



US008257707B2

(12) **United States Patent**  
**Smith et al.**(10) **Patent No.:** **US 8,257,707 B2**  
(45) **Date of Patent:** **Sep. 4, 2012**(54) **IL-18 RECEPTOR ANTIGEN BINDING  
PROTEINS**(75) Inventors: **Dirk E. Smith**, Bainbridge Island, WA  
(US); **John E. Sims**, Seattle, WA (US);  
**Jeffrey T. McGrew**, Woodinville, WA  
(US); **Marek Z. Kubin**, Bainbridge  
Island, WA (US); **Duncan Cochrane**,  
Granta Park (GA); **Louise Conroy**,  
Granta Park (GA)(73) Assignee: **Amgen Inc.**, Thousand Oaks, CA (US)(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.(21) Appl. No.: **12/670,112**(22) PCT Filed: **Jul. 24, 2008**  
(Under 37 CFR 1.47)(86) PCT No.: **PCT/US2008/071047**  
§ 371 (c)(1),  
(2), (4) Date: **Aug. 16, 2010**(87) PCT Pub. No.: **WO2009/015284**  
PCT Pub. Date: **Jan. 29, 2009**(65) **Prior Publication Data**

US 2011/0014201 A1 Jan. 20, 2011

**Related U.S. Application Data**(60) Provisional application No. 60/951,691, filed on Jul.  
24, 2007, provisional application No. 60/951,692,  
filed on Jul. 24, 2007, provisional application No.  
61/073,142, filed on Jun. 17, 2008.(51) **Int. Cl.**  
**A61K 39/395** (2006.01)(52) **U.S. Cl.** ..... **424/133.1**; 424/141.1; 424/143.1;  
530/388.1; 530/350; 536/23.1; 435/69.1;  
435/235.1; 435/325; 435/6.1(58) **Field of Classification Search** ..... None  
See application file for complete search history.(56) **References Cited**

## U.S. PATENT DOCUMENTS

5,776,731 A 7/1998 Parnet  
6,090,918 A 7/2000 Parnet  
6,451,760 B1 9/2002 Parnet  
6,492,496 B1 \* 12/2002 Parnet et al. .... 530/388.22  
6,589,764 B1 7/2003 Sims  
6,600,022 B1 7/2003 Torigoe et al.  
6,664,077 B1 12/2003 Sims  
6,692,740 B2 2/2004 Sims  
6,693,171 B2 2/2004 Sims  
7,001,599 B2 2/2006 Parnet  
7,169,581 B2 1/2007 Sims  
7,235,238 B2 6/2007 Parnet  
7,270,817 B2 9/2007 Sims  
7,625,710 B2 12/2009 Parnet  
7,704,945 B2 4/2010 Sims7,838,248 B2 11/2010 Sims  
7,993,845 B2 8/2011 Sims  
8,034,343 B2 10/2011 Parnet  
2002/0098185 A1 7/2002 Sims et al.  
2003/0059937 A1 3/2003 Ruben et al.  
2004/0018198 A1 1/2004 Gudas et al.  
2004/0023336 A1 2/2004 Heavner et al.  
2005/0147610 A1 7/2005 Ghayur et al.  
2007/0025992 A1 2/2007 Takayama et al.

## FOREIGN PATENT DOCUMENTS

EP 1808446 A1 7/2007  
WO 97/31010 A1 8/1997  
WO 99/37772 A1 7/1999  
WO 99/37773 A1 7/1999  
WO 01/007480 A2 2/2001  
WO 01/27279 A1 4/2001  
WO 02/092008 A2 11/2002  
WO 03/057821 A2 7/2003

## OTHER PUBLICATIONS

Nishida et al., "Cloning and expression of a single-chain Fv fragment  
specific for the human interleukin 18 receptor," Hybridoma  
17(6):577-581, 1998.Vermot-Desroches et al., "Monoclonal antibodies specific for the  
IL-18 receptor," Cell Immunol, 236:101-104, 2005.Caldas et al., Humanization of the anti-CD18 antibody 6.7: an unex-  
pected effect of a framework residue in binding to antigen, Mol  
Immunol 39(15):941-952, 2003.Casset et al., "A peptide mimetic of an anti-CD4 monoclonal anti-  
body by rational design," Biochem Biophys Res Commun  
307(1):198-205, 2003.Chien et al., "Significant structural and functional change of an  
antigen-binding site by a distant amino acid substitution: proposal of  
a structural mechanism," Proc Natl Acad Sci USA 86(14):5532-  
5536, 1989.De Pascalis et al., "Grafting of 'abbreviated' complementarity-deter-  
mining regions containing specificity-determining residues essential  
for ligand contact to engineer a less immunogenic humanized  
monoclonal antibody," J Immunol 169(6):3076-3084, 2002.Guisti et al., "Somatic diversification of S107 from an  
antiphosphocholine to an anti-DNA autoantibody is due to a single  
base change in its heavy chain variable region," Proc Natl Acad Sci  
USA 84(9):2926-2930, 1987.Gussow and Seemann, "Humanization of monoclonal antibodies,"  
Methods Enzymol 203:99-121, 1991.Holm et al., "Functional mapping and single chain construction of the  
anti-cytokeratin 8 monoclonal antibody TS1," Mol Immunol  
44(6):1075-1084, 2007.MacCallum et al., "Antibody-antigen interactions: contact analysis  
and binding site topography," J Mol Biol 262(5):732-745, 1996.

(Continued)

*Primary Examiner* — Dong Jiang(74) *Attorney, Agent, or Firm* — Thomas J. Wrona(57) **ABSTRACT**Provided herein are IL-18 receptor antigen binding proteins  
and polynucleotides encoding the same. Expression vectors  
and host cells comprising the same for production of the  
antigen binding proteins are also provided. In addition, pro-  
vided are compositions and methods for diagnosing and treat-  
ing diseases mediated by IL-18 receptor.**21 Claims, 34 Drawing Sheets**

OTHER PUBLICATIONS

Mariuzza et al., "The structural basis of antigen-antibody recognition," *Annu Rev Biophys Biophys Chem* 16:139-159, 1987.

Rudikoff et al., "Single amino acid substitution altering antigen-binding specificity," *Proc Natl Acad Sci USA* 79(6):1979-1983, 1982.

Wu et al., "Humanization of a murine monoclonal antibody by simultaneous optimization of framework and CDR residues," *J Mol Biol* 294(1):151-162, 1999.

R&D Systems online catalog, "Human IL-18 R $\alpha$ /IL-1 R5 Antibody," Jan. 1, 2007, retrieved from the Internet: URL:<http://www.rndsystems.com/pdf/mab840.pdf> [retrieved on Feb. 3, 2012].

Tokmadzic, V.S. et al., "IL-18 is present at the maternal-fetal interface and enhances cytotoxic activity of decidual lymphocytes," *AJRI* 48:191-200 (2002).

Wu, C. et al., "IL-18 receptor  $\beta$ -induced changes in the presentation of IL-18 binding sites affect ligand binding and signal transduction," *J Immunol* 170:5571-5577 (2003).

Imaoka H. et al., "Interleukin-18 production and pulmonary function in COPD," *Eur Respir J* 31:287-297 (2008).

\* cited by examiner

**AM<sub>H</sub>1 polynucleotide sequence (SEQ ID:35)**

caggtgcagctggtgcagctctgggggagggcgtggtccagcctgggaggtccctgagactc	60
Q V Q L V Q S G G G V V Q P G R S L R L	
tcctgtgcagcgtctggattcaccttcagcggttatggcatgcactgggtccgccaggct	120
S C A A S G F T F S G Y G M H W V R Q A	
ccaggcaaggggctggagtggtggcagtaatatcaaatgatggaagtaagaaatattat	180
P G K G L E W V A V I S N D G S K K Y Y	
tcagactccgtgaagggccgattcacctctccagagacaattccaaaaacacgctgtat	240
S D S V K G R F T I S R D N S K N T L Y	
ctgcagatgaacagcctgagagctgaggacacggctgtatattactgtgcgaaagggctcc	300
L Q M N S L R A E D T A V Y Y C A K G S	
agttccatatggctgaccagtcctggaccactgggggcaggggaccacggtcaccgctc	360
S S I W L T Q S L D H W G Q G T T V T V	
tcctca	366
S S	

**AM<sub>H</sub>1 amino acid sequence (SEQ ID:1)**

QVQLVQSGGGVVQPGRSLRLSCAASGFTFSGYGMHWVRQAPGKGLEWVAVISNDGSKKYY	60
SDSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAKSSSIWLTQSLDHWGQGTTVTV	120
SS	122

**AM<sub>H</sub>2 polynucleotide sequence (SEQ ID:36)**

gaggtgcagctggtggagctctgggggagggcgtggtccagcctgggaggtccctgagactc	60
E V Q L V E S G G G V V Q P G R S L R L	
tcctgtgcagcgtctggattcaccttcagcggttatggcatgcactgggtccgccaggct	120
S C A A S G F T F S G Y G M H W V R Q A	
ccaggcaaggggctggagtggtggcagtaatatcaaatgatggaagtaagaaatattat	180
P G K G L E W V A V I S N D G S K K Y Y	
tcagactccgtgaagggccgattcacctctccagagacaattccaaaaacacgctgtat	240
S D S V K G R F T I S R D N S K N T L Y	
ctgcagatgaacagcctgagagctgaggacacggctgtatattactgtgcgaaagggctcc	300
L Q M N S L R A E D T A V Y Y C A K G S	
agttccatatggctgaccagtcctggaccactgggggcaggggaccacggtcaccgctc	360
S S I W L T Q S L D H W G Q G T T V T V	
tcctca	366
S S	

**AM<sub>H</sub>2 amino acid sequence (SEQ ID NO:2)**

EVQLVESGGGVVQPGRSLRLSCAASGFTFSGYGMHWVRQAPGKGLEWVAVISNDGSKKYY	60
SDSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCAKSSSIWLTQSLDHWGQGTTVTV	120
SS	122

FIGURE 1A

**AM<sub>H</sub>3 polynucleotide sequence (SEQ ID 37)**

cagggtgcagctggtgcagtcctgggggagggcgtggtccagcctgggaggtccctgagactc	60
Q V Q L V Q S G G G V V Q P G R S L R L	
tcctgtgcagcgtctggattcaccttcagcggttatggcatgcaactgggtccgccaggct	120
S C A A S G F T F S G Y G M H W V R Q A	
ccaggcaaggggctggagtggtggcagtaatatcaaatgatggaagtaagaaatattat	180
P G K G L E W V A V I S N D G S K K Y Y	
tcagactccgtgaagggccgattcaccatctccagagacaattccaaaaacacgctgtat	240
S D S V K G R F T I S R D N S K N T L Y	
ctgcagatgaacagcctgagagctgaggacacggctgtatattactgtgcgaaaggggtcc	300
L Q M N S L R A E D T A V Y Y C A K G S	
agttccatatggctgtcgcagtcctggacggctgggggcaggggaccacggtcaccgtc	360
S S I W L S Q S L D G W G Q G T T V T V	
tcctca	366
S S	

**AM<sub>H</sub>3 amino acid sequence (SEQ ID NO:3)**

QVQLVQSGGGVVQPGRSLRLS CAASGFTFSGYGMHWVRQAPGKGLEWVAVISNDGSKKYY	60
SDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK GSSSIWLSQSLD GWGQGT TVTV	120
SS	122

**AM<sub>H</sub>4 polynucleotide sequence (SEQ ID:38)**

gagggtgcagctggtggagtcctgggggagggcgtggtccagcctgggaggtccctgagactc	60
E V Q L V E S G G G V V Q P G R S L R L	
tcctgtgcagcgtctggattcaccttcagcggttatggcatgcaactgggtccgccaggct	120
S C A A S G F T F S G Y G M H W V R Q A	
ccaggcaaggggctggagtggtggcagtaatatcaaatgatggaagtaagaaatattat	180
P G K G L E W V A V I S N D G S K K Y Y	
tcagactccgtgaagggccgattcaccatctccagagacaattccaaaaacacgctgtat	240
S D S V K G R F T I S R D N S K N T L Y	
ctgcagatgaacagcctgagagctgaggacacggctgtatattactgtgcgaaaggggtcc	300
L Q M N S L R A E D T A V Y Y C A K G S	
agttccatatggctgtcgcagtcctggacggctgggggcaggggaccacggtcaccgtc	360
S S I W L S Q S L D G W G Q G T T V T V	
tcctca	366
S S	

**AM<sub>H</sub>4 amino acid sequence (SEQ ID:4)**

EVQLVESGGGVVQPGRSLRLS CAASGFTFSGYGMHWVRQAPGKGLEWVAVISNDGSKKYY	60
SDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK GSSSIWLSQSLD GWGQGT TVTV	120
SS	122

FIGURE 1B

**AM<sub>H</sub>5 polynucleotide sequence (SEQ ID:39)**

caggtgcagctggtgcagctctgggggagggcgtggtccagcctgggaggtccctgagactc	60
Q V Q L V Q S G G G V V Q P G R S L R L	
tcctgtgcagcgtctggattcaccttcagcggttatggcatgcactgggtccgccaggct	120
S C A A S G F T F S G Y G M H W V R Q A	
ccaggcaaggggctggagtggtggcagtaatatcaaatgatggaagtaagaaatattat	180
P G K G L E W V A V I S N D G S K K Y Y	
tcagactccgtgaagggccgattcaccatctccagagacaattccaaaaacacgctgtat	240
S D S V K G R F T I S R D N S K N T L Y	
ctgcagatgaacagcctgagagctgaggacacggctgtatattactgtgcgaaaggggcc	300
L Q M N S L R A E D T A V Y Y C A K G S	
agttccatattggctgacctcggccctgaacctgtgggggaggggaccacggtcaccgctc	360
S S I W L T S A L N L W G Q G T T V T V	
tcctca	366
S S	

**AM<sub>H</sub>5 amino acid sequence (SEQ ID:5)**

QVQLVQSGGGVVQPGRSLRLS CAASGFTFSGYGMHWVRQAPGKGLEWVAVISNDGSKKYY	60
SDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAKGSSSIWLTSALNLWGQGT TVTV	120
SS	122

**AM<sub>H</sub>6 polynucleotide sequence (SEQ ID:40)**

gaggtgcagctggtggagctctgggggagggcgtggtccagcctgggaggtccctgagactc	60
E V Q L V E S G G G V V Q P G R S L R L	
tcctgtgcagcgtctggattcaccttcagcggttatggcatgcactgggtccgccaggct	120
S C A A S G F T F S G Y G M H W V R Q A	
ccaggcaaggggctggagtggtggcagtaatatcaaatgatggaagtaagaaatattat	180
P G K G L E W V A V I S N D G S K K Y Y	
tcagactccgtgaagggccgattcaccatctccagagacaattccaaaaacacgctgtat	240
S D S V K G R F T I S R D N S K N T L Y	
ctgcagatgaacagcctgagagctgaggacacggctgtatattactgtgcgaaaggggcc	300
L Q M N S L R A E D T A V Y Y C A K G S	
agttccatattggctgacctcggccctgaacctgtgggggaggggaccacggtcaccgctc	360
S S I W L T S A L N L W G Q G T T V T V	
tcctca	366
S S	

**AM<sub>H</sub>6 polynucleotide sequence (SEQ ID:6)**

EVQLVESGGGVVQPGRSLRLS CAASGFTFSGYGMHWVRQAPGKGLEWVAVISNDGSKKYY	60
SDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAKGSSSIWLTSALNLWGQGT TVTV	120
SS	122

FIGURE 1C

**AM<sub>H</sub>7 polynucleotide sequence (SEQ ID:41)**

gaggtgcagctggtggagtctgggggagggcgtggtccagcctgggaggtccctgagactc	60
E V Q L V E S G G G V V Q P G R S L R L	
tcctgtgcagcgtctggattcaccttcagcggttatggcatgcactgggtccgccaggct	120
S C A A S G F T F S G Y G M H W V R Q A	
ccaggcaaggggctggagtgggtggcagtaatatcaaagtatggaagtaagaaatattat	180
P G K G L E W V A V I S N D G S K K Y Y	
tcagactccgtgaagggccgattcaccatctccagagacaattccaaaaacacgctgtat	240
S D S V K G R F T I S R D N S K N T L Y	
ctgcagatgaacagcctgagagctgaggacacggctatatattactgtgcgaaagggctcc	300
L Q M N S L R A E D T A I Y Y C A K G S	
agttccatattggttcggggagaccgttgactactgggggcaggggaccacg	351
S S I W F G E T V D Y W G Q G T T	

**AM<sub>H</sub>7 amino acid sequence (SEQ ID:7)**

EVQLVESGGGVVQPGRSLRLSCAASGFTFSGYGMHWVRQAPGKGLEWVAVISNDGSKKYY	60
SDSVKGRFTISRDNKNTLYLQMNSLRAEDTAIYYCAKGSIIWFGETVDYWGQGT	117

**AM<sub>H</sub>8 polynucleotide sequence (SEQ ID:42)**

caggtgcagctggtgcagctctggggctgaggtgaagaagcctggggcctcagtgaaggtc	60
Q V Q L V Q S G A E V K K P G A S V K V	
tcctgcaaggtttccggatacacctcactgaattatccatgcactgggtgacagagct	120
S C K V S G Y T L T E L S M H W V R Q A	
cctggaaaagggcttgagtggatgggaggttttgatcgtgaagatgatgaaacaatccac	180
P G K G L E W M G G F D R E D D E T I H	
gcacagaagttccagggcagagtcaccatgaccgaggacacatctacagacacagcctac	240
A Q K F Q G R V T M T E D T S T D T A Y	
atggaactgagcagcctgcgatctgaggacacggccgtttattactgtgcaacagatcct	300
M E L S S L R S E D T A V Y Y C A T D L	
atgggtgtggggcgatttttgatccagcactggggccaggggacactgggtcaccgtctcc	360
M V W G D F W I Q H W G Q G T L V T V S	
tca	363
S	

**AM<sub>H</sub>8 polynucleotide sequence (SEQ ID:8)**

QVQLVQSGAEVKKPGASVKVSKVSGYTLTELSMHWVRQAPGKGLEWMGGFDREDETIH	60
AQKFQGRVTMTEDTSTDTAYMELSSLRSEDTAVYYCATDLMVWGDVFIQHWGQGLVTVS	120
S	121

FIGURE 1D

**AM<sub>H</sub>9 polynucleotide sequence (SEQ ID:43)**

caggtgcagctggtgcagtcctggggctgaggtgaagaagcctggggcctcagtgaaggtc	60
Q V Q L V Q S G A E V K K P G A S V K V	
tcctgcaagggttccggatacacccctcactgaattatccatgcactgggtgacagggct	120
S C K V S G Y T L T E L S M H W V R Q A	
cctggaaaagggttgagtggatgggaggttttgatcgtgaagatgatgaaacaatccac	180
P G K G L E W M G G F D R E D D E T I H	
gcacagaagttccagggcagagtcacatgaccgaggacacatctacagacacagcctac	240
A Q K F Q G R V T M T E D T S T D T A Y	
atggaactgagcagcctgcatctgaggacacggccgtttattactgtgcaacagatctt	300
M E L S S L R S E D T A V Y Y C A T D L	
atgggtgtggggcgatttttggatccagcactggggccaggggacactggtcaccgtctcc	360
M V W G D F W I Q H W G Q G T L V T V S	
tca	363
S	

**AM<sub>H</sub>9 amino acid sequence (SEQ ID:9)**

QVQLVQSGAEVKKPGASVKVSCKVSGYTLTELSMHWVRQAPGKGLEWMGGFDREDETIH	60
AQKFQGRVTMTEDTSTDTAYMELSSLRSEDTAVYYCATDLMVWGDFWIQHWGQGLVTVS	120
S	121

**AM<sub>H</sub>10 polynucleotide sequence (SEQ ID:44)**

caggtgcagctggtgcagtcctggggctgaggtgaagaagcctggggcctcagtgaaggtc	60
Q V Q L V Q S G A E V K K P G A S V K V	
tcctgcaagggttccggatacacccctcactgaattatccatgcactgggtgacagggct	120
S C K V S G Y T L T E L S M H W V R Q A	
cctggaaaagggttgagtggatgggaggttttgatcgtgaagatgatgaaacaatccac	180
P G K G L E W M G G F D R E D D E T I H	
gcacagaagttccagggcagagtcacatgaccgaggacacatctacagacacagcctac	240
A Q K F Q G R V T M T E D T S T D T A Y	
atggaactgagcagcctgcatctgaggacacggccgtttattactgtgcaacagatctt	300
M E L S S L R S E D T A V Y Y C A T D L	
atggcctgggactaccgcccacccagcactggggccaggggacactggtcaccgtctcc	360
M A W D Y P P I Q H W G Q G T L V T V S	
tca	363
S	

**AM<sub>H</sub>10 amino acid sequence (SEQ ID:10)**

QVQLVQSGAEVKKPGASVKVSCKVSGYTLTELSMHWVRQAPGKGLEWMGGFDREDETIH	60
AQKFQGRVTMTEDTSTDTAYMELSSLRSEDTAVYYCATDLMWDYPPIQHWGQGLVTVS	120
S	121

FIGURE 1E

**AM<sub>H</sub>11 polynucleotide sequence (SEQ ID:45)**

caggtgcagctggtgcagtcctggggctgaggtgaagaagcctggggcctcagtgaaggtc	60
Q V Q L V Q S G A E V K K P G A S V K V	
tcctgcaagggttccggatacaccctcactgaattatccatgcactgggtgacagggct	120
S C K V S G Y T L T E L S M H W V R Q A	
cctggaaaagggttgagtggatgggaggttttgatcgtgaagatgatgaaacaatccac	180
P G K G L E W M G G F D R E D D E T I H	
gcacagaagttccagggcagagtcacccatgaccgaggacacatctacagacacagcctac	240
A Q K F Q G R V T M T E D T S T D T A Y	
atggaactgagcagcctgcatctgaggacacggccggtttattactgtgcaacagatcct	300
M E L S S L R S E D T A V Y Y C A T D L	
atgggtgtggaacttccccccatccagcactggggccaggggacactggtcaccgctctcc	360
M V W N F P P I Q H W G Q G T L V T V S	
tca	363
S	

**AM<sub>H</sub>11 amino acid sequence (SEQ ID:11)**

QVQLVQSGAEVKKPGASVKVSKVSGYTLTELSMHWVRQAPGKGLEWMGGFDREDDETIH	60
AQKFQGRVTMTEDTSTDTAYMELSSLRSEDTAVYYCATDLMVWNFPPIQHWGQGLVTVS	120
S	121

**AM<sub>J</sub>12 polynucleotide sequence (SEQ ID:46)**

caggtgcagctggtgcagtcctggggctgaggtgaagaagcctggggcctcagtgaaggtc	60
Q V Q L V Q S G A E V K K P G A S V K V	
tcctgcaagggttccggatacaccctcactgaattatccatgcactgggtgacagggct	120
S C K V S G Y T L T E L S M H W V R Q A	
cctggaaaagggttgagtggatgggaggttttgatcgtgaagatgatgaaacaatccac	180
P G K G L E W M G G F D R E D D E T I H	
gcacagaagttccagggcagagtcacccatgaccgaggacacatctacagacacagcctac	240
A Q K F Q G R V T M T E D T S T D T A Y	
atggaactgagcagcctgcatctgaggacacggccggtttattactgtgcaacagatcct	300
M E L S S L R S E D T A V Y Y C A T D L	
atgggtgtggggcgatttttgatccagcactggggcaaggggacaatg	348
M V W G D F W I Q H W G K G T M	

**AM<sub>H</sub>12 amino acid sequence (SEQ ID:12)**

QVQLVQSGAEVKKPGASVKVSKVSGYTLTELSMHWVRQAPGKGLEWMGGFDREDDETIH	60
AQKFQGRVTMTEDTSTDTAYMELSSLRSEDTAVYYCATDLMVWGDVFIQHWGKGTM	111

FIGURE 1F



**AM<sub>H</sub>13 polynucleotide sequence (SEQ ID:47)**

gaggtgcagctggtggagtctgggggaggcttggtacagcctgggggggtccctgagactc	60
E V Q L L E S G G G L V Q P G G S L R L	
tcctgtgcagcctctggattcacctttagcagctatgccatgagctgggtccgccaggct	120
S C A A S G F T F S S Y A M S W V R Q A	
ccaggaaggggctggagtgggtctcagctattagtggtagtggtgggtggcacatactac	180
P G K G L E W V S A I S G S G G G T Y Y	
gcagactccgtgaagggccggtccaccatctccagagacaattccaagaacacgctgtat	240
A D S V K G R F T I S R D N S K N T L Y	
ctgcaaatgaacagcctgagagccgaggacacggccgtgtattactgtgogagaattcgg	300
L Q M N S L R A E D T A V Y Y C A R I R	
ggcgactaccggacggacatctggggccaggggaaccacggtcaccgtctcctca	354
G D Y R T D I W G Q G T T V T V S S	

**AM<sub>H</sub>13 amino acid sequence (SEQ ID:13)**

EVQLLESGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSAISGSGGGTYY	60
ADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARIRGDYRTDIWGQTTVTVSS	118

**AM<sub>H</sub>14 polynucleotide sequence (SEQ ID:48)**

gaggtgcagctggtggagtctgggggaggcttggtacagcctgggggggtccctgagactc	60
E V Q L L E S G G G L V Q P G G S L R L	
tcctgtgcagcctctggattcacctttagcagctatgccatgagctgggtccgccaggct	120
S C A A S G F T F S S Y A M S W V R Q A	
ccaggaaggggctggagtgggtctcagctattagtggtagtggtgggtggcacatactac	180
P G K G L E W V S A I S G S G G G T Y Y	
gcagactccgtgaagggccggtccaccatctccagagacaattccaagaacacgctgtat	240
A D S V K G R S T I S R D N S K N T L Y	
ctgcaaatgaacagcctgagagccgaggacacggccgtgtattactgtgogagaattcgg	300
L Q M N S L R A E D T A V Y Y C A R I R	
ggggactaccggacggacatctggggccgggggaaccctgggtcaccgtctcctca	354
G D Y R T D I W G R G T L V T V S S	

**AM<sub>H</sub>14 amino acid sequence (SEQ ID:14)**

EVQLLESGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSAISGSGGGTYY	60
ADSVKGRSTISRDN SKNTLYLQMNSLRAEDTAVYYCARIRGDYRTDIWGRGTLVTVSS	118

**FIGURE 1G**

**AM<sub>H</sub>15 polynucleotide sequence (SEQ ID:49)**

gagggtgcagctggtggagctctgggggaggcttggcacagcctgggggggtccctgagactc	60
E V Q L L E S G G G L A Q P G G S L R L	
tcctgtgcagcctctgggttcacctttagcagctatgccatgagctgggtccgccaggct	120
S C A A S G F T F S S Y A M S W V R Q A	
ccagggaaggggctggagtggtctcagctattagtggttagtggttagcacatactac	180
P G K G L E W V S A I S G S G G S T Y Y	
gcagactccgtgaaggccgggttcaccatctccagagacaattccaagaacacgctgtat	240
A D S V K G R F T I S R D N S K N T L Y	
ctgcaaatgaacagcctgagagccgaggacacggccgtgtattactgtgagagagttcgg	300
L Q M N S L R A E D T A V Y Y C A R V R	
ggggactaccggacggacatctggggccggggaaccctgggtcaccgtctcctca	354
G D Y R T D I W G R G T L V T V S S	

**AM<sub>H</sub>15 amino acid sequence (SEQ ID:15)**

EVQLLESGGGLAQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEWVSAISGSGGSTYY	60
ADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARVRGDIWGRGTLVTVSS	118

**AM<sub>H</sub>16 polynucleotide sequence (SEQ ID:50)**

gagggtgcagctggtggagctctgggggaggcttggtagcagcctgggggggtccctgagactc	60
E V Q L L E S G G G L V Q P G G S L R L	
tcctgtgcagcctctagattcacctttagcagctatgccatgagctgggtccgccaggct	120
S C A A S R F T F S S Y A M S W V R Q A	
ccagggaaggggctggagtggtctcagctattagtggttagtggttagcacatactac	180
P G K G L E W V S A I S G S G G S T Y Y	
gcagactccgtgaaggccgggttcaccatctccagagacaattccaagaacacgctgtat	240
A D S V K G R F T I S R D N S K N T L Y	
ctgcaaatgaacagcctgagagccgaggacacggccgtgtattactgtgagagagttcgg	300
L Q M N S L R A E D T A V Y Y C A R V R	
ggggactaccggacggacatctggggccggggaaccctgggtcaccgtctcctca	354
G D Y R T D I W G R G T L V T V S S	

**AM<sub>H</sub>16 amino acid sequence (SEQ ID:16)**

EVQLLESGGGLVQPGGSLRLSCAASRFTFSSYAMSWVRQAPGKGLEWVSAISGSGGSTYY	60
ADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARVRGDIWGRGTLVTVSS	118

FIGURE 1H

**AM<sub>H</sub>17 polynucleotide sequence (SEQ ID:51)**

gaggtgcagctggtggagtctgggggaggcttgggtacagcctgggggggccctgagactc	60
E V Q L L E S G G G L V Q P G G S L R L	
tcctgtgcagcctctagattcacctttagcagctatgccatgagctgggtccgccaggct	120
S C A A S R F T F S S Y A M S W V R Q A	
ccaggaaggggctggagtgggtctcagctattagtggtagtggtggtagcacatactac	180
P G K G L E W V S A I S G S G G S T Y Y	
gcagactccgtgaagggccggttcaccatctccagagacaattccaagaacacgctgtat	240
A D S V K G R F T I S R D N S K N T L Y	
ctgcaaataaacagcctgagagccgaggacacggccgtgtattactgtgcgagagttcgg	300
L Q M N S L R A E D T A V Y Y C A R V R	
ggcatatacgggtatggacgtctggggccggggaacctg	339
G I Y G M D V W G R G T L	

**AM<sub>H</sub>17 amino acid sequence (SEQ ID:17)**

EVQLLESGLVQPGGSLRLSCAASRFTFSSYAMSWVRQAPGKGLEWVSAISGSGGSTYY	60
ADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARVRGIYGMVWGRGTL	113

**AM<sub>L</sub>1 polynucleotide sequence (SEQ ID:52)**

cagcctgtgctgactcagccccctcagtgtccgtgtccccaggacagactgccagcatc	60
Q P V L T Q P P S V S V S P G Q T A S I	
acctgctctggagataaattgggggataaatatgcttctctgggtatcagcagaagccaggc	120
T C S G D K L G D K Y A S W Y Q Q K P G	
aagtcccctgtactgggtcatctatcaagattccaatcggccctcagggatccctgagcga	180
K S P V L V I Y Q D S N R P S G I P E R	
ttctctggctccaactctgggaacacagccactctgaccatcagcgggacccaggctagg	240
F S G S N S G N T A T L T I S G T Q A R	
gatgaggctgactattactgtcaggcgtgggacagcagcactgcacgggtgttcggcgga	300
D E A D Y Y C Q A W D S S T A S V F G G	
gggaccaag	309
G T K	

**AM<sub>L</sub>1 amino acid sequence (SEQ ID:18)**

QPVLTPPSVSVSPGQTASITCSGDKLGDKYASWYQQKPGKSPVLVIYQDSNRPSGIPER	60
FSGNSGNTATLTIISGTQARDEADYYCQAWDSSTASVFGGGTK	103

FIGURE 11

**AM<sub>1</sub>2 polynucleotide sequence (SEQ ID:53)**

cagcctgtgctgactcagccccctcagtgtccgtgtccccaggacagactgccagcatc	60
Q P V L T Q P P S V S V S P G Q T A S I	
acctgctctggagataaattgggggataaatatgcttctctggatcagcagaagccaggc	120
T C S G D K L G D K Y A S W Y Q Q K P G	
aagtccccctgtactggatcatctatcaagattccaatcggccctcagggatccctgagcga	180
K S P V L V I Y Q D S N R P S G I P E R	
ttctctggctccaactctgggaacacagccactctgaccatcagcgggacccaggctagg	240
F S G S N S G N T A T L T I S G T Q A R	
gatgaggctgactattactgtcaggcgtggaccactccttgcagcacagggtcggcgga	300
D E A D Y Y C Q A W D H S L Q H R F G G	
gggaccaaggtcaccgtcctaggt	324
G T K V T V L G	

**AM<sub>1</sub>2 amino acid sequence (SEQ ID:19)**

QPVLTPPSVSVSPGQTASITCSGDKLGDKYASWYQQKPGKSPVLVIYQDSNRPSGIPER	60
FSGSNSGNTATLTI SGTQARDEADYYCQAWDHS LQHRFGGGTKVTVLG	108

**AM<sub>1</sub>3 polynucleotide sequence (SEQ ID:54)**

cagcctgtgctgactcagccccctcagtgtccgtgtccccaggacagactgccagcatc	60
Q P V L T Q P P S V S V S P G Q T A S I	
acctgctctggagataaattgggggataaatatgcttctctggatcagcagaagccaggc	120
T C S G D K L G D K Y A S W Y Q Q K P G	
cagaccctgtactggatcatctatcaagattccaatcggccctcagggatccctgagcga	180
Q T P V L V I Y Q D S N R P S G I P E R	
ttctctggctccaactccgggaacacagccactctgaccatcagcgggacccaggctagg	240
F S G S N S G N T A T L T I S G T Q A R	
gatgaggctgactattactgtcaggcgtggaccagcgcctgaactcgcaggtcggcgga	300
D E A D Y Y C Q A W T S A L N S Q F G G	
gggaccaaggtcaccgtcctaggt	324
G T K V T V L G	

**AM<sub>1</sub>3 amino acid sequence (SEQ ID:20)**

QPVLTPPSVSVSPGQTASITCSGDKLGDKYASWYQQKPGQTPVLVIYQDSNRPSGIPER	60
FSGSNSGNTATLTI SGTQARDEADYYCQAWTSALNSQFGGGTKVTVLG	108

FIGURE 1J

**AM<sub>1</sub>4 polynucleotide sequence (SEQ ID:55)**

cagcctgtgctgactcagccccctcagtgtccgtgtccccaggacagactgccagcatc	60
Q P V L T Q P P S V S V S P G Q T A S I	
acctgctctggagataaattgggggataaatatgcttcctgggtatcagcagaagccaggc	120
T C S G D K L G D K Y A S W Y Q Q K P G	
cagtccccctgtactgggtcatctatcaagattccaatcggccctcagggatccctgagcga	180
Q S P V L V I Y Q D S N R P S G I P E R	
ttctctggctccaactctggggacacagccactctgaccatcagcgggacccaggctagg	240
F S G S N S G D T A T L T I S G T Q A R	
gatgaggctgactattactgtcaggcgtggacgcactccctcagcacgttggttcggcgga	300
D E A D Y Y C Q A W T H S L S T L F G G	
gggaccaaggtcaccgtcctaggt	324
G T K V T V L G	

**AM<sub>1</sub>4 amino acid sequence (SEQ ID:21)**

QPVLTPPSVSVSPGQTASITCSGDKLGDKYASWYQQKPGQSPVLVIYQDSNRPSGIPER	60
FSGSNSGDTATLTI SGTQARDEADYYCQAWTHSLSTLFGGGTKVTVLG	108

**AM<sub>1</sub>5 polynucleotide sequence (SEQ ID:56)**

tcctatgagctgactcagccccctcagtgtccgtgtccccaggacagactgccagcatc	60
S Y E L T Q P P S V S V S P G Q T A S I	
acctgctctggagataaattgggggataaatatgcttcctgggtatcagcagaagccaggc	120
T C S G D K L G D K Y A S W Y Q Q K P G	
cagtccccctgtactgggtcatctatcaagattccaatcggccctcagggatccctgagcga	180
Q S P V L V I Y Q D S N R P S G I P E R	
ttctctggctccaactctgggaacacagccactctgaccatcagcgggacccaggctatg	240
F S G S N S G N T A T L T I S G T Q A M	
gatgaggctgactattactgtcaggcgtggacccacagcctgagcacgttggttcggcgga	300
D E A D Y Y C Q A W T H S L S T L F G G	
gggaccaagctgaccgtcctaggt	324
G T K L T V L G	

**AM<sub>1</sub>5 amino acid sequence (SEQ ID:22)**

SYELTPPSVSVSPGQTASITCSGDKLGDKYASWYQQKPGQSPVLVIYQDSNRPSGIPER	60
FSGSNSGNTATLTI SGTQAMDEADYYCQAWTHSLSTLFGGGTKLTVLG	108

FIGURE 1K

**AM<sub>1</sub>6 polynucleotide sequence (SEQ ID:57)**

cagtctgtgctgactcagccaccctcagcgtctgggacccccgggcagaggggtcaccatc	60
Q S V L T Q P P S A S G T P G Q R V T I	
tcttggtctggaaggaactccaacatcggaagttatactgtaacctggtaccagcagctc	120
S C S G R N S N I G S Y T V T W Y Q Q L	
ccaggaacggcccccaactcctcatctatagtaatagtcagcggccctcaggggtccct	180
P G T A P K L L I Y S N S Q R P S G V P	
gaccgattctcaggtccaagtctggcacctcagcctccttggccatcagtgggctccag	240
D R F S G S K S G T S A S L A I S G L Q	
tctgaagatgaggctgattattactgtgcagcatgggatgacagcctgaatggccccggtg	300
S E D E A D Y Y C A A W D D S L N G P V	
ttcggcggaggaccacag	318
F G G G T K	

**AM<sub>1</sub>6 amino acid sequence (SEQ ID:23)**

QSVLTQPPSASGTPGQRVTISCSGRNSNIGSYTVTWTYQQLPGTAPKLLIYSNSQRPSGVF	60
DRFSGSKSGTSASLAISGLQSEDEADYYCAAWDDSLNGPVFVGGGTK	106

**AM<sub>1</sub>7 polynucleotide sequence (SEQ ID:58)**

cagtctgtgctgactcagccaccctcagcgtctgggacccccgggcagaggggtcaccatc	60
Q S V L T Q P P S A S G T P G Q R V T I	
tcttggtctggaaggaactccaacatcggaagttatactgtaacctggtaccagcagctc	120
S C S G R N S N I G S Y T V T W Y Q Q L	
ccaggaacggcccccaactcctcatctatagtaatagtcagcggccctcaggggtccct	180
P G T A P K L L I Y S N S Q R P S G V P	
gaccgattctcaggtccaagtctggcacctcagcctccttggccatcagtgggctccag	240
D R F S G S K S G T S A S L A I S G L Q	
tctgaagatgaggctgattattactgtgtggtgtgggatgacgtgctgaatggccccggtg	300
S E D E A D Y Y C V V W D D V L N G P V	
ttcggcggaggaccacagctgaccgtcctaggt	333
F G G G T K L T V L G	

**AM<sub>1</sub>7 amino acid sequence (SEQ ID:24)**

QSVLTQPPSASGTPGQRVTISCSGRNSNIGSYTVTWTYQQLPGTAPKLLIYSNSQRPSGVF	60
DRFSGSKSGTSASLAISGLQSEDEADYYCVVWDDVNLNGPVFVGGGTKLTVLG	111

**FIGURE 1L**

**AM<sub>1</sub>8 polynucleotide sequence (SEQ ID:59)**

cagtcctgtgctgactcagccaccctcagcgtctgggacccccgggcagagggtcaccatc	60
Q S V L T Q P P S A S G T P G Q R V T I	
tcttgttctggaaggaactccaacatcggaagttatactgtaacctggtaccagcagctc	120
S C S G R N S N I G S Y T V T W Y Q Q L	
ccaggaacggcccccaactcctcatctatagtaatagtcagcggccctcaggggtccct	180
P G T A P K L L I Y S N S Q R P S G V P	
gaccgattctcaggtccaagtctggcacctcagcctccttggccatcagtgggctccag	240
D R F S G S K S G T S A S L A I S G L Q	
tctgaagatgaggctgattattactgtgtcgtgtgggatgacaagctgaatggccccggtg	300
S E D E A D Y Y C V V W D D K L N G P V	
ttcggcggaggaccgaagctgaccgtcctaggt	333
F G G G T K L T V L G	

**AM<sub>1</sub>8 polynucleotide sequence (SEQ ID:25)**

QSVLTQPPSASGTPGQRVTISCSGRNSNIGSYTVTWTYQQLPGTAPKLLIYSNSQRPSGVP	60
DRFSGSKSGTSASLAISGLQSEDEADYYCVVWDDKLNGLPVFGGGTKLTVLG	111

**AM<sub>1</sub>9 polynucleotide sequence (SEQ ID:60)**

cagtcctgtgctgactcagccaccctcagcgtctgggacccccgggcagagggtcaccatc	60
Q S V L T Q P P S A S G T P G Q R V T I	
tcttgttctggaaggaactccaacatcggaagttatactgtaacctggtaccagcagctc	120
S C S G R N S N I G S Y T V T W Y Q Q L	
ccaggaacggcccccaactcctcatctatagtaatagtcagcggccctcaggggtccct	180
P G T A P K L L I Y S N S Q R P S G V P	
gaccgattctcaggtccaagtctggcacctcagcctccttggccatcagtgggctccag	240
D R F S G S K S G T S A S L A I S G L Q	
tctgaagatgaggctgattattactgtgtgggtgtgggacgagatcctgaatggccccggtg	300
S E D E A D Y Y C V V W D E I L N G P V	
ttcggcggaggaccgaagctgaccgtcctaggt	333
F G G G T K L T V L G	

**AM<sub>1</sub>9 amino acid sequence (SEQ ID:26)**

QSVLTQPPSASGTPGQRVTISCSGRNSNIGSYTVTWTYQQLPGTAPKLLIYSNSQRPSGVP	60
DRFSGSKSGTSASLAISGLQSEDEADYYCVVWDEILNGLPVFGGGTKLTVLG	111

FIGURE 1M

**AM<sub>1</sub>10 polynucleotide sequence (SEQ ID:61)**

tcgtctgagctgactcaggaccctgctgtgtctgtggccttgggacagacagtcaggatc	60
S S E L T Q D P A V S V A L G Q T V R I	
acatgccaaaggagacagcctcagaagctattatgcaagctggtaccagcagaagccagga	120
T C Q G D S L R S Y Y A S W Y Q Q K P G	
caggccccctgtacttgtcatctctgctaaaaacaaccggccctcagggatcccagaccga	180
Q A P V L V I S A K N N R P S G I P D R	
ttctctggctccagctcaggaaacacagcttccttgaccatcactggggctcaggcgga	240
F S G S S S G N T A S L T I T G A Q A E	
gatgaagctgactattactgtaactcccgggacagcagtaacctgtggtattcggcgga	300
D E A D Y Y C N S R D S S N H V V F G G	
gggaccaag	309
G T K	

**AM<sub>1</sub>10 amino acid sequence (SEQ ID:27)**

SSELTQDPAVSVVALGQTVRITCQGD SLRSYYASWYQQKPGQAPV LVISAKNNRPSGIPDR	60
FSGSSGNTASLTITGAQAEDEADYYCNSRDSSNHVVFVGGGTK	103

**AM<sub>1</sub>11 polynucleotide sequence (SEQ ID:62)**

cagtctgtgctgactcagccaccctcagcgtctgggacccccgggcagagggtcaccatc	60
Q S V L T Q P P S A S G T P G Q R V T I	
tcttgttctggaaggaactccaacatcggaagttatactgtaacctggtaccagcagctc	120
S C S G R N S N I G S Y T V T W Y Q Q L	
ccaggaacggcccccaactcctcatctatagtaatagtcagcggccctcaggggtccct	180
P G T A P K L L I Y S N S Q R P S G V P	
gaccgattctcaggctccaagtctggcacctcagcctccttggccatcagtgggctccag	240
D R F S G S K S G T S A S L A I S G L Q	
tctgaagatgaggctgattattactgtctcgtgtgggacgacgtcctgaatggccccggtg	300
S E D E A D Y Y C L V W D D V L N G P V	
ttcggcggagggaccaagctgaccgtcctaggt	333
F G G G T K L T V L G	

**AM<sub>1</sub>11 amino acid sequence (SEQ ID:28)**

QSVLTQPPSASGTPGQRV TISCSGRNSNIGSYTVT WYQQLPGTAPKLLIYSNSQRPSGVP	60
DRFSGSKSGTSASLAISGLQSEDEADYYCLVWDDV L N G P V F G G G T K L T V L G	111

FIGURE 1N



**AM<sub>L</sub>12 polynucleotide sequence (SEQ ID:63)**

tcgtctgagctgactcaggaccctgctgtgtctgtggccttgggacagacagtcaggatc	60
S S E L T Q D P A V S V A L G Q T V R I	
acatgccaaaggagacagcctcagaagctattatgcaagctggtaccagcagaagccagga	120
T C Q G D S L R S Y Y A S W Y Q Q K P G	
caggccctgtacttgtcatctctgctaaaaacaaccggccctcagggatcccagaccga	180
Q A P V L V I S A K N N R P S G I P D R	
ttctctggctccagctcaggaaacacagcttccttgaccatcactggggctcaggcggaa	240
F S G S S S G N T A S L T I T G A Q A E	
gatgaggctgactattactgtgctcccggaaacggctggaaccatgtggtattcggcgga	300
D E A D Y Y C A S R N G W N H V V F G G	
gggaccaagctgaccgtcctaggt	324
G T K L T V L G	

**AM<sub>L</sub>12 amino acid sequence (SEQ ID:29)**

SSELTQDPAVSVALGQTVRITCQGDSLRSYYASWYQQKPGQAPVLVISAANNRPSGIPDR	60
FSGSSSGNTASLTITGAQAEADYYCASRNGWNHVVFGGGTKLTVLG	108

**AM<sub>L</sub>13 polynucleotide sequence (SEQ ID:64)**

tcgtctgagctgactcaggaccctgctgtgtctgtggccttgggacagacagtcagggtc	60
S S E L T Q D P A V S V A L G Q T V R V	
acatgccaaaggagacagcctcagaagctattatgcaagctggtaccagcagaagccagga	120
T C Q G D S L R S Y Y A S W Y Q Q K P G	
caggccctgtacttgtcatctctgctaaaaacaaccggccctcagggatcccagaccga	180
Q A P V L V I S A K N N R P S G I P D R	
ttctctggctccagctcaggaaacacagcttccttgaccatcactggggctcaggcggaa	240
F S G S S S G N T A S L T I T G A Q A E	
gatgaggctgactattactgtgctcccggaaacggctggaaccatgtggtattcggcgga	300
D E A D Y Y C A S R N G W N H V V F G G	
gggaccaagctgaccgtcctaggt	324
G T K L T V L G	

**AM<sub>L</sub>13 amino acid sequence (SEQ ID:30)**

SSELTQDPAVSVALGQTVRVTCQGDLSRSYYASWYQQKPGQAPVLVISAANNRPSGIPDR	60
FSGSSSGNTASLTITGAQAEADYYCASRNGWNHVVFGGGTKLTVLG	108

FIGURE 10

**AM<sub>1</sub>14 polynucleotide sequence (SEQ ID:65)**

tcgtctgagctgactcaggaccctgctgtgtctgtggccttgggacagacagtcaggatc	60
S S E L T Q D P A V S V A L G Q T V R -I	
acatgccaaaggagacagcctcagaagctattatgcaagctggtaccagcagaagccagga	120
T C Q G D S L R S Y Y A S W Y Q Q K P G	
caggcccctgtacttgtcatctctgctaaaaacaaccggccctcagggatcccagaccga	180
Q A P V L V I S A K N N R P S G I P D R	
ttctctggctccagctcaggaaacacagcttccttgaccatcactggggctcaggcggaa	240
F S G S S S G N T A S L T I T G A Q A E	
gatgaggctgactattactgtgctcccggacggctggaaccatgtggtattcggcgga	300
D E A D Y Y C A S R N G W N H V V F G G	
gggaccaagctgaccgtcctaggt	324
G T K L T V L G	

**AM<sub>1</sub>14 amino acid sequence (SEQ ID:31)**

SSELTQDPAVSVVALGQTVRITCQGDSLRSYYASWYQQKPGQAPVLVISAKNNRPSGIPDR	60
FSGSSGNTASLTITGAQAEDEADYYCASRNGWNHVVFGGGTKLTVLG	108

**AM<sub>1</sub>15 polynucleotide sequence (SEQ ID:66)**

tcgtctgagctgactcaggaccctgctgtgtctgtggccttgggacagacagtcagggtc	60
S S E L T Q D P A V S V A L G Q T V R V	
acatgccaaaggagacagcctcagaagctattatgcaagctggtaccagcagaagccagga	120
T C Q G D S L R S Y Y A S W Y Q Q K P G	
caggcccctgtacttgtcatctctgctaaaaacaaccggccctcagggatcccagaccga	180
Q A P V L V I S A K N N R P S G I P D R	
ttctctggctccagctcaggaaacacagcttccttgaccatcactggggctcaggcggaa	240
F S G S S S G N T A S L T I T G A Q A E	
gatgaggctgactattactgtgctcccggacggctggaaccatgtggtattcggcgga	300
D E A D Y Y C A S R N G W N H V V F G G	
gggaccaagctgaccgtcctaggt	324
G T K L T V L G	

**AM<sub>1</sub>15 amino acid sequence (SEQ ID:32)**

SSELTQDPAVSVVALGQTVRVITCQGDSLRSYYASWYQQKPGQAPVLVISAKNNRPSGIPDR	60
FSGSSGNTASLTITGAQAEDEADYYCASRNGWNHVVFGGGTKLTVLG	120

FIGURE 1P

**AM<sub>1</sub>16 polynucleotide sequence (SEQ ID:67)**

tcgtctgagctgactcaggaccctgctgtgtctgtggccttgggacagacagtcaggatc	60
S S E L T Q D P A V S V A L G Q T V R I	
acatgccaaaggagacagcctcagaagctattatgcaagctggtaccagcagaagccagga	120
T C Q G D S L R S Y Y A S W Y Q Q K P G	
caggccccctgtacttgtcatctctgctaaaaacaaccggccctcagggatcccagaccga	180
Q A P V L V I S A K N N R P S G I P D R	
ttctctggctccagctcaggaaacacagcttccttgaccatcactggggctcaggcggaa	240
F S G S S S G N T A S L T I T G A Q A E	
gatgaggctgactattactgtgcgacccggaacggctggaaccatgtggtattcggcgga	300
D E A D Y Y C A T R N G W N H V V F G G	
gggaccaagctgaccgtcctaggt	324
G T K L T V L G	

**AM<sub>1</sub>16 amino acid sequence (SEQ ID:33)**

SSELTQDPAVSVALGQTVRITCQGDLSRSYYASWYQQKPGQAPV LVI SAKNNRPSGI PDR	60	
FSGSSGNTASLTITGAQAEDEADYYCATRNGWNHVVFGGGTKLTVLG		108

**AM<sub>1</sub>17 polynucleotide sequence (SEQ ID:68)**

tcgtctgagctgactcaggaccctgctgtgtctgtggccttgggacagacagtcagggtc	60
S S E L T Q D P A V S V A L G Q T V R V	
acatgccaaaggagacagcctcagaagctattatgcaagctggtaccagcagaagccagga	120
T C Q G D S L R S Y Y A S W Y Q Q K P G	
caggccccctgtacttgtcatctctgctaaaaacaaccggccctcagggatcccagaccga	180
Q A P V L V I S A K N N R P S G I P D R	
ttctctggctccagctcaggaaacacagcttccttgaccatcactggggctcaggcggaa	240
F S G S S S G N T A S L T I T G A Q A E	
gatgaggctgactattactgtgcgacccggaacggctggaaccatgtggtattcggcgga	300
D E A D Y Y C A T R N G W N H V V F G G	
gggaccaagctgaccgtcctaggt	324
G T K L T V L G	

**AM<sub>1</sub>17 amino acid sequence (SEQ ID:34)**

SSELTQDPAVSVALGQTVRVTCQGDLSRSYYASWYQQKPGQAPV LVI SAKNNRPSGI PDR	60	
FSGSSGNTASLTITGAQAEDEADYYCATRNGWNHVVFGGGTKLTVLG		108

**FIGURE 1Q**

AM<sub>H</sub>1 amino acid sequence (SEQ ID:1)

QVQLVQSGGGVVPGRSLRLSCAASGFTFSGYGMHWVRQAPGKGLEWVAVISNDGSKKYYSDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAKGSSEIWLIOSLDHWGQGTITVTVSS  
CDR1 CDR2 CDR3

AM<sub>H</sub>2 amino acid sequence (SEQ ID NO:2)

EVQLVESGGGVVPGRSLRLSCAASGFTFSGYGMHWVRQAPGKGLEWVAVISNDGSKKYYSDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAKGSSEIWLIOSLDHWGQGTITVTVSS  
CDR1 CDR2 CDR3

AM<sub>H</sub>3 amino acid sequence (SEQ ID NO:3)

QVQLVQSGGGVVPGRSLRLSCAASGFTFSGYGMHWVRQAPGKGLEWVAVISNDGSKKYYSDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAKGSSEIWLIOSLDHWGQGTITVTVSS  
CDR1 CDR2 CDR3

AM<sub>H</sub>4 amino acid sequence (SEQ ID:4)

EVQLVESGGGVVPGRSLRLSCAASGFTFSGYGMHWVRQAPGKGLEWVAVISNDGSKKYYSDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAKGSSEIWLIOSLDHWGQGTITVTVSS  
CDR1 CDR2 CDR3

AM<sub>H</sub>5 amino acid sequence (SEQ ID:5)

QVQLVQSGGGVVPGRSLRLSCAASGFTFSGYGMHWVRQAPGKGLEWVAVISNDGSKKYYSDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAKGSSEIWLIOSLDHWGQGTITVTVSS  
CDR1 CDR2 CDR3

AM<sub>H</sub>6 polynucleotide sequence (SEQ ID:6)

EVQLVESGGGVVPGRSLRLSCAASGFTFSGYGMHWVRQAPGKGLEWVAVISNDGSKKYYSDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAKGSSEIWLIOSLDHWGQGTITVTVSS  
CDR1 CDR2 CDR3

AM<sub>H</sub>7 amino acid sequence (SEQ ID:7)

EVQLVESGGGVVPGRSLRLSCAASGFTFSGYGMHWVRQAPGKGLEWVAVISNDGSKKYYSDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAKGSSEIWLIOSLDHWGQGTITVTVSS  
CDR1 CDR2 CDR3

FIGURE 2A

**AM<sub>H</sub>8 polynucleotide sequence (SEQ ID:8)**

QVQLVQSGAEVKKPGASVKVSCKVSGYTLT<sup>CDR1</sup>ELSMHWVRQAPGKGLEW<sup>CDR2</sup>MGGFDREDELIHAOKFOGRVTMTEDTSDTAYMELSSLRSEDTAVYYCATDLMVWGDFMIQHMGQGLVTVSS<sup>CDR3</sup>

**AM<sub>H</sub>9 amino acid sequence (SEQ ID:9)**

QVQLVQSGAEVKKPGASVKVSCKVSGYTLT<sup>CDR1</sup>ELSMHWVRQAPGKGLEW<sup>CDR2</sup>MGGFDREDELIHAOKFOGRVTMTEDTSDTAYMELSSLRSEDTAVYYCATDLMVWGDFMIQHMGQGLVTVSS<sup>CDR3</sup>

**AM<sub>H</sub>10 amino acid sequence (SEQ ID:10)**

QVQLVQSGAEVKKPGASVKVSCKVSGYTLT<sup>CDR1</sup>ELSMHWVRQAPGKGLEW<sup>CDR2</sup>MGGFDREDELIHAOKFOGRVTMTEDTSDTAYMELSSLRSEDTAVYYCATDLMANDYFPFIQHMGQGLVTVSS<sup>CDR3</sup>

**AM<sub>H</sub>11 amino acid sequence (SEQ ID:11)**

QVQLVQSGAEVKKPGASVKVSCKVSGYTLT<sup>CDR1</sup>ELSMHWVRQAPGKGLEW<sup>CDR2</sup>MGGFDREDELIHAOKFOGRVTMTEDTSDTAYMELSSLRSEDTAVYYCATDLMVMNPEPFIQHMGQGLVTVSS<sup>CDR3</sup>

**AM<sub>H</sub>12 amino acid sequence (SEQ ID:12)**

QVQLVQSGAEVKKPGASVKVSCKVSGYTLT<sup>CDR1</sup>ELSMHWVRQAPGKGLEW<sup>CDR2</sup>MGGFDREDELIHAOKFOGRVTMTEDTSDTAYMELSSLRSEDTAVYYCATDLMVWGDFMIQHMGKGT<sup>CDR3</sup>

**AM<sub>H</sub>13 amino acid sequence (SEQ ID:13)**

EVQLLESGGGLVQPGGSLRLSCAASGFTT<sup>CDR1</sup>FSSYAMSNVRQAPGKGLEWVSAISGSGGGTYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARIRGDIYRIDINGQGITVTVSS<sup>CDR2</sup><sup>CDR3</sup>

**AM<sub>H</sub>14 amino acid sequence (SEQ ID:14)**

EVQLLESGGGLVQPGGSLRLSCAASGFTT<sup>CDR1</sup>FSSYAMSNVRQAPGKGLEWVSAISGSGGGTYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARIEGDIYRIDINGRGTLLVTVSS<sup>CDR2</sup><sup>CDR3</sup>

FIGURE 2B

AM<sub>H</sub>15 amino acid sequence (SEQ ID:15)

EVQLLESGGGLAQPGGSLRLSCAAAGFTFSYAMSNVRQAPKGLLEWVSAISGSGSIIYADSVKGRFTISRDN SKNTLYLQMN SLRAEDTAVYYCARVGGDYRTDIWGRTLVTVSS  
CDR1 CDR2 CDR3

AM<sub>H</sub>16 amino acid sequence (SEQ ID:16)

EVQLLESGGGLVQPGGSLRLSCAAARFTFSYAMSNVRQAPFGKLEWVSAISGSGSIIYADSVKGRFTISRDN SKNTLYLQMN SLRAEDTAVYYCARVGGDYRTDIWGRTLVTVSS  
CDR1 CDR2 CDR3

AM<sub>H</sub>17 amino acid sequence (SEQ ID:17)

EVQLLESGGGLVQPGGSLRLSCAAARFTFSYAMSNVRQAPFGKLEWVSAISGSGSIIYADSVKGRFTISRDN SKNTLYLQMN SLRAEDTAVYYCARVGGIYGMVWNGRGTLL  
CDR1 CDR2 CDR3

AM<sub>L</sub>1 amino acid sequence (SEQ ID:18)

QPVLTPPPSVSPGQTASITCSGDKLGDKYASWYQQKPGKSPVLVIYQDSNRPSGI PERFGSNGNTATLTI SGTQARDEADYYQAMDSIIASVFGGGTK  
CDR1 CDR2 CDR3

AM<sub>L</sub>2 amino acid sequence (SEQ ID:19)

QPVLTPPPSVSPGQTASITCSGDKLGDKYASWYQQKPGKSPVLVIYQDSNRPSGI PERFGSNGNTATLTI SGTQARDEADYYQAMDHSLQHRFGGGTKVTVLG  
CDR1 CDR2 CDR3

AM<sub>L</sub>3 amino acid sequence (SEQ ID:20)

QPVLTPPPSVSPGQTASITCSGDKLGDKYASWYQQKPGQT FVLVIYQDSNRPSGI PERFGSNGNTATLTI SGTQARDEADYYQAMTSALNSQFGGGTKVTVLG  
CDR1 CDR2 CDR3

AM<sub>L</sub>4 amino acid sequence (SEQ ID:21)

QPVLTPPPSVSPGQTASITCSGDKLGDKYASWYQQKPGQS FVLVIYQDSNRPSGI PERFGSNGSDTATLTI SGTQARDEADYYQAWTHSLSLIFGGGTKVTVLG  
CDR1 CDR2 CDR3

FIGURE 2C

**AM<sub>L</sub>5 amino acid sequence (SEQ ID:22)**

SYELTQPPSVSVSPGQTASI TCSGDKLGDKYASWYQQKPGQSFVLVI YQDSNRPSGI PERFSGNSGNTATLTI SGTQANDEADYYCQANTHSLSLSTLFGGGTKLTVLG  
CDR1 CDR2 CDR3

**AM<sub>L</sub>6 amino acid sequence (SEQ ID:23)**

QSVLTQPPSASGTPGQRVTI SC~~SG~~RNSNIGSYITVWYQQLPGTAPKLLIYNSORPSGVDPDRFSGSKSGTSASLAI SGLQSEDEADYYCAAWDDSLNGPVPVFGGGTK  
CDR1 CDR2 CDR3

**AM<sub>L</sub>7 amino acid sequence (SEQ ID:24)**

QSVLTQPPSASGTPGQRVTI SC~~SG~~RNSNIGSYITVWYQQLPGTAPKLLIYNSORPSGVDPDRFSGSKSGTSASLAI SGLQSEDEADYYCVVWDDVLLNGPVPVFGGGTKLTVLG  
CDR1 CDR2 CDR3

**AM<sub>L</sub>8 polynucleotide sequence (SEQ ID:25)**

QSVLTQPPSASGTPGQRVTI SC~~SG~~RNSNIGSYITVWYQQLPGTAPKLLIYNSORPSGVDPDRFSGSKSGTSASLAI SGLQSEDEADYYCVVWDDVLLNGPVPVFGGGTKLTVLG  
CDR1 CDR2 CDR3

**AM<sub>L</sub>9 amino acid sequence (SEQ ID:26)**

QSVLTQPPSASGTPGQRVTI SC~~SG~~RNSNIGSYITVWYQQLPGTAPKLLIYNSORPSGVDPDRFSGSKSGTSASLAI SGLQSEDEADYYCVVWDEILLNGPVPVFGGGTKLTVLG  
CDR1 CDR2 CDR3

**AM<sub>L</sub>10 amino acid sequence (SEQ ID:27)**

SSELTDQPAVSVLQQTVRTITCGESLRSYASWYQQKPGQAFVLVISA~~KN~~RPSGITPDRFSGSSGNTASLTI TGAQAEDEADYYCNSRDS~~SN~~HVYVFGGGTK  
CDR1 CDR2 CDR3

**AM<sub>L</sub>11 amino acid sequence (SEQ ID:28)**

QSVLTQPPSASGTPGQRVTI SC~~SG~~RNSNIGSYITVWYQQLPGTAPKLLIYNSORPSGVDPDRFSGSKSGTSASLAI SGLQSEDEADYYC~~V~~WDDVLLNGPVPVFGGGTKLTVLG  
CDR1 CDR2 CDR3

FIGURE 2D

AM<sub>1</sub>,12 amino acid sequence (SEQ ID:29)

SSELTQDPAVSVALGQTVRITCOGDSLRSYYASWYQKFGQAFVLVISA<sup>CDR1</sup>AKNNRPSGI<sup>CDR2</sup>PDRFSGSSSGNTASLITGAQAEDY<sup>CDR3</sup>CASRNGMNHVVF<sup>CDR3</sup>GGGTKLTVLG

AM<sub>1</sub>,13 amino acid sequence (SEQ ID:30)

SSELTQDPAVSVALGQTVRVTCOGDSLRSYYASWYQKFGQAFVLVISA<sup>CDR1</sup>AKNNRPSGI<sup>CDR2</sup>PDRFSGSSSGNTASLITGAQAEDY<sup>CDR3</sup>CASRNGMNHVVF<sup>CDR3</sup>GGGTKLTVLG

AM<sub>1</sub>,14 amino acid sequence (SEQ ID:31)

SSELTQDPAVSVALGQTVRITCOGDSLRSYYASWYQKFGQAFVLVISA<sup>CDR1</sup>AKNNRPSGI<sup>CDR2</sup>PDRFSGSSSGNTASLITGAQAEDY<sup>CDR3</sup>CASRNGMNHVVF<sup>CDR3</sup>GGGTKLTVLG

AM<sub>1</sub>,15 amino acid sequence (SEQ ID:32)

SSELTQDPAVSVALGQTVRVTCOGDSLRSYYASWYQKFGQAFVLVISA<sup>CDR1</sup>AKNNRPSGI<sup>CDR2</sup>PDRFSGSSSGNTASLITGAQAEDY<sup>CDR3</sup>CASRNGMNHVVF<sup>CDR3</sup>GGGTKLTVLG

AM<sub>1</sub>,16 amino acid sequence (SEQ ID:33)

SSELTQDPAVSVALGQTVRITCOGDSLRSYYASWYQKFGQAFVLVISA<sup>CDR1</sup>AKNNRPSGI<sup>CDR2</sup>PDRFSGSSSGNTASLITGAQAEDY<sup>CDR3</sup>CATRNGMNHVVF<sup>CDR3</sup>GGGTKLTVLG

AM<sub>1</sub>,17 amino acid sequence (SEQ ID:34)

SSELTQDPAVSVALGQTVRVTCOGDSLRSYYASWYQKFGQAFVLVISA<sup>CDR1</sup>AKNNRPSGI<sup>CDR2</sup>PDRFSGSSSGNTASLITGAQAEDY<sup>CDR3</sup>CATRNGMNHVVF<sup>CDR3</sup>GGGTKLTVLG

FIGURE 2E



Heavy Chain

1 70

AM <sub>H</sub> 1	SEQ ID NO: 1	QVQLVQSGGG VVQPGRSRLR SCAASGFTFS GYGMHWVRQA PGKGLEWVAV ISNDGSKKYY SDSVKGRFTI	CDR1	CDR2
AM <sub>H</sub> 2	SEQ ID NO: 2	EVQLVESGGG VVQPGRSRLR SCAASGFTFS GYGMHWVRQA PGKGLEWVAV ISNDGSKKYY SDSVKGRFTI		
AM <sub>H</sub> 3	SEQ ID NO: 3	QVQLVQSGGG VVQPGRSRLR SCAASGFTFS GYGMHWVRQA PGKGLEWVAV ISNDGSKKYY SDSVKGRFTI		
AM <sub>H</sub> 4	SEQ ID NO: 4	EVQLVESGGG VVQPGRSRLR SCAASGFTFS GYGMHWVRQA PGKGLEWVAV ISNDGSKKYY SDSVKGRFTI		
AM <sub>H</sub> 5	SEQ ID NO: 5	QVQLVQSGGG VVQPGRSRLR SCAASGFTFS GYGMHWVRQA PGKGLEWVAV ISNDGSKKYY SDSVKGRFTI		
AM <sub>H</sub> 6	SEQ ID NO: 6	EVQLVESGGG VVQPGRSRLR SCAASGFTFS GYGMHWVRQA PGKGLEWVAV ISNDGSKKYY SDSVKGRFTI		
AM <sub>H</sub> 15	SEQ ID NO: 15	EVQLLESVGG LAQPGGSLRL SCAASGFTFS SYAMSWVRQA PGKGLEWVSA ISGSGGSTYY ADSVKGRFTI		
AM <sub>H</sub> 16	SEQ ID NO: 16	EVQLLESVGG LVQPGGSLRL SCAASRFTFS SYAMSWVRQA PGKGLEWVSA ISGSGGSTYY ADSVKGRFTI		
AM <sub>H</sub> 13	SEQ ID NO: 13	EVQLLESVGG LVQPGGSLRL SCAASGFTFS SYAMSWVRQA PGKGLEWVSA ISGSGGSTYY ADSVKGRFTI		
AM <sub>H</sub> 14	SEQ ID NO: 14	EVQLLESVGG LVQPGGSLRL SCAASGFTFS SYAMSWVRQA PGKGLEWVSA ISGSGGSTYY ADSVKGRFTI		
AM <sub>H</sub> 8	SEQ ID NO: 8	QVQLVQSGAE VVKPGASVKV SKVSGYTLT ELSMHWVRQA PKGLEWVSA ISGSGGSTYY AOKFQGRVTM		
AM <sub>H</sub> 9	SEQ ID NO: 9	QVQLVQSGAE VVKPGASVKV SKVSGYTLT ELSMHWVRQA PKGLEWVSA ISGSGGSTYY AOKFQGRVTM		
AM <sub>H</sub> 10	SEQ ID NO: 10	QVQLVQSGAE VVKPGASVKV SKVSGYTLT ELSMHWVRQA PKGLEWVSA ISGSGGSTYY AOKFQGRVTM		
AM <sub>H</sub> 11	SEQ ID NO: 11	QVQLVQSGAE VVKPGASVKV SKVSGYTLT ELSMHWVRQA PKGLEWVSA ISGSGGSTYY AOKFQGRVTM		
AM <sub>H</sub> 7	SEQ ID NO: 7	EVQLVESGGG VVQPGRSRLR SCAASGFTFS GYGMHWVRQA PGKGLEWVAV ISNDGSKKYY SDSVKGRFTI		
AM <sub>H</sub> 12	SEQ ID NO: 12	QVQLVQSGAE VVKPGASVKV SKVSGYTLT ELSMHWVRQA PKGLEWVSA ISGSGGSTYY AOKFQGRVTM		
AM <sub>H</sub> 17	SEQ ID NO: 17	EVQLLESVGG LVQPGGSLRL SCAASRFTFS SYAMSWVRQA PGKGLEWVSA ISGSGGSTYY ADSVKGRFTI		

71 122

AM <sub>H</sub> 1	SEQ ID NO: 1	SRDNSKNTLY LQMNSLRAED TAVYICAKGS SSIWLTQSLE HWGQGITVTV SS.....	CDR3
AM <sub>H</sub> 2	SEQ ID NO: 2	SRDNSKNTLY LQMNSLRAED TAVYICAKGS SSIWLTQSLE HWGQGITVTV SS.....	
AM <sub>H</sub> 3	SEQ ID NO: 3	SRDNSKNTLY LQMNSLRAED TAVYICAKGS SSIWLTQSLE HWGQGITVTV SS.....	
AM <sub>H</sub> 4	SEQ ID NO: 4	SRDNSKNTLY LQMNSLRAED TAVYICAKGS SSIWLTQSLE HWGQGITVTV SS.....	
AM <sub>H</sub> 5	SEQ ID NO: 5	SRDNSKNTLY LQMNSLRAED TAVYICAKGS SSIWLTQSLE HWGQGITVTV SS.....	
AM <sub>H</sub> 6	SEQ ID NO: 6	SRDNSKNTLY LQMNSLRAED TAVYICAKGS SSIWLTQSLE HWGQGITVTV SS.....	
AM <sub>H</sub> 15	SEQ ID NO: 15	SRDNSKNTLY LQMNSLRAED TAVYICARV GDY...RTD IWGRGTLVTV SS.....	
AM <sub>H</sub> 16	SEQ ID NO: 16	SRDNSKNTLY LQMNSLRAED TAVYICARV GDY...RTD IWGRGTLVTV SS.....	
AM <sub>H</sub> 13	SEQ ID NO: 13	SRDNSKNTLY LQMNSLRAED TAVYICARV GDY...RTD IWGRGTLVTV SS.....	
AM <sub>H</sub> 14	SEQ ID NO: 14	SRDNSKNTLY LQMNSLRAED TAVYICARV GDY...RTD IWGRGTLVTV SS.....	
AM <sub>H</sub> 8	SEQ ID NO: 8	TEDTSTDYAY MELSLRSED TAVYICATDL M.VWGDFWVQ HWGQGITVTV SS.....	
AM <sub>H</sub> 9	SEQ ID NO: 9	TEDTSTDYAY MELSLRSED TAVYICATDL M.VWGDFWVQ HWGQGITVTV SS.....	
AM <sub>H</sub> 10	SEQ ID NO: 10	TEDTSTDYAY MELSLRSED TAVYICATDL M.VWGDFWVQ HWGQGITVTV SS.....	
AM <sub>H</sub> 11	SEQ ID NO: 11	TEDTSTDYAY MELSLRSED TAVYICATDL M.VWGDFWVQ HWGQGITVTV SS.....	
AM <sub>H</sub> 7	SEQ ID NO: 7	SRDNSKNTLY LQMNSLRAED TAVYICAKGS SSIWLTQSLE HWGQGITVTV SS.....	
AM <sub>H</sub> 12	SEQ ID NO: 12	TEDTSTDYAY MELSLRSED TAVYICATDL M.VWGDFWVQ HWGQGITVTV SS.....	
AM <sub>H</sub> 17	SEQ ID NO: 17	SRDNSKNTLY LQMNSLRAED TAVYICARV GDY...RTD IWGRGTLVTV SS.....	

FIGURE 3A

Light Chain

1 70

AM <sub>L</sub> 17	SEQ ID NO:34	SSELTQDP	AV	SVALGQTV	RV	TCQG..	DSL	RYASWYQ	QK	PGQAPV	LVIS	AKNRP	SGIP	DRFSG	SSGN
AM <sub>L</sub> 16	SEQ ID NO:33	SSELTQDP	AV	SVALGQTV	RV	TCQG.,	DSL	RYASWYQ	QK	PGQAPV	LVIS	AKNRP	SGIP	DRFSG	SSGN
AM <sub>L</sub> 15	SEQ ID NO:32	SSELTQDP	AV	SVALGQTV	RV	TCQG..	DSL	RYASWYQ	QK	PGQAPV	LVIS	AKNRP	SGIP	DRFSG	SSGN
AM <sub>L</sub> 13	SEQ ID NO:30	SSELTQDP	AV	SVALGQTV	RV	TCQG..	DSL	RYASWYQ	QK	PGQAPV	LVIS	AKNRP	SGIP	DRFSG	SSGN
AM <sub>L</sub> 14	SEQ ID NO:31	SSELTQDP	AV	SVALGQTV	RV	TCQG..	DSL	RYASWYQ	QK	PGQAPV	LVIS	AKNRP	SGIP	DRFSG	SSGN
AM <sub>L</sub> 12	SEQ ID NO:29	SSELTQDP	AV	SVALGQTV	RV	TCQG..	DSL	RYASWYQ	QK	PGQAPV	LVIS	AKNRP	SGIP	DRFSG	SSGN
AM <sub>L</sub> 3	SEQ ID NO:20	QPVLTP	PPSV	SVSPGQ	TASI	TCSG..	DKL	RYASWYQ	QK	PGQSPV	LVIY	QDSNR	PSGIP	ERFSG	NSGN
AM <sub>L</sub> 4	SEQ ID NO:21	QPVLTP	PPSV	SVSPGQ	TASI	TCSG..	DKL	RYASWYQ	QK	PGQSPV	LVIY	QDSNR	PSGIP	ERFSG	NSGN
AM <sub>L</sub> 2	SEQ ID NO:19	QPVLTP	PPSV	SVSPGQ	TASI	TCSG..	DKL	RYASWYQ	QK	PGKSPV	LVIY	QDSNR	PSGIP	ERFSG	NSGN
AM <sub>L</sub> 5	SEQ ID NO:22	SYELTP	PPSA	SGTPGQ	RVTI	TCSG..	DKL	RYASWYQ	QK	PGQSPV	LVIY	QDSNR	PSGIP	ERFSG	NSGN
AM <sub>L</sub> 11	SEQ ID NO:26	QSVLTQ	PPSA	SGTPGQ	RVTI	TCSG..	DKL	RYASWYQ	QK	PGTAPK	LVIY	SNSQR	PSGVP	DRFSG	SKSGT
AM <sub>L</sub> 7	SEQ ID NO:24	QSVLTQ	PPSA	SGTPGQ	RVTI	TCSG..	DKL	RYASWYQ	QK	PGTAPK	LVIY	SNSQR	PSGVP	DRFSG	SKSGT
AM <sub>L</sub> 9	SEQ ID NO:26	QSVLTQ	PPSA	SGTPGQ	RVTI	TCSG..	DKL	RYASWYQ	QK	PGTAPK	LVIY	SNSQR	PSGVP	DRFSG	SKSGT
AM <sub>L</sub> 8	SEQ ID NO:25	QSVLTQ	PPSA	SGTPGQ	RVTI	TCSG..	DKL	RYASWYQ	QK	PGTAPK	LVIY	SNSQR	PSGVP	DRFSG	SKSGT
AM <sub>L</sub> 10	SEQ ID NO:27	SSELTQ	DP	AV	SVALGQ	TV	RV	TCQG..	DSL	RYASWYQ	QK	PGQAPV	LVIS	AKNRP	SGIP
AM <sub>L</sub> 1	SEQ ID NO:18	QPVLTP	PPSV	SVSPGQ	TASI	TCSG..	DKL	RYASWYQ	QK	PGKSPV	LVIY	QDSNR	PSGIP	ERFSG	NSGN
AM <sub>L</sub> 6	SEQ ID NO:23	QSVLTQ	PPSA	SGTPGQ	RVTI	TCSG..	DKL	RYASWYQ	QK	PGTAPK	LVIY	SNSQR	PSGVP	DRFSG	SKSGT

51 129

AM <sub>L</sub> 17	SEQ ID NO:34	TASLTIT	GAQ	AEDEADY	YCA	TRNGWNH	.VV	FGGGTKL	TVL	G.....	.....	.....	.....	.....	.....
AM <sub>L</sub> 16	SEQ ID NO:33	TASLTIT	GAQ	AEDEADY	YCA	TRNGWNH	.VV	FGGGTKL	TVL	G.....	.....	.....	.....	.....	.....
AM <sub>L</sub> 15	SEQ ID NO:32	TASLTIT	GAQ	AEDEADY	YCA	SRNGWNH	.VV	FGGGTKL	TVL	G.....	.....	.....	.....	.....	.....
AM <sub>L</sub> 13	SEQ ID NO:30	TASLTIT	GAQ	AEDEADY	YCA	SRNGWNH	.VV	FGGGTKL	TVL	G.....	.....	.....	.....	.....	.....
AM <sub>L</sub> 14	SEQ ID NO:31	TASLTIT	GAQ	AEDEADY	YCA	SRNGWNH	.VV	FGGGTKL	TVL	G.....	.....	.....	.....	.....	.....
AM <sub>L</sub> 12	SEQ ID NO:29	TASLTIT	GAQ	AEDEADY	YCA	SRNGWNH	.VV	FGGGTKL	TVL	G.....	.....	.....	.....	.....	.....
AM <sub>L</sub> 3	SEQ ID NO:20	TATLTIS	GTQ	ARDEADY	YCA	AMTSA	LN.SQ	FGGGTKV	TVL	G.....	.....	.....	.....	.....	.....
AM <sub>L</sub> 4	SEQ ID NO:21	TATLTIS	GTQ	ARDEADY	YCA	AWTHSL	.TL	FGGGTKV	TVL	G.....	.....	.....	.....	.....	.....
AM <sub>L</sub> 2	SEQ ID NO:19	TATLTIS	GTQ	ARDEADY	YCA	AWTHSL	.TL	FGGGTKV	TVL	G.....	.....	.....	.....	.....	.....
AM <sub>L</sub> 5	SEQ ID NO:22	TATLTIS	GTQ	AMDEADY	YCA	AWTHSL	.TL	FGGGTKL	TVL	G.....	.....	.....	.....	.....	.....
AM <sub>L</sub> 11	SEQ ID NO:28	SASLAIS	GLQ	SEDEADY	YCA	VMDPVL	NGPV	FGGGTKL	TVL	G.....	.....	.....	.....	.....	.....
AM <sub>L</sub> 7	SEQ ID NO:24	SASLAIS	GLQ	SEDEADY	YCA	VMDPVL	NGPV	FGGGTKL	TVL	G.....	.....	.....	.....	.....	.....
AM <sub>L</sub> 9	SEQ ID NO:26	SASLAIS	GLQ	SEDEADY	YCA	VMDPVL	NGPV	FGGGTKL	TVL	G.....	.....	.....	.....	.....	.....
AM <sub>L</sub> 8	SEQ ID NO:25	SASLAIS	GLQ	SEDEADY	YCA	VMDPVL	NGPV	FGGGTKL	TVL	G.....	.....	.....	.....	.....	.....
AM <sub>L</sub> 10	SEQ ID NO:27	TASLTIT	GAQ	AEDEADY	YCA	SRDSS	.NHV	FGGGTK	.....	.....	.....	.....	.....	.....	.....
AM <sub>L</sub> 1	SEQ ID NO:18	TATLTIS	GTQ	ARDEADY	YCA	AWDSS	.TASV	FGGGTK	.....	.....	.....	.....	.....	.....	.....
AM <sub>L</sub> 6	SEQ ID NO:23	SASLAIS	GLQ	SEDEADY	YCA	AWDSS	.TASV	FGGGTK	.....	.....	.....	.....	.....	.....	.....

FIGURE 3B



MNCRELPLTLWVLISVSTAESCTSRPHITVVEGEPFYLKHCSCSLAHEIETTTKSWYKSSGSGQEHVELNPR  
SSSRIALHDCVLEFWPVELNDTGSYFFQMKNYTQKWKLVIRRNKHSCFTERQVTSKIVEVKKFFQITCE  
NSYYQTLVNSTSLYKNCKKLLLENNKNPTIKKNAEFEDQGYSCVHFLHHNGKLFNITKTFNITIVEDRSNI  
VPVLLGPKLNHVAVELGKNVRLNCSALL**NEEDVIYWMFGEENGSDPNIHEEKEMRIMTPEGKWHASKVL**  
RIENIGESNLNVLYNCTVASTGGTDTKSFILVRKADMADIPGHVFTRGMIIAVLILVAVVCLVTVCVIYRVDLV  
LFYRHLTRRDETLTDGKTYDAFVSYLKECRPENGEETHFAVEILPRVLEKHFGYKLCIFERDVPPGAVVD  
EIHSLIEKSRRLIIVLSKSYMSNEVRYELESGLHEALVERKIKIILIEFTPVTDFLFPQSLKLLKSHRVLKWK  
DKSLSYNSRFWKNLLYLMPAKTVKPGRDEPEVLPVLS

**Amino acids of the human IL-18 receptor sequence that contribute to the antigen binding epitope are shown in bold (Amino Acids 243-271). Specific amino acids that were experimentally determined to influence binding are in bold.**

**FIGURE 5**

**AM<sub>H</sub>6 (SEQ ID NO:74)**

GTCGACGCCGCCACCATGGGGTCAACCGCCATCCTTGGCCTCCTCCTGGCTGT  
CCTGCAGGGAGGGCGCGCCGAGGTGCAGCTGGTGGAGTCTGGGGGAGGCGT  
GGTCCAGCCTGGGAGGTCCCTGAGACTCTCCTGTGCAGCGTCTGGATTCACCT  
TCAGCGGTTATGGCATGCACTGGGTCCGCCAGGCTCCAGGCAAGGGGGCTGGA  
GTGGGTGGCAGTAATATCAAATGATGGAAGTAAGAAATATTATTCAGACTCC  
GTGAAGGGCCGATTCACCATCTCCAGAGACAATTCCAAAAACACGCTGTATC  
TGCAGATGAACAGCCTGAGAGCTGAGGACACGGCTGTATATTACTGTGCGAA  
AGGGTCCAGTTCATATGGCTGACCTCGGCCCTGAACCTGTGGGGGCAGGGG  
ACCACGGTCACCGTCTCCTCAGCTAGCACCAAGGGCCCATCGGTCTTCCCCCT  
GGCGCCCTGCTCCAGGAGCACCTCCGAGAGCACAGCGGCCCTGGGCTGCCTG  
GTCAAGGACTACTTCCCCGAACCGGTGACGGTGTCTGTTGAACTCAGGCGCTC  
TGACCAGCGGCGTGCACACCTTCCCAGCTGTCCTACAGTCCTCAGGACTCTAC  
TCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAACTTCGGCACCCAGACCT  
ACACCTGCAACGTAGATCACAAGCCCAGCAACACCAAGGTGGACAAGACAG  
TTGAGCGCAAATGTTGTGTCGAGTGCCACCGTGCCAGCACCACCTGTGGC  
AGGACCGTCAGTCTTCTTCCCCCAAACCAAGGACACCCTCATGATCT  
CCCGGACCCCTGAGGTACAGTGGTGGTGGTGGACGTGAGCCACGAAGACCC  
CGAGGTCCAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAG  
ACAAAGCCACGGGAGGAGCAGTTCAACAGCACGTTCCGTGTGGTCAGCGTCC  
TCACCGTTGTGCACCAGGACTGGCTGAACGGCAAGGAGTACAAGTGCAAGGT  
CTCCAACAAAGGCCTCCAGCCCCCATCGAGAAAACCATCTCCAAAACCAAA  
GGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAG  
ATGACCAAGAACCAGGTGACCTGACCTGCCTGGTCAAAGGCTTCTACCCCA  
GCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGGAGAACAACTACA  
AGACCACACCTCCCATGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAG  
CTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCG  
TGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTC  
TCCGGGTAAATGAGCGGCCGC

**AM<sub>H</sub>6 (SEQ ID NO:73)**

MGSTAILGLLLAVLQGGRA<sup>^</sup>EVQLVESGGGVVQPGRSLRLSCAASGFTFSG  
YGMHWVRQAPGKLEWVAVISNDGSKKYYSDSVKGRFTISRDNKNTLYL  
QMNSLRAEDTAVYYCAKGSSSIWLTSALNLWGQGTTVTVSSASTKGPSVF  
PLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQS  
SGLYSLSVTVTPSSNFGTQTYTCNVDHKPSNTKVDKTVRKCCVECPPC  
PAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVD  
GVEVHNAKTKPREEQFNSTFRVVSVLTVVHQDWLNGKEYKCKVSNKGLPA  
PIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVE  
WESNGQPENNYKTTTPMLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHE  
ALHNHYTQKSLSLSPGK

**FIGURE 6**

**AM<sub>L</sub>12 (SEQ ID NO:76)**

GTCGACGTTTAAACGCCGCCACCAATGGAGACAGACACACTCCTGCTATGGGT  
ACTGCTGCTCTGGGTTCCAGGTTCCACTGGTTCGTCTGAGCTGACTCAGGACC  
CTGCTGTGTCTGTGGCCTTGGGACAGACAGTCAGGATCACATGCCAAGGAGA  
CAGCCTCAGAAGCTATTATGCAAGCTGGTACCAGCAGAAGCCAGGACAGGCC  
CCTGTACTIONTGTATCTCTGCTAAAAACAACCGGCCCTCAGGGATCCCAGACC  
GATTCTCTGGCTCCAGCTCAGGAAACACAGCTTCCTTGACCATCACTGGGGCT  
CAGGCGGAAGATGAGGCTGACTATTACTGTGCGTCCCGGAACGGCTGGAACC  
ATGTGGTATTCGGCGGAGGGACCAAGCTGACCGTCCTAGGCCAACCGAAAGC  
GGCGCCCTCGGTCACTCTGTTCCCGCCCTCCTCTGAGGAGCTTCAAGCCAACA  
AGGCCACACTGGTGTGTCTCATAAGTGACTTCTACCCGGGAGCCGTGACAGT  
GGCCTGGAAGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGGAGACCACCAC  
ACCCTCCAAACAAAGCAACAACAAGTACGCGGCCAGCAGCTATCTGAGCCTG  
ACGCCTGAGCAGTGGAAGTCCCACAGAAGCTACAGCTGCCAGGTCACGCATG  
AAGGGAGCACCGTGGAGAAGACAGTGGCCCTACAGAATGTTTCATAGGCGG  
CCGC

**AM<sub>L</sub>12 (SEQ ID NO:75)**

METDTLLLWVLLLWVPGSTG^SSELTQDPAVSVALGQTVRITCQGDSLRSYYA  
SWYQQKPGQAPVLVISAKNNRPSGIPDRFSGSSSGNTASLTITGAQAEDEADY  
YCASRNGWNHVVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLI  
SDFYPGAVTVAWKADSSPVKAGVETTTPSKQSNKYAASSYLSLTPEQWKS  
HRSYSCQVTHEGSTVEKTVAPTECS

**FIGURE 7**

**AM<sub>H</sub>4 (SEQ ID NO:78)**

ATGGGGTCAACCGCCATCCTTGGCCTCCTCCTGGCTGTCCTGCAGGGAGGGC  
GCGCCGAGGTGCAGCTGGTGGAGTCTGGGGGAGGCGTGGTCCAGCCTGGGA  
GGTCCCTGAGACTCTCCTGTGCAGCGTCTGGATTCACCTTCAGCGGTTATGGC  
ATGCACTGGGTCCGCCAGGCTCCAGGCAAGGGGCTGGAGTGGGTGGCAGTAA  
TATCAAATGATGGAAGTAAGAAATATTATTCAGACTCCGTGAAGGGCCGATT  
CACCATCTCCAGAGACAATCCAAAAACACGCTGTATCTGCAGATGAACAGC  
CTGAGAGCTGAGGACACGGCTGTATATTACTGTGCGAAAGGGTCCAGTTCCA  
TATGGCTGTCGCAGTCCCTGGACGGCTGGGGGCAGGGGACCACGGTCACCGT  
CTCCTCAGCTAGCACCAAGGGCCCATCGGTCTTCCCCCTGGCGCCCTGCTCCA  
GGAGCACCTCCGAGAGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTT  
CCCCGAACCGGTGACGGTGTCTGGAAGTCAAGGCGCTCTGACCAGCGGCGTG  
CACACCTTCCCAGCTGTCTACAGTCCCTCAGGACTCTACTCCCTCAGCAGCGT  
GGTGACCGTGCCCTCCAGCAACTTCGGCACCCAGACCTACACCTGCAACGTA  
GATCACAAGCCCAGCAACACCAAGGTGGACAAGACAGTTGAGCGCAAATGT  
TGTGTGAGTGCCACCGTGCCAGCACCACCTGTGGCAGGACCGTCAGTCT  
TCCTCTTCCCCCAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAG  
GTCACGTGCGTGGTGGTGGACGTGAGCCACGAAGACCCCGAGGTCCAGTTCA  
ACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCACGGG  
AGGAGCAGTTCAACAGCACGTTCCGTGTGGTCAGCGTCCCTACCGTTGTGCA  
CCAGGACTGGCTGAACGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGG  
CCTCCCAGCCCCATCGAGAAAACCATCTCCAAAACCAAAGGGCAGCCCCGA  
GAACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACCAAGAACC  
AGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCTACCCAGCGACATCGCCGT  
GGAGTGGGAGAGCAATGGGCAGCCGGAGAACAATAAGACCACACCTCC  
CATGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACA  
AGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGC  
TCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAATGA

**AM<sub>H</sub>4 (SEQ ID NO:77)**

MGSTAILGLLLAVLQGGRA<sup>^</sup>EVQLVESGGGVVQPGRSLRLSCAASGFTFSG  
YGMHWVRQAPGKGLEWVAVISNDGSKKYSDSVKGRFTISRDN SKNTLYL  
QMNSLRAEDTAVYYCAKGS SSIWLSQSLDGWGQGTTVTVSSASTKGPSVF  
PLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQS  
SGLYSLSSVVTVPSSNFGTQTYTCNV DHKPSNTKVDKTV ERKCCVECPPC  
PAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVD  
GVEVHNAKTKPREEQFNSTFRVVS VLTVVHQDWLNGKEYKCKVSNKGLPA  
PIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVE  
WESNGQPENNYKTTTPMLDSDGSFFLYSKLTVDKSRWQQGNV FSCSVMHE  
ALHNHYTQKSLSLSPGK

**FIGURE 8**

**AM<sub>L</sub>14 (SEQ ID NO:80)**

ATGGAGACAGACACACTCCTGCTATGGGTACTGCTGCTCTGGGTTCAGGTTCCAGGTTCCAGGTTCCGTTCTGAGCTGACTCAGGACCCTGCTGTGTCTGTGGCCTTGGGACAGACAGTCAGGATCACATGCCAAGGAGACAGCCTCAGAAGCTATTATGCAAGCTGGTACCAGCAGAAGCCAGGACAGGCCCTGTACTTGTCATCTCTGCTAAAACAACCGGCCCTCAGGGATCCCAGACCGATTCTCTGGCTCCAGCTCAGGAAACACAGCTTCCTTGACCATCACTGGGGCTCAGGCGGAAGATGAGGCTGACTATTACTGTGCGTCCCGGAACGGCTGGAACCATGTGGTATTCGGCGGAGGGACC AAGCTGACCGTCCTAGGCCAACCGAAAGCGGGCCCTCGGTCACTCTGTTCCCGCCCTCCTCTGAGGAGCTTCAAGCCAACAAGGCCACACTGGTGTGTCTCAT AAGTGACTTCTACCCGGGAGCCGTGACAGTGGCCTGGAAGGCAGATAGCAGC CCGTCAAGGCGGGAGTGGAGACCACACCCCTCCAAACAAAGCAACAAC AAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCAGTGGAAGTCCC ACAGAAGCTACAGCTGCCAGGTCACGCATGAAGGGAGCACCGTGGAGAAGA CAGTGGCCCCTACAGAATGTTTCATAGGCGGCCGC

**AM<sub>L</sub>14 (SEQ ID NO:79)**

METDTLLLWVLLLWVPGSTG<sup>^</sup>SSELTQDPAVSVALGQTVRITCQGDSLRSYYA SWYQQKPGQAPVLVISAKNNRPSGIPDRFSGSSSGNTASLTITGAQAEDEADY YCASRNGWNHVVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLI SDFYPGAVTVAWKADSSPVKAGVETTTPSKQSNKYAASSYLSLTPEQWKS HRSYSCQVTHEGSTVEKTVAPTECS

**FIGURE 9**



**AM<sub>H</sub>9 (SEQ ID NO:82)**

ATGGGGTCAACCGCCATCCTTGGCCTCCTCCTGGCTGTCCTGCAGGGAGGGC  
GCGCCAGGTGCAGCTGGTGCAGTCTGGGGCTGAGGTGAAGAAGCCTGGGGC  
CTCAGTGAAGGTCTCCTGCAAGGTTTCCGGATACACCCTCACTGAATTATCCA  
TGCACTGGGTGCGACAGGCTCCTGGAAAAGGGCTTGAGTGGATGGGAGGTTT  
TGATCGTGAAGATGATGAAACAATCCACGCACAGAAGTTCCAGGGCAGAGTC  
ACCATGACCGAGGACACATCTACAGACACAGCCTACATGGAAGTGAAGCAGCC  
TGCGATCTGAGGACACGGCCGTTTATTACTGTGCAACAGATCTTATGGTGTGG  
GGCGATTTTTGGATCCAGCACTGGGGCCAGGGGACACTGGTCACCGTCTCCT  
CAGCTAGCACCAAGGGCCCATCGGTCTTCCCCCTGGCGCCCTGCTCCAGGAG  
CACCTCCGAGAGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCC  
GAACCGGTGACGGTGTCTGGAAGTCAAGGCGCTCTGACCAGCGGCGTGCACA  
CCTTCCCAGCTGTCCTACAGTCTCAGGACTCTACTCCCTCAGCAGCGTGGTG  
ACCGTGCCCTCCAGCAACTTCGGCACCCAGACCTACACCTGCAACGTAGATC  
ACAAGCCCAGCAACACCAAGGTGGACAAGACAGTTGAGCGCAAATGTTGTGT  
CGAGTGCCACCGTGCCAGCACCACCTGTGGCAGGACCGTCAGTCTTCTCT  
TCCCCCAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTCAC  
GTGCGTGGTGGTGGACGTGAGCCACGAAGACCCCGAGGTCCAGTTCAACTGG  
TACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCACGGGAGGAG  
CAGTTCAACAGCACGTTCCGTGTGGTCAAGCTCCTCACCGTTGTGCACCAGG  
ACTGGCTGAACGGCAAGGAGTACAAGTGAAGGTCTCCAACAAAGGCCT  
CCCAGCCCCATCGAGAAAACCATCTCCAAAACCAAAGGGCAGCCCCGAGA  
ACCACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACCAAGAACCAG  
GTCAGCCTGACCTGCCTGGTCAAAGGCTTCTACCCCAGCGACATCGCCGTGG  
AGTGGGAGAGCAATGGGCAGCCGGAGAACAATAAGACCACACCTCCCA  
TGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAG  
AGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTC  
TGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAATGA

**AM<sub>H</sub>9 (SEQ ID NO:81)**

MGSTAILGLLLAVLQGGRA^QVQLVQSGAEVKKPGASVKVSCKVSGYTLTE  
LSMHWVRQAPGKGLEWMGGFDREDETIHAQKFQGRVTMTEDTSTDYAYM  
ELSSLRSEDVAVYYCATDLMVWGFVWQHWGQGLVTVSSASTKGPSVFP  
LAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS  
GLYSLSSVTVPSNFGTQTYTCNVDHKPSNTKVDKTKVERKCCVECP  
APPVAGPSVFLFPPKPKDTLMISRTPETVTCVVVDVSHEDPEVQFNWYVDG  
VEVHNAKTKPREEQFNSTFRVVSVLTVVHQQDWLNGKEYKCKVSNKGLPAP  
IEKTIKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEW  
ESNGQPENNYKTTTPMLDSDGSFFLYSKLTVDKSRWQQGNVVFSCSVMHEA  
LHNHYTQKSLSLSPGK

**FIGURE 10**



**AM<sub>H</sub>11 (SEQ ID NO:86)**

ATGGGGTCAACCGCCATCCTTGGCCTCCTCCTGGCTGTCCTGCAGGGAGGGC  
GCGCCCAGGTGCAGCTGGTGCAGTCTGGGGCTGAGGTGAAGAAGCCTGGGGC  
CTCAGTGAAGGTCTCCTGCAAGGTTTCCGGATACACCCTCACTGAATTATCCA  
TGCACTGGGTGCGACAGGCTCCTGGAAAAGGGCTTGAGTGGATGGGAGGTTT  
TGATCGTGAAGATGATGAAACAATCCACGCACAGAAGTTCCAGGGCAGAGTC  
ACCATGACCGAGGACACATCTACAGACACAGCCTACATGGAAGTGAAGCAGCC  
TGCGATCTGAGGACACGGCCGTTTATTACTGTGCAACAGATCTTATGGTGTGG  
AACTTCCCCCCCATCCAGCACTGGGGCCAGGGGACACTGGTCACCGTCTCCT  
CAGCTAGCACCAAGGGCCCATCGGTCTTCCCCCTGGCGCCCTGCTCCAGGAG  
CACCTCCGAGAGCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCC  
GAACCGGTGACGGTGTCTGGAAGTCAAGGCGCTCTGACCAGCGGCGTGCACA  
CCTTCCCAGCTGTCTACAGTCTCAGGACTCTACTCCCTCAGCAGCGTGGTG  
ACCGTGCCCTCCAGCAACTTCGGCACCCAGACCTACACCTGCAACGTAGATC  
ACAAGCCCAGCAACACCAAGGTGGACAAGACAGTTGAGCGCAAATGTTGTGT  
CGAGTGCCACCGTGCCAGCACCACTGTGGCAGGACCGTCAGTCTTCTCT  
TCCCCCAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTCAC  
GTGCGTGGTGGTGGACGTGAGCCACGAAGACCCCGAGGTCCAGTTCAACTGG  
TACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCACGGGAGGAG  
CAGTTCAACAGCACGTTCCGTGTGGTCAAGCGTCTCACCCTGTCACCAGG  
ACTGGCTGAACGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAGGCCTCCC  
AGCCCCATCGAGAAAACCATCTCCAAAACCAAGGGCAGCCCCGAGAACC  
ACAGGTGTACACCCTGCCCCCATCCCGGGAGGAGATGACCAAGAACCAGGTC  
AGCCTGACCTGCCTGGTCAAAGGCTTCTACCCAGCGACATCGCCGTGGAGT  
GGGAGAGCAATGGGCAGCCGGAGAACAATAAGACCACACCTCCCATGC  
TGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAG  
CAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTG  
CACAACTACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAATGA

**AM<sub>H</sub>11 (SEQ ID NO:85)**

MGSTAILGLLLAVLQGGRA<sup>^</sup>QVQLVQSGAEVKKPGASVKVSKVSGYTLTE  
LSMHWVRQAPGKGLEWMGGFDREDETIHAQKFQGRVTMTEDTSTDYAYM  
ELSSLRSEDVAVYYCATDLMVWNFPPIQHWGQGLVTVSSASTKGPSVFP  
LAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS  
GLYSLSSVTVPSNFGTQTYTCNVDPKPSNTKVDKTKVERKCCVECP  
APPVAGPSVFLFPPKPKDTLMISRTPPEVTCVVVDVSHEDPEVQFNWYVDG  
VEVHNAKTKPREEQFNSTFRVVSVLTVVHQQDWLNGKEYKCKVSNKGLPAP  
IEKTIKTKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEW  
ESNGQPENNYKTTTPMLDSDGSFFLYSKLTVDKSRWQQGNVFSQVMSHEA  
LHNHYTQKSLSLSPGK

**FIGURE 12**

**AM<sub>L</sub>7 (SEQ ID NO:88)**

ATGGAGACAGACACTCCTGCTATGGGTACTGCTGCTCTGGGTTCCAGGTTCC  
CACTGGTGCTAGCCAGTCTGTGCTGACTCAGCCACCCTCAGCGTCTGGGACCC  
CCGGGCAGAGGGTCACCATCTCTTGTCTGGAAGGAACTCCAACA'ICGGAAG  
TTATACTGTAACCTGGTACCAGCAGCTCCCAGGAACGGCCCCCAAACCTCCTC  
ATCTATAGTAATAGTCAGCGGCCCTCAGGGGTCCCTGACCGATTCTCAGGCTC  
CAAGTCTGGCACCTCAGCCTCCTTGGCCATCAGTGGGCTCCAGTCTGAAGAT  
GAGGCTGATTACTGTGTGGTGTGGGATGACGTGCTGAATGGCCCCGGTGT  
CGGCGGAGGGACCAAGCTGACCGTCCTAGGCCAACCGAAAGCGGCGCCCTC  
GGTCACTCTGTTCCCGCCCTCCTCTGAGGAGCTTCAAGCCAACAAGGCCACA  
CTGGTGTGTCTCATAAGTGACTTCTACCCGGGAGCCGTGACAGTGGCCTGGA  
AGGCAGATAGCAGCCCCGTCAAGGCGGGAGTGGAGACCACCACACCCTCCA  
AACAAAGCAACAACAAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTG  
AGCAGTGGAAGTCCCACAGAAGCTACAGCTGCCAGGTACGCATGAAGGGA  
GCACCGTGGAGAAGACAGTGGCCCCCTACAGAATGTTCATAG

**AM<sub>L</sub>7 (SEQ ID NO:87)**

METDTLLLWVLLLWVPGSTG^ASQSVLTQPPSASGTPGQRVTISCSGRNSN  
IGSYTVTWYQQLPGTAPKLLIYSNSQRPSGVPDRFSGSKSGTSASLAISG  
LQSEDEADYYCVVWDDVLNGPVFGGGTKLTVLGQPKAAPSVTLFPPSSEE  
LQANKATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTPSKQSNNKYAA  
SSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS

**FIGURE 13**

## IL-18 RECEPTOR ANTIGEN BINDING PROTEINS

This application is a National Stage application under 35 U.S.C. §371 of International Application No. PCT/US2008/071047 (which designated the United States), having an international filing date of Jul. 24, 2008, which claims the priority benefit of U.S. Provisional Patent Application Ser. No. 61/073,142, filed Jun. 17, 2008, U.S. Provisional Patent Application Ser. No. 60/951,692, filed Jul. 24, 2007, and U.S. Provisional Patent Application Ser. No. 60/951,691, filed Jul. 24, 2007 each of which is hereby incorporated by reference in its entirety.

### REFERENCE TO THE SEQUENCE LISTING

The present application is being filed along with a Sequence Listing in electronic format. The Sequence Listing is provided as a file entitled Sequence Listing.txt, created Jan. 21, 2010, which is 134 KB in size. The information in the electronic format of the Sequence Listing is incorporated herein by reference in its entirety.

### I. FIELD OF THE INVENTION

Provided herein are IL-18 receptor antigen binding proteins and polynucleotides encoding the same. Expression vectors and host cells comprising the same for production of the antigen binding proteins are also provided. In addition, provided are compositions and methods for diagnosing and treating diseases mediated by IL-18 receptor.

### II. BACKGROUND

IL-18 is a proinflammatory cytokine that belongs to the IL-1 family of ligands. Okamura et al., 1995, *Nature* 378:88-91. Also referred to as IFN- $\gamma$ -inducing factor, IL-18 is a cytokine that plays an important role in the TH1 response, primarily by its ability to induce IFN- $\gamma$  production in T cells and natural killer cells. IL-18 is related to the IL-1 family in both structure and function. In terms of structure, IL-18 and IL-1 $\beta$  share significant primary amino acid sequences and are both folded as  $\beta$ -sheet polypeptides. In terms of function, IL-18 induces gene expression and synthesis of IL-1, TNF, Fas ligand, and several cytokines.

The activity of IL-18 is transduced through a signal transducing pathway initiated by its forming of a IL-18 receptor (IL-18R) complex. The IL-18R includes a binding chain termed  $\alpha$ -IL-18 receptor ( $\alpha$ -IL-18R), a member of the IL-1R family previously identified as the IL-1R-related protein (IL-1Rrp), and a  $\beta$ -IL-18 receptor ( $\beta$ -IL-18R), also a member of the IL-1R family and previously identified as AcPL; both chains are required for signaling. Born et al., 1998, *J. Biol. Chem.* 273:29445-50. The IL-18/IL-18R complex recruits IL-1R-activating kinase and TNF receptor-associated factor-6, which phosphorylates nuclear factor kappaB (NFkappaB)-inducing kinase with subsequent activation of NFkappaB. IL-18 participates in both innate and acquired immunity. Dinarello, 1999, *J. Allergy Clin. Immunol.* 103:11-24.

Increased levels of IL-18 and/or involvement of IL-18 mediated signals in pathogenesis have been demonstrated in a variety of human disease states, including autoimmune diseases (WO2004/002519; WO2005/063290; WO2004/034988; Mallet et al., 2002, *Circ. Res.* 91:441-448), hepatic diseases (Finitto et al., 2004, *Liver* 53:392-400; Tsutsui et al., 2000, *Immunological Reviews* 174:192-209; Ludwiczek et al., 2002, *J. Clinical Immunology* 22:331-337), pancreatic

diseases, and cardiovascular diseases (Gerdes et al, 2002, *J. Exp. Med.* 195:245-257; WO03/080104; WO02/060479; WO01/85201; Raeburn et al., 2002, *Am. J. Physiol. Heart Circ. Physiol.* 283:H650-H657). Accordingly, it is desirable to generate new agents capable of modulating the IL-18/IL-18 receptor interaction.

### III. SUMMARY

Provided herein are  $\alpha$ - and  $\beta$ -IL-18 receptor (also referred to herein collectively as "IL-18 receptor" or "IL-18R") antigen binding proteins and polynucleotides that encode them. The IL-18 receptor antigen binding proteins inhibit, interfere with, or modulate at least one of the biological responses mediated by IL-18 and as such can be useful for ameliorating the effects of IL-18 mediated diseases or disorders. Also provided are expression systems, including cell lines, for the production of  $\alpha$ - and  $\beta$ -IL-18 receptor antigen binding proteins and methods for diagnosing and treating diseases associated with aberrant IL-18 activity.

In one embodiment, antigen binding proteins bind the  $\alpha$ - and  $\beta$ -IL-18 receptor, and comprise (a) a scaffold structure; and (b) at least one complementary determining region (CDR), selected from the CDRH regions of any of SEQ ID NOs:89-139 or the CDRL regions of any of SEQ ID NOs:140-190. In this embodiment, of particular use are antigen binding proteins with a CDRH3 or CDRL3 region of SEQ ID NO:91, 94, 97, 100, 103, 106, 109, 112, 115, 118, 121, 124, 127, 130, 133, 136, 139 or SEQ ID NO:142, 145, 148, 151, 154, 157, 160, 163, 166, 169, 172, 175, 178, 181, 184, 187, 190, respectively. Additional embodiments utilize antigen binding proteins with one CDR selected from the CDRH regions of any of SEQ ID NOs:89-139 and a CDRL region of any of SEQ ID NOs:140-190 (e.g., the antigen binding protein has two CDR regions, one heavy and one light; again, in a specific embodiment the antigen binding proteins have both a CDRH3 and a CDRL3 region).

The antigen binding proteins can bind to an IL-18 receptor  $\alpha$ - or  $\beta$ -chain having the amino acid sequence of SEQ ID NO:69 or SEQ ID NO:71, respectively.

Described herein are antigen binding proteins that comprise a heavy chain amino acid sequence that comprises at least one CDR selected from the group consisting of: (a) a CDRH1 of any of SEQ ID NOs:89, 92, 95, 98, 101, 104, 107, 110, 113, 116, 119, 122, 125, 128, 131, 134, 137; (b) a CDRH2 of any of SEQ ID NOs:90, 93, 96, 99, 102, 105, 108, 111, 114, 117, 120, 123, 126, 129, 132, 135, 138; and (c) a CDRH3 of any of SEQ ID NOs:91, 94, 97, 100, 103, 106, 109, 112, 115, 118, 121, 124, 127, 130, 133, 136, 139; and/or a light chain amino acid sequence that comprises at least one CDR selected from the group consisting of: (a) a CDRL1 of any of SEQ ID NOs:140, 143, 146, 149, 152, 155, 158, 161, 164, 167, 170, 173, 176, 179, 182, 185, 188; (b) a CDRL2 of any of SEQ ID NOs:141, 144, 147, 150, 153, 156, 159, 162, 165, 168, 171, 174, 177, 180, 183, 186, 189; and (c) a CDRL3 of any of SEQ ID NOs:142, 145, 148, 151, 154, 157, 160, 163, 166, 169, 172, 175, 178, 181, 184, 187, 190.

In certain aspects, the antigen binding protein comprises a heavy chain amino acid sequence having a CDRH1, a CDRH2, and a CDRH3 of any of SEQ ID NOs:89-139, and/or a light chain amino acid sequence that comprises a CDRL1, a CDRL2, and a CDRL3 of any of SEQ ID NOs:140-190. Preferred antigen binding proteins comprise a heavy chain amino acid sequence selected from the group consisting of SEQ ID NOs:1-17 and/or a light chain amino acid sequence selected from the group consisting of SEQ ID NOs:18-34. Preferred CDRH3s include those set forth in any of SEQ ID

NOs:91, 94, 97, 100, 103, 106, 109, 112, 115, 118, 121, 124, 127, 130, 133, 136, 139. Preferred CDRL3s include those set forth in any of SEQ ID NOs:142, 145, 148, 151, 154, 157, 160, 163, 166, 169, 172, 175, 178, 181, 184, 187, 190.

In certain aspects, the antigen binding protein comprises one or more IgG heavy or light chains, including those of the IgG1-, IgG2-IgG3- or IgG4-type. Preferred IgG heavy chains include, but are not limited to, those set forth in SEQ ID NO:73, 77, 81, and 85. Preferred IgG light chains include, but are not limited to, those set forth in SEQ ID NO:75, 79, 83, and 87.

As described herein, antigen binding proteins that bind to amino acid residues 250-253 and 267-271 of a three dimensional structure formed by mature  $\alpha$ -IL-18 receptor (SEQ ID NO:69) are particularly useful in blocking the interaction of IL-18 with IL-18 receptor.

An antigen binding protein can be a monoclonal antibody, a human antibody, a recombinant antibody, a chimeric antibody, a humanized antibody, a bispecific antibody, or a fragment thereof. Antibody fragments include, but are not limited to, a minibody, a domain antibody, a F(ab) fragment, a F(ab') fragment, a F(ab')<sub>2</sub> fragment, a Fv fragment, a scFv fragment, a Fd fragment, a diabody, or a single chain antibody molecule.

In other aspects, provided herein are isolated nucleic acids encoding one or more IL-18 receptor antigen binding proteins. Such nucleic acids can be comprised within a vector and operably linked to a control sequence. Also, provided herein are host cells transformed with such isolated nucleic acids.

Additionally, provided herein are pharmaceutical compositions comprising an IL-18 receptor antigen binding protein and a pharmaceutically acceptable carrier. Such pharmaceutical compositions are useful in methods for preventing or treating a condition associated with IL-18 receptor in a patient, which comprise administering an effective amount thereof to the patient. Diseases and conditions associated with IL-18 receptor include inflammatory and autoimmune diseases (such as psoriasis, rheumatoid arthritis, juvenile idiopathic arthritis, Still's disease, ankylosing spondylitis, osteoarthritis, ulcerative arthritis, celiac disease, psoriatic arthritis, chronic obstructive pulmonary disease, asthma, particularly chronic severe asthma, acute respiratory distress syndrome, sepsis, Alzheimer disease, lupus, allergic rhinitis, idiopathic thrombocytopenic purpura, transplantation, atopic dermatitis, type II diabetes, Crohn's disease, inflammatory bowel disease, multiple sclerosis, autoimmune hepatitis, HIV, atopic dermatitis, myasthenia gravis, sarcoidosis), a hepatic disease (such as hepatitis), a pancreatic disease (such as chronic pancreatitis or acute pancreatitis), and a cardiovascular disease (such as acute heart attacks, atheromatous plaque rupture, post-ischemic heart failure, reperfusion injury, vascular inflammation, chronic heart failure, atherosclerosis, cardiovascular complications of rheumatoid arthritis, and atherogenesis).

Further provided herein are methods of inhibiting the binding of IL-18 to IL-18 receptor comprising contacting an IL-18 receptor with an IL-18 receptor antigen binding protein. Upon binding IL-18 receptor, the IL-18 receptor antigen binding protein will prevent or block binding of the receptor to IL-18.

#### IV. DESCRIPTION OF THE DRAWINGS

FIGS. 1A, 1B, 1C, 1D, 1E, 1F, 1G, 1H, 1I, 1J, 1K, 1L, 1M, 1N, 1O, 1P, and 1Q depict nucleic acid and amino acid sequences of VH and VL variable domains of  $\alpha$ - and  $\beta$ -IL-18 receptor antigen binding proteins.

FIGS. 2A, 2B, 2C, 2D, and 2E show the CDR1, CDR2, and CDR3 regions of various heavy and light chain variable regions of antigen binding proteins. The amino acid sequences of the various heavy and light chain regions are identified in SEQ ID NOs:1-34. The sequences of the individual CDRs are identified in SEQ ID NOs:89-190.

FIGS. 3A and 3B depict an alignment of the amino acid sequences of heavy and light chain variable sequences of  $\alpha$ - and  $\beta$ -IL-18 receptor antigen binding proteins. The CDR1, CDR2 and CDR3 regions are highlighted in grey.

FIG. 4 depicts a chart showing various possible combinations of heavy chain variable regions and light chain variable region sequences. Shown are dimers of each one heavy and one light chain variable region. As naturally occurring antibodies typically are tetramers, an antibody may comprise a combination of two of the depicted dimers.

FIG. 5 depicts the portions of the  $\alpha$ -IL-18 receptor amino acid sequences that form the epitope for a specific antigen binding protein embodiment.

FIG. 6 depicts the complete AM<sub>H</sub>6 heavy chain nucleotide and amino acid sequences (SEQ ID NOs:74 and 73, respectively). The arrow indicates the cleavage site of the leader sequence.

FIG. 7 depicts the complete AM<sub>L</sub>12 light chain nucleotide and amino acid sequences (SEQ ID NOs:76 and 75, respectively). The arrow indicates the cleavage site of the leader sequence.

FIG. 8 depicts the complete AM<sub>H</sub>4 heavy chain nucleotide and amino acid sequences (SEQ ID NOs:78 and 77, respectively). The arrow indicates the cleavage site of the leader sequence.

FIG. 9 depicts the complete AM<sub>L</sub>14 light chain nucleotide and amino acid sequences (SEQ ID NOs:80 and 79, respectively). The arrow indicates the cleavage site of the leader sequence.

FIG. 10 depicts the complete AM<sub>H</sub>9 heavy chain nucleotide and amino acid sequences (SEQ ID NOs:82 and 81, respectively). The arrow indicates the cleavage site of the leader sequence.

FIG. 11 depicts the complete AM<sub>L</sub>9 light chain nucleotide and amino acid sequences (SEQ ID NOs:84 and 83, respectively). The arrow indicates the cleavage site of the leader sequence.

FIG. 12 depicts the complete AM<sub>H</sub>11 heavy chain nucleotide and amino acid sequences (SEQ ID NOs:86 and 85, respectively). The arrow indicates the cleavage site of the leader sequence.

FIG. 13 depicts the complete AM<sub>L</sub>7 light chain nucleotide and amino acid sequences (SEQ ID NOs:88 and 87, respectively). The arrow indicates the cleavage site of the leader sequence.

#### V. DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described.

Standard techniques may be used for recombinant DNA, oligonucleotide synthesis, tissue culture and transformation, protein purification etc. Enzymatic reactions and purification techniques may be performed according to the manufacturer's specifications or as commonly accomplished in the art or as described herein. The following procedures and techniques may be generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed

throughout the specification. See, e.g., Sambrook et al., 2001, *Molecular Cloning: A Laboratory Manual*, 3<sup>rd</sup> ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., which is incorporated herein by reference for any purpose. Unless specific definitions are provided, the nomenclature used in connection with, and the laboratory procedures and techniques of, analytical chemistry, organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art. Standard techniques may be used for chemical synthesis, chemical analyses, pharmaceutical preparation, formulation, and delivery and treatment of patients.

#### A. General Overview

Provided herein are antigen binding proteins that bind an  $\alpha$ - or  $\beta$ -IL-18 receptor; the amino acid sequence of the human  $\alpha$ - and  $\beta$ -IL-18 receptor are depicted in SEQ ID NOs:69 and 71, respectively. The antigen binding proteins of the invention comprise a scaffold structure with one or more complementarity-determining region (CDRs) as depicted in FIGS. 2A-2E, 3A and 3B, namely the CDRH1, CDRH2, CDRH3, CDRL1, CDRL2 and CDRL3 portion of SEQ ID NOs:1-34 (see, also SEQ ID NOs:89-292, depicting the amino acid sequences of the various CDRs). In certain embodiments, the scaffold structure of the antigen binding proteins is based on antibodies (including, but not limited to, monoclonal antibodies, human antibodies, murine antibodies, chimeric antibodies, humanized antibodies, bispecific antibodies), antibody fragments (such as minibodies, domain antibodies, F(ab) fragments, F(ab') fragments, F(ab)<sub>2</sub> fragments, F(ab')<sub>2</sub> fragments, Fv fragments, scFv fragments, Fd fragments), synthetic antibodies (sometimes referred to herein as "antibody mimetics"), antibody fusions (sometimes referred to as "antibody conjugates"), including Fc fusions. The various structures are further described and defined hereinbelow.

$\alpha$ - and  $\beta$ -IL-18 receptor antigen binding proteins are useful in treating conditions associated with IL-18 activity, including TH1-driven autoimmune diseases (WO2004/002519; WO2005/063290; WO2004/034988; Mallet et al., 2002, *Circ. Res.* 91:441-448), hepatic diseases (Finitto et al., 2004, *Liver* 53:392-400; Tsutsui et al., 2000, *Immunological Reviews* 174:192-209; Ludwiczek et al., 2002, *J. Clinical Immunology* 22:331-337), pancreatic diseases (Yoshida et al., 1998, *Anticancer Res.* 18:333-5), and cardiovascular diseases (Gerdes et al., 2002, *J. Exp. Med.* 195:245-257; WO03/080104; WO02/060479; WO01/85201; Raeburn et al., 2002, *Am. J. Physiol. Heart Circ. Physiol.* 283:H650-H657), as is further described below. Other uses for antigen binding proteins include, for example, diagnosis of IL-18 associated diseases or conditions and screening assays to determine the presence or absence of the  $\alpha$ - or  $\beta$ -IL-18 receptor. Also provided are  $\alpha$ - or  $\beta$ -IL-18 receptor antigen binding proteins, particularly antigen binding proteins that include at least one complementarity determining region (CDR) including heavy and/or light CDRs, as more fully described below, and combinations thereof.

The antigen binding proteins of the invention interfere with, block or modulate the interaction between IL-18 and the IL-18 receptor. In some embodiments, the antigen binding proteins interrupt the IL-18 pathway, thereby decreasing at least one biological activity of IL-18, including, but not limited to, induction of IFN- $\gamma$  production, induction of killer cell formation, and enhancement of cytotoxicity of killer cells. As demonstrated in the Examples herein, antigen binding proteins that reduce IL-18 induced production of IFN- $\gamma$  by KG cells include those comprising AM<sub>H</sub>8 and AM<sub>L</sub>11, AM<sub>H</sub>9 and AM<sub>L</sub>9, AM<sub>H</sub>10 and AM<sub>L</sub>8, AM<sub>H</sub>11 and AM<sub>L</sub>7, AM<sub>H</sub>15 and AM<sub>L</sub>3, AM<sub>H</sub>13 and AM<sub>L</sub>4, AM<sub>H</sub>13 and AM<sub>L</sub>5, AM<sub>H</sub>16 and

AM<sub>L</sub>2, AM<sub>H</sub>2 and AM<sub>L</sub>16, AM<sub>H</sub>2 and AM<sub>L</sub>17, AM<sub>H</sub>1 and AM<sub>L</sub>16, AM<sub>H</sub>1 and AM<sub>L</sub>17, AM<sub>H</sub>4 and AM<sub>L</sub>14, AM<sub>H</sub>4 and AM<sub>L</sub>15, AM<sub>H</sub>3 and AM<sub>L</sub>14, AM<sub>H</sub>3 and AM<sub>L</sub>15, AM<sub>H</sub>6 and AM<sub>L</sub>12, AM<sub>H</sub>6 and AM<sub>L</sub>13, AM<sub>H</sub>5 and AM<sub>L</sub>12, and AM<sub>H</sub>5 and AM<sub>L</sub>13.

The antigen binding proteins of the invention thus may serve to identify conditions related to IL-18 or IL-18 receptor induced immune responses. In addition, the antigen binding proteins may be utilized to regulate and/or suppress IL-18 or IL-18 receptor mediated immune responses, as such having efficacy in the treatment and prevention of various diseases caused by excessive immune responses, e.g., inflammatory diseases. Accordingly, the  $\alpha$ - and  $\beta$ -IL-18 receptor antigen binding proteins of the present invention can be used for the diagnosis, prevention or treatment of diseases or conditions associated with the IL-18 and IL-18 receptor mediated signal transduction pathway.

#### B. IL-18 Receptor Antigen Binding Proteins

In one aspect, antigen binding proteins that bind an  $\alpha$ - or  $\beta$ -IL-18 receptor are provided. By "antigen binding protein" as used herein is meant a protein that specifically binds a specified antigen. In specific embodiments, the antigen is a human  $\alpha$ - or  $\beta$ -IL-18 receptor.

By "protein," as used herein, is meant at least two covalently attached amino acids, which includes proteins, polypeptides, oligopeptides and peptides. In some embodiments, the two or more covalently attached amino acids are attached by a peptide bond. The protein may be made up of naturally occurring amino acids and peptide bonds, for example when the protein is made recombinantly using expression systems and host cells, as outlined below. Alternatively, the protein may include synthetic amino acids (e.g., homophenylalanine, citrulline, ornithine, and norleucine), or peptidomimetic structures, i.e., "peptide or protein analogs", such as peptoids (see, Simon et al., 1992, *Proc. Natl. Acad. Sci. U.S.A.* 89:9367, incorporated by reference herein), which can be resistant to proteases or other physiological and/or storage conditions. Such synthetic amino acids may be incorporated in particular when the antigen binding protein is synthesized in vitro by conventional methods well known in the art. In addition, any combination of peptidomimetic, synthetic and naturally occurring residues/structures can be used. "Amino acid" also includes imino acid residues such as proline and hydroxyproline. The amino acid "R group" or "side chain" may be in either the (L)- or the (D)-configuration. In a specific embodiment, the amino acids are in the (L)- or (D)-configuration.

In certain aspects, the invention provides recombinant antigen binding proteins that bind an IL-18 receptor, in some embodiments a human IL-18 receptor. In this context, a "recombinant protein" is a protein made using recombinant techniques, i.e., through the expression of a recombinant nucleic acid as described herein. Methods and techniques for the production of recombinant proteins are well known in the art.

In some embodiments, the antigen binding proteins are isolated proteins or substantially pure proteins. An "isolated" protein is unaccompanied by at least some of the material with which it is normally associated in its natural state, preferably constituting at least about 5%, more preferably at least about 50% by weight of the total protein in a given sample. A "substantially pure" protein comprises at least about 75% by weight of the total protein, with at least about 80% being preferred, and at least about 90% being particularly preferred. The definition includes the production of an antigen binding protein from one organism in a different organism or host cell. Alternatively, the protein may be made at a significantly

higher concentration than is normally seen, through the use of an inducible promoter or high expression promoter, such that the protein is made at increased concentration levels.

The antigen binding proteins can specifically bind to an IL-18 receptor, preferably a human IL-18 receptor. “Specifically binds” as used herein means the equilibrium dissociation constant is at least  $10^{-6}$  M, preferably  $10^{-7}$  to  $10^{-10}$  M, more preferably  $<10^{-8}$  to  $<10^{-10}$  M, even more preferably  $<10^{-9}$  to  $<10^{-10}$  M. In a specific embodiment, the antigen binding protein binds to a human IL-18 receptor having the amino acid sequence of SEQ ID NO:69 or 71. An epitope in the  $\alpha$ - or  $\beta$ -IL-18 receptor to which preferred antigen binding proteins specifically bind is detailed below.

In embodiments where the antigen binding protein is used for therapeutic applications, an important characteristic of an IL-18 receptor antigen binding protein is whether it can inhibit, interfere with or modulate one or more biological activities of an IL-18 receptor. In this case, an antigen binding protein binds specifically and/or substantially inhibits binding of IL-18 to its receptor when an excess of antibody reduces the quantity of IL-18 bound to IL-18 receptor, or vice versa, by at least about 20%, 40%, 60%, 80%, 85%, or more (for example by measuring binding in an in vitro competitive binding assay). IL-18 receptor has many distinct biological effects, which can be measured in many different assays in different cell types. The ability of an IL-18 receptor antigen binding protein to inhibit, interfere with, or modulate the biological activity of IL-18 can be determined, for example, by measuring the inhibition of IFN- $\gamma$  release in KG1 cells, as described in Example 4 or using a similar assay in which the ability of an antigen binding protein to inhibit IFN- $\gamma$  release is measured.

Not every antigen binding protein that specifically binds to an antigen can block antigen binding to its normal ligand and thus inhibit or modulate the biological effects of the antigen. As is known in the art, such an effect can depend on what portion of the antigen the antigen binding protein binds to, and on both the absolute and the relative concentrations of the antigen and the antigen binding protein, in this case, an IL-18 receptor and the IL-18 receptor antigen binding protein. To be considered capable of inhibiting or modulating the biological activity of an IL-18 receptor as meant herein, an antigen binding protein may be able, for example, to inhibit the release of IFN- $\gamma$  observed in the presence of IL-18, as measured in the KG1 cell assay of Example 4 or a similar assay, by at least about 20%, 40%, 60%, 80%, 85%, 90%, 95%, 99%, or more when the IL-18 concentration is within a range, for example, at about  $EC_{80}$  or  $EC_{90}$ , where the effects of an agent that inhibits its biological activity can be readily apparent. An  $EC_{80}$ , as meant herein, is the amount of IL-18 required for 80% of the maximal effect of IL-18 to be observed. If the IL-18 concentration is well above  $EC_{90}$ , effects of an inhibiting agent may be less apparent due to the excess of IL-18. The concentration of an antigen binding protein required to inhibit, interfere with or modulate the biological activity of IL-18 receptor can vary widely and may depend upon how tightly the antibody binds to IL-18 receptor. For example, one molecule or less of an antigen binding protein per molecule of IL-18 may be sufficient to inhibit, interfere with or modulate biological activity in the KG1 cell assay. In some embodiments, a ratio of IL-18 receptor/antibody of about 1,000:1 to about 1:1,000, including about 2:1, 1:1, 1:2, 1:4, 1:6, 1:8, 1:10, 1:20, 1:40, 1:60, 1:100, 1:500, 1:1,000 or more may be required to inhibit, interfere with or modulate the biological activity of IL-18 receptor when the IL-18 concentration is

from about  $EC_{50}$  to about  $EC_{90}$ . Ratios of IL-18 receptor antigen binding protein between these values are also possible.

As a general structure, the antigen binding proteins of the invention comprise (a) a scaffold, and (b) one or a plurality of CDRs, regions that are determinative to antigen binding specificity and affinity. A “complementary determining region” or “CDR,” as used herein, refers to a binding protein region that constitutes the major surface contact points for antigen binding. One or more CDRs are embedded in the scaffold structure of the antigen binding protein. The scaffold structure of the antigen binding proteins may be the framework of an antibody, or fragment or variant thereof, or may be completely synthetic in nature. The various scaffold structures of antigen binding proteins are further described hereinbelow.

#### 1. CDRs

An antigen binding protein may have six CDRs (as typically does each “arm” of a naturally occurring antibody), for example one heavy chain CDR1 (“CDRH1”), one heavy chain CDR2 (“CDRH2”), one heavy chain CDR3 (“CDRH3”), one light chain CDR1 (“CDRL1”), one light chain CDR2 (“CDRL2”), one light chain CDR3 (“CDRL3”). The term “naturally occurring” as used throughout the specification in connection with biological materials such as polypeptides, nucleic acids, host cells, and the like, refers to materials which are found in nature. In naturally occurring antibodies, a CDRH1 typically comprises about five (5) to about seven (7) amino acids, CDRH2 typically comprises about sixteen (16) to about nineteen (19) amino acids, and CDRH3 typically comprises about three (3) to about twenty five (25) amino acids. CDRL1 typically comprises about ten (10) to about seventeen (17) amino acids, CDRL2 typically comprises about seven (7) amino acids, and CDRL3 typically comprises about seven (7) to about ten (10) amino acids. Preferred CDRs are depicted in FIGS. 2A-2E, 3A, and 3B.

The structure and properties of CDRs within a naturally occurring antibody are described further in this Section hereinbelow. Briefly, in a traditional antibody scaffold, the CDRs are embedded within a framework in the heavy and light chain variable region where they constitute the regions responsible for antigen binding and recognition. A variable region comprises at least three heavy or light chain CDRs, see, supra (Kabat et al., 1991, *Sequences of Proteins of Immunological Interest*, Public Health Service N.I.H., Bethesda, Md.; see also Chothia and Lesk, 1987, *J. Mol. Biol.* 196:901-917; Chothia et al., 1989, *Nature* 342: 877-883), within a framework region (designated framework regions 1-4, FR1, FR2, FR3, and FR4, by Kabat et al., 1991, supra; see also Chothia and Lesk, 1987, supra). See, infra. The CDRs provided by the present invention, however, may not only be used to define the antigen binding domain of a traditional antibody structure, but may be embedded in a variety of other scaffold structures, as described herein.

In certain embodiments, one or more CDRs of an antigen binding protein are each independently selected from the CDRH regions of any of SEQ ID NOs:89-139 and the CDRL regions of any of SEQ ID NOs:140-190. Thus, in one embodiment, the invention provides an antigen binding protein that binds an  $\alpha$ - or  $\beta$ -IL-18 receptor, wherein said antigen binding protein comprises (a) a scaffold structure (as described below); and (b) at least one CDR selected from the CDRH regions of any of SEQ ID NOs:89-139 and the CDRL regions of any of SEQ ID NOs:140-190. In this embodiment, of particular use are antigen binding proteins with a CDRH3 or CDRL3 region. Additional embodiments utilize antigen binding proteins with one CDR selected from the CDRH



regions of any of SEQ ID NOs:89-139 and a CDRL region of any of SEQ ID NOs:140-190 (e.g., the antigen binding protein has two CDR regions, one CDRH and one CDHL, a specific embodiment are antigen binding proteins with both a CDRH3 and a CDRL3 region).

As will be appreciated by those in the art, particularly useful embodiments may contain one, two, three, four, five or six of independently selected CDRs of SEQ ID NOs:89-190. However, as will be appreciated by those in the art, specific embodiments generally utilize combinations of CDRs that are non-repetitive, e.g., antigen binding proteins are generally not made with two CDRH2 regions, etc.

In some embodiments, antigen binding proteins are generated that comprise a CDRH3 region and a CDRL3 region, particularly with the CDRH3 region being selected from a CDRH3 region of any of SEQ ID NOs:91, 94, 97, 100, 103, 106, 109, 112, 115, 118, 121, 124, 127, 130, 133, 136, 139 and the CDRL3 region being selected from a CDRL3 region of any of SEQ ID NOs:142, 145, 148, 151, 154, 157, 160, 163, 166, 169, 172, 175, 178, 181, 184, 187, 190. Particular combinations are depicted in FIG. 4.

In additional embodiments, antigen binding proteins are utilized that comprise a CDRH1, a CDRH2, and a CDRH3 region independently selected from SEQ ID NOs:89-139. In more specific embodiments, of particular use may be antigen binding proteins of this type that have all three CDRH regions selected from the same variable region of any of SEQ ID NOs:1-17.

In further embodiments, antigen binding proteins are utilized that comprise a CDRL1, a CDRL2, and a CDRL3 region independently selected from SEQ ID NOs:140-190. In more specific embodiments, of particular use are antigen binding proteins of this type that have all three CDRL regions selected from the same variable region of any of SEQ ID NOs:18-34.

In an additional embodiment, the antigen binding protein comprises a CDRH1, CDRH2, and CDRH3 region independently selected from SEQ ID NOs:89-139, again, in one embodiment with all three regions selected from the same SEQ ID NO, and a CDRL1, CDRL2, and CDRL3 region independently selected from SEQ ID NOs:140-190, again, in one embodiment with all three regions selected from the same variable region of any of SEQ ID NOs:1-34.

In yet another aspect of the invention provides for an antigen binding protein that binds the  $\alpha$ - or  $\beta$ IL-18 receptor where the isolated antigen binding protein comprises a heavy chain amino acid sequence that comprises a CDRH1, a CDRH2, or a CDRH3, each selected from any of SEQ ID NOs:89-139, or a fragment thereof, or a light chain amino acid sequence that comprises a CDRL1, a CDRL2, or a CDRL3, each selected from any of SEQ ID NOs:140-190, or a fragment thereof. A heavy or light chain variable region "fragment," as used herein includes at least one CDR and at least a portion of a framework region of an antibody framework of SEQ ID NOs:1-34, said portion comprising at least one amino acid.

In yet another aspect, the invention provides for an antigen binding protein that binds an  $\alpha$ - or  $\beta$ IL-18 receptor where the isolated antigen binding protein comprises a heavy chain amino acid sequence that comprises a CDRH1, a CDRH2, and a CDRH3, each independently selected from any of SEQ ID NOs:89-139, or a light chain amino acid sequence that comprises a CDRL1, a CDRL2, and a CDRL3, each independently selected from any of SEQ ID NOs:140-190. In a specific embodiment, the CDRs are from the same contiguous heavy chain amino acid sequence of SEQ ID NOs:1-17 or from the same contiguous light chain amino acid sequence of SEQ ID NOs:18-34.

A further aspect of the invention provides for an isolated antigen binding protein that binds an  $\alpha$ - or  $\beta$ IL-18 receptor where the isolated antigen binding protein comprises a heavy chain amino acid sequence that comprises a CDRH1, a CDRH2, and a CDRH3, each independently selected from any of SEQ ID NOs:89-139, and a light chain amino acid sequence that comprises a CDRL1, a CDRL2, and a CDRL3, each independently selected from any of SEQ ID NOs:140-190. In a specific embodiment, the heavy chain CDRs are from the same contiguous heavy chain amino acid sequence of SEQ ID NO:1-17 and the light chain CDRs are from the same contiguous light chain amino acid sequence of SEQ ID NO:18-34.

An additional aspect of the invention provides for an isolated antigen binding protein that binds an  $\alpha$ - or  $\beta$ IL-18 receptor where the isolated antigen binding protein comprises a heavy chain amino acid sequence of any of SEQ ID NOs:1-17, or a light chain amino acid sequence of any of SEQ ID NOs:18-34.

A further aspect of the invention provides for an isolated antigen binding protein that binds an  $\alpha$ - or  $\beta$ IL-18 receptor where the isolated antigen binding protein comprises a heavy chain amino acid sequence of any of SEQ ID NOs:1-17, and a light chain amino acid sequence of any of SEQ ID NOs:18-34. It is noted that the any the heavy chain sequences of SEQ ID NOs:1-17 can be mixed and matched with any of the light chain sequences of SEQ ID NOs:18-34. The resulting possible combinations are depicted in FIG. 4. Shown are dimers of a combination of each one heavy and one light chain variable region. As most antibodies are tetramers, an antigen binding protein of the invention may comprise any combination of any two of the depicted dimers thus including both hetero- and homo-tetramers, with homo-tetramers (e.g., two identical dimers) being specific.

In again a further aspect the antigen binding protein of the invention comprises any of the sequences depicted in SEQ ID NOs:73-88.

TABLE 1 provides a brief description of the sequences as they relate to their sequence identification numbers. The CDRs within the variable regions of the invention are identified in FIGS. 2A-2E, 3A and 3B.

TABLE 1

Brief Description Of Sequence Listings	
Brief Description	Sequence Identification Number
Amino acid sequence encoding the heavy chain variable region AM <sub>H</sub> 1	SEQ ID NO: 1
Amino acid sequence encoding the heavy chain variable region AM <sub>H</sub> 2	SEQ ID NO: 2
Amino acid sequence encoding the heavy chain variable region AM <sub>H</sub> 3	SEQ ID NO: 3
Amino acid sequence encoding the heavy chain variable region AM <sub>H</sub> 4	SEQ ID NO: 4
Amino acid sequence encoding the heavy chain variable region AM <sub>H</sub> 5	SEQ ID NO: 5



TABLE 1-continued

Brief Description Of Sequence Listings	
Brief Description	Sequence Identification Number
Amino acid sequence of complete heavy chain of AM <sub>H</sub> 9	SEQ ID NO: 81
Nucleotide sequence of complete heavy chain of AM <sub>H</sub> 9	SEQ ID NO: 82
Amino acid sequence of complete light chain of AM <sub>L</sub> 9	SEQ ID NO: 83
Nucleotide sequence of complete light chain of AM <sub>L</sub> 9	SEQ ID NO: 84
Amino acid sequence of complete heavy chain of AM <sub>H</sub> 11	SEQ ID NO: 85
Nucleotide sequence of complete heavy chain of AM <sub>H</sub> 11	SEQ ID NO: 86
Amino acid sequence of complete light chain of AM <sub>L</sub> 7	SEQ ID NO: 87
Nucleotide sequence of complete light chain of AM <sub>L</sub> 7	SEQ ID NO: 88
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 1	SEQ ID NO: 89, 90, 91
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 2	SEQ ID NO: 92, 93, 94
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 3	SEQ ID NO: 95, 96, 97
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 4	SEQ ID NO: 98, 99, 100
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 5	SEQ ID NO: 101, 102, 103
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 6	SEQ ID NO: 104, 105, 106
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 7	SEQ ID NO: 107, 108, 109
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 8	SEQ ID NO: 110, 111, 112
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 9	SEQ ID NO: 113, 114, 115
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 10	SEQ ID NO: 116, 117, 118
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 11	SEQ ID NO: 119, 120, 121
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 12	SEQ ID NO: 122, 123, 124
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 13	SEQ ID NO: 125, 126, 127
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 14	SEQ ID NO: 128, 129, 130
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 15	SEQ ID NO: 131, 132, 133
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 16	SEQ ID NO: 134, 135, 136
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 17	SEQ ID NO: 137, 138, 139
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 1	SEQ ID NO: 140, 141, 142
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 2	SEQ ID NO: 143, 144, 145
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 3	SEQ ID NO: 146, 147, 148
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 4	SEQ ID NO: 149, 150, 151
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 5	SEQ ID NO: 152, 153, 154
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 6	SEQ ID NO: 155, 156, 157
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 7	SEQ ID NO: 158, 159, 160
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 8	SEQ ID NO: 161, 162, 163
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 9	SEQ ID NO: 164, 165, 166
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 10	SEQ ID NO: 167, 168, 169
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 11	SEQ ID NO: 170, 171, 172
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 12	SEQ ID NO: 173, 174, 175
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 13	SEQ ID NO: 176, 177, 178
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 14	SEQ ID NO: 179, 180, 181
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 15	SEQ ID NO: 182, 183, 184
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 16	SEQ ID NO: 185, 186, 187

TABLE 1-continued

Brief Description Of Sequence Listings	
Brief Description	Sequence Identification Number
Amino acid sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 17	SEQ ID NO: 188, 189, 190
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 1	SEQ ID NO: 191, 192, 193
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 2	SEQ ID NO: 194, 195, 196
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 3	SEQ ID NO: 197, 198, 199
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 4	SEQ ID NO: 200, 201, 202
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 5	SEQ ID NO: 203, 204, 205
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 6	SEQ ID NO: 206, 207, 208
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 7	SEQ ID NO: 209, 210, 211
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 8	SEQ ID NO: 212, 213, 214
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 9	SEQ ID NO: 215, 216, 217
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 10	SEQ ID NO: 218, 219, 220
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 11	SEQ ID NO: 221, 222, 223
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 12	SEQ ID NO: 224, 225, 226
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 13	SEQ ID NO: 227, 228, 229
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 14	SEQ ID NO: 230, 231, 232
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 15	SEQ ID NO: 233, 234, 235
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 16	SEQ ID NO: 236, 237, 238
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of heavy chain variable region AM <sub>H</sub> 17	SEQ ID NO: 239, 240, 241
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 1	SEQ ID NO: 242, 243, 244
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 2	SEQ ID NO: 245, 246, 247
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 3	SEQ ID NO: 248, 249, 250
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 4	SEQ ID NO: 251, 252, 253
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 5	SEQ ID NO: 254, 255, 256
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 6	SEQ ID NO: 257, 258, 259
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 7	SEQ ID NO: 260, 261, 262
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 8	SEQ ID NO: 263, 264, 265
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 9	SEQ ID NO: 266, 267, 268
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 10	SEQ ID NO: 269, 270, 271
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 11	SEQ ID NO: 272, 273, 274
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 12	SEQ ID NO: 275, 276, 277
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 13	SEQ ID NO: 278, 279, 280
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 14	SEQ ID NO: 281, 282, 283
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 15	SEQ ID NO: 284, 285, 286
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 16	SEQ ID NO: 287, 288, 289
Nucleotide sequence encoding CDR1, CDR 2, CDR 3, respectively, of light chain variable region AM <sub>L</sub> 17	SEQ ID NO: 290, 291, 292

## 2. Scaffolds

As noted herein, the antigen binding proteins can comprise a scaffold structure into which the CDR(s) are grafted. The scaffold structure may be based on antibodies (including, but not limited to, monoclonal antibodies, human antibodies, murine antibodies, chimeric antibodies, humanized antibodies, bispecific antibodies), antibody fragments (such as minibodies, domain antibodies, F(ab) fragments, F(ab') fragments, F(ab)<sub>2</sub> fragments, F(ab')<sub>2</sub> fragments, Fv fragments, scFv fragments, Fd fragments), synthetic antibodies (sometimes referred to herein as "antibody mimetics"), antibody fusions (sometimes referred to as "antibody conjugates"), including Fc fusions. Some embodiments include the use of human scaffold components. The invention as such at least encompasses any of the below described scaffolds comprising one or several of the CDRs as identified in SEQ ID NOs:89-190, preferably of SEQ ID NOs:89-189, that can bind to and/or inhibit the biological activity of IL-18 receptor. In some embodiments, the scaffold comprises one or several heavy chain variable regions as identified in SEQ ID NOs:1-17, and or one or several light chain variable regions as identified in any of SEQ ID NOs:18-34. In some embodiments, the scaffold comprises an IgG chain as identified in any of SEQ ID NOs:77-88.

In one embodiment, the scaffold into which one or several CDRs are grafted is an antibody. As used herein, the term "antibody" refers to a multimeric protein having a traditional antibody structure, comprising at least two, more typically four polypeptide chains. An antibody binds specifically to an antigen and may be able to inhibit or modulate the biological activity of the antigen. In certain embodiments, antibodies are produced by recombinant DNA techniques. In additional embodiments, antibodies are produced by enzymatic or chemical cleavage of naturally occurring antibodies.

Traditional antibody structural units typically comprise a tetramer. Each tetramer is typically composed of two identical pairs of polypeptide chains, each pair having one "light" (typically having a molecular weight of about 25 kDa) and one "heavy" chain (typically having a molecular weight of about 50-70 kDa). The amino-terminal portion of each chain includes a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The carboxy-terminal portion of each chain defines a constant region primarily responsible for effector function. Human light chains are classified as kappa and lambda light chains. Heavy chains are classified as mu, delta, gamma, alpha, or epsilon, and define the antibody's isotype as IgM, IgD, IgG, IgA, and IgE, respectively. IgG has several subclasses, including, but not limited to IgG1, IgG2, IgG3, and IgG4. IgM has subclasses, including, but not limited to, IgM1 and IgM2.

Within light and heavy chains, the variable and constant regions are joined by a "J" region of about twelve (12) or more amino acids, with the heavy chain also including a "D" region of about ten (10) more amino acids. See, generally, Paul, W., ed., 1989, *Fundamental Immunology* Ch. 7, 2nd ed. Raven Press, N.Y. The variable regions of each light/heavy chain pair form the antibody binding site.

Some naturally occurring antibodies, for example found in camels and llamas, are dimers consisting of two heavy chain and include no light chains. Muldermans et al., 2001, *J. Biotechnol.* 74:277-302; Desmyter et al., 2001, *J. Biol. Chem.* 276:26285-26290. Crystallographic studies of a camel antibody have revealed that the CDR3 regions form a surface that interacts with the antigen and thus is critical for antigen binding like in the more typical tetrameric antibodies.

The variable regions of the heavy and light chains typically exhibit the same general structure of relatively conserved

framework regions (FR) joined by three hypervariable regions, also called complementarity determining regions or CDRs. The CDRs are the hypervariable regions of an antibody (or antigen binding protein, as outlined herein), which are responsible for antigen recognition and binding. The CDRs from the two chains of each pair are aligned by the framework regions, enabling binding to a specific epitope. From N-terminal to C-terminal, both light and heavy chains comprise the domains FR1, CDR1, FR2, CDR2, FR3, CDR3 and FR4. The assignment of amino acids to each domain is in accordance with the definitions of Kabat Sequences of Proteins of Immunological Interest. Chothia et al., 1987, *J. Mol. Biol.* 196:901-917; Chothia et al., 1989, *Nature* 342:878-883.

CDRs constitute the major surface contact points for antigen binding. See, e.g., Chothia and Lesk, 1987, *J. Mol. Biol.* 196:901-917. Further, CDR3 of the light chain and, especially, CDR3 of the heavy chain may constitute the most important determinants in antigen binding within the light and heavy chain variable regions. See, e.g., Chothia and Lesk, 1987, *supra*; Desiderio et al., 2001, *J. Mol. Biol.* 310:603-615; Xu and Davis, 2000, *Immunity* 13:37-45; Desmyter et al., 2001, *J. Biol. Chem.* 276:26285-26290; and Muyldermans, 2001, *J. Biotechnol.* 74:277-302. In some antibodies, the heavy chain CDR3 appears to constitute the major area of contact between the antigen and the antibody. Desmyter et al., 2001, *supra*. In vitro selection schemes in which CDR3 alone is varied can be used to vary the binding properties of an antibody. Muyldermans, 2001, *supra*; Desiderio et al., 2001, *supra*.

Naturally occurring antibody chains typically include a signal sequence, which directs the antibody chain into the cellular pathway for protein secretion and which is not present in the mature antibody. A polynucleotide encoding an antibody chain may encode a naturally occurring signal sequence or a heterologous signal sequence as described below.

In one embodiment, the antigen binding protein is a monoclonal antibody, with from one (1) to six (6) of the depicted CDRs of any of SEQ ID NOs:89-190, as outlined herein. The antibodies of the invention may be of any type including IgM, IgG (including IgG1, IgG2, IgG3, IgG4), IgD, IgA, or IgE antibody. In specific embodiment, the antigen binding protein is an IgG type antibody. In an even more specific embodiment, the antigen binding protein is an IgG2 type antibody.

In some embodiments, for example when the antigen binding protein is an antibody with complete heavy and light chains, the CDRs are all from the same species, e.g., human. In some embodiments, however, the scaffold components can be a mixture from different species. As such, if the antigen binding protein is an antibody, such antibody may be a chimeric antibody and/or a humanized antibody. In general, both "chimeric antibodies" refer to antibodies that combine regions from more than one species. For example, "chimeric antibodies" traditionally comprise variable region(s) from a mouse (or rat, in some cases) and the constant region(s) from a human.

For example in embodiments wherein the antigen binding protein contains less than six CDRs from the sequences outlined above, additional CDRs may be either from other species (e.g., murine CDRs), or may be different human CDRs than those depicted in the sequences. For example, human CDRH3 and CDRL3 regions from the appropriate sequences identified herein may be used, with CDRH1, CDRH2, CDRL1 and CDRL2 being optionally selected from alternate species, or different human antibody sequences, or combina-

tions thereof. For example, the CDRs of the invention can replace the CDR regions of commercially relevant chimeric or humanized antibodies.

Specific embodiments of the invention utilize scaffold components of the antigen binding proteins that are human components.

“Humanized antibodies” generally refer to non-human antibodies that have had the variable-domain framework regions swapped for sequences found in human antibodies. Generally, in a humanized antibody, the entire antibody, except the CDRs, is encoded by a polynucleotide of human origin or is identical to such an antibody except within its CDRs. The CDRs, some or all of which are encoded by nucleic acids originating in a non-human organism, are grafted into the beta-sheet framework of a human antibody variable region to create an antibody, the specificity of which is determined by the engrafted CDRs. The creation of such antibodies is described in, e.g., WO 92/11018, Jones, 1986, *Nature* 321:522-525, Verhoeyen et al., 1988, *Science* 239: 1534-1536. Humanized antibodies can also be generated using mice with a genetically engineered immune system. Roque et al., 2004, *Biotechnol. Prog.* 20:639-654. In the present invention, the identified CDRs are human, and thus both humanized and chimeric antibodies in this context include some non-human CDRs; for example, humanized antibodies may be generated that comprise the CDRH3 and CDRL3 regions, with one or more of the other CDR regions being of a different special origin.

In one embodiment, the IL-18 antigen binding protein is a multispecific antibody, and notably a bispecific antibody, also sometimes referred to as “diabodies”. These are antibodies that bind to two (or more) different antigens. Diabodies can be manufactured in a variety of ways known in the art (Holliger and Winter, 1993, *Current Opinion Biotechnol.* 4:446-449), e.g., prepared chemically or from hybrid hybridomas.

In one embodiment, the IL-18 antigen binding protein is a fully human antibody, i.e., an antibody fully composed of human components. In this embodiment, as outlined above, specific structures comprise complete heavy and light chains depicted comprising the CDR regions depicted in FIGS. 2A-2E, 3A and 3B. Additional embodiments utilize one or more of the CDRs of the invention, with the other CDRs, framework regions, J and D regions, constant regions, etc., coming from other human antibodies. For example, the CDRs of the invention can replace the CDRs of any number of human antibodies, particularly commercially relevant antibodies.

In one embodiment, the IL-18 antigen binding protein is an antibody fragment, that is a fragment of any of the antibodies outlined herein that retain binding specificity to an  $\alpha$ - or  $\beta$ -18 receptor.

Specific antibody fragments include, but are not limited to, (i) the Fab fragment consisting of VL, VH, CL and CH1 domains, (ii) the Fab' fragment consisting of VL, VH, CL and CH1 domains plus the heavy chain hinge region; (iii) the Fd fragment consisting of the VH and CH1 domains, (iii) the Fv fragment consisting of the VL and VH domains of a single antibody; (iv) the dAb fragment (Ward et al., 1989, *Nature* 341:544-546) which consists of a single variable, (v) isolated CDR regions, (vi) F(ab')<sub>2</sub> fragments, a bivalent fragment comprising two linked Fab' fragments (vii) single chain Fv molecules (scFv), wherein a VH domain and a VL domain are linked by a peptide linker which allows the two domains to associate to form an antigen binding site (Bird et al., 1988, *Science* 242:423-426, Huston et al., 1988, *Proc. Natl. Acad. Sci. U.S.A.* 85:5879-5883), (viii) bispecific single chain Fv dimers (PCT/US92/09965) and (ix) “diabodies” or “triabod-

ies”, multivalent or multispecific fragments constructed by gene fusion (Tomlinson et al., 2000, *Methods Enzymol.* 326: 461-479; WO94/13804; Holliger et al., 1993, *Proc. Natl. Acad. Sci. U.S.A.* 90:6444-6448). The antibody fragments may be modified. For example, the molecules may be stabilized by the incorporation of disulphide bridges linking the VH and VL domains (Reiter et al., 1996, *Nature Biotech.* 14:1239-1245). Again, as outlined herein, the non-CDR components of these fragments are preferably human sequences. In one embodiment, the IL-18 antigen binding protein is a minibody. Minibodies are minimized antibody-like proteins comprising a scFv joined to a CH3 domain. Hu et al., 1996, *Cancer Res.* 56:3055-3061.

In one embodiment, the IL-18 antigen binding protein is a domain antibody; see for example U.S. Pat. No. 6,248,516. Domain antibodies (dAbs) are functional binding domains of antibodies, corresponding to the variable regions of either the heavy (VH) or light (VL) chains of human antibodies dAbs have a molecular weight of approximately 13 kDa, or less than one-tenth the size of a full antibody. dAbs are well expressed in a variety of hosts including bacterial, yeast, and mammalian cell systems. In addition, dAbs are highly stable and retain activity even after being subjected to harsh conditions, such as freeze-drying or heat denaturation. See, for example, U.S. Pat. Nos. 6,291,158; 6,582,915; 6,593,081; 6,172,197; US Serial No. 2004/0110941; European Patent 0368684; U.S. Pat. No. 6,696,245, WO04/058821, WO04/003019 and WO03/002609.

In one embodiment, the IL-18 antigen binding protein is an antibody fusion protein or an antibody fragment fusion, such as an Fc fusion (sometimes collectively referred to herein as an “antibody conjugate”). The conjugate partner can be proteinaceous or non-proteinaceous; the latter generally being generated using functional groups on the antigen binding protein (see the discussion on covalent modifications of the antigen binding proteins) and on the conjugate partner. For example linkers are known in the art; for example, homo- or hetero-bifunctional linkers as are well known (see, 1994 Pierce Chemical Company catalog, technical section on cross-linkers, pages 155-200, incorporated herein by reference).

Suitable conjugates include, but are not limited to, labels as described below, drugs and cytotoxic agents including, but not limited to, cytotoxic drugs (e.g., chemotherapeutic agents) or toxins or active fragments of such toxins. Suitable toxins and their corresponding fragments include diphtheria A chain, exotoxin A chain, ricin A chain, abrin A chain, curcin, crotin, phenomycin, enomycin and the like. Cytotoxic agents also include radiochemicals made by conjugating radioisotopes to antigen binding proteins, or binding of a radionuclide to a chelating agent that has been covalently attached to the antigen binding protein. Additional embodiments utilize calicheamicin, auristatins, geldanamycin and maytansine.

In one embodiment, the IL-18 antigen binding protein is an antibody analog, sometimes referred to as “synthetic antibodies.” For example, a variety of recent work utilizes either alternative protein scaffolds or artificial scaffolds with grafted CDRs. Such scaffolds include, but are not limited to, mutations introduced to stabilize the three-dimensional structure of the binding protein as well as wholly synthetic scaffolds consisting for example of biocompatible polymers. See, for example, Korndorfer et al., 2003, *Proteins: Structure, Function, and Bioinformatics*, Volume 53, Issue 1:121-129. Roque et al., 2004, *Biotechnol. Prog.* 20:639-654. In addition, peptide antibody mimetics (“PAMs”) can be used, as well as scaffolds based on antibody mimetics utilizing fibronectin components as a scaffold. Alternate scaffolds that may be

used to produce an IL-18 antigen binding protein are reviewed in Hey et al., 2005, *Trends Biotechnol.* 23:514-22 and Binz et al., *Nature Biotechnology* 23:1257-68 (both incorporated herein by reference in their entirety).

### 3. CDR Variants

Also included within the invention are variants of the CDRH1, CDRH2, CDRH3, CDRL1, CDRL2 and CDRL3 amino acid sequences depicted in SEQ ID NOs:89-190. Thus variant CDRs are included within the definition of CDR as used herein. These variants fall into one or more of three classes: substitutional, insertional or deletional variants, with the former being specific.

As it is known in the art, a number of different programs can be used to identify the degree of sequence identity or similarity a protein or nucleic acid has to a known sequence.

For amino acid sequences, sequence identity and/or similarity is determined by using standard techniques known in the art, including, but not limited to, the local sequence identity algorithm of Smith and Waterman, 1981, *Adv. Appl. Math.* 2:482, the sequence identity alignment algorithm of Needleman and Wunsch, 1970, *J. Mol. Biol.* 48:443, the search for similarity method of Pearson and Lipman, 1988, *Proc. Nat. Acad. Sci. U.S.A.* 85:2444, computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Drive, Madison, Wis.), the Best Fit sequence program described by Devereux et al., 1984, *Nucl. Acid Res.* 12:387-395, preferably using the default settings, or by inspection. Preferably, percent identity is calculated by FastDB based upon the following parameters: mismatch penalty of 1; gap penalty of 1; gap size penalty of 0.33; and joining penalty of 30, "Current Methods in Sequence Comparison and Analysis," Macromolecule Sequencing and Synthesis, Selected Methods and Applications, pp 127-149 (1988), Alan R. Liss, Inc.

An example of a useful algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments. It can also plot a tree showing the clustering relationships used to create the alignment. PILEUP uses a simplification of the progressive alignment method of Feng & Doolittle, 1987, *J. Mol. Evol.* 35:351-360; the method is similar to that described by Higgins and Sharp, 1989, *CABIOS* 5:151-153. Useful PILEUP parameters including a default gap weight of 3.00, a default gap length weight of 0.10, and weighted end gaps.

Another example of a useful algorithm is the BLAST algorithm, described in: Altschul et al., 1990, *J. Mol. Biol.* 215:403-410; Altschul et al., 1997, *Nucleic Acids Res.* 25:3389-3402; and Karin et al., 1993, *Proc. Natl. Acad. Sci. U.S.A.* 90:5873-5787. A particularly useful BLAST program is the WU-BLAST-2 program which was obtained from Altschul et al., 1996, *Methods in Enzymology* 266:460-480. WU-BLAST-2 uses several search parameters, most of which are set to the default values. The adjustable parameters are set with the following values: overlap span=1, overlap fraction=0.125, word threshold (T)=II. The HSP S and HSP S2 parameters are dynamic values and are established by the program itself depending upon the composition of the particular sequence and composition of the particular database against which the sequence of interest is being searched; however, the values may be adjusted to increase sensitivity.

An additional useful algorithm is gapped BLAST as reported by Altschul et al., 1993, *Nucl. Acids Res.* 25:3389-3402. Gapped BLAST uses BLOSUM-62 substitution scores; threshold T parameter set to 9; the two-hit method to trigger ungapped extensions, charges gap lengths of k a cost of 10+k;  $X_{10}$  set to 16, and  $X_g$  set to 40 for database search

stage and to 67 for the output stage of the algorithms. Gapped alignments are triggered by a score corresponding to about 22 bits.

Generally, the amino acid homology, similarity, or identity between individual variant CDRs are at least 80% to the sequences depicted herein, and more typically with preferably increasing homologies or identities of at least 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, and almost 100%.

In a similar manner, "percent (%) nucleic acid sequence identity" with respect to the nucleic acid sequence of the binding proteins identified herein is defined as the percentage of nucleotide residues in a candidate sequence that are identical with the nucleotide residues in the coding sequence of the antigen binding protein. A specific method utilizes the BLASTN module of WU-BLAST-2 set to the default parameters, with overlap span and overlap fraction set to 1 and 0.125, respectively.

Generally, the nucleic acid sequence homology, similarity, or identity between the nucleotide sequences encoding individual variant CDRs and the nucleotide sequences depicted herein are at least 60%, and more typically with preferably increasing homologies or identities of at least 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, and almost 100%.

Homology between nucleotide sequences is often defined by their ability to hybridize to each other. The term "selectively hybridize" referred to herein means to detectably and specifically bind. Polynucleotides, oligonucleotides and fragments thereof in accordance with the invention selectively hybridize to nucleic acid strands under hybridization and wash conditions that minimize appreciable amounts of detectable binding to nonspecific nucleic acids. High stringency conditions can be used to achieve selective hybridization conditions as known in the art and discussed herein.

High stringency conditions are known in the art; see, for example Sambrook et al., 2001, supra, and *Short Protocols in Molecular Biology*, Second Edition, Ausubel et al. eds., John Wiley & Sons, 1992, both of which are hereby incorporated by reference. Stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures. An extensive guide to the hybridization of nucleic acids is found in Tijssen, *Techniques In Biochemistry and Molecular Biology—Hybridization with Nucleic Acid Probes*, "Overview of principles of hybridization and the strategy of nucleic acid assays" (1993).

Generally, stringent conditions are selected to be about 5-10° C. lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength and pH. The  $T_m$  is the temperature (under defined ionic strength, pH and nucleic acid concentration) at which 50% of the probes complementary to the target hybridize to the target sequence at equilibrium (as the target sequences are present in excess, at  $T_m$ , 50% of the probes are occupied at equilibrium). Stringent conditions will be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30° C. for short probes (e.g., 10 to 50 nucleotides) and at least about 60° C. for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide.

In another embodiment, less stringent hybridization conditions are used; for example, moderate or low stringency

conditions may be used, as are known in the art; see, Sambrook et al., 2001, supra; Ausubel et al., 1992, supra, and Tijssen, 1993, supra.

The variants according to the invention are ordinarily prepared by site specific mutagenesis of nucleotides in the DNA encoding the antigen binding protein, using cassette or PCR mutagenesis or other techniques well known in the art, to produce DNA encoding the variant, and thereafter expressing the recombinant DNA in cell culture as outlined herein. However, antigen binding protein fragments comprising variant CDRs having up to about 100-150 residues may be prepared by in vitro synthesis using established techniques. The variants typically exhibit the same qualitative biological activity as the naturally occurring analogue, e.g., binding to IL-18 receptor and inhibiting signaling, although variants can also be selected which have modified characteristics as will be more fully outlined below.

Thus, a "variant CDR" is one with the specified homology, similarity, or identity to the parent CDR of the invention, and shares biological function, including, but not limited to, at least 90, 91, 92, 93, 94, 95, 96, 97, 98% or 99% of the specificity and/or activity of the parent CDR. For example, the variants typically will bind to the same IL-18 receptor epitope outlined below, with a similar inhibition of IL-18 receptor signaling.

While the site or region for introducing an amino acid sequence variation is predetermined, the mutation per se need not be predetermined. For example, in order to optimize the performance of a mutation at a given site, random mutagenesis may be conducted at the target codon or region and the expressed antigen binding protein CDR variants screened for the optimal combination of desired activity. Techniques for making substitution mutations at predetermined sites in DNA having a known sequence are well known, for example, M13 primer mutagenesis and PCR mutagenesis. Screening of the mutants is done using assays of antigen binding protein activities, such as IL-18 receptor binding.

Amino acid substitutions are typically of single residues; insertions usually will be on the order of from about one (1) to about twenty (20) amino acid residues, although considerably larger insertions may be tolerated. Deletions range from about one (1) to about twenty (20) amino acid residues, although in some cases deletions may be much larger.

Substitutions, deletions, insertions or any combination thereof may be used to arrive at a final derivative or variant. Generally these changes are done on a few amino acids to minimize the alteration of the molecule, particularly the immunogenicity and specificity of the antigen binding protein. However, larger changes may be tolerated in certain circumstances. When small alterations in the characteristics of the CDR of the antigen binding protein are desired, substitutions are generally made in accordance with the following chart depicted as TABLE 2.

TABLE 2

Original Residue	Exemplary Substitutions
Ala	Ser
Arg	Lys
Asn	Gln, His
Asp	Glu
Cys	Ser
Gln	Asn
Glu	Asp
Gly	Pro
His	Asn, Gln
Ile	Leu, Val

TABLE 2-continued

Original Residue	Exemplary Substitutions
Leu	Ile, Val
Lys	Arg, Gln, Glu
Met	Leu, Ile
Phe	Met, Leu, Tyr
Ser	Thr
Thr	Ser
Trp	Tyr
Tyr	Trp, Phe
Val	Ile, Leu

Substantial changes in function or immunological identity are made by selecting substitutions that are less conservative than those shown in TABLE 2. For example, substitutions may be made which more significantly affect: the structure of the polypeptide backbone in the area of the alteration, for example the alpha-helical or beta-sheet structure; the charge or hydrophobicity of the molecule at the target site; or the bulk of the side chain. The substitutions which in general are expected to produce the greatest changes in the polypeptide's properties are those in which (a) a hydrophilic residue, e.g., seryl or threonyl, is substituted for (or by) a hydrophobic residue, e.g., leucyl, isoleucyl, phenylalanyl, valyl or alanyl; (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue having an electropositive side chain, e.g., lysyl, arginyl, or histidyl, is substituted for (or by) an electronegative residue, e.g., glutamyl or aspartyl; or (d) a residue having a bulky side chain, e.g., phenylalanine, is substituted for (or by) one not having a side chain, e.g., glycine.

The variants typically exhibit the same qualitative biological activity and will elicit the same immune response as the naturally-occurring analogue, although variants also are selected to modify the characteristics of the antigen binding protein proteins as needed. Alternatively, the variant may be designed such that the biological activity of the antigen binding protein is altered. For example, glycosylation sites may be altered or removed as discussed herein.

#### 4. VH And VL Variants

As outlined above, in some embodiments the invention provides antigen binding proteins comprising, or consisting of, a heavy chain variable region of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, or 17 and/or a light chain variable region of SEQ ID NO:18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, or 34, respectively, or fragments thereof as defined above. Thus, in those embodiments, the antigen binding protein comprises not only at least one CDR or variant thereof depicted in SEQ ID NOs:1-34, but also at least part of a depicted framework sequence. In addition, the invention encompasses variants of such heavy chain variable sequences or light chain variable sequences.

A "variant VH" or "variant heavy chain variable region," and a "variant VL" or "variant light chain variable region" generally shares an amino acid homology, similarity, or identity of at least 80% with those depicted herein, and more typically with preferably increasing homologies or identities of at least 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% and almost 100%. The nucleic acid sequence homology, similarity, or identity between the nucleotide sequences encoding individual variant VHs and VLs and the nucleic acid sequences depicted herein are at least 60% with those depicted herein, and more typically with preferably increasing homologies or identities of at least 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% and almost 100%. In addition, a "variant



VH” or “variant heavy chain variable region,” and a “variant VL” or “variant light chain variable region” typically shares the biological function, including, but not limited to, at least 90, 91, 92, 93, 94, 95, 96, 97, 98% or 99% of the specificity and/or activity of the parent CDR. For example, the variants typically will bind to the same IL-18 receptor epitopes outlined below, with a similar inhibition of IL-18 receptor signaling.

Methods of generating variants, as well as methods of determining sequence homology, similarity, and identity, are outlined supra, see Section V.B.1.

In some embodiments, constant region variants may also be included. Preferred constant region variants include those that alter a biological function of the antibody containing the variation. For example, the antibody may contain a variation that alters the antibody’s ability to activate complement or induce antibody-dependent cellular cytotoxicity (ADCC). Such variants may include those that result in an alteration of the glycosylation of the antibody.

#### C. IL-18 Receptor and IL-18 Receptor Epitopes

By “IL-18 receptor” or “IL-18R” herein is meant the cell surface receptor that binds to a ligand, including, but not limited to, IL-18 and as a result initiates a signal transduction pathway within the cell. The IL-18 receptor complex is made up of an IL-18 binding chain termed “ $\alpha$ -IL-18 receptor” ( $\alpha$ -IL-18R) or “IL-18R $\alpha$  chain,” and a signaling chain, termed  $\beta$ -IL-18 receptor “ $\beta$ -IL-18 receptor” ( $\beta$ -IL-18R”) or “IL-18R $\beta$  chain.” As used herein, the term “IL-18 receptor” collectively refers to both the  $\alpha$ - and the  $\beta$ -IL-18 receptor.

The antigen binding proteins disclosed herein bind to the IL-18R $\alpha$  chain, the human amino acid sequence of which is depicted in SEQ ID NO:69 (its nucleic acid sequence is depicted in SEQ ID NO:70), or the IL-18R $\beta$  chain, the human amino acid sequence of which is depicted in SEQ ID NO:71 (its nucleic acid sequence is depicted in SEQ ID NO:72). In a specific embodiment, the IL-18 receptor is human, although in some cases, other species may be used. In addition, as described below, IL-18 receptor proteins may also include fragments.

As is described below, binding of antigen binding proteins to specific epitopes is specific.

By “epitope”, “antigenic determinant”, and grammatical equivalents herein are meant a region of an antigen, e.g., IL-18 receptor, which can be specifically bound by an antigen binding protein. As the skilled artisan will appreciate, an epitope can be linear or conformational. “Linear epitope” refers to an epitope comprising a sequence of at least about five (5) and not more than about twenty (20) amino acids connected in a linear fashion, which amino acids, by themselves or as part of a larger sequence, bind to an antigen binding protein of the invention. “Conformational epitope” refers to an epitope whose three dimensional, secondary and/or tertiary structure can be a substantial aspect of antibody binding. Generally but not uniformly, amino acids that comprise a conformational epitope do not comprise a linear sequence of a protein’s primary structure. Thus, a conformational epitope may be shared by proteins having non-homologous linear amino acid sequences. Without being bound by theory, a conformational epitope can be shared because the tertiary structure recognized by an antibody can be shared between two or more amino acid sequences. In one embodiment, suitable IL-18 receptor epitopes include any which are recognized by the antigen binding proteins of the present invention.

The invention provides antigen binding proteins recognizing and binding to a conformational epitope in the third Ig domain of human IL-18R $\alpha$ , in particular, the region defined

by amino acid residues 243-271, made up by amino acid residues 250-253 (i.e., residues MFGE) and amino acid residues 267-271 (i.e., residues MRIMT) of SEQ ID NO:69. The amino acid structure of this epitope is depicted in FIG. 5.

Antigen binding proteins that bind to this epitope are particularly effective at blocking the interaction of IL-18 with the IL-18R. Methods of determining the binding epitope of an antigen binding protein are well known in the art and one such method is described in Example 4 herein.

Example 4 demonstrates that certain human IL-18R antigen binding proteins had significantly reduced ability to bind the human IL-18R $\alpha$  when residues within the epitope defined by amino acids 243-271, e.g., 250-253 or 267-271, were changed to the corresponding mouse residues. Thus, provided herein are antigen binding proteins that bind human IL-18R but such binding is reduced when residues 250-253 of the human IL-18R $\alpha$  chain are substituted with the corresponding mouse amino acids. Also provided herein are antigen binding proteins that bind human IL-18R but such binding is reduced when residues 267-271 of the human IL-18R $\alpha$  chain are substituted with the corresponding mouse amino acids.

#### D. Covalent Modifications of Antigen Binding Protein

Covalent modifications of antigen binding proteins are included within the scope of this invention, and are generally, but not always, done post-translationally. For example, several types of covalent modifications of the antigen binding protein are introduced into the molecule by reacting specific amino acid residues of the antigen binding protein with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C-terminal residues.

Cysteinyll residues most commonly are reacted with  $\alpha$ -haloacetates (and corresponding amines), such as chloroacetic acid or chloroacetamide, to give carboxymethyl or carboxyamidomethyl derivatives. Cysteinyll residues also are derivatized by reaction with bromotrifluoroacetone,  $\alpha$ -bromo- $\beta$ -(5-imidazolyl)propionic acid, chloroacetyl phosphate, N-alkylmaleimides, 3-nitro-2-pyridyl disulfide, methyl 2-pyridyl disulfide, p-chloromercuribenzoate, 2-chloromercuri-4-nitrophenol, or chloro-7-nitrobenzo-2-oxa-1,3-diazole.

Histidyl residues are derivatized by reaction with diethylpyrocarbonate at pH 5.5-7.0 because this agent is relatively specific for the histidyl side chain. Para-bromophenacyl bromide also is useful; the reaction is preferably performed in 0.1M sodium cacodylate at pH 6.0.

Lysinyll and amino terminal residues are reacted with succinic or other carboxylic acid anhydrides. Derivatization with these agents has the effect of reversing the charge of the lysinyll residues. Other suitable reagents for derivatizing alpha-amino-containing residues include imidoesters such as methyl picolinimidate; pyridoxal phosphate; pyridoxal; chloroborohydride; trinitrobenzenesulfonic acid; O-methylisourea; 2,4-pentanedione; and transaminase-catalyzed reaction with glyoxylate.

Arginyll residues are modified by reaction with one or several conventional reagents, among them phenylglyoxal, 2,3-butanedione, 1,2-cyclohexanedione, and ninhydrin. Derivatization of arginine residues requires that the reaction be performed in alkaline conditions because of the high pK<sub>a</sub> of the guanidine functional group. Furthermore, these reagents may react with the groups of lysine as well as the arginine epsilon-amino group.

The specific modification of tyrosyl residues may be made, with particular interest in introducing spectral labels into tyrosyl residues by reaction with aromatic diazonium compounds or tetranitromethane. Most commonly, N-acetylimidazole and tetranitromethane are used to form O-acetyl

tyrosyl species and 3-nitro derivatives, respectively. Tyrosyl residues are iodinated using  $^{125}\text{I}$  or  $^{131}\text{I}$  to prepare labeled proteins for use in radioimmunoassay, the chloramine T method described above being suitable.

Carboxyl side groups (aspartyl or glutamyl) are selectively modified by reaction with carbodiimides ( $\text{R}'\text{—N}=\text{C}=\text{N—R}'$ ), where R and R' are optionally different alkyl groups, such as 1-cyclohexyl-3-(2-morpholinyl-4-ethyl) carbodiimide or 1-ethyl-3-(4-azonia-4,4-dimethylpentyl) carbodiimide. Furthermore, aspartyl and glutamyl residues are converted to asparaginyl and glutaminyl residues by reaction with ammonium ions.

Derivatization with bifunctional agents is useful for crosslinking antigen binding proteins to a water-insoluble support matrix or surface for use in a variety of methods. Commonly used crosslinking agents include, e.g., 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), and bifunctional maleimides such as bis-N-maleimido-1,8-octane. Derivatizing agents such as methyl-3-[(p-azidophenyl)dithio]propioimidate yield photoactivatable intermediates that are capable of forming crosslinks in the presence of light. Alternatively, reactive water-insoluble matrices such as cyanogen bromide-activated carbohydrates and the reactive substrates described in U.S. Pat. Nos. 3,969,287; 3,691,016; 4,195,128; 4,247,642; 4,229,537; and 4,330,440 are employed for protein immobilization.

Glutamyl and asparaginyl residues are frequently deamidated to the corresponding glutamyl and aspartyl residues, respectively. Alternatively, these residues are deamidated under mildly acidic conditions. Either form of these residues falls within the scope of this invention.

Other modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the  $\alpha$ -amino groups of lysine, arginine, and histidine side chains (T. E. Creighton, *Proteins: Structure and Molecular Properties*, W. H. Freeman & Co., San Francisco, pp. 79-86 [1983]), acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

### 1. Glycosylation

Another type of covalent modification of the antigen binding protein included within the scope of this invention comprises altering the glycosylation pattern of the protein. As is known in the art, glycosylation patterns can depend on both the sequence of the protein (e.g., the presence or absence of particular glycosylation amino acid residues, discussed below), or the host cell or organism in which the protein is produced. Particular expression systems are discussed below.

Glycosylation of polypeptides is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tri-peptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tri-peptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-acetyl-galactosamine, galactose, or xylose, to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

Addition of glycosylation sites to the antigen binding protein is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tri-peptide sequences (for N-linked glycosylation

sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the starting sequence (for O-linked glycosylation sites). For ease, the antigen binding protein amino acid sequence is preferably altered through changes at the DNA level, particularly by mutating the DNA encoding the target polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids.

Another means of increasing the number of carbohydrate moieties on the antigen binding protein is by chemical or enzymatic coupling of glycosides to the protein. These procedures are advantageous in that they do not require production of the protein in a host cell that has glycosylation capabilities for N- and O-linked glycosylation. Depending on the coupling mode used, the sugar(s) may be attached to (a) arginine and histidine, (b) free carboxyl groups, (c) free sulfhydryl groups such as those of cysteine, (d) free hydroxyl groups such as those of serine, threonine, or hydroxyproline, (e) aromatic residues such as those of phenylalanine, tyrosine, or tryptophan, or (f) the amide group of glutamine. These methods are described in WO 87/05330 published Sep. 11, 1987, and in Aplin and Wriston, 1981, *CRC Crit. Rev. Biochem.*, pp. 259-306.

Removal of carbohydrate moieties present on the starting antigen binding protein may be accomplished chemically or enzymatically. Chemical deglycosylation requires exposure of the protein to the compound trifluoromethanesulfonic acid, or an equivalent compound. This treatment results in the cleavage of most or all sugars except the linking sugar (N-acetylglucosamine or N-acetylgalactosamine), while leaving the polypeptide intact. Chemical deglycosylation is described by Hakimuddin et al., 1987, *Arch. Biochem. Biophys.* 259:52 and by Edge et al., 1981, *Anal. Biochem.* 118: 131. Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura et al., 1987, *Meth. Enzymol.* 138:350. Glycosylation at potential glycosylation sites may be prevented by the use of the compound tunicamycin as described by Duskin et al., 1982, *J. Biol. Chem.* 257:3105. Tunicamycin blocks the formation of protein-N-glycoside linkages.

### 2. PEGylation

Another type of covalent modification of the antigen binding protein comprises linking the antigen binding protein to various nonproteinaceous polymers, including, but not limited to, various polyols such as polyethylene glycol, polypropylene glycol or polyoxyalkylenes, in the manner set forth in U.S. Pat. No. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337. In addition, as is known in the art, amino acid substitutions may be made in various positions within the antigen binding protein to facilitate the addition of polymers such as PEG.

### 3. Labels And Effector Groups

In some embodiments, the covalent modification of the antigen binding proteins of the invention comprises the addition of one or more labels.

The term "labeling group" means any detectable label. Examples of suitable labelling groups include, but are not limited to, the following: radioisotopes or radionuclides (e.g.,  $^3\text{H}$ ,  $^{14}\text{C}$ ,  $^{15}\text{N}$ ,  $^{35}\text{S}$ ,  $^{90}\text{Y}$ ,  $^{99}\text{Tc}$ ,  $^{111}\text{In}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ ), fluorescent groups (e.g., FITC, rhodamine, lanthanide phosphors), enzymatic groups (e.g., horseradish peroxidase,  $\beta$ -galactosidase, luciferase, alkaline phosphatase), chemiluminescent groups, biotinyl groups, or predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags). In some embodiments, the

labelling group is coupled to the antigen binding protein via spacer arms of various lengths to reduce potential steric hindrance. Various methods for labelling proteins are known in the art and may be used in performing the present invention.

The term “effector group” means any group coupled to an antigen binding protein that acts as a cytotoxic agent. Examples for suitable effector groups are radioisotopes or radionuclides (e.g., <sup>3</sup>H, <sup>14</sup>C, <sup>15</sup>N, <sup>35</sup>S, <sup>90</sup>Y, <sup>99</sup>Tc, <sup>111</sup>In, <sup>125</sup>I, <sup>131</sup>I). Other suitable groups include toxins, therapeutic groups, or chemotherapeutic groups. Examples of suitable groups include calicheamicin, auristatins, geldanamycin and maytansine. In some embodiments, the effector group is coupled to the antigen binding protein via spacer arms of various lengths to reduce potential steric hindrance.

In general, labels fall into a variety of classes, depending on the assay in which they are to be detected: a) isotopic labels, which may be radioactive or heavy isotopes; b) magnetic labels (e.g., magnetic particles); c) redox active moieties; d) optical dyes; enzymatic groups (e.g. horseradish peroxidase,  $\beta$ -galactosidase, luciferase, alkaline phosphatase); e) biotinylated groups; and f) predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags, etc.). In some embodiments, the labelling group is coupled to the antigen binding protein via spacer arms of various lengths to reduce potential steric hindrance. Various methods for labelling proteins are known in the art and may be used in performing the present invention.

Specific labels include optical dyes, including, but not limited to, chromophores, phosphors and fluorophores, with the latter being specific in many instances. Fluorophores can be either “small molecule” fluo- res, or proteinaceous fluo- res.

By “fluorescent label” is meant any molecule that may be detected via its inherent fluorescent properties. Suitable fluorescent labels include, but are not limited to, fluorescein, rhodamine, tetramethylrhodamine, eosin, erythrosin, coumarin, methyl-coumarins, pyrene, Malacite green, stilbene, Lucifer Yellow, Cascade BlueJ, Texas Red, IAEDANS, EDANS, BODIPY FL, LC Red 640, Cy 5, Cy 5.5, LC Red 705, Oregon green, the Alexa-Fluor dyes (Alexa Fluor 350, Alexa Fluor 430, Alexa Fluor 488, Alexa Fluor 546, Alexa Fluor 568, Alexa Fluor 594, Alexa Fluor 633, Alexa Fluor 660, Alexa Fluor 680), Cascade Blue, Cascade Yellow and R-phycoerythrin (PE) (Molecular Probes, Eugene, Oreg.), FITC, Rhodamine, and Texas Red (Pierce, Rockford, Ill.), Cy5, Cy5.5, Cy7 (Amersham Life Science, Pittsburgh, Pa.). Suitable optical dyes, including fluorophores, are described in Molecular Probes Handbook by Richard P. Haugland, hereby expressly incorporated by reference.

Suitable proteinaceous fluorescent labels also include, but are not limited to, green fluorescent protein, including a *Renilla*, *Ptilosarcus*, or *Aequorea* species of GFP (Chalfie et al., 1994, *Science* 263:802-805), EGFP (Clontech Laboratories, Inc., Genbank Accession Number U55762), blue fluorescent protein (BFP, Quantum Biotechnologies, Inc. 1801 de Maisonneuve Blvd. West, 8th Floor, Montreal, Quebec, Canada H3H 1J9; Stauber, 1998, *Biotechniques* 24:462-471; Heim et al., 1996, *Curr. Biol.* 6:178-182), enhanced yellow fluorescent protein (EYFP, Clontech Laboratories, Inc.), luciferase (Ichiki et al., 1993, *J. Immunol.* 150:5408-5417),  $\beta$  galactosidase (Nolan et al., 1988, *Proc. Natl. Acad. Sci. U.S.A.* 85:2603-2607) and *Renilla* (WO92/15673, WO95/07463, WO98/14605, WO98/26277, WO99/49019, U.S. Pat. Nos. 5,292,658, 5,418,155, 5,683,888, 5,741,668, 5,777,079,

5,804,387, 5,874,304, 5,876,995, 5,925,558). All of the above-cited references are expressly incorporated herein by reference.

E. Polynucleotides Encoding IL-18 Receptor Antigen Binding Proteins

In certain aspects, the invention provides nucleic acid molecules encoding the IgGs, variable regions and CDRs of SEQ ID NOs:1-34, 73, 75, 77, 79, 81, 83, 85, 87, 89-190. In one embodiment, the nucleic acids have the nucleotide sequence of any of SEQ ID NOs:35-68, 74, 76, 78, 80, 82, 84, 86, 88, and 191-292.

As described herein, a variable region or CDR nucleic acid encodes a variable region or CDR protein, respectively. By “nucleic acid” herein is meant any nucleic acid, including both DNA and RNA. Nucleic acids of the present invention are typically polynucleic acids; that is, polymers of individual nucleotides that are covalently joined by 3', 5' phosphodiester bonds.

Depending on its use, the nucleic acid may be double stranded, single stranded, or contain portions of both double stranded or single stranded sequence. As will be appreciated by those in the art, the depiction of a single strand (“Watson”) also defines the sequence of the other strand (“Crick”); thus the nucleic acid sequences depicted in SEQ ID NOs:35-68 also include the complement of these sequences. By the term “recombinant nucleic acid” herein is meant nucleic acid, originally formed in vitro, in general, by the manipulation of nucleic acid by endonucleases, in a form not normally found in nature. Thus an isolated antigen binding protein nucleic acid, in a linear form, or an expression vector formed in vitro by ligating DNA molecules that are not normally joined, are both considered recombinant for the purposes of this invention. It is understood that once a recombinant nucleic acid is made and reintroduced into a host cell or organism, it will replicate non-recombinantly, i.e., using the in vivo cellular machinery of the host cell rather than in vitro manipulations; however, such nucleic acids, once produced recombinantly, although subsequently replicated non-recombinantly, are still considered recombinant for the purposes of the invention.

As will be appreciated by those in the art, due to the degeneracy of the genetic code, an extremely large number of nucleic acids may be made, all of which encode the CDRs (and heavy and light chains or other components of the antigen binding protein) of the present invention. Thus, having identified a particular amino acid sequence, such as SEQ ID NOs:1-34, those skilled in the art could make any number of different nucleic acids, by simply modifying the sequence of one or more codons in a way which does not change the amino acid sequence of the encoded protein.

F. Methods of Producing Antigen Binding Proteins

The present invention also provides expression systems and constructs in the form of plasmids, expression vectors, transcription or expression cassettes which comprise at least one polynucleotide as above. In addition, the invention provides host cells comprising such expression systems or constructs.

Typically, expression vectors used in any of the host cells will contain sequences for plasmid maintenance and for cloning and expression of exogenous nucleotide sequences. Such sequences, collectively referred to as “flanking sequences” in certain embodiments will typically include one or more of the following nucleotide sequences: a promoter, one or more enhancer sequences, an origin of replication, a transcriptional termination sequence, a complete intron sequence containing a donor and acceptor splice site, a sequence encoding a leader sequence for polypeptide secretion, a ribosome binding site, a polyadenylation sequence, a polylinker region for inserting

the nucleic acid encoding the polypeptide to be expressed, and a selectable marker element. Each of these sequences is discussed below.

Optionally, the vector may contain a "tag"-encoding sequence, i.e., an oligonucleotide molecule located at the 5' or 3' end of the IL-18 receptor antigen binding protein coding sequence; the oligonucleotide sequence encodes polyHis (such as hexaHis), or another "tag" such as FLAG, HA (hemagglutinin influenza virus), or myc, for which commercially available antibodies exist. This tag is typically fused to the polypeptide upon expression of the polypeptide, and can serve as a means for affinity purification or detection of the IL-18 receptor antigen binding protein from the host cell. Affinity purification can be accomplished, for example, by column chromatography using antibodies against the tag as an affinity matrix. Optionally, the tag can subsequently be removed from the purified IL-18 receptor antigen binding protein by various means such as using certain peptidases for cleavage.

Flanking sequences may be homologous (i.e., from the same species and/or strain as the host cell), heterologous (i.e., from a species other than the host cell species or strain), hybrid (i.e., a combination of flanking sequences from more than one source), synthetic or native. As such, the source of a flanking sequence may be any prokaryotic or eukaryotic organism, any vertebrate or invertebrate organism, or any plant, provided that the flanking sequence is functional in, and can be activated by, the host cell machinery.

Flanking sequences useful in the vectors of this invention may be obtained by any of several methods well known in the art. Typically, flanking sequences useful herein will have been previously identified by mapping and/or by restriction endonuclease digestion and can thus be isolated from the proper tissue source using the appropriate restriction endonucleases. In some cases, the full nucleotide sequence of a flanking sequence may be known. Here, the flanking sequence may be synthesized using the methods described herein for nucleic acid synthesis or cloning.

Whether all or only a portion of the flanking sequence is known, it may be obtained using polymerase chain reaction (PCR) and/or by screening a genomic library with a suitable probe such as an oligonucleotide and/or flanking sequence fragment from the same or another species. Where the flanking sequence is not known, a fragment of DNA containing a flanking sequence may be isolated from a larger piece of DNA that may contain, for example, a coding sequence or even another gene or genes. Isolation may be accomplished by restriction endonuclease digestion to produce the proper DNA fragment followed by isolation using agarose gel purification, Qiagen® column chromatography (Chatsworth, Calif.), or other methods known to the skilled artisan. The selection of suitable enzymes to accomplish this purpose will be readily apparent to one of ordinary skill in the art.

An origin of replication is typically a part of those prokaryotic expression vectors purchased commercially, and the origin aids in the amplification of the vector in a host cell. If the vector of choice does not contain an origin of replication site, one may be chemically synthesized based on a known sequence, and ligated into the vector. For example, the origin of replication from the plasmid pBR322 (New England Biolabs, Beverly, Mass.) is suitable for most gram-negative bacteria, and various viral origins (e.g., SV40, polyoma, adenovirus, vesicular stomatitis virus (VSV), or papillomaviruses such as HPV or BPV) are useful for cloning vectors in mammalian cells. Generally, the origin of replication component is not needed for mammalian expression vectors (for

example, the SV40 origin is often used only because it also contains the virus early promoter).

A transcription termination sequence is typically located 3' to the end of a polypeptide coding region and serves to terminate transcription. Usually, a transcription termination sequence in prokaryotic cells is a G-C rich fragment followed by a poly-T sequence. While the sequence is easily cloned from a library or even purchased commercially as part of a vector, it can also be readily synthesized using methods for nucleic acid synthesis such as those described herein.

A selectable marker gene encodes a protein necessary for the survival and growth of a host cell grown in a selective culture medium. Typical selection marker genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, tetracycline, or kanamycin for prokaryotic host cells; (b) complement auxotrophic deficiencies of the cell; or (c) supply critical nutrients not available from complex or defined media. Specific selectable markers are the kanamycin resistance gene, the ampicillin resistance gene, and the tetracycline resistance gene. Advantageously, a neomycin resistance gene may also be used for selection in both prokaryotic and eukaryotic host cells.

Other selectable genes may be used to amplify the gene that will be expressed. Amplification is the process wherein genes that are required for production of a protein critical for growth or cell survival are reiterated in tandem within the chromosomes of successive generations of recombinant cells. Examples of suitable selectable markers for mammalian cells include dihydrofolate reductase (DHFR) and promoterless thymidine kinase genes. Mammalian cell transformants are placed under selection pressure wherein only the transformants are uniquely adapted to survive by virtue of the selectable gene present in the vector. Selection pressure is imposed by culturing the transformed cells under conditions in which the concentration of selection agent in the medium is successively increased, thereby leading to the amplification of both the selectable gene and the DNA that encodes another gene, such as an antigen binding protein antibody that binds to IL-18 receptor polypeptide. As a result, increased quantities of a polypeptide such as an IL-18 receptor antigen binding protein are synthesized from the amplified DNA.

A ribosome-binding site is usually necessary for translation initiation of mRNA and is characterized by a Shine-Dalgarno sequence (prokaryotes) or a Kozak sequence (eukaryotes). The element is typically located 3' to the promoter and 5' to the coding sequence of the polypeptide to be expressed.

In some cases, such as where glycosylation is desired in a eukaryotic host cell expression system, one may manipulate the various pre- or prosequences to improve glycosylation or yield. For example, one may alter the peptidase cleavage site of a particular signal peptide, or add prosequences, which also may affect glycosylation. The final protein product may have, in the -1 position (relative to the first amino acid of the mature protein) one or more additional amino acids incident to expression, which may not have been totally removed. For example, the final protein product may have one or two amino acid residues found in the peptidase cleavage site, attached to the amino-terminus. Alternatively, use of some enzyme cleavage sites may result in a slightly truncated form of the desired polypeptide, if the enzyme cuts at such area within the mature polypeptide.

Expression and cloning vectors of the invention will typically contain a promoter that is recognized by the host organism and operably linked to the molecule encoding the IL-18 receptor antigen binding protein. Promoters are untranscribed sequences located upstream (i.e., 5') to the start codon

of a structural gene (generally within about 100 to 1000 bp) that control transcription of the structural gene. Promoters are conventionally grouped into one of two classes: inducible promoters and constitutive promoters. Inducible promoters initiate increased levels of transcription from DNA under their control in response to some change in culture conditions, such as the presence or absence of a nutrient or a change in temperature. Constitutive promoters, on the other hand, uniformly transcribe gene to which they are operably linked, that is, with little or no control over gene expression. A large number of promoters, recognized by a variety of potential host cells, are well known. A suitable promoter is operably linked to the DNA encoding heavy chain or light chain comprising an IL-18 receptor antigen binding protein of the invention by removing the promoter from the source DNA by restriction enzyme digestion and inserting the desired promoter sequence into the vector.

Suitable promoters for use with yeast hosts are also well known in the art. Yeast enhancers are advantageously used with yeast promoters. Suitable promoters for use with mammalian host cells are well known and include, but are not limited to, those obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, retroviruses, hepatitis-B virus and most preferably Simian Virus 40 (SV40). Other suitable mammalian promoters include heterologous mammalian promoters, for example, heat-shock promoters and the actin promoter.

Additional promoters which may be of interest include, but are not limited to: SV40 early promoter (Benoist and Chambon, 1981, *Nature* 290:304-310); CMV promoter (Thornsen et al., 1984, *Proc. Natl. Acad. U.S.A.* 81:659-663); the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., 1980, *Cell* 22:787-797); herpes thymidine kinase promoter (Wagner et al., 1981, *Proc. Natl. Acad. Sci. U.S.A.* 78:1444-1445); promoter and regulatory sequences from the metallothionein gene Prinster et al., 1982, *Nature* 296:39-42); and prokaryotic promoters such as the beta-lactamase promoter (Villa-Kamaroff et al., 1978, *Proc. Natl. Acad. Sci. U.S.A.* 75:3727-3731); or the tac promoter (DeBoer et al., 1983, *Proc. Natl. Acad. Sci. U.S.A.* 80:21-25). Also of interest are the following animal transcriptional control regions, which exhibit tissue specificity and have been utilized in transgenic animals: the elastase I gene control region that is active in pancreatic acinar cells (Swift et al., 1984, *Cell* 38:639-646; Ornitz et al., 1986, *Cold Spring Harbor Symp. Quant. Biol.* 50:399-409; MacDonald, 1987, *Hepatology* 7:425-515); the insulin gene control region that is active in pancreatic beta cells (Hanahan, 1985, *Nature* 315:115-122); the immunoglobulin gene control region that is active in lymphoid cells (Grosschedl et al., 1984, *Cell* 38:647-658; Adames et al., 1985, *Nature* 318:533-538; Alexander et al., 1987, *Mol. Cell. Biol.* 7:1436-1444); the mouse mammary tumor virus control region that is active in testicular, breast, lymphoid and mast cells (Leder et al., 1986, *Cell* 45:485-495); the albumin gene control region that is active in liver (Pinkert et al., 1987, *Genes and Devel.* 1:268-276); the alpha-feto-protein gene control region that is active in liver (Krumlauf et al., 1985, *Mol. Cell. Biol.* 5:1639-1648; Hammer et al., 1987, *Science* 253:53-58); the alpha 1-antitrypsin gene control region that is active in liver (Kelsey et al., 1987, *Genes and Devel.* 1:161-171); the beta-globin gene control region that is active in myeloid cells (Mogram et al., 1985, *Nature* 315:338-340; Kollias et al., 1986, *Cell* 46:89-94); the myelin basic protein gene control region that is active in oligodendrocyte cells in the brain (Readhead et al., 1987, *Cell* 48:703-712); the myosin light chain-2 gene control region that is

active in skeletal muscle (Sani, 1985, *Nature* 314:283-286); and the gonadotropic releasing hormone gene control region that is active in the hypothalamus (Mason et al., 1986, *Science* 234:1372-1378).

An enhancer sequence may be inserted into the vector to increase transcription of DNA encoding light chain or heavy chain comprising an IL-18 receptor antigen binding protein of the invention by higher eukaryotes. Enhancers are cis-acting elements of DNA, usually about 10-300 bp in length, that act on the promoter to increase transcription. Enhancers are relatively orientation and position independent, having been found at positions both 5' and 3' to the transcription unit. Several enhancer sequences available from mammalian genes are known (e.g., globin, elastase, albumin, alpha-feto-protein and insulin). Typically, however, an enhancer from a virus is used. The SV40 enhancer, the cytomegalovirus early promoter enhancer, the polyoma enhancer, and adenovirus enhancers known in the art are exemplary enhancing elements for the activation of eukaryotic promoters. While an enhancer may be positioned in the vector either 5' or 3' to a coding sequence, it is typically located at a site 5' from the promoter. A sequence encoding an appropriate native or heterologous signal sequence (leader sequence or signal peptide) can be incorporated into an expression vector, to promote extracellular secretion of the antibody. The choice of signal peptide or leader depends on the type of host cells in which the antibody is to be produced, and a heterologous signal sequence can replace the native signal sequence. Examples of signal peptides that are functional in mammalian host cells include the following: the signal sequence for interleukin-7 (IL-7) described in U.S. Pat. No. 4,965,195; the signal sequence for interleukin-2 receptor described in Cosman et al., 1984, *Nature* 312:768; the interleukin-4 receptor signal peptide described in EP Patent No. 0367 566; the type I interleukin-1 receptor signal peptide described in U.S. Pat. No. 4,968,607; the type II interleukin-1 receptor signal peptide described in EP Patent No. 0 460 846.

Expression vectors of the invention may be constructed from a starting vector such as a commercially available vector. Such vectors may or may not contain all of the desired flanking sequences. Where one or more of the flanking sequences described herein are not already present in the vector, they may be individually obtained and ligated into the vector. Methods used for obtaining each of the flanking sequences are well known to one skilled in the art.

After the vector has been constructed and a nucleic acid molecule encoding light chain, a heavy chain, or a light chain and a heavy chain comprising an IL-18 receptor antigen binding sequence has been inserted into the proper site of the vector, the completed vector may be inserted into a suitable host cell for amplification and/or polypeptide expression. The transformation of an expression vector for an IL-18 receptor antigen binding protein into a selected host cell may be accomplished by well known methods including transfection, infection, calcium phosphate co-precipitation, electroporation, microinjection, lipofection, DEAE-dextran mediated transfection, or other known techniques. The method selected will in part be a function of the type of host cell to be used. These methods and other suitable methods are well known to the skilled artisan, and are set forth, for example, in Sambrook et al., 2001, *supra*.

A host cell, when cultured under appropriate conditions, synthesizes an IL-18 receptor antigen binding protein that can subsequently be collected from the culture medium (if the host cell secretes it into the medium) or directly from the host cell producing it (if it is not secreted). The selection of an appropriate host cell will depend upon various factors, such as

desired expression levels, polypeptide modifications that are desirable or necessary for activity (such as glycosylation or phosphorylation) and ease of folding into a biologically active molecule.

Mammalian cell lines available as hosts for expression are well known in the art and include, but are not limited to, immortalized cell lines available from the American Type Culture Collection (ATCC), including but not limited to Chinese hamster ovary (CHO) cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (e.g., Hep G2), and a number of other cell lines. In certain embodiments, cell lines may be selected through determining which cell lines have high expression levels and constitutively produce antigen binding proteins with IL-18 receptor binding properties. In another embodiment, a cell line from the B cell lineage that does not make its own antibody but has a capacity to make and secrete a heterologous antibody can be selected.

#### G. Use of IL-18 Receptor Antigen Binding Proteins for Diagnostic and Therapeutic Purposes

Antigen binding proteins of the invention are useful for detecting IL-18 receptor in biological samples and identification of cells or tissues that produce IL-18 receptor protein. Antigen binding proteins of the invention that specifically bind to IL-18 receptor may be used in treatment of IL-18 receptor mediated diseases in a patient in need thereof. For one, the IL-18 receptor antigen binding proteins of the invention can be used in diagnostic assays, e.g., binding assays to detect and/or quantify IL-18 receptor expressed in a tissue or cell. In addition, the IL-18 receptor antigen binding protein of the invention can be used to inhibit IL-18 receptor from forming a complex with its ligand, e.g., IL-18, thereby modulating the biological activity of IL-18 receptor in a cell or tissue. Antigen binding proteins that bind to IL-18 receptor thus may modulate and/or block interaction with other binding compounds and as such may have therapeutic use in ameliorating IL-18 receptor mediated diseases. In specific embodiments, IL-18 receptor antigen binding proteins may block IL-18 binding to its receptor, which may result in disruption of the IL-18 receptor induced signal transduction cascade.

##### 1. Indications

Increased levels of IL-18 and/or involvement of IL-18 mediated signals in disease pathogenesis have been demonstrated in a variety of conditions and diseases. The antigen binding proteins of the present invention thus serve to regulate or suppress an immune response and have efficacy in the treatment and prevention of various diseases caused by an excessive immune response (see, WO2004/002519; WO2005/063290; WO2004/034988; Mallat et al., 2002, *Circ. Res.* 91:441-448). Accordingly, the IL-18 receptor antigen binding proteins of the present invention can be used for the diagnosis, prevention or treatment of diseases or conditions associated with the IL-18.

A disease or condition associated with IL-18 means any disease or condition whose onset in a patient is caused by or prevented by the interaction of IL-18 with the IL-18 receptor. The severity of the disease or condition can also be increased or decreased by the interaction of IL-18 with the IL-18 receptor. For example, IL-18 is associated with autoimmune diseases (WO2004/002519; WO2005/063290; WO2004/034988; Mallat et al., 2002, *Circ. Res.* 91:441-448), hepatic disease (Finitto et al., 2004, *Liver* 53:392-400; Tsutsui et al., 2000, *Immunological Reviews* 174:192-209; Ludwiczek et al., 2002, *J. Clinical Immunology* 22:331-337), pancreatic disease and cardiovascular diseases (Gerdes et al., 2002, *J. Exp. Med.* 195:245-257; WO03/080104; WO02/060479;

WO01/85201; Raeburn et al., 2002, *Am. J. Physiol. Heart Circ. Physiol.* 283:H650-H657).

Examples of autoimmune diseases that are associated with IL-18 include psoriasis, inflammatory arthritis such as rheumatoid arthritis (WO2005/063290; Cannetti et al., 2003, *J. Immunol.* 171:1009-1015; Charles et al., 1999, *J. Immunol.* 163: 1521-1528; Cunnane et al., 2000, *Online J. Rheumatol.* 27: 58-63; Yoshimoto, 1998, *J. Immunol.* 161: 3400-3407), lupus (WO2005/063290), Type I diabetes, Type II diabetes, Crohn's disease (Niederer, 1997, *Online NLM*), inflammatory bowel disease (WO2004/002519), multiple sclerosis, autoimmune hepatitis (Tsutsui et al., 2000, *supra*), primary biliary cirrhosis (PBC), acquired immune deficiency syndrome (AIDS), atopic dermatitis (Konishi et al., 2002, *Proc. Natl. Acad. Sci. U.S.A.* 99:11340-11345), myasthenia gravis, and sarcoidosis.

In rheumatoid arthritis, elevated levels of mature IL-18 have been demonstrated in patient sera and synovial fluid. In some studies, IL-18 levels were shown to correlate with disease activity and response to disease modifying treatment. Extremely elevated serum levels of IL-18 have consistently been measured in systemic Juvenile Idiopathic Arthritis and the closely related Adult-Onset Still's Disease. WO2005/063290; Cannetti et al., 2003, *J. Immunol.* 171:1009-1015; Charles et al., 1999, *J. Immunol.* 163: 1521-1528; Cunnane et al., 2000, *Online J. Rheumatol.* 27: 58-63; Yoshimoto, 1998, *J. Immunol.* 161: 3400-3407.

Other forms of arthritis that are associated with IL-18 include for example ankylosing spondylitis, back pain, carpal deposition syndrome, Ehlers-Danlos-Syndrome, gout, juvenile arthritis, lupus erythematosus, myositis, osteogenesis imperfecta, osteoporosis, polyarthritis; polymyositis, psoriatic arthritis, Reiter's syndrome, scleroderma, arthritis with bowel disease, Behcets's disease, children's arthritis, degenerative joint disease, fibromyalgia, infectious arthritis, Lyme disease, Marian syndrome, osteoarthritis, osteonecrosis, Pagets Disease, Polymyalgia rheumatica, pseudogout, reflex sympathetic dystrophy, rheumatoid arthritis, rheumatism, Sjogren's syndrome, familial adenomatous polyposis and the like. Dai et al., 2004, *Arthritis Rheum.* 50:432-443; Kawashima et al., 2004, *Online Arthritis Res. Ther.* 6:R39-R45; Myers et al., 2004, *Rheumatology* 43:272-276; Wei et al., 2001, *American Association Of Immunologists*, pp. 517-521.

Elevated levels of IL-18 have also been found in patients with Crohn's disease when compared with patients with ulcerative colitis or non-inflammatory intestinal conditions. Both intestinal epithelial cells and lamina propria mononuclear cells have been identified as the source of increased IL-18 production in situ. Crohn's disease lesions have been shown to be infiltrated with IL-18R expressing cells. Niederer, 1997, *Online NLM*.

IL-18 has also been implicated as being associated with ulcerative colitis and Coeliac Disease.

Central Nervous System (CNS) lesions, cerebrospinal fluid, and sera from patients with Multiple Sclerosis have been shown to contain increased levels of IL-18 message or protein. Within lesions, microglia and macrophages are thought to be the source of IL-18. IL-18 cannot be detected in control tissue biopsies from individuals with non-inflammatory CNS diseases. Particularly high levels of IL-18 has been found in the patient subset with relapsing-remitting disease; and IL-18 levels have been found to increase during relapses compared to periods of remission. Huang et al., 2004, *Mult. Scler.* 10:482-7; Karni et al., 2002, *J. Neuroimmunol.* 125:

134-40; Losy et al., 2001, *Acta Neurol. Scand.* 104:171-3; Nicoletti et al., 2001, *Neurology* 57:342-4; Fassbender et al., 1999, *Neurology* 53:1104-6.

In patients with psoriasis, serum levels of IL-18 were reported to be increased, correlating with the extent of skin lesions and PASI score. Overexpression of both IL-18 and IL-18R mRNA has been demonstrated in lesional skin compared with non-lesional or normal skin controls. Documented overexpression of IFN- $\gamma$  and TNF- $\alpha$  in psoriatic skin is consistent with biological activities exerted by IL-18. Arican et al., 2005, *Mediators Inflamm.* 2005:273-9; Piskin et al., 2004, *Exp. Dermatol.* 13:764-72; Companjen et al., 2004 *Eur. Cytokine Netw.* 15:210-6; Pietrzak et al., 2003, *Acta Derm. Venereol.* 83:262-5.

Various other autoimmune diseases have been associated with increased levels of IL-18 either in diseased tissue or in the serum. These include Systemic Lupus Erythematosus, atopic dermatitis, myasthenia gravis, type I diabetes, and sarcoidosis. IL-18 may also be involved in asthma, Alzheimer's Disease, allergic rhinitis, Idiopathic Thrombocytopenic Purpura (ITP), transplantation and GvHD.

IL-18 has also been implicated in a liver or hepatic diseases and in conditions associated with liver damage or injury. Liver damage or injury may have diverse causes. It may be due to viral or bacterial infections, alcohol abuse, immunological disorders, or cancer, for example. Liver injury also includes damages of the bile ducts, and damage to the liver in conditions such as alcoholic hepatitis, liver cirrhosis, viral hepatitis, primary biliary cirrhosis, and alcohol-related hepatic necro-inflammation. Finitto et al., 2004, *Liver* 53:392-400; Tsutsui et al., 2000, *Immunological Reviews* 174:192-209; Ludwiczek et al., 2002, *J. Clinical Immunology* 22:331-337.

Hepatic diseases that are associated with IL-18 include hepatitis C and hepatitis B. IL-18 has been implicated in the pathogenesis of both autoimmune and infectious hepatitis. It is thought to contribute to hepatocyte death via upregulation of proapoptotic molecules, including FasL. It has been suggested that the beneficial effect of interferon-alpha in hepatitis C may be mediated through reduced levels of IL-18. In contrast, IL-18 administration had a beneficial effect in a transgenic model of hepatitis B, improving viral clearance through increased NK and CTL activity. Finitto et al., 2004, *Liver* 53:392-400; Tsutsui et al., 2000, *Immunological Reviews* 174:192-209; Ludwiczek et al., 2002, *J. Clinical Immunology* 22:331-337.

Apart from Hepatitis B and C virus, at least four other viruses causing virus-associated hepatitis have been discovered so far, called Hepatitis A, D, E and G-Virus.

IL-18 is also associated with cardiovascular disease, including atheromatous plaque rupture, post-ischemic heart failure, reperfusion injury, atherosclerosis, chronic heart failure, cardiovascular complications of rheumatoid arthritis and other cardiovascular disorders. IL-18 is thought to markedly depress cardiac output in the setting of sepsis or endotoxin shock. IL-18 is an important link between inflammatory processes and atherogenesis, which is particularly relevant given the accumulating evidence for large excess mortality from cardiovascular causes in patients with chronic inflammatory conditions, including RA and lupus. IL-18 levels have been shown to be a strong independent predictor of death from cardiac events (with better predictive power than CRP levels). Gerdes et al, 2002, *J. Exp. Med.* 195:245-257; WO03/080104; WO02/060479; WO01/85201; Raeburn et al., 2002, *Am. J. Physiol. Heart Circ. Physiol.* 283:H650-H657.

IL-18 may also be associated with pulmonary diseases such as, for example, Chronic Obstructed Pulmonary Disease (COPD), chronic severe asthma and Acute Respiratory Distress Syndrome (ARDS).

## 2. Diagnostic Methods

The antigen binding proteins of the invention can be used for diagnostic purposes to detect, diagnose, or monitor diseases and/or conditions associated with IL-18 or the IL-18 receptor. The invention provides for the detection of the presence of the IL-18 receptor in a sample using classical immunohistological methods known to those of skill in the art (e.g., Tijssen, 1993, *Practice and Theory of Enzyme Immunoassays*, vol 15 (Eds R. H. Burdon and P. H. van Knippenberg, Elsevier, Amsterdam); Zola, 1987, *Monoclonal Antibodies: A Manual of Techniques*, pp. 147-158 (CRC Press, Inc.); Jalkanen et al., 1985, *J. Cell. Biol.* 101:976-985; Jalkanen et al., 1987, *J. Cell Biol.* 105:3087-3096). The detection of the IL-18 receptor can be performed in vivo or in vitro.

Diagnostic applications provided herein include use of the antigen binding proteins to detect expression of the IL-18 receptor and binding of the ligands to the IL-18 receptor. Examples of methods useful in the detection of the presence of the IL-18 receptor include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA).

For diagnostic applications, the antigen binding protein typically will be labeled with a detectable labeling group. Suitable labeling groups include, but are not limited to, the following: radioisotopes or radionuclides (e.g.,  $^3\text{H}$ ,  $^{14}\text{C}$ ,  $^{15}\text{N}$ ,  $^{35}\text{S}$ ,  $^{90}\text{Y}$ ,  $^{99}\text{Tc}$ ,  $^{111}\text{In}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ ), fluorescent groups (e.g., FITC, rhodamine, lanthanide phosphors), enzymatic groups (e.g., horseradish peroxidase,  $\beta$ -galactosidase, luciferase, alkaline phosphatase), chemiluminescent groups, biotinyl groups, or predetermined polypeptide epitopes recognized by a secondary reporter (e.g., leucine zipper pair sequences, binding sites for secondary antibodies, metal binding domains, epitope tags). In some embodiments, the labelling group is coupled to the antigen binding protein via spacer arms of various lengths to reduce potential steric hindrance. Various methods for labelling proteins are known in the art and may be used in performing the present invention.

One aspect of the invention provides for identifying a cell or cells that express the IL-18 receptor. In a specific embodiment, the antigen binding protein is labeled with a labeling group and the binding of the labeled antigen binding protein to the IL-18 receptor is detected. In a further specific embodiment, the binding of the antigen binding protein to the IL-18 receptor is detected in vivo. In a further specific embodiment, the antigen binding protein-IL-18 receptor is isolated and measured using techniques known in the art. See, for example, Harlow and Lane, 1988, *Antibodies: A Laboratory Manual*, New York: Cold Spring Harbor (ed. 1991 and periodic supplements); John E. Coligan, ed., 1993, *Current Protocols In Immunology* New York: John Wiley & Sons.

Another aspect of the invention provides for detecting the presence of a test molecule that competes for binding to the IL-18 receptor with the antigen binding proteins of the invention. An example of one such assay would involve detecting the amount of free antigen binding protein in a solution containing an amount of IL-18 receptor in the presence or absence of the test molecule. An increase in the amount of free antigen binding protein (i.e., the antigen binding protein not bound to the IL-18 receptor) would indicate that the test molecule is capable of competing for IL-18 receptor binding with the antigen binding protein. In one embodiment, the antigen binding protein is labeled with a labeling group.

Alternatively, the test molecule is labeled and the amount of free test molecule is monitored in the presence and absence of an antigen binding protein.

### 3. Methods of Treatment: Pharmaceutical Formulations, Routes of Administration

In some embodiments, the invention provides pharmaceutical compositions comprising a therapeutically effective amount of one or a plurality of the antigen binding proteins of the invention together with a pharmaceutically acceptable diluent, carrier, solubilizer, emulsifier, preservative, and/or adjuvant. In addition, the invention provides methods of treating a patient by administering such pharmaceutical composition. The term "patient" includes human and animal subjects.

Preferably, acceptable formulation materials are nontoxic to recipients at the dosages and concentrations employed. In specific embodiments, pharmaceutical compositions comprising a therapeutically effective amount of IL-18 receptor antigen binding proteins are provided.

In certain embodiments, acceptable formulation materials preferably are nontoxic to recipients at the dosages and concentrations employed. In certain embodiments, the pharmaceutical composition may contain formulation materials for modifying, maintaining or preserving, for example, the pH, osmolarity, viscosity, clarity, color, isotonicity, odor, sterility, stability, rate of dissolution or release, adsorption or penetration of the composition. In such embodiments, suitable formulation materials include, but are not limited to, amino acids (such as glycine, glutamine, asparagine, arginine or lysine); antimicrobials; antioxidants (such as ascorbic acid, sodium sulfite or sodium hydrogen-sulfite); buffers (such as borate, bicarbonate, Tris-HCl, citrates, phosphates or other organic acids); bulking agents (such as mannitol or glycine); chelating agents (such as ethylenediamine tetraacetic acid (EDTA)); complexing agents (such as caffeine, polyvinylpyrrolidone, beta-cyclodextrin or hydroxypropyl-beta-cyclodextrin); fillers; monosaccharides; disaccharides; and other carbohydrates (such as glucose, mannose or dextrans); proteins (such as serum albumin, gelatin or immunoglobulins); coloring, flavoring and diluting agents; emulsifying agents; hydrophilic polymers (such as polyvinylpyrrolidone); low molecular weight polypeptides; salt-forming counterions (such as sodium); preservatives (such as benzalkonium chloride, benzoic acid, salicylic acid, thimerosal, phenethyl alcohol, methylparaben, propylparaben, chlorhexidine, sorbic acid or hydrogen peroxide); solvents (such as glycerin, propylene glycol or polyethylene glycol); sugar alcohols (such as mannitol or sorbitol); suspending agents; surfactants or wetting agents (such as pluronics, PEG, sorbitan esters, polysorbates such as polysorbate 20, polysorbate, triton, tromethamine, lecithin, cholesterol, tyloxapal); stability enhancing agents (such as sucrose or sorbitol); tonicity enhancing agents (such as alkali metal halides, preferably sodium or potassium chloride, mannitol sorbitol); delivery vehicles; diluents; excipients and/or pharmaceutical adjuvants. See, REMINGTON'S PHARMACEUTICAL SCIENCES, 18<sup>th</sup> Edition, (A. R. Genrmo, ed.), 1990, Mack Publishing Company.

In certain embodiments, the optimal pharmaceutical composition will be determined by one skilled in the art depending upon, for example, the intended route of administration, delivery format and desired dosage. See, for example, REMINGTON'S PHARMACEUTICAL SCIENCES, supra. In certain embodiments, such compositions may influence the physical state, stability, rate of in vivo release and rate of in vivo clearance of the antigen binding proteins of the invention. In certain embodiments, the primary vehicle or carrier in

a pharmaceutical composition may be either aqueous or non-aqueous in nature. For example, a suitable vehicle or carrier may be water for injection, physiological saline solution or artificial cerebrospinal fluid, possibly supplemented with other materials common in compositions for parenteral administration. Neutral buffered saline or saline mixed with serum albumin are further exemplary vehicles. In specific embodiments, pharmaceutical compositions comprise Tris buffer of about pH 7.0-8.5, or acetate buffer of about pH 4.0-5.5, and may further include sorbitol or a suitable substitute therefor. In certain embodiments of the invention, IL-18 receptor antigen binding protein compositions may be prepared for storage by mixing the selected composition having the desired degree of purity with optional formulation agents (REMINGTON'S PHARMACEUTICAL SCIENCES, supra) in the form of a lyophilized cake or an aqueous solution. Further, in certain embodiments, the IL-18 receptor antigen binding protein product may be formulated as a lyophilizate using appropriate excipients such as sucrose.

The pharmaceutical compositions of the invention can be selected for parenteral delivery. Alternatively, the compositions may be selected for inhalation or for delivery through the digestive tract, such as orally. Preparation of such pharmaceutically acceptable compositions is within the skill of the art.

The formulation components are present preferably in concentrations that are acceptable to the site of administration. In certain embodiments, buffers are used to maintain the composition at physiological pH or at a slightly lower pH, typically within a pH range of from about 5 to about 8.

When parenteral administration is contemplated, the therapeutic compositions for use in this invention may be provided in the form of a pyrogen-free, parenterally acceptable aqueous solution comprising the desired IL-18 receptor antigen binding protein in a pharmaceutically acceptable vehicle. A particularly suitable vehicle for parenteral injection is sterile distilled water in which the IL-18 receptor antigen binding protein is formulated as a sterile, isotonic solution, properly preserved. In certain embodiments, the preparation can involve the formulation of the desired molecule with an agent, such as injectable microspheres, bio-erodible particles, polymeric compounds (such as polylactic acid or polyglycolic acid), beads or liposomes, that may provide controlled or sustained release of the product which can be delivered via depot injection. In certain embodiments, hyaluronic acid may also be used, having the effect of promoting sustained duration in the circulation. In certain embodiments, implantable drug delivery devices may be used to introduce the desired antigen binding protein.

Pharmaceutical compositions of the invention can be formulated for inhalation. In these embodiments, IL-18 receptor antigen binding proteins are advantageously formulated as a dry, inhalable powder. In specific embodiments, IL-18 receptor antigen binding protein inhalation solutions may also be formulated with a propellant for aerosol delivery. In certain embodiments, solutions may be nebulized. Pulmonary administration and formulation methods therefore are further described in International Patent Application No. PCT/US94/001875, which is incorporated by reference and describes pulmonary delivery of chemically modified proteins. It is also contemplated that formulations can be administered orally. IL-18 receptor antigen binding proteins that are administered in this fashion can be formulated with or without carriers customarily used in the compounding of solid dosage forms such as tablets and capsules. In certain embodiments, a capsule may be designed to release the active portion of the formulation at the point in the gastrointestinal tract when



bioavailability is maximized and pre-systemic degradation is minimized. Additional agents can be included to facilitate absorption of the IL-18 receptor antigen binding protein. Diluents, flavorings, low melting point waxes, vegetable oils, lubricants, suspending agents, tablet disintegrating agents, and binders may also be employed.

A pharmaceutical composition of the invention is preferably provided to comprise an effective quantity of one or a plurality of IL-18 receptor antigen binding proteins in a mixture with non-toxic excipients that are suitable for the manufacture of tablets. By dissolving the tablets in sterile water, or another appropriate vehicle, solutions may be prepared in unit-dose form. Suitable excipients include, but are not limited to, inert diluents, such as calcium carbonate, sodium carbonate or bicarbonate, lactose, or calcium phosphate; or binding agents, such as starch, gelatin, or acacia; or lubricating agents such as magnesium stearate, stearic acid, or talc.

Additional pharmaceutical compositions will be evident to those skilled in the art, including formulations involving IL-18 receptor antigen binding proteins in sustained- or controlled-delivery formulations. Techniques for formulating a variety of other sustained- or controlled-delivery means, such as liposome carriers, bio-erodible microparticles or porous beads and depot injections, are also known to those skilled in the art. See, for example, International Patent Application No. PCT/US93/00829, which is incorporated by reference and describes controlled release of porous polymeric microparticles for delivery of pharmaceutical compositions. Sustained-release preparations may include semipermeable polymer matrices in the form of shaped articles, e.g., films, or microcapsules. Sustained release matrices may include polyesters, hydrogels, polylactides (as disclosed in U.S. Pat. No. 3,773,919 and European Patent Application Publication No. EP 058481, each of which is incorporated by reference), copolymers of L-glutamic acid and gamma ethyl-L-glutamate (Sidman et al., 1983, *Biopolymers* 2:547-556), poly (2-hydroxyethyl-methacrylate) (Langer et al., 1981, *J. Biomed. Mater. Res.* 15:167-277 and Langer, 1982, *Chem. Tech.* 12:98-105), ethylene vinyl acetate (Langer et al., 1981, supra) or poly-D(-)-3-hydroxybutyric acid (European Patent Application Publication No. EP 133,988). Sustained release compositions may also include liposomes that can be prepared by any of several methods known in the art. See, e.g., Eppstein et al., 1985, *Proc. Natl. Acad. Sci. U.S.A.* 82:3688-3692; European Patent Application Publication Nos. EP 036,676; EP 088,046 and EP 143,949, incorporated by reference.

Pharmaceutical compositions used for in vivo administration are typically provided as sterile preparations. Sterilization can be accomplished by filtration through sterile filtration membranes. When the composition is lyophilized, sterilization using this method may be conducted either prior to or following lyophilization and reconstitution. Compositions for parenteral administration can be stored in lyophilized form or in a solution. Parenteral compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

Once the pharmaceutical composition has been formulated, it may be stored in sterile vials as a solution, suspension, gel, emulsion, solid, crystal, or as a dehydrated or lyophilized powder. Such formulations may be stored either in a ready-to-use form or in a form (e.g., lyophilized) that is reconstituted prior to administration. The invention also provides kits for producing a single-dose administration unit. The kits of the invention may each contain both a first container having a dried protein and a second container having an aqueous formulation. In certain embodiments of this inven-

tion, kits containing single and multi-chambered pre-filled syringes (e.g., liquid syringes and lysosyringes) are provided.

The therapeutically effective amount of an IL-18 receptor antigen binding protein-containing pharmaceutical composition to be employed will depend, for example, upon the therapeutic context and objectives. One skilled in the art will appreciate that the appropriate dosage levels for treatment will vary depending, in part, upon the molecule delivered, the indication for which the IL-18 receptor antigen binding protein is being used, the route of administration, and the size (body weight, body surface or organ size) and/or condition (the age and general health) of the patient. In certain embodiments, the clinician may titer the dosage and modify the route of administration to obtain the optimal therapeutic effect. A typical dosage may range from about 0.1  $\mu\text{g}/\text{kg}$  to up to about 30 mg/kg or more, depending on the factors mentioned above. In specific embodiments, the dosage may range from 0.1  $\mu\text{g}/\text{kg}$  up to about 30 mg/kg, optionally from 1  $\mu\text{g}/\text{kg}$  up to about 30 mg/kg or from 10  $\mu\text{g}/\text{kg}$  up to about 5 mg/kg.

Dosing frequency will depend upon the pharmacokinetic parameters of the particular IL-18 receptor antigen binding protein in the formulation used. Typically, a clinician administers the composition until a dosage is reached that achieves the desired effect. The composition may therefore be administered as a single dose, or as two or more doses (which may or may not contain the same amount of the desired molecule) over time, or as a continuous infusion via an implantation device or catheter. Further refinement of the appropriate dosage is routinely made by those of ordinary skill in the art and is within the ambit of tasks routinely performed by them. Appropriate dosages may be ascertained through use of appropriate dose-response data. In certain embodiments, the antigen binding proteins of the invention can be administered to patients throughout an extended time period. Chronic administration of an antigen binding protein of the invention minimizes the adverse immune or allergic response commonly associated with antigen binding proteins that are not fully human, for example an antibody raised against a human antigen in a non-human animal, for example, a non-fully human antibody or non-human antibody produced in a non-human species.

The route of administration of the pharmaceutical composition is in accord with known methods, e.g., orally, through injection by intravenous, intraperitoneal, intracerebral (intraparenchymal), intracerebroventricular, intramuscular, intraocular, intraarterial, intraportal, or intralesional routes; by sustained release systems or by implantation devices. In certain embodiments, the compositions may be administered by bolus injection or continuously by infusion, or by implantation device.

The composition also may be administered locally via implantation of a membrane, sponge or another appropriate material onto which the desired molecule has been absorbed or encapsulated. In certain embodiments, where an implantation device is used, the device may be implanted into any suitable tissue or organ, and delivery of the desired molecule may be via diffusion, timed-release bolus, or continuous administration.

It also may be desirable to use IL-18 receptor antigen binding protein pharmaceutical compositions according to the invention ex vivo. In such instances, cells, tissues or organs that have been removed from the patient are exposed to IL-18 receptor antigen binding protein pharmaceutical compositions after which the cells, tissues and/or organs are subsequently implanted back into the patient.

In particular, IL-18 receptor antigen binding proteins can be delivered by implanting certain cells that have been geneti-

cally engineered, using methods such as those described herein, to express and secrete the polypeptide. In certain embodiments, such cells may be animal or human cells, and may be autologous, heterologous, or xenogeneic. In certain embodiments, the cells may be immortalized. In other embodiments, in order to decrease the chance of an immunological response, the cells may be encapsulated to avoid infiltration of surrounding tissues. In further embodiments, the encapsulation materials are typically biocompatible, semi-permeable polymeric enclosures or membranes that allow the release of the protein product(s) but prevent the destruction of the cells by the patient's immune system or by other detrimental factors from the surrounding tissues.

All references cited within the body of the instant specification are hereby expressly incorporated by reference in their entirety.

The following examples, including the experiments conducted and the results achieved, are provided for illustrative purposes only and are not to be construed as limiting the invention.

## EXAMPLES

### A. Example 1

#### Production of IgG2 and IgG4 Versions of Anti-IL-18 Receptor Antibodies Using pVE414N Transient Expression Constructs

The following example describes the generation of transient expression constructs used to produce IgG2 and IgG4 versions of various anti-IL18 receptor binding proteins, and experimental approaches to test their binding characteristics and potency.

##### 1. Generation of Constructs

Expression constructs for transient expression of IgG4 versions of AM<sub>H</sub>9/AM<sub>L</sub>9, AM<sub>H</sub>11/AM<sub>L</sub>7, AM<sub>H</sub>3/AM<sub>L</sub>14, and AM<sub>H</sub>6/AM<sub>L</sub>12 were generated by subcloning the polynucleotide sequences of SEQ ID NOs:74, 76, 78, 80, 82, 84, 86 and 88 into a transient expression vector. IgG2 versions of the same variable regions were generated by subcloning the polynucleotide portions encoding the variable regions of those IgGs into a separate transient expression vector.

##### 2. Transient Roller Bottle Transfections

Eight roller bottle transfections into CosPKB cell line for each of the antibodies were performed. The titers for the IgG2's were as follows:

IgG2	Titer
AMH9/AML9	33.5
AMH11/AML7	35.1
AMH3/AML14	42.9
AMH6/AML12	41.5

##### 3. Assays for Potency and Cross Reactivity of Various Antibodies

**KG-1 IFN $\gamma$  Release Assay.** The various IgG constructs were tested to determine their inhibitory activity in an in vitro Interferon- $\gamma$  (IFN $\gamma$ ) release assay. The IFN $\gamma$  release assay works on the principle that human myelomonocytic KG-1 cells that express the endogenous IL-18R release IFN $\gamma$  in response to IL-18.

Briefly, reagents such as affinity-purified scFv are pre-incubated with KG-1 cells in a 96-well tissue culture plate. IL-18 (+TNF $\alpha$ ) is added to the cells to induce IFN $\gamma$  release.

TNF $\alpha$  is added with the IL-18 to increase the IFN $\gamma$  response and therefore makes the assay more sensitive by allowing a lower concentration of IL-18 to be used. This is at least in part, likely due to TNF $\alpha$  induced upregulation of IL-18R surface expression.

After a defined incubation period, the cell supernatants are harvested and analyzed for IFN $\gamma$  content using ELISA. Test compounds that inhibit IL-18R mediated signaling can be assessed in this assay by showing a reduction in IFN $\gamma$  release.

KG-1 cells were obtained from the European Collection of Cell Cultures (ECACC, 86111306). The cells were propagated in supplemented Iscove's Dulbecco modified medium (IMDM), and were maintained at  $1-2 \times 10^6$  cells/ml. Recombinant human IL-18 was obtained from Peprotech (200-18) and the recombinant human TNF $\alpha$  was purchased from R&D Systems (210-TA). The amount of IFN $\gamma$  released in response to 1 nM IL-18 (+1.1 nM TNF $\alpha$ ) was monitored in each experiment, and ranged from approximately 250 to 4000 pg/ml.

The KG-1 assay was carried out in 96 flat bottom well cell culture plates (Costar). Test solutions of antibody (in duplicate) were used neat (or diluted to the required concentration in Dulbecco's PBS) in a volume of 50  $\mu$ l. The antibodies were then titrated in a 6-9 point  $\frac{1}{3}$  dilution series (using KG-1 medium), followed by the addition of 50  $\mu$ l KG-1 cells with gentle mixing. A "no antibody" control with only IL-18 and a "cells only" control were always included. After incubation of the antibody/cell mixture for 30-60 minutes at 37 $^\circ$  C. with 5% CO<sub>2</sub>, 100  $\mu$ l of IL-18+TNF $\alpha$  diluted in KG-1 medium was added with gentle mixing. The final concentration of IMAC-purified scFv typically ranged between 25-200  $\mu$ g/ml. Three reference inhibitors were used in all assays. The first two were monoclonal antibodies against the two different chains of the IL-18, RP1 (R&D systems, MAB840) and AcPL (R&D Systems, MAB1181). These antibodies were used as described above for scFv, except that the final starting concentration for the dilution series was 10-20  $\mu$ g/ml. In addition, a recombinant IL-18BP/Fc chimera (R&D Systems, 119-BP) was used to neutralize IL-18. In this case, 50  $\mu$ l of KG-1 medium was added to the cell culture plate, followed by 50  $\mu$ l of cells and the cells were incubated as above for the antibodies. In a separate 96 well "U" bottom cell culture plate, the IL-18BP/Fc was titrated in a 6-9 point  $\frac{1}{2}$  dilution series (using KG-1 medium) in a volume of 60  $\mu$ l/well. An equal volume of IL-18+TNF $\alpha$  was added to the IL-18BP/Fc dilution series. After incubation of the IL-18BP/Fc/IL-18 mixture for 30-60 minutes at 37 $^\circ$  C. with 5% CO<sub>2</sub>, 100  $\mu$ l/well was added to the KG-1 cell plate with gentle mixing. The final starting concentration of the IL-18BP/Fc for the dilution series was 1 mg/ml. The final concentration of the KG-1 cells was  $1.5 \times 10^6$  cells/ml (i.e.  $3 \times 10^6$ /well) and IL-18 was used at a final concentration of 1 nM (+1.1 nM TNF $\alpha$ ).

IL-18 was also titrated to determine the EC<sub>50</sub> of the IFN $\gamma$  response. The IL-18 was titrated in a 6-10 point  $\frac{1}{2}$  dilution series (using KG-1 medium), with the TNF $\alpha$  held constant at 1.1 nM. In this case 50 ml of KG-1 medium was added to the cell culture plate, followed by 50 ml of cells. The cell plate was incubated for 30-60 minutes at 37 $^\circ$  C. with 5% CO<sub>2</sub>, then 100 ml of the titrated IL-18 was added with gentle mixing. The final concentration of IL-18 started at 5 nM for the dilution series. Typically, the EC<sub>50</sub> for IL-18 (+TNF $\alpha$ ) was in the range of 0.5-1 nM, although minimum and maximum EC<sub>50</sub> values of down to 0.3 nM or up to 5 nM were occasionally seen.

The KG-1 cell plates were incubated at 37 $^\circ$  C. with 5% CO<sub>2</sub> overnight. The cell supernatants were harvested by removal of 180 ml medium into a clean "U" bottom 96-well

plate, which was then sealed and centrifuged at 1200 rpm for 5 minutes. 160 ml of the cell free supernatant was then transferred to another clean “U” bottom 96-well plate, and the clarified supernatants tested immediately or frozen at  $-20^{\circ}\text{C}$ .

The amount of IFN $\gamma$  in the KG-1 cell supernatants was determined by a standard sandwich ELISA based assay, using a time-resolved fluorometric readout. FLUORONUNC flat bottom 96-well plates (NUNC, 437958) were coated using 100 ml/well of the IFN $\gamma$  specific monoclonal capture antibody (R&D Systems, MAB2851) at 4 mg/ml and left at  $+4^{\circ}\text{C}$  overnight. The plates were rinsed with Dulbecco’s PBS, then non-specific protein binding was blocked by the addition of 200 ml/well of 3% milk powder in PBS and incubation at RT (RT) for 1-2 hours. The recombinant human IFN $\gamma$  standard (R&D Systems, 285-IF-100) was diluted to 16,000 pg/ml in reagent diluent (0.1% BSA, 0.05% Tween-20, 20 mM Tris, 150 mM NaCl; pH 7.2-7.4), then titrated in a 12 point  $\frac{1}{2}$  dilution series. The blocking buffer was removed from the capture antibody coated plates, and 100 ml/well of the IFN $\gamma$  standard or clarified cell supernatants was added. Reagent diluent was added for the “blank” control. After incubation for 1-2 hours at RT, the plates were washed 3 $\times$  with PBS containing 0.1% Tween-20. The biotinylated anti-human IFN $\gamma$  polyclonal detection antibody (R&D Systems, BAF285) was diluted to 100 ng/ml in reagent diluent supplemented with 2% normal goat serum, and 100 ml/well was added. After incubation for 1 hour at RT, the plates were washed 3 $\times$  with PBS containing 0.1% Tween-20. Streptavidin-Europium (Perkin-Elmer, 4001-0010) was diluted 1/1000 in DELFIA assay buffer (Perkin-Elmer, 4002-0010), and 100 ml/well was added. After 30-60 minutes incubation at RT, the plates were washed 7 $\times$  with DELFIA wash buffer (Perkin-Elmer, 1244-114). DELFIA Enhancement solution (Perkin-Elmer, 4001-0010) was added at 100 ml/well and the plates were left for at least 10 min at RT. The resulting fluorescent signal was measured using dissociation-enhanced time-resolved fluorometry using the Victor2 V plate reader (PerkinElmer).

The average value for the ELISA “blank” control was subtracted from the results for the IFN $\gamma$  standard, while the average value for the “cells alone” control was subtracted for the cell supernatant results. GraphPad PRISM (GraphPad Software, Inc.) was used to calculate the IFN $\gamma$  standard curve using non-linear regression (with a variable slope). The concentration of IFN $\gamma$  in the cell supernatants was then determined by using an “unknown X from Y” output for the IFN $\gamma$  standard curve. The EC50 for IFN $\gamma$  release from KG-1 cells in response to IL-18 was calculated using non-linear regression (with a variable slope) and constraining the top and bottom as necessary. Inhibition of IFN $\gamma$  release from KG-1 cells by test compound was normalised as a percentage of the average value of the “no antibody” IL-18 alone control, using the fluorescent counts data. The IC<sub>50</sub> values for test compounds could then be calculated using non-linear regression (with a variable slope), and constraining the bottom and top to 0 and 100% respectively.

The IgG2 versions of antibodies AM<sub>H</sub>9/AM<sub>L</sub>9, AM<sub>H</sub>11/AM<sub>L</sub>7, AM<sub>H</sub>3/AM<sub>L</sub>14, and AM<sub>H</sub>6/AM<sub>L</sub>12 have at least equivalent potency as the original IgG4 versions in an assay measuring their effect on the IFN- $\gamma$  production by KG1 cells. Furthermore, the IgG2 versions have at least equivalent potency as the original IgG4s in an assay measuring INF- $\gamma$  production by human NK cells.

The IgG2 versions of the antibodies have equivalent potency as the original IgG4s in an assay measuring their effect on IL-18 induced INF- $\gamma$  production by cynomolgus

PBMC#010182 cells. This confirms that the conversion to IgG2 did not affect the cyno cross reactivity of the tested antibodies.

#### 4. Assays for Specificity

IgGs of various antibodies were analyzed for cross-reactivity by ELISA against a panel of proteins.

Test antigens were coated onto Protein Immobiliser 96-well plates (Exiqon, Prd#10203-111-60) at 1  $\mu\text{g}/\text{ml}$  in PBS (Dulbecco’s w/o Ca and Mg, Invitrogen, Cat#14190-086) in duplicate, 50  $\mu\text{l}$  per well, overnight at  $4^{\circ}\text{C}$ .

Plates were washed three times with 300  $\mu\text{l}$  PBS-Tween (0.1%) (PBS-T) and three times with 300  $\mu\text{l}$  PBS per well using a 96-well plate washer (BIO-TEK, ELX405UV). To the washed plates, 300  $\mu\text{l}$  of 3% Marvel PBS (MPBS) was added per well as a blocking agent. Plates were blocked at room temperature (RT) for 1 hour.

Plates were washed three times with PBS-T and three times with PBS as previously stated.

Antibodies (huIgG<sub>4</sub>) were diluted to 0.5  $\mu\text{g}/\text{ml}$  in 3% MPBS. 50  $\mu\text{l}$  of diluted huIgG<sub>4</sub> were added per well. Plates were incubated at RT for 1 hour. Plates were washed as previously stated.

Primary detection antibody (Monoclonal anti-human IgG4 clone HP-6025 biotin conjugate, Sigma, Cat#B-3648) was diluted 1:15,000 in 3% MPBS and added to plates at 50  $\mu\text{l}$  per well. Incubation with primary detection antibody was at RT for 1 hour. Plates were washed previously stated.

Secondary detection antibody (ExtrAvidin peroxidase conjugate, Sigma, Cat#E-2886), was diluted 1:1,000 in 3% MPBS and added to plates at 50  $\mu\text{l}$  per well. Incubation with secondary detection antibody was at RT for 30 minutes. Plates were washed previously stated.

50  $\mu\text{l}$  per well Tetramethyl-benzidine (TMB) (Liquid substrate for ELISA, Sigma, Cat#T-0440) was added and incubated at RT for 10 minutes. To stop the enzyme color reaction, 50  $\mu\text{l}$  0.5 M H<sub>2</sub>SO<sub>4</sub> per well was added.

Plates were read at 450 nm on a 96-well plate reader (Victor<sup>2</sup> V plate reader (PerkinElmer).

The specific anti-IL-18 receptor IgG4 antibodies were positive against human and cynomolgus IL-18 receptor protein only. There was no cross-reactivity with other species. An IgG2 version of an above antibody had the same cross-reactivity properties, i.e., it cross-reacted with cynomolgus IL-18 receptor only.

#### B. Example 2

##### Characterization of the Binding Affinity of an IL-18 Receptor Antibody

This Example provides an exemplary method of determining the binding affinity of an IL-18 receptor antigen binding protein to cell surface-expressed human IL-18R $\alpha$ . An IL-18 receptor antibody was iodinated using [<sup>125</sup>I]-Bolton-Hunter Reagent (diiodinated; PerkinElmer Life Sciences, Inc., Boston, Mass.). One millicurie of [<sup>125</sup>I]-Bolton-Hunter Reagent supplied in anhydrous benzene was evaporated to dryness under a nitrogen stream. Five microliters of IL-18 receptor antibody (16 micrograms) was diluted with an equal volume of borate buffered saline and then added to the dried [<sup>125</sup>I]-Bolton-Hunter Reagent in its original vial. After an overnight incubation at  $4^{\circ}\text{C}$ ., 10 microliters of PBS, 0.1% gelatin was added and the entire sample transferred to an equilibrated 2 mL P6 column (BioRad; Hercules, Calif.) where iodinated IL-18 receptor antibody was separated from free <sup>125</sup>I by gel filtration with 0.1% gelatin as a carrier protein. Fractions containing iodinated antibody were pooled, diluted to a con-

centration of 100 nM in binding media (RPMI 1640 containing 2.5% bovine albumin, Fraction V, 20 mM Hepes, and 0.2% sodium azide, pH 7.2), and stored at 4° C. A specific activity of  $3.5 \times 10^{15}$  cpm/mmol was calculated based on an initial protein concentration of antibody by amino acid analysis, and a recovery of 70% from a control experiment in which an aliquot of iodinated antibody was put through the iodination protocol with omission of [ $^{125}$ I]-Bolton-Hunter Reagent.

#### 1. Direct Equilibrium Binding

KG-1 cells were stimulated for 20 hours at 37° C. in 5% CO<sub>2</sub> in IMDM medium containing 20% fetal calf serum in the presence of 20 ng/mL of human TNF $\alpha$ . The stimulated KG-1 cells ( $5 \times 10^5$  cells/150 microliters final) were washed twice with PBS, and then incubated with a range of concentrations of iodinated antibody. Nonspecific binding was measured at a single concentration of iodinated antibody (~350 pM, in triplicate) in the presence of a 1000-fold molar excess of unlabeled antibody, and assumed to be a linear function of the concentration of iodinated antibody present. All reagents were diluted in binding media containing sodium azide (0.2%) to inhibit potential internalization of iodinated antibody by the cells.

Cells were incubated in 96-well round-bottom microtiter plates at 37° C., 5% CO<sub>2</sub> on a miniorbital shaker. After 4 hours, two 60 microliter aliquots of each mixture were transferred to chilled 400 microliter polyethylene centrifuge tubes containing 200 microliters phthalate oil (1.5 parts dibutylphthalate: 1 part dioctylphthalate) and spun for 1.5 minutes in a 4° C. tabletop microfuge (Sorvall, Asheville, N.C.) at 10,000 rpm to separate cell associated iodinated antibody from free iodinated antibody. The oil tubes were cut, and each cell pellet and supernatant was collected in individual 12x75 mm glass tubes and loaded on a COBRA gamma counter (Packard Instrument Company; Boston, Mass.) for cpm measurements. Cpm from duplicate aliquots for each well were averaged for analysis. Data were fit to a simple 1-site binding equation via nonlinear regression in Prism version 3.03 (GraphPad Software, Inc; San Diego, Calif.).

The iodinated antibody bound to stimulated KG-1 cells at 37° C. with a  $K_D$  of 81 pM and ~4700 sites/cell.

#### 2. Competition Assay

Stimulated and washed KG-1 cells ( $5 \times 10^5$  cells/150 microliters final) were incubated with a single concentration of iodinated antibody (15.6 pM) and varying concentrations of unlabeled antibody in binding media. Nonspecific binding was determined in the presence of a 1000-fold molar excess of unlabeled antibody. Iodinated and unlabeled antibody were mixed just prior to the addition of cells, i.e., there was no pre-incubation step.

Cells were incubated in 96-well round-bottom microtiter plates at 37° C., 5% CO<sub>2</sub> on a miniorbital shaker. After 4 hours, two 60 microliter aliquots of each mixture were transferred to chilled 400 microliter polyethylene centrifuge tubes containing 200 microliters phthalate oil (1.5 parts dibutylphthalate: 1 part dioctylphthalate) and spun for 1.5 minutes in a 4° C. tabletop microfuge (Sorvall, Asheville, N.C.) at 10,000 rpm to separate cell associated iodinated antibody from free iodinated antibody. The oil tubes were cut, and each cell pellet and supernatant was collected in individual 12x75 mm glass tubes and loaded on a COBRA gamma counter (Packard Instrument Company; Boston, Mass.) for cpm measurements. Cpm from duplicate aliquots for each well were averaged for analysis. Data were fit to a single competitive inhibition equation via nonlinear regression using the  $K_d$  value of 81 pM in Prism.

The  $K_d$  of unlabeled antibody binding was 53 pM.

### C. Example 3

#### Characterization of the Activity of Anti-IL-18 Receptor Antibodies

A IFN $\gamma$  release assay was performed as described in Example 1, Section 3 for various IgG constructs, as described below.

#### 1. Comparison of the Inhibition of INF $\gamma$ Release by KG1 Cells with AM<sub>H</sub>2/AM<sub>L</sub>16, AM<sub>H</sub>2/AM<sub>L</sub>17, AM<sub>H</sub>1/AM<sub>L</sub>16, and AM<sub>H</sub>1/AM<sub>L</sub>17 Constructs

Inhibition of INF- $\gamma$  release by KG1 cells when treated with AM<sub>H</sub>/AM<sub>L</sub> antigen binding protein constructs AM<sub>H</sub>2/AM<sub>L</sub>16, AM<sub>H</sub>2/AM<sub>L</sub>17, AM<sub>H</sub>1/AM<sub>L</sub>16, and AM<sub>H</sub>1/AM<sub>L</sub>17 was compared with control treatment with IL-18 binding protein (BP). The AM<sub>H</sub>/AM<sub>L</sub> antigen binding proteins were significantly more efficacious at inhibiting IFN $\gamma$  release, all demonstrating an ED<sub>50</sub> between 6 and 10 pM compared to an ED<sub>50</sub> for IL-18 BP of 520 pM.

#### 2. Comparison of the Inhibition of INF $\gamma$ Release by KG1 Cells with AM<sub>H</sub>4/AM<sub>L</sub>14, AM<sub>H</sub>4/AM<sub>L</sub>15, AM<sub>H</sub>3/AM<sub>L</sub>14, and AM<sub>H</sub>3/AM<sub>L</sub>15 Constructs

Inhibition of INF- $\gamma$  release by KG1 cells when treated with AM<sub>H</sub>/AM<sub>L</sub> antigen binding protein constructs AM<sub>H</sub>4/AM<sub>L</sub>14, AM<sub>H</sub>4/AM<sub>L</sub>15, AM<sub>H</sub>3/AM<sub>L</sub>14, and AM<sub>H</sub>3/AM<sub>L</sub>15 was compared with control treatment with IL-18 binding protein (BP). The AM<sub>H</sub>/AM<sub>L</sub> antigen binding proteins were significantly more efficacious at inhibiting IFN $\gamma$  release, all demonstrating an ED<sub>50</sub> between 3 and 7 pM compared to an ED<sub>50</sub> for IL-18 BP of 520 pM.

#### 3. Comparison of the Inhibition Of INF $\gamma$ Release by KG1 Cells with AM<sub>H</sub>6/AM<sub>L</sub>12, AM<sub>H</sub>6/AM<sub>L</sub>13, AM<sub>H</sub>5/AM<sub>L</sub>12, and AM<sub>H</sub>5/AM<sub>L</sub>13 Constructs

Inhibition of INF- $\gamma$  release by KG1 cells when treated with AM<sub>H</sub>/AM<sub>L</sub> antigen binding protein constructs AM<sub>H</sub>6/AM<sub>L</sub>12, AM<sub>H</sub>6/AM<sub>L</sub>13, AM<sub>H</sub>5/AM<sub>L</sub>12, and AM<sub>H</sub>5/AM<sub>L</sub>13 was compared with control treatment with IL-18 binding protein (BP). The AM<sub>H</sub>/AM<sub>L</sub> antigen binding proteins were significantly more efficacious at inhibiting IFN $\gamma$  release, all demonstrating an ED<sub>50</sub> between 2.9 and 11.3 pM compared to an ED<sub>50</sub> for IL-18 BP of 520 pM.

#### 4. Inhibition of INF $\gamma$ Release by KG1 Cells by AM<sub>H</sub>8/AM<sub>L</sub>11, AM<sub>H</sub>9/AM<sub>L</sub>9, AM<sub>H</sub>10/AM<sub>L</sub>8, and AM<sub>H</sub>11/AM<sub>L</sub>7 IgGs

Inhibition of INF- $\gamma$  release by KG1 cells with AM<sub>H</sub>/AM<sub>L</sub> IgG antigen binding protein constructs comprising combinations of AM<sub>H</sub>8/AM<sub>L</sub>11, AM<sub>H</sub>9/AM<sub>L</sub>9, AM<sub>H</sub>10/AM<sub>L</sub>8, and AM<sub>H</sub>11/AM<sub>L</sub>7 IgGs was compared with control treatment with IL-18 binding protein (BP). The AM<sub>H</sub>/AM<sub>L</sub> antigen binding proteins were significantly more efficacious at inhibiting IFN $\gamma$  release, all demonstrating an ED<sub>50</sub> between 3 and 45 pM compared to an ED<sub>50</sub> for IL-18 BP of 520 pM.

#### 5. Inhibition of INF $\gamma$ Release by KG1 Cells by AM<sub>H</sub>15/AM<sub>L</sub>3, AM<sub>H</sub>13/AM<sub>L</sub>4, AM<sub>H</sub>13/AM<sub>L</sub>5, and AM<sub>H</sub>16/AM<sub>L</sub>2 IgGs

Inhibition of INF- $\gamma$  release by KG1 cells with AM<sub>H</sub>/AM<sub>L</sub> IgG antigen binding protein constructs comprising combinations of AM<sub>H</sub>15/AM<sub>L</sub>3, AM<sub>H</sub>13/AM<sub>L</sub>4, AM<sub>H</sub>13/AM<sub>L</sub>5, and AM<sub>H</sub>16/AM<sub>L</sub>2 IgGs was compared with control treatment with IL-18 binding protein (BP). The AM<sub>H</sub>/AM<sub>L</sub> antigen binding proteins were significantly more efficacious at inhib-

iting IFN $\gamma$  release, all demonstrating an ED<sub>50</sub> between 17 and 320 pM compared to an ED<sub>50</sub> for IL-18 BP of 520 pM.

#### D. Example 4

##### Identification of the Binding Epitope of the Described IL-18 Receptor Antibodies

Experiments were carried out to determine the amino acids residues present in human IL-18 receptor (IL-18R) that are important for binding to one or more of the IL-18 receptor binding proteins. An exemplary antibody bound with high affinity to human IL-18R but did not bind the mouse ortholog of IL-18R. Experiments were directed towards examining the amino acids that differ between human and mouse IL-18R. This was coupled with analysis of a three dimensional computational model of the IL-18R to determine which of those amino acids lie on the surface of the receptor and are therefore more likely to interact with the antibody. Candidate amino acids were examined for their contribution to antibody recognition by using site-specific mutagenesis to change the particular amino acids from the human sequence to mouse sequence and then testing the mutant IL-18R molecules for

binding to antibody. The relative ability of antibody to bind the different mutations was assessed using antibody dose-response curves and subsequent determination of EC<sub>50</sub> concentrations (the concentration of antibody required for 50% of the maximal binding signal).

A region on the surface of human IL-18R was identified that is particularly important for antibody binding and thus contributes to the epitope. This region contains amino acids 243-271 (shown in bold in FIG. 5). When specific amino acids in this region were mutated to mouse sequence, antibody binding was diminished (effects on antibody binding reported in TABLE 3). Amino acids 250-253 (MFGE mutant) and 267-271 (MRIMT mutation) had the most profound influence on antibody binding however neither completely influenced antibody binding alone (see, underlined amino acid residues). When the receptor was mutated at all four of the specific sites tested, antibody binding was virtually abolished. Binding of a control anti-IL-18R receptor antibody was not significantly affected by the mutations indicating that the overall structure of the receptor had not been disrupted by the mutations. Amino acids 243-271 encode one of the predicted IL-18 contact points and thus this epitope is consistent with the antibody's ability to block IL-18 binding to the receptor.

TABLE 3

Results of Antibody Binding Assay with IL-18R and Mutated IL-18R (the mouse amino acids corresponding to each mutation is provided in parenthesis)			
	Exemplary Ab binding: avg. EC50 (pM)	Fold decrease in binding ability relative to huIL18R	Control Ab binding: avg. EC50 (pM)
huIL18R	15.7 +/- 9.0	0	7.9 +/- 4.7
EEDV mutation (KDDL)	25.3 +/- 11.9	1.6	6.7 +/- 4.4
MFGE mutation (SIRK)	57.9 +/- 26.7	3.7	5.9 +/- 3.7
MRIMT mutation (TTTWI)	177.5 +/- 58.9	11.3	8.5 +/- 5.1
STGGT (NEEAI)	15.7 +/- 8.5	0	10.0 +/- 6.2
EEDV + MRIMT + STGGT	2615.1	2287.3	7.9 +/- 5.1
EEDV + MRIMT + MFGE + STGGT	too low to measure	N/A	10.3 +/- 5.5
Mouse IL18R	does not bind	N/A	N/A

Human IL18R was mutated to mouse IL18R sequence at the indicated residues and recombinant mutated receptors with and avidin tag were immobilized on biotin-coated beads. Immobilized receptor was then used to determine relative antibody binding ability in a soluble binding assay. Binding was also performed using an anti-huIL-18R antibody that recognizes a distinct epitope from the exemplary antibody. Binding experiments were all performed at least two times. The average EC50 represents the concentration of antibody required to achieve 50% of maximal binding.

#### SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 292

<210> SEQ ID NO 1

<211> LENGTH: 122

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

Gln Val Gln Leu Val Gln Ser Gly Gly Gly Val Val Gln Pro Gly Arg  
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Gly Tyr  
20 25 30

-continued

---

Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
                   35                                  40                                  45

Ala Val Ile Ser Asn Asp Gly Ser Lys Lys Tyr Tyr Ser Asp Ser Val  
           50                                  55                                  60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
   65                                  70                                  75                                  80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
                                   85                                  90                                  95

Ala Lys Gly Ser Ser Ser Ile Trp Leu Thr Gln Ser Leu Asp His Trp  
                                   100                                  105                                  110

Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
           115                                  120

<210> SEQ ID NO 2  
 <211> LENGTH: 122  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg  
 1                                  5                                  10                                  15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Gly Tyr  
           20                                  25                                  30

Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
           35                                  40                                  45

Ala Val Ile Ser Asn Asp Gly Ser Lys Lys Tyr Tyr Ser Asp Ser Val  
           50                                  55                                  60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
   65                                  70                                  75                                  80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
                                   85                                  90                                  95

Ala Lys Gly Ser Ser Ser Ile Trp Leu Thr Gln Ser Leu Asp His Trp  
                                   100                                  105                                  110

Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
           115                                  120

<210> SEQ ID NO 3  
 <211> LENGTH: 122  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

Gln Val Gln Leu Val Gln Ser Gly Gly Gly Val Val Gln Pro Gly Arg  
 1                                  5                                  10                                  15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Gly Tyr  
           20                                  25                                  30

Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
           35                                  40                                  45

Ala Val Ile Ser Asn Asp Gly Ser Lys Lys Tyr Tyr Ser Asp Ser Val  
           50                                  55                                  60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
   65                                  70                                  75                                  80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
                                   85                                  90                                  95

Ala Lys Gly Ser Ser Ser Ile Trp Leu Ser Gln Ser Leu Asp Gly Trp  
                                   100                                  105                                  110

-continued

Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 115 120

<210> SEQ ID NO 4  
 <211> LENGTH: 122  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg  
 1 5 10 15  
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Gly Tyr  
 20 25 30  
 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
 35 40 45  
 Ala Val Ile Ser Asn Asp Gly Ser Lys Lys Tyr Tyr Ser Asp Ser Val  
 50 55 60  
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
 65 70 75 80  
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
 85 90 95  
 Ala Lys Gly Ser Ser Ser Ile Trp Leu Ser Gln Ser Leu Asp Gly Trp  
 100 105 110  
 Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 115 120

<210> SEQ ID NO 5  
 <211> LENGTH: 122  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

Gln Val Gln Leu Val Gln Ser Gly Gly Gly Val Val Gln Pro Gly Arg  
 1 5 10 15  
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Gly Tyr  
 20 25 30  
 Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
 35 40 45  
 Ala Val Ile Ser Asn Asp Gly Ser Lys Lys Tyr Tyr Ser Asp Ser Val  
 50 55 60  
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
 65 70 75 80  
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
 85 90 95  
 Ala Lys Gly Ser Ser Ser Ile Trp Leu Thr Ser Ala Leu Asn Leu Trp  
 100 105 110  
 Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 115 120

<210> SEQ ID NO 6  
 <211> LENGTH: 122  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg  
 1 5 10 15  
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Gly Tyr  
 20 25 30

-continued

---

Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
                   35                                  40                                  45

Ala Val Ile Ser Asn Asp Gly Ser Lys Lys Tyr Tyr Ser Asp Ser Val  
           50                                  55                                  60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
  65                                  70                                  75                                  80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
                                   85                                  90                                  95

Ala Lys Gly Ser Ser Ser Ile Trp Leu Thr Ser Ala Leu Asn Leu Trp  
                                   100                                  105                                  110

Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
           115                                  120

<210> SEQ ID NO 7  
 <211> LENGTH: 117  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln Pro Gly Arg  
 1                                  5                                  10                                  15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Gly Tyr  
           20                                  25                                  30

Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
           35                                  40                                  45

Ala Val Ile Ser Asn Asp Gly Ser Lys Lys Tyr Tyr Ser Asp Ser Val  
           50                                  55                                  60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
  65                                  70                                  75                                  80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Ile Tyr Tyr Cys  
                                   85                                  90                                  95

Ala Lys Gly Ser Ser Ser Ile Trp Phe Gly Glu Thr Val Asp Tyr Trp  
                                   100                                  105                                  110

Gly Gln Gly Thr Thr  
           115

<210> SEQ ID NO 8  
 <211> LENGTH: 121  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala  
 1                                  5                                  10                                  15

Ser Val Lys Val Ser Cys Lys Val Ser Gly Tyr Thr Leu Thr Glu Leu  
           20                                  25                                  30

Ser Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Met  
           35                                  40                                  45

Gly Gly Phe Asp Arg Glu Asp Asp Glu Thr Ile His Ala Gln Lys Phe  
           50                                  55                                  60

Gln Gly Arg Val Thr Met Thr Glu Asp Thr Ser Thr Asp Thr Ala Tyr  
  65                                  70                                  75                                  80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys  
                                   85                                  90                                  95

Ala Thr Asp Leu Met Val Trp Gly Asp Phe Trp Ile Gln His Trp Gly  
           100                                  105                                  110



-continued

Gln Gly Thr Leu Val Thr Val Ser Ser  
 115 120

<210> SEQ ID NO 9  
 <211> LENGTH: 121  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala  
 1 5 10 15  
 Ser Val Lys Val Ser Cys Lys Val Ser Gly Tyr Thr Leu Thr Glu Leu  
 20 25 30  
 Ser Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Met  
 35 40 45  
 Gly Gly Phe Asp Arg Glu Asp Asp Glu Thr Ile His Ala Gln Lys Phe  
 50 55 60  
 Gln Gly Arg Val Thr Met Thr Glu Asp Thr Ser Thr Asp Thr Ala Tyr  
 65 70 75 80  
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys  
 85 90 95  
 Ala Thr Asp Leu Met Val Trp Gly Asp Phe Trp Ile Gln His Trp Gly  
 100 105 110  
 Gln Gly Thr Leu Val Thr Val Ser Ser  
 115 120

<210> SEQ ID NO 10  
 <211> LENGTH: 121  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala  
 1 5 10 15  
 Ser Val Lys Val Ser Cys Lys Val Ser Gly Tyr Thr Leu Thr Glu Leu  
 20 25 30  
 Ser Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Met  
 35 40 45  
 Gly Gly Phe Asp Arg Glu Asp Asp Glu Thr Ile His Ala Gln Lys Phe  
 50 55 60  
 Gln Gly Arg Val Thr Met Thr Glu Asp Thr Ser Thr Asp Thr Ala Tyr  
 65 70 75 80  
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys  
 85 90 95  
 Ala Thr Asp Leu Met Ala Trp Asp Tyr Pro Pro Ile Gln His Trp Gly  
 100 105 110  
 Gln Gly Thr Leu Val Thr Val Ser Ser  
 115 120

<210> SEQ ID NO 11  
 <211> LENGTH: 121  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 11

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala  
 1 5 10 15  
 Ser Val Lys Val Ser Cys Lys Val Ser Gly Tyr Thr Leu Thr Glu Leu  
 20 25 30

-continued

---

Ser Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Met  
                   35                                  40  45

Gly Gly Phe Asp Arg Glu Asp Asp Glu Thr Ile His Ala Gln Lys Phe  
       50  55  60

Gln Gly Arg Val Thr Met Thr Glu Asp Thr Ser Thr Asp Thr Ala Tyr  
  65                                  70                                  75                                  80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys  
                                   85                                  90                                  95

Ala Thr Asp Leu Met Val Trp Asn Phe Pro Pro Ile Gln His Trp Gly  
                   100                                  105                                  110

Gln Gly Thr Leu Val Thr Val Ser Ser  
       115                                  120

<210> SEQ ID NO 12  
 <211> LENGTH: 116  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 12

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala  
 1                  5                                  10                                  15

Ser Val Lys Val Ser Cys Lys Val Ser Gly Tyr Thr Leu Thr Glu Leu  
                   20                                  25                                  30

Ser Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Met  
                   35                                  40                                  45

Gly Gly Phe Asp Arg Glu Asp Asp Glu Thr Ile His Ala Gln Lys Phe  
       50  55  60

Gln Gly Arg Val Thr Met Thr Glu Asp Thr Ser Thr Asp Thr Ala Tyr  
  65                                  70                                  75                                  80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys  
                                   85                                  90                                  95

Ala Thr Asp Leu Met Val Trp Gly Asp Phe Trp Ile Gln His Trp Gly  
                   100                                  105                                  110

Lys Gly Thr Met  
       115

<210> SEQ ID NO 13  
 <211> LENGTH: 118  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 13

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly  
 1                  5                                  10                                  15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr  
                   20                                  25                                  30

Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
                   35                                  40                                  45

Ser Ala Ile Ser Gly Ser Gly Gly Gly Thr Tyr Tyr Ala Asp Ser Val  
       50                                  55                                  60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
  65                                  70                                  75                                  80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
                                   85                                  90                                  95

Ala Arg Ile Arg Gly Asp Tyr Arg Thr Asp Ile Trp Gly Gln Gly Thr  
                   100                                  105                                  110

-continued

Thr Val Thr Val Ser Ser  
115

<210> SEQ ID NO 14  
<211> LENGTH: 118  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 14

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly  
1 5 10 15  
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr  
20 25 30  
Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
35 40 45  
Ser Ala Ile Ser Gly Ser Gly Gly Gly Thr Tyr Tyr Ala Asp Ser Val  
50 55 60  
Lys Gly Arg Ser Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
65 70 75 80  
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
85 90 95  
Ala Arg Ile Arg Gly Asp Tyr Arg Thr Asp Ile Trp Gly Arg Gly Thr  
100 105 110  
Leu Val Thr Val Ser Ser  
115

<210> SEQ ID NO 15  
<211> LENGTH: 118  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 15

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Ala Gln Pro Gly Gly  
1 5 10 15  
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr  
20 25 30  
Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
35 40 45  
Ser Ala Ile Ser Gly Ser Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val  
50 55 60  
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
65 70 75 80  
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
85 90 95  
Ala Arg Val Arg Gly Asp Tyr Arg Thr Asp Ile Trp Gly Arg Gly Thr  
100 105 110  
Leu Val Thr Val Ser Ser  
115

<210> SEQ ID NO 16  
<211> LENGTH: 118  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 16

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly  
1 5 10 15  
Ser Leu Arg Leu Ser Cys Ala Ala Ser Arg Phe Thr Phe Ser Ser Tyr  
20 25 30

-continued

Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
                   35                                  40                                  45

Ser Ala Ile Ser Gly Ser Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val  
   50  55                                  60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
  65                                  70                                  75                                  80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
                                   85                                  90                                  95

Ala Arg Val Arg Gly Asp Tyr Arg Thr Asp Ile Trp Gly Arg Gly Thr  
                           100                                  105                                  110

Leu Val Thr Val Ser Ser  
           115

<210> SEQ ID NO 17  
 <211> LENGTH: 113  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 17

Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly  
 1                                  5                                  10                                  15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Arg Phe Thr Phe Ser Ser Tyr  
                           20                                  25                                  30

Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val  
                   35                                  40                                  45

Ser Ala Ile Ser Gly Ser Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val  
   50  55                                  60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr  
  65                                  70                                  75                                  80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys  
                                   85                                  90                                  95

Ala Arg Val Arg Gly Ile Tyr Gly Met Asp Val Trp Gly Arg Gly Thr  
                           100                                  105                                  110

Leu

<210> SEQ ID NO 18  
 <211> LENGTH: 103  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 18

Gln Pro Val Leu Thr Gln Pro Pro Ser Val Ser Val Ser Pro Gly Gln  
 1                                  5                                  10                                  15

Thr Ala Ser Ile Thr Cys Ser Gly Asp Lys Leu Gly Asp Lys Tyr Ala  
                           20                                  25                                  30

Ser Trp Tyr Gln Gln Lys Pro Gly Lys Ser Pro Val Leu Val Ile Tyr  
                   35                                  40                                  45

Gln Asp Ser Asn Arg Pro Ser Gly Ile Pro Glu Arg Phe Ser Gly Ser  
   50                                  55                                  60

Asn Ser Gly Asn Thr Ala Thr Leu Thr Ile Ser Gly Thr Gln Ala Arg  
  65                                  70                                  75                                  80

Asp Glu Ala Asp Tyr Tyr Cys Gln Ala Trp Asp Ser Ser Thr Ala Ser  
                           85                                  90                                  95

Val Phe Gly Gly Gly Thr Lys  
           100

-continued

<210> SEQ ID NO 19  
 <211> LENGTH: 108  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 19

Gln Pro Val Leu Thr Gln Pro Pro Ser Val Ser Val Ser Pro Gly Gln  
 1 5 10 15  
 Thr Ala Ser Ile Thr Cys Ser Gly Asp Lys Leu Gly Asp Lys Tyr Ala  
 20 25 30  
 Ser Trp Tyr Gln Gln Lys Pro Gly Lys Ser Pro Val Leu Val Ile Tyr  
 35 40 45  
 Gln Asp Ser Asn Arg Pro Ser Gly Ile Pro Glu Arg Phe Ser Gly Ser  
 50 55 60  
 Asn Ser Gly Asn Thr Ala Thr Leu Thr Ile Ser Gly Thr Gln Ala Arg  
 65 70 75 80  
 Asp Glu Ala Asp Tyr Tyr Cys Gln Ala Trp Asp His Ser Leu Gln His  
 85 90 95  
 Arg Phe Gly Gly Gly Thr Lys Val Thr Val Leu Gly  
 100 105

<210> SEQ ID NO 20  
 <211> LENGTH: 108  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 20

Gln Pro Val Leu Thr Gln Pro Pro Ser Val Ser Val Ser Pro Gly Gln  
 1 5 10 15  
 Thr Ala Ser Ile Thr Cys Ser Gly Asp Lys Leu Gly Asp Lys Tyr Ala  
 20 25 30  
 Ser Trp Tyr Gln Gln Lys Pro Gly Gln Thr Pro Val Leu Val Ile Tyr  
 35 40 45  
 Gln Asp Ser Asn Arg Pro Ser Gly Ile Pro Glu Arg Phe Ser Gly Ser  
 50 55 60  
 Asn Ser Gly Asn Thr Ala Thr Leu Thr Ile Ser Gly Thr Gln Ala Arg  
 65 70 75 80  
 Asp Glu Ala Asp Tyr Tyr Cys Gln Ala Trp Thr Ser Ala Leu Asn Ser  
 85 90 95  
 Gln Phe Gly Gly Gly Thr Lys Val Thr Val Leu Gly  
 100 105

<210> SEQ ID NO 21  
 <211> LENGTH: 108  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 21

Gln Pro Val Leu Thr Gln Pro Pro Ser Val Ser Val Ser Pro Gly Gln  
 1 5 10 15  
 Thr Ala Ser Ile Thr Cys Ser Gly Asp Lys Leu Gly Asp Lys Tyr Ala  
 20 25 30  
 Ser Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Val Leu Val Ile Tyr  
 35 40 45  
 Gln Asp Ser Asn Arg Pro Ser Gly Ile Pro Glu Arg Phe Ser Gly Ser  
 50 55 60  
 Asn Ser Gly Asp Thr Ala Thr Leu Thr Ile Ser Gly Thr Gln Ala Arg  
 65 70 75 80

-continued

Asp Glu Ala Asp Tyr Tyr Cys Gln Ala Trp Thr His Ser Leu Ser Thr  
85 90 95

Leu Phe Gly Gly Gly Thr Lys Val Thr Val Leu Gly  
100 105

<210> SEQ ID NO 22  
<211> LENGTH: 108  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 22

Ser Tyr Glu Leu Thr Gln Pro Pro Ser Val Ser Val Ser Pro Gly Gln  
1 5 10 15

Thr Ala Ser Ile Thr Cys Ser Gly Asp Lys Leu Gly Asp Lys Tyr Ala  
20 25 30

Ser Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Val Leu Val Ile Tyr  
35 40 45

Gln Asp Ser Asn Arg Pro Ser Gly Ile Pro Glu Arg Phe Ser Gly Ser  
50 55 60

Asn Ser Gly Asn Thr Ala Thr Leu Thr Ile Ser Gly Thr Gln Ala Met  
65 70 75 80

Asp Glu Ala Asp Tyr Tyr Cys Gln Ala Trp Thr His Ser Leu Ser Thr  
85 90 95

Leu Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly  
100 105

<210> SEQ ID NO 23  
<211> LENGTH: 106  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 23

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln  
1 5 10 15

Arg Val Thr Ile Ser Cys Ser Gly Arg Asn Ser Asn Ile Gly Ser Tyr  
20 25 30

Thr Val Thr Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu  
35 40 45

Ile Tyr Ser Asn Ser Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser  
50 55 60

Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Gln  
65 70 75 80

Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Ser Leu  
85 90 95

Asn Gly Pro Val Phe Gly Gly Gly Thr Lys  
100 105

<210> SEQ ID NO 24  
<211> LENGTH: 111  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 24

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln  
1 5 10 15

Arg Val Thr Ile Ser Cys Ser Gly Arg Asn Ser Asn Ile Gly Ser Tyr  
20 25 30

Thr Val Thr Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu  
35 40 45

-continued

---

Ile Tyr Ser Asn Ser Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser  
 50 55 60

Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Gln  
 65 70 75 80

Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Val Val Trp Asp Asp Val Leu  
 85 90 95

Asn Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly  
 100 105 110

<210> SEQ ID NO 25  
 <211> LENGTH: 111  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 25

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln  
 1 5 10 15

Arg Val Thr Ile Ser Cys Ser Gly Arg Asn Ser Asn Ile Gly Ser Tyr  
 20 25 30

Thr Val Thr Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu  
 35 40 45

Ile Tyr Ser Asn Ser Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser  
 50 55 60

Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Gln  
 65 70 75 80

Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Val Val Trp Asp Asp Lys Leu  
 85 90 95

Asn Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly  
 100 105 110

<210> SEQ ID NO 26  
 <211> LENGTH: 111  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 26

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln  
 1 5 10 15

Arg Val Thr Ile Ser Cys Ser Gly Arg Asn Ser Asn Ile Gly Ser Tyr  
 20 25 30

Thr Val Thr Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu  
 35 40 45

Ile Tyr Ser Asn Ser Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser  
 50 55 60

Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Gln  
 65 70 75 80

Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Val Val Trp Asp Glu Ile Leu  
 85 90 95

Asn Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly  
 100 105 110

<210> SEQ ID NO 27  
 <211> LENGTH: 103  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 27

Ser Ser Glu Leu Thr Gln Asp Pro Ala Val Ser Val Ala Leu Gly Gln

-continued

---

```

1           5           10           15
Thr Val Arg Ile Thr Cys Gln Gly Asp Ser Leu Arg Ser Tyr Tyr Ala
                20                25                30
Ser Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Val Leu Val Ile Ser
                35                40                45
Ala Lys Asn Asn Arg Pro Ser Gly Ile Pro Asp Arg Phe Ser Gly Ser
                50                55                60
Ser Ser Gly Asn Thr Ala Ser Leu Thr Ile Thr Gly Ala Gln Ala Glu
65                70                75                80
Asp Glu Ala Asp Tyr Tyr Cys Asn Ser Arg Asp Ser Ser Asn His Val
                85                90                95
Val Phe Gly Gly Gly Thr Lys
                100

```

```

<210> SEQ ID NO 28
<211> LENGTH: 111
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 28

```

```

Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
1           5           10           15
Arg Val Thr Ile Ser Cys Ser Gly Arg Asn Ser Asn Ile Gly Ser Tyr
                20                25                30
Thr Val Thr Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
35                40                45
Ile Tyr Ser Asn Ser Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
50                55                60
Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Gln
65                70                75                80
Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Leu Val Trp Asp Asp Val Leu
85                90                95
Asn Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
100                105                110

```

```

<210> SEQ ID NO 29
<211> LENGTH: 108
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 29

```

```

Ser Ser Glu Leu Thr Gln Asp Pro Ala Val Ser Val Ala Leu Gly Gln
1           5           10           15
Thr Val Arg Ile Thr Cys Gln Gly Asp Ser Leu Arg Ser Tyr Tyr Ala
20                25                30
Ser Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Val Leu Val Ile Ser
35                40                45
Ala Lys Asn Asn Arg Pro Ser Gly Ile Pro Asp Arg Phe Ser Gly Ser
50                55                60
Ser Ser Gly Asn Thr Ala Ser Leu Thr Ile Thr Gly Ala Gln Ala Glu
65                70                75                80
Asp Glu Ala Asp Tyr Tyr Cys Ala Ser Arg Asn Gly Trp Asn His Val
85                90                95
Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
100                105

```

```

<210> SEQ ID NO 30

```



-continued

---

<211> LENGTH: 108  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 30

Ser Ser Glu Leu Thr Gln Asp Pro Ala Val Ser Val Ala Leu Gly Gln  
 1 5 10 15  
 Thr Val Arg Val Thr Cys Gln Gly Asp Ser Leu Arg Ser Tyr Tyr Ala  
 20 25 30  
 Ser Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Val Leu Val Ile Ser  
 35 40 45  
 Ala Lys Asn Asn Arg Pro Ser Gly Ile Pro Asp Arg Phe Ser Gly Ser  
 50 55 60  
 Ser Ser Gly Asn Thr Ala Ser Leu Thr Ile Thr Gly Ala Gln Ala Glu  
 65 70 75 80  
 Asp Glu Ala Asp Tyr Tyr Cys Ala Ser Arg Asn Gly Trp Asn His Val  
 85 90 95  
 Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly  
 100 105

<210> SEQ ID NO 31  
 <211> LENGTH: 108  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 31

Ser Ser Glu Leu Thr Gln Asp Pro Ala Val Ser Val Ala Leu Gly Gln  
 1 5 10 15  
 Thr Val Arg Ile Thr Cys Gln Gly Asp Ser Leu Arg Ser Tyr Tyr Ala  
 20 25 30  
 Ser Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Val Leu Val Ile Ser  
 35 40 45  
 Ala Lys Asn Asn Arg Pro Ser Gly Ile Pro Asp Arg Phe Ser Gly Ser  
 50 55 60  
 Ser Ser Gly Asn Thr Ala Ser Leu Thr Ile Thr Gly Ala Gln Ala Glu  
 65 70 75 80  
 Asp Glu Ala Asp Tyr Tyr Cys Ala Ser Arg Asn Gly Trp Asn His Val  
 85 90 95  
 Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly  
 100 105

<210> SEQ ID NO 32  
 <211> LENGTH: 108  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 32

Ser Ser Glu Leu Thr Gln Asp Pro Ala Val Ser Val Ala Leu Gly Gln  
 1 5 10 15  
 Thr Val Arg Val Thr Cys Gln Gly Asp Ser Leu Arg Ser Tyr Tyr Ala  
 20 25 30  
 Ser Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Val Leu Val Ile Ser  
 35 40 45  
 Ala Lys Asn Asn Arg Pro Ser Gly Ile Pro Asp Arg Phe Ser Gly Ser  
 50 55 60  
 Ser Ser Gly Asn Thr Ala Ser Leu Thr Ile Thr Gly Ala Gln Ala Glu  
 65 70 75 80  
 Asp Glu Ala Asp Tyr Tyr Cys Ala Ser Arg Asn Gly Trp Asn His Val

-continued

---

85	90	95	
Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly			
100	105		
<210> SEQ ID NO 33 <211> LENGTH: 108 <212> TYPE: PRT <213> ORGANISM: Homo sapiens  <400> SEQUENCE: 33			
Ser Ser Glu Leu Thr Gln Asp Pro Ala Val Ser Val Ala Leu Gly Gln			
1	5	10	15
Thr Val Arg Ile Thr Cys Gln Gly Asp Ser Leu Arg Ser Tyr Tyr Ala			
	20	25	30
Ser Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Val Leu Val Ile Ser			
	35	40	45
Ala Lys Asn Asn Arg Pro Ser Gly Ile Pro Asp Arg Phe Ser Gly Ser			
	50	55	60
Ser Ser Gly Asn Thr Ala Ser Leu Thr Ile Thr Gly Ala Gln Ala Glu			
65	70	75	80
Asp Glu Ala Asp Tyr Tyr Cys Ala Thr Arg Asn Gly Trp Asn His Val			
	85	90	95
Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly			
100	105		

<210> SEQ ID NO 34  
 <211> LENGTH: 108  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 34

Ser Ser Glu Leu Thr Gln Asp Pro Ala Val Ser Val Ala Leu Gly Gln			
1	5	10	15
Thr Val Arg Val Thr Cys Gln Gly Asp Ser Leu Arg Ser Tyr Tyr Ala			
	20	25	30
Ser Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Val Leu Val Ile Ser			
	35	40	45
Ala Lys Asn Asn Arg Pro Ser Gly Ile Pro Asp Arg Phe Ser Gly Ser			
	50	55	60
Ser Ser Gly Asn Thr Ala Ser Leu Thr Ile Thr Gly Ala Gln Ala Glu			
65	70	75	80
Asp Glu Ala Asp Tyr Tyr Cys Ala Thr Arg Asn Gly Trp Asn His Val			
	85	90	95
Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly			
100	105		

<210> SEQ ID NO 35  
 <211> LENGTH: 366  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 35

caggtgcagc tgggtgcagtc tgggggagggc gtggtccagc ctgggaggtc cctgagactc	60
tcctgtgcag cgtctggatt caccttcagc gggtatggca tgcactgggt ccgccaggct	120
ccaggcaagg ggctggagtg ggtggcagta atatcaaatg atggaagtaa gaaatattat	180
tcagactccg tgaagggccg attcaccatc tccagagaca attccaaaaa cacgctgtat	240
ctgcagatga acagcctgag agctgaggac acggctgtat attactgtgc gaaaggggtcc	300

-continued

---

```

agttccatat ggctgacca gtcctggac cactgggggc aggggaccac ggtcacgctc 360
tcctca 366

```

```

<210> SEQ ID NO 36
<211> LENGTH: 366
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 36

```

```

gaggtgcagc tgggtggagtc tgggggaggc gtggtccagc ctgggaggtc cctgagactc 60
tcctgtgcag cgtctggatt caccttcagc gggtatggca tgcactgggt ccgccaggct 120
ccaggcaagg ggctggagtg ggtggcagta atatcaaatg atggaagtaa gaaatattat 180
tcagactccg tgaagggccg attcaccatc tccagagaca attccaaaaa cacgctgtat 240
ctgcagatga acagcctgag agctgaggac acggctgtat attactgtgc gaaagggctc 300
agttccatat ggctgacca gtcctggac cactgggggc aggggaccac ggtcacgctc 360
tcctca 366

```

```

<210> SEQ ID NO 37
<211> LENGTH: 366
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 37

```

```

caggtgcagc tgggtgcagtc tgggggaggc gtggtccagc ctgggaggtc cctgagactc 60
tcctgtgcag cgtctggatt caccttcagc gggtatggca tgcactgggt ccgccaggct 120
ccaggcaagg ggctggagtg ggtggcagta atatcaaatg atggaagtaa gaaatattat 180
tcagactccg tgaagggccg attcaccatc tccagagaca attccaaaaa cacgctgtat 240
ctgcagatga acagcctgag agctgaggac acggctgtat attactgtgc gaaagggctc 300
agttccatat ggctgtcgca gtcctggac ggctgggggc aggggaccac ggtcacgctc 360
tcctca 366

```

```

<210> SEQ ID NO 38
<211> LENGTH: 366
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 38

```

```

gaggtgcagc tgggtggagtc tgggggaggc gtggtccagc ctgggaggtc cctgagactc 60
tcctgtgcag cgtctggatt caccttcagc gggtatggca tgcactgggt ccgccaggct 120
ccaggcaagg ggctggagtg ggtggcagta atatcaaatg atggaagtaa gaaatattat 180
tcagactccg tgaagggccg attcaccatc tccagagaca attccaaaaa cacgctgtat 240
ctgcagatga acagcctgag agctgaggac acggctgtat attactgtgc gaaagggctc 300
agttccatat ggctgtcgca gtcctggac ggctgggggc aggggaccac ggtcacgctc 360
tcctca 366

```

```

<210> SEQ ID NO 39
<211> LENGTH: 366
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 39

```

```

caggtgcagc tgggtgcagtc tgggggaggc gtggtccagc ctgggaggtc cctgagactc 60

```

-continued

---

```

tcctgtgcag cgtctggatt caccttcagc ggttatggca tgcactgggt ccgccaggct 120
ccaggcaagg ggctggagtg ggtggcagta atatcaaagc atggaagtaa gaaatattat 180
tcagactccg tgaagggccg attcaccatc tccagagaca attccaaaaa cacgctgtat 240
ctgcagatga acagcctgag agctgaggac acggctgtat attactgtgc gaaagggctc 300
agttccatat ggctgacctc ggccctgaac ctgtgggggc aggggaccac ggtcaccgtc 360
tcctca 366

```

```

<210> SEQ ID NO 40
<211> LENGTH: 366
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 40

```

```

gaggtgcagc tgggtggagt cgggggaggc gtggtccagc ctgggaggtc cctgagactc 60
tcctgtgcag cgtctggatt caccttcagc ggttatggca tgcactgggt ccgccaggct 120
ccaggcaagg ggctggagtg ggtggcagta atatcaaagc atggaagtaa gaaatattat 180
tcagactccg tgaagggccg attcaccatc tccagagaca attccaaaaa cacgctgtat 240
ctgcagatga acagcctgag agctgaggac acggctgtat attactgtgc gaaagggctc 300
agttccatat ggctgacctc ggccctgaac ctgtgggggc aggggaccac ggtcaccgtc 360
tcctca 366

```

```

<210> SEQ ID NO 41
<211> LENGTH: 351
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 41

```

```

gaggtgcagc tgggtggagt cgggggaggc gtggtccagc ctgggaggtc cctgagactc 60
tcctgtgcag cgtctggatt caccttcagc ggttatggca tgcactgggt ccgccaggct 120
ccaggcaagg ggctggagtg ggtggcagta atatcaaagc atggaagtaa gaaatattat 180
tcagactccg tgaagggccg attcaccatc tccagagaca attccaaaaa cacgctgtat 240
ctgcagatga acagcctgag agctgaggac acggctatat attactgtgc gaaagggctc 300
agttccatat ggttcgggga gaccgttgac tactgggggc aggggaccac g 351

```

```

<210> SEQ ID NO 42
<211> LENGTH: 363
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 42

```

```

caggtgcagc tgggtgcagt cggggctgag gtgaagaagc ctggggcctc agtgaaggct 60
tcctgcaagg tttccggata caccctcact gaattatcca tgcactgggt gcgacaggct 120
cctggaaaag ggcttgagtg gatgggaggt tttgatcgtg aagatgatga aacaatccac 180
gcacagaagt tccagggcag agtcaccatg accgaggaca catctacaga cacagcctac 240
atggaactga gcagcctgag atctgaggac acggccgctt attactgtgc aacagatctt 300
atggtgtggg gcgatttttg gatccagcac tggggccagg ggacactggt caccgtctcc 360
tca 363

```

```

<210> SEQ ID NO 43
<211> LENGTH: 363

```

-continued

---

```

<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 43

caggtgcagc tgggtgcagtc tggggctgag gtgaagaagc ctggggcctc agtgaaggtc      60
tcttgcaagg tttccggata caccctcact gaattatcca tgcactgggt gcgacaggct      120
cctggaaaag ggcttgagtg gatgggaggt tttgatcgtg aagatgatga aacaatccac      180
gcacagaagt tccagggcag agtcaccatg accgaggaca catctacaga cacagcctac      240
atggaactga gcagcctgcg atctgaggac acggccgttt attactgtgc aacagatcct      300
atggtgtggg gcgatttttg gatccagcac tggggccagg ggacactggt caccgtctcc      360
tca                                                                                   363

```

```

<210> SEQ ID NO 44
<211> LENGTH: 363
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 44

caggtgcagc tgggtgcagtc tggggctgag gtgaagaagc ctggggcctc agtgaaggtc      60
tcttgcaagg tttccggata caccctcact gaattatcca tgcactgggt gcgacaggct      120
cctggaaaag ggcttgagtg gatgggaggt tttgatcgtg aagatgatga aacaatccac      180
gcacagaagt tccagggcag agtcaccatg accgaggaca catctacaga cacagcctac      240
atggaactga gcagcctgcg atctgaggac acggccgttt attactgtgc aacagatcct      300
atggcctggg actaccgcc catccagcac tggggccagg ggacactggt caccgtctcc      360
tca                                                                                   363

```

```

<210> SEQ ID NO 45
<211> LENGTH: 363
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 45

caggtgcagc tgggtgcagtc tggggctgag gtgaagaagc ctggggcctc agtgaaggtc      60
tcttgcaagg tttccggata caccctcact gaattatcca tgcactgggt gcgacaggct      120
cctggaaaag ggcttgagtg gatgggaggt tttgatcgtg aagatgatga aacaatccac      180
gcacagaagt tccagggcag agtcaccatg accgaggaca catctacaga cacagcctac      240
atggaactga gcagcctgcg atctgaggac acggccgttt attactgtgc aacagatcct      300
atggtgtgga acttcccccc catccagcac tggggccagg ggacactggt caccgtctcc      360
tca                                                                                   363

```

```

<210> SEQ ID NO 46
<211> LENGTH: 348
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 46

caggtgcagc tgggtgcagtc tggggctgag gtgaagaagc ctggggcctc agtgaaggtc      60
tcttgcaagg tttccggata caccctcact gaattatcca tgcactgggt gcgacaggct      120
cctggaaaag ggcttgagtg gatgggaggt tttgatcgtg aagatgatga aacaatccac      180
gcacagaagt tccagggcag agtcaccatg accgaggaca catctacaga cacagcctac      240
atggaactga gcagcctgcg atctgaggac acggccgttt attactgtgc aacagatcct      300

```

-continued

---

atggtgtggg gcgatttttg gatccagcac tggggcaagg ggacaatg 348

<210> SEQ ID NO 47  
 <211> LENGTH: 354  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 47

gaggtgcagc tgttgagtc tgggggaggc ttggtacagc ctggggggtc cctgagactc 60  
 tcctgtgcag cctctggatt cacctttagc agctatgcca tgagctgggt ccgccaggct 120  
 ccaggggaagg ggctggagtg ggtctcagct attagtggta gtggtggtgg cacatactac 180  
 gcagactccg tgaagggccg gttcaccatc tccagagaca attccaagaa cacgctgtat 240  
 ctgcaaatga acagcctgag agccgaggac acggccgtgt attactgtgc gagaattcgg 300  
 ggcgactacc ggacggacat ctggggccag ggaaccacgg tcaccgtctc ctca 354

<210> SEQ ID NO 48  
 <211> LENGTH: 354  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 48

gaggtgcagc tgttgagtc tgggggaggc ttggtacagc ctggggggtc cctgagactc 60  
 tcctgtgcag cctctggatt cacctttagc agctatgcca tgagctgggt ccgccaggct 120  
 ccaggggaagg ggctggagtg ggtctcagct attagtggta gtggtggtgg cacatactac 180  
 gcagactccg tgaagggccg gtccaccatc tccagagaca attccaagaa cacgctgtat 240  
 ctgcaaatga acagcctgag agccgaggac acggccgtgt attactgtgc gagaattcgg 300  
 ggggactacc ggacggacat ctggggcccg ggaaccctgg tcaccgtctc ctca 354

<210> SEQ ID NO 49  
 <211> LENGTH: 354  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 49

gaggtgcagc tgttgagtc tgggggaggc ttggcacagc ctggggggtc cctgagactc 60  
 tcctgtgcag cctctgggtt cacctttagc agctatgcca tgagctgggt ccgccaggct 120  
 ccaggggaagg ggctggagtg ggtctcagct attagtggta gtggtggttag cacatactac 180  
 gcagactccg tgaagggccg gttcaccatc tccagagaca attccaagaa cacgctgtat 240  
 ctgcaaatga acagcctgag agccgaggac acggccgtgt attactgtgc gagagtccgg 300  
 ggggactacc ggacggacat ctggggcccg ggaaccctgg tcaccgtctc ctca 354

<210> SEQ ID NO 50  
 <211> LENGTH: 354  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 50

gaggtgcagc tgttgagtc tgggggaggc ttggtacagc ctggggggtc cctgagactc 60  
 tcctgtgcag cctctagatt cacctttagc agctatgcca tgagctgggt ccgccaggct 120  
 ccaggggaagg ggctggagtg ggtctcagct attagtggta gtggtggttag cacatactac 180  
 gcagactccg tgaagggccg gttcaccatc tccagagaca attccaagaa cacgctgtat 240  
 ctgcaaatga acagcctgag agccgaggac acggccgtgt attactgtgc gagagtccgg 300

-continued

---

ggggactacc ggacggacat ctggggccgg ggaaccctgg tcaccgtctc ctca 354

<210> SEQ ID NO 51  
 <211> LENGTH: 339  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 51

gaggtgcagc tgttgagtc tgggggaggc ttgtacagc ctggggggtc cctgagactc 60  
 tcctgtgcag cctctagatt cacctttagc agctatgcca tgagctgggt ccgccaggct 120  
 ccaggggaagg ggctggagtg ggtctcagct attagtggta gtgggtgtag cacatactac 180  
 gcagactccg tgaagggccg gttcaccatc tccagagaca attccaagaa cacgctgtat 240  
 ctgcaaatga acagcctgag agccgaggac acggccgtgt attactgtgc gagagttcgg 300  
 ggcataatcg gtatggacgt ctggggccgg ggaaccctg 339

<210> SEQ ID NO 52  
 <211> LENGTH: 309  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 52

cagcctgtgc tgactcagcc cccctcagtg tccgtgtccc caggacagac tgccagcatc 60  
 acctgctctg gagataaatt gggggataaa tatgcttctt ggtatcagca gaagccaggc 120  
 aagtcccctg tactggatcat ctatcaagat tccaatcggc cctcagggat ccctgagcga 180  
 ttctctggct ccaactctgg gaacacagcc actctgacca tcagcgggac ccaggctagg 240  
 gatgaggctg actattactg tcaggcgtgg gacagcagca ctgcatcggg gttcggcgga 300  
 gggaccaag 309

<210> SEQ ID NO 53  
 <211> LENGTH: 324  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 53

cagcctgtgc tgactcagcc cccctcagtg tccgtgtccc caggacagac tgccagcatc 60  
 acctgctctg gagataaatt gggggataaa tatgcttctt ggtatcagca gaagccaggc 120  
 aagtcccctg tactggatcat ctatcaagat tccaatcggc cctcagggat ccctgagcga 180  
 ttctctggct ccaactctgg gaacacagcc actctgacca tcagcgggac ccaggctagg 240  
 gatgaggctg actattactg tcaggcgtgg gaccactcct tgcagcacag gttcggcgga 300  
 gggaccaagg tcaccgtcct aggt 324

<210> SEQ ID NO 54  
 <211> LENGTH: 324  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 54

cagcctgtgc tgactcagcc cccctcagtg tccgtgtccc caggacagac tgccagcatc 60  
 acctgctctg gagataaatt gggggataaa tatgcttctt ggtatcagca gaagccaggc 120  
 cagaccctg tactggatcat ctatcaagat tccaatcggc cctcagggat ccctgagcga 180  
 ttctctggct ccaactccgg gaacacagcc actctgacca tcagcgggac ccaggctagg 240  
 gatgaggctg actattactg tcaggcgtgg accagcggcc tgaactcga gttcggcgga 300

-continued

---

gggaccaagg tcaccgtcct aggt 324

<210> SEQ ID NO 55  
 <211> LENGTH: 324  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 55

cagcctgtgc tgactcagcc cccctcagtg tccgtgtccc caggacagac tgccagcatc 60  
 acctgctctg gagataaatt gggggataaa tatgcttctt ggtatcagca gaagccaggc 120  
 cagtcccctg tactggatcat ctatcaagat tccaatcggc cctcagggat ccctgagcga 180  
 ttctctggct ccaactctgg ggacacagcc actctgacca tcagcgggac ccaggctagg 240  
 gatgaggctg actattactg tcaggcgtgg acgcaactccc tcagcacggt gttcggcgga 300  
 gggaccaagg tcaccgtcct aggt 324

<210> SEQ ID NO 56  
 <211> LENGTH: 324  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 56

tcctatgagc tgactcagcc cccctcagtg tccgtgtccc caggacagac tgccagcatc 60  
 acctgctctg gagataaatt gggggataaa tatgcttctt ggtatcagca gaagccaggc 120  
 cagtcccctg tactggatcat ctatcaagat tccaatcggc cctcagggat ccctgagcga 180  
 ttctctggct ccaactctgg gaacacagcc actctgacca tcagcgggac ccaggctatg 240  
 gatgaggctg actattactg tcaggcgtgg acccacagcc tgagcacggt gttcggcgga 300  
 gggaccaagc tgaccgtcct aggt 324

<210> SEQ ID NO 57  
 <211> LENGTH: 318  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 57

cagtctgtgc tgactcagcc accctcagcg tctgggaccc ccgggcagag ggtcaccatc 60  
 tcttgctctg gaaggaactc caacatcgga agttatactg taacctggta ccagcagctc 120  
 ccaggaacgg cccccaaact cctcatctat agtaatagtc agcggccctc aggggtccct 180  
 gaccgattct caggctccaa gtctggcacc tcagcctcct tggccatcag tgggctccag 240  
 tctgaagatg aggctgatta ttactgtgca gcatgggatg acagcctgaa tggcccgggtg 300  
 ttcggcggag ggaccaag 318

<210> SEQ ID NO 58  
 <211> LENGTH: 333  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 58

cagtctgtgc tgactcagcc accctcagcg tctgggaccc ccgggcagag ggtcaccatc 60  
 tcttgctctg gaaggaactc caacatcgga agttatactg taacctggta ccagcagctc 120  
 ccaggaacgg cccccaaact cctcatctat agtaatagtc agcggccctc aggggtccct 180  
 gaccgattct caggctccaa gtctggcacc tcagcctcct tggccatcag tgggctccag 240  
 tctgaagatg aggctgatta ttactgtgtg gtgtgggatg acgtgctgaa tggcccgggtg 300



-continued

---

```
ttcggcggag ggaccaagct gaccgtccta ggt 333
```

```
<210> SEQ ID NO 59
<211> LENGTH: 333
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
```

```
<400> SEQUENCE: 59
```

```
cagtctgtgc tgactcagcc accctcagcg tctgggaccc ccgggcagag ggtcaccatc 60
tcttgttctg gaaggaactc caacatcgga agttatactg taacctggta ccagcagctc 120
ccaggaacgg cccccaaact cctcatctat agtaatagtc agcggccctc aggggtccct 180
gaccgattct caggctccaa gtctggcacc tcagcctcct tggccatcag tgggctccag 240
tctgaagatg aggctgatta ttactgtgtc gtgtgggatg acaagctgaa tggcccggtg 300
ttcggcggag ggaccaagct gaccgtccta ggt 333
```

```
<210> SEQ ID NO 60
<211> LENGTH: 333
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
```

```
<400> SEQUENCE: 60
```

```
cagtctgtgc tgactcagcc accctcagcg tctgggaccc ccgggcagag ggtcaccatc 60
tcttgttctg gaaggaactc caacatcgga agttatactg taacctggta ccagcagctc 120
ccaggaacgg cccccaaact cctcatctat agtaatagtc agcggccctc aggggtccct 180
gaccgattct caggctccaa gtctggcacc tcagcctcct tggccatcag tgggctccag 240
tctgaagatg aggctgatta ttactgtgtg gtgtgggacg agatcctgaa tggcccggtg 300
ttcggcggag ggaccaagct gaccgtccta ggt 333
```

```
<210> SEQ ID NO 61
<211> LENGTH: 309
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
```

```
<400> SEQUENCE: 61
```

```
tcgtctgagc tgactcagga cctgtctgtg tctgtggcct tgggacagac agtcaggatc 60
acatgccaag gagacagcct cagaagctat tatgcaagct ggtaccagca gaagccagga 120
caggcccttg tacttgtcat ctctgctaaa aacaaccggc cctcagggat cccagaccga 180
ttctctggct ccagctcagg aaacacagct tccttgacca tcactggggc tcaggcggaa 240
gatgaagctg actattactg taactcccgg gacagcagta accatgtggt attcggcgga 300
gggaccaag 309
```

```
<210> SEQ ID NO 62
<211> LENGTH: 333
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
```

```
<400> SEQUENCE: 62
```

```
cagtctgtgc tgactcagcc accctcagcg tctgggaccc ccgggcagag ggtcaccatc 60
tcttgttctg gaaggaactc caacatcgga agttatactg taacctggta ccagcagctc 120
ccaggaacgg cccccaaact cctcatctat agtaatagtc agcggccctc aggggtccct 180
gaccgattct caggctccaa gtctggcacc tcagcctcct tggccatcag tgggctccag 240
tctgaagatg aggctgatta ttactgtctc gtgtgggacg acgtcctgaa tggcccggtg 300
```

-continued

ttcggcggag ggaccaagct gaccgtccta ggt 333

<210> SEQ ID NO 63  
 <211> LENGTH: 324  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 63

tcgtctgagc tgactcagga cctgctgtg tctgtggcct tgggacagac agtcaggatc 60  
 acatgccaaag gagacagcct cagaagctat tatgcaagct ggtaccagca gaagccagga 120  
 caggccccctg tacttgatcat ctctgctaaa aacaaccggc cctcagggat cccagaccga 180  
 ttctctggct ccagctcagg aaacacagct tccttgacca tcaactggggc tcaggcggaa 240  
 gatgaggctg actattactg tgcgtcccgg aacggctgga accatgtggt attcggcgga 300  
 gggaccaagc tgaccgtcct aggt 324

<210> SEQ ID NO 64  
 <211> LENGTH: 324  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 64

tcgtctgagc tgactcagga cctgctgtg tctgtggcct tgggacagac agtcagggtc 60  
 acatgccaaag gagacagcct cagaagctat tatgcaagct ggtaccagca gaagccagga 120  
 caggccccctg tacttgatcat ctctgctaaa aacaaccggc cctcagggat cccagaccga 180  
 ttctctggct ccagctcagg aaacacagct tccttgacca tcaactggggc tcaggcggaa 240  
 gatgaggctg actattactg tgcgtcccgg aacggctgga accatgtggt attcggcgga 300  
 gggaccaagc tgaccgtcct aggt 324

<210> SEQ ID NO 65  
 <211> LENGTH: 324  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 65

tcgtctgagc tgactcagga cctgctgtg tctgtggcct tgggacagac agtcaggatc 60  
 acatgccaaag gagacagcct cagaagctat tatgcaagct ggtaccagca gaagccagga 120  
 caggccccctg tacttgatcat ctctgctaaa aacaaccggc cctcagggat cccagaccga 180  
 ttctctggct ccagctcagg aaacacagct tccttgacca tcaactggggc tcaggcggaa 240  
 gatgaggctg actattactg tgcgtcccgg aacggctgga accatgtggt attcggcgga 300  
 gggaccaagc tgaccgtcct aggt 324

<210> SEQ ID NO 66  
 <211> LENGTH: 324  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 66

tcgtctgagc tgactcagga cctgctgtg tctgtggcct tgggacagac agtcagggtc 60  
 acatgccaaag gagacagcct cagaagctat tatgcaagct ggtaccagca gaagccagga 120  
 caggccccctg tacttgatcat ctctgctaaa aacaaccggc cctcagggat cccagaccga 180  
 ttctctggct ccagctcagg aaacacagct tccttgacca tcaactggggc tcaggcggaa 240  
 gatgaggctg actattactg tgcgtcccgg aacggctgga accatgtggt attcggcgga 300

-continued

gggaccaagc tgaccgtcct aggt 324

<210> SEQ ID NO 67  
<211> LENGTH: 324  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 67

tcgtctgagc tgactcagga ccttgctgtg tctgtggcct tgggacagac agtcaggatc 60  
 acatgccaaag gagacagcct cagaagctat tatgcaagct ggtaccagca gaagccagga 120  
 caggccccctg tacttgtcat ctctgctaaa aacaaccggc cctcagggat cccagaccga 180  
 ttctctggct ccagctcagg aaacacagct tccttgacca tcaactggggc tcaggcggaa 240  
 gatgaggctg actattactg tgcgacccgg aacggctgga accatgtggt attcggcgga 300  
 gggaccaagc tgaccgtcct aggt 324

<210> SEQ ID NO 68  
<211> LENGTH: 324  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 68

tcgtctgagc tgactcagga ccttgctgtg tctgtggcct tgggacagac agtcagggtc 60  
 acatgccaaag gagacagcct cagaagctat tatgcaagct ggtaccagca gaagccagga 120  
 caggccccctg tacttgtcat ctctgctaaa aacaaccggc cctcagggat cccagaccga 180  
 ttctctggct ccagctcagg aaacacagct tccttgacca tcaactggggc tcaggcggaa 240  
 gatgaggctg actattactg tgcgacccgg aacggctgga accatgtggt attcggcgga 300  
 gggaccaagc tgaccgtcct aggt 324

<210> SEQ ID NO 69  
<211> LENGTH: 541  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 69

Met Asn Cys Arg Glu Leu Pro Leu Thr Leu Trp Val Leu Ile Ser Val  
 1 5 10 15  
 Ser Thr Ala Glu Ser Cys Thr Ser Arg Pro His Ile Thr Val Val Glu  
 20 25 30  
 Gly Glu Pro Phe Tyr Leu Lys His Cys Ser Cys Ser Leu Ala His Glu  
 35 40 45  
 Ile Glu Thr Thr Thr Lys Ser Trp Tyr Lys Ser Ser Gly Ser Gln Glu  
 50 55 60  
 His Val Glu Leu Asn Pro Arg Ser Ser Ser Arg Ile Ala Leu His Asp  
 65 70 75 80  
 Cys Val Leu Glu Phe Trp Pro Val Glu Leu Asn Asp Thr Gly Ser Tyr  
 85 90 95  
 Phe Phe Gln Met Lys Asn Tyr Thr Gln Lys Trp Lys Leu Asn Val Ile  
 100 105 110  
 Arg Arg Asn Lys His Ser Cys Phe Thr Glu Arg Gln Val Thr Ser Lys  
 115 120 125  
 Ile Val Glu Val Lys Lys Phe Phe Gln Ile Thr Cys Glu Asn Ser Tyr  
 130 135 140  
 Tyr Gln Thr Leu Val Asn Ser Thr Ser Leu Tyr Lys Asn Cys Lys Lys  
 145 150 155 160

-continued

---

Leu Leu Leu Glu Asn Asn Lys Asn Pro Thr Ile Lys Lys Asn Ala Glu  
 165 170 175  
 Phe Glu Asp Gln Gly Tyr Tyr Ser Cys Val His Phe Leu His His Asn  
 180 185 190  
 Gly Lys Leu Phe Asn Ile Thr Lys Thr Phe Asn Ile Thr Ile Val Glu  
 195 200 205  
 Asp Arg Ser Asn Ile Val Pro Val Leu Leu Gly Pro Lys Leu Asn His  
 210 215 220  
 Val Ala Val Glu Leu Gly Lys Asn Val Arg Leu Asn Cys Ser Ala Leu  
 225 230 235 240  
 Leu Asn Glu Glu Asp Val Ile Tyr Trp Met Phe Gly Glu Glu Asn Gly  
 245 250 255  
 Ser Asp Pro Asn Ile His Glu Glu Lys Glu Met Arg Ile Met Thr Pro  
 260 265 270  
 Glu Gly Lys Trp His Ala Ser Lys Val Leu Arg Ile Glu Asn Ile Gly  
 275 280 285  
 Glu Ser Asn Leu Asn Val Leu Tyr Asn Cys Thr Val Ala Ser Thr Gly  
 290 295 300  
 Gly Thr Asp Thr Lys Ser Phe Ile Leu Val Arg Lys Ala Asp Met Ala  
 305 310 315 320  
 Asp Ile Pro Gly His Val Phe Thr Arg Gly Met Ile Ile Ala Val Leu  
 325 330 335  
 Ile Leu Val Ala Val Val Cys Leu Val Thr Val Cys Val Ile Tyr Arg  
 340 345 350  
 Val Asp Leu Val Leu Phe Tyr Arg His Leu Thr Arg Arg Asp Glu Thr  
 355 360 365  
 Leu Thr Asp Gly Lys Thr Tyr Asp Ala Phe Val Ser Tyr Leu Lys Glu  
 370 375 380  
 Cys Arg Pro Glu Asn Gly Glu Glu His Thr Phe Ala Val Glu Ile Leu  
 385 390 395 400  
 Pro Arg Val Leu Glu Lys His Phe Gly Tyr Lys Leu Cys Ile Phe Glu  
 405 410 415  
 Arg Asp Val Val Pro Gly Gly Ala Val Val Asp Glu Ile His Ser Leu  
 420 425 430  
 Ile Glu Lys Ser Arg Arg Leu Ile Ile Val Leu Ser Lys Ser Tyr Met  
 435 440 445  
 Ser Asn Glu Val Arg Tyr Glu Leu Glu Ser Gly Leu His Glu Ala Leu  
 450 455 460  
 Val Glu Arg Lys Ile Lys Ile Ile Leu Ile Glu Phe Thr Pro Val Thr  
 465 470 475 480  
 Asp Phe Thr Phe Leu Pro Gln Ser Leu Lys Leu Leu Lys Ser His Arg  
 485 490 495  
 Val Leu Lys Trp Lys Ala Asp Lys Ser Leu Ser Tyr Asn Ser Arg Phe  
 500 505 510  
 Trp Lys Asn Leu Leu Tyr Leu Met Pro Ala Lys Thr Val Lys Pro Gly  
 515 520 525  
 Arg Asp Glu Pro Glu Val Leu Pro Val Leu Ser Glu Ser  
 530 535 540

&lt;210&gt; SEQ ID NO 70

&lt;211&gt; LENGTH: 1626

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 70

-continued

---

```

atgaattgta gagaattacc cttgaccctt tgggtgctta tatctgtaag cactgcagaa    60
tcttgtactt cacgtcccca cattactgtg gttgaagggg aacctttcta tctgaaacat    120
tgctcgtggt cacttgca ca tgagattgaa acaaccacca aaagctggta caaaagcagt    180
ggatcacagg aacatgtgga gctgaaccca aggagttcct cgagaattgc tttgcatgat    240
tgtgttttgg agttttggcc agttgagttg aatgacacag gatcttactt tttccaaatg    300
aaaaattata ctcaaaaatg gaaattaaat gtcatcagaa gaaataaaca cagctgtttc    360
actgaaagac aagtaactag taaaattgtg gaagttaaaa aattttttca gataacctgt    420
gaaaacagtt actatcaaac actgggtcaac agcacatcat tgtataagaa ctgtaaaaag    480
ctactactgg agaacaataa aaacccaacg ataaagaaga acgccgagtt tgaagatcag    540
gggtattact cctgcgtgca tttccttcat cataatggaa aactatttaa tatcaccaaa    600
accttcaata taacaatagt ggaagatcgc agtaatatag ttccggttct tcttgacca    660
aagcttaacc atgttgcaagt ggaattagga aaaaacgtaa ggctcaactg ctctgctttg    720
ctgaatgaag aggatgtaat ttattggatg ttcggggaag aaaatggatc ggatccta    780
atacatgaag agaaagaaat gagaattatg actccagaag gcaaatggca tgcttcaaaa    840
gtattgagaa ttgaaaatat tggtgaaagc aatctaaatg ttttatataa ttgcaactgt    900
gccagcacgg gaggcacaga caccaaaagc ttcactcttg tgagaaaagc agacatggct    960
gatatcccag gccacgtctt cacaagagga atgatcatag ctgttttgat cttgggtggca 1020
gtagtgtgcc tagtgactgt gtgtgtcatt tatagagttg acttggttct atttataga 1080
catttaacga gaagagatga aacattaaca gatggaaaaa catatgatgc ttttgtgtct 1140
tacctaaaag aatgccgacc tgaaaatgga gaggagcaca cctttgctgt ggagattttg 1200
cccagggtgt tggagaaaca ttttgggtat aagttagca tatttgaaag ggatgtagtg 1260
cctggaggag ctgttggtga tgaaatccac tcaactgatag agaaaagccg aagactaatc 1320
attgtcctaa gtaaaagtta tatgtctaag gaggtcaggt atgaactga aagtggactc 1380
catgaagcat tgggtgaaag aaaaattaa ataacttaa ttgaattac acctgttact 1440
gacttcacat tcttgcccca atcactaaag cttttgaaat ctacagagt tctgaagtgg 1500
aaggccgata aatctctttc ttataactca aggttctgga agaaccttct ttacttaatg 1560
cctgcaaaaa cagtcaagcc aggtagagac gaaccggaag tcttgctgt tctttccgag 1620
tcttaa 1626

```

&lt;210&gt; SEQ ID NO 71

&lt;211&gt; LENGTH: 599

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 71

```

Met Leu Cys Leu Gly Trp Ile Phe Leu Trp Leu Val Ala Gly Glu Arg
 1             5             10             15
Ile Lys Gly Phe Asn Ile Ser Gly Cys Ser Thr Lys Lys Leu Leu Trp
      20             25             30
Thr Tyr Ser Thr Arg Ser Glu Glu Glu Phe Val Leu Phe Cys Asp Leu
      35             40             45
Pro Glu Pro Gln Lys Ser His Phe Cys His Arg Asn Arg Leu Ser Pro
      50             55             60
Lys Gln Val Pro Glu His Leu Pro Phe Met Gly Ser Asn Asp Leu Ser
      65             70             75             80

```



-continued

Lys Phe Cys Tyr Phe Gln Glu Pro Glu Ser Leu Pro His Leu Val Lys  
 515 520 525

Lys Ala Leu Arg Val Leu Pro Thr Val Thr Trp Arg Gly Leu Lys Ser  
 530 535 540

Val Pro Pro Asn Ser Arg Phe Trp Ala Lys Met Arg Tyr His Met Pro  
 545 550 555 560

Val Lys Asn Ser Gln Gly Phe Thr Trp Asn Gln Leu Arg Ile Thr Ser  
 565 570 575

Arg Ile Phe Gln Trp Lys Gly Leu Ser Arg Thr Glu Thr Thr Gly Arg  
 580 585 590

Ser Ser Gln Pro Lys Glu Trp  
 595

<210> SEQ ID NO 72  
 <211> LENGTH: 1800  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 72

atgctctggt tgggctggat atttctttgg cttgttgacg gagagcgaat taaaggattt 60  
 aatatttcag gttgttccac aaaaaaactc ctttgacat attctacaag gagtgaagag 120  
 gaatttgctt tattttgtga tttaccagag ccacagaaat cacatttctg ccacagaaat 180  
 cgactctcac caaaacaagt ccttgagcac ctgcccttca tgggtagtaa cgacctatct 240  
 gatgtccaat ggtaccaaca accttcgaat ggagatccat tagaggacat taggaaaagc 300  
 tatcctcaca tcattcagga caaatgtacc cttcactttt tgaccccagg ggtgaataat 360  
 tctgggtcat atattttag acccaagatg attaagagcc cctatgatgt agcctggtgt 420  
 gtcaagatga ttttagaagt taagccccag acaaatgcat cctgtgagta ttccgcatca 480  
 cataagcaag acctacttct tgggagcact ggctctatct cttgccccag tctcagctgc 540  
 caaagtgatg cacaaagtcc agcggtaacc tggtagaaga atggaaaact cctctctgtg 600  
 gaaaggagca accgaatcgt agtggatgaa gtttatgact atcaccaggg cacatatgta 660  
 tgtgattaca ctcagtcgga tactgtgagt tcgtggacag tcagagctgt tgttcaagtg 720  
 agaaccattg tgggagacac taaactcaaa ccagatattc tggatcctgt cgaggacaca 780  
 ctggaagtag aacttgaaa gcctttaact attagctgca aagcacgatt tggctttgaa 840  
 agggctctta accctgtcat aaaatggtac atcaaagatt ctgacctaga gtgggaagtc 900  
 tcagtacctg aggcgaaaag tattaatcc actttaaagg atgaaatcat tgagcgtaat 960  
 atcatcttgg aaaaagtcac tcagcgtgat cttcgcagga agtttgtttg ctttgtccag 1020  
 aactccattg gaaacacaac ccagtccgtc caactgaaag aaaagagagg agtggtgctc 1080  
 ctgtacatcc tgcttgccac catcgggacc ctggtggccg tgctggcgcc gagtgcctc 1140  
 ctctacaggc actggattga aatagtctg ctgtaccgga cctaccagag caaggatcag 1200  
 acgcttgggg ataaaaagga ttttgatgct ttcgtatcct atgcaaatg gagctctttt 1260  
 ccaagtgagg ccacttcac tctgagtga gaacacttg ccctgagcct atttcctgat 1320  
 gttttagaaa acaaatatgg atatagcctg tgtttgcttg aaagagatgt ggctccagga 1380  
 ggagtgtatg cagaagacat tgtgagcatt attaagagaa gcagaagagg aatatttatc 1440  
 ttgagcccca actatgtcaa tggaccagc atctttgaac tacaagcagc agtgaatctt 1500  
 gccttgatg atcaaacact gaaactcatt ttaattaagt tctgttactt ccaagagcca 1560  
 gagtctctac ctcatctcgt gaaaaaagct ctcagggttt tgcccacagt tacttggaga 1620

-continued

ggcttaaaat cagttctcc caattctagg ttctgggcca aaatgegeta ccacatgcct 1680  
 gtgaaaaact ctcaggatt cacgtggaac cagctcagaa ttacctctag gatttttcag 1740  
 tggaaaggac tcagtagaac agaaaccact gggaggagct cccagcctaa ggaatggtga 1800

<210> SEQ ID NO 73  
 <211> LENGTH: 467  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 73

Met Gly Ser Thr Ala Ile Leu Gly Leu Leu Leu Ala Val Leu Gln Gly  
 1 5 10 15  
 Gly Arg Ala Glu Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln  
 20 25 30  
 Pro Gly Arg Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe  
 35 40 45  
 Ser Gly Tyr Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu  
 50 55 60  
 Glu Trp Val Ala Val Ile Ser Asn Asp Gly Ser Lys Lys Tyr Tyr Ser  
 65 70 75 80  
 Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn  
 85 90 95  
 Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val  
 100 105 110  
 Tyr Tyr Cys Ala Lys Gly Ser Ser Ser Ile Trp Leu Thr Ser Ala Leu  
 115 120 125  
 Asn Leu Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr  
 130 135 140  
 Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser  
 145 150 155 160  
 Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu  
 165 170 175  
 Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His  
 180 185 190  
 Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser  
 195 200 205  
 Val Val Thr Val Pro Ser Ser Asn Phe Gly Thr Gln Thr Tyr Thr Cys  
 210 215 220  
 Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys Thr Val Glu  
 225 230 235 240  
 Arg Lys Cys Cys Val Glu Cys Pro Pro Cys Pro Ala Pro Pro Val Ala  
 245 250 255  
 Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met  
 260 265 270  
 Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His  
 275 280 285  
 Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val  
 290 295 300  
 His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Phe  
 305 310 315 320  
 Arg Val Val Ser Val Leu Thr Val Val His Gln Asp Trp Leu Asn Gly  
 325 330 335  
 Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ala Pro Ile  
 340 345 350



-continued

Glu Lys Thr Ile Ser Lys Thr Lys Gly Gln Pro Arg Glu Pro Gln Val  
 355 360 365

Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser  
 370 375 380

Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu  
 385 390 395 400

Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro  
 405 410 415

Met Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val  
 420 425 430

Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met  
 435 440 445

His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser  
 450 455 460

Pro Gly Lys  
 465

<210> SEQ ID NO 74  
 <211> LENGTH: 1427  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 74

gtcgcgccc ccaccatggg gtcaaccgcc atccttgccc tctcctggc tgcctgcag 60  
 ggagggcgcg ccgaggtgca gctggtggag tctgggggag gcgtggtcca gcctgggagg 120  
 tccctgagac tctcctgtgc agcgtctgga ttcacctca gcggttatgg catgcactgg 180  
 gtccgccagg ctccaggcaa ggggctggag tgggtggcag taatatcaa tgatggaagt 240  
 aagaaatatt attcagactc cgtgaagggc cgattcacca tctccagaga caattccaaa 300  
 aacacgctgt atctgcagat gaacagcctg agagctgagg acacggctgt atattactgt 360  
 gcgaaagggc ccagttccat atggctgacc tggccctga acctgtgggg gcaggggacc 420  
 acggtcaccg tctcctcagc tagcaccaag ggccatcgg tcttccccct ggcgcctgc 480  
 tccaggagca cctccgagag cacagcggcc ctgggctgcc tggtaagga ctacttcccc 540  
 gaaccggtga cgggtgctgt gaactcaggc gctctgacca gcggcgtgca caccttccca 600  
 gctgtcctac agtcctcagg actctactcc ctacgcagcg tggtgaccgt gccctccagc 660  
 aacttcggca cccagacctc cacctgcaac gtagatcaca agcccagcaa caccaaggtg 720  
 gacaagacag ttgagcgcaa atgttggtgc gtagtcccac cgtgcccagc accacctgtg 780  
 gcaggaccgt cagtcttctt cttcccccca aaaccaagg acacctcat gatctcccg 840  
 accctgagg tcacgtgctg ggtggtggac gtgagccacg aagacccga ggtccagttc 900  
 aactggtacg tggacggcgt ggaggtgcat aatgccaaaga caaagccagc ggaggagcag 960  
 ttcaacagca cgttccgtgt ggtcagcgtc ctaccgctg tgcaccagga ctggctgaac 1020  
 ggcaaggagt acaagtcaa ggtctccaac aaaggcctcc cagccccat cgagaaaacc 1080  
 atctccaaaa ccaaaggca gccccgagaa ccacaggtgt acacctgcc cccatcccg 1140  
 gaggagatga ccaagaacca ggtcagcctg acctgcctgg tcaaaggctt ctaccccagc 1200  
 gacatcgccg tggagtggga gagcaatggg cagccggaga acaactacaa gaccacacct 1260  
 cccatgctgg actccgacgg ctcttcttct ctctacagca agctcacctg ggacaagagc 1320  
 aggtggcagc aggggaacgt cttctcatgc tccgtgatgc atgaggctct gcacaaccac 1380  
 tacacgcaga agagcctctc cctgtctccg ggtaaatgag cggccgc 1427

-continued

<210> SEQ ID NO 75  
 <211> LENGTH: 233  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 75

Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro  
 1 5 10 15  
 Gly Ser Thr Gly Ser Ser Glu Leu Thr Gln Asp Pro Ala Val Ser Val  
 20 25 30  
 Ala Leu Gly Gln Thr Val Arg Ile Thr Cys Gln Gly Asp Ser Leu Arg  
 35 40 45  
 Ser Tyr Tyr Ala Ser Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Val  
 50 55 60  
 Leu Val Ile Ser Ala Lys Asn Asn Arg Pro Ser Gly Ile Pro Asp Arg  
 65 70 75 80  
 Phe Ser Gly Ser Ser Ser Gly Asn Thr Ala Ser Leu Thr Ile Thr Gly  
 85 90 95  
 Ala Gln Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ser Arg Asn Gly  
 100 105 110  
 Trp Asn His Val Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly  
 115 120 125  
 Gln Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu  
 130 135 140  
 Glu Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe  
 145 150 155 160  
 Tyr Pro Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val  
 165 170 175  
 Lys Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys  
 180 185 190  
 Tyr Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser  
 195 200 205  
 His Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu  
 210 215 220  
 Lys Thr Val Ala Pro Thr Glu Cys Ser  
 225 230

<210> SEQ ID NO 76  
 <211> LENGTH: 733  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 76

gtcgacgttt aaacgccgc accatggaga cagacacact cctgctatgg gtactgctgc 60  
 tctgggttcc aggttccact ggttcgtctg agctgactca ggaccctgct gtgtctgtgg 120  
 ccttgggaca gacagtcagg atcacatgcc aaggagacag cctcagaagc tattatgcaa 180  
 gctggtacca gcagaagcca ggacaggccc ctgtacttgt catctctgct aaaaacaacc 240  
 ggcctcagg gatccagac cgattctctg gctccagctc aggaacaca gcttccttga 300  
 ccatcactgg ggctcaggcg gaagatgagg ctgactatta ctgtgctgct cggaacggct 360  
 ggaaccatgt ggtattcggc ggagggacca agctgaccgt cctaggccaa ccgaaagcgg 420  
 cgccctcggc cactctgttc ccgccctcct ctgaggagct tcaagccaac aaggccacac 480  
 tgggtgtgtct cataagtgc ttctaccgg gagccgtgac agtggcctgg aaggcagata 540

-continued

---

```

gcagccccgt caaggcggga gtggagacca ccacaccctc caaacaagc aacaacaagt 600
acgcgccag cagctatctg agcctgacgc ctgagcagtg gaagtccac agaagctaca 660
gctgccaggt cacgcatgaa gggagcaccg tggagaagac agtggcccct acagaatggt 720
cataggcggc cgc 733

```

```

<210> SEQ ID NO 77
<211> LENGTH: 467
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 77

```

```

Met Gly Ser Thr Ala Ile Leu Gly Leu Leu Leu Ala Val Leu Gln Gly
1          5          10          15
Gly Arg Ala Glu Val Gln Leu Val Glu Ser Gly Gly Gly Val Val Gln
20          25          30
Pro Gly Arg Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe
35          40          45
Ser Gly Tyr Gly Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
50          55          60
Glu Trp Val Ala Val Ile Ser Asn Asp Gly Ser Lys Lys Tyr Tyr Ser
65          70          75          80
Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn
85          90          95
Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val
100         105         110
Tyr Tyr Cys Ala Lys Gly Ser Ser Ser Ile Trp Leu Ser Gln Ser Leu
115         120         125
Asp Gly Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr
130         135         140
Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser
145         150         155         160
Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu
165         170         175
Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His
180         185         190
Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser
195         200         205
Val Val Thr Val Pro Ser Ser Asn Phe Gly Thr Gln Thr Tyr Thr Cys
210         215         220
Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys Thr Val Glu
225         230         235         240
Arg Lys Cys Cys Val Glu Cys Pro Pro Cys Pro Ala Pro Pro Val Ala
245         250         255
Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met
260         265         270
Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His
275         280         285
Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val
290         295         300
His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Phe
305         310         315         320
Arg Val Val Ser Val Leu Thr Val Val His Gln Asp Trp Leu Asn Gly
325         330         335
Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ala Pro Ile

```

-continued

340				345				350							
Glu	Lys	Thr	Ile	Ser	Lys	Thr	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val
		355					360					365			
Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser
	370					375					380				
Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu
385					390					395					400
Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro
				405					410					415	
Met	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val
			420					425					430		
Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met
		435					440					445			
His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser
	450					455					460				
Pro	Gly	Lys													
465															

&lt;210&gt; SEQ ID NO 78

&lt;211&gt; LENGTH: 1404

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 78

```

atgggggtcaa cgcctatcct tggcctcctc ctggctgtcc tgcagggagg gcgcgccgag    60
gtgcagctgg tggagtctgg gggaggcgtg gtccagcctg ggaggtcct gagactctcc    120
tgtgcagcgt ctggattcac cttcagcggg tatggcatgc actgggtccg ccaggctcca    180
ggcaaggggc tggagtgggt ggcagtaata tcaaatgatg gaagtaagaa atattattca    240
gactccgtga agggccgatt caccatctcc agagacaatt ccaaaaacac gctgtatctg    300
cagatgaaca gcctgagagc tgaggacacg gctgtatatt actgtgcgaa aggggtccagt    360
tccatattgg tgtcgcagtc cctggacggc tgggggcagg ggaccacggt caccgtctcc    420
tcagctagca ccaagggccc atcggctctc ccctggcgc cctgctccag gagcacctcc    480
gagagcacag cggccctggg ctgcctggtc aaggactact tccccgaacc ggtgacgggtg    540
tcgtggaact caggcgtctt gaccagcggc gtgcacacct tcccagctgt cctacagtcc    600
tcaggactct actcctcag cagcgtgggt accgtgcctt ccagcaactt cggcaccag    660
acctacacct gcaacgtaga tcacaagccc agcaacacca aggtggacaa gacagttgag    720
cgaaaatggt gtgtcgagtg cccaccgtgc ccagcaccac ctgtggcagg accgtcagtc    780
ttcctcttcc ccccaaaacc caaggacacc ctcatgatct cccggacccc tgaggtcacg    840
tgcgtgggtg tggacgtgag ccacgaagac cccgaggtcc agttcaactg gtacgtggac    900
ggcgtggagg tgcataatgc caagacaaaag ccacgggagg agcagttcaa cagcagttc    960
cgtgtgggtca gcgtcctcac cgttgtgcac caggactggc tgaacggcaa ggagtacaag   1020
tgcaaggtct ccaacaaagg cctcccagcc cccatcgaga aaaccatctc caaaaccaa   1080
gggcagcccc gagaaccaca ggtgtacacc ctgccccat cccgggagga gatgaccaag   1140
aaccaggtca gcctgacctg cctgggtcaaa ggcttctacc ccagcgacat cgcctggag   1200
tgggagagca atgggcagcc ggagaacaac tacaagacca cacctccat gctggactcc   1260
gacggctcct tcttcteta cagcaagctc accgtggaca agagcaggtg gcagcagggg   1320
aacgtcttct catgctccgt gatgcatgag gctctgcaca accactacac gcagaagagc   1380

```

-continued

ctctccctgt ctccgggtaa atga

1404

&lt;210&gt; SEQ ID NO 79

&lt;211&gt; LENGTH: 233

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 79

Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro  
1 5 10 15Gly Ser Thr Gly Ser Ser Glu Leu Thr Gln Asp Pro Ala Val Ser Val  
20 25 30Ala Leu Gly Gln Thr Val Arg Ile Thr Cys Gln Gly Asp Ser Leu Arg  
35 40 45Ser Tyr Tyr Ala Ser Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Val  
50 55 60Leu Val Ile Ser Ala Lys Asn Asn Arg Pro Ser Gly Ile Pro Asp Arg  
65 70 75 80Phe Ser Gly Ser Ser Ser Gly Asn Thr Ala Ser Leu Thr Ile Thr Gly  
85 90 95Ala Gln Ala Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ser Arg Asn Gly  
100 105 110Trp Asn His Val Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly  
115 120 125Gln Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu  
130 135 140Glu Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe  
145 150 155 160Tyr Pro Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val  
165 170 175Lys Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys  
180 185 190Tyr Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser  
195 200 205His Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu  
210 215 220Lys Thr Val Ala Pro Thr Glu Cys Ser  
225 230

&lt;210&gt; SEQ ID NO 80

&lt;211&gt; LENGTH: 710

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 80

atggagacag acacactcct gctatgggta ctgtgctct gggttccagg ttccactggt 60

tcgtctgagc tgactcagga cctgctgtg tctgtggcct tgggacagac agtcaggatc 120

acatgccaag gagacagcct cagaagctat tatgcaagct ggtaccagca gaagccagga 180

caggccctg tacttgcat ctctgctaaa aacaaccggc cctcaggat cccagaccga 240

ttctctggct ccagctcagg aaacacagct tccttgacca tcaactggggc tcaggcggaa 300

gatgaggctg actattactg tgcgtcccgg aacggctgga accatgtggt attcggcgga 360

gggaccaagc tgaccgtcct aggccaaccg aaagcggcgc cctcggtcac tctgttcccg 420

ccctcctctg aggagcttca agccaacaag gccacactgg tgtgtctcat aagtgacttc 480

taccgggag ccgtgacagt ggctggaag gcagatagca gcccgtcaa ggccgggagtg 540

-continued

---

```

gagaccacca caccctccaa acaaagcaac aacaagtacg cggccagcag ctatctgagc 600
ctgacgcctg agcagtggaa gtcccacaga agctacagct gccaggtcac gcatgaaggg 660
agcaccgtgg agaagacagt ggcccctaca gaatgttcat aggeggccgc 710

```

```

<210> SEQ ID NO 81
<211> LENGTH: 466
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 81

```

```

Met Gly Ser Thr Ala Ile Leu Gly Leu Leu Leu Ala Val Leu Gln Gly
1           5           10           15
Gly Arg Ala Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys
20           25           30
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Val Ser Gly Tyr Thr Leu
35           40           45
Thr Glu Leu Ser Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
50           55           60
Glu Trp Met Gly Gly Phe Asp Arg Glu Asp Asp Glu Thr Ile His Ala
65           70           75           80
Gln Lys Phe Gln Gly Arg Val Thr Met Thr Glu Asp Thr Ser Thr Asp
85           90           95
Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val
100          105          110
Tyr Tyr Cys Ala Thr Asp Leu Met Val Trp Gly Asp Phe Trp Ile Gln
115          120          125
His Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys
130          135          140
Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu
145          150          155          160
Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro
165          170          175
Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr
180          185          190
Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val
195          200          205
Val Thr Val Pro Ser Ser Asn Phe Gly Thr Gln Thr Tyr Thr Cys Asn
210          215          220
Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys Thr Val Glu Arg
225          230          235          240
Lys Cys Cys Val Glu Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly
245          250          255
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
260          265          270
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
275          280          285
Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
290          295          300
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Phe Arg
305          310          315          320
Val Val Ser Val Leu Thr Val Val His Gln Asp Trp Leu Asn Gly Lys
325          330          335
Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ala Pro Ile Glu
340          345          350

```

-continued

Lys Thr Ile Ser Lys Thr Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr  
           355                          360                          365  
 Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu  
           370                          375                          380  
 Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp  
 385                          390                          395                          400  
 Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Met  
                           405                          410                          415  
 Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp  
                           420                          425                          430  
 Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His  
           435                          440                          445  
 Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro  
           450                          455                          460  
 Gly Lys  
 465

&lt;210&gt; SEQ ID NO 82

&lt;211&gt; LENGTH: 1401

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 82

```

atgggggtcaa cgcctcctc tggcctcctc ctggctgtcc tgcagggagg gcgcgcccag    60
gtgcagctgg tgcagtctgg ggctgaggtg aagaagcctg gggcctcagt gaaggtctcc    120
tgcaaggttt cgggatacac cctcactgaa ttatccatgc actgggtgcg acaggctcct    180
ggaaaagggc ttgagtggat gggaggtttt gatcgtgaag atgatgaaac aatccacgca    240
cagaagttcc agggcagagt caccatgacc gaggacacat ctacagacac agcctacatg    300
gaactgagca gcctgcgatc tgaggacacg gccgtttatt actgtgcaac agatcttatg    360
gtgtggggcg atttttggat ccagcactgg ggccagggga cactggtcac cgtctcctca    420
gctagcacca agggcccac ggtcttcccc ctggcgcctt gctccaggag cacctccgag    480
agcacagcgg cctgggctg cctgggtcaag gactacttcc ccgaaccggt gacgggtgctg    540
tggaaactcag gcgctctgac cagcggcgtg cacaccttcc cagctgtcct acagtcctca    600
ggactctact ccctcagcag cgtgggtgacc gtgccctcca gcaacttcg caccagacc    660
tacacctgca acgtagatca caagcccagc aacaccaagg tggacaagac agttgagcgc    720
aaatgttggtg tcgagtgcc accgtgcca gcaccacctg tggcaggacc gtcagtcttc    780
ctcttcccc caaaacccaa ggacaccctc atgatctccc ggaccctga ggtcacgtgc    840
gtggtggtgg acgtgagcca cgaagacccc gaggtccagt tcaactggta cgtggacggc    900
gtggagggtgc ataatgccaa gacaaagcca cgggaggagc agttcaacag cacgttccgt    960
gtggtcagcg tcctcacctg tgtgcaccag gactggctga acggcaagga gtacaagtgc   1020
aaggtctcca acaaaggcct cccagcccc atcgagaaaa ccatctcca aaccaaaggg   1080
cagccccgag aaccacaggt gtacaccctg ccccatccc gggaggagat gaccaagaac   1140
caggctcagc tgacctgctt ggtcaaaggc ttctaccca gcgacatcgc cgtggagtgg   1200
gagagcaatg ggcagccgga gaacaactac aagaccacac ctccatgct ggactccgac   1260
ggctccttct tcctctacag caagctcacc gtggacaaga gcaggtggca gcaggggaac   1320
gtcttctcat gctccgtgat gcatgaggct ctgcacaacc actacacgca gaagagcctc   1380
tcctgtctc cgggtaaatg a                                     1401

```

-continued

<210> SEQ ID NO 83  
 <211> LENGTH: 238  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 83

Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro  
 1 5 10 15  
 Gly Ser Thr Gly Ala Ser Gln Ser Val Leu Thr Gln Pro Pro Ser Ala  
 20 25 30  
 Ser Gly Thr Pro Gly Gln Arg Val Thr Ile Ser Cys Ser Gly Arg Asn  
 35 40 45  
 Ser Asn Ile Gly Ser Tyr Thr Val Thr Trp Tyr Gln Gln Leu Pro Gly  
 50 55 60  
 Thr Ala Pro Lys Leu Leu Ile Tyr Ser Asn Ser Gln Arg Pro Ser Gly  
 65 70 75 80  
 Val Pro Asp Arg Phe Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu  
 85 90 95  
 Ala Ile Ser Gly Leu Gln Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Val  
 100 105 110  
 Val Trp Asp Glu Ile Leu Asn Gly Pro Val Phe Gly Gly Gly Thr Lys  
 115 120 125  
 Leu Thr Val Leu Gly Gln Pro Lys Ala Ala Pro Ser Val Thr Leu Phe  
 130 135 140  
 Pro Pro Ser Ser Glu Glu Leu Gln Ala Asn Lys Ala Thr Leu Val Cys  
 145 150 155 160  
 Leu Ile Ser Asp Phe Tyr Pro Gly Ala Val Thr Val Ala Trp Lys Ala  
 165 170 175  
 Asp Ser Ser Pro Val Lys Ala Gly Val Glu Thr Thr Thr Pro Ser Lys  
 180 185 190  
 Gln Ser Asn Asn Lys Tyr Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro  
 195 200 205  
 Glu Gln Trp Lys Ser His Arg Ser Tyr Ser Cys Gln Val Thr His Glu  
 210 215 220  
 Gly Ser Thr Val Glu Lys Thr Val Ala Pro Thr Glu Cys Ser  
 225 230 235

<210> SEQ ID NO 84  
 <211> LENGTH: 717  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 84

atggagacag acacactcct gctatgggta ctgctgctct gggttccagg ttccactggt 60  
 gctagccagt ctgtgctgac tcagccaccc tcagcgtctg ggacccccgg gcagagggtc 120  
 accatctctt gttctggaag gaactccaac atcggaagtt atactgtaac ctggtaccag 180  
 cagctcccag gaacggcccc caaactcctc atctatagta atagtcagcg gcctcaggg 240  
 gtccctgacc gattctcagg ctccaagtct ggcacctcag cctccttggc catcagtggg 300  
 ctccagtctg aagatgaggc tgattattac tgtgtgggtg gggacgagat cctgaatggc 360  
 ccggtgttcg gcggaggac caagctgacc gtccatggcc aaccgaaagc ggccgacctg 420  
 gtcactctgt tcccgcctc ctctgaggag cttcaagcca acaaggccac actggtgtgt 480  
 ctcataagtg acttctaccc gggagccgtg acagtggcct ggaaggcaga tagcagcccc 540



-continued

---

```

gtcaaggcgg gagtggagac caccacaccc tccaaacaaa gcaacaacaa gtacgcggcc 600
agcagctatc tgagcctgac gctgagcag tggaaagccc acagaagcta cagctgccag 660
gtcacgcatg aaggagcac cgtggagaag acagtggccc ctacagaatg ttcatag 717

```

&lt;210&gt; SEQ ID NO 85

&lt;211&gt; LENGTH: 466

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 85

```

Met Gly Ser Thr Ala Ile Leu Gly Leu Leu Leu Ala Val Leu Gln Gly
1           5           10           15
Gly Arg Ala Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys
20           25           30
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Val Ser Gly Tyr Thr Leu
35           40           45
Thr Glu Leu Ser Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu
50           55           60
Glu Trp Met Gly Gly Phe Asp Arg Glu Asp Asp Glu Thr Ile His Ala
65           70           75           80
Gln Lys Phe Gln Gly Arg Val Thr Met Thr Glu Asp Thr Ser Thr Asp
85           90           95
Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val
100          105          110
Tyr Tyr Cys Ala Thr Asp Leu Met Val Trp Asn Phe Pro Pro Ile Gln
115          120          125
His Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys
130          135          140
Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu
145          150          155          160
Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro
165          170          175
Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr
180          185          190
Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val
195          200          205
Val Thr Val Pro Ser Ser Asn Phe Gly Thr Gln Thr Tyr Thr Cys Asn
210          215          220
Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys Thr Val Glu Arg
225          230          235          240
Lys Cys Cys Val Glu Cys Pro Pro Cys Pro Ala Pro Pro Val Ala Gly
245          250          255
Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
260          265          270
Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
275          280          285
Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
290          295          300
Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Phe Arg
305          310          315          320
Val Val Ser Val Leu Thr Val Val His Gln Asp Trp Leu Asn Gly Lys
325          330          335
Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ala Pro Ile Glu
340          345          350

```

-continued

---

Lys Thr Ile Ser Lys Thr Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr  
 355 360 365

Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu  
 370 375 380

Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp  
 385 390 395 400

Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Met  
 405 410 415

Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp  
 420 425 430

Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His  
 435 440 445

Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro  
 450 455 460

Gly Lys  
 465

&lt;210&gt; SEQ ID NO 86

&lt;211&gt; LENGTH: 1401

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 86

```

atgggggtcaa cgcctcctc tggcctcctc ctggctgtcc tgcagggagg ggcgccccag   60
gtgcagctgg tgcagtctgg ggctgaggtg aagaagcctg gggcctcagt gaaggtctcc   120
tgcaaggttt cggatacac cctcactgaa ttatccatgc actgggtgcg acaggctcct   180
ggaaaagggc ttgagtggat gggaggtttt gatcgtgaag atgatgaaac aatccacgca   240
cagaagttcc agggcagagt caccatgacc gaggacacat ctacagacac agcctacatg   300
gaactgagca gcctgcgatc tgaggacacg gccgtttatt actgtgcaac agatcttatg   360
gtgtggaact tccccccat ccagcactgg ggccagggga cactggtcac cgtctcctca   420
gctagcacca agggcccatc ggtcttcccc ctggcgccct gctccaggag cacctccgag   480
agcacagcgg ccctgggctg cctgggtcaag gactacttcc ccgaaccggt gacgggtgctg   540
tggaaactcag gcgctctgac cagcggcgtg cacaccttcc cagctgtcct acagtctctca   600
ggactctact ccctcagcag cgtgggtgacc gtgccctcca gcaacttcgg caccagacc   660
tacacctgca acgtagatca caagcccagc aacaccaagg tggacaagac agttgagcgc   720
aaatgttggtg tcgagtgcc accgtgccc gcaccacctg tggcaggacc gtcagtcttc   780
ctcttcccc caaaaccaa ggacaccctc atgatctccc ggaccctga ggtcacgtgc   840
gtggtggtgg acgtgagcca cgaagacccc gaggtccagt tcaactggta cgtggacggc   900
gtggaggtgc ataatgcaa gacaaagcca cgggaggagc agttcaacag cacgttccgt   960
gtggtcagcg tcctcaccgt tgtgcaccag gactggctga acggcaagga gtacaagtgc  1020
aaggtctcca acaaaggcct cccagcccc atcgagaaaa ccatctcaa aaccaaaggg  1080
cagccccgag aaccacaggt gtacaccctg cccccatccc gggaggagat gaccaagaac  1140
caggtcagcc tgacctgct ggtcaaaggc ttctaccca gcgacatcgc cgtggagtgg  1200
gagagcaatg ggcagccgga gaacaactac aagaccacac ctcccatgct ggactccgac  1260
ggctccttct tcctctacag caagctcacc gtggacaaga gcaggtggca gcaggggaac  1320
gtcttctcat gctcctgat gcatgaggct ctgcacaacc actacacgca gaagagcctc  1380
tcctgtctc cgggtaaatg a                                     1401

```

-continued

<210> SEQ ID NO 87  
 <211> LENGTH: 238  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens  
  
 <400> SEQUENCE: 87  
  
 Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro  
 1 5 10 15  
 Gly Ser Thr Gly Ala Ser Gln Ser Val Leu Thr Gln Pro Pro Ser Ala  
 20 25 30  
 Ser Gly Thr Pro Gly Gln Arg Val Thr Ile Ser Cys Ser Gly Arg Asn  
 35 40 45  
 Ser Asn Ile Gly Ser Tyr Thr Val Thr Trp Tyr Gln Gln Leu Pro Gly  
 50 55 60  
 Thr Ala Pro Lys Leu Leu Ile Tyr Ser Asn Ser Gln Arg Pro Ser Gly  
 65 70 75 80  
 Val Pro Asp Arg Phe Ser Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu  
 85 90 95  
 Ala Ile Ser Gly Leu Gln Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Val  
 100 105 110  
 Val Trp Asp Asp Val Leu Asn Gly Pro Val Phe Gly Gly Gly Thr Lys  
 115 120 125  
 Leu Thr Val Leu Gly Gln Pro Lys Ala Ala Pro Ser Val Thr Leu Phe  
 130 135 140  
 Pro Pro Ser Ser Glu Glu Leu Gln Ala Asn Lys Ala Thr Leu Val Cys  
 145 150 155 160  
 Leu Ile Ser Asp Phe Tyr Pro Gly Ala Val Thr Val Ala Trp Lys Ala  
 165 170 175  
 Asp Ser Ser Pro Val Lys Ala Gly Val Glu Thr Thr Thr Pro Ser Lys  
 180 185 190  
 Gln Ser Asn Asn Lys Tyr Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro  
 195 200 205  
 Glu Gln Trp Lys Ser His Arg Ser Tyr Ser Cys Gln Val Thr His Glu  
 210 215 220  
 Gly Ser Thr Val Glu Lys Thr Val Ala Pro Thr Glu Cys Ser  
 225 230 235

<210> SEQ ID NO 88  
 <211> LENGTH: 717  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
  
 <400> SEQUENCE: 88  
  
 atggagacag acacactcct gctatgggta ctgctgctct gggttccagg ttccactggt 60  
 gctagccagt ctgtgctgac tcagccaccc tcagcgtctg ggacccccgg gcagagggtc 120  
 accatctctt gttctggaag gaactccaac atcggaagtt atactgtaac ctggtaccag 180  
 cagctcccag gaacggcccc caaactcctc atctatagta atagtcagcg gccctcaggg 240  
 gtccctgacc gattctcagg ctccaagtct ggcacctcag cctccttggc catcagtggg 300  
 ctccagtctg aagatgaggc tgattattac tgtgtggtgt gggatgacgt gctgaatggc 360  
 ccggtgttcg gcggagggac caagctgacc gtcctaggcc aaccgaaagc ggcgccctcg 420  
 gtcactctgt tcccgcctc ctctgaggag ctcaagcca acaaggccac actggtgtgt 480  
 ctcataagtg acttctaccc gggagccgtg acagtggcct ggaaggcaga tagcagcccc 540  
 gtcaaggcgg gactggagac caccacaccc tccaacaaa gcaacaaca gtacgcggcc 600

-continued

agcagctatc tgagcctgac gcctgagcag tggaagtccc acagaagcta cagctgccag 660

gtcacgcatg aaggagcac cgtggagaag acagtggccc ctacagaatg ttcatag 717

<210> SEQ ID NO 89  
 <211> LENGTH: 5  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 89

Gly Tyr Gly Met His  
 1 5

<210> SEQ ID NO 90  
 <211> LENGTH: 18  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 90

Val Ile Ser Asn Asp Gly Ser Lys Lys Tyr Tyr Ser Asp Ser Val Lys  
 1 5 10 15

Gly Arg

<210> SEQ ID NO 91  
 <211> LENGTH: 12  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 91

Ser Ser Ser Ile Trp Leu Thr Gln Ser Leu Asp His  
 1 5 10

<210> SEQ ID NO 92  
 <211> LENGTH: 5  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 92

Gly Tyr Gly Met His  
 1 5

<210> SEQ ID NO 93  
 <211> LENGTH: 18  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 93

Val Ile Ser Asn Asp Gly Ser Lys Lys Tyr Tyr Ser Asp Ser Val Lys  
 1 5 10 15

Gly Arg

<210> SEQ ID NO 94  
 <211> LENGTH: 12  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 94

Ser Ser Ser Ile Trp Leu Thr Gln Ser Leu Asp His  
 1 5 10

<210> SEQ ID NO 95  
 <211> LENGTH: 5  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

-continued

&lt;400&gt; SEQUENCE: 95

Gly Tyr Gly Met His  
1 5

&lt;210&gt; SEQ ID NO 96

&lt;211&gt; LENGTH: 18

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 96

Val Ile Ser Asn Asp Gly Ser Lys Lys Tyr Tyr Ser Asp Ser Val Lys  
1 5 10 15

Gly Arg

&lt;210&gt; SEQ ID NO 97

&lt;211&gt; LENGTH: 12

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 97

Ser Ser Ser Ile Trp Leu Ser Gln Ser Leu Asp Gly  
1 5 10

&lt;210&gt; SEQ ID NO 98

&lt;211&gt; LENGTH: 5

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 98

Gly Tyr Gly Met His  
1 5

&lt;210&gt; SEQ ID NO 99

&lt;211&gt; LENGTH: 18

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 99

Val Ile Ser Asn Asp Gly Ser Lys Lys Tyr Tyr Ser Asp Ser Val Lys  
1 5 10 15

Gly Arg

&lt;210&gt; SEQ ID NO 100

&lt;211&gt; LENGTH: 12

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 100

Ser Ser Ser Ile Trp Leu Ser Gln Ser Leu Asp Gly  
1 5 10

&lt;210&gt; SEQ ID NO 101

&lt;211&gt; LENGTH: 5

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 101

Gly Tyr Gly Met His  
1 5

&lt;210&gt; SEQ ID NO 102

&lt;211&gt; LENGTH: 18

&lt;212&gt; TYPE: PRT

-continued

---

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 102

Val Ile Ser Asn Asp Gly Ser Lys Lys Tyr Tyr Ser Asp Ser Val Lys  
 1                   5                   10                   15

Gly Arg

<210> SEQ ID NO 103

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 103

Ser Ser Ser Ile Trp Leu Ser Gln Ser Leu Asp Gly  
 1                   5                   10

<210> SEQ ID NO 104

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 104

Gly Tyr Gly Met His  
 1                   5

<210> SEQ ID NO 105

<211> LENGTH: 18

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 105

Val Ile Ser Asn Asp Gly Ser Lys Lys Tyr Tyr Ser Asp Ser Val Lys  
 1                   5                   10                   15

Gly Arg

<210> SEQ ID NO 106

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 106

Ser Ser Ser Ile Trp Leu Thr Ser Ala Leu Asn Leu  
 1                   5                   10

<210> SEQ ID NO 107

<211> LENGTH: 5

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 107

Gly Tyr Gly Met His  
 1                   5

<210> SEQ ID NO 108

<211> LENGTH: 18

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 108

Val Ile Ser Asn Asp Gly Ser Lys Lys Tyr Tyr Ser Asp Ser Val Lys  
 1                   5                   10                   15

Gly Arg

-continued

---

<210> SEQ ID NO 109  
 <211> LENGTH: 12  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 109

Ser Ser Ser Ile Trp Phe Gly Glu Thr Val Asp Tyr  
 1                   5                   10

<210> SEQ ID NO 110  
 <211> LENGTH: 5  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 110

Glu Leu Ser Met His  
 1                   5

<210> SEQ ID NO 111  
 <211> LENGTH: 18  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 111

Gly Phe Asp Arg Glu Asp Asp Glu Thr Ile His Ala Gln Lys Phe Gln  
 1                   5                   10                   15

Gly Arg

<210> SEQ ID NO 112  
 <211> LENGTH: 11  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 112

Leu Met Val Trp Gly Asp Phe Trp Ile Gln His  
 1                   5                   10

<210> SEQ ID NO 113  
 <211> LENGTH: 5  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 113

Glu Leu Ser Met His  
 1                   5

<210> SEQ ID NO 114  
 <211> LENGTH: 18  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 114

Gly Phe Asp Arg Glu Asp Asp Glu Thr Ile His Ala Gln Lys Phe Gln  
 1                   5                   10                   15

Gly Arg

<210> SEQ ID NO 115  
 <211> LENGTH: 11  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 115

Leu Met Val Trp Gly Asp Phe Trp Ile Gln His  
 1                   5                   10

-continued

<210> SEQ ID NO 116  
 <211> LENGTH: 5  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 116

Glu Leu Ser Met His  
 1 5

<210> SEQ ID NO 117  
 <211> LENGTH: 18  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 117

Gly Phe Asp Arg Glu Asp Asp Glu Thr Ile His Ala Gln Lys Phe Gln  
 1 5 10 15

Gly Arg

<210> SEQ ID NO 118  
 <211> LENGTH: 11  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 118

Leu Met Ala Trp Asp Tyr Pro Pro Ile Gln His  
 1 5 10

<210> SEQ ID NO 119  
 <211> LENGTH: 5  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 119

Glu Leu Ser Met His  
 1 5

<210> SEQ ID NO 120  
 <211> LENGTH: 18  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 120

Gly Phe Asp Arg Glu Asp Asp Glu Thr Ile His Ala Gln Lys Phe Gln  
 1 5 10 15

Gly Arg

<210> SEQ ID NO 121  
 <211> LENGTH: 11  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 121

Leu Met Val Trp Asn Phe Pro Pro Ile Gln His  
 1 5 10

<210> SEQ ID NO 122  
 <211> LENGTH: 5  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 122

Glu Leu Ser Met His  
 1 5



-continued

<210> SEQ ID NO 123  
 <211> LENGTH: 18  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 123

Gly Phe Asp Arg Glu Asp Asp Glu Thr Ile His Ala Gln Lys Phe Gln  
 1                   5                   10                   15

Gly Arg

<210> SEQ ID NO 124  
 <211> LENGTH: 11  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 124

Leu Met Val Trp Gly Asp Phe Trp Ile Gln His  
 1                   5                   10

<210> SEQ ID NO 125  
 <211> LENGTH: 5  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 125

Ser Tyr Ala Met Ser  
 1                   5

<210> SEQ ID NO 126  
 <211> LENGTH: 18  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 126

Ala Ile Ser Gly Ser Gly Gly Gly Thr Tyr Tyr Ala Asp Ser Val Lys  
 1                   5                   10                   15

Gly Arg

<210> SEQ ID NO 127  
 <211> LENGTH: 8  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 127

Arg Gly Asp Tyr Arg Thr Asp Ile  
 1                   5

<210> SEQ ID NO 128  
 <211> LENGTH: 5  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 128

Ser Tyr Ala Met Ser  
 1                   5

<210> SEQ ID NO 129  
 <211> LENGTH: 18  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 129

Ala Ile Ser Gly Ser Gly Gly Gly Thr Tyr Tyr Ala Asp Ser Val Lys

-continued

---

 1                    5                    10                    15

Gly Arg

<210> SEQ ID NO 130  
 <211> LENGTH: 8  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 130

Arg Gly Asp Tyr Arg Thr Asp Ile  
 1                    5

<210> SEQ ID NO 131  
 <211> LENGTH: 5  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 131

Ser Tyr Ala Met Ser  
 1                    5

<210> SEQ ID NO 132  
 <211> LENGTH: 11  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 132

Ala Ile Ser Gly Ser Gly Gly Ser Thr Tyr Tyr  
 1                    5                    10

<210> SEQ ID NO 133  
 <211> LENGTH: 8  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 133

Arg Gly Asp Tyr Arg Thr Asp Ile  
 1                    5

<210> SEQ ID NO 134  
 <211> LENGTH: 5  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 134

Ser Tyr Ala Met Ser  
 1                    5

<210> SEQ ID NO 135  
 <211> LENGTH: 11  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 135

Ala Ile Ser Gly Ser Gly Gly Ser Thr Tyr Tyr  
 1                    5                    10

<210> SEQ ID NO 136  
 <211> LENGTH: 8  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 136

Arg Gly Asp Tyr Arg Thr Asp Ile  
 1                    5

-continued

---

<210> SEQ ID NO 137  
<211> LENGTH: 5  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 137

Ser Tyr Ala Met Ser  
1 5

<210> SEQ ID NO 138  
<211> LENGTH: 11  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 138

Ala Ile Ser Gly Ser Gly Gly Ser Thr Tyr Tyr  
1 5 10

<210> SEQ ID NO 139  
<211> LENGTH: 9  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 139

Val Arg Gly Ile Tyr Gly Met Asp Val  
1 5

<210> SEQ ID NO 140  
<211> LENGTH: 11  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 140

Ser Gly Asp Lys Leu Gly Asp Lys Tyr Ala Ser  
1 5 10

<210> SEQ ID NO 141  
<211> LENGTH: 7  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 141

Gln Asp Ser Asn Arg Pro Ser  
1 5

<210> SEQ ID NO 142  
<211> LENGTH: 10  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 142

Gln Ala Trp Asp Ser Ser Thr Ala Ser Val  
1 5 10

<210> SEQ ID NO 143  
<211> LENGTH: 11  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 143

Ser Gly Asp Lys Leu Gly Asp Lys Tyr Ala Ser  
1 5 10

<210> SEQ ID NO 144

-continued

---

<211> LENGTH: 7  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 144

Gln Asp Ser Asn Arg Pro Ser  
 1 5

<210> SEQ ID NO 145  
 <211> LENGTH: 10  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 145

Gln Ala Trp Asp His Ser Leu Gln His Arg  
 1 5 10

<210> SEQ ID NO 146  
 <211> LENGTH: 11  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 146

Ser Gly Asp Lys Leu Gly Asp Lys Tyr Ala Ser  
 1 5 10

<210> SEQ ID NO 147  
 <211> LENGTH: 7  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 147

Gln Asp Ser Asn Arg Pro Ser  
 1 5

<210> SEQ ID NO 148  
 <211> LENGTH: 10  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 148

Gln Ala Trp Thr Ser Ala Leu Asn Ser Gln  
 1 5 10

<210> SEQ ID NO 149  
 <211> LENGTH: 11  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 149

Ser Gly Asp Lys Leu Gly Asp Lys Tyr Ala Ser  
 1 5 10

<210> SEQ ID NO 150  
 <211> LENGTH: 7  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 150

Gln Asp Ser Asn Arg Pro Ser  
 1 5

<210> SEQ ID NO 151  
 <211> LENGTH: 10  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

-continued

---

&lt;400&gt; SEQUENCE: 151

Gln Ala Trp Thr His Ser Leu Ser Thr Leu  
1                   5                   10

&lt;210&gt; SEQ ID NO 152

&lt;211&gt; LENGTH: 11

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 152

Ser Gly Asp Lys Leu Gly Asp Lys Tyr Ala Ser  
1                   5                   10

&lt;210&gt; SEQ ID NO 153

&lt;211&gt; LENGTH: 7

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 153

Gln Asp Ser Asn Arg Pro Ser  
1                   5

&lt;210&gt; SEQ ID NO 154

&lt;211&gt; LENGTH: 10

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 154

Gln Ala Trp Thr His Ser Leu Ser Thr Leu  
1                   5                   10

&lt;210&gt; SEQ ID NO 155

&lt;211&gt; LENGTH: 13

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 155

Ser Gly Arg Asn Ser Asn Ile Gly Ser Tyr Thr Val Thr  
1                   5                   10

&lt;210&gt; SEQ ID NO 156

&lt;211&gt; LENGTH: 7

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 156

Ser Asn Ser Gln Arg Pro Ser  
1                   5

&lt;210&gt; SEQ ID NO 157

&lt;211&gt; LENGTH: 11

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 157

Ala Ala Trp Asp Asp Ser Leu Asn Gly Pro Val  
1                   5                   10

&lt;210&gt; SEQ ID NO 158

&lt;211&gt; LENGTH: 13

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 158

-continued

---

Ser Gly Arg Asn Ser Asn Ile Gly Ser Tyr Thr Val Thr  
1 5 10

<210> SEQ ID NO 159  
<211> LENGTH: 7  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 159

Ser Asn Ser Gln Arg Pro Ser  
1 5

<210> SEQ ID NO 160  
<211> LENGTH: 11  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 160

Val Val Trp Asp Asp Val Leu Asn Gly Pro Val  
1 5 10

<210> SEQ ID NO 161  
<211> LENGTH: 13  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 161

Ser Gly Arg Asn Ser Asn Ile Gly Ser Tyr Thr Val Thr  
1 5 10

<210> SEQ ID NO 162  
<211> LENGTH: 7  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 162

Ser Asn Ser Gln Arg Pro Ser  
1 5

<210> SEQ ID NO 163  
<211> LENGTH: 11  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 163

Val Val Trp Asp Asp Lys Leu Asn Gly Pro Val  
1 5 10

<210> SEQ ID NO 164  
<211> LENGTH: 13  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 164

Ser Gly Arg Asn Ser Asn Ile Gly Ser Tyr Thr Val Thr  
1 5 10

<210> SEQ ID NO 165  
<211> LENGTH: 7  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 165

Ser Asn Ser Gln Arg Pro Ser  
1 5

-continued

<210> SEQ ID NO 166  
 <211> LENGTH: 11  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 166

Val Val Trp Asp Glu Ile Leu Asn Gly Pro Val  
 1 5 10

<210> SEQ ID NO 167  
 <211> LENGTH: 11  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 167

Gln Gly Asp Ser Leu Arg Ser Tyr Tyr Ala Ser  
 1 5 10

<210> SEQ ID NO 168  
 <211> LENGTH: 7  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 168

Ala Lys Asn Asn Arg Pro Ser  
 1 5

<210> SEQ ID NO 169  
 <211> LENGTH: 10  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 169

Asn Ser Arg Asp Ser Ser Asn His Val Val  
 1 5 10

<210> SEQ ID NO 170  
 <211> LENGTH: 13  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 170

Ser Gly Arg Asn Ser Asn Ile Gly Ser Tyr Thr Val Thr  
 1 5 10

<210> SEQ ID NO 171  
 <211> LENGTH: 7  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 171

Ser Asn Ser Gln Arg Pro Ser  
 1 5

<210> SEQ ID NO 172  
 <211> LENGTH: 11  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 172

Leu Val Trp Asp Asp Val Leu Asn Gly Pro Val  
 1 5 10

<210> SEQ ID NO 173  
 <211> LENGTH: 11

-continued

---

<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 173  
  
Gln Gly Asp Ser Leu Arg Ser Tyr Tyr Ala Ser  
1                   5                   10

<210> SEQ ID NO 174  
<211> LENGTH: 7  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 174

Ala Lys Asn Asn Arg Pro Ser  
1                   5

<210> SEQ ID NO 175  
<211> LENGTH: 10  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 175

Ala Ser Arg Asn Gly Trp Asn His Val Val  
1                   5                   10

<210> SEQ ID NO 176  
<211> LENGTH: 11  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 176

Gln Gly Asp Ser Leu Arg Ser Tyr Tyr Ala Ser  
1                   5                   10

<210> SEQ ID NO 177  
<211> LENGTH: 7  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 177

Ala Lys Asn Asn Arg Pro Ser  
1                   5

<210> SEQ ID NO 178  
<211> LENGTH: 10  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 178

Ala Ser Arg Asn Gly Trp Asn His Val Val  
1                   5                   10

<210> SEQ ID NO 179  
<211> LENGTH: 11  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens  
  
<400> SEQUENCE: 179

Gln Gly Asp Ser Leu Arg Ser Tyr Tyr Ala Ser  
1                   5                   10

<210> SEQ ID NO 180  
<211> LENGTH: 7  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens



-continued

---

<400> SEQUENCE: 180

Ala Lys Asn Asn Arg Pro Ser  
1 5

<210> SEQ ID NO 181

<211> LENGTH: 10

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 181

Ala Ser Arg Asn Gly Trp Asn His Val Val  
1 5 10

<210> SEQ ID NO 182

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 182

Gln Gly Asp Ser Leu Arg Ser Tyr Tyr Ala Ser  
1 5 10

<210> SEQ ID NO 183

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 183

Ala Lys Asn Asn Arg Pro Ser  
1 5

<210> SEQ ID NO 184

<211> LENGTH: 10

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 184

Ala Ser Arg Asn Gly Trp Asn His Val Val  
1 5 10

<210> SEQ ID NO 185

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 185

Gln Gly Asp Ser Leu Arg Ser Tyr Tyr Ala Ser  
1 5 10

<210> SEQ ID NO 186

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 186

Ala Lys Asn Asn Arg Pro Ser  
1 5

<210> SEQ ID NO 187

<211> LENGTH: 10

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 187

Ala Thr Arg Asn Gly Trp Asn His Val Val



-continued

---

&lt;400&gt; SEQUENCE: 195

gtaatataca atgatggaag taagaaatat tattcagact ccgtgaaggg ccga 54

&lt;210&gt; SEQ ID NO 196

&lt;211&gt; LENGTH: 36

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 196

tccagttcca tatggctgac ccagtcctg gaccac 36

&lt;210&gt; SEQ ID NO 197

&lt;211&gt; LENGTH: 15

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 197

ggttatggca tgcac 15

&lt;210&gt; SEQ ID NO 198

&lt;211&gt; LENGTH: 54

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 198

gtaatataca atgatggaag taagaaatat tattcagact ccgtgaaggg ccga 54

&lt;210&gt; SEQ ID NO 199

&lt;211&gt; LENGTH: 36

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 199

tccagttcca tatggctgtc gcagtcctg gacggc 36

&lt;210&gt; SEQ ID NO 200

&lt;211&gt; LENGTH: 15

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 200

ggttatggca tgcac 15

&lt;210&gt; SEQ ID NO 201

&lt;211&gt; LENGTH: 54

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 201

gtaatataca atgatggaag taagaaatat tattcagact ccgtgaaggg ccga 54

&lt;210&gt; SEQ ID NO 202

&lt;211&gt; LENGTH: 36

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 202

tccagttcca tatggctgtc gcagtcctg gacggc 36

&lt;210&gt; SEQ ID NO 203

&lt;211&gt; LENGTH: 15

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

-continued

---

<400> SEQUENCE: 203  
ggttatggca tgcac 15

<210> SEQ ID NO 204  
<211> LENGTH: 54  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 204  
gtaatatcaa atgatggaag taagaaatat tattcagact ccgtgaaggg ccga 54

<210> SEQ ID NO 205  
<211> LENGTH: 36  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 205  
tccagttcca tatggctgac ctggccctg aacctg 36

<210> SEQ ID NO 206  
<211> LENGTH: 15  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 206  
ggttatggca tgcac 15

<210> SEQ ID NO 207  
<211> LENGTH: 54  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 207  
gtaatatcaa atgatggaag taagaaatat tattcagact ccgtgaaggg ccga 54

<210> SEQ ID NO 208  
<211> LENGTH: 36  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 208  
tccagttcca tatggctgac ctggccctg aacctg 36

<210> SEQ ID NO 209  
<211> LENGTH: 15  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 209  
ggttatggca tgcac 15

<210> SEQ ID NO 210  
<211> LENGTH: 54  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 210  
gtaatatcaa atgatggaag taagaaatat tattcagact ccgtgaaggg ccga 54

<210> SEQ ID NO 211  
<211> LENGTH: 36  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

-continued

---

&lt;400&gt; SEQUENCE: 211

tccagttcca tatggttcgg ggagaccgtt gactac 36

&lt;210&gt; SEQ ID NO 212

&lt;211&gt; LENGTH: 15

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 212

gaattatcca tgcac 15

&lt;210&gt; SEQ ID NO 213

&lt;211&gt; LENGTH: 54

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 213

ggttttgatc gtgaagatga tgaacaatc cacgcacaga agttccaggg caga 54

&lt;210&gt; SEQ ID NO 214

&lt;211&gt; LENGTH: 33

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 214

cttatggtgt ggggcatatt ttggatccag cac 33

&lt;210&gt; SEQ ID NO 215

&lt;211&gt; LENGTH: 15

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 215

gaattatcca tgcac 15

&lt;210&gt; SEQ ID NO 216

&lt;211&gt; LENGTH: 54

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 216

ggttttgatc gtgaagatga tgaacaatc cacgcacaga agttccaggg caga 54

&lt;210&gt; SEQ ID NO 217

&lt;211&gt; LENGTH: 33

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 217

cttatggtgt ggggcatatt ttggatccag cac 33

&lt;210&gt; SEQ ID NO 218

&lt;211&gt; LENGTH: 15

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 218

gaattatcca tgcac 15

&lt;210&gt; SEQ ID NO 219

&lt;211&gt; LENGTH: 54

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

-continued

---

<400> SEQUENCE: 219  
 ggttttgatc gtgaagatga tgaacaatc cacgcacaga agttccagg caga 54

<210> SEQ ID NO 220  
 <211> LENGTH: 33  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 220  
 cttatggcct gggactacc gccatccag cac 33

<210> SEQ ID NO 221  
 <211> LENGTH: 15  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 221  
 gaattatcca tgcac 15

<210> SEQ ID NO 222  
 <211> LENGTH: 54  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 222  
 ggttttgatc gtgaagatga tgaacaatc cacgcacaga agttccagg caga 54

<210> SEQ ID NO 223  
 <211> LENGTH: 33  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 223  
 cttatggtgt ggaacttccc cccatccag cac 33

<210> SEQ ID NO 224  
 <211> LENGTH: 15  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 224  
 gaattatcca tgcac 15

<210> SEQ ID NO 225  
 <211> LENGTH: 54  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 225  
 ggttttgatc gtgaagatga tgaacaatc cacgcacaga agttccagg caga 54

<210> SEQ ID NO 226  
 <211> LENGTH: 33  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 226  
 cttatggtgt ggggcgattt ttgatccag cac 33

<210> SEQ ID NO 227  
 <211> LENGTH: 15  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

-continued

---

<400> SEQUENCE: 227  
agctatgccca tgagc 15

<210> SEQ ID NO 228  
<211> LENGTH: 54  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 228  
gctattagtg gtagtggtgg tggcacatac tacgcagact ccgtgaaggg ccgg 54

<210> SEQ ID NO 229  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 229  
cggggcgact accggacgga catc 24

<210> SEQ ID NO 230  
<211> LENGTH: 15  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 230  
agctatgccca tgagc 15

<210> SEQ ID NO 231  
<211> LENGTH: 54  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 231  
gctattagtg gtagtggtgg tggcacatac tacgcagact ccgtgaaggg ccgg 54

<210> SEQ ID NO 232  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 232  
cggggggact accggacgga catc 24

<210> SEQ ID NO 233  
<211> LENGTH: 15  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 233  
agctatgccca tgagc 15

<210> SEQ ID NO 234  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 234  
gctattagtg gtagtggtgg tagcacatac tac 33

<210> SEQ ID NO 235  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

-continued

---

<400> SEQUENCE: 235  
cggggggact accggacgga catc 24

<210> SEQ ID NO 236  
<211> LENGTH: 15  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 236  
agctatgcca tgagc 15

<210> SEQ ID NO 237  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 237  
gctattagtg gtagtgggtg tagcacatac tac 33

<210> SEQ ID NO 238  
<211> LENGTH: 24  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 238  
cggggggact accggacgga catc 24

<210> SEQ ID NO 239  
<211> LENGTH: 15  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 239  
agctatgcca tgagc 15

<210> SEQ ID NO 240  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 240  
gctattagtg gtagtgggtg tagcacatac tac 33

<210> SEQ ID NO 241  
<211> LENGTH: 27  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 241  
gttcggggca tatacgggtat ggacgtc 27

<210> SEQ ID NO 242  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 242  
tctggagata aattggggga taaatatgct tcc 33

<210> SEQ ID NO 243  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens



-continued

---

<400> SEQUENCE: 243  
caagattcca atcggcctc a 21

<210> SEQ ID NO 244  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 244  
caggcgtggg acagcagcac tgcacggtg 30

<210> SEQ ID NO 245  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 245  
tctggagata aattggggga taaatatgct tcc 33

<210> SEQ ID NO 246  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 246  
caagattcca atcggcctc a 21

<210> SEQ ID NO 247  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 247  
caggcgtggg accactcctt gcagcacagg 30

<210> SEQ ID NO 248  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 248  
tctggagata aattggggga taaatatgct tcc 33

<210> SEQ ID NO 249  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 249  
caagattcca atcggcctc a 21

<210> SEQ ID NO 250  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 250  
caggcgtgga ccagcgcctt gaactcgcag 30

<210> SEQ ID NO 251  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

-continued

---

<400> SEQUENCE: 251  
tctggagata aattggggga taaatagct tcc 33

<210> SEQ ID NO 252  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 252  
caagattcca atcgccctc a 21

<210> SEQ ID NO 253  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 253  
caggcgtgga cgcactcct cagcacgttg 30

<210> SEQ ID NO 254  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 254  
tctggagata aattggggga taaatagct tcc 33

<210> SEQ ID NO 255  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 255  
caagattcca atcgccctc a 21

<210> SEQ ID NO 256  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 256  
caggcgtgga cccacagcct gagcacgttg 30

<210> SEQ ID NO 257  
<211> LENGTH: 39  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 257  
tctggaagga actccaacat cggaagttat actgtaacc 39

<210> SEQ ID NO 258  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 258  
agtaatagtc agcggccctc a 21

<210> SEQ ID NO 259  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

-continued

---

&lt;400&gt; SEQUENCE: 259

gcagcatggg atgacagcct gaatggcccg gtg

33

&lt;210&gt; SEQ ID NO 260

&lt;211&gt; LENGTH: 39

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 260

tctggaagga actccaacat cggaagttat actgtaacc

39

&lt;210&gt; SEQ ID NO 261

&lt;211&gt; LENGTH: 21

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 261

agtaatagtc agcggccctc a

21

&lt;210&gt; SEQ ID NO 262

&lt;211&gt; LENGTH: 33

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 262

gtggtgtggg atgacgtgct gaatggcccg gtg

33

&lt;210&gt; SEQ ID NO 263

&lt;211&gt; LENGTH: 39

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 263

tctggaagga actccaacat cggaagttat actgtaacc

39

&lt;210&gt; SEQ ID NO 264

&lt;211&gt; LENGTH: 21

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 264

agtaatagtc agcggccctc a

21

&lt;210&gt; SEQ ID NO 265

&lt;211&gt; LENGTH: 33

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 265

gtcgtgtggg atgacaagct gaatggcccg gtg

33

&lt;210&gt; SEQ ID NO 266

&lt;211&gt; LENGTH: 39

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 266

tctggaagga actccaacat cggaagttat actgtaacc

39

&lt;210&gt; SEQ ID NO 267

&lt;211&gt; LENGTH: 21

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

-continued

---

<400> SEQUENCE: 267  
agtaatagtc agcggcctc a 21

<210> SEQ ID NO 268  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 268  
gtggtgtggg acgagatcct gaatggcccg gtg 33

<210> SEQ ID NO 269  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 269  
caaggagaca gcctcagaag ctattatgca agc 33

<210> SEQ ID NO 270  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 270  
gctaaaaaca accggcctc a 21

<210> SEQ ID NO 271  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 271  
tgtaactccc gggacagcag taaccatgtg gta 33

<210> SEQ ID NO 272  
<211> LENGTH: 39  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 272  
tctggaagga actccaacat cggaagttat actgtaacc 39

<210> SEQ ID NO 273  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 273  
agtaatagtc agcggcctc a 21

<210> SEQ ID NO 274  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 274  
ctcgtgtggg acgacgtcct gaatggcccg gtg 33

<210> SEQ ID NO 275  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

-continued

---

<400> SEQUENCE: 275  
caaggagaca gcctcagaag ctattatgca agc 33

<210> SEQ ID NO 276  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 276  
gctaaaaaca accggcctc a 21

<210> SEQ ID NO 277  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 277  
gcgtcccga acggctggaa ccatgtggta 30

<210> SEQ ID NO 278  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 278  
caaggagaca gcctcagaag ctattatgca agc 33

<210> SEQ ID NO 279  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 279  
gctaaaaaca accggcctc a 21

<210> SEQ ID NO 280  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 280  
gcgtcccga acggctggaa ccatgtggta 30

<210> SEQ ID NO 281  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 281  
caaggagaca gcctcagaag ctattatgca agc 33

<210> SEQ ID NO 282  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 282  
gctaaaaaca accggcctc a 21

<210> SEQ ID NO 283  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

-continued

---

<400> SEQUENCE: 283  
gcgtcccgga acggctggaa ccatgtggta 30

<210> SEQ ID NO 284  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 284  
caaggagaca gcctcagaag ctattatgca agc 33

<210> SEQ ID NO 285  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 285  
gctaaaaaca accggcctc a 21

<210> SEQ ID NO 286  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 286  
gcgtcccgga acggctggaa ccatgtggta 30

<210> SEQ ID NO 287  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 287  
caaggagaca gcctcagaag ctattatgca agc 33

<210> SEQ ID NO 288  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 288  
gctaaaaaca accggcctc a 21

<210> SEQ ID NO 289  
<211> LENGTH: 30  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 289  
gcgaccgga acggctggaa ccatgtggta 30

<210> SEQ ID NO 290  
<211> LENGTH: 33  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 290  
caaggagaca gcctcagaag ctattatgca agc 33

<210> SEQ ID NO 291  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

-continued

&lt;400&gt; SEQUENCE: 291

gctaaaaaca accggcctc a

21

&lt;210&gt; SEQ ID NO 292

&lt;211&gt; LENGTH: 30

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 292

gcgaccggga acggctggaa ccatgtggta

30

What is claimed is:

**1.** An isolated antigen binding protein that binds a human IL-18 receptor,

wherein said antigen binding protein comprises:

an antibody heavy chain comprising (a) a CDRH1 of SEQ ID NO:104; (b) a CDRH2 of SEQ ID NO:105; and (c) a CDRH3 of SEQ ID NO:106; and

an antibody light chain comprising (a) a CDRL1 of SEQ ID NO:173; (b) a CDRL2 of SEQ ID NO:174; and (c) a CDRL3 of SEQ ID NO:175.

**2.** The antigen binding protein of claim 1, wherein said antigen binding protein comprises an antibody heavy chain of SEQ ID NO:6 or an antibody light chain of SEQ ID NO:29.

**3.** The antigen binding protein of claim 1, wherein said antigen binding protein comprises an antibody heavy chain of SEQ ID NO:6 and an antibody light chain of SEQ ID NO:29.

**4.** The antigen binding protein of claim 1, wherein said antigen binding protein is a monoclonal antibody, a recombinant antibody, a bispecific antibody, or a fragment thereof.

**5.** The antigen binding protein of claim 4, wherein said antibody fragment is a minibody, a domain antibody, a F(ab) fragment, a F(ab') fragment, a F(ab')<sub>2</sub> fragment, a Fv fragment, a scFv fragment, a Fd fragment, a diabody, or a single chain antibody molecule.

**6.** The antigen binding protein of claim 1, wherein said antigen binding protein is of an IgG1-, IgG2-IgG3- or IgG4-type.

**7.** The antigen binding protein of claim 6, wherein said antigen binding protein is of an IgG2-type.

**8.** A pharmaceutical composition comprising the antigen binding protein of claim 1, and pharmaceutically acceptable carrier.

**9.** An isolated antigen binding protein comprising an IgG heavy chain of SEQ ID NO:73, or an IgG light chain of SEQ ID NO:75.

**10.** An isolated antigen binding protein comprising an IgG heavy chain of SEQ ID NO:73 and an IgG light chain of SEQ ID NO:75.

**11.** A pharmaceutical composition comprising the antigen binding protein of claim 10, and pharmaceutically acceptable carrier.

**12.** An isolated nucleic acid molecule encoding:

a) an antibody heavy chain comprising (a) a CDRH1 of SEQ ID NO:104; (b) a CDRH2 of SEQ ID NO:105; and (c) a CDRH3 of SEQ ID NO:106;

b) an antibody light chain comprising (a) a CDRL1 of SEQ ID NO:173; (b) a CDRL2 of SEQ ID NO:174; and (c) a CDRL3 of SEQ ID NO:175; or

c) the antibody heavy chain of a) and antibody light chain of b).

**13.** The isolated nucleic acid molecule of claim 12, wherein said nucleic acid molecule is operably linked to a control sequence.

**14.** A vector comprising a nucleic acid molecule of claim 13.

**15.** An isolated host cell transformed with the vector of claim 14.

**16.** A vector comprising a nucleic acid molecule of claim 12.

**17.** An isolated host cell transformed with the vector of claim 16.

**18.** An isolated host cell comprising:

a) a nucleic acid encoding an antibody heavy chain comprising (a) a CDRH1 of SEQ ID NO:104; (b) a CDRH2 of SEQ ID NO:105; and (c) a CDRH3 of SEQ ID NO:106; and

b) a nucleic acid encoding an antibody light chain comprising (a) a CDRL1 of SEQ ID NO:173; (b) a CDRL2 of SEQ ID NO:174; and (c) a CDRL3 of SEQ ID NO:175.

**19.** The isolated host cell of claim 18, wherein said host cell comprises a nucleic acid encoding an IgG heavy chain of SEQ ID NO:73 and a nucleic acid encoding an IgG light chain of SEQ ID NO:75.

**20.** A process for preparing an isolated antigen binding protein comprising an antibody heavy chain comprising (a) a CDRH1 of SEQ ID NO:104; (b) a CDRH2 of SEQ ID NO:105; and (c) a CDRH3 of SEQ ID NO:106; and

an antibody light chain (a) a CDRL1 of SEQ ID NO:173; (b) a CDRL2 of SEQ ID NO:174; and (c) a CDRL3 of SEQ ID NO:175, said process comprising the step of culturing a host cell that secretes said binding protein, and isolating said binding protein.

**21.** The process of claim 20, wherein the host cell comprises a nucleic acid encoding an IgG heavy chain of SEQ ID NO:73 and a nucleic acid encoding an IgG light chain of SEQ ID NO:75.

\* \* \* \* \*

UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

PATENT NO. : 8,257,707 B2  
APPLICATION NO. : 12/670112  
DATED : September 4, 2012  
INVENTOR(S) : Dirk E. Smith et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

On the Title Page

Item (75) Inventors: “Duncan Cochrane, Granta Park (GA); Louise Conroy, Granta Park (GA)”  
should read -- Duncan Cochrane, Cambridge (UK); Louise Conroy,  
Cambridge (UK) --.

In the Claims

Col. 181,  
Claim 4, lines 33-34: “a monoclonal antibody, a recombinant antibody” should read  
-- a monoclonal antibody, a human antibody, a recombinant antibody, a  
chimeric antibody, a humanized antibody, --.

Col. 181,  
Claim 11, line 54: “the antigen” should read -- an antigen --.

Signed and Sealed this  
Fifteenth Day of July, 2014



Michelle K. Lee  
*Deputy Director of the United States Patent and Trademark Office*