



US008318159B2

(12) **United States Patent**
Adam et al.(10) **Patent No.:** **US 8,318,159 B2**
(45) **Date of Patent:** **Nov. 27, 2012**

- (54) **ANTI-IGF ANTIBODIES**
- (75) Inventors: **Paul Adam**, Vienna (AT); **Eric Borges**, Maria Enzersdorf (AT)
- (73) Assignee: **Boehringer Ingelheim International GmbH**, Ingelheim am Rhein (DE)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 68 days.
- (21) Appl. No.: **12/636,195**
- (22) Filed: **Dec. 11, 2009**
- (65) **Prior Publication Data**
US 2010/0150940 A1 Jun. 17, 2010
- (30) **Foreign Application Priority Data**
Dec. 12, 2008 (EP) 08171554
- (51) **Int. Cl.**
A61K 39/395 (2006.01)
C07K 16/26 (2006.01)
- (52) **U.S. Cl.** **424/130.1**; 530/387.9; 530/388.24; 424/133.1; 424/139.1; 424/140.1; 424/145.1
- (58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,773,919	A	11/1973	Boswell et al.
4,342,566	A	8/1982	Theofilopoulos et al.
6,342,221	B1	1/2002	Thorpe et al.
6,696,245	B2	2/2004	Winter et al.
6,991,790	B1	1/2006	Lam et al.
7,020,563	B1	3/2006	Bentley et al.
7,037,498	B2	5/2006	Cohen et al.
7,060,268	B2	6/2006	Andy et al.
7,438,911	B2	10/2008	Shitara et al.
7,498,415	B2	3/2009	Shitara et al.
7,749,966	B2	7/2010	Raso
2006/0165695	A1	7/2006	Shitara et al.
2007/0196376	A1	8/2007	Raeber et al.
2009/0016967	A1	1/2009	Schnapp et al.
2010/0099147	A1	4/2010	Hariharan et al.
2010/0196395	A1	8/2010	Adam et al.

FOREIGN PATENT DOCUMENTS

CA	2473039	A1	7/2003
CA	2483848	A1	11/2003
CA	2536288	A1	3/2005
CA	2540133	A1	3/2005
CA	2540138	A1	3/2005
EP	0123228	A2	10/1984
EP	0292656	A1	11/1988
EP	0492552	A1	7/1992
EP	0700994	A1	3/1996
EP	1505075	A1	2/2005
JP	2003310275	A	11/2003
WO	8500831	A1	2/1985
WO	8911297	A1	11/1989
WO	9000562	A1	1/1990
WO	9429348	A2	12/1994
WO	9525794	A1	9/1995
WO	9928347	A1	6/1999

WO	02053596	A2	7/2002
WO	02056910	A1	7/2002
WO	03002609	A2	1/2003
WO	03050531	A2	6/2003
WO	03059951	A2	7/2003
WO	03093317	A1	11/2003
WO	03100008	A2	12/2003
WO	03106621	A2	12/2003
WO	2004003019	A2	1/2004
WO	2004058821	A2	7/2004
WO	2004071529	A2	8/2004
WO	2004083248	A1	9/2004
WO	2005005635	A2	1/2005
WO	2005016970	A2	2/2005
WO	2005018671	A1	3/2005
WO	2005027970	A1	3/2005
WO	2005028515	A1	3/2005
WO	2005058967	A2	6/2005
WO	2005061541	A1	7/2005
WO	2006008639	A1	1/2006
WO	2006069202	A2	6/2006
WO	2006125640	A2	11/2006
WO	2007012614	A2	2/2007
WO	2007042309	A2	4/2007
WO	2007070432	A2	6/2007
WO	2007115814	A2	10/2007
WO	2007118214	A2	10/2007

(Continued)

OTHER PUBLICATIONS

Sell, Christian, et al; Effect of a Null Mutation of the Insulin-Like Growth Factor I Receptor Gene on Growth and Transformation of Mouse Embryo Fibroblasts; *Molecular and Cellular Biology* (1994) vol. 14, No. 6 pp. 3604-3612.

Sell, Christian, et al; Simian Virus 40 Large Tumor Antigen is Unable to Transform Mouse Embryonic Fibroblasts lacking Type 1 Insulin-Like Growth Factor Receptor; *Proc. Natl. Acad. Sci. USA* vol. 90 pp. 11217-11221, 1993.

Shukla, Abhinav, A., et al; Downstream Processing of Monoclonal Antibodies—Application of Platform Approaches; *Journal of Chromatography* (2007) vol. 848 pp. 28-39.

Srinivasan, Mythily, et al; Immunomodulatory Peptides From IgSF Proteins: A Review; *Current Protein and Peptide Science* (2005) vol. 6, No. 2 pp. 185-196.

Strumberg, Dirk; Preclinical and Clinical Development of the Oral Multikinase Inhibitor Sorafenib in Cancer Treatment; *Drugs of Today* (2005) vol. 41, No. 12 pp. 773-784.

Takanami, Iwao, et al; Insulin-Like Growth Factor-II as a Prognostic Factor in Pulmonary Adenocarcinoma; *Journal of Surgical Oncology* (1996) vol. 61 pp. 205-208.

Tsai, J. F., et al; Serum Insulin-Like Growth Factor-II as a Serologic Marker of Small Hepatocellular Carcinoma; *Scandinavian Journal of Gastroenterology* (2005) vol. 40 pp. 68-75.

(Continued)

Primary Examiner — Marianne P Allen(74) *Attorney, Agent, or Firm* — Michael P. Morris; Edouard G. Lebel(57) **ABSTRACT**

Antibody molecules, in particular fully human antibodies that bind to human IGF-1 and cross-react with IGF-2 such that binding of IGF-1 and IGF-2 to the IGF-1 receptor is prevented and IGF-1 receptor-mediated signaling is inhibited. The antibodies do not bind to insulin and thus do not affect the mitogenic properties of insulin that are mediated by its binding to the insulin receptors. The antibodies are useful for the treatment of hyperproliferative diseases, in particular cancer.

13 Claims, 25 Drawing Sheets

FOREIGN PATENT DOCUMENTS

WO	2007126876	A2	11/2007
WO	2007141626	A1	12/2007
WO	2008005469	A2	1/2008
WO	2008079324	A1	7/2008
WO	2008079849	A2	7/2008
WO	2008098917	A2	8/2008
WO	2008108986	A2	9/2008
WO	2008115470	A2	9/2008
WO	2008116103	A2	9/2008
WO	2008144345	A2	11/2008
WO	2008144720	A2	11/2008
WO	2008152422	A2	12/2008
WO	2008155387	A2	12/2008
WO	WO 2008/155387	*	12/2008
WO	2009005673	A1	1/2009
WO	2009006336	A1	1/2009
WO	2009016164	A1	2/2009
WO	2009017679	A2	2/2009
WO	2009019117	A1	2/2009
WO	2009021054	A2	2/2009
WO	2009032145	A1	3/2009
WO	2009032782	A2	3/2009
WO	2009039457	A1	3/2009
WO	2009045361	A2	4/2009
WO	2009045389	A2	4/2009
WO	2009079587	A2	6/2009
WO	2009120767	A1	10/2009
WO	2009126304	A1	10/2009
WO	2009137378	A2	11/2009
WO	2009137758	A2	11/2009
WO	2009149185	A2	12/2009
WO	2010034441	A1	4/2010
WO	2010036767	A1	4/2010
WO	2010045315	A1	4/2010
WO	2010048123	A2	4/2010
WO	2010052344	A2	5/2010
WO	2010062896	A1	6/2010
WO	2010066868	A2	6/2010
WO	2010069858	A1	6/2010
WO	2010075511	A1	7/2010

OTHER PUBLICATIONS

Wang, Zheng, et al; Expression of IGF-II in Early Experimental Hepatocellular Carcinomas and its Significance in Early Diagnosis; *World Journal of Gastroenterology* (2003) vol. 9 pages 267-270.

Woodson, Karen, et al; Loss of Insulin-Like Growth Factor-II Imprinting and the Presence of Screen-Detected Colorectal Adenomas in Women; *Journal of the National Cancer Institute* (2004) vol. 96, No. 5 pages 407-410.

Yao, Xiaoming, et al; A Methylated Oligonucleotide Inhibits IGF2 Expression and Enhances Survival in a Model of Hepatocellular Carcinoma; *The Journal of Clinical Investigation* (2003) vol. 111, No. 2 pp. 265-273.

Yao, Xiaoming, et al; A Novel Orthotopic Tumor Model to Study Growth Factors and Oncogenes in Hepatocarcinogenesis; *Clinical Cancer Research* (2003) vol. 9 pp. 2719-2726.

Yelton, Dale, E. et al; Affinity Maturation of the BR96 Anti-Carcinoma Antibody by Codon-Based Mutagenesis; *The American Association of Immunologists* (1995) vol. 155 pp. 1994-2004.

Zapata, Gerardo, et al; Engineering Linear F(ab')₂ Fragments for Efficient Production in *Escherichia coli* and Enhanced Antiproliferative Activity; *Protein Engineering* (1995) vol. 8, No. 10 pp. 1057-1062.

Zhao, Ronghua, et al; Positive Correlation of Insulin-Like Growth Factor-II with Proliferating Cell Index in Patients with Colorectal Neoplasia; *Cancer Epidemiology, Biomarkers and Prevention* (2005) vol. 14 pp. 1819-1822.

Barbas, Carlos F., et al; In Vitro evolution of a Neutralizing Human Antibody to Human Immunodeficiency Virus Type 1 to Enhance Affinity and Broaden Strain Cross-Reactivity; *Proc. Natl. Acad. Sci. USA* (1994) vol. 91 pp. 3809-1813.

Burtrum, Douglas, et al; A Fully Human Monoclonal Antibody to the Insulin-Like Growth Factor I Receptor Blocks Ligand-dependent Signaling and Inhibits Human Tumor Growth in Vivo; *Cancer Research* (2003) vol. 63 pp. 8912-8921.

Cascieri, Margaret, A., et al; Identification of the Insulin-Like Growth Factor I (IGF I) Epitopes Recognized by Monoclonal and Polyclonal Antibodies to IGF I; *Endocrinology* (1990) vol. 126, No. 6 pp. 2773-2777.

Chen, Jian-Wen, et al; Free Rather than Total Circulating Insulin-Like Growth Factor-I Determines the Feedback on Growth Hormone Release in Normal Subjects; *The Journal of Clinical Endocrinology & Metabolism* (2005) vol. 90, No. 1 pp. 366-371.

Chothia, Cyrus, et al; Canonical Structures for the Hypervariable Regions of Immunoglobulins; *Journal Molecular Biology* (1987) vol. 196 pp. 901-917.

Cruz-Correa, M., et al; IGF2 Loss of Imprinting: A Potential Heritable Risk Factor for Colorectal Cancer; *Gastroenterology* (2004) vol. 126 pp. 1190-1201.

Cui, Hengmi, et al; Loss of IGF2 Imprinting: A Potential Marker of Colorectal Cancer Risk; *Science* (2003) vol. 299 pp. 1753-1755.

Dufner, Almut, et al; Ribosomal S6 Kinase Signaling and the Control of Translation; *Experimental Cell Research* (1999) vol. 253 pp. 100-109.

Feng, Yang, et al; Novel Human Monoclonal Antibodies to Insulin-Like Growth Factor (IGF)-II That Potently Inhibit the IGF Receptor Type I Signal Transduction Function; *Molecular Cancer Therapy* (2006) vol. 5, No. 1 pp. 114-120.

Frasca, F., et al; Insulin Receptor Isoform A, a Newly Recognized, High-Affinity Insulin-Like Growth Factor II Receptor in Fetal and Cancer Cells; *Molecular and Cellular Biology* (1999) vol. 19, No. 5 pp. 3278-3288.

Freier, S., et al; Expression of the Insulin-Like Growth Factors and their Receptors in Adenocarcinoma of the Colon; *Gut* (1999) vol. 44 pp. 704-708.

Fukuzawa, Ryuji, et al; High Frequency of Inactivation of the Imprinted H19 gene in "Sporadic" Hepatoblastoma; *International Journal of Cancer* (1999) vol. 82 pp. 490-497.

Goetsch, Liliane, et al; A Recombinant Humanized Anti-Insulin-Like Growth Factor Receptor Type I Antibody (h7C10) Enhances the Antitumor Activity of Vinorelbine and Anti-Epidermal Growth Factor Receptor Therapy Against Human Cancer Xenografts; *International Journal of Cancer* (2005) vol. 113 pp. 316-328.

Goya, Masato, et al; Growth Inhibition of Human Prostate Cancer Cells in Human Adult Bone implanted into Nonobese Diabetic/Severe Combined Immunodeficient Mice by a Ligand-Specific Antibody to Human Insulin-Like Growth Factors; *Cancer Research, American Association for Cancer Research* (2004) vol. 64, No. 17 pp. 6252-6258.

Haenel, Cornelia et al; Characterization of High-Affinity Antibodies by Electrochemiluminescence-Based Equilibrium Titration; *Analytical Biochemistry* (2005) vol. 339 pp. 182-184.

Hassan, A. Bassim., et al; Insulin-Like Growth factor II Supply Modifies growth of Intestinal Adenoma in ApcMin/+Mice; *Cancer Research* (2000) vol. 60 pp. 1070-1076.

Hawkins, Robert E., et al; Selection of Phage Antibodies by Binding Affinity Mimicking Affinity Maturation; *Journal Mol. Biol.* (1992) vol. 226 pp. 889-896.

Jackson, Jeffrey R., et al; In Vitro Antibody Maturation: Improvement of High Affinity, Neutralizing Antibody Against IL-1b; *Journal of Immunology* (1995) vol. 154, No. 7 pp. 3310-3319.

Jerome L, et al; Deregulation of the IGF Axis in Cancer: Epidemiological Evidence and Potential Therapeutic Interventions; *Endocrine-Related Cancer* (2003) vol. 10 pp. 561-578.

Kipriyanov, Sergey M., et al; Generation and Production of Engineered Antibodies; *Molecular Biotechnology* (2004) vol. 26 pp. 39-60.

Knappik, Achim, et al; Fully Synthetic Human Combinatorial Antibody Libraries (HuCAL) Based on Modular Consensus Frameworks and CDRs Randomized with Trinucleotides; *Journal Molecular Biology* (2000) vol. 296 pp. 57-86.

Kolb, E. Anders, et al; Initial Testing (Stage 1) of a Monoclonal Antibody (SCH 717454) Against the IGF-1 Receptor by the Pediatric Preclinical Testing Program; *Pediatr Blood Cancer* (2008) vol. 50 pp. 1190-1197.

Kulik, George et al; Antiapoptotic Signalling by the Insulin-Like Growth Factor I receptor, Phosphatidylinositol 3-Kinase, and AKT; *Molecular and Cellular Biology* (1997) vol. 17, No. 3 pp. 1595-1606.

- Leroith, Derek; The Insulin-Like Growth Factor System; *Experimental Diab. Res.* (2003) vol. 4 pp. 205-212.
- Li, Shu-Rui, et al; Differential Expression Patterns of the Insulin-Like Growth Factor 2 Gene in Human Colorectal Cancer; *Tumor Biology* (2004) vol. 25 pp. 62-68.
- Lin, Yvonne, S., et al; Preclinical Pharmacokinetics, Interspecies scaling, and Tissue Distribution of a Humanized Monoclonal Antibody Against Vascular endothelial growth Factor; *The Journal of Pharmacology and Experimental Therapeutics* (1999) vol. 288 pp. 371-378.
- Lowman, Henry, B., et al; Selecting High-Affinity Binding Proteins by Monovalent Phage Display; *Biochemistry* (1991) vol. 30, No. 45 pp. 10832-10837.
- Lund, Per, et al; Autocrine Inhibition of Chemotherapy Response in Human Liver Tumor Cells by Insulin-Like Growth Factor-II; *Cancer Letters* (2004) vol. 206 pp. 85-96.
- Manara, Maria C., et al; Preclinical in Vivo Study of New Insulin-Like Growth Factor-I Receptor-Specific Inhibitor in Ewing's Sarcoma; *Clinical Cancer Research* (2007) vol. 13, No. 4 pp. 1322-1330.
- Manes, Santos, et al; Functional Epitope Mapping of Insulin-Like Growth Factor I (IGF-I) by Anti-IGF-I Monoclonal Antibodies; *Endocrinology* (1997) vol. 138, No. 3 pp. 905-915.
- Manes, Santos, et al; Physical Mapping of Human Insulin-Like Growth Factor-I Using Specific Monoclonal Antibodies; *Journal of endocrinology* (1997) vol. 154 pp. 293-302.
- Marks, James D., et al; By-Passing Immunization: Building High Affinity Human Antibodies by Chain Shuffling; *Bio/Technology* (1992) vol. 10 pp. 779-783.
- Miyamoto, Shin'ichi, et al; Blockade of Paracrine Supply of Insulin-Like Growth Factors Using Neutralizing Antibodies Suppresses the Liver Metastasis of Human Colorectal Cancers; *Clin Cancer Research* (2005) vol. 11, No. 9 pp. 3494-3502.
- Moorehead, Roger A., et al; Transgenic Overexpression of IGF-II Induces Spontaneous Lung Tumors: A Model for Human Lung Adenocarcinoma; *Oncogene* (2003) vol. 22 pp. 853-857.
- Nagy, Zoltan A., et al; Fully Human, HLA-DR-Specific Monoclonal Antibodies efficiently Induce Programmed Death of Malignant Lymphoid Cells; *Nature Medicine* (2002) vol. 8, Issue 8 pp. 801-807.
- Ng, Irene OL, et al; "Hepatocellular Carcinoma Expression of Insulin-Like Growth Factor II mRNA in Hepatocellular Carcinoma" *Journal of Gastroenterology and Hepatology* (1998) vol. 13, p. 152-157.
- Pandini, Giuseppe, et al.; "Insulin/Insulin-like Growth Factor I Hybrid Receptors Have Different Biological Characteristics Depending on the Insulin Receptor Isoform Involved"; *The Journal of Biological Chemistry*; (2002) V. 277, Issue: 42, pp. 39684-39695.
- Pollak, M.N., et al; Pharmacodynamic Properties of the Anti-IGF-IR Monoclonal Antibody CP-751,871 in Cancer Patients; *American Society of Clinical Oncology (ASCO)* vol. 25, No. 18Sp. 3587, 2007.
- Pollak, Michael N. et al.; "Insulin-Like Growth Factors and Neoplasia" *Nature Reviews Cancer*, (2004) vol. 4, pp. 505-518.
- Quinn Kathryn A., et al; insulin-Like Growth Factor Expression in Human Cancer Cell Lines; *The Journal of Biological Chemistry* (1996) vol. 271, No. 19 pp. 11477-11483.
- Rauchenberger, Robert, et al; Human Combinatorial Fab Library Yielding Specific and Functional Antibodies Against the Human Fibroblast Growth Factor Receptor 3*; *The Journal of Biological Chemistry* (2003) vol. 278, No. 40 pp. 38194-38205.
- Renehan, Andrew, G., et al; Circulating Insulin-Like Growth Factor II and Colorectal Adenomas; *The Journal of Clinical Endocrinology and Metabolism* (2000) vol. 85, No. 9 pp. 3402-3408.
- Renehan, Andrew, G., et al; Elevated Serum Insulin-Like Growth Factor (IGF)-II and IGF Binding Protein-2 in Patients with Colorectal Cancer; *British Journal of Cancer* (2000) vol. 83, No. 10 pp. 1344-1350.
- Reverts, Hilde, et al; Nanobodies as Novel Agents for Cancer Therapy; *Experts Opin. Biol. Ther.* (2005) vol. 5, No. 1 pp. 111-124.
- Rubin, Raphael, et al; Biology of Disease: Insulin-Like Growth Factor-I Receptor; *Laboratory Investigation* (1995) vol. 73, No. 3 pp. 311-331.
- Rusell, William, E., et al; Inhibition of the Mitogenic Effects of Plasma by a Monoclonal Antibody to Somatomedin C; *Proc. Natl. Acad. Sci. USA* (1984) vol. 81 pp. 2389-2392.
- Schier, Robert, et al; Identification of Functional and Structural Amino-Acid residues by Parsimonious Mutagenesis; *Gene* (1996) vol. 169 pp. 147-155.
- Scotlandi, Katia, et al; Insulin-like Growth Factor I Receptor-Mediated Circuit in Ewing's Sarcoma/Peripheral Neuroectodermal Tumor: A Possible Therapeutic Target; *Cancer Research* (1996) vol. 56 pp. 4570-4574.
- Reinberg, Steven "Rare Gene Mutation Plays Role in Longevity" *Healthday News*, published by US News & World Report, Mar. 4, 2008; pp. 1-3.
- Green, Larry L. "Antibody engineering via genetic engineering of the mouse: XenoMouse strains are a vehicle for the facile generation of therapeutic human monoclonal antibodies" *Journal of Immunological Methods*; 231, (1991) 11-23.
- Jirtle Randy L. "IGF2 Loss of Imprinting: A Potential Heritable Risk Factor for Colorectal Cancer"; *Gastroenterology* (2004) vol. 126 pp. 1190-1201.
- Krebs, Barbara, et al "High-throughput generation and engineering of recombinant human antibodies" *Journal of Immunological Methods* 254 (2001) pp. 67-84.
- Casset, Florence et al. "A peptide mimetic of an anti-CD4 monoclonal antibody by rational design" *Biochemical and Biophysical Research Communications*, (2003) vol. 307, pp. 198-205.
- Chen, Yvonne, et al. "Selection and Analysis of an Optimized Anti-VEGF Antibody: Crystal Structure of an Affinity-matured Fab in Complex with Antigen" *Journal of Molecular Biology*, (1999) vol. 293, pp. 865-881.
- De Pascalis, Roberto, et al. "Grafting of "Abbreviated" Complementarity-Determining Regions Containing Specificity-Determining Residues Essential for Ligand Contact to Engineer a Less Immunogenic Humanized Monoclonal Antibody" *The Journal of Immunology* (2002) vol. 169 pp. 3076-3084.
- Green, Larry L. "Antibody engineering via genetic engineering of the mouse: XenoMouse strains are a vehicle for the facile generation of therapeutic human monoclonal antibodies" *Journal of Immunological Methods*, (1999) vol. 231, pp. 11-23.
- MacCallum, Robert M. et al. "Antibody-antigen Interactions: Contact Analysis and Binding Site Topography" *Journal of Molecular Biology* (1996) vol. 262, pp. 732-745.
- Rudikoff, Stuart et al. "Single Amino Acid Substitution Altering Antigen-Binding Specificity" *Proc. Natl. Acad. Sci. USA* (1982) vol. 79, pp. 1979-1983.
- Wu, Henry, et al "Humanization of a Murine Monoclonal Antibody by Simultaneous Optimization of Framework and CDR Residues" (1999) *Journal of Molecular Biology* vol. 294, pp. 151-162.

* cited by examiner




Fig. 1A




Fig. 1B




Fig. 1C




Fig. 1D




Fig. 1E




Fig. 1F




Fig. 1G




Fig. 2A




Fig. 2B




Fig. 3A




Fig. 3B




Fig. 4A




Fig. 4B




Fig. 4C




Fig. 4D




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9





Fig. 10



GPETLCGAELVDALQFVCGDRGFYFNKPTGYGSSRRAPQ
FGIVDECCFRSCDLRRLEMYCAPLKPAKSA

Fig. II

60814

VH3 Amino Acid Sequence:

QVELVESGGGLVQPGGSLRLSCLAAAGFTFSNYWMHWVRQAPKGLEWVSGISGWSSWYYADSVKGRFT
 ISRDNSKNITLYLQMNLSLRAEDTAVYYCARFGIDAYTKVYFDYWGQGLVTVSS

VH3 DNA Sequence:

CAGGTGGAATTGGTGGAAAGCGCGGCCCTGGTGCAACCGGGCCAGCCCTGAGCTGCGCGGCCCTCCGGATT
 TACCTTTTCATAATTATTGGATGCATTGGGTGCCCCAGCCCTGGGAAGGCTCCGAGTGGGTGAGCGGTATCTCTGGTT
 GGTCTAGCTGGACCTATATATGCGGATAGCGTGAAGGCCGTTTACCATTTACCGTGATAATTCGAAAACACCCCTGTAT
 CTGCAAAATGAACAGCCCTGCGTGGGAAGATACGGCCGTGATTAATTGCGCGCCTTTTGGTATTGATGCTTATACCTAAGGT
 TTATTTGATTATTGGGCCAAGGCACCCTGGTGACGGTTAGCTCA

60814

VA3 Amino Acid Sequence:

DIELTQPPSVVAPGQTARISCSGDNIPLKYVSWYQOKPGQAPVLVIHDDNKRPSGIPERFSGNSGN
 TATLTIISGTQAEDEADYYCSSWDTLDFNVFGGTKLTVLG (Q)

VA3 DNA Sequence:

GATATCGAACTGACCCAGCCCTTCAGTGAGCGTTGCACCAGGTCAGACCGCGGTATCTCGTGTAGCGCGGATAATAT
 TCCTCTTAAGTATGTTCTTGGTACCAGCAGAAACCCGGCAGCCAGTTCTTGTGATTCATGATGATAATAAGCGTC
 CCTCAGGCATCCCGGAACGCTTTAGCGGATCCAACAGCGGCAACCCGACCCCTGACCATTAGCGGCACCTCAGGCGGAA
 GACGAAGCGGATTATTATTGCTCTTCTTGGGATACTCTTGATATTTTAAATGFTTGGCGGCGCACGAAGTTAACCGT
 CCTAGGT

Fig. 12A

60819

VH3 Amino Acid Sequence:

QVELVESGGGLVQPGGSLRLSLCAASGFTFSNYMHWVRQAPGKLEWVSGISGWSSTYYADSVKGRFT
ISRDNSKNTLLYLQMNLSLRAEDTAVYYCARFGIDAYTKVYFDYWGQGTLLVTVSS

VH3 DNA Sequence:

CAGGTGGAATTGGTGAAAGCGCGGCCCTGGTGCAACCGGGCGGAGCCCTGCGTCTGAGCTGCCGGCCCTCCGGATT
TACCTTTTCTAATTATTGGATGCAATGGGTGGCCAGCCCTGGGAAGGCTCGAGTGGGTGAGCGGTATCTCTGGTT
GGCTAGCTGGACCTATTATGCGGATAGCGTGAAGGCCGTTTACCAATTCACGTGATAAATTCGAAAACAACCCCTGTAT
CTGCAAAATGAACAGCCCTGCGTGCGGAAGATACGGCCGTTTATTTGGTATTGCGCCGTTTGGTATTCATGCTTATATACTAAGGT
TTATTTTGATTATTGGGGCCCAAGCCACCTGGTGACGGTTAGCTCA

60819

Vλ3 Amino Acid Sequence:

DIELTQPPSVSAPGQTARISCSGDNIPKLYVSWYQQKPGQAPVLVHDDNKRPSGIPERFSGSNSGNTATLTIS
GTQAEDEADYYCCSYDYFPKFFVFGGKLTVLG (Q)

Vλ3 DNA Sequence:

GATATCGAACTGACCCAGCCGCTTCAGTGAGCGTTGCACCAGGTCAGACCCGCGTATCTCGTGTAGCGCGGATAATATF
TCCCTCTTAAGTATGTTCTTGGTACCAGCAAACCCGGCAGGCCAGTTCTTGTGATTCATGATGATAATAAGCGTC
CCTCAGGCATCCCGGAACGCTTTAGCGGATCCAAACAGCGCAACCCGACCCCTGACCATTAGCGGCACTCAGCGGAA
GACGAAAGCGGATTATTATTGCCAGTCTTATGATTATTTCCCTAAGTTTGTGTTGGCGCGGCACGAAAGTTAACCCGT
CCTAGGT

Fig. 12B

60833

VH3 Amino Acid Sequence:

QVELVESGGGLVQPGGSLRLSCAASGFTFTSYW
 NSKNILYLQMNSLRAEDTAVYYCARNMYTHFI

VH3 DNA Sequence:

CAGGTGGAATTGGTGGAAAGCGGGCGGCCCTGGTGT
 TACCTTTACTTCTTATGGATGTCTTGGGTGCGCCA
 ATGTAAGCTTTACCTATTATGCCGGATAGCGTGAAAG
 CTGCAAATGAACAGCCTGCCTGCCGGAAGATACGGCC
 GGGCCAAGGCACCCCTGGTGACGGTTAGCTCA

60833

Vλ1 Amino Acid Sequence:

DIVLTQPPSVSGAPGQRVTISCSGSSNSVSWYQQLPGTAPKLLIYDNSKRPSGVPDFRFSKSGTSASLAI
 TGLQSEDEADYYCQSRDITYGYWVFGGGTKLTVLG (Q)

Vλ1 DNA Sequence:

GATATCGTGTGACCCAGCCGCTTCAGTGAGTGGCGCACCCAGGTCAGCGTGTGACCATCTCGTGTAGCGGCAGCAG
 CAACATTGGTTCTAATTCTGTGTTGGTACCAGCAGTTGCCCGGGACGGCCGAAACTTCTGATTTATGATAATTCTA
 AGCGTCCCTCAGGCGTCCGGATCGTATTAGCGGATCCAAAGCGGCACCCAGCGGAGCCTTGCATTTACGGCCCTGCAA
 AGCGAAGACGAAGCGGATTAATTATTGCCAGTCTCGTGATACTTATGGTTATTGCGGTGTTGGCGGGCCACGAAAGTT
 AACCGTCCCTAGGT

Fig. 12C

1

ANTI-IGF ANTIBODIES

This application claims the benefit of European application number EP 08171554.2 filed Dec. 12, 2008, the contents of which are incorporated herein their entirety.

The present invention relates to the therapy of hyperproliferative diseases, in particular to the therapy of cancers.

BACKGROUND OF THE INVENTION

Insulin-like growth factor-1 (IGF-1; a 70 amino-acid polypeptide) and insulin-like growth factor-2 (IGF-2; a 67 amino-acid polypeptide) are 7.5-kD soluble factors present in serum that can potentially stimulate the growth of many mammalian cells (reviewed by Pollack et al., 2004). On secretion into the bloodstream the IGFs form complexes with the IGF-BPs which protect them from proteolytic degradation in the serum en route to their target tissues and prevents their association with the IGF receptors. IGFs are also known to be secreted in an autocrine or paracrine manner in target tissues themselves. This is known to occur during normal fetal development where the IGFs play a key role in the growth of tissues, bone and organs. It is also seen in many cancer tissues where there is thought to be paracrine signaling between tumour cells and stromal cells or autocrine IGF production by the tumour cells themselves (reviewed by LeRoith D, 2003).

IGF-1 and IGF-2 are able to bind to the IGF-1 receptor (IGF-1R) expressed on many normal tissues, which functionally is a 460 kD heterotetramer consisting of a dimerised alpha- and beta-subunit, with similar affinities (Rubin et al., 1995). IGF-2 can also bind to the IGF-2 receptor, which is thought to prevent IGF-2 from binding and signaling through the IGF-1R. In this respect the IGF-2R has been demonstrated to be a tumour suppressor protein. The IGF-1R is structurally similar to the insulin receptor which exists in two forms, IR-A and IR-B, which differ by an alternatively spliced 12 amino acid exon deletion in the extracellular domain of IR-A. IR-B is the predominant IR isoform expressed in most normal adult tissues where it acts to mediate the effects of insulin on metabolism. IR-A on the other hand is known to be highly expressed in developing fetal tissues but not in adult normal tissues. Recent studies have also shown that IR-A, but not IR-B, is highly expressed in some cancers. The exon deletion in IR-A has no impact on insulin binding but does cause a small conformational change that allows IGF-2 to bind with much higher affinity than for IR-B (Frasca et al., 1999; Pandini et al., 2002). Thus, because of its expression in cancer tissues and increased propensity for IGF-2 binding, IR-A may be as important as IGF-1R in mediating the mitogenic effects of IGF-2 in cancer.

Binding of the IGFs to IGF-1R triggers a complex intracellular signaling cascade which results in activation of proteins that stimulate proliferation and survival (reviewed by Pollack et al., 2004).

Unlike the EGFR and Her2neu receptors there is no known amplification of the IGF-1R or IR-A receptors in cancers indicating that receptor activation is controlled by the presence of active ligand. There is a very large body of scientific, epidemiological and clinical literature implicating a role for the IGFs in the development, progression and metastasis of many different cancer types (reviewed by Jerome et al., 2003; and Pollack et al., 2004).

For example, in colorectal cancer the expression of IGF-2 mRNA and protein is elevated in clinical colorectal tumour specimens compared with adjacent normal tissue (Freier et al., 1999; Li et al., 2004). There is also a positive correlation of elevated IGF serum levels with proliferating cell index in

2

patients with colorectal neoplasia (Zhao et al., 2005). In addition, elevated circulating levels of IGF-2 correlate with an increased risk of developing colorectal cancers and adenomas (Renahan et al., 2000a) and b); Hassan et al., 2000). Loss of parental imprinting (LOI) of the IGF-2 gene, an epigenetic alteration that results in elevated IGF-2 expression, is a heritable molecular trait that has recently been identified in patients with colorectal and other tumour types. Loss of IGF-2 imprinting has been shown to be associated with a five-fold risk of colorectal neoplasia (Cui et al., 2003; Cruz-Correa et al., 2004) and adenomas (Woodson et al., 2004). Antibodies targeting the alpha-subunit of the IGF-1R which block IGF binding and internalize the receptor have been shown to delay the growth of the xenografted colon cancer-derived cell lines such as COLO 205 (Burtrum et al., 2003).

Elevated levels of IGFs are associated with a poor prognosis in human pulmonary adenocarcinomas (Takanami et al., 1996) and IGFs are expressed and secreted by many SCLC- and NSCLC-derived cell lines (Quinn et al., 1996). Transgenic over-expression of IGF-2 induces spontaneous lung tumours in a murine model (Moorhead et al., 2003). In terms of hepatocellular carcinoma (HCC), human clinical specimens and animal models of HCC express higher levels of IGF mRNA and protein than corresponding normal tissues and this has been correlated with increased tumour growth (Wang et al., 2003; Ng et al., 1998). IGF-2 has also been shown to be a serological marker of HCC with elevated levels in the serum of HCC patients compared with controls (Tsai et al., 2005).

Many childhood solid tumours such as Ewing's sarcoma and rhabdomyosarcoma appear to be particularly dependent on the IGF signaling pathway for their growth (Scotlandi et al., 1996). LOI of the IGF-2 gene has been implicated as a primary genetic event in the development for embryonal rhabdomyosarcoma (Fukuzawa et al., 1999). Autocrine IGF signaling is also thought to strongly influence the growth of Ewing's sarcoma in cases where the type-1 EWS-FLI1 chimeric transcription factor is expressed through a chromosomal translocation resulting in elevated expression of target genes including the IGF ligands and IGF-1R, and reduced expression of IGF-BP-3. Antibodies and small molecule compounds targeting the IGF-1R have been shown to reduce the growth of xenografted pediatric solid tumour derived cell lines (Kolb et al., 2008; Manara et al., 2007).

Using IGF ligand-specific antibodies it has been demonstrated that the growth of human prostate cancer cells in adult human bone implanted into SCID mice can be inhibited (Goya et al., 2004). In addition, it was demonstrated that the same IGF ligand antibodies could block the paracrine supply of IGF and suppress the liver metastasis of human colorectal cancer cells in a murine xenograft system (Miyamoto et al., 2005).

There is also considerable evidence suggesting that the IGF signaling system reduces the sensitivity of cancers to chemotherapeutic agents and radiation. One of the earliest findings in this respect was the demonstration that IGF-1R knock-out mouse embryos are refractory to transformation by viruses, oncogenes and over-expressed growth factor receptors (Sell et al., 1993; Sell et al., 1994) and that over-expression of IGF-1R protects cells from UV irradiation and gamma radiation-induced apoptosis (Kulik et al., 1997). Furthermore, using liver tumour cell lines that secrete large amounts of IGF-2, it was found that neutralization of IGF-2 significantly increased response to chemotherapeutic agents such as cisplatin and etoposide in vitro, especially at lower, cytostatic doses, suggesting that IGF-2 can reduce the susceptibility to chemotherapeutic agents (Lund et al., 2004). Consistent with these findings it has been demonstrated that antibodies tar-

getting the IGF-1R increase the susceptibility of tumour xenografts to growth inhibition by chemotherapeutic drugs and radiation (Goetsch et al., 2005).

A number of antibodies that show cross-reactive binding to human IGF-1 and human IGF-2 have been reported. Antibody sm1.2 was raised against human IGF-1 and shows 40% cross-reactivity to human IGF-2 and was shown to inhibit the proliferation of a mouse fibroblast cell line BALB/c3T3 which was stimulated with 20 ng/ml human IGF-1 (Russell et al., 1984). In a study designed to functionally epitope map IGF-1 by raising monoclonal antibodies to whole IGF-1 protein and portions of the protein a number of antibodies were identified that cross reacted with IGF-2 (Manes et al., 1997). The percent cross-reactivity with IGF-2 ranged from 0 to 800% and several antibodies were identified which were equally IGF-1 and IGF-2 reactive. KM1486 is a rat monoclonal antibody that cross-reacts with human IGF-1 and IGF-2 and it was demonstrated that KM1486 can inhibit growth of human prostate cancer cells in human adult bone implanted into nonobese diabetic/severe combined immunodeficient mice (Goya et al., 2004). In addition, it was demonstrated that KM1486 suppresses the liver metastasis of human colorectal cancers (Miyamoto et al., 2005). KM1486 has also been described in WO 03/093317, JP 2003-310275, WO 2005/018671, WO 2005/028515, and WO 2005/027970.

For the treatment of human disease an antibody with a fully human sequence is highly desirable in order to minimize the risk of generating a human anti-antibody reaction and neutralizing antibodies that will rapidly eliminate the administered antibody from the body and thereby reduce the therapeutic effect. As such, and given the roles of IGF-1 and IGF-2 dependent signaling in the development and progression of cancers, it has become desirable to obtain fully human antibodies. WO 2007/070432 describes fully human antibodies that co-neutralise the mitogenic effects of both ligands.

It was an object of the invention to provide alternative human anti-IGF antibodies with high affinities.

It was a further object of the invention to provide human anti-IGF antibodies with high affinity to IGF-1.

It was a further object of the invention to provide human anti-IGF antibodies with high affinity to IGF-1 and to IGF-2.

It was a further object of the invention to provide human anti-IGF antibodies with adequate relative affinities to IGF-1 and to IGF-2.

It was a further object of the invention to provide human anti-IGF antibodies with a higher affinity to IGF-1 than to IGF-2.

It was a further object of the invention to provide human anti-IGF antibodies with high IGF-1 neutralisation potency.

It was a further object of the invention to provide human anti-IGF antibodies with high IGF-1 and IGF-2 neutralisation potency.

It was a further object of the invention to provide human anti-IGF antibodies with high solubility and stability.

It was a further object of the invention to obtain antibodies that do not affect binding of insulin to its receptor.

The clinical development of therapeutic agents is supported by pharmacodynamic biomarkers of drug activity. Clinical studies with antibodies targeting the IGF-1R have demonstrated that an increase in total serum IGF-1 levels may be a useful pharmacodynamic marker for these agents (Pollack et al., 2007). The reason for the increase in total serum IGF-1 levels is likely due to a feedback mechanism involving pituitary growth hormone (GH) secretion which releases both IGF-1 and IGF-BPs from the liver. Indeed, in humans it has been demonstrated that free or bioactive IGF-1, which repre-

sents only around 1% of total IGF-1 levels, determines the feedback response (Chen et al., 2005).

It was therefore a further object of the invention to provide, for the treatment of diseases in whose development and/or progression the IGFs are causally involved, a therapy that is accompanied by a biomarker that allows the pharmacological monitoring of the effectiveness of the therapy.

In the experiments of the present invention, it could be demonstrated that total serum IGF-1 levels are elevated upon application of the anti-IGF antibodies of the invention. Thus, total IGF-1 levels are a useful pharmacodynamic marker for the effectiveness of the therapy with an anti-IGF antibody. It is therefore highly advantageous that the antibodies of the invention are cross-reactive with IGFs from a suitable animal species, e.g. mouse or rat, such that a pharmacodynamic effect can already be tested pre-clinically.

“Total IGF-1 levels” refers to the combined amount of IGF-1 in plasma or serum comprising the amount of IGF-1 bound to serum binding proteins plus the free (unbound) IGF-1.

Therefore, in a further aspect, the present invention relates to a method for determining the effectiveness of a treatment of a cancer patient with an antibody molecule that binds to IGF-1 and IGF-2. In such method, in a first step, the level of total IGF-1 is measured in a biological sample of the patient, e.g. serum or plasma. Next, the antibody molecule is administered and then, after a period of time sufficient to allow the therapeutic antibody to exert its effect, the level of total IGF-1 is again determined. The amount of increase in the level of total IGF-1 compared to the level of total IGF-1 measured in the first step, indicates to which extent the patient responds to said anti-IGF antibody molecule. This method is preferably used for monitoring therapies in which the antibodies of the invention are administered.

BRIEF DESCRIPTION OF THE FIGURES

FIGS. 1A-1G show an ELISA binding titration of IgG1 antibodies designated 60814, 60819 and 60833 to human IGF-1 (FIG. 1A), mouse IGF-1 (FIG. 1B), rat IGF-1 (FIG. 1C), human IGF-2 (FIG. 1D), mouse IGF-2 (FIG. 1E), rat IGF-2 (FIG. 1F), and human insulin (FIG. 1G).

FIG. 2 shows typical titrations of antibody 60833 neutralising IGF-1 (20 ng/mL) (FIG. 2A) and IGF-2 (100 ng/mL) (FIG. 2B) induced phosphorylation of the IGF-1R using a cell based ELISA.

FIG. 3A shows a typical titration of antibody 60833 neutralising IGF-2 (100 ng/mL) induced IR-A phosphorylation. FIG. 3B shows a typical titration of antibody 60833 neutralising human serum (20%) induced phosphorylation of the IGF-1R. Both assays are performed using cell based ELISAs.

FIGS. 4A-4D show the effect of antibodies 60814 and 60819 on IGF-1 (FIGS. 4A and 4C) and IGF-2 (FIGS. 4B and 4D) stimulated MCF-7 (FIGS. 4A and 4B) and COLO 205 (FIGS. 4C and 4D) cell proliferation.

FIG. 5 shows the effect of antibodies 60819 and 60833 on the proliferation of the Ewing's sarcoma-derived cell line TC-71 in 10% growth medium.

FIG. 6 shows the effect of antibody 60819 on murine total serum IGF-1 levels 24 hours following the administration of single doses of 25, 12.5, 6.25, 3.13 mg/kg. 0 mg/kg represents the total serum IGF-1 levels prior to antibody treatment.

FIG. 7 shows the effect of antibody 60819 on rat total plasma IGF-1 levels 24 hours following the administration of single doses of 30, 100, 200 mg/kg by a 10 minute intravenous infusion. 0 mg/kg represents the total serum IGF-1 levels prior to antibody treatment.

FIG. 8 demonstrates the effect of antibody 60819 and rapamycin, alone or in combination, on the proliferation of the Ewing's sarcoma-derived cell line SK-ES-1 in 10% FCS containing growth medium.

FIG. 9 shows the effect of antibody 60819 and rapamycin, alone or in combination, on the phosphorylation of AKT and levels of PTEN.

FIG. 10 demonstrates the effect of antibody 60819 and erlotinib/Tarceva, alone or in combination, on the proliferation of the NSCLC-derived cell line A-549 in 10% FCS containing growth medium.

FIG. 11 shows the 3D structure of human IGF-1 where the amino acids bound by antibody 60833 are highlighted (dark grey). The linear amino acid sequence of human IGF-1 where the amino acids that interact with antibody 60833 are underlined is shown underneath (SEQ ID NO: 43).

FIG. 12 shows the amino acid and DNA sequences of the variable chains of antibodies 60814 (A, SEQ ID NOs: 7 and 8, and SEQ ID NOs: 9 and 10), 60819 (B, SEQ ID NOs: 17 and 18, and SEQ ID NOs: 19 and 20), and 60833 (C, SEQ ID NOs: 27 and 28, and SEQ ID NOs: 29 and 30); CDRs are in bold letters.

BRIEF DESCRIPTION OF INVENTION

In one aspect, the present invention relates to an isolated human antibody molecule, which

- a) binds to human IGF-1 and IGF-2 such that
 - i) binding of IGF-1 and IGF-2 to the IGF-1 receptor is prevented and
 - ii) IGF-1 receptor-mediated signaling is inhibited,
- b) binds to mouse and rat IGF-1 and IGF-2,
- c) does not bind to human insulin;

wherein said antibody molecule is selected from the group comprising

- i) an antibody molecule that has heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:1 (CDR1), SEQ ID NO:2 (CDR2) and SEQ ID NO:3 (CDR3) and that has light chain CDRs comprising the amino acid sequences of SEQ ID NO:4 (CDR1), SEQ ID NO:5 (CDR2) and SEQ ID NO:6 (CDR3);
- ii) an antibody molecule that has heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:11 (CDR1), SEQ ID NO:12 (CDR2) and SEQ ID NO:13 (CDR3) and that has light chain CDRs comprising the amino acid sequences of SEQ ID NO:14 (CDR1), SEQ ID NO:15 (CDR2) and SEQ ID NO:16 (CDR3);
- iii) an antibody molecule that has heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:21 (CDR1), SEQ ID NO:22 (CDR2) and SEQ ID NO:23 (CDR3) and that has light chain CDRs comprising the amino acid sequences of SEQ ID NO:24 (CDR1), SEQ ID NO:25 (CDR2) and SEQ ID NO:26 (CDR3).

In another aspect, the present invention relates to an anti-IGF antibody molecule, wherein said antibody molecule has heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:1 (CDR1), SEQ ID NO:2 (CDR2) and SEQ ID NO:3 (CDR3) and light chain CDRs comprising the amino acid sequences of SEQ ID NO:4 (CDR1), SEQ ID NO:5 (CDR2) and SEQ ID NO:6 (CDR3).

In another aspect, the present invention relates to an anti-IGF antibody molecule, wherein said antibody molecule has heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:11 (CDR1), SEQ ID NO:12 (CDR2) and SEQ ID NO:13 (CDR3) and light chain CDRs comprising the amino acid sequences of SEQ ID NO:14 (CDR1), SEQ ID NO:15 (CDR2) and SEQ ID NO:16 (CDR3).

In another aspect, the present invention relates to an anti-IGF antibody molecule, wherein said antibody molecule has heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:21 (CDR1), SEQ ID NO:22 (CDR2) and SEQ ID NO:23 (CDR3) and has light chain CDRs comprising the amino acid sequences of SEQ ID NO:24 (CDR1), SEQ ID NO:25 (CDR2) and SEQ ID NO:26 (CDR3).

In another aspect, the present invention relates to anti-IGF antibody molecules having heavy and light chains or CDRs having amino acid sequences as depicted in FIGS. 12A-C.

In another aspect, the present invention relates to an anti-IGF antibody molecule, wherein said antibody molecule binds to a nonlinear epitope within IGF-1 comprising the amino acid sequences LCGAELVDALQFVCGDR (SEQ ID NO:41) and CCFRSCDLRRLEM (SEQ ID NO:42) of human IGF-1 (SEQ ID NO:43). In a preferred embodiment, said antibody molecule makes contact with at least 8 amino acids within the amino acid sequence LCGAELVDALQFVCGDR (SEQ ID NO:41), and at least 10 amino acids within amino acid sequence CCFRSCDLRRLEM (SEQ ID NO:42) of human IGF-1 (SEQ ID NO:43). In a further preferred embodiment, such anti-IGF antibody molecule makes contact with Leu (5), Cys (6), Glu (9), Leu (10), Asp (12), Ala (13), Phe (16), Val (17), Arg (21), Cys (47), Cys (48), Phe (49), Ser (51), Cys (52), Asp (53), Leu (54), Arg (55), Leu (57), and Glu (58) of human IGF-1 (SEQ ID NO:43), as determined by X-ray crystallography. A respective method is disclosed in Example 9 herein. Preferably, said antibody molecule has heavy chain CDRs comprising the amino acid sequences of SEQ ID NO:21 (CDR1), SEQ ID NO:22 (CDR2) and SEQ ID NO:23 (CDR3) and has light chain CDRs comprising the amino acid sequences of SEQ ID NO:24 (CDR1), SEQ ID NO:25 (CDR2) and SEQ ID NO:26 (CDR3).

Binding of the antibody is defined as the interaction that occurs via the non-covalent bonds that hold the antigen (or a protein or a fragment thereof that is structurally similar) to the antibody combining site, i.e. the region of the immunoglobulin that combines with the determinant of an appropriate antigen (or a structurally similar protein).

Affinity (i.e. the interaction between a single antigen-binding site on an antibody and a single epitope) is expressed by the association constant $K_A = k_{ass}/k_{diss}$ or the dissociation constant $K_D = k_{diss}/k_{ass}$.

In one aspect according to a), the antibody binds to each IGF protein with an affinity, as determined by surface plasmon resonance analysis, with a K_D value ranging from 0.02 nM to 20 nM, e.g. 0.2 nM to 2 nM, for example, with an affinity of 0.05, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, or 1.0 nM. Based on this property, neutralization of IGF functional signaling is achieved.

In one aspect according to c), the antibody does not bind to human insulin at concentrations that are at least 100-fold higher than the minimum concentration required for binding to human IGF-1 or IGF-2.

In another aspect, the property of the anti-IGF antibody molecule defined in c) is characterized by the fact that the affinity of the anti-IGF antibody molecule to IGF-1 and IGF-2, respectively, is at least 100-fold, and even more than 1000-fold, as compared to its affinity to insulin. Even though at very high doses, e.g. more than 100 mg/kg, weak binding may not be completely excluded, the anti-IGF antibody molecule does not bind to insulin at therapeutic doses.

In one embodiment, the antibody molecules of the invention do not affect the mitogenic properties of human insulin that are mediated by its binding to the insulin receptor. (In general, a mitogenic property is defined as the ability of a

compound to encourage a cell to commence cell division, triggering mitosis, e.g. in the case of insulin, its ability to promote cell growth).

In another embodiment, in addition to its ability to inhibit IGF signaling mediated via the IGF-1 receptor, an antibody of the invention also has the ability to inhibit IGF-2 signaling mediated via the insulin receptor IR-A.

The antibodies of the invention have a surprisingly high neutralisation potency towards IGF-1 and IGF-2. Furthermore, they have an unexpected higher potency and binding affinity towards IGF-1 than towards IGF-2. They have high solubility and stability, they are free of undesirable glycosylation or hydrolysis motifs in the variable domain, and have a long half-life in the circulation.

DETAILED DESCRIPTION OF THE INVENTION

In the following, an antibody molecule of the invention, which binds to human IGF-1 and IGF-2, is termed "anti-IGF antibody molecule".

The term "anti-IGF antibody molecule" encompasses human anti-IGF antibodies, anti-IGF antibody fragments, anti-IGF antibody-like molecules and conjugates with any of the above mentioned antibody molecules. Antibodies include, in the meaning of the present invention, but are not limited to, monoclonal, chimerized monoclonal, and bi- or multispecific antibodies. The term "antibody" shall encompass complete immunoglobulins as they are produced by lymphocytes and for example present in blood sera, monoclonal antibodies secreted by hybridoma cell lines, polypeptides produced by recombinant expression in host cells, which have the binding specificity of immunoglobulins or monoclonal antibodies, and molecules which have been derived from such immunoglobulins, monoclonal antibodies, or polypeptides by further processing while retaining their binding specificity.

In particular, the term "antibody molecule" includes fully human complete immunoglobulins comprising two heavy chains and two light chains, preferably.

In a further aspect, the antibody molecule is an anti-IGF antibody-fragment that has an antigen binding region. To obtain antibody fragments, e.g. Fab fragments, digestion can be accomplished by means of routine techniques, e.g. using papain or pepsin. Examples of papain digestion are described in WO 94/29348 and U.S. Pat. No. 4,342,566. Papain digestion of antibodies typically produces two identical antigen binding fragments, so-called Fab fragments, each with a single antigen binding site, and a residual Fc fragment. Pepsin treatment yields an F(ab')₂ fragment that has two antigen combining sites and is still capable of cross-linking the antigen. Antibody fragments can also be generated by molecular biology methods producing the respective coding DNA fragments.

Fab fragments also contain the constant domains of the light chain and the first constant domain (CH₁) of the heavy chain. Fab' fragments differ from Fab fragments in that they contain additional residues at the carboxy terminus of the heavy chain CH₁ domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. F(ab')₂ antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them.

Antigen-binding antibody fragments or antibody-like molecules, including single-chain antibodies and linear antibodies as described in Zapata et al., 1995, may comprise, on a single polypeptide, the variable region(s) alone or in combi-

nation with the entirety or a portion of the following: constant domain of the light chain, CH1, hinge region, CH2, and CH3 domains, e.g. a so-called "SMIP" ("Small Modular Immunopharmaceutical"), which is an antibody like molecule employing a single polypeptide chain as its binding domain Fv, which is linked to single-chain hinge and effector domains devoid of the constant domain CH1 (WO 02/056910). SMIPs can be prepared as monomers or dimers, but they do not assume the dimer-of-dimers structure of traditional antibodies. Also included in the invention are antigen-binding fragments comprising any combination of variable region(s) with a constant domain region of a light chain, VH1, CH1, hinge region, CH2, and CH3 domains.

The antibody fragments or antibody-like molecules may contain all or only a portion of the constant region as long as they exhibit specific binding to the relevant portion of the IGF-1/IGF-2 antigen. The choice of the type and length of the constant region depends, if no effector functions like complement fixation or antibody dependent cellular toxicity are desired, mainly on the desired pharmacological properties of the antibody protein. The antibody molecule will typically be a tetramer consisting of two light chain/heavy chain pairs, but may also be dimeric, i.e. consisting of a light chain/heavy chain pair, e.g. a Fab or Fv fragment, or it may be a monomeric single chain antibody (scFv).

The anti-IGF antibody-like molecules may also be single domain antibodies (e.g. the so-called "nanobodies"), which harbour an antigen-binding site in a single Ig-like domain (described e.g. in WO 03/050531, and by Revets et al., 2005). Other examples for antibody-like molecules are immunoglobulin super family antibodies (IgSF; Srinivasan and Roeske, 2005), or CDR-containing or CDR-grafted molecules or "Domain Antibodies" (dAbs). dAbs are functional binding units of antibodies, corresponding to the variable regions of either the heavy (VH) or light (VL) chains of human antibodies. Domain Antibodies have a molecular weight of approximately 13 kDa, or less than one-tenth the size of a full antibody. A series of large and highly functional libraries of fully human VH and VL dAbs has been developed. dAbs are also available for "dual targeting", i.e. dAbs that bind, in addition to IGF-1/IGF-2, to a further target in one molecule. dAb libraries, selection and screening methods, dAb formats for dual targeting and for conferring extended serum half life are described in e.g. U.S. Pat. No. 6,696,245, WO 04/058821, WO 04/003019, and WO 03/002609.

In general, antibody fragments and antibody-like molecules are well expressed in bacterial, yeast, and mammalian cell systems.

In a preferred embodiment, an antibody molecule of the invention, as defined above in i), has a variable heavy chain comprising the amino acid sequence of SEQ ID NO:8 and a variable light chain comprising the amino acid sequence of SEQ ID NO:10 (this sequence may contain, at its C-terminus, an additional Gln. This amino acid position may either be considered the C-terminal end of the variable region, according to the Kabat numbering, or alternatively, and in line with the sequences in the sequence listing, it may represent the first amino acid of the constant light chain, see SEQ ID NO:34).

Preferably, an antibody with the variable heavy chain comprising the amino acid sequence of SEQ ID NO:8 and a variable light chain comprising the amino acid sequence of SEQ ID NO:10 has an IgG1 constant heavy chain region. Preferably, such antibody has an Igλ constant light chain region. Preferably, such antibody is the antibody designated 60814, which has a heavy chain constant region which comprises the amino acid sequence of SEQ ID NO:32 and a light chain constant region which comprises the amino acid

sequence of SEQ ID NO:34. The complete amino acid sequences of the antibody designated 60814 are depicted in SEQ ID NO:35 (heavy chain) and SEQ ID NO:36 (light chain).

In another preferred embodiment, an antibody molecule of the invention, as defined above in ii), has a variable heavy chain comprising the amino acid sequence of SEQ ID NO:18 and a variable light chain comprising the amino acid sequence of SEQ ID NO:20 (this sequence may contain, at its C-terminus, an additional Gln. This amino acid position may either be considered the C-terminal end of the variable region, according to the Kabat numbering, or alternatively, and in line with the sequences in the sequence listing, it may represent the first amino acid of the constant light chain, see SEQ ID NO:34).

Preferably, an antibody with the variable heavy chain comprising the amino acid sequence of SEQ ID NO:18 and a variable light chain comprising the amino acid sequence of SEQ ID NO:20 has an IgG1 constant heavy chain region. Preferably, such antibody has an Ig λ constant light chain region. Preferably, such antibody is the antibody designated 60819, which has a heavy chain constant region which comprises the amino acid sequence of SEQ ID NO:32 and a light chain constant region which comprises the amino acid sequence of SEQ ID NO:34. The complete amino acid sequences of the antibody designated 60819 are depicted in SEQ ID NO:37 (heavy chain) and SEQ ID NO:38 (light chain).

In another preferred embodiment, an antibody of the invention, as defined above in iii), has a variable heavy chain comprising the amino acid sequence of SEQ ID NO:28 and a variable light chain comprising the amino acid sequence of SEQ ID NO:30 (this sequence may contain, at its C-terminus, an additional Gln. This amino acid position may either be considered the C-terminal end of the variable region, according to the Kabat numbering, or alternatively, and in line with the sequences in the sequence listing, it may represent the first amino acid of the constant light chain, see SEQ ID NO:34).

Preferably, an antibody with the variable heavy chain comprising the amino acid sequence of SEQ ID NO:28 and a variable light chain comprising the amino acid sequence of SEQ ID NO:30 has an IgG1 constant heavy chain region. Preferably, such antibody has an Ig λ constant light chain region. Preferably, such antibody is the antibody designated 60833, which has a heavy chain constant region which comprises the amino acid sequence of SEQ ID NO:32 and a light chain constant region which comprises the amino acid sequence of SEQ ID NO:34. The complete amino acid sequences of the antibody designated 60833 are depicted in SEQ ID NO:39 (heavy chain) and SEQ ID NO:40 (light chain).

The cross-reactivity of the antibodies of the invention with mouse and rat IGF-1 allows to examine their endocrine effects, e.g. the effect on the growth hormone pathway, in these species. Cross-reactivity with the rat IGFs is particularly advantageous because the rat is an excellent animal model that is preferably used in drug development to study toxicological effects.

The observed pharmacodynamic effect of the antibodies on total IGF-1 levels, likely due to removal of the free IGF-1, which results in feedback regulation through the growth hormone pathway resulting in increased secretion of IGF-1 by the liver, is a useful pharmacodynamic marker. The availability of such marker in animal species, which allows determination of a dose/effect relationship early in drug development, facilitates the preparation of Phase I clinical studies where, in addition to PK analysis, the pharmacodynamic response on total IGF-1 levels in patients are monitored.

The anti-IGF antibody molecule of the invention may also be a variant of an antibody as defined by the amino acid sequences shown in the sequence listing. Thus, the invention also embodies antibodies that are variants of these polypeptides, which have the features a) to c) defined above. Using routinely available technologies, the person skilled in the art will be able to prepare, test and utilize functional variants of the antibodies 60814, 60819 and 60833. Examples are variant antibodies with at least one position in a CDR and/or framework altered, variant antibodies with single amino acid substitutions in the framework region where there is a deviation from the germline sequence, antibodies with conservative amino substitutions, antibodies that are encoded by DNA molecules that hybridize, under stringent conditions, with the DNA molecules presented in the sequence listing encoding antibody variable chains of 60814, 60819 or 60833, functionally equivalent codon-optimized variants of 60814, 60819 and 60833.

A variant may also be obtained by using an antibody of the invention as starting point for optimization and diversifying one or more amino acid residues, preferably amino acid residues in one or more CDRs, and by screening the resulting collection of antibody variants for variants with improved properties. Particularly preferred is diversification of one or more amino acid residues in CDR3 of the variable light chain, CDR3 of the variable heavy chain, CDR1 of the variable light and/or CDR2 of the variable heavy chain. Diversification can be done by methods known in the art, e.g. the so-called TRIM technology referred to in WO 2007/042309.

Given the properties of individual amino acids, rational substitutions can be performed to obtain antibody variants that conserve the overall molecular structure of antibody 60814, 60819 or 60833. Amino acid substitutions, i.e., "conservative substitutions", may be made, for instance, on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the respective amino acid. The skilled person is familiar with commonly practiced amino acid substitutions, as described e.g. in WO 2007/042309, and methods for obtaining thus modified antibodies. Given the genetic code and recombinant and synthetic DNA techniques, DNA molecules encoding variant antibodies with one or more conservative amino acid exchanges can be routinely designed and the respective antibodies readily obtained.

Preferred antibody variants have a sequence identity in the variable regions of at least 60%, more preferably, at least 70% or 80%, still more preferably at least 90% and most preferably at least 95%. Preferred antibodies also have a sequence similarity in the variable regions of at least 80%, more preferably 90% and most preferably 95%.

("Sequence identity" between two polypeptide sequences indicates the percentage of amino acids that are identical between the sequences. "Sequence similarity" indicates the percentage of amino acids that either are identical or that represent conservative amino acid substitutions.)

In a further embodiment, the anti-IGF antibody molecule of the invention is an "affinity matured" antibody.

An "affinity matured" anti-IGF antibody is an anti-IGF antibody derived from a parent anti-IGF antibody, e.g. 60814, 60819 or 60833, that has one or more alterations in one or more CDRs or in which one or more complete CDRs have been replaced, which results in an improvement in the affinity for the antigens, compared to the respective parent antibody. One of the procedures for generating such antibody mutants involves phage display (Hawkins et al., 1992; and Lowman et al., 1991). Briefly, several hypervariable region sites (e.g. 6-7 sites) are mutated to generate all possible amino substitutions

at each site. The antibody mutants thus generated are displayed in a monovalent fashion from filamentous phage particles as fusions to the gene III product of M13 packaged within each particle. The phage-displayed mutants are then screened for their biological activity (e.g. binding affinity) as herein disclosed.

Affinity matured antibodies may also be produced by methods as described, for example, by Marks et al., 1992, (affinity maturation by variable heavy chain (VH) and variable light chain (VL) domain shuffling), or Barbas et al., 1994; Shier et al., 1995; Yelton et al., 1995; Jackson et al., 1995; and Hawkins et al., 1992, (random mutagenesis of CDR and/or framework residues). Preferred affinity matured antibodies will have very high affinities, e.g. low picomolar, for the target antigen.

The present invention also relates to DNA molecules that encode an anti-IGF antibody molecule of the invention. These sequences include, but are not limited to, those DNA molecules encoding antibodies 60814, 60819 and 60833 as shown in the sequence listing: SEQ ID NO:7 and SEQ ID NO:9, respectively, encoding the variable heavy and light chain, respectively, of antibody 60814; SEQ ID NO:17 and SEQ ID NO:19, encoding the variable heavy and light chain, respectively, of antibody 60819; SEQ ID NO:27 and SEQ ID NO:29, encoding the variable heavy and light chain, respectively, of antibody 60833.

The sequences shown in SEQ ID NO:9, SEQ ID NO:19 and 29, encoding the variable light chains, may, at their 3' end, contain an additional codon for Gln.

Accordingly, the present invention also relates to nucleic acid molecules that hybridize to the DNA molecules set forth in the sequence listing under high stringency binding and washing conditions, as defined in WO 2007/042309, where such nucleic molecules encode an antibody or functional fragment thereof that has properties equivalent or superior to antibody 60814, 60819 or 60833. Preferred molecules (from an mRNA perspective) are those that have at least 75% or 80% (preferably at least 85%, more preferably at least 90% and most preferably at least 95%) homology or sequence identity with one of the DNA molecules described herein.

Yet another class of DNA variants that are within the scope of the invention may be defined with reference to the polypeptide they encode. These DNA molecules deviate with respect to their sequence from those depicted in the sequence listing (SEQ ID NOs:7, 17 and 27, or 9, 19, 29, respectively), but encode, due to the degeneracy of the genetic code, antibodies with the identical amino acid sequences of antibodies 60814, 60819 or 60833, respectively. By way of example, in view of expressing antibodies 60814, 60819 or 60833 in eukaryotic cells, the last nine nucleotides, respectively, that encode the last three amino acids of the variable light chains, can be designed to match codon usage in eukaryotic cells. If it is desired to express the antibodies in *E. coli*, these sequences can be changed to match *E. coli* codon usage.

Variants of DNA molecules of the invention can be constructed in several different ways, as described in WO 2007/042309.

For producing the recombinant anti-IGF antibody molecules of the invention, the DNA molecules (cDNA and/or genomic DNA) encoding full-length light chain (in the case of antibody 60814, a sequence comprising SEQ ID NO:9 and SEQ ID NO:33) and heavy chain (in the case of antibody 60814, the sequence comprising SEQ ID NO:7 and SEQ ID NO:31), or fragments thereof, are inserted into expression vectors such that the sequences are operatively linked to transcriptional and/or translational control sequences. In the case of antibody 60819, the sequences are those of SEQ ID

NO:19 and SEQ ID NO:33, and SEQ ID NO:17 and SEQ ID NO:31, respectively, in the case of antibody 60833, the sequences are those of SEQ ID NO:29 and SEQ ID NO:33, and SEQ ID NO:27 and SEQ ID NO:31, respectively.

For manufacturing the antibodies of the invention, the skilled artisan may choose from a great variety of expression systems well known in the art, e.g. those reviewed by Kipriyanow and Le Gall, 2004.

In another aspect, the present invention relates to an expression vector containing a DNA molecule comprising the nucleotide sequence encoding the variable heavy chain and/or the variable light chain of an antibody molecule as described above. Preferably, such an expression vector of containing a DNA molecule comprising the nucleotide sequence of SEQ ID NO:7 and/or SEQ ID NO:9, or comprising the sequence of SEQ ID NO:17 and/or SEQ ID NO:19, or comprising the sequence of SEQ ID NO:27 and/or SEQ ID NO:29. Preferably, such an expression vector additionally comprises a DNA molecule encoding the constant heavy chain and/or the constant light chain, respectively, linked to the DNA molecule encoding the variable heavy chain and/or the variable light chain, respectively.

Expression vectors include plasmids, retroviruses, cosmids, EBV derived episomes, and the like. The expression vector and expression control sequences are selected to be compatible with the host cell. The antibody light chain gene and the antibody heavy chain gene can be inserted into separate vectors. In certain embodiments, both DNA sequences are inserted into the same expression vector. Convenient vectors are those that encode a functionally complete human CH (constant heavy) or CL (constant light) immunoglobulin sequence, with appropriate restriction sites engineered so that any VH (variable heavy) or VL (variable light) sequence can be easily inserted and expressed, as described above. In the case of the antibodies with the variable regions of 60814, 60819 and 60833, the constant chain is usually kappa or lambda for the antibody light chain, for the antibody heavy chain, it can be, without limitation, any IgG isotype (IgG1, IgG2, IgG3, IgG4) or other immunoglobulins, including allelic variants.

The recombinant expression vector may also encode a signal peptide that facilitates secretion of the antibody chain from a host cell. The DNA encoding the antibody chain may be cloned into the vector such that the signal peptide is linked in-frame to the amino terminus of the mature antibody chain DNA. The signal peptide may be an immunoglobulin signal peptide or a heterologous peptide from a non-immunoglobulin protein. Alternatively, the DNA sequence encoding the antibody chain may already contain a signal peptide sequence.

In addition to the antibody chain DNA sequences, the recombinant expression vectors carry regulatory sequences including promoters, enhancers, termination and polyadenylation signals and other expression control elements that control the expression of the antibody chains in a host cell. Examples for promoter sequences (exemplified for expression in mammalian cells) are promoters and/or enhancers derived from CMV (such as the CMV Simian Virus 40 (SV40) promoter/enhancer), adenovirus, (e. g., the adenovirus major late promoter (AdMLP)), polyoma and strong mammalian promoters such as native immunoglobulin and actin promoters. Examples for polyadenylation signals are BGH polyA, SV40 late or early polyA; alternatively, 3'UTRs of immunoglobulin genes etc. can be used.

The recombinant expression vectors may also carry sequences that regulate replication of the vector in host cells (e. g. origins of replication) and selectable marker genes.

Nucleic acid molecules encoding the heavy chain or an antigen-binding portion thereof and/or the light chain or an antigen-binding portion thereof of an anti-IGF antibody, and vectors comprising these DNA molecules can be introduced into host cells, e.g. bacterial cells or higher eukaryotic cells, e.g. mammalian cells, according to transfection methods well known in the art, including liposome-mediated transfection, polycation-mediated transfection, protoplast fusion, micro-injections, calcium phosphate precipitation, electroporation or transfer by viral vectors.

Preferably, the DNA molecules encoding the heavy chain and the light chain are present on two vectors which are co-transfected into the host cell, preferably a mammalian cell.

In a further aspect, the present invention relates to a host cell carrying one or more expression vectors as described before, preferably a mammalian cell.

Mammalian cell lines available as hosts for expression are well known in the art and include, inter alia, Chinese hamster ovary (CHO) cells, NSO, SP2/0 cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human carcinoma cells (e. g., Hep G2 and A-549 cells), 3T3 cells or the derivatives/progenies of any such cell line. Other mammalian cells, including but not limited to human, mice, rat, monkey and rodent cells lines, or other eukaryotic cells, including but not limited to yeast, insect and plant cells, or prokaryotic cells such as bacteria may be used. The anti-IGF antibody molecules of the invention are produced by culturing the host cells for a period of time sufficient to allow for expression of the antibody molecule in the host cells.

Thus, in a further aspect, the present invention relates to a method for producing an antibody molecule as described, comprising transfecting a mammalian host cell with one or more vectors as described, cultivating the host cell and recovering and purifying the antibody. In another embodiment, the present invention relates to a method for producing an antibody as described above, comprising obtaining a mammalian host cell comprising one or more vectors as described, and cultivating the host cell. In another embodiment, the method further comprises recovering and purifying the antibody.

Antibody molecules are preferably recovered from the culture medium as a secreted polypeptide or it can be recovered from host cell lysates if for example expressed without a secretory signal. It is necessary to purify the antibody molecules using standard protein purification methods used for recombinant proteins and host cell proteins in a way that substantially homogenous preparations of the antibody are obtained. By way of example, state-of-the art purification methods useful for obtaining the anti-IGF antibody molecule of the invention include, as a first step, removal of cells and/or particulate cell debris from the culture medium or lysate. The antibody is then purified from contaminant soluble proteins, polypeptides and nucleic acids, for example, by fractionation on immunoaffinity or ion-exchange columns, ethanol precipitation, reverse phase HPLC, Sephadex chromatography, chromatography on silica or on a cation exchange resin. As a final step in the process for obtaining an anti-IGF antibody molecule preparation, the purified antibody molecule may be dried, e.g. lyophilized, as described below for therapeutic applications.

In one embodiment, the anti-IGF antibody molecule of the invention may be purified by a sequence of state-of-the art purifications steps comprising affinity chromatography (recombinant Protein A), low pH viral inactivation, depth filtration, cation exchange chromatography, anion exchange chromatography, nanofiltration, and 30 kD ultra/diafiltration (Shukla et al., 2007).

In a further aspect, the present invention relates to an antibody molecule as described above for use in medicine.

In a further aspect, the present invention relates to a pharmaceutical composition containing, as the active ingredient, an anti-IGF antibody molecule, preferably a full antibody, of the invention.

To be used in therapy, the anti-IGF antibody molecule is included into pharmaceutical compositions appropriate to facilitate administration to animals or humans. Typical formulations of the anti-IGF antibody molecule can be prepared by mixing the anti-IGF antibody molecule with physiologically acceptable carriers, excipients or stabilizers, in the form of lyophilized or otherwise dried formulations or aqueous solutions or aqueous or non-aqueous suspensions. Carriers, excipients, modifiers or stabilizers are nontoxic at the dosages and concentrations employed. They include buffer systems such as phosphate, citrate, acetate and other anorganic or organic acids and their salts; antioxidants including ascorbic acid and methionine; preservatives such as octadecyltrimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone or polyethylene glycol (PEG); amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, oligosaccharides or polysaccharides and other carbohydrates including glucose, mannose, sucrose, trehalose, dextrans or dextrans; chelating agents such as EDTA; sugar alcohols such as, mannitol or sorbitol; salt-forming counter-ions such as sodium; metal complexes (e.g., Zn-protein complexes); and/or ionic or non-ionic surfactants such as TWEEN™ (polysorbates), PLURONICS™ or fatty acid esters, fatty acid ethers or sugar esters. Also organic solvents can be contained in the antibody formulation such as ethanol or isopropanol. The excipients may also have a release-modifying or absorption-modifying function.

The anti-IGF antibody molecules may also be dried (freeze-dried, spray-dried, spray-freeze dried, dried by near or supercritical gases, vacuum dried, air-dried), precipitated or crystallized or entrapped in microcapsules that are prepared, for example, by coacervation techniques or by interfacial polymerization using, for example, hydroxymethylcellulose or gelatin and poly-(methylmethacrylate), respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules), in macroemulsions or precipitated or immobilized onto carriers or surfaces, for example by pcmc technology (protein coated microcrystals). Such techniques are disclosed in Remington, 2005.

Naturally, the formulations to be used for in vivo administration must be sterile; sterilization may be accomplished by conventional techniques, e.g. by filtration through sterile filtration membranes.

It may be useful to increase the concentration of the anti-IGF antibody to come to a so-called high concentration liquid formulation (HCLF); various ways to generate such HCLFs have been described.

The anti-IGF antibody molecule may also be contained in a sustained-release preparation. Such preparations include solid, semi-solid or liquid matrices of hydrophobic or hydrophilic polymers, and may be in the form of shaped articles, e.g., films, sticks or microcapsules and may be applied via an application device. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxy-

ethyl-methacrylate or sucrose acetate butyrate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated antibodies remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37° C., resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S—S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilization (e.g. as described in WO 89/011297) from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

Formulations that may also be used for the anti-IGF antibody molecule of the invention are described in U.S. Pat. Nos. 7,060,268 and 6,991,790.

The IGF antibody molecule can be incorporated also in other application forms, such as dispersions, suspensions or liposomes, tablets, capsules, powders, sprays, transdermal or intradermal patches or creams with or without permeation enhancing devices, wafers, nasal, buccal or pulmonary formulations, or may be produced by implanted cells or—after gene therapy—by the individual's own cells.

An anti-IGF antibody molecule may also be derivatized with a chemical group such as polyethylene glycol (PEG), a methyl or ethyl group, or a carbohydrate group. These groups may be useful to improve the biological characteristics of the antibody, e.g., to increase serum half-life or to increase tissue binding.

The preferred mode of application is parenteral, by infusion or injection (intravenous, intramuscular, subcutaneous, intraperitoneal, intradermal), but other modes of application such as by inhalation, transdermal, intranasal, buccal, oral, may also be applicable.

In a preferred embodiment, the pharmaceutical composition of the invention contains the anti-IGF-antibody, e.g. antibody 60814, 60819 or 60833, in a concentration of 10 mg/ml and further comprises 25 mM Na citrate pH 6, 115 mM NaCl, 0.02% Tween® (polysorbate 20).

In another embodiment, the pharmaceutical composition of the invention is an aqueous solution which contains the anti-IGF-antibody, e.g. antibody 60814, 60819 or 60833, in a concentration of 10 mg/ml, and further comprises 25 mM histidine HCl pH 6, 38.8 g/L mannitol, 9.70 g/L sucrose, and 0.02% Tween® (polysorbate 20).

For intravenous infusion, the pharmaceutical composition of the invention may be diluted with a physiological solution, e.g. with 0.9% sodium chloride or G5 solution.

The pharmaceutical composition may be freeze-dried and reconstituted with water for injection (WFI) before use.

For the prevention or treatment of disease, the appropriate dosage of antibody will depend on the type of disease to be treated, the severity and course of the disease, whether the antibody is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the antibody, and the discretion of the attending

physician. The antibody is suitably administered to the patient at one time or over a series of treatments.

Depending on the type and severity of the disease, about 1 μ g/kg to 20 mg/kg (e.g. 0.1-15 mg/kg) of antibody is an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion, e.g. infusion over 1 hour. A typical treatment schedule usually involves administration of the antibody once every week to once every three weeks with doses ranging from about 0.1 μ g/kg to ca. 20 mg/kg or more, depending on the factors mentioned above. For example, a weekly dose could be 5, 10, or 15 mg/kg. Progress of this therapy is easily monitored by conventional techniques and assays.

The “therapeutically effective amount” of the antibody to be administered is the minimum amount necessary to prevent, ameliorate, or treat a disease or disorder.

The anti-IGF antibody molecule of the invention and pharmaceutical compositions containing it are useful for the treatment of hyperproliferative disorders.

In certain embodiments, the hyperproliferative disorder is cancer.

Cancers are classified in two ways: by the type of tissue in which the cancer originates (histological type) and by primary site, or the location in the body, where the cancer first developed. The most common sites in which cancer develops include the skin, lung, breast, prostate, colon and rectum, cervix and uterus.

The anti-IGF antibody molecules of the invention are useful in the treatment of a variety of cancers, including but not limited to the following:

- AIDS-related cancer such as Kaposi's sarcoma;
- bone related cancer such as Ewing's family of tumours and osteosarcoma;
- brain related cancer such as adult brain tumour, childhood brain stem glioma, childhood cerebellar astrocytoma, childhood cerebral astrocytoma/malignant glioma, childhood ependymoma, childhood medulloblastoma, childhood supratentorial primitive neuroectodermal tumours, childhood visual pathway and hypothalamic glioma and other childhood brain tumours;
- breast cancer;
- digestive/gastrointestinal related cancer such as anal cancer, extrahepatic bile duct cancer, gastrointestinal carcinoid tumour, gastrointestinal stroma tumour (GIST), cholangiocarcinoma, colon cancer, esophageal cancer, gallbladder cancer, adult primary liver cancer (hepatocellular carcinoma, hepatoblastoma) childhood liver cancer, pancreatic cancer, rectal cancer, small intestine cancer and stomach (gastric) cancer;
- endocrine related cancer such as adrenocortical carcinoma, gastrointestinal carcinoid tumour, islet cell carcinoma (endocrine pancreas), parathyroid cancer, pheochromocytoma, pituitary tumour and thyroid cancer;
- eye related cancer such as intraocular melanoma, and retinoblastoma;
- genitourinary related cancer such as bladder cancer, kidney (renal cell) cancer, penile cancer, prostate cancer, transitional cell renal pelvis and ureter cancer, testicular cancer, urethral cancer, Wilms' tumour and other childhood kidney tumours;
- germ cell related cancer such as childhood extracranial germ cell tumour, extragonadal germ cell tumour, ovarian germ cell tumour and testicular cancer;
- gynecologic cancer such as cervical cancer, endometrial cancer, gestational trophoblastic tumour, ovarian epithe-

lial cancer, ovarian germ cell tumour, ovarian low malignant potential tumour, uterine sarcoma, vaginal cancer and vulvar cancer;

head and neck related cancer such as hypopharyngeal cancer, laryngeal cancer, lip and oral cavity cancer, metastatic squamous neck cancer with occult primary, nasopharyngeal cancer, oropharyngeal cancer, paranasal sinus and nasal cavity cancer, parathyroid cancer and salivary gland cancer;

hematologic/blood related cancer such as leukemias, such as adult acute lymphoblastic leukemia, childhood acute lymphoblastic leukemia, adult acute myeloid leukemia, childhood acute myeloid leukemia, chronic lymphocytic leukemia, chronic myelogenous leukemia and hairy cell leukemia; and lymphomas, such as AIDS-related lymphoma, cutaneous T-cell lymphoma, adult Hodgkin's lymphoma, childhood Hodgkin's lymphoma, Hodgkin's lymphoma during pregnancy, mycosis fungoides, adult non-Hodgkin's lymphoma, childhood non-Hodgkin's lymphoma, non-Hodgkin's lymphoma during pregnancy, primary central nervous system lymphoma, Sezary syndrome, cutaneous T-cell lymphoma and Waldenström's macroglobulinemia and other hematologic/blood related cancer such as chronic myeloproliferative disorders, multiple myeloma/plasma cell neoplasm, myelodysplastic syndromes and myelodysplastic/myeloproliferative diseases;

musculoskeletal related cancer such as Ewing's family of tumours, osteosarcoma, malignant fibrous histiocytoma of bone, childhood rhabdomyosarcoma, adult soft tissue sarcoma, childhood soft tissue sarcoma and uterine sarcoma; hemangiosarcomas and angiosarcoma;

neurologic related cancer such as adult brain tumour, childhood brain tumour, brain stem glioma, cerebellar astrocytoma, cerebral astrocytoma/malignant glioma, ependymoma, medulloblastoma, supratentorial primitive neuroectodermal tumours, visual pathway and hypothalamic glioma and other brain tumours such as neuroblastoma, pituitary tumour and primary central nervous system lymphoma;

respiratory/thoracic related cancer such as non-small cell lung cancer, small cell lung cancer, malignant mesothelioma, thymoma and thymic carcinoma;

skin related cancer such as cutaneous T-cell lymphoma, Kaposi's sarcoma, melanoma, Merkel cell carcinoma and skin cancer;

Small blue round cell tumours.

In particular, the anti-IGF antibody molecules of the invention and pharmaceutical compositions containing them are beneficial in the treatment of cancers of the hematopoietic system including leukemias, lymphomas and myelomas, cancers of the gastrointestinal tract including esophageal, gastric, colorectal, pancreatic, liver and gall bladder and bile duct cancer; kidney, prostate and bladder cancer; gynecological cancers including breast, ovarian, cervical and endometrial cancer; skin and head and neck cancers including malignant melanomas; pediatric cancers like Wilms' tumour, neuroblastoma and Ewing' sarcoma; brain cancers like glioblastoma; sarcomas like osteosarcoma, soft tissue sarcoma, rhabdomyosarcoma, hemangiosarcoma; lung cancer, mesothelioma and thyroid cancer.

In a preferred aspect of the invention, the anti-IGF antibody molecules of the invention and pharmaceutical compositions containing them are beneficial in the treatment of non-small cell lung cancer (NSCLC), in particular locally advanced or metastatic NSCLC (stage IIIB/IV). In this context, the anti-IGF antibody molecules of the invention can be combined

with platinum-based chemotherapy, in particular paclitaxel/carboplatin or gemcitabine/cisplatin platinum doublet therapy.

In a further preferred aspect of the invention, the anti-IGF antibody molecules of the invention and pharmaceutical compositions containing them are beneficial in the treatment of hepatocellular carcinoma, in particular locally advanced or hepatocellular carcinoma (stage III/IV). In this context, the anti-IGF antibody molecules of the invention can be combined with sorafenib (Strumberg D., 2005).

In another embodiment, the anti-IGF antibody molecules and pharmaceutical compositions containing them are useful for non-cancerous hyperproliferative disorders such as, without limitation, psoriasis and restenosis after angioplasty. In addition, based on the recent observation (Reinberg, 2008) that a gene mutation that decreases the activity of IGF-1 has a positive effect on longevity, the antibodies of the invention have the potential to be useful, when applied to adults, in therapies to slow aging and prevent age-related diseases.

Thus, in a further aspect, the present invention relates to the use of an antibody molecule as described above for the preparation of a medicament for the treatment of a cancerous disease outlined above.

In another aspect, the present invention relates to a pharmaceutical composition as described above for the treatment of a cancerous disease as outlined before.

In another aspect, the present invention relates to a method for treating a patient suffering from a cancerous disease as outlined above, comprising administering to said patient an effective amount of a pharmaceutical composition as described herein.

Depending on the disorder to be treated, the anti-IGF antibody molecule of the invention may be used on its own or in combination with one or more additional therapeutic agents, in particular selected from DNA damaging agents or therapeutically active compounds that inhibit angiogenesis, signal transduction pathways or mitotic checkpoints in cancer cells.

The additional therapeutic agent may be administered simultaneously with, optionally as a component of the same pharmaceutical preparation, or before or after administration of the anti-IGF antibody molecule.

In certain embodiments, the additional therapeutic agent may be, without limitation, one or more inhibitors selected from the group of inhibitors of EGFR, VEGFR, HER2-neu, AuroraA, AuroraB, PLK and PI3 kinase, FGFR, PDGFR, Raf, KSP or PDK1.

Further examples of additional therapeutic agents are inhibitors of CDK, Akt, src/bcr-abl, cKit, cMet/HGF, c-Myc, Flt3, HSP90, hedgehog antagonists, inhibitors of JAK/STAT, Mek, mTor, NFkappaB, the proteasome, Rho, an inhibitor of wnt signaling or an ubiquitination pathway inhibitor.

Examples for Aurora inhibitors are, without limitation, PHA-739358, AZD-1152, AT-9283, CYC-116, R-763, VX-667, MLN-8045, PF-3814735, SNS-314, VX-689, GSK-1070916, TTP-607, PHA-680626, MLN-8237 and ENMD-2076.

An example for a PLK inhibitor is GSK-461364.

Examples for raf inhibitors are BAY-73-4506 (also a VEGFR inhibitor), PLX-4032, RAF-265 (also a VEGFR inhibitor), sorafenib (also a VEGFR inhibitor), XL-281, and Nevavar (also an inhibitor of the VEGFR).

Examples for KSP inhibitors are ispinesib, ARRY-520, AZD-4877, CK-1122697, GSK-246053A, GSK-923295, MK-0731, SB-743921, LY-2523355, and EMD-534085.

Examples for a src and/or bcr-abl inhibitors are dasatinib, AZD-0530, bosutinib, XL-228 (also an IGF-1R inhibitor), nilotinib (also a PDGFR and cKit inhibitor), imatinib (also a

cKit inhibitor), NS-187, KX2-391, AP-24534 (also an inhibitor of EGFR, FGFR, Tie2, Flt3), KM-80 and LS-104 (also an inhibitor of Flt3, Jak2).

An example for a PDK1 inhibitor is AR-12.

An example for a Rho inhibitor is BA-210.

Examples for PI3 kinase inhibitors are PX-866, PX-867, BEZ-235 (also an mTor inhibitor), XL-147, XL-765 (also an mTor inhibitor), BGT-226, CDC-0941, GSK-1059615.

Examples for inhibitors of cMet or HGF are XL-184 (also an inhibitor of VEGFR, cKit, Flt3), PF-2341066, MK-2461, XL-880 (also an inhibitor of VEGFR), MGCD-265 (also an inhibitor of VEGFR, Ron, Tie2), SU-11274, PHA-665752, AMG-102, AV-299, ARQ-197, MetMAb, CGEN-241, BMS-777607, JNJ-38877605, PF-4217903, SGX-126, CEP-17940, AMG-458, INCB-028060, and E-7050.

An example for a c-Myc inhibitor is CX-3543.

Examples for Flt3 inhibitors are AC-220 (also an inhibitor of cKit and PDGFR), KW-2449, LS-104 (also an inhibitor of bcr-abl and Jak2), MC-2002, SB-1317, lestaurtinib (also an inhibitor of VEGFR, PDGFR, PKC), TG-101348 (also an inhibitor of JAK2), XL-999 (also an inhibitor of cKit, FGFR, PDGFR and VEGFR), sunitinib (also an inhibitor of PDGFR, VEGFR and cKit), and tandutinib (also an inhibitor of PDGFR, and cKit).

Examples for HSP90 inhibitors are, tanespimycin, alvespimycin, IPI-504, STA-9090, MEDI-561, AUY-922, CNF-2024, and SNX-5422.

Examples for JAK/STAT inhibitors are CYT-997 (also interacting with tubulin), TG-101348 (also an inhibitor of Flt3), and XL-019.

Examples for Mek inhibitors are ARRY-142886, AS-703026, PD-325901, AZD-8330, ARRY-704, RDEA-119, and XL-518.

Examples for mTor inhibitors are rapamycin, temsirolimus, deforolimus (which also acts as a VEGF inhibitor), everolimus (a VEGF inhibitor in addition), XL-765 (also a PI3 kinase inhibitor), and BEZ-235 (also a PI3 kinase inhibitor).

Examples for Akt inhibitors are perifosine, GSK-690693, RX-0201, and triciribine.

Examples for cKit inhibitors are masitinib, OSI-930 (also acts as a VEGFR inhibitor), AC-220 (also an inhibitor of Flt3 and PDGFR), tandutinib (also an inhibitor of Flt3 and PDGFR), axitinib (also an inhibitor of VEGFR and PDGFR), sunitinib (also an inhibitor of Flt3, PDGFR, VEGFR), and XL-820 (also acts as a VEGFR- and PDGFR inhibitor), imatinib (also a bcr-abl inhibitor), nilotinib (also an inhibitor of bcr-abl and PDGFR).

Examples for hedgehog antagonists are IPI-609, CUR-61414, GDC-0449, IPI-926, and XL-139.

Examples for CDK inhibitors are seliciclib, AT-7519, P-276, ZK-CDK (also inhibiting VEGFR2 and PDGFR), PD-332991, R-547, SNS-032, PHA-690509, PHA-848125, and SCH-727965.

Examples for proteasome inhibitors/NFkappaB pathway inhibitors are bortezomib, carfilzomib, NPI-0052, CEP-18770, MLN-2238, PR-047, PR-957, AVE-8680, and SPC-839.

An example for an ubiquitination pathway inhibitor is HBX-41108.

Examples for anti-angiogenic agents are inhibitors of the FGFR, PDGFR and VEGF(R), and thalidomides, such agents being selected from, without limitation, BIBF 1120 (Vargatef®), bevacizumab, motesanib, CDP-791, SU-14813, telatinib, KRN-951, ZK-CDK (also an inhibitor of CDK), ABT-869, BMS-690514, RAF-265, IMC-KDR, IMC-18F1, IMiDs, thalidomide, CC-4047, lenalidomide, ENMD-0995,

IMC-D11, Ki-23057, brivanib, cediranib, 1B3, CP-868596, IMC-3G3, R-1530 (also an inhibitor of Flt1), sunitinib (also an inhibitor of cKit and Flt3), axitinib (also an inhibitor of cKit), lestaurtinib (also an inhibitor of Flt3 and PKC), vatalanib, tandutinib (also an inhibitor of Flt3 and cKit), pazopanib, PF-337210, aflibercept, E-7080, CHIR-258, sorafenib tosylate (also an inhibitor of Raf), vandetanib, CP-547632, OSI-930, AEE-788 (also an inhibitor of EGFR and Her2), BAY-57-9352 (also an inhibitor of Raf), BAY-73-4506 (also an inhibitor of Raf), XL-880 (also an inhibitor of cMet), XL-647 (also an inhibitor of EGFR and EphB4), XL-820 (also an inhibitor of cKit), nilotinib (also an inhibitor of cKit and bcr-abl), CYT-116, PTC-299, BMS-584622, CEP-11981, dovitinib, CY-2401401, and ENMD-2976.

The additional therapeutic agent may also be selected from EGFR inhibitors, it may be a small molecule EGFR inhibitor or an anti-EGFR antibody. Examples for anti-EGFR antibodies, without limitation, are cetuximab, panitumumab, nimotuzumab, zalutumumab; examples for small molecule EGFR inhibitors are gefitinib, erlotinib and vandetanib (also an inhibitor of the VEGFR). Another example for an EGFR modulator is the EGF fusion toxin.

Further EGFR and/or Her2 inhibitors useful for combination with an anti-IGF antibody molecule of the invention are BIBW 2992 (Tovok®), lapatinib, trastuzumab, pertuzumab, XL-647, neratinib, BMS-599626 ARRY-334543, AV-412, mAB-806, BMS-690514, JNJ-26483327, AEE-788 (also an inhibitor of VEGFR), AZD-8931, ARRY-380 ARRY-333786, IMC-11F8, Zemab, TAK-285, AZD-4769.

Other agents that may be advantageously combined in a therapy with the anti-IGF antibody molecule of the invention are tositumumab and ibritumomab tiuxetan (two radiolabelled anti-CD20 antibodies); ofatumumab, rituximab, LY-2469298, ocrelizumab, TRU-015, PRO-131921, FBT-A05, veltuzumab, R-7159 (CD20 inhibitors), alemtuzumab (an anti-CD52 antibody), denosumab, (an osteoclast differentiation factor ligand inhibitor), galiximab (a CD80 antagonist), zanolimumab (a CD4 antagonist), SGN40 (a CD40 ligand receptor modulator), XmAb-5485, Chi Lob 7/4, luca-tumumab, CP-870893 (CD40 inhibitors), CAT-8015, epratuzumab, Y90-epratuzumab, inotuzumab ozogamicin (CD22 inhibitors), lumiliximab (a CD23 inhibitor), TRU-016 (a CD37 inhibitor), MDX-1342, SAR-3419, MT-103 (CD19 inhibitors), or mapatumumab, tigatuzumab, lexatumumab, Apomab, AMG-951 and AMG-655 (TRAIL receptor modulators).

Other chemotherapeutic drugs that may be used in combination with the anti-IGF antibody molecules of the present invention are selected from, but not limited to hormones, hormonal analogues and antihormonals (e.g. tamoxifen, toremifene, raloxifene, fulvestrant, megestrol acetate, flutamide, nilutamide, bicalutamide, cyproterone acetate, finasteride, buserelin acetate, fludrocortisone, fluoxymesterone, medroxyprogesterone, octreotide, arzoxifene, pasireotide, vapreotide), aromatase inhibitors (e.g. anastrozole, letrozole, liarozole, exemestane, atamestane, formestane), LHRH agonists and antagonists (e.g. goserelin acetate, leuprolide, abarelix, cetrorelix, deslorelin, histrelin, triptorelin), antimitabolites (e.g. antifolates like methotrexate, pemetrexed, pyrimidine analogues like 5-fluorouracil, capecitabine, decitabine, nelarabine, and gemcitabine, purine and adenosine analogues such as mercaptopurine thioguanine, cladribine and pentostatin, cytarabine, fludarabine); antitumour antibiotics (e.g. anthracyclines like doxorubicin, daunorubicin, epirubicin and idarubicin, mitomycin-C, bleomycin dactinomycin, plicamycin, mitoxantrone, pixantrone, streptozocin); platinum derivatives (e.g. cisplatin, oxaliplatin, carboplatin,

lobaplatin, satraplatin); alkylating agents (e.g. estramustine, meclorethamine, melphalan, chlorambucil, busulphan, dacarbazine, cyclophosphamide, ifosfamide, hydroxyurea, temozolomide, nitrosoureas such as carmustine and lomustine, thiotepa); antimetabolic agents (e.g. vinca alkaloids like vinblastine, vindesine, vinorelbine, vinflunine and vincristine; and taxanes like paclitaxel, docetaxel and their formulations, larotaxel; simotaxel, and epothilones like ixabepilone, patupilone, ZK-EPO); topoisomerase inhibitors (e.g. epipodophyllotoxins like etoposide and etopophos, teniposide, amsacrine, topotecan, irinotecan) and miscellaneous chemotherapeutics such as amifostine, anagrelide, interferone alpha, procarbazine, mitotane, and porfimer, bexarotene, celecoxib.

In one aspect, the anti-IGF antibody molecules of the invention are used in combination with platinum-based chemotherapy, for example in combination with paclitaxel/carboplatin or gemcitabine/cisplatin platinum doublet therapy. In one embodiment, such combination therapy may be repeated several times, for examples 6 cycles (q3 weeks). This treatment may be followed by further repeated treatment (e.g. 6 cycles q3 weeks) with anti-IGF antibody molecule alone. This regimen can be used e.g. in the treatment of NSCLC. In another aspect, the anti-IGF antibody molecules of the invention are used in combination with sorafenib. In one embodiment, the anti-IGF antibody molecule may be administered repeatedly in intervals of 1-3 weeks, e.g. for 12 cycles, in combination with continuous administration of sorafenib. This regimen can be used e.g. in the treatment of hepatocellular carcinoma.

The anti-IGF antibody molecules of the invention, e.g. when used at lower concentrations, may also be combined with agents that target the IGF-1R. Such agents include antibodies that bind to IGF-1R (e.g. CP-751871, AMG-479, IMC-A12, MK-0646, AVE-1642, R-1507, BBB-022, SCH-717454, rhu Mab IGFR and novel chemical entities that target the kinase domain of the IGF1-R (e.g. OSI-906 or BMS-554417, XL-228, BMS-754807).

The anti-IGF antibody molecules of the invention may also be used in combination with other therapies including surgery, radiotherapy, endocrine therapy, biologic response modifiers, hyperthermia and cryotherapy and agents to attenuate any adverse effect, e.g. antiemetics and, in a preferred embodiment, antidiabetics, e.g. metformin.

The anti-IGF antibody molecules of the invention are also useful in diagnosis of cancers where elevated serum levels of IGF-1 and/or IGF-2 correlate with development or progression of disease, e.g. for determining elevated IGF-2 levels due to loss of imprinting (LOI), an epigenetic alteration affecting the insulin-like growth factor II gene (IGF2). In certain embodiments, an antibody for diagnostic applications, e.g. for detection of IGF-1 in human tissue sections by immunohistological staining, is a chimeric antibody that is derived from a human antibody. In such antibody, the constant regions, or parts thereof, have been replaced by the respective sequences from an antibody of another species, e.g. mouse. By using such chimeric antibody as a primary antibody, the secondary antibody, e.g. a goat antibody which specifically reacts with the murine Fc portion, will specifically recognize the murine sequences of the chimeric primary antibody and not bind to the Fc portions of the other human immunoglobulin molecules that are present in the human tissue sample. Thus, undesired background staining is avoided.

The antibodies of the invention, by blocking IGF-1 and IGF-2 mediated signal transduction, may also be useful for the control of body weight and adipose tissue formation. To

this end, the antibodies of the invention are administered alone or in combination with other anti-obesity drugs.

Materials & Methods

5 Selection of High Affinity Fully Human Antibodies that Bind IGF-1

Selection of specific Fab fragment clones from the human combinatorial antibody library (HuCAL Gold) (Knappik et al., 2000) that bind human IGF-1 with low nanomolar affinity is performed essentially as described by Rauchenberger et al., 2003, in three panning cycles. In order to identify Fab fragments with improved affinity to human IGF-1, several of these 'parental' Fab clones are subjected to 'in vitro affinity maturation' essentially as described by Nagy et al., 2002. The

L-CDR3 (light chain CDR3) and H-CDR2 (heavy chain CDR2) sequences of each clone are separately diversified by substituting the parental sequence with approximately 10^8 L-CDR3 and H-CDR2 cassettes from HuCAL (Knappik et al., 2000). Phages are prepared from the resultant 'maturation libraries' and each library is subjected to solution panning on human IGF-1. In order to select the highest affinity human IGF-1 binders, the solution panning is performed under normal and increased stringency washing conditions according to methods known in the art, with antigen reduction, and with and without blocking by human insulin. The panning outputs after three phage panning rounds are subcloned into a Fab expression vector and the affinity of each Fab for human IGF-1 determined by an electrochemiluminescence-based equilibrium titration technology developed by BioVeris (Witney, Oxfordshire, UK) essentially as described by Haenel et al., 2005. The Fab clones with the best IGF-1 affinities are sequenced, then converted into human IgG1 antibodies as described by Krebs et al., 2001, with subnanomolar affinity to human IGF-1 without any change in specificity compared with the parental antibodies.

35 Cloning and Recombinant Expression of IgG1 Antibodies

Variable heavy chain regions (VH) and variable light chain regions (VL) are excised from the Fab expression vectors by restriction enzyme digestion and ligated into compatible restriction enzyme sites of pcDNA3.1 based plasmids containing the human IgG1 heavy chain and human Ig λ light chain constant regions respectively. EndoFree plasmid preparations (Qiagen) are prepared and the heavy and light chain plasmids are co-transfected into HEK293 freestyle cells (Invitrogen) at a concentration of 1 mg/L of each plasmid according to the supplier's protocol. After 72 hours the supernatant is harvested and the IgG concentration determined by ELISA. Antibody is purified on a modified protein A column (GE Healthcare), eluted into a citrate buffer and then dialysed to a concentration of 2.5 mg/ml in PBS. Alternatively, a CHO cell line stably integrated with the antibody expression plasmids is generated and used to produce the antibodies.

Surface Plasmon Resonance Analysis for Determining Affinity Constants

a) Antibody Capture Method

55 The sensor chip is coated with approximately 1000 RU of the reference antibody in flow cell 1 and approximately 1000 RU of a rabbit-anti-human Fc-gamma-specific antibody in flow cell 2 using the coupling reagents from an amine coupling kit. A target of 1000 RU is set in the surface preparation wizard of the Biacore 3000 software at a flow rate of 5 μ l/min. Running buffer used is HBS-EP. The affinity measurements are made using the following parameters: 20 μ l/min flow (HCB running buffer); 25° C. detection temperature; Fc1, Fc2 flow paths; Fc1, Fc2 detection; anti-IGF-huMAb-capturing: 3 min of a 1 μ g/ml solution; 5 min IGF-Ag-association; 5 min IGF-Ag-dissociation; regeneration: 30 sec pulse with 50 mM HCl. The IGF antigens are diluted to 500, 250, 125, 62.5

and 31.3 nM in running buffer (HCB) and the different anti-
gen dilutions are run singly over Fc1 and Fc2 with random
order. Blank runs using running buffer only are run in-be-
tween. A blank run curve is subtracted from each binding
curve before affinity analysis. Data evaluation is performed
using the BIAevaluation software (version 4.1, Biacore,
Freiburg, Germany). The dissociation and association phases
of the kinetics are fitted separately. For the separate fit of the
 k_{diss} values a time-frame of the initial 200-300 seconds in the
dissociation phase is used (range of steady decrease of sig-
nal). For the separate fit of the k_{ass} values, initial time frames
of approx 100 seconds are used (range of steady increase of
signal) and for calculation the individual k_{diss} values are used
with the 1:1 Langmuir association model. The average values
with the standard deviations of the kinetic data together with
the corresponding dissociation (K_D) and association (K_A)
constants are calculated.

b) IGF Coating Method

The determination of binding constants of IGF antibodies
to IGF ligands when the sensor chip is coated with IGF
ligands is essentially performed as described above except
that the sensor chip is coated with 35.1 $\mu\text{g}/\text{mm}^2$ and 38.5
 $\mu\text{g}/\text{mm}^2$ IGF-1 and IGF-2 respectively. The antibodies are
then flowed over the chip at the following concentrations: 50,
25, 12.5, 6.25, 3.12 nM.

Measurement of Binding to Human, Murine and Rat IGFs
and to Human Insulin in Immunosorbent Assays

Fully human IgG1 antibodies that bound with high affinity
to IGF-1 are also tested for binding to human IGF-1 in direct
immunosorbent assays (ELISA). Assays are performed by
coating human IGF-1 (R&D Systems, No. 291-G1) to
96-well Maxisorb plates at a concentration of 0.5 $\mu\text{g}/\text{ml}$ over-
night at 4° C. (100 $\mu\text{l}/\text{well}$). Coating buffer alone is used as a
control for unspecific binding. Wells are then washed once
with washing buffer (1 \times TBS-T) and residual binding sites are
blocked with 200 μl blocking buffer for 1 hour at room tem-
perature on an orbital shaker followed by a further wash
cycle. Serial three-fold dilutions of each test antibody in
blocking buffer are prepared directly on the coated plates.
Typical concentrations used are 50, 16.6, 5.6, 1.8, 0.6, 0.2,
and 0.07 ng/ml. Blocking buffer alone is used as a positive
control. The plates are then incubated for 2 hours at room
temperature with agitation. After three wash cycles 100
 $\mu\text{l}/\text{well}$ of HRPO-conjugated anti-human IgG secondary
reagent (Jackson ImmunoResearch Inc.) diluted in blocking
buffer is added to all wells. After 2 hours incubation at room
temperature with agitation the plates are washed three-times
and 100 $\mu\text{l}/\text{well}$ of TMB substrate solution (equal amounts of
solution A and B) are pipetted into all wells. The plates are
incubated for 10-20 min at RT with agitation and then the
reaction is stopped by addition of 100 $\mu\text{l}/\text{well}$ 1 M phosphoric
acid. The absorbance is measured at a wavelength of 450 nm
(reference 650 nm).

Binding of the fully human IGF-1 binding IgG1 antibodies
to mouse IGF-1 (R&D Systems, No. 791-MG), rat IGF-1
(IBT, No. RU100), human IGF-2 (GroPep, No. FM001),
mouse IGF-2 (R&D Systems, No. 792-MG), rat IGF-2 (IBT,
No. AAU100), and human insulin (Roche) is also tested as
described above for human IGF-1 (except that the concentra-
tion of human insulin used for coating is 3 $\mu\text{g}/\text{ml}$).

In Vitro Cell Proliferation Assays for Determining Neutral-
ization Potency

The MCF-7 breast cancer derived cell line (ATCC, HTB-
22) and COLO 205 colon cancer-derived cell line (ATCC
#CCL-222) are plated in 96-well plates at a cell density of
1000 cells per well in serum-free RPMI medium. 10 ng/ml of
either IGF-1 or IGF-2 is added in the presence or absence of

a humanized isotype control antibody that does not bind
IGF-1 or IGF-2, or antibodies 60814, 60819, and 60833 at
concentrations of 12, 37, 111, 333, 1000 and 3000 ng/ml.
Cells are cultured for 5 days then the relative cell number in
each well determined using the CellTiter-Glo luminescent
cell viability assay (Promega). Luminescence
(LU=Luminescence Units) is recorded using a XFluor
GENios Pro 4.

Ewing's Sarcoma-Derived Cell Line Growth Assay

The Ewing's sarcoma-derived cell lines TC-71 (ATCC
#ACC516) and SK-ES-1 (ATCC #HTB86) are plated in
96-well plates at a density of 1000 cells per well in DMEM
medium containing 1 \times NEAA, 1 \times sodium pyruvate, 1 \times
glutamax and 10% fetal calf serum (FCS) and incubated
overnight at 37° C. and 5% CO₂ in a humidified atmosphere.
The following day a serial dilution of test antibody, human-
ized isotype control antibody (a humanized IgG1 antibody
targeted to CD44-v6) that does not bind IGF-1 or IGF-2,
rapamycin, or a combination of rapamycin and test antibody,
are added to the cells. The typical concentrations used are 30,
10, 3.3, 1.1, 0.37, and 0.12 $\mu\text{g}/\text{ml}$ (or 100, 10, 1, 0.1, 0.01,
0.001 nM rapamycin and test antibody for combination stud-
ies) and each dilution is performed in triplicate wells. The
cells plus antibody are then incubated for 120 hours after
which time the relative cell number in each well is determined
using the CellTiter-Glo luminescent cell viability assay
(Promega). Luminescence (LU=Luminescence Units) is
recorded using a XFluor GENios Pro 4 and for data analysis
the mean value from triplicate wells is taken and fitted by
iterative calculations using a sigmoidal curve analysis pro-
gram (Graph Pad Prism) with variable Hill slope.

Western Blot Analysis of Phosphorylated AKT and PTEN Levels

SK-ES-1 cells are plated in 6-well plates in medium con-
taining 10% fetal bovine serum and after overnight incuba-
tion they are treated with either 100 nM isotype control anti-
body (a humanized IgG1 antibody targeted to CD44-v6) that
does not bind IGF-1 or IGF-2, 100 nM 60819, 100 nM rapa-
mycin, or a combination of 100 nM 60819 and 100 nM
rapamycin. 24 hours later the cells are lysed and the cell lysate
frozen after the protein concentration is determined by Brad-
ford assay. Western blotting is performed by applying 30 μg
of protein lysates to an SDS PAGE gel (BioRad) and the gel
blotted on a Citerian gel blotting sandwich. Western blots are
incubated overnight with a rabbit anti-beta actin (control)
antibody, a rabbit anti-PTEN antibody (Cell Signaling
#9559), or a rabbit anti-phospho-pAKT antibody (Cell Sig-
naling #4060), at 1:5000 (anti-beta actin), 1:1000 (anti-
PTEN), or 1:2000 (anti-phosphoAKT) dilutions in 1% milk
powder. Following washing in TBS an anti-rabbit IgG
HRPO-conjugated secondary antibody (Amersham) is
applied for 1 hour and after further washes in TBS antibody
reactivity is detected by ECL and captured on Hyperfilm
(Amersham).

In Vitro Combination of Anti-IGF Antibody with EGFR Inhibitor in NSCLC-Derived Cell Line

The NSCLC-derived cell line A-549 (ATCC #CCL-185) is
plated in 96-well plates at a density of 1000 cells per well in
RPMI 1640 medium containing 2 mM L-glutamine and 10%
fetal bovine serum and incubated overnight at 37° C. and 5%
CO₂ in a humidified atmosphere. The following day a serial
dilution of test IGF antibody, erlotinib/Tarceva, or a combi-
nation of test IGF antibody and erlotinib are added to the
cells. The typical concentrations of the test IGF antibody used
are 30000, 10000, 3333, 1111, 370, 123, 41, 14 ng/mL, and
the typical concentration of erlotinib used are 20000, 6667,
2222, 741, 247, 82, 27, 9 nM, and each dilution is performed

in triplicate wells. The cells are then incubated for 120 hours after which time the relative cell number in each well is determined using the CellTiter-Glo luminescent cell viability assay (Promega). Luminescence (LU=Luminescence Units) is recorded using a XFluor GENios Pro 4 and for data analysis the mean value from triplicate wells is taken and fitted by iterative calculations using a sigmoidal curve analysis program (Graph Pad Prism) with variable Hill slope.

Determination of the Effect on Total Murine and Total Rat Serum IGF-1 Levels

Single intravenous (bolus) administrations of 25, 12.5, 6.25, and 3.13 mg/kg of test IGF antibody are given to female athymic NMRI nude mice, 6-8 weeks old (n=5). Single 10 minute intravenous administrations of 30, 100, 200 mg/kg of antibody 60819 are given to male and female Wistar Han rats, 6-8 weeks old (n=4 male, 4 female). Prior to antibody treatment and 24 hours post administration a blood sample is taken, serum collected, and total murine or rat IGF-1 levels determined using the OCTEIA rat/mouse total IGF-1 immunocytometric assay. The assay is performed according to the manufacturer's instructions, absorbance is measured at 450 nm and evaluated using the SoftMax Pro software. A standard curve is used to determine the serum concentration of total IGF-1 in ng/ml. Statistical analysis is performed using the GraphPad Prism software.

Cell Based IGF-1R Phosphorylation ELISA

Mouse fibroblast cell lines recombinantly expressing human IGF-1R or human IR-A are maintained in DMEM supplemented with 10% heat inactivated FCS, 1 mM sodium pyrovate, 0.075% sodium bicarbonate, MEM NEAA, and 0.3 µg/ml puromycin at 37° C. and 5% CO₂ in a humidified incubator. Cells are detached with trypsin/EDTA, resuspended in growth medium and diluted to 100,000 cells/mL. 100 µL (10,000 cells) are seeded in wells of a sterile 96-well plate and incubated overnight in a humidified incubator at 37° C. and 5% CO₂. The cells are then starved with 100 µL/well assay medium (DMEM supplemented with 0.5% heat inactivated FCS; 1 mM sodium pyrovate, 0.075% sodium bicarbonate, and MEM NEAA) and incubated overnight as before. A range of test antibody concentrations prepared in assay medium is added to the cells, all samples are prepared in triplicate to determine the standard deviation for each assay condition. An IGF-1R antibody, αIR-3 (Calbiochem, No. GR11L) is also tested in these experiments. IGF-1 (20 ng/mL final concentration), IGF-2 (100 ng/mL final concentration), or human serum (20% final concentration) is then added and the plates incubated for 30 min in the humidified incubator. Cells are fixed by replacing the growth medium with 4% formaldehyde in PBS for 20 min at RT. After two wash cycles with 300 µL/well wash buffer (PBS with 0.1% Triton X-100) for 5 min (with agitation) the cells are quenched with 100 µL/well 1.2 wt % hydrogen peroxide in wash buffer for 30 minutes at RT. Cells are washed again with 300 µL/well washing buffer and blocked with 100 µL/well blocking buffer (5% BSA in wash buffer) for 60 min at RT with agitation. Blocking buffer is removed and 50 µL/well primary phospho-IGF-1 receptor β (tyr1135/1136)/insulin receptor β (tyr1150/1151) antibody (Cell Signaling, No. 3024) diluted 1:1000 in blocking buffer is added. Plates are incubated overnight at 4° C. with agitation then washed three times as above and 50 µL/well anti-rabbit IgG goat immunoglobulins conjugated with horseradish peroxidase (Dako, No. P0448) diluted 1:500 in blocking buffer is added. After a 60 min incubation at RT with agitation the wells are washed twice with washing buffer as above and once with 300 µL PBS. 100 µL/well TMB substrate solution (Bender MedSystems, No. BMS406.1000) is added to the wells and incubated for 10 min with agitation,

following this the reaction is stopped by adding 100 µL/well 1 M phosphoric acid and the absorbance read using a photometer (OD 450 nm, OD 650 nm as reference) Inhibition of IGF-1R or IR-A phosphorylation IC₅₀ values are determined by graphical analysis.

Fab-IGF-1 Co-crystallisation and Structure Determination

Monoclonal antibodies are prepared in a buffer of 100 mM Na-phosphate (pH 7.0) prior to papain digestion. Papain (Sigma Aldrich, P #3125) is activated in digestion buffer (phosphate buffer containing 10 mM cysteine hydrochloride, 4 mM EDTA, pH 7.0) following the manufacturer's instructions. IgG antibody is mixed with the activated papain (ratio enzyme:IgG=1:100) and the reaction is incubated at 37° C. on a rotor shaker overnight. Digestion is stopped by adding iodacetamid to a final concentration of 30 mM. To separate the Fab fragment from Fc fragments, Fc cleavage products and intact Mab, the digestion mix is loaded onto a Protein A MabSelect column equilibrated with phosphate buffer. The column is washed with column volumes of PBS, and the Fab fragment is collected in the flow-through and wash fractions. The Fc fragment and intact Mab are eluted from the column with 100 mM citrate buffer (pH 3.0) and subsequent size exclusion chromatography of the Fab fragment is performed using a HiLoad Superdex 75 column The column is run at 0.5 mL per min with 20 mM triethanolamine, 130 mM NaCl, pH 8.0. The protein concentration of Fab fragments is determined by measuring absorbance at 280 nm Quality of Fab fragments is analysed by Western Blotting and ELISA.

Fab-IGF-1 complex is generated by adding a 2-fold molar excess of the recombinant IGF-1 (Gropep; Receptor Grade) to the purified Fab which is then incubated overnight on a rotor shaker at 4° C. Concentration of the complex to (15 mg/mL) and removal of unbound IGF-1 is performed using an Amicon-Ultra device. Crystallization of the Fab:IGF-1 complex is carried out using various techniques such as hanging drop, sitting drop, and seeding. In one embodiment, the crystal is precipitated by contacting the solution with a reservoir that reduces the solubility of the proteins due to presence of precipitants, i.e., reagents that induce precipitation. Screening of various conditions lead to a suitable buffer system manipulated by addition of a precipitant and additives. The concentration of the precipitants is preferably between 5-50% w/v. The pH of the buffer is preferably about 3 to about 6. The concentration of the protein in the solution is preferably that of super-saturation to allow precipitation. The temperature during crystallization is preferably between 4 and 25° C.

The three dimensional structure of Fab:IGF-1 complex as defined by atomic coordinates is obtained from the X-ray diffraction pattern of the crystal and the electron density map derived there from. The diffraction of the crystals is better than 2 Å resolution. The crystals preferably have the space group P3221 (number 154) and unit cell dimensions of approximately=70 Å, b=70 Å, c=195 Å; and γ=120°. The method for determining the three dimensional structure is molecular replacement which involves use of the structure of a closely related molecule or receptor ligand complex. Model building and refining is done in several iterative steps to final R-factors (R and R_{free}) of 21 and 23% respectively.

Determination of Pharmacokinetic Parameters in Rats

Wistar rats are given five intravenous bolus administrations of 18, 52, and 248 mg/kg antibody every 72 hours. At various time points a blood sample is taken and the human antibody concentration in the plasma is determined by sandwich ELISA. This allowed the mean pharmacokinetic parameters

of the antibody to be calculated on the first day of dosing and half-life is calculated after the last day of dosing (with $t(n) = 1008$ hours).

EXAMPLE 1

Selection of High Affinity Antibodies that Bind IGF-1

In order to identify Fab fragments with improved affinity to human IGF-1, several 'parental' Fab clones that are identified to bind IGF-1 with low nanomolar affinity are subjected to 'in vitro affinity maturation' where the L-CDR3 and H-CDR2 sequences of each clone are separately diversified by substituting the parental sequence with a library of new L-CDR3 and H-CDR2 sequences. The resultant 'maturation libraries' are subjected to solution panning on human IGF-1 and the clones with the best affinity are selected for conversion into IgG1 antibodies and tested further. The three antibodies with the best human IGF-1 affinities are 60814, 60819, and 60833 which had affinities (K_D) of 180, 190, and 130 pM respectively (shown in Table 1) as determined by an electrochemiluminescence-based equilibrium titration method.

TABLE 1

IGF-1 BINDING SUMMARY	
Antibody	Affinity (pM)
60814	180
60819	190
60833	130

The antibodies are also tested for their binding to human, murine, and rat IGF-1 and IGF-2, and human insulin, in immunosorbent assays. This demonstrated that 60814, 60819, and 60833 show comparable cross-reactive binding with mouse and rat IGF-1, and human, murine and rat IGF-2, but no reactivity to human insulin (at the highest concentration tested, 50 ng/ml) (FIGS. 1A-1G).

Affinity constants for binding of the antibodies to human, mouse, and rat IGF-1 and IGF-2 is also determined by surface plasmon resonance (Biacore) analysis. The method involves capturing the antibodies on the sensor and flowing the IGF antigens over the captured antibodies, thus overcoming any avidity effect that could occur if the IGF antigens are coated onto the sensor and the antibodies added. The affinity constants using this method for antibody 60833 are shown in Table 2 where it can be seen that the measured K_D values for human IGF-1 and human IGF-2 are 0.07 nM and 0.9 nM respectively.

TABLE 2

AFFINITY CONSTANTS OF ANTIBODY 60833 FOR HUMAN, MOUSE, AND RAT IGF-1 AND IGF-2 DETERMINED BY SURFACE PLASMON RESONANCE (ANTIBODY CAPTURE METHOD)			
Antigen	K_{on} [$M^{-1}s^{-1}$]	K_{off} [s^{-1}]	K_D [nM]
Human IGF-1	4.74×10^6	3.01×10^{-4}	0.07
Mouse IGF-1	1.00×10^6	3.23×10^{-4}	0.33
Rat IGF-1	3.81×10^6	2.53×10^{-4}	0.07
Human IGF-2	3.97×10^6	3.53×10^{-3}	0.913
Mouse IGF-2	8.68×10^5	1.1×10^{-2}	13.4
Rat IGF-2	2.56×10^6	6.13×10^{-3}	2.41

EXAMPLE 2

Inhibition of IGF Signalling

The first signalling event which occurs following binding of IGFs to the IGF-1R is the phosphorylation of the IGF-1R. A cell-based ELISA assay is used to measure the inhibition of IGF induced IGF-1R phosphorylation by the antibody 60833. The potency and effectiveness (of up to 15 μ g/mL (100 nM)) of 60833 in neutralising recombinant bioactive IGF-1 and IGF-2 induced IGF-1R phosphorylation is determined As shown in Table 3 and example FIG. 2 60833 potently and effectively inhibits IGF-1 (FIG. 2A) and IGF-2 (FIG. 2B) induced signalling. In the same assay the IGF-1R targeted mAb α IR3 is much less potent and effective with respect to IGF-1 induced signalling, and displays a very weak effect on IGF-2 induced signalling.

A similar cell based IR-A phosphorylation ELISA is used to demonstrate that 60833 can also inhibit IGF-2 signalling via IR-A. As shown in Table 4 and example FIG. 3A, 60833 potently and effectively inhibits IGF-2 induced IR-A phosphorylation. In contrast, α IR3, which cannot bind IR-A, shows no inhibitory effect.

The level of IGF bioactivity in human serum or plasma samples can also be measured using the IGF-1R phosphorylation cell based ELISA. This is used to determine the potency and effectiveness (up to 15 μ g/mL (100 nM)) of 60833 in neutralising human serum IGF bioactivity. As shown in Table 3 and example FIG. 3B 60833 potently and effectively inhibits IGF bioactivity in human serum.

TABLE 3

EFFECT OF 60833 ON IGF-1R PHOSPHORYLATION			
IGF-1R Phosphorylation Stimulus	Inhibitor	IC ₅₀ (μ g/mL)	% Remaining Phosphorylation at 15 μ g/mL (100 nM) Inhibitor
IGF-1 (20 ng/mL)	60833	0.09	0
	α IR3	1.16	35
	Control IgG	>15	108
IGF-2 (100 ng/mL)	60833	1.12	7
	α IR3	>15	76
	Control IgG	>15	108
Human Serum Pooled from Healthy Donors (20%)	60833	0.25	5
	α IR3	>15	120
	Control IgG	>15	110

TABLE 4

EFFECT OF 60833 ON IR-A PHOSPHORYLATION			
IR-A Phosphorylation Stimulus	Inhibitor	IC ₅₀ (μ g/mL)	% Remaining Phosphorylation at 15 μ g/mL (100 nM) Inhibitor
IGF-2 (100 ng/mL)	60833	0.82	6
	α IR3	>15	115
	Control IgG	>15	109

EXAMPLE 3

Effects on IGF-1 and IGF-2-induced Cell Proliferation

The effects of antibodies 60814, 60819, and 60833 on IGF-1 and IGF-2 induced MCF-7 (breast cancer derived) and COLO 205 (colon cancer derived) cell line proliferation is determined Examples of the effects of antibodies 60814 and 60819 are shown in FIGS. 4A-D. All three antibodies show a

29

dose dependent inhibition of IGF-1 (FIGS. 4A and 4C) and IGF-2 (FIGS. 4B and 4D) induced MCF-7 (FIGS. 4A and 4B) and COLO 205 (FIGS. 4C and 4D) cell proliferation. The concentration of each antibody required to inhibit 50% of the IGF-1 or IGF-2 induced proliferation of each cell line is shown in Table 5.

TABLE 5

INHIBITION OF IGF-1 AND IGF-2 INDUCED PROLIFERATION OF THE MCF-7 AND COLO 205 CANCER CELL LINES				
Cell Line	Stimulation	IC ₅₀ (ng/ml)		
		60814	60819	60833
MCF7	IGF-1	24.1	54.0	38.6
MCF7	IGF-2	78.2	40.8	81.2
COLO-205	IGF-1	135.0	216.9	165.1
COLO 205	IGF-2	576.1	100.8	632.3

EXAMPLE 4

Effects on Proliferation of Ewing's Sarcoma-derived Cell Lines

The effect of antibodies 60819 and 60833 on the proliferation of the Ewing's sarcoma-derived cell line TC-71 grown in medium containing 10% FCS is shown in FIG. 5. Relative to a humanized IgG1 isotype control antibody, that does not bind IGF-1 or IGF-2, both 60819 and 60833 show a dose-dependent inhibition of TC-71 cell proliferation.

EXAMPLE 5

Effect on Total Murine and Rat IGF-1 Levels

Neutralization of active IGF-1 with an IGF targeted antibody may be expected to result in an endocrine feedback through the GH pathway which results in elevated total serum IGF-1 levels. Antibodies 60814, 60819, and 60833 are cross-reactive with mouse and rat IGF-1 which allows any pharmacodynamic effect on total serum IGF-1 levels to be measured in these species. As shown in FIGS. 6 and 7, administration of antibody 60819 to mice (FIG. 6) and rats (FIG. 7) results in a dose dependent elevation of serum total murine and rat IGF-1 levels 24 hours post administration. This represents a useful pharmacodynamic marker of the activity of these antibodies which can be tested during clinical development in humans.

EXAMPLE 6

Effect of Combination of IGF Ligand Targeting Antibodies and Rapamycin on Ewing's Sarcoma-derived Cell Line Proliferation and Intracellular Signaling

The effect of antibody 60819 and the mTOR inhibitor rapamycin, alone or in combination, on the proliferation of the Ewing's sarcoma-derived cell line SK-ES-1 is shown in FIG. 8. There is a dose dependent inhibition of proliferation with both antibody 60819 and rapamycin alone, with both single agents achieving around 60% proliferation inhibition at 100 nM. Combination of equivalent doses of both antibody 60819 and rapamycin demonstrated an additive effect on the inhibition of cell proliferation with approximately 95% inhibition when 100 nM doses are combined.

IGF-induced cell proliferation is mediated via a chain of intracellular protein phosphorylation events. One protein whose phosphorylation is increased by IGF stimulation is AKT. FIG. 9 demonstrates the effect of antibody 60819 and rapamycin, alone or in combination, on the phosphorylation

30

of AKT in SK-ES-1 cells 24 hours following treatment using 100 nM doses. Compared with proliferating untreated cells which show phosphorylation of AKT, 100 nM antibody 60819 inhibited AKT phosphorylation. Conversely, 100 nM rapamycin treatment resulted in higher levels of phosphorylated AKT than the control which is thought to be due to a compensatory feedback mechanism following mTOR inhibition. However, when 100 nM rapamycin and 100 nM antibody 60819 are combined the phosphorylation of AKT is inhibited. This suggests that the compensatory feedback which leads to phosphorylated AKT upon rapamycin treatment is due to elevation of the IGF ligands and these are inhibited by antibody 60819. FIG. 9 also demonstrates that both antibody 60819 and rapamycin, alone or in combination, do not affect the total levels of PTEN.

EXAMPLE 7

Effect of Combination of an IGF Ligand Targeting Antibody and an EGFR Inhibitor on NSCLC-derived Cell Line Proliferation

The effect of antibody 60819 and the EGFR inhibitor erlotinib/tarceva, alone or in combination, on the proliferation of the NSCLC-derived cell line A-549 is shown in FIG. 10. In this model, there is only a small effect of antibody 60819 alone on cell proliferation whilst tarceva shows a dose dependent effect with around 60% cell proliferation inhibition at the highest dose tested (20 μ M). However, when antibody 60819 and tarceva are combined there is a more potent and effective inhibition of cell proliferation indicative of a synergistic effect.

EXAMPLE 8

Pharmacokinetic Properties in Wistar Rats

The mean pharmacokinetic parameters of antibody 60833 in Wistar rats on the first day of dosing with 18, 52, and 248 mg/kg is shown in Table 6. Terminal half-life was calculated after the last day of dosing (with $t(n)=1008$ hours), the average terminal half-life for all three dose levels is 221 hr (9.2 days).

TABLE 6

	60833 Dose (mg/kg)		
	18	52	248
C(max) [mg/mL]	0.531	1.70	5.56
AUC(0-72 h) [mg · h/mL]	15.5	40.2	120
CL [(mL/day)/kg]	22.9	28.7	37.3
V(ss) [mL/kg]	68.1	76.3	65.4
$t_{1/2\phi}$ [hr]	210	197	255

ϕ = after last day of dosing with $t(n) = 1008$ hr

Example 9

Fab-IGF-1 Co-Crystallisation and Structure Determination to Identify Antibody Binding Sites on IGF-1

To definitively determine the residues on human IGF-1 that interact with the IGF antibodies the Fab and IGF-1 were co-crystallised and the structure of the interaction determined with better than 2 Å resolution. The residues on IGF-1 that are contacted by antibody (Fab) 60833 are shown in Table 7. In total 19 residues on IGF-1 make contact with 15 CDR resi-

dues on 60833. Of these 19 IGF-1 residues 17 are identical in human IGF-2 when the human IGF-1 and IGF-2 amino acid sequences are aligned (listed in Table 7). FIG. 11 shows the 3D structure of IGF-1 with the amino acids that are bound by 60833 highlighted, the linear amino acid sequence of human IGF-1 is also shown with the interacting amino acids underlined.

TABLE 7

RESIDUES IN HUMAN IGF-1 THAT MAKE CONTACTS WITH RESIDUES OF 60833 FAB		
IGF-1 residues in contact with 60833	Contact residues on 60833 (CDR)	Homologous residue on IGF-2
Leu (L) 5	Tyr (Y) 54; (HCDR 2)	Leu (L) 8
Cys (C) 6	Ser (S) 56; (HCDR 2)	Cys (C) 9
Glu (E) 9	Thr (T) 52; (HCDR 2)	Glu (E) 12
	Ser (S) 53; (HCDR 2)	
	Tyr (Y) 54; (HCDR 2)	
	Gly (G) 55; (HCDR 2)	
	Ser (S) 56; (HCDR 2)	
Leu (L) 10	Phe (F) 57; (HCDR 2)	Leu (L) 13
Asp (D) 12	Trp (W) 33; (HCDR 1)	Asp (D) 15
Ala (A) 13	Trp (W) 33; (HCDR 1)	—
Phe (F) 16	Trp (W) 33; (HCDR 1)	Phe (F) 19
	Arg (R) 92; (LCDR 3)	
	Tyr (Y) 98; (LCDR 3)	
	Trp (W) 99; (LCDR 3)	
	Tyr (Y) 101; (HCDR 3)	
Val (V) 17	Arg (R) 92; (LCDR 3)	Val (V) 20
	Tyr (Y) 98; (LCDR 3)	
Arg (R) 21	Tyr (Y) 95; (LCDR 3)	Arg (R) 24
Cys (C) 47	Ser (S) 56; (HCDR 2)	Cys (C) 46
	Phe (F) 57; (HCDR 2)	
Cys (C) 48	Ser (S) 56; (HCDR 2)	Cys (C) 47
Phe (F) 49	Tyr (Y) 54; (HCDR 2)	Phe (F) 48
	Gly (G) 55; (HCDR 2)	
	Ser (S) 56; (HCDR 2)	
Ser (S) 51	Gly (G) 55; (HCDR 2)	Ser (S) 50
	Ser (S) 56; (HCDR 2)	
	Thr (T) 58; (HCDR 2)	
Cys (C) 52	Ser (S) 56; (HCDR 2)	Cys (C) 51
	Phe (F) 57; (HCDR 2)	
	Thr (T) 58; (HCDR 2)	
Asp (D) 53	Phe (F) 57; (HCDR 2)	Asp (D) 52
	Thr (T) 58; (HCDR 2)	
Leu (L) 54	Trp (W) 33; (HCDR 1)	Leu (L) 53
	Phe (F) 57; (HCDR 2)	
	Thr (T) 58; (HCDR 2)	
	Tyr (Y) 98; (LCDR 3)	
Arg (R) 55	Lys (K) 65; (HCDR 2)	—
	Gly (G) 96; (LCDR 3)	
	Tyr (Y) 98; (LCDR 3)	
Leu (L) 57	Phe (F) 57; (HCDR 2)	Leu (L) 56
Glu (E) 58	Tyr (Y) 95; (LCDR 3)	Glu (E) 57
	Gly (G) 96; (LCDR 3)	
	Tyr (Y) 98; (LCDR 3)	
19 residues on IGF-1 involved in contact with 60833	15 residues on 60833 involved in contacts with IGF-1: HCDR 1: 1 residues HCDR 2: 8 residues HCDR 3: 1 residues LCDR 1: — LCDR 2: — LCDR 3: 5 residues	

REFERENCES

Barbas, et al., 1994, Proc. Nat. Acad. Sci, USA 91:3809-3813.
 Burtrum et al., Cancer Res. 63: 8912-21, 2003).
 Chen et al., J. Clin. Endocrinol. 90: 366-371, 2005.
 Cui et al., Science 299: 1753-55, 2003.
 Cruz-Correa et al., Gastroenterology 126: 1190-3, 2004.
 Dufner and Thomas, Exp. Cell Res. 253: 100-109, 1999.

Frasca et al., Mol. Cell. Biol. 19: 3278-88, 1999.
 Freier et al., Gut, May;44(5): 704-08, 1999;
 Fukuzawa et al., Int. J. Cancer 82: 490-497, 1999.
 Goetsch et al., Int. J. Cancer 113: 316-28, 2005.
 5 Goya et al., Cancer Res. 64: 6252-58, 2004.
 Haenel et al., Anal. Biochem. 339: 182-184, 2005.
 Hassan et al., Cancer Res. 60: 1070-6, 2000
 Hawkins et al., 1992, J. Mol. Biol. 226(3): 889 896.
 Jackson et al., 1995, J. Immunol. 154(7):3310-9.
 10 Jerome et al., End. Rel. Cancer 10: 561-578, 2003.
 Kabat et al., Sequences of Proteins of Immunological Interest (5th Ed.). NIH Publication No. 91 3242. U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health, Bethesda, Md. (1991).
 15 Kipriyanow and Le Gall, Molecular Biotechnology 26: 39-60, 2004.
 Knappik et al., J. Mol. Biol. 296: 57-86, 2000.
 Kolb et al. Pediatr. Blood Cancer 50: 1190-1197, 2008.
 Krebs, B. et al., J. Immunol. Meth. 245: 67 84, 2001.
 20 Kulik et al., Mol. Cell. Biol. 17: 1595-606, 1997.
 LeRoith D, Experimental Diab. Res. 4: 205-212, 2003.
 Li et al., Tumour Biol. 25: 62-8, 2004.
 Lowman et al., Biochemistry 30(45): 10832-10837, 1991.
 Lund et al., Cancer Lett. 206: 85-96, 2004.
 25 Manara et al., Clin. Cancer Res. 13: 1322-1330, 2007.
 Manes et al., Endocrinology 138: 905-915, 1997.
 Marks et al., 1992, Biotechnology 10:779-783.
 Miyamoto et al., Clin. Cancer Res. 11: 3494-3502, 2005.
 Moorhead et al., Oncogene 22: 853-7, 2003.
 30 Nagy et al., Nature Med. 8(8): 801-807, 2002.
 Ng et al., J. Gastroenterol. Hepatol. 13: 152-7, 1998.
 Pandini et al., J. Biol. Chem. 277: 39684-95, 2002.
 Pollack et al., Nature Rev. Can. 4: 505-518, 2004.
 Pollack et al., American Society for Clinical Oncology (ASCO), Annual Meeting 2007, abstract 3587.
 35 Quinn et al., J. Biol. Chem. 271: 11477-83, 1996.
 Rauchenberger, R. et al., J. Biol. Chem. 278: 38194-38205, 2003.
 Reinberg, U.S. News World Report, Mar. 5, 2008.
 40 Remington: "The Science and Practice of Pharmacy", 2005, 21st edition, Hendrickson Randy, Editor; Advanced Concepts Institute, University of The Sciences in Philadelphia, 600 S. 43rd Street, Philadelphia, Pa. 19104, USA; 215-895-1184.
 45 Renehan et al., Br. J. Cancer 83: 1344-50, 2000a).
 Renehan et al., J. Clin. Endocrinol. Metab. 85: 3402-8, 2000b).
 Revets et al., Expert Opin Biol Ther. 5(1):111-24, 2005.
 Rubin et al., Lab. Invest. 73: 311-31, 1995.
 50 Russell et al., Proc. Natl. Acad. Sci USA 81: 2389-2392, 1984.
 Scotlandi et al., Cancer Res. 56: 4570-4574, 1996.
 Sell et al., Natl. Acad. Sci. USA 90: 11217-21, 1993.
 Sell et al., Mol. Cell. Biol. 14: 3604-12, 1994.
 55 Shier et al., 1995, Gene 169:147-155.
 Shukla et al., 2007, J. Chromatography B, 848(1): 28-39
 Srinivasan, M. and Roeske, R W., Curr Protein Pept Sci. 2005, April;6(2):185-96.
 Strumberg D., 2005, Drugs Today (Barc). 2005 Dec;41(12): 773-84
 60 Takanami et al., J. Surg. Oncol. 61: 205-8, 1996.
 Tsai et al., Scand. J. Gastroenterol. 40: 68-75, 2005.
 Wang et al., World J. Gastroenterol. 9: 267-70, 2003.
 Woodson et al., J. Natl. Cancer Inst. 96: 407-10, 2004.
 65 Yao et al., Clin. Cancer Res. 9: 2719-26, 2003a).
 Yao et al., J. Clin. Invest. 111: 265-273, 2003b).
 Yelton et al., 1995, Immunol. 155:1994-2004.

Zapata et al., Protein Eng. 8(10): 1057-1062., 1995.
 Zhao et al., Cancer Epidemiol. Biomarkers Prev. 14: 1819-22,
 2005.
 WO 89/011297
 WO 94/29348
 WO 02/056910
 WO 03/002609
 WO 03/050531
 WO 03/093317
 WO 04/003019
 WO 04/058821

WO 2005/018671
 WO 2005/027970
 WO 2005/028515
 WO 2007/042309
 5 WO 2007/070432
 JP 2003-310275
 U.S. Pat. No. 4,342,566
 U.S. Pat. No. 3,773,919
 U.S. Pat. No. 6,696,245
 10 U.S. Pat. No. 6,991,790
 U.S. Pat. No. 7,060,268

 SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 43

<210> SEQ ID NO 1
 <211> LENGTH: 5
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

Asn Tyr Trp Met His
 1 5

<210> SEQ ID NO 2
 <211> LENGTH: 17
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

Gly Ile Ser Gly Trp Ser Ser Trp Thr Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> SEQ ID NO 3
 <211> LENGTH: 13
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

Phe Gly Ile Asp Ala Tyr Thr Lys Val Tyr Phe Asp Tyr
 1 5 10

<210> SEQ ID NO 4
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

Ser Gly Asp Asn Ile Pro Leu Lys Tyr Val Ser
 1 5 10

<210> SEQ ID NO 5
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

Asp Asp Asn Lys Arg Pro Ser
 1 5

<210> SEQ ID NO 6
 <211> LENGTH: 11
 <212> TYPE: PRT

-continued

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

Ser Ser Trp Asp Thr Leu Asp Ile Phe Asn Val
 1 5 10

<210> SEQ ID NO 7

<211> LENGTH: 366

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (1)..(366)

<400> SEQUENCE: 7

cag gtg gaa ttg gtg gaa agc ggc ggc ggc ctg gtg caa ccg ggc ggc 48
 Gln Val Glu Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

agc ctg cgt ctg agc tgc gcg gcc tcc gga ttt acc ttt tct aat tat 96
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Tyr
 20 25 30

tgg atg cat tgg gtg cgc caa gcc cct ggg aag ggt ctc gag tgg gtg 144
 Trp Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

agc ggt atc tct ggt tgg tct agc tgg acc tat tat gcg gat agc gtg 192
 Ser Gly Ile Ser Gly Trp Ser Ser Trp Thr Tyr Tyr Ala Asp Ser Val
 50 55 60

aaa ggc cgt ttt acc att tca cgt gat aat tcg aaa aac acc ctg tat 240
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

ctg caa atg aac agc ctg cgt gcg gaa gat acg gcc gtg tat tat tgc 288
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

gcg cgt ttt ggt att gat gct tat act aag gtt tat ttt gat tat tgg 336
 Ala Arg Phe Gly Ile Asp Ala Tyr Thr Lys Val Tyr Phe Asp Tyr Trp
 100 105 110

ggc caa ggc acc ctg gtg acg gtt agc tca 366
 Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 115 120

<210> SEQ ID NO 8

<211> LENGTH: 122

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

Gln Val Glu Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Tyr
 20 25 30

Trp Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ser Gly Ile Ser Gly Trp Ser Ser Trp Thr Tyr Tyr Ala Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Phe Gly Ile Asp Ala Tyr Thr Lys Val Tyr Phe Asp Tyr Trp
 100 105 110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser

-continued

115 120

<210> SEQ ID NO 9
 <211> LENGTH: 327
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(327)

<400> SEQUENCE: 9

gat atc gaa ctg acc cag ccg cct tca gtg agc gtt gca cca ggt cag 48
 Asp Ile Glu Leu Thr Gln Pro Pro Ser Val Ser Val Ala Pro Gly Gln
 1 5 10 15

acc gcg cgt atc tcg tgt agc ggc gat aat att cct ctt aag tat gtt 96
 Thr Ala Arg Ile Ser Cys Ser Gly Asp Asn Ile Pro Leu Lys Tyr Val
 20 25 30

tct tgg tac cag cag aaa ccc ggg cag gcg cca gtt ctt gtg att cat 144
 Ser Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Val Leu Val Ile His
 35 40 45

gat gat aat aag cgt ccc tca ggc atc ccg gaa cgc ttt agc gga tcc 192
 Asp Asp Asn Lys Arg Pro Ser Gly Ile Pro Glu Arg Phe Ser Gly Ser
 50 55 60

aac agc ggc aac acc gcg acc ctg acc att agc ggc act cag gcg gaa 240
 Asn Ser Gly Asn Thr Ala Thr Leu Thr Ile Ser Gly Thr Gln Ala Glu
 65 70 75 80

gac gaa gcg gat tat tat tgc tct tct tgg gat act ctt gat att ttt 288
 Asp Glu Ala Asp Tyr Tyr Cys Ser Ser Trp Asp Thr Leu Asp Ile Phe
 85 90 95

aat gtg ttt ggc ggc ggc acg aag tta acc gtc cta ggt 327
 Asn Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
 100 105

<210> SEQ ID NO 10
 <211> LENGTH: 109
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

Asp Ile Glu Leu Thr Gln Pro Pro Ser Val Ser Val Ala Pro Gly Gln
 1 5 10 15

Thr Ala Arg Ile Ser Cys Ser Gly Asp Asn Ile Pro Leu Lys Tyr Val
 20 25 30

Ser Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Val Leu Val Ile His
 35 40 45

Asp Asp Asn Lys Arg Pro Ser Gly Ile Pro Glu Arg Phe Ser Gly Ser
 50 55 60

Asn Ser Gly Asn Thr Ala Thr Leu Thr Ile Ser Gly Thr Gln Ala Glu
 65 70 75 80

Asp Glu Ala Asp Tyr Tyr Cys Ser Ser Trp Asp Thr Leu Asp Ile Phe
 85 90 95

Asn Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
 100 105

<210> SEQ ID NO 11
 <211> LENGTH: 5
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 11

Asn Tyr Trp Met His
 1 5

-continued

<210> SEQ ID NO 12
 <211> LENGTH: 17
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 12

Gly Ile Ser Gly Trp Ser Ser Trp Thr Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> SEQ ID NO 13
 <211> LENGTH: 13
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 13

Phe Gly Ile Asp Ala Tyr Thr Lys Val Tyr Phe Asp Tyr
 1 5 10

<210> SEQ ID NO 14
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 14

Ser Gly Asp Asn Ile Pro Leu Lys Tyr Val Ser
 1 5 10

<210> SEQ ID NO 15
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 15

Asp Asp Asn Lys Arg Pro Ser
 1 5

<210> SEQ ID NO 16
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 16

Gln Ser Tyr Asp Tyr Phe Pro Lys Phe Val Val
 1 5 10

<210> SEQ ID NO 17
 <211> LENGTH: 366
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(366)

<400> SEQUENCE: 17

cag gtg gaa ttg gtg gaa agc ggc ggc ggc ctg gtg caa ccg ggc ggc 48
 Gln Val Glu Leu Val Glu Ser Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

agc ctg cgt ctg agc tgc gcg gcc tcc gga ttt acc ttt tct aat tat 96
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Tyr
 20 25 30

tgg atg cat tgg gtg cgc caa gcc cct ggg aag ggt ctc gag tgg gtg 144
 Trp Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

-continued

```

aac agc ggc aac acc gcg acc ctg acc att agc ggc act cag gcg gaa      240
Asn Ser Gly Asn Thr Ala Thr Leu Thr Ile Ser Gly Thr Gln Ala Glu
65                      70                      75                      80

```

```

gac gaa gcg gat tat tat tgc cag tct tat gat tat ttt cct aag ttt      288
Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Tyr Phe Pro Lys Phe
85                      90                      95

```

```

gtt gtg ttt ggc ggc ggc acg aag tta acc gtc cta ggt      327
Val Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
100                      105

```

```

<210> SEQ ID NO 20
<211> LENGTH: 109
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 20

```

```

Asp Ile Glu Leu Thr Gln Pro Pro Ser Val Ser Val Ala Pro Gly Gln
1                      5                      10                      15

```

```

Thr Ala Arg Ile Ser Cys Ser Gly Asp Asn Ile Pro Leu Lys Tyr Val
20                      25                      30

```

```

Ser Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Val Leu Val Ile His
35                      40                      45

```

```

Asp Asp Asn Lys Arg Pro Ser Gly Ile Pro Glu Arg Phe Ser Gly Ser
50                      55                      60

```

```

Asn Ser Gly Asn Thr Ala Thr Leu Thr Ile Ser Gly Thr Gln Ala Glu
65                      70                      75                      80

```

```

Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Tyr Phe Pro Lys Phe
85                      90                      95

```

```

Val Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
100                      105

```

```

<210> SEQ ID NO 21
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 21

```

```

Ser Tyr Trp Met Ser
1                      5

```

```

<210> SEQ ID NO 22
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 22

```

```

Ser Ile Thr Ser Tyr Gly Ser Phe Thr Tyr Tyr Ala Asp Ser Val Lys
1                      5                      10                      15

```

```

Gly

```

```

<210> SEQ ID NO 23
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 23

```

```

Asn Met Tyr Thr His Phe Asp Ser
1                      5

```

```

<210> SEQ ID NO 24
<211> LENGTH: 13
<212> TYPE: PRT

```

-continued

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 24

Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn Ser Val Ser
 1 5 10

<210> SEQ ID NO 25

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 25

Asp Asn Ser Lys Arg Pro Ser
 1 5

<210> SEQ ID NO 26

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 26

Gln Ser Arg Asp Thr Tyr Gly Tyr Tyr Trp Val
 1 5 10

<210> SEQ ID NO 27

<211> LENGTH: 351

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (1)..(351)

<400> SEQUENCE: 27

cag gtg gaa ttg gtg gaa agc ggc ggc ggc ctg gtg caa ccg ggc ggc 48
 Gln Val Glu Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

agc ctg cgt ctg agc tgc gcg gcc tcc gga ttt acc ttt act tct tat 96
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Thr Ser Tyr
 20 25 30

tgg atg tct tgg gtg cgc caa gcc cct ggg aag ggt ctc gag ctt gtg 144
 Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Leu Val
 35 40 45

agc tct atc act tct tat ggt agc ttt acc tat tat gcg gat agc gtg 192
 Ser Ser Ile Thr Ser Tyr Gly Ser Phe Thr Tyr Tyr Ala Asp Ser Val
 50 55 60

aaa ggc cgt ttt acc att tca cgt gat aat tcg aaa aac acc ctg tat 240
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

ctg caa atg aac agc ctg cgt gcg gaa gat acg gcc gtg tat tat tgc 288
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

gcg cgt aat atg tat act cat ttt gat tct tgg ggc caa ggc acc ctg 336
 Ala Arg Asn Met Tyr Thr His Phe Asp Ser Trp Gly Gln Gly Thr Leu
 100 105 110

gtg acg gtt agc tca 351
 Val Thr Val Ser Ser
 115

<210> SEQ ID NO 28

<211> LENGTH: 117

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 28

-continued

Gln Val Glu Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Thr Ser Tyr
 20 25 30
 Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Leu Val
 35 40 45
 Ser Ser Ile Thr Ser Tyr Gly Ser Phe Thr Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Asn Met Tyr Thr His Phe Asp Ser Trp Gly Gln Gly Thr Leu
 100 105 110
 Val Thr Val Ser Ser
 115

<210> SEQ ID NO 29
 <211> LENGTH: 333
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(333)

<400> SEQUENCE: 29

gat atc gtg ctg acc cag ccg cct tca gtg agt ggc gca cca ggt cag 48
 Asp Ile Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
 1 5 10 15
 cgt gtg acc atc tcg tgt agc ggc agc agc agc aac att ggt tct aat 96
 Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn
 20 25 30
 tct gtg tct tgg tac cag cag ttg ccc ggg acg gcg ccg aaa ctt ctg 144
 Ser Val Ser Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
 35 40 45
 att tat gat aat tct aag cgt ccc tca ggc gtg ccg gat cgt ttt agc 192
 Ile Tyr Asp Asn Ser Lys Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
 50 55 60
 gga tcc aaa agc ggc acc agc gcg agc ctt gcg att acg ggc ctg caa 240
 Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Leu Gln
 65 70 75 80
 agc gaa gac gaa gcg gat tat tat tgc cag tct cgt gat act tat ggt 288
 Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Arg Asp Thr Tyr Gly
 85 90 95
 tat tat tgg gtg ttt ggc ggc ggc acg aag tta acc gtc cta ggt 333
 Tyr Tyr Trp Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
 100 105 110

<210> SEQ ID NO 30
 <211> LENGTH: 111
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 30

Asp Ile Val Leu Thr Gln Pro Pro Ser Val Ser Gly Ala Pro Gly Gln
 1 5 10 15
 Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn
 20 25 30
 Ser Val Ser Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
 35 40 45

-continued

Ile Tyr Asp Asn Ser Lys Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
 50 55 60
 Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Thr Gly Leu Gln
 65 70 75 80
 Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Arg Asp Thr Tyr Gly
 85 90 95
 Tyr Tyr Trp Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
 100 105 110

<210> SEQ ID NO 31
 <211> LENGTH: 993
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(993)

<400> SEQUENCE: 31

gcc tcc acc aag ggt cca tcc gtc ttc ccc ctg gca ccc tcc tcc aag 48
 Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
 1 5 10 15
 agc acc tct ggg ggc aca gcg gcc ctg ggc tgc ctg gtc aag gac tac 96
 Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20 25 30
 ttc ccc gaa ccg gtg acg gtg tcc tgg aac tca ggc gcc ctg acc agc 144
 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35 40 45
 ggc gtg cac acc ttc ccg gct gtc cta cag tcc tca gga ctc tac tcc 192
 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50 55 60
 ctc agc agc gtg gtg acc gtg ccc tcc agc agc ttg ggc acc cag acc 240
 Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
 65 70 75 80
 tac atc tgc aac gtg aat cac aag ccc agc aac acc aag gtg gac aag 288
 Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
 85 90 95
 aaa gtt gag ccc aaa tct tgt gac aaa act cac aca tgc cca ccg tgc 336
 Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
 100 105 110
 cca gca cct gaa ctc ctg ggg gga ccg tca gtc ttc ctc ttc ccc cca 384
 Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
 115 120 125
 aaa ccc aag gac acc ctc atg atc tcc cgg acc cct gag gtc aca tgc 432
 Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
 130 135 140
 gtg gtg gtg gac gtg agc cac gaa gac cct gag gtc aag ttc aac tgg 480
 Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
 145 150 155 160
 tac gtg gac ggc gtg gag gtg cat aat gcc aag aca aag ccg cgg gag 528
 Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
 165 170 175
 gag cag tac aac agc acg tac cgg gtg gtc agc gtc ctc acc gtc ctg 576
 Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
 180 185 190
 cac cag gac tgg ctg aat ggc aag gag tac aag tgc aag gtc tcc aac 624
 His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
 195 200 205
 aaa gcc ctc cca gcc ccc atc gag aaa acc atc tcc aaa gcc aaa ggg 672
 Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
 210 215 220
 cag ccc cga gaa cca cag gtg tac acc ctg ccc cca tcc cgg gat gag 720

-continued

Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
 245 250 255

Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
 260 265 270

Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
 275 280 285

Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
 290 295 300

Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
 305 310 315 320

Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 325 330

<210> SEQ ID NO 33
 <211> LENGTH: 318
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(318)

<400> SEQUENCE: 33

cag ccc aag gct gcc ccc tcg gtc act ctg ttc ccg ccc tcc tct gag 48
 Gln Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu
 1 5 10 15

gag ctt caa gcc aac aag gcc aca ctg gtg tgt ctc ata agt gac ttc 96
 Glu Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe
 20 25 30

tac ccg gga gcc gtg aca gtg gcc tgg aag gga gat agc agc ccc gtc 144
 Tyr Pro Gly Ala Val Thr Val Ala Trp Lys Gly Asp Ser Ser Pro Val
 35 40 45

aag gcg gga gtg gag acc acc aca ccc tcc aaa caa agc aac aac aag 192
 Lys Ala Gly Val Glu Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys
 50 55 60

tac gcg gcc agc agc tat ctg agc ctg acg cct gag cag tgg aag tcc 240
 Tyr Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser
 65 70 75 80

cac aga agc tac agc tgc cag gtc acg cat gaa ggg agc acc gtg gag 288
 His Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu
 85 90 95

aag aca gtg gcc cct aca gaa tgt tca tag 318
 Lys Thr Val Ala Pro Thr Glu Cys Ser
 100 105

<210> SEQ ID NO 34
 <211> LENGTH: 105
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 34

Gln Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu
 1 5 10 15

Glu Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe
 20 25 30

Tyr Pro Gly Ala Val Thr Val Ala Trp Lys Gly Asp Ser Ser Pro Val
 35 40 45

Lys Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys
 50 55 60

Tyr Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser
 65 70 75 80

-continued

His Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu
 85 90 95

 Lys Thr Val Ala Pro Thr Glu Cys Ser
 100 105

 <210> SEQ ID NO 35
 <211> LENGTH: 452
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 35

 Gln Val Glu Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Tyr
 20 25 30

 Trp Met His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

 Ser Gly Ile Ser Gly Trp Ser Ser Trp Thr Tyr Tyr Ala Asp Ser Val
 50 55 60

 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

 Ala Arg Phe Gly Ile Asp Ala Tyr Thr Lys Val Tyr Phe Asp Tyr Trp
 100 105 110

 Gly Gln Gly Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro
 115 120 125

 Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr
 130 135 140

 Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr
 145 150 155 160

 Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro
 165 170 175

 Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr
 180 185 190

 Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn
 195 200 205

 His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser
 210 215 220

 Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu
 225 230 235 240

 Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu
 245 250 255

 Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser
 260 265 270

 His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu
 275 280 285

 Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr
 290 295 300

 Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn
 305 310 315 320

 Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro
 325 330 335

 Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln
 340 345 350

-continued

Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val
 355 360 365

Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val
 370 375 380

Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro
 385 390 395 400

Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr
 405 410 415

Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val
 420 425 430

Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu
 435 440 445

Ser Pro Gly Lys
 450

<210> SEQ ID NO 36
 <211> LENGTH: 214
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 36

Asp Ile Glu Leu Thr Gln Pro Pro Ser Val Ser Val Ala Pro Gly Gln
 1 5 10 15

Thr Ala Arg Ile Ser Cys Ser Gly Asp Asn Ile Pro Leu Lys Tyr Val
 20 25 30

Ser Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Val Leu Val Ile His
 35 40 45

Asp Asp Asn Lys Arg Pro Ser Gly Ile Pro Glu Arg Phe Ser Gly Ser
 50 55 60

Asn Ser Gly Asn Thr Ala Thr Leu Thr Ile Ser Gly Thr Gln Ala Glu
 65 70 75 80

Asp Glu Ala Asp Tyr Tyr Cys Ser Ser Trp Asp Thr Leu Asp Ile Phe
 85 90 95

Asn Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln Pro Lys
 100 105 110

Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu Glu Leu Gln
 115 120 125

Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe Tyr Pro Gly
 130 135 140

Ala Val Thr Val Ala Trp Lys Gly Asp Ser Ser Pro Val Lys Ala Gly
 145 150 155 160

Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys Tyr Ala Ala
 165 170 175

Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser His Arg Ser
 180 185 190

Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu Lys Thr Val
 195 200 205

Ala Pro Thr Glu Cys Ser
 210

<210> SEQ ID NO 37
 <211> LENGTH: 452
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 37

Gln Val Glu Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly

-continued

1	5	10	15
Ser Leu Arg	Leu Ser Cys Ala Ala	Ser Gly Phe Thr Phe	Ser Asn Tyr
	20	25	30
Trp Met His	Trp Val Arg Gln Ala Pro	Gly Lys Gly Leu Glu Trp Val	
	35	40	45
Ser Gly Ile	Ser Gly Trp Ser Ser Trp Thr Tyr Tyr	Ala Asp Ser Val	
	50	55	60
Lys Gly Arg	Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr		
	65	70	75
Leu Gln Met	Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys		
	85	90	95
Ala Arg Phe	Gly Ile Asp Ala Tyr Thr Lys Val Tyr Phe Asp Tyr Trp		
	100	105	110
Gly Gln Gly	Thr Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro		
	115	120	125
Ser Val Phe	Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr		
	130	135	140
Ala Ala Leu	Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr		
	145	150	155
Val Ser Trp	Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro		
	165	170	175
Ala Val Leu	Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr		
	180	185	190
Val Pro Ser	Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn		
	195	200	205
His Lys Pro	Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro Lys Ser		
	210	215	220
Cys Asp Lys	Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu		
	225	230	235
Gly Gly Pro	Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu		
	245	250	255
Met Ile Ser	Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser		
	260	265	270
His Glu Asp	Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu		
	275	280	285
Val His Asn	Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr		
	290	295	300
Tyr Arg Val	Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn		
	305	310	315
Gly Lys Glu	Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro		
	325	330	335
Ile Glu Lys	Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln		
	340	345	350
Val Tyr Thr	Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val		
	355	360	365
Ser Leu Thr	Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val		
	370	375	380
Glu Trp Glu	Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro		
	385	390	395
Pro Val Leu	Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr		
	405	410	415
Val Asp Lys	Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val		
	420	425	430

-continued

Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu
 435 440 445

Ser Pro Gly Lys
 450

<210> SEQ ID NO 38
 <211> LENGTH: 214
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 38

Asp Ile Glu Leu Thr Gln Pro Pro Ser Val Ser Val Ala Pro Gly Gln
 1 5 10 15

Thr Ala Arg Ile Ser Cys Ser Gly Asp Asn Ile Pro Leu Lys Tyr Val
 20 25 30

Ser Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Val Leu Val Ile His
 35 40 45

Asp Asp Asn Lys Arg Pro Ser Gly Ile Pro Glu Arg Phe Ser Gly Ser
 50 55 60

Asn Ser Gly Asn Thr Ala Thr Leu Thr Ile Ser Gly Thr Gln Ala Glu
 65 70 75 80

Asp Glu Ala Asp Tyr Tyr Cys Gln Ser Tyr Asp Tyr Phe Pro Lys Phe
 85 90 95

Val Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly Gln Pro Lys
 100 105 110

Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu Glu Leu Gln
 115 120 125

Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe Tyr Pro Gly
 130 135 140

Ala Val Thr Val Ala Trp Lys Gly Asp Ser Ser Pro Val Lys Ala Gly
 145 150 155 160

Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys Tyr Ala Ala
 165 170 175

Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser His Arg Ser
 180 185 190

Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu Lys Thr Val
 195 200 205

Ala Pro Thr Glu Cys Ser
 210

<210> SEQ ID NO 39
 <211> LENGTH: 447
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 39

Gln Val Glu Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Thr Ser Tyr
 20 25 30

Trp Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Leu Val
 35 40 45

Ser Ser Ile Thr Ser Tyr Gly Ser Phe Thr Tyr Tyr Ala Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys

-continued

85				90				95							
Ala	Arg	Asn	Met	Tyr	Thr	His	Phe	Asp	Ser	Trp	Gly	Gln	Gly	Thr	Leu
			100					105					110		
Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu
		115					120					125			
Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys
	130					135					140				
Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser
145					150					155					160
Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser
				165					170					175	
Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser
			180					185					190		
Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn
		195					200					205			
Thr	Lys	Val	Asp	Lys	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His
	210					215					220				
Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val
225					230					235					240
Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr
				245					250					255	
Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu
			260					265					270		
Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys
		275					280					285			
Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser
	290					295					300				
Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys
305					310					315					320
Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile
				325					330					335	
Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro
			340					345					350		
Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu
		355					360					365			
Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn
	370					375					380				
Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser
385					390					395					400
Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg
			405						410					415	
Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu
			420					425					430		
His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys	
	435						440					445			

<210> SEQ ID NO 40

<211> LENGTH: 216

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 40

Asp	Ile	Val	Leu	Thr	Gln	Pro	Pro	Ser	Val	Ser	Gly	Ala	Pro	Gly	Gln
1				5					10					15	

Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ser Asn

-continued

20				25				30							
Ser	Val	Ser	Trp	Tyr	Gln	Gln	Leu	Pro	Gly	Thr	Ala	Pro	Lys	Leu	Leu
		35					40					45			
Ile	Tyr	Asp	Asn	Ser	Lys	Arg	Pro	Ser	Gly	Val	Pro	Asp	Arg	Phe	Ser
	50					55					60				
Gly	Ser	Lys	Ser	Gly	Thr	Ser	Ala	Ser	Leu	Ala	Ile	Thr	Gly	Leu	Gln
65					70					75					80
Ser	Glu	Asp	Glu	Ala	Asp	Tyr	Tyr	Cys	Gln	Ser	Arg	Asp	Thr	Tyr	Gly
				85					90					95	
Tyr	Tyr	Trp	Val	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Thr	Val	Leu	Gly	Gln
			100						105				110		
Pro	Lys	Ala	Ala	Pro	Ser	Val	Thr	Leu	Phe	Pro	Pro	Ser	Ser	Glu	Glu
		115					120						125		
Leu	Gln	Ala	Asn	Lys	Ala	Thr	Leu	Val	Cys	Leu	Ile	Ser	Asp	Phe	Tyr
	130					135					140				
Pro	Gly	Ala	Val	Thr	Val	Ala	Trp	Lys	Gly	Asp	Ser	Ser	Pro	Val	Lys
145					150					155					160
Ala	Gly	Val	Glu	Thr	Thr	Thr	Pro	Ser	Lys	Gln	Ser	Asn	Asn	Lys	Tyr
				165					170					175	
Ala	Ala	Ser	Ser	Tyr	Leu	Ser	Leu	Thr	Pro	Glu	Gln	Trp	Lys	Ser	His
			180						185				190		
Arg	Ser	Tyr	Ser	Cys	Gln	Val	Thr	His	Glu	Gly	Ser	Thr	Val	Glu	Lys
		195					200						205		
Thr	Val	Ala	Pro	Thr	Glu	Cys	Ser								
	210					215									

<210> SEQ ID NO 41
 <211> LENGTH: 17
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 41

Leu	Cys	Gly	Ala	Glu	Leu	Val	Asp	Ala	Leu	Gln	Phe	Val	Cys	Gly	Asp
1				5					10					15	

Arg

<210> SEQ ID NO 42
 <211> LENGTH: 13
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 42

Cys	Cys	Phe	Arg	Ser	Cys	Asp	Leu	Arg	Arg	Leu	Glu	Met
1				5					10			

<210> SEQ ID NO 43
 <211> LENGTH: 70
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 43

Gly	Pro	Glu	Thr	Leu	Cys	Gly	Ala	Glu	Leu	Val	Asp	Ala	Leu	Gln	Phe
1				5					10					15	

Val	Cys	Gly	Asp	Arg	Gly	Phe	Tyr	Phe	Asn	Lys	Pro	Thr	Gly	Tyr	Gly
			20					25					30		

Ser	Ser	Ser	Arg	Arg	Ala	Pro	Gln	Thr	Gly	Ile	Val	Asp	Glu	Cys	Cys
		35					40					45			

Phe	Arg	Ser	Cys	Asp	Leu	Arg	Arg	Leu	Glu	Met	Tyr	Cys	Ala	Pro	Leu
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

