

US008367062B2

(12) **United States Patent**
Lombardo et al.

(10) **Patent No.:** US 8,367,062 B2
(45) **Date of Patent:** Feb. 5, 2013

(54) **GLYCOPEPTIDES DERIVED FROM PANCREATIC STRUCTURES, ANTIBODIES AND APPLICATIONS THEREOF IN DIAGNOSTICS AND THERAPEUTICS**

(75) Inventors: **Dominique Lombardo**, Marseilles (FR); **Eric Mas**, Marseilles (FR); **Marie-Odile Sadoulet**, Marseilles (FR); **Laurence Panicot-Dubois**, Canohes (FR); **Jean-Paul Bernard**, Marseilles (FR)

(73) Assignee: **Universite d'Aix-Marseille**, Marseilles Cedex (FR)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 583 days.

(21) Appl. No.: **12/488,793**

(22) Filed: **Jun. 22, 2009**

(65) **Prior Publication Data**

US 2009/0285823 A1 Nov. 19, 2009

Related U.S. Application Data

(62) Division of application No. 10/593,859, filed as application No. PCT/FR2005/000771 on Mar. 30, 2005, now Pat. No. 7,557,193.

(30) **Foreign Application Priority Data**

Mar. 31, 2004 (FR) 04 03378
Dec. 16, 2004 (FR) 04 13428

(51) **Int. Cl.**
A61K 39/395 (2006.01)

(52) **U.S. Cl.** 424/133.1; 530/388.1

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

6,180,370 B1 1/2001 Queen et al.

FOREIGN PATENT DOCUMENTS

WO WO 94/20610 9/2004

OTHER PUBLICATIONS

Takeda et al. (J. Cancer Res. Clin. Oncol., vol. 118(5), pp. 377-385, May 1992).*

Lombardo, D. "Bile Salt-Dependent Lipase: Its Pathophysiological Implications", *Biochimica et Biophysica Acta*, May 8, 2001, pp. 1-28, vol. 1533.

Mas, E. et al. "The Oncofetal J28 Epitope Involves Fucosylated O-Linked Oligosaccharide Structures of the Fetoacinar Pancreatic Protein", *Glycobiology*, 1997, pp. 745-752, vol. 7, No. 6.

Panicot, L. et al. "The Formation of the Oncofetal J28 Glycotope Involves Core-2 β 6-N-Acetylglucosaminyltransferase and α 3/4-Fusosyltransferase Activities", *Glycobiology*, 1999, pp. 935-946, vol. 9, No. 9.

Reue, K. et al. "cDNA Cloning of Carboxyl Ester Lipase from Human Pancreas Reveals a Unique Proline-Rich Repeat Unit", *Journal of Lipid Research*, 1991, pp. 267-276, vol. 32.

Panicot-Dubois, L. et al. "Monoclonal Antibody 16D10 to the C-Terminal Domain of the Feta-Acinar Pancreatic Protein Binds to Membrane of Human Pancreatic Tumoral SOJ-6 Cells and Inhibits the Growth of Tumor Xenografts", *Neoplasia*, 2004, pp. 713-724, vol. 6, No. 6.

Takeda, Y. et al. "Radioimmunolocalization of the monoclonal antibody J28 in early transformation stages in *N*-nitrosobis(2-hydroxypropyl)amine-induced pancreatic tumors in the Syrian golden hamster", *J. Cancer Research Clinical Oncology*, 1992, pp. 377-385, vol. 118, No. 5.

Holmes, S. et al. "Monoclonal Antibodies", *Hybridoma*, 2000, p. 503, vol. 19, No. 6.

Thirion, S. et al. "Mono- and bispecific single-chain antibody fragments for cancer therapy", *Eur. J. Cancer Prev.*, 1996, pp. 507-511, vol. 5, No. 6, abstract only.

* cited by examiner

Primary Examiner — Laura B Goddard

Assistant Examiner — Meera Natarajan

(74) *Attorney, Agent, or Firm* — Saliwanchik, Lloyd & Eisenschenk

(57) **ABSTRACT**

Glycopeptide comprising 1 to 40 repeated C-terminal polypeptides, with 11 amino acids, of BSDL or FAPP, whereby said polypeptides are glycosylated and bear glycosylated epitopes giving rise to a specific immunological reaction with induced antibodies in a patient suffering from type 1 diabetes and/or purified from biological fluids of human or animal origin or recombinant and produced by expression in a standard host cell comprising an enzymatic material necessary for priming a glycosylation, said host cell being genetically modified such as to comprise a gene coding for said polypeptides and a gene coding for one or more enzymes selected from among glycosyltransferases, anti-glycopeptide antibodies and the applications thereof in therapeutics and diagnostics.

11 Claims, 4 Drawing Sheets






FIGURE 1



Exp. 1



Exp. 2

FIGURE 2




FIGURE 3




FIGURE 4

**GLYCOPEPTIDES DERIVED FROM
PANCREATIC STRUCTURES, ANTIBODIES
AND APPLICATIONS THEREOF IN
DIAGNOSTICS AND THERAPEUTICS**

CROSS-REFERENCE TO RELATED
APPLICATIONS

This application is a divisional of U.S. Ser. No. 10/593,859, filed Sep. 22, 2006, now U.S. Pat. No. 7,557,193, which is the U.S. national stage application of International Patent Application No. PCT/FR2005/000771, filed Mar. 30, 2005, the disclosures of which are hereby incorporated by reference in their entirety, including all figures, tables and amino acid or nucleic acid sequences.

The invention relates to glycopeptides derived from pancreatic structures, antibodies and applications thereof in diagnostics and therapeutics.

More particularly, the invention relates to the obtaining of natural C-terminal glycopeptides of BSDL and/or FAPP carrying various glycosylated epitopes (alone or combined) recognized by induced antibodies in a patient suffering from type I diabetes and purified from biological fluids of human or animal origin as well as to said recombinant glycopeptides produced by biological engineering from cell lines containing cDNAs coding for the C-terminal peptides of FAPP and/or BSDL and cDNAs coding for the different glycosyltransferases required to build said glycopeptides. The invention also relates to monoclonal antibodies recognizing the recombinant and/or natural C-terminal glycopeptides of BSDL and/or FAPP, and to the use of said glycopeptides and said monoclonal antibodies as immunogenic agents, diagnostic agents and therapeutic agents that can be used for the diagnosis or the preventive or curative treatment of pancreatic pathologies such as diabetes and cancer of the exocrine pancreas or any other pathologies as well as in curative or preventive protocols of said pathologies.

Cancer of the exocrine pancreas, which accounts for over 20% of digestive tract cancers, is one of the most aggressive. In France for example, 4,000 new cases are diagnosed each year. Furthermore, its frequency is rising markedly in many regions of the world. Survival rates do not exceed 20% at 1 year and 3% at 5 years and mean survival is 3 to 4 months after diagnosis. The cancer is diagnosed late and progresses very rapidly, mainly through the formation of peritoneal and hepatic metastases. In addition, the deep anatomic location of the tumor, the absence of sensitive and specific early biological markers and its asymptomatic nature result in a diagnosis that always occurs late. Currently there are no effective therapies for exocrine pancreatic cancer, which is refractory to chemotherapy and radiation.

It would therefore be desirable to have specific markers of pancreatic cancer or other pancreatic pathologies available so as to diagnose these diseases and have products to provide effective treatments.

Now, after extensive research, and in a surprising manner, the applicants have discovered specific markers of pancreatic pathologies, and in particular of pancreatic cancer, expressed at the surface of the pancreatic tumor cell.

The applicants have discovered circulating antibodies directed against glycosylated antigenic structures of the C-terminal peptides of bile salt dependent lipase (BSDL), in diabetic patients.

The applicants have also discovered that monoclonal and polyclonal antibodies anti-glycosylated epitope expressed in the C-terminal part of human fetoacinar pancreatic protein

(FAPP) specifically recognized human pancreatic tumor cell lines, but did not recognize tumor cell lines which were not of pancreatic origin.

The applicants have also discovered that monoclonal and polyclonal antibodies anti-glycosylated epitope expressed in the C-terminal part of human fetoacinar pancreatic protein (FAPP) specifically recognized human pancreatic tumor tissue, but did not recognize normal pancreatic tissue.

The applicants have further discovered that the antibodies anti-glycosylated epitope expressed in the C-terminal part of BSDL or FAPP could detect BSDL- and FAPP-derived glycopeptides in urine. In particular, the antibodies anti-glycosylated epitope expressed in the C-terminal part of FAPP could detect glycopeptides in urine allowing to identify a subject with a pancreatic pathology, particularly pancreatic cancer.

More particularly, therefore, the applicants have discovered that compounds having a glycopeptide structure whose peptide part is based on the repeated C-terminal sequences of BSDL, a digestive lipolytic enzyme present in normal pancreatic secretions, or based on the repeated C-terminal sequences of FAPP (an oncofetal form of BSDL) constituted such specific markers of pancreatic pathologies.

Indeed, BSDL and FAPP comprise repeated C-terminal peptide sequences of 11 amino acids, comprising a generally invariant part with 7 amino acids having the sequence Ala Pro Pro Val Pro Pro Thr and a glycosylation site. Said generally invariant part is flanked on either side by a glycine often substituted by a glutamic acid and contains the amino acids Asp and Ser on the N-terminal side.

The applicants have also discovered that the compounds having a glycopeptide structure which could be prepared by expression and secretion by a host cell, for example from Chinese hamster ovary (CHO), comprising a gene construct including a DNA molecule coding for one or more repeated sequences of the C-terminal peptide, particularly recombinant of BSDL, for example all or part of the 16 repeated sequences and also comprising a gene construct such as a DNA molecule coding for at least one enzyme with α -mannosyltransferase activity, in particular selected in the group consisting of Core 2 β (1-6) N-acetylglucosaminyltransferase, fucosyltransferase FUT3 which has α (1-3) and α (1-4) fucosyltransferase activity, or fucosyltransferase FUT7 which only has α (1-3) fucosyltransferase activity, constituted said specific markers of pancreatic cancer.

Thus the present application has as object a glycopeptide, particularly recombinant, possibly isolated or purified, comprising from 1 to 40 repeated C-terminal polypeptides, composed of 11 amino acids, of BSDL or FAPP, said polypeptides being glycosylated and carrying glycosylated epitopes giving rise to a specific immunological reaction with induced antibodies in a patient with type I diabetes and

or else purified from biological fluids of human or animal origin

or else recombinant and being produced by expression in a conventional host cell comprising an enzymatic machinery necessary for priming a glycosylation, said host cell being genetically modified so as to comprise a gene coding for said polypeptide and a gene coding for one or more enzymes selected from glycosyltransferases and in particular from Core2 β (1-6) N-acetylglucosaminyltransferase (abbreviated C2GnT), α (1-3) galactosyltransferase, fucosyltransferase 3 (abbreviated FUT3) and fucosyltransferase 7 (abbreviated FUT7).

In other preferred conditions of implementation of the invention, the aforementioned glycopeptide is essentially constituted of 1 to 40 glycosylated repeated C-terminal pep-

tides, with 11 amino acids of BSDL or FAPP, and in particular exclusively constituted of said glycosylated peptides. Hence, the invention therefore relates to isolated, purified or recombinant glycopeptides comprising, or essentially consisting of, repetitions of the repeated C-terminal peptide sequence of 11 amino acids, preferably: D-S-G/E-A-P-P-V-P-P-T-G/E (SEQ ID No 14). Said glycopeptides can contain 1 to 40 repetitions. In a preferred embodiment, said glycopeptides comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 or 40 repetitions. Preferably, the glycopeptides comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 or 40 repetitions. In a particular embodiment, the glycopeptides comprise 1, 2, 3, 4, 5, 7, 8, 9, 10, 11, 12, 13, 14, 15, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 or 40 repetitions. In another alternative embodiment, the glycopeptides comprise between 1 and 15 repetitions (for example, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, or 15 repetitions), preferably between 2 and 10 repetitions (for example, 2, 3, 4, 5, 6, 7, 8, 9, or 10 repetitions). In an additional embodiment, the glycopeptides comprise between 17 and 40 repetitions (for example, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39 or 40 repetitions). In preferred conditions of carrying out the invention, the aforementioned glycopeptide comprises from 1 to 40, preferably from 4 to 25 and more particularly from 6 to 16 repeated C-terminal polypeptides. In a particular embodiment, the glycopeptide comprises, or essentially consists of, 6 repetitions.

Preferably, the glycopeptides according to the invention are glycosylated by one or more enzymes having ose-transferase activity selected in the group consisting of Core 2 β (1-6) N-acetylglucosaminyltransferase (C2GnT), fucosyltransferase FUT3 which has α (1-3) and α (1-4) fucosyltransferase activity, or fucosyltransferase FUT7 which only has α (1-3) fucosyltransferase activity. In a particular embodiment, the glycopeptides according to the invention have been glycosylated by the enzymes C2GnT and FUT3. In an alternative embodiment, the glycopeptides according to the invention have been glycosylated by the enzymes C2GnT and FUT7. The invention also encompasses glycopeptides which have been glycosylated by the enzymes C2GnT, FUT3 and FUT7. In a preferred embodiment, the glycopeptides according to the invention have additionally been glycosylated by α (1-3)galactosyltransferase (GT). Thus, when the glycopeptide according to the invention is recombinant, the host cell which produces it comprises a gene coding for said polypeptide and a gene coding for one or more enzymes selected from glycosyltransferases and in particular from among C2GnT, FUT3 and FUT7. In a preferred embodiment, said host cell additionally comprises a gene coding for α (1-3)galactosyltransferase (GT).

In a particularly preferred embodiment, the invention relates to a recombinant, isolated or purified glycopeptide comprising 1 to 40 repetitions of the peptide sequence described in SEQ ID No 14 and glycosylated by one or more enzymes having ose-transferase activity selected in the group consisting of Core 2 β (1-6) N-acetylglucosaminyltransferase (C2GnT), fucosyltransferase FUT3 which has α (1-3) and α (1-4) fucosyltransferase activity, or fucosyltransferase FUT7 which has α (1-3) fucosyltransferase activity, said glycopeptide additionally being glycosylated by α (1-3)galactosyltransferase (GT).

“Isolated glycopeptides” is understood to mean that said compounds are separated from their natural environment.

Thus, said compounds are separated from some or all of the other components of their natural environment.

“Purified glycopeptides” is understood to mean that said compounds are enriched in a mixture by at least approximately a factor of 1, preferably by at least 2, 3, 4 or 5-fold. The term “purified” does not necessarily mean that the compounds are absolutely pure. For instance, said compounds can have a purity of 30%, 40%, 50%, 60%, 70%, 80%, 90% or 100%. Preferably, however, said compounds generate only single bands in polyacrylamide gel electrophoresis.

Any suitable separation and/or purification method by which to obtain them can be used. In a particular embodiment, the glycopeptides are purified from a biological fluid from a human or animal subject. The biological fluid can be selected in the group consisting of serum, urine, pancreatic juices and milk secretions. Preferably, the biological fluid is urine. In a preferred embodiment, the subject is a patient who may be or who is suffering from a pancreatic disease, preferably pancreatic cancer. In another embodiment, the glycopeptides are purified from a cell culture. In an additional embodiment, the glycopeptides are purified from a tissue sample, preferably pancreatic tissue and in particular a pancreatic tissue tumor.

In the spirit of the application, the term “glycotype” refers to glycosylated epitopes. The glycosylated epitopes can involve one or more glycosylated chains. The term “mAb” (“Acm” in French) refers to a monoclonal antibody, the term “pAb” (“Acp” in French) to a polyclonal antibody.

In a particular embodiment, the invention relates to a glycopeptide according to the invention which can give rise to a specific immunological reaction with an antibody according to the invention. Preferably, the glycopeptide can give rise to a specific immunological reaction with one or more antibodies selected from J28 and 16D10. Optionally, the glycopeptide will give rise to a specific immunological reaction with J28. Alternatively, it will give rise to a specific immunological reaction with 16D10.

The host cells classically used in genetic engineering naturally contain enzymes, for example α -3 N-acetylgalactosaminyltransferase which adds an N-acetylgalactosamine generally on a serine or threonine residue of a peptide. For example this is the case for widely used host cells such as C127, COS, CHO or CHO-K1 which contain such enzymes.

In the present application and in that which follows, the term “conventional host cell” denotes a cell normally used for the production of glycopeptides, preferably a human cell or an animal cell such as C127, COS cells, or insect cells such as *Spodoptera frugiperda* ovary cells and particularly a CHO or CHO-K1 cell. Such a suitable conventional host cell can be selected in particular on the basis of its genetic machinery and its abilities to generate various glycosylated epitopes. To make said selection, an enzymatic assay can be carried out to detect the presence in particular of α -3 N-acetylglucosaminyltransferase in the host cell that one wishes to use.

The glycosyltransferases which are used can be of animal or preferably human origin.

In preferred conditions of implementation of the invention, the genetically modified conventional host cell comprises at least one gene coding for said polypeptide and at least one gene coding for Core2 β (1-6) N-acetylglucosaminyltransferase (C2GnT) and fucosyltransferase 3 (FUT3).

In other preferred conditions of implementation of the invention, particularly for applications in pancreatic cancer, the genetically modified conventional host cell comprises at least one gene coding for said polypeptide and at least one gene coding for Core2 β (1-6) N-acetylglucosaminyltransferase (C2GnT), fucosyltransferase 3 (FUT3) and α (1-3)galactosyltransferase (GT).

In still other preferred conditions of implementation of the invention, particularly for applications in type I diabetes, the genetically modified conventional host cell comprises at least one gene coding for said polypeptide and at least one gene coding for glycosyltransferase Core2 β (1-6) N-acetylglucosaminyltransferase (C2GnT) and fucosyltransferase 7 (FUT7).

All or part of the repeated C-terminal peptides can be glycosylated. Preferably, all of the repeated sequences will be glycosylated, generally 16 sequences, more specifically 6 sequences and in particular 2 sequences will be glycosylated.

The glycosylations can be due to one or more N-acetylglucosamine, N-acetylgalactosamine, sialic acid, glucose, galactose, fucose.

In yet other preferred conditions of implementation of the invention, the size of an aforementioned glycopeptide estimated by polyacrylamide gel electrophoresis (SDS-PAGE) is advantageously comprised between 20 and 120 kDa, preferably comprised between 20 and 85 kDa, more particularly comprised between 45 and 83 kDa.

In particular, the glycopeptides named C2-F3-16R, C2-F3-GT-6R and more particularly C2-F3-6R and C2-F7-16R are considered. C2-F3-16R designates a glycopeptide having 16 repetitions glycosylated by C2GnT and FUT3. C2-F3-GT-6R designates a glycopeptide having 6 repetitions glycosylated by C2GnT, FUT3 and GT. C2-F3-6R designates a glycopeptide having 6 repetitions glycosylated by C2GnT and FUT3. C2-F7-16R designates a glycopeptide having 16 repetitions glycosylated by C2GnT and FUT7. Preferably, the glycopeptides according to the invention are selected in the group consisting of C2-F3-6R, C2-F3-GT-6R, C2-F7-6R, C2-F7-GT-6R, C2-F3-F7-6R, and C2-F3-F7-GT-6R.

A cell line producing the glycopeptide C2-F7-16R was deposited with the Collection Nationale de Culture de Microorganismes (CNCM), Institut Pasteur, 28, rue du Dr Roux, 75724 Paris Cédex 15, France on 16 Mar. 2004 under the number I-3189.

The person skilled in the art knows that a peptide can undergo several structural modifications without altering function.

The peptide can contain one or more modifications in its amino acids such as deletions, substitutions, additions or functionalizations such as acylation of amino acids, in so far as said modifications do not affect the immunologic characteristics thereof. In particular, the modification or modifications can be substitutions by a conservative amino acid (that is, having similar physicochemical characteristics). For example, in general, the substitution of a leucine residue by an isoleucine residue does not alter such properties. Generally, the modifications must concern less than 40%, in particular less than 30%, preferably less than 20% and more particularly less than 10% of the amino acids of the peptide. It is important that the modified peptide does not be denatured such as can be done for example by a physical treatment such as heat, so as to preserve the conformational sites thereof so that the antibodies induced by the modified derivatives will be active towards the native structure.

In general, with respect to the modifications, the homology or similarity between the aforementioned modified glycopeptide and the aforementioned native glycopeptide, as well as to the modes of use, or of coupling the immunogenic compound according to the invention to an immunogenic protein like tetanus toxoid, reference can be made in particular to WO-A-86/06 414 or to EP-A-0.220.273 or else to PCT/US.86/00831, equivalent.

The glycopeptides which are the object of the invention can be prepared in particular from pancreatic juices, serum, milk,

urine or amniotic fluid of human or animal origin. In a preferred embodiment, the glycopeptides according to the invention are isolated and/or purified from urine of human or animal origin. Preferably, the urine is of human origin. In a preferred embodiment, the urine is from a patient who may be or who is suffering from a pancreatic pathology. Preferably, the pancreatic pathology is pancreatic cancer. Those skilled in the art are familiar with the techniques that can be used to purify and/or isolate the inventive glycopeptides. In particular, the glycopeptides can be purified by using for example, but without being limited to said methods, centrifugations, ultrafiltrations, concentration steps, liquid chromatography on exclusion or affinity columns, reverse phase chromatography or cation or anion exchange chromatography, electrophoresis or a combination of said techniques. In particular, the antibodies specific of the glycopeptides according to the invention such as J28 and 16D10 can be used for an affinity purification. An example of a purification protocol is described in Experiment 7. Thus, the invention relates to a method of preparation of a glycopeptide according to the invention comprising collecting the urine of a subject and purifying said glycopeptide from the urine.

The invention more particularly relates to glycopeptides according to the invention which are purified and/or isolated from the urine of a patient who may be or who is suffering from a pancreatic pathology, in particular pancreatic cancer. As described hereinbelow, said purified glycopeptides can be used for preventive or therapeutic treatment by immunotherapy. For instance, they can be used to prepare a pharmaceutical and/or vaccine composition.

The inventive glycopeptides can be recombinant and prepared as follows.

A suitable conventional host cell line like CHO-K1 is transfected with a plasmid comprising the complementary DNA of the desired transferase or by plasmids comprising the complementary DNAs of the desired transferases, for example the glycosyltransferase Core2 β (1-6) N-acetylglucosaminyltransferase (C2GnT), fucosyltransferase 3 (FUT3), fucosyltransferase 7 (FUT7) or α (1-3)galactosyltransferase (GT), by using lipofectamine for example.

The resulting cell line is transfected with a plasmid enabling the expression and preferably the secretion of the recombinant peptides, for example a pSecTag plasmid containing the cDNA coding for an inventive glycopeptide, for example the C-terminal domain of BSDL or FAPP. SEQ ID Nos 6, 8, 10 and 12 are examples of sequences coding for a glycopeptide according to the invention.

The recombinant glycopeptide obtained by culturing the cell line containing the desired cDNAs is then advantageously purified for example on a polyacrylamide gel or by liquid and/or affinity chromatography and if desired it is analyzed by SDS-PAGE for identification.

Of course it is possible to transfer the genes required to prepare the inventive glycopeptides into the host cell by any other suitable method.

In preferred conditions of implementation of the aforementioned method, a suitable cell line like CHO-K1 is transfected with a plasmid comprising the complementary DNA of glycosyltransferase Core2 β (1-6) N-acetylglucosaminyltransferase (C2GnT), fucosyltransferase 3 (FUT3), and a glycopeptide according to the invention, for example the C-terminal polypeptide domain of FAPP or in particular of BSDL, preferably in that order.

In an advantageous manner, the CHO cell line and in particular the CHO-K1 cell line is used as host cell.

In an advantageous manner, the plasmids pSecTag, pcDNA-3 or pBK-CMV are used as plasmid.

More generally, the invention also has as object a method of preparation of a glycopeptide such as described hereinabove, characterized in that one transfers into a conventional host cell comprising an enzymatic machinery necessary for priming a glycosylation, a gene coding for the polypeptide according to the invention and at least one gene coding for one or more enzymes selected from glycosyltransferases, in conditions enabling the expression and optionally the secretion of the desired glycopeptide, and one isolates the expected glycopeptide.

The glycopeptides according to the invention, and in particular the natural or derivative glycopeptides of FAPP or BSDL, can be prepared from proteins isolated from biological fluids or tissues of any human or animal origin whatsoever, particularly sera, urine, pancreatic juices, milk secretions, tissue or cell homogenates, and the like, by chromatographic methods such as described by Lombardo et al. 1978, *Biochim. Biophys. Acta*, 527: 142-149, or by Abouakil et al. 1988, *Biochim. Biophys. Acta*, 961: 299-308, or by Wang & Johnson, 1983, *Anal. Biochem.*, 133: 457-461 or else by Blackberg & Hernell, 1981, *Eur J. Biochem.* 116: 221-225. The homogeneous proteins are then hydrolyzed either by enzymatic methods involving any enzyme particularly proteolytic, or by chemical methods.

The natural glycopeptides can be isolated by chromatographic methods (gel filtration, affinity and immunoaffinity, ion exchange, etc.) as described in particular in Mas et al. 1997, *Glycobiology*, 7: 745-752, or Wang et al. 1995, *Biochemistry*, 34: 10639-10644, or by any other method.

Said glycopeptides can also be purified, isolated either directly from biological fluids or tissues as indicated earlier or subsequent to the aforementioned enzymatic or chemical treatments of said biological fluids or tissues.

The glycosylation of said natural glycopeptides of BSDL or FAPP can be modified by any chemical or enzymatic routes (use of native, soluble recombinant, membranar in particular, of glycosidases or glycosyltransferases) in order to obtain the glycan structures hereinbelow used in particular for the diagnosis of the pathologies indicated hereinabove and hereinbelow or for vaccination against said pathologies, or for obtaining or producing antibodies which are the object of the invention.

In addition, it is possible to modify the peptide sequence of the natural or recombinant glycopeptides of the invention.

The natural and particularly recombinant, isolated or purified glycopeptides which are object of the invention have very interesting properties. In particular, they exhibit remarkable immunogenic properties, specific of pancreatic pathologies.

Said properties are illustrated hereinbelow in the experimental section. They justify the use of the aforementioned glycopeptides as medicament.

This is why the invention also has as object the aforementioned glycopeptides, for the use thereof in a method of therapeutic treatment of the human or animal body, that is to say, as medicament.

The medicaments according to the invention can be used for example in immunotherapy (immunization, vaccination) preventive and/or curative treatment of pancreatic cancer, breast cancer, for cell-based immunotherapy: autologous vaccination by activation of the patient's immune system (dendritic or other cells, etc.), for the diagnosis of certain pathologies such as type I diabetes by detection of circulating antibodies directed against said structures in patients afflicted with a diabetic pathology and more generally with any other pathology requiring the use of one or more of said glycopeptides.

They can also be used in other biological systems such as inflammation, the formation of metastases and the inhibition of invasion by pathogens.

Thus, the invention relates to a pharmaceutical or vaccine composition comprising a glycopeptide according to the invention. In particular, the invention relates to a pharmaceutical or vaccine composition comprising a glycopeptide comprising 1 to 40 repetitions of the peptide sequence described in SEQ ID No 14 and glycosylated by one or more enzymes having ose-transferase activity selected in the group consisting of Core 2 β (1-6) N-acetylglucosaminyltransferase (C2GnT), fucosyltransferase FUT3 which has α (1-3) and α (1-4) fucosyltransferase activity, or fucosyltransferase FUT7 which has α (1-3) fucosyltransferase activity. Optionally, said glycopeptide is glycosylated by the enzymes C2GnT and FUT3. Optionally, said glycopeptide is glycosylated by the enzymes C2GnT and FUT7. Optionally, said glycopeptide is glycosylated by the enzymes C2GnT, FUT3 and FUT7. Preferably, said glycopeptide is further glycosylated by α (1-3)galactosyltransferase (GT). In a preferred embodiment, said glycopeptide comprises between 1 and 15 of said repetitions. Said glycopeptide can be recombinant or purified from a biological fluid. Preferably, said biological fluid is urine, in particular urine from a subject suffering from a pancreatic pathology, particularly pancreatic cancer. In a particular embodiment, said glycopeptide is loaded on antigen-presenting cells, preferably dendritic cells.

The invention also relates to the use of a glycopeptide or a pharmaceutical or vaccine composition according to the invention as medicament. In particular, the invention relates to the use of a glycopeptide or a pharmaceutical or vaccine composition according to the invention for preparing a medicament intended for the preventive or curative treatment of a disease selected in the group consisting of a cancer, an inflammatory disorder, a vascular pathology or an infection by a pathogen. Preferably, the disease is a breast or pancreas cancer. In a preferred embodiment, the invention relates to the use of a glycopeptide or a pharmaceutical or vaccine composition according to the invention for preparing a medicament intended for the treatment or prevention of a pancreatic pathology, in particular pancreatic cancer. In a preferred embodiment, the glycopeptide is obtained by purification from the urine of a subject suffering from a pancreatic pathology, preferably pancreatic cancer.

Within an adjuvant formulation, the immunogen, in this case a glycopeptide according to the invention, can be included in particular in a water-in-oil emulsion, by using ISA 51 for example.

A vaccine preparation containing the immunogenic glycopeptide can be administered in a suitable pharmaceutical formulation to induce an immune response of the systemic type by the intramuscular (im), subcutaneous (sc), intradermal (id) route or of the mucosal type by the intranasal, oral, vaginal or rectal route.

A vaccine preparation containing the immunogenic glycopeptide can also contain other immunogens.

A systemic pharmaceutical preparation, administered by the sc, im, id route, can be a water-in-oil emulsion containing the immunogenic glycopeptide, or an immunogen-embedded calcium phosphate suspension, or aluminium hydroxide adsorbing the immunogen.

The vaccine preparations can be formulated for the intranasal route in the form of a gel with carbopol as excipient, nose drops or spray and for the oral route in the form of gastroresistant capsules, gastroresistant granules or sugar coated tablets.

The usual dose, which varies according to the subject and the causal pathology, can be for example from 1 to 100 mg of a glycopeptide according to the invention, in particular of the glycopeptide described in example 4 hereinbelow, administered by the systemic route in humans, every two weeks for 10 to 20 weeks.

The invention also has as object pharmaceutical compositions which contain at least one aforementioned glycopeptide, as active ingredient.

In said compositions, the active ingredient is advantageously present at physiologically effective doses; in particular the aforementioned compositions contain an effective vaccine dose of at least one aforementioned active ingredient.

As medicaments, the glycopeptides described hereinabove can be incorporated in pharmaceutical compositions intended for the digestive or parenteral route.

For example, said pharmaceutical compositions can be solids or liquids and be supplied as pharmaceutical formulations commonly used in human medicine, such as for example simple or sugar-coated tablets, capsules, granules, caramels, suppositories, particularly injectable preparations; they are prepared by the usual methods. The active ingredient(s) can be formulated therein with excipients usually employed in said pharmaceutical compositions, such as talc, gum arabic, lactose, starch, magnesium stearate, cocoa butter, vehicles which are aqueous or not, animal or vegetable fats, paraffin derivatives, glycols, various wetting agents, dispersives or emulsifiers, preservatives.

The invention also has as object a method of preparation of a composition such as described hereinabove, characterized in that the active ingredient(s) are mixed with acceptable excipients, particularly pharmaceutically acceptable, according to established methods.

The invention further has as object the use of a glycopeptide such as described hereinabove, for preparing a medicament intended for the curative or preventive treatment of pancreatic cancer or breast cancer or other pathologies such as a vascular pathology or angiogenesis.

The invention further relates to a method of preventive or curative treatment of a disease selected in the group consisting of a cancer, an inflammatory disorder, a vascular pathology or an infection by a pathogen in a subject, comprising administering an effective dose of a glycopeptide according to the invention.

In a preferred embodiment of cancer treatment, in particular prostate cancer, the glycopeptide can be loaded on an allogeneic antigen-presenting cell. Preferably, the antigen-presenting cell is a dendritic cell. In a preferred manner, the antigen-presenting cell is autologous, that is, it has previously been removed from the recipient subject or it is derived from stem cells originating from the recipient subject. In an alternative, the antigen-presenting cell originates from a sample taken from an allogeneic subject, that is to say, compatible with the recipient. Thus, in this embodiment, the pharmaceutical or vaccine composition according to the invention comprises antigen-presenting cells loaded or "pulsed" with a glycopeptide according to the invention. The invention therefore relates to the use of a pharmaceutical or vaccine composition comprising antigen-presenting cells loaded or "pulsed" with a glycopeptide according to the invention as medicament, in particular as medicament for the treatment or prevention of breast or pancreatic cancer, preferably pancreatic cancer. In a preferred embodiment, the glycopeptide is obtained by purification from the urine of a subject suffering from a pancreatic pathology, preferably pancreatic cancer. In a most desirable embodiment, the glycopeptide is obtained by purification from the urine of a treated subject. Preferably, the subject is

human. The invention also relates to a method of curative or preventive treatment of a pancreatic pathology, preferably pancreatic cancer, or breast cancer, comprising: providing the glycopeptides according to the invention; and administering said glycopeptides, optionally loaded on antigen-presenting cells, said administration allowing to treat, attenuate or prevent a pancreatic pathology, preferably pancreatic cancer, or breast cancer. In a preferred embodiment, the treated disease is pancreatic cancer. Said glycopeptides can be recombinant or isolated from a biological sample, preferably a biological fluid. In a preferred embodiment, said glycopeptides are isolated and/or purified from the urine of a subject suffering from a pancreatic pathology, preferably pancreatic cancer, or breast cancer. Preferably, said glycopeptides are isolated and/or purified from the urine of a treated subject. Thus, the step of providing the glycopeptides can comprise the collection of the biological fluid, preferably urine, and the purification of said glycopeptides.

The properties of the glycopeptides described hereinabove also justify their use in diagnosis, particularly in immunoenzymatic methods.

This is why the invention also has as object a composition for the diagnosis in vitro of the presence of an antibody directed against a glycopeptide described hereinabove, in a human biological sample, characterized in that it contains at least one glycopeptide such as described hereinabove.

The invention also has as object a method for the detection in vitro of the presence of induced antibodies in a human subject suffering from an aforementioned pathology, in a human biological sample such as serum and, more particularly, for the diagnosis in vitro of pancreatic cancer or a diabetic pathology and originating from the person who is the object of the diagnosis, characterized in that said biological sample is contacted with an antigen recognized by an induced antibody in the patient such as defined hereinabove or hereinbelow, in conditions allowing the formation of an immunological complex between said antigen and said antibody and in that the immunological complex possibly formed between said antibody and antigen is detected.

The invention also has as object the necessary materials or kit for the detection of induced antibodies in a human subject suffering from an aforementioned pathology, in a biological sample, particularly a possible carrier of said antibodies, characterized in that it comprises:

at least one glycopeptide such as described hereinabove means for detecting the immunological complex resulting from the immunological reaction between the antigen and said biological sample.

The glycopeptides, particularly recombinant, isolated or purified which are the object of the invention can also be used in the diagnosis of certain pathologies such as type I diabetes by detection of circulating antibodies directed against these structures in patients with such diabetic pathology.

In particular, the glycopeptides named C2-F7-16R are used for the diagnosis of type I diabetes.

As the glycopeptides, particularly recombinant, isolated or purified which are the object of the invention possess very interesting antigenic properties, they also enable the production of antibodies particularly monoclonal.

Hence the application also has as object an antibody particularly monoclonal giving rise to a specific immunological reaction with a glycopeptide such as described hereinabove, with the exception of antibody J28 and any other antibody meeting the hereinabove definition and which can be described in the literature. The monoclonal antibodies can be from any class, in particular IgM, IgD, IgE or IgA. Preferably, the monoclonal antibodies are IgG. An antibody according to

the invention can be modified, regardless of the initial class thereof, to an antibody from another immunoglobulin class, for example by known molecular biology methods (Lewis et al., 1992, *Hum. Antibodies Hybridomas*, 3: 146-52; Lewis et al., 1993, *J. Immunol.*, 151: 2829-38; Hall et al., 1994, *Cancer Res.*, 54: 5178-85; Shepherd & Dean, *Monoclonal Antibodies*, Oxford University Press, 2000, 479 pages). These same monoclonal antibodies can be humanized by genetic recombination or by any other method in order to conserve their recognition site for said recombinant or natural glycopeptides so as to use them in the conditions indicated earlier.

In particular an antibody obtained by immunization of a mammal with the aid of a glycopeptide such as described hereinabove, is chosen.

More particularly, the invention relates to the antibodies described in the examples and in particular the 16D10 antibodies of the IgM type which can be produced by the hybridoma deposited with the Collection Nationale de Culture de Microorganismes (CNCM), Institut Pasteur, 28, rue du Dr Roux, 75724 Paris Cédex 15, France on 16 Mar. 2004 under the number I-3188 and 8H8 of the IgG type.

Preferably, the invention relates to a monoclonal antibody selected in the group consisting of the monoclonal antibody 16D10, a fragment or derivative thereof, and an antibody which essentially binds to the same epitope as monoclonal antibody 16D10. Optionally, said antibody is humanized, chimeric or human. Preferably, the antibody is of the IgG type. Optionally, the antibody is a single chain antibody.

An antibody fragment or derivative is understood to mean an entity conserving substantially the same specificity as the antibody in question. A fragment thereof can be a Fab, Fab', F(ab)2 and sFv fragment (Blazar et al., 1997, *Journal of Immunology* 159: 5821-5833 and Bird et al., 1988, *Science* 242: 423-426). Fv, Fab, or F(ab)2 fragments according to the invention can be obtained by classical enzymatic digestion methods. A derivative thereof can be a chimeric, humanized or single chain scFv antibody. Said derivatives can be obtained by classical genetic engineering methods. Polynucleotides coding for variable regions of antibody 16D10 can be obtained for example by cloning said variable regions from a 16D10-antibody-producing hybridoma cDNA library. They can also be totally or partially prepared by nucleic acid synthesis, based on the nucleotide sequences of said variable regions. For example polynucleotides coding for the CDRs of 16D10 can be synthesized, and incorporated into suitable regions of an other antibody, in particular an antibody of human origin and/or of the IgG type, by known techniques of CDR grafting, such as those described by Routledge et al. ("Reshaping antibodies for therapy", in *Protein Engineering of Antibody Molecules for Prophylactic and Therapeutic Applications in Man*, 13-44, Academic Titles, Nottingham, England (1993)) or by Roguska et al., 1996, *Protein Engineering*, 9 (10): 895-904).

The invention also has as object any nucleic acid molecule coding for a protein comprising the CDRs of antibody 16D10, and any recombinant vector, in particular any expression vector, comprising said nucleic acid molecule, any host cell comprising said nucleic acid or said vector.

The invention also has as object a method of preparation of an anti-glycopeptide antibody such as described hereinabove, characterized in that a mammal is immunized with the aid of said glycopeptide or with the aid of a mixture of FAPP and BSDL isolated from any human or animal biological tissue or fluid and in particular normal and pathological human pancreatic juices and the expected antibodies recognizing the C-terminal domain of said proteins are then recovered and purified if so desired.

The application equally has as object a method of preparation of an anti-glycopeptide antibody such as described hereinabove characterized in that one clones B lymphocytes originating from a subject immunized with the aid of a natural or recombinant glycopeptide such as described hereinabove or with the aid of a mixture of FAPP and BSDL such as described hereinabove and transformed by Epstein-Barr virus (EBV), then recovers the expected antibodies secreted by said transformed B lymphocytes.

In particular, the monoclonal antibodies object of the invention can give rise to a specific immunological reaction with a glycopeptide such as described hereinabove.

In a particular embodiment of the invention, the antibody essentially binds to the same epitope as the monoclonal antibody 16D10 or J28 (respectively produced by hybridoma 16D10 or J28). Preferably, said antibody is a monoclonal antibody. Preferably, the invention relates to the monoclonal antibody 16D10 (produced by hybridoma 16D10), but it shall be understood that the invention also relates to monoclonal antibodies which can specifically compete with antibody 16D10. Likewise, in a particular embodiment, the monoclonal antibodies according to the invention can specifically compete with antibody J28.

The term "essentially binds to the same epitope or determinant as" an antibody of interest means that the antibody "competes" with said antibody of interest. The term "essentially binds to the same epitope or determinant as" an antibody of interest does not mean that the two antibodies have exactly the same epitope. However, in a particular embodiment, the two antibodies can have the same epitope. The term "essentially binds to the same epitope or determinant as" monoclonal antibody 16D10 means that the antibody "competes" with antibody 16D10. Generally, an antibody which "essentially binds to the same epitope or determinant as" an antibody of interest (e.g., antibody 16D10) means that the antibody "competes" with said antibody of interest for a glycopeptide according to the invention, in particular one or more proteins selected from BSDL or FAPP or a fragment thereof, preferably a C-terminal part of BSDL or FAPP as described herein, even more preferably a part comprising one or more repetitions of 11 amino acids of BSDL or FAPP, or a part or fragment of same. In other examples, an antibody essentially binds to the same epitope or determinant of BSDL or FAPP when the antibody "competes" with the antibody of interest for binding to BSDL or FAPP.

The term "essentially binds to the same epitope or determinant as" an antibody of interest means that the antibody "competes" with said antibody of interest for any BSDL and/or FAPP molecule to which said antibody of interest specifically binds. The term "essentially binds to the same epitope or determinant as" monoclonal antibody 16D10 means that the antibody "competes" with said monoclonal antibody 16D10 for any BSDL and/or FAPP molecule to which said monoclonal antibody 16D10 specifically binds. For example, an antibody which essentially binds to the same epitope or determinant as monoclonal antibodies 16D10 or J28 "competes" with said antibodies 16D10 or J28 for binding to BSDL or FAPP, respectively.

The identification of one or more antibodies which essentially bind(s) to the same epitope as the monoclonal antibodies described herein can be carried out by using any immunological screening method in which a competition between antibodies can be evaluated. Many assays are routinely carried out and are well known to those skilled in the art (see for example U.S. Pat. No. 5,660,827 granted 26 Aug. 1997, which is specifically incorporated herein by reference). The determination of the epitope to which an antibody binds is not

necessary to identify an antibody which binds, or which essentially binds, to the same epitope as the monoclonal antibody described herein.

For example, when the candidate antibodies to be studied have been obtained from different animal sources, or possibly have different Ig isotypes, a simple competition assay can be employed wherein the control antibody (antibody 16D10 for example) and the candidate antibody are mixed (or pre-adsorbed) and contacted with a sample containing either BSDL or FAPP, both known to bind to antibody 16D10. Protocols based on ELISA, radioimmunological assays, western blot analyses, or the use of a BIACORE analysis (as described, for example, in the Examples) are suitable for use in simple competition tests.

In particular embodiments, it is possible to first prepare mixtures of control antibodies (for example, antibody 16D10) with variable amounts of candidate antibodies (for example, approximately 1:10 or approximately 1:100) before contacting with the BSDL or FAPP antigen sample. In another embodiment, the control antibody and the variable amounts of candidate antibodies can simply be mixed at the time of contact with the BSDL or FAPP antigen sample. So long as one can distinguish bound antibodies from free antibodies (for example, by using separation methods or by washing to remove unbound antibodies) and distinguish antibody 16D10 from the candidate antibodies (for example, by using secondary antibodies specific of the species or the isotype or by labelling antibody 16D10 with a detectable tag), it will be possible to determine if the candidate antibodies reduce the binding of antibody 16D10 to the BSDL or FAPP antigen, thereby indicating whether the candidate antibody essentially recognizes the same epitope as antibody 16D10. The binding of the control antibody in the presence of a completely irrelevant antibody can serve as an upper control value. The lower control value can be obtained by incubating the labelled control antibody (16D10) with unlabelled antibodies recognizing exactly the same epitope (16D10), in this way inducing a competition which reduces the binding of the labelled antibody. In one test, a significant reduction in the reactivity of a labelled antibody in the presence of a candidate antibody indicates that the candidate antibody essentially recognizes the same epitope, that is to say, it cross-reacts with the labelled antibody (16D10). Any candidate antibody which reduces the binding of antibody 16D10 to each of the antigens BSDL or FAPP by at least approximately 20%, 30%, 40%, 50%, preferably by at least about 60%, or even more preferably by at least about 70% (for example, 65-100%), at a ratio of 16D10 antibody to candidate antibody of between about 1:10 and about 1:100, is considered to be an antibody which essentially binds to the same epitope or determinant as antibody 16D10. Preferably, said candidate antibodies reduce the binding of antibody 16D10 to the BSDL or FAPP antigen by at least about 90% (for example, about 95%). Preferably, antibody 16D10 also reduces the binding of the candidate antibody to the BSDL or FAPP antigen when the binding is evaluated in the same way, although the degree of reduction in binding may be different.

The competition can be evaluated by a flow cytometry assay for example. In such assay, cells carrying a BSDL or FAPP antigen, for example cells transfected with BSDL or FAPP, are first incubated with antibody 16D10, for example, then with the candidate antibody labelled with a fluorochrome or with biotin. The antibody is considered to compete with antibody 16D10 if the binding observed after preincubation with a saturating amount of 16D10 is about 30%, preferably about 40% about 50%, about 80%, or more (for example, about 90%) of the binding observed (as measured

by fluorescence) with the antibody without preincubation with 16D10. Alternatively, an antibody is considered to compete with antibody 16D10 when the binding observed with labelled 16D10 antibody (labelled with a fluorochrome or biotin) in cells preincubated with a saturating amount of the candidate antibody is about 80%, preferably about 50%, about 40%, about 30%, or less (for example, about 20%) of the binding observed without preincubation with the antibody.

In an advantageous manner, a simple competition assay in which a candidate antibody is preadsorbed and applied at a saturating concentration on a surface on which a BSDL or FAPP antigen has been immobilized can also be used. The surface in the simple competition assay is preferably a BIACORE chip (or another support compatible with surface plasmon resonance analysis). The control antibody (for example, 16D10) is then contacted with a surface at a saturating concentration of BSDL or FAPP and binding of the control antibody to the surface carrying BSDL or FAPP is measured. Said binding of the control antibody is compared with the binding of the control antibody to the surface carrying BSDL or FAPP in the absence of the candidate antibody. In an assay test, a significant reduction in binding of the control antibody to the surface carrying BSDL or FAPP in the presence of the candidate antibody indicates that the candidate antibody essentially recognizes the same epitope as the control antibody in such a way that the candidate antibody cross-reacts with the control antibody. Any candidate antibody which reduces the binding of the control antibody (such as antibody 16D10) to the BSDL or FAPP antigen by at least about 30% or preferably about 40% can be considered an antibody which essentially binds to the same epitope as the control antibody (for example, 16D10). Preferably, said candidate antibody reduces the binding of the control antibody (for example, 16D10) to the BSDL or FAPP antigen by at least about 50% (for example, at least about 60%, at least about 70%, or more). It shall be understood that the order between the control and candidate antibodies can be reversed, that is to say, that the control antibody can be the first to bind to the surface and that the candidate antibody is subsequently contacted with the surface in a competition test. Preferably, the antibody displaying the highest affinity for BSDL or FAPP is bound to the surface carrying BSDL or FAPP first, since it is hoped that the reduction in binding observed with the second antibody (presuming that the antibodies cross-react) will be of greater amplitude. Other examples of such tests are described in the examples and in Saunal and Regenmortel, (1995, J. Immunol. Methods 183: 33-41, the content thereof being incorporated herein by reference).

Said properties are illustrated in the experimental section which follows. They justify the use of the monoclonal antibodies described hereinabove and in particular of the monoclonal antibodies 8H8, 16D10, 14H10 and other monoclonal antibodies directed against the C-terminal part of BSDL or FAPP, for the diagnosis, prognosis and/or prediction of several pancreatic pathologies including pancreatic cancer, pancreatitis and type I diabetes, but also breast cancer or cardiovascular diseases. The diagnostic information can be obtained from serum and/or urine assays.

The antibody 16D10 has particularly interesting properties. In fact, it is specific of pancreatic cancer. It recognizes neither normal tissues nor other types of tumor tissues. The epitope recognized by antibody 16D10 is different from that recognized by antibody J28 because these two antibodies do not compete. These results are described in example 4.

Thus, the invention relates to a method enabling the detection, preferably in vitro, of a pancreatic pathology, in particu-

lar pancreatic cancer, comprising contacting a biological sample with an antibody according to the invention and detecting the formation of immunological complexes resulting from the immunological reaction between said antibody and said biological sample. Preferably, the antibody is the antibody 16D10, a fragment or derivative thereof, or an antibody which essentially binds to the same epitope or determinant as the former. Preferably, the biological sample is a sample of pancreatic tissue (biopsy) or a biological fluid. The biological sample can be a tissue slice (immunohistochemistry) or cells contained in the sample or derived by culturing the sample (immunocytochemistry). The biological fluid can be selected in the group consisting of serum, urine, pancreatic juices and milk. Preferably, the biological fluid is urine. The complex can be detected directly by labelling the antibody according to the invention or indirectly by adding a molecule which reveals the presence of the antibody according to the invention (secondary antibody, streptavidin/biotin tag, etc.). For example, labelling can be accomplished by coupling the antibody with radioactive or fluorescent tags. These methods are well known to those skilled in the art.

The invention also relates to the use of an antibody according to the invention for preparing a diagnostic composition that can be used for detecting a pancreatic pathology in vivo or in vitro. Preferably, the antibody is the antibody 16D10, a fragment or derivative thereof, or an antibody which essentially binds to the same epitope or determinant as the former. Preferably, the pancreatic pathology is pancreatic cancer.

In a preferred embodiment, the invention relates to a method enabling the detection of a pancreatic pathology in a subject, in particular pancreatic cancer, comprising recovering the urine of the subject, contacting said urine with an antibody according to the invention, and detecting the formation of immunological complexes resulting from the immunological reaction between said antibody and said urine. Preferably, the antibody is the antibody 16D10, a fragment or derivative thereof, or an antibody which essentially binds to the same epitope or determinant as the former. The antibody can also be the antibody J28, a fragment or derivative thereof, or an antibody which essentially binds to the same epitope or determinant as the former. Preferably, the pancreatic pathology is pancreatic cancer. The complex can be detected directly by labelling the antibody according to the invention or indirectly by adding a molecule which reveals the presence of the antibody according to the invention (secondary antibody, streptavidin/biotin tag, etc.). These methods are well known to those skilled in the art. Optionally, the method can comprise intermediate steps of treating the urine sample before incubation with said antibody. For example, said steps can comprise concentration of the urine, steps of glycopeptide enrichment or purification, and the like.

The invention also relates to the use of an antibody according to the invention for the diagnosis of a pancreatic pathology, in particular pancreatic cancer in a subject. Preferably, the antibody is antibody 16D10, a fragment or derivative thereof, or an antibody which essentially binds to the same epitope or determinant as the former. The antibody can also be antibody J28 a fragment or derivative thereof, or an antibody which essentially binds to the same epitope or determinant as the former.

The invention relates to a diagnostic kit for a pancreatic pathology, in particular pancreatic cancer, comprising an antibody according to the invention. Said kit can additionally comprise means by which to detect the immunological complex resulting from the immunological reaction between the biological sample and said antibody, in particular reagents enabling the detection of said antibody. In a particular

embodiment, the kit comprises the antibody 16D10, a fragment or derivative thereof, or an antibody which essentially binds to the same epitope or determinant as the former.

These properties justify the use of the monoclonal antibodies described hereinabove for developing new therapeutic protocols for cancer of the exocrine pancreas such as targeting tumor cells with said antibodies coupled with drugs (chemotherapy) or radioactive elements (radiotherapy and radio-diagnosis) or else with genes modifying the outcome or behavior of the neoplastic cells (gene therapy).

The antibodies coupled with radioactive elements can be used in radiolocalization (immunoscintigraphy) of primary tumors and metastases (secondary tumors).

Said diagnostic information can be acquired in serum and/or urine assays.

In particular, the invention relates to antibodies according to the invention, in particular monoclonal, coupled with an antitumoral active substance, which can be used in the treatment of pancreatic cancer to target and destroy pancreatic tumor cells.

Similarly, radioactive elements (or others) can be coupled with said antibodies and allow a precise localization of primary tumors and metastases. Said antibodies can also make it possible to clearly and accurately differentiate normal tissue from tumor tissue.

These same properties justify the use of said monoclonal antibodies alone, combined or coupled with a diagnostic or therapeutic molecule to develop new passive immunotherapy protocols for the aforementioned pathologies.

Thus the invention also has as object a composition for the in vitro diagnosis of a pathology such as described hereinabove, in a human biological sample, characterized in that it contains at least one monoclonal antibody such as described hereinabove, in particular coupled with at least one radioactive element. Depending on the nature of the radioactive element, said composition can be used curatively to irradiate a region of the pancreas.

As a diagnostic composition, the aforementioned antibodies can be mixed with acceptable excipients.

The invention also has as object a kit for detecting the presence of a glycopeptide such as described hereinabove appearing in a human subject afflicted with a diabetic or neoplastic pathology in a human biological sample originating from the individual to be diagnosed, characterized in that it comprises:

at least one antibody such as described hereinabove, means for detecting the immunological complex resulting from the immunological reaction between the antibody and the biological sample.

The preferred conditions of employing the glycopeptides and antibodies described hereinabove also apply to the other objects of the invention mentioned earlier, in particular to the diagnostic methods.

The invention also relates to a pharmaceutical composition comprising an antibody according to the invention. Preferably, the antibody is the antibody 16D10, a fragment or derivative thereof, or an antibody which essentially binds to the same epitope or determinant as the former. The antibody can also be the antibody J28, a fragment or derivative thereof, or an antibody which essentially binds to the same epitope or determinant as the former. The composition can also comprise a pharmaceutically acceptable support. In a particular embodiment, said antibody is coupled with an antitumoral substance.

The invention further relates to the use of an antibody according to the invention or of a pharmaceutical composition comprising such antibody as medicament. In particular,

the invention relates to the use of an antibody according to the invention or of a pharmaceutical composition comprising such antibody for preparing a medicament intended for the preventive or curative treatment of a pancreatic pathology, preferably pancreatic cancer, or breast cancer. Preferably, the antibody is the antibody 16D10, a fragment or derivative thereof, or an antibody which essentially binds to the same epitope or determinant as the former. The antibody can also be the antibody J28, a fragment or derivative thereof, or an antibody which essentially binds to the same epitope or determinant as the former. In a particular embodiment, said antibody is coupled with an antitumoral substance.

The invention relates to a method of preventive or curative treatment of a subject suffering from a pancreatic pathology, in particular pancreatic cancer, or breast cancer, comprising administering to said subject an effective amount of an antibody according to the invention, said administration resulting in a decrease or disappearance of the pancreatic pathology, in particular pancreatic cancer, or breast cancer, in the subject. Preferably, the antibody is the antibody 16D10, a fragment or derivative thereof, or an antibody which essentially binds to the same epitope or determinant as the former. The antibody can also be the antibody J28, a fragment or derivative thereof, or an antibody which essentially binds to the same epitope or determinant as the former. In a particular embodiment, said antibody is coupled with an antitumoral substance.

In another protocol the usual dose, which varies according to the subject being treated and the causal disease, can be for example from 1 to 10 mg of the monoclonal antibody described in example 18 hereinbelow per kilogram of body weight administered systemically in humans, once a week for two weeks.

The invention is also directed at providing new products (glycopeptides, antibodies, etc.) as well as other better, more effective or purer products than those described in the prior art.

The invention is illustrated in the following examples.

LEGENDS OF FIGURES

FIG. 1. Study comparing the reactivity of mAbJ28 and mAb16D10 in two pancreatic tumor tissues (PDAC=ADK1 and PDAC4=ADK4).

FIG. 2. Competition test between antibodies mAbJ28 and mAb16D10 for glycopeptide J28.

FIG. 3. Competition test between antibodies mAbJ28 and mAb16D10 in SOJ-6 cells.

FIG. 4. Comparative immunodetection analysis in urine from healthy subjects (samples 29 to 36) and subjects with pancreatic cancer (samples 1 to 5) with the aid of polyclonal antibody pAbL64, monoclonal antibody mAb8H8 and monoclonal antibody mAbJ28. Urine proteins were separated on a 10% polyacrylamide gel then electrotransferred to a nitrocellulose membrane. The membranes were incubated with pAbL64, mAb8H8 or mAbJ28. The left lanes (Std) correspond to the molecular weight markers.

EXAMPLES

Preparation of Recombinant Glycopeptides

Recombinant glycopeptides of BSDL (SEQ ID No 9) and FAPP (SEQ ID No 13), were expressed in different strains derived from the well-known cell line CHO-K1, a Chinese hamster ovarian tumor cell line with fibroblast morphology, which can be obtained in particular from the *American Type*

Culture Collection ATCC under the number CRL61 (Buck et al., 1958, *J. Exp. Med.*, 108: 945-955).

The CHO-K1 cell line was chosen because it does not show detectable C2GnT glycosyltransferase, $\alpha(1-2)$ fucosyltransferase, $\alpha(1-3)$ fucosyltransferase, $\alpha(1-4)$ fucosyltransferase or $\alpha(1-3)$ galactosyltransferase activity in standard assay conditions.

Preparation of Plasmids (Cloned cDNA)

The cloned cDNAs enabling the expression of the different glycopeptide components were obtained by reverse transcription, carried out with a polydT polynucleotide probe (18 bases) hybridizing with the polyA tail of messenger RNA extracted from tissue samples of healthy human pancreas or the human pancreatic tumor cell line such as for example the cell line SOJ-6 [Fujii et al., 1990, *Hum. Cell.*, 3: 31-36] originating from a human pancreatic adenocarcinoma, according to the method described by Chirgwin et al. [1979, *Biochemistry*, 18: 5294-5299].

Preparation 1: Plasmid pSec-FAPP

The amplification of cDNA coding for fetoacinar pancreatic protein (FAPP) (SEQ ID No 10) was reverse transcribed from 5 μ g of total RNA extracted from the human pancreatic tumor line SOJ-6 mentioned earlier (45 minutes at 55° C., reverse transcription kit, Sigma, St Louis, Mo., USA) and followed by a PCR using the nucleotide primers C-ter/FAPP-BSDL and N-ter/FAPP, defined from the cDNA sequence coding for human BSDL (SEQ ID No 6) in the conditions described hereinbelow. The polymerase chain reaction was carried out with an "Advantage-GC cDNA PCR" kit (Clontech) enabling amplification of nucleotide sequences particularly rich in guanine and cytosine. The following program was employed: one cycle of 2 minutes at 94° C. followed by 35 cycles of 1 minute at 94° C. (denaturation), 1 minute at 52° C. (primer annealing), and 4 minutes at 68° C. (extension), followed by one cycle of 10 minutes at 68° C., in a Robocycler Gradient 96 thermocycler (Stratagene).

The primers had the following sequences:

N-ter/FAPP (SEQ ID No 1):

5'-TTCGTAagcttGCGAAGCTGGGCGCCGTGTACAGAA-3';

C-ter/FAPP-BSDL (SEQ ID No 2):

5'-TTTCGTgaattcACGCTAAAACCTAATGACTGCAGCATCTG-3'.

The C-ter/FAPP-BSDL primer (SEQ ID No 2) which was used comprises a termination codon so as to eliminate translation of the c-myc epitope as well as that of the 6 \times -histidine tag carried by the commercial vector.

The cDNA (nucleotides 1 to 2169) amplified in this manner (SEQ ID No 12) does not comprise a signal peptide.

The resulting cDNA was then cloned into plasmid pSec-Tag (Invitrogen, Leek, the Netherlands).

Preparation 2: Plasmid pSec-16R

The cDNA coding for the C-terminal part of BSDL (from nucleotide 1089: Phe364 to nucleotide 2169: stop codon; SEQ ID Nos 8 and 9) from RNA extracted from normal pancreas was amplified by PCR using the primers C-ter/FAPP-BSDL (primer SEQ ID No 2, see Preparation 1) and N-ter-Ct.

The N-ter-Ct primer had the following sequence:

N-ter-Ct (SEQ ID No 3):
5'-CGTCTAaagcttTTTGATGTCTACACCGAGTCC-3'.

The polymerase chain reaction was carried out in the conditions described earlier (see Preparation 1).

The resulting cDNA was then cloned into plasmid pSec-Tag.

Preparation 3: Plasmid pSec-6R

The cDNA coding for the C-terminal part of FAPP (from nucleotide 1089: Phe364 to nucleotide 1839: stop codon; SEQ ID Nos 12 and 13) present in the plasmid pSec-FAPP was amplified by PCR using nucleotide primers No 2 (C-ter/FAPP-BSDL) and No 3 (N-ter-Ct) described hereinabove.

The polymerase chain reaction was carried out in the conditions described in preparation 1.

The resulting cDNA was then cloned into plasmid pSec-Tag.

Preparation 4: Plasmid pBK- α GT

The cDNA coding for $\alpha(1-3)$ galactosyltransferase (from nucleotide -10 to nucleotide +1122: stop codon) from RNA extracted from mouse heart (Balb/c mice) was amplified by PCR using the primers Cter/ α Gal and Nter/ α Gal defined from the cDNA sequence coding for mouse $\alpha(1-3)$ galactosyltransferase. These primers had the following sequence:

Nter/ α Gal (SEQ ID No 4):
5'-AAAAAgaattcGGAGAAAATAATGAAT-3'.

Cter/ α Gal (SEQ ID No 5):
5'-AAAAagggccACAAAGTCAGACATTAT-3'.

The polymerase chain reaction was carried out with the "Taq PCR Master Mix" kit (Qiagen, Courtaboeuf, France) according to the following program: one cycle of 2 minutes at 94° C. followed by 35 cycles of 1 minute at 94° C. (denaturation), 1 minute at 54° C. (primer annealing), and 2 minutes at 74° C. (extension), then one cycle of 10 minutes at 74° C., in a Gene Amp PCR System 2400 thermocycler (Perkin Elmer).

The resulting cDNA was then cloned into plasmid pBK-CMV (Stratagene, La Jolla, Calif., USA).

Preparation 5: Plasmid pcDNA3-C2 β 6GnT-flag

The cDNA coding for glycosyltransferase Core2 $\beta(1-6)$ N-acetylglucosaminyltransferase described by Panicot et al. in *Glycobiology*, 1999, 9: 935-946 was cloned into the vector pCDNA-3 which at its 3' end comprises a sequence encoding the flag epitope. Expression of the flag epitope allows detection of C2GnT expression in the transformed cells.

The following steps were common to the plasmid preparations:

a) Ligation

The different PCR products were analyzed on a 1% agarose gel, purified using the "GeneClean" kit (Bio 101), then subcloned into the shuttle vector pCR2.1 TOPO and sequenced (Euro Sequences Gènes Service, Paris, France) with the universal M13 and reverse M13 primers. The PCR products were then released from said shuttle vector by the action of the restriction enzymes HindIII and EcoRI and cloned into the eukaryote expression and secretion plasmid pSec-Tag to yield plasmids pSec-FAPP and pSec-16R (16R for C-terminal domain of BSDL) (SEQ ID No 8). The complementary DNA coding for the C-terminal part of FAPP, obtained by PCR, was directly cloned after purification and HindIII/EcoRI digestion into the pSec-Tag plasmid to yield plasmid pSec-6R (6R for C-terminal of domain FAPP) (SEQ ID No 12). Sequencing

was carried out with the universal T7 primer and the BGH reverse primer synthesized for this purpose by Euro Sequences Gènes Service.

The PCR product coding for $\alpha(1-3)$ galactosyltransferase was released with the restriction enzymes EcoRI and ApaI, purified, then directly ligated into the EcoRI/ApaI sites of the vector pBK-CMV to yield plasmid pBK- α GT.

b) Bacterial Transformation

A 5 μ l aliquot of the ligation product was contacted with 50 μ l of competent bacterial cells (*Escherichia coli*, strain TOP10F') according to the protocol described by Hanahan [Hanahan, 1983, J. Mol. Biol., 166: 557-580]. Two microliters of 0.5 M β -mercaptoethanol were added and the sample was kept on ice for 30 minutes, then heat-shocked at 42° C. for 30 seconds, and immediately put back on ice. After 2 minutes, the sample was diluted in 450 μ l of SOC medium (Life Technologies). The bacterial suspension was incubated for 1 hour at 37° C. with shaking. The bacteria were then spread, using glass beads, on Petri dishes containing Luria-Bertagni agar supplemented with 50 μ g/ml ampicillin.

c) Plasmid Purification

The bacterial colonies which appeared after 18 hours of culture on selective agar medium (50 μ g/ml ampicillin) were picked and inoculated in 2 ml of Luria-Bertagni liquid medium containing 50 μ g/ml ampicillin. The cultures were incubated for 8 hours at 37° C. with shaking, then 1.5 ml of the bacterial suspension was centrifuged at 5,000 g for 5 minutes. The bacterial pellet was taken up in 100 μ l of bacterial membrane destabilizing buffer (50 mM Tris-HCl pH 8, 10 mM EDTA, Ribonuclease A 100 μ g/ml). The bacterial suspension was lysed by addition of 200 μ l of alkaline lysis buffer (200 mM NaOH, 1% SDS) and the pH of the preparation was neutralized with 150 μ l of 3 M potassium acetate pH 5.5. The alkaline lysis and neutralization steps require a 5-minute incubation on ice. Cell debris, denatured proteins and chromosomal DNA were eliminated by centrifugation at 10,000 g for 10 minutes at 4° C. The supernatant was then extracted with phenol-chloroform by 1:2 dilution in phenol/chloroform/isoamyl alcohol (25/24/1 V/V/V) stabilized to pH 8 with 100 mM Tris-HCl. The aqueous and organic phases were separated by centrifugation at 10,000 g for 2 minutes at room temperature. The upper aqueous phase was recovered and diluted 1:3 in absolute ethanol (-20° C.). Plasmid DNA was precipitated by centrifugation at 12,000 g for 15 minutes at 4° C., then washed twice with 70% ethanol. Plasmid DNA was then resuspended in 20 μ l of ultra-pure water and stored at -20° C.

Preparation of Cell Clones

Preparation 6: Cell Clone CHO-C2 and General Points Regarding the Obtaining of Intermediate Cell Clones

A CHO-K1 cell line was transfected with plasmid pcDNA3-C2 β 6GnT-flag comprising the cDNA coding for the glycosyltransferase Core2 $\beta(1-6)$ N-acetylglucosaminyltransferase (C2GnT), which at its 3' end comprises a sequence coding for the flag epitope. Expression of the flag epitope allows detection of C2GnT expression in the transformed cells.

To this effect, CHO-K1 cells were grown in HAM-F12 medium. When the cells reached 60 to 80% confluence, the culture medium was removed, the cell surface washed three times with Opti-MEM medium then covered with 200 μ l of transfection solution diluted in Opti-MEM medium without fetal calf serum. The transfection solution contained the aforementioned plasmid and a lipofectamine liposome suspension (Life Technologies).

The cells were kept like this for 16 hours at which time the transfection supernatant was replaced with 2 ml of complete

HAM-F12 medium containing a suitable selective antibiotic (neomycin, zeocin or hygromycin) for a period of at least four weeks. Clones resistant to the action of the antibiotic were isolated and then amplified.

The CHO-K1 cell clone selected in this manner with neomycin (250 µg/ml) was named CHO-C2.

Said clone CHO-C2 was then transfected with plasmid pSec-16R as described below for preparation 7. The clone resulting from transformation of CHO-C2 cells with plasmid pSEC-16R was selected with zeocin (1 mg/ml) and named CHO-C2-16R.

CHO-K1 cells were also cotransfected with plasmid pCDM7-FUT3 coding for fucosyltransferase 3 described by Kukowska-Lallo et al. (1990, *Genes Dev.* 4: 1288-1303), and plasmid pMamNeo according to the method described hereinabove. The CHO-F3 clone resulting from transformation of CHO-K1 cells with plasmid pCDM7-FUT3 was selected on the one hand with neomycin (250 µg/ml) and on the other hand for its high expression of sialyl Lewis x motifs. Said expression was detected by indirect immunofluorescence with the aid of the antibody CSLEX-1 (Becton-Dickinson). In this manner clone CHO-F3 was obtained.

Said clone CHO-F3 was then transfected with plasmid pSec-16R as described below for preparation 7. The clone resulting from transformation of CHO-F3 cells with plasmid pSec-16R and selection with zeocin (1 mg/ml) was named CHO-F3-16R.

CHO-K1 cells were also transfected with plasmid pCDM8-FUT7 coding for fucosyltransferase 7 described by Lowe et al. (1991, *J. Biol. Chem.* 266: 17467-17477) instead of plasmid pCDM7-FUT3, then with plasmid pSec16R. After selection with zeocin and neomycin in the conditions described earlier, the resulting clone was named CHO-F7-16R.

To prepare the intermediate clones expressing the C-terminal peptide of FAPP (6R), the aforementioned method was employed by transfecting the cell lines or cell clones CHO-K1, CHO-C2 and CHO-F3 with plasmid pSec-6R instead of plasmid pSec16R (see method described in preparation 7). However, only the glycosyltransferases C2GnT and fucosyltransferase 3 required to build the J28 glycosylated epitope were used during preparation of the cell lines.

Three intermediate cell clones CHO-6R, CHO-C2-6R and CHO-F3-6R were thus obtained after antibiotic selection in the conditions described hereinabove.

Preparation 7: Cell Clone CHO-16R

CHO-K1 cells were transfected with plasmid pSec-16R, which corresponds to plasmid pSec-Tag (Invitrogen, Leek, the Netherlands) into which the cDNA coding for the C-terminal domain of BSDL was cloned (see preparation 2). This was carried out as for the preparation of clone CHO-C2 in preparation 6.

The pSec-Tag plasmid carries a sequence coding for the V-J2C region of the mouse immunoglobulin κ light chain. Said sequence directs the protein encoded by the cDNA cloned in the plasmid to the secretory pathway of the eukaryotic cell in which it has been transfected.

The clone resulting from transformation of CHO-K1 cells with plasmid pSec-16R was named CHO-16R and was selected for expression of the corresponding (glyco)peptide [(glyco)peptide 16R].

Preparation 8: Cell Clone CHO-C2-F3-16R

Clone CHO-C2 from preparation 6 was transfected with plasmid pCDM7-FUT3 and with plasmid pLSVHg (R & D Systems) conferring hygromycin resistance, to yield cell clone CHO-C2-F3.

The CHO-C2-F3 cell line was transfected with plasmid pSec-16R as in preparation 7. The clone resulting from transformation of CHO-C2-F3 cells with plasmid pSec-16R was named CHO-C2-F3-16R.

CHO-C2-F3-16R expresses sialyl Lewis x motifs and shows reactivity towards antibody pAbL64.

Preparation 9: Cell Clone CHO-C2-F7-16R

This was obtained as in preparation 8, but by using plasmid pCDM8-FUT7 instead of plasmid pCDM7-FUT3.

The clone resulting from transformation of CHO-C2-F7 cells with plasmid pSec-16R was named CHO-C2-F7-16R and the corresponding cell line was deposited with the Collection Nationale de Culture de Microorganismes (CNCM), Institut Pasteur, 28, rue du Dr Roux, 75724 Paris Cédex 15, France on 16 Mar. 2004 under the number I-3189.

The clones CHO-C2-F3-16R and CHO-C2-F7-16R from preparations 8 and 9 express sialyl Lewis x motifs and show reactivity towards antibody pAbL64.

Preparation 10: Cell Clone CHO-C2-F3-6R

Clone CHO-C2-F3 from preparation 8 was transfected with plasmid pSec-6R (see preparation 3) as described for preparations 6 and 7 to yield clone CHO-C2-F3-6R. This clone expresses a peptide bearing among others a glycosylated epitope recognized by monoclonal antibody J28.

Preparation 11: Cell clone CHO-C2-F3-GT-6R

Clone CHO-C2-F3-6R from preparation 10 was transfected with plasmid pBK-αGT (see preparation 4) as described earlier to yield clone CHO-C2-F3-GT-6R bearing the glycosylated epitope recognized by monoclonal antibody J28 as well as the glycosylated epitope αGal recognized by natural circulating antibodies in human blood.

Example 1

Recombinant Glycopeptide Having the Size and the Characteristics of Those Corresponding to the C-Terminal End of BSDL (CHO-16R)

The recombinant glycopeptide was produced by routine culture of cell clone CHO-16R (preparation 7) for 16 hours at 37° C. in a humid, 5% CO₂ atmosphere in Opti-MEM medium (Invitrogen) containing 2 mM L-glutamine, 100 U/ml penicillin, 100 mg/ml streptomycin and 0.1% fongizone supplemented with suitable selective antibiotics. The cells were seeded at a density of 2×10⁴ cells/cm².

The culture medium was changed every 48 hours. When the cells reached confluence, they were washed in phosphate buffer (PBS) without CaCl₂ and without MgCl₂, then detached with trypsin/0.05% EDTA, sedimented by low speed centrifugation (400 g for 3 min) and resuspended in 5 ml of culture medium.

Intracellular immunofluorescence and flow cytometry were used to check that the cell clones expressed the recombinant glycopeptide of interest.

The recombinant glycopeptide was purified by liquid chromatography followed by affinity chromatography and analyzed by SDS-PAGE.

It could be seen on the gel that the secreted peptide resolved into two different molecular weight bands of approximately 78 kDa and 83 kDa.

The 83 kDa band probably corresponds to the glycosylated peptide whereas the 78 kDa band would either be a non-glycosylated form or a form with less glycosylation.

It therefore appears that the aforementioned clone expressed a glycopeptide having the size and the characteristics of those corresponding to the C-terminal end of BSDL.

23

Example 2

Recombinant Glycopeptide Having the Size and the Characteristics of Those Corresponding to the C-Terminal End of BSDL (CHO-C2-16R)

The method described in example 1 was employed, but by culturing the cell clone CHO-C2-16R (preparation 6).

The recombinant glycopeptide obtained was purified by liquid chromatography followed by affinity chromatography and analyzed by SDS-PAGE.

It could be seen on the gel that the secreted peptide resolved into two different molecular weight bands of approximately 78 kDa and 83 kDa.

It therefore appears that the aforementioned clone expressed a glycopeptide having the size and the characteristics of those corresponding to the C-terminal end of BSDL.

Example 3

Recombinant Glycopeptide Having the Size and the Characteristics of Those Corresponding to the C-Terminal End of BSDL (CHO-F3-16R)

The method described in example 1 was employed, but by culturing the cell clone CHO-F3-16R (preparation 6).

The recombinant glycopeptide obtained was purified by liquid chromatography followed by affinity chromatography and analyzed by SDS-PAGE.

It could be seen on the gel that the secreted peptide resolved into two different molecular weight bands of approximately 78 kDa and 83 kDa.

It therefore appears that the aforementioned clone expressed a glycopeptide having the size and the characteristics of those corresponding to the C-terminal end of BSDL.

Example 4

Recombinant Glycopeptide Having the Size and the Characteristics of Those Corresponding to the C-Terminal End of BSDL (CHO-C2-F3-16R)

The method described in example 1 was employed, but by culturing the cell clone CHO-C2-F3-16R (preparation 8).

The recombinant glycopeptide obtained was purified by liquid chromatography followed by affinity chromatography and analyzed by SDS-PAGE.

It could be seen on the gel that the secreted peptide resolved into two different molecular weight bands of approximately 78 kDa and 83 kDa.

It therefore appears that the aforementioned clone expressed a glycopeptide having the size and the characteristics of those corresponding to the C-terminal end of BSDL.

Example 5

Recombinant Glycopeptide Having the Size and the Characteristics of Those Corresponding to the C-Terminal End of BSDL (CHO-F7-16R)

The method described in example 1 was employed, but by culturing the cell clone CHO-F7-16R (preparation 6).

The recombinant glycopeptide obtained was purified by liquid chromatography followed by affinity chromatography and analyzed by SDS-PAGE.

Only one band of molecular weight 83 kDa was seen on the gel.

24

Example 6

Recombinant Glycopeptide Having the Size and the Characteristics of Those Corresponding to the C-Terminal End of BSDL (CHO-C2-F7-16R)

The method described in example 1 was employed, but by culturing the cell clone CHO-C2-F7-16R (preparation 9).

The recombinant glycopeptide obtained was purified by liquid chromatography followed by affinity chromatography and analyzed by SDS-PAGE.

It could be seen on the gel that the secreted peptide resolved into two different molecular weight bands of approximately 78 kDa and 83 kDa.

It therefore appears that the aforementioned clone expressed a glycopeptide having the size and the characteristics of those corresponding to the C-terminal end of BSDL.

It therefore appears that all the aforementioned clones (16R) expressed glycopeptides having the size and the characteristics of those corresponding to the C-terminal end of BSDL.

Examples 7 to 13

Recombinant Glycopeptides Having the Size and the Characteristics of Those Corresponding to the C-Terminal End of FAPP

The recombinant polypeptides produced by clones CHO-K1 (control), CHO-6R (Ex 7), CHO-C2-6R (Ex 8), CHO-F3-6R (Ex 9), CHO-C2-F3-6R (Ex 10), CHO-GT-6R (Ex 11), CHO-C2-GT-6R (Ex 12) and CHO-C2-F3-GT-6R (Ex 13), were purified on a polyacrylamide gel or else by liquid and/or affinity chromatography and analyzed by immunodetection using antibody pAbL64 described by Abouakil et al., 1988, *Biochim. Biophys. Acta*, 961: 299-308, and mAbJ28. The mouse monoclonal antibody mAbJ28, specific of the fucosylated J28 glycopeptide derived from O-glycosylation of the oncofetal form of FAPP, is described by Panicot et al. in *Glycobiology*, 1999, 9: 935-946.

The anti-BSDL polyclonal antibody pAbL64 was obtained by immunizing rabbits after purifying the antigen from human pancreatic juices. Said antibody perfectly recognizes FAPP, the oncofetal form of BSDL.

The polyclonal antibody specific of the BSDL protein core, named anti-peptide, was obtained by immunizing rabbits with a synthetic peptide corresponding to the peptide sequence of BSDL comprised between the serine residue in position 346 and glutamine in position 370 coupled to hemocyanin.

Said two antibodies were purified on a protein A-Sepharose 4B column (Pharmacia). The anti-peptide was subjected to a second purification step by affinity chromatography on a column of keyhole limpet hemocyanin (KLH) coupled to Sepharose gel to eliminate anti-KLH antibodies.

The recombinant peptides produced by the different clones of the 6R type were loaded at 100 µg per lane on a 10% polyacrylamide gel, separated, then electrotransferred to a nitrocellulose membrane. The membranes were then incubated in the presence of antibody pAbL64 on the one hand, and antibody mAbJ28 on the other hand, and developed.

In both cases, the antibodies employed bound specifically to polypeptides having a size of 50 kDa, corresponding to the

expected size of the C-terminal glycopeptides of FAPP, except in the case of the non-transfected CHO-K1 cell line used as negative control.

Examples 14 to 19

Anti-Glycopeptide Monoclonal Antibodies

Recombinant anti-glycopeptide antibodies having the size and the characteristics of those corresponding to the C-terminal end of FAPP were prepared as follows:

Stage A—Immunization of Mice.

Six-week-old male Balb/c mice were immunized according to the following protocol:

Day 0: Intraperitoneal injection of 25 µg of a mixture of FAPP and BSDL purified from normal and pathological human pancreatic juices (hereinbelow named BSDL-FAPP antigen) in a 50:50 emulsion (150 µl NaCl/150 µl complete Freund's adjuvant).

Day 15: Intraperitoneal challenge with 25 µg of BSDL-FAPP antigen in a 50:50 emulsion (150 µl NaCl/150 µl incomplete Freund's adjuvant).

Day 30: Intraperitoneal challenge with 25 µg of BSDL-FAPP antigen in a 50:50 emulsion (150 µl NaCl/150 µl incomplete Freund's adjuvant).

Day 110: Intraperitoneal challenge with 20 µg of BSDL-FAPP antigen in a 50:50 emulsion (150 µl NaCl/150 µl incomplete Freund's adjuvant).

Day 140: Intraperitoneal challenge with 20 µg of BSDL-FAPP antigen in a 50:50 emulsion (150 µl NaCl/150 µl incomplete Freund's adjuvant).

Day 215: Intraperitoneal challenge with 20 µg of BSDL-FAPP antigen in a 50:50 emulsion (150 µl NaCl/150 µl incomplete Freund's adjuvant).

Day 244: Intravenous injection of 10 µg of BSDL-FAPP antigen in 100 µl of sterile NaCl.

Day 247: Cell fusion.

Stage B—Cell Fusion According to the Protocol of Kohler and Milstein.

a) At day 247, the selected mouse was sacrificed and the spleen removed and ground up. The spleen cells were washed in RPMI 1640 medium. P3.X63.Ag8 653 myeloma cells, previously grown in RPMI 1640 medium containing 20% fetal calf serum (FCS), 1% glutamine, 1% nonessential amino acids and 1% sodium pyruvate were also washed in the same medium.

At the same time, peritoneal macrophages were collected by peritoneal lavage of non-immunized Balb/c mice with RPMI.

For hybridoma formation, the spleen cells and myeloma cells were mixed in a tube at a ratio of 5 spleen cells for 1 myeloma cell. After centrifugation, the cell pellet was resuspended in 800 µl of 50% polyethylene glycol 1500 in 75 mM Hepes buffer pH 7.5. After 1 minute of contact at 37° C., 20 ml of RPMI 1640 medium were slowly added to the fused cells.

b) The initial culture was carried out on 96-well microtiteration plates in the presence of RPMI medium containing 20% fetal calf serum (FCS) and supplemented with 5×10^{-3} M hypoxanthine, 2×10^{-5} M aminopterin and 8×10^{-4} M thymidine. Next, 5×10^3 peritoneal macrophages followed by 10^5 fused cells were deposited in each well.

Stage C—Cloning and Subcloning.

Each hybridoma selected by the method described in stage D hereinbelow resulted from cloning by a limit dilution technique in which 10, 5, 2, 1 and 0.5 cells were statistically distributed into microwells containing peritoneal macroph-

ages. Two subclonings were thus carried out, each clone and subclone having been replicated then frozen in 90% FCS and 10% dimethylsulfoxide (DMSO). Subclones from the last generation were then expanded in vivo to obtain ascites fluid in the Balb/c mice, followed by immunoglobulin purification on a protein A column.

Stage D—Hybrid Cell Selection Method.

Selection was carried out by a liquid phase ELISA method using culture supernatant. Five micrograms of BSDL-FAPP antigen in 100 µl of bicarbonate buffer pH 8.5 were deposited in the wells of a 96-well ELISA plate. The plate was activated for 12 h at 4° C. and saturated for 2 hours with 300 µl of 1 mg/ml bovine albumin. The plates were then washed and incubated with 100 µl of cell culture supernatants potentially containing antibodies directed against the BSDL-FAPP antigen. After a 2-hour incubation, the plates were washed and incubated for 2 hours with alkaline phosphatase-labelled secondary antibodies. At the end of the incubation, the plates were again washed and incubated with para-nitrophenylphosphate (100 µl, 1 mg/ml in 0.2 M Tris/HCl buffer pH 8.2 and 1 mM CaCl_2). After 1 hour at 37° C., the plates were read on a microplate reader at 410 nm.

About 15 antibodies were selected for their response in the ELISA test. Additional analyses by Western blot (immunoprinting) using FAPP as immunogenic protein enabled the selection of hybridomas 7B4 (Example 14), 11D7 (Example 15), 14H9 (Example 16), 14H10 (Example 17), 16D10 (Example 18) and 8H8 (Example 19).

Cells producing the IgM antibody 16D10 were deposited with the Collection Nationale de Culture de Microorganismes (CNCM), Institut Pasteur, 28, rue du Dr Roux, 75724 Paris Cédex 15, France on 16 Mar. 2004 under the number I-3188.

Example 20

Preparation of Membrane Glycopeptides of Natural Origin

Membrane glycopeptides carrying epitopes recognized by the monoclonal antibodies described in examples 14 to 19 were prepared in the following manner: pancreatic tumor cells were cultivated on a plastic support then detached therefrom with non-enzymatic dissociation solution (Sigma). The cell pellet obtained by centrifugation (2 min at 1000 rpm) was sonicated (2×15 sec) and again centrifuged (20 min, 14,000 rpm, 4° C.). The pellet suspended in phosphate buffer corresponds to membrane glycopeptides carrying epitopes recognized by the monoclonal antibodies described in examples 14 to 19.

Example 21

A vaccine having the following composition was prepared: membrane glycopeptides from example 20 isolated from

HaPT-1 pancreatic cells, 20 µg
ALU-gel-ser adjuvant (Serva) 150 µl
excipient including water for injections 150 µl

Vaccination was carried out by intraperitoneal injection two weeks then one week before transplanting the tumor cells into the flank of the animals. The result of said vaccination was compared to that of a placebo (injection of isotonic solution).

Example 22

A 0.5 mg/ml injectable isotonic solution of antibodies from example 18 (16D10) was prepared.

Control Example 1

Recombinant Glycopeptide Having the Size and the Characteristics of Those Corresponding to the C-terminal End of BSDL

The method described in example 1 was employed, but by cultivating the cell clone CHO-K1.

The gel did not show any bands at molecular weight 78 kDa or 83 kDa.

PHARMACOLOGICAL STUDY

Experiment 1

Use of Recombinant C-Terminal Glycopeptides of BSDL and FAPP in Cellular Immunotherapy of Exocrine Pancreatic Cancer

Operating Protocol:

Hamsters (90-100 g) were divided into several groups and vaccinated with HaPT-1 cells inactivated by 30 minutes of UV exposure. The hamsters were inoculated with the glycopeptides from example 20 and inactivated cells mixed with ALU-gel-ser adjuvant by intraperitoneal injection once a week for two weeks. Two weeks after the last inoculation, the hamsters received an ectopic transplantation of HaPT-1 cells (subcutaneous, on the flank).

Tumor growth was monitored weekly and tumor volume was compared with the control group vaccinated with a placebo (PBS alone).

Results:

The tumor growth curve was significantly slower in hamsters immunized with the glycopeptides from example 20 (antigens represented in large part by structures recognized by monoclonal antibodies J28 and 16D10 (Example 18), directed against the C-terminal domain of BSDL and/or FAPP).

At the end of the experiment, tumor volume in the group vaccinated with inactivated cells was 80% lower than in the control group.

The hamsters were sacrificed and pancreatic and tumor tissue samples were removed for immunohistochemical analysis. Monoclonal antibodies J28 and 16D10 and polyclonal antibody L64 were used to test for expression of the 16D10 and J28 epitopes (recognized by monoclonal antibodies J28 and 16D10, respectively) and the FAPP protein core.

Examination of the tumor tissue slices revealed a decrease in the expression of the 16D10 and J28 epitopes (smaller decrease for the latter), between the group vaccinated with inactivated cells and the control group.

Therefore, the C-terminal glycopeptides of BSDL and FAPP according to the invention can be used in cellular immunotherapy of exocrine pancreatic cancer.

Experiment 2

Use of the Glycopeptides According to the Invention in the Diagnosis of Diabetic Pathologies

Operating Protocol:

A series of experiments carried out by the ELISA method on sera from patients suffering from various pancreatic and other pathologies showed that BSDL, used as antigen to activate the ELISA plates, was specifically recognized by antibodies present in approximately 80% of the sera from patients with type I diabetes whereas less than 10% of the sera from

control patients were reactive. Sera from patients with other pathologies such as pancreatitis, pancreatic cancer or Graves' disease were not reactive (Panicot et al. 1999, Diabetes, 48: 2316-2323).

The recombinant glycopeptides from examples 2, 5 and 6 were tested by Western blot (or immuno-imprinting).

Results:

BSDL, and in particular the C-terminal domain thereof, was recognized by sera from diabetic patients whereas BSDL was not recognized by sera from normal control patients.

The recombinant glycopeptides from examples 2 and 6 were recognized by sera from diabetic (type I) patients.

The control sera (from healthy patients) did not recognize any of said three glycopeptides.

Therefore, the natural and recombinant C-terminal glycopeptides of the invention described in the examples can be used for the diagnosis of diabetic pathologies.

Experiment 3

Use of Monoclonal Antibodies Directed Against the C-Terminal Domain of FAPP and/or BSDL in the Diagnosis of Pancreatic Cancer

Operating Protocol:

Human pancreatic tumor cell lines SOJ-6 and Bx-PC-3 and human non-pancreatic tumor cell lines Caco-2 and Hep-G2 were treated with monoclonal antibodies J28, 16D10 (Example 18), and 8H8 (Example 19) and with polyclonal antibody (pAb) L64 specific of the C-terminal domain of BSDL and FAPP. The respective antigen-antibody complexes were detected with the aid of a fluorescent rabbit or mouse fluorescein conjugated N-isothiocyanate anti-IgG antibody. FACS analysis was used to detect the presence of antigens associated with the C-terminal domain of FAPP and BSDL at their surface.

Results:

The monoclonal antibodies J28 and 16D10 (Example 18), and the polyclonal antibody (pAb) L64 specifically recognized the surface of the human pancreatic tumor cell lines SOJ-6 and Bx-PC-3 and did not recognize the surface of the human non-pancreatic tumor cell lines which were tested, such as Caco-2 and Hep-G2 cells.

The monoclonal antibodies J28 and 16D10 directed against the glycosylated epitopes expressed specifically on the C-terminal domain of FAPP and BSDL are therefore capable of distinguishing pancreatic tumor cells from non-pancreatic tumor cells.

Therefore, the monoclonal antibodies directed against the glycosylated epitopes expressed specifically on the C-terminal domain of FAPP and BSDL can be used in the diagnosis of pancreatic cancer.

In particular said antibodies can determine whether a secondary tumor site (metastasis) is of pancreatic origin or from a primary tumor of another origin.

Experiment 4

Targeting of Pancreatic Neoplastic Cells by the Monoclonal Antibodies of the Invention

A—Study on Human Pancreatic Tissue Sections.

Operating Protocol:

Tissue expression of the glycosylated epitopes recognized by the monoclonal antibodies directed against the C-terminal domain of FAPP and/or BSDL was tested in immunohis-

tochemical studies carried out with the various monoclonal antibodies of the invention on normal and cancerous human pancreatic tissue sections.

In a first step, pancreatic tissue was obtained from four patients with pancreatic cancer and from five normal pancreases. The tissue sections were incubated with the antibodies from examples 18 (16D10) and 19 (8H8). In a second step, seven human pancreatic tumor tissue samples and five non-tumoral tissues (diagnosed by a pathologist) were sliced into 5- μ m thick sections, dried in acetone at 4° C. and rehydrated in Tris/HCl buffer pH 7.6. The sections were then incubated either for 1 h at 25° C. (mAbJ28 and mAb8H8, example 19) or overnight at 4° C. (mAb 16D10, example 18), the antibodies being diluted 1:50 in Dako diluent (EnVision System, Dako, Copenhagen, Denmark). The sections were then incubated with a suitable alkaline phosphatase-labelled secondary antibody. The antigen-antibody complex was then detected with the chromogenic substrate Fast-Red. In the case of quantitative analysis by confocal laser scanning microscopy, the suitable secondary antibody was labelled with streptavidin-fluorescein. The sections were then examined under a microscope or with a confocal laser fluorescence microscope.

Results:

In the first experiment, the monoclonal antibody from example 18 (16D10) specifically recognized neoplastic cells from four pancreatic tumor tissue samples.

No reaction was observed with the five normal pancreatic tissues.

The monoclonal antibody from example 19 (8H8) preferentially recognized normal pancreatic tissue.

In the second experiment, out of all seven tumor tissue samples studied, seven were recognized by mAb16D10 (example 18) and five by mAbJ28. The immunohistochemical results are presented in FIG. 1 and Tables 1 and 2. The mAb16D10 (example 18) showed very high specificity for tumor tissue whereas mAb8H8 (example 19) only recognized normal pancreatic tissue. mAbJ28 showed intermediate specificity since it preferentially recognized tumor tissue, but also some normal tissue. Table 2 presents the confocal laser microscopy results defining the fluorescence intensity on the antibody-labelled tissue sections. It should be noted that none of these antibodies recognized the other tumor tissues tested: liver, lung, stomach, colon, esophagus and thyroid.

The monoclonal antibodies of the invention coupled to an antitumoral active substance can be used in the treatment of pancreatic cancer to target and destroy pancreatic tumor cells.

Similarly, radioactive (or other) elements can be coupled to said antibodies and enable a precise localization of primary tumors and metastases. Said antibodies can also afford a clear and sharp differentiation between normal tissue and tumor tissue.

TABLE 1

Semi-quantitative immunohistochemical studies: results obtained with antibodies mAb8H8, mAbJ28 and mAb16D10			
	Label		
	mAb8H8	mAbJ28	mAb16D10
Control 1			
Acini	4	1	0
Canals	0	0	0
Control 2			
Acini	4	1	0
Canals	0	0	0

TABLE 1-continued

Semi-quantitative immunohistochemical studies: results obtained with antibodies mAb8H8, mAbJ28 and mAb16D10			
	Label		
	mAb8H8	mAbJ28	mAb16D10
Control 3			
Acini	4	1	0
Canals	0	0	0
Control 4			
Acini	4	1	0
Canals	0	0	0
Control 5			
Acini	4	1	0
Canals	0	0	0
PDAC1	0	0	4
PDAC2	0	3	4
PDAC3	0	2	4
PDAC4	0	0	4
PDAC5	1	3	4
PDAC6	0	2	4
PDAC7*	0	1	3
CCA*	0	0	0
ECA*	0	0	0
LiCa*	0	0	0
LuCa	0	0	0
SCA*	0	0	0
TCA*	0	0	0

C, control; PDAC, pancreatic adenocarcinoma; LuCa, lung carcinoma; CCA, colon carcinoma; ECA, esophageal carcinoma; LiCa, liver carcinoma; SCA, stomach carcinoma; TCA, thyroid carcinoma. Tissues were obtained from the BioChain Institute (USA). The number of labelled cells is expressed semi-quantitatively as follows: 0, no labelling; 1, from 1 to 10% of cells were labelled; 2, from 10 to 30% of cells were labelled; 3, from 30 to 40% of cells were labelled and 4: >40% of cells were labelled.

TABLE 2

Fluorescence intensity on sections labelled with the antibodies of the invention.			
Case	Fluorescence intensity		
	mAb8H8	mAbJ28	mAb16D10
Control 1	92.33 \pm 2.80	19.11 \pm 8.75	31.41 \pm 1.19
Control 2	138.72 \pm 3.36	14.54 \pm 6.57	8.16 \pm 5.16
Control 3	110.95 \pm 2.71	17.41 \pm 8.09	0
Control 4	109.04 \pm 4.58	26.49 \pm 12.01	0
Control 5	114.14 \pm 1.26	35.56 \pm 16.30	0
PDAC1	0	53.12 \pm 2.12	124.02 \pm 2.43
PDAC 2	36.47 \pm 0.35	87.88 \pm 4.28	129.84 \pm 6.35
PDAC 3	0	89.60 \pm 2.84	123.27 \pm 6.16
PDAC 4	17.06 \pm 7.64	50.61 \pm 2.36	143.95 \pm 2.71
PDAC 5	40.42 \pm 0.22	88.92 \pm 5.48	135.06 \pm 4.03
PDAC 6	0	59.97 \pm 1.64	129.30 \pm 1.91
PDAC 7*	18.65 \pm 8.38	72.74 \pm 4.84	132.34 \pm 2.75

C, control; PDAC, pancreatic adenocarcinoma. Tissues were obtained from the BioChain Institute (USA). In each case the mean \pm standard error of labelling on six different regions of each tissue slice are shown. Significant differences were observed between control and pancreatic tumor tissue for mAb16D10 ($P < 0.001$), mAb8H8 ($P < 0.001$) and mAbJ28 ($P < 0.05$).

B—ELISA Competition Studies Between mAbJ28 and mAb16D10 (Example 18) on the Recombinant Glycopeptide Carrying the J28 Glycotope.

Operating Protocol:

The glycosyltransferases involved in formation of the oncofetal glycotipe J28 were characterized and said glycotipe was reproduced ex vivo by genetic engineering. It was reconstructed in CHO cells from the recombinant C-terminal peptide of FAPP (composed of 6 repeated sequences) and two glycosyltransferases, α (1-3/4) fucosyltransferase FUT3 and Core 2 β (1-6) N-acetylglucosaminyltransferase required to build said structure. The recombinant C-terminal glycopep-

tion of FAPP carrying the J28 glyco-type (example 10) was secreted in the culture supernatant of this cell line.

Competition experiments were carried out with an ELISA test in the presence of culture supernatant containing the recombinant C-terminal glycopeptide of FAPP carrying the J28 glyco-type (example 10), biotinylated mAbJ28 and increasing concentrations of mAb16D10 (example 18) as competitor with the aim of demonstrating the specificity of said antibodies.

The principle of the protocol was as follows:

1. The bottom of the wells was coated with concentrated culture supernatant containing the recombinant J28 glycopeptide diluted 1:10, 1:20, 1:50, 1:100.

2. Biotinylated mAbJ28 was added at 10 ng per well (Exp. 1) or 5 ng per well (Exp. 2) together with increasing concentrations of purified mAb16D10.

3. Biotinylated mAbJ28 was recognized with the aid of an alkaline phosphatase-conjugated anti-biotin antibody (anti-biotin Ab control on mAb26D10=0).

Results:

FIG. 2 presenting the results of two experiments shows that there was no significant loss in the reactivity of mAbJ28 for its glyco-type coated at the bottom of the wells in the presence of increasing concentrations of mAb16D10 (example 18) used as competitor (even with a four-fold excess).

C—FACS Competition Study Between Antibodies mAbJ28 and mAb16D10 on human pancreatic tumor cell line SOJ-6.

Operating Protocol:

The monoclonal antibodies mAbJ28 and mAb16D10 specifically recognized human pancreatic tumor cell lines SOJ-6, Bx-PC-3 and did not recognize the human non-pancreatic tumor cell lines tested.

Preliminary competition experiments by FACS were carried out on the human pancreatic tumor cell line SOJ-6 in the presence of monoclonal antibody mAbJ28 coupled to a fluorochrome (Alexa 488) and increasing concentrations of monoclonal antibody mAb16D10 (example 18) used as competitor in the form of ascites fluid.

The principle of the protocol was as follows:

1. Human pancreatic tumor SOJ-6 cells were detached, fixed and saturated with BSA at 4° C.

2. Fluorescent-labelled monoclonal antibody mAbJ28 was added at 2 ng per sample in the presence of increasing concentrations of ascites fluid containing mAb16D10 at 1:100, 1:40, 1:20, 1:10, 1:4 and 1:2 dilution.

3—The cells were analyzed by FACS.

Results:

The results are shown in FIG. 3. Despite a very low labeling of the cells (due to the first experiments with Alexa-488-labelled mAbJ28), no significant loss of reactivity of antibody mAbJ28 for its glyco-type was observed in the presence of increasing concentrations of antibody mAb16D10 used as competitor.

Experiment 5

Use of the Monoclonal Antibodies of the Invention in the Treatment of Exocrine Pancreatic Cancer

Experimental Protocol:

Two groups of six-week-old nude mice (NMRI mice) received an intraperitoneal inoculation on Day -1 either with monoclonal antibody 16D10 from example 18 at a dose of 150 µl of a 0.5 mg/mg isotonic antibody solution or with an isotonic placebo solution.

At Day 0, 2×10^6 SOJ-6 human pancreatic tumor cells were subcutaneously injected into the right flank. The animals were then inoculated with monoclonal antibody 16D10 or isotonic solution at Days +1, +3, +6, +8 and +10 (in the quantities indicated hereinabove).

The volume of the palpable subcutaneous tumors was measured in each animal at weeks 1, 2, 3, 4, 5, 6, 7, 8, 9 and 10.

Results:

The tumor growth rate in animals treated with the monoclonal antibody from example 18 was 10 times lower than that of placebo-treated animals.

At the end of the experiment (10 weeks post-inoculation), the median volume of tumors removed from control animals was 1.25 versus a markedly lower median value of 0.15 cm³ in the treated animals.

The monoclonal antibodies directed against the C-terminal domain of FAPP and/or BSDL and in particular the monoclonal antibody from example 18 can therefore be used in therapy for pancreatic cancer.

Experiment 6

Use of the Monoclonal Antibodies Directed Against the C-terminal Domain of FAPP and/or BSDL in the Urinary Diagnosis of Pancreatic Cancer

Operating Protocol:

A series of experiments carried out by Western blot, liquid chromatography and mass spectrometric characterization led to the identification of intact BSDL in the urine of healthy patients. In light of these findings, the different glycosylated epitopes J28, 16D10 carried by the C-terminal domain of BSDL/FAPP and which appear in pancreatic adenocarcinoma, might also be present in the urine of patients suffering from this cancer.

Results:

FIG. 4 showing the initial results obtained in a preliminary Western blot experiment reveals the presence of the glycosylated J28 epitope carried by BSDL/FAPP in urine samples from patients with pancreatic adenocarcinoma, and the absence thereof in urine samples from healthy individuals.

FIG. 4 shows that both healthy subjects and cancer patients had one or two bands displaying immunoreactivity towards pAbL64 and mAb8H8 (example 19) located at 110 kDa. On the other hand, the two groups displayed different reactivity towards mAbJ28, a mAbJ28-immunoreactive band at 110 kDa being seen in cancer patients but not in healthy subjects.

Experiment 7

Use of Urinary C-terminal Glycopeptides of BSDL and FAPP in Cellular Immunotherapy of Exocrine Pancreatic Cancer

Operating Protocol:

Preliminary results showed that the glycosylated J28 epitope carried by BSDL/FAPP was present in urine samples from patients with pancreatic adenocarcinoma tested at the present time whereas it was absent in urine samples from healthy individuals. The presence of said tumor marker in urine, which can be obtained in large quantities without the use of invasive methods, can serve as a source of antigen in order to develop different active immunotherapy protocols such as described by Ramanathan et al., *Cancer Immunol Immunoth*, 2004, 54: 254-64; O'Neill et al., *Blood*, 2004, 104: 2235-46; Vlad et al., *J Exp Med*, 2002, 196: 1435-46; Schmidt et al., *Cancer Res*, 2003, 63: 8962-67. One such

avenue is to develop autologous cell immunotherapy protocols (as described by Yamanaka et al., *Br J Cancer*, 2003, 89: 1172-9; Svane et al., *Cancer Immunol Immunother*, 2004, 53: 633-41; Yu et al., *Cancer Res*, 2004, 64: 4973-9) using dendritic cells from patients who have their pancreatic tumor resected. The patient's dendritic cells are then loaded ex vivo with his own antigens, that is to say, with the C-terminal glycopeptide(s) of BSDL/FAPP carrying the J28 glyco-
 5 tope(s) (and perhaps also the 16D10 glycotope) obtained from the urine, before being reinjected in the patient to induce an immune response directed against residual tumor foci which express said same antigens.

Said strategy of human dendritic cell activation, currently under evaluation using as tumor antigen the recombinant C-terminal glycopeptide of FAPP carrying the glycosylated J28 epitope (example 10), can also be carried out by using the urinary C-terminal glycopeptide of BSDL/FAPP carrying said glycosylated epitopes. The purification protocols (different steps of centrifugation, ultrafiltration and concentration, combined with liquid chromatography on exclusion and affinity columns) developed in order to purify and characterize BSDL in the urine of healthy individuals are used to purify BSDL/FAPP carrying the J28 glycotope (and also that of 16D10) from urine samples of patients with pancreatic adenocarcinoma. The protocol is as follows: Native urine is

centrifuged to eliminate cellular debris and partially desalted by ultrafiltration on a Centripreps® YM-10, then completely desalted on a Sephadex® G-25 column. The recovered fractions are lyophilized, then taken up in 10 mM Tris-HCl elution buffer, pH 7.8, 400 mM NaCl, and loaded on Sephacryl™ S-200 gel to separate the proteins according to their molecular mass. Fractions containing BSDL/FAPP carrying the J28 glycotope (and/or the 16D10 glycotope) are collected, dialyzed and lyophilized. The lyophilizates obtained after molecular sieving are rehydrated in PBS, and BSDL/FAPP carrying the J28 glycotope (and perhaps also the 16D10 glycotope) is purified either by immunoaffinity chromatography on an agarose-mAbJ28 column or on a heparin Sepharose column. Purified BSDL/FAPP carrying the J28 glycotope (and/or the 16D10 glycotope) is digested with trypsin or cyanogen bromide to obtain the C-terminal glycopeptide of BSDL/FAPP carrying said glycotopes, as described elsewhere to isolate the C-terminal domain of BSDL/FAPP (Mas et al., 1997, *Glycobiology*, 7: 745-752). The dendritic cells of patients with pancreatic adenocarcinoma are cultured in the presence of said glycopeptide carrying the J28 glycotope (and/or the 16D10 glycotope) purified from their own urine. The presence of the glycopeptides at the surface of the pulsed mature dendritic cells will be checked by confocal microscopy before the cells are reinjected in the patient to induce an anti-tumor response.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 14

<210> SEQ ID NO 1

<211> LENGTH: 36

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

ttcgtaagct tgcgaagctg ggcgccgtgt acagaa 36

<210> SEQ ID NO 2

<211> LENGTH: 42

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

tttcgtgaat tcacgctaaa acctaatgac tgcaggcatc tg 42

<210> SEQ ID NO 3

<211> LENGTH: 33

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

cgtctaaagc tttttgatgt ctacaccgag tcc 33

<210> SEQ ID NO 4

<211> LENGTH: 27

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

aaaaagaatt cggagaaaat aatgaat 27

<210> SEQ ID NO 5

<211> LENGTH: 28

-continued

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

aaaaagggcc cacaaagtca gacattat 28

<210> SEQ ID NO 6

<211> LENGTH: 2350

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (17)..(2251)

<400> SEQUENCE: 6

ggccaccag aggctg atg ctc acc atg ggg cgc ctg caa ctg gtt gtg ttg 52
 Met Leu Thr Met Gly Arg Leu Gln Leu Val Val Leu
 1 5 10

ggc ctc acc tgc tgc tgg gca gtg gcg agt gcc gcg aag ctg ggc gcc 100
 Gly Leu Thr Cys Cys Trp Ala Val Ala Ser Ala Ala Lys Leu Gly Ala
 15 20 25

gtg tac aca gaa ggt ggg ttc gtg gaa ggc gtc aat aag aag ctc ggc 148
 Val Tyr Thr Glu Gly Gly Phe Val Glu Gly Val Asn Lys Lys Leu Gly
 30 35 40

ctc ctg ggt gac tct gtg gac atc ttc aag ggc atc ccc ttc gca gct 196
 Leu Leu Gly Asp Ser Val Asp Ile Phe Lys Gly Ile Pro Phe Ala Ala
 45 50 55 60

ccc acc aag gcc ctg gaa aat cct cag cca cat cct ggc tgg caa ggg 244
 Pro Thr Lys Ala Leu Glu Asn Pro Gln Pro His Pro Gly Trp Gln Gly
 65 70 75

acc ctg aag gcc aag aac ttc aag aag aga tgc ctg cag gcc acc atc 292
 Thr Leu Lys Ala Lys Asn Phe Lys Lys Arg Cys Leu Gln Ala Thr Ile
 80 85 90

acc cag gac agc acc tac ggg gat gaa gac tgc ctg tac ctc aac att 340
 Thr Gln Asp Ser Thr Tyr Gly Asp Glu Asp Cys Leu Tyr Leu Asn Ile
 95 100 105

tgg gtg ccc cag ggc agg aag caa gtc tcc cgg gac ctg ccc gtt atg 388
 Trp Val Pro Gln Gly Arg Lys Gln Val Ser Arg Asp Leu Pro Val Met
 110 115 120

atc tgg atc tat gga ggc gcc ttc ctc atg ggg tcc ggc cat ggg gcc 436
 Ile Trp Ile Tyr Gly Gly Ala Phe Leu Met Gly Ser Gly His Gly Ala
 125 130 135 140

aac ttc ctc aac aac tac ctg tat gac ggc gag gag atc gcc aca cgc 484
 Asn Phe Leu Asn Asn Tyr Leu Tyr Asp Gly Glu Glu Ile Ala Thr Arg
 145 150 155

gga aac gtc atc gtg gtc acc ttc aac tac cgt gtc ggc ccc ctt ggg 532
 Gly Asn Val Ile Val Val Thr Phe Asn Tyr Arg Val Gly Pro Leu Gly
 160 165 170

ttc ctc agc act ggg gac gcc aat ctg cca ggt aac tat ggc ctt cgg 580
 Phe Leu Ser Thr Gly Asp Ala Asn Leu Pro Gly Asn Tyr Gly Leu Arg
 175 180 185

gat cag cac atg gcc att gct tgg gtg aag agg aat atc gcg gcc ttc 628
 Asp Gln His Met Ala Ile Ala Trp Val Lys Arg Asn Ile Ala Ala Phe
 190 195 200

ggg ggg gac ccc aac aac atc acg ctc ttc ggg gag tct gct gga ggt 676
 Gly Gly Asp Pro Asn Asn Ile Thr Leu Phe Gly Glu Ser Ala Gly Gly
 205 210 215 220

gcc agc gtc tct ctg cag acc ctc tcc ccc tac aac aag ggc ctc atc 724
 Ala Ser Val Ser Leu Gln Thr Leu Ser Pro Tyr Asn Lys Gly Leu Ile
 225 230 235

cgg cga gcc atc agc cag agc ggc gtg gcc ctg agt ccc tgg gtc atc 772
 Arg Arg Ala Ile Ser Gln Ser Gly Val Ala Leu Ser Pro Trp Val Ile

-continued

240	245	250	
cag aaa aac cca ctc ttc tgg gcc	aaa aag gtg gct gag aag gtg ggt		820
Gln Lys Asn Pro Leu Phe Trp Ala Lys Lys Val Ala Glu Lys Val Gly			
255	260	265	
tgc cct gtg ggt gat gcc gcc agg atg gcc cag tgt ctg aag gtt act			868
Cys Pro Val Gly Asp Ala Ala Arg Met Ala Gln Cys Leu Lys Val Thr			
270	275	280	
gat ccc cga gcc ctg acg ctg gcc tat aag gtg ccg ctg gca ggc ctg			916
Asp Pro Arg Ala Leu Thr Leu Ala Tyr Lys Val Pro Leu Ala Gly Leu			
285	290	295	300
gag tac ccc atg ctg cac tat gtg ggc ttc gtc cct gtc att gat gga			964
Glu Tyr Pro Met Leu His Tyr Val Gly Phe Val Pro Val Ile Asp Gly			
305	310	315	
gac ttc atc ccc gct gac ccg atc aac ctg tac gcc aac gcc gcc gac			1012
Asp Phe Ile Pro Ala Asp Pro Ile Asn Leu Tyr Ala Asn Ala Ala Asp			
320	325	330	
atc gac tat ata gca ggc acc aac aac atg gac ggc cac atc ttc gcc			1060
Ile Asp Tyr Ile Ala Gly Thr Asn Asn Met Asp Gly His Ile Phe Ala			
335	340	345	
agc atc gac atg cct gcc atc aac aag ggc aac aag aaa gtc acg gag			1108
Ser Ile Asp Met Pro Ala Ile Asn Lys Gly Asn Lys Lys Val Thr Glu			
350	355	360	
gag gac ttc tac aag ctg gtc agt gag ttc aca atc acc aag ggg ctc			1156
Glu Asp Phe Tyr Lys Leu Val Ser Glu Phe Thr Ile Thr Lys Gly Leu			
365	370	375	380
aga ggc gcc aag acg acc ttt gat gtc tac acc gag tcc tgg gcc cag			1204
Arg Gly Ala Lys Thr Thr Phe Asp Val Tyr Thr Glu Ser Trp Ala Gln			
385	390	395	
gac cca tcc cag gag aat aag aag aag act gtg gtg gac ttt gag acc			1252
Asp Pro Ser Gln Glu Asn Lys Lys Lys Thr Val Val Asp Phe Glu Thr			
400	405	410	
gat gtc ctc ttc ctg gtg ccc acc gag att gcc cta gcc cag cac aga			1300
Asp Val Leu Phe Leu Val Pro Thr Glu Ile Ala Leu Ala Gln His Arg			
415	420	425	
gcc aat gcc aag agt gcc aag acc tac gcc tac ctg ttt tcc cat ccc			1348
Ala Asn Ala Lys Ser Ala Lys Thr Tyr Ala Tyr Leu Phe Ser His Pro			
430	435	440	
tct cgg atg ccc gtc tac ccc aaa tgg gtg ggg gcc gac cat gca gat			1396
Ser Arg Met Pro Val Tyr Pro Lys Trp Val Gly Ala Asp His Ala Asp			
445	450	455	460
gac att cag tac gtt ttc ggg aag ccc ttc gcc acc ccc acg ggc tac			1444
Asp Ile Gln Tyr Val Phe Gly Lys Pro Phe Ala Thr Pro Thr Gly Tyr			
465	470	475	
cgg ccc caa gac agg aca gtc tct aag gcc atg atc gcc tac tgg acc			1492
Arg Pro Gln Asp Arg Thr Val Ser Lys Ala Met Ile Ala Tyr Trp Thr			
480	485	490	
aac ttt gcc aaa aca ggg gac ccc aac atg ggc gac tcg gct gtg ccc			1540
Asn Phe Ala Lys Thr Gly Asp Pro Asn Met Gly Asp Ser Ala Val Pro			
495	500	505	
aca cac tgg gaa ccc tac act acg gaa aac agc ggc tac ctg gag atc			1588
Thr His Trp Glu Pro Tyr Thr Thr Glu Asn Ser Gly Tyr Leu Glu Ile			
510	515	520	
acc aag aag atg ggc agc agc tcc atg aag cgg agc ctg aga acc aac			1636
Thr Lys Lys Met Gly Ser Ser Ser Met Lys Arg Ser Leu Arg Thr Asn			
525	530	535	540
ttc ctg cgc tac tgg acc ctc acc tat ctg gcg ctg ccc aca gtg acc			1684
Phe Leu Arg Tyr Trp Thr Leu Thr Tyr Leu Ala Leu Pro Thr Val Thr			
545	550	555	
gac cag gag gcc acc cct gtg ccc ccc aca ggg gac tcc gag gcc act			1732
Asp Gln Glu Ala Thr Pro Val Pro Pro Thr Gly Asp Ser Glu Ala Thr			

-continued

560	565	570	
ccc gtg ccc ccc acg ggt gac tcc gag acc gcc ccc gtg ccg ccc acg Pro Val Pro Pro Thr Gly Asp Ser Glu Thr Ala Pro Val Pro Pro Thr 575 580 585			1780
ggt gac tcc ggg gcc ccc ccc gtg ccg ccc acg ggt gac tcc ggg gcc Gly Asp Ser Gly Ala Pro Pro Val Pro Pro Thr Gly Asp Ser Gly Ala 590 595 600			1828
ccc ccc gtg ccg ccc acg ggt gac tcc ggg gcc ccc ccc gtg ccg ccc Pro Pro Val Pro Pro Thr Gly Asp Ser Glu Thr Ala Pro Pro Val Pro Pro 605 610 615 620			1876
acg ggt gac tcc ggg gcc ccc ccc gtg ccg ccc acg ggt gac tcc ggg Thr Gly Asp Ser Gly Ala Pro Pro Val Pro Pro Thr Gly Asp Ser Gly 625 630 635			1924
gcc ccc ccc gtg ccg ccc acg ggt gac tcc ggg gcc ccc ccc gtg ccg Ala Pro Pro Val Pro Pro Thr Gly Asp Ser Glu Thr Ala Pro Pro Val Pro 640 645 650			1972
ccc acg ggt gac tcc ggc gcc ccc ccc gtg ccg ccc acg ggt gac gcc Pro Thr Gly Asp Ser Gly Ala Pro Pro Val Pro Pro Thr Gly Asp Ala 655 660 665			2020
ggg ccc ccc ccc gtg ccg ccc acg ggt gac tcc ggc gcc ccc ccc gtg Gly Pro Pro Pro Val Pro Pro Thr Gly Asp Ser Glu Thr Ala Pro Pro Val 670 675 680			2068
ccg ccc acg ggt gac tcc ggg gcc ccc ccc gtg acc ccc acg ggt gac Pro Pro Thr Gly Asp Ser Gly Ala Pro Pro Val Thr Pro Thr Gly Asp 685 690 695 700			2116
tcc gag acc gcc ccc gtg ccg ccc acg ggt gac tcc ggg gcc ccc cct Ser Glu Thr Ala Pro Val Pro Pro Thr Gly Asp Ser Gly Ala Pro Pro 705 710 715			2164
gtg ccc ccc acg ggt gac tct gag gct gcc cct gtg ccc ccc aca gat Val Pro Pro Thr Gly Asp Ser Glu Ala Ala Pro Val Pro Pro Thr Asp 720 725 730			2212
gac tcc aag gaa gct cag atg cct gca gtc att agg ttt tagcgtccca Asp Ser Lys Glu Ala Gln Met Pro Ala Val Ile Arg Phe 735 740 745			2261
tgagccttg tatcaagagg ccacaagagt gggaccccag gggctcccct cccatcttga			2321
gctcttctg aataaagcct catacccct			2350

<210> SEQ ID NO 7

<211> LENGTH: 745

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

Met Leu Thr Met Gly Arg Leu Gln Leu Val Val Leu Gly Leu Thr Cys 1 5 10 15
Cys Trp Ala Val Ala Ser Ala Ala Lys Leu Gly Ala Val Tyr Thr Glu 20 25 30
Gly Gly Phe Val Glu Gly Val Asn Lys Lys Leu Gly Leu Leu Gly Asp 35 40 45
Ser Val Asp Ile Phe Lys Gly Ile Pro Phe Ala Ala Pro Thr Lys Ala 50 55 60
Leu Glu Asn Pro Gln Pro His Pro Gly Trp Gln Gly Thr Leu Lys Ala 65 70 75 80
Lys Asn Phe Lys Lys Arg Cys Leu Gln Ala Thr Ile Thr Gln Asp Ser 85 90 95
Thr Tyr Gly Asp Glu Asp Cys Leu Tyr Leu Asn Ile Trp Val Pro Gln 100 105 110
Gly Arg Lys Gln Val Ser Arg Asp Leu Pro Val Met Ile Trp Ile Tyr

-continued

115					120					125					
Gly	Gly	Ala	Phe	Leu	Met	Gly	Ser	Gly	His	Gly	Ala	Asn	Phe	Leu	Asn
	130					135					140				
Asn	Tyr	Leu	Tyr	Asp	Gly	Glu	Glu	Ile	Ala	Thr	Arg	Gly	Asn	Val	Ile
145					150					155					160
Val	Val	Thr	Phe	Asn	Tyr	Arg	Val	Gly	Pro	Leu	Gly	Phe	Leu	Ser	Thr
				165					170					175	
Gly	Asp	Ala	Asn	Leu	Pro	Gly	Asn	Tyr	Gly	Leu	Arg	Asp	Gln	His	Met
			180					185					190		
Ala	Ile	Ala	Trp	Val	Lys	Arg	Asn	Ile	Ala	Ala	Phe	Gly	Gly	Asp	Pro
		195					200					205			
Asn	Asn	Ile	Thr	Leu	Phe	Gly	Glu	Ser	Ala	Gly	Gly	Ala	Ser	Val	Ser
	210					215					220				
Leu	Gln	Thr	Leu	Ser	Pro	Tyr	Asn	Lys	Gly	Leu	Ile	Arg	Arg	Ala	Ile
225					230					235					240
Ser	Gln	Ser	Gly	Val	Ala	Leu	Ser	Pro	Trp	Val	Ile	Gln	Lys	Asn	Pro
				245					250					255	
Leu	Phe	Trp	Ala	Lys	Lys	Val	Ala	Glu	Lys	Val	Gly	Cys	Pro	Val	Gly
			260					265					270		
Asp	Ala	Ala	Arg	Met	Ala	Gln	Cys	Leu	Lys	Val	Thr	Asp	Pro	Arg	Ala
		275					280					285			
Leu	Thr	Leu	Ala	Tyr	Lys	Val	Pro	Leu	Ala	Gly	Leu	Glu	Tyr	Pro	Met
	290					295					300				
Leu	His	Tyr	Val	Gly	Phe	Val	Pro	Val	Ile	Asp	Gly	Asp	Phe	Ile	Pro
305					310					315					320
Ala	Asp	Pro	Ile	Asn	Leu	Tyr	Ala	Asn	Ala	Ala	Asp	Ile	Asp	Tyr	Ile
				325				330					335		
Ala	Gly	Thr	Asn	Asn	Met	Asp	Gly	His	Ile	Phe	Ala	Ser	Ile	Asp	Met
			340					345					350		
Pro	Ala	Ile	Asn	Lys	Gly	Asn	Lys	Lys	Val	Thr	Glu	Glu	Asp	Phe	Tyr
		355					360					365			
Lys	Leu	Val	Ser	Glu	Phe	Thr	Ile	Thr	Lys	Gly	Leu	Arg	Gly	Ala	Lys
	370					375					380				
Thr	Thr	Phe	Asp	Val	Tyr	Thr	Glu	Ser	Trp	Ala	Gln	Asp	Pro	Ser	Gln
385					390					395					400
Glu	Asn	Lys	Lys	Lys	Thr	Val	Val	Asp	Phe	Glu	Thr	Asp	Val	Leu	Phe
				405					410					415	
Leu	Val	Pro	Thr	Glu	Ile	Ala	Leu	Ala	Gln	His	Arg	Ala	Asn	Ala	Lys
			420					425					430		
Ser	Ala	Lys	Thr	Tyr	Ala	Tyr	Leu	Phe	Ser	His	Pro	Ser	Arg	Met	Pro
		435					440					445			
Val	Tyr	Pro	Lys	Trp	Val	Gly	Ala	Asp	His	Ala	Asp	Asp	Ile	Gln	Tyr
	450					455					460				
Val	Phe	Gly	Lys	Pro	Phe	Ala	Thr	Pro	Thr	Gly	Tyr	Arg	Pro	Gln	Asp
465					470					475					480
Arg	Thr	Val	Ser	Lys	Ala	Met	Ile	Ala	Tyr	Trp	Thr	Asn	Phe	Ala	Lys
				485					490					495	
Thr	Gly	Asp	Pro	Asn	Met	Gly	Asp	Ser	Ala	Val	Pro	Thr	His	Trp	Glu
			500					505					510		
Pro	Tyr	Thr	Thr	Glu	Asn	Ser	Gly	Tyr	Leu	Glu	Ile	Thr	Lys	Lys	Met
		515					520						525		
Gly	Ser	Ser	Ser	Met	Lys	Arg	Ser	Leu	Arg	Thr	Asn	Phe	Leu	Arg	Tyr
	530					535					540				

-continued

Trp Thr Leu Thr Tyr Leu Ala Leu Pro Thr Val Thr Asp Gln Glu Ala
 545 550 555 560
 Thr Pro Val Pro Pro Thr Gly Asp Ser Glu Ala Thr Pro Val Pro Pro
 565 570 575
 Thr Gly Asp Ser Glu Thr Ala Pro Val Pro Pro Thr Gly Asp Ser Gly
 580 585 590
 Ala Pro Pro Val Pro Pro Thr Gly Asp Ser Gly Ala Pro Pro Val Pro
 595 600 605
 Pro Thr Gly Asp Ser Gly Ala Pro Pro Val Pro Pro Thr Gly Asp Ser
 610 615 620
 Gly Ala Pro Pro Val Pro Pro Thr Gly Asp Ser Gly Ala Pro Pro Val
 625 630 635 640
 Pro Pro Thr Gly Asp Ser Gly Ala Pro Pro Val Pro Pro Thr Gly Asp
 645 650 655
 Ser Gly Ala Pro Pro Val Pro Pro Thr Gly Asp Ala Gly Pro Pro Pro
 660 665 670
 Val Pro Pro Thr Gly Asp Ser Gly Ala Pro Pro Val Pro Pro Thr Gly
 675 680 685
 Asp Ser Gly Ala Pro Pro Val Thr Pro Thr Gly Asp Ser Glu Thr Ala
 690 695 700
 Pro Val Pro Pro Thr Gly Asp Ser Gly Ala Pro Pro Val Pro Pro Thr
 705 710 715 720
 Gly Asp Ser Glu Ala Ala Pro Val Pro Pro Thr Asp Asp Ser Lys Glu
 725 730 735
 Ala Gln Met Pro Ala Val Ile Arg Phe
 740 745

<210> SEQ ID NO 8
 <211> LENGTH: 1083
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(1077)

<400> SEQUENCE: 8

ttt gat gtc tac acc gag tcc tgg gcc cag gac cca tcc cag gag aat 48
 Phe Asp Val Tyr Thr Glu Ser Trp Ala Gln Asp Pro Ser Gln Glu Asn
 1 5 10 15
 aag aag aag act gtg gtg gac ttt gag acc gat gtc ctc ttc ctg gtg 96
 Lys Lys Lys Thr Val Val Asp Phe Glu Thr Asp Val Leu Phe Leu Val
 20 25 30
 ccc acc gag att gcc cta gcc cag cac aga gcc aat gcc aag agt gcc 144
 Pro Thr Glu Ile Ala Leu Ala Gln His Arg Ala Asn Ala Lys Ser Ala
 35 40 45
 aag acc tac gcc tac ctg ttt tcc cat ccc tct cgg atg ccc gtc tac 192
 Lys Thr Tyr Ala Tyr Leu Phe Ser His Pro Ser Arg Met Pro Val Tyr
 50 55 60
 ccc aaa tgg gtg ggg gcc gac cat gca gat gac att cag tac gtt ttc 240
 Pro Lys Trp Val Gly Ala Asp His Ala Asp Asp Ile Gln Tyr Val Phe
 65 70 75 80
 ggg aag ccc ttc gcc acc ccc acg ggc tac cgg ccc caa gac agg aca 288
 Gly Lys Pro Phe Ala Thr Pro Thr Gly Tyr Arg Pro Gln Asp Arg Thr
 85 90 95
 gtc tct aag gcc atg atc gcc tac tgg acc aac ttt gcc aaa aca ggg 336
 Val Ser Lys Ala Met Ile Ala Tyr Trp Thr Asn Phe Ala Lys Thr Gly
 100 105 110
 gac ccc aac atg ggc gac tcg gct gtg ccc aca cac tgg gaa ccc tac 384
 Asp Pro Asn Met Gly Asp Ser Ala Val Pro Thr His Trp Glu Pro Tyr

-continued

115	120	125	
act acg gaa aac agc ggc tac ctg gag atc acc aag aag atg ggc agc Thr Thr Glu Asn Ser Gly Tyr Leu Glu Ile Thr Lys Lys Met Gly Ser 130 135 140			432
agc tcc atg aag cgg agc ctg aga acc aac ttc ctg cgc tac tgg acc Ser Ser Met Lys Arg Ser Leu Arg Thr Asn Phe Leu Arg Tyr Trp Thr 145 150 155 160			480
ctc acc tat ctg gcg ctg ccc aca gtg acc gac cag gag gcc acc cct Leu Thr Tyr Leu Ala Leu Pro Thr Val Thr Asp Gln Glu Ala Thr Pro 165 170 175			528
gtg ccc ccc aca ggg gac tcc gag gcc act ccc gtg ccc ccc acg ggt Val Pro Pro Thr Gly Asp Ser Glu Ala Thr Pro Val Pro Pro Thr Gly 180 185 190			576
gac tcc gag acc gcc ccc gtg ccg ccc acg ggt gac tcc ggg gcc ccc Asp Ser Glu Thr Ala Pro Val Pro Pro Thr Gly Asp Ser Gly Ala Pro 195 200 205			624
ccc gtg ccg ccc acg ggt gac tcc ggg gcc ccc ccc gtg ccg ccc acg Pro Val Pro Pro Thr Gly Asp Ser Gly Ala Pro Pro Val Pro Pro Thr 210 215 220			672
ggt gac tcc ggg gcc ccc ccc gtg ccg ccc acg ggt gac tcc ggg gcc Gly Asp Ser Gly Ala Pro Pro Val Pro Pro Thr Gly Asp Ser Gly Ala 225 230 235 240			720
ccc ccc gtg ccg ccc acg ggt gac tcc ggg gcc ccc ccc gtg ccg ccc Pro Pro Val Pro Pro Thr Gly Asp Ser Gly Ala Pro Pro Val Pro Pro 245 250 255			768
acg ggt gac tcc ggg gcc ccc ccc gtg ccg ccc acg ggt gac tcc ggc Thr Gly Asp Ser Gly Ala Pro Pro Val Pro Pro Thr Gly Asp Ser Gly 260 265 270			816
gcc ccc ccc gtg ccg ccc acg ggt gac gcc ggg ccc ccc ccc gtg ccg Ala Pro Pro Val Pro Pro Thr Gly Asp Ala Gly Pro Pro Pro Val Pro 275 280 285			864
ccc acg ggt gac tcc ggc gcc ccc ccc gtg ccg ccc acg ggt gac tcc Pro Thr Gly Asp Ser Gly Ala Pro Pro Val Pro Pro Thr Gly Asp Ser 290 295 300			912
ggg gcc ccc ccc gtg acc ccc acg ggt gac tcc gag acc gcc ccc gtg Gly Ala Pro Pro Val Thr Pro Thr Gly Asp Ser Glu Thr Ala Pro Val 305 310 315 320			960
ccg ccc acg ggt gac tcc ggg gcc ccc cct gtg ccc ccc acg ggt gac Pro Pro Thr Gly Asp Ser Gly Ala Pro Pro Val Pro Pro Thr Gly Asp 325 330 335			1008
tct gag gct gcc cct gtg ccc ccc aca gat gac tcc aag gaa gct cag Ser Glu Ala Ala Pro Val Pro Pro Thr Asp Asp Ser Lys Glu Ala Gln 340 345 350			1056
atg cct gca gtc att agg ttt tagcgt Met Pro Ala Val Ile Arg Phe 355			1083

<210> SEQ ID NO 9

<211> LENGTH: 359

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

Phe Asp Val Tyr Thr Glu Ser Trp Ala Gln Asp Pro Ser Gln Glu Asn
1 5 10 15

Lys Lys Lys Thr Val Val Asp Phe Glu Thr Asp Val Leu Phe Leu Val
20 25 30

Pro Thr Glu Ile Ala Leu Ala Gln His Arg Ala Asn Ala Lys Ser Ala
35 40 45

Lys Thr Tyr Ala Tyr Leu Phe Ser His Pro Ser Arg Met Pro Val Tyr

-continued

50	55	60
Pro Lys Trp Val Gly Ala Asp His Ala Asp Asp Ile Gln Tyr Val Phe 65 70 75 80		
Gly Lys Pro Phe Ala Thr Pro Thr Gly Tyr Arg Pro Gln Asp Arg Thr 85 90 95		
Val Ser Lys Ala Met Ile Ala Tyr Trp Thr Asn Phe Ala Lys Thr Gly 100 105 110		
Asp Pro Asn Met Gly Asp Ser Ala Val Pro Thr His Trp Glu Pro Tyr 115 120 125		
Thr Thr Glu Asn Ser Gly Tyr Leu Glu Ile Thr Lys Lys Met Gly Ser 130 135 140		
Ser Ser Met Lys Arg Ser Leu Arg Thr Asn Phe Leu Arg Tyr Trp Thr 145 150 155 160		
Leu Thr Tyr Leu Ala Leu Pro Thr Val Thr Asp Gln Glu Ala Thr Pro 165 170 175		
Val Pro Pro Thr Gly Asp Ser Glu Ala Thr Pro Val Pro Pro Thr Gly 180 185 190		
Asp Ser Glu Thr Ala Pro Val Pro Pro Thr Gly Asp Ser Gly Ala Pro 195 200 205		
Pro Val Pro Pro Thr Gly Asp Ser Gly Ala Pro Pro Val Pro Pro Thr 210 215 220		
Gly Asp Ser Gly Ala Pro Pro Val Pro Pro Thr Gly Asp Ser Gly Ala 225 230 235 240		
Pro Pro Val Pro Pro Thr Gly Asp Ser Gly Ala Pro Pro Val Pro Pro 245 250 255		
Thr Gly Asp Ser Gly Ala Pro Pro Val Pro Pro Thr Gly Asp Ser Gly 260 265 270		
Ala Pro Pro Val Pro Pro Thr Gly Asp Ala Gly Pro Pro Pro Val Pro 275 280 285		
Pro Thr Gly Asp Ser Gly Ala Pro Pro Val Pro Pro Thr Gly Asp Ser 290 295 300		
Gly Ala Pro Pro Val Thr Pro Thr Gly Asp Ser Glu Thr Ala Pro Val 305 310 315 320		
Pro Pro Thr Gly Asp Ser Gly Ala Pro Pro Val Pro Pro Thr Gly Asp 325 330 335		
Ser Glu Ala Ala Pro Val Pro Pro Thr Asp Asp Ser Lys Glu Ala Gln 340 345 350		
Met Pro Ala Val Ile Arg Phe 355		

<210> SEQ ID NO 10
 <211> LENGTH: 1839
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(1836)

<400> SEQUENCE: 10

gcg aag ctg ggc gcc gtg tac aca gaa ggt ggg ttc gtg gaa ggc gtc Ala Lys Leu Gly Ala Val Tyr Thr Glu Gly Gly Phe Val Glu Gly Val 1 5 10 15	48
aat aag aag ctc ggc ctc ctg ggt gac tct gtg gac atc ttc aag ggc Asn Lys Lys Leu Gly Leu Leu Gly Asp Ser Val Asp Ile Phe Lys Gly 20 25 30	96
atc ccc ttc gca gct ccc acc aag gcc ctg gaa aat cct cag cca cat Ile Pro Phe Ala Ala Pro Thr Lys Ala Leu Glu Asn Pro Gln Pro His	144

-continued

35	40	45	
cct ggc tgg caa ggg acc	ctg aag gcc aag aac ttc	aag aag aga tgc	192
Pro Gly Trp Gln Gly Thr	Leu Lys Ala Lys Asn Phe	Lys Lys Arg Cys	
50	55	60	
ctg cag gcc acc atc acc	cag gac agc acc tac ggg	gat gaa gac tgc	240
Leu Gln Ala Thr Ile Thr	Gln Asp Ser Thr Tyr Gly	Asp Glu Asp Cys	
65	70	75 80	
ctg tac ctc aac att tgg	gtg ccc cag ggc agg aag	caa gtc tcc cgg	288
Leu Tyr Leu Asn Ile Trp	Val Pro Gln Gly Arg Lys	Gln Val Ser Arg	
85	90	95	
gac ctg ccc gtt atg atc	tgg atc tat gga ggc gcc	ttc ctc atg ggg	336
Asp Leu Pro Val Met Ile	Trp Ile Tyr Gly Gly Ala	Phe Leu Met Gly	
100	105	110	
tcc ggc cat ggg gcc aac	ttc ctc aac aac tac ctg	tat gac ggc gag	384
Ser Gly His Gly Ala Asn	Phe Leu Asn Asn Tyr Leu	Tyr Asp Gly Glu	
115	120	125	
gag atc gcc aca cgc gga	aac gtc atc gtg gtc acc	ttc aac tac cgt	432
Glu Ile Ala Thr Arg Gly	Asn Val Ile Val Val Thr	Phe Asn Tyr Arg	
130	135	140	
gtc ggc ccc ctt ggg ttc	ctc agc act ggg gac gcc	aat ctg cca ggt	480
Val Gly Pro Leu Gly Phe	Leu Ser Thr Gly Asp Ala	Asn Leu Pro Gly	
145	150	155 160	
aac tat ggc ctt cgg gat	cag cac atg gcc att gct	tgg gtg aag agg	528
Asn Tyr Gly Leu Arg Asp	Gln His Met Ala Ile Ala	Trp Val Lys Arg	
165	170	175	
aat atc gcg gcc ttc ggg	ggg gac ccc aac aac atc	acg ctc ttc ggg	576
Asn Ile Ala Ala Phe Gly	Gly Asp Pro Asn Asn Ile	Thr Leu Phe Gly	
180	185	190	
gag tct gct gga ggt gcc	agc gtc tct ctg cag act	ctc tcc ccc tac	624
Glu Ser Ala Gly Gly Ala	Ser Val Ser Leu Gln Thr	Leu Ser Pro Tyr	
195	200	205	
aac aag ggc ctc atc cgg	cga gcc atc agc cag agc	ggc gtg gcc ctg	672
Asn Lys Gly Leu Ile Arg	Arg Ala Ile Ser Gln Ser	Gly Val Ala Leu	
210	215	220	
agt ccc tgg gtc atc cag	aaa aac cca ctc ttc	tgg gcc aaa aag gtg	720
Ser Pro Trp Val Ile Gln	Lys Asn Pro Leu Phe Trp	Ala Lys Lys Val	
225	230	235 240	
gct gag aag gtg ggt tgc	cct gtg ggt gat gcc gcc	agg atg gcc cag	768
Ala Glu Lys Val Gly Cys	Pro Val Gly Asp Ala Ala	Arg Met Ala Gln	
245	250	255	
tgt ctg aag gtt act gat	ccc cga gcc ctg acg ctg	gcc tat aag gtg	816
Cys Leu Lys Val Thr Asp	Pro Arg Ala Leu Thr Leu	Ala Tyr Lys Val	
260	265	270	
ccg ctg gca ggc ctg gag	tac ccc atg ctg cac tat	gtg ggc ttc gtc	864
Pro Leu Ala Gly Leu Glu	Tyr Pro Met Leu His Tyr	Val Gly Phe Val	
275	280	285	
cct gtc att gat gga gac	ttc atc ccc gct gac ccg	atc aac ctg tac	912
Pro Val Ile Asp Gly Asp	Phe Ile Pro Ala Asp Pro	Ile Asn Leu Tyr	
290	295	300	
gcc aac gcc gcc gac atc	gac tat ata gca ggc acc	aac aac atg gac	960
Ala Asn Ala Ala Asp Ile	Asp Tyr Ile Ala Gly Thr	Asn Asn Met Asp	
305	310	315 320	
ggc cac atc ttc gcc agc	atc gac atg cct gcc atc	aac aag ggc aac	1008
Gly His Ile Phe Ala Ser	Ile Asp Met Pro Ala Ile	Asn Lys Gly Asn	
325	330	335	
aag aaa gtc acg gag gag	gac ttc tac aag ctg gtc	agt gag ttc aca	1056
Lys Lys Val Thr Glu Glu	Asp Phe Tyr Lys Leu Val	Ser Glu Phe Thr	
340	345	350	
atc acc aag ggg ctc aga	ggc gcc aag acg acc ttt	gat gtc tac acc	1104
Ile Thr Lys Gly Leu Arg	Gly Ala Lys Thr Thr Phe	Asp Val Tyr Thr	

-continued

355	360	365	
gag tcc tgg gcc cag gac cca tcc cag gag aat aag aag aag act gtg			1152
Glu Ser Trp Ala Gln Asp Pro Ser Gln Glu Asn Lys Lys Lys Thr Val			
370	375	380	
gtg gac ttt gag acc gat gtc ctc ttc ctg gtg ccc acc gag att gcc			1200
Val Asp Phe Glu Thr Asp Val Leu Phe Leu Val Pro Thr Glu Ile Ala			
385	390	395	400
cta gcc cag cac aga gcc aat gcc aag agt gcc aag acc tac gcc tac			1248
Leu Ala Gln His Arg Ala Asn Ala Lys Ser Ala Lys Thr Tyr Ala Tyr			
405	410	415	
ctg ttt tcc cat ccc tct cgg atg ccc gtc tac ccc aaa tgg gtg ggg			1296
Leu Phe Ser His Pro Ser Arg Met Pro Val Tyr Pro Lys Trp Val Gly			
420	425	430	
gcc gac cat gca gat gac att cag tac gtt ttc ggg aag ccc ttc gcc			1344
Ala Asp His Ala Asp Asp Ile Gln Tyr Val Phe Gly Lys Pro Phe Ala			
435	440	445	
acc ccc acg ggc tac cgg ccc caa gac agg aca gtc tct aag gcc atg			1392
Thr Pro Thr Gly Tyr Arg Pro Gln Asp Arg Thr Val Ser Lys Ala Met			
450	455	460	
atc gcc tac tgg acc aac ttt gcc aaa aca ggg gac ccc aac atg ggc			1440
Ile Ala Tyr Trp Thr Asn Phe Ala Lys Thr Gly Asp Pro Asn Met Gly			
465	470	475	480
gac tcg gct gtg ccc aca cac tgg gaa ccc tac act acg gaa aac agc			1488
Asp Ser Ala Val Pro Thr His Trp Glu Pro Tyr Thr Thr Glu Asn Ser			
485	490	495	
ggc tac ctg gag atc acc aag aag atg ggc agc agc tcc atg aag cgg			1536
Gly Tyr Leu Glu Ile Thr Lys Lys Met Gly Ser Ser Ser Met Lys Arg			
500	505	510	
agc ctg aga acc aac ttc ctg cgc tac tgg acc ctc acc tat ctg gcg			1584
Ser Leu Arg Thr Asn Phe Leu Arg Tyr Trp Thr Leu Thr Tyr Leu Ala			
515	520	525	
ctg ccc aca gtg acc gac cag gag gcc acc cct gtg ccc ccc aca ggg			1632
Leu Pro Thr Val Thr Asp Gln Glu Ala Thr Pro Val Pro Pro Thr Gly			
530	535	540	
gac tcc gag gcc act ccc gtg ccc ccc acg ggt gac tcc gag acc gcc			1680
Asp Ser Glu Ala Thr Pro Val Pro Pro Thr Gly Asp Ser Glu Thr Ala			
545	550	555	560
ccc gtg ccg ccc acg ggc gac tcc ggg gcc ccc ccc gtg ccg ccc acg			1728
Pro Val Pro Pro Thr Gly Asp Ser Gly Ala Pro Pro Val Pro Pro Thr			
565	570	575	
ggt gac tcc ggg gcc ccc cct gtg ccc ccc acg ggt gac tct gag gct			1776
Gly Asp Ser Gly Ala Pro Pro Val Pro Pro Thr Gly Asp Ser Glu Ala			
580	585	590	
gcc cct gtg ccc ccc aca ggt gac tcc aag gaa gct cag atg cct gca			1824
Ala Pro Val Pro Pro Thr Gly Asp Ser Lys Glu Ala Gln Met Pro Ala			
595	600	605	
gtc att agg ttt tag			1839
Val Ile Arg Phe			
610			

<210> SEQ ID NO 11

<211> LENGTH: 612

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 11

Ala Lys Leu Gly Ala Val Tyr Thr Glu Gly Gly Phe Val Glu Gly Val
1 5 10 15

Asn Lys Lys Leu Gly Leu Leu Gly Asp Ser Val Asp Ile Phe Lys Gly
20 25 30

-continued

Ile	Pro	Phe	Ala	Ala	Pro	Thr	Lys	Ala	Leu	Glu	Asn	Pro	Gln	Pro	His
		35					40					45			
Pro	Gly	Trp	Gln	Gly	Thr	Leu	Lys	Ala	Lys	Asn	Phe	Lys	Lys	Arg	Cys
	50					55				60					
Leu	Gln	Ala	Thr	Ile	Thr	Gln	Asp	Ser	Thr	Tyr	Gly	Asp	Glu	Asp	Cys
65					70					75					80
Leu	Tyr	Leu	Asn	Ile	Trp	Val	Pro	Gln	Gly	Arg	Lys	Gln	Val	Ser	Arg
				85					90					95	
Asp	Leu	Pro	Val	Met	Ile	Trp	Ile	Tyr	Gly	Gly	Ala	Phe	Leu	Met	Gly
			100					105					110		
Ser	Gly	His	Gly	Ala	Asn	Phe	Leu	Asn	Asn	Tyr	Leu	Tyr	Asp	Gly	Glu
		115					120					125			
Glu	Ile	Ala	Thr	Arg	Gly	Asn	Val	Ile	Val	Val	Thr	Phe	Asn	Tyr	Arg
	130					135					140				
Val	Gly	Pro	Leu	Gly	Phe	Leu	Ser	Thr	Gly	Asp	Ala	Asn	Leu	Pro	Gly
145					150					155					160
Asn	Tyr	Gly	Leu	Arg	Asp	Gln	His	Met	Ala	Ile	Ala	Trp	Val	Lys	Arg
				165					170					175	
Asn	Ile	Ala	Ala	Phe	Gly	Gly	Asp	Pro	Asn	Asn	Ile	Thr	Leu	Phe	Gly
			180					185					190		
Glu	Ser	Ala	Gly	Gly	Ala	Ser	Val	Ser	Leu	Gln	Thr	Leu	Ser	Pro	Tyr
		195					200					205			
Asn	Lys	Gly	Leu	Ile	Arg	Arg	Ala	Ile	Ser	Gln	Ser	Gly	Val	Ala	Leu
	210					215					220				
Ser	Pro	Trp	Val	Ile	Gln	Lys	Asn	Pro	Leu	Phe	Trp	Ala	Lys	Lys	Val
225					230					235					240
Ala	Glu	Lys	Val	Gly	Cys	Pro	Val	Gly	Asp	Ala	Ala	Arg	Met	Ala	Gln
				245					250					255	
Cys	Leu	Lys	Val	Thr	Asp	Pro	Arg	Ala	Leu	Thr	Leu	Ala	Tyr	Lys	Val
			260					265					270		
Pro	Leu	Ala	Gly	Leu	Glu	Tyr	Pro	Met	Leu	His	Tyr	Val	Gly	Phe	Val
		275					280					285			
Pro	Val	Ile	Asp	Gly	Asp	Phe	Ile	Pro	Ala	Asp	Pro	Ile	Asn	Leu	Tyr
	290					295					300				
Ala	Asn	Ala	Ala	Asp	Ile	Asp	Tyr	Ile	Ala	Gly	Thr	Asn	Asn	Met	Asp
305					310					315					320
Gly	His	Ile	Phe	Ala	Ser	Ile	Asp	Met	Pro	Ala	Ile	Asn	Lys	Gly	Asn
				325					330					335	
Lys	Lys	Val	Thr	Glu	Glu	Asp	Phe	Tyr	Lys	Leu	Val	Ser	Glu	Phe	Thr
			340					345					350		
Ile	Thr	Lys	Gly	Leu	Arg	Gly	Ala	Lys	Thr	Thr	Phe	Asp	Val	Tyr	Thr
		355					360					365			
Glu	Ser	Trp	Ala	Gln	Asp	Pro	Ser	Gln	Glu	Asn	Lys	Lys	Lys	Thr	Val
		370				375					380				
Val	Asp	Phe	Glu	Thr	Asp	Val	Leu	Phe	Leu	Val	Pro	Thr	Glu	Ile	Ala
385					390					395					400
Leu	Ala	Gln	His	Arg	Ala	Asn	Ala	Lys	Ser	Ala	Lys	Thr	Tyr	Ala	Tyr
				405					410					415	
Leu	Phe	Ser	His	Pro	Ser	Arg	Met	Pro	Val	Tyr	Pro	Lys	Trp	Val	Gly
			420					425					430		
Ala	Asp	His	Ala	Asp	Asp	Ile	Gln	Tyr	Val	Phe	Gly	Lys	Pro	Phe	Ala
		435					440					445			
Thr	Pro	Thr	Gly	Tyr	Arg	Pro	Gln	Asp	Arg	Thr	Val	Ser	Lys	Ala	Met
						455					460				

-continued

Ile Ala Tyr Trp Thr Asn Phe Ala Lys Thr Gly Asp Pro Asn Met Gly
 465 470 475 480

Asp Ser Ala Val Pro Thr His Trp Glu Pro Tyr Thr Thr Glu Asn Ser
 485 490 495

Gly Tyr Leu Glu Ile Thr Lys Lys Met Gly Ser Ser Ser Met Lys Arg
 500 505 510

Ser Leu Arg Thr Asn Phe Leu Arg Tyr Trp Thr Leu Thr Tyr Leu Ala
 515 520 525

Leu Pro Thr Val Thr Asp Gln Glu Ala Thr Pro Val Pro Pro Thr Gly
 530 535 540

Asp Ser Glu Ala Thr Pro Val Pro Pro Thr Gly Asp Ser Glu Thr Ala
 545 550 555 560

Pro Val Pro Pro Thr Gly Asp Ser Gly Ala Pro Pro Val Pro Pro Thr
 565 570 575

Gly Asp Ser Gly Ala Pro Pro Val Pro Pro Thr Gly Asp Ser Glu Ala
 580 585 590

Ala Pro Val Pro Pro Thr Gly Asp Ser Lys Glu Ala Gln Met Pro Ala
 595 600 605

Val Ile Arg Phe
 610

<210> SEQ ID NO 12
 <211> LENGTH: 750
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(747)

<400> SEQUENCE: 12

ttt gat gtc tac acc gag tcc tgg gcc cag gac cca tcc cag gag aat 48
 Phe Asp Val Tyr Thr Glu Ser Trp Ala Gln Asp Pro Ser Gln Glu Asn
 1 5 10 15

aag aag aag act gtg gtg gac ttt gag acc gat gtc ctc ttc ctg gtg 96
 Lys Lys Lys Thr Val Val Asp Phe Glu Thr Asp Val Leu Phe Leu Val
 20 25 30

ccc acc gag att gcc cta gcc cag cac aga gcc aat gcc aag agt gcc 144
 Pro Thr Glu Ile Ala Leu Ala Gln His Arg Ala Asn Ala Lys Ser Ala
 35 40 45

aag acc tac gcc tac ctg ttt tcc cat ccc tct cgg atg ccc gtc tac 192
 Lys Thr Tyr Ala Tyr Leu Phe Ser His Pro Ser Arg Met Pro Val Tyr
 50 55 60

ccc aaa tgg gtg ggg gcc gac cat gca gat gac att cag tac gtt ttc 240
 Pro Lys Trp Val Gly Ala Asp His Ala Asp Asp Ile Gln Tyr Val Phe
 65 70 75 80

ggg aag ccc ttc gcc acc ccc acg ggc tac cgg ccc caa gac agg aca 288
 Gly Lys Pro Phe Ala Thr Pro Thr Gly Tyr Arg Pro Gln Asp Arg Thr
 85 90 95

gtc tct aag gcc atg atc gcc tac tgg acc aac ttt gcc aaa aca ggg 336
 Val Ser Lys Ala Met Ile Ala Tyr Trp Thr Asn Phe Ala Lys Thr Gly
 100 105 110

gac ccc aac atg ggc gac tcg gct gtg ccc aca cac tgg gaa ccc tac 384
 Asp Pro Asn Met Gly Asp Ser Ala Val Pro Thr His Trp Glu Pro Tyr
 115 120 125

act acg gaa aac agc ggc tac ctg gag atc acc aag aag atg ggc agc 432
 Thr Thr Glu Asn Ser Gly Tyr Leu Glu Ile Thr Lys Lys Met Gly Ser
 130 135 140

agc tcc atg aag cgg agc ctg aga acc aac ttc ctg cgc tac tgg acc 480
 Ser Ser Met Lys Arg Ser Leu Arg Thr Asn Phe Leu Arg Tyr Trp Thr

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 8,367,062 B2
APPLICATION NO. : 12/488793
DATED : February 5, 2013
INVENTOR(S) : Dominique Lombardo et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

In the Specification

Column 18,

Line 49, “5'-TTTCGTgaattcACGCTAAAACCTAATGACTGCAGCATCTG-3'” should
read -- 5'-TTTCGTgaattcACGCTAAAACCTAATGACTGCAGGCATCTG-3' --.

Column 20,

Line 13, “0.5 M 62 -mercaptoethanol” should read --0.5 M β -mercaptoethanol--.


Column 25,

Line 27, “(150 pi” should read --(150 μ l--.

Column 31,

Line 20, “mAb26D10=0).” should read --mAb16D10=0).--.

Signed and Sealed this
Twenty-sixth Day of November, 2013



Margaret A. Focarino
Commissioner for Patents of the United States Patent and Trademark Office