

US008568730B2

(12) **United States Patent**
Wilkes(10) **Patent No.:** **US 8,568,730 B2**
(45) **Date of Patent:** **Oct. 29, 2013**(54) **COMPOSITIONS FOR USE IN THE
TREATMENT OF CHRONIC OBSTRUCTIVE
PULMONARY DISEASES AND ASTHMA**(75) Inventor: **David S. Wilkes**, Indianapolis, IN (US)(73) Assignee: **Indiana University Research &
Technology Corporation**, Indianapolis,
IN (US)(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.(21) Appl. No.: **13/265,847**(22) PCT Filed: **Apr. 22, 2010**(86) PCT No.: **PCT/US2010/032007**§ 371 (c)(1),
(2), (4) Date: **Oct. 21, 2011**(87) PCT Pub. No.: **WO2010/124058**PCT Pub. Date: **Oct. 28, 2010**(65) **Prior Publication Data**

US 2012/0052081 A1 Mar. 1, 2012

Related U.S. Application Data(60) Provisional application No. 61/266,048, filed on Dec.
2, 2009, provisional application No. 61/171,705, filed
on Apr. 22, 2009.(51) **Int. Cl.**
A61K 39/00 (2006.01)(52) **U.S. Cl.**
USPC **424/184.1**; 514/12.2; 514/21.2; 514/17.2(58) **Field of Classification Search**
None
See application file for complete search history.(56) **References Cited**

U.S. PATENT DOCUMENTS

3,169,094	A	2/1965	Wretling	
4,237,224	A	12/1980	Cohen et al.	
4,554,101	A	11/1985	Hopp	
4,683,195	A	7/1987	Mullis et al.	
4,683,202	A	7/1987	Mullis	
4,751,180	A	6/1988	Cousens et al.	
4,800,159	A	1/1989	Mullis et al.	
4,883,750	A	11/1989	Whiteley et al.	
4,935,233	A	6/1990	Bell et al.	
5,633,234	A	5/1997	August et al.	
5,837,458	A	11/1998	Minshull et al.	
6,288,118	B1 *	9/2001	Nieman et al.	514/572
7,348,005	B2	3/2008	Wilkes	
7,759,075	B2 *	7/2010	Wilkes et al.	435/7.21
2012/0052081	A1	3/2012	Wilkes	

FOREIGN PATENT DOCUMENTS

EP	320308	6/1989
EP	329822	8/1989
GB	2202328	9/1988

WO	WO 87/06270	10/1987
WO	WO 88/10315	12/1988
WO	WO 89/06700	7/1989
WO	WO 89/09284	10/1989
WO	WO 02/053092	* 10/2002
WO	WO 2007/120947	10/2007
WO	WO/2007120947	* 10/2007

OTHER PUBLICATIONS

Goodnow CC., *Lancet*. Jun. 30, 2001;357(9274):2115-21.*
 Skyler, J.S., et al. *Diabetes Care*. 2005;28:1068-1076.*
 Pozzilli, P., et al. *Diabetol*. 2000;43:1000-1004.*
 Dong, V.M., et al. *Ped. Transplant.*. 1999;161:181-189.*
 Bell, J.J. et al. *J. Immunol*. 2008;180:1508-1516.*
 Kraus, T.A., and Mayer, L. *Curr. Opin. Gastroenterol*. 2005;21:692-696.*
 Schroeder, R.A., et al. *J. Surg. Sci. Res*. 2003;111:109-119.*
 Marketletter, Sep. 13, 1999, 2 pages.*
 International Search Report and Written Opinion for International Application No. PCT/US2010/032007, mailed Oct. 13, 2010.
 International Preliminary Report on Patentability for International Application No. PCT/US2010/032007, dated Oct. 25, 2011.
 Alpan et al., "The Role of Dendritic Cells, B Cells, and M Cells in Gut-Oriented Immune Responses," *J. Immunol*. 166:4843-52, 2001.
 Altschul et al., "Gapped BLAST and PSI-BLAST: a new generation of protein database search programs," *Nucleic Acids Research*, 25:3389-3402, 1997.
 Altschul et al., "Basic Local Alignment Search Tool," *J. Mol. Biol.*, 215:403-410, 1990.
 Bettelli et al., "Reciprocal developmental pathways for the generation of pathogenic effector T_H17 and regulatory T cells," *Nature*, 11:441(7090):235-8, 2006.
 Bitter et al., "Expression and Secretion Vectors for Yeast," *Methods Enzymol.*, 153:516-544, 1987.
 Broglie, R. et al., "Light-Regulated Expression of a Pea Ribulose-1,5-Bisphosphate Carboxylase Small Subunit Gene in Transformed Plant Cells," *Science*, 224:838-843, 1984.
 Burlingham et al., "IL-17-dependent cellular immunity to collagen type V predisposes to obliterative bronchiolitis in human lung transplants," *J. Clin. Invest*. 117:3498-3506, 2007.
 Chen et al., "Peripheral deletion of antigen-reactive T cells in oral tolerance," *Nature*, 376:177-180, 1995.
 Chen et al., "Regulatory T-cell clones induced by oral tolerance: suppression of autoimmune encephalomyelitis," *Science*, 265:1237-1240 1994.
 Chiang et al., "Type V(A-B) collagen induces platelet aggregation," *J. Lab. Clin. Med.*, 95:99-107, 1980.
 Colbere-Garapin, F. et al, "A New Dominant Hybrid Selective Marker for Higher Eukaryotic Cells," *J. Mol. Biol.*, 150:1-14, 1981.
 Coruzzi, G. et al., "Tissue specific and light-regulated expression of a pea nuclear gene encoding the small subunit of ribulose-1,5-bisphosphate carboxylase," *Embo J*. 3:1671-1680, 1984.
 Cremer et al., "Type XI collagen-induced arthritis in the Lewis rat: characterization of cellular and humoral immune responses to native types XI, V, and II collagen and constituent α -chains," *J. Immunol*. 153:824-832, 1994.
 Danzer et al., "Cytokine interactions in human mixed lymphocyte culture," *Transplantation*, 57(11):1638-1642, 1994.

(Continued)

Primary Examiner — Maher Haddad
 (74) *Attorney, Agent, or Firm* — Cooley LLP

(57) **ABSTRACT**

The present invention provides compounds and methods for treating or preventing pulmonary diseases include COPD and asthma. In particular, the present invention provides for compounds comprising type V collagen, or tolerizing fragments thereof, for the treatment of COPD and asthma.

7 Claims, 6 Drawing Sheets

(56)

References Cited

OTHER PUBLICATIONS

- Dayhoff, M.O. (ed.) Atlas of Protein Sequence and Structure, National Biomedical Research Foundation, Washington DC, 5(3):345-358, 1978.
- Deavin et al., "Statistical Comparison of Established T-Cell Epitope Predictors Against a Large Database of Human and Murine Antigens," *Mol. Immunol.* 33(2):145-155, 1996.
- DeMeester et al., "The Bimodal Expression of Tumor Necrosis Factor- α in Association with Rat Lung Reimplantation and Allograft Rejection," *J. Immunol.*, 150(6):2494-2505, 1993.
- Engelhard, E. K. et al., "The insect tracheal system: A conduit for the systemic spread of *Autographa californica* M nuclear polyhedrosis virus," *Proc. Natl. Acad. Sci.*, 91:3224-3227, 1994.
- Faria et al., "Oral tolerance: mechanisms and therapeutic applications," *Adv. Immunol.*, 73:153-264, 1999.
- Fedoseyeva et al., "De Novo Autoimmunity to Cardiac Myosin After Heart Transplantation and Its Contribution to the Rejection Process," *J. Immunol.*, 162:6836-42, 1999.
- Fouser et al., "Th17 cytokines and their emerging roles in inflammation and autoimmunity," *Immunol Rev.*, 226:87-102, 2008.
- Garrovillo et al., "Indirect allorecognition in acquired thymic tolerance: induction of donor-specific tolerance to rat cardiac allografts by allopeptide-pulsed host dendritic cells," *Transplantation*, 68:1827-1834, 1999.
- Hancock et al., "Oral, but not intravenous, alloantigen prevents accelerated allograft rejection by selective intragraft Th2 cell activation," *Transplantation*, 55:1112-1118, 1993.
- Hanson et al., "The human $\alpha 2(XI)$ collagen gene (COL11A2) maps to the centromeric of the major histocompatibility complex on chromosome 6," *Genomics*, 5:925-931, 1989.
- Hartman, S. C. And Mulligan, R. C. "Two dominant-acting selectable markers for gene transfer studies in mammalian cells," *Proc. Natl. Acad. Sci.*, 85:8047-51, 1988.
- Hein, J., "Unified Approach to Alignment and Phylogenesis," *Methods in Enzymology*, 183:626-645, Academic Press, Inc., San Diego, CA, 1990.
- Henikoff et al., "Amino Acid substitution matrices from protein blocks," *Proc. Natl. Acad. Sci.*, 89:10915, 1992.
- Higgins, D.G. And Sharp, P.M., "Fast and sensitive multiple sequence alignments on a microcomputer," *CABIOS*, 5:151-153, 1989.
- Hirt et al., "Development of obliterative bronchiolitis after allogeneic rat lung transplantation: Implication of acute rejection and the time point of treatment," *J. Heart Lung Transplant.*, 18:542-548, 1999.
- Hobbs, S. or Murry, L. E. in McGraw Hill Yearbook of Science and Technology, McGraw Hill, New York, N.Y.; pp. 191-196, 1992.
- Horn, T. et al., "Synthesis of oligonucleotides on cellulose. Part II: design and synthetic strategy to the synthesis of 22 oligodeoxynucleotides coding for Gastric Inhibitory Polypeptide (GIP)," *Nucl. Acids Res. Symp. Ser.*, 7:225-232, 1980.
- Huang et al.; "Stable mixed chimerism and tolerance using a nonmyeloablative preparative regimen in a large-animal model," *J. Clin. Invest.*, 105:173-181, 2000.
- Ishido et al., "Induction of donor-specific hyporesponsiveness and prolongation of cardiac allograft survival by jejunal administration of donor splenocytes," *Transplantation*, 68:1377-1382, 1999.
- Iyer et al., "Characterization and biologic significance of immunosuppressive peptide D2702.75-84(E α V) binding protein," *J. Bio. Chem.*, 273(5):2692-2697, 1998.
- Jeffery, P. K., "Remodling in asthma and chronic obstructive lung disease," *American Journal of Respiratory and Critical Care Medicine*, 164(10-Pt.2):S28-S38, 2001.
- Joo et al., "T-cell mediated responses in a murine model of orthotopic corneal transplantation," *Invest. Ophthalmol. Vis. Sci.*, 36:1530-1540, 1995.
- Kang, et al., "Cutting Edge: Immunosuppressant as Adjuvant for Tolerogenic Immunization," *J. Immunol.*, 180: 5172-5176, 2008.
- Kapp J. A. and Bucy R. P., "CD8⁺ suppressor T cells resurrected," *Hum Immunol.*, 69(11):715-20, 2008.
- Konomi et al., "Localization of Type V Collagen and Type IV Collagen in Human Cornea, Lung, and Skin," *Am. J. Pathol.*, 116:417-426, 1984.
- Krensky et al., "HLA-derived peptides as novel immunosuppressives," *Nephrol. Dial. Transplant.*, 12:865-878, 1997.
- Kroll, D. J. et al., "A Multifunctional Prokaryotic Protein Expression System: Overproduction, Affinity Purification, and Selective Detection," *DNA and Cell Biol.*, 12:441-453, 1993.
- Logan, J. And Shenk, T., "Adenovirus tripartite leader sequence enhances of mRNAs late after infection," *Proc. Natl. Acad. Sci.*, 81:3655-3659, 1984.
- Lowry et al., "Immune Mechanisms in Organ Allograft Rejection. VI. Delayed-Type Hypersensitivity and Lymphotoxin in Experimental Renal Allograft Rejection," *Transplantation*, 40:183-188, 1985.
- Lowy, I., "Isolation of Transforming DNA: Cloning the Hamster aprt Gene," *Cell*, 22:817-23, 1980.
- Maddox, D. E. et al., "Elevated Serum Levels in Human Pregnancy of a Molecule Immunochemically Similar to Eosinophil Granule Major Basic Protein," *J. Exp. Med.*, 158:1211-1216, 1983.
- Madri et al., "Isolation and Tissue Localization of Type AB Collagen From Normal Lung Parenchyma Human Pathology," 11:353-366, 1980.
- Madri et al., "Collagen Polymorphism in the Lung, An Immunochemical Study of Pulmonary Fibrosis," *Am. J. Pathol.*, 94:323-332, 1979.
- Marck et al., "Lung Transplantation in the Rat. III. Functional Studies in Iso- and Allografts," *J. Surgical Res.*, 35:149-158, 1983.
- Matsumura et al., "Assessment of Pathological Changes Associated with Chronic Allograft Rejection and Tolerance in Two Experimental Models of Rat Lung Transplantation," *Transplantation*, 59:1509-1517, 1995.
- Merrifield, "Solid Phase Peptide Synthesis, I. The Synthesis of a Tetrapeptide," *J. Am. Chem. Soc.*, 85:2149-2154, 1963.
- Mizobuchi et al., "Differential Expression of Smad7 Transcripts Identifies the CD4+CD45RC^{high} Regulatory T Cells That Mediate Type V Collagen-Induced Tolerance to Lung Allografts," *J. Immunol.* 171:1140-1147, 2003.
- Moore, "Update in Asthma 2007," *Am J Respir Crit Care Med.*, 177(10):1068-1073, 2008.
- Morphy et a., "Matrix Metalloproteinase Inhibitors: Current Status," *Current Medicinal Chemistry*, 2:743-762, 1995.
- Morris et al., "Type XI Collagen is a Heterotrimer with the Composition (1 α , 2 α , 3 α) Retaining Non-triple-helical Domains," *J. Biological Chem.*, 262:11345-11350, 1987.
- Murphy et al., "Inhibition of allorecognition by a human class II MHC-derived peptide through the induction of apoptosis," *J. Clin. Invest.*, 103:859-867, 1999.
- Murphy et al.; Genetic construction, expression, and melanoma-selective cytotoxicity of a diphtheria toxin-related a-melanocyte-stimulating hormone fusion protein, *Proc. Natl. Acad. Sci.*, 83:8258-8262, 1986.
- Myers, E.W. And Muller W., "Optimal alignments in linear space," *CABIOS*, 4:(1):11-17, 1988.
- Needleman et al., "A General Method Applicable to the Search for Similarities in the Amino Acid Sequence of Two Proteins," *J. Mol. Biol.*, 48:443-453, 1970.
- Nöbner et al., "HLA-derived Peptides which Inhibit T Cell Function Bind to Members of the Heat-Shock Protein 70 Family," *J. Exp. Med.*, 183:339-348, 1996.
- Oluwole et al., "Induction of Transplantation Intolerance to Rat Cardiac Allografts by Intrathymic Inoculation of Allogeneic Soluble Peptides," *Transplantation*, 56(6):1523-1527, 1993.
- Palmans, E. et al., "Repeated allergen exposure changes collagen composition in airways of sensitised Brown Norway rats," *The European Respiratory Journal*, 20(2):280-285, 2002.
- Parker et al., "Scheme for Ranking Potential HLA-A2 Binding Peptides Based on Independent Binding of Individual Peptide Side-Chains," *J. Immunol.*, 152:163-175, 1994.
- Pearson et al., "Improved tools for biological sequence comparison," *Proc. Natl. Acad. Sci.*, 85:2444-2448, 1988.
- Porath, J. et al., "Immobilized Metal Ion Affinity Chromatography," *Prot. Exp. Purif.* 3:263-281, 1992.

(56)

References Cited

OTHER PUBLICATIONS

- Porter et al., "Recent developments in matrix metalloproteinase inhibitors," *Exp. Opin. Ther. Patents*, 5(12):1287-1296, 1995.
- Prop et al., "Lung Allograft Rejection in the Rat. I. Accelerated Rejection Caused by Graft Lymphocytes," *Transplantation*, 40:25-30, 1985.
- Prop et al., "Lung Allograft Rejection in the Rat. II. Specific immunological Properties of Lung Grafts," *Transplantation*, 40(2):126-131, 1985.
- Rammensee et al., "MHC ligands and peptide motifs: first listing" *Immunogenetics*, 41:178-228, 1995.
- Rhodes, C. A. et al., "Transformation of Maize by Electroporation of Embryos," *Methods Mol. Biol.* 55:121-131, 1995.
- Roberge, J. Y. et al., "A Strategy for a Convergent Synthesis of N-Linked Glycopeptides on a Solid Support," *Science*, 269:202-204, 1995.
- Robinson, "Comparison of Labeled Trees with Valency Three," *J. Comb. Theory*, 11:105-119, 1971.
- Rothbard et al., "A sequence pattern common to T cell epitopes," *The EMBO Journal*, 7:93-100, 1988.
- Saitou, N. And Nei, M., "The Neighbor-joining Method: A New Method for Reconstructing Phylogenetic Trees," *Mol. Biol. Evol.*, 4:406-425, 1987.
- Sayegh et al., "Mechanisms of T Cell Recognition of Alloantigen," *Transplantation*, 57(9):1295-1302, 1994.
- Sayegh et al., "Novel immunotherapeutic strategies using MHC derived peptides," *Kidney Int. Suppl.*, 53:S13-20, 1996.
- Sayegh et al., "Induction of immunity and oral tolerance with polymorphic class II major histocompatibility complex allopeptides in the rat," *Proc. Natl. Acad. Sci.*, 89:7762-7766, 1992.
- Sayegh et al., "Down-regulation of the immune response to histocompatibility antigens and prevention of sensitization by skin allografts by orally administered alloantigen," *Transplantation*, 53:163-166, 1992.
- Scharf, D. et al., "Heat Stress Promoters and Transcription Factors," *Results Probl. Cell Differ.*, 20:125-162, 1994.
- Sekine et al., "Role of Passenger Leukocytes in Allograft Rejection—Effect of Depletion Donor Alveolar Macrophages on the Local Production of TNF-alpha, T Helper 1/Helper 2 Cytokines, IgG Subclasses, and Pathology in a Rat Model of Lung Transplantation," *J. Immunol.*, 159:4084-4093, 1997.
- Seyer et al., "Covalent Structure of Collagen: Amino Acid Sequence of Three Cyanogen Bromide-Derived Peptides from Human Alpha 1(V) Collagen Chain," *Arch. Biochem. Biophys.*, 271(1):120-129, 1989.
- Sivasai et al., "Indirect Recognition of donor HLA class I Peptides in Lung Transplant Recipients with Bronchiolitis Obliterans Syndrome," *Transplantation*, 67(8):1094-1098, 1999.
- Smith, Jr. et al., "Interaction of proteoglycans with pericellular (1 alpha, 2 alpha, 3 alpha) collagens of cartilage," *J. Biol. Chem.*, 260:10761-10767, 1985.
- Smith and Waterman, "Comparison of Biosequences Add," *APL. Math* 2:482-489, 1981.
- Strober et al., "Tolerance and Immunity in the Mucosal Immune System," *Res. Immunol.*, 148:489-599, 1997.
- Takamatsu, N., "Expression of bacterial chloramphenicol acetyltransferase gene in tobacco plants mediated by TMV-RNA," *EMBO J.*, 6:307-311, 1987.
- Trulock, "Lung transplantation," *Am. J. Respir. Crit. Care Med.*, 155:789-818, 1997.
- Vanbuskirk et al., "Patterns of allosensitization in allograft recipients: long-term allograft acceptance is associated with active alloantibody production in conjunction with active inhibition of allereactive delayed-type hypersensitivity," *Transplantation*, 65:1115-1123, 1998.
- Van Heeke, G. et al., "Expression of Human Asparagine in *Escherichia coli*," *J. Biol. Chem.*, 264:5503-5509, 1989.
- Vignola et al., "Tissue Remodeling as a Feature of Persistent Asthma," *Journal of Allergy and Clinical Immunology*, 105(6):1041-1053, 2000.
- Vogel et al., "Sensing extracellular matrix: An update on discoidin domain receptor function," *Cellular Signalling*, 18(8):1108-16, 2006.
- Westra et al., "A Paradox in Heart and Lung Rejection," *Transplantation*, 49:826-828, 1990.
- Weiner, "Oral tolerance: immune mechanisms and treatment of autoimmune diseases," *Imm. Today*, 7:335-44, 1997.
- Whitacre et al., "Oral Tolerance in Experimental Autoimmune Encephalomyelitis. III. Evidence for Clonal Anergy," *J. Immunol.*, 147:2155-2163, 1991.
- Wigler, M. et al., "Transformation of mammalian cells with an amplifiable dominant-acting gene," *Proc. Natl. Acad. Sci.*, 77:3567-70, 1980.
- Wigler, M. et al., "Transfer of Purified Herpes Virus Thymidine Kinase Gene to Cultured Mouse Cells," *Cell*, 11:223-32, 1977.
- Wilbur et al., "Rapid similarity searches of nucleic acid and protein data banks," *Proc. Natl. Acad. Sci.*, 80:726-730, 1983.
- Wilkes et al., "Allogeneic Bronchoalveolar Lavage Cells Induce the Histology and Immunology of Lung Allograft Rejection in Recipient Murine Lungs. Role of ICAM-1 on Donor cells," *Transplantation*, 67(6):890-896, 1999.
- Wilkes et al., "Allogeneic Bronchoalveolar Lavage Cells Induce the Histology of Acute Lung Allograft Rejection, and Deposition of IgG2a in Recipient Murine Lungs," *J. Immunol.*, 155:2775-2783, 1995.
- Wilkes et al., "Instillation of allogeneic lung macrophages and dendritic cells cause differential effects on local IFN-gamma production, lymphocytic bronchitis, and vasculitis in recipient murine lungs," *J. Leukoc. Biol.*, 64:578-586, 1998.
- Wilkes et al., "Cell-Mediated Immunity to Collagen V in Lung Transplant Recipients: Correlation with Collagen V Release into BAL fluid," *J. Heart Lung Transplant.*, 20:167, 2001.
- Wilson et al., "Shared amino acid sequences between major histocompatibility complex class II glycoproteins, type XI collagen and *Proteus mirabilis* in rheumatoid arthritis," *Ann. Rheum. Dis.*, 54:216-220, 1995.
- Winter, J. et al., "The Expression of Heat Shock Protein and Cognate Genes During Plant Development," *Results Probl. Cell Differ.*, 17:85-105, 1991.
- Woessner, Jr., "The Determination of Hydroxyproline in Tissue and Protein Samples Containing Small Proportions of this Imino Acid," *Arch. Biochem. Biophys.*, 93:440-447, 1961.
- Yagyu et al., "Comparison of mononuclear cell populations in bronchoalveolar lavage fluid in acute rejection after lung transplantation and Mycoplasma infection in rats," *J. Heart Transplant.*, 9:516-525, 1990.
- Yamagami et al., "Suppression of Allograft Rejection with anti-[alpha][beta] T Cell Receptor Antibody in Rat Corneal Transplantation," *Transplantation*, 67:600-604, 1999.
- Yasufuku et al., "Prevention of Bronchiolitis Obliterans in Rat Lung allografts by Type V Collagen-Induced Oral Tolerance," *Transplantation*, 73:500-505, 2002.
- Yasufuku et al., "Oral Tolerance Induction by Type V Collagen Downregulates Lung Allograft Rejection," *Am. J. Respir. Cell Mol. Biol.*, 25:26-34, 2001.
- Yoshino et al., "Suppression of Antigen-Induced Arthritis in Lewis Rats by Oral Administration of Type II Collagen," *Arthritis Rheum.*, 38:1092-1096, 1995.
- Yousem et al., "Revision of the 1990 working formulation for the classification of pulmonary allograft rejection: Lung rejection study group," *J. Heart Lung Transplant.*, 15:1-15, 1996.
- Zheng et al., "CTLA4 Signals Are Required to Optimally Induce Allograft Tolerance with Combined Donor-Specific Transfusion and Anti-CD154 Monoclonal Antibody Treatment," *J. Immunol.*, 162:4983-4990, 1999.

* cited by examiner

Serum anti-col(V) antibodies are increased in patients with COPD

Circulating col(V)ab in human volunteers




FIG. 1




FIG. 2




FIG. 3

Lung Mononuclear Cell Transcript Levels




FIG. 4A




FIG. 4B

Lung Mononuclear Cell Transcript Levels




FIG. 4C




FIG. 4D

Lung Mononuclear Cell Transcript Levels




FIG. 4E

**COMPOSITIONS FOR USE IN THE
TREATMENT OF CHRONIC OBSTRUCTIVE
PULMONARY DISEASES AND ASTHMA**

CROSS-REFERENCE TO RELATED
APPLICATIONS

This is a U.S. national phase application of International PCT Patent Application No. PCT/US2010/032007, which was filed on Apr. 22, 2010, which claims the benefit under 35 U.S.C. 119(e) of U.S. Provisional Patent Application No. 61/171,705, filed Apr. 22, 2009, and U.S. Provisional Patent Application No. 61/266,048 filed Dec. 2, 2009, which are incorporated herein by reference in their entireties.

STATEMENT REGARDING SEQUENCE
LISTING

The Sequence Listing associated with this application is provided in text format in lieu of a paper copy, and is hereby incorporated by reference into the specification. The name of the text file containing the Sequence Listing is IURT_004_02US_ST25.txt. The text file is 67 KB, was created on Sep. 10, 2012, and is being submitted electronically via EFS-Web.

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to methods for the treatment of chronic obstructive pulmonary disease (COPD) and asthma. In particular the invention relates to the treatment of COPD and asthma by administering type V Collagen (colV) or tolerogenic fragments thereof.

2. Description of the Related Art

Chronic obstructive pulmonary disease (COPD) is a group of diseases of the lungs in which the airways become narrowed. This leads to a limitation of the flow of air to and from the lungs causing shortness of breath. In contrast to asthma, the limitation of airflow is poorly reversible and usually gets progressively worse over time. COPD is also known as chronic obstructive lung disease (COLD), chronic obstructive airway disease (COAD), chronic airflow limitation (CAL) and chronic obstructive respiratory disease. The term "COPD" includes two main conditions—emphysema and chronic obstructive bronchitis.

In emphysema, the walls between many of the air sacs are damaged, causing them to lose their shape and become floppy. This damage also can destroy the walls of the air sacs, leading to fewer and larger air sacs instead of many tiny ones.

In chronic obstructive bronchitis, the lining of the airways is constantly irritated and inflamed. This causes the lining to thicken. Lots of thick mucus forms in the airways, making it hard to breathe.

Most people who have COPD have both emphysema and chronic obstructive bronchitis. Thus, the general term "COPD" is more accurate.

Chronic bronchitis and emphysema are most commonly caused by smoking; approximately 90% of patients with COPD are or were smokers. Although approximately 50% of smokers develop chronic bronchitis, only 15% of smokers develop disabling airflow obstruction. Certain other mammals, particularly horses, suffer from COPD as well.

The airflow obstruction associated with COPD is progressive, may be accompanied by airway hyperreactivity, and may be partially reversible. Non-specific airway hyper-

sponsiveness may also play a role in the development of COPD and may be predictive of an accelerated rate of decline in lung function in smokers.

COPD is a significant cause of death and disability. It is currently the fourth leading cause of death in the United States and Europe. Treatment guidelines advocate early detection and implementation of smoking cessation programs to help reduce morbidity and mortality due to the disease. However, early detection and diagnosis has been difficult for a number of reasons.

COPD takes years to develop and smokers often deny any ill effects from smoking, attributing the early warning signs of increased breathlessness as a sign of age. Similarly, acute episodes of bronchitis often are not recognized by the general practitioner as early signs of COPD. Many patients exhibit features of more than one disease (e.g. chronic bronchitis or asthmatic bronchitis) making precise diagnosis a challenge, particularly in early disease. Also, many patients do not seek medical help until they are experiencing more severe symptoms associated with reduced lung function, such as dyspnea, persistent cough, and sputum production. As a consequence, the vast majority of patients are not diagnosed or treated until they are in a more advanced stage of disease.

Asthma is a heterogeneous disorder of the airways that afflicts millions of people. Airway inflammation, hyperresponsiveness, and obstruction characterize the condition. The disease often causes spasms of the bronchial smooth muscle system, and affects both the upper and lower respiratory tracts. There are several forms of asthma, characterized by varying degrees of severity. Mild asthma, for example, is defined as brief episodes of wheezing, with or without dyspnea or cough. Moderately severe asthma is defined as wheezing and dyspnea, and can be with or without cough and expectoration, but generally interferes with daily activities and/or sleeping. Severe asthma is characterized by incapacitation due to dyspnea, and the afflicted patient typically is unable to eat or sleep normally, is very anxious, and is often exhausted. A condition known as status asthmaticus is the most severe form of asthma, and generally requires intensive hospital care, and may even prove fatal. The disease may occur as a result of both allergic and nonallergic mechanisms.

While there are several treatments available for relieving the symptoms and discomfort associated with asthma, there are no cures. Moreover, the current treatments often cause side effects that exacerbate the discomfort and precipitate other debilitating conditions. Mild asthma generally is treated with beta-adrenergic drugs, as well as antihistamines, especially in the case of children, to prevent or abort sporadic episodes. Moderately severe and severe asthma are generally treated with adrenergic agents and bronchodilators, as well as corticosteroids. Other actions caused by antiasthmatic agents which limit their widespread use include headache, fatigue, dry mouth, nervousness, and in some cases addiction and substance abuse. Recent advances in the understanding of the pathogenesis and treatment of asthma is discussed more fully in *Am J Respir Crit. Care Med.* 2008 May 15; 177(10):1068-73.

Because asthma is so prevalent in both children and adults, there is an ongoing need for agents that can treat the disease, or at least relieve the symptoms that accompany the disease, without causing undesirable side effects. Likewise, there is an ongoing need for compositions and methods for treating COPD. The present invention provides compositions and methods for the treatment of COPD and asthma and other advantages as described in the detailed description.

BRIEF SUMMARY OF THE INVENTION

One aspect of the present invention provides a method for treating chronic obstructive pulmonary disease comprising

administering to a COPD patient a therapeutically effective amount of type V collagen, or a tolerogenic fragment thereof. In one embodiment of the methods described herein, the COPD patient has emphysema and/or chronic obstructive bronchitis. In another embodiment of the invention, the type V collagen or tolerogenic fragment thereof is administered orally, and may be administered at a dose of between 0.1 mg and 0.5 mg. In a further embodiment, the type V collagen or tolerogenic fragment thereof is administered intravenously, by intrapulmonary instillation, by inhalation, or intramuscularly. In certain embodiments, a combination of different routes may also be used. In yet another embodiment of the invention, the methods may further comprise administering to the COPD patient a bronchodilator, a corticosteroid of other known treatment for COPD.

Another aspect of the invention provides a method for treating asthma comprising administering to an asthma patient a therapeutically effective amount of type V collagen, or a tolerogenic fragment thereof. In one embodiment of the methods, the type V collagen or tolerogenic fragment thereof is administered orally, and may be administered at a dose of between 0.1 mg and 0.5 mg. In further embodiments, the type V collagen or tolerogenic fragment thereof is administered intravenously, by intrapulmonary instillation, by inhalation, or intramuscularly. In a yet a further embodiment, the method further comprises administering to the asthma patient a corticosteroid, a bronchodilator and/or a leukotriene modifier, or other known treatment for asthma.

Another aspect of the invention provides a method for preventing the development of chronic obstructive pulmonary disease in a subject at risk for developing chronic obstructive pulmonary disease comprising administering to the subject a therapeutically effective amount of type V collagen, or a tolerogenic fragment thereof. In one embodiment, the type V collagen or tolerogenic fragment thereof is administered orally and may be administered at a dose of between 0.1 mg and 0.5 mg. In further embodiments, the type V collagen or tolerogenic fragment thereof is administered intravenously, by intrapulmonary instillation, by inhalation, or intramuscularly and may be administered by a combination of these routes.

A further aspect of the invention provides a method for preventing the development or worsening of asthma in a subject at risk for developing asthma comprising administering to the subject a therapeutically effective amount of type V collagen, or a tolerogenic fragment thereof. In one embodiment, the type V collagen or tolerogenic fragment thereof is administered orally and in certain embodiments, may be administered at a dose of between 0.1 mg and 0.5 mg. In a further embodiment, the type V collagen or tolerogenic fragment thereof is administered intravenously, by intrapulmonary instillation, by inhalation, intramuscularly, or by a combination of one or more of these routes.

One aspect of the present invention provides a method for identifying a COPD or asthma patient as a candidate for collagen V tolerance therapy comprising, contacting at least a portion of a sample of blood from the patient with collagen V or an antigenic fragment thereof; and measuring the level of antibodies that bind to the collagen V or antigenic fragment thereof (i.e., measuring the level of Type V collagen-specific antibodies); wherein the presence of antibodies bound to the collagen V is indicative of COPD or asthma. In this regard, collagen V-specific antibody level may be used in conjunction with other clinical factors as described herein in diagnosing COPD or asthma. In one embodiment, the collagen V or antigenic fragment thereof is conjugated to a bead. In a further embodiment, the measuring comprises contacting the

antibodies that bind to the collagen V or antigenic fragment thereof with a fluorescently labeled anti-IgG antibody; and detecting by flow cytometry the amount of fluorescently labeled anti-IgG antibody bound to the antibodies bound to the collagen V.

Another aspect of the present invention provides a method for identifying an individual at risk for developing COPD or asthma, comprising: contacting at least a portion of a sample of blood from the individual with collagen V or an antigenic fragment thereof; and measuring the level of antibodies that bind to the collagen V or an antigenic fragment thereof (i.e., measuring the level of Type V collagen-specific antibodies); wherein the presence of antibodies that bind to the collagen V is associated with a higher risk than would be expected in an individual with no antibodies that bind to the collagen V. In one embodiment, the collagen V or antigenic fragment thereof is conjugated to a bead. In another embodiment, the measuring comprises contacting the antibodies that bind to the collagen V or antigenic fragment thereof with a fluorescently labeled anti-IgG antibody; and detecting by flow cytometry the amount of fluorescently labeled anti-IgG antibody bound to the antibodies bound to the collagen V.

In certain embodiments of the methods for diagnosing or measuring risk for developing COPD or asthma, the anti-IgG antibody used in the methods detects all IgG subtypes. In further embodiments, the anti-IgG antibody specifically detects the IgG1 subtype, or the IgG2 subtype, or the IgG3 subtype, or the IgG4 subtype. In this regard, a switch from one subtype to another subtype may occur during the course of disease and may indicate worsening of disease. Therefore, an increase in one subtype over time may indicate worsening of disease.

A further aspect of the invention provides a method for monitoring the progression of COPD or asthma in an individual comprising, contacting at least a portion of a first sample of blood from the individual with collagen V or an antigenic fragment thereof; measuring the level of antibodies that bind to the collagen V or antigenic fragment thereof in the first sample of blood; contacting at least a portion of a second sample of blood from the individual taken at a later time point, with collagen V or an antigenic fragment thereof; measuring the level of antibodies that bind to the collagen V or antigenic fragment thereof in the second sample of blood; and comparing the level of antibodies that bind to the collagen V or antigenic fragment thereof in the second sample of blood to the level of antibodies that bind to the collagen V or antigenic fragment thereof in the first sample of blood; wherein an increase in the level of antibodies bound to the collagen V in the second sample as compared to the first sample is indicative of worsening of COPD or asthma and a decrease in the level of antibodies bound to the collagen V in the second sample as compared to the first sample is indicative of amelioration of COPD or asthma. Other clinical indicators of COPD and asthma may be used in conjunction with the methods provided herein. In certain embodiments, an increase in anti-collagen V antibodies of a particular IgG subtype (e.g., IgG1, IgG2, IgG3, or IgG4) is indicative of a progression of COPD or asthma. In certain embodiments, the collagen V or antigenic fragment thereof is conjugated to a bead. In another embodiment, the measuring comprises contacting the antibodies that bind to the collagen V or antigenic fragment thereof with a fluorescently labeled anti-IgG antibody; and detecting by flow cytometry the amount of fluorescently labeled anti-IgG antibody bound to the antibodies bound to the collagen V. In certain embodiments of the methods for monitoring progression the anti-IgG antibody detects all IgG

subtypes. In other embodiments, the anti-IgG antibody specifically detects the IgG1, IgG2, IgG3 or IgG4 subtype.

These and other aspects of the invention will be evident upon reference to the following detailed description and attached drawings.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a bar graph showing elevated anti-type V collagen antibodies in COPD patients.

FIG. 2 is a bar graph showing elevated anti-type V collagen antibodies in asthma patients.

FIG. 3 shows prevention of ovalbumin-induced airway hyper-responsiveness in mice following intravenous administration of collagen V. n=5 in all groups except col V where n=2-5 for each data point.

FIGS. 4A-4E are bar graphs showing induction of IFN- γ transcripts in lung mononuclear cells by intravenous administration of col(v). Quantitative PCR was performed for the IL-4, IL-5, IL-13, IFN- γ and IL-10 cytokines shown. Only Col(V) IV induced IFN- γ transcripts in lung mononuclear cells. Data represent lung mononuclear cells of RNA pooled from 5 mice in each group.

DETAILED DESCRIPTION OF THE INVENTION

Type V Collagen

Collagen protein is made of polypeptide chains composed of a repeated sequence of amino acids primarily consisting of hydroxyproline (Hyp), glycine (Gly), and proline (Pro). Collagen is one of the most predominant proteins found in the human body, comprising about 80-85% of the extracellular matrix (ECM) in the dermal layer of normal (non-wounded) skin tissue.

Collagens are classified into several types based on sequence identity and function. Types I, II, & III collagen molecules make up the main fibers of most animal extracellular structures. Type I forms about 90% of the body's collagen and is the primary component of bone, skin and tendons. Type II makes up the major fibers of cartilage. Collagen fibers are arranged in rigid plates in bones, in parallel bundles in tendons, and in a dense meshwork in cartilage. Type I and lesser amounts of type III make up tendons and skin. Type IV collagen molecules make up very fine, unstriated fibers present in basal laminae. Type V Collagen (colV) is a minor collagen present in the lung (Madri and Furthmayr, Human Pathology, 11:353-366, 1980) and is located in the peribronchiolar connective tissues (Madri and Furthmayr, Am. J. Pathol., 94:323-332, 1979), alveolar interstitium (Konomi et al., 1984), and capillary basement membranes (Madri and Furthmayr, 1979, Supra). Over a dozen other collagen types are known but are less well characterized.

Collagen polypeptide chains are characterized by a core helical domain made up of repeating glycine-X-Y triplets and globular N-terminal and C-terminal domains. Three such chains are wound around one another in a superhelix to generate an individual ropelike collagen molecule.

Previous work has demonstrated that autoimmunity to colV is associated with chronic allograft dysfunction (including obliterative bronchiolitis, bronchiolitis obliterans syndrome (BOS)), lung allograft rejection and with risk of developing IPF (see e.g., U.S. Pat. No. 7,348,005 and WO 2007/120947). Furthermore, this work showed that administration of colV induced tolerance to alloantigens and to colV (see e.g., WO 2007/120947; FIG. 10). However, prior to the present invention, no association with autoimmunity to colV had been shown in either asthma or COPD. In fact, previous

findings had shown that patients with COPD do not have anti-col(V) DTH responses which were significantly different from that seen in normal subjects with no known lung disease (see WO 2007/120947, Example 2; FIG. 3). Therefore, the present invention is surprising in that, despite a previous observation that patients with COPD do not have elevated anti-colV DTH responses as compared to normal controls, it has now been found that COPD patients have elevated anti-colV antibodies (see e.g., Example 1, FIG. 1).

Thus, the present invention relates to inducing tolerance to colV in COPD and asthma patients, or in subjects at risk for developing these diseases.

ColV polynucleotide and polypeptide sequences are known to the skilled person and are available in public databases. Illustrative colV polynucleotides and polypeptides of the present invention include, but are not limited to *Homo sapien* collagen, type V, alpha 1 (COL5A1), mRNA NCBI Reference Sequence: NM_000093.3 version GI:89276750 (SEQ ID NO:1); alpha 1 type V collagen preproprotein [*Homo sapiens*]: Accession NP_000084, version GI:89276751 (SEQ ID NO:2); *Homo sapien* collagen, type V, alpha 2 (COL5A2), mRNA; accession NM_000393, version GI:89363016 (SEQ ID NO:3); alpha 2 type V collagen preproprotein [*Homo sapiens*]; accession NP_000384, version GI:89363017 (SEQ ID NO:4); *Homo sapien* collagen, type V, alpha 3 (COL5A3), mRNA; accession NM_015719, NM_015719.3, GI:110735434 (SEQ ID NO:5); collagen, type V, alpha 3 preproprotein [*Homo sapien*]; accession NP_056534, version NP_056534.2, GI:110735435 (SEQ ID NO:6).

As would be recognized by the skilled person, the preprocollagen is processed in a cell into procollagen which is exported from the cell and eventually formed into collagen fibrils and fibers. Thus, the present invention specifically contemplates procollagen and other processed or mature forms of collagen proteins described herein. In this regard, for example, amino acids 1-26 of SEQ ID NO:4 corresponds to the signal peptide which is cleaved during processing, amino acids 27-1229 is the collagen alpha-2(V) chain, and amino acids 1230-1499 correspond to the c-terminal propeptide. These positions within the sequences specifically disclosed herein would be recognized by the skilled person and are available through various public databases where annotation of sequences is provided. It should also be noted that certain amino acids of the collagen proteins are modified during processing (e.g., proline to hydroxyproline). Mature, modified forms of the type V collagen chains, particularly alpha-2 chains, are specifically contemplated herein. As noted elsewhere, the type V collagen and alpha chains thereof for use in the present invention may be purified from a variety of sources or produced recombinantly.

As used herein, the term "polypeptide" is used in its conventional meaning, i.e., as a sequence of amino acids. The polypeptides are not limited to a specific length of the product; thus, peptides, oligopeptides, and proteins are included within the definition of polypeptide, and such terms may be used interchangeably herein unless specifically indicated otherwise. This term also does not refer to or exclude post-expression modifications of the polypeptide, for example, glycosylations, acetylations, phosphorylations and the like, as well as other modifications known in the art, both naturally occurring and non-naturally occurring. A polypeptide may be an entire protein, or a subsequence thereof. Particular polypeptides of interest in the context of this invention are amino acid subsequences comprising tolerogenic fragments.

The present invention, in another aspect, provides polypeptide fragments comprising at least about 5, 10, 15, 20, 25, 50,

or 100 contiguous amino acids, or more, including all intermediate lengths, of a collagen polypeptide compositions set forth herein, such as those set forth in SEQ ID NOs: 2, 4 or 6, or those encoded by a polynucleotide sequence set forth in a sequence of SEQ ID NOs: 1, 3 or 5.

In another aspect, the present invention provides variants of the polypeptide compositions described herein. Polypeptide variants generally encompassed by the present invention will typically exhibit at least about 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% or more identity (determined as described below), along its length, to a polypeptide sequences set forth herein.

In one embodiment, the polypeptide fragments and variants provided by the present invention are immunologically tolerogenic as described herein.

A polypeptide "variant," as the term is used herein, is a polypeptide that typically differs from a polypeptide specifically disclosed herein in one or more substitutions, deletions, additions and/or insertions. Such variants may be naturally occurring or may be synthetically generated, for example, by modifying one or more of the above polypeptide sequences of the invention and evaluating their tolerogenic activity as described herein and/or using any of a number of techniques well known in the art.

In many instances, a variant will contain conservative substitutions. A "conservative substitution" is one in which an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydrophobic nature of the polypeptide to be substantially unchanged. Modifications may be made in the structure of the polynucleotides and polypeptides of the present invention and still obtain a functional molecule that encodes a variant or derivative polypeptide with desirable characteristics, e.g., with tolerogenic characteristics. When it is desired to alter the amino acid sequence of a polypeptide to create an equivalent, or even an improved, tolerogenic variant or portion of a polypeptide of the invention, one skilled in the art will typically change one or more of the codons of the encoding DNA sequence according to Table 1.

For example, certain amino acids may be substituted for other amino acids in a protein structure without appreciable loss of interactive binding capacity with structures such as, for example, antigen-binding regions of antibodies or binding sites on substrate molecules. Since it is the interactive capacity and nature of a protein that defines that protein's biological functional activity, certain amino acid sequence substitutions can be made in a protein sequence, and, of course, its underlying DNA coding sequence, and nevertheless obtain a protein with like properties. It is thus contemplated that various changes may be made in the peptide sequences of the disclosed compositions, or corresponding DNA sequences which encode said peptides without appreciable loss of their tolerogenic utility or activity.

TABLE 1

Amino Acids		Codons	
Alanine	Ala A	GCA GCC GCG GCU	
Cysteine	Cys C	UGC UGU	
Aspartic acid	Asp D	GAC GAU	
Glutamic acid	Glu E	GAA GAG	
Phenylalanine	Phe F	UUC UUU	

TABLE 1-continued

Amino Acids		Codons	
Glycine	Gly G	GGA GGC GGG GGU	
Histidine	His H	CAC CAU	
Isoleucine	Ile I	AUA AUC AUU	
Lysine	Lys K	AAA AAG	
Leucine	Leu L	UUA UUG CUA CUC CUG CUU	
Methionine	Met M	AUG	
Asparagine	Asn N	AAC AAU	
Proline	Pro P	CCA CCC CCG CCU	
Glutamine	Gln Q	CAA CAG	
Arginine	Arg R	AGA AGG CGA CGC CGG CGU	
Serine	Ser S	AGC AGU UCA UCC UCG UCU	
Threonine	Thr T	ACA ACC ACG ACU	
Valine	Val V	GUA GUC GUG GUU	
Tryptophan	Trp W	UGG	
Tyrosine	Tyr Y	UAC UAU	

In making such changes, the hydrophobic index of amino acids may be considered. The importance of the hydrophobic amino acid index in conferring interactive biologic function on a protein is generally understood in the art (Kyte and Doolittle, 1982, incorporated herein by reference). It is accepted that the relative hydrophobic character of the amino acid contributes to the secondary structure of the resultant protein, which in turn defines the interaction of the protein with other molecules, for example, enzymes, substrates, receptors, DNA, antibodies, antigens, and the like. Each amino acid has been assigned a hydrophobic index on the basis of its hydrophobicity and charge characteristics (Kyte and Doolittle, 1982). These values are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (-0.4); threonine (-0.7); serine (-0.8); tryptophan (-0.9); tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate (-3.5); glutamine (-3.5); aspartate (-3.5); asparagine (-3.5); lysine (-3.9); and arginine (-4.5).

It is known in the art that certain amino acids may be substituted by other amino acids having a similar hydrophobic index or score and still result in a protein with similar biological activity, i.e., still obtain a biological functionally equivalent protein. In making such changes, the substitution of amino acids whose hydrophobic indices are within ± 2 is preferred, those within ± 1 are particularly preferred, and those within ± 0.5 are even more particularly preferred. It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity. U.S. Pat. No. 4,554,101 (specifically incorporated herein by reference in its entirety), states that the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with a biological property of the protein.

As detailed in U.S. Pat. No. 4,554,101, the following hydrophilicity values have been assigned to amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0 \pm 1); glutamate (+3.0 \pm 1); serine (+0.3); asparagine (+0.2);

glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5±1); alanine (-0.5); histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5); tryptophan (-3.4). It is understood that an amino acid can be substituted for another having a similar hydrophilicity value and still obtain a biologically equivalent, and in particular, an immunologically equivalent protein. In such changes, the substitution of amino acids whose hydrophilicity values are within ±2 is preferred, those within ±1 are particularly preferred, and those within ±0.5 are even more particularly preferred.

As outlined above, amino acid substitutions are generally therefore based on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size, and the like. Exemplary substitutions that take various of the foregoing characteristics into consideration are well known to those of skill in the art and include: arginine and lysine; glutamate and aspartate; serine and threonine; glutamine and asparagine; and valine, leucine and isoleucine.

In addition, any polynucleotide may be further modified to increase stability in vivo. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends; the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages in the backbone; and/or the inclusion of nontraditional bases such as inosine, queosine and wybutosine, as well as acetyl-methyl-, thio- and other modified forms of adenine, cytidine, guanine, thymine and uridine.

Amino acid substitutions may further be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity and/or the amphipathic nature of the residues. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values include leucine, isoleucine and valine; glycine and alanine; asparagine and glutamine; and serine, threonine, phenylalanine and tyrosine. Other groups of amino acids that may represent conservative changes include: (1) ala, pro, gly, glu, asp, gln, asn, ser, thr; (2) cys, ser, tyr, thr; (3) val, ile, leu, met, ala, phe; (4) lys, arg, his; and (5) phe, tyr, trp, his. A variant may also, or alternatively, contain nonconservative changes. In a preferred embodiment, variant polypeptides differ from a native sequence by substitution, deletion or addition of five amino acids or fewer. Variants may also (or alternatively) be modified by, for example, the deletion or addition of amino acids that have minimal influence on the tolerogenicity, secondary structure and hydrophobic nature of the polypeptide.

As noted above, polypeptides may comprise a signal (or leader) sequence at the N-terminal end of the protein, which co-translationally or post-translationally directs transfer of the protein. The polypeptide may also be conjugated to a linker or other sequence for ease of synthesis, purification or identification of the polypeptide (e.g., poly-His), or to enhance binding of the polypeptide to a solid support. For example, a polypeptide may be conjugated to an immunoglobulin Fc region.

When comparing polypeptide sequences, two sequences are said to be "identical" if the sequence of amino acids in the two sequences is the same when aligned for maximum correspondence, as described below. Comparisons between two sequences are typically performed by comparing the sequences over a comparison window to identify and compare local regions of sequence similarity. A "comparison window" as used herein, refers to a segment of at least about 20 contiguous positions, usually 30 to about 75, 40 to about

50, in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned.

Optimal alignment of sequences for comparison may be conducted using the Megalign program in the Lasergene suite of bioinformatics software (DNASTAR, Inc., Madison, Wis.), using default parameters. This program embodies several alignment schemes described in the following references: Dayhoff, M. O., (1978) *A model of evolutionary change in proteins—Matrices for detecting distant relationships*. In Dayhoff, M. O. (ed.) *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, Washington D.C. Vol. 5, Suppl. 3, pp. 345-358; Hein J. (1990) *Unified Approach to Alignment and Phylogenesis*, pp. 626-645 *Methods in Enzymology* vol. 183, Academic Press, Inc., San Diego, Calif.; Higgins, D. G. and Sharp, P. M., *CABIOS* 5:151-153 (1989); Myers, E. W. and Muller W., *CABIOS* 4:11-17 (1988); Robinson, E. D., *Comb. Theor* 11:105 (1971); Saitou, N. Nei, M., *Mol. Biol. Evol.* 4:406-425 (1987); Sneath, P. H. A. and Sokal, R. R., *Numerical Taxonomy—the Principles and Practice of Numerical Taxonomy*, Freeman Press, San Francisco, Calif. (1973); Wilbur, W. J. and Lipman, D. J., *Proc. Natl. Acad., Sci. USA* 80:726-730 (1983).

Alternatively, optimal alignment of sequences for comparison may be conducted by the local identity algorithm of Smith and Waterman, *Add. APL. Math* 2:482 (1981), by the identity alignment algorithm of Needleman and Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity methods of Pearson and Lipman, *Proc. Natl. Acad. Sci. USA* 85: 2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, Wis.), or by inspection.

One preferred example of algorithms that are suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al., *Nucl. Acids Res.* 25:3389-3402 (1977), and Altschul et al., *J. Mol. Biol.* 215:403-410 (1990), respectively. BLAST and BLAST 2.0 can be used, for example with the parameters described herein, to determine percent sequence identity for the polynucleotides and polypeptides of the invention. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information. For amino acid sequences, a scoring matrix can be used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment.

In one preferred approach, the "percentage of sequence identity" is determined by comparing two optimally aligned sequences over a window of comparison of at least 20 positions, wherein the portion of the polypeptide sequence in the comparison window may comprise additions or deletions (i.e., gaps) of 20 percent or less, usually 5 to 15 percent, or 10 to 12 percent, as compared to the reference sequences (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of

matched positions by the total number of positions in the reference sequence (i.e., the window size) and multiplying the results by 100 to yield the percentage of sequence identity.

Within other illustrative embodiments, a polypeptide may be a fusion polypeptide that comprises multiple polypeptides as described herein, or that comprises at least one polypeptide as described herein and an unrelated sequence, such as a His tag for purification, or a targeting peptide. A fusion partner may, for example, assist in expressing the protein (an expression enhancer) at higher yields than the native recombinant protein. Other fusion partners may be selected so as to increase the solubility of the polypeptide or to enable the polypeptide to be targeted to desired intracellular compartments. Still further fusion partners include affinity tags, which facilitate purification of the polypeptide. In certain embodiments, a fusion partner increases the tolerogenicity of the polypeptide or increases its uptake by cells. In further embodiments, a fusion partner comprises an immune response enhancer.

Fusion polypeptides may generally be prepared using standard techniques, including chemical conjugation. Preferably, a fusion polypeptide is expressed as a recombinant polypeptide, allowing the production of increased levels, relative to a non-fused polypeptide, in an expression system. Briefly, DNA sequences encoding the polypeptide components may be assembled separately, and ligated into an appropriate expression vector. The 3' end of the DNA sequence encoding one polypeptide component is ligated, with or without a peptide linker, to the 5' end of a DNA sequence encoding the second polypeptide component so that the reading frames of the sequences are in phase. This permits translation into a single fusion polypeptide that retains the biological activity of both component polypeptides.

A peptide linker sequence may be employed to separate the first and second polypeptide components by a distance sufficient to ensure that each polypeptide folds into its secondary and tertiary structures. Such a peptide linker sequence is incorporated into the fusion polypeptide using standard techniques well known in the art. Suitable peptide linker sequences may be chosen based on the following factors: (1) their ability to adopt a flexible extended conformation; (2) their inability to adopt a secondary structure that could interact with functional epitopes on the first and second polypeptides; and (3) the lack of hydrophobic or charged residues that might react with the polypeptide functional epitopes. Preferred peptide linker sequences contain Gly, Asn and Ser residues. Other near neutral amino acids, such as Thr and Ala may also be used in the linker sequence. Amino acid sequences which may be usefully employed as linkers include those disclosed in Maratea et al., *Gene* 40:39-46, 1985; Murphy et al., *Proc. Natl. Acad. Sci. USA* 83:8258-8262, 1986; U.S. Pat. No. 4,935,233 and U.S. Pat. No. 4,751,180. The linker sequence may generally be from 1 to about 50 amino acids in length. Linker sequences are not required when the first and second polypeptides have non-essential N-terminal amino acid regions that can be used to separate the functional domains and prevent steric interference.

The ligated DNA sequences are operably linked to suitable transcriptional or translational regulatory elements. The regulatory elements responsible for expression of DNA are located only 5' to the DNA sequence encoding the first polypeptides. Similarly, stop codons required to end translation and transcription termination signals are only present 3' to the DNA sequence encoding the second polypeptide.

One embodiment of the invention involves fusion polypeptides, and the polynucleotides encoding them, wherein the fusion partner comprises a targeting signal capable of direct-

ing a polypeptide to the endosomal/lysosomal compartment, as described in U.S. Pat. No. 5,633,234. A tolerogenic polypeptide of the invention, when fused with this targeting signal, will associate more efficiently with MHC class II molecules and thereby provide enhanced in vivo stimulation of appropriate CD4⁺ T cells specific for the polypeptide.

Polypeptides of the invention are prepared using any of a variety of well known synthetic and/or recombinant techniques, the latter of which are further described below. Polypeptides, portions and other variants generally less than about 150 amino acids can be generated by synthetic means, using techniques well known to those of ordinary skill in the art. In one illustrative example, such polypeptides are synthesized using any of the commercially available solid-phase techniques, such as the Merrifield solid-phase synthesis method, where amino acids are sequentially added to a growing amino acid chain. See Merrifield, *J. Am. Chem. Soc.* 85:2149-2146, 1963. Equipment for automated synthesis of polypeptides is commercially available from suppliers such as Perkin Elmer/Applied BioSystems Division (Foster City, Calif.), and may be operated according to the manufacturer's instructions.

In general, polypeptide compositions (including fusion polypeptides) of the invention are isolated. An "isolated" polypeptide is one that is removed from its original environment. For example, a naturally-occurring protein or polypeptide is isolated if it is separated from some or all of the coexisting materials in the natural system. Preferably, such polypeptides are also purified, e.g., are at least about 90% pure, more preferably at least about 95% pure and most preferably at least about 99% pure.

ColV protein may be purified from a variety of sources or may be purchased from a commercial source (Collaborative Biomedical Products/Becton, Dickinson and Company, Franklin Lakes, N.J. USA). In order to practice some embodiments it may be necessary to obtain pure or partially pure collagen or a tolerogenic fragment, epitope or antigenic portion thereof. These materials for example type V collagen or fragments thereof can be readily obtained by a variety of means including but not limited to animal sources, human cadavers, or recombinant means. Additional methods include partial digests of collagen such as type V collagen. In this regard, Human type V collagen may be extracted from human placenta or other sources and purified by differential NaCl precipitation (Seyer and Kang, 1989). For example, placental tissues are minced, washed, and suspended in 0.5 M acetic acid containing 0.2 M NaCl, and digested by pepsin at 4° C. Supernatants are aspirated from centrifuged specimens, the pellet collected and the extraction procedure repeated. The supernatants are combined from the two digests, and col(V) was purified from the supernatants by differential NaCl precipitation from 0.5 M acetic acid (Smith et al., 1985; Seyer and Kang, 1989). The type V collagen is generally soluble in 0.7 M NaCl and precipitated in 1.2 M NaCl.

For those embodiments where it is required to purify $\alpha(V)$ chains, the cycle of solubilization in acetic acid and NaCl precipitation may be repeated until a type V preparation with an α -chain ratio $\alpha1(V)/\alpha2(V)$ of approximately 2 is obtained as determined by SDS-polyacrylamide gel electrophoresis (Smith et al., 1985), or other appropriate method known to the skilled artisan. Separation of $\alpha1(V)$ from $\alpha2(V)$ may be achieved by chromatography on DEAE-cellulose (Seyer and Kang, 1989) or other methods known to the skilled person, such as those described in Protein Purification Protocols, Ed. Shawn Doonan, Humana Press, 1996). The $\alpha1(V)$ and $\alpha2(V)$ chains may be eluted from the column, and purity confirmed by SDS-polyacrylamide gel electrophoresis as previously

reported (Smith, Jr. et al, 1985). Intact col(V), or $\alpha 1(V)$ and $\alpha 2(V)$ chains may be diluted in PBS (0.5 mg/ml) or other appropriate buffer until use.

The present invention, in certain embodiments, provides polynucleotides encoding the collagen proteins of the present invention. Illustrative polynucleotides are those set forth in SEQ ID NOs:1, 3 and 5, and fragments thereof that encode a tolerogenic fragment of a collagen protein as described herein.

The terms "DNA" and "polynucleotide" are used essentially interchangeably herein to refer to a DNA molecule that has been isolated free of total genomic DNA of a particular species. "Isolated", as used herein, means that a polynucleotide is substantially away from other coding sequences, and that the DNA molecule does not contain large portions of unrelated coding DNA, such as large chromosomal fragments or other functional genes or polypeptide coding regions. Of course, this refers to the DNA molecule as originally isolated, and does not exclude genes or coding regions later added to the segment by the hand of man.

As will be understood by those skilled in the art, the polynucleotide compositions of this invention can include genomic sequences, extra-genomic and plasmid-encoded sequences and smaller engineered gene segments that express, or may be adapted to express, proteins, polypeptides, peptides and the like. Such segments may be naturally isolated, or modified synthetically by the hand of man.

Polynucleotides compositions of the present invention may be identified, prepared and/or manipulated using any of a variety of well established techniques (see generally, Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, N.Y., 1989; Ausubel et al. (2001 *Current Protocols in Molecular Biology*, Greene Publ. Assoc. Inc. & John Wiley & Sons, Inc., NY, N.Y.) and other like references).

Many template dependent processes are available to amplify a target sequences of interest present in a sample. One of the best known amplification methods is the polymerase chain reaction (PCRTM) which is described in detail in U.S. Pat. Nos. 4,683,195, 4,683,202 and 4,800,159, each of which is incorporated herein by reference in its entirety.

Any of a number of other template dependent processes, many of which are variations of the PCRTM amplification technique, are readily known and available in the art. Illustratively, some such methods include the ligase chain reaction (referred to as LCR), described, for example, in Eur. Pat. Appl. Publ. No. 320,308 and U.S. Pat. No. 4,883,750; Qbeta Replicase, described in PCT Intl. Pat. Appl. Publ. No. PCT/US87/00880; Strand Displacement Amplification (SDA) and Repair Chain Reaction (RCR). Still other amplification methods are described in Great Britain Pat. Appl. No. 2 202 328, and in PCT Intl. Pat. Appl. Publ. No. PCT/US89/01025. Other nucleic acid amplification procedures include transcription-based amplification systems (TAS) (PCT Intl. Pat. Appl. Publ. No. WO 88/10315), including nucleic acid sequence based amplification (NASBA) and 3SR. Eur. Pat. Appl. Publ. No. 329,822 describes a nucleic acid amplification process involving cyclically synthesizing single-stranded RNA ("ss-RNA"), ssDNA, and double-stranded DNA (dsDNA). PCT Intl. Pat. Appl. Publ. No. WO 89/06700 describes a nucleic acid sequence amplification scheme based on the hybridization of a promoter/primer sequence to a target single-stranded DNA ("ssDNA") followed by transcription of many RNA copies of the sequence. Other amplification methods such as "RACE" (Frohman, 1990), and "one-sided PCR" (Ohara, 1989) are also well-known to those of skill in the art.

An amplified portion of a polynucleotide of the present invention may be used to isolate a full length gene from a suitable library (e.g., a tumor cDNA library) using well known techniques.

As will be also recognized by the skilled artisan, polynucleotides of the invention may be single-stranded (coding or antisense) or double-stranded, and may be DNA (genomic, cDNA or synthetic) or RNA molecules. RNA molecules may include HnRNA molecules, which contain introns and correspond to a DNA molecule in a one-to-one manner, and mRNA molecules, which do not contain introns. Additional coding or non-coding sequences may, but need not, be present within a polynucleotide of the present invention, and a polynucleotide may, but need not, be linked to other molecules and/or support materials.

Polynucleotides may comprise a native sequence (i.e., an endogenous sequence that encodes a polypeptide/protein of the invention or a portion thereof) or may comprise a sequence that encodes a variant or derivative, preferably a tolerogenic variant or derivative, of such a sequence.

In other related embodiments, the present invention provides polynucleotide variants having substantial identity to the sequences disclosed herein in SEQ ID NOs: 1, 3 and 5, for example those comprising at least 70% sequence identity, preferably at least 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% or higher, sequence identity compared to a polynucleotide sequence of this invention using the methods described herein, (e.g., BLAST analysis using standard parameters, as described below). One skilled in this art will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning and the like.

Typically, polynucleotide variants will contain one or more substitutions, additions, deletions and/or insertions, preferably such that the tolerogenic activity of the polypeptide encoded by the variant polynucleotide is not substantially diminished relative to a polypeptide encoded by a polynucleotide sequence specifically set forth herein. The term "variants" should also be understood to encompass homologous genes of xenogenic origin.

In additional embodiments, the present invention provides polynucleotide fragments comprising or consisting of various lengths of contiguous stretches of sequence identical to or complementary to one or more of the sequences disclosed herein. For example, polynucleotides are provided by this invention that comprise or consist of at least about 10, 15, 20, 30, 40, 50, 75, 100, 150, 200, 300, 400, 500 or 1000 or more contiguous nucleotides of one or more of the sequences disclosed herein as well as all intermediate lengths there between. It will be readily understood that "intermediate lengths", in this context, means any length between the quoted values, such as 16, 17, 18, 19, etc.; 21, 22, 23, etc.; 30, 31, 32, etc.; 50, 51, 52, 53, etc.; 100, 101, 102, 103, etc.; 150, 151, 152, 153, etc.; including all integers through 200-500; 500-1,000, and the like. A polynucleotide sequence as described here may be extended at one or both ends by additional nucleotides not found in the native sequence. This additional sequence may consist of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 nucleotides at either end of the disclosed sequence or at both ends of the disclosed sequence.

In another embodiment of the invention, polynucleotide compositions are provided that are capable of hybridizing under moderate to high stringency conditions to a polynucleotide sequence provided herein, or a fragment thereof, or a

complementary sequence thereof. Hybridization techniques are well known in the art of molecular biology. For purposes of illustration, suitable moderately stringent conditions for testing the hybridization of a polynucleotide of this invention with other polynucleotides include prewashing in a solution of 5×SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0); hybridizing at 50° C.-60° C., 5×SSC, overnight; followed by washing twice at 65° C. for 20 minutes with each of 2×, 0.5× and 0.2×SSC containing 0.1% SDS. One skilled in the art will understand that the stringency of hybridization can be readily manipulated, such as by altering the salt content of the hybridization solution and/or the temperature at which the hybridization is performed. For example, in another embodiment, suitable highly stringent hybridization conditions include those described above, with the exception that the temperature of hybridization is increased, e.g., to 60-65° C. or 65-70° C.

In one embodiment, such polynucleotide variants encode polypeptides that have a level of tolerogenic activity of at least about 50%, preferably at least about 70%, and more preferably at least about 90%, 95%, 96%, 97%, 98%, 99%, or more, of that for a polypeptide sequence specifically set forth herein.

The polynucleotides of the present invention, or fragments thereof, regardless of the length of the coding sequence itself, may be combined with other DNA sequences, such as promoters, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, other coding segments, and the like, such that their overall length may vary considerably. It is therefore contemplated that a nucleic acid fragment of almost any length may be employed, with the total length preferably being limited by the ease of preparation and use in the intended recombinant DNA protocol. For example, illustrative polynucleotide segments with total lengths of about 10,000, about 5000, about 3000, about 2,000, about 1,000, about 500, about 200, about 100, about 50 base pairs in length, and the like, (including all intermediate lengths) are contemplated to be useful in many implementations of this invention.

When comparing polynucleotide sequences, two sequences are said to be “identical” if the sequence of nucleotides in the two sequences is the same when aligned for maximum correspondence, as described below. Comparisons between two sequences are typically performed by comparing the sequences over a comparison window to identify and compare local regions of sequence similarity. A “comparison window” as used herein, refers to a segment of at least about 20 contiguous positions, usually 30 to about 75, 40 to about 50, in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned.

Optimal alignment of sequences for comparison may be conducted using the Megalign program in the Lasergene suite of bioinformatics software (DNASTAR, Inc., Madison, Wis.), using default parameters. This program embodies several alignment schemes described in the following references: Dayhoff, M. O. (1978) A model of evolutionary change in proteins—Matrices for detecting distant relationships. In Dayhoff, M. O. (ed.) *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, Washington D.C. Vol. 5, Suppl. 3, pp. 345-358; Hein J., *Unified Approach to Alignment and Phylogenesis*, pp. 626-645 (1990); *Methods in Enzymology* vol. 183, Academic Press, Inc., San Diego, Calif.; Higgins, D. G. and Sharp, P. M., *CABIOS* 5:151-153 (1989); Myers, E. W. and Muller W., *CABIOS* 4:11-17 (1988); Robinson, E. D., *Comb. Theor* 11:105 (1971); Saitou, N. Nei, M., *Mol. Biol. Evol.* 4:406-425 (1987); Sneath, P. H. A. and Sokal, R. R., *Numerical Tax-*

onomy—the Principles and Practice of Numerical Taxonomy, Freeman Press, San Francisco, Calif. (1973); Wilbur, W. J. and Lipman, D. J., *Proc. Natl. Acad., Sci. USA* 80:726-730 (1983).

Alternatively, optimal alignment of sequences for comparison may be conducted by the local identity algorithm of Smith and Waterman, *Add. APL. Math* 2:482 (1981), by the identity alignment algorithm of Needleman and Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity methods of Pearson and Lipman, *Proc. Natl. Acad. Sci. USA* 85: 2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, Wis.), or by inspection.

One preferred example of algorithms that are suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al., *Nucl. Acids Res.* 25:3389-3402 (1977), and Altschul et al., *J. Mol. Biol.* 215:403-410 (1990), respectively. BLAST and BLAST 2.0 can be used, for example with the parameters described herein, to determine percent sequence identity for the polynucleotides of the invention. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information. In one illustrative example, cumulative scores can be calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always >0) and N (penalty score for mismatching residues; always <0). Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, and expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff and Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915 (1989)) alignments, (B) of 50, expectation (E) of 10, M=5, N=-4 and a comparison of both strands.

Preferably, the “percentage of sequence identity” is determined by comparing two optimally aligned sequences over a window of comparison of at least 20 positions, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) of 20 percent or less, usually 5 to 15 percent, or 10 to 12 percent, as compared to the reference sequences (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid bases occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the reference sequence (i.e., the window size) and multiplying the results by 100 to yield the percentage of sequence identity.

It will be appreciated by those of ordinary skill in the art that, as a result of the degeneracy of the genetic code, there are many nucleotide sequences that encode a polypeptide as described herein. Some of these polynucleotides bear minimal homology to the nucleotide sequence of any native gene. Nonetheless, polynucleotides that vary due to differences in codon usage are specifically contemplated by the present invention. Further, alleles of the genes comprising the polynucleotide sequences provided herein are within the scope of the present invention. Alleles are endogenous genes that are

altered as a result of one or more mutations, such as deletions, additions and/or substitutions of nucleotides. The resulting mRNA and protein may, but need not, have an altered structure or function. Alleles may be identified using standard techniques (such as hybridization, amplification and/or data-

base sequence comparison). Therefore, in another embodiment of the invention, a mutagenesis approach, such as site-specific mutagenesis, is employed for the preparation of tolerogenic variants and/or derivatives of the polypeptides described herein. By this approach, specific modifications in a polypeptide sequence can be made through mutagenesis of the underlying polynucleotides that encode them. These techniques provides a straightforward approach to prepare and test sequence variants, for example, incorporating one or more of the foregoing considerations, by introducing one or more nucleotide sequence changes into the polynucleotide.

Site-specific mutagenesis allows the production of mutants through the use of specific oligonucleotide sequences which encode the DNA sequence of the desired mutation, as well as a sufficient number of adjacent nucleotides, to provide a primer sequence of sufficient size and sequence complexity to form a stable duplex on both sides of the deletion junction being traversed. Mutations may be employed in a selected polynucleotide sequence to improve, alter, decrease, modify, or otherwise change the properties of the polynucleotide itself, and/or alter the properties, activity, composition, stability, or primary sequence of the encoded polypeptide.

In certain embodiments of the present invention, the inventors contemplate the mutagenesis of the disclosed polynucleotide sequences to alter one or more properties of the encoded polypeptide, such as the tolerogenicity of a polypeptide. The techniques of site-specific mutagenesis are well-known in the art, and are widely used to create variants of both polypeptides and polynucleotides. For example, site-specific mutagenesis is often used to alter a specific portion of a DNA molecule. In such embodiments, a primer comprising typically about 14 to about 25 nucleotides or so in length is employed, with about 5 to about 10 residues on both sides of the junction of the sequence being altered.

As will be appreciated by those of skill in the art, site-specific mutagenesis techniques have often employed a phage vector that exists in both a single stranded and double stranded form. Typical vectors useful in site-directed mutagenesis include vectors such as the M13 phage. These phage are readily commercially-available and their use is generally well-known to those skilled in the art. Double-stranded plasmids are also routinely employed in site directed mutagenesis that eliminates the step of transferring the gene of interest from a plasmid to a phage.

In general, site-directed mutagenesis in accordance herewith is performed by first obtaining a single-stranded vector or melting apart of two strands of a double-stranded vector that includes within its sequence a DNA sequence that encodes the desired peptide. An oligonucleotide primer bearing the desired mutated sequence is prepared, generally synthetically. This primer is then annealed with the single-stranded vector, and subjected to DNA polymerizing enzymes such as *E. coli* polymerase I Klenow fragment, in order to complete the synthesis of the mutation-bearing strand. Thus, a heteroduplex is formed wherein one strand encodes the original non-mutated sequence and the second strand bears the desired mutation. This heteroduplex vector is then used to transform appropriate cells, such as *E. coli* cells, and clones are selected which include recombinant vectors bearing the mutated sequence arrangement.

The preparation of sequence variants of the selected peptide-encoding DNA segments using site-directed mutagenesis provides a means of producing potentially useful species and is not meant to be limiting as there are other ways in which sequence variants of peptides and the DNA sequences encoding them may be obtained. For example, recombinant vectors encoding the desired peptide sequence may be treated with mutagenic agents, such as hydroxylamine, to obtain sequence variants. Specific details regarding these methods and protocols are found in the teachings of Maloy et al., 1994; Segal, 1976; Prokop and Bajpai, 1991; Kuby, 1994; and Maniatis et al., 1982, each incorporated herein by reference, for that purpose.

As used herein, the term "oligonucleotide directed mutagenesis procedure" refers to template-dependent processes and vector-mediated propagation which result in an increase in the concentration of a specific nucleic acid molecule relative to its initial concentration, or in an increase in the concentration of a detectable signal, such as amplification. As used herein, the term "oligonucleotide directed mutagenesis procedure" is intended to refer to a process that involves the template-dependent extension of a primer molecule. The term template dependent process refers to nucleic acid synthesis of an RNA or a DNA molecule wherein the sequence of the newly synthesized strand of nucleic acid is dictated by the well-known rules of complementary base pairing (see, for example, Watson, 1987). Typically, vector mediated methodologies involve the introduction of the nucleic acid fragment into a DNA or RNA vector, the clonal amplification of the vector, and the recovery of the amplified nucleic acid fragment. Examples of such methodologies are provided by U.S. Pat. No. 4,237,224, specifically incorporated herein by reference in its entirety.

In another approach for the production of polypeptide variants of the present invention, recursive sequence recombination, as described in U.S. Pat. No. 5,837,458, may be employed. In this approach, iterative cycles of recombination and screening or selection are performed to "evolve" individual polynucleotide variants of the invention having, for example, enhanced tolerogenic activity.

In other embodiments of the invention, polynucleotide sequences or fragments thereof which encode polypeptides of the invention, or fusion proteins or functional equivalents thereof, may be used in recombinant DNA molecules to direct expression of a polypeptide in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences that encode substantially the same or a functionally equivalent amino acid sequence may be produced and these sequences may be used to clone and express a given polypeptide.

As will be understood by those of skill in the art, it may be advantageous in some instances to produce polypeptide-encoding nucleotide sequences possessing non-naturally occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce a recombinant RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

Moreover, the polynucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter polypeptide encoding sequences for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the gene product. For example, DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucle-

otide sequences. In addition, site-directed mutagenesis may be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, or introduce mutations, and so forth.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences may be ligated to a heterologous sequence to encode a fusion protein. For example, to screen peptide libraries for inhibitors of polypeptide activity, it may be useful to encode a chimeric protein that can be recognized by a commercially available antibody. A fusion protein may also be engineered to contain a cleavage site located between the polypeptide-encoding sequence and the heterologous protein sequence, so that the polypeptide may be cleaved and purified away from the heterologous moiety.

Sequences encoding a desired polypeptide may be synthesized, in whole or in part, using chemical methods well known in the art (see Caruthers, M. H. et al. (1980) *Nucl. Acids Res. Symp. Ser.* 215-223, Horn, T. et al. (1980) *Nucl. Acids Res. Symp. Ser.* 225-232). Alternatively, the protein itself may be produced using chemical methods to synthesize the amino acid sequence of a polypeptide, or a portion thereof. For example, peptide synthesis can be performed using various solid-phase techniques (Roberge, J. Y. et al. (1995) *Science* 269:202-204) and automated synthesis may be achieved, for example, using the ABI 431A Peptide Synthesizer (Perkin Elmer, Palo Alto, Calif.).

A newly synthesized peptide may be substantially purified by preparative high performance liquid chromatography (e.g., Creighton, T. (1983) *Proteins, Structures and Molecular Principles*, WH Freeman and Co., New York, N.Y.) or other comparable techniques available in the art. The composition of the synthetic peptides may be confirmed by amino acid analysis or sequencing (e.g., the Edman degradation procedure). Additionally, the amino acid sequence of a polypeptide, or any part thereof, may be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

In order to express a desired polypeptide, the nucleotide sequences encoding the polypeptide, or functional equivalents, may be inserted into appropriate expression vector, i.e., a vector which contains the necessary elements for the transcription and translation of the inserted coding sequence. Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding a polypeptide of interest and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. Such techniques are described, for example, in Sambrook, J. et al. (1989) *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Press, Plainview, N.Y., and Ausubel et al. (2001-2008) *Current Protocols in Molecular Biology*, Greene Publ. Assoc. Inc. & John Wiley & Sons, Inc., NY, N.Y.).

A variety of expression vector/host systems may be utilized to contain and express polynucleotide sequences. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with virus expression vectors (e.g., baculovirus); plant cell systems transformed with virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems.

The “control elements” or “regulatory sequences” present in an expression vector are those non-translated regions of the vector—enhancers, promoters, 5' and 3' untranslated regions—which interact with host cellular proteins to carry out transcription and translation. Such elements may vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the pBLUESCRIPT phagemid (Stratagene, La Jolla, Calif.) or pSPORT1 plasmid (Gibco BRL, Gaithersburg, Md.) and the like may be used. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are generally preferred. If it is necessary to generate a cell line that contains multiple copies of the sequence encoding a polypeptide, vectors based on SV40 or EBV may be advantageously used with an appropriate selectable marker.

In bacterial systems, any of a number of expression vectors may be selected depending upon the use intended for the expressed polypeptide. For example, when large quantities are needed, for example for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified may be used. Such vectors include, but are not limited to, the multifunctional *E. coli* cloning and expression vectors such as pBLUESCRIPT (Stratagene), in which the sequence encoding the polypeptide of interest may be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced; pIN vectors (Van Heeke, G. and S. M. Schuster (1989) *J. Biol. Chem.* 264: 5503-5509); and the like. pGEX Vectors (Promega, Madison, Wis.) may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems may be designed to include heparin, thrombin, or factor XA protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

In the yeast, *Saccharomyces cerevisiae*, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH may be used. For reviews, see Ausubel et al. (supra) and Grant et al. (1987) *Methods Enzymol.* 153:516-544.

In cases where plant expression vectors are used, the expression of sequences encoding polypeptides may be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV may be used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) *EMBO J.* 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used (Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984) *Science* 224:838-843; and Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105). These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. Such techniques are described in a number of generally available reviews (see, for example, Hobbs, S. or Murry, L. E. in McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York, N.Y.; pp. 191-196).

An insect system may also be used to express a polypeptide of interest. For example, in one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a

vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. The sequences encoding the polypeptide may be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of the polypeptide-encoding sequence will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses may then be used to infect, for example, *S. frugiperda* cells or *Trichoplusia* larvae in which the polypeptide of interest may be expressed (Engelhard, E. K. et al. (1994) *Proc. Natl. Acad. Sci.* 91:3224-3227).

In mammalian host cells, a number of viral-based expression systems are generally available. For example, in cases where an adenovirus is used as an expression vector, sequences encoding a polypeptide of interest may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain a viable virus which is capable of expressing the polypeptide in infected host cells (Logan, J. and Shenk, T. (1984) *Proc. Natl. Acad. Sci.* 81:3655-3659). In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells.

Specific initiation signals may also be used to achieve more efficient translation of sequences encoding a polypeptide of interest. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding the polypeptide, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a portion thereof, is inserted, exogenous translational control signals including the ATG initiation codon should be provided. Furthermore, the initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used, such as those described in the literature (Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162).

In addition, a host cell strain may be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be used to facilitate correct insertion, folding and/or function. Different host cells such as CHO, COS, HeLa, MDCK, HEK293, and WI38, which have specific cellular machinery and characteristic mechanisms for such post-translational activities, may be chosen to ensure the correct modification and processing of the foreign protein.

For long-term, high-yield production of recombinant proteins, stable expression is generally preferred. For example, cell lines which stably express a polynucleotide of interest may be transformed using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resis-

tant clones of stably transformed cells may be proliferated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler, M. et al. (1977) *Cell* 11:223-32) and adenine phosphoribosyltransferase (Lowy, I. et al. (1990) *Cell* 22:817-23) genes which can be employed in tk⁻ or aprt⁻ cells, respectively. Also, antimitabolite, antibiotic or herbicide resistance can be used as the basis for selection; for example, dhfr which confers resistance to methotrexate (Wigler, M. et al. (1980) *Proc. Natl. Acad. Sci.* 77:3567-70); npt, which confers resistance to the aminoglycosides, neomycin and G-418 (Colbere-Garapin, F. et al. (1981) *J. Mol. Biol.* 150:1-14); and als or pat, which confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively (Murry, supra). Additional selectable genes have been described, for example, trpB, which allows cells to utilize indole in place of tryptophan, or hisD, which allows cells to utilize histinol in place of histidine (Hartman, S. C. and R. C. Mulligan (1988) *Proc. Natl. Acad. Sci.* 85:8047-51). The use of visible markers has gained popularity with such markers as anthocyanins, beta-glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, being widely used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system (Rhodes, C. A. et al. (1995) *Methods Mol. Biol.* 55:121-131).

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, its presence and expression may need to be confirmed. For example, if the sequence encoding a polypeptide is inserted within a marker gene sequence, recombinant cells containing sequences can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a polypeptide-encoding sequence under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

Alternatively, host cells that contain and express a desired polynucleotide sequence may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassay or immunoassay techniques which include, for example, membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein.

A variety of protocols for detecting and measuring the expression of polynucleotide-encoded products, using either polyclonal or monoclonal antibodies specific for the product are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on a given polypeptide may be preferred for some applications, but a competitive binding assay may also be employed. These and other assays are described, among other places, in Hampton, R. et al. (1990; *Serological Methods, a Laboratory Manual*, APS Press, St Paul, Minn.) and Maddox, D. E. et al. (1983; *J. Exp. Med.* 158:1211-1216).

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides include oligolabeling, nick translation, end-labeling or PCR amplification using a labeled nucleotide. Alternatively, the sequences, or any portions

thereof may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits. Suitable reporter molecules or labels, which may be used include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with a polynucleotide sequence of interest may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a recombinant cell may be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides of the invention may be designed to contain signal sequences which direct secretion of the encoded polypeptide through a prokaryotic or eukaryotic cell membrane. Other recombinant constructions may be used to join sequences encoding a polypeptide of interest to nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Wash.). The inclusion of cleavable linker sequences such as those specific for Factor XA or enterokinase (Invitrogen, San Diego, Calif.) between the purification domain and the encoded polypeptide may be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing a polypeptide of interest and a nucleic acid encoding 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification on IMIAC (immobilized metal ion affinity chromatography) as described in Porath, J. et al. (1992, *Prot. Exp. Purif.* 3:263-281) while the enterokinase cleavage site provides a means for purifying the desired polypeptide from the fusion protein. A discussion of vectors which contain fusion proteins is provided in Kroll, D. J. et al. (1993; *DNA Cell Biol.* 12:441-453).

In addition to recombinant production methods, polypeptides of the invention, and fragments thereof, may be produced by direct peptide synthesis using solid-phase techniques (Merrifield J. (1963) *J. Am. Chem. Soc.* 85:2149-2154). Protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer). Alternatively, various fragments may be chemically synthesized separately and combined using chemical methods to produce the full length molecule.

Tolerance

Immunological tolerance is defined as immune unresponsiveness to an antigen, usually an antigen implicated in causing disease. Although tolerance may be induced by administering antigens by different routes, oral tolerance refers to the oral administration of the antigen, which has resulted in suppression of disease activity in several animal models including experimental autoimmune encephalomyelitis—a rodent model of multiple sclerosis, myasthenia gravis, uveitis, insulin dependent diabetes, and collagen-induced arthritis (Faria and Weiner 1999). Early results from clinical trials in humans suggest that oral tolerance is effective in autoimmune uveitis,

diabetes, nickel allergy, and possibly multiple sclerosis (Faria and Weiner 1999; Duda et al. 2000). There are few studies reporting oral tolerance induction in organ transplantation (Sayegh et al. 1996; Hancock et al. 1993; Ishido et al. 1999; Sayegh et al. 1992a; Sayegh et al. 1992b). In each report, tolerance was induced by feeding donor MHC-derived peptides or feeding allogeneic cells prior to transplantation (Sayegh et al. 1996; Hancock et al. 1993; Ishido et al. 1999; Sayegh et al. 1992a; Sayegh et al. 1992b). These techniques were effective in preventing rejection of cardiac and corneal allografts (Sayegh et al. 1996; Hancock et al. 1993; Ishido et al. 1999; Sayegh et al. 1992a; Sayegh et al. 1992b; Faria and Weiner 1999). In addition to diminished disease activity, immune suppression induced by oral tolerance in these studies was also quantitated by down regulation of delayed type hypersensitivity (DTH) responses to target antigens, as well as diminished cellular and humoral immunity (Faria and Weiner 1999; Mayer 2000; Garside and Mowat 1997).

There are three mechanisms by which oral (and other routes of administration) tolerance down regulates antigen-specific immune responses: 1. active suppression of antigen specific cells, 2. clonal anergy of antigen specific cells, and 3. clonal deletion of antigen specific cells (Faria and Weiner 1999, Miller et al. 1991; Chen et al. 1994; Chen et al. 1995). Although all three mechanisms can be operative simultaneously in response to oral tolerance, active suppression and clonal anergy are the key mechanisms of immune suppression induced by oral tolerance (Faria and Weiner 1999).

Active suppression describes the regulation of one lymphocyte subset by another in an antigen-specific manner. Depending on the antigen and disease state, the suppressor cells may be CD4+ and/or CD8+ T-lymphocytes which migrate from peripheral lymphoid tissues, such as spleen and peripheral lymph nodes, to sites of disease activity. Adoptive transfer of these cells to naive recipients has confirmed the role of these cells in active suppression in rodent models of ovalbumin-induced hypersensitivity, and multiple sclerosis. In vitro evidence of active suppression is demonstrated by data showing that tolerized lymphocytes from animals can suppress proliferation of other antigen-specific T-lymphocytes across a transwell cell culture system (Faria and Weiner 1999; Miller et al. 1991).

Clonal anergy refers to unresponsiveness of antigen-specific T-lymphocytes, which is characterized by diminished proliferation after exposure to an antigen, and is involved in oral tolerance in several animal models. Anergy could be the result of production of soluble suppressive factors by CD4+ or CD8+ T-lymphocytes themselves, other T-lymphocytes or cells in the local environment, or as result of decreased expression of appropriate costimulatory molecules (Faria and Weiner 1999). Clonal deletion refers to the elimination of antigen-specific T-lymphocytes, but has been reported rarely as a mechanism of oral tolerance to an antigen (Chen et al. 1995).

The soluble mediators that suppress the immune response during oral tolerance are derived mainly from regulatory or suppressor T-lymphocytes (Faria and Weiner 1999). There are five types of T-lymphocytes described by the cytokines they produce: Th1-type that produce interleukin-2 (IL-2) and gamma interferon (γ IFN); Th2-type that produce IL-4 and IL-10; Th3-type that produce high levels of transforming growth factor beta (TGF- β), alone, or in conjunction with very low levels of IL-4, IL-10, or γ IFN; Tr1 cells that produce high levels of IL-10 in conjunction with low levels of TGF- β (Faria and Weiner et al. 1999; Mayer 2000; Garside and Mowat 1997; Groux et al. 1997); and Th17 cells that produce IL-17 (see e.g., *Immunol Rev.* 2008, 226:87-102; *Nature.*

2006 May 11; 441(7090):235-8). Since Th3, Th2, and Tr1-T-lymphocytes have been shown to be the major mediators of active suppression induced by oral tolerance, then TGF- β , IL-4 and IL-10 are believed to be key cytokines in this process (Teng et al. 1998; Shi et al. 1999b). A report from Barone et al., and others showing that oral tolerance induction occurred in the absence of these cytokines suggests that other mediators or cells could suppress the immune response (Barone et al. 1998; Shi et al. 1999a).

Although studies of oral tolerance have focused on T-lymphocyte-derived cytokines that suppress immune responses, nitric oxide, which is not produced by T-lymphocytes, is known to be a potent suppressor of alloimmune responses (Garside and Mowat 1997). These data and others showing that nitric oxide modulates apoptosis, which is involved in the rejection response (Meyer et al. 1998; Kallio et al. 1997; Shiraishi et al. 1997; Shiraishi et al. 1995; Medot-Pirenne et al. 1999), suggests that nitric oxide could be a mediator of oral tolerance and prevent the rejection response. TGF- β , is a potent inducer of nitric oxide synthesis, and is a key mediator of active suppression in oral tolerance (Faria and Weiner 1999; Meyer et al. 1998; Vodovotz et al. 1998; Vodovitz et al. 1999). Therefore, immunosuppression induced by TGF- β in the tolerized host could be mediated, in part, by nitric oxide. However, production of nitric oxide in response to oral tolerance is unknown.

Antigen-specific T-lymphocyte activation induced by APCs requires bi-directional interaction between the T-lymphocyte and APC. Initially, APCs present MHC molecules that bind to the T-cell-receptor which stimulates upregulated expression of CD40-ligand (CD40L) on T-lymphocytes. CD40L, in turn, binds to its receptor, CD40, on the APC. Signaling through CD40 induces the expression of CD80 and CD86 on the APC which, upon binding to their receptor, CD28, on the T-lymphocyte, results in co-stimulation and subsequent T-lymphocyte activation (Liu et al. 1999; Li et al. 1999; Lederman and Siciu-Foca 1999). Although studies of oral tolerance induction have focused on T-lymphocyte function, a recent study from Taams et al (1998), reported that tolerance induction may affect function of APCs, with similar data from other investigators (Wu et al. 1998; Finkelman et al. 1996; Viney et al. 1998). For example, a report from Wu, et al (1998), showing that expression of CD80 is decreased on APCs from the lymph nodes and spleens of orally tolerized mice suggests that ineffective APCs could contribute to impaired T-lymphocyte activation in tolerized recipients. Furthermore, studies in vitro showing that suppressor T-lymphocytes inhibit expression of CD86 in APCs highlights another mechanism of how tolerance induces impaired APC function (Liu et al. 1999; Li et al. 1999; Lederman and Siciu-Foca 1999).

Administration of col(V) prevents proliferative responses to alloantigens, in addition to preventing proliferative responses to itself, and prevents the development of acute rejection pathology in recipient lungs (see e.g., U.S. Pat. No. 7,348,005; WO 2007/120947). Thus, col(V) may induce anergy to donor alloantigens and to itself; or alternatively, the lack of proliferative responses to donor antigens and to colV may be due to clonal deletion of alloantigen-specific lung lymphocytes; or yet alternatively may result from suppressor cell activity.

Without being bound by theory, tolerance induction by colV may be induced through linked suppression (see, e.g., Hum Immunol. 2008 November; 69(11):715-20). Also, it is thought that differential binding of colV to the collagen receptors on lymphocytes, leading to differential activation

signals may be involved (see e.g., Cell Signal. 2006 August; 18(8):1108-16). For example, only collagen V induced IL-17 signaling in T cells.

Oral administration of antigens is an effective method of inducing peripheral T-cell tolerance. This phenomenon, often referred to as oral tolerance, has been well studied in various models of autoimmune diseases in animals including encephalomyelitis, uveitis, diabetes, myasthenia gravis, and arthritis. However, the mechanisms for inducing tolerance are not completely understood. All of the known mechanisms for tolerance induction, including clonal anergy, clonal deletion, and regulation by IL-4, IL-10, or TGF-beta-mediated active suppression may have a role in oral tolerance (Faria and Weiner, 1999). Generally, higher doses of antigen are reported to induce anergy or clonal deletion (Chen et al., 1995; Whitacre et al., 1991), whereas low doses induce cytokine regulation and active suppression (Faria and Weiner, 1999; Chen et al., 1994). In the animal model of cardiac transplantation, oral administration of allogeneic splenocytes has been shown to be effective in tolerance induction by bypassing Th1 activation and selectively stimulating induction of Th-2 derived inhibitory cytokines such as IL-4 (Hancock et al., 1993; Ishido et al., 1999).

Thus, oral tolerance is a method of downregulating an immune response in a subject by orally administering an antigen (i.e. by feeding) to the subject. Oral tolerance is characterized by decreased levels of systemic antibody production, as well as decreased delayed type hypersensitivity responses (DTH), T cell proliferation, cytotoxic responses and graft rejection (Alpan et al. 2001. J. Immunol. 166:4843-52; Chen et al. 1995. Nature 376:177-80; Weiner. 1997. Imm. Today. 7:335-44; Sayegh et al. 1992. Transplantation. 53:163-6).

Other routes for inducing tolerance are also contemplated herein, in particular by intramuscular, subcutaneous, intradermal and intravenous injection. Intradermal injections are contemplated herein where low doses tend to induce tolerance and high doses induce an immune response.

Studies of tolerance have focused primarily on the effect of the tolerizing antigen on T-lymphocyte function, and the role of T-lymphocytes in suppressing immune activation (Faria and Weiner 1999; Mayer 2000; Garside and Mowat 1997). However, immune responses to any antigen require interactions between APCs and T-lymphocytes, and the T-lymphocyte may affect APC function (Liu et al. 1999; Li et al. 1999; Lederman and Siciu-Foca 1999). Therefore, down-regulated antigen presentation by APCs from tolerized hosts could contribute to tolerance induction either indirectly as a result of interactions with suppressor T-lymphocytes, or possibly as a result of direct effects of the tolerizing antigen on the APC.

Thus, in certain aspects of the invention, the present invention provides methods for restoring or reinforcing self-tolerance to colV in COPD and asthma patients. One embodiment is a method of treating COPD or asthma by administering colV either by oral therapy (Yasufuku et al, 2001; Yasufuku, et al, 2002) interstitially into the lung or by other desensitization strategies on a dosage regimen designed to increase the patient tolerance for collagens including, but not necessarily limited to colV and antigenic components and variants thereof.

By "tolerogenic fragment" is meant a fragment that can induce tolerance to the full-length protein of which it is a fragment (e.g., full-length type V collagen, and tolerogenic fragments thereof). In certain embodiments, a tolerogenic fragment can induce tolerance to the full-length type V collagen at least as well as the full-length type V collagen protein can and in certain embodiments may be more effective than

the full-length collagen protein at inducing tolerance. However, in certain embodiments, a tolerogenic fragment induces tolerance to the full-length type V collagen but may not induce tolerance as effectively as the full-length type V collagen protein. Such tolerogenic fragments may still be useful in the present invention particularly where said tolerogenic fragments have other advantageous properties, such as ease of preparation or purification as compared to the full-length protein. As would be recognized by the skilled person, a variety of known assays can be used to assess induction of tolerance, including measuring delayed-type hypersensitivity (DTH) responses, measuring cytokine productions by ELISA or other methods, T cell proliferation or cytotoxicity assays, B cell proliferation assays, antibody production, and the like. Such assays are known in the art and are described, for example, in *Current Protocols in Immunology*, Edited by: John E. Coligan, Ada M. Kruisbeek, David H. Margulies, Ethan M. Shevach, Warren Strober (2001 John Wiley & Sons, NY, N.Y.); Ausubel et al. (2001 *Current Protocols in Molecular Biology*, Greene Publ. Assoc. Inc. & John Wiley & Sons, Inc., NY, N.Y.); U.S. Pat. No. 7,348,005; and elsewhere.

Thus, a tolerogenic fragment is a fragment of a tolerogenic polypeptide such as type V collagen, or any one or more of the alpha chains thereof, that itself is immunologically tolerogenic (i.e., induces tolerance) with regard to the specific B-cells and/or T-cells that recognize the polypeptide via their surface receptors (e.g., B cell antibody receptor or T cell receptor). Tolerogenic fragments may generally be identified using well known techniques, such as those summarized in Paul, *Fundamental Immunology*, 3rd ed., (Raven Press, 1993) and references cited therein. Such techniques include screening polypeptides for the ability to reduce T cell and/or B cell reactivity. As used herein, antisera and antibodies are "antigen-specific" if they specifically bind to an antigen (i.e., they react specifically with the protein in an ELISA or other immunoassay, and do not react detectably with unrelated proteins). Such antisera and antibodies may be prepared as described herein, and using well-known techniques.

In one embodiment, a tolerogenic fragment of a polypeptide of the present invention is a portion that induces B cell and/or T cell tolerance at a level that is not substantially less than the tolerogenic activity of the full-length polypeptide (e.g., in an appropriate assay such as antibody production, which may be measured by ELISA, and/or T cell reactivity assay (T cell proliferation or cytokine production assay). Preferably, the level of tolerogenic activity of the tolerogenic portion is at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, or greater than about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% of the tolerogenic activity of the full-length polypeptide. In some instances, tolerogenic fragments will be identified that have a level of tolerogenic activity greater than that of the corresponding full-length polypeptide, e.g., having greater than about 100%, 110%, 120%, 130%, 140% or 150% or more tolerogenic activity.

In certain embodiments, tolerogenic fragments may be identified using computer analysis, such as the Tsites program (see Rothbard and Taylor, *EMBO J.* 7:93-100, 1988; Deavin et al., *Mol. Immunol.* 33:145-155, 1996), which searches for peptide motifs that have the potential to elicit Th responses. CTL peptides with motifs appropriate for binding to murine and human class I or class II MHC may be identified according to BIMAS (Parker et al., *J. Immunol.* 152:163, 1994) and other HLA peptide binding prediction analyses. Alternatively, portions that bind to a particular MHC molecule can be identified by using defined peptide binding motifs such as those described in Rammensee et al., *Immunogenetics* 41:178-228, 1995. To confirm peptide binding to

murine and human class I or class II MHC molecules, peptide binding assays known in the art may be used. To confirm immunogenicity or tolerogenicity, a peptide may be tested using an HLA A2 or other transgenic mouse model and/or an in vitro stimulation assay using dendritic cells, fibroblasts or peripheral blood cells.

It should be noted that in certain embodiments, a tolerogenic fragment of the invention is also an immunogenic fragment. In this regard, as would be recognized by the skilled artisan, highly immunogenic fragments, such as immunodominant epitopes of proteins like colV, may be tolerogenic when administered correctly, e.g., generally in low doses over extended periods of time. Thus, the present invention also contemplates the identification and use of immunogenic fragments of colV where such immunogenic fragments may be used to induce tolerance. In this regard, the level of immunogenic activity of an immunogenic portion is at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, or greater than about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% of the immunogenic activity of the full-length polypeptide. In some instances, immunogenic fragments will be identified that have a level of immunogenic activity greater than that of the corresponding full-length polypeptide, e.g., having greater than about 100%, 110%, 120%, 130%, 140% or 150% or more immunogenic activity.

The same analyses may be used to identify immunogenic fragments that are used to identify tolerogenic fragments, including using computer analysis, such as the Tsites program (see Rothbard and Taylor, *EMBO J.* 7:93-100, 1988; Deavin et al., *Mol. Immunol.* 33:145-155, 1996), which searches for peptide motifs that have the potential to elicit Th responses. CTL peptides with motifs appropriate for binding to murine and human class I or class II MHC may be identified according to BIMAS (Parker et al., *J. Immunol.* 152:163, 1994) and other HLA peptide binding prediction analyses. Alternatively, portions that bind to a particular MHC molecule can be identified by using defined peptide binding motifs such as those described in Rammensee et al., *Immunogenetics* 41:178-228, 1995. To confirm peptide binding to murine and human class I or class II MHC molecules, peptide binding assays known in the art may be used. To confirm immunogenicity or tolerogenicity, a peptide may be tested using an HLA A2 or other transgenic mouse model and/or an in vitro stimulation assay using dendritic cells, fibroblasts or peripheral blood cells.

Intact type V collagen or any one or more of its component alpha chains that have tolerogenic or immunogenic activity, are contemplated for use in the methods of the present invention. As such, a tolerogenic fragment or immunogenic fragment of collagen V may refer to a fragment of intact type V collagen or may refer to a tolerogenic or immunogenic fragment of any one of the component alpha chains. In certain embodiments, the colV as used herein may comprise the collagen molecule composed of the three alpha chains. In a further embodiment, colV may comprise any one or more of the alpha chains, such as those set forth in SEQ ID NOs:2, 4 or 6, encoded by the polynucleotides set forth in SEQ ID NOs:1, 3, or 5, or a tolerogenic fragment or an immunogenic fragment thereof.

In certain embodiments, two or more tolerogenic or immunogenic fragments may be used concurrently, either administered separately, mixed in a composition, or as a fusion protein. In this regard, any number of tolerogenic fragments or immunogenic fragments may be used to induce tolerance to colV, such as 2, 3, 4, 5, 6, 7, 8, 9, 10 or more tolerogenic or immunogenic fragments, either in a composition as separate fragments or as a fusion protein, with or without linkers. In

certain embodiments, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 or more fragments may be used in the methods of the present invention.

Detecting the presence of antibodies to collagen in accordance with some embodiments may be accomplished using any of a number of immunoassay procedures, such as by ELISA procedures. A wide range of immunoassay techniques is available as can be seen by reference to standard immunoassay textbooks these include, but are not limited to singlesite and two-site or "sandwich" assays of the non-competitive types, as well as the traditional competitive binding assays.

Sandwich assays are among the most useful and commonly used antibody based assay methods and may be used to practice various embodiments. A number of variations of the sandwich assay technique exist, and all are intended to be encompassed by various embodiments. Briefly, in a typical assay to detect antibodies in a sample, an unlabelled antigen is immobilized on a solid substrate and the sample to be tested is contacted with the bound antigen molecule. After a suitable period of incubation, i.e. for a period of time sufficient to allow formation of an antibody-antigen complex, a second antibody such as anti-human IgG, labeled with a reporter molecule capable of producing a detectable signal, is then added and incubated, allowing time sufficient for the formation of an antibody-antigen-labeled antibody. Any unreacted material is washed away, and the presence of the antibody to be detected in the sample is determined by observation of a signal produced by the reporter molecule. The results may either be qualitative, e.g., by simple observation of the visible signal, or may be quantitated by comparing the signal generated by a sample of interest with a control sample containing known amounts of antibody to be detected. Variations on this assay include a simultaneous assay, in which both the sample and labeled antibody are added simultaneously to the bound antigen. These techniques are well known, including any minor variations as will be readily apparent to those in the art. In the typical sandwich assay, antigen is immobilized, for example by being either covalently or passively bound to a solid surface. In some embodiments the solid surface is typically glass or a polymer, the most commonly used polymers being cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride or polypropylene. The solid supports may be in the form of tubes, beads, discs, or microplates, or any other surface suitable for conducting an immunoassay. Various binding processes are well-known in the art and generally consist of crosslinking, covalent binding or physical adsorption of the antigen to a given surface. The immobilized antigen is then washed in preparation for the addition of the test sample. An aliquot of the sample to be tested is then contacted with the immobilized antigen and incubated for a period of time sufficient (e.g. 2-40 minutes) and under suitable conditions (e.g. 25° C.) to allow binding of any antibody to collagen present in the sample. The actual length of contact time, buffer conditions, temperatures and the like are readily adjustable parameters and are typically readily arrived at for a given test. Following the incubation period, the immobilized antigen including any bound antibody is washed and dried, and incubated with a second antibody specific for the bound antibody, for example anti-human IgG. The second antibody is linked to a reporter molecule which is used to indicate the binding of the second antibody to the antibody-immobilized antigen complex.

One particular method for measuring antibodies to colV is described in WO 2007/120947. Specifically, this bead assay detects antibodies to type V collagen as may be present in serum and/or lung lavage fluid from patients that have an

autoimmune response type V collagen. Type V collagen-coated beads along with other necessary reagents are provided for this assay. Briefly, a typical assay is as follows: 1) Streptavidin-coated beads (e.g., such as those from Polyscience, Warrington, Pa.) are washed with sterile PBS. Beads are suspended in an appropriate volume of PBS with human type V collagen. 2) A positive control is generated by following the same procedures in 1 above, using rabbit antibody to human collagen V antibody (Bioten) (Abeam, Cambridge, Mass.). 3) For each assay, conjugated beads are washed in PBS, and incubated in PBS plus serum of lung lavage fluid. The beads are then washed with PBS containing 10% FCS. 4) The beads are then suspended in sterile PBS+10% FBS and incubated at room temperature with secondary antibody. Typically, anti-human IgG antibody conjugated with R-PE is used (Sigma, Saint Louis) although as would be understood by the skilled artisan, other suitable antibodies are available. In this regard, anti-human IgG1-, IgG2-, IgG3-, or IgG4-specific antibodies may be used in certain embodiments in order to detect switching from one subtype to another during the course of disease. The beads are washed in PBS containing 10% FCS, suspended in PBS/FCS solution and analyzed using a flow cytometer. For the positive control, known amounts of anti-colV antisera or antibody may be added to the bead assay.

Typically, antibodies to colV found in COPD and asthma patients are IgG but other classes of antibodies may also be present, such as IgM. Further, in certain embodiments of the present invention, the subtype of IgG antibodies to type V collagen as may be present in serum and/or lung lavage fluid from patients changes through the course of the disease. In this regard, the IgG subtype switching may occur during the course of a disease and certain subtypes may be indicative of worsening disease. Thus, any one or more of the IgG subtypes may be present during the course of disease, e.g. IgG1, IgG2, IgG3, or IgG4, or any combination thereof. The present invention provides for methods for detecting Type V collagen-specific IgG1, IgG2, IgG3, IgG4 subtype antibodies using the bead assay as described herein. As would be recognized by the skilled person, IgG subtype-specific antibodies are commercially available and may be used in the methods described herein. In certain embodiments, an increase in IgG1 indicates worsening of disease. In another embodiment, a switch to IgG2 subtype indicates worsening of disease. In further embodiment, a switch to IgG3 subtype indicates worsening of disease. In yet an additional embodiment, a switch to IgG4 subtype indicates worsening of disease.

Compositions, Pharmaceutical Compositions and Methods of Use

Administration of the tolerogenic compounds or compositions of the invention, or their pharmaceutically acceptable salts, in pure form or in an appropriate pharmaceutical composition, can be carried out via any of the accepted modes of administration of agents for serving similar utilities. As noted elsewhere herein, one route contemplated herein for inducing tolerance is oral administration. However, any of a variety of other routes may also be used, in particular including intravenous, intramuscular, intradermal, subcutaneous injection, and other routes. The pharmaceutical compositions of the invention can be prepared by combining a compound of the invention with an appropriate pharmaceutically acceptable carrier, diluent or excipient, and may be formulated into preparations in solid, semi-solid, liquid or gaseous forms, such as tablets, capsules, powders, granules, ointments, solutions, suppositories, injections, inhalants, gels, microspheres, and aerosols. In addition, other pharmaceutically active ingredients (including other immunosuppressive agents) and/

or suitable excipients such as salts, buffers and stabilizers may, but need not, be present within the composition.

Administration may be achieved by a variety of different routes, including oral, parenteral, nasal, intravenous, intradermal, subcutaneous or topical. Preferred modes of administration depend upon the nature of the condition to be treated or prevented. An amount that, following administration, reduces, inhibits, prevents or delays the onset of an anti-colV immune response or clinical indication of such a response is considered effective.

In certain embodiments, the amount administered is sufficient to result in reduced immune activity as described elsewhere herein (e.g., T cell response, B cell response, anti-colV antibody level, and the like). The precise dosage and duration of treatment is a function of the disease being treated and may be determined empirically using known testing protocols or by testing the compositions in model systems known in the art and extrapolating therefrom. Controlled clinical trials may also be performed. Dosages may also vary with the severity of the condition to be alleviated. A pharmaceutical composition is generally formulated and administered to exert a therapeutically useful effect while minimizing undesirable side effects. The composition may be administered one time, or may be divided into a number of smaller doses to be administered at intervals of time. For any particular subject, specific dosage regimens may be adjusted over time according to the individual need.

The compositions of the present invention may be administered alone or in combination with other known treatments, such as immunosuppressive regimens, radiation therapy, chemotherapy, transplantation, oral collagen therapy, immunotherapy, hormone therapy, photodynamic therapy, etc.

Typical routes of administering these and related pharmaceutical compositions thus include, without limitation, oral, topical, transdermal, inhalation, intrapulmonary instillation, parenteral, sublingual, buccal, rectal, vaginal, and intranasal. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques. Compositions of the invention are formulated so as to allow the active ingredients contained therein to be bioavailable upon administration of the composition to a patient. Compositions that will be administered to a subject or patient take the form of one or more dosage units, where for example, a tablet may be a single dosage unit, and a container of a compound of the invention in aerosol form may hold a plurality of dosage units. Actual methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see *Remington: The Science and Practice of Pharmacy*, 20th Edition (Philadelphia College of Pharmacy and Science, 2000). The composition to be administered will, in any event, contain a therapeutically effective amount of a compound of the invention, or a pharmaceutically acceptable salt thereof, for treatment of a disease or condition of interest in accordance with the teachings of this invention.

A pharmaceutical composition of the invention may be in the form of a solid or liquid. In one aspect, the carrier(s) are particulate, so that the compositions are, for example, in tablet or powder form. The carrier(s) may be liquid, with the compositions being, for example, an oral oil, injectable liquid or an aerosol, which is useful in, for example, inhalatory administration.

In certain embodiments, the therapeutic compound(s) are directly administered as a pressurized aerosol or nebulized formulation to the patient's lungs via inhalation. Such formulations may contain any of a variety of known aerosol propellants useful for endopulmonary and/or intranasal inhalation

administration. In addition, water may be present, with or without any of a variety of cosolvents, surfactants, stabilizers (e.g., antioxidants, chelating agents, inert gases and buffers). For compositions to be administered from multiple dose containers, antimicrobial agents are typically added. Such compositions are also generally filtered and sterilized, and may be lyophilized to provide enhanced stability and to improve solubility.

When intended for oral administration, the pharmaceutical composition is preferably in either solid or liquid form, where semi-solid, semi-liquid, suspension and gel forms are included within the forms considered herein as either solid or liquid.

As a solid composition for oral administration, the pharmaceutical composition may be formulated into a powder, granule, compressed tablet, pill, capsule, chewing gum, wafer or the like form. Such a solid composition will typically contain one or more inert diluents or edible carriers. In addition, one or more of the following may be present: binders such as carboxymethylcellulose, ethyl cellulose, microcrystalline cellulose, gum tragacanth or gelatin; excipients such as starch, lactose or dextrans, disintegrating agents such as alginate, sodium alginate, Primogel, corn starch and the like; lubricants such as magnesium stearate or Sterotex; glidants such as colloidal silicon dioxide; sweetening agents such as sucrose or saccharin; a flavoring agent such as peppermint, methyl salicylate or orange flavoring; and a coloring agent.

When the pharmaceutical composition is in the form of a capsule, for example, a gelatin capsule, it may contain, in addition to materials of the above type, a liquid carrier such as polyethylene glycol or oil.

The pharmaceutical composition may be in the form of a liquid, for example, an elixir, syrup, solution, emulsion or suspension. The liquid may be for oral administration or for delivery by injection, as two examples. When intended for oral administration, preferred composition contain, in addition to the present compounds, one or more of a sweetening agent, preservatives, dye/colorant and flavor enhancer. In a composition intended to be administered by injection, one or more of a surfactant, preservative, wetting agent, dispersing agent, suspending agent, buffer, stabilizer and isotonic agent may be included.

The liquid pharmaceutical compositions of the invention, whether they be solutions, suspensions or other like form, may include one or more of the following adjuvants: sterile diluents such as water for injection, saline solution, preferably physiological saline, Ringer's solution, isotonic sodium chloride, fixed oils such as synthetic mono or diglycerides which may serve as the solvent or suspending medium, polyethylene glycols, glycerin, propylene glycol or other solvents; antibacterial agents such as benzyl alcohol or methyl paraben; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic. Physiological saline is a preferred adjuvant. An injectable pharmaceutical composition is preferably sterile.

A liquid pharmaceutical composition of the invention intended for either parenteral or oral administration should contain an amount of a compound of the invention such that a suitable dosage will be obtained. Typically, this amount is at least 0.01% of a compound of the invention in the composition. When intended for oral administration, this amount may be varied to be between 0.1 and about 70% of the weight of the

composition. Certain oral pharmaceutical compositions contain between about 4% and about 75% of the compound of the invention. Certain pharmaceutical compositions and preparations according to the present invention are prepared so that a parenteral dosage unit contains between 0.01 to 10% by weight of the compound prior to dilution of the invention.

The pharmaceutical composition of the invention may be intended for topical administration, in which case the carrier may suitably comprise a solution, emulsion, ointment or gel base. The base, for example, may comprise one or more of the following: petrolatum, lanolin, polyethylene glycols, bee wax, mineral oil, diluents such as water and alcohol, and emulsifiers and stabilizers. Thickening agents may be present in a pharmaceutical composition for topical administration. If intended for transdermal administration, the composition may include a transdermal patch or iontophoresis device. Topical formulations may contain a concentration of the compound of the invention from about 0.1 to about 10% w/v (weight per unit volume).

The pharmaceutical composition of the invention may be intended for rectal administration, in the form, for example, of a suppository, which will melt in the rectum and release the drug. The composition for rectal administration may contain an oleaginous base as a suitable nonirritating excipient. Such bases include, without limitation, lanolin, cocoa butter and polyethylene glycol.

The pharmaceutical composition of the invention may include various materials, which modify the physical form of a solid or liquid dosage unit. For example, the composition may include materials that form a coating shell around the active ingredients. The materials that form the coating shell are typically inert, and may be selected from, for example, sugar, shellac, and other enteric coating agents. Alternatively, the active ingredients may be encased in a gelatin capsule.

The pharmaceutical composition of the invention in solid or liquid form may include an agent that binds to the compound of the invention and thereby assists in the delivery of the compound. Suitable agents that may act in this capacity include a monoclonal or polyclonal antibody, a protein or a liposome.

The pharmaceutical composition of the invention may consist of dosage units that can be administered as an aerosol. The term aerosol is used to denote a variety of systems ranging from those of colloidal nature to systems consisting of pressurized packages. Delivery may be by a liquefied or compressed gas or by a suitable pump system that dispenses the active ingredients. Aerosols of compounds of the invention may be delivered in single phase, bi-phasic, or tri-phasic systems in order to deliver the active ingredient(s). Delivery of the aerosol includes the necessary container, activators, valves, subcontainers, and the like, which together may form a kit. One of ordinary skill in the art, without undue experimentation may determine preferred aerosols.

The pharmaceutical compositions of the invention may be prepared by methodology well known in the pharmaceutical art. For example, a pharmaceutical composition intended to be administered by injection can be prepared by combining a compound of the invention with sterile, distilled water so as to form a solution. A surfactant may be added to facilitate the formation of a homogeneous solution or suspension. Surfactants are compounds that non-covalently interact with the compound of the invention so as to facilitate dissolution or homogeneous suspension of the compound in the aqueous delivery system.

The compounds of the invention, or their pharmaceutically acceptable salts, are administered in a therapeutically effective amount, which will vary depending upon a variety of

factors including the activity of the specific compound employed; the metabolic stability and length of action of the compound; the age, body weight, general health, sex, and diet of the patient; the mode and time of administration; the rate of excretion; the drug combination; the severity of the particular disorder or condition; and the subject undergoing therapy. Generally, a therapeutically effective daily dose is (for a 70 kg mammal) from about 0.001 mg/kg (i.e., 0.07 mg) to about 100 mg/kg (i.e., 7.0 g); preferably a therapeutically effective dose is (for a 70 kg mammal) from about 0.01 mg/kg (i.e., 0.7 mg) to about 50 mg/kg (i.e., 3.5 g); more preferably a therapeutically effective dose is (for a 70 kg mammal) from about 1 mg/kg (i.e., 70 mg) to about 25 mg/kg (i.e., 1.75 g).

In certain embodiments, the dose of colV administered orally is from 0.001 mg to 500 mg per day. In one particular embodiment, the oral dose of colV as described herein is from 0.01 mg to 50 mg per day. In a further embodiment, the oral dose of colV as described herein is from 0.1 mg to 0.5 mg per day. In one embodiment, the oral dose of colV is 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.07, 0.08, 0.09, or 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3.0, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9, 4.0, 4.1, 4.2, 4.3, 4.4, 4.5, 4.6, 4.7, 4.8, 4.9, or 5.0 mg per day. In another embodiment, the oral dose of colV may be 5.1, 5.2, 5.3, 5.4, 5.5, 5.6, 5.7, 5.8, 5.9, 6.0, 6.1, 6.2, 6.3, 6.4, 6.5, 6.6, 6.7, 6.8, 6.9, 7.0, 7.1, 7.2, 7.3, 7.4, 7.5, 7.6, 7.7, 7.8, 7.9, or 8.0 mg per day. In certain embodiments, the dose may given in a single dose, or may be given in multiple doses over the course of the day, for example in 2, 3 or 4 doses per day for a total of a particular mg/day dose.

As described elsewhere herein, in certain embodiments, a therapeutically effective dose of colV as used herein is a dose sufficient to induce tolerance to colV measured using any of a variety of methods as described herein. In certain embodiments, induction of tolerance to colV results in a decrease in serum anti-colV antibodies as measured using the methods described herein, such as an ELISA. In a further embodiment, a therapeutically effective dose of colV as used herein is a dose sufficient to induce T cell tolerance to colV as measured using any of a variety of methods as described herein, such as cytokine release assays, intracellular cytokine staining and flow cytometry, ELISPOT, and the like. Functional T cell assays, such as proliferation of cytotoxicity assays may also be used.

Compounds of the invention, or pharmaceutically acceptable salts thereof, may also be administered simultaneously with, prior to, or after administration of one or more other therapeutic agents. Such combination therapy includes administration of a single pharmaceutical dosage formulation which contains a compound of the invention and one or more additional active agents, as well as administration of the compound of the invention and each active agent in its own separate pharmaceutical dosage formulation. For example, a compound of the invention and the other active agent can be administered to the patient together in a single oral dosage composition such as a tablet or capsule, or each agent administered in separate oral dosage formulations. Where separate dosage formulations are used, the compounds of the invention and one or more additional active agents can be administered at essentially the same time, i.e., concurrently, or at separately staggered times, i.e., sequentially; combination therapy is understood to include all these regimens.

The compounds of the present invention may be administered to an individual afflicted with a disease or disorder as described herein, such as COPD and severe and persistent asthma. For in vivo use for the treatment of human disease,

the compounds described herein are generally incorporated into a pharmaceutical composition prior to administration. A pharmaceutical composition comprises one or more of the compounds described herein in combination with a physiologically acceptable carrier or excipient as described elsewhere herein. To prepare a pharmaceutical composition, an effective amount of one or more of the compounds is mixed with any pharmaceutical carrier(s) or excipient known to those skilled in the art to be suitable for the particular mode of administration. A pharmaceutical carrier may be liquid, semi-liquid or solid. Solutions or suspensions used for parenteral, intradermal, subcutaneous or topical application may include, for example, a sterile diluent (such as water), saline solution, fixed oil, polyethylene glycol, glycerin, propylene glycol or other synthetic solvent; antimicrobial agents (such as benzyl alcohol and methyl parabens); antioxidants (such as ascorbic acid and sodium bisulfite) and chelating agents (such as ethylenediaminetetraacetic acid (EDTA)); buffers (such as acetates, citrates and phosphates). If administered intravenously, suitable carriers include physiological saline or phosphate buffered saline (PBS), and solutions containing thickening and solubilizing agents, such as glucose, polyethylene glycol, polypropylene glycol and mixtures thereof.

The compounds described herein may be prepared with carriers that protect it against rapid elimination from the body, such as time release formulations or coatings. Such carriers include controlled release formulations, such as, but not limited to, implants and microencapsulated delivery systems, and biodegradable, biocompatible polymers, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, polyorthoesters, polylactic acid and others known to those of ordinary skill in the art.

In certain embodiments, adjuvants which assist in inducing tolerance include dexamethasone (see e.g., Y. Kang, et al., *J. Immunol.* 2008, 180: 5172-5176), lipopolysaccharides (LPS) and cholera toxin β -subunit and may be added to the formulations. Certain other tolerogenic carriers are also contemplated for use with the colV compositions of the present invention. Such carriers include mineral oil carriers such as incomplete, Freund's adjuvant (IFA) or complete Freund's adjuvant (CFA). IFA is an emulsion of mineral oil. CFA is a preparation of mineral oil containing various amounts of killed organisms of *Mycobacterium*. However, IFA and CFA are not allowed for human use because the mineral oil is not metabolizable and cannot be degraded by the body.

In certain embodiments, fat emulsions, which have been in use for many years for intravenous nutrition of human patients, can also act as a vehicle for tolerogenic polypeptide therapy using the polypeptides of the present invention. Two examples of such emulsions are the available commercial fat emulsions known as Intralipid and Lipofundin. "Intralipid" is a registered trademark of Kabi Pharmacia, Sweden, for a fat emulsion for intravenous nutrition, described in U.S. Pat. No. 3,169,094. "Lipofundin" is a registered trademark of B. Braun Melsungen, Germany. Both contain soybean oil as fat (100 or 200 g in 1,000 ml distilled water: 10% or 20%, respectively). Egg-yolk phospholipids are used as emulsifiers in Intralipid (12 g/l distilled water) and egg-yolk lecithin in Lipofundin (12 g/l distilled water). Isotonicity results from the addition of glycerol (25 g/l) both in Intralipid and Lipofundin. It is believed that these vehicles are actually biologically active carriers which when complexed with the suspected auto-antigen, promote a TH1 to TH2 shift of the autoimmune T cells. In certain embodiments, such a vehicle

is a fat emulsion comprising 10-20% triglycerides of plant and/or animal origin, 1.2-2.4% phospholipids of plant and/or animal origin, 2.25-4.5% osmo-regulator, 0-0.05% anti-oxidant, and sterile water to 100%.

In certain embodiments, colV or tolerogenic fragments thereof may be linked to the diphtheria toxin receptor to enhance GI uptake.

The tolerogenic compositions of the present invention may be used to treat asthma and COPD and other pulmonary diseases associated with autoimmunity to colV. Thus, in one embodiment, the present invention provides a method for treating chronic obstructive pulmonary disease comprising administering to a COPD patient a therapeutically effective amount of type V collagen, or a tolerogenic fragment thereof, or an immunogenic fragment thereof administered in a dose effective to induce tolerance. In this regard, a COPD patient may have emphysema or chronic obstructive bronchitis.

Another aspect of the invention provides a method for treating asthma comprising administering to an asthma patient a therapeutically effective amount of type V collagen, or a tolerogenic fragment thereof, or an immunogenic fragment thereof administered in a dose effective to induce tolerance. In this regard, the asthma patient may have severe, persistent asthma. In another embodiment, the present invention provides methods for reducing the severity of asthma in an asthma patient.

The present invention further provides, a method for preventing the development of chronic obstructive pulmonary disease in a subject at risk for developing chronic obstructive pulmonary disease comprising administering to the subject a therapeutically effective amount of type V collagen, or a tolerogenic fragment thereof.

As noted elsewhere herein, intact colV or any one or more of its component α chains may be administered, or a tolerogenic fragment of any of the aforementioned molecules. Further, also as noted elsewhere herein, an immunogenic fragment of intact colV or any one or more of its component α chains may be administered in any of the methods described herein, using a dose effective to induce tolerance to colV. Such a dose will vary depending upon a variety of factors including the activity of the specific proteins or fragments employed; the metabolic stability and length of action of these compounds; the age, body weight, general health, sex, and diet of the patient; the mode and time of administration; the rate of excretion; the drug combination; the severity of the particular disorder or condition; and the subject undergoing therapy. Thus, such tolerogenic doses may be determined empirically using known testing protocols or by testing the compositions in model systems known in the art and extrapolating therefrom. Controlled clinical trials may also be performed. Dosages may also vary with the severity of the condition to be alleviated.

As would be readily appreciated by the skilled artisan, a variety of factors can be assessed to determine the effectiveness of the compounds and methods of the invention for treating, preventing, or reducing the severity of COPD or asthma. Such factors include typical clinical symptoms of asthma or COPD which can be assessed by the skilled clinician. These symptoms may include, but are not limited to: for asthma: wheezing; chest tightness or pain; rapid heart rate; sweating; peak flow rates as measured by a peak flow meter; frequent cough, especially at night; loss of breath easily or shortness of breath; feeling very tired or weak when exercising; wheezing or coughing after exercise; feeling tired, easily upset, grouchy, or moody; decreases or changes in lung function as measured on a peak flow meter; signs of a cold, or allergies (sneezing, runny nose, cough, nasal congestion, sore

throat, and headache); trouble sleeping. Such factors include typical clinical symptoms of COPD such as, but not limited to the amount of sputum produced; thickness or stickiness of sputum; sputum color or the presence of blood in the sputum; severity of shortness of breath, cough and/or wheezing; ankle swelling; forgetfulness, confusion, slurring of speech and sleepiness; trouble sleeping; using more pillows or sleeping in a chair instead of a bed to avoid shortness of breath; an unexplained increase or decrease in weight; fatigue and lack of energy that is persistent; a lack of sexual drive; morning headaches, dizzy spells, restlessness.

The compositions and methods of the present invention can be used in conjunction with other known treatments for COPD and asthma, such as, but not limited to corticosteroids (e.g., prednisone, fluticasone, methylprednisolone), bronchodilators (e.g., short- and long-acting β 2-agonists, theophylline, pirbuterol, ephedrine, albuterol, salmeterol, levalbuterol, clenbuterol ipratropium bromide), and leukotriene modifiers (e.g., montelukast, zafirlukast, zileuton).

REFERENCES

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference. Chen, Inobe, Marks, Gonnella, Kuchroo, Weiner, "Peripheral deletion of antigen-reactive T cells in oral tolerance," *Nature*, 376:177-180, 1995. Chen, Kuchroo, Inobe, Hafler, Weiner, "Regulatory T-cell clones induced by oral tolerance: suppression of autoimmune encephalomyelitis," *Science*, 265:1237-1240, 1994. Chiang, Mainardi, Seyer, "Type V(A-B) collagen induces platelet aggregation," *J. Lab. Clin. Med.*, 95:99-107, 1980. Cremer, Ye, Terato, Owens, Seyer, Kang, "Type XI collagen-induced arthritis in the Lewis rat: characterization of cellular and humoral immune responses to native types XI, V, and II collagen and constituent α -chains," *J. Immunol.* 153:824-832, 1994. Danzer, Kirchner, Rink, "Cytokine interactions in human mixed lymphocyte culture," *Transplantation*, 57(11):1638-1642, 1994. DeMeester, Rolfe, Kunkel, Swiderski, Lincoln, Deeb, Strieter, "The bimodal expression of tumor necrosis factor- α in association with rat lung reimplantation and allograft rejection," *J. Immunol.*, 150(6):2494-2505, 1993. Faria and Weiner, "Oral tolerance: mechanisms and therapeutic applications," *Adv. Immunol.*, 73:153-264, 1999. Fedoseyeva, Zhang, Orr, Levin, Buncke, Benichou, "De novo autoimmunity to cardiac myosin after heart transplantation and its contribution to the rejection process," *J. Immunol.*, 162:6836-42, 1999. Garrovillo, Ali, Oluwole, "Indirect allorecognition in acquired thymic tolerance: induction of donor-specific tolerance to rat cardiac allografts by allopeptide-pulsed host dendritic cells," *Transplantation*, 68:1827-1834, 1999. Hancock, Sayegh, Kwok, Weiner, Carpenter, "Oral, but not intravenous, alloantigen prevents accelerated allograft rejection by selective intragraft Th2 cell activation," *Transplantation*, 55:1112-1118, 1993. Hanson, Gorman, Oui, Cheah, Solomon, Trowsdale, "The human α 2(XI) collagen gene (COL11A2) maps to the centromeric of the major histocompatibility complex on chromosome 6," *Genomics*, 5:925-931, 1989. Hirt, You, Moller, Boeke, Starke, Spranger, Wottge, "Development of obliterative bronchiolitis after allogeneic rat lung transplantation: Implication of acute rejection and the time point of treatment," *J. Heart Lung Transplant.*, 18:542-548, 1999. Huang, Fuchimoto, Scheier-Dolberg, Murphy, Neville, Sachs, J. Clin. Invest., 105:173-181, 2000. Ishido, Matsuoka, Matsuno, Nakagawa, Tanaka, "Induction of donor-specific hyporesponsiveness and prolongation of cardiac allograft survival

by jejunal administration of donor splenocytes," *Transplantation*, 68:1377-1382, 1999. Iyer, Woo, Cornejo, Gao, McCoubrey, Maines, Buelow, "Characterization and biologic significance of immunosuppressive peptide D2702.75-84 (E α V) binding protein," *J. Bio. Chem.*, 273(5):2692-2697, 1998. Joo, Pepose, Stuart, "T-cell mediated responses in a murine model of orthotopic corneal transplantation," *Invest. Ophthalmol. Vis. Sci.*, 36:1530-1540, 1995. Konomi, Hayashi, Nakayasu, Arima, "Localization of type V collagen and type IV collagen in human cornea, lung, and skin," *Am. J. Pathol.*, 116:417-426, 1984. Krensky and Clayberger, "HLA-derived peptides as novel immunosuppressives," *Nephrol. Dial. Transplant.*, 12:865-878, 1997. Lowry, Marghesco, Blackburn, "Immune mechanisms in organ allograft rejection. VI. Delayed-type hypersensitivity and lymphotoxin in experimental renal allograft rejection," *Transplantation.*, 40:183-188, 1985. Madri and Furthmayr, "Collagen polymorphism in the lung," *Human Pathology*, 11:353-366, 1980. Madri and Furthmayr, "Isolation and tissue localization of type AB2 collagen from normal lung parenchyma," *Am. J. Pathol.*, 94:323-332, 1979. Marck, Prop, Widevuur, "Lung transplantation in the rat. III. Functional studies in iso- and allografts," *J. Surgical Res.*, 35:149-158, 1983. Matsumura, Marchevsky, Zuo, Kass, Matloff, Jordan, "Assessment of pathological changes associated with chronic allograft rejection and tolerance in two experimental models of rat lung transplantation," *Transplantation.*, 59:1509-1517, 1995. Morris and Bachinger, "Type XI collagen is a heterotrimer with the composition (1 α ,2 α ,3 α) retaining non-triple helical domains. *J. Biological Chem.*, 262:11345-11350, 1987. Murphy, Magee, Alexander, Waaga, Snoeck, Vella, Carpenter, Sayagh, "Inhibition of allorecognition by a human class II MHC-derived peptide through the induction of apoptosis," *J. Clin. Invest.*, 103:859-867, 1999. Nosner, Goldberg, Nafitzger, Lyu, Clayberger, Krensky, "HLA-derived peptides which inhibit T cell function bind to members of the heat-shock protein 70 family," *J. Exp. Med.*, 183:339-348, 1996. Oluwole, Chowdhury, Jin, Hardy, "Induction of transplantation intolerance to rat cardiac allografts by intrathymic inoculation of allogeneic soluble peptides," *Transplantation*, 56(6):1523-1527, 1993. Prop, Nieuwenhuis, Wildevuur, "Lung allograft rejection in the rat. I. Accelerated rejection caused by graft lymphocytes," *Transplantation*, 40:25-30, 1985. Prop, Wildevuur, Nieuwenhuis, "Lung allograft rejection in the rat. II. Specific immunological properties of lung grafts," *Transplantation*, 40:126-131, 1985. Sayagh, Watschinger, Carpenter, "Mechanisms of T cell recognition of alloantigen," *Transplantation*, 57:(9)1295-1302, 1994. Sayegh and Krensky, "Novel immunotherapeutic strategies using MHC derived peptides," *Kidney Int. Suppl.* 53:S13-20, 1996. Sayegh, Khoury, Hancock, Weiner, Carpenter, "Induction of immunity and oral tolerance with polymorphic class II major histocompatibility complex allopeptides in the rat," *Proc. Natl. Acad. Sci.*, 89: 7762-7766, 1992. Sayegh, Zhang, Hancock, Kwok, Carpenter, Weiner, "Down-regulation of the immune response to histocompatibility antigens and prevention of sensitization by skin allografts by orally administered alloantigen," *Transplantation*, 53:163-166, 1992. Sekine, Nowen, Heidler, Van Rooijen, Brown, Cummings, Wilkes, "Role of passenger leukocytes in allograft rejection—Effect of depletion donor alveolar Macrophages on the local production of TNF-alpha, T helper 1/Thelper 2 cytokines, IgG subclasses, and pathology in a rat model of lung transplantation," *J. Immunol*, 159:4084-4093, 1997. Seyer and Kang, "Covalent structure of collagen: amino acid sequence of three cyanogen bromide-derived peptides from human alpha 1(V) collagen chain. *Arch. Biochem. Biophys.* 271(1): 120-129,

1989. SivaSai, Smith, Poindexter, Sundaresan, Trulock, Lynch, Cooper, Patterson, Mohanakumar, "Indirect recognition of donor HLA class I peptides in lung transplant recipients with bronchiolitis obliterans syndrome," *Transplantation*, 67(8):1094-1098, 1999. Smith Jr, Williams, Brandt, "Interaction of proteoglycans with pericellular (1 alpha, 2 alpha, 3 alpha) collagens of cartilage," *J. Biol. Chem.*, 260:10761-10767, 1985. Stark and Ostrow, *Training Manual Series, Laboratory Animal Technician, American Association for Laboratory Animal Science*, 181-182, 1990. Strober and Coffman, "Tolerance and immunity in the mucosal immune system," *Res. Immunol.*, 148:489-599, 1997. Trulock, "Lung transplantation," *Am. J. Respir. Crit. Care Med.*, 155:789-818, 1997. VanBuskirk, Wakely, Sirak, Orosz, "Patterns of allosensitization in allograft recipients: long-term allograft acceptance is associated with active alloantibody production in conjunction with active inhibition of alloreactive delayed-type hypersensitivity," *Transplantation.*, 65:1115-1123, 1998. Westra, Prop, Kuijpers, "A paradox in heart and lung rejection," *Transplantation*, 49:826-828, 1990. Whitacre, Gienapp, Orosz, Bitar, "Oral tolerance in experimental autoimmune encephalomyelitis. III. Evidence for clonal anergy," *J. Immunol.*, 147:2155-2163, 1991. Wilkes, Bowman, Cummings, Heidler, "Allogeneic bronchoalveolar lavage cells induce the histology and immunology of lung allograft rejection in recipient murine lungs. Role of ICAM-1 on donor cells," *Transplantation*, 67(6):890-896, 1999. Wilkes, Heidler, Bowen, Quinlan, Doyle, Cummings, Doerschuk, "Allogeneic bronchoalveolar lavage cells induce the histology of acute lung allograft rejection, and deposition of IgG2a in recipient murine lungs," *J. Immunol.*, 155:2775-2783, 1995. Wilkes, Thompson, Cummings, Bragg, Heidler, "Instillation of allogeneic lung macrophages and dendritic cells cause differential effects on local IFN- γ production, lymphocytic bronchitis, and vasculitis in recipient murine lungs," *J. Leukoc. Biol.* 64:578-586, 1998. Wilson, Ebringer, Ahmadi, Wrigglesworth, Tiwana, Fielder, Binder, Ettelaie, Cunningham, Joannou, Bansal, "Shared amino acid sequences between major histocompatibility complex class II glycoproteins, type XI collagen and *Proteus mirabilis* in rheumatoid arthritis," *Ann. Rheum. Dis.*, 54:216-220, 1995. Woessner Jr., "The determination of hydroxyproline in tissue and protein samples containing small proportions of this imino acid," *Arch. Biochem. Biophys.* 93:440-447, 1961. Yagyu, Steinhoff, Schafers, Dammenhayn, Haverich, Borst, "Comparison of mononuclear cell populations in bronchoalveolar lavage fluid in acute rejection after lung transplantation and *Mycoplasma* infection in rats," *J. Heart Transplant.*, 9:516-525, 1990. Yamagami, Tsuru, Ohkawa, Endo, Isobe, "Suppression of allograft rejection with anti-alpha beta T cell receptor antibody in rat corneal transplantation," *Transplantation*, 67:600-604, 1999. Yoshino, Quattrocchi, Weiner, "Suppression of antigen-induced arthritis in Lewis rats by oral administration of type II collagen," *Arthritis Rheum.* 38:1092-1096, 1995. Yousem, Berry, Cagle, Chamberlain, Husain, Hruban, Marchevsky, Ohori, Ritter, Stewart, Tazelaar, "Revision of the 1990 working formulation for the classification of pulmonary allograft rejection: Lung rejection study group," *J. Heart Lung Transplant*, 15:1-15, 1996. Zheng, Markees, Hancock, Li, Greine, Li, Mordes, Sayegh, Rossini, Strom, "CTLA4 signals are required to optimally induce allograft tolerance with combined donor-specific transfusion and anti-CD154 monoclonal antibody treatment," *J. Immunol.*, 162:4983-4990, 1999.

EXAMPLES

Example 1

Anti-Collagen V Antibodies are Increased in COPD Patients

Plasma was obtained from normal volunteers (non smoking adults, age 18-55), and volunteers with documented emphysema. Levels of anti-colV antibodies in COPD patients and control healthy subjects were detected by the flow cytometry bead assay as described in WO 2007/120947.

Briefly, 1) Streptavidin-coated beads (5 μ m, binding capacity 10-20 μ g/ 1×10^7 beads (Polyscience, Warrington, Pa.)) were washed two times with sterile PBS. Beads (1×10^7) were suspended in 100 μ l of PBS with 40 μ g of human Type V collagen and incubated for 60 minutes at 4° C. 2) A positive control was generated by following the same procedures in 1 above, using 20 μ m of rabbit antibody to human collagen V antibody (biotin) (Abeam, Cambridge, Mass.). 3) For each assay, 1×10^6 conjugated beads were washed two times in PBS, and incubated in 100 μ l PBS plus 50 μ l serum. After incubating for 30-minutes at room temperature, the beads were washed three times with PBS containing 10% FCS. 4) The beads were suspended in 100 μ l of sterile PBS+10% FBS and incubated for about 30 minutes at room temperature with secondary antibody. Typically, 5 μ l of anti-human IgG antibody conjugated with R-PE was used (Sigma, Saint Louis). The beads were washed three times in PBS containing 10% FCS, suspended in 300 μ l of PBS/FCS solution and analyzed using a flow cytometer.

As shown in FIG. 1, anti-type V collagen antibodies are significantly elevated in COPD patients (N=16) as compared to controls (N=42).

Example 2

Anti-Collagen V Antibodies are Increased in Asthma Patients

Plasma was obtained from normal volunteers (non smoking adults, age 18-55), and volunteers with documented chronic asthma. Levels of anti-colV antibodies in the asthma patients and control volunteer subjects were measured using the bead assay as outlined in Example 1.

As shown in FIG. 2, elevated levels of anti-type V collagen antibodies were found in 8 of 20 asthmatics.

Example 3

Intravenous Collagen V Prevents Ovalbumin-Induced Airway Hyper-Responsiveness in Mice

This Example demonstrates that intravenous administration of collagen V prevents ovalbumin-induced airway hyper-responsiveness in this well-established murine asthma model.

Balb/c mice were injected via tail vein with 100 μ g col (V), alone, or col(V) mixed in complete Freund's adjuvant, (CFA), or PBS, or CFA, alone. Seven days later mice received an IP injection of ovalbumin in alum and this was repeated seven days later. Seven days after the last ova/alum injection, mice in each group were challenged with increasing doses of aerosolized ova followed by measurements of airways resistance (PenH).

The results showed that col(V), alone, abrogated ova-induced airway hyperresponsiveness (see FIG. 3). Further experiments were conducted which confirm these results. These results are summarized below in Table 2.

TABLE 2

Col(V) alone abrogates ova-induced airway hyperresponsiveness							
Mg/ml	PBS			Col(V)			2-WAY
	PenH Units			PenH Units			ANOVA
methacholine	Mean	SEM	N	Mean	SEM	N	P value
Baseline	0.419913	0.021817	6	0.564856	0.092752	6	>0.05
Saline	0.466943	0.034769	6	0.519667	0.073273	6	>0.05
10	1.413495	0.189206	6	1.615039	0.172871	6	>0.05
25	2.448044	0.438264	6	2.914945	0.510278	6	>0.05
50	4.463914	0.319439	6	3.147791	0.531683	6	>0.05
100	5.121152	0.327822	6	3.661426	0.544324	6	>0.05
200	6.064105	0.368829	6	3.387585	0.46153	6	<0.001
300	6.825607	0.499452	6	3.700204	0.657257	6	<0.001

Example 4

Intravenous Col(V) Induces IFN- γ Transcripts in Lung Mononuclear Cells

This example shows intravenous injection of col(V) alone induced a TH1 response in lung mononuclear cells, characterized by induction of IFN- γ transcripts. Balb/c mice were injected via tail vein with 100 ug col(V) alone, CFA alone at the base of the tail, colV plus CFA the base of the tale or PBS i.v. alone. Seven days later mice received an IP injection of ovalbumin in alum and this was repeated seven days later. Seven days after the last ova/alum injection, mice in each group were challenged with increasing doses of aerosolized methacholine followed by measurements of airways resistance in response to methacholine challenge (PenH). RNA was extracted from mononuclear cells isolated from the lung parenchyma of mice in each treatment group and quantitative PCR performed for determining expression levels of IL-4, IL-5, IL-13, IFN- γ and IL-10 (see FIG. 4; data represent lung mononuclear cells of RNA pooled from 5 mice in each group).

The results of the experiment show that Col(V) alone administered intravenously induced IFN- γ transcripts in lung mononuclear cells. IFN- γ is antagonistic to IL-13 and IL4, two cytokines thought to play a key role in asthma pathogenesis. Therefore, without being bound by theory, induction of this TH1 response which counteracts the effects of IL4/IL13 may play a part in the colV-mediated protective effect in ova-induced asthma.

All of the above U.S. patents, U.S. patent application publications, U.S. patent applications, foreign patents, foreign patent applications and non-patent publications referred to in this specification and/or listed in the Application Data Sheet, are incorporated herein by reference, in their entirety.

From the foregoing it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 6

<210> SEQ ID NO 1

<211> LENGTH: 8439

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

cgcaactctcc gtccccgcgg ctggcgcagg acctcactcg agcggagcgc ccacggggag      60
cgggtcgcgg ggcggcggcg gcgaggagga ggcgagaagg agttggagga ggaggaggag      120
gaggcgaggg cgagctagcc cagcgggggtc ccggccgccc cgcgggccaa agtcgagccc      180
tcccgcccggt gggcgagcgc gccagccgcc ccttcagaa cagccgccgc cacaagaag      240
aacgggggggt gccgaggtcc ccatgacctc ctaaagtggg gcggtccctg ctgagtgcgc      300
tgcccgggccc gtgaccgcg cccctgtgcg tcccgcgcg cctccgagcg ccctgtgcg      360
ccccggcccc cgccccgccc gcatggacgt ccatacccgc tggaaagcgc gcagcgcgct      420
ccgcccgggc gccccgctgc tgcctccgct gctgctgctg ctgctgtggg cgccgctcc      480
gagccgcgca gctcagccag cagatctcct gaaggttcta gattttcaca acttgctga      540
tggataaaca aagacaacag gcttttgcgc cagcggcga tcttccaaag gcccgatgt      600

```


-continued

cgcttacaga	gtcaccaaag	acgcgcagct	cagcgcaccc	accaagcagc	tgtaccctgc	660
gtctgcattt	cccaggagact	tctccatcct	aacaactgtg	aaagccaaga	aaggcagcca	720
ggccttctctg	gtctccatct	acaacgagca	gggtatccag	cagattgggc	tggagctggg	780
ccgctctccc	gtcttctct	acgaggacca	cacggggaag	cctggccccg	aagactaccc	840
cctcttccgg	ggcatcaacc	tgtcagatgg	caagtggcac	agaattgctc	tcagcgtcca	900
caagaaaaat	gtcaccttga	tcctcgactg	taaaaaaag	accaccaa	tcctcgaccg	960
cagcgaccac	cccattgatc	acatcaatgg	catcatcgtg	tttggcacc	ggatcctgga	1020
tgaggaggtg	tttgagggtg	acatccagca	gctgctcttt	gtctcggacc	accgggcagc	1080
ttatgattac	tgtgagcact	acagcctga	ctgtgacacc	gcagtacctg	acaccccaca	1140
gtcgcaggac	cccaatccag	atgaatatta	cacggaagga	gacggcgagg	gtgagaccta	1200
ttactacgaa	taccctact	acgaagacc	cgaagacct	gggaaggagc	ccacccccag	1260
caagaagccc	gtggaagctg	ccaaagaaac	cacagaggtc	cccaggagc	tgacccccgac	1320
ccccacggaa	gctgctccca	tgctgaaac	cagtgaaggg	gctgggaagg	aagaggacgt	1380
cggcatcggg	gactatgact	acgtgccag	tgaggactac	tacacgcct	caccgtatga	1440
tgacctcacc	tatggcgagg	gggaggagaa	ccccgaccag	cccacagacc	caggcgctgg	1500
ggccgaaatt	cccaccagca	ccgccgacac	ctccaactcc	tccaatccag	ctccgcctcc	1560
aggggaaggt	gcggatgact	tggaggggga	gttcaactgag	gaaacgatcc	ggaaccttga	1620
cgagaactac	tacgaccct	actacgacc	caccagctcc	ccgtcggaga	tcgggccggg	1680
aatccggcg	aaccaggata	ccatctatga	agggattgga	ggacctcggg	gcgagaaagg	1740
ccaaaagggg	gaaccagcga	ttatcgagcc	gggatgctc	atcgagggcc	cgctggccc	1800
agaaggcccc	gcgggtcttc	ccggacctcc	aggaacctg	ggccccactg	gccaagtcgg	1860
ggacctgga	gaaaggggcc	cccctggacg	cccaggcctt	cctggggccg	atggcctgcc	1920
cggctctcca	ggaacctatc	tcattgctgc	cttccggttt	ggaggtggcg	gcgatgcggg	1980
ctccaaaggc	cccattggtct	cagcccagga	gtcccaggcg	caagccattc	tccagcaggc	2040
caggttggca	ctgaggggac	cagctggccc	gatgggtctc	acagggagac	ctggccctgt	2100
gggtccccct	gggagcggag	gtttgaaggg	cgagccggga	gacgtggggc	ctcagggtcc	2160
tcgaggtgtg	caaggcccgc	ctggtccggc	cgggaagccc	ggaagacggg	gtcgggctgg	2220
gagtgatgga	gccagaggaa	tgctggaca	aactggcccc	aagggtgacc	ggggtttcga	2280
cggcctggct	gggttgccag	gcgagaaggg	ccacaggggt	gacctggtc	cttccggccc	2340
accaggacct	ccgggagacg	atggagaaa	gggtgacgac	ggagaagttg	ggcccagggg	2400
gctgcctggg	gagcccgggc	cacgtggtct	gcttgggccc	aagggcccc	caggtcctcc	2460
cggacctccc	ggtgtcacgg	gtatggacgg	ccagccgggg	ccaaaaggaa	atgtgggtcc	2520
ccagggagag	cctggcccc	caggacagca	gggtaatcca	ggcgcccagg	gtcttccagg	2580
ccccagggt	gcaattggtc	ctccaggaga	aaagggtccc	ttggggaaac	caggccttcc	2640
aggaatgccc	ggtgctgacg	gacccccggg	acacctggc	aaagaaggcc	ctccaggaga	2700
gaaaggaggt	cagggtccac	ctggccccc	gggtccgatt	ggctaccag	gtcctcgagg	2760
agtcaagggg	gccgatggca	tccgtggtct	gaagggcaca	aagggcgaga	aggggtgaaga	2820
cggcttctct	gggtttaaag	gagacatggg	catcaaggg	gatcggggg	agatcggccc	2880
acccggtccc	aggggagaag	atggcctga	aggcccaaag	ggtcgaggag	gtcccaatgg	2940
tgacccccgt	cctctgggac	cccctgggga	gaagggaaa	ctcggagtcc	cagggttacc	3000

-continued

agggtatcca	ggaagacaag	gaccaaaggg	ctctattgga	ttccctggat	ttcctggcgc	3060
caatggagag	aagggcggca	gggggacccc	tggaaagcca	ggaccgcggg	ggcagcgagg	3120
cccaacgggt	ccgaggggtg	aaagaggccc	ccggggcatc	actgggaagc	ctggccccaa	3180
gggcaactcc	ggaggtgacg	gcccagctgg	ccctcctggg	gaacggggac	ccaatggacc	3240
ccaaggaccc	acaggatttc	ctggaccaa	gggccccct	ggccctccag	gcaaggatgg	3300
actcccagga	cacctggac	agagaggcga	gactggtttc	caaggcaaga	ccggccctcc	3360
aggccccccc	ggcgtggtcg	gccctcaggg	tcccacggga	gaaacgggcc	caatgggtga	3420
gcgtagccac	cctgggcccc	ctggaccccc	cggtgaacag	gggcttccgg	gccttgctgg	3480
aaaagaaggg	acgaaggtg	accagggccc	tgcaggcctc	cctgggaaag	atggccctcc	3540
aggattacgt	ggtttcctg	gggaccgagg	gcttctggg	ccagtgggag	ctcttgact	3600
gaaaggcaat	gaagggcccc	ctggcccc	aggccctcg	ggatctccag	gggagagagg	3660
tccagctgga	gccgctggc	ccatcggaa	tccagggaga	cctgggcccc	agggaccccc	3720
agggccggca	ggagagaaag	gggctcctgg	cgagaaaggc	ccacaaggcc	cagctggccg	3780
agacggcttc	caggggctg	tggggctccc	gggtccagct	ggccctgtgg	gtccccctgg	3840
agaagacgga	gataagggag	agatcgggga	gccggggcag	aaaggaagca	aggggggaca	3900
aggagaacag	ggcctcctg	ggcctacagg	tcctcaaggc	cccatcggac	agccaggccc	3960
ctctggagct	gacggcgagc	cggggcctcg	gggccagcag	ggccttttcg	ggcagaaagg	4020
tgatgaaggt	cccagaggct	ttcctggacc	ccctgggcca	gtggggctgc	agggtttgcc	4080
aggacctcca	ggcgagaag	gtgagacagg	agacgtgggc	cagatgggcc	ccccgggtcc	4140
ccctggcccc	cgaggaccct	ccggagctcc	aggtgctgat	ggcccacaag	gtcccccagg	4200
tggaatagga	aaccctggtg	cagtgggaga	gaagggcgag	cctggcgaag	caggtgagcc	4260
tggccttccg	ggagaaggcg	gccccccggg	acccaaagga	gaaaggggag	agaagggcga	4320
gtcaggccct	tcaggtgctg	ccggaccccc	tggacccaaa	ggccctcccg	gagatgatgg	4380
tcccaaaggc	agccctggcc	cagtgggttt	tcctggagat	cctggccccc	ccggagagcc	4440
tggccccgcg	ggtcaagatg	gtccccctgg	tgacaaagga	gatgatggtg	aaccggggca	4500
gacgggatcc	cccggcccta	ctggtgaacc	aggtccatcg	gggcctccag	gaaaaagggg	4560
tccccaggc	cccgcaggcc	ccgaaggcag	acagggagag	aaaggggcca	agggagaagc	4620
cggttgga	ggccctcctg	ggaagactgg	ccccatcggc	ccccaggggg	cccctgggaa	4680
gcccggaccg	gatggccttc	gagggatccc	tggccctgtg	ggagaacaag	gtctcccagg	4740
atccccaggc	ccggacggtc	ccccggccc	catgggtccc	ccaggacttc	ccggcctcaa	4800
aggagattct	ggtcccaaag	gtgaaaaggg	tcattccaggc	ctgatcgggc	tcattcggctc	4860
tccgggtgaa	cagggtgaga	agggcgaccg	tggctctcct	ggcccccagg	gctcctccgg	4920
tcctaagggg	gaacagggtg	tactgggtcc	ttctggcccc	attgggcctc	ctgggcccc	4980
tggcctgccg	ggtccgctg	gtccaaaagg	tgtaagggc	tcctcgggtc	caactggccc	5040
gaagggtgag	gcaggccacc	caggaccccc	aggccccccg	ggcccccccg	gagaggtcat	5100
ccagccccctg	ccaatccagg	catccaggac	gcggcggaac	atcgacgcca	gccagctgct	5160
ggacgacggg	aatggcgaga	actacgtgga	ctacgcggac	ggcatggaag	agatcttcgg	5220
ctctctcaac	tctctgaagc	tggagattga	gcagatgaaa	cggccccctg	gcacgcagca	5280
gaacccccgc	cgcacctgca	aggacctgca	gctctgccac	cccgaacttc	cagatgggtga	5340
atactgggtc	gatcctaacc	aaggatgctc	cagggatcc	ttcaaggttt	actgcaactt	5400

-continued

cacagccggg	gggtcgacat	gcgtcttccc	tgacaagaag	tccgaagggg	ccagaatcac	5460
ttcttgcccc	aaagaaaacc	cgggctcctg	gttcagtga	ttcaagcgtg	ggaaactgct	5520
ctcctatgtg	gacgccgagg	gcaaccctgt	gggtgtggta	cagatgacct	tctgcccgtt	5580
gctgagcgcc	tctgcccacc	agaacgtcac	ctaccactgc	taccagtcag	tggcctggca	5640
ggacgcagcc	acgggcagct	acgacaaggc	cctccgcttc	ctgggctcca	acgacgagga	5700
gatgtcctat	gacaacaacc	cctacatccg	cgccctggtg	gacggctgtg	ctaccaagaa	5760
aggctaccag	aagacggttc	tggagatcga	cacccccaaa	gtggagcagg	tgccatcgt	5820
ggacatcatg	ttcaatgact	tcggtgaagc	gtcacagaaa	tttggatttg	aagtggggcc	5880
ggcttgcttc	atgggctagg	agccgccgag	cccgggctcc	cgagagcaac	ctcgtgacct	5940
cagcatgcca	ttcgttcgtg	agtgtcccgt	gcacgtcctg	accctggaca	gtgaaggctt	6000
ctccctcccc	tcccacctga	cttcatctac	gcctcggcac	cacggggtgt	gggaccccag	6060
cccggagaga	acagagggaa	ggagccgcgc	ccccacctgg	agctgaatca	catgacctag	6120
ctgcacccca	gcgctggggc	ccgccccacg	ctctgtccac	accacgcgc	cccgggagcg	6180
ggcccatgcc	tccagcccc	cagctcggcc	gacccatcct	gttcgtgaat	aggtctcagg	6240
ggttggggga	gggactgcca	gatttggaca	ctatatTTTT	ttctaaattc	aacttgaaga	6300
tgtgtatttc	ccctgacctt	caaaaaatgt	tccaaggtaa	gcctcgtaaa	ggtcatccca	6360
ccatcaccaa	agcctccgtt	tttaacaacc	tccaacacga	tccatttaga	ggccaaatgt	6420
cattctgcag	gtgccttccc	gatggattaa	aggtgcttat	gtttttgtga	gttttaagta	6480
aatatttgta	ttgtattggt	ataaatgtta	agtgtgcctg	gctttcaatc	atgcacggaa	6540
accagtctc	agtcccacgg	acagaatggg	cgaggcatgg	attctggggt	gcagtaccgt	6600
tctgattaga	aataggaagt	ctccccaccc	ccgcccctggc	caagaacgtg	caataaattg	6660
gaagtttgcc	ccggggcagc	aagaatttat	gctgccattg	aaaagcaggt	accagtgccc	6720
cttttcagac	agtttttgat	tcgctctaga	cttttttttt	ttttaatagg	gaaaaaattt	6780
gataattttc	ttttttctac	atgcacttaa	gactaaaaca	caggtttggg	ttaattttat	6840
ttgcttccct	tttccgcttt	tcttcccgca	gagcctgatg	ggagaatgtc	cagggcaggg	6900
aaaccacatt	ttttgtaggt	gataactcaa	tgaaaattgg	tgcttatttt	ttacacttct	6960
ctcttggtgc	tctcttggtg	tgctatctat	ctgttttaag	gtctccttga	aggcgcactg	7020
gggacctggg	ccatgcctcg	ttctccctgc	ttcttttata	ctgttattgc	ctccacagtc	7080
tgttgccaag	gactctaaga	tcaatgcacg	tcactttcct	ttccactggg	caggatagcc	7140
aagcacactc	cctcctgcgc	tctcccgcgc	cggtgcgctc	actcccaggg	gctggtatga	7200
ggactggggt	gtgcctactt	gatttgaaaa	cacacacaag	caataaaaag	cctcttctctg	7260
cattgtctgt	ggtgtgacca	tagcagatta	tatttggttc	ctgaatgttt	gtggtgctaa	7320
tttctgtggt	tgttccaagc	cgttcagtca	tgccatgcgc	tgctcggta	gatggagtaa	7380
tgtacaatga	actccatgag	tctctccagg	gctgcctgca	gcacgtcttt	tccaagtagc	7440
ctatttggat	tcccatctca	aatgtcctgg	atgcgagcgt	cagcggctcc	agagctcggg	7500
gcgggtgagg	tcccctttgg	ggaacccttt	cctggccatc	gaggtcgggg	ggctgccgtc	7560
tgtgggcagg	aggacctgag	gggcagccag	gaaaggcgat	ctcttcaactg	tgaaaagtgtg	7620
cccgggtgca	gcgctttttc	cttctaccat	gggaaatgca	ggctggggcc	ttggggtgag	7680
cctgcggggc	tctggtgctg	tccccgacct	ccaccaccac	cagaatgcag	ttccagctta	7740
ggaagccaca	aacaagccac	ccaggaggaa	caaaacaccg	ccagcgtgga	ttttccaaat	7800

-continued

```

ttccctggaa agtaagtctc gctcttgcca aagaaaagtc tggcttgag agtctctgga 7860
gccaggatg ccagcatgtg ccaatgactg tcaccttcat ctcttcaaaa gaaaagccat 7920
agccgaggac tgtcccgcga cccccgtgga ctgctctag gtcattgat tctgttttca 7980
tttctcatcc catccaattt gtcttttct cctgtcattt tcttctctg tggtccttc 8040
aaagttgta taattgtac tgaacttcaa aatgtgtccc gttctccca gaccactcta 8100
gccacagtat attgcaataa aattacttct tatatttgca gaaattcttt tgggtgtaatt 8160
ttatttttct ctctcaatat atataattgg acaaacgctg gcaaaaagaa aaaaatggta 8220
agcaaaaaac ccaagataaa gtttcgagga catcaggcct tttgaaatac aatgtcaaat 8280
gacacattgt acggtttcaa aaaatccgct agacatgtca taagttttaa ctgtaatgcc 8340
caggaaagga tatcttaaaa tattctaaac ttgtgtaaca aaggaataat taactgtaat 8400
agtttttcaa taaatcgagt tgggtgtttc caccgtaaa 8439

```

<210> SEQ ID NO 2

<211> LENGTH: 1838

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

```

Met Asp Val His Thr Arg Trp Lys Ala Arg Ser Ala Leu Arg Pro Gly
 1             5             10             15

Ala Pro Leu Leu Pro Pro Leu Leu Leu Leu Leu Trp Ala Pro Pro
          20             25             30

Pro Ser Arg Ala Ala Gln Pro Ala Asp Leu Leu Lys Val Leu Asp Phe
          35             40             45

His Asn Leu Pro Asp Gly Ile Thr Lys Thr Thr Gly Phe Cys Ala Thr
 50             55             60

Arg Arg Ser Ser Lys Gly Pro Asp Val Ala Tyr Arg Val Thr Lys Asp
65             70             75             80

Ala Gln Leu Ser Ala Pro Thr Lys Gln Leu Tyr Pro Ala Ser Ala Phe
          85             90             95

Pro Glu Asp Phe Ser Ile Leu Thr Thr Val Lys Ala Lys Lys Gly Ser
100            105            110

Gln Ala Phe Leu Val Ser Ile Tyr Asn Glu Gln Gly Ile Gln Gln Ile
115            120            125

Gly Leu Glu Leu Gly Arg Ser Pro Val Phe Leu Tyr Glu Asp His Thr
130            135            140

Gly Lys Pro Gly Pro Glu Asp Tyr Pro Leu Phe Arg Gly Ile Asn Leu
145            150            155            160

Ser Asp Gly Lys Trp His Arg Ile Ala Leu Ser Val His Lys Lys Asn
165            170            175

Val Thr Leu Ile Leu Asp Cys Lys Lys Lys Thr Thr Lys Phe Leu Asp
180            185            190

Arg Ser Asp His Pro Met Ile Asp Ile Asn Gly Ile Ile Val Phe Gly
195            200            205

Thr Arg Ile Leu Asp Glu Glu Val Phe Glu Gly Asp Ile Gln Gln Leu
210            215            220

Leu Phe Val Ser Asp His Arg Ala Ala Tyr Asp Tyr Cys Glu His Tyr
225            230            235            240

Ser Pro Asp Cys Asp Thr Ala Val Pro Asp Thr Pro Gln Ser Gln Asp
245            250            255

Pro Asn Pro Asp Glu Tyr Tyr Thr Glu Gly Asp Gly Glu Gly Glu Thr
260            265            270

```


-continued

Tyr Tyr Tyr Glu Tyr Pro Tyr Tyr Glu Asp Pro Glu Asp Leu Gly Lys
 275 280 285
 Glu Pro Thr Pro Ser Lys Lys Pro Val Glu Ala Ala Lys Glu Thr Thr
 290 300
 Glu Val Pro Glu Glu Leu Thr Pro Thr Pro Thr Glu Ala Ala Pro Met
 305 310 315 320
 Pro Glu Thr Ser Glu Gly Ala Gly Lys Glu Glu Asp Val Gly Ile Gly
 325 330 335
 Asp Tyr Asp Tyr Val Pro Ser Glu Asp Tyr Tyr Thr Pro Ser Pro Tyr
 340 345 350
 Asp Asp Leu Thr Tyr Gly Glu Gly Glu Glu Asn Pro Asp Gln Pro Thr
 355 360 365
 Asp Pro Gly Ala Gly Ala Glu Ile Pro Thr Ser Thr Ala Asp Thr Ser
 370 375 380
 Asn Ser Ser Asn Pro Ala Pro Pro Pro Gly Glu Gly Ala Asp Asp Leu
 385 390 395 400
 Glu Gly Glu Phe Thr Glu Glu Thr Ile Arg Asn Leu Asp Glu Asn Tyr
 405 410 415
 Tyr Asp Pro Tyr Tyr Asp Pro Thr Ser Ser Pro Ser Glu Ile Gly Pro
 420 425 430
 Gly Met Pro Ala Asn Gln Asp Thr Ile Tyr Glu Gly Ile Gly Gly Pro
 435 440 445
 Arg Gly Glu Lys Gly Gln Lys Gly Glu Pro Ala Ile Ile Glu Pro Gly
 450 455 460
 Met Leu Ile Glu Gly Pro Pro Gly Pro Glu Gly Pro Ala Gly Leu Pro
 465 470 475 480
 Gly Pro Pro Gly Thr Met Gly Pro Thr Gly Gln Val Gly Asp Pro Gly
 485 490 495
 Glu Arg Gly Pro Pro Gly Arg Pro Gly Leu Pro Gly Ala Asp Gly Leu
 500 505 510
 Pro Gly Pro Pro Gly Thr Met Leu Met Leu Pro Phe Arg Phe Gly Gly
 515 520 525
 Gly Gly Asp Ala Gly Ser Lys Gly Pro Met Val Ser Ala Gln Glu Ser
 530 535 540
 Gln Ala Gln Ala Ile Leu Gln Gln Ala Arg Leu Ala Leu Arg Gly Pro
 545 550 555
 Ala Gly Pro Met Gly Leu Thr Gly Arg Pro Gly Pro Val Gly Pro Pro
 565 570 575
 Gly Ser Gly Gly Leu Lys Gly Glu Pro Gly Asp Val Gly Pro Gln Gly
 580 585 590
 Pro Arg Gly Val Gln Gly Pro Pro Gly Pro Ala Gly Lys Pro Gly Arg
 595 600 605
 Arg Gly Arg Ala Gly Ser Asp Gly Ala Arg Gly Met Pro Gly Gln Thr
 610 615 620
 Gly Pro Lys Gly Asp Arg Gly Phe Asp Gly Leu Ala Gly Leu Pro Gly
 625 630 635 640
 Glu Lys Gly His Arg Gly Asp Pro Gly Pro Ser Gly Pro Pro Gly Pro
 645 650 655
 Pro Gly Asp Asp Gly Glu Arg Gly Asp Asp Gly Glu Val Gly Pro Arg
 660 665 670
 Gly Leu Pro Gly Glu Pro Gly Pro Arg Gly Leu Leu Gly Pro Lys Gly
 675 680 685
 Pro Pro Gly Pro Pro Gly Pro Pro Gly Val Thr Gly Met Asp Gly Gln

-continued

690			695			700									
Pro 705	Gly	Pro	Lys	Gly	Asn 710	Val	Gly	Pro	Gln	Gly	Glu	Pro	Gly	Pro	Pro 720
Gly	Gln	Gln	Gly	Asn 725	Pro	Gly	Ala	Gln	Gly	Leu	Pro	Gly	Pro	Gln 735	Gly
Ala	Ile	Gly	Pro	Pro	Gly	Glu	Lys	Gly	Pro	Leu	Gly	Lys	Pro	Gly	Leu
Pro	Gly	Met 755	Pro	Gly	Ala	Asp	Gly	Pro	Pro	Gly	His	Pro	Gly	Lys	Glu
Gly	Pro	Pro	Gly	Glu	Lys	Gly	Gly	Gln	Gly	Pro	Pro	Gly	Pro	Gln	Gly
Pro	Ile	Gly	Tyr	Pro	Gly	Pro	Arg	Gly	Val	Lys	Gly	Ala	Asp	Gly	Ile 800
Arg	Gly	Leu	Lys	Gly	Thr	Lys	Gly	Glu	Lys	Gly	Glu	Asp	Gly	Phe	Pro
Gly	Phe	Lys	Gly	Asp	Met	Gly	Ile	Lys	Gly	Asp	Arg	Gly	Glu	Ile	Gly
Pro	Pro	Gly	Pro	Arg	Gly	Glu	Asp	Gly	Pro	Glu	Gly	Pro	Lys	Gly	Arg
Gly	Gly	Pro	Asn	Gly	Asp	Pro	Gly	Pro	Leu	Gly	Pro	Pro	Gly	Glu	Lys
Gly	Lys	Leu	Gly	Val	Pro	Gly	Leu	Pro	Gly	Tyr	Pro	Gly	Arg	Gln	Gly 880
Pro	Lys	Gly	Ser	Ile	Gly	Phe	Pro	Gly	Phe	Pro	Gly	Ala	Asn	Gly	Glu 895
Lys	Gly	Gly	Arg	Gly	Thr	Pro	Gly	Lys	Pro	Gly	Pro	Arg	Gly	Gln	Arg
Gly	Pro	Thr	Gly	Pro	Arg	Gly	Glu	Arg	Gly	Pro	Arg	Gly	Ile	Thr	Gly
Lys	Pro	Gly	Pro	Lys	Gly	Asn	Ser	Gly	Gly	Asp	Gly	Pro	Ala	Gly	Pro
Pro	Gly	Glu	Arg	Gly	Pro	Asn	Gly	Pro	Gln	Gly	Pro	Thr	Gly	Phe	Pro 960
Gly	Pro	Lys	Gly	Pro	Pro	Gly	Pro	Pro	Gly	Lys	Asp	Gly	Leu	Pro	Gly 975
His	Pro	Gly	Gln	Arg	Gly	Glu	Thr	Gly	Phe	Gln	Gly	Lys	Thr	Gly	Pro 990
Pro	Gly	Pro	Pro	Gly	Val	Val	Gly	Pro	Gln	Gly	Pro	Thr	Gly	Glu	Thr
Gly	Pro	Met	Gly	Glu	Arg	Gly	His	Pro	Gly	Pro	Pro	Gly	Pro	Pro	Gly
Glu	Gln	Gly	Leu	Pro	Gly	Leu	Ala	Gly	Lys	Glu	Gly	Thr	Lys	Gly	Asp 1040
Pro	Gly	Pro	Ala	Gly	Leu	Pro	Gly	Lys	Asp	Gly	Pro	Pro	Gly	Leu	Arg 1055
Gly	Phe	Pro	Gly	Asp	Arg	Gly	Leu	Pro	Gly	Pro	Val	Gly	Ala	Leu	Gly
Leu	Lys	Gly	Asn	Glu	Gly	Pro	Pro	Gly	Pro	Pro	Gly	Pro	Ala	Gly	Ser 1085
Pro	Gly	Glu	Arg	Gly	Pro	Ala	Gly	Ala	Ala	Gly	Pro	Ile	Gly	Ile	Pro 1100
Gly	Arg	Pro	Gly	Pro	Gln	Gly	Pro	Pro	Gly	Pro	Ala	Gly	Glu	Lys	Gly 1120

-continued

Ala Pro Gly Glu Lys Gly Pro Gln Gly Pro Ala Gly Arg Asp Gly Leu
1125 1130 1135

Gln Gly Pro Val Gly Leu Pro Gly Pro Ala Gly Pro Val Gly Pro Pro
1140 1145 1150

Gly Glu Asp Gly Asp Lys Gly Glu Ile Gly Glu Pro Gly Gln Lys Gly
1155 1160 1165

Ser Lys Gly Asp Lys Gly Glu Gln Gly Pro Pro Gly Pro Thr Gly Pro
1170 1175 1180

Gln Gly Pro Ile Gly Gln Pro Gly Pro Ser Gly Ala Asp Gly Glu Pro
1185 1190 1195 1200

Gly Pro Arg Gly Gln Gln Gly Leu Phe Gly Gln Lys Gly Asp Glu Gly
1205 1210 1215

Pro Arg Gly Phe Pro Gly Pro Pro Gly Pro Val Gly Leu Gln Gly Leu
1220 1225 1230

Pro Gly Pro Pro Gly Glu Lys Gly Glu Thr Gly Asp Val Gly Gln Met
1235 1240 1245

Gly Pro Pro Gly Pro Pro Gly Pro Arg Gly Pro Ser Gly Ala Pro Gly
1250 1255 1260

Ala Asp Gly Pro Gln Gly Pro Pro Gly Gly Ile Gly Asn Pro Gly Ala
1265 1270 1275 1280

Val Gly Glu Lys Gly Glu Pro Gly Glu Ala Gly Glu Pro Gly Leu Pro
1285 1290 1295

Gly Glu Gly Gly Pro Pro Gly Pro Lys Gly Glu Arg Gly Glu Lys Gly
1300 1305 1310

Glu Ser Gly Pro Ser Gly Ala Ala Gly Pro Pro Gly Pro Lys Gly Pro
1315 1320 1325

Pro Gly Asp Asp Gly Pro Lys Gly Ser Pro Gly Pro Val Gly Phe Pro
1330 1335 1340

Gly Asp Pro Gly Pro Pro Gly Glu Pro Gly Pro Ala Gly Gln Asp Gly
1345 1350 1355 1360

Pro Pro Gly Asp Lys Gly Asp Asp Gly Glu Pro Gly Gln Thr Gly Ser
1365 1370 1375

Pro Gly Pro Thr Gly Glu Pro Gly Pro Ser Gly Pro Pro Gly Lys Arg
1380 1385 1390

Gly Pro Pro Gly Pro Ala Gly Pro Glu Gly Arg Gln Gly Glu Lys Gly
1395 1400 1405

Ala Lys Gly Glu Ala Gly Leu Glu Gly Pro Pro Gly Lys Thr Gly Pro
1410 1415 1420

Ile Gly Pro Gln Gly Ala Pro Gly Lys Pro Gly Pro Asp Gly Leu Arg
1425 1430 1435 1440

Gly Ile Pro Gly Pro Val Gly Glu Gln Gly Leu Pro Gly Ser Pro Gly
1445 1450 1455

Pro Asp Gly Pro Pro Gly Pro Met Gly Pro Pro Gly Leu Pro Gly Leu
1460 1465 1470

Lys Gly Asp Ser Gly Pro Lys Gly Glu Lys Gly His Pro Gly Leu Ile
1475 1480 1485

Gly Leu Ile Gly Pro Pro Gly Glu Gln Gly Glu Lys Gly Asp Arg Gly
1490 1495 1500

Leu Pro Gly Pro Gln Gly Ser Ser Gly Pro Lys Gly Glu Gln Gly Ile
1505 1510 1515 1520

Thr Gly Pro Ser Gly Pro Ile Gly Pro Pro Gly Pro Pro Gly Leu Pro
1525 1530 1535

Gly Pro Pro Gly Pro Lys Gly Ala Lys Gly Ser Ser Gly Pro Thr Gly
1540 1545 1550

-continued

Pro Lys Gly Glu Ala Gly His Pro Gly Pro Pro Gly Pro Pro Gly Pro
 1555 1560 1565

Pro Gly Glu Val Ile Gln Pro Leu Pro Ile Gln Ala Ser Arg Thr Arg
 1570 1575 1580

Arg Asn Ile Asp Ala Ser Gln Leu Leu Asp Asp Gly Asn Gly Glu Asn
 1585 1590 1595 1600

Tyr Val Asp Tyr Ala Asp Gly Met Glu Glu Ile Phe Gly Ser Leu Asn
 1605 1610 1615

Ser Leu Lys Leu Glu Ile Glu Gln Met Lys Arg Pro Leu Gly Thr Gln
 1620 1625 1630

Gln Asn Pro Ala Arg Thr Cys Lys Asp Leu Gln Leu Cys His Pro Asp
 1635 1640 1645

Phe Pro Asp Gly Glu Tyr Trp Val Asp Pro Asn Gln Gly Cys Ser Arg
 1650 1655 1660

Asp Ser Phe Lys Val Tyr Cys Asn Phe Thr Ala Gly Gly Ser Thr Cys
 1665 1670 1675 1680

Val Phe Pro Asp Lys Lys Ser Glu Gly Ala Arg Ile Thr Ser Trp Pro
 1685 1690 1695

Lys Glu Asn Pro Gly Ser Trp Phe Ser Glu Phe Lys Arg Gly Lys Leu
 1700 1705 1710

Leu Ser Tyr Val Asp Ala Glu Gly Asn Pro Val Gly Val Val Gln Met
 1715 1720 1725

Thr Phe Leu Arg Leu Leu Ser Ala Ser Ala His Gln Asn Val Thr Tyr
 1730 1735 1740

His Cys Tyr Gln Ser Val Ala Trp Gln Asp Ala Ala Thr Gly Ser Tyr
 1745 1750 1755 1760

Asp Lys Ala Leu Arg Phe Leu Gly Ser Asn Asp Glu Glu Met Ser Tyr
 1765 1770 1775

Asp Asn Asn Pro Tyr Ile Arg Ala Leu Val Asp Gly Cys Ala Thr Lys
 1780 1785 1790

Lys Gly Tyr Gln Lys Thr Val Leu Glu Ile Asp Thr Pro Lys Val Glu
 1795 1800 1805

Gln Val Pro Ile Val Asp Ile Met Phe Asn Asp Phe Gly Glu Ala Ser
 1810 1815 1820

Gln Lys Phe Gly Phe Glu Val Gly Pro Ala Cys Phe Met Gly
 1825 1830 1835

<210> SEQ ID NO 3
 <211> LENGTH: 6930
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

gaccgttgct tggcagacac tggatgggta tgagcctgaa caagctgaaa aggggcagga 60
 aaagaagtgg aggcagcatt cttcctatth aaagctgcat cgcttgaaaa aagttttcgc 120
 agactgtgct ggagctggtg ctgaaaaagg gggtttgag aggctgcctt ggggctggtg 180
 ctgaaagaag agcccacagc tgacttcatg gtgctacaat aacctcagaa tctacttttc 240
 actctcagga gaaccacat gtctaattatt tagacatgat ggcaaactgg gcggaagcaa 300
 gacctctcct cattcttatt gttttattag ggcaatttgt ctcaataaaa gccaggaag 360
 aagacgagga tgaaggatat ggtgaagaaa tagcctgcac tcagaatggc cagatgtact 420
 taaacagggg catttggaag cctgcccctt gtcagatctg tgtctgtgac aatggagcca 480
 ttctctgtga caagatagaa tgccaggatg tgctggactg tgccgaccct gtaacgcccc 540

-continued

ctggggaatg	ctgtcctgtc	tggtcacaaa	cacctggagg	tggcaataca	aatdddggta	600
gaggaagaaa	gggacaaaag	ggagaaccag	gattagtgcc	tggtgtaaca	ggcatatcgtg	660
gtcgtccagg	accggcagga	cctccaggat	cacagggacc	aagaggagag	cgagggccaa	720
aaggaagacc	tggccctcgt	ggacctcagg	gaattgatgg	agaaccaggt	gttctctggtc	780
aacctggtgc	tccaggacct	cctggacatc	cgtdccaccc	aggacctgat	ggcttgagca	840
ggccgttttc	agctcaaatg	gctgggttgg	atgaaaaatc	tggacttggg	agtcaagtag	900
gactaatgcc	tggctctgtg	ggctctgttg	gcccaggggg	accacagggg	ttacaaggac	960
agcaaggtgg	tgcaggacct	acaggacctc	ctggtgaacc	tggatgatcct	ggaccaatgg	1020
gtccgattgg	ttcacgtgga	ccagagggcc	ctctggtaa	acctggggaa	gatggtgaac	1080
ctggcagaaa	tggaaatcct	ggtgaagtgg	gatttgacgg	atctccggga	gctcgtggat	1140
ttctggggc	tcctggtctt	ccaggtctga	agggtcaccg	aggacacaaa	ggtcttgaag	1200
gccctaaagg	tgaagttgga	gcacctgggt	ccaaggggga	agctggcccc	actgggtccaa	1260
tgggtgccat	gggtcctctg	ggctccaggg	gaatgccagg	agagagaggg	agacttgggc	1320
cacaggtgct	tcctggacaa	cgaggtgcac	atggtatgcc	tggaaaacct	ggaccaatgg	1380
gtcctcttgg	gataccaggc	tcttctgggt	ttccaggaaa	tcctggaatg	aagggagaag	1440
caggtcctac	aggggcgcga	ggccctgaag	gtcctcaggg	gcagagaggt	gaaactgggc	1500
ccccaggtcc	agttggctct	ccaggtcttc	ctggtgcaat	aggaactgat	ggtactcctg	1560
gtgcccagg	cccaacgggc	tctccgggta	cctctgggtc	tcctgggtca	gcagggcctc	1620
ctggatctcc	aggacctcag	ggtagcactg	gtcctcaggg	aattcagggc	caaccgggtg	1680
atccaggagt	tccaggtttc	aaaggagaag	ctggcccaaa	aggggaacca	gggccacatg	1740
gtattcaggg	tccgataggc	ccaccgggtg	aagaaggcaa	aagaggtccc	agaggtgacc	1800
caggaacagt	tggctctcca	gggccagtgg	gagaaagggg	tgctcctggc	aatcgtgggt	1860
ttccaggctc	tgatggttta	cctgggccaa	agggtgctca	aggagaacgg	ggtcctgtag	1920
gttcttcagg	acccaaagga	agccaggggg	atccaggacg	tccaggggaa	cctgggcttc	1980
caggtgctcg	gggtttgaca	ggaaatcctg	gtgttcaagg	tcctgaagga	aaacttgac	2040
ctttgggtgc	gccaggggaa	gatggccgtc	caggtcctcc	aggctccata	ggaatcagag	2100
ggcagcccgg	gagcatgggc	cttccaggcc	ccaaaggtag	cagtgggtgac	cctgggaaac	2160
ctggagaagc	aggaaatgct	ggagttcctg	ggcagagggg	agctcctgga	aaagatgggtg	2220
aagtgggtcc	ttctggtcct	gtgggcccgc	cggtctagc	tggtgaaaga	ggagaacaag	2280
gacctccagg	ccccacaggt	tttcaggggc	ttcctgggtc	tccagggcct	cctggagaag	2340
gtggaaaacc	aggtgatcaa	ggtgttctctg	gagatcccgg	agcagttggc	ccgttaggac	2400
ctagaggaga	acgaggaaat	cctggggaaa	gaggagaacc	tgggataact	ggactccctg	2460
gtgagaaggg	aatggctgga	ggacatggtc	ctgatggccc	aaaaggcagt	ccaggtccat	2520
ctgggacccc	tggagataca	ggcccaccag	gtcttcaagg	tatgccggga	gaaagaggaa	2580
ttgcaggaac	tcctggcccc	aaggggtgaca	gaggtggcat	aggagaaaaa	ggtgctgaag	2640
gcacagctgg	aatgatgggt	gcaagaggtc	ttccagggtc	tttgggcctc	ccaggtccgg	2700
caggtcctac	tggagaaaag	ggtgaacctg	gtcctcaggg	tttagttggc	cctcctggct	2760
cccggggcaa	tcctggttct	cgaggtgaaa	atgggccaac	tggagctgtt	ggttttgccg	2820
gaccccaggg	tcctgacgga	cagcctggag	taaaagggtga	acctggagag	ccaggacaga	2880
aggagatgc	tggttctcct	ggaccacaag	gttagcagg	atcccctggc	cctcatggtc	2940

-continued

ctaattggtgt	tcttggacta	aaaggtggtc	gaggaacca	aggtecgct	ggtgctacag	3000
gatttcctgg	ttctgctggc	agagttggac	ctccaggccc	tgctggagct	ccaggacctg	3060
cgggacctct	aggggaacct	gggaaggagg	gacctccagg	tcttctggg	gacctggct	3120
ctcatgggag	tgtgggagat	cgaggaccag	ctggccccc	tggtggccca	ggagacaaag	3180
gggacctcag	agaagatggg	caacctggtc	cagatggccc	ccctggtcca	gctggaacga	3240
ccgggcagag	aggaattggt	ggcatgcctg	ggcaactggg	agagagaggc	atgcccggcc	3300
taccaggccc	agcgggaaca	ccaggaaaag	taggaccaac	tggtgcaaca	ggagataaag	3360
gtccacctgg	acctgtgggg	ccccaggct	ccaatggtcc	tgtaggggaa	cctggaccag	3420
aaggtccagc	tggcaatgat	ggtaccccag	gacgggatgg	tgctggtgga	gaactgggtg	3480
atcgtggaga	ccctgggccc	gcaggtctgc	caggctctca	gggtgcccct	ggaactcctg	3540
gcctgtggg	tgctccagga	gatgcaggac	aaagaggaga	tccgggttct	cggggtccta	3600
taggaccacc	tggtcgagct	gggaaactg	gattacctgg	acccaagga	cctcgtgggtg	3660
aaaaggtga	tcatggagac	cgaggcgaca	gaggtcagaa	gggccacaga	ggctttactg	3720
gtcttcaggg	tcttctggc	cctcctggtc	caaatggtga	acaaggaagt	gctggaatcc	3780
ctggaccatt	tggcccaaga	ggtcctccag	gccagttgg	tccttcaggt	aaagaaggaa	3840
accctgggcc	acttgggcca	attggacctc	caggtgtacg	aggcagtgt	ggagaagcag	3900
gacctgaggg	ccctcctggt	gagcctggcc	cacctggccc	tccgggtccc	cctggccacc	3960
ttacagctgc	tcttggggat	atcatggggc	actatgatga	aagcatgcca	gatccacttc	4020
ctgagtttac	tgaagatcag	gcggtcctctg	atgacaaaa	caaacggac	ccaggggttc	4080
atgctacctc	gaagtcactc	agtagtcaga	ttgaaacct	gcgcagcccc	gatggctcga	4140
aaaagcacc	agcccgcacg	tgtgatgacc	taaagctttg	ccattccgca	aagcagagtg	4200
gtgaatactg	gattgatcct	aaccaaggat	ctgttgaaga	tgcaatcaa	gtttactgca	4260
acatggaaac	aggagaaaca	tgtatttcag	caaacctac	cagtgtacca	cgtaaacct	4320
ggtgggcccag	taaactctct	gacaataaac	ctgtttggt	tggtcttgat	atgaacagag	4380
ggtctcagtt	cgcttatgga	gaccaccaat	cacctaatc	agccattact	cagatgactt	4440
ttttgcgct	tttatcaaaa	gaagcctccc	agaacatcac	ttacatctgt	aaaaacagtg	4500
taggatacat	ggacgatcaa	gctaagaacc	tcaaaaaagc	tgtggttctc	aaaggggcaa	4560
atgacttaga	tatcaaagca	gagggaaata	ttagattccg	gtatatcggt	cttcaagaca	4620
cttgctctaa	gcggaatgga	aatgtgggca	agactgtctt	tgaatataga	acacagaatg	4680
tggcacgctt	gcccatacata	gatcttgctc	ctgtggatgt	tggcggcaca	gaccaggaat	4740
tcggcggtga	aattgggcca	gtttgttttg	tgtaaagtaa	gccaagacac	atcgacaatg	4800
agcaccacca	tcaatgacca	ccgccattca	caagaacttt	gactgtttga	agttgatcct	4860
gagactcttg	aagtaatggc	tgatcctgca	tcagcattgt	atatatggtc	ttaagtgcct	4920
ggcctcctta	tccttcagaa	tatttatttt	acttacaatc	ctcaagtttt	aattgatttt	4980
aaatattttt	caatacaaca	gtttaggttt	aagatgacca	atgacaatga	ccacctttgc	5040
agaaagttaa	ctgattgaat	aaataaatct	ccgttttctt	caatttattt	cagtgtaatg	5100
aaaaagttag	ttagtattta	tgaggaaatt	cttcttctctg	gcaggtagct	taaagagtgg	5160
ggtatataga	gccacaacac	atgtttattt	tgcttgctg	cagttgaaaa	atagaaatta	5220
gtgccctttt	gtgacctctc	attccaagat	tgtcaattaa	aatgagttt	aaaatgttta	5280
acttgtgatc	gagacctaca	tgcattgctt	gatattgtgt	aactataata	gagactcttt	5340

-continued

```

aaggagaatc ttaaaaaaaaa aaaaacgttt ctcaactgtct taaatagaat ttttaaatag 5400
tatatatcca gtggcatttt ggagaacaaa gtgaatttac ttcgacttct taaatttttg 5460
taaaagacta taagttaga catctttctc attcaaattt aaagatatct ttctcctctt 5520
gatcaatcta tcaatattga tagaagtcac actagtatat accatttaat acatttacac 5580
tttcttattt aagaagatat tgaatgcaaa ataattgaca tatagaactt tacaaacata 5640
tgtccaagga ctctaaattg agactcttcc acatgtacaa tctcatcatc ctgaagccta 5700
taatgaagaa aaagatctag aaactgagtt gtggagctga ctctaatcaa atgtgatgat 5760
tggattaga ccatttggcc tttgaacttt cataggaaaa atgacceaac atttcttagc 5820
atgagctacc tcatctctag aagctgggat ggacttacta ttcttgttta tatttttagat 5880
actgaaaggt gctatgcttc tgttattatt ccaagactgg agataggcag ggctaaaaag 5940
gtattattat ttttccttta atgatggtgc taaaattctt cctataaaat tccttaaaaa 6000
taaagatggt ttaatcacta ccatttgtaa aacataactg ttagacttcc cgtttctgaa 6060
agaaagagca tcgttccaat gcttggtcac tgttcctctg tcatactgta tctggaatgc 6120
tttgaatac ttgcatgctt cttagaccag aacatgtagg tccccttggt tctcaatact 6180
tttttttct taattgcatt tgttggctct attttaattt ttttcttta aaataaacag 6240
ctgggaccat cccaaaagac aagccatgca tacaactttg gtcatgtatc tctgcaaagc 6300
atcaaattaa atgcacgctt ttgtcatgtc agtggttttt gttttgtgaa attcctttga 6360
ccatattaga tctatttcat ttccaatagt gaaaaggaga tgtggtggtta tactttgttt 6420
gccatttggt taaaagatac aacggatacc ttctatcatg tatgtactgg cttataaatg 6480
aaaatctatc tacaacatta cccacaaagg caacatgaca ccaattatca ctgcctctgc 6540
ccttaaaaat gtcagagtag tattattgat aaaaagggca agcaatagat ttttcatgac 6600
tgaataaact gtaataataa aacatattgtc tcaaagtgtg tcacatatga atttagccta 6660
attgttttca gtttcattct caatatttag ttacaacat cattttcccc taaactggtt 6720
atattttgac ctgtatatct taaatttgag tatttatatg cctaaataga tgtgtgagtt 6780
ttgtttgact tccaagtcca aactataaga ttatataagt tcatatagat gaatcagaaa 6840
tatgtggtaa tactattaag tcacaaacac taacaatttc caactataga aataacagtt 6900
cttatttggg ttttgggaat gctaccaata 6930

```

```

<210> SEQ ID NO 4
<211> LENGTH: 1499
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 4

```

```

Met Met Ala Asn Trp Ala Glu Ala Arg Pro Leu Leu Ile Leu Ile Val
 1           5           10           15
Leu Leu Gly Gln Phe Val Ser Ile Lys Ala Gln Glu Glu Asp Glu Asp
 20           25           30
Glu Gly Tyr Gly Glu Glu Ile Ala Cys Thr Gln Asn Gly Gln Met Tyr
 35           40           45
Leu Asn Arg Asp Ile Trp Lys Pro Ala Pro Cys Gln Ile Cys Val Cys
 50           55           60
Asp Asn Gly Ala Ile Leu Cys Asp Lys Ile Glu Cys Gln Asp Val Leu
 65           70           75           80
Asp Cys Ala Asp Pro Val Thr Pro Pro Gly Glu Cys Cys Pro Val Cys
 85           90           95

```


-continued

Ser Gln Thr Pro Gly Gly Gly Asn Thr Asn Phe Gly Arg Gly Arg Lys
 100 105 110

Gly Gln Lys Gly Glu Pro Gly Leu Val Pro Val Val Thr Gly Ile Arg
 115 120 125

Gly Arg Pro Gly Pro Ala Gly Pro Pro Gly Ser Gln Gly Pro Arg Gly
 130 135 140

Glu Arg Gly Pro Lys Gly Arg Pro Gly Pro Arg Gly Pro Gln Gly Ile
 145 150 155 160

Asp Gly Glu Pro Gly Val Pro Gly Gln Pro Gly Ala Pro Gly Pro Pro
 165 170 175

Gly His Pro Ser His Pro Gly Pro Asp Gly Leu Ser Arg Pro Phe Ser
 180 185 190

Ala Gln Met Ala Gly Leu Asp Glu Lys Ser Gly Leu Gly Ser Gln Val
 195 200 205

Gly Leu Met Pro Gly Ser Val Gly Pro Val Gly Pro Arg Gly Pro Gln
 210 215 220

Gly Leu Gln Gly Gln Gln Gly Gly Ala Gly Pro Thr Gly Pro Pro Gly
 225 230 235 240

Glu Pro Gly Asp Pro Gly Pro Met Gly Pro Ile Gly Ser Arg Gly Pro
 245 250 255

Glu Gly Pro Pro Gly Lys Pro Gly Glu Asp Gly Glu Pro Gly Arg Asn
 260 265 270

Gly Asn Pro Gly Glu Val Gly Phe Ala Gly Ser Pro Gly Ala Arg Gly
 275 280 285

Phe Pro Gly Ala Pro Gly Leu Pro Gly Leu Lys Gly His Arg Gly His
 290 295 300

Lys Gly Leu Glu Gly Pro Lys Gly Glu Val Gly Ala Pro Gly Ser Lys
 305 310 315 320

Gly Glu Ala Gly Pro Thr Gly Pro Met Gly Ala Met Gly Pro Leu Gly
 325 330 335

Pro Arg Gly Met Pro Gly Glu Arg Gly Arg Leu Gly Pro Gln Gly Ala
 340 345 350

Pro Gly Gln Arg Gly Ala His Gly Met Pro Gly Lys Pro Gly Pro Met
 355 360 365

Gly Pro Leu Gly Ile Pro Gly Ser Ser Gly Phe Pro Gly Asn Pro Gly
 370 375 380

Met Lys Gly Glu Ala Gly Pro Thr Gly Ala Arg Gly Pro Glu Gly Pro
 385 390 395 400

Gln Gly Gln Arg Gly Glu Thr Gly Pro Pro Gly Pro Val Gly Ser Pro
 405 410 415

Gly Leu Pro Gly Ala Ile Gly Thr Asp Gly Thr Pro Gly Ala Lys Gly
 420 425 430

Pro Thr Gly Ser Pro Gly Thr Ser Gly Pro Pro Gly Ser Ala Gly Pro
 435 440 445

Pro Gly Ser Pro Gly Pro Gln Gly Ser Thr Gly Pro Gln Gly Ile Arg
 450 455 460

Gly Gln Pro Gly Asp Pro Gly Val Pro Gly Phe Lys Gly Glu Ala Gly
 465 470 475 480

Pro Lys Gly Glu Pro Gly Pro His Gly Ile Gln Gly Pro Ile Gly Pro
 485 490 495

Pro Gly Glu Glu Gly Lys Arg Gly Pro Arg Gly Asp Pro Gly Thr Val
 500 505 510

Gly Pro Pro Gly Pro Val Gly Glu Arg Gly Ala Pro Gly Asn Arg Gly

-continued

515					520					525					
Phe	Pro	Gly	Ser	Asp	Gly	Leu	Pro	Gly	Pro	Lys	Gly	Ala	Gln	Gly	Glu
	530					535					540				
Arg	Gly	Pro	Val	Gly	Ser	Ser	Gly	Pro	Lys	Gly	Ser	Gln	Gly	Asp	Pro
545					550					555					560
Gly	Arg	Pro	Gly	Glu	Pro	Gly	Leu	Pro	Gly	Ala	Arg	Gly	Leu	Thr	Gly
				565					570					575	
Asn	Pro	Gly	Val	Gln	Gly	Pro	Glu	Gly	Lys	Leu	Gly	Pro	Leu	Gly	Ala
			580					585					590		
Pro	Gly	Glu	Asp	Gly	Arg	Pro	Gly	Pro	Pro	Gly	Ser	Ile	Gly	Ile	Arg
		595					600					605			
Gly	Gln	Pro	Gly	Ser	Met	Gly	Leu	Pro	Gly	Pro	Lys	Gly	Ser	Ser	Gly
	610					615					620				
Asp	Pro	Gly	Lys	Pro	Gly	Glu	Ala	Gly	Asn	Ala	Gly	Val	Pro	Gly	Gln
625					630					635					640
Arg	Gly	Ala	Pro	Gly	Lys	Asp	Gly	Glu	Val	Gly	Pro	Ser	Gly	Pro	Val
				645					650					655	
Gly	Pro	Pro	Gly	Leu	Ala	Gly	Glu	Arg	Gly	Glu	Gln	Gly	Pro	Pro	Gly
			660					665					670		
Pro	Thr	Gly	Phe	Gln	Gly	Leu	Pro	Gly	Pro	Pro	Gly	Pro	Pro	Gly	Glu
		675					680					685			
Gly	Gly	Lys	Pro	Gly	Asp	Gln	Gly	Val	Pro	Gly	Asp	Pro	Gly	Ala	Val
	690					695					700				
Gly	Pro	Leu	Gly	Pro	Arg	Gly	Glu	Arg	Gly	Asn	Pro	Gly	Glu	Arg	Gly
705					710					715					720
Glu	Pro	Gly	Ile	Thr	Gly	Leu	Pro	Gly	Glu	Lys	Gly	Met	Ala	Gly	Gly
				725					730					735	
His	Gly	Pro	Asp	Gly	Pro	Lys	Gly	Ser	Pro	Gly	Pro	Ser	Gly	Thr	Pro
			740					745					750		
Gly	Asp	Thr	Gly	Pro	Pro	Gly	Leu	Gln	Gly	Met	Pro	Gly	Glu	Arg	Gly
		755					760					765			
Ile	Ala	Gly	Thr	Pro	Gly	Pro	Lys	Gly	Asp	Arg	Gly	Gly	Ile	Gly	Glu
	770					775				780					
Lys	Gly	Ala	Glu	Gly	Thr	Ala	Gly	Asn	Asp	Gly	Ala	Arg	Gly	Leu	Pro
785						790				795					800
Gly	Pro	Leu	Gly	Pro	Pro	Gly	Pro	Ala	Gly	Pro	Thr	Gly	Glu	Lys	Gly
				805					810					815	
Glu	Pro	Gly	Pro	Arg	Gly	Leu	Val	Gly	Pro	Pro	Gly	Ser	Arg	Gly	Asn
			820					825					830		
Pro	Gly	Ser	Arg	Gly	Glu	Asn	Gly	Pro	Thr	Gly	Ala	Val	Gly	Phe	Ala
		835					840					845			
Gly	Pro	Gln	Gly	Pro	Asp	Gly	Gln	Pro	Gly	Val	Lys	Gly	Glu	Pro	Gly
	850					855					860				
Glu	Pro	Gly	Gln	Lys	Gly	Asp	Ala	Gly	Ser	Pro	Gly	Pro	Gln	Gly	Leu
865					870					875					880
Ala	Gly	Ser	Pro	Gly	Pro	His	Gly	Pro	Asn	Gly	Val	Pro	Gly	Leu	Lys
				885					890					895	
Gly	Gly	Arg	Gly	Thr	Gln	Gly	Pro	Pro	Gly	Ala	Thr	Gly	Phe	Pro	Gly
			900					905					910		
Ser	Ala	Gly	Arg	Val	Gly	Pro	Pro	Gly	Pro	Ala	Gly	Ala	Pro	Gly	Pro
		915						920				925			
Ala	Gly	Pro	Leu	Gly	Glu	Pro	Gly	Lys	Glu	Gly	Pro	Pro	Gly	Leu	Arg
	930						935					940			

-continued

Gly Asp Pro Gly Ser His Gly Arg Val Gly Asp Arg Gly Pro Ala Gly
 945 950 955 960
 Pro Pro Gly Gly Pro Gly Asp Lys Gly Asp Pro Gly Glu Asp Gly Gln
 965 970 975
 Pro Gly Pro Asp Gly Pro Pro Gly Pro Ala Gly Thr Thr Gly Gln Arg
 980 985 990
 Gly Ile Val Gly Met Pro Gly Gln Arg Gly Glu Arg Gly Met Pro Gly
 995 1000 1005
 Leu Pro Gly Pro Ala Gly Thr Pro Gly Lys Val Gly Pro Thr Gly Ala
 1010 1015 1020
 Thr Gly Asp Lys Gly Pro Pro Gly Pro Val Gly Pro Pro Gly Ser Asn
 1025 1030 1035 1040
 Gly Pro Val Gly Glu Pro Gly Pro Glu Gly Pro Ala Gly Asn Asp Gly
 1045 1050 1055
 Thr Pro Gly Arg Asp Gly Ala Val Gly Glu Arg Gly Asp Arg Gly Asp
 1060 1065 1070
 Pro Gly Pro Ala Gly Leu Pro Gly Ser Gln Gly Ala Pro Gly Thr Pro
 1075 1080 1085
 Gly Pro Val Gly Ala Pro Gly Asp Ala Gly Gln Arg Gly Asp Pro Gly
 1090 1095 1100
 Ser Arg Gly Pro Ile Gly Pro Pro Gly Arg Ala Gly Lys Arg Gly Leu
 1105 1110 1115 1120
 Pro Gly Pro Gln Gly Pro Arg Gly Asp Lys Gly Asp His Gly Asp Arg
 1125 1130 1135
 Gly Asp Arg Gly Gln Lys Gly His Arg Gly Phe Thr Gly Leu Gln Gly
 1140 1145 1150
 Leu Pro Gly Pro Pro Gly Pro Asn Gly Glu Gln Gly Ser Ala Gly Ile
 1155 1160 1165
 Pro Gly Pro Phe Gly Pro Arg Gly Pro Pro Gly Pro Val Gly Pro Ser
 1170 1175 1180
 Gly Lys Glu Gly Asn Pro Gly Pro Leu Gly Pro Ile Gly Pro Pro Gly
 1185 1190 1195 1200
 Val Arg Gly Ser Val Gly Glu Ala Gly Pro Glu Gly Pro Pro Gly Glu
 1205 1210 1215
 Pro Gly Pro Pro Gly Pro Pro Gly Pro Pro Gly His Leu Thr Ala Ala
 1220 1225 1230
 Leu Gly Asp Ile Met Gly His Tyr Asp Glu Ser Met Pro Asp Pro Leu
 1235 1240 1245
 Pro Glu Phe Thr Glu Asp Gln Ala Ala Pro Asp Asp Lys Asn Lys Thr
 1250 1255 1260
 Asp Pro Gly Val His Ala Thr Leu Lys Ser Leu Ser Ser Gln Ile Glu
 1265 1270 1275 1280
 Thr Met Arg Ser Pro Asp Gly Ser Lys Lys His Pro Ala Arg Thr Cys
 1285 1290 1295
 Asp Asp Leu Lys Leu Cys His Ser Ala Lys Gln Ser Gly Glu Tyr Trp
 1300 1305 1310
 Ile Asp Pro Asn Gln Gly Ser Val Glu Asp Ala Ile Lys Val Tyr Cys
 1315 1320 1325
 Asn Met Glu Thr Gly Glu Thr Cys Ile Ser Ala Asn Pro Ser Ser Val
 1330 1335 1340
 Pro Arg Lys Thr Trp Trp Ala Ser Lys Ser Pro Asp Asn Lys Pro Val
 1345 1350 1355 1360
 Trp Tyr Gly Leu Asp Met Asn Arg Gly Ser Gln Phe Ala Tyr Gly Asp
 1365 1370 1375

-continued

His Gln Ser Pro Asn Thr Ala Ile Thr Gln Met Thr Phe Leu Arg Leu
 1380 1385 1390

 Leu Ser Lys Glu Ala Ser Gln Asn Ile Thr Tyr Ile Cys Lys Asn Ser
 1395 1400 1405

 Val Gly Tyr Met Asp Asp Gln Ala Lys Asn Leu Lys Lys Ala Val Val
 1410 1415 1420

 Leu Lys Gly Ala Asn Asp Leu Asp Ile Lys Ala Glu Gly Asn Ile Arg
 1425 1430 1435 1440

 Phe Arg Tyr Ile Val Leu Gln Asp Thr Cys Ser Lys Arg Asn Gly Asn
 1445 1450 1455

 Val Gly Lys Thr Val Phe Glu Tyr Arg Thr Gln Asn Val Ala Arg Leu
 1460 1465 1470

 Pro Ile Ile Asp Leu Ala Pro Val Asp Val Gly Gly Thr Asp Gln Glu
 1475 1480 1485

 Phe Gly Val Glu Ile Gly Pro Val Cys Phe Val
 1490 1495

<210> SEQ ID NO 5
 <211> LENGTH: 6192
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

gcgagtgact gcaccgagcc cgagaagtcg ccgccccccg cagccgcccc gactgggttcc 60
 ccgccttgcc cgtgggcccc gccgggatgg ggaaccgccc ggacctgggc cagccgcggg 120
 ccggtctctg cctgctcctg gccgcgctgc agcttctgcc ggggacgcag gccgatcctg 180
 tggatgtect gaaggccctg ggtgtgcagg gaggccaggc tgggggtccc gaggggcctg 240
 gcttctgtcc ccagaggact ccagaggggtg accgggcatt cagaattggc caggccagca 300
 cgctcggcat ccccacgtgg gaactctttc cagaaggcca ctttcctgag aacttctcct 360
 tgetgateac cttgeggga cagccagcca atcagtctgt cctgctgtcc atttatgatg 420
 aaaggggtgc ccggcagttg ggctggcac tggggccagc gctgggtctc ctaggtgacc 480
 cttccgccc cctccccag caggtcaacc tcacagatgg caggtggcac cgtgtggccc 540
 tcagcataga tggtagatg gtgaccctgg tagctgactg tgaagctcag ccccctgttt 600
 tgggccatgg cccccgcttc atcagcatag ctggactcac tgtgctgggg acccaggacc 660
 ttggggaaaa gactttcgag ggagacattc aggagctgct gataagccca gatcctcagg 720
 ctgccttcca ggcttctgag cggctacctc ccgactgtga caacctggca ccggcagcca 780
 cagtggctcc ccagggtgaa ccagaaacct ctcgtcctcg gcggaagggg aagggaaaag 840
 ggaggaagaa agggcgaggt cgcaagggga agggcaggaa aaagaacaag gaaatttggg 900
 cctcaagtcc acctcctgac tccgagaga accagacctc cactgacatc cccaagacag 960
 agactccagc tccaaatctg cctccgacct ccacgccttt ggtcgtcacc tccactgtga 1020
 ctactggact caatgccag atcctagaga ggagcttggg ccctgacagt ggaaccgagc 1080
 tggggaccct ggagaccaag gcagccaggg aggatgaaga aggagatgat tccaccatgg 1140
 gccctgactt ccgggcagca gaatatccat ctcggactca gttccagatc tttcctgggtg 1200
 ctggagagaa aggagcaaaa ggagagcccc cagtgattga aaaggggag cagtttgagg 1260
 gacctccagg agccccagga cccaagggg tggttgcccc ctcaggccct cccggcccc 1320
 caggattccc tggcgaccct ggtccaccgg gccctgctgg cctcccagga atccccggca 1380
 ttgatgggat ccgaggccca ccgggactg tgatcatgat gccgttccag tttgcaggcg 1440

-continued

gctcctttaa	aggccccca	gtctcattcc	agcaggccca	ggctcaggca	gttctgcagc	1500
agactcagct	ctctatgaaa	ggccccctg	gtccagtggg	gctcactggg	cgcccaggcc	1560
ctgtgggtct	ccccgggat	ccaggtctga	aaggagagga	gggagcagaa	gggccacagg	1620
gtccccgagg	cctgcaggga	cctcatggac	cccctggccg	agtgggcaag	atgggccgcc	1680
ctggagcaga	tggagctcgg	ggcctcccag	gggacactgg	acctaagggt	gatcgtggct	1740
tcgatggcct	ccctgggctg	cctgggtgaga	agggccaaag	gggtgacttt	ggccatgtgg	1800
ggcaaccocg	tccccagga	gaggatgggt	agaggggagc	agagggacct	ccagggccca	1860
ctggccaggc	tggggagccg	ggccacagag	gactgcttgg	ccccagaggc	tctcctggcc	1920
ccacgggtcg	cccgggtgtg	actggaattg	atggtgctcc	tggtgccaaa	ggcaatgtgg	1980
gtcctccagg	agaaccaggc	cctccgggac	agcagggaaa	ccatgggtcc	cagggactcc	2040
ccggtcccca	gggactcatt	ggcactcctg	gggagaaggg	tccccctgga	aaccaggaa	2100
ttccaggcct	cccaggatcc	gatggccctc	tgggtcacc	aggacatgag	ggccccacgg	2160
gagagaaagg	ggctcagggt	ccaccagggt	cggcaggccc	tccgggctat	cctggacctc	2220
ggggagtga	gggcacttca	ggcaaccggg	gcctccaggg	ggagaaaggc	gagaaggag	2280
aggacggctt	cccaggcttc	aagggcgatg	tggggctcaa	aggtgatcag	gggaaaccgc	2340
gagctccagg	tccccgggga	gaggatgggt	ctgaggggcc	gaaggggag	gcggggcagg	2400
ctggcgagga	ggggcccca	ggctcagctg	gggagaaggg	caagcttggg	gtgccaggcc	2460
tcccaggtta	tccaggacgc	cctggacctc	aggatctat	tggatttccc	ggccccctgg	2520
gacctatagg	agagaaaggg	aagtcgggaa	agacagggca	gccaggcctg	gaaggagagc	2580
ggggaccacc	aggttcccgt	ggagagaggg	ggcaaccggg	tgccacaggg	caaccaggcc	2640
ccaagggcga	tgtgggcca	gatggagccc	ctgggatccc	tggagaaaag	ggcctcctg	2700
gtctgcaagg	ccctccagga	ttccctgggc	caaagggccc	ccctggcac	caaggtaaag	2760
atgggagacc	agggaccct	ggacagagag	gagaactggg	cttccaaggt	cagacaggcc	2820
cgctggacc	agctggtgtc	ttaggcctc	agggaaagac	aggagaagtg	ggacctctag	2880
gtgaaagggg	gcctccaggc	ccccctggac	ctcctggtga	acaaggtctt	cctggcctgg	2940
aaggcagaga	gggggccaag	ggggaactgg	gaccaccagg	acccttggg	aaagaagggc	3000
cagctggact	caggggcttt	cccggcccca	aagggggccc	tggggaccgc	ggacctactg	3060
gcttaaaggg	tgataagggc	ccccaggggc	ccgtgggggc	caatggctcc	cctggtgagc	3120
gcggtccttt	gggcccagca	ggaggcattg	gacttccctg	ccaaagtggc	agcgaaggcc	3180
ccgttggccc	tgcaggcaag	aaggggtccc	ggggagaacg	tggccccct	ggccccactg	3240
gcaaagatgg	gatcccaggg	cccctggggc	ctctgggacc	ccctggagct	gctgggcctt	3300
ctggcgagga	aggggacaag	ggggatgtgg	gtgcccccg	acacaagggg	agtaaaggcg	3360
ataaaggaga	cgcgggcca	cctggacaac	cagggatcag	gggtcctgca	ggacaccag	3420
gtccccggg	agcagacggg	gctcaggggc	gccggggacc	cccaggcctc	tttgggcaga	3480
aaggagatga	cggagtcaga	ggctttgtgg	gggtgattgg	ccctcctgga	ctgcaggggc	3540
tgccaggccc	tccgggagag	aaaggggagg	tcggagacgt	cgggtccatg	ggtcccatg	3600
gagctccagg	tcctcgggg	ccccagggcc	ccactggatc	agagggcact	ccagggctgc	3660
ctggaggagt	tggtcagcca	ggcgccgtgg	gtgagaaggg	tgagcgaggg	gacgctggag	3720
accagggccc	tccaggagcc	ccaggcatcc	cggggcccaa	gggagacatt	ggtgaaaagg	3780
gggactcagg	cccatctgga	gctgctggac	ccccaggcaa	gaaaggtccc	cctggagagg	3840

-continued

atggagccaa	agggagcgtg	ggccccacgg	ggctgcccgg	agatctaggg	ccccaggag	3900
accctggagt	ttcaggcata	gatggttccc	caggggagaa	gggagaccct	ggtgatgttg	3960
ggggaccggg	tccgctgga	gcttctgggg	agccccggcg	ccccgggccc	cccggcaaga	4020
ggggtccttc	aggccacatg	ggtcgagaag	gcagagaagg	ggagaaaggt	gccaaagggg	4080
agccaggfcc	tgatggggcc	ccaggaggga	cgggtccaat	gggggctaga	gggccccctg	4140
gacgtgtggg	gcctgagggt	cttcgaggga	tccctggccc	tgtgggtgaa	ccaggcctcc	4200
tgggagcccc	tggacagatg	ggccctcctg	gccccctggg	gccctctggc	ctcccagggc	4260
tgaagggaga	cactggcccc	aagggggaaa	agggccacat	tggattgatc	ggtctcattg	4320
gccccccggg	agaagctggt	gagaaaggag	atcaggggtt	gccaggcgtg	cagggacccc	4380
ctgggtcccaa	gggagaccct	ggtccccctg	gtcccattgg	ctctctgggc	cacctggggc	4440
ccccagggtg	ggcgggccc	ctaggacaga	aaggctcaaa	agggtctccg	gggtccatgg	4500
gccccctggg	agacactgga	cctgcaggcc	caccaggccc	cccgggtgcc	cctgccgagc	4560
tgcatgggct	gcgcaggcgc	cggcgcttcg	tcccagtcct	gcttccagtc	gtggagggcg	4620
gcctggaggga	ggtgctggcc	tcgctcacat	cgctgagctt	ggagctggag	cagctgcggc	4680
gtcctcccgg	cactgctggg	cgccccgggc	tcgtgtgcca	cgagctgcac	cgcaaccacc	4740
cgcacctgcc	tgatggggaa	tactggattg	accccaacca	gggctgcgcg	cgggactcgt	4800
tcagggtttt	ttgcaacttc	acggcggggg	gagagacctg	cctctatccc	gacaagaagt	4860
ttgagatcgt	gaaattggcc	tcctgggtcca	aggaaaagcc	tggaggctgg	tatagcacat	4920
tccgtcgagg	gaagaagttc	tcctacgtgg	acgccgacgg	gtccccagtg	aatgtcgtgc	4980
agctgaactt	cctgaaactg	ctgagtgcc	cagctcgcca	gaacttcacc	tactcctgcc	5040
agaatgcagc	tgccctggctg	gacgaagcca	cgggtgacta	cagccactcc	gcccgttccc	5100
ttggcaccaa	tggagaggag	ctgtctttca	accagacgac	agcagccact	gtcagcgtcc	5160
cccaggatgg	ctgcccggctc	cggaaaggac	agacgaagac	ccttttcgaa	ttcagctctt	5220
ctcgagcggg	atttctgccc	ctgtgggatg	tggcggccac	tgactttggc	cagacgaacc	5280
aaaagtttgg	gtttgaactg	ggccccgtct	gcttcagcag	ctgagagtgt	ccgggggtggg	5340
agggaccatg	agggagcccc	agaatggggg	gcatttggtg	ctgaggcttt	gaagccaccg	5400
tatttttcgt	tacctgtgac	tatggagcca	atgggatgtg	acttcgctca	tcacggacag	5460
tcattccttc	tcctttccag	ggtgctgggg	gctgggggtc	cctggcccaa	gggtccagcc	5520
tcctctcacc	ccattccagg	tggcactactg	cagtctggct	ctttctcccc	tcctcccca	5580
cccaagcctc	acctccccac	cccttgaacc	cccatgcaat	gagcttctaa	ctcagagctg	5640
atgaacaaaa	gcccccccac	ccccaatgcc	tgctcctca	ctcctccgct	gctgcccttc	5700
acaccttttg	gtgctacccc	tcccagaggt	taagcaactgg	atgtctcctg	atcccaggct	5760
gggaccocct	ccccacccc	ctttgatcct	ttctacttcc	acggtgaaag	gactgaggtc	5820
ggactacaga	gggaagaggg	acttcctctg	actgggttgt	gtttcttttc	ctgcctcagc	5880
ccagctctgc	aatcccctc	cccctgcccc	ccacctcccc	aggctcacct	tgccatgcca	5940
ggtggtttgg	ggaccaagat	ggtggggggg	tgaatcagga	tcctaatggt	gctgccctat	6000
ttatacctgg	gtctgtatta	aaagggaaaag	tccccctgt	tgtagatttc	atctgcttcc	6060
tccttaggga	aggctgggat	atgatgagag	attccagccc	aagcctggcc	ccccaccgcc	6120
aggccatagg	gcataatttg	catctcaaat	ctgagaataa	actgatgaac	tgtgaaaaaa	6180
aaaaaaaaaa	aa					6192

-continued

<210> SEQ ID NO 6
 <211> LENGTH: 1745
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 6

 Met Gly Asn Arg Arg Asp Leu Gly Gln Pro Arg Ala Gly Leu Cys Leu
 1 5 10 15

 Leu Leu Ala Ala Leu Gln Leu Leu Pro Gly Thr Gln Ala Asp Pro Val
 20 25 30

 Asp Val Leu Lys Ala Leu Gly Val Gln Gly Gly Gln Ala Gly Val Pro
 35 40 45

 Glu Gly Pro Gly Phe Cys Pro Gln Arg Thr Pro Glu Gly Asp Arg Ala
 50 55 60

 Phe Arg Ile Gly Gln Ala Ser Thr Leu Gly Ile Pro Thr Trp Glu Leu
 65 70 75 80

 Phe Pro Glu Gly His Phe Pro Glu Asn Phe Ser Leu Leu Ile Thr Leu
 85 90 95

 Arg Gly Gln Pro Ala Asn Gln Ser Val Leu Leu Ser Ile Tyr Asp Glu
 100 105 110

 Arg Gly Ala Arg Gln Leu Gly Leu Ala Leu Gly Pro Ala Leu Gly Leu
 115 120 125

 Leu Gly Asp Pro Phe Arg Pro Leu Pro Gln Gln Val Asn Leu Thr Asp
 130 135 140

 Gly Arg Trp His Arg Val Ala Val Ser Ile Asp Gly Glu Met Val Thr
 145 150 155 160

 Leu Val Ala Asp Cys Glu Ala Gln Pro Pro Val Leu Gly His Gly Pro
 165 170 175

 Arg Phe Ile Ser Ile Ala Gly Leu Thr Val Leu Gly Thr Gln Asp Leu
 180 185 190

 Gly Glu Lys Thr Phe Glu Gly Asp Ile Gln Glu Leu Leu Ile Ser Pro
 195 200 205

 Asp Pro Gln Ala Ala Phe Gln Ala Cys Glu Arg Tyr Leu Pro Asp Cys
 210 215 220

 Asp Asn Leu Ala Pro Ala Ala Thr Val Ala Pro Gln Gly Glu Pro Glu
 225 230 235 240

 Thr Pro Arg Pro Arg Arg Lys Gly Lys Gly Lys Gly Arg Lys Lys Gly
 245 250 255

 Arg Gly Arg Lys Gly Lys Gly Arg Lys Lys Asn Lys Glu Ile Trp Thr
 260 265 270

 Ser Ser Pro Pro Pro Asp Ser Ala Glu Asn Gln Thr Ser Thr Asp Ile
 275 280 285

 Pro Lys Thr Glu Thr Pro Ala Pro Asn Leu Pro Pro Thr Pro Thr Pro
 290 295 300

 Leu Val Val Thr Ser Thr Val Thr Thr Gly Leu Asn Ala Thr Ile Leu
 305 310 315 320

 Glu Arg Ser Leu Asp Pro Asp Ser Gly Thr Glu Leu Gly Thr Leu Glu
 325 330 335

 Thr Lys Ala Ala Arg Glu Asp Glu Glu Gly Asp Asp Ser Thr Met Gly
 340 345 350

 Pro Asp Phe Arg Ala Ala Glu Tyr Pro Ser Arg Thr Gln Phe Gln Ile
 355 360 365

 Phe Pro Gly Ala Gly Glu Lys Gly Ala Lys Gly Glu Pro Ala Val Ile
 370 375 380

-continued

Glu Lys Gly Gln Gln Phe Glu Gly Pro Pro Gly Ala Pro Gly Pro Gln
 385 390 395 400
 Gly Val Val Gly Pro Ser Gly Pro Pro Gly Pro Pro Gly Phe Pro Gly
 405 410 415
 Asp Pro Gly Pro Pro Gly Pro Ala Gly Leu Pro Gly Ile Pro Gly Ile
 420 425 430
 Asp Gly Ile Arg Gly Pro Pro Gly Thr Val Ile Met Met Pro Phe Gln
 435 440 445
 Phe Ala Gly Gly Ser Phe Lys Gly Pro Pro Val Ser Phe Gln Gln Ala
 450 455 460
 Gln Ala Gln Ala Val Leu Gln Gln Thr Gln Leu Ser Met Lys Gly Pro
 465 470 475 480
 Pro Gly Pro Val Gly Leu Thr Gly Arg Pro Gly Pro Val Gly Leu Pro
 485 490 495
 Gly His Pro Gly Leu Lys Gly Glu Glu Gly Ala Glu Gly Pro Gln Gly
 500 505 510
 Pro Arg Gly Leu Gln Gly Pro His Gly Pro Pro Gly Arg Val Gly Lys
 515 520 525
 Met Gly Arg Pro Gly Ala Asp Gly Ala Arg Gly Leu Pro Gly Asp Thr
 530 535 540
 Gly Pro Lys Gly Asp Arg Gly Phe Asp Gly Leu Pro Gly Leu Pro Gly
 545 550 555 560
 Glu Lys Gly Gln Arg Gly Asp Phe Gly His Val Gly Gln Pro Gly Pro
 565 570 575
 Pro Gly Glu Asp Gly Glu Arg Gly Ala Glu Gly Pro Pro Gly Pro Thr
 580 585 590
 Gly Gln Ala Gly Glu Pro Gly Pro Arg Gly Leu Leu Gly Pro Arg Gly
 595 600 605
 Ser Pro Gly Pro Thr Gly Arg Pro Gly Val Thr Gly Ile Asp Gly Ala
 610 615 620
 Pro Gly Ala Lys Gly Asn Val Gly Pro Pro Gly Glu Pro Gly Pro Pro
 625 630 635 640
 Gly Gln Gln Gly Asn His Gly Ser Gln Gly Leu Pro Gly Pro Gln Gly
 645 650 655
 Leu Ile Gly Thr Pro Gly Glu Lys Gly Pro Pro Gly Asn Pro Gly Ile
 660 665 670
 Pro Gly Leu Pro Gly Ser Asp Gly Pro Leu Gly His Pro Gly His Glu
 675 680 685
 Gly Pro Thr Gly Glu Lys Gly Ala Gln Gly Pro Pro Gly Ser Ala Gly
 690 695 700
 Pro Pro Gly Tyr Pro Gly Pro Arg Gly Val Lys Gly Thr Ser Gly Asn
 705 710 715 720
 Arg Gly Leu Gln Gly Glu Lys Gly Glu Lys Gly Glu Asp Gly Phe Pro
 725 730 735
 Gly Phe Lys Gly Asp Val Gly Leu Lys Gly Asp Gln Gly Lys Pro Gly
 740 745 750
 Ala Pro Gly Pro Arg Gly Glu Asp Gly Pro Glu Gly Pro Lys Gly Gln
 755 760 765
 Ala Gly Gln Ala Gly Glu Glu Gly Pro Pro Gly Ser Ala Gly Glu Lys
 770 775 780
 Gly Lys Leu Gly Val Pro Gly Leu Pro Gly Tyr Pro Gly Arg Pro Gly
 785 790 795 800
 Pro Lys Gly Ser Ile Gly Phe Pro Gly Pro Leu Gly Pro Ile Gly Glu

-continued

805				810				815							
Lys	Gly	Lys	Ser	Gly	Lys	Thr	Gly	Gln	Pro	Gly	Leu	Glu	Gly	Glu	Arg
			820					825					830		
Gly	Pro	Pro	Gly	Ser	Arg	Gly	Glu	Arg	Gly	Gln	Pro	Gly	Ala	Thr	Gly
		835					840					845			
Gln	Pro	Gly	Pro	Lys	Gly	Asp	Val	Gly	Gln	Asp	Gly	Ala	Pro	Gly	Ile
850						855					860				
Pro	Gly	Glu	Lys	Gly	Leu	Pro	Gly	Leu	Gln	Gly	Pro	Pro	Gly	Phe	Pro
865					870				875						880
Gly	Pro	Lys	Gly	Pro	Pro	Gly	His	Gln	Gly	Lys	Asp	Gly	Arg	Pro	Gly
			885					890					895		
His	Pro	Gly	Gln	Arg	Gly	Glu	Leu	Gly	Phe	Gln	Gly	Gln	Thr	Gly	Pro
			900					905					910		
Pro	Gly	Pro	Ala	Gly	Val	Leu	Gly	Pro	Gln	Gly	Lys	Thr	Gly	Glu	Val
		915					920					925			
Gly	Pro	Leu	Gly	Glu	Arg	Gly	Pro	Pro	Gly	Pro	Pro	Gly	Pro	Pro	Gly
	930					935					940				
Glu	Gln	Gly	Leu	Pro	Gly	Leu	Glu	Gly	Arg	Glu	Gly	Ala	Lys	Gly	Glu
945					950					955					960
Leu	Gly	Pro	Pro	Gly	Pro	Leu	Gly	Lys	Glu	Gly	Pro	Ala	Gly	Leu	Arg
				965					970						975
Gly	Phe	Pro	Gly	Pro	Lys	Gly	Gly	Pro	Gly	Asp	Pro	Gly	Pro	Thr	Gly
		980						985					990		
Leu	Lys	Gly	Asp	Lys	Gly	Pro	Pro	Gly	Pro	Val	Gly	Ala	Asn	Gly	Ser
		995					1000					1005			
Pro	Gly	Glu	Arg	Gly	Pro	Leu	Gly	Pro	Ala	Gly	Gly	Ile	Gly	Leu	Pro
	1010					1015						1020			
Gly	Gln	Ser	Gly	Ser	Glu	Gly	Pro	Val	Gly	Pro	Ala	Gly	Lys	Lys	Gly
1025					1030					1035					1040
Ser	Arg	Gly	Glu	Arg	Gly	Pro	Pro	Gly	Pro	Thr	Gly	Lys	Asp	Gly	Ile
				1045					1050					1055	
Pro	Gly	Pro	Leu	Gly	Pro	Leu	Gly	Pro	Pro	Gly	Ala	Ala	Gly	Pro	Ser
			1060					1065					1070		
Gly	Glu	Glu	Gly	Asp	Lys	Gly	Asp	Val	Gly	Ala	Pro	Gly	His	Lys	Gly
	1075						1080					1085			
Ser	Lys	Gly	Asp	Lys	Gly	Asp	Ala	Gly	Pro	Pro	Gly	Gln	Pro	Gly	Ile
	1090					1095					1100				
Arg	Gly	Pro	Ala	Gly	His	Pro	Gly	Pro	Pro	Gly	Ala	Asp	Gly	Ala	Gln
1105					1110					1115					1120
Gly	Arg	Arg	Gly	Pro	Pro	Gly	Leu	Phe	Gly	Gln	Lys	Gly	Asp	Asp	Gly
				1125					1130					1135	
Val	Arg	Gly	Phe	Val	Gly	Val	Ile	Gly	Pro	Pro	Gly	Leu	Gln	Gly	Leu
			1140					1145					1150		
Pro	Gly	Pro	Pro	Gly	Glu	Lys	Gly	Glu	Val	Gly	Asp	Val	Gly	Ser	Met
		1155					1160					1165			
Gly	Pro	His	Gly	Ala	Pro	Gly	Pro	Arg	Gly	Pro	Gln	Gly	Pro	Thr	Gly
						1175					1180				
Ser	Glu	Gly	Thr	Pro	Gly	Leu	Pro	Gly	Gly	Val	Gly	Gln	Pro	Gly	Ala
1185					1190					1195					1200
Val	Gly	Glu	Lys	Gly	Glu	Arg	Gly	Asp	Ala	Gly	Asp	Pro	Gly	Pro	Pro
				1205					1210					1215	
Gly	Ala	Pro	Gly	Ile	Pro	Gly	Pro	Lys	Gly	Asp	Ile	Gly	Glu	Lys	Gly
			1220					1225					1230		

-continued

Asp Ser Gly Pro Ser Gly Ala Ala Gly Pro Pro Gly Lys Lys Gly Pro
 1235 1240 1245

Pro Gly Glu Asp Gly Ala Lys Gly Ser Val Gly Pro Thr Gly Leu Pro
 1250 1255 1260

Gly Asp Leu Gly Pro Pro Gly Asp Pro Gly Val Ser Gly Ile Asp Gly
 1265 1270 1275 1280

Ser Pro Gly Glu Lys Gly Asp Pro Gly Asp Val Gly Gly Pro Gly Pro
 1285 1290 1295

Pro Gly Ala Ser Gly Glu Pro Gly Ala Pro Gly Pro Pro Gly Lys Arg
 1300 1305 1310

Gly Pro Ser Gly His Met Gly Arg Glu Gly Arg Glu Gly Glu Lys Gly
 1315 1320 1325

Ala Lys Gly Glu Pro Gly Pro Asp Gly Pro Pro Gly Arg Thr Gly Pro
 1330 1335 1340

Met Gly Ala Arg Gly Pro Pro Gly Arg Val Gly Pro Glu Gly Leu Arg
 1345 1350 1355 1360

Gly Ile Pro Gly Pro Val Gly Glu Pro Gly Leu Leu Gly Ala Pro Gly
 1365 1370 1375

Gln Met Gly Pro Pro Gly Pro Leu Gly Pro Ser Gly Leu Pro Gly Leu
 1380 1385 1390

Lys Gly Asp Thr Gly Pro Lys Gly Glu Lys Gly His Ile Gly Leu Ile
 1395 1400 1405

Gly Leu Ile Gly Pro Pro Gly Glu Ala Gly Glu Lys Gly Asp Gln Gly
 1410 1415 1420

Leu Pro Gly Val Gln Gly Pro Pro Gly Pro Lys Gly Asp Pro Gly Pro
 1425 1430 1435 1440

Pro Gly Pro Ile Gly Ser Leu Gly His Pro Gly Pro Pro Gly Val Ala
 1445 1450 1455

Gly Pro Leu Gly Gln Lys Gly Ser Lys Gly Ser Pro Gly Ser Met Gly
 1460 1465 1470

Pro Arg Gly Asp Thr Gly Pro Ala Gly Pro Pro Gly Pro Pro Gly Ala
 1475 1480 1485

Pro Ala Glu Leu His Gly Leu Arg Arg Arg Arg Arg Phe Val Pro Val
 1490 1495 1500

Pro Leu Pro Val Val Glu Gly Gly Leu Glu Glu Val Leu Ala Ser Leu
 1505 1510 1515 1520

Thr Ser Leu Ser Leu Glu Leu Glu Gln Leu Arg Arg Pro Pro Gly Thr
 1525 1530 1535

Ala Glu Arg Pro Gly Leu Val Cys His Glu Leu His Arg Asn His Pro
 1540 1545 1550

His Leu Pro Asp Gly Glu Tyr Trp Ile Asp Pro Asn Gln Gly Cys Ala
 1555 1560 1565

Arg Asp Ser Phe Arg Val Phe Cys Asn Phe Thr Ala Gly Gly Glu Thr
 1570 1575 1580

Cys Leu Tyr Pro Asp Lys Lys Phe Glu Ile Val Lys Leu Ala Ser Trp
 1585 1590 1595 1600

Ser Lys Glu Lys Pro Gly Gly Trp Tyr Ser Thr Phe Arg Arg Gly Lys
 1605 1610 1615

Lys Phe Ser Tyr Val Asp Ala Asp Gly Ser Pro Val Asn Val Val Gln
 1620 1625 1630

Leu Asn Phe Leu Lys Leu Leu Ser Ala Thr Ala Arg Gln Asn Phe Thr
 1635 1640 1645

Tyr Ser Cys Gln Asn Ala Ala Ala Trp Leu Asp Glu Ala Thr Gly Asp
 1650 1655 1660

