

US008728481B2

(12) **United States Patent**
Banchereau et al.

(10) **Patent No.:** **US 8,728,481 B2**
(45) **Date of Patent:** **May 20, 2014**

(54) **AGENTS THAT ENGAGE
ANTIGEN-PRESENTING CELLS THROUGH
DENDRITIC CELL ASIALOGLYCOPROTEIN
RECEPTOR (DC-ASGPR)**

(75) Inventors: **Jacques F. Banchereau**, Montclair, NJ
(US); **SangKOn Oh**, Baltimore, MD
(US); **Gerard Zurawski**, Midlothian,
TX (US); **Sandra Zurawski**,
Midlothian, TX (US); **Dapeng Li**,
Dallas, TX (US)

(73) Assignee: **Baylor Research Institute**, Dallas, TX
(US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **13/551,198**

(22) Filed: **Jul. 17, 2012**

(65) **Prior Publication Data**

US 2012/0282281 A1 Nov. 8, 2012

Related U.S. Application Data

(62) Division of application No. 12/025,010, filed on Feb.
2, 2008, now Pat. No. 8,236,934.

(60) Provisional application No. 60/888,036, filed on Feb.
2, 2007.

(51) **Int. Cl.**
A61K 39/00 (2006.01)
A61K 39/395 (2006.01)
A01N 63/00 (2006.01)

(52) **U.S. Cl.**
USPC **424/178.1**; 424/173.1; 424/93.71

(58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,791,932 A 2/1974 Schuurs 435/7.8
3,949,064 A 4/1976 Bornstein 436/527
4,174,384 A 11/1979 Ullman 436/537
4,554,101 A 11/1985 Hopp 530/324
5,945,308 A 8/1999 Tang 435/69.1
6,046,158 A 4/2000 Ariizumi 514/21.2
6,277,959 B1 8/2001 Valladeau 530/350
6,410,241 B1 6/2002 Sykes 435/6.13
6,451,995 B1 9/2002 Cheung 536/23.53
6,541,011 B2 4/2003 Punnonen 424/204.1
6,738,985 B2 5/2004 Hahn 2/181
7,129,039 B2 10/2006 Ariizumi 435/4
7,179,595 B2 2/2007 Li 435/6.16
7,592,003 B2 9/2009 Nagai 424/93.7
7,666,596 B2 2/2010 Halloran 435/6.16
7,786,267 B2 8/2010 Flamar 530/387.1
8,236,934 B2 8/2012 Banchereau et al.
8,449,888 B2 5/2013 Zurawski et al.
2004/0126357 A1 7/2004 Segal 424/85.1
2004/0192892 A1* 9/2004 Valladeau et al. 530/350

2004/0258688 A1 12/2004 Hawiger et al.
2005/0037001 A1 2/2005 Germeraad 424/141.1
2005/0064509 A1 3/2005 Bradbury 435/7.1
2005/0106700 A1 5/2005 Kanaya 435/226
2006/0257412 A1 11/2006 Bowdish 424/178.1
2009/0186025 A1 7/2009 Colaco 424/134.1
2012/0039916 A1 2/2012 Zurawski et al.
2012/0121592 A1 5/2012 Oh et al.
2012/0231023 A1 9/2012 Zurawski et al.
2012/0237513 A1 9/2012 Zurawski et al.
2012/0244155 A1 9/2012 Lecine et al.
2012/0282281 A1 11/2012 Banchereau et al.
2012/0301465 A1 11/2012 Dutartre et al.
2012/0315269 A1 12/2012 Klechevsky et al.

FOREIGN PATENT DOCUMENTS

EP 1441030 A1 7/2004
WO 97/14789 A2 4/1997
WO 00/63151 A1 10/2000
WO WO 00/63251 10/2000
WO 03/073827 A2 9/2003
WO WO 2006/004663 1/2006
WO WO 2006004663 A2 * 1/2006
WO 2006/107617 A2 10/2006
WO 2008/097817 A2 8/2008

OTHER PUBLICATIONS

Karni et al., 2006. J. Immunol. vol. 177: 4196-4202.*
Craig et al., 2006, J. Biotech. vol. 121: 165-173.*
Wiley, K.N., et al., "The In-Vitro Inhibition of Rat Alloantigen Pre-
sentation by Immunotoxins—Implications for Allografting," Clin.
Exp. Immunol., (1989), 76:132-137.
Wykes, M. N., et al., "Dendritic cell-B-cell interaction: dendritic cells
provide B cells with CD40-independent proliferation signals and
CD40-dependent survival signals," Immunology (2000), 100:1-3.
Wykes, M. N., et al., "Dendritic cells and follicular dendritic cells
express a novel ligand for CD38 which influences their maturation
and antibody responses," Immunology (2004), 113:318-327.
Zhu, Zhenping, et al., "Inhibition of Tumor Growth and Metastasis by
Targeting Tumor-Associated Angiogenesis with Antagonists to the
Receptors of Vascular Endothelial Growth Factor," Investigational
New Drugs, 1999, 17:195-212.
Ahmad-Nejad, et al., "Bacterial CpG-DNA and Lipopolysaccharides
Activate Toll-Like Receptors at Distinct Cellular Compartments,"
European J. Immunol., (2002), 32:1958-1968.
Arce, Ignacio et al., "The human C-type lectin CLECSF18 is a novel
monocyte/macrophage endocytic receptor", European Journal of
Immunology, 2004, vol. 34(1), pp. 210-220.
Asea, A., et al., "HSP70 Stimulates Cytokine Production Through a
CD14-Dependant Pathway, Demonstrating its Dual Role as a Chap-
erone and Cytokine," Nat Med, (2000), 6:435-442.
Balazs, F., et al., "Blood Dendritic Cells Interact with Splenic Mar-
ginal Zone B Cells to Initiate T-Independent Immune Responses,"
Immunity, (2002), 17:341-352.

(Continued)

Primary Examiner — Amy Juedes

(74) *Attorney, Agent, or Firm* — Fulbright & Jaworski LLP

(57) **ABSTRACT**

The present invention includes compositions and methods for
making and using anti DC-ASGPR antibodies that can, e.g.,
activate DCs and other cells.

9 Claims, 12 Drawing Sheets

(56)

References Cited

OTHER PUBLICATIONS

- Banchereau, J., et al., "Dendritic Cells and the Control of Immunity," *Nature*, (1998), 392:245-252.
- Banchereau, J., et al., "Immunobiology of Dendritic Cells," *Annu Rev Immunol*, (2000), 18:767-811.
- Banchereau, J., et al., "Dendritic Cells as Vectors for Thereapy," *Cell*, (2001), 106(3):271-4.
- Banchereau J., et al., "Immune and Clinical Responses in Patients with Metastatic Melanoma to CD34(+) Progenitor-Derived Dendritic Cell Vaccine," *Cancer Res*, (2001), 61(17):6451-8.
- Barak, Y. et al., "Matching Fusion Protein Systems for Affinity Analysis of Two Interacting Families of Proteins: the Cohesin-Dockerin Interaction", *J. Mo. Recognt*, 2005:18:491-501.
- Barrios, et al., "Length of the Antibody Heavy Chain Complementarity Determining Region 3 as a Specificity-Determining Factor", *J. Mol. Recognt*. 2004: 17:332-338.
- Bates, E. E., et al., "APCs Express DCIR, a Novel C-Type Lectin Surface Receptor Containing an Immunoreceptor Tyrosine-Based Inhibitory Motif," *J Immunol*, (1999), 163(4):1973-83.
- Bausinger, H., et al., "Endotoxin-Free Heat-Shock Protein 70 Fails to Induce APC Activation," *Eur J Immunol* (2002), 32:3708-3713.
- Bendsten, J.D., et al., "Improved Prediction of Signal Peptides: SignalP 3.0." *J Mol Biol*, (2004), 340(4):783-95.
- Berard, F. et al., "Cross-Priming of Naïve CD8 T Cells Against Melanoma Antigens Using Dendritic Cells Loaded with Killed Allogeneic Melanoma Cells," *J Exp Med*, (Dec. 2000), 192(11):1535-44.
- Bergtold, A., et al., "Cell Surface Recycling of Internalized Antigen Permits Dendritic Cell Priming of B Cells," *Immunity* (2005), 23:503-514.
- Bernasconi, N. L., et al., "Maintenance of Serological Memory by Polyclonal Activation of Human Memory B Cells," *Science* (2002), 298:2199-2202.
- Beutler, B., et al., "Genetic Analysis of Host Resistance: Toll-Like Receptor Signaling and Immunity at Large," *Annu Rev Immunol* (2006), 24:353-389.
- Bonifaz, Laura, et al., "Efficient Targeting of Protein Antigen to the Dendritic Cell Receptor DEC-205 in the Steady State Leads to Antigen Presentation on Major Histocompatibility Complex Class I Products and Peripheral CD8 T Cell Tolerance," *J. Exp. Med.*, Dec. 16, 2002, vol. 196, No. 12, pp. 1627-1638.
- Bonifaz, L.C., et al., "In Vivo Targeting of Antigens to Maturing Dendritic Cells via the DEC-205 Receptor Improves T Cell Vaccination." *J Exp Med*, (Mar. 2004), 199(6):815-24.
- Brown, G. D., "Dectin-1: a signalling non-TLR pattern-recognition receptor," *Nat Rev Immunol* (2006), 6:33-43.
- Cambi, A., et al., "The C-type lectin DC-SIGN (CD209) is an antigenuptake receptor for *Candida albicans* on dendritic cells," *Eur J Immunol* (2003), 33:532-538.
- Carter, et al., "Preferential Induction of CD4+ T Cell Responses through In Vivo Targeting of Antigen to Dendritic Cell-Associated C-Type Lectin-1", *J. of Immunol.*, vol. 177, No. 4, 2006, pp. 2276-2284.
- Carvalho, A.L., et al., "Cellulosome Assembly Revealed by the Crystal Structure of the Cohesin-Dockerin Complex." *Proc Natl Acad Sci USA*, (2003), 100(24):13809-14.
- Chu. et al., "CpG Oligodeoxynucleotides Down-Regulate Macrophage Class II MHC Antigen Processing", *J Immunol* 1999:163:1188-1194.
- Colonna, M., et al., "Molecular Characterization of Two Novel C-Type Lectin-Like Receptors, one of which is Selectively Expressed in Human Dendritic Cells," *Eur J Immunol* (2000), 30:697-704.
- Cooper, A. M., et al., "Mice Lacking Bioactive IL-12 Can Generate Protective, Antigen-Specific Cellular Responses to Mycobacterial Infection Only if the IL-12 p40 Subunit Is Present," *J Immunol* (2002), 168:1322-1327.
- Craxton, A., et al., "Macrophage and Dendritic Cell-Dependent Regulation of Human B-Cell Proliferation Requires the TNF Family Ligand BAFF," *Blood*, (2003), 101:4464-4471.
- Deinste, Y., G. et al., "Involvement of LOX-1 in Dendritic Cell-Mediated Antigen Cross-Presentation." *Immunity* (2002), 17(3):353-62.
- Deyev, S.M. et al., "Design of multivalent complexes using the barnase-barstar module", *Nature Biotechnology*, vol. 21, No. 12, Dec. 2003, pp. 1486-1492.
- Ding, et al., "A Scaffoldin of the *Bacteriodes cellulosolvens* Cellulosome That Contains 11 Type II Cohesins", *J. Bacteriology* 182(17):4915-4925, Sep. 2000.
- Ding, et al., "The Bacterial Scaffoldin: Structure, Function and Potential Applications in the Nanosciences", *Genetic Engineering*, vol. 25:209-225, (2003).
- D'Ostiani, C. F., et al., "Dendritic Cells Discriminate between Yeasts and Hyphae of the Fungus *Candida albicans*: Implications for Initiation of T Helper Cell Immunity In Vitro and In Vivo," *J Exp Med* (2000), 191:1661-1673.
- Dubois, B., et al., "Dendritic cells directly modulate B cell growth and differentiation," *J Leukoc Biol* (1999), 66:224-230.
- Extended European Search Report for Application No. 08714180.0, dated Sep. 29, 2011, 14 pages.
- Extended European Search Report for Application No. 08728767.8, dated Aug. 24, 2010, 11 pages.
- Extended European Search Report for Application No. 08728868.4, dated Aug. 3, 2010, 10 pages.
- Extended European Search Report for Application No. 08799678.1, dated Feb. 5, 2010, 12 pages.
- Fierobe, H-P et al., "Design and Production of Active Cellulosome Chimeras," *J Biol Chem* (2001), 276:21257-21261.
- Figdor, C. G., et al., "C-Type Lectin Receptors on Dendritic Cells and Langerhans Cells," *Nat Rev Immunol*, (2002), 2(2): 77-84.
- Flamar, A.L. et al., "Antibody and Antigen Complexes Based on Cohesin and Dockerin Interaction are Versatile Tools for Eliciting and Monitoring Specific Immune Responses", *Clinical Immunology*, US LnkD-DOI:10.1016/J. Clin.2009.03.019, vol. 131, Jan. 1, 2009, pp. S9.
- Flomes, L.M. et al., "Identification of Lectin-Like Receptors Expressed by Antigen Presenting Cells and Neutrophils and their Mapping to a Novel Gene Complex," *Immunogenetics*, (2004), vol. 56, pp. 506-517.
- Fradin, C., et al., "β-1,2-Linked Oligomannosides from *Candida albicans* Bind to a 32-Kilodalton Macrophage Membrane Protein Homologous to the Mammalian Lectin Galectin-3," *Infect Immun* (2000), 68:4391-4398.
- Frankel, A.F., "Increased Sophistication of Immunotoxins," *Clin Can Res*, (2002), 8:942-944.
- Gantner, B. N., et al., "Collaborative Induction of Inflammatory Responses by Dectin-1 and Toll-like Receptor 2," *J Exp Med* (2003), 197:1107-1117.
- Geijtenbeek, T. B., et al., "DC-SIGN—ICAM-2 Interaction Mediates Dendritic Cell Trafficking," *Nat Immunol* (2000), 1:353-357.
- Geijtenbeek, T. B., et al., "Identification of DC-SIGN, a Novel Dendritic Cell—Specific ICAM-3 Receptor that Supports Primary Immune Responses," *Cell* (2000), 100:575-585.
- Geijtenbeek, T. B., et al., "Mycobacteria Target DC-SIGN to Suppress Dendritic Cell Function," *J Exp Med* (2003), 197:7-17.
- Geijtenbeek, T.B., et al., "Self-and Nonself-Recognition by C-type Lectins on Dendritic Cells." *Annu Rev Immunol* (2004), 22:33-54.
- Gross, J. A., et al., "TACI and BCMA are Receptors for a TNF Homologue Implicated in B-cell Autoimmune Disease," *Nature* (2000), 404:995-999.
- Grueneback, et al., "Human (h) Dectin-1 is a Member of C-Type-Lectin-Like Receptor Family that was Shown to be the Major Receptor for Fungal Beta-Glucans . . ." Abstract from 49th Annual Meeting of the American Society of Hematology, Dec. 2007, XP002563205, 2 pages.
- Hayashida, K., et al., "Lectin-Like Oxidized LDL Receptor-1 (LOX-1) Supports Adhesion of Mononuclear Leukocytes and a Monocyte-Like Cell Line THP-1 Cells Under Static and Flow Conditions," *FEBS Letters*, (2002), 511:133-138.
- Huang, et al., "Cloning and Characterization of a Novel ITIM Containing Lectin-like Immunoreceptor LLIR and Its Two Transmembrane Region Deletion Variants", *Biochem. Biophys. Res. Commun.*, 281(1):131-140, (2001).

(56)

References Cited

OTHER PUBLICATIONS

- International Search Report and Written Opinion for PCT/US2008/052714, dated Jul. 8, 2008, 8 pages.
- International Search Report and Written Opinion for PCT/US2008/52850, dated Sep. 24, 2008, 9 pages.
- International search Report and Written Opinion for PCT/US08/52865, dated Aug. 13, 2008, 8 pages.
- International Search Report and Written Opinion for PCT/US2008/054785, dated Sep. 25, 2008, 12 pages.
- International Search Report and Written Opinion for PCT/US2008/054792, dated Aug. 13, 2008, 10 pages.
- International Search Report and Written Opinion for PCT/US2008/054798, dated Dec. 8, 2008, 13 pages.
- Jeannin, P., et al., "Complexity and Complementarity of Outer Membrane Protein A Recognition by Cellular and Humoral Innate Immunity Receptors," *Immunity* (2005), 22:551-560.
- Jego, G., et al., "Plasmacytoid Dendritic Cells Induce Plasma Cell Differentiation through Type I Interferon and Interleukin 6," *Immunity* (2003), 19:225-234.
- Jindou, S. et al., "Interaction between a Type-II Dockerin Domain and a Type-II Cohesin Domain from *Clostridium thermocellum* Cellulosome", *Biosci. Biotechnol. Biochem.* 68(4):924-926,(2004).
- Kakutani, M., et al., "A platelet—endothelium interaction mediated by lectin-like oxidized low-density lipoprotein receptor-1," *PNAS* (2000), 97:360-364.
- Karpol, A. et al., "Engineering a Reversible, High-Affinity System for Efficient Protein Purification Based on the Cohesin-Dockerin Interaction", *J. Mol. Recognit.* 2009, vol. 22:91-98.
- Kikuchi, T., et al., "Dendritic Cells Genetically Modified to Express CD40 Ligand and Pulsed with Antigen can Initiate Antigen-Specific Humoral Immunity Independent of CD4+ T Cells," *Nat Med.* (2000), 6:1154-1159.
- Klimka et al., "Human Anti-CD30 Recombinant Antibodies by Guided Phage Antibody Selection Using Cell Planning", *Brit. J. Cancer*, vol. 83, 252-260, 2000.
- Kobrin, et al., "A V Region Mutation in a Phosphocholine-Binding Monoclonal Antibody Results in Loss of Antigen Binding", *J. Immunology*, 146:2017-2020 (1991).
- Latz, et al., "TLR9 Signals After Translocating from the ER to CpG DNA in the Lysosome", *Nat. Immunol.*, (2004), vol. 5:190-198.
- Lee, S. J., et al., "Mannose Receptor-Mediated Regulation of Serum Glycoprotein Homeostasis," *Science* (2002), 295:1898-1901.
- Maclennan, C., et al., "Dendritic Cells, BAFF, and APRIL: Innate Players in Adaptive Antibody Responses," *Immunity* (2002), 17:235-238.
- Maeda, N., et al., "The Cell Surface Receptor DC-SIGN Discriminates between *Mycobacterium* Species through Selective Recognition of the Mannose Caps on Lipoarabinomannan," *J Biol Chem*, (2003), 278:5513-5513.
- Marks et al., "By-Passing Immunization: Human Antibodies from V-gene Libraries Displayed on Phage", *J. Mol. Biol.*, (1991), 222:581-597.
- Mechaly, A. et al., "Cohesin-Dockerin Interaction in Cellulosome Assembly", *J. Biological Chemistry*, Mar. 30, 2001, vol. 276, No. 13, pp. 9883-9888.
- Mellman, I. et al., "Dendritic Cells: Specialized and Regulated Antigen Processing Machines," *Cell*, (2001), 106 (3):255-8.
- Moore, P. A., et al., "BLyS: Member of the Tumor Necrosis Factor Family and B Lymphocyte Stimulator," *Science* (1999), 285:260-263.
- Moriwaki, H., et al., "Expression of Lectin-Like Oxidized Low Density Lipoprotein Receptor-1 in Human and Murine Macrophages: Upregulated Expression by TNF- α ," *FEBS Letters*, (1998), 440:29-32.
- Neidhardt-Berard, E.M., et al., "Dendritic Cells Loaded with Killed Breast Cancer Cells Induce Differentiation of Tumor-Specific Cytotoxic T Lymphocytes." *Breast Cancer Res*, (2004), 6(4):R322-8.
- Netea, M. G., et al., "CD40/CD40 Ligand Interactions in the Host Defense Against Disseminated *Candida albicans* Infection: the Role of Macrophage-Derived Nitric Oxide," *Eur J Immunol*, (2002), 32:1455-1463.
- Palucka, A.K., et al., "Human Dendritic Cell Subsets in NOD/SCID Mice Engrafted with CD34+ Hematopoietic progenitors," *Blood*, (2003), 102(9):3302-10.
- Proudfoot et al., "Receptor-Mediated Delivery of Antigens to Dendritic Cells: Anticancer Applications", *Molec. Pharm.*, vol. 4, 2007, pp. 58-72.
- Pyz, E., et al., "C-Type Lectin-Like Receptors on Myeloid Cells," *Ann Med*, (2006), 38:242-251.
- Qi, H., et al., "Extrafollicular Activation of Lymph Node B Cells by Antigen-Bearing Dendritic Cells," *Science* (2006), 312:1672-1676.
- Ramakrishna, V., et al., "Mannose Receptor Targeting of Tumor Antigen PMEL17 to Human Dendritic Cells Directs Antimelanoma T Cell Responses via Multiple HLA Molecules." *J Immunol*, (2004), 172(5):2845-52.
- Ramakrishna, et al., "Toll-like Receptor Activation Enhances Cell-Mediated Immunity Inducted by an Antibody Vaccine Targeting Human Dendritic Cells", *Journal of Translational Medicine*, vol. 5:5, pp. 1-14, Jan. 2007.
- Reddy, M.P., et al., "Elimination of FC Receptor-Dependent Effector Functions of a Modified IgG4 Monoclonal Antibody to Human CD4," *J Immunol*, (2000), 164(4):1925-33.
- Rossi, E.A. et al., "Development of New Multivalent-bispecific Agents for Pretargeting Tumor Localization and Therapy", *Clinical Cancer Research*, vol. 9, Sep. 1, 2003, pp. 3886S-3896S.
- Ruprecht, C. R., et al., "Toll-like Receptor Stimulation as a Third Signal Required for Activation of Human Naive B Cells," *Eur J Immunol*, (2006), 36:810-816.
- Saeland, S., "Langerin and the DC Asialoglycoprotein Receptor: Two Closely Related Endocytic Type-II Lectins with Divergent Functions in Dendritic Cells," *Journal of Investigative Dermatology*, vol. 117, No. 4, Oct. 2001, p. 1003.
- Schaft, Neils, et al., "Dendritic Cell Vaccination and Other Strategies to TLP the Balance of the Immune System," *Cancer Immunol. Immunother.* (2008), 57:913-928.
- Shortman, Ken, et al., "Mouse and Human Dendritic Cell Subtypes," *Nature Reviews, Immunology*, vol. 2, Mar. 2002, pp. 454-465.
- Smith, Daniel C., et al., "Lack of Dendritic Cell Maturation by the Plant Toxin Ricin," *European J. Immunol.*, (2004), 34:2149-2157.
- Spooner, RA, et al., "Expression of Immunoglobulin Heavy Chain-Ricin a Chain Fusions in Mammalian Cells," *Mol. Immunol.*, Feb. 1994, 31(2):117-125.
- Starovasnik, Melissa A., et al., "Structural Mimicry of a Native Protein by a Minimized Binding Domain," *Proc. Nat. Acad. Sci. USA*, Sep. 1997, vol. 94, pp. 10080-10085.
- Steinman, Ralph M., et al., "Active Immunization Against Cancer with Dendritic Cells: The Near Future," *Int. J. Cancer*, (2001), 94, 459-473.
- Tacke, Paul J., et al., "Effective Induction of Naive and Recall T-Cell Responses by Targeting Antigen to Human Dendritic Cells via a Humanized Anti-DC-SIGN Antibody," *Blood*, May 5, 2005, 106:1278-1285.
- Tailleux, L., et al., "DC-SIGN Is the Major *Mycobacterium tuberculosis* Receptor on Human Dendritic Cells," *J Exp Med* (2003), 197:121-127.
- Tan, et al., "Mannose Receptor-Mediated Uptake of Antigens Strongly Enhances HLA Class II-Restricted Antigen Presentation by Cultured Dendritic Cells," *Eur. J. Immunol.*, 1997, 27:2426-2435.
- Trombetta, E. Sergio, et al., "Cell Biology of Antigen Processing in Vitro and In Vivo," *Annu. Rev. Immunol.*, 2005, 23:975-1028.
- Trumpfheller, Christine, et al., "Intensified and Protective CD4+ T Cell Immunity in Mice with Anti-Dendritic Cell HIV Gag Fusion Antibody Vaccine," *Mar. 20, 2006*, vol. 203, No. 3, pp. 607-617.
- Valladeau, J., et al., "Immature Human Dendritic Cells Express Asialoglycoprotein Receptor Isoforms for Efficient Receptor-Mediated Endocytosis," *J Immunol* (2001), 167:5767-5774.
- Wang, Y., et al., "CD40 Is a Cellular Receptor Mediating Mycobacterial Heat Shock Protein 70 Stimulation of CC-Chemokines," *Immunity* (2001), 15:971-983.

(56)

References Cited

OTHER PUBLICATIONS

Accession No. NP_498023, Nov. 2008, pp. 1-2.
Balch, et al., *J Biol Chem.* 273(29): 18656-64, 1998.
Bayer, et al., *Trends Biotechnol.* 12: 379-386, 1994.
Burgdorf, et al., *J Immunol.* 176:6770-6, 2006.
Daly, et al., *Anal Lett.* 34:1799-1827, 2001.
Engering, et al., *Trends Immunol.* 23(10): 480-5, 2002.
Janeway, et al., *Immunobiol.* 3:3-3: 4, 1997.
Li, et al., *Clin Exp Immunol.* 162:251-61, 2010.
Luft, et al., *Internat Immunol.* 14(4):367-380, 2002.

Ma, "Genome-Wide Analysis of Human Peripheral Leukocyte Gene Expression," PhD Thesis, p. 51 & 60, 2003.
Office Communication in Israeli Patent Application No. 224539 mailed Apr. 23, 2013.
Riemer, et al., *Mol Immunol.* 42: 1121-4, 2005.
Rocca-Serra, et al., *Nature.* 304:353-355, 1983.
Rose, et al., *J. Immunol.* MRC1 mannose receptor, C type 1 [*Homo sapiens*], 1999, <http://www.ncbi.nlm.nih.gov/gene/4360>.
Chu, et al., *J. Immunol.* 163: 1188-94, 1999.
Stancovski, et al., *PNAS.* 88:8691-5, 1991.
Zymosan, Dectin-1 Review, "A Major Receptor in Antifungal Immunity", 2008.

* cited by examiner




FIG. 1A




FIG. 1B




FIG. 1C





FIG. 1D

FIG. 1E






FIG. 2A





FIG. 2B




FIG. 2C





FIG. 2D




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 3D




FIG. 4A




FIG. 4B




FIG. 4C




FIG. 4D




FIG. 5




FIG. 6




FIG. 7




FIG. 8A




FIG. 8B




FIG. 8C




FIG. 8D

1

**AGENTS THAT ENGAGE
ANTIGEN-PRESENTING CELLS THROUGH
DENDRITIC CELL ASIALOGLYCOPROTEIN
RECEPTOR (DC-ASGPR)**

CROSS-REFERENCE TO RELATED
APPLICATIONS

This application is a divisional of U.S. patent application Ser. No. 12/025,010 filed Feb. 2, 2008 which claims priority to U.S. Provisional Application Ser. No. 60/888,036, filed Feb. 2, 2007, the contents of which are incorporated by reference herein in its entirety.

STATEMENT OF FEDERALLY FUNDED
RESEARCH

This invention was made with U.S. Government support under Contract No. 1U19AI057234-0100003 awarded by the NIH. The government has certain rights in this invention.

TECHNICAL FIELD OF THE INVENTION

The present invention relates in general to the field of agents that engage antigen-presenting cells through dendritic cell asialoglycoprotein receptor (DC-ASGPR).

REFERENCE TO A SEQUENCE LISTING

The present application includes a Sequence Listing filed separately as required by 37 CFR 1.821-1.825.

BACKGROUND OF THE INVENTION

Without limiting the scope of the invention, its background is described in connection with antigen presentation.

Dendritic Cells play a pivotal role in controlling the interface of innate and acquired immunity by providing soluble and intercellular signals, followed by recognition of pathogens. These functions of DCs are largely dependent on the expression of specialized surface receptors, 'pattern recognition receptors' (PRRs), represented, most notably, by toll-like receptors (TLRs) and C-type lectins or lectin-like receptors (LLRs) (1-3).

In the current paradigm, a major role of TLRs is to alert DCs to produce interleukin 12 (IL-12) and other inflammatory cytokines for initiating immune responses. C-type LLRs operate as constituents of the powerful antigen capture and uptake mechanism of macrophages and DCs (1). Compared to TLRs, however, LLRs might have broader ranges of biological functions that include cell migrations (4), intercellular interactions (5). These multiple functions of LLRs might be due to the facts that LLRs, unlike TLRs, can recognize both self and non-self. However, the complexity of LLRs, including the redundancy of a number of LLRs expressed in immune cells, has been one of the major obstacles to understand the detailed functions of individual LLRs. In addition, natural ligands for most of these receptors remain unidentified. Nonetheless, evidence from recent studies suggests that LLRs, in collaboration with TLRs, may contribute to the activation of immune cells during microbial infections (6-14).

Valladeau et al. (The Journal of Immunology, 2001, 167: 5767-5774) described a novel LLR receptor on immature human Dendritic Cells related to hepatic Asialoglycoprotein Receptor and demonstrated that it efficiently mediated endocytosis. DC-ASGPR mRNA was observed predomi-

2

nantly in immune tissues—in DC and granulocytes, but not in T, B, or NK cells, or monocytes. DC-ASGPR species were restricted to the CD14-derived DC obtained from CD34-derived progenitors, while absent from the CD1a-derived subset. Both monocyte-derived DC and tonsillar interstitial-type DC expressed DC-ASGPR protein, while Langerhans-type cells did not. Furthermore, DC-ASGPR was a feature of immaturity, as expression was lost upon CD40 activation. In agreement with the presence of tyrosine-based and dileucine motifs in the intracytoplasmic domain, mAb against DC-ASGPR was rapidly internalized by DC at 37° C. Finally, intracellular DC-ASGPR was localized to early endosomes, suggesting that the receptor recycles to the cell surface following internalization of ligand. These findings identified DC-ASGPR/human macrophage lectin as a feature of immature DC, and as another lectin important for the specialized Ag-capture function of DC.

SUMMARY OF THE INVENTION

While DC-ASGPR is known to be capable of directing the internalization of surrogate antigen into human DC, the invention uses novel biological activities of DC-ASGPR to effect particularly desirable changes in the immune system, some in the context of antigen uptake (e.g., vaccination), others through the unique action of DC-ASGPR effectors (alone or in concert with other immune regulatory molecules) capable of eliciting signaling through this receptor on DC, B cells, and monocytes. The invention disclosure reveals means of developing unique agents capable of activating cells bearing DC-ASGPR, as well as the effect of the resulting changes in cells receiving these signals regards action on other cells in the immune system. These effects (either alone, or in concert with other signals (i.e., co-stimulation)) are highly predictive of therapeutic outcomes for certain disease states or for augmenting protective outcomes in the context of vaccination.

The present invention includes compositions and methods for increasing the effectiveness of antigen presentation by a DC-ASGPR-expressing antigen presenting cell by isolating and purifying a DC-ASGPR-specific antibody or fragment thereof to which a targeted agent is attached that forms an antibody-antigen complex, wherein the agent is processed and presented by, e.g., a dendritic cell, that has been contacted with the antibody-agent complex. In one embodiment, the antigen presenting cell is a dendritic cell and the DC-ASGPR-specific antibody or fragment thereof is bound to one half of a Coherin/Dockerin pair. The DC-ASGPR-specific antibody or fragment thereof may also be bound to one half of a Coherin/Dockerin pair and an antigen is bound to the complementary half of the Coherin/Dockerin pair to form a complex. Non-limiting examples agents include one or more peptides, proteins, lipids, carbohydrates, nucleic acids and combinations thereof.

The agent may one or more cytokine selected from interleukins, transforming growth factors (TGFs), fibroblast growth factors (FGFs), platelet derived growth factors (PDGFs), epidermal growth factors (EGFs), connective tissue activated peptides (CTAPs), osteogenic factors, and biologically active analogs, fragments, and derivatives of such growth factors, B/T-cell differentiation factors, B/T-cell growth factors, mitogenic cytokines, chemotactic cytokines, colony stimulating factors, angiogenesis factors, IFN- α , IFN- β , IFN- γ , IL1, IL2, IL3, IL4, IL5, IL6, IL7, IL8, IL9, IL10, IL11, IL12, IL13, IL14, IL15, IL16, IL17, IL18, etc., leptin, myostatin, macrophage stimulating protein, platelet-derived growth factor, TNF- α , TNF- β , NGF, CD40L, CD137L/4-1BBL, human lymphotoxin- β , G-CSF, M-CSF, GM-CSF,

PDGF, IL-1 α , IL1- β , IP-10, PF4, GRO, 9E3, erythropoietin, endostatin, angiostatin, VEGF, transforming growth factor (TGF) supergene family include the beta transforming growth factors (for example TGF- β 1, TGF- β 2, TGF- β 3); bone morphogenetic proteins (for example, BMP-1, BMP-2, BMP-3, BMP-4, BMP-5, BMP-6, BMP-7, BMP-8, BMP-9); heparin-binding growth factors (fibroblast growth factor (FGF), epidermal growth factor (EGF), platelet-derived growth factor (PDGF), insulin-like growth factor (IGF)); Inhibins (for example, Inhibin A, Inhibin B); growth differentiating factors (for example, GDF-1); and Activins (for example, Activin A, Activin B, Activin AB). In another embodiment, the agent comprises an antigen that is a bacterial, viral, fungal, protozoan or cancer protein.

The present invention also includes compositions and methods for increasing the effectiveness of antigen presentation by dendritic cells comprising binding a DC-ASGPR-specific antibody or fragment thereof to which an antigen is attached that forms an antibody-antigen complex, wherein the antigen is processed and presented by a dendritic cell that has been contacted with the antibody-antigen complex. Another embodiment is the use of antibodies or other specific binding molecules directed to DC-ASGPR for delivering antigens to antigen-presenting cells for the purpose of eliciting protective or therapeutic immune responses. The use of antigen-targeting reagents specific to DC-ASGPR for vaccination via the skin; antigen-targeting reagents specific to DC-ASGPR in association with co-administered or linked adjuvant for vaccination or use for antigen-targeting (vaccination) purposes of specific antigens which can be expressed as recombinant antigen-antibody fusion proteins.

Another embodiment includes a method for increasing the effectiveness of dendritic cells by isolating patient dendritic cells; exposing the dendritic cells to activating amounts of anti-DC-ASGPR antibodies or fragments thereof and antigen to form antigen-loaded, activated dendritic cells; and reintroducing the antigen-loaded, activated dendritic cells into the patient. The antigen may be a bacterial, viral, fungal, protozoan or cancer protein. The present invention also includes an anti-DC-ASGPR immunoglobulin or portion thereof that is secreted from mammalian cells and an antigen bound to the immunoglobulin. The immunoglobulin is bound to one half of a cohesin/dockerin domain, or it may also include a complementary half of the cohesin-dockerin binding pair bound to an antigen that forms a complex with the modular rAb carrier, or a complementary half of the cohesin-dockerin binding pair that is a fusion protein with an antigen. The antigen specific domain may be a full length antibody, an antibody variable region domain, an Fab fragment, a Fab' fragment, an F(ab)₂ fragment, and Fv fragment, and Fabc fragment and/or a Fab fragment with portions of the Fc domain. The anti-DC-ASGPR immunoglobulin may also be bound to a toxin selected from wherein the toxin is selected from the group consisting of a radioactive isotope, metal, enzyme, botulin, tetanus, ricin, cholera, diphtheria, aflatoxins, perfringens toxin, mycotoxins, shigatoxin, staphylococcal enterotoxin B, T2, seguitoxin, saxitoxin, abrin, cyanoginosin, alphatoxin, tetrodotoxin, aconotoxin, snake venom and spider venom. The antigen may be a fusion protein with the immunoglobulin or bound chemically covalently or not.

The present invention also includes compositions and methods for increasing the effectiveness of dendritic cells by isolating patient dendritic cells, exposing the dendritic cells to activating amounts of anti-DC-ASGPR antibodies or fragments thereof and antigen to form antigen-loaded, activated dendritic cells; and reintroducing the antigen-loaded, activated dendritic cells into the patient. The agents may be used

to engage DC-ASGPR, alone or with co-activating agents, to activate antigen-presenting cells for therapeutic or protective applications, to bind DC-ASGPR and/or activating agents linked to antigens, alone or with co-activating agents, for protective or therapeutic vaccination. Another use of is the development of specific antibody V-region sequences capable of binding to and activating DC-ASGPR, for use as anti-DC-ASGPR agents linked to toxic agents for therapeutic purposes in the context of diseases known or suspected to result from inappropriate activation of immune cells via DC-ASGPR and as a vaccine with a DC-ASGPR-specific antibody or fragment thereof to which an antigen is attached that forms an antibody-antigen complex, wherein the antigen is processed and presented by a dendritic cell that has been contacted with the antibody-antigen complex.

BRIEF DESCRIPTION OF THE DRAWINGS

For a more complete understanding of the features and advantages of the present invention, reference is now made to the detailed description of the invention along with the accompanying figures and in which:

FIGS. 1A to 1E demonstrate signaling through lectin-like receptor DC-ASGPR activates DCs, resulting in increased levels of costimulatory molecules as well as cytokines and chemokines FIG. 1A shows three day and six day GM/IL-4 DCs were stained with FITC-labeled goat anti-mouse IgG followed by mouse monoclonal anti-human DC-ASGPR, antibody. FIG. 1B shows six day GM/IL-4 DCs were cultured in plates coated with the anti-DC-ASGPR or control mAbs (1-2 ug/ml) for 16-18 h. Cells were stained with anti-CD86 and HLA-DR antibodies labeled with fluorescent dyes. Open and filled bars in the histograms represent cells activated with isotype control mAbs and anti-lectin mAbs, respectively. FIG. 1C shows six day GM/IL-4 DCs were cultured in plates coated with the mAbs for 12 h, and subjected to RNA isolation and Affymetrix Gene Chip analysis, as described in Methods. Fold increases of gene expression by anti-lectin mAbs were compared with the gene expression levels in DCs stimulated with control mAbs. FIG. 1D shows the cytokines and chemokines in the culture supernatants from the experiment shown in FIG. 1B were measured by Luminex. FIG. 1E shows six day GM/IL-4 DCs were cultured in plates coated with mAbs in the presence or absence of 50 ng/ml soluble CD40L, for 16-18 h, and then stained with anti-CD83 antibodies. Cytokines and chemokines in the culture supernatants from the experiment shown in FIG. 1E were measured by Luminex. Results shown are representative of three independent experiments using cells from different normal donors.

FIGS. 2A to 2D shows that DC-ASGPR expressed on DCs, contributes to enhanced humoral immune responses. Six day GM/IL-4 DCs, 5×10^3 /well, were incubated in 96 well plates coated with anti-DC-ASGPR or control mAb for 16-18 h, and then 1×10^5 autologous CD19⁺ B cells stained with CFSE were co-cultured in the presence of 20 units/ml IL-2 and 50 nM CpG. FIG. 2A is a FACS of day six cells stained with fluorescently labeled antibodies. CD3⁺ and 7-AAD⁺ cells were gated out. CD38⁺ and CFSE⁻ cells were purified by FACS sorter and Giemsa staining was performed. FIG. 2B are culture supernatants on day thirteen were analyzed for total IgM, IgG, and IgM by sandwich ELISA. FIG. 1C shows DCs pulsed with 5 multiplicity of infection (moi) of heat-inactivated influenza virus (PR8), and cultured with B cells. Culture supernatant was analyzed for influenza-specific immunoglobulins (Igs) on day thirteen. FIG. 1D shows DC cultured

5

with anti-DC-ASGPR or control mAb were stained for cell surface APRIL expression and the supernatants assayed for soluble APRIL.

FIGS. 3A to 3D shows the cell surface expression of DC-ASGPR on B cells contribute to B cell activation and immunoglobulin production. FIG. 3A are PBMCs from buffy coats were stained with anti-CD19, anti-CD3, and anti-DC-ASGPR or control mAb. CD19⁺ and CD3⁺ cells were gated and the expression levels of the molecules on CD19⁺ B cells were measured by flow cytometry. FIG. 3B are CD19⁺ B cells from buffy coats were cultured in plates coated with the mAbs for 12 h, and subjected to RNA isolation and Affymetrix Gene Chip analysis as described in Methods. Fold increases of gene expression by anti-DC-ASGPR mAb were compared to the gene expression levels in CD19⁺ B cells stimulated with control mAb. FIG. 3C shows CD19⁺ B cells were cultured in plates coated with the mAbs for 16-18 h, and then culture supernatants were analyzed for cytokines and chemokines by Luminex. FIG. 3D shows 1×10⁵ CD19⁺ B cells were cultured in plates coated with the mAbs for thirteen days. Total Ig levels were measured by ELISA. Data are representative of two repeat experiments using cells from three different normal donors.

FIGS. 4A to 4D shows that the proliferation of purified allogeneic T cells was significantly enhanced by DCs stimulated with mAb specific for DC-ASGPR.

FIG. 5 shows that certain anti-DC-ASGPR mAbs can activate DC. GM-CSF/IL-4. DC were incubated for 24 hrs with one of a panel of 12 pure anti-ASGPR mAbs. Cells were then tested for expression of cell surface CD86 (a DC activation marker) and supernatants were assayed for secreted cytokines. Three mAbs (36, 38, 43) from the anti-ASGPR mAb panel activated DC.

FIG. 6 shows that different antigens can be expressed in the context of a DC-ASGPR rAb. Such an anti-DC-ASGPR rAb.Doc protein can be simply mixed with any Cohesin.fusion protein to assemble a stable non-covalent [rAb.Doc:Coh.fusion] complex that functions just as a rAb.fusion protein.

FIG. 7—GM-CSF/IFN α DCs (5,000/well) were loaded with 10 or 1 nM anti-DC-ASGPR.Doc:Coh.Flu M1, or hIgG4.Doc:Coh.Flu M1 complexes. After 6 h, autologous CD8⁺ T cells (200,000/well) were added into the cultures. At day 8, the CD8⁺ T cells were analyzed for expansion of cells bearing TCR specific for a HLA-A201 immunodominant peptide. The inner boxes indicate the percentage of tetramer-specific CD8⁺ T cells.

FIGS. 8A-8D demonstrated the cross reactivity of the different antibodies with monkey ASGPR.

DETAILED DESCRIPTION OF THE INVENTION

While the making and using of various embodiments of the present invention are discussed in detail below, it should be appreciated that the present invention provides many applicable inventive concepts that can be embodied in a wide variety of specific contexts. The specific embodiments discussed herein are merely illustrative of specific ways to make and use the invention and do not delimit the scope of the invention.

To facilitate the understanding of this invention, a number of terms are defined below. Terms defined herein have meanings as commonly understood by a person of ordinary skill in the areas relevant to the present invention. Terms such as “a”, “an” and “the” are not intended to refer to only a singular entity, but include the general class of which a specific example may be used for illustration. The terminology herein

6

is used to describe specific embodiments of the invention, but their usage does not delimit the invention, except as outlined in the claims.

Dendritic cells (DCs) are antigen-presenting cells that play a key role in regulating antigen-specific immunity (Mellman and Steinman 2001), (Banchereau, Briere et al. 2000), (Cella, Sallusto et al. 1997). DCs capture antigens, process them into peptides, and present these to T cells. Therefore delivering antigens directly to DC is a focus area for improving vaccines. One such example is the development of DC-based vaccines using ex-vivo antigen-loading of autologous DCs that are then re-administered to patients (Banchereau, Schuler-Thurner et al. 2001), (Steinman and Dhodapkar 2001). Another strategy to improve vaccine efficacy is specific targeting to DC of antigen conjugated to antibodies against internalizing DC-specific receptors. The potential of targeting DC for vaccination is highlighted by key mouse studies. In vivo, targeting with an anti-LOX-1 mAb coupled to ovalbumin (OVA) induced a protective CD8⁺ T cell response, via exogenous antigen cross-presentation toward the MHC class I pathway (Delneste, Magistrelli et al. 2002). Also, OVA conjugated to anti-DEC205 mAb in combination with a CD40L maturation stimulus enhanced the MHC class I-restricted presentation by DCs in vivo and led to the durable formation of effector memory CD8⁺ T cells (Bonifaz, Bonnyay et al. 2004). Both these studies showed dramatic dose-sparing (i.e., strong immune-responses at very low antigen doses) and suggested broader responses than normally seen with other types of OVA immunization. Recent work with targeting of HIV gag antigen to DC via DEC205 has extended these concepts to a clinically relevant antigen and confirmed the tenets of targeting antigen to DC—dramatic dose-sparing, protective responses from a single vaccination, and expansion of antigen-specific T cells in both the CD8 and CD4 compartments (Trumpfheller, Finke et al. 2006).

The present invention provides for the complexing of multiple antigens or proteins (engineered, expressed, and purified independently from the primary mAb) in a controlled, multivariable fashion, to one single primary recombinant mAb. Presently, there are methods for engineering site-specific biotinylation sites that provide for the addition of different proteins (each engineered separately linked to streptavidin) to the one primary mAb. However, the present invention provides for addition to the primary mAb of multiple combinations, in fixed equimolar ratios and locations, of separately engineered proteins.

As used herein, the term “modular rAb carrier” is used to describe a recombinant antibody system that has been engineered to provide the controlled modular addition of diverse antigens, activating proteins, or other antibodies to a single recombinant monoclonal antibody (mAb). The rAb may be a monoclonal antibody made using standard hybridoma techniques, recombinant antibody display, humanized monoclonal antibodies and the like. The modular rAb carrier can be used to, e.g., target (via one primary recombinant antibody against an internalizing receptor, e.g., a human dendritic cell receptor) multiple antigens and/or antigens and an activating cytokine to dendritic cells (DC). The modular rAb carrier may also be used to join two different recombinant mAbs end-to-end in a controlled and defined manner.

The antigen binding portion of the “modular rAb carrier” may be one or more variable domains, one or more variable and the first constant domain, an Fab fragment, a Fab' fragment, an F(ab)₂ fragment, and Fv fragment, and Fabc fragment and/or a Fab fragment with portions of the Fc domain to which the cognate modular binding portions are added to the amino acid sequence and/or bound. The antibody for use in

the modular rAb carrier can be of any isotype or class, subclass or from any source (animal and/or recombinant).

In one non-limiting example, the modular rAb carrier is engineered to have one or more modular cohesin-dockerin protein domains for making specific and defined protein complexes in the context of engineered recombinant mAbs. The mAb is a portion of a fusion protein that includes one or more modular cohesin-dockerin protein domains carboxy from the antigen binding domains of the mAb. The cohesin-dockerin protein domains may even be attached post-translationally, e.g., by using chemical cross-linkers and/or disulfide bonding.

The term "antigen" as used herein refers to a molecule that can initiate a humoral and/or cellular immune response in a recipient of the antigen. Antigen may be used in two different contexts with the present invention: as a target for the antibody or other antigen recognition domain of the rAb or as the molecule that is carried to and/or into a cell or target by the rAb as part of a dockerin/cohesin-molecule complement to the modular rAb carrier. The antigen is usually an agent that causes a disease for which a vaccination would be advantageous treatment. When the antigen is presented on MHC, the peptide is often about 8 to about 25 amino acids. Antigens include any type of biologic molecule, including, for example, simple intermediary metabolites, sugars, lipids and hormones as well as macromolecules such as complex carbohydrates, phospholipids, nucleic acids and proteins. Common categories of antigens include, but are not limited to, viral antigens, bacterial antigens, fungal antigens, protozoal and other parasitic antigens, tumor antigens, antigens involved in autoimmune disease, allergy and graft rejection, and other miscellaneous antigens.

The modular rAb carrier is able to carry any number of active agents, e.g., antibiotics, anti-infective agents, antiviral agents, anti-tumoral agents, antipyretics, analgesics, anti-inflammatory agents, therapeutic agents for osteoporosis, enzymes, cytokines, anticoagulants, polysaccharides, collagen, cells, and combinations of two or more of the foregoing active agents. Examples of antibiotics for delivery using the present invention include, without limitation, tetracycline, aminoglycosides, penicillins, cephalosporins, sulfonamide drugs, chloramphenicol sodium succinate, erythromycin, vancomycin, lincomycin, clindamycin, nystatin, amphotericin B, amantidine, idoxuridine, p-amino salicylic acid, isoniazid, rifampin, antinomycin D, mithramycin, daunomycin, adriamycin, bleomycin, vinblastine, vincristine, procarbazine, imidazole carboxamide, and the like.

Examples of anti-tumor agents for delivery using the present invention include, without limitation, doxorubicin, Daunorubicin, taxol, methotrexate, and the like. Examples of antipyretics and analgesics include aspirin, Motrin®, Ibuprofen®, naprosyn, acetaminophen, and the like.

Examples of anti-inflammatory agents for delivery using the present invention include, without limitation, include NSAIDS, aspirin, steroids, dexamethasone, hydrocortisone, prednisolone, Diclofenac Na, and the like.

Examples of therapeutic agents for treating osteoporosis and other factors acting on bone and skeleton include for delivery using the present invention include, without limitation, calcium, alendronate, bone GLa peptide, parathyroid hormone and its active fragments, histone H4-related bone formation and proliferation peptide and mutations, derivatives and analogs thereof.

Examples of enzymes and enzyme cofactors for delivery using the present invention include, without limitation, pancrease, L-asparaginase, hyaluronidase, chymotrypsin, trypsin, tPA, streptokinase, urokinase, pancreatin, colla-

nase, trypsinogen, chymotrypsinogen, plasminogen, streptokinase, adenyl cyclase, superoxide dismutase (SOD), and the like.

Examples of cytokines for delivery using the present invention include, without limitation, interleukins, transforming growth factors (TGFs), fibroblast growth factors (FGFs), platelet derived growth factors (PDGFs), epidermal growth factors (EGFs), connective tissue activated peptides (CTAPs), osteogenic factors, and biologically active analogs, fragments, and derivatives of such growth factors. Cytokines may be B/T-cell differentiation factors, B/T-cell growth factors, mitogenic cytokines, chemotactic cytokines, colony stimulating factors, angiogenesis factors, IFN- α , IFN- β , IFN- γ , IL1, IL2, IL3, IL4, IL5, IL6, IL7, IL8, IL9, IL10, IL11, IL12, IL13, IL14, IL15, IL16, IL17, IL18, etc., leptin, myostatin, macrophage stimulating protein, platelet-derived growth factor, TNF- α , TNF- β , NGF, CD40L, CD137L/4-1BBL, human lymphotoxin- β , G-CSF, M-CSF, GM-CSF, PDGF, IL-1 α , IL-1- β , IP-10, PF4, GRO, 9E3, erythropoietin, endostatin, angiostatin, VEGF or any fragments or combinations thereof. Other cytokines include members of the transforming growth factor (TGF) supergene family include the beta transforming growth factors (for example TGF- β 1, TGF- β 2, TGF- β 3); bone morphogenetic proteins (for example, BMP-1, BMP-2, BMP-3, BMP-4, BMP-5, BMP-6, BMP-7, BMP-8, BMP-9); heparin-binding growth factors (for example, fibroblast growth factor (FGF), epidermal growth factor (EGF), platelet-derived growth factor (PDGF), insulin-like growth factor (IGF)); Inhibins (for example, Inhibin A, Inhibin B); growth differentiating factors (for example, GDF-1); and Activins (for example, Activin A, Activin B, Activin AB).

Examples of growth factors for delivery using the present invention include, without limitation, growth factors that can be isolated from native or natural sources, such as from mammalian cells, or can be prepared synthetically, such as by recombinant DNA techniques or by various chemical processes. In addition, analogs, fragments, or derivatives of these factors can be used, provided that they exhibit at least some of the biological activity of the native molecule. For example, analogs can be prepared by expression of genes altered by site-specific mutagenesis or other genetic engineering techniques.

Examples of anticoagulants for delivery using the present invention include, without limitation, include warfarin, heparin, Hirudin, and the like. Examples of factors acting on the immune system include for delivery using the present invention include, without limitation, factors which control inflammation and malignant neoplasms and factors which attack infective microorganisms, such as chemotactic peptides and bradykinins.

Examples of viral antigens include, but are not limited to, e.g., retroviral antigens such as retroviral antigens from the human immunodeficiency virus (HIV) antigens such as gene products of the gag, pol, and env genes, the Nef protein, reverse transcriptase, and other HIV components; hepatitis viral antigens such as the S, M, and L proteins of hepatitis B virus, the pre-S antigen of hepatitis B virus, and other hepatitis, e.g., hepatitis A, B, and C, viral components such as hepatitis C viral RNA; influenza viral antigens such as hemagglutinin and neuraminidase and other influenza viral components; measles viral antigens such as the measles virus fusion protein and other measles virus components; rubella viral antigens such as proteins E1 and E2 and other rubella virus components; rotaviral antigens such as VP7sc and other rotaviral components; cytomegaloviral antigens such as envelope glycoprotein B and other cytomegaloviral antigen com-

ponents; respiratory syncytial viral antigens such as the RSV fusion protein, the M2 protein and other respiratory syncytial viral antigen components; herpes simplex viral antigens such as immediate early proteins, glycoprotein D, and other herpes simplex viral antigen components; varicella zoster viral antigens such as gpI, gpII, and other varicella zoster viral antigen components; Japanese encephalitis viral antigens such as proteins E, M-E, M-E-NS1, NS1, NS1-NS2A, 80% E, and other Japanese encephalitis viral antigen components; rabies viral antigens such as rabies glycoprotein, rabies nucleoprotein and other rabies viral antigen components. See Fundamental Virology, Second Edition, eds. Fields, B. N. and Knipe, D. M. (Raven Press, New York, 1991) for additional examples of viral antigens.

Antigenic targets that may be delivered using the rAb-DC/DC-antigen vaccines of the present invention include genes encoding antigens such as viral antigens, bacterial antigens, fungal antigens or parasitic antigens. Viruses include picornavirus, coronavirus, togavirus, flavivirus, rhabdovirus, paramyxovirus, orthomyxovirus, bunyavirus, arenavirus, reovirus, retrovirus, papillomavirus, parvovirus, herpesvirus, poxvirus, hepadnavirus, and spongiform virus. Other viral targets include influenza, herpes simplex virus 1 and 2, measles, dengue, smallpox, polio or HIV. Pathogens include trypanosomes, tapeworms, roundworms, helminthes, and malaria. Tumor markers, such as fetal antigen or prostate specific antigen, may be targeted in this manner. Other examples include: HIV env proteins and hepatitis B surface antigen. Administration of a vector according to the present invention for vaccination purposes would require that the vector-associated antigens be sufficiently non-immunogenic to enable long term expression of the transgene, for which a strong immune response would be desired. In some cases, vaccination of an individual may only be required infrequently, such as yearly or biennially, and provide long term immunologic protection against the infectious agent. Specific examples of organisms, allergens and nucleic and amino sequences for use in vectors and ultimately as antigens with the present invention may be found in U.S. Pat. No. 6,541, 011, relevant portions incorporated herein by reference, in particular, the tables that match organisms and specific sequences that may be used with the present invention.

Bacterial antigens for use with the rAb vaccine disclosed herein include, but are not limited to, e.g., bacterial antigens such as pertussis toxin, filamentous hemagglutinin, pertactin, FIM2, FIM3, adenylate cyclase and other pertussis bacterial antigen components; diphtheria bacterial antigens such as diphtheria toxin or toxoid and other diphtheria bacterial antigen components; tetanus bacterial antigens such as tetanus toxin or toxoid and other tetanus bacterial antigen components; streptococcal bacterial antigens such as M proteins and other streptococcal bacterial antigen components; gram-negative bacilli bacterial antigens such as lipopolysaccharides and other gram-negative bacterial antigen components, *Mycobacterium tuberculosis* bacterial antigens such as mycolic acid, heat shock protein 65 (HSP65), the 30 kDa major secreted protein, antigen 85A and other mycobacterial antigen components; *Helicobacter pylori* bacterial antigen components; pneumococcal bacterial antigens such as pneumolysin, pneumococcal capsular polysaccharides and other pneumococcal bacterial antigen components; *haemophilus influenzae* bacterial antigens such as capsular polysaccharides and other *haemophilus influenzae* bacterial antigen components; anthrax bacterial antigens such as anthrax protective antigen and other anthrax bacterial antigen components; rickettsiae bacterial antigens such as rompA and other rickettsiae bacterial antigen component. Also included with the bacterial antigens

described herein are any other bacterial, mycobacterial, mycoplasmal, rickettsial, or chlamydial antigens. Partial or whole pathogens may also be: *haemophilus influenzae*; *Plasmodium falciparum*; *neisseria meningitidis*; *streptococcus pneumoniae*; *neisseria gonorrhoeae*; *salmonella* serotype typhi; *shigella*; *vibrio cholerae*; Dengue Fever; Encephalitides; Japanese Encephalitis; Lyme disease; *Yersinia pestis*; west nile virus; yellow fever; tularemia; hepatitis (viral; bacterial); RSV (respiratory syncytial virus); HPIV 1 and HPIV 3; adenovirus; small pox; allergies and cancers.

Fungal antigens for use with compositions and methods of the invention include, but are not limited to, e.g., candida fungal antigen components; histoplasma fungal antigens such as heat shock protein 60 (HSP60) and other histoplasma fungal antigen components; cryptococcal fungal antigens such as capsular polysaccharides and other cryptococcal fungal antigen components; coccidiodes fungal antigens such as spherule antigens and other coccidiodes fungal antigen components; and tinea fungal antigens such as trichophytin and other coccidiodes fungal antigen components.

Examples of protozoal and other parasitic antigens include, but are not limited to, e.g., *plasmodium falciparum* antigens such as merozoite surface antigens, sporozoite surface antigens, circumsporozoite antigens, gametocyte/gamete surface antigens, blood-stage antigen pf 155/RESA and other plasmodial antigen components; toxoplasma antigens such as SAG-1, p30 and other toxoplasmal antigen components; schistosomae antigens such as glutathione-S-transferase, paramyosin, and other schistosomal antigen components; *leishmania major* and other leishmaniae antigens such as gp63, lipophosphoglycan and its associated protein and other leishmanial antigen components; and *trypanosoma cruzi* antigens such as the 75-77 kDa antigen, the 56 kDa antigen and other trypanosomal antigen components.

Antigen that can be targeted using the rAb of the present invention will generally be selected based on a number of factors, including: likelihood of internalization, level of immune cell specificity, type of immune cell targeted, level of immune cell maturity and/or activation and the like. Examples of cell surface markers for dendritic cells include, but are not limited to, MHC class I, MHC Class II, B7-2, CD18, CD29, CD31, CD43, CD44, CD45, CD54, CD58, CD83, CD86, CMRF-44, CMRF-56, DCIR and/or ASPGR and the like; while in some cases also having the absence of CD2, CD3, CD4, CD8, CD14, CD15, CD16, CD 19, CD20, CD56, and/or CD57. Examples of cell surface markers for antigen presenting cells include, but are not limited to, MHC class I, MHC Class II, CD40, CD45, B7-1, B7-2, IFN- γ receptor and IL-2 receptor, ICAM-1 and/or Fc γ receptor. Examples of cell surface markers for T cells include, but are not limited to, CD3, CD4, CD8, CD 14, CD20, CD11b, CD16, CD45 and HLA-DR.

Target antigens on cell surfaces for delivery includes those characteristic of tumor antigens typically will be derived from the cell surface, cytoplasm, nucleus, organelles and the like of cells of tumor tissue. Examples of tumor targets for the antibody portion of the present invention include, without limitation, hematological cancers such as leukemias and lymphomas, neurological tumors such as astrocytomas or glioblastomas, melanoma, breast cancer, lung cancer, head and neck cancer, gastrointestinal tumors such as gastric or colon cancer, liver cancer, pancreatic cancer, genitourinary tumors such cervix, uterus, ovarian cancer, vaginal cancer, testicular cancer, prostate cancer or penile cancer, bone tumors, vascular tumors, or cancers of the lip, nasopharynx, pharynx and oral cavity, esophagus, rectum, gall bladder, biliary tree, larynx, lung and bronchus, bladder, kidney, brain

and other parts of the nervous system, thyroid, Hodgkin's disease, non-Hodgkin's lymphoma, multiple myeloma and leukemia.

Examples of antigens that may be delivered alone or in combination to immune cells for antigen presentation using the present invention include tumor proteins, e.g., mutated oncogenes; viral proteins associated with tumors; and tumor mucins and glycolipids. The antigens may be viral proteins associated with tumors would be those from the classes of viruses noted above. Certain antigens may be characteristic of tumors (one subset being proteins not usually expressed by a tumor precursor cell), or may be a protein which is normally expressed in a tumor precursor cell, but having a mutation characteristic of a tumor. Other antigens include mutant variant(s) of the normal protein having an altered activity or subcellular distribution, e.g., mutations of genes giving rise to tumor antigens.

Specific non-limiting examples of tumor antigens include: CEA, prostate specific antigen (PSA), HER-2/neu, BAGE, GAGE, MAGE 1-4, 6 and 12, MUC (Mucin) (e.g., MUC-1, MUC-2, etc.), GM2 and GD2 gangliosides, ras, myc, tyrosinase, MART (melanoma antigen), Pmel 17(gp100), GnT-V intron V sequence (N-acetylglucosaminyltransferase V intron V sequence), Prostate Ca psm, PRAME (melanoma antigen), β -catenin, MUM-1-B (melanoma ubiquitous mutated gene product), GAGE (melanoma antigen) 1, BAGE (melanoma antigen) 2-10, c-ERB2 (Her2/neu), EBNA (Epstein-Barr Virus nuclear antigen) 1-6, gp75, human papilloma virus (HPV) E6 and E7, p53, lung resistance protein (LRP), Bcl-2, and Ki-67. In addition, the immunogenic molecule can be an autoantigen involved in the initiation and/or propagation of an autoimmune disease, the pathology of which is largely due to the activity of antibodies specific for a molecule expressed by the relevant target organ, tissue, or cells, e.g., SLE or MG. In such diseases, it can be desirable to direct an ongoing antibody-mediated (i.e., a Th2-type) immune response to the relevant autoantigen towards a cellular (i.e., a Th1-type) immune response. Alternatively, it can be desirable to prevent onset of or decrease the level of a Th2 response to the autoantigen in a subject not having, but who is suspected of being susceptible to, the relevant autoimmune disease by prophylactically inducing a Th1 response to the appropriate autoantigen. Autoantigens of interest include, without limitation: (a) with respect to SLE, the Smith protein, RNP ribonucleoprotein, and the SS-A and SS-B proteins; and (b) with respect to MG, the acetylcholine receptor. Examples of other miscellaneous antigens involved in one or more types of autoimmune response include, e.g., endogenous hormones such as luteinizing hormone, follicular stimulating hormone, testosterone, growth hormone, prolactin, and other hormones.

Antigens involved in autoimmune diseases, allergy, and graft rejection can be used in the compositions and methods of the invention. For example, an antigen involved in any one or more of the following autoimmune diseases or disorders can be used in the present invention: diabetes, diabetes mellitus, arthritis (including rheumatoid arthritis, juvenile rheumatoid arthritis, osteoarthritis, psoriatic arthritis), multiple sclerosis, myasthenia gravis, systemic lupus erythematosus, autoimmune thyroiditis, dermatitis (including atopic dermatitis and eczematous dermatitis), psoriasis, Sjogren's Syndrome, including keratoconjunctivitis sicca secondary to Sjogren's Syndrome, alopecia areata, allergic responses due to arthropod bite reactions, Crohn's disease, aphthous ulcer, iritis, conjunctivitis, keratoconjunctivitis, ulcerative colitis, asthma, allergic asthma, cutaneous lupus erythematosus, scleroderma, vaginitis, proctitis, drug eruptions, leprosy reversal reactions, erythema nodosum leprosum, autoim-

mune uveitis, allergic encephalomyelitis, acute necrotizing hemorrhagic encephalopathy, idiopathic bilateral progressive sensorineural hearing loss, aplastic anemia, pure red cell anemia, idiopathic thrombocytopenia, polychondritis, Wegener's granulomatosis, chronic active hepatitis, Stevens-Johnson syndrome, idiopathic sprue, lichen planus, Crohn's disease, Graves ophthalmopathy, sarcoidosis, primary biliary cirrhosis, uveitis posterior, and interstitial lung fibrosis. Examples of antigens involved in autoimmune disease include glutamic acid decarboxylase 65 (GAD 65), native DNA, myelin basic protein, myelin proteolipid protein, acetylcholine receptor components, thyroglobulin, and the thyroid stimulating hormone (TSH) receptor. Examples of antigens involved in allergy include pollen antigens such as Japanese cedar pollen antigens, ragweed pollen antigens, rye grass pollen antigens, animal derived antigens such as dust mite antigens and feline antigens, histocompatibility antigens, and penicillin and other therapeutic drugs. Examples of antigens involved in graft rejection include antigenic components of the graft to be transplanted into the graft recipient such as heart, lung, liver, pancreas, kidney, and neural graft components. The antigen may be an altered peptide ligand useful in treating an autoimmune disease.

As used herein, the term "epitope(s)" refer to a peptide or protein antigen that includes a primary, secondary or tertiary structure similar to an epitope located within any of a number of pathogen polypeptides encoded by the pathogen DNA or RNA. The level of similarity will generally be to such a degree that monoclonal or polyclonal antibodies directed against such polypeptides will also bind to, react with, or otherwise recognize, the peptide or protein antigen. Various immunoassay methods may be employed in conjunction with such antibodies, such as, for example, Western blotting, ELISA, RIA, and the like, all of which are known to those of skill in the art. The identification of pathogen epitopes, and/or their functional equivalents, suitable for use in vaccines is part of the present invention. Once isolated and identified, one may readily obtain functional equivalents. For example, one may employ the methods of Hopp, as taught in U.S. Pat. No. 4,554,101, incorporated herein by reference, which teaches the identification and preparation of epitopes from amino acid sequences on the basis of hydrophilicity. The methods described in several other papers, and software programs based thereon, can also be used to identify epitopic core sequences (see, for example, Jameson and Wolf, 1988; Wolf et al., 1988; U.S. Pat. No. 4,554,101). The amino acid sequence of these "epitopic core sequences" may then be readily incorporated into peptides, either through the application of peptide synthesis or recombinant technology.

The preparation of vaccine compositions that includes the nucleic acids that encode antigens of the invention as the active ingredient, may be prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid prior to infection can also be prepared. The preparation may be emulsified, encapsulated in liposomes. The active immunogenic ingredients are often mixed with carriers which are pharmaceutically acceptable and compatible with the active ingredient.

The term "pharmaceutically acceptable carrier" refers to a carrier that does not cause an allergic reaction or other untoward effect in subjects to whom it is administered. Suitable pharmaceutically acceptable carriers include, for example, one or more of water, saline, phosphate buffered saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the vaccine can contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, and/or adjuvants which enhance the

effectiveness of the vaccine. Examples of adjuvants that may be effective include but are not limited to: aluminum hydroxide, N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-nor-muramyl-L-alanyl-D-isoglutamine, MTP-PE and RIBI, which contains three components extracted from bacteria, monophosphoryl lipid A, trehalose dimycolate and cell wall skeleton (MPL+TDM+CWS) in a 2% squalene/Tween 80 emulsion. Other examples of adjuvants include DDA (dimethyldioctadecylammonium bromide), Freund's complete and incomplete adjuvants and QuilA. In addition, immune modulating substances such as lymphokines (e.g., IFN- γ , IL-2 and IL-12) or synthetic IFN- γ inducers such as poly I:C can be used in combination with adjuvants described herein.

Pharmaceutical products that may include a naked polynucleotide with a single or multiple copies of the specific nucleotide sequences that bind to specific DNA-binding sites of the apolipoproteins present on plasma lipoproteins as described in the current invention. The polynucleotide may encode a biologically active peptide, antisense RNA, or ribozyme and will be provided in a physiologically acceptable administrable form. Another pharmaceutical product that may spring from the current invention may include a highly purified plasma lipoprotein fraction, isolated according to the methodology, described herein from either the patients blood or other source, and a polynucleotide containing single or multiple copies of the specific nucleotide sequences that bind to specific DNA-binding sites of the apolipoproteins present on plasma lipoproteins, prebound to the purified lipoprotein fraction in a physiologically acceptable, administrable form.

Yet another pharmaceutical product may include a highly purified plasma lipoprotein fraction which contains recombinant apolipoprotein fragments containing single or multiple copies of specific DNA-binding motifs, prebound to a polynucleotide containing single or multiple copies of the specific nucleotide sequences, in a physiologically acceptable administrable form. Yet another pharmaceutical product may include a highly purified plasma lipoprotein fraction which contains recombinant apolipoprotein fragments containing single or multiple copies of specific DNA-binding motifs, prebound to a polynucleotide containing single or multiple copies of the specific nucleotide sequences, in a physiologically acceptable administrable form.

The dosage to be administered depends to a great extent on the body weight and physical condition of the subject being treated as well as the route of administration and frequency of treatment. A pharmaceutical composition that includes the naked polynucleotide prebound to a highly purified lipoprotein fraction may be administered in amounts ranging from 1 μ g to 1 mg polynucleotide and 1 μ g to 100 mg protein.

Administration of an rAb and rAb complexes a patient will follow general protocols for the administration of chemotherapeutics, taking into account the toxicity, if any, of the vector. It is anticipated that the treatment cycles would be repeated as necessary. It also is contemplated that various standard therapies, as well as surgical intervention, may be applied in combination with the described gene therapy.

Where clinical application of a gene therapy is contemplated, it will be necessary to prepare the complex as a pharmaceutical composition appropriate for the intended application. Generally this will entail preparing a pharmaceutical composition that is essentially free of pyrogens, as well as any other impurities that could be harmful to humans or animals. One also will generally desire to employ appropriate salts and buffers to render the complex stable and allow for complex uptake by target cells.

Aqueous compositions of the present invention may include an effective amount of the compound, dissolved or dispersed in a pharmaceutically acceptable carrier or aqueous medium. Such compositions can also be referred to as inocula. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients also can be incorporated into the compositions. The compositions of the present invention may include classic pharmaceutical preparations. Dispersions also can be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

Disease States. Depending on the particular disease to be treated, administration of therapeutic compositions according to the present invention will be via any common route so long as the target tissue is available via that route in order to maximize the delivery of antigen to a site for maximum (or in some cases minimum) immune response. Administration will generally be by orthotopic, intradermal, subcutaneous, intramuscular, intraperitoneal or intravenous injection. Other areas for delivery include: oral, nasal, buccal, rectal, vaginal or topical. Topical administration would be particularly advantageous for treatment of skin cancers. Such compositions would normally be administered as pharmaceutically acceptable compositions that include physiologically acceptable carriers, buffers or other excipients.

Vaccine or treatment compositions of the invention may be administered parenterally, by injection, for example, either subcutaneously or intramuscularly. Additional formulations which are suitable for other modes of administration include suppositories, and in some cases, oral formulations or formulations suitable for distribution as aerosols. In the case of the oral formulations, the manipulation of T-cell subsets employing adjuvants, antigen packaging, or the addition of individual cytokines to various formulation that result in improved oral vaccines with optimized immune responses. For suppositories, traditional binders and carriers may include, for example, polyalkylene glycols or triglycerides; such suppositories may be formed from mixtures containing the active ingredient in the range of 0.5% to 10%, preferably 1%-2%. Oral formulations include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, and the like. These compositions take the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations or powders and contain 10%-95% of active ingredient, preferably 25-70%.

The antigen encoding nucleic acids of the invention may be formulated into the vaccine or treatment compositions as neutral or salt forms. Pharmaceutically acceptable salts include the acid addition salts (formed with free amino groups of the peptide) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or with organic acids such as acetic, oxalic, tartaric, maleic, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine, and the like.

Vaccine or treatment compositions are administered in a manner compatible with the dosage formulation, and in such amount as will be prophylactically and/or therapeutically effective. The quantity to be administered depends on the

subject to be treated, including, e.g., capacity of the subject's immune system to synthesize antibodies, and the degree of protection or treatment desired. Suitable dosage ranges are of the order of several hundred micrograms active ingredient per vaccination with a range from about 0.1 mg to 1000 mg, such as in the range from about 1 mg to 300 mg, and preferably in the range from about 10 mg to 50 mg. Suitable regimens for initial administration and booster shots are also variable but are typified by an initial administration followed by subsequent inoculations or other administrations. Precise amounts of active ingredient required to be administered depend on the judgment of the practitioner and may be peculiar to each subject. It will be apparent to those of skill in the art that the therapeutically effective amount of nucleic acid molecule or fusion polypeptides of this invention will depend, inter alia, upon the administration schedule, the unit dose of antigen administered, whether the nucleic acid molecule or fusion polypeptide is administered in combination with other therapeutic agents, the immune status and health of the recipient, and the therapeutic activity of the particular nucleic acid molecule or fusion polypeptide.

The compositions can be given in a single dose schedule or in a multiple dose schedule. A multiple dose schedule is one in which a primary course of vaccination may include, e.g., 1-10 separate doses, followed by other doses given at subsequent time intervals required to maintain and or reinforce the immune response, for example, at 1-4 months for a second dose, and if needed, a subsequent dose(s) after several months. Periodic boosters at intervals of 1-5 years, usually 3 years, are desirable to maintain the desired levels of protective immunity. The course of the immunization can be followed by in vitro proliferation assays of peripheral blood lymphocytes (PBLs) co-cultured with ESAT6 or ST-CF, and by measuring the levels of IFN- γ released from the primed lymphocytes. The assays may be performed using conventional labels, such as radionucleotides, enzymes, fluorescent labels and the like. These techniques are known to one skilled in the art and can be found in U.S. Pat. Nos. 3,791,932, 4,174,384 and 3,949,064, relevant portions incorporated by reference.

The modular rAb carrier and/or conjugated rAb carrier-(cohesion/dockerin and/or dockerin-cohesin)-antigen complex (rAb-DC/DC-antigen vaccine) may be provided in one or more "unit doses" depending on whether the nucleic acid vectors are used, the final purified proteins, or the final vaccine form is used. Unit dose is defined as containing a predetermined-quantity of the therapeutic composition calculated to produce the desired responses in association with its administration, i.e., the appropriate route and treatment regimen. The quantity to be administered, and the particular route and formulation, are within the skill of those in the clinical arts. The subject to be treated may also be evaluated, in particular, the state of the subject's immune system and the protection desired. A unit dose need not be administered as a single injection but may include continuous infusion over a set period of time. Unit dose of the present invention may conveniently be described in terms of DNA/kg (or protein/Kg) body weight, with ranges between about 0.05, 0.10, 0.15, 0.20, 0.25, 0.5, 1, 10, 50, 100, 1,000 or more mg/DNA or protein/kg body weight are administered. Likewise the amount of rAb-DC/DC-antigen vaccine delivered can vary from about 0.2 to about 8.0 mg/kg body weight. Thus, in particular embodiments, 0.4 mg, 0.5 mg, 0.8 mg, 1.0 mg, 1.5 mg, 2.0 mg, 2.5 mg, 3.0 mg, 4.0 mg, 5.0 mg, 5.5 mg, 6.0 mg, 6.5 mg, 7.0 mg and 7.5 mg of the vaccine may be delivered to an individual in vivo. The dosage of rAb-DC/DC-antigen vaccine to be administered depends to a great extent on the

weight and physical condition of the subject being treated as well as the route of administration and the frequency of treatment. A pharmaceutical composition that includes a naked polynucleotide prebound to a liposomal or viral delivery vector may be administered in amounts ranging from 1 μ g to 1 mg polynucleotide to 1 μ g to 100 mg protein. Thus, particular compositions may include between about 1 μ g, 5 μ g, 10 μ g, 20 μ g, 30 μ g, 40 μ g, 50 μ g, 60 μ g, 70 μ g, 80 μ g, 100 μ g, 150 μ g, 200 μ g, 250 μ g, 500 μ g, 600 μ g, 700 μ g, 800 μ g, 900 μ g or 1,000 μ g polynucleotide or protein that is bound independently to 1 μ g, 5 μ g, 10 μ g, 20 μ g, 30 μ g, 40 μ g, 50 μ g, 60 μ g, 70 μ g, 80 μ g, 100 μ g, 150 μ g, 200 μ g, 250 μ g, 500 μ g, 600 μ g, 700 μ g, 800 μ g, 900 μ g, 1 mg, 1.5 mg, 5 mg, 10 mg, 20 mg, 30 mg, 40 mg, 50 mg, 60 mg, 70 mg, 80 mg, 90 mg or 100 mg vector.

The present invention was tested in an in vitro cellular system that measures immune stimulation of human Flu-specific T cells by dendritic cells to which Flu antigen has been targeted. The results shown herein demonstrate the specific expansion of such antigen specific cells at doses of the antigen which are by themselves ineffective in this system.

The present invention may also be used to make a modular rAb carrier that is, e.g., a recombinant humanized mAb (directed to a specific human dendritic cell receptor) complexed with protective antigens from Ricin, Anthrax toxin, and *Staphylococcus* B enterotoxin. The potential market for this entity is vaccination of all military personnel and stored vaccine held in reserve to administer to large population centers in response to any biothreat related to these agents. The invention has broad application to the design of vaccines in general, both for human and animal use. Industries of interest include the pharmaceutical and biotechnology industries.

The present invention includes compositions and methods, including vaccines, that specifically target (deliver) antigens to antigen-presenting cells (APCs) for the purpose of eliciting potent and broad immune responses directed against the antigen. These compositions evoke protective or therapeutic immune responses against the agent (pathogen or cancer) from which the antigen was derived. In addition the invention creates agents that are directly, or in concert with other agents, therapeutic through their specific engagement of a receptor called DC-ASGPR that is expressed on antigen-presenting cells.

The novel recombinant humanized mAb (directed to the specific human dendritic cell receptor DC-ASGPR) fused through the antibody (Ab) heavy chain to antigens known or suspected to encode protective antigens. These include as examples for vaccination against various agents—hemagglutinins from Influenza H5N1; HIV gag from attenuated toxins from Ricin, Anthrax toxin, and *Staphylococcus* B enterotoxin; 'strings' of antigenic peptides from melanoma antigens, etc. The present invention may be used as a preventative or therapeutic vaccination for at risk or infected patients. The invention has broad application for vaccination against many diseases and cancers, both for human and animal use. Industries that can use the present invention include the pharmaceutical and biotechnological.

The present invention can be used to target antigens to APC for vaccination purposes. It is not known which antigen internalizing receptor will be best suited for this purpose. The invention describes particularly advantageous features of DC-ASGPR as for this purpose. Furthermore, the invention shows that engaging DC-ASGPR can be beneficial in the sense of activating the immune system with highly predicted significant therapeutic benefit.

The present invention includes the development of high affinity monoclonal antibodies against human DC-ASGPR.

Receptor ectodomain.hIgG (human IgG1Fc) and AP (human placental alkaline phosphatase) fusion proteins were produced for immunization of mice and screening of mAbs, respectively. An expression construct for hDCIR ectodomain.IgG was described previously (Bates, Fournier et al. 1999) and used the mouse SLAM (mSLAM) signal peptide to direct secretion (Bendtsen, Nielsen et al. 2004). An expression vector for hDCIR ectodomain.AP was generated using PCR to amplify AP residues 133-1581 (gb|BC009647) while adding a proximal in-frame Xho I site and a distal TGA stop codon and Not I site. This Xho I-Not I fragment replaced the IgG coding sequence in the above hDCIR ectodomain.IgG vector. DC-ASGPR ectodomain constructs in the same Ig and AP vector series contained inserts encoding (bp 484-1251, gi|53832017). DC-ASGPR fusion proteins were produced using the FreeStyle™ 293 Expression System (Invitrogen) according to the manufacturer's protocol (1 mg total plasmid DNA with 1.3 ml 293 Fectin reagent/L of transfection). For rAb production, equal amounts of vector encoding the H and L chain were co-transfected. Transfected cells are cultured for 3 days, the culture supernatant was harvested and fresh media added with continued incubation for two days. The pooled supernatants were clarified by filtration. Receptor ectodomain.hIgG was purified by HiTrap protein A affinity chromatography with elution by 0.1 M glycine pH 2.7 and then dialyzed versus PBS. rAbs (recombinant antibodies described later) were purified similarly, by using HiTrap MabSelect™ columns. Mouse mAbs were generated by conventional cell fusion technology. Briefly, 6-week-old BALB/c mice were immunized intraperitoneally with 20 µg of receptor ectodomain.hIgGFc fusion protein with Ribi adjuvant, then boosts with 20 µg antigen 10 days and 15 days later. After 3 months, the mice were boosted again three days prior to taking the spleens. Alternately, mice were injected in the footpad with 1-10 µg antigen in Ribi adjuvant every 3-4 days over a 30-40 day period. 3-4 days after a final boost, draining lymph nodes were harvested. B cells from spleen or lymph node cells were fused with SP2/O-Ag 14 cells (Shulman, Wilde et al. 1978) using conventional techniques. ELISA was used to screen hybridoma supernatants against the receptor ectodomain fusion protein compared to the fusion partner alone, or versus the receptor ectodomain fused to AP (Bates, Fournier et al. 1999). Positive wells were then screened in FACS using 293F cells transiently transfected with expression plasmids encoding full-length receptor cDNAs. Selected hybridomas were single cell cloned and expanded in CEL-Line flasks (Intergra). Hybridoma supernatants were mixed with an equal volume of 1.5 M glycine, 3 M NaCl, 1xPBS, pH 7.8 and tumbled with MabSelect resin. The resin was washed with binding buffer and eluted with 0.1 M glycine, pH 2.7. Following neutralization with 2 M Tris, mAbs were dialyzed versus PBS.

Characterization of purified anti-DC-ASGPR monoclonal antibodies by direct ELISA. the relative affinities of several anti-DC-ASGPR mAbs by ELISA were determined (i.e., DC-ASGPR.Ig protein is immobilized on the microplate surface and the antibodies are tested in a dose titration series for their ability to bind to DC-ASGPR.Ig (as detected by an anti-mouse IgG.HRP conjugate reagent. In this example, PAB42 and PAB44 show higher affinity binding than other mAbs. The same mAbs fail to bind significantly to human Ig bound to the microplate surface. This shows that the mAbs react to the DC-ASGPR ectodomain part of the DC-ASGPR.Ig fusion protein (data not shown).

Characterization of purified anti-DC-ASGPR monoclonal antibodies by indirect ELISA. Next, the relative affinities of several anti-DC-ASGPR mAbs were determined by ELISA

(i.e., anti-DC-ASGPR mAb is immobilized on the microplate surface and tested in a dose titration series for their ability to bind to DC-ASGPR.AP reagent. It was found that the supernatants from the hybridomas listed as: PAB42, PAB44 and PAB54 show higher affinity binding than other mAbs (data not shown).

Characterization of anti-DC-ASGPR mAbs by FACS. The panel of mAbs was also tested by FACS versus 293F cells transfected with expression plasmid directing synthesis of cell surface DC-ASGPR. Mean fluorescence intensity of the signal was subtracted from the analogous signal versus non-transfected 293F cells. By this criterion, the mAbs are able to bind to specifically to the surface of cells bearing DC-ASGPR. Some mAbs, e.g., 37A7 appear particularly advantageous in this regard (data not shown).

FIGS. 1A to 1D shows that signaling through DC-ASGPR activates DCs. DCs are the primary immune cells that determine the results of immune responses, either induction or tolerance, depending on their activation (15). The role of LLRs in DC activation is not clear yet. Therefore, we tested whether triggering the LLR DC-ASGPR can result in the activation of DCs. Both three and six day in vitro cultured GM/IL-4 DCs express LOX-1, ASGPR, and CLEC-6 (FIG. 1A). Six day DCs were stimulated with mAb specific to DC-ASGPR, and data in FIG. 1B show that signals through DC-ASGPR could activate DCs, resulting in the increased expression of CD86 and HLA-DR. Triggering DC-ASGPR on DCs also resulted in the increased production of IL-6, MCP-1, IL-12p40, and IL-8 from DCs (FIG. 1C). Other cytokines and chemokines, TNF α , IP-10, MIP-1a, and IL-10, were also significantly increased (data not shown) by signaling through DC-ASGPR, suggesting that DC-ASGPR can deliver cellular signals to activate DCs. Consistently, DCs stimulated with DC-ASGPR specific mAb expressed increased levels of multiple genes, including co-stimulatory molecules as well as chemokine and cytokine-related genes (FIG. 1D). The possible contribution of LLRs in TLR2 and TLR4-mediated immune cell activation has been described previously (13, 16). We observed that signals through DC-ASGPR could synergize with signal through CD40 for a further activation of DCs (FIG. 1E). This is important because LLRs could serve as co-stimulatory molecules during in vivo DC activation. Taken together, data in FIG. 1 prove that signaling through DC-ASGPR can activate DCs and that DC-ASGPR serves as a co-stimulatory molecule for the activation of DCs. DC-ASGPR engagement during CD40-CD40L interaction results in dramatically increased production of IL-12p70.

DCs stimulated through DC-ASGPR induce potent humoral immune responses. DCs play an important role in humoral immune responses by providing signals for both T-dependent and T-independent B cell responses (19-22) and by transferring antigens to B cells (23, 24). In addition to DCs, signaling through TLR9 as a third signal is necessary for efficient B cell responses (25, 26).

Therefore, we tested the role of DC-ASGPR in DCs-mediated humoral immune responses in the presence of TLR9 ligand, CpG. Six day GM/IL-4 DCs were stimulated with anti-DC-ASGPR mAb, and then purified B cells were co-cultured. As shown in FIG. 2A, DCs activated with anti-DC-ASGPR mAb resulted in remarkably enhanced B cell proliferation (CFSE dilution) and plasma cell differentiation (CD38⁺CD20⁻), compared to DCs stimulated with control mAb. CD38⁺CD20⁻ B cells have a typical morphology of plasma cells, but they do not express CD138. The majority of proliferating cells did not express CCR2, CCR4, CCR6, or CCR7. The amounts of total immunoglobulins (Igs) produced

were measured by ELISA (FIG. 2B). Consistent with the data in FIG. 2A, B cells cultured with anti-DC-ASGPR-stimulated DCs resulted in significantly increased production of total IgM, IgG, and IgA. In addition to the total Igs, we also observed that DCs activated by triggering DC-ASGPR are more potent than DCs stimulated with control mAb for the production of influenza-virus-specific IgM, IgG, and IgA (FIG. 2C) by B cells, suggesting that DC-ASGPR-mediated DC activation contributes to both total and antigen specific humoral immune responses. We tested the role of DC-ASGPR in ex vivo antigen presenting cells (APCs) in humoral immune responses. Parts of APCs in PBMCs, including CD19⁺ and CD14⁺ cells, express DC-ASGPR (Supplementary FIG. 2). PBMCs from buffy coats were cultured in the plates coated with anti-DC-ASGPR mAb, and the total Igs and B cell proliferation were measured. Consistent with the data generated from DCs (FIG. 2A), APCs stimulated through DC-ASGPR resulted in enhanced B cell proliferation and plasma cell differentiation in the absence (upper panels in FIG. 2d) or presence (lower panels in FIG. 2D) of TLR9 ligand. The total IgM, IgG, and IgA were also significantly increased when PBMCs were cultured in the plates coated with mAb against DC-ASGPR (FIG. 2e). As shown in FIG. 1, DCs activated by signaling through DC-ASGPR have matured phenotypes and produce large amounts of inflammatory cytokines and chemokines, and both matured DC phenotypes and soluble factors from DCs could contribute to the enhanced B cells responses (FIG. 2). However, DC-derived B lymphocyte stimulator protein (BLyS, BAFF) and a proliferation-inducing ligand (APRIL) are also important molecules by which DCs can directly regulate human B cell proliferation and function (27-30). Therefore, we tested whether signals through DC-ASGPR could alter the expression levels of BLyS and APRIL. Data in FIG. 2d show that DCs stimulated through DC-ASGPR expressed increased levels of intracellular APRIL as well as APRIL secreted, but not BLyS (not shown). Expression levels of BLyS and APRIL receptors on B cells in the mixed cultures were measured, but there was no significant change (not shown).

DC-ASGPR contributes to B cell activation and Ig production. CD19⁺ B cells express DC-ASGPR (FIG. 3A). Therefore, we tested the role of DC-ASGPR in B cell activation. Data in FIG. 3B show that B cells stimulated through DC-ASGPR produced significantly higher amounts of chemokines. In addition to IL-8 and MIP-1a, slight increases in IL-6 and TNF α were also observed when B cells were stimulated with the anti-DC-ASGPR mAb, compared to control mAb. Genes related to cell activation were also up-regulated (FIG. 3C). B cells produced IgM, IgG, and IgA when they were stimulated through DC-ASGPR (FIG. 3D), suggesting that DC-ASGPR could play an important role in the maintenance of normal immunoglobulin levels in vivo. However, signaling through DC-ASGPR alone did not induce significant B cell proliferation.

Role of DC-ASGPR in T cell responses. DCs stimulated through DC-ASGPR express enhanced levels of co-stimulatory molecules and produce increased amounts of cytokines and chemokines (see FIG. 1), suggesting that DC-ASGPR contributes to cellular immune responses as well as humoral immune responses. This was tested by a mixed lymphocyte reaction (MLR). Proliferation of purified allogeneic T cells was significantly enhanced by DCs stimulated with mAb specific for DC-ASGPR (FIG. 4A). DCs activated through DC-ASGPR could also prime Mart-1-specific CD8 T cells more efficiently than DC stimulated with control mAb (upper panels in FIG. 4B). More importantly, signaling through DC-ASGPR permitted DCs to cross-prime Mart-1 peptides to

CD8 T cells (lower panels in FIG. 4B). This indicates that DC-ASGPR plays an important role in enhancing DC function, resulting in better priming and cross-priming of antigens to CD8 T cells. The role of DC-ASGPR expressed on the mixture of APCs in PBMCs in activation of T cell responses is shown in FIG. 4C where PBMCs stimulated with mAb to DC-ASGPR resulted in an increased frequency of Flu M1 tetramer specific CD8 T cells compared to DCs stimulated with control mAb. This enhanced antigen specific CD8 T cell response was supported by the data in FIG. 4D, showing that DCs stimulated through DC-ASGPR significantly increase CD4 T cell proliferation.

Antibodies and tetramers—Antibodies (Abs) for surface staining of DCs and B cells, including isotype control Abs, were purchased from BD Biosciences (CA). Abs for ELISA were purchased from Bethyl (TX). Anti-BLyS and anti-APRIL were from PeproTech (NJ). Tetramers, HLA-A*0201-GILGFVFTL (SEQ ID NO.: 1) (Flu M1) and HLA-A*0201-ELAGIGILTV (SEQ ID NO.: 2) (Mart-1), were purchased from Beckman Coulter (CA).

Cells and cultures—Monocytes (1×10^6 /ml) from normal donors were cultured in Cellgenics (France) media containing GM-CSF (100 ng/ml) and IL-4 (50 ng/ml) (R&D, CA). For day three and day six, DCs, the same amounts of cytokines were supplemented into the media on day one and day three, respectively. B cells were purified with a negative isolation kit (BD). CD4 and CD8 T cells were purified with magnetic beads coated with anti-CD4 or CD8 (Milteniy, CA). PBMCs were isolated from Buffy coats using Percoll™ gradients (GE Healthcare UK Ltd, Buckinghamshire, UK) by density gradient centrifugation. For DC activation, 1×10^5 DCs were cultured in the mAb-coated 96-well plate for 16-18 h. mAbs (1-2 μ g/well) in carbonate buffer, pH 9.4, were incubated for at least 3 h at 37° C. Culture supernatants were harvested and cytokines/chemokines were measured by Luminex (Biorad, CA). For gene analysis, DCs were cultured in the plates coated with mAbs for 8 h. In some experiments, soluble 50 ng/ml of CD40L (R&D, CA) or 50 nM CpG (InVivogen, CA) was added into the cultures. In the DCs and B cell co-cultures, 5×10^3 DCs resuspended in RPMI 1640 with 10% FCS and antibiotics (Biosource, CA) were first cultured in the plates coated with mAbs for at least 6 h, and then 1×10^5 purified autologous B cells labeled with CFSE (Molecular Probes, OR) were added. In some experiments, DCs were pulsed with 5 moi (multiplicity of infection) of heat-inactivated influenza virus (A/PR/8 H1N1) for 2 h, and then mixed with B cells. For the DCs and T cell co-cultures, 5×10^3 DCs were cultured with 1×10^5 purified autologous CD8 T cells or mixed allogeneic T cells. Allogeneic T cells were pulsed with 1 μ Ci/well ³[H]-thymidine for the final 18 h of incubation, and then cpm were measured by a μ -counter (Wallac, MN). 5×10^5 PBMCs/well were cultured in the plates coated with mAbs. The frequency of Mart-1 and Flu M1 specific CD8 T cells was measured by staining cells with anti-CD8 and tetramers on day ten and day seven of the cultures, respectively. 10 μ M of Mart-1 peptide (ELAGIGILTV) (SEQ ID NO.: 2) and 20 nM of recombinant protein containing Mart-1 peptides (see below) were added to the DC and CD8 T cell cultures. 20 nM purified recombinant Flu M1 protein (see below) was add to the PBMC cultures.

Monoclonal antibodies—Mouse mAbs were generated by conventional technology. Briefly, six-week-old BALB/c mice were immunized i.p. with 20 μ g of receptor ectodomain.hIgG-Fc fusion protein with Ribi adjuvant, then boosts with 20 μ g antigen ten days and fifteen days later. After three months, the mice were boosted again three days prior to taking the spleens. Alternately, mice were injected in the footpad with

1-10 µg antigen in Ribi adjuvant every three to four days over a thirty to forty day period. Three to four days after a final boost, draining lymph nodes were harvested. B cells from spleen or lymph node cells were fused with SP2/O-Ag 14 cells. Hybridoma supernatants were screened to analyze Abs to the receptor ectodomain fusion protein compared to the fusion partner alone, or the receptor ectodomain fused to alkaline phosphatase (44). Positive wells were then screened in FACS using 293F cells transiently transfected with expression plasmids encoding full-length receptor cDNAs. Selected hybridomas were single cell cloned and expanded in CEL-Line flasks (Integra, CA). Hybridoma supernatants were mixed with an equal volume of 1.5 M glycine, 3 M NaCl, 1×PBS, pH 7.8 and tumbled with MabSelect resin. The resin was washed with binding buffer and eluted with 0.1 M glycine, pH 2.7. Following neutralization with 2 M Tris, mAbs were dialyzed versus PBS.

ELISA—Sandwich ELISA was performed to measure total IgM, IgG, and IgA as well as flu-specific immunoglobulins (Igs). Standard human serum (Bethyl) containing known amounts of Igs and human AB serum were used as standard for total Igs and flu-specific Igs, respectively. Flu specific Ab titers, units, in samples were defined as dilution factor of AB serum that shows an identical optical density. The amounts of BAFF and BlyS were measured by ELISA kits (Bender MedSystem, CA).

RNA purification and gene analysis—Total RNA extracted with RNeasy columns (Qiagen), and analyzed with the 2100 Bioanalyser (Agilent). Biotin-labeled cRNA targets were prepared using the Illumina totalprep labeling kit (Ambion) and hybridized to Sentrix Human6 BeadChips (46K transcripts). These microarrays consist of 50mer oligonucleotide probes attached to 3 µm beads which are lodged into microwells etched at the surface of a silicon wafer. After staining with Streptavidin-Cy3, the array surface is imaged using a sub-micron resolution scanner manufactured by Illumina (Beadstation 500x). A gene expression analysis software program, GeneSpring, Version 7.1 (Agilent), was used to perform data analysis. Expression and purification of recombinant Flu M1 and MART-1 proteins—PCR was used to amplify the ORF of Influenza A/Puerto Rico/8/34/Mount Sinai (H1N1) M1 gene while incorporating an Nhe I site distal to the initiator codon and a Not I site distal to the stop codon. The digested fragment was cloned into pET-28b(+) (Novagen), placing the M1 ORF in-frame with a His6 tag, thus encoding His.Flu M1 protein. A pET28b (+) derivative encoding an N-terminal 169 residue cohesin domain from *C. thermocellum* (unpublished) inserted between the Nco I and Nhe I sites expressed Coh.His. For expression of Cohesin-Flex-hMART-1-PeptideA-His, the sequence GACACCACCGAGGCCCGCCACCCCCACCCCCCGTGACCACCCCCACCACCACCGA CCGGAAGGGCACCACCGCCGAGGAGCTGGC-CGGC ATCGGCATCCTGACCGTGATCC TGGGCGGCAAGCGGACCAACAACAGCAC-CCCCACCAAGGGCGAATTCTGCAGATA TCCATCACTGGCGGCCG (SEQ ID NO.: 3) (encoding DTTEAR-HPHPVTTPTTDRKGTAEELAGIGILTV ILGGKRTNNSPTKGEFCRYPSHWR P (SEQ ID NO.: 4)—the marked residues are the immunodominant HLA-A2-restricted peptide and the underlined residues surrounding the peptide are from MART-1) was inserted between the Nhe I and Xho I sites of the above vector. The proteins were expressed in *E. coli* strain BL21 (DE3) (Novagen) or T7 Express (NEB), grown in LB at 37° C. with selection for kanamycin resistance (40 µg/ml) and shaking at 200 rounds/min to mid log phase growth when 120 mg/L IPTG was added. After three hours, the cells were harvested by centrifu-

gation and stored at -80° C. *E. coli* cells from each 1 L fermentation were resuspended in 30 ml ice-cold 50 mM Tris, 1 mM EDTA pH 8.0 (buffer B) with 0.1 ml of protease inhibitor Cocktail II (Calbiochem, CA). The cells were sonicated on ice 2×5 min at setting 18 (Fisher Sonic Dismembrator 60) with a 5 min rest period and then spun at 17,000 r.p.m. (Sorvall SA-600) for 20 min at 4° C. For His.Flu M1 purification the 50 ml cell lysate supernatant fraction was passed through 5 ml Q Sepharose beads and 6.25 ml 160 mM Tris, 40 mM imidazole, 4 M NaCl pH 7.9 was added to the Q Sepharose flow through. This was loaded at 4 ml/min onto a 5 ml HiTrap chelating HP column charged with Ni⁺⁺. The column-bound protein was washed with 20 mM NaPO₄, 300 mM NaCl pH 7.6 (buffer D) followed by another wash with 100 mM H₃COONa pH 4.0. Bound protein was eluted with 100 mM H₃COONa pH 4.0. The peak fractions were pooled and loaded at 4 ml/min onto a 5 ml HiTrap S column equilibrated with 100 mM H₃COONa pH 5.5, and washed with the equilibration buffer followed by elution with a gradient from 0-1 M NaCl in 50 mM NaPO₄ pH 5.5. Peak fractions eluting at about 500 mM NaCl were pooled. For Coh.Flu M1.His purification, cells from 2 L of culture were lysed as above. After centrifugation, 2.5 ml of Triton X114 was added to the supernatant with incubation on ice for 5 min. After further incubation at 25° C. for 5 min, the supernatant was separated from the Triton X114 following centrifugation at 25° C. The extraction was repeated and the supernatant was passed through 5 ml of Q Sepharose beads and 6.25 ml 160 mM Tris, 40 mM imidazole, 4 M NaCl pH 7.9 was added to the Q Sepharose flow through. The protein was then purified by Ni⁺⁺ chelating chromatography as described above and eluted with 0-500 mM imidazole in buffer D.

Only particular anti-DC-ASGPR mAbs have DC activation properties—The invention discloses that DC activation is not a general property of anti-DC-ASGPR antibodies, rather only certain anti-DC-ASGPR mAbs have this function. FIG. 5 shows that only certain mAbs activate DCS through the DC-ASGPR, which must be characterized by screening against actual DCs.

Particular sequences corresponding to the L and H variable regions of anti-DC-ASGPR mAbs—The invention encompasses particular amino acid sequences shown below corresponding to anti-DC-ASGPR monoclonal antibodies that are desirable components (in the context of e.g., humanized recombinant antibodies) of therapeutic or protective products. The following are such sequences in the context of chimeric mouse V region-human C region recombinant antibodies. [mAnti-ASGPR_49C11_7H-LV-hIgG4H-C] is DVQLQESGPDLVKPSQSLSLTCTVTGYSITSGYSW-HWIRQFPGNKLEWMGYILFSGSTNYNPSLKSRSITR-DTSKNQFFLQLNSVTTEDTATYFCARSNYGSFASW-GQGTLVTVSAAKTTGPSVFPLAPCSRSTSESTAALG-CLVKDYFPEPVTVSWNSGALTSQVHTFPAVLQSSGLYSLSSVVTVPSSSLGKTYTCNVDH-KPSNTKVDKRVESKYGPPCPPAPEFEGGPPSVF LFP-PKPKDTLMISRTPEVTCVVVDVSDQEDPE-VQFNWYVDGVEVHNAKTKPREEQFNST YRVVSVLTVLHQDWLNGKEYKCK-VSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEE MTKNQVSLTCLVKGFYPSDIA-VEWESNGQPENNYKTTTPVLDSDGSFF-LYSRLTVDKSR WQEGNVFSCSVMHEALHNHYTQK-SLSLSLGKAS (SEQ ID NO.: 5). The above sequence is a chimera between the H chain V-region of the mAb 49C11

23

(shown underlined) and the C region of hIgG4. [mAnti-ASGPR_49C11_7K-LV-hIgGK-C] is the corresponding L chain chimera QIVLTQSPAIMSASPGE-
 KVTMTCSASSSVSHMHWYQKSGTSPKRWIYDTS-
 RLASGVPARFSGSGSGTSYSLSISSME-
 AEDAATYYCQQWSSHPWSFGGGT-
 KLEIKRTVAAPSVFIFPP SDEQLKSGTASVVCLLNN-
 FYPREAKVQWKVDNALQSGNSQESVTEQDSKDY-
 SLSST LTLSKADYEKHKVY-
 ACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO.: 6). [mAnti-ASGPR_4G2.2_Hv-V-hIgG4H-C] is -
 QIQLVQSGPELKKPGETVKISCKASGY-
 TFTNYGMNWVKQVPGKGLRWMGWMDFTEPT-
 YADDFKGRFAFSLETSASTAYLQINSLKNEDTATYFC-
 ARGGILRLNYFDYWGGTTLTVSSAKTKGPSVFPLA-
 PCSRSTSESTAALGCLVKDYFPEP-
 VTVSWNSGALTSVHTFPA VLQSSGLYSLSSV-
 VTPSSSLGTKTYTCNVDPKPSNTKVD-
 KRVESKYGPPCPPCPAPEFE
 GGPSVFLFPPPKDITLMISRTPEVTCV-
 VVDVSQEDPEVQFNWYVDGVEVHNAKTKPRE
 EQFNSTYRVVSVLTVLHQDWL-
 NGKEYKCKVSNKGLPSSIEK-
 TISKAKGQPREPQVYTLPSQEEMTKNQVSLT-
 CLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDG-
 SFFLYSRLT VDКСRWQEGNVFSCSVMHEALH-
 NHYTQKSLSLGKAS (SEQ ID NO.: 7). [mAnti-AS-
 GPR_4G2.2_Kv-V-hIgGK-C] is -DIQMTQS-
 SSSFSVSLGDRVTITCKASEDIYNRLGWYQQKPGNA-
 PRLISGATSLETGVPSRFSGSGSGKDYALSITSLQTE-
 DLATYYCQQCWTSPTTFGGGTKLEIKRTVAAPSVFI-
 FPP SDEQLKSGTASVVCLLNN-
 FYPREAKVQWKVDNALQSGNSQESVTE-
 QDSKDYSLSS LTLSKADYEKHKVY-
 ACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO.: 8). [mAnti-ASGPR_5F10H-LV-hIgG4H-C] is -
 EVQLQSGPELVKPGASVKMSCKASGYTFDYDM-
 KWVKQSHGKSLEWIGDINPNYGDTFYNQKFEGKA-
 TLTVDKSSRTAYMQLNSLTSEDSAVYYCGRGDYGY-
 FDVWGAGTTVTVSSAKTKGPSVFPLAPCSRSTSESTA-
 ALGCLVKDYFPEPVTVSWNSGALTSVHTFPAV
 LQSSGLYSLSSVTVPSSSLGTKTYTCN-
 VDHPKPSNTKVDKRVESKYGPPCPPCPAPEFEG GPSV-
 FLFPPPKDITLMISRTPEVTCVVVD-
 VSQEDPEVQFNWYVDGVEVHNAKTKPREE
 QFNSTYRVVSVLTVLHQDWLNGKEYKCK-
 VSNKGLPSSIEKTISKAKGQPREPQVYTLPP SQEEMT-
 KNQVSLTCLVKGFYPSDIA-
 VEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTV
 DKSRWQEGNVFSCSVMHEALHNHYTQK-
 SLLSLGKAS (SEQ ID NO.: 9). [mAnti-ASGPR_5F10K-
 LV-hIgGK-C] is -DIVMTQS-
 HKFMSTVGDVRSITCKASQDVGTAVAWYQQKPGQ-
 SPKLLIYWASTRHTGVPDRFTGSGSGTDFTLINNV-
 QSEDLADYFCQQYSSNPYMFGGGTKLEIKRTVAAP-
 SVFI FPPSDEQLKSGTASVVCLLNN-
 FYPREAKVQWKVDNALQSGNSQESVTE-
 QDSKDYSL SSSLTLSKADYEKHKVY-
 ACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO.: 10). [mAnti-ASGPR1H11H-V-hIgG4H-C] is -
 QLQQSGPELVKPGASVKISCKTSGYTFTEYTMHWV-
 RQSHGKSLEWIGGINPINGGPTYNQKFEGKATLTV-
 DKSSSTAYMELRSLTSEDSAVYYCARWDYGSRDVM-
 DYWGQTSVTVSSAKTKGPSVFPLAPCSRSTSESTA-
 ALGCLVKDYFPEPVTVSWNSGALTSVHTFPAV
 LQSSGLYSLSSVTVPSSSLGTKTYTCN-
 VDHPKPSNTKVDKRVESKYGPPCPPCPAPEFEG GPSV-

24

FLFPPPKDITLMISRTPEVTCVVVD-
 VSQEDPEVQFNWYVDGVEVHNAKTKPREE
 QFNSTYRVVSVLTVLHQDWLNGKEYKCK-
 VSNKGLPSSIEKTISKAKGQPREPQVYTLPP SQEEMT-
 5 KNQVSLTCLVKGFYPSDIA-
 VEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTV
 DKSRWQEGNVFSCSVMHEALHNHYTQK-
 SLLSLGKAS (SEQ ID NO.: 11). [mAnti-ASGPR1H11K-
 LV-hIgGK-C] is -NIVMTQSPKSM-
 10 SVGERVTLSCASENVGTYVSWYQQRPEQSPKLLIY-
 GASNRYTGVPDRFTGSGSATDFTLTSSVQAEDLAD-
 YHCGQTYSYIFTFGSGTKLEIKRTVAAPSVFIFP PSD-
 EQLKSGTASVVCLLNN-
 FYPREAKVQWKVDNALQSGNSQESVTE-
 15 QDSKDYSLSS
 LTLSKADYEKHKVY-
 ACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO.: 12). The
 invention envisions these V-region sequences and related
 sequences modified by those well versed in the art to e.g.,
 20 enhance affinity for DC-ASGPR and/or integrated into
 human V-region framework sequences to be engineered into
 expression vectors to direct the expression of protein forms
 that can bind to DC-ASGPR on antigen presenting cells.
 Engineered recombinant anti-DC-ASGPR recombinant
 25 antibody—antigen fusion proteins ((rAb.antigen) are effica-
 cious prototype vaccines in vitro—Expression vectors can be
 constructed with diverse protein coding sequence e.g., fused
 in-frame to the H chain coding sequence. For example, anti-
 gens such as Influenza HAS, Influenza M1, HIV gag, or
 immuno-dominant peptides from cancer antigens, or cytok-
 30 ines, can be expressed subsequently as rAb.antigen or rAb-
 .cytokine fusion proteins, which in the context of this inven-
 tion, can have utility derived from using the anti-DC-ASGPR
 V-region sequence to bring the antigen or cytokine (or toxin)
 35 directly to the surface of the antigen presenting cell bearing
 DC-ASGPR. This permits internalization of e.g., antigen—
 sometimes associated with activation of the receptor and
 ensuing initiation of therapeutic or protective action (e.g., via
 initiation of a potent immune response, or via killing of the
 40 targeted cell. An exemplative prototype vaccine based on this
 concept could use a H chain vector such as [mAnti-ASGPR_5F10H-LV-hIgG4H-C-Flex-FluHA5-1-6xHis] or -EVQ-
 LQQSGPELVKPGASVKMSCKASGYTFDYDMKW-
 VKQSHGKSLEWIGDINPNYGDTFYNQKFEGKATLT-
 45 VDKSSRTAYMQLNSLTSEDSAVYYCGRGDYGYFDV-
 WGAGTTVTVSSAKTKGPSVFPLAPCSRSTSESTAAL-
 GCLVKDYFPEPVTVSWNSGALTSVHTFPAV LQSS-
 GLYSLSSVTVPSSSLGTKTYTCNVDPK-
 PSNTKVDKRVESKYGPPCPPCPAPEFEG GPSVFLFP-
 50 PKPKDITLMISRTPEVTCVVVDVSQEDPEVQFNWYV-
 DGVEVHNAKTKPREE QFNSTYRVVSVLTVLHQDWL-
 NGKEYKCKVSNKGLPSSIEK-
 TISKAKGQPREPQVYTLPP SQEEMTKNQVSLT-
 CLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSD-
 55 GSFFLYSRLTV DKSRWQEGNVFSCSVMHEALH-
 NHYTQKSLSLGKASDTTEPATPTTPVTTDQICIGYH
 ANNSTEQVDTIMEKNVTVTHAQDILE-
 KKHNGKLCDLGDKPLILRDCSVAGWLL GNPMC-
 DEFINVPESYIVEKANPVNDLCYPGD-
 60 FNDYEELKHLISRINHFEKIIPK
 SSWSHEASLGVSACPYQGGKSSFFRN-
 VWLIKKNSTYPTIKRSYNNTNQEDLLVL WGIHHP-
 NDAAEQTKLYQNPTTYISVGTSTLNQR-
 LVPRIATRSKVNQSGRMEFFW
 65 TILKPNDAINFESNGNFIAP-
 EYAYKIVKKGDSTIMKSELEYGNCNTKC-
 QTPMGAINS SMPFHNIHPLTIGECPKYVKS NRLV-

25

LAHHHHHH (SEQ ID NO.: 13). The above sequence corresponds to the chimeric H chain shown already fused via a flexible linker sequence (shown italicized) to HA-1 domain of avian Flu HAS (shown in bold). This can be co-expressed with the corresponding L chain chimeric sequence already shown above. Similarly, the sequence [mAnti-ASGPR_49C11_7H-LV-hIgG4H-C-Dockerin]-

(SEQ ID NO.: 14)
 DVQLQESGPDLVKPSQSLSLTCTVTGYSITSGYSWHWIRQFPGNKLEWVG
 YILFSGSTNYNPSLKSRISTRDTSKNQFFLQNLNSVTTEDTATYFCARSN
 YGSFASWGOGLTVTVSAAKTKGPSVFPLAPCSRSTSESTAALGCLVKDYF
 PEPVTVSWNSGALTSKVHTFPAVLQSSGLYSLSSVTVPSSSLGKTKYTC
 NVDHKPSNTKVDKRVESKYGPPCPAPEFEGGSPVFLFPPKPKDMLMI
 SRTPEVTCVVVDVSDQEDPEVFQFNWYVDGVEVHNAKTKPREEQFNSTYRVV
 SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTIKAKGQPREPQVYTLPP
 SQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGS
 FFLYRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSLSLGLKASNSEPC
 NEVLYGDEVMDGKVNSTDLTLLKRYVLYKAVSTLPSKAEKNAVDNRDGRV
 NSSDVTLISRYLIRVTEKLP

can be used to express via co-transfection of the corresponding L chain sequence already shown above a rAb.Dockerin fusion protein.

FIG. 5 shows that certain anti-DC-ASGPR mAbs can activate DC. GM-CSF/IL-4. DC were incubated for 24 hrs with one of a panel of 12 pure anti-ASGPR mAbs. Cells were then tested for expression of cell surface CD86 (a DC activation marker) and supernatants were assayed for secreted cytokines. Three mAbs (36, 38, 43) from the anti-ASGPR mAb panel activated DC.

FIG. 6 shows that different antigens can be expressed in the context of a DC-ASGPR rAb. Such an anti-DC-ASGPR rAb.Doc protein can be simply mixed with any Cohesin.fusion protein to assemble a stable non-covalent [rAb.Doc:Coh.fusion] complex that functions just as a rAb.fusion protein. FIG.

26

6 shows that such a [rAb.Doc:Coh.fusion] complex can focus antigen to the surface of cells expressing DC-ASGPR. The figure also shows anti-DC-ASGPR.Doc:Coh.Flu M1 complexes deliver Flu M1 to the surface of 293F cells transfected with DC-ASGPR cDNA. 1 µg/ml of anti-DC-ASGPR.Doc rAb (shown as the peak on the right) or control hIgG4.Doc rAb (shown as the peak on the left) were incubated with biotinylated Coh.Flu M1 (2 µg/ml) for 1 hr at R.T. transfected 293F cells were added and incubation continued for 20 min on ice. Cells were then washed and stained with PE-labeled streptavidin. Cells were then analyzed for PE fluorescence.

Anti-DC-ASGPR rAb complexed to Flu M1 via Dockerin: Cohesin interaction targets the antigen to human DCs and results in the expansion of Flu M1-specific CD8+ T cells—the potential utility of anti-DC-ASGPR rAbs as devices to deliver antigen to e.g., DC is shown in the figure below. FIG. 7 shows the dramatic expansion of Flu M1-specific CD8+ cells is highly predictive of potency of such an agent as a vaccine directed to eliciting protective immune responses against Flu M1.

FIGS. 8A-8D demonstrated the cross reactivity of the different antibodies with monkey ASGPR. For pIRES_ASGPR-mon (monkey) was cloned by inserting the PCR product into NheI-NotI sites of pIRES vector. The sequence of final product is base on clone 5S10. Most other clones are either similar to this with one aa difference or identical to this. However, one clone, 5S1, has an A deletion near the 3' end, which generated a shortened and different C' terminus and maybe used as a second variant. To clone the monkey ASGPR, the following oligos were used: DC-ASGPR_MoN: gaattcgtagcCACCATGACATATGAAAACCTCCAAGACTTGGAGAGTGAGGAGAAAGTCCAAGGGG (SEQ ID NO.: 15); and DC-ASGPR_Mo: CGAATTCGCGGCCGCTCAGTGACTCTCCTGGCTGGCCTGGGTCAGACCAGCCTCGC AGACCC (SEQ ID NO.: 16), which is a reverse complement of GGGTCTGCGAGGCTGGTCTGACCCAGGC-CAGCCAGGAGAGTCACTGAGCGGCCGC GAATTCG (SEQ ID NO.: 17). Sequence comparisons indicate the likely regions of overlap and, hence, the cross-reactivity, as is known to those of skill in the art.

The following table demonstrated the binding of the DC-ASGPR 334998 200 ug/ml 12.05.07 cfg#558 anti-Human IgG PE

Glycan number	Glycan name	Avg w/o Max & Min	StDev w/o Max & Min	SEM w/o Max & Min	% CV
82	GalNAcα1-3(Fucα1-2)Galβ1-4GlcNAcβ-Sp8	52930	10265	5132	19
210	Neu5Acα2-3(GalNAcβ1-4)Galβ1-4GlcNAcβ-Sp8	49937	4969	2484	10
86	GalNAcα1-3Galβ-Sp8	49067	4672	2336	10
89	GalNAcα1-3(Fucα1-2)Galβ-Sp8	47375	5453	2726	12
84	GalNAcα1-3(Fucα1-2)Galβ-Sp8	46555	6618	3309	14
209	Neu5Acα2-3(GalNAcβ1-4)Galβ1-4GlcNAcβ-Sp0	46169	2121	1060	5
175	GlcNAcβ1-6GalNAcα-Sp8	44809	1939	969	4
301	GalNAcα1-3(Fucα1-2)Galβ-Sp18	44147	6003	3002	14
211	Neu5Acα2-3(GalNAcβ1-4)Galβ1-4Glcβ-Sp0	43603	3517	1759	8
10	α-GalNAc-Sp8	43514	2476	1238	6
128	Galβ1-3GalNAcβ1-4(Neu5Acα2-3)Galβ1-4Glcβ-Sp0	43152	13339	6669	31
151	Galβ1-4GlcNAcβ1-6GalNAcα-Sp8	42871	2466	1233	6
92	GalNAcβ1-4GlcNAcβ-Sp0	42845	3394	1697	8
93	GalNAcβ1-4GlcNAcβ-Sp8	41764	7340	3670	18
87	GalNAcα1-4(Fucα1-2)Galβ1-4GlcNAcβ-Sp8	41584	2925	1462	7
79	GalNAcα1-3(Fucα1-2)Galβ1-3GlcNAcβ-Sp0	41406	14134	7067	34
20	β-GalNAc-Sp8	40803	2388	1194	6
206	Neu5Acα2-8Neu5Acα2-3(GalNAcβ1-4)Galβ1-4Glcβ-Sp0	38720	2736	1368	7
242	Neu5Acα2-6GalNAcα-Sp8	37500	1934	967	5
91	GalNAcβ1-4(Fucα1-3)GlcNAcβ-Sp0	37286	5046	2523	14

-continued

Glycan number	Glycan name	Avg w/o Max & Min	StDev w/o Max & Min	SEM w/o Max & Min	% CV
204	Neu5Acα2-8Neu5Acα2-8Neu5Acα2-3(GalNAcβ1-4)Galβ1-4Glcβ-Sp0	37237	995	497	3
203	NeuAcα2-8NeuAcα2-8NeuAcα2-8NeuAcα2-3(GalNAcβ1-4)Galβ1-4Glcβ-Sp0	36746	2399	1200	7
243	Neu5Acα2-6GalNAcβ1-4GlcNAcβ-Sp0	36375	1661	830	5
59	Fucα1-2Galβ1-3GalNAcβ1-4(Neu5Acα2-3)Galβ1-4Glcβ-Sp0	35701	6903	3452	19
90	GalNAcβ1-3Galα1-4Galβ1-4GlcNAcβ-Sp0	34350	760	380	2
83	GalNAcα1-3(Fucα1-2)Galβ1-4Glcβ-Sp0	28846	9844	4922	34
302	GalNAcβ1-3Galβ-Sp8	28745	15727	7864	55
300	GalNAcα-Sp15	18125	18847	9424	104
127	Galβ1-3GalNAcβ1-3Galα1-4Galβ1-4Glcβ-Sp0	17999	9798	4899	54
85	GalNAcα1-3GalNAcβ-Sp8	12643	10843	5422	86
173	GlcNAcβ1-4GlcNAcβ1-4GlcNAcβ-Sp8	8673	940	470	11
81	GalNAcα1-3(Fucα1-2)Galβ1-4GlcNAcβ-Sp0	7672	12937	6469	169
30	[3OSO3]Galβ1-4(6OSO3)Glcβ-Sp8	7394	292	146	4
120	Galβ1-3(Galβ1-4GlcNAcβ1-6)GalNAcα-Sp8	5664	1311	655	23
80	GalNAcα1-3(Fucα1-2)Galβ1-4(Fucα1-3)GlcNAcβ-Sp0	5444	907	454	17
147	Galβ1-4GlcNAcβ1-3Galβ1-4GlcNAcβ-Sp0	4927	410	205	8
29	[3OSO3]Galβ1-4(6OSO3)Glcβ-Sp0	4871	908	454	19
101	Galα1-3GalNAcα-Sp8	4815	3163	1581	66
214	Neu5Acα2-3GalNAcα-Sp8	4109	569	284	14
287	[3OSO3][4OSO3]Galβ1-4GlcNAcβ-SpSp0	3959	1646	823	42
40	[4OSO3]Galβ1-4GlcNAcβ-Sp8	3848	673	337	17
45	[6OSO3]Galβ1-4[6OSO3]Glcβ-Sp8	3790	993	497	26
166	GlcNAcβ1-3Galβ1-4GlcNAcβ1-3Galβ1-4GlcNAcβ-Sp0	3720	435	218	12
227	Neu5Acα2-3Galβ1-4[6OSO3]GlcNAcβ-Sp8	3576	793	397	22
218	NeuAcα2-3Galβ1-3(Fucα1-4)GlcNAcβ1-3Galβ1-4(Fucα1-3)GlcNAcβ Sp0	3360	104	52	3
240	Neu5Acα2-3Galβ1-4Glcβ-Sp8	3313	976	488	29
149	Galβ1-4GlcNAcβ1-3Galβ1-4Glcβ-Sp8	3233	263	132	8
244	Neu5Acα2-6Galβ1-4[6OSO3]GlcNAcβ-Sp8	3195	757	379	24
270	Fucα1-2Galβ1-4[6OSO3]GlcNAc-Sp8	3161	2563	1282	81
42	[6OSO3]Galβ1-4Glcβ-Sp0	3084	529	264	17
271	Fucα1-2[6OSO3]Galβ1-4[6OSO3]Glc-Sp0	3063	377	188	12
172	(GlcNAcβ1-4)5β-Sp8	3032	1058	529	35
47	[6OSO3]GlcNAcβ-Sp8	3008	159	80	5
143	Neu5Acα2-3Galβ1-4GlcNAcβ1-2Manα1-3(Neu5Acα2-3Galβ1-4GlcNAcβ1-2Manα1-6)Manβ1-4GlcNAcβ1-4GlcNAcβ-Sp12	3008	309	155	10
265	[3OSO3]Galβ1-4(Fucα1-3)(6OSO3)Glc-Sp0	2995	1841	921	61
139	Galβ1-4[6OSO3]Glcβ-Sp0	2988	1070	535	36
27	[3OSO3][6OSO3]Galβ1-4GlcNAcβ-Sp0	2930	317	158	11
273	Fucα1-2-Galβ1-4[6OSO3]Glc-Sp0	2919	495	247	17
319	Neu5Acα2-6Galβ1-4GlcNAcβ1-2Manα1-3(Galβ1-4GlcNAcβ1-2Manα1-6)Manβ1-4GlcNAcβ1-4GlcNAcβ-Sp12	2730	993	497	36
35	[3OSO3]Galβ1-4[6OSO3]GlcNAcβ-Sp8	2722	516	258	19
28	[3OSO3]Galβ1-4Glcβ-Sp8	2674	197	98	7
38	[3OSO3]Galβ-Sp8	2652	1680	840	63
253	Neu5Acα2-8Neu5Acα2-3Galβ1-4Glcβ-Sp0	2631	1136	568	43
289	6-H2PO3Glcβ-Sp10	2611	674	337	26
26	[3OSO3][6OSO3]Galβ1-4[6OSO3]GlcNAcβ-Sp0	2550	153	76	6
266	[3OSO3]Galβ1-4(Fucα1-3)Glc-Sp0	2529	444	222	18
54	Neu5Acα2-6Galβ1-4GlcNAcβ1-2Manα1-3(Neu5Acα2-6Galβ1-4GlcNAcβ1-2Manα1-6)Manβ1-4GlcNAcβ1-4GlcNAcβ-Sp8	2476	300	150	12
303	GlcAβ1-3GlcNAcβ-Sp8	2463	130	65	5
32	[3OSO3]Galβ1-3GalNAcα-Sp8	2461	622	311	25
53	Neu5Acα2-6Galβ1-4GlcNAcβ1-2Manα1-3(Neu5Acα2-6Galβ1-4GlcNAcβ1-2Manα1-6)Manβ1-4GlcNAcβ1-4GlcNAcβ-Sp13	2455	283	142	12
181	Glcβ1-6Glcβ-Sp8	2455	154	77	6
267	[3OSO3]Galβ1-4[Fucα1-3][6OSO3]GlcNAc-Sp8	2447	1065	532	44
293	Galβ1-3(Neu5Acα2-3Galβ1-4(Fucα1-3)GlcNAcβ1-6)GalNAc-Sp14	2359	648	324	27
202	Neu5Acα2-3Galβ1-3GalNAcα-Sp8	2349	928	464	40
163	GlcNAcβ1-3Galβ1-3GalNAcα-Sp8	2347	375	188	16
1	Neu5Acα2-8Neu5Acα-Sp8	2339	1539	769	66
31	[3OSO3]Galβ1-3(Fucα1-4)GlcNAcβ-Sp8	2332	319	160	14
230	Neu5Acα2-3Galβ1-4(Fucα1-3)GlcNAcβ-Sp0	2306	164	82	7
286	[3OSO3]Galβ1-4[6OSO3]GlcNAcβ-Sp0	2290	472	236	21
318	Neu5Acα2-3Galβ1-4GlcNAcβ1-2Manα1-3(Neu5Acα2-6Galβ1-4GlcNAcβ1-2Manα1-6)Manβ1-4GlcNAcβ1-4GlcNAcβ-Sp12	2262	246	123	11
199	Neu5Acα2-6Galβ1-4GlcNAcβ1-2Manα1-3(Neu5Acα2-3Galβ1-4GlcNAcβ1-2Manα1-6)Manβ1-4GlcNAcβ1-4GlcNAcβ-Sp12	2217	138	69	6

-continued

Glycan number	Glycan name	Avg w/o Max & Min	StDev w/o Max & Min	SEM w/o Max & Min	% CV
39	[4OSO3][6OSO3]Galβ1-4GlcNAcβ-Sp0	2215	619	310	28
77	Fuca1-4GlcNAcβ-Sp8	2207	83	42	4
285	Neu5Acα2-3Galβ1-4GlcNAcβ1-3Galβ1-3GlcNAcβ-Sp0	2193	1679	839	77
262	Neu5Gcα2-6GalNAcα-Sp0	2192	734	367	33
216	Neu5Acα2-3Galβ1-3(6OSO3)GlcNAc-Sp8	2163	1062	531	49
43	[6OSO3]Galβ1-4Glcβ-Sp8	2149	700	350	33
297	Galβ1-4GlcNAcβ1-3(GlcNAcβ1-6)Galβ1-4GlcNAc-Sp0	2141	983	491	46
224	NeuAcα2-3Galβ1-3GlcNAcβ1-3Galβ1-4GlcNAcβ-Sp0	2133	1208	604	57
3	Neu5Acα2-8Neu5Acα2-8Neu5Acβ-Sp8	2117	611	306	29
171	(GlcNAcβ1-4)6β-Sp8	2112	302	151	14
316	Neu5Acα2-3Galβ1-3(Neu5Acα2-6)GalNAc-Sp14	2105	1171	585	56
15	α-Neu5Ac-Sp11	2099	250	125	12
52	Galβ1-4GlcNAcβ1-2Manα1-3(Galβ1-4GlcNAcβ1-2Manα1-6)Manβ1-4GlcNAcβ1-4GlcNAcβ-Sp13	2092	429	215	21
268	[3OSO3]Galβ1-4[Fuca1-3]GlcNAc-Sp0	2085	955	477	46
313	Manα1-2Manα1-2Manα1-3(Manα1-2Manα1-6(Manα1-3)Manα1-6)Manα-Sp9	2020	812	406	40
225	Neu5Acα2-3Galβ1-3GlcNAcβ-Sp0	2019	1052	526	52
36	[3OSO3]Galβ1-4GlcNAcβ-Sp0	2012	389	194	19
263	Neu5Gcα2-6Galβ1-4GlcNAcβ-Sp0	1999	664	332	33
141	Galβ1-4GalNAcα1-3(Fuca1-2)Galβ1-4GlcNAcβ-Sp8	1968	772	386	39
274	Galβ1-3(Fuca1-4)GlcNAcβ1-3Galβ1-3(Fuca1-4)GlcNAcβ-Sp0	1961	78	39	4
275	Galβ1-3-(Galβ1-4GlcNAcβ1-6)GalNAc-Sp14	1953	409	205	21
7	α-D-Gal-Sp8	1925	636	318	33
41	6-H2PO3Manα-Sp8	1919	223	111	12
247	Neu5Acα2-6Galβ1-4GlcNAcβ1-3Galβ1-4(Fuca1-3)GlcNAcβ1-3Galβ1-4(Fuca1-3)GlcNAcβ-Sp0	1914	169	85	9
311	Manα1-6Manβ-Sp10	1906	522	261	27
205	Neu5Acα2-8Neu5Acα2-8Neu5Acα2-3Galβ1-4Glcβ-Sp0	1902	222	111	12
280	Galβ1-4[Fuca1-3][6OSO3]GlcNAc-Sp0	1881	982	491	52
152	Galβ1-4GlcNAcβ-Sp0	1868	924	462	49
113	Galα1-6Glcβ-Sp8	1864	321	161	17
115	Galβ1-3(Fuca1-4)GlcNAcβ1-3Galβ1-4(Fuca1-3)GlcNAcβ-Sp0	1855	338	169	18
251	Neu5Acα2-6Galβ-Sp8	1842	316	158	17
116	Galβ1-3(Fuca1-4)GlcNAcβ1-3Galβ1-4GlcNAcβ-Sp0	1836	798	399	43
194	Manα1-2Manα1-2Manα1-3(Manα1-2Manα1-3(Manα1-2Manα1-6)Manα1-6)Manβ1-4GlcNAcβ1-4GlcNAcβ-Sp12	1829	176	88	10
33	[3OSO3]Galβ1-3GlcNAcβ-Sp8	1812	889	445	49
272	Fuca1-2-(6OSO3)-Galβ1-4Glc-Sp0	1805	86	43	5
207	Neu5Acα2-8Neu5Acα2-8Neu5Acα-Sp8	1804	454	227	25
74	Fuca1-2Galβ-Sp8	1796	648	324	36
213	Neu5Acα2-3(Neu5Acα2-6)GalNAcα-Sp8	1768	312	156	18
234	Neu5Acα2-3Galβ1-4GlcNAcβ1-3Galβ1-4(Fuca1-3)GlcNAc-Sp0	1767	178	89	10
50	Manα1-3(Manα1-6)Manβ1-4GlcNAcβ1-4GlcNAcβ-Sp13	1759	553	277	31
111	Galα1-4Galβ1-4Glcβ-Sp0	1740	635	318	36
291	Galα1-3GalNAcα-Sp16	1738	1090	545	63
296	Galβ1-4GlcNAcβ1-3(Galβ1-4GlcNAcβ1-6)Galβ1-4GlcNAc-Sp0	1726	850	425	49
154	Galβ1-4Glcβ-Sp0	1725	457	229	27
56	Fuca1-2Galβ1-3GalNAcβ1-3Galα1-4Galβ1-4Glcβ-Sp9	1719	384	192	22
66	Fuca1-2Galβ1-4(Fuca1-3)GlcNAcβ1-3Galβ1-4(Fuca1-3)GlcNAcβ1-3Galβ1-4(Fuca1-3)GlcNAcβ-Sp0	1703	224	112	13
299	Galβ1-4GlcNAcβ1-6Galβ1-4GlcNAcβ-Sp0	1658	820	410	49
44	[6OSO3]Galβ1-4GlcNAcβ-Sp8	1632	242	121	15
237	Neu5Acα2-3Galβ1-4GlcNAcβ-Sp8	1632	1049	524	64
233	Neu5Acα2-3Galβ1-4(Fuca1-3)GlcNAcβ1-3Galβ1-4GlcNAcβ-Sp8	1620	862	431	53
192	Manα1-6(Manα1-2Manα1-3)Manα1-6(Manα2Manα1-3)Manβ1-4GlcNAcβ1-4GlcNAcβ-Sp12	1608	903	452	56
64	Fuca1-2Galβ1-3GlcNAcβ-Sp8	1602	625	313	39
62	Fuca1-2Galβ1-3GlcNAcβ1-3Galβ1-4Glcβ-Sp8	1580	417	208	26
148	Galβ1-4GlcNAcβ1-3Galβ1-4Glcβ-Sp0	1568	617	308	39
295	Galβ1-4GlcNAcβ1-2Manα1-3(Neu5Acα2-6Galβ1-4GlcNAcβ1-2Manα1-6)Manβ1-4GlcNAcβ1-4GlcNAcβ-Sp12	1556	190	95	12
137	Galβ1-4(Fuca1-3)GlcNAcβ1-4Galβ1-4(Fuca1-3)GlcNAcβ-Sp0	1552	1313	656	85
17	β-D-Gal-Sp8	1544	871	435	56
168	GlcNAcβ1-4MDPLys	1542	345	172	22
254	Neu5Acβ2-6GalNAcα-Sp8	1541	688	344	45
231	Neu5Acα2-3Galβ1-4(Fuca1-3)GlcNAcβ-Sp8	1534	257	129	17
125	Galβ1-3GalNAcα-Sp8	1483	1025	512	69
269	Fuca1-2[6OSO3]Galβ1-4GlcNAc-Sp0	1473	191	96	13
182	G-ol-Sp8	1471	264	132	18
37	[3OSO3]Galβ1-4GlcNAcβ-Sp8	1462	1187	593	81
229	Neu5Acα2-3Galβ1-4(Fuca1-3)GlcNAcβ1-3Galβ1-4(Fuca1-3)GlcNAcβ1-3Galβ1-4(Fuca1-3)GlcNAcβ-Sp0	1451	333	167	23
315	Neu5Acα2-3Galβ1-3(Neu5Acα2-3Galβ1-4GlcNAcβ1-6)GalNAc-Sp14	1448	1476	738	102

-continued

Glycan number	Glycan name	Avg w/o Max & Min	StDev w/o Max & Min	SEM w/o Max & Min	% CV
65	Fuc α 1-2Gal β 1-4(Fuc α 1-3)GlcNAc β 1-3Gal β 1-4(Fuc α 1-3)GlcNAc β -Sp0	1442	748	374	52
164	GlcNAc β 1-3Gal β 1-4GlcNAc β -Sp0	1436	1332	666	93
305	GlcNAc β 1-2Man α 1-3(GlcNAc β 1-2Man α 1-6)Man β 1-4GlcNAc β 1-4GlcNAc β -Sp12	1428	288	144	20
304	GlcNAc β 1-2Man α 1-3(Neu5Ac α 2-6Gal β 1-4GlcNAc β 1-2Man α 1-6)Man β 1-4GlcNAc β 1-4GlcNAc β -Sp12	1428	499	249	35
145	Gal β 1-4GlcNAc β 1-3Gal β 1-4(Fuc α 1-3)GlcNAc β 1-3Gal β 1-4(Fuc α 1-3)GlcNAc β -Sp0	1422	323	162	23
117	Gal β 1-3(Fuc α 1-4)GlcNAc-Sp0	1407	681	341	48
193	Man α 1-2Man α 1-6(Man α 1-3)Man α 1-6(Man α 2Man α 2Man α 1-3)Man β 1-4GlcNAc β 1-4GlcNAc β -Sp12	1404	285	142	20
19	β -D-Man-Sp8	1389	635	317	46
176	GlcNAc β 1-6Gal β 1-4GlcNAc β -Sp8	1383	1000	500	72
232	Neu5Ac α 2-3Gal β 1-4(Fuc α 1-3)GlcNAc β 1-3Gal β -Sp8	1355	374	187	28
219	Neu5Ac α 2-3Gal β 1-3(Neu5Ac α 2-3Gal β 1-4)GlcNAc β -Sp8	1350	753	377	56
123	Gal β 1-3(Neu5Ac β 2-6)GalNAc α -Sp8	1350	852	426	63
276	Gal β 1-3(GlcNAc β 1-6)GalNAc-Sp14	1345	353	176	26
208	Neu5Ac α 2-3(6-O-Su)Gal β 1-4(Fuc α 1-3)GlcNAc β -Sp8	1341	642	321	48
55	Fuc α 1-2Gal β 1-3GalNAc β 1-3Gal α -Sp9	1331	466	233	35
257	Neu5Gc α 2-3Gal β 1-3(Fuc α 1-4)GlcNAc β -Sp0	1315	108	54	8
201	Fuc α 1-3(Gal β 1-4)GlcNAc β 1-2Man α 1-3(Fuc α 1-3(Gal β 1-4)GlcNAc β 1-2Man α 1-6)Man β 1-4GlcNAc β 1-4GlcNAc β -Sp20	1294	289	144	22
97	Gal α 1-3(Fuc α 1-2)Gal β 1-4GlcNAc-Sp0	1282	583	291	45
150	Gal β 1-4GlcNAc β 1-6(Gal β 1-3)GalNAc α -Sp8	1265	778	389	62
60	Fuc α 1-2Gal β 1-3GalNAc β 1-4(Neu5Ac α 2-3)Gal β 1-4Glc β -Sp9	1261	738	369	59
317	Neu5Ac α 2-3Gal β 1-3GalNAc-Sp14	1239	780	390	63
23	β -GlcN(Gc)-Sp8	1219	436	218	36
279	Gal β 1-3GlcNAc β 1-3Gal β 1-3GlcNAc β -Sp0	1219	570	285	47
190	Man α 1-2Man α 1-3(Man α 1-2Man α 1-6)Man α -Sp9	1217	1305	653	107
178	Glc α 1-4Glc α -Sp8	1216	560	280	46
146	Gal β 1-4GlcNAc β 1-3Gal β 1-4GlcNAc β 1-3Gal β 1-4GlcNAc β -Sp0	1211	1315	658	109
292	Gal β 1-3GalNAc α -Sp16	1198	370	185	31
221	Neu5Ac α 2-3Gal β 1-3(Neu5Ac α 2-6)GalNAc α -Sp8	1194	238	119	20
99	Gal α 1-3(Fuc α 1-2)Gal β -Sp8	1189	767	383	64
309	HOOC(CH ₃)CH-3-O-GlcNAc β 1-4GlcNAc β -Sp10	1186	1108	554	93
248	Neu5Ac α 2-6Gal β 1-4GlcNAc β 1-3Gal β 1-4GlcNAc β -Sp0	1181	334	167	28
107	Gal α 1-3Gal β -Sp8	1148	688	344	60
236	Neu5Ac α 2-3Gal β 1-4GlcNAc β -Sp0	1148	441	220	38
320	Neu5Ac α 2-6Gal β 1-4GlcNAc β 1-2Man α 1-3(GlcNAc β 1-2Man α 1-6)Man β 1-4GlcNAc β 1-4GlcNAc β -Sp12	1142	55	27	5
197	Man α 1-6(Man α 1-3)Man α 1-6(Man α 2Man α 1-3)Man β 1-4GlcNAc β 1-4GlcNAc β -Sp12	1134	200	100	18
185	GlcA β 1-3Gal β -Sp8	1133	470	235	42
34	[3OSO ₃]Gal β 1-4(Fuc α 1-3)GlcNAc β -Sp8	1117	980	490	88
109	Gal α 1-4Gal β 1-4GlcNAc β -Sp0	1094	499	250	46
235	Neu5Ac α 2-3Gal β 1-4GlcNAc β 1-3Gal β 1-4GlcNAc β 1-3Gal β 1-4GlcNAc β -Sp0	1092	1077	539	99
228	Neu5Ac α 2-3Gal β 1-4(Fuc α 1-3)(6OSO ₃)GlcNAc β -Sp8	1090	771	385	71
184	GlcA β -Sp8	1072	476	238	44
282	Gal β 1-4(Fuc α 1-3)GlcNAc β 1-3Gal β 1-3(Fuc α 1-4)GlcNAc β -Sp0	1062	239	120	23
2	Neu5Ac α 2-8Neu5Ac β -Sp17	1060	84	42	8
174	GlcNAc β 1-6(Gal β 1-3)GalNAc α -Sp8	1039	913	456	88
261	Neu5Gc α 2-3Gal β 1-4Glc β -Sp0	1034	440	220	43
18	β -D-Glc-Sp8	1024	335	167	33
217	Neu5Ac α 2-3Gal β 1-3(Fuc α 1-4)GlcNAc β -Sp8	1023	646	323	63
260	Neu5Gc α 2-3Gal β 1-4GlcNAc β -Sp0	1020	208	104	20
104	Gal α 1-3Gal β 1-3GlcNAc β -Sp0	1017	297	149	29
245	Neu5Ac α 2-6Gal β 1-4GlcNAc β -Sp0	1010	394	197	39
14	α -Neu5Ac-Sp8	998	1046	523	105
283	Gal β 1-4GlcNAc β 1-3Gal β 1-3GlcNAc β -Sp0	978	514	257	53
156	GlcNAc α 1-3Gal β 1-4GlcNAc β -Sp8	969	276	138	29
310	Man α 1-3(Man α 1-6)Man β 1-4GlcNAc β 1-4GlcNAc β -Sp12	965	238	119	25
183	GlcA α -Sp8	960	463	232	48
138	Gal β 1-4(Fuc α 1-3)GlcNAc β 1-4Gal β 1-4(Fuc α 1-3)GlcNAc β 1-4Gal β 1-4(Fuc α 1-3)GlcNAc β -Sp0	948	595	297	63
96	Gal α 1-3(Fuc α 1-2)Gal β 1-4(Fuc α 1-3)GlcNAc β -Sp0	948	260	130	27
6	Neu5Ac α 2-6Gal β 1-4GlcNAc β 1-2Man α 1-3(Neu5Ac α 2-6Gal β 1-4GlcNAc β 1-2Man α 1-6)Man β 1-4GlcNAc β 1-4GlcNAc β -Sp12	943	351	176	37
306	GlcNAc β 1-3Man-Sp10	938	153	77	16
121	Gal β 1-3(GlcNAc β 1-6)GalNAc α -Sp8	936	748	374	80
258	Neu5Gc α 2-3Gal β 1-3GlcNAc β -Sp0	932	375	188	40
246	Neu5Ac α 2-6Gal β 1-4GlcNAc β -Sp8	931	635	317	68
200	Man β 1-4GlcNAc β -Sp0	920	322	161	35
78	Fuc β 1-3GlcNAc β -Sp8	911	464	232	51
94	Gal α 1-2Gal β -Sp8	911	393	197	43

-continued

Glycan number	Glycan name	Avg w/o Max & Min	StDev w/o Max & Min	SEM w/o Max & Min	% CV
256	Galβ1-4GlcNAcβ1-2Manα1-3(Neu5Acα2-6Galβ1-4GlcNAcβ1-2Manα1-6)Manβ1-4GlcNAcβ1-4GlcNAcβ-Sp21	909	428	214	47
95	Galα1-3(Fuca1-2)Galβ1-3GlcNAcβ-Sp0	908	245	123	27
8	α-D-Glc-Sp8	904	417	209	46
103	Galα1-3Galβ1-4(Fuca1-3)GlcNAcβ-Sp8	893	445	222	50
118	Galβ1-3(Fuca1-4)GlcNAc-Sp8	890	624	312	70
9	α-D-Man-Sp8	881	403	201	46
16	β-Neu5Ac-Sp8	876	935	468	107
119	Galβ1-3(Fuca1-4)GlcNAcβ-Sp8	872	283	141	32
278	Galβ1-3GalNAc-Sp14	851	144	72	17
187	KDNα2-3Galβ1-3GlcNAcβ-Sp0	839	386	193	46
69	Fuca1-2Galβ1-4GlcNAcβ1-3Galβ1-4GlcNAc-Sp0	837	328	164	39
76	Fuca1-3GlcNAcβ-Sp8	836	276	138	33
108	Galα1-4(Fuca1-2)Galβ1-4GlcNAcβ-Sp8	819	58	29	7
212	NeuAcα2-3(NeuAcα2-3Galβ1-3GalNAcβ1-4)Galβ1-4Glcβ-Sp0	818	1442	721	176
132	Galβ1-3GlcNAcβ1-3Galβ1-4Glcβ-Sp10	816	353	176	43
105	Galα1-3Galβ1-4GlcNAcβ-Sp8	806	184	92	23
308	GlcNAcβ1-4GlcNAcβ-Sp12	796	360	180	45
160	GlcNAcβ1-3(GlcNAcβ1-6)Galβ1-4GlcNAcβ-Sp8	794	416	208	52
284	Neu5Acα2-3Galβ1-3GlcNAcβ1-3Galβ1-3GlcNAcβ-Sp0	777	491	245	63
188	KDNα2-3Galβ1-4GlcNAcβ-Sp0	774	320	160	41
215	Neu5Acα2-3GalNAcβ1-4GlcNAcβ-Sp0	762	252	126	33
294	Galβ1-3Galβ1-4GlcNAcβ-Sp8	746	255	128	34
196	Manα1-3(Manα1-2Manα1-2Manα1-6)Manα-Sp9	744	177	88	24
189	Manα1-2Manα1-2Manα1-3Manα-Sp9	743	207	103	28
25	GlcNAcβ1-3(GlcNAcβ1-4)(GlcNAcβ1-6)GlcNAc-Sp8	735	270	135	37
131	Galβ1-3GlcNAcβ1-3Galβ1-4GlcNAcβ-Sp0	728	290	145	40
277	Galβ1-3(Neu5Acα2-3Galβ1-4GlcNAcβ1-6)GalNAc-Sp14	722	324	162	45
136	Galβ1-4(Fuca1-3)GlcNAcβ-Sp8	718	93	46	13
70	Fuca1-2Galβ1-4GlcNAcβ1-3Galβ1-4GlcNAcβ1-3Galβ1-4GlcNAcβ-Sp0	713	861	430	121
110	Galα1-4Galβ1-4GlcNAcβ-Sp8	712	183	91	26
129	Galβ1-3GalNAcβ1-4Galβ1-4Glcβ-Sp8	702	224	112	32
71	Fuca1-2Galβ1-4GlcNAcβ-Sp0	686	160	80	23
169	GlcNAcβ1-4(GlcNAcβ1-6)GalNAcα-Sp8	686	229	115	33
122	Galβ1-3(Neu5Acα2-6)GalNAcα-Sp8	679	157	79	23
106	Galα1-3Galβ1-4Glcβ-Sp0	678	137	69	20
255	Neu5Acβ2-6Galβ1-4GlcNAcβ-Sp8	671	153	76	23
130	Galβ1-3Galβ-Sp8	668	285	143	43
144	Galβ1-4GlcNAcβ1-3GalNAcα-Sp8	663	227	113	34
13	α-L-Rha-Sp8	662	245	123	37
22	β-GlcNAc-Sp8	655	313	157	48
72	Fuca1-2Galβ1-4GlcNAcβ-Sp8	646	95	47	15
157	GlcNAcα1-6Galβ1-4GlcNAcβ-Sp8	644	323	162	50
307	GlcNAcβ1-4GlcNAcβ-Sp10	640	336	168	53
180	Glcβ1-4Glcβ-Sp8	608	316	158	52
191	Manα1-2Manα1-3Manα-Sp9	607	104	52	17
134	Galβ1-3GlcNAcβ-Sp8	603	103	51	17
21	β-GlcNAc-Sp0	595	285	142	48
24	(Galβ1-4GlcNAcβ)2-3,6-GalNAcα-Sp8	590	240	120	41
223	NeuAcα2-3Galβ1-3GalNAcβ1-3Galα1-4Galβ1-4Glcβ-Sp0	580	191	95	33
162	GlcNAcβ1-3Galβ-Sp8	577	435	217	75
135	Galβ1-4(Fuca1-3)GlcNAcβ-Sp0	561	139	70	25
249	Neu5Acα2-6Galβ1-4Glcβ-Sp0	560	377	189	67
48	9NAcNeu5Acα-Sp8	556	470	235	85
158	GlcNAcβ1-2Galβ1-3GalNAcα-Sp8	550	417	208	76
264	Neu5Gcα-Sp8	550	305	152	55
46	NeuAcα2-3[6OSO3]Galβ1-4GlcNAcβ-Sp8	545	363	182	67
68	Fuca1-2Galβ1-4(Fuca1-3)GlcNAcβ-Sp8	541	208	104	38
222	Neu5Acα2-3Galβ-Sp8	526	277	139	53
298	Galβ1-4GlcNAcα1-6Galβ1-4GlcNAcβ-Sp0	494	335	167	68
98	Galα1-3(Fuca1-2)Galβ1-4Glcβ-Sp0	482	112	56	23
312	Manα1-6(Manα1-3)Manα1-6(Manα1-3)Manβ-Sp10	453	292	146	64
133	Galβ1-3GlcNAcβ-Sp0	452	165	82	36
57	Fuca1-2Galβ1-3(Fucβ1-4)GlcNAcβ-Sp8	450	268	134	60
114	Galβ1-2Galβ-Sp8	449	324	162	72
198	Manα1-6(Manα1-3)Manα1-6(Manα1-3)Manβ1-4GlcNAcβ1-4GlcNAcβ-Sp12	448	204	102	45
161	GlcNAcβ1-3GalNAcα-Sp8	442	156	78	35
281	Galβ1-4[Fuca1-3][6OSO3]Glc-Sp0	439	144	72	33
259	Neu5Gcα2-3Galβ1-4(Fuca1-3)GlcNAcβ-Sp0	433	357	179	83
67	Fuca1-2Galβ1-4(Fuca1-3)GlcNAcβ-Sp0	420	94	47	22
12	α-L-Fuc-Sp9	410	303	151	74
159	GlcNAcβ1-3(GlcNAcβ1-6)GalNAcα-Sp8	407	88	44	22
75	Fuca1-3GlcNAcβ-Sp8	399	182	91	46
239	Neu5Acα2-3Galβ1-4Glcβ-Sp0	395	156	78	39
290	Galα1-3(Fuca1-2)Galβ-Sp18	389	246	123	63

-continued

Glycan number	Glycan name	Avg w/o Max & Min	StDev w/o Max & Min	SEM w/o Max & Min	% CV
11	α -L-Fuc-Sp8	387	231	115	60
51	GlcNAc β 1-2Man α 1-3(GlcNAc β 1-2Man α 1-6)Man β 1-4GlcNAc β 1-4GlcNAc β -Sp13	383	164	82	43
5	Gal β 1-3GlcNAc β 1-2Man α 1-3(Gal β 1-3GlcNAc β 1-2Man α 1-6)Man β 1-4GlcNAc β 1-4GlcNAc β -Sp19	381	529	265	139
63	Fuc α 1-2Gal β 1-3GlcNAc β -Sp0	362	187	93	52
241	Gal β 1-4GlcNAc β 1-2Man α 1-3(Fuc α 1-3(Gal β 1-4)GlcNAc β 1-2Man α 1-6)Man β 1-4GlcNAc β 1-4GlcNAc β -Sp20	352	68	34	19
155	Gal β 1-4Glc β -Sp8	315	105	53	33
126	Gal β 1-3GalNAc β -Sp8	288	265	132	92
195	Man α 1-3(Man α 1-6)Man α -Sp9	269	92	46	34
88	GalNAc β 1-3GalNAc α -Sp8	262	107	54	41
252	Neu5Ac α 2-8Neu5Ac α -Sp8	260	214	107	82
167	GlcNAc β 1-3Gal β 1-4Glc β p-Sp0	257	129	64	50
140	Gal β 1-4[6OSO3]Glc β -Sp8	256	345	172	135
177	Glc α 1-4Glc β -Sp8	246	113	57	46
179	Glc α 1-6Glc α 1-6Glc β -Sp8	225	380	190	168
314	Man α 1-2Man α 1-2Man α 1-3(Man α 1-2Man α 1-6(Man α 1-2Man α 1-3)Man α 1-6)Man α -Sp9	221	329	165	149
238	Neu5Ac α 2-3Gal β 1-4GlcNAc β 1-3Gal β 1-4GlcNAc β -Sp0	212	200	100	94
220	Neu5Ac α 2-3Gal β 1-3[6OSO3]GalNAc α -Sp8	210	153	77	73
142	Gal β 1-4GalNAc β 1-3(Fuc α 1-2)Gal β 1-4GlcNAc β -Sp8	204	126	63	62
61	Fuc α 1-2Gal β 1-3GlcNAc β 1-3Gal β 1-4Glc β -Sp10	196	67	34	34
102	Gal α 1-3GalNAc β -Sp8	188	198	99	105
170	GlcNAc β 1-4Gal β 1-4GlcNAc β -Sp8	184	127	64	69
124	Gal β 1-3(Neu5Ac α 2-6)GlcNAc β 1-4Gal β 1-4Glc β -Sp10	173	146	73	84
100	Gal α 1-3(Gal α 1-4)Gal β 1-4GlcNAc β -Sp8	168	112	56	66
186	GlcA β 1-6Gal β -Sp8	158	171	86	108
4	Neu5Gc β 2-6Gal β 1-4GlcNAc-Sp8	152	96	48	63
73	Fuc α 1-2Gal β 1-4Glc β -Sp0	148	205	103	139
49	9NAcNeu5Ac α 2-6Gal β 1-4GlcNAc β -Sp8	146	159	79	108
58	Fuc α 1-2Gal β 1-3GalNAc α -Sp8	136	171	86	126
250	Neu5Ac α 2-6Gal β 1-4Glc β -Sp8	122	144	72	119
112	Gal α 1-4GlcNAc β -Sp8	115	82	41	72
165	GlcNAc β 1-3Gal β 1-4GlcNAc β -Sp8	84	68	34	81
226	Neu5Ac α 2-3Gal β 1-3GlcNAc β -Sp8	76	85	42	112
288	[6OSO3]Gal β 1-4[6OSO3]GlcNAc β -Sp0	72	130	65	180
153	Gal β 1-4GlcNAc β -Sp8	48	58	29	120

It is contemplated that any embodiment discussed in this specification can be implemented with respect to any method, kit, reagent, or composition of the invention, and vice versa. Furthermore, compositions of the invention can be used to achieve methods of the invention.

It will be understood that particular embodiments described herein are shown by way of illustration and not as limitations of the invention. The principal features of this invention can be employed in various embodiments without departing from the scope of the invention. Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, numerous equivalents to the specific procedures described herein. Such equivalents are considered to be within the scope of this invention and are covered by the claims.

All publications and patent applications mentioned in the specification are indicative of the level of skill of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

The use of the word “a” or “an” when used in conjunction with the term “comprising” in the claims and/or the specification may mean “one,” but it is also consistent with the meaning of “one or more,” “at least one,” and “one or more than one.” The use of the term “or” in the claims is used to mean “and/or” unless explicitly indicated to refer to alternatives only or the alternatives are mutually exclusive, although the disclosure supports a definition that refers to only alter-

natives and “and/or.” Throughout this application, the term “about” is used to indicate that a value includes the inherent variation of error for the device, the method being employed to determine the value, or the variation that exists among the study subjects.

As used in this specification and claim(s), the words “comprising” (and any form of comprising, such as “comprise” and “comprises”), “having” (and any form of having, such as “have” and “has”), “including” (and any form of including, such as “includes” and “include”) or “containing” (and any form of containing, such as “contains” and “contain”) are inclusive or open-ended and do not exclude additional, unrecited elements or method steps.

The term “or combinations thereof” as used herein refers to all permutations and combinations of the listed items preceding the term. For example, “A, B, C, or combinations thereof” is intended to include at least one of: A, B, C, AB, AC, BC, or ABC, and if order is important in a particular context, also BA, CA, CB, CBA, BCA, ACB, BAC, or CAB. Continuing with this example, expressly included are combinations that contain repeats of one or more item or term, such as BB, AAA, MB, BBC, AAABCCCC, CBBAAA, CABABB, and so forth. The skilled artisan will understand that typically there is no limit on the number of items or terms in any combination, unless otherwise apparent from the context.

All of the compositions and/or methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have

been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the compositions and/or methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

REFERENCES

1. C. G. Figdor, Y. van Kooyk, G. J. Adema, *Nat Rev Immunol* 2, 77 (February, 2002).
2. E. Pyz, A. S. Marshall, S. Gordon, G. D. Brown, *Ann Med* 38, 242 (2006).
3. G. D. Brown, *Nat Rev Immunol* 6, 33 (January, 2006).
4. T. B. Geijtenbeek et al., *Nat Immunol* 1, 353 (October, 2000).
5. T. B. Geijtenbeek et al., *Cell* 100, 575 (Mar. 3, 2000).
6. C. F. d'Ostiani et al., *J Exp Med* 191, 1661 (May 15, 2000).
7. C. Fradin, D. Poulain, T. Jouault, *Infect Immun* 68, 4391 (August, 2000).
8. A. Cambi et al., *Eur J Immunol* 33, 532 (February, 2003).
9. M. G. Netea, J. W. Meer, I. Verschueren, B. J. Kullberg, *Eur J Immunol* 32, 1455 (May, 2002).
10. S. J. Lee et al., *Science* 295, 1898 (Mar. 8, 2002).
11. N. Maeda et al., *J Biol Chem* 278, 5513 (Feb. 21, 2003).
12. L. Tailleux et al., *J Exp Med* 197, 121 (Jan. 6, 2003).
13. T. B. Geijtenbeek et al., *J Exp Med* 197, 7 (Jan. 6, 2003).
14. A. M. Cooper et al., *J Immunol* 168, 1322 (Feb. 1, 2002).
15. J. Banchereau et al., *Annu Rev Immunol* 18, 767 (2000).
16. P. Jeannin et al., *Immunity* 22, 551 (May, 2005).
17. B. N. Gantner, R. M. Simmons, S. J. Canavera, S. Akira, D. M. Underhill, *J Exp Med* 197, 1107 (May 5, 2003).
18. H. Moriwaki et al., *FEBS Lett* 440, 29 (Nov. 27, 1998).
19. M. Wykes, G. MacPherson, *Immunology* 100, 1 (May, 2000).
20. M. Balazs, F. Martin, T. Zhou, J. Kearney, *Immunity* 17, 341 (September, 2002).
21. T. Kikuchi, S. Worgall, R. Singh, M. A. Moore, R. G. Crystal, *Nat Med* 6, 1154 (October, 2000).
22. B. Dubois et al., *J Leukoc Biol* 66, 224 (August, 1999).
23. H. Qi, J. G. Egen, A. Y. Huang, R. N. Germain, *Science* 312, 1672 (Jun. 16, 2006).
24. A. Bergtold, D. D. Desai, A. Gavhane, R. Clynes, *Immunity* 23, 503 (November, 2005).
25. C. R. Ruprecht, A. Lanzavecchia, *Eur J Immunol* 36, 810 (April, 2006).
- 10 26. N. L. Bernasconi, E. Traggiai, A. Lanzavecchia, *Science* 298, 2199 (Dec. 13, 2002).
27. P. A. Moore et al., *Science* 285, 260 (Jul. 9, 1999).
28. J. A. Gross et al., *Nature* 404, 995 (Apr. 27, 2000).
29. A. Craxton, D. Magaletti, E. J. Ryan, E. A. Clark, *Blood* 101, 4464 (Jun. 1, 2003).
30. J. MacLennan, C. Vinuesa, *Immunity* 17, 235 (September, 2002).
31. J. Banchereau, R. M. Steinman, *Nature* 392, 245 (Mar. 19, 1998).
- 20 32. J. Banchereau, P. de Paoli, A. Valle, E. Garcia, F. Rousset, *Science* 251, 70 (Jan. 4, 1991).
33. B. Beutler et al., *Annu Rev Immunol* 24, 353 (2006).
34. K. Hayashida, N. Kume, M. Minami, T. Kita, *FEBS Lett* 511, 133 (Jan. 30, 2002).
- 25 35. M. Kakutani, T. Masaki, T. Sawamura, *Proc Natl Acad Sci USA* 97, 360 (Jan. 4, 2000).
36. M. Colonna, J. Samaridis, L. Angman, *Eur J Immunol* 30, 697 (February, 2000).
- 30 37. J. Valladeau et al., *J Immunol* 167, 5767 (Nov. 15, 2001).
38. G. Jego et al., *Immunity* 19, 225 (August, 2003).
39. M. N. Wykes, L. Beattie, G. G. Macpherson, D. N. Hart, *Immunology* 113, 318 (November, 2004).
40. Y. Delneste et al., *Immunity* 17, 353 (September, 2002).
- 35 41. Y. Wang et al., *Immunity* 15, 971 (December, 2001).
42. A. Asea et al., *Nat Med* 6, 435 (April, 2000).
43. H. Bausinger et al., *Eur J Immunol* 32, 3708 (December, 2002).
44. E. E. Bates et al., *J Immunol* 163, 1973 (Aug. 15, 1999).

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 17

<210> SEQ ID NO 1
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically Synthesized Peptide

<400> SEQUENCE: 1

Gly Ile Leu Gly Phe Val Phe Thr Leu
 1 5

<210> SEQ ID NO 2
 <211> LENGTH: 10
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically Synthesized Peptide

<400> SEQUENCE: 2

Glu Leu Ala Gly Ile Gly Ile Leu Thr Val
 1 5 10

-continued

<210> SEQ ID NO 3
 <211> LENGTH: 186
 <212> TYPE: DNA
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically Synthesized Oligonucleotide

 <400> SEQUENCE: 3

 gacaccaccg aggcccgcca cccccacccc cccgtgacca cccccaccac caccgaccgg 60
 aagggcacca ccgccgagga gctggccggc atcgcatcc tgaccgtgat cctggggcggc 120
 aagcggacca acaacagcac ccccaccaag ggccaattct gcagatatcc atcacactgg 180
 cggccg 186

<210> SEQ ID NO 4
 <211> LENGTH: 61
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically Synthesized Peptide

 <400> SEQUENCE: 4

 Asp Thr Thr Glu Ala Arg His Pro His Pro Pro Val Thr Thr Pro Thr
 1 5 10 15
 Thr Asp Arg Lys Gly Thr Thr Ala Glu Glu Leu Ala Gly Ile Gly Ile
 20 25 30
 Leu Thr Val Ile Leu Gly Gly Lys Arg Thr Asn Asn Ser Thr Pro Thr
 35 40 45
 Lys Gly Glu Phe Cys Arg Tyr Pro Ser His Trp Arg Pro
 50 55 60

<210> SEQ ID NO 5
 <211> LENGTH: 446
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically Synthesized Peptide

 <400> SEQUENCE: 5

 Asp Val Gln Leu Gln Glu Ser Gly Pro Asp Leu Val Lys Pro Ser Gln
 1 5 10 15
 Ser Leu Ser Leu Thr Cys Thr Val Thr Gly Tyr Ser Ile Thr Ser Gly
 20 25 30
 Tyr Ser Trp His Trp Ile Arg Gln Phe Pro Gly Asn Lys Leu Glu Trp
 35 40 45
 Met Gly Tyr Ile Leu Phe Ser Gly Ser Thr Asn Tyr Asn Pro Ser Leu
 50 55 60
 Lys Ser Arg Ile Ser Ile Thr Arg Asp Thr Ser Lys Asn Gln Phe Phe
 65 70 75 80
 Leu Gln Leu Asn Ser Val Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys
 85 90 95
 Ala Arg Ser Asn Tyr Gly Ser Phe Ala Ser Trp Gly Gln Gly Thr Leu
 100 105 110
 Val Thr Val Ser Ala Ala Lys Thr Thr Gly Pro Ser Val Phe Pro Leu
 115 120 125
 Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys
 130 135 140
 Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser
 145 150 155 160

-continued

Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser
 165 170 175
 Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser
 180 185 190
 Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn
 195 200 205
 Thr Lys Val Asp Lys Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro
 210 215 220
 Pro Cys Pro Ala Pro Glu Phe Glu Gly Gly Pro Ser Val Phe Leu Phe
 225 230 235 240
 Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val
 245 250 255
 Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe
 260 265 270
 Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro
 275 280 285
 Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr
 290 295 300
 Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val
 305 310 315 320
 Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala
 325 330 335
 Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln
 340 345 350
 Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly
 355 360 365
 Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro
 370 375 380
 Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser
 385 390 395 400
 Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu
 405 410 415
 Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His
 420 425 430
 Tyr Thr Gln Lys Ser Leu Ser Leu Ser Leu Gly Lys Ala Ser
 435 440 445

<210> SEQ ID NO 6
 <211> LENGTH: 213
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically Synthesized Peptide

<400> SEQUENCE: 6

Gln Ile Val Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Pro Gly
 1 5 10 15
 Glu Lys Val Thr Met Thr Cys Ser Ala Ser Ser Ser Val Ser His Met
 20 25 30
 His Trp Tyr Gln Gln Lys Ser Gly Thr Ser Pro Lys Arg Trp Ile Tyr
 35 40 45
 Asp Thr Ser Arg Leu Ala Ser Gly Val Pro Ala Arg Phe Ser Gly Ser
 50 55 60
 Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile Ser Ser Met Glu Ala Glu
 65 70 75 80

-continued

Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser His Pro Trp Ser
85 90 95

Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala Pro
100 105 110

Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly Thr
115 120 125

Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala Lys
130 135 140

Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln Glu
145 150 155 160

Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser Ser
165 170 175

Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr Ala
180 185 190

Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser Phe
195 200 205

Asn Arg Gly Glu Cys
210

<210> SEQ ID NO 7
 <211> LENGTH: 449
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically Synthesized Peptide

<400> SEQUENCE: 7

Gln Ile Gln Leu Val Gln Ser Gly Pro Glu Leu Lys Lys Pro Gly Glu
1 5 10 15

Thr Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
20 25 30

Gly Met Asn Trp Val Lys Gln Val Pro Gly Lys Gly Leu Arg Trp Met
35 40 45

Gly Trp Met Asp Thr Phe Thr Gly Glu Pro Thr Tyr Ala Asp Asp Phe
50 55 60

Lys Gly Arg Phe Ala Phe Ser Leu Glu Thr Ser Ala Ser Thr Ala Tyr
65 70 75 80

Leu Gln Ile Asn Ser Leu Lys Asn Glu Asp Thr Ala Thr Tyr Phe Cys
85 90 95

Ala Arg Gly Gly Ile Leu Arg Leu Asn Tyr Phe Asp Tyr Trp Gly Gln
100 105 110

Gly Thr Thr Leu Thr Val Ser Ser Ala Lys Thr Lys Gly Pro Ser Val
115 120 125

Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala
130 135 140

Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser
145 150 155 160

Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val
165 170 175

Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro
180 185 190

Ser Ser Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp His Lys
195 200 205

Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Ser Lys Tyr Gly Pro
210 215 220

Pro Cys Pro Pro Cys Pro Ala Pro Glu Phe Glu Gly Gly Pro Ser Val

-continued

225				230						235				240	
Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr
				245					250					255	
Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	Gln	Glu	Asp	Pro	Glu
			260					265					270		
Val	Gln	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys
		275					280					285			
Thr	Lys	Pro	Arg	Glu	Glu	Gln	Phe	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser
	290					295					300				
Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys
305					310					315					320
Cys	Lys	Val	Ser	Asn	Lys	Gly	Leu	Pro	Ser	Ser	Ile	Glu	Lys	Thr	Ile
				325					330					335	
Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro
			340					345					350		
Pro	Ser	Gln	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu
		355					360					365			
Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn
370						375					380				
Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser
385				390						395					400
Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Arg	Leu	Thr	Val	Asp	Lys	Ser	Arg
			405						410					415	
Trp	Gln	Glu	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu
			420					425					430		
His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Leu	Gly	Lys	Ala
		435					440					445			

Ser

<210> SEQ ID NO 8
 <211> LENGTH: 214
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically Synthesized Peptide

<400> SEQUENCE: 8

Asp	Ile	Gln	Met	Thr	Gln	Ser	Ser	Ser	Ser	Phe	Ser	Val	Ser	Leu	Gly
1				5					10					15	
Asp	Arg	Val	Thr	Ile	Thr	Cys	Lys	Ala	Ser	Glu	Asp	Ile	Tyr	Asn	Arg
			20					25					30		
Leu	Gly	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Asn	Ala	Pro	Arg	Leu	Leu	Ile
		35					40					45			
Ser	Gly	Ala	Thr	Ser	Leu	Glu	Thr	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly
	50					55					60				
Ser	Gly	Ser	Gly	Lys	Asp	Tyr	Ala	Leu	Ser	Ile	Thr	Ser	Leu	Gln	Thr
65				70						75				80	
Glu	Asp	Leu	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	Cys	Trp	Thr	Ser	Pro	Tyr
				85					90					95	
Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Arg	Thr	Val	Ala	Ala
			100					105					110		
Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	Asp	Glu	Gln	Leu	Lys	Ser	Gly
		115					120					125			
Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	Asn	Asn	Phe	Tyr	Pro	Arg	Glu	Ala
	130					135					140				

-continued

Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
 145 150 155 160

Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
 165 170 175

Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
 180 185 190

Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
 195 200 205

Phe Asn Arg Gly Glu Cys
 210

<210> SEQ ID NO 9
 <211> LENGTH: 445
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically Synthesized Peptide

<400> SEQUENCE: 9

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
 1 5 10 15

Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30

Tyr Met Lys Trp Val Lys Gln Ser His Gly Lys Ser Leu Glu Trp Ile
 35 40 45

Gly Asp Ile Asn Pro Asn Tyr Gly Asp Thr Phe Tyr Asn Gln Lys Phe
 50 55 60

Glu Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Arg Thr Ala Tyr
 65 70 75 80

Met Gln Leu Asn Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95

Gly Arg Gly Asp Tyr Gly Tyr Phe Asp Val Trp Gly Ala Gly Thr Thr
 100 105 110

Val Thr Val Ser Ser Ala Lys Thr Lys Gly Pro Ser Val Phe Pro Leu
 115 120 125

Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys
 130 135 140

Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser
 145 150 155 160

Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser
 165 170 175

Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser
 180 185 190

Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn
 195 200 205

Thr Lys Val Asp Lys Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro
 210 215 220

Pro Cys Pro Ala Pro Glu Phe Glu Gly Gly Pro Ser Val Phe Leu Phe
 225 230 235 240

Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val
 245 250 255

Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe
 260 265 270

Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro
 275 280 285

Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr

-continued

290	295	300
Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val 305	310	315
Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala 325	330	335
Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln 340	345	350
Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly 355	360	365
Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro 370	375	380
Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser 385	390	395
Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu 405	410	415
Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His 420	425	430
Tyr Thr Gln Lys Ser Leu Leu Ser Leu Gly Lys Ala Ser 435	440	445

<210> SEQ ID NO 10

<211> LENGTH: 214

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Chemically Synthesized Peptide

<400> SEQUENCE: 10

Asp Ile Val Met Thr Gln Ser His Lys Phe Met Ser Thr Ser Val Gly 1	5	10	15
Asp Arg Val Ser Ile Thr Cys Lys Ala Ser Gln Asp Val Gly Thr Ala 20	25	30	
Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile 35	40	45	
Tyr Trp Ala Ser Thr Arg His Thr Gly Val Pro Asp Arg Phe Thr Gly 50	55	60	
Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Asn Asn Val Gln Ser 65	70	75	80
Glu Asp Leu Ala Asp Tyr Phe Cys Gln Gln Tyr Ser Ser Asn Pro Tyr 85	90	95	
Met Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala 100	105	110	
Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly 115	120	125	
Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala 130	135	140	
Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln 145	150	155	160
Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser 165	170	175	
Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr 180	185	190	
Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser 195	200	205	
Phe Asn Arg Gly Glu Cys 210			

-continued

<210> SEQ ID NO 11
 <211> LENGTH: 447
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically Synthesized Peptide

 <400> SEQUENCE: 11

 Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala Ser Val
 1 5 10 15

 Lys Ile Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Glu Tyr Thr Met
 20 25 30

 His Trp Val Arg Gln Ser His Gly Lys Ser Leu Glu Trp Ile Gly Gly
 35 40 45

 Ile Asn Pro Ile Asn Gly Gly Pro Thr Tyr Asn Gln Lys Phe Lys Gly
 50 55 60

 Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr Met Glu
 65 70 75 80

 Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys Ala Arg
 85 90 95

 Trp Asp Tyr Gly Ser Arg Asp Val Met Asp Tyr Trp Gly Gln Gly Thr
 100 105 110

 Ser Val Thr Val Ser Ser Ala Lys Thr Lys Gly Pro Ser Val Phe Pro
 115 120 125

 Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly
 130 135 140

 Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn
 145 150 155 160

 Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln
 165 170 175

 Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser
 180 185 190

 Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser
 195 200 205

 Asn Thr Lys Val Asp Lys Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys
 210 215 220

 Pro Pro Cys Pro Ala Pro Glu Phe Glu Gly Gly Pro Ser Val Phe Leu
 225 230 235 240

 Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu
 245 250 255

 Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln
 260 265 270

 Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys
 275 280 285

 Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu
 290 295 300

 Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys
 305 310 315 320

 Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys
 325 330 335

 Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser
 340 345 350

 Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys
 355 360 365

-continued

Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln
 370 375 380

Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly
 385 390 395 400

Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln
 405 410 415

Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn
 420 425 430

His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Leu Gly Lys Ala Ser
 435 440 445

<210> SEQ ID NO 12
 <211> LENGTH: 214
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically Synthesized Peptide

<400> SEQUENCE: 12

Asn Ile Val Met Thr Gln Ser Pro Lys Ser Met Ser Met Ser Val Gly
 1 5 10 15

Glu Arg Val Thr Leu Ser Cys Lys Ala Ser Glu Asn Val Gly Thr Tyr
 20 25 30

Val Ser Trp Tyr Gln Gln Arg Pro Glu Gln Ser Pro Lys Leu Leu Ile
 35 40 45

Tyr Gly Ala Ser Asn Arg Tyr Thr Gly Val Pro Asp Arg Phe Thr Gly
 50 55 60

Ser Gly Ser Ala Thr Asp Phe Thr Leu Thr Ile Ser Ser Val Gln Ala
 65 70 75 80

Glu Asp Leu Ala Asp Tyr His Cys Gly Gln Thr Tyr Ser Tyr Ile Phe
 85 90 95

Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys Arg Thr Val Ala Ala
 100 105 110

Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
 115 120 125

Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
 130 135 140

Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
 145 150 155 160

Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
 165 170 175

Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
 180 185 190

Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
 195 200 205

Phe Asn Arg Gly Glu Cys
 210

<210> SEQ ID NO 13
 <211> LENGTH: 780
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically Synthesized Peptide

<400> SEQUENCE: 13

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
 1 5 10 15

-continued

Thr Glu Pro Ala Thr Pro Thr Thr Pro Val Thr Thr Asp Gln Ile Cys
 450 455 460
 Ile Gly Tyr His Ala Asn Asn Ser Thr Glu Gln Val Asp Thr Ile Met
 465 470 475 480
 Glu Lys Asn Val Thr Val Thr His Ala Gln Asp Ile Leu Glu Lys Lys
 485 490 495
 His Asn Gly Lys Leu Cys Asp Leu Asp Gly Val Lys Pro Leu Ile Leu
 500 505 510
 Arg Asp Cys Ser Val Ala Gly Trp Leu Leu Gly Asn Pro Met Cys Asp
 515 520 525
 Glu Phe Ile Asn Val Pro Glu Trp Ser Tyr Ile Val Glu Lys Ala Asn
 530 535 540
 Pro Val Asn Asp Leu Cys Tyr Pro Gly Asp Phe Asn Asp Tyr Glu Glu
 545 550 555 560
 Leu Lys His Leu Leu Ser Arg Ile Asn His Phe Glu Lys Ile Gln Ile
 565 570 575
 Ile Pro Lys Ser Ser Trp Ser Ser His Glu Ala Ser Leu Gly Val Ser
 580 585 590
 Ser Ala Cys Pro Tyr Gln Gly Lys Ser Ser Phe Phe Arg Asn Val Val
 595 600 605
 Trp Leu Ile Lys Lys Asn Ser Thr Tyr Pro Thr Ile Lys Arg Ser Tyr
 610 615 620
 Asn Asn Thr Asn Gln Glu Asp Leu Leu Val Leu Trp Gly Ile His His
 625 630 635 640
 Pro Asn Asp Ala Ala Glu Gln Thr Lys Leu Tyr Gln Asn Pro Thr Thr
 645 650 655
 Tyr Ile Ser Val Gly Thr Ser Thr Leu Asn Gln Arg Leu Val Pro Arg
 660 665 670
 Ile Ala Thr Arg Ser Lys Val Asn Gly Gln Ser Gly Arg Met Glu Phe
 675 680 685
 Phe Trp Thr Ile Leu Lys Pro Asn Asp Ala Ile Asn Phe Glu Ser Asn
 690 695 700
 Gly Asn Phe Ile Ala Pro Glu Tyr Ala Tyr Lys Ile Val Lys Lys Gly
 705 710 715 720
 Asp Ser Thr Ile Met Lys Ser Glu Leu Glu Tyr Gly Asn Cys Asn Thr
 725 730 735
 Lys Cys Gln Thr Pro Met Gly Ala Ile Asn Ser Ser Met Pro Phe His
 740 745 750
 Asn Ile His Pro Leu Thr Ile Gly Glu Cys Pro Lys Tyr Val Lys Ser
 755 760 765
 Asn Arg Leu Val Leu Ala His His His His His His
 770 775 780

<210> SEQ ID NO 14

<211> LENGTH: 521

<212> TYPE: PRT

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Chemically Synthesized Peptide

<400> SEQUENCE: 14

Asp Val Gln Leu Gln Glu Ser Gly Pro Asp Leu Val Lys Pro Ser Gln
 1 5 10 15
 Ser Leu Ser Leu Thr Cys Thr Val Thr Gly Tyr Ser Ile Thr Ser Gly
 20 25 30

-continued

Tyr Ser Trp His Trp Ile Arg Gln Phe Pro Gly Asn Lys Leu Glu Trp
 35 40 45
 Met Gly Tyr Ile Leu Phe Ser Gly Ser Thr Asn Tyr Asn Pro Ser Leu
 50 55 60
 Lys Ser Arg Ile Ser Ile Thr Arg Asp Thr Ser Lys Asn Gln Phe Phe
 65 70 75 80
 Leu Gln Leu Asn Ser Val Thr Thr Glu Asp Thr Ala Thr Tyr Phe Cys
 85 90 95
 Ala Arg Ser Asn Tyr Gly Ser Phe Ala Ser Trp Gly Gln Gly Thr Leu
 100 105 110
 Val Thr Val Ser Ala Ala Lys Thr Lys Gly Pro Ser Val Phe Pro Leu
 115 120 125
 Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys
 130 135 140
 Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser
 145 150 155 160
 Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser
 165 170 175
 Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser
 180 185 190
 Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn
 195 200 205
 Thr Lys Val Asp Lys Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro
 210 215 220
 Pro Cys Pro Ala Pro Glu Phe Glu Gly Gly Pro Ser Val Phe Leu Phe
 225 230 235 240
 Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val
 245 250 255
 Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe
 260 265 270
 Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro
 275 280 285
 Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr
 290 295 300
 Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val
 305 310 315 320
 Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala
 325 330 335
 Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln
 340 345 350
 Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly
 355 360 365
 Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro
 370 375 380
 Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser
 385 390 395 400
 Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu
 405 410 415
 Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His
 420 425 430

8. The method of claim 1, wherein the antibody or fragment thereof comprises an immunoglobulin light chain selected from the group consisting of SEQ ID NO:6, 8, 10, or 12.

9. A method for increasing the effectiveness of dendritic cells comprising:

isolating dendritic cells from an individual;

exposing the dendritic cells to an effective amount of a complex of an antigen and a DC-ASGPR-specific antibody or DC-ASGPR-specific fragment thereof, wherein the antibody or fragment thereof comprises an immunoglobulin heavy chain selected from the group consisting of SEQ ID NO: 5, 7, 9, or 11, and wherein the antibody or fragment thereof targets a human DC-ASGPR; and reintroducing the cells into the individual.

* * * * *

5

10

15