



US008735125B2

(12) **United States Patent**
Dicosimo et al.

(10) **Patent No.:** **US 8,735,125 B2**
(45) **Date of Patent:** **May 27, 2014**

(54) **PERHYDROLASE VARIANT PROVIDING
IMPROVED SPECIFIC ACTIVITY**

(71) Applicant: **E. I. du Pont de Nemours and
Company**, Wilmington, DE (US)

(72) Inventors: **Robert Dicosimo**, Chadds Ford, PA
(US); **Mark Scott Payne**, Wilmington,
DE (US); **John Edward Gavagan**,
Wilmington, DE (US)

(73) Assignee: **E. I. du Pont de Nemours and
Company**, Wilmington, DE (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 81 days.

(21) Appl. No.: **13/657,601**

(22) Filed: **Oct. 22, 2012**

(65) **Prior Publication Data**

US 2013/0102669 A1 Apr. 25, 2013

Related U.S. Application Data

(60) Provisional application No. 61/551,233, filed on Oct.
25, 2011.

(51) **Int. Cl.**
C12N 9/18 (2006.01)
A01N 37/16 (2006.01)
C12N 15/63 (2006.01)

(52) **U.S. Cl.**
USPC **435/197**; 435/196; 435/132; 435/188;
424/76.1; 530/350

(58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,974,082 A 8/1976 Weyn
4,585,150 A 4/1986 Beacham et al.
5,116,575 A 5/1992 Badertscher et al.
5,296,161 A 3/1994 Wiersema et al.
5,338,676 A 8/1994 Mitsushima et al.
5,364,554 A 11/1994 Stanislawski et al.
5,398,846 A 3/1995 Corba et al.
5,528,152 A 6/1996 Hinoshita et al.
5,552,018 A 9/1996 Devenyns
5,624,634 A 4/1997 Brougham et al.
5,683,724 A 11/1997 Hei et al.
5,932,532 A 8/1999 Ghosh et al.
6,183,807 B1 2/2001 Gutzmann et al.

6,210,639 B1 4/2001 Vlass et al.
6,319,888 B2 11/2001 Wei et al.
6,391,840 B1 5/2002 Thompson et al.
6,518,307 B2 2/2003 McKenzie et al.
6,545,047 B2 4/2003 Gutzmann et al.
6,995,125 B2 2/2006 Dasque et al.
7,550,420 B2 6/2009 DiCosimo et al.
7,723,083 B2 5/2010 DiCosimo et al.
7,910,347 B1 3/2011 DiCosimo et al.
7,923,233 B1 4/2011 DiCosimo et al.
7,927,854 B1 4/2011 DiCosimo et al.
7,932,072 B1 4/2011 DiCosimo et al.
7,951,566 B2 5/2011 DiCosimo et al.
7,960,528 B1 6/2011 DiCosimo et al.
7,964,378 B2 6/2011 DiCosimo et al.
2003/0026846 A1 2/2003 Hei et al.
2005/0008526 A1 1/2005 Bianchetti et al.
2005/0139608 A1 6/2005 Muehlhausen et al.
2008/0176299 A1 7/2008 DiCosimo et al.
2010/0041752 A1 2/2010 DiCosimo et al.
2010/0086510 A1 4/2010 DiCosimo et al.
2010/0086621 A1 4/2010 DiCosimo et al.
2010/0087529 A1 4/2010 DiCosimo et al.
2011/0236335 A1 9/2011 DiCosimo et al.
2011/0236336 A1 9/2011 DiCosimo et al.
2011/0236337 A1 9/2011 DiCosimo et al.
2011/0236338 A1 9/2011 DiCosimo et al.
2011/0236339 A1 9/2011 DiCosimo et al.

FOREIGN PATENT DOCUMENTS

EP 0807156 B1 11/1997
EP 1040222 B1 2/2008
WO WO 0061713 A1 10/2000
WO WO 2007070609 6/2007

OTHER PUBLICATIONS

Gilbert et al., Recent Advances in Carbohydrate Bioengineering, The
Royal Society of Chemistry, Cambridge, pp. 3-12 (1999).
Mitsushima et al Gene Cloning, Nucleotide Sequence, and Expres-
sion., Appl. Env. Microbiol. 61(6):2224-2229 (1995).
Vincent et al., Multifunctional Xylooligosaccharide/Cephalosporin
C Deacetylase . . . , J. Mol. Biol., 330:593-606 (2003).

Primary Examiner — Kagne H Gebreyesus

(57) **ABSTRACT**

An acetyl xylan esterase variant having perhydrolytic activity
is provided for producing peroxy-carboxylic acids from car-
boxylic acid esters and a source of peroxygen. More specifi-
cally, a *Thermotoga maritima* acetyl xylan esterase gene was
modified using error-prone PCR and site-directed mutagen-
esis to create an enzyme catalyst characterized by an increase
in specific activity. The variant acetyl xylan esterase may be
used to produce peroxy-carboxylic acids suitable for use in a
variety of applications such as cleaning, disinfecting, sanitiz-
ing, bleaching, wood pulp processing, and paper pulp pro-
cessing applications.

17 Claims, No Drawings

1

PERHYDROLASE VARIANT PROVIDING
IMPROVED SPECIFIC ACTIVITYCROSS-REFERENCE TO RELATED
APPLICATION

This application claims the benefit of U.S. Provisional Application No. 61/551,233 filed Oct. 25, 2011, which is incorporated by reference in its entirety.

TECHNICAL FIELD

The invention relates to the field of peroxycarboxylic acid biosynthesis and enzyme catalysis. More specifically, an enzyme catalyst comprising a variant enzyme having perhydrolytic activity is provided having an increase in specific activity. Methods of using the present enzyme catalyst to produce peroxycarboxylic acids are also provided.

BACKGROUND

Peroxycarboxylic acid compositions can be effective antimicrobial agents. Methods of using peroxycarboxylic acids to clean, disinfect, and/or sanitize hard surfaces, textiles, meat products, living plant tissues, and medical devices against undesirable microbial growth have been described (U.S. Pat. No. 6,545,047; U.S. Pat. No. 6,183,807; U.S. Pat. No. 6,518,307; U.S. Patent Application Publication No. 2003-0026846; and U.S. Pat. No. 5,683,724). Peroxycarboxylic acids have also been used in various bleaching applications including, but not limited to, wood pulp bleaching/delignification and laundry care applications (European Patent 1040222B1; U.S. Pat. No. 5,552,018; U.S. Pat. No. 3,974,082; U.S. Pat. No. 5,296,161; and U.S. Pat. No. 5,364,554). The desired efficacious concentration of peroxycarboxylic acid may vary according to the product application (for example, ca. 500 ppm to 1000 ppm for medical instrument disinfection, ca. 30 ppm to 80 ppm for laundry bleaching or disinfection applications) in 1 min to 5 min reaction time at neutral pH.

Enzymes structurally classified as members of family 7 of the carbohydrate esterases (CE-7) have been employed as perhydrolases to catalyze the reaction of hydrogen peroxide (or alternative peroxide reagent) with alkyl esters of carboxylic acids in water at a basic to acidic pH range (from ca. pH 11.5 to ca. pH 5) to produce an efficacious concentration of a peroxycarboxylic acid for such applications as disinfection (such as medical instruments, hard surfaces, textiles), bleaching (such as wood pulp or paper pulp processing/delignification, textile bleaching and laundry care applications), and other laundry care applications such as destaining, deodorizing, and sanitization (U.S. Pat. Nos. 7,964,378; 7,951,566; and 7,723,083 and Published U.S. Patent Application Nos. 2008-0176299, and 2010-0041752 to DiCosimo et al.). The CE-7 enzymes have been found to have high specific activity for perhydrolysis of esters, particularly acetyl esters of alcohols, diols, glycerols and phenols, and acetyl esters of mono-, di- and polysaccharides.

Published U.S. Patent Application No. 2010-0087529 to DiCosimo et al. describes several variant CE-7 perhydrolases derived from several *Thermotoga* sp. having higher perhydrolytic specific activity and/or improved selectivity for perhydrolysis when used to prepare peroxycarboxylic acid from carboxylic acid esters. Two of the variants described in Published U.S. Patent Application No. 2010-0087529, *Thermotoga maritima* C277S and *Thermotoga maritima* C277T, exhibited a significant improvement in specific activity relative to the *T. maritima* wild-type enzyme.

2

Thermotoga maritima variants having higher peracid stability were also reported by DiCosimo et al. in U.S. Pat. Nos. 7,927,854; 7,923,233; 7,932,072; 7,910,347, and 7,960,528. Each variant was characterized as having an increased peracetic acid formation to peracetic acid hydrolysis ratio (PAAF:PAAH) when compared to the *T. maritima* wild-type perhydrolase or the *T. maritima* C277S variant perhydrolase.

Several variants of the *Thermotoga maritima* C277S perhydrolase have been identified which have higher specific activity for the perhydrolysis of esters when compared to the specific activity of the *Thermotoga maritima* C277S (U.S. Patent Application Nos. 2011-0236335, 2011-0236336, 2011-0236337, 2011-0236338, and 2011-0236339 to DiCosimo et al.). However, there remains a need to identify additional variants having an increase in perhydrolytic specific activity.

The problem to be solved is to provide an enzyme catalyst comprising a CE-7 perhydrolase having higher specific activity for the perhydrolysis of esters when compared to the specific activity of the *Thermotoga maritima* C277T perhydrolase.

SUMMARY

Nucleic acid molecules encoding the *Thermotoga maritima* acetyl xylan esterase variant C277T were mutated to create libraries of variant enzymes having perhydrolytic activity. Several perhydrolase variants were identified having an increase in specific activity when compared to the parent enzyme from which they were derived (i.e., the *Thermotoga maritima* C277T perhydrolase) under the same assay conditions.

In one embodiment, an isolated nucleic acid molecule encoding a polypeptide having perhydrolytic activity is provided selected from the group consisting of:

- (a) a polynucleotide encoding a polypeptide having perhydrolytic activity, said polypeptide comprising the amino acid sequence of SEQ ID NO: 18;
- (b) a polynucleotide comprising the nucleic acid sequence of SEQ ID NO: 44; and
- (c) a polynucleotide fully complementary to the polynucleotide of (a) or (b).

In other embodiments, a vector, a recombinant DNA construct, and a recombinant host cell comprising the present polynucleotide are also provided.

In another embodiment, a method for transforming a cell is provided comprising transforming a cell with the above nucleic acid molecule.

In another embodiment, an isolated polypeptide having perhydrolytic activity is provided comprising the amino acid sequence of SEQ ID NO: 18.

In one embodiment, the variant polypeptide having perhydrolytic activity is characterized by an increase in specific activity (as determined by an increased amount of peroxycarboxylic acid produced) when compared to the specific activity of the *Thermotoga maritima* C277T variant (Published U.S. Patent Application No. 2010-0087529 to DiCosimo et al.) under identical reaction conditions. In a preferred aspect, the relative increase in activity is measured under simulated laundry conditions (i.e., in the presence of a laundry care detergent formulation).


3

In another embodiment, a process for producing a peroxy-carboxylic acid is also provided comprising:

(a) providing a set of reaction components comprising:

(1) at least one substrate selected from the group consisting of:

(i) one or more esters having the structure



wherein


X=an ester group of the formula $R_6-C(O)O$;

$R_6=C1$ to $C7$ linear, branched or cyclic hydrocarbyl moiety, optionally substituted with hydroxyl groups or $C1$ to $C4$ alkoxy groups, wherein R_6 optionally comprises one or more ether linkages for $R_6=C2$ to $C7$;

R_5 =a $C1$ to $C6$ linear, branched, or cyclic hydrocarbyl moiety or a five-membered cyclic heteroaromatic moiety or six-membered cyclic aromatic or heteroaromatic moiety optionally substituted with hydroxyl groups; wherein each carbon atom in R_5 individually comprises no more than one hydroxyl group or no more than one ester or carboxylic acid group; wherein R_5 optionally comprises one or more ether linkages;


m =is an integer ranging from 1 to the number of carbon atoms in R_5 ; and wherein said esters have solubility in water of at least 5 ppm at 25° C.;

(ii) one or more glycerides having the structure



wherein $R_1=C1$ to $C21$ straight chain or branched chain alkyl optionally substituted with an hydroxyl or a $C1$ to $C4$ alkoxy group and R_3 and R_4 are individually H or $R_1C(O)$;

(iii) one or more esters of the formula:



wherein R_1 is a $C1$ to $C7$ straight chain or branched chain alkyl optionally substituted with an hydroxyl or a $C1$ to $C4$ alkoxy group and R_2 is a $C1$ to $C10$ straight chain or branched chain alkyl, alkenyl, alkynyl, aryl, alkylaryl, alkylheteroaryl, heteroaryl, $(CH_2CH_2O)_n$, or $(CH_2CH(CH_3)-O)_nH$ and n is 1 to 10;

(iv) one or more acylated monosaccharides, acylated disaccharides, or acylated polysaccharides; and

(v) any combination of (i) through (iv);

(2) a source of peroxygen; and

(3) an enzyme catalyst comprising a polypeptide having perhydrolytic activity, said polypeptide comprising the amino acid sequence of SEQ ID NO: 18;

(b) combining the set of reaction components under suitable reaction conditions whereby peroxy-carboxylic acid is produced; and

(c) optionally diluting the peroxy-carboxylic acid produced in step (b).

4


In another embodiment, a process is provided further comprising a step (d) wherein the peroxy-carboxylic acid produced in step (b) or step (c) is contacted with a hard surface, an article of clothing or an inanimate object whereby the hard surface, article of clothing or inanimate object is disinfected, sanitized, bleached, destained, deodorized or any combination thereof.

In another embodiment, a composition is provided comprising:

(a) a set of reaction components comprising:

(1) at least one substrate selected from the group consisting of:

(i) one or more esters having the structure



wherein

X=an ester group of the formula $R_6-C(O)O$;


$R_6=C1$ to $C7$ linear, branched or cyclic hydrocarbyl moiety, optionally substituted with hydroxyl groups or $C1$ to $C4$ alkoxy groups, wherein R_6 optionally comprises one or more ether linkages for $R_6=C2$ to $C7$;

R_5 =a $C1$ to $C6$ linear, branched, or cyclic hydrocarbyl moiety or a five-membered cyclic heteroaromatic moiety or six-membered cyclic aromatic or heteroaromatic moiety optionally substituted with hydroxyl groups; wherein each carbon atom in R_5 individually comprises no more than one hydroxyl group or no more than one ester group or carboxylic acid group; wherein R_5 optionally comprises one or more ether linkages;

m =is an integer ranging from 1 to the number of carbon atoms in R_5 ; and


wherein said esters have solubility in water of at least 5 ppm at 25° C.;

(ii) one or more glycerides having the structure



wherein $R_1=C1$ to $C21$ straight chain or branched chain alkyl optionally substituted with an hydroxyl or a $C1$ to $C4$ alkoxy group and R_3 and R_4 are individually H or $R_1C(O)$;

(iii) one or more esters of the formula:



wherein R_1 is a $C1$ to $C7$ straight chain or branched chain alkyl optionally substituted with an hydroxyl or a $C1$ to $C4$ alkoxy group and R_2 is a $C1$ to $C10$ straight chain or branched chain alkyl, alkenyl, alkynyl, aryl, alkylaryl, alkylheteroaryl, heteroaryl, $(CH_2CH_2O)_n$, or $(CH_2CH(CH_3)-O)_nH$ and n is 1 to 10;

(iv) one or more acylated monosaccharides, acylated disaccharides, or acylated polysaccharides; and

(v) any combination of (i) through (iv);

5

- (2) a source of peroxygen; and
 (3) an enzyme catalyst comprising a polypeptide, said polypeptide comprising perhydrolytic activity having the amino acid sequence of SEQ ID NO: 18; and
 (b) at least one peroxy-carboxylic acid formed upon combining the set of reaction components of (a).

The present process produces the desired peroxy-carboxylic acid upon combining the reaction components. The reaction components may remain separated until use.

In a further aspect, a peroxy-carboxylic acid generation and delivery system is provided comprising:

- (a) a first compartment comprising
 (1) an enzyme catalyst comprising a polypeptide having perhydrolytic activity, said polypeptide comprising the amino acid sequence of SEQ ID NO: 18;
 (2) at least one substrate selected from the group consisting of:
 (i) one or more esters having the structure

$[X]_mR_5$

wherein

X=an ester group of the formula $R_6-C(O)O$;


$R_6=C1$ to $C7$ linear, branched or cyclic hydrocarbyl moiety, optionally substituted with hydroxyl groups or $C1$ to $C4$ alkoxy groups, wherein R_6 optionally comprises one or more ether linkages for $R_6=C2$ to $C7$;

$R_5=a$ $C1$ to $C6$ linear, branched, or cyclic hydrocarbyl moiety or a five-membered cyclic heteroaromatic moiety or six-membered cyclic aromatic or heteroaromatic moiety optionally substituted with hydroxyl groups; wherein each carbon atom in R_5 individually comprises no more than one hydroxyl group or no more than one ester group or carboxylic acid group; wherein R_5 optionally comprises one or more ether linkages;

m =is an integer ranging from 1 to the number of carbon atoms in R_5 ; and


wherein said esters have solubility in water of at least 5 ppm at $25^\circ C$;

- (ii) one or more glycerides having the structure



wherein $R_1=C1$ to $C21$ straight chain or branched chain alkyl optionally substituted with an hydroxyl or a $C1$ to $C4$ alkoxy group and R_3 and R_4 are individually H or $R_1C(O)$;

- (iii) one or more esters of the formula:



wherein R_1 is a $C1$ to $C7$ straight chain or branched chain alkyl optionally substituted with an hydroxyl or a $C1$ to $C4$ alkoxy group and R_2 is a $C1$ to $C10$ straight chain or branched chain alkyl,

6

alkenyl, alkynyl, aryl, alkylaryl, alkylheteroaryl, heteroaryl, $(CH_2CH_2O)_n$, or $(CH_2CH(CH_3)-O)_nH$ and n is 1 to 10;

- (iv) one or more acylated monosaccharides, acylated disaccharides, or acylated polysaccharides; and
 (v) any combination of (i) through (iv); and

(3) an optional buffer; and

(b) a second compartment comprising

(1) source of peroxygen;

(2) a peroxide stabilizer; and

(3) an optional buffer.

In a further embodiment, a laundry care composition is provided comprising a polypeptide comprising the amino acid sequence of SEQ ID NO: 18.

In a further embodiment, a personal care composition is provided comprising a polypeptide comprising the amino acid sequence of SEQ ID NO: 18.

BRIEF DESCRIPTION OF THE BIOLOGICAL SEQUENCES

The following sequences comply with 37 C.F.R. §§1.821-1.825 ("Requirements for Patent Applications Containing Nucleotide Sequences and/or Amino Acid Sequence Disclosures—the Sequence Rules") and are consistent with World Intellectual Property Organization (WIPO) Standard ST.25 (2009) and the sequence listing requirements of the European Patent Convention (EPC) and the Patent Cooperation Treaty (PCT) Rules 5.2 and 49.5(a-bis), and Section 208 and Annex C of the Administrative Instructions. The symbols and format used for nucleotide and amino acid sequence data comply with the rules set forth in 37 C.F.R. §1.822.

SEQ ID NO: 1 is the nucleic acid sequence of the codon-optimized coding region encoding the wild-type *Thermotoga maritima* acetyl xylan esterase having perhydrolytic activity.

SEQ ID NO: 2 is the amino acid sequence of the wild-type *Thermotoga maritima* acetyl xylan esterase having perhydrolytic activity.

SEQ ID NOs: 3 and 4 are the nucleic acid sequences of primers used to prepare the C277T variant acetyl xylan esterase.

SEQ ID NO: 5 is the amino acid sequence of the C277T variant acetyl xylan esterase having perhydrolytic activity (Published U.S. Patent Application No. 2010-0087529 to DiCosimo et al.).

SEQ ID NO: 6 is the nucleic acid sequence of the plasmid pSW202/C277T.

SEQ ID NOs: 7 and 8 are the nucleic acid sequences of primers used for error-prone PCR.

SEQ ID NO: 9 is the amino acid sequence of the 007C7 variant acetyl xylan esterase.

SEQ ID NO: 10 is the amino acid sequence of the 007A11 variant acetyl xylan esterase.

SEQ ID NO: 11 is the amino acid sequence of the 007A12 variant acetyl xylan esterase.

SEQ ID NO: 12 is the amino acid sequence of the 007H10 variant acetyl xylan esterase.

SEQ ID NO: 13 is the amino acid sequence of the 007A5 variant acetyl xylan esterase.

SEQ ID NO: 14 is the amino acid sequence of the 007A8 variant acetyl xylan esterase.

SEQ ID NO: 15 is the amino acid sequence of the 007B12 variant acetyl xylan esterase.

SEQ ID NO: 16 is the amino acid sequence of the 007B2 variant acetyl xylan esterase.

SEQ ID NO: 17 is the amino acid sequence of the 007B3 variant acetyl xylan esterase.

SEQ ID NO: 18 is the amino acid sequence of the 007B5 variant acetyl xylan esterase.

SEQ ID NO: 19 is the amino acid sequence of the 007B7 variant acetyl xylan esterase.

SEQ ID NO: 20 is the amino acid sequence of the 007C4 variant acetyl xylan esterase.

SEQ ID NO: 21 is the amino acid sequence of the 007C3 variant acetyl xylan esterase.

SEQ ID NO: 22 is the amino acid sequence of the 007C8 variant acetyl xylan esterase.

SEQ ID NO: 23 is the amino acid sequence of the 007D1 variant acetyl xylan esterase.

SEQ ID NO: 24 is the amino acid sequence of the 007D11 variant acetyl xylan esterase.

SEQ ID NO: 25 is the amino acid sequence of the 007D12 variant acetyl xylan esterase.

SEQ ID NO: 26 is the amino acid sequence of the 007D2 variant acetyl xylan esterase.

SEQ ID NO: 27 is the amino acid sequence of the 007D5 variant acetyl xylan esterase.

SEQ ID NO: 28 is the amino acid sequence of the 007D6 variant acetyl xylan esterase.

SEQ ID NO: 29 is the amino acid sequence of the 007D8 variant acetyl xylan esterase.

SEQ ID NO: 30 is the amino acid sequence of the 007E11 variant acetyl xylan esterase.

SEQ ID NO: 31 is the amino acid sequence of the 007E6 variant acetyl xylan esterase.

SEQ ID NO: 32 is the amino acid sequence of the 007E9 variant acetyl xylan esterase.

SEQ ID NO: 33 is the amino acid sequence of the 007F10 variant acetyl xylan esterase.

SEQ ID NO: 34 is the amino acid sequence of the 007G11 variant acetyl xylan esterase.

SEQ ID NO: 35 is the amino acid sequence of the 007G5 variant acetyl xylan esterase.

SEQ ID NO: 36 is the amino acid sequence of the 007G6 variant acetyl xylan esterase.

SEQ ID NO: 37 is the amino acid sequence of the 007G8 variant acetyl xylan esterase.

SEQ ID NO: 38 is the nucleic acid sequence encoding the 007C7 variant acetyl xylan esterase.

SEQ ID NO: 39 is the nucleic acid sequence encoding the 007A11 variant acetyl xylan esterase.

SEQ ID NO: 40 is the nucleic acid sequence encoding the 007A12 variant acetyl xylan esterase.

SEQ ID NO: 41 is the nucleic acid sequence encoding the 007A5 variant acetyl xylan esterase.

SEQ ID NO: 42 is the nucleic acid sequence encoding the 007A8 variant acetyl xylan esterase.

SEQ ID NO: 43 is the nucleic acid sequence encoding the 007B2 variant acetyl xylan esterase.

SEQ ID NO: 44 is the nucleic acid sequence encoding the 007B5 variant acetyl xylan esterase.

SEQ ID NO: 45 is the nucleic acid sequence encoding the 007D1 variant acetyl xylan esterase.

SEQ ID NO: 46 is the nucleic acid sequence encoding the 007D5 variant acetyl xylan esterase.

SEQ ID NO: 47 is the nucleic acid sequence encoding the 007E9 variant acetyl xylan esterase.

DETAILED DESCRIPTION

A nucleic acid molecule encoding the *Thermotoga maritima* C277T variant acetyl xylan esterase was mutated to create a library of variant perhydrolases. Several perhydrolase variants were identified exhibiting an increase in specific

activity when compared to the specific activity of the *Thermotoga maritima* C277T perhydrolase having amino acid sequence SEQ ID NO: 5.

Compositions and methods are provided comprising the variant perhydrolase enzyme having amino acid sequence SEQ ID NO: 18.

In this disclosure, a number of terms and abbreviations are used. The following definitions apply unless specifically stated otherwise.

As used herein, the articles “a”, “an”, and “the” preceding an element or component of the invention are intended to be nonrestrictive regarding the number of instances (i.e., occurrences) of the element or component. Therefore “a”, “an” and “the” should be read to include one or at least one, and the singular word form of the element or component also includes the plural unless the number is obviously meant to be singular.

The term “comprising” means the presence of the stated features, integers, steps, or components as referred to in the claims, but does not preclude the presence or addition of one or more other features, integers, steps, components or groups thereof. The term “comprising” is intended to include embodiments encompassed by the terms “consisting essentially of” and “consisting of”. Similarly, the term “consisting essentially of” is intended to include embodiments encompassed by the term “consisting of”.

As used herein, the term “about” modifying the quantity of an ingredient or reactant employed refers to variation in the numerical quantity that can occur, for example, through typical measuring and liquid handling procedures used for making concentrates or use solutions in the real world; through inadvertent error in these procedures; through differences in the manufacture, source, or purity of the ingredients employed to make the compositions or carry out the methods; and the like. The term “about” also encompasses amounts that differ due to different equilibrium conditions for a composition resulting from a particular initial mixture. Whether or not modified by the term “about”, the claims include equivalents to the quantities.

Where present, all ranges are inclusive and combinable. For example, when a range of “1 to 5” is recited, the recited range should be construed as including ranges “1 to 4”, “1 to 3”, “1-2”, “1-2 & 4-5”, “1-3 & 5”, and the like.

As used herein, the term “multi-component system” will refer to a system of enzymatically generating peroxy-carboxylic acid wherein the components remain separated until use. As such, the multi-component system will include at least one first component that remains separated from at least one second component. The first and second components are separated in different compartments until use (i.e., using first and second compartments). The design of the multi-component systems will often depend on the physical form of the components to be combined and are described in more detail below.

As used herein, the term “peroxy-carboxylic acid” is synonymous with peracid, peroxyacid, peroxy acid, percarboxylic acid and peroxy acid.

As used herein, the term “peracetic acid” is abbreviated as “PAA” and is synonymous with peroxyacetic acid, ethaneperoxy acid and all other synonyms of CAS Registry Number 79-21-0.

As used herein, the term “monoacetin” is synonymous with glycerol monoacetate, glycerin monoacetate, and glyceryl monoacetate.

As used herein, the term “diacetin” is synonymous with glycerol diacetate; glycerin diacetate, glyceryl diacetate, and all other synonyms of CAS Registry Number 25395-31-7.

As used herein, the term “triacetin” is synonymous with glycerin triacetate; glycerol triacetate; glyceryl triacetate; 1,2,3-triacetoxypropane; 1,2,3-propanetriol triacetate; and all other synonyms of CAS Registry Number 102-76-1.

As used herein, the term “monobutylin” is synonymous with glycerol monobutyrate, glycerin monobutyrate, and glyceryl monobutyrate.

As used herein, the term “dibutylin” is synonymous with glycerol dibutyrate and glyceryl dibutyrate.

As used herein, the term “tributylin” is synonymous with glycerol tributyrate; 1,2,3-tributyrylglycerol; and all other synonyms of CAS Registry Number 60-01-5.

As used herein, the term “monopropionin” is synonymous with glycerol monopropionate, glycerin monopropionate, and glyceryl monopropionate.

As used herein, the term “dipropionin” is synonymous with glycerol dipropionate and glyceryl dipropionate.

As used herein, the term “tripropionin” is synonymous with glyceryl tripropionate, glycerol tripropionate; 1,2,3-tripropionylglycerol; and all other synonyms of CAS Registry Number 139-45-7.

As used herein, the term “ethyl acetate” is synonymous with acetic ether, acetoxyethane, ethyl ethanoate, acetic acid ethyl ester, ethanoic acid ethyl ester, ethyl acetic ester and all other synonyms of CAS Registry Number 141-78-6.

As used herein, the term “ethyl lactate” is synonymous with lactic acid ethyl ester and all other synonyms of CAS Registry Number 97-64-3.

As used herein, the terms “acylated sugar” and “acylated saccharide” refer to mono-, di- and polysaccharides comprising at least one acyl group, where the acyl group is selected from the group consisting of straight chain aliphatic carboxylates having a chain length from C2 to C8. Examples include, but are not limited to, glucose pentaacetate, galactose pentaacetate, sucrose octaacetate, sorbitol hexaacetate, tetraacetylxylofuranose, α -D-glucopyranose pentaacetate, α -D-mannopyranose pentaacetate, acetylated xylan, acetylated xylan fragments, β -D-ribofuranose-1,2,3,5-tetraacetate, tri-O-acetyl-D-galactal, and tri-O-acetyl-glucal.

As used herein, the terms “hydrocarbyl”, “hydrocarbyl group”, and “hydrocarbyl moiety” mean a straight chain, branched or cyclic arrangement of carbon atoms connected by single, double, or triple carbon to carbon bonds and/or by ether linkages, and substituted accordingly with hydrogen atoms. Such hydrocarbyl groups may be aliphatic and/or aromatic. Examples of hydrocarbyl groups include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, t-butyl, cyclopropyl, cyclobutyl, pentyl, cyclopentyl, methylcyclopentyl, hexyl, cyclohexyl, benzyl, and phenyl. In one embodiment, the hydrocarbyl moiety is a straight chain, branched or cyclic arrangement of carbon atoms connected by single carbon to carbon bonds and/or by ether linkages, and substituted accordingly with hydrogen atoms.

As used herein, the term “aromatic” refers to an organic compound or moiety characterized by increased chemical stability resulting from the delocalization of electrons in a ring system containing usually multiple conjugated double bonds. Planar monocyclic conjugated rings having delocalized electrons should be aromatic if they have $(4n+2)$ π electrons. Examples of aromatic compounds may include derivatives of benzene (such as 2-, 3- or 4-acetoxybenzoic acid). In one embodiment, the ester substrate may be 4-acetoxybenzoic acid.

As used herein, the term “heterocyclic” refers to an organic compound or moiety with a ring structure having one or more atoms other than carbon in at least one of its rings.

As used herein, the term “heteroaromatic” refers to an organic compound or moiety with a ring structure that is both heterocyclic and aromatic, wherein the ring comprises at least one of the heteroatoms oxygen, nitrogen, or sulfur. Examples of heteroaromatic moieties may include pyridine, pyrrole, furan, and thiophene moieties.

As used herein, the terms “monoesters” and “diesters” of 1,2-ethanediol, 1,2-propanediol, 1,3-propanediol, 1,2-butanediol, 1,3-butanediol, 2,3-butanediol, 1,4-butanediol, 1,2-pentanediol, 2,5-pentanediol, 1,6-pentanediol, 1,2-hexanediol, 2,5-hexanediol, 1,6-hexanediol, refer to said compounds comprising at least one ester group of the formula $RC(O)O$, wherein R is a C1 to C7 linear hydrocarbyl moiety.

As used herein, the terms “suitable enzymatic reaction formulation”, “components suitable for generation of a peroxycarboxylic acid”, “suitable reaction components”, “reaction components”, “reaction formulation”, and “suitable aqueous reaction formulation” refer to the materials and water in which the reactants and the enzyme catalyst comprising the present variant polypeptide having perhydrolytic activity come into contact to form the desired peroxycarboxylic acid. The components of the reaction formulation are provided herein and those skilled in the art appreciate the range of component variations suitable for this process. In one embodiment, the enzymatic reaction formulation produces peroxycarboxylic acid in situ upon combining the reaction components. As such, the reaction components may be provided as a multi-component system wherein one or more of the reaction components remains separated until use. The design of systems and means for separating and combining multiple active components are known in the art and generally will depend upon the physical form of the individual reaction components. For example, multiple active fluids (liquid-liquid) systems typically use multi-chamber dispenser bottles or two-phase systems (U.S. Patent Application Publication No. 2005-0139608; U.S. Pat. No. 5,398,846; U.S. Pat. No. 5,624,634; U.S. Pat. No. 6,391,840; E.P. Patent 080715661; U.S. Patent Application Publication No. 2005-0008526; and PCT Publication No. WO 00/61713A1) such as found in some bleaching applications wherein the desired bleaching agent is produced upon mixing the reactive fluids. Multi-component formulations and multi-component generation systems to enzymatically produce peroxycarboxylic acids from carboxylic acid esters are described by DiCosimo et al. in published U.S. Patent Application Nos. 2010-0086510 and 2010-0086621, respectively. Other forms of multi-component systems used to generate peroxycarboxylic acid may include, but are not limited to, those designed for one or more solid components or combinations of solid-liquid components, such as powders used in many commercially available bleaching compositions (e.g., U.S. Pat. No. 5,116,575), multi-layered tablets (e.g., U.S. Pat. No. 6,210,639), water dissolvable packets having multiple compartments (e.g., U.S. Pat. No. 6,995,125) and solid agglomerates that react upon the addition of water (e.g., U.S. Pat. No. 6,319,888).

As used herein, the term “substrate” or “carboxylic acid ester substrate” will refer to the reaction components enzymatically perhydrolyzed using the present enzyme catalyst in the presence of a suitable source of peroxygen, such as hydrogen peroxide. In one embodiment, the substrate comprises at least one ester group capable of being enzymatically perhydrolyzed using the enzyme catalyst, whereby a peroxycarboxylic acid is produced.

As used herein, the term “perhydrolysis” is defined as the reaction of a selected substrate with a source of hydrogen peroxide to form a peroxycarboxylic acid. Typically, inorganic peroxide is reacted with the selected substrate in the

presence of a catalyst to produce the peroxy-carboxylic acid. As used herein, the term “chemical perhydrolysis” includes perhydrolysis reactions in which a substrate (such as a peroxy-carboxylic acid precursor) is combined with a source of hydrogen peroxide wherein peroxy-carboxylic acid is formed in the absence of an enzyme catalyst. As used herein, the term “enzymatic perhydrolysis” refers to a reaction of a selected substrate with a source of hydrogen peroxide to form a peroxy-carboxylic acid, wherein the reaction is catalyzed by an enzyme catalyst having perhydrolysis activity.

As used herein, the term “perhydrolase activity” refers to the enzyme catalyst activity per unit mass (for example, milligram) of protein, dry cell weight, or immobilized catalyst weight.

As used herein, “one unit of enzyme activity” or “one unit of activity” or “U” is defined as the amount of perhydrolase activity required for the production of 1 μ mol of peroxy-carboxylic acid product (such as peracetic acid) per minute at a specified temperature. “One unit of enzyme activity” may also be used herein to refer to the amount of peroxy-carboxylic acid hydrolysis activity required for the hydrolysis of 1 μ mol of peroxy-carboxylic acid (e.g., peracetic acid) per minute at a specified temperature.

The present variant CE-7 carbohydrate esterase is characterized by an increase in specific activity when compared to the perhydrolase from which it was derived (*Thermotoga maritima* C277T; Published U.S. Patent Application No. 2010-0087529) under the same reaction conditions. As used herein, the “fold increase” in specific activity is measured relative to the specific activity of the parent perhydrolase from which the variant was derived (the *Thermotoga maritima* C277T perhydrolase (SEQ ID NO: 5) under the same reaction conditions). In one embodiment, the fold increase in specific activity of the variant polypeptide (i.e., variant perhydrolase) relative to the parent perhydrolase (*Thermotoga maritima* C277T, SEQ ID NO: 5) is at least 1.01, 1.05, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 3.0, 4.0, 5.0, 6.0, 7.0, 8.0, 9.0, or 10-fold when compared under identical reaction/assay conditions.

As used herein, “identical assay conditions” or “same assay conditions” refer to the conditions used to measure the peracid formation (i.e., perhydrolysis of a carboxylic acid ester substrate) specific activity of the variant polypeptide relative to the respective specific activity of the polypeptide from which it is derived. The assay conditions used to measure the respective specific activities should be as close to identical as possible such that only the structure of the polypeptide having perhydrolytic activity varies.

As used herein, the terms “enzyme catalyst” and “perhydrolase catalyst” refer to a catalyst comprising an enzyme (i.e., a polypeptide) having perhydrolysis activity and may be in the form of a whole microbial cell, permeabilized microbial cell(s), one or more cell components of a microbial cell extract, partially purified enzyme, or purified enzyme. The enzyme catalyst may also be chemically modified (for example, by pegylation or by reaction with cross-linking reagents). The perhydrolase catalyst may also be immobilized on a soluble or insoluble support using methods well-known to those skilled in the art; see for example, *Immobilization of Enzymes and Cells*; Gordon F. Bickerstaff, Editor; Humana Press, Totowa, N.J., USA; 1997.

The present enzyme catalyst comprises a variant polypeptide having perhydrolytic activity and is structurally classified as a member of the carbohydrate family esterase family 7 (CE-7 family) of enzymes (see Coutinho, P. M., Henrissat, B. “Carbohydrate-active enzymes: an integrated database approach” in *Recent Advances in Carbohydrate Bioengineer-*

ing, H. J. Gilbert, G. Davies, B. Henrissat and B. Svensson eds., (1999) The Royal Society of Chemistry, Cambridge, pp. 3-12.). The CE-7 family of enzymes has been demonstrated to be particularly effective for producing peroxy-carboxylic acids from a variety of carboxylic acid ester substrates when combined with a source of peroxygen (See PCT publication No. WO2007/070609 and U.S. Patent Application Publication No. 2008-0176299 and U.S. Pat. Nos. 7,951,566 and 7,723,083 to DiCosimo et al.; each herein incorporated by reference in their entireties). The CE-7 enzyme family includes cephalosporin C deacetylases (CAHs; E.C. 3.1.1.41) and acetyl xylan esterases (AXEs; E.C. 3.1.1.72). Members of the CE-7 enzyme family share a conserved signature motif (Vincent et al., *J. Mol. Biol.*, 330:593-606 (2003)).

As used herein, the terms “signature motif” and “CE-7 signature motif”, refer to conserved structures shared among a family of enzymes having a perhydrolytic activity.

As used herein, “structurally classified as a CE-7 enzyme”, “structurally classified as a carbohydrate esterase family 7 enzyme”, “structurally classified as a CE-7 carbohydrate esterase”, and “CE-7 perhydrolase” will be used to refer to enzymes having perhydrolysis activity that are structurally classified as a CE-7 carbohydrate esterase based on the presence of the CE-7 signature motif (Vincent et al., supra). The “signature motif” for CE-7 esterases comprises three conserved motifs (residue position numbering relative to reference sequence SEQ ID NO: 2; the wild-type *Thermotoga maritima* acetyl xylan esterase):

- a) Arg118-Gly119-Gln120;
- b) Gly186-Xaa187-**Ser188**-Gln189-Gly190;
and
- c) **His303**-Glu304.

Typically, the Xaa at amino acid residue position 187 is glycine, alanine, proline, tryptophan, or threonine. Two of the three amino acid residues belonging to the catalytic triad are in bold. In one embodiment, the Xaa at amino acid residue position 187 is selected from the group consisting of glycine, alanine, proline, tryptophan, and threonine.

Further analysis of the conserved motifs within the CE-7 carbohydrate esterase family indicates the presence of an additional conserved motif (LXD at amino acid positions 272-274 of SEQ ID NO: 2) that may be used to further define a member of the CE-7 carbohydrate esterase family. In a further embodiment, the signature motif defined above includes a fourth conserved motif defined as:

Leu272-Xaa273-Asp274.

The Xaa at amino acid residue position 273 is typically isoleucine, valine, or methionine. The fourth motif includes the aspartic acid residue (bold) belonging to the catalytic triad (Ser188-Asp274-His303).

As used herein, the terms “cephalosporin C deacetylase” and “cephalosporin C acetyl hydrolase” refer to an enzyme (E.C. 3.1.1.41) that catalyzes the deacetylation of cephalosporins such as cephalosporin C and 7-aminocephalosporanic acid (Mitsushima et al., *Appl. Environ. Microbiol.*, 61(6): 2224-2229 (1995); U.S. Pat. No. 5,528,152; and U.S. Pat. No. 5,338,676). Enzymes classified as cephalosporin C deacetylases have been shown to often have significant perhydrolytic activity (U.S. Pat. No. 7,951,566 and U.S. Patent Application Publication No. 2008-0176299 to DiCosimo et al.).

As used herein, “acetyl xylan esterase” refers to an enzyme (E.C. 3.1.1.72; AXEs) that catalyzes the deacetylation of

acetylated xylans and other acetylated saccharides. Enzymes classified as acetyl xylan esterases have been shown to have significant perhydrolytic activity (U.S. Pat. Nos. 7,951,566 and 7,723,083 and U.S. Patent Application Publication No. 2008-0176299; each to DiCosimo et al.).

As used herein, the term “*Thermotoga maritima*” refers to a bacterial cell reported to have acetyl xylan esterase activity (GENBANK®NP_227893.1). In one aspect, the *Thermotoga maritima* strain is *Thermotoga maritima* MSB8. The amino acid sequence of the wild-type enzyme having perhydrolytic activity from *Thermotoga maritima* is provided as SEQ ID NO: 2.

As used herein, the terms “variant”, “variant polypeptide”, and “variant enzyme catalyst” refer to an enzyme catalyst comprising at least one polypeptide (i.e., a perhydrolytic activity wherein the polypeptide comprises at least one amino acid change relative to the enzyme/polypeptide from which it was derived (i.e., *Thermotoga maritima* C277T perhydrolytic activity). Several variant polypeptides are provided herein having perhydrolytic activity and are characterized by an increase in specific activity relative to the *Thermotoga maritima* C277T acetyl xylan esterase having amino acid sequence SEQ ID NO: 5.

For a particular variant perhydrolytic, amino acid substitutions are specified with reference to the wild type *Thermotoga maritima* amino acid sequence (SEQ ID NO: 2). The wild-type amino acid (denoted by the standard single letter abbreviation) is followed by the amino acid residue position of SEQ ID NO: 2 followed by the amino acid of the variant (also denoted by the standard single letter abbreviation). For example, “C277T” describes a change in SEQ ID NO: 2 at amino acid residue position 277 where cysteine was changed to threonine. The variant polypeptide may be comprised of multiple point substitutions. For example, R261 S/I264F/C277T refers to a variant polypeptide having three point substitutions: 1) a change at amino acid residue position 261 where an arginine was changed to a serine, 2) a change at residue position 264 where an isoleucine was changed to a phenylalanine, and 3) a change at position 277 where a cysteine was changed to a threonine.

The term “amino acid” refers to the basic chemical structural unit of a protein or polypeptide. The following abbreviations are used herein to identify specific amino acids:

Amino Acid	Three-Letter Abbreviation	One-Letter Abbreviation
Alanine	Ala	A
Arginine	Arg	R
Asparagine	Asn	N
Aspartic acid	Asp	D
Cysteine	Cys	C
Glutamine	Gln	Q
Glutamic acid	Glu	E
Glycine	Gly	G
Histidine	His	H
Isoleucine	Ile	I
Leucine	Leu	L
Lysine	Lys	K
Methionine	Met	M
Phenylalanine	Phe	F
Proline	Pro	P
Serine	Ser	S
Threonine	Thr	T
Tryptophan	Trp	W
Tyrosine	Tyr	Y
Valine	Val	V
Any amino acid (or as defined herein)	Xaa	X

As used herein, the term “biological contaminants” refers to one or more unwanted and/or pathogenic biological entities including, but not limited to, microorganisms, spores, viruses, prions, and mixtures thereof. The present enzyme can be used to produce an efficacious concentration of at least one peroxycarboxylic acid useful to reduce and/or eliminate the presence of the viable biological contaminants. In a preferred embodiment, the biological contaminant is a viable pathogenic microorganism.

As used herein, the term “disinfect” refers to the process of destruction of or prevention of the growth of biological contaminants. As used herein, the term “disinfectant” refers to an agent that disinfects by destroying, neutralizing, or inhibiting the growth of biological contaminants. Typically, disinfectants are used to treat inanimate objects or surfaces. As used herein, the term “antiseptic” refers to a chemical agent that inhibits the growth of disease-carrying microorganisms. In one aspect of the embodiment, the biological contaminants are pathogenic microorganisms.

As used herein, the term “sanitary” means of or relating to the restoration or preservation of health, typically by removing, preventing or controlling an agent that may be injurious to health. As used herein, the term “sanitize” means to make sanitary. As used herein, the term “sanitizer” refers to a sanitizing agent. As used herein the term “sanitization” refers to the act or process of sanitizing.

As used herein, the term “virucide” refers to an agent that inhibits or destroys viruses, and is synonymous with “viricide”. An agent that exhibits the ability to inhibit or destroy viruses is described as having “virucidal” activity. Peroxycarboxylic acids can have virucidal activity. Typical alternative virucides known in the art which may be suitable for use with the present invention include, for example, alcohols, ethers, chloroform, formaldehyde, phenols, beta propiolactone, iodine, chlorine, mercury salts, hydroxylamine, ethylene oxide, ethylene glycol, quaternary ammonium compounds, enzymes, and detergents.

As used herein, the term “biocide” refers to a chemical agent, typically broad spectrum, which inactivates or destroys microorganisms. A chemical agent that exhibits the ability to inactivate or destroy microorganisms is described as having “biocidal” activity. Peroxycarboxylic acids can have biocidal activity. Typical alternative biocides known in the art, which may be suitable for use in the present invention include, for example, chlorine, chlorine dioxide, chloroisocyanurates, hypochlorites, ozone, acrolein, amines, chlorinated phenolics, copper salts, organo-sulphur compounds, and quaternary ammonium salts.

As used herein, the phrase “minimum biocidal concentration” refers to the minimum concentration of a biocidal agent that, for a specific contact time, will produce a desired lethal, irreversible reduction in the viable population of the targeted microorganisms. The effectiveness can be measured by the \log_{10} reduction in viable microorganisms after treatment. In one aspect, the targeted reduction in viable microorganisms after treatment is at least a 3- \log_{10} reduction, more preferably at least a 4- \log_{10} reduction, and most preferably at least a 5- \log_{10} reduction. In another aspect, the minimum biocidal concentration is at least a 6- \log_{10} reduction in viable microbial cells.

As used herein, the terms “peroxygen source” and “source of peroxygen” refer to compounds capable of providing hydrogen peroxide at a concentration of about 0.5 mM or more when in an aqueous solution including, but not limited to, hydrogen peroxide, hydrogen peroxide adducts (e.g., urea-hydrogen peroxide adduct (carbamide peroxide)), perborates, and percarbonates, such as sodium percarbonate. As

described herein, the concentration of the hydrogen peroxide provided by the peroxygen compound in the aqueous reaction formulation is initially at least 0.5 mM or more upon combining the reaction components. In one embodiment, the hydrogen peroxide concentration in the aqueous reaction formulation is at least 1 mM. In another embodiment, the hydrogen peroxide concentration in the aqueous reaction formulation is at least 10 mM. In another embodiment, the hydrogen peroxide concentration in the aqueous reaction formulation is at least 100 mM. In another embodiment, the hydrogen peroxide concentration in the aqueous reaction formulation is at least 200 mM. In another embodiment, the hydrogen peroxide concentration in the aqueous reaction formulation is 500 mM or more. In yet another embodiment, the hydrogen peroxide concentration in the aqueous reaction formulation is 1000 mM or more. The molar ratio of the hydrogen peroxide to enzyme substrate, such as triglyceride, (H₂O₂:substrate) in the aqueous reaction formulation may be from about 0.002 to 20, preferably about 0.1 to 10, and most preferably about 0.5 to 5.

As used herein, the term "benefit agent" refers to a material that promotes or enhances a useful advantage, a favorable/desirable effect or benefit. In one embodiment, a process is provided whereby a benefit agent, such as a composition comprising a peroxycarboxylic acid, is applied to a textile or article of clothing to achieve a desired benefit, such as disinfecting, bleaching, destaining, deodorizing, and any combination thereof. In another embodiment, the present variant polypeptide having perhydrolytic activity may be used to produce a peracid-based benefit agent for use in personal care products (such as hair care products, skin care products, nail care products or oral care products). In one embodiment, a personal care product is provided comprising the variant perhydrolytic activity having amino acid sequence SEQ ID NO: 18. The personal care products are formulated to provide a safe and efficacious concentration of the desired peracid benefit agent. Variant Polypeptides Having an Increase in Specific Activity.

The present variant polypeptides were derived from the *Thermotoga maritima* C277T acetyl xylan esterase that has been previously demonstrated to have significant perhydrolytic activity for producing peroxycarboxylic acids from carboxylic acid esters and a source of peroxygen, such as hydrogen peroxide (U.S. Patent Application Publication No. 2008-0176299 and 2010-0087529, each to DiCosimo et al.).

Libraries of variant polypeptides were created from the C277T *Thermotoga maritima* perhydrolytic activity (SEQ ID NO: 5) and assayed for an increase in the specific activity for producing peroxycarboxylic acids from carboxylic acid ester substrates. The assay conditions used to measure the respective specific activities should be as close to identical as possible such that only the structure of the polypeptide having perhydrolytic activity varies. In one embodiment, reactions used to measure specific activity are run at ca. 25° C. in reactions containing 1 mM triacetin, 4 mM hydrogen peroxide and approximately 1.0 µg/mL of heat-treated extract supernatant total protein from *E. coli* strain KLP18 expressing the C277T perhydrolytic activity or variant perhydrolytic activity, in the presence of 2 mg/mL of a laundry detergent (powder) having a formulation generally comprising surfactants (sodium dodecylbenzenesulfonate, C12-15 Pareth-5, C12-15 Pareth-7, sodium stearate, and stearic acid), builders (sodium carbonate, zeolite, sodium silicate, and citric acid), binders (cellulose, PEG-75, dextrin, and sucrose), bulking agents (sodium sulfate, sodium chloride, sodium bicarbonate, and calcium carbonate), structurants (sodium acrylic acid/MA copolymer, and sodium polyacrylate), sequestrants (tetrasodium etidronate and calcium sodium EDTMP), optical


brighteners (disodium anilinomorpholinotriazinyl-aminostilbenesulfonate), stabilizing agents, anti-redeposition agents, antifoaming agents, and softness extenders at pH 10.7 (see Example 5).

In another embodiment, the reactions to measure specific activity are conducted under simulated laundry care conditions at ca. 25° C. using 2 mg/mL of a laundry detergent (powder), 0.75 mM triacetin, 1.4 mM hydrogen peroxide and 0.5 µg/mL to 4 µg/mL of heat-treated extract supernatant total soluble protein from *E. coli* strain KLP18 expressing the *T. maritima* C277T perhydrolytic activity or the present variant perhydrolytic activity at pH 10.7 (see Example 6).

Suitable Reaction Conditions for the Enzyme-Catalyzed Preparation of Peroxycarboxylic Acids from Carboxylic Acid Esters and Hydrogen Peroxide

A process is provided to produce an aqueous formulation comprising at least one peroxycarboxylic acid by reacting carboxylic acid esters and an inorganic peroxide (such as, e.g., hydrogen peroxide, sodium perborate or sodium percarbonate) in the presence of an enzyme catalyst having perhydrolytic activity, wherein the enzyme catalyst comprises, in one embodiment, a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NOs: 9, 10, 11, 13, 14, 16, 18, 23, 27, and 32. In a further embodiment, the polypeptide has the amino acid sequence of SEQ ID NO: 18.

In one embodiment, suitable substrates include one or more esters provided by the following formula:



wherein X=an ester group of the formula R₆C(O)O

R₆=C1 to C7 linear, branched or cyclic hydrocarbyl moiety, optionally substituted with hydroxyl groups or C1 to C4 alkoxy groups, wherein R₆ optionally comprises one or more ether linkages for R₆=C2 to C7;

R₅=a C1 to C6 linear, branched, or cyclic hydrocarbyl moiety or a five-membered cyclic heteroaromatic moiety or six-membered cyclic aromatic or heteroaromatic moiety optionally substituted with hydroxyl groups; wherein each carbon atom in R₅ individually comprises no more than one hydroxyl group or no more than one ester group or carboxylic acid group; wherein R₅ optionally comprises one or more ether linkages;


m=is an integer ranging from 1 to the number of carbon atoms in R₅; and

wherein said esters have solubility in water of at least 5 ppm at 25° C.

In another embodiment, R₆ is C1 to C7 linear hydrocarbyl moiety, optionally substituted with hydroxyl groups or C1 to C4 alkoxy groups, optionally comprising one or more ether linkages. In a further preferred embodiment, R₆ is C2 to C7 linear hydrocarbyl moiety, optionally substituted with hydroxyl groups, and/or optionally comprising one or more ether linkages.

In one embodiment, the suitable substrate may include 2-acetoxybenzoic acid, 3-acetoxybenzoic acid, 4-acetoxybenzoic acid or mixtures thereof.


In another embodiment, suitable substrates also include one or more glycerides of the formula:



17

wherein R_1 =C1 to C21 straight chain or branched chain alkyl optionally substituted with an hydroxyl or a C1 to C4 alkoxy group and R_3 and R_4 are individually H or $R_1C(O)$. In one embodiment, the suitable substrate is a glyceride of the above formula wherein R_1 =C1 to C7 straight chain or branched chain alkyl optionally substituted with an hydroxyl or a C1 to C4 alkoxy group and R_3 and R_4 are individually H or $R_1C(O)$.

In another aspect, suitable substrates may also include one or more esters of the formula:



wherein R_1 is a C1 to C7 straight chain or branched chain alkyl optionally substituted with an hydroxyl or a C1 to C4 alkoxy group and R_2 is a C1 to C10 straight chain or branched chain alkyl, alkenyl, alkynyl, aryl, alkylaryl, alkylheteroaryl, heteroaryl, $(CH_2CH_2O)_n$, or $(CH_2CH(CH_3)-O)_n$, H and n is 1 to 10.

Suitable substrates may also include one or more acylated saccharides selected from the group consisting of acylated mono-, di-, and polysaccharides. In another embodiment, the acylated saccharides are selected from the group consisting of acetylated xylan; fragments of acetylated xylan; acetylated xylose (such as xylose tetraacetate); acetylated glucose (such as α -D-glucose pentaacetate; β -D-glucose pentaacetate); β -D-galactose pentaacetate; sorbitol hexaacetate; sucrose octaacetate; β -D-ribofuranose-1,2,3,5-tetraacetate; tri-O-acetyl-D-galactal; tri-O-acetyl-D-glucal; tetraacetylxylofuranose, α -D-glucopyranose pentaacetate, α -D-mannopyranose pentaacetate, and acetylated cellulose. In a preferred embodiment, the acetylated saccharide is selected from the group consisting of β -D-ribofuranose-1,2,3,5-tetraacetate; tri-O-acetyl-D-galactal; tri-O-acetyl-D-glucal; sucrose octaacetate and acetylated cellulose.

In another embodiment, suitable substrates are selected from the group consisting of: monoacetin; diacetin; triacetin; monopropionin; dipropionin; tripropionin; monobutyryn; dibutyryn; tributyrin; glucose pentaacetate; xylose tetraacetate; acetylated xylan; acetylated xylan fragments; β -D-ribofuranose-1,2,3,5-tetraacetate; tri-O-acetyl-D-galactal; tri-O-acetyl-D-glucal; monoesters or diesters of 1,2-ethanediol; 1,2-propanediol; 1,3-propanediol; 1,2-butanediol; 1,3-butanediol; 2,3-butanediol; 1,4-butanediol; 1,2-pentanediol; 2,5-pentanediol; 1,6-pentanediol; 1,2-hexanediol; 2,5-hexanediol; 1,6-hexanediol; and mixtures thereof.

In another embodiment, the carboxylic acid ester is selected from the group consisting of monoacetin, diacetin, triacetin, and combinations thereof. In another embodiment, the substrate is a C1 to C6 polyol comprising one or more ester groups. In a preferred embodiment, one or more of the hydroxyl groups on the C1 to C6 polyol are substituted with one or more acetoxy groups (such as 1,3-propanediol diacetate, 1,4-butanediol diacetate, etc.). In a further embodiment, the substrate is propylene glycol diacetate (PGDA), ethylene glycol diacetate (EGDA), or a mixture thereof.

In another embodiment, suitable substrates are selected from the group consisting of ethyl acetate; methyl lactate; ethyl lactate; methyl glycolate; ethyl glycolate; methyl methoxyacetate; ethyl methoxyacetate; methyl 3-hydroxybutyrate; ethyl 3-hydroxybutyrate; triethyl 2-acetyl citrate; glucose pentaacetate; gluconolactone; glycerides (mono-, di-, and triglycerides) such as monoacetin, diacetin, triacetin, monopropionin, dipropionin (glyceryl dipropionate), tripro-

18

pionin (1,2,3-tripropionylglycerol), monobutyryn, dibutyryn (glyceryl dibutyrate), tributyrin (1,2,3-tributyrylglycerol); acetylated saccharides; and mixtures thereof.

In a further embodiment, suitable substrates are selected from the group consisting of monoacetin, diacetin, triacetin, monopropionin, dipropionin, tripropionin, monobutyryn, dibutyryn, tributyrin, ethyl acetate, and ethyl lactate. In yet another aspect, the substrate is selected from the group consisting of diacetin, triacetin, ethyl acetate, and ethyl lactate. In a most preferred embodiment, the suitable substrate comprises triacetin.

The carboxylic acid ester is present in the aqueous reaction formulation at a concentration sufficient to produce the desired concentration of peroxy-carboxylic acid upon enzyme-catalyzed perhydrolysis. The carboxylic acid ester need not be completely soluble in the aqueous reaction formulation, but has sufficient solubility to permit conversion of the ester by the perhydrolyase catalyst to the corresponding peroxy-carboxylic acid. The carboxylic acid ester is present in the aqueous reaction formulation at a concentration of 0.0005 wt % to 40 wt % of the aqueous reaction formulation, preferably at a concentration of 0.005 wt % to 20 wt % of the aqueous reaction formulation, and more preferably at a concentration of 0.01 wt % to 10 wt % of the aqueous reaction formulation. The wt % of carboxylic acid ester may optionally be greater than the solubility limit of the carboxylic acid ester, such that the concentration of the carboxylic acid ester is at least 0.0005 wt % in the aqueous reaction formulation that is comprised of water, enzyme catalyst, and source of peroxide, where the remainder of the carboxylic acid ester remains as a second separate phase of a two-phase aqueous/organic reaction formulation. Not all of the added carboxylic acid ester must immediately dissolve in the aqueous reaction formulation, and after an initial mixing of all reaction components, additional continuous or discontinuous mixing is optional.

The peroxy-carboxylic acids produced by the present reaction components may vary depending upon the selected substrates, so long as the present enzyme catalyst is used. In one embodiment, the peroxy-carboxylic acid produced is peracetic acid, perpropionic acid, perbutyric acid, peroctanoic acid, perlactic acid, perglycolic acid, permethoxyacetic acid, per- β -hydroxybutyric acid, or mixtures thereof.

The peroxygen source may include, but is not limited to, hydrogen peroxide, hydrogen peroxide adducts (e.g., urea-hydrogen peroxide adduct (carbamide peroxide)), perborate salts and percarbonate salts. Alternatively, hydrogen peroxide can be generated in situ by the reaction of a substrate and oxygen catalyzed by an enzyme having oxidase activity (including, but not limited to, glucose oxidase, galactose oxidase, sorbitol oxidase, hexose oxidase, alcohol oxidase, glycerol oxidase, monoamine oxidase, glycolate oxidase, lactate oxidase, pyruvate oxidase, oxalate oxidase, choline oxidase, cholesterol oxidase, pyranose oxidase, carbon/alcohol oxidase, L-amino acid oxidase, glycine oxidase, glutamate oxidase, lysine oxidase, and uricase). The concentration of peroxygen compound in the aqueous reaction formulation may range from 0.0017 wt % to about 50 wt %, preferably from 0.017 wt % to about 40 wt %, more preferably from 0.17 wt % to about 30 wt %.

Many perhydrolyase catalysts (such as whole cells, permeabilized whole cells, and partially purified whole cell extracts) have been reported to have catalase activity (EC 1.11.1.6). Catalases catalyze the conversion of hydrogen peroxide into oxygen and water. In one aspect, the enzyme catalyst having perhydrolyase activity lacks catalase activity. In another aspect, the enzyme catalyst having perhydrolyase

activity has a sufficiently-low catalase activity that the presence of said catalase activity does not significantly interfere with perhydrolase-catalyzed peroxy-carboxylic acid production. In another aspect, a catalase inhibitor is added to the aqueous reaction formulation. Examples of catalase inhibitors include, but are not limited to, sodium azide and hydroxylamine sulfate. One of skill in the art can adjust the concentration of catalase inhibitor as needed. The concentration of the catalase inhibitor typically ranges from 0.1 mM to about 1 M; preferably about 1 mM to about 50 mM; more preferably from about 1 mM to about 20 mM. In one aspect, sodium azide concentration typically ranges from about 20 mM to about 60 mM while hydroxylamine sulfate is concentration is typically about 0.5 mM to about 30 mM, preferably about 10 mM.

The catalase activity in a host cell can be down-regulated or eliminated by disrupting expression of the gene(s) responsible for the catalase activity using well known techniques including, but not limited to, transposon mutagenesis, RNA antisense expression, targeted mutagenesis, and random mutagenesis. In a preferred embodiment, the gene(s) encoding the endogenous catalase activity are down-regulated or disrupted (i.e., “knocked-out”). As used herein, a “disrupted” gene is one where the activity and/or function of the protein encoded by the modified gene is no longer present. Means to disrupt a gene are well-known in the art and may include, but are not limited to, insertions, deletions, or mutations to the gene so long as the activity and/or function of the corresponding protein is no longer present. In a further preferred embodiment, the production host is an *E. coli* production host comprising a disrupted catalase gene selected from the group consisting of *katG* and *katE* (see U.S. Pat. No. 7,951,566 to DiCosimo et al.). In another embodiment, the production host is an *E. coli* strain comprising a down-regulation and/or disruption in both *katG* and *katE* catalase genes. An *E. coli* strain comprising a double-knockout of *katG* and *katE* has been prepared and is described as *E. coli* strain KLP18 (U.S. Pat. No. 7,951,566 to DiCosimo et al.).

The concentration of the catalyst in the aqueous reaction formulation depends on the specific catalytic activity of the catalyst, and is chosen to obtain the desired rate of reaction. The weight of catalyst in perhydrolysis reactions typically ranges from 0.0001 mg to 50 mg per mL of total reaction volume, preferably from 0.0005 mg to 10 mg per mL, more preferably from 0.0010 mg to 2.0 mg per mL. The catalyst may also be immobilized on a soluble or insoluble support using methods well-known to those skilled in the art; see for example, *Immobilization of Enzymes and Cells*; Gordon F. Bickerstaff, Editor; Humana Press, Totowa, N.J., USA; 1997. The use of immobilized catalysts permits the recovery and reuse of the catalyst in subsequent reactions. The enzyme catalyst may be in the form of whole microbial cells, permeabilized microbial cells, microbial cell extracts, partially-purified or purified enzymes, and mixtures thereof.

In one aspect, the concentration of peroxy-carboxylic acid generated by the combination of chemical perhydrolysis and enzymatic perhydrolysis of the carboxylic acid ester is sufficient to provide an effective concentration of peroxy-carboxylic acid for disinfection, bleaching, sanitization, deodorizing or destaining at a desired pH. In another aspect, the peroxy-carboxylic acid is generated at a safe and efficacious concentration suitable for use in a personal care product to be applied to the hair, skin, nails or tissues of the oral cavity, such as tooth enamel, tooth pellicle or the gums. In another aspect, the present methods provide combinations of enzymes and enzyme substrates to produce the desired effective concentration of peroxy-carboxylic acid, where, in the absence of

added enzyme, there is a significantly lower concentration of peroxy-carboxylic acid produced. Although there may be some chemical perhydrolysis of the enzyme substrate by direct chemical reaction of inorganic peroxide with the enzyme substrate, there may not be a sufficient concentration of peroxy-carboxylic acid generated to provide an effective concentration of peroxy-carboxylic acid in the desired applications, and a significant increase in total peroxy-carboxylic acid concentration is achieved by the addition of an appropriate perhydrolase catalyst to the aqueous reaction formulation.

In one aspect of the invention, the concentration of peroxy-carboxylic acid generated (e.g. peracetic acid) by the enzymatic perhydrolysis is at least about 2 ppm, preferably at least 20 ppm, preferably at least 100 ppm, more preferably at least about 200 ppm peroxy-carboxylic acid, more preferably at least 300 ppm, more preferably at least 500 ppm, more preferably at least 700 ppm, more preferably at least about 1000 ppm peroxy-carboxylic acid, more preferably at least about 2000 ppm peroxy-carboxylic acid, most preferably at least 10,000 ppm peroxy-carboxylic acid within 5 minutes more preferably within 1 minute of initiating the enzymatic perhydrolysis reaction. In a second aspect of the invention, the concentration of peroxy-carboxylic acid generated (e.g. peracetic acid) by the enzymatic perhydrolysis is at least about 2 ppm, preferably at least 20 ppm, preferably at least 30 ppm, more preferably at least about 40 ppm peroxy-carboxylic acid, more preferably at least 50 ppm, more preferably at least 60 ppm, more preferably at least 70 ppm, more preferably at least about 80 ppm peroxy-carboxylic acid, most preferably at least 100 ppm peroxy-carboxylic acid within 5 minutes, more preferably within 1 minute, of initiating the enzymatic perhydrolysis reaction (i.e., time measured from combining the reaction components to form the formulation).

The aqueous formulation comprising the peroxy-carboxylic acid may be optionally diluted with diluent comprising water, or a solution predominantly comprised of water, to produce a formulation with the desired lower target concentration of peroxy-carboxylic acid. In one aspect, the reaction time required to produce the desired concentration (or concentration range) of peroxy-carboxylic acid is about 20 minutes or less, preferable about 5 minutes or less, most preferably about 1 minute or less.

In other aspects, the surface or inanimate object contaminated with a concentration of a biological contaminant(s) is contacted with the peroxy-carboxylic acid formed in accordance with the processes described herein within about 1 minute to about 168 hours of combining said reaction components, or within about 1 minute to about 48 hours, or within about 1 minute to 2 hours of combining said reaction components, or any such time interval therein.

In another aspect, the peroxy-carboxylic acid formed in accordance with the processes describe herein is used in a laundry care application wherein the peroxy-carboxylic acid is contacted with clothing or a textile to provide a benefit, such as disinfecting, bleaching, destaining, deodorizing and/or a combination thereof. The peroxy-carboxylic acid may be used in a variety of laundry care products including, but not limited to, laundry or textile pre-wash treatments, laundry detergents or additives, stain removers, bleaching compositions, deodorizing compositions, and rinsing agents. In one embodiment, the present process to produce a peroxy-carboxylic acid for a target surface is conducted in situ.

In the context of laundry care applications, the term “contacting an article of clothing or textile” means that the article of clothing or textile is exposed to a formulation disclosed herein. To this end, there are a number of formats the formulation may be used to treat articles of clothing or textiles

including, but not limited to, liquid, solids, gel, paste, bars, tablets, spray, foam, powder, or granules and can be delivered via hand dosing, unit dosing, dosing from a substrate, spraying and automatic dosing from a laundry washing or drying machine. Granular compositions can also be in compact form; liquid compositions can also be in a concentrated form.

When the formulations disclosed herein are used in a laundry washing machine, the formulation can further contain components typical to laundry detergents. For example, typical components include, but are not limited to, surfactants, bleaching agents, bleach activators, additional enzymes, suds suppressors, dispersants, lime-soap dispersants, soil suspension and anti-redeposition agents, softening agents, corrosion inhibitors, tarnish inhibitors, germicides, pH adjusting agents, non-builder alkalinity sources, chelating agents, organic and/or inorganic fillers, solvents, hydrotropes, optical brighteners, dyes, and perfumes. For example, a laundry detergent powder was used in Example 5 and 6 comprising a formulation generally comprising surfactants (sodium dodecylbenzenesulfonate, C12-15 Pareth-5, C12-15 Pareth-7, sodium stearate, and stearic acid), builders (sodium carbonate, zeolite, sodium silicate, and citric acid), binders (cellulose, PEG-75, dextrin, and sucrose), bulking agents (sodium sulfate, sodium chloride, sodium bicarbonate, and calcium carbonate), structurants (sodium acrylic acid/MA copolymer, and sodium polyacrylate), sequestrants (tetrasodium etidronate and calcium sodium EDTMP), optical brighteners (disodium anilinomorpholinotriazinyl-aminostilbene-sulfonate), stabilizing agents, anti-redeposition agents, antifoaming agents, and softness extenders. The formulations disclosed herein can also be used as detergent additive products in solid or liquid form. Such additive products are intended to supplement or boost the performance of conventional detergent compositions and can be added at any stage of the cleaning process.

In connection with the present systems and methods for laundry care where the peracid is generated for one or more of bleaching, stain removal, and odor reduction, the concentration of peracid generated (e.g., peracetic acid) by the perhydrolysis of at least one carboxylic acid ester may be at least about 2 ppm, preferably at least 20 ppm, more preferably at least 40 ppm, and even more preferably at least about 100 ppm peracid. In connection with the present systems and methods for laundry care where the peracid is generated for disinfection or sanitization, the concentration of peracid generated (e.g., peracetic acid) by the perhydrolysis of at least one carboxylic acid ester may be at least about 40 ppm, more preferably at least 80 ppm, and most preferably at least 100 ppm peracid within 10 minutes, preferably within 5 minutes, and most preferably within 1 minute of initiating the perhydrolysis reaction. The product formulation comprising the peracid may be optionally diluted with water, or a solution predominantly comprised of water, to produce a formulation with the desired lower concentration of peracid. In one aspect of the present methods and systems, the reaction time required to produce the desired concentration of peracid is not greater than about two hours, preferably not greater than about 30 minutes, more preferably not greater than about 10 minutes, even more preferably not greater than about 5 minutes, and most preferably in about 1 minute or less.

The temperature of the reaction is chosen to control both the reaction rate and the stability of the enzyme catalyst activity. The temperature of the reaction may range from just above the freezing point of the aqueous reaction formulation (approximately 0° C.) to about 85° C., with a preferred range of reaction temperature of from about 5° C. to about 75° C.

The pH of the aqueous reaction formulation while enzymatically producing peroxy-carboxylic acid is maintained at a pH ranging from about 5.0 to about 11.5, preferably about 6.5 to about 11.0, and yet even more preferably about 7.5 to about 11.0. In one embodiment, the pH of the aqueous reaction formulation ranges from about 10.5 to about 11.0 for at least 30 minutes after combining the reaction components. The pH of the aqueous reaction formulation may be adjusted or controlled by the addition or incorporation of a suitable buffer, including, but not limited to, phosphate, pyrophosphate, bicarbonate, acetate, or citrate. In one embodiment, the buffer is selected from a phosphate buffer, a bicarbonate buffer, or a buffer formed by the combination of hard water (tap water to simulate laundry care applications) and percarbonate (from sodium percarbonate used to generate hydrogen peroxide). The concentration of buffer, when employed, is typically from 0.1 mM to 1.0 M, preferably from 1 mM to 300 mM, most preferably from 10 mM to 100 mM. In another aspect of the present invention, no buffer is added to the reaction mixture while enzymatically producing peroxy-carboxylic acid.

In yet another aspect, the enzymatic perhydrolysis aqueous reaction formulation may contain an organic solvent that acts as a dispersant to enhance the rate of dissolution of the carboxylic acid ester in the aqueous reaction formulation. Such solvents include, but are not limited to, propylene glycol methyl ether, acetone, cyclohexanone, diethylene glycol butyl ether, tripropylene glycol methyl ether, diethylene glycol methyl ether, propylene glycol butyl ether, dipropylene glycol methyl ether, cyclohexanol, benzyl alcohol, isopropanol, ethanol, propylene glycol, and mixtures thereof.

In another aspect, the enzymatic perhydrolysis product may contain additional components that provide desirable functionality. These additional components include, but are not limited to, buffers, detergent builders, thickening agents, emulsifiers, surfactants, wetting agents, corrosion inhibitors (e.g., benzotriazole), enzyme stabilizers, and peroxide stabilizers (e.g., metal ion chelating agents). Many of the additional components are well known in the detergent industry (see, for example, U.S. Pat. No. 5,932,532; hereby incorporated by reference). Examples of emulsifiers include, but are not limited to, polyvinyl alcohol or polyvinylpyrrolidone. Examples of thickening agents include, but are not limited to, LAPONITE® RD (synthetic layered silicate), corn starch, PVP, CARBOWAX® (polyethylene glycol and/or methoxy-polyethylene glycol), CARBOPOL® (acrylates crosspolymer), CABOSIL® (synthetic amorphous fumed silicon dioxide), polysorbate 20, PVA, and lecithin. Examples of buffering systems include, but are not limited to, sodium phosphate monobasic/sodium phosphate dibasic; sulfamic acid/triethanolamine; citric acid/triethanolamine; tartaric acid/triethanolamine; succinic acid/triethanolamine; and acetic acid/triethanolamine. Examples of surfactants include, but are not limited to, a) non-ionic surfactants such as block copolymers of ethylene oxide or propylene oxide, ethoxylated or propoxylated linear and branched primary and secondary alcohols, and aliphatic phosphine oxides; b) cationic surfactants such as quaternary ammonium compounds, particularly quaternary ammonium compounds having a C8-C20 alkyl group bound to a nitrogen atom additionally bound to three C1-C2 alkyl groups; c) anionic surfactants such as alkane carboxylic acids (e.g., C8-C20 fatty acids), alkyl phosphonates, alkane sulfonates (e.g., sodium dodecylsulfate "SDS") or linear or branched alkyl benzene sulfonates, alkene sulfonates; and d) amphoteric and zwitterionic surfactants such as aminocarboxylic acids, aminodicarboxylic acids, alkybetaines, and mixtures thereof. Additional components may include fragrances, dyes, stabilizers of hydrogen

peroxide (e.g., metal chelators such as 1-hydroxyethylidene-1,1-diphosphonic acid (DEQUEST® 2010, Solutia Inc., St. Louis, Mo.) and ethylenediaminetetraacetic acid (EDTA)), TURPINAL® SL (etidronic acid), DEQUEST® 0520 (phosphonate), DEQUEST® 0531 (phosphonate), stabilizers of enzyme activity (e.g., polyethylene glycol (PEG)), and detergent builders.

In another aspect, the enzymatic perhydrolysis product may be pre-mixed to generate the desired concentration of peroxycarboxylic acid prior to contacting the surface or inanimate object to be disinfected.

In another aspect, the enzymatic perhydrolysis product is not pre-mixed to generate the desired concentration of peroxycarboxylic acid prior to contacting the surface or inanimate object to be disinfected, but instead the components of the aqueous reaction formulation that generate the desired concentration of peroxycarboxylic acid are contacted with the surface or inanimate object to be disinfected and/or bleached or destained, generating the desired concentration of peroxycarboxylic acid. In some embodiments, the components of the aqueous reaction formulation combine or mix at the locus. In some embodiments, the reaction components are delivered or applied to the locus and subsequently mix or combine to generate the desired concentration of peroxycarboxylic acid.

Production of Peroxycarboxylic Acids Using a Perhydrolase Catalyst

The peroxycarboxylic acids, once produced, are quite reactive and may decrease in concentration over extended periods of time, depending on variables that include, but are not limited to, temperature and pH. As such, it may be desirable to keep the various reaction components separated, especially for liquid formulations. In one aspect, the hydrogen peroxide source is separate from either the substrate or the perhydrolase catalyst, preferably from both. This can be accomplished using a variety of techniques including, but not limited to, the use of multicompartiment chambered dispensers (U.S. Pat. No. 4,585,150) and at the time of use physically combining the perhydrolase catalyst with a source of peroxygen (such as hydrogen peroxide) and the present substrates to initiate the aqueous enzymatic perhydrolysis reaction. The perhydrolase catalyst may optionally be immobilized within the body of reaction chamber or separated (e.g., filtered, etc.) from the reaction product comprising the peroxycarboxylic acid prior to contacting the surface and/or object targeted for treatment. The perhydrolase catalyst may be in a liquid matrix or in a solid form (e.g., powder or tablet) or embedded within a solid matrix that is subsequently mixed with the substrates to initiate the enzymatic perhydrolysis reaction. In a further aspect, the perhydrolase catalyst may be contained within a dissolvable or porous pouch that may be added to the aqueous substrate matrix to initiate enzymatic perhydrolysis. In yet a further aspect, the perhydrolase catalyst may comprise the contents contained within a separate compartment of a dissolvable or porous pouch that has at least one additional compartment for the containment contents comprising the ester substrate and/or source of peroxide. In an additional further aspect, a powder comprising the enzyme catalyst is suspended in the substrate (e.g., triacetin), and at time of use is mixed with a source of peroxygen in water.

Method for Determining the Concentration of Peroxycarboxylic Acid and Hydrogen Peroxide.

A variety of analytical methods can be used in the present method to analyze the reactants and products including, but not limited to, titration, high performance liquid chromatography (HPLC), gas chromatography (GC), mass spectrometry (MS), capillary electrophoresis (CE), the HPLC analyti-

cal procedure described by U. Karst et al. (*Anal. Chem.*, 69(17):3623-3627 (1997)), and the 2,2'-azino-bis(3-ethylbenzothiazoline)-6-sulfonate (ABTS) assay (see U. Pinkernell et al., *The Analyst* 122:567-571(1997), S. Minning, et al., *Analytica Chimica Acta* 378:293-298 (1999) and WO 2004/058961 A1) as described in U.S. Pat. No. 7,951,566.

Determination of Minimum Biocidal Concentration of Peroxycarboxylic Acids

The method described by J. Gabrielson et al. (*J. Microbiol. Methods* 50: 63-73 (2002)) can be employed for determination of the Minimum Biocidal Concentration (MBC) of peroxycarboxylic acids, or of hydrogen peroxide and enzyme substrates. The assay method is based on XTT reduction inhibition, where XTT (2,3-bis[2-methoxy-4-nitro-5-sulfophenyl]-5-[(phenylamino)carbonyl]-2H-tetrazolium, inner salt, monosodium salt) is a redox dye that indicates microbial respiratory activity by a change in optical density (OD) measured at 490 nm or 450 nm. However, there are a variety of other methods available for testing the activity of disinfectants and antiseptics including, but not limited to, viable plate counts, direct microscopic counts, dry weight, turbidity measurements, absorbance, and bioluminescence (see, for example Brock, Semour S., *Disinfection, Sterilization, and Preservation*, 5th edition, Lippincott Williams & Wilkins, Philadelphia, Pa., USA; 2001).

Uses of Enzymatically Prepared Peroxycarboxylic Acid Compositions

The enzyme catalyst-generated peroxycarboxylic acid produced according to the present method can be used in a variety of hard surface/inanimate object applications for reduction of concentrations of biological contaminants, such as decontamination of medical instruments (e.g., endoscopes), textiles (such as garments and carpets), food preparation surfaces, food storage and food-packaging equipment, materials used for the packaging of food products, chicken hatcheries and grow-out facilities, animal enclosures, and spent process waters that have microbial and/or virucidal activity. The enzyme-generated peroxycarboxylic acids may be used in formulations designed to inactivate prions (e.g., certain proteases) to additionally provide biocidal activity (see U.S. Pat. No. 7,550,420 to DiCosimo et al.).

In one aspect, the peroxycarboxylic acid composition is useful as a disinfecting agent for non-autoclavable medical instruments and food packaging equipment. As the peroxycarboxylic acid-containing formulation may be prepared using GRAS (generally recognized as safe) or food-grade components (enzyme, enzyme substrate, hydrogen peroxide, and buffer), the enzyme-generated peroxycarboxylic acid may also be used for decontamination of animal carcasses, meat, fruits and vegetables, or for decontamination of prepared foods. The enzyme-generated peroxycarboxylic acid may be incorporated into a product whose final form is a powder, liquid, gel, film, solid or aerosol. The enzyme-generated peroxycarboxylic acid may be diluted to a concentration that still provides an efficacious decontamination.

The compositions comprising an efficacious concentration of peroxycarboxylic acid can be used to disinfect surfaces and/or objects contaminated (or suspected of being contaminated) with biological contaminants, such as pathogenic microbial contaminants, by contacting the surface or object with the products produced by the present processes. As used herein, "contacting" refers to placing a disinfecting composition comprising an effective concentration of peroxycarboxylic acid in contact with the surface or inanimate object suspected of contamination with a biological contaminant for a period of time sufficient to clean and disinfect. Contacting includes spraying, treating, immersing, flushing, pouring on

or in, mixing, combining, painting, coating, applying, affixing to and otherwise communicating a peroxycarboxylic acid solution or composition comprising an efficacious concentration of peroxycarboxylic acid, or a solution or composition that forms an efficacious concentration of peroxycarboxylic acid, with the surface or inanimate object suspected of being contaminated with a concentration of a biological contaminant. The disinfectant compositions may be combined with a cleaning composition to provide both cleaning and disinfection. Alternatively, a cleaning agent (e.g., a surfactant or detergent) may be incorporated into the formulation to provide both cleaning and disinfection in a single composition.

The compositions comprising an efficacious concentration of peroxycarboxylic acid can also contain at least one additional antimicrobial agent, combinations of prion-degrading proteases, a virucide, a sporicide, or a biocide. Combinations of these agents with the peroxycarboxylic acid produced by the claimed processes can provide for increased and/or synergistic effects when used to clean and disinfect surfaces and/or objects contaminated (or suspected of being contaminated) with biological contaminants. Suitable antimicrobial agents include carboxylic esters (e.g., p-hydroxy alkyl benzoates and alkyl cinnamates); sulfonic acids (e.g., dodecylbenzene sulfonic acid); iodo-compounds or active halogen compounds (e.g., elemental halogens, halogen oxides (e.g., NaOCl, HOCl, HOBr, ClO₂), iodine, interhalides (e.g., iodine monochloride, iodine dichloride, iodine trichloride, iodine tetrachloride, bromine chloride, iodine monobromide, or iodine dibromide), polyhalides, hypochlorite salts, hypochlorous acid, hypobromite salts, hypobromous acid, chloro- and bromo-hydantoin, chlorine dioxide, and sodium chlorite); organic peroxides including benzoyl peroxide, alkyl benzoyl peroxides, ozone, singlet oxygen generators, and mixtures thereof; phenolic derivatives (e.g., o-phenyl phenol, o-benzyl-p-chlorophenol, tert-amyl phenol and C₁-C₆ alkyl hydroxy benzoates); quaternary ammonium compounds (e.g., alkyldimethylbenzyl ammonium chloride, dialkyldimethyl ammonium chloride and mixtures thereof); and mixtures of such antimicrobial agents, in an amount sufficient to provide the desired degree of microbial protection. Effective amounts of antimicrobial agents include about 0.001 wt % to about 60 wt % antimicrobial agent, about 0.01 wt % to about 15 wt % antimicrobial agent, or about 0.08 wt % to about 2.5 wt % antimicrobial agent.

In one aspect, the peroxycarboxylic acids formed by the process can be used to reduce the concentration of viable biological contaminants (such as a microbial population) when applied on and/or at a locus. As used herein, a "locus" comprises part or all of a target surface suitable for disinfecting or bleaching. Target surfaces include all surfaces that can potentially be contaminated with biological contaminants. Non-limiting examples include equipment surfaces found in the food or beverage industry (such as tanks, conveyors, floors, drains, coolers, freezers, equipment surfaces, walls, valves, belts, pipes, drains, joints, crevasses, combinations thereof, and the like); building surfaces (such as walls, floors and windows); non-food-industry related pipes and drains, including water treatment facilities, pools and spas, and fermentation tanks; hospital or veterinary surfaces (such as walls, floors, beds, equipment (such as endoscopes), clothing worn in hospital/veterinary or other healthcare settings, including clothing, scrubs, shoes, and other hospital or veterinary surfaces); restaurant surfaces; bathroom surfaces; toilets; clothes and shoes; surfaces of barns or stables for livestock, such as poultry, cattle, dairy cows, goats, horses and pigs; hatcheries for poultry or for shrimp; and pharmaceutical or biopharmaceutical surfaces (e.g., pharmaceutical or biop-

harmaceutical manufacturing equipment, pharmaceutical or biopharmaceutical ingredients, pharmaceutical or biopharmaceutical excipients). Additional hard surfaces include food products, such as beef, poultry, pork, vegetables, fruits, seafood, combinations thereof, and the like. The locus can also include water absorbent materials such as linens or other textiles. The locus also includes harvested plants or plant products including seeds, corms, tubers, fruit, and vegetables, growing plants, and especially crop growing plants, including cereals, leaf vegetables and salad crops, root vegetables, legumes, berried fruits, citrus fruits, and hard fruits.

Non-limiting examples of hard surface materials are metals (e.g., steel, stainless steel, chrome, titanium, iron, copper, brass, aluminum, and alloys thereof), minerals (e.g., concrete), polymers and plastics (e.g., polyolefins, such as polyethylene, polypropylene, polystyrene, poly(meth)acrylate, polyacrylonitrile, polybutadiene, poly(acrylonitrile, butadiene, styrene), poly(acrylonitrile, butadiene), acrylonitrile butadiene; polyesters such as polyethylene terephthalate; and polyamides such as nylon). Additional surfaces include brick, tile, ceramic, porcelain, wood, wood pulp, paper, vinyl, linoleum, and carpet.

The peroxycarboxylic acids formed by the present process may be used to provide a benefit to an article of clothing or a textile including, but not limited to, disinfecting, sanitizing, bleaching, destaining, and deodorizing. The peroxycarboxylic acids formed by the present process may be used in any number of laundry care products including, but not limited to, textile pre-wash treatments, laundry detergents, laundry detergents or additives, stain removers, bleaching compositions, deodorizing compositions, and rinsing agents, to name a few.

The peroxycarboxylic acids formed by the present process can be used in one or more steps of the wood pulp or paper pulp bleaching/delignification process, particularly where peracetic acid is used (for example, see EP1040222 B1 and U.S. Pat. No. 5,552,018 to Devenyns, J.).

Recombinant Microbial Expression

The genes and gene products of the instant sequences may be produced in heterologous host cells, particularly in the cells of microbial hosts. Preferred heterologous host cells for expression of the instant genes and nucleic acid molecules are microbial hosts that can be found within the fungal or bacterial families and which grow over a wide range of temperature, pH values, and solvent tolerances. For example, it is contemplated that any of bacteria, yeast, and filamentous fungi may suitably host the expression of the present nucleic acid molecules. The perhydrolase may be expressed intracellularly, extracellularly, or a combination of both intracellularly and extracellularly, where extracellular expression renders recovery of the desired protein from a fermentation product more facile than methods for recovery of protein produced by intracellular expression. Transcription, translation and the protein biosynthetic apparatus remain invariant relative to the cellular feedstock used to generate cellular biomass; functional genes will be expressed regardless. Examples of host strains include, but are not limited to, bacterial, fungal or yeast species such as *Aspergillus*, *Trichoderma*, *Saccharomyces*, *Pichia*, *Phaffia*, *Kluyveromyces*, *Candida*, *Hansenula*, *Yarrowia*, *Salmonella*, *Bacillus*, *Acinetobacter*, *Zymomonas*, *Agrobacterium*, *Erythrobacter*, *Chlorobium*, *Chromatium*, *Flavobacterium*, *Cytophaga*, *Rhodobacter*, *Rhodococcus*, *Streptomyces*, *Brevibacterium*, *Corynebacteria*, *Mycobacterium*, *Deinococcus*, *Escherichia*, *Erwinia*, *Pantoea*, *Pseudomonas*, *Sphingomonas*, *Methylobacter*, *Methylobacter*, *Methylococcus*, *Methylosinus*, *Methylomicrobium*, *Methylocystis*, *Alcaligenes*, *Synechocys-*

tis, *Synechococcus*, *Anabaena*, *Thiobacillus*, *Methanobacterium*, *Klebsiella*, and *Myxococcus*. In one embodiment, bacterial host strains include *Escherichia*, *Bacillus*, and *Pseudomonas*. In a preferred embodiment, the bacterial host cell is *Bacillus subtilis* or *Escherichia coli*.

Industrial Production

A variety of culture methodologies may be applied to produce the perhydrolase catalyst. Large-scale production of a specific gene product over expressed from a recombinant microbial host may be produced by batch, fed-batch or continuous culture methodologies. Batch and fed-batch culturing methods are common and well known in the art and examples may be found in Thomas D. Brock in *Biotechnology: A Textbook of Industrial Microbiology*, Second Edition, Sinauer Associates, Inc., Sunderland, Mass. (1989) and Deshpande, Mukund V., *Appl. Biochem. Biotechnol.*, 36:227 (1992).

In one embodiment, commercial production of the desired perhydrolase catalyst is accomplished with a continuous culture. Continuous cultures are an open system where a defined culture media is added continuously to a bioreactor and an equal amount of conditioned media is removed simultaneously for processing. Continuous cultures generally maintain the cells at a constant high liquid phase density where cells are primarily in log phase growth. Alternatively, continuous culture may be practiced with immobilized cells where carbon and nutrients are continuously added and valuable products, by-products or waste products are continuously removed from the cell mass. Cell immobilization may be performed using a wide range of solid supports composed of natural and/or synthetic materials.

Recovery of the desired perhydrolase catalysts from a batch or fed-batch fermentation, or continuous culture may be accomplished by any of the methods that are known to those skilled in the art. For example, when the enzyme catalyst is produced intracellularly, the cell paste is separated from the culture medium by centrifugation or membrane filtration, optionally washed with water or an aqueous buffer at a desired pH, then a suspension of the cell paste in an aqueous buffer at a desired pH is homogenized to produce a cell extract containing the desired enzyme catalyst. The cell extract may optionally be filtered through an appropriate filter aid such as celite or silica to remove cell debris prior to a heat-treatment step to precipitate undesired protein from the enzyme catalyst solution. The solution containing the desired enzyme catalyst may then be separated from the precipitated cell debris and protein produced during the heat-treatment step by membrane filtration or centrifugation, and the resulting partially-purified enzyme catalyst solution concentrated by additional membrane filtration, then optionally mixed with an appropriate excipient (for example, maltodextrin, trehalose, sucrose, lactose, sorbitol, mannitol, phosphate buffer, citrate buffer, or mixtures thereof) and spray-dried to produce a solid powder comprising the desired enzyme catalyst. Alternatively, the resulting partially-purified enzyme catalyst solution prepared as described above may be optionally concentrated by additional membrane filtration, and the partially-purified enzyme catalyst solution or resulting enzyme concentrate is then optionally mixed with one or more stabilizing agents (e.g., glycerol, sorbitol, propylene glycol, 1,3-propanediol, polyols, polymeric polyols, polyvinylalcohol or mixtures thereof), one or more salts (e.g., sodium chloride, sodium sulfate, potassium chloride, potassium sulfate, or mixtures thereof), and one or more biocides, and maintained as an aqueous solution until used.

When an amount, concentration, or other value or parameter is given either as a range, preferred range, or a list of upper preferable values and lower preferable values, this is to

be understood as specifically disclosing all ranges formed from any pair of any upper range limit or preferred value and any lower range limit or preferred value, regardless of whether ranges are separately disclosed. Where a range of numerical values is recited herein, unless otherwise stated, the range is intended to include the endpoints thereof, and all integers and fractions within the range. It is not intended that the scope be limited to the specific values recited when defining a range.

General Methods

The following examples are provided to demonstrate different embodiments. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the methods disclosed herein, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the presently disclosed methods.

All reagents and materials were obtained from DIFCO Laboratories (Detroit, Mich.), GIBCO/BRL (Gaithersburg, Md.), TCI America (Portland, Oreg.), Roche Diagnostics Corporation (Indianapolis, Ind.) or Sigma-Aldrich Chemical Company (St. Louis, Mo.), unless otherwise specified.

The following abbreviations in the specification correspond to units of measure, techniques, properties, or compounds as follows: "sec" or "s" means second(s), "min" means minute(s), "h" or "hr" means hour(s), "μL" means microliter(s), "mL" means milliliter(s), "L" means liter(s), "mM" means millimolar, "M" means molar, "mmol" means millimole(s), "ppm" means part(s) per million, "wt" means weight, "wt %" means weight percent, "g" means gram(s), "μg" means microgram(s), "ng" means nanogram(s), "g" means gravity, "HPLC" means high performance liquid chromatography, "dd H₂O" means distilled and deionized water, "dcw" means dry cell weight, "ATCC" or "ATCC®" means the American Type Culture Collection (Manassas, Va.), "U" means unit(s) of perhydrolase activity, "rpm" means revolution(s) per minute, "EDTA" means ethylenediaminetetraacetic acid, "IPTG" means isopropyl-β-D-thio-galactoside, "DTT" means dithiothreitol, "BCA" means bicinchoninic acid, and "ABTS" means 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid).

Example 1

Construction of a Random Mutagenesis Library of *Thermotoga maritima* Acetyl Xylan Esterase C277T Variant

The coding sequence of a *Thermotoga maritima* acetyl xylan esterase (GENBANK® accession #NP_227893.1) was synthesized using codons optimized for expression in *E. coli* (DNA 2.0, Menlo Park, Calif.) and cloned into pUC19 between PstI and XbaI to create the plasmid known as pSW202 (U.S. Patent Application Publication No. 2008-0176299). The codon-optimized sequence is provided as SEQ ID NO: 1 encoding the wild-type *Thermotoga maritima* acetyl xylan esterase provided as SEQ ID NO: 2.

A codon change was made in the gene using primer pairs identified as SEQ ID NO: 3 and SEQ ID NO: 4, and the QUIKCHANGE® site-directed mutagenesis kit (Stratagene, La Jolla, Calif.), according to the manufacturer's instructions,

resulting in the amino acid change C277T (SEQ ID NO: 5), to create the plasmid known as pSW202/C277T (SEQ ID NO: 6). Plasmid pSW202/C277T served as a template for error-prone PCR using primers identified as SEQ ID NO: 7 and SEQ ID NO: 8, and the GENEMORPH® II random mutagenesis kit (Stratagene), according to the manufacturer's recommendations. The resulting PCR product was digested with Pst1 and Xba1 and ligated with pUC19 also digested with Pst1 and Xba1. *E. coli* KLP18 (see U.S. Patent Application Publication No. 2008-0176299) was transformed with the ligation mixture and plated onto LB plates supplemented with 0.1 mg ampicillin/mL. Nucleotide sequencing of a random sample indicated a mutation frequency of 2-8 changes per PCR product.

Example 2

Screening of *Thermotoga maritima* Error-Prone PCR Library for Increased Enzyme Activity in the Presence of Laundry Detergent

Colonies were picked (automated) and placed into 96-well "master plates" containing 0.1 mL LB media supplemented with 0.1 mg ampicillin/mL and grown 16-18 h at 37° C. and 80% humidity. From each well of the master plates, 0.003 mL of culture was transferred to 96-well "induction plates" containing 0.3 mL LB media supplemented with 0.1 mg ampicillin/mL and 0.5 mM IPTG, which were incubated for 16-18 h with shaking at 37° C. and 80% humidity. Separately, 0.1 mL of 50% glycerol was added to each well of the master plates, which were stored at -80° C. as stocks. Induction plates were centrifuged, decanted, and cell pellets frozen at -20° C. Cell pellets were thawed and 0.075 mL of 25.2 mg/mL CELLYTIC™ Express (Sigma Aldrich, St. Louis,

Mo.) was added to each cell pellet to lyse the cells. After incubation for 30 minutes with shaking at ambient temperature, 0.525 mL of distilled/deionized water was added to each well. The cell lysates were further diluted 1:1 with distilled/deionized water, after which 0.010 mL was transferred to 96-well "assay plates" containing 0.050 mL of 4 g/L of a laundry detergent (powder) having a formulation generally comprising surfactants (sodium dodecylbenzenesulfonate, C12-15 Pareth-5, C12-15 Pareth-7, sodium stearate, and stearic acid), builders (sodium carbonate, zeolite, sodium silicate, and citric acid), binders (cellulose, PEG-75, dextrin, and sucrose), bulking agents (sodium sulfate, sodium chloride, sodium bicarbonate, and calcium carbonate), structurants (sodium acrylic acid/MA copolymer, and sodium polyacrylate), sequestrants (tetrasodium etidronate and calcium sodium EDTMP), optical brighteners (disodium anilino-morpholinotriazinyl-aminostilbenesulfonate), stabilizing agents, anti-redeposition agents, antifoaming agents, and softness extenders and 0.030 mL of 33 mM hydrogen peroxide. To each well of the assay plates was then added 0.010 mL of 50 mM triacetin. To each well of the assay plate was added 0.1 mL of "stop buffer" (100 mM o-phenylenediamine and 500 mM sodium dihydrogen phosphate, pH 2.0). The plates were incubated for 30 minutes at 30° C., after which absorbance at 458 nm was read. *T. maritima* WT (codon optimized gene (SEQ ID NO: 1) encoding the wild type enzyme (SEQ ID NO: 2)) and *T. maritima* C277T (SEQ ID NO: 5) were incorporated into each plate as controls. Screening approximately 7500 colonies resulted in the identification of numerous "hits" demonstrating activity significantly greater than *T. maritima* C277T (SEQ ID NO: 5). Nucleotide sequencing was used to determine the amino acid changes in the perhydrolase enzyme of these hits (Table 1).

TABLE 1

T. maritima perhydrolase variants identified in primary screen as having improved perhydrolytic activity relative to the C277T perhydrolase control when screened in the presence of a laundry detergent.

Variant ID	Amino acid changes relative to <i>T. maritima</i> WT sequence SEQ ID NO: 2						Amino acid sequence (SEQ ID NO:)
007C7	F253S	C277T					9
007A11	F161L	C277T					10
007A12	E17D	K77R	F174L	C277T			11
007H10	D25G	C277T	G307S				12
007A5	N257D	C277T					13
007A8	Q67R	C277T					14
007B12	L32P	C277T					15
007B2	R15H	Q176H	L208Q	C277T			16
007B3	P16L	F38S	T53N	R159C	V183D	C277T	17
007B5	A266G	C277T					18
007B7	H100N	W105S	C277T				19
007C4	R261S	I264F	C277T				20
007C3	D178N	C277T					21
007C8	M47V	E134D	L150Q	Y252F	C277T		22
007D1	V270A	C277T					23
007D11	E55K	C277T					24
007D12	R261H	C277T					25
007D2	A206T	C277T					26
007D5	C277T	R296P					27
007D6	F44S	C277T					28
007D8	E9K	G255C	C277T				29
007E11	M47K	R158H	Q179K	C277T			30
007E6	E17V	R261H	C277T				31
007E9	C215S	C277T					32
007F10	E244D	V270I	C277T				33
007G11	V137L	C277T					34
007G5	V256I	C277T					35
007G6	F61L	C277T					36
007G8	K82R	E180K	C277T				37

TABLE 1-continued

<i>T. maritima</i> perhydrolase variants identified in primary screen as having improved perhydrolytic activity relative to the C277T perhydrolase control when screened in the presence of a laundry detergent.		
Variant ID	Amino acid changes relative to <i>T. maritima</i> WT sequence SEQ ID NO: 2	Amino acid sequence (SEQ ID NO:)
Control (C277T)	C277T	5

Example 3

Production of Variant *Thermotoga maritima* Perhydrolases

Variant strains identified in the screens were grown in LB media at 37° C. with shaking up to OD_{600 nm}=0.4-0.5, at which time IPTG was added to a final concentration of 1 mM, and incubation continued for 2-3 h. Cells were harvested by centrifugation and SDS-PAGE was performed to confirm expression of the perhydrolase enzyme at 15-30% of total soluble protein.

Example 4

Preparation of Heat-Treated Cell Extracts Containing Semi-Purified Variant *Thermotoga maritima* Perhydrolases

Cell cultures (prepared as described in Example 3) were harvested by centrifugation at 5,000×g for 15 minutes then resuspended (20% w/v) in 50 mM phosphate buffer pH 7.0 supplemented with 1.0 mM DTT. Resuspended cells were passed through a French pressure cell twice to ensure >95% cell lysis. Lysed cells were centrifuged for 30 minutes at 12,000×g, and the supernatant was heated at 75° C. for 20 minutes, followed by quenching in an ice bath for 2 minutes. Precipitated protein was removed by centrifugation for 10 minutes at 11,000×g. The resulting heat-treated extract supernatants were analyzed by SDS-PAGE and BCA protein assay, and stored frozen at -80° C.

Example 5

Comparison of *Thermotoga maritima* Perhydrolase Variants vs. *Thermotoga maritima* C277T Perhydrolase Under Simulated Laundry Care Application Conditions

Nine error-prone PCR (epPCR)-generated variant perhydrolases were evaluated in duplicate reactions at 25° C. for

peracetic acid (PAA) production under laundry application conditions (10 mL total volume, 1 mM triacetin (TA), 4 mM H₂O₂, 1.0 µg/mL of heat-treated extract total soluble protein containing an error-prone PCR (epPCR)-generated variant perhydrolase prepared as described in Example 4) using 2 g/L of a laundry detergent (powder) having a formulation generally comprising surfactants (sodium dodecylbenzenesulfonate, C12-15 Pareth-5, C12-15 Pareth-7, sodium stearate, and stearic acid), builders (sodium carbonate, zeolite, sodium silicate, and citric acid), binders (cellulose, PEG-75, dextrin, and sucrose), bulking agents (sodium sulfate, sodium chloride, sodium bicarbonate, and calcium carbonate), structurants (sodium acrylic acid/MA copolymer, and sodium polyacrylate), sequestrants (tetrasodium etidronate and calcium sodium EDTMP), optical brighteners (disodium anilino-morpholinotriazinyl-aminostilbenesulfonate), stabilizing agents, anti-redeposition agents, antifoaming agents, and softness extenders, and compared to PAA production using the same concentration of C277T variant perhydrolase (Published U.S. Patent Application No. 2010-0087529 to DiCosimo et al., SEQ ID NO: 5). Peracid concentrations were determined using an ABTS assay as described by U. Pinkernell et al., *The Analyst* 122:567-571(1997). The results of the various comparisons under simulated laundry care conditions are provided in Table 2. Under these reaction conditions, variants 007C7 (F253S/C277T, SEQ ID NO: 9), 007A11 (F161L/C277T, SEQ ID NO: 10), 007A12 (E17D/K77R/F174L/C277T, SEQ ID NO: 11), 007A5 (N257D/C277T, SEQ ID NO: 13), 007B2 (R15H/Q176H/L208Q/C277T, SEQ ID NO: 16), 007B5 (A266G/C277T, SEQ ID NO: 18), 007D1 (V270A/C277T, SEQ ID NO: 23), 007D5 (C277T/R296P, SEQ ID NO: 27) and 007E9 (C215S/C277T, SEQ ID NO: 32) show a measurable improvement in specific activity for peracetic acid production when compared to the *T. maritima* C277T perhydrolase (SEQ ID NO: 5) under the specified conditions with added powder detergent at pH 10.7.

TABLE 2

Peracetic acid (PAA) production at 5 minutes and 10 minutes from perhydrolase variants vs. <i>T. maritima</i> C277T perhydrolase at 25° C., 1.0 µg/mL heat-treated extract total soluble protein and 2.0 mg/mL laundry detergent powder.									
Variant ID	SEQ ID NO:	Samples	Triacetin (mM)	H ₂ O ₂ (mM)	Total soluble protein (µg/mL)	detergent (mg/mL)	Initial pH	PAA (ppm); 5 min.	PAA (ppm); 10 min
		no perhydrolase	1	4	0	2	10.7	10.5	14.1
Control	5	C277T	1	4	1.0	2	10.7	18.6	23.1
Control	5	C277T	1	4	1.0	2	10.7	18.4	23.6
007C7	9	F253S/C277T	1	4	1.0	2	10.7	26.2	32.1
007C7	9	F253S/C277T	1	4	1.0	2	10.7	25.5	32.2
007A11	10	F161L/C277T	1	4	1.0	2	10.7	21.6	27.7
007A11	10	F161L/C277T	1	4	1.0	2	10.7	22.5	26.3

TABLE 2-continued

Peracetic acid (PAA) production at 5 minutes and 10 minutes from perhydrolase variants vs. <i>T. maritima</i> C277T perhydrolase at 25° C., 1.0 µg/mL heat-treated extract total soluble protein and 2.0 mg/mL laundry detergent powder.									
Variant ID	SEQ ID NO:	Samples	Triacetin (mM)	H ₂ O ₂ (mM)	Total soluble protein (µg/mL)	detergent powder (mg/mL)	Initial pH	PAA (ppm); 5 min.	PAA (ppm); 10 min
007A12	11	E17D/K77R/F174L/C277T	1	4	1.0	2	10.7	21.1	27.1
007A12	11	E17D/K77R/F174L/C277T	1	4	1.0	2	10.7	22.2	28.8
007A5	13	N257D/C277T	1	4	1.0	2	10.7	20.5	29.0
007A5	13	N257D/C277T	1	4	1.0	2	10.7	21.1	25.3
007B2	16	R15H/Q176H/L208Q/C277T	1	4	1.0	2	10.7	22.7	29.0
007B2	16	R15H/Q176H/L208Q/C277T	1	4	1.0	2	10.7	24.3	30.8
007B5	18	A266G/C277T	1	4	1.0	2	10.7	23.2	29.4
007B5	18	A266G/C277T	1	4	1.0	2	10.7	23.6	29.5
007D1	23	V270A/C277T	1	4	1.0	2	10.7	23.3	29.9
007D1	23	V270A/C277T	1	4	1.0	2	10.7	22.6	28.7
007D5	27	C277T/R296P	1	4	1.0	2	10.7	26.5	33.6
007D5	27	C277T/R296P	1	4	1.0	2	10.7	26.5	32.9
007E9	32	C215S/C277T	1	4	1.0	2	10.7	20.7	27.9
007E9	32	C215S//C277T	1	4	1.0	2	10.7	20.2	27.0

Example 6

Comparison of a *Thermotoga maritima* Perhydrolase Variant vs. *Thermotoga maritima* C277T Perhydrolase Under Simulated Laundry Care Application Conditions

An error-prone PCR (epPCR)-generated variant perhydrolase was evaluated for peracetic acid (PAA) production at 25° C. under laundry application conditions (10 mL total volume, 0.75 mM triacetin (TA), 1.4 mM H₂O₂, 0.5 µg/mL or 4.0 µg/mL of heat-treated extract total soluble protein containing an error-prone PCR (epPCR)-generated variant perhydrolase prepared as described in Example 4) using 2 g/L of a laundry detergent (powder) having a formulation generally comprising surfactants (sodium dodecylbenzenesulfonate, C12-15 Pareth-5, C12-15 Pareth-7, sodium stearate, and stearic acid), builders (sodium carbonate, zeolite, sodium silicate, and citric acid), binders (cellulose, PEG-75, dextrin, and sucrose),

bulking agents (sodium sulfate, sodium chloride, sodium bicarbonate, and calcium carbonate), structurants (sodium acrylic acid/MA copolymer, and sodium polyacrylate), sequestrants (tetrasodium etidronate and calcium sodium EDTMP), optical brighteners (disodium anilinomorpholinotriazinyl-aminostilbenesulfonate), stabilizing agents, anti-redeposition agents, antifoaming agents, and softness extenders, and compared to PAA production using the same concentration of C277T variant perhydrolase (Published U.S. Patent Application No. 2010-0087529 to DiCosimo et al., SEQ ID NO: 5). Peracid concentrations were determined using an ABTS assay as described by U. Pinkernell et al., *The Analyst* 122:567-571(1997). The results of the various comparisons under simulated laundry care conditions are provided in Table 3. Under these reaction conditions, variant 007A8 (Q67R/C277T, SEQ ID NO: 14) showed a measurable improvement in specific activity for peracetic acid production when compared to the *T. maritima* C277T perhydrolase (SEQ ID NO: 5) under the specified conditions with added powder detergent at pH 10.7.

TABLE 3

Peracetic acid (PAA) production at 3 minutes and 10 minutes from variant 007A8 vs. <i>T. maritima</i> C277T at 25° C., 0.5 µg/mL or 4.0 µg/mL heat-treated extract total soluble protein and 2.0 mg/mL laundry detergent powder.									
Variant ID	SEQ ID NO:	Samples	Triacetin (mM)	H ₂ O ₂ (mM)	Total soluble protein (µg/mL)	detergent powder (mg/mL)	Initial pH	PAA (ppm); 3 min.	PAA (ppm); 10 min.
		no perhydrolase	0.75	1.4	0	2	10.7	2.9	5.5
Control	5	C277T	0.75	1.4	0.5	2	10.7	5.6	11.5
Control	5	C277T	0.75	1.4	0.5	2	10.7	5.8	11.2
007A8	14	Q67R/C277T	0.75	1.4	0.5	2	10.7	6.9	16.0
007A8	14	Q67R/C277T	0.75	1.4	0.5	2	10.7	7.3	16.2
Control	5	C277T	0.75	1.4	4.0	2	10.7	17.1	23.7
Control	5	C277T	0.75	1.4	4.0	2	10.7	14.7	23.7
007A8	14	Q67R/C277T	0.75	1.4	4.0	2	10.7	22.9	22.1
007A8	14	Q67R/C277T	0.75	1.4	4.0	2	10.7	25.3	24.4

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 47

<210> SEQ ID NO 1

<211> LENGTH: 978

<212> TYPE: DNA

<213> ORGANISM: *Thermotoga maritima*

<400> SEQUENCE: 1

```

atggccttct tcgatttacc actcgaagaa ctgaagaaat atcgctcaga gcggtacgaa      60
gagaaagact tcgatgagtt ctgggaagag aactcgcag agagcgaaaa gttcccctta      120
gaccccgctct tcgagaggat ggagtctcac ctcaaacag tcaagcgta cgatgtcacc      180
ttctccggat acaggggaca gaggatcaaa ggggtggctcc ttgttccaaa actggaagaa      240
gaaaaacttc cctgcgttgt gcagtacata ggatacaacg gtggaagagg attccctcac      300
gactggctgt tctggccttc tatgggttac atatgtttcg tcatggatac tcgaggtcag      360
ggaagcggct ggctgaaagg agacacaccg gattaccctg aggggtcccgt tgaccctcag      420
tatccaggat tcatgacaag aggaatactg gatcccagaa cttactacta cagacgagtc      480
ttcacggacg ctgtcagagc cgttgaagct gctgcttctt ttctcaggt agatcaagaa      540
agaatcgtga tagctggagg cagtcagggt ggcggaatag cccttgcggt gagcgctctc      600
tcaaagaaag caaaggctct tctgtgcgat gtgccgttcc tgtgtcactt cagaagagca      660
gtacagcttg tggatacgca tccatacgcg gagatcacga actttctaaa gaccacaga      720
gacaaggaag aatcgtggt caggactctt tcctatttcg atggagtga cttcgcagcc      780
agagcgaaga tcctgcgct gttttctgtg ggtctcatgg acaacatttg tcctccttca      840
acggttttcg ctgcctacaa ttactacgct ggaccgaagg aatcagaat ctatccgtac      900
aacaaccacg agggaggagg ctctttccaa gcggttgaac aggtgaaatt cttgaaaaaa      960
ctatttgaga aaggctaa                                     978

```

<210> SEQ ID NO 2

<211> LENGTH: 325

<212> TYPE: PRT

<213> ORGANISM: *Thermotoga maritima*

<400> SEQUENCE: 2

```

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
1           5           10           15
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
20           25           30
Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
35           40           45
Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
50           55           60
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
65           70           75           80
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
85           90           95
Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
100          105          110
Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
115          120          125
Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
130          135          140

```

-continued

Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
 145 150 155 160

Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
 165 170 175

Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
 180 185 190

Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
 195 200 205

Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
 210 215 220

Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
 225 230 235 240

Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
 245 250 255

Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
 260 265 270

Met Asp Asn Ile Cys Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
 275 280 285

Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
 290 295 300

Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
 305 310 315 320

Leu Phe Glu Lys Gly
 325

<210> SEQ ID NO 3
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer

<400> SEQUENCE: 3

tcggacaaca tcacctcctt cta

23

<210> SEQ ID NO 4
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer

<400> SEQUENCE: 4

tagaaggagg tgagatggtg tcc

23

<210> SEQ ID NO 5
 <211> LENGTH: 325
 <212> TYPE: PRT
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 5

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
 1 5 10 15

Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
 20 25 30

Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
 35 40 45

-continued

Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
50 55 60

Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
65 70 75 80

Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
85 90 95

Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
100 105 110

Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
115 120 125

Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
130 135 140

Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
145 150 155 160

Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
165 170 175

Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
180 185 190

Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
195 200 205

Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
210 215 220

Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
225 230 235 240

Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
245 250 255

Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
260 265 270

Met Asp Asn Ile Thr Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
275 280 285

Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
290 295 300

Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
305 310 315 320

Leu Phe Glu Lys Gly
325

<210> SEQ ID NO 6
 <211> LENGTH: 3674
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: plasmid

<400> SEQUENCE: 6

gacgaaaggg cctcgtgata cgctatattt tatagggttaa tgtcatgata ataatggttt 60
 cttagacgtc aggtggcact tttcggggaa atgtgcgagg aaccctatt tgtttatttt 120
 tctaaataca ttcaaatatg tatccgctca tgagacaata accctgataa atgcttcaat 180
 aatattgaaa aaggaagagt atgagtattc aacatttccg tgtcgccctt attccctttt 240
 ttgcggcatt ttgccttccct gtttttgctc acccagaaac gctggtgaaa gtaaaagatg 300
 ctgaagatca gttgggtgca cgagtgggtt acatcgaact ggatctcaac agcggtaaga 360
 tccttgagag ttttegcccc gaagaacgtt ttccaatgat gagcactttt aaagttctgc 420
 tatgtggcgc ggtattatcc cgtattgacg ccgggcaaga gcaactcggc cgccgcatac 480

-continued

actattctca	gaatgacttg	gttgagtact	caccagtcac	agaaaagcat	cttacggatg	540
gcatgacagt	aagagaatta	tgcagtgctg	ccataacccat	gagtgataac	actgcgccca	600
acttacttct	gacaacgatc	ggaggaccga	aggagctaac	cgcttttttg	cacaacatgg	660
gggatcatgt	aactcgctt	gatcgttggg	aaccggagct	gaatgaagcc	ataccaaacg	720
acgagcgtga	caccacgatg	cctgtagcaa	tggcaacaac	gttgcgcaa	ctattaactg	780
gcgaactact	tactctagct	tcccggcaac	aattaataga	ctggatggag	gcggataaag	840
ttgcaggacc	acttctgctc	tggcccttc	cggtggctg	gtttattgct	gataaatctg	900
gagccggtga	gctggttct	cgcggtatca	ttgcagcact	ggggccagat	ggtaagccct	960
cccgtatcgt	agttatctac	acgacgggga	gtcaggcaac	tatggatgaa	cgaaatagac	1020
agatcgctga	gataggtgcc	tactgatta	agcattggta	actgtcagac	caagtttact	1080
catatatact	ttagattgat	ttaaaacttc	atthttaatt	taaaaggatc	taggtgaaga	1140
tcctttttga	taatctcatg	acaaaatcc	ctaacgtga	gttttcgctc	caactgagcgt	1200
cagaccccg	agaaaagatc	aaaggatctt	cttgagatcc	ttttttctg	cgcgtaatct	1260
gctgcttgca	aacaaaaaaa	ccaccgctac	cagcggtggt	ttgtttgccg	gatcaagagc	1320
taccaactct	ttttccgaag	gtaactggct	tcagcagagc	gcagatacca	aatactgtcc	1380
ttctagtgt	gccgtagtta	ggccaccact	tcaagaactc	tgtagcaccg	cctacatacc	1440
tcgctctgct	aatcctgtta	ccagtggctg	ctgccagtgg	cgataagtcg	tgtcttaccg	1500
ggttgactc	aagacgatag	ttaccggata	aggcgcagcg	gtcgggctga	acggggggtt	1560
cgtgcacaca	gcccagcttg	gagcgaacga	cctacaccga	actgagatac	ctacagcgtg	1620
agctatgaga	aagcgccacg	cttcccgaag	ggagaaaggc	ggacaggtat	ccggtaaagc	1680
gcagggctcg	aacaggagag	cgcacgaggg	agcttccagg	gggaaacgcc	tggtatcttt	1740
atagtcctgt	cgggtttcgc	caactctgac	ttgagcgtcg	atthttgtga	tgctcgtcag	1800
ggggcgag	cctatgaaa	aacgccagca	acgcggcctt	tttacggttc	ctggcctttt	1860
gctggccttt	tgctcacatg	ttctttcctg	cgttatcccc	tgattctgtg	gataaccgta	1920
ttaccgcctt	tgagtgagct	gataccgctc	gccgcagccg	aacgaccgag	cgcagcagct	1980
cagtgagcga	ggaagcggaa	gagcgcccaa	tacgcaaacc	gcctctcccc	gcgcggtggc	2040
cgattcatta	atgcagctgg	cacgacaggt	ttcccactg	gaaagcgggc	agtgagcga	2100
acgcaattaa	tgtgagttag	ctcactcatt	aggcaacca	ggctttacac	tttatgcttc	2160
cggtcgtat	gttgtgtgga	attgtgagcg	gataacaatt	tcacacagga	aacagctatg	2220
accatgatta	cgccaagctt	gcatgcctgc	agtaaggagg	aataggacat	ggcgttcttc	2280
gacctgcctc	tggaagaact	gaagaaatac	cgtccagagc	gttacgaaga	gaaggacttc	2340
gacgagttct	gggaggaaac	tctggcggag	agcgaagagt	ttccgctgga	cccagtgctc	2400
gagcgtatgg	aatctcacct	gaaaaccgtg	gaggcatatg	acgttacttt	ttctggttac	2460
cgtggccagc	gtatcaaagg	ctggctgctg	gttccgaaac	tggaggaaga	aaaactgccg	2520
tgcgtagttc	agtacatcgg	ttacaacggg	ggccgtggct	ttccgcacga	ttggctgttc	2580
tggccgtcta	tgggctacat	ttgcttcgct	atggatactc	gtggtcaggg	ttccggctgg	2640
ctgaaaggcg	atactccgga	ttatccggag	ggcccggtag	acccgcagta	ccctggcttc	2700
atgacgcgtg	gtattctgga	tccgcgtacc	tattactatc	gccgcgtttt	taccgatgca	2760
gttcgtgccg	tagaggccgc	ggcttctttc	cctcaggttg	accaggagcg	tattgttatac	2820
gctggtggct	cccagggtgg	cggcatcgcc	ctggcggtag	ctgcgctgag	caagaaagct	2880

-continued

```

aaggcactgc tgtgtgacgt cccgttctctg tgtcacttcc gtcgcgctgt tcagctggta 2940
gatacccatc cgtacgcgga gattactaac ttctgaaaa ctcaccgcga caaagaagaa 3000
atcgttttcc gcaccctgtc ctatttcgac ggcgttaact tcgcggctcg tgcaaaaatt 3060
ccggcactgt tctctgttgg tctgatggac aacatcacc ctccttctac cgttttcgcg 3120
gcatataact attatgcggg tccgaaagaa atccgtatct atccgtacaa caaccacgaa 3180
ggcggtggtg gctttcaggc tgttgaacaa gtgaaattcc tgaagaaact gtttgagaag 3240
ggctaatact gaggatcccc gggtagcgag ctgcaattca ctggccgctc ttttacaacg 3300
tcgtgactgg gaaaaccctg gcgttacca acttaatcgc cttgcagcac atcccccttt 3360
cgccagctgg cgtaatagcg aagaggcccc caccgatcgc ctttcccaac agttgcgag 3420
cctgaatggc gaatggcgcc tgatgcggta ttttctcctt acgcatctgt gcggtatttc 3480
acaccgata tgggtgactc tcagtacaat ctgctctgat gccgcatagt taagccagcc 3540
ccgacaccgc ccaacaccgc ctgacgcgcc ctgacgggct tgtctgctcc cggcatccgc 3600
ttacagacaa gctgtgaccg tctccgggag ctgcatgtgt cagaggtttt caccgtcatc 3660
accgaaacgc gcga 3674

```

```

<210> SEQ ID NO 7
<211> LENGTH: 49
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

```

```
<400> SEQUENCE: 7
```

```
taactgcagt aaggaggaat aggacatggc gttcttcgac ctgcctctg 49
```

```

<210> SEQ ID NO 8
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

```

```
<400> SEQUENCE: 8
```

```
tgatctagat tagcccttct caaacagttt ctttcagg 38
```

```

<210> SEQ ID NO 9
<211> LENGTH: 325
<212> TYPE: PRT
<213> ORGANISM: ARTIFICIAL SEQUENCE
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

```

```
<400> SEQUENCE: 9
```

```

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
1           5           10           15
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
20           25           30
Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
35           40           45
Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
50           55           60
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
65           70           75           80
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
85           90           95

```

-continued

Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
 100 105 110
 Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
 115 120 125
 Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
 130 135 140
 Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
 145 150 155 160
 Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
 165 170 175
 Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
 180 185 190
 Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
 195 200 205
 Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
 210 215 220
 Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
 225 230 235 240
 Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Ser Asp Gly Val
 245 250 255
 Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
 260 265 270
 Met Asp Asn Ile Thr Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
 275 280 285
 Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
 290 295 300
 Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
 305 310 315 320
 Leu Phe Glu Lys Gly
 325

<210> SEQ ID NO 10
 <211> LENGTH: 325
 <212> TYPE: PRT
 <213> ORGANISM: ARTIFICIAL SEQUENCE
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 10

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
 1 5 10 15
 Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
 20 25 30
 Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
 35 40 45
 Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
 50 55 60
 Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
 65 70 75 80
 Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
 85 90 95
 Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
 100 105 110
 Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
 115 120 125
 Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe

-continued

130	135	140																		
Met	Thr	Arg	Gly	Ile	Leu	Asp	Pro	Arg	Thr	Tyr	Tyr	Tyr	Arg	Arg	Val					
145					150					155					160					
Leu	Thr	Asp	Ala	Val	Arg	Ala	Val	Glu	Ala	Ala	Ala	Ser	Phe	Pro	Gln					
				165					170					175						
Val	Asp	Gln	Glu	Arg	Ile	Val	Ile	Ala	Gly	Gly	Ser	Gln	Gly	Gly	Gly					
			180					185					190							
Ile	Ala	Leu	Ala	Val	Ser	Ala	Leu	Ser	Lys	Lys	Ala	Lys	Ala	Leu	Leu					
		195					200					205								
Cys	Asp	Val	Pro	Phe	Leu	Cys	His	Phe	Arg	Arg	Ala	Val	Gln	Leu	Val					
210						215					220									
Asp	Thr	His	Pro	Tyr	Ala	Glu	Ile	Thr	Asn	Phe	Leu	Lys	Thr	His	Arg					
225					230					235					240					
Asp	Lys	Glu	Glu	Ile	Val	Phe	Arg	Thr	Leu	Ser	Tyr	Phe	Asp	Gly	Val					
				245					250					255						
Asn	Phe	Ala	Ala	Arg	Ala	Lys	Ile	Pro	Ala	Leu	Phe	Ser	Val	Gly	Leu					
			260					265					270							
Met	Asp	Asn	Ile	Thr	Pro	Pro	Ser	Thr	Val	Phe	Ala	Ala	Tyr	Asn	Tyr					
		275					280						285							
Tyr	Ala	Gly	Pro	Lys	Glu	Ile	Arg	Ile	Tyr	Pro	Tyr	Asn	Asn	His	Glu					
	290					295					300									
Gly	Gly	Gly	Ser	Phe	Gln	Ala	Val	Glu	Gln	Val	Lys	Phe	Leu	Lys	Lys					
305					310					315					320					
Leu	Phe	Glu	Lys	Gly																
				325																

<210> SEQ ID NO 11

<211> LENGTH: 325

<212> TYPE: PRT

<213> ORGANISM: ARTIFICIAL SEQUENCE

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 11

Met	Ala	Phe	Phe	Asp	Leu	Pro	Leu	Glu	Glu	Leu	Lys	Lys	Tyr	Arg	Pro					
1				5					10					15						
Asp	Arg	Tyr	Glu	Glu	Lys	Asp	Phe	Asp	Glu	Phe	Trp	Glu	Glu	Thr	Leu					
			20					25					30							
Ala	Glu	Ser	Glu	Lys	Phe	Pro	Leu	Asp	Pro	Val	Phe	Glu	Arg	Met	Glu					
			35				40					45								
Ser	His	Leu	Lys	Thr	Val	Glu	Ala	Tyr	Asp	Val	Thr	Phe	Ser	Gly	Tyr					
			50			55					60									
Arg	Gly	Gln	Arg	Ile	Lys	Gly	Trp	Leu	Leu	Val	Pro	Arg	Leu	Glu	Glu					
65				70						75				80						
Glu	Lys	Leu	Pro	Cys	Val	Val	Gln	Tyr	Ile	Gly	Tyr	Asn	Gly	Gly	Arg					
				85					90					95						
Gly	Phe	Pro	His	Asp	Trp	Leu	Phe	Trp	Pro	Ser	Met	Gly	Tyr	Ile	Cys					
			100					105					110							
Phe	Val	Met	Asp	Thr	Arg	Gly	Gln	Gly	Ser	Gly	Trp	Leu	Lys	Gly	Asp					
		115					120					125								
Thr	Pro	Asp	Tyr	Pro	Glu	Gly	Pro	Val	Asp	Pro	Gln	Tyr	Pro	Gly	Phe					
		130				135					140									
Met	Thr	Arg	Gly	Ile	Leu	Asp	Pro	Arg	Thr	Tyr	Tyr	Tyr	Arg	Arg	Val					
145					150					155					160					
Phe	Thr	Asp	Ala	Val	Arg	Ala	Val	Glu	Ala	Ala	Ala	Ser	Leu	Pro	Gln					
				165					170					175						

-continued

Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
 180 185 190
 Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
 195 200 205
 Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
 210 215 220
 Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
 225 230 235 240
 Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
 245 250 255
 Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
 260 265 270
 Met Asp Asn Ile Thr Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
 275 280 285
 Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
 290 295 300
 Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
 305 310 315 320
 Leu Phe Glu Lys Gly
 325

<210> SEQ ID NO 12
 <211> LENGTH: 325
 <212> TYPE: PRT
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 12

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
 1 5 10 15
 Glu Arg Tyr Glu Glu Lys Asp Phe Gly Glu Phe Trp Glu Glu Thr Leu
 20 25 30
 Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
 35 40 45
 Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
 50 55 60
 Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
 65 70 75 80
 Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
 85 90 95
 Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
 100 105 110
 Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
 115 120 125
 Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
 130 135 140
 Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
 145 150 155 160
 Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
 165 170 175
 Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
 180 185 190
 Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
 195 200 205

-continued

Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
 210 215 220
 Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
 225 230 235 240
 Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
 245 250 255
 Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
 260 265 270
 Met Asp Asn Ile Thr Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
 275 280 285
 Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
 290 295 300
 Gly Gly Ser Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
 305 310 315 320
 Leu Phe Glu Lys Gly
 325

<210> SEQ ID NO 13
 <211> LENGTH: 325
 <212> TYPE: PRT
 <213> ORGANISM: ARTIFICIAL SEQUENCE
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 13

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
 1 5 10 15
 Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
 20 25 30
 Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
 35 40 45
 Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
 50 55 60
 Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
 65 70 75 80
 Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
 85 90 95
 Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
 100 105 110
 Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
 115 120 125
 Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
 130 135 140
 Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
 145 150 155 160
 Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
 165 170 175
 Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
 180 185 190
 Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
 195 200 205
 Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
 210 215 220
 Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
 225 230 235 240
 Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val

-continued

	245		250		255										
Asp	Phe	Ala	Ala	Arg	Ala	Lys	Ile	Pro	Ala	Leu	Phe	Ser	Val	Gly	Leu
		260						265					270		
Met	Asp	Asn	Ile	Thr	Pro	Pro	Ser	Thr	Val	Phe	Ala	Ala	Tyr	Asn	Tyr
		275					280						285		
Tyr	Ala	Gly	Pro	Lys	Glu	Ile	Arg	Ile	Tyr	Pro	Tyr	Asn	Asn	His	Glu
	290					295					300				
Gly	Gly	Gly	Ser	Phe	Gln	Ala	Val	Glu	Gln	Val	Lys	Phe	Leu	Lys	Lys
305					310					315					320
Leu	Phe	Glu	Lys	Gly											
				325											

<210> SEQ ID NO 14

<211> LENGTH: 325

<212> TYPE: PRT

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 14

Met	Ala	Phe	Phe	Asp	Leu	Pro	Leu	Glu	Glu	Leu	Lys	Lys	Tyr	Arg	Pro
1				5					10					15	
Glu	Arg	Tyr	Glu	Glu	Lys	Asp	Phe	Asp	Glu	Phe	Trp	Glu	Glu	Thr	Leu
			20					25					30		
Ala	Glu	Ser	Glu	Lys	Phe	Pro	Leu	Asp	Pro	Val	Phe	Glu	Arg	Met	Glu
		35					40					45			
Ser	His	Leu	Lys	Thr	Val	Glu	Ala	Tyr	Asp	Val	Thr	Phe	Ser	Gly	Tyr
	50					55					60				
Arg	Gly	Arg	Arg	Ile	Lys	Gly	Trp	Leu	Leu	Val	Pro	Lys	Leu	Glu	Glu
65					70					75				80	
Glu	Lys	Leu	Pro	Cys	Val	Val	Gln	Tyr	Ile	Gly	Tyr	Asn	Gly	Gly	Arg
				85					90					95	
Gly	Phe	Pro	His	Asp	Trp	Leu	Phe	Trp	Pro	Ser	Met	Gly	Tyr	Ile	Cys
			100					105					110		
Phe	Val	Met	Asp	Thr	Arg	Gly	Gln	Gly	Ser	Gly	Trp	Leu	Lys	Gly	Asp
		115					120					125			
Thr	Pro	Asp	Tyr	Pro	Glu	Gly	Pro	Val	Asp	Pro	Gln	Tyr	Pro	Gly	Phe
	130					135					140				
Met	Thr	Arg	Gly	Ile	Leu	Asp	Pro	Arg	Thr	Tyr	Tyr	Tyr	Arg	Arg	Val
145					150					155					160
Phe	Thr	Asp	Ala	Val	Arg	Ala	Val	Glu	Ala	Ala	Ala	Ser	Phe	Pro	Gln
			165						170					175	
Val	Asp	Gln	Glu	Arg	Ile	Val	Ile	Ala	Gly	Gly	Ser	Gln	Gly	Gly	Gly
		180						185					190		
Ile	Ala	Leu	Ala	Val	Ser	Ala	Leu	Ser	Lys	Lys	Ala	Lys	Ala	Leu	Leu
	195						200					205			
Cys	Asp	Val	Pro	Phe	Leu	Cys	His	Phe	Arg	Arg	Ala	Val	Gln	Leu	Val
	210					215					220				
Asp	Thr	His	Pro	Tyr	Ala	Glu	Ile	Thr	Asn	Phe	Leu	Lys	Thr	His	Arg
225					230					235				240	
Asp	Lys	Glu	Glu	Ile	Val	Phe	Arg	Thr	Leu	Ser	Tyr	Phe	Asp	Gly	Val
				245					250					255	
Asn	Phe	Ala	Ala	Arg	Ala	Lys	Ile	Pro	Ala	Leu	Phe	Ser	Val	Gly	Leu
		260						265					270		
Met	Asp	Asn	Ile	Thr	Pro	Pro	Ser	Thr	Val	Phe	Ala	Ala	Tyr	Asn	Tyr
	275						280					285			

-continued

Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
 290 295 300

Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
 305 310 315 320

Leu Phe Glu Lys Gly
 325

<210> SEQ ID NO 15

<211> LENGTH: 325

<212> TYPE: PRT

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 15

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
 1 5 10 15

Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Pro
 20 25 30

Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
 35 40 45

Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
 50 55 60

Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
 65 70 75 80

Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
 85 90 95

Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
 100 105 110

Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
 115 120 125

Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
 130 135 140

Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
 145 150 155 160

Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
 165 170 175

Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
 180 185 190

Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
 195 200 205

Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
 210 215 220

Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
 225 230 235 240

Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
 245 250 255

Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
 260 265 270

Met Asp Asn Ile Thr Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
 275 280 285

Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
 290 295 300

Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
 305 310 315 320

-continued

Leu Phe Glu Lys Gly
325

<210> SEQ ID NO 16
<211> LENGTH: 325
<212> TYPE: PRT
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 16

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr His Pro
1 5 10 15
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
20 25 30
Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
35 40 45
Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
50 55 60
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
65 70 75 80
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
85 90 95
Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
100 105 110
Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
115 120 125
Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
130 135 140
Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
145 150 155 160
Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro His
165 170 175
Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
180 185 190
Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Gln
195 200 205
Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
210 215 220
Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
225 230 235 240
Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
245 250 255
Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
260 265 270
Met Asp Asn Ile Thr Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
275 280 285
Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
290 295 300
Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
305 310 315 320
Leu Phe Glu Lys Gly
325

<210> SEQ ID NO 17
<211> LENGTH: 325
<212> TYPE: PRT

-continued

```

<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 17
Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Leu
1          5          10          15
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
20          25          30
Ala Glu Ser Glu Lys Ser Pro Leu Asp Pro Val Phe Glu Arg Met Glu
35          40          45
Ser His Leu Lys Asn Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
50          55          60
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
65          70          75          80
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
85          90          95
Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
100         105         110
Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
115         120         125
Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
130         135         140
Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Cys Val
145         150         155         160
Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
165         170         175
Val Asp Gln Glu Arg Ile Asp Ile Ala Gly Gly Ser Gln Gly Gly Gly
180         185         190
Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
195         200         205
Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
210         215         220
Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
225         230         235         240
Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
245         250         255
Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
260         265         270
Met Asp Asn Ile Thr Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
275         280         285
Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
290         295         300
Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
305         310         315         320
Leu Phe Glu Lys Gly
325

```

```

<210> SEQ ID NO 18
<211> LENGTH: 325
<212> TYPE: PRT
<213> ORGANISM: ARTIFICIAL SEQUENCE
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

```

```

<400> SEQUENCE: 18

```

```

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro

```

-continued

1	5	10	15
Glu Arg Tyr	Glu Glu Lys Asp Phe Asp	Glu Phe Trp	Glu Glu Thr Leu
	20	25	30
Ala Glu Ser	Glu Lys Phe Pro Leu Asp	Pro Val Phe	Glu Arg Met Glu
	35	40	45
Ser His Leu	Lys Thr Val Glu Ala Tyr Asp	Val Thr Phe	Ser Gly Tyr
	50	55	60
Arg Gly Gln	Arg Ile Lys Gly Trp Leu Leu	Val Pro Lys	Leu Glu Glu
	65	70	75
Glu Lys Leu	Pro Cys Val Val Gln Tyr	Ile Gly Tyr	Asn Gly Gly Arg
	85	90	95
Gly Phe Pro	His Asp Trp Leu Phe Trp	Pro Ser Met	Gly Tyr Ile Cys
	100	105	110
Phe Val Met	Asp Thr Arg Gly Gln Gly	Ser Gly Trp	Leu Lys Gly Asp
	115	120	125
Thr Pro Asp	Tyr Pro Glu Gly Pro Val	Asp Pro Gln	Tyr Pro Gly Phe
	130	135	140
Met Thr Arg	Gly Ile Leu Asp Pro Arg	Thr Tyr Tyr	Tyr Arg Arg Val
	145	150	155
Phe Thr Asp	Ala Val Arg Ala Val Glu	Ala Ala Ala	Ser Phe Pro Gln
	165	170	175
Val Asp Gln	Glu Arg Ile Val Ile Ala	Gly Gly Ser	Gln Gly Gly Gly
	180	185	190
Ile Ala Leu	Ala Val Ser Ala Leu Ser	Lys Lys Ala	Lys Ala Leu Leu
	195	200	205
Cys Asp Val	Pro Phe Leu Cys His Phe	Arg Arg Ala	Val Gln Leu Val
	210	215	220
Asp Thr His	Pro Tyr Ala Glu Ile Thr	Asn Phe Leu	Lys Thr His Arg
	225	230	235
Asp Lys Glu	Glu Ile Val Phe Arg Thr	Leu Ser Tyr	Phe Asp Gly Val
	245	250	255
Asn Phe Ala	Ala Arg Ala Lys Ile Pro	Gly Leu Phe	Ser Val Gly Leu
	260	265	270
Met Asp Asn	Ile Thr Pro Pro Ser Thr	Val Phe Ala	Ala Tyr Asn Tyr
	275	280	285
Tyr Ala Gly	Pro Lys Glu Ile Arg Ile	Tyr Pro Tyr	Asn Asn His Glu
	290	295	300
Gly Gly Gly	Ser Phe Gln Ala Val Glu	Gln Val Lys	Phe Leu Lys Lys
	305	310	315
Leu Phe Glu	Lys Gly		
	325		

<210> SEQ ID NO 19

<211> LENGTH: 325

<212> TYPE: PRT

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 19

Met Ala Phe	Phe Asp Leu Pro Leu	Glu Glu Leu Lys	Lys Tyr Arg Pro
	5	10	15
Glu Arg Tyr	Glu Lys Asp Phe Asp	Glu Phe Trp	Glu Glu Thr Leu
	20	25	30
Ala Glu Ser	Glu Lys Phe Pro Leu Asp	Pro Val Phe	Glu Arg Met Glu
	35	40	45

-continued

Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
 50 55 60
 Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
 65 70 75 80
 Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
 85 90 95
 Gly Phe Pro Asn Asp Trp Leu Phe Ser Pro Ser Met Gly Tyr Ile Cys
 100 105 110
 Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
 115 120 125
 Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
 130 135 140
 Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
 145 150 155 160
 Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
 165 170 175
 Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
 180 185 190
 Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
 195 200 205
 Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
 210 215 220
 Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
 225 230 235 240
 Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
 245 250 255
 Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
 260 265 270
 Met Asp Asn Ile Thr Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
 275 280 285
 Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
 290 295 300
 Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
 305 310 315 320
 Leu Phe Glu Lys Gly
 325

<210> SEQ ID NO 20
 <211> LENGTH: 325
 <212> TYPE: PRT
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 20

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
 1 5 10 15
 Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
 20 25 30
 Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
 35 40 45
 Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
 50 55 60
 Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
 65 70 75 80

-continued

Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
 85 90 95
 Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
 100 105 110
 Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
 115 120 125
 Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
 130 135 140
 Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
 145 150 155 160
 Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
 165 170 175
 Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
 180 185 190
 Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
 195 200 205
 Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
 210 215 220
 Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
 225 230 235 240
 Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
 245 250 255
 Asn Phe Ala Ala Ser Ala Lys Phe Pro Ala Leu Phe Ser Val Gly Leu
 260 265 270
 Met Asp Asn Ile Thr Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
 275 280 285
 Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
 290 295 300
 Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
 305 310 315 320
 Leu Phe Glu Lys Gly
 325

<210> SEQ ID NO 21
 <211> LENGTH: 325
 <212> TYPE: PRT
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 21

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
 1 5 10 15
 Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
 20 25 30
 Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
 35 40 45
 Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
 50 55 60
 Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
 65 70 75 80
 Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
 85 90 95
 Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
 100 105 110
 Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp

-continued

115					120					125					
Thr	Pro	Asp	Tyr	Pro	Glu	Gly	Pro	Val	Asp	Pro	Gln	Tyr	Pro	Gly	Phe
130					135					140					
Met	Thr	Arg	Gly	Ile	Leu	Asp	Pro	Arg	Thr	Tyr	Tyr	Tyr	Arg	Arg	Val
145					150					155					160
Phe	Thr	Asp	Ala	Val	Arg	Ala	Val	Glu	Ala	Ala	Ala	Ser	Phe	Pro	Gln
				165					170					175	
Val	Asn	Gln	Glu	Arg	Ile	Val	Ile	Ala	Gly	Gly	Ser	Gln	Gly	Gly	Gly
			180					185					190		
Ile	Ala	Leu	Ala	Val	Ser	Ala	Leu	Ser	Lys	Lys	Ala	Lys	Ala	Leu	Leu
		195					200					205			
Cys	Asp	Val	Pro	Phe	Leu	Cys	His	Phe	Arg	Arg	Ala	Val	Gln	Leu	Val
	210					215					220				
Asp	Thr	His	Pro	Tyr	Ala	Glu	Ile	Thr	Asn	Phe	Leu	Lys	Thr	His	Arg
225					230					235					240
Asp	Lys	Glu	Glu	Ile	Val	Phe	Arg	Thr	Leu	Ser	Tyr	Phe	Asp	Gly	Val
				245					250					255	
Asn	Phe	Ala	Ala	Arg	Ala	Lys	Ile	Pro	Ala	Leu	Phe	Ser	Val	Gly	Leu
			260					265					270		
Met	Asp	Asn	Ile	Thr	Pro	Pro	Ser	Thr	Val	Phe	Ala	Ala	Tyr	Asn	Tyr
		275					280						285		
Tyr	Ala	Gly	Pro	Lys	Glu	Ile	Arg	Ile	Tyr	Pro	Tyr	Asn	Asn	His	Glu
	290					295					300				
Gly	Gly	Gly	Ser	Phe	Gln	Ala	Val	Glu	Gln	Val	Lys	Phe	Leu	Lys	Lys
305					310					315					320
Leu	Phe	Glu	Lys	Gly											
				325											

<210> SEQ ID NO 22

<211> LENGTH: 325

<212> TYPE: PRT

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 22

Met	Ala	Phe	Phe	Asp	Leu	Pro	Leu	Glu	Glu	Leu	Lys	Lys	Tyr	Arg	Pro
1				5					10					15	
Glu	Arg	Tyr	Glu	Glu	Lys	Asp	Phe	Asp	Glu	Phe	Trp	Glu	Glu	Thr	Leu
			20					25					30		
Ala	Glu	Ser	Glu	Lys	Phe	Pro	Leu	Asp	Pro	Val	Phe	Glu	Arg	Val	Glu
			35				40					45			
Ser	His	Leu	Lys	Thr	Val	Glu	Ala	Tyr	Asp	Val	Thr	Phe	Ser	Gly	Tyr
	50					55					60				
Arg	Gly	Gln	Arg	Ile	Lys	Gly	Trp	Leu	Leu	Val	Pro	Lys	Leu	Glu	Glu
65					70					75					80
Glu	Lys	Leu	Pro	Cys	Val	Val	Gln	Tyr	Ile	Gly	Tyr	Asn	Gly	Gly	Arg
				85					90					95	
Gly	Phe	Pro	His	Asp	Trp	Leu	Phe	Trp	Pro	Ser	Met	Gly	Tyr	Ile	Cys
			100					105					110		
Phe	Val	Met	Asp	Thr	Arg	Gly	Gln	Gly	Ser	Gly	Trp	Leu	Lys	Gly	Asp
		115					120					125			
Thr	Pro	Asp	Tyr	Pro	Asp	Gly	Pro	Val	Asp	Pro	Gln	Tyr	Pro	Gly	Phe
	130					135					140				
Met	Thr	Arg	Gly	Ile	Gln	Asp	Pro	Arg	Thr	Tyr	Tyr	Tyr	Arg	Arg	Val
145					150					155					160

-continued

Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
 195 200 205

Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
 210 215 220

Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
 225 230 235 240

Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
 245 250 255

Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Ala Gly Leu
 260 265 270

Met Asp Asn Ile Thr Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
 275 280 285

Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
 290 295 300

Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
 305 310 315 320

Leu Phe Glu Lys Gly
 325

<210> SEQ ID NO 24
 <211> LENGTH: 325
 <212> TYPE: PRT
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 24

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
 1 5 10 15

Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
 20 25 30

Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
 35 40 45

Ser His Leu Lys Thr Val Lys Ala Tyr Asp Val Thr Phe Ser Gly Tyr
 50 55 60

Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
 65 70 75 80

Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
 85 90 95

Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
 100 105 110

Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
 115 120 125

Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
 130 135 140

Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
 145 150 155 160

Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
 165 170 175

Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
 180 185 190

Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
 195 200 205

Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
 210 215 220

Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg

-continued

```

225          230          235          240
Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
          245          250          255

Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
          260          265          270

Met Asp Asn Ile Thr Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
          275          280          285

Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
          290          295          300

Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
305          310          315          320

Leu Phe Glu Lys Gly
          325

```

```

<210> SEQ ID NO 25
<211> LENGTH: 325
<212> TYPE: PRT
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

```

```

<400> SEQUENCE: 25

```

```

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
1          5          10          15

Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
          20          25          30

Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
          35          40          45

Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
50          55          60

Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
65          70          75          80

Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
          85          90          95

Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
          100          105          110

Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
          115          120          125

Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
          130          135          140

Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
145          150          155          160

Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
          165          170          175

Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
          180          185          190

Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
          195          200          205

Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
          210          215          220

Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
225          230          235          240

Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
          245          250          255

Asn Phe Ala Ala His Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
          260          265          270

```

-continued

Met Asp Asn Ile Thr Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
 275 280 285

Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
 290 295 300

Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
 305 310 315 320

Leu Phe Glu Lys Gly
 325

<210> SEQ ID NO 26
 <211> LENGTH: 325
 <212> TYPE: PRT
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 26

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
 1 5 10 15

Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
 20 25 30

Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
 35 40 45

Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
 50 55 60

Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
 65 70 75 80

Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
 85 90 95

Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
 100 105 110

Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
 115 120 125

Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
 130 135 140

Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
 145 150 155 160

Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
 165 170 175

Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
 180 185 190

Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Thr Leu Leu
 195 200 205

Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
 210 215 220

Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
 225 230 235 240

Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
 245 250 255

Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
 260 265 270

Met Asp Asn Ile Thr Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
 275 280 285

Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
 290 295 300

-continued

Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
305 310 315 320

Leu Phe Glu Lys Gly
325

<210> SEQ ID NO 27

<211> LENGTH: 325

<212> TYPE: PRT

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 27

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
1 5 10 15

Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
20 25 30

Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
35 40 45

Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
50 55 60

Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
65 70 75 80

Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
85 90 95

Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
100 105 110

Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
115 120 125

Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
130 135 140

Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
145 150 155 160

Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
165 170 175

Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
180 185 190

Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
195 200 205

Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
210 215 220

Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
225 230 235 240

Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
245 250 255

Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
260 265 270

Met Asp Asn Ile Thr Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
275 280 285

Tyr Ala Gly Pro Lys Glu Ile Pro Ile Tyr Pro Tyr Asn Asn His Glu
290 295 300

Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
305 310 315 320

Leu Phe Glu Lys Gly
325

-continued

```

<210> SEQ ID NO 28
<211> LENGTH: 325
<212> TYPE: PRT
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 28

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
1          5          10          15
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
20          25          30
Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Ser Glu Arg Met Glu
35          40          45
Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
50          55          60
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
65          70          75          80
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
85          90          95
Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
100         105         110
Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
115         120         125
Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
130         135         140
Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
145         150         155         160
Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
165         170         175
Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
180         185         190
Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
195         200         205
Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
210         215         220
Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
225         230         235         240
Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
245         250         255
Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
260         265         270
Met Asp Asn Ile Thr Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
275         280         285
Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
290         295         300
Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
305         310         315         320
Leu Phe Glu Lys Gly
325

```

```

<210> SEQ ID NO 29
<211> LENGTH: 325
<212> TYPE: PRT
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

```

-continued

<400> SEQUENCE: 29

Met Ala Phe Phe Asp Leu Pro Leu Lys Glu Leu Lys Lys Tyr Arg Pro
 1 5 10 15
 Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
 20 25 30
 Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
 35 40 45
 Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
 50 55 60
 Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
 65 70 75 80
 Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
 85 90 95
 Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
 100 105 110
 Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
 115 120 125
 Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
 130 135 140
 Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
 145 150 155 160
 Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
 165 170 175
 Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
 180 185 190
 Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
 195 200 205
 Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
 210 215 220
 Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
 225 230 235 240
 Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Cys Val
 245 250 255
 Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
 260 265 270
 Met Asp Asn Ile Thr Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
 275 280 285
 Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
 290 295 300
 Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
 305 310 315 320
 Leu Phe Glu Lys Gly
 325

<210> SEQ ID NO 30

<211> LENGTH: 325

<212> TYPE: PRT

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 30

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
 1 5 10 15
 Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
 20 25 30

-continued

Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
 65 70 75 80
 Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
 85 90 95
 Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
 100 105 110
 Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
 115 120 125
 Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
 130 135 140
 Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
 145 150 155 160
 Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
 165 170 175
 Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
 180 185 190
 Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
 195 200 205
 Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
 210 215 220
 Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
 225 230 235 240
 Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
 245 250 255
 Asn Phe Ala Ala His Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
 260 265 270
 Met Asp Asn Ile Thr Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
 275 280 285
 Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
 290 295 300
 Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
 305 310 315 320
 Leu Phe Glu Lys Gly
 325

<210> SEQ ID NO 32
 <211> LENGTH: 325
 <212> TYPE: PRT
 <213> ORGANISM: ARTIFICIAL SEQUENCE
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 32

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
 1 5 10 15
 Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
 20 25 30
 Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
 35 40 45
 Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
 50 55 60
 Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
 65 70 75 80
 Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
 85 90 95
 Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys

-continued

100					105					110					
Phe	Val	Met	Asp	Thr	Arg	Gly	Gln	Gly	Ser	Gly	Trp	Leu	Lys	Gly	Asp
		115					120					125			
Thr	Pro	Asp	Tyr	Pro	Glu	Gly	Pro	Val	Asp	Pro	Gln	Tyr	Pro	Gly	Phe
	130					135					140				
Met	Thr	Arg	Gly	Ile	Leu	Asp	Pro	Arg	Thr	Tyr	Tyr	Tyr	Arg	Arg	Val
145					150					155					160
Phe	Thr	Asp	Ala	Val	Arg	Ala	Val	Glu	Ala	Ala	Ala	Ser	Phe	Pro	Gln
			165						170					175	
Val	Asp	Gln	Glu	Arg	Ile	Val	Ile	Ala	Gly	Gly	Ser	Gln	Gly	Gly	Gly
		180						185					190		
Ile	Ala	Leu	Ala	Val	Ser	Ala	Leu	Ser	Lys	Lys	Ala	Lys	Ala	Leu	Leu
		195					200					205			
Cys	Asp	Val	Pro	Phe	Leu	Ser	His	Phe	Arg	Arg	Ala	Val	Gln	Leu	Val
	210					215					220				
Asp	Thr	His	Pro	Tyr	Ala	Glu	Ile	Thr	Asn	Phe	Leu	Lys	Thr	His	Arg
225					230					235					240
Asp	Lys	Glu	Glu	Ile	Val	Phe	Arg	Thr	Leu	Ser	Tyr	Phe	Asp	Gly	Val
				245					250					255	
Asn	Phe	Ala	Ala	Arg	Ala	Lys	Ile	Pro	Ala	Leu	Phe	Ser	Val	Gly	Leu
			260					265					270		
Met	Asp	Asn	Ile	Thr	Pro	Pro	Ser	Thr	Val	Phe	Ala	Ala	Tyr	Asn	Tyr
		275					280					285			
Tyr	Ala	Gly	Pro	Lys	Glu	Ile	Arg	Ile	Tyr	Pro	Tyr	Asn	Asn	His	Glu
	290					295					300				
Gly	Gly	Gly	Ser	Phe	Gln	Ala	Val	Glu	Gln	Val	Lys	Phe	Leu	Lys	Lys
305					310					315					320
Leu	Phe	Glu	Lys	Gly											
				325											

<210> SEQ ID NO 33
 <211> LENGTH: 325
 <212> TYPE: PRT
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 33

Met	Ala	Phe	Phe	Asp	Leu	Pro	Leu	Glu	Glu	Leu	Lys	Lys	Tyr	Arg	Pro
1				5					10					15	
Glu	Arg	Tyr	Glu	Glu	Lys	Asp	Phe	Asp	Glu	Phe	Trp	Glu	Glu	Thr	Leu
			20					25					30		
Ala	Glu	Ser	Glu	Lys	Phe	Pro	Leu	Asp	Pro	Val	Phe	Glu	Arg	Met	Glu
		35					40					45			
Ser	His	Leu	Lys	Thr	Val	Glu	Ala	Tyr	Asp	Val	Thr	Phe	Ser	Gly	Tyr
	50					55					60				
Arg	Gly	Gln	Arg	Ile	Lys	Gly	Trp	Leu	Leu	Val	Pro	Lys	Leu	Glu	Glu
65					70					75				80	
Glu	Lys	Leu	Pro	Cys	Val	Val	Gln	Tyr	Ile	Gly	Tyr	Asn	Gly	Gly	Arg
				85					90					95	
Gly	Phe	Pro	His	Asp	Trp	Leu	Phe	Trp	Pro	Ser	Met	Gly	Tyr	Ile	Cys
			100					105					110		
Phe	Val	Met	Asp	Thr	Arg	Gly	Gln	Gly	Ser	Gly	Trp	Leu	Lys	Gly	Asp
		115					120						125		
Thr	Pro	Asp	Tyr	Pro	Glu	Gly	Pro	Val	Asp	Pro	Gln	Tyr	Pro	Gly	Phe
						135					140				

-continued

Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
 145 150 155 160

Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
 165 170 175

Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
 180 185 190

Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
 195 200 205

Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
 210 215 220

Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
 225 230 235 240

Asp Lys Glu Asp Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
 245 250 255

Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Ile Gly Leu
 260 265 270

Met Asp Asn Ile Thr Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
 275 280 285

Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
 290 295 300

Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
 305 310 315 320

Leu Phe Glu Lys Gly
 325

<210> SEQ ID NO 34
 <211> LENGTH: 325
 <212> TYPE: PRT
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 34

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
 1 5 10 15

Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
 20 25 30

Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
 35 40 45

Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
 50 55 60

Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
 65 70 75 80

Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
 85 90 95

Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
 100 105 110

Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
 115 120 125

Thr Pro Asp Tyr Pro Glu Gly Pro Leu Asp Pro Gln Tyr Pro Gly Phe
 130 135 140

Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
 145 150 155 160

Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
 165 170 175

-continued

Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
 180 185 190

Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
 195 200 205

Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
 210 215 220

Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
 225 230 235 240

Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
 245 250 255

Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
 260 265 270

Met Asp Asn Ile Thr Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
 275 280 285

Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
 290 295 300

Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
 305 310 315 320

Leu Phe Glu Lys Gly
 325

<210> SEQ ID NO 35
 <211> LENGTH: 325
 <212> TYPE: PRT
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 35

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
 1 5 10 15

Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
 20 25 30

Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
 35 40 45

Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
 50 55 60

Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
 65 70 75 80

Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
 85 90 95

Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
 100 105 110

Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
 115 120 125

Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
 130 135 140

Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
 145 150 155 160

Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
 165 170 175

Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
 180 185 190

Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
 195 200 205

Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val

-continued

210	215	220			
Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg					
225	230	235			240
Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Ile					
	245	250			255
Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu					
	260	265			270
Met Asp Asn Ile Thr Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr					
	275	280			285
Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu					
	290	295			300
Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys					
305	310	315			320
Leu Phe Glu Lys Gly					
	325				

<210> SEQ ID NO 36
 <211> LENGTH: 325
 <212> TYPE: PRT
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 36

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro					
1	5	10			15
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu					
	20	25			30
Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu					
	35	40			45
Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Leu Ser Gly Tyr					
	50	55			60
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu					
65	70	75			80
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg					
	85	90			95
Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys					
	100	105			110
Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp					
	115	120			125
Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe					
	130	135			140
Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val					
145	150	155			160
Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln					
	165	170			175
Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly					
	180	185			190
Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu					
	195	200			205
Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val					
	210	215			220
Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg					
225	230	235			240
Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val					
	245	250			255

-continued

Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
 260 265 270

Met Asp Asn Ile Thr Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
 275 280 285

Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
 290 295 300

Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
 305 310 315 320

Leu Phe Glu Lys Gly
 325

<210> SEQ ID NO 37
 <211> LENGTH: 325
 <212> TYPE: PRT
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 37

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
 1 5 10 15

Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
 20 25 30

Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
 35 40 45

Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
 50 55 60

Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
 65 70 75 80

Glu Arg Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
 85 90 95

Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
 100 105 110

Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
 115 120 125

Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
 130 135 140

Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
 145 150 155 160

Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
 165 170 175

Val Asp Gln Lys Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
 180 185 190

Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
 195 200 205

Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
 210 215 220

Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
 225 230 235 240

Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
 245 250 255

Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
 260 265 270

Met Asp Asn Ile Thr Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
 275 280 285

-continued

Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
 290 295 300

Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
 305 310 315 320

Leu Phe Glu Lys Gly
 325

<210> SEQ ID NO 38
 <211> LENGTH: 978
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 38

```

atggcggttct tgcacctgcc tctggaagaa ctgaagaaat accgtccaga gcgttacgaa    60
gagaaggact tgcacgagtt ctgggaggaa actctggcgg agagcgaaaa gtttccgctg    120
gaccagtggt tgcagcgtat ggaatctcac ctgaaaaccg tggaggcata tgacgttact    180
ttttctgggt accgtggcca gcgtatcaaa ggctggctgc tggttccgaa actggaggaa    240
gaaaaactgc cgtgcgtagt tcagtacatc ggttacaacg gtggccgtgg ctttccgcac    300
gattggctgt tctggccgtc tatgggctac atttgcttcg tcatggatac tctgggtcag    360
ggttccggct ggctgaaagg cgatactccg gattatccgg agggcccggg agaccgcag    420
taccctggct tcatgacgcg tggattctg gatccgcgta cctattacta tcgccgcggt    480
ttaccgatg cagttcgtgc cgtagaggcc gcgcttctt tccctcaggt tgaccaggag    540
cgtattgta tcgctgggtg cctccagggt ggccgcatcg cctggcgggt atctgcgctg    600
agcaagaaag ctaaggcact gctgtgtgac gtcccgttc tgtgtcactt ccgtcgcgct    660
gttcagctgg tagataccca tccgtacgcg gagattacta acttctgaa aactcaccgc    720
gacaaagaag aaatcgtttt ccgcaccctg tcctattccg acggcgtaa cttcgcggct    780
cgtgcaaaaa ttccggcact gttctctggt ggtctgatgg acaacatcac cctccttct    840
accgttttct cggcatataa ctattatgcg ggtccgaaag aaatccgtat ctatccgtac    900
aacaaccacg aaggcgggtg tagctttcag gctgttgaac aagtgaatt cctgaagaaa    960
ctgtttgaga agggctaa                                     978

```

<210> SEQ ID NO 39
 <211> LENGTH: 978
 <212> TYPE: DNA
 <213> ORGANISM: ARTIFICIAL SEQUENCE
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 39

```

atggcggttct tgcacctgcc tctggaagaa ctgaagaaat accgtccaga gcgttacgaa    60
gagaaggact tgcacgagtt ctgggaggaa actctggcgg agagcgaaaa gtttccgctg    120
gaccagtggt tgcagcgtat ggaatctcac ctgaaaaccg tggaggcata tgacgttact    180
ttttctgggt accgtggcca gcgtatcaaa ggctggctgc tggttccgaa actggaggaa    240
gaaaaactgc cgtgcgtagt tcagtacatc ggttacaacg gtggccgtgg ctttccgcac    300
gattggctgt tctggccgtc tatgggctac atttgcttcg tcatggatac tctgggtcag    360
ggttccggct ggctgaaagg cgatactccg gattatccgg agggcccggg agaccgcag    420
taccctggct tcatgacgcg tggattctg gatccgcgta cctattacta tcgccgcggt    480
ttaaccgatg cagttcgtgc cgtagaggcc gcgcttctt tccctcaggt tgaccaggag    540

```

-continued

```

cgtattgtta tcgctggtgg ctcccagggt ggccgcatcg ccctggcggt atctgcgctg 600
agcaagaaag ctaaggcact gctgtgtgac gtcccgttcc tgtgtcactt ccgtcgcgct 660
gttcagctgg tagataccca tccgtacgcg gagattacta acttctgaa aactcaccgc 720
gacaaagaag aaatcgtttt ccgcaccctg tcctatttcg acggcgtaa cttcgcggct 780
cgtgcaaaaa ttccggcact gttctctggt ggtctgatgg acaacatcac ccctccttct 840
accgttttcg cggcatataa ctattatgcg ggtccgaaag aaatccgtat ctatccgtac 900
aacaaccacg aaggcgggtg tagctttcag gctgttgaac aagtgaatt cctgaagaaa 960
ctgtttgaga agggctaa 978

```

```

<210> SEQ ID NO 40
<211> LENGTH: 978
<212> TYPE: DNA
<213> ORGANISM: ARTIFICIAL SEQUENCE
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

```

```

<400> SEQUENCE: 40

```

```

atggcgttct tcgacctgcc tctggaagaa ctgaagaaat accgtccaga tcgttacgaa 60
gagaaggact tcgacgggtt ctgggaggaa actctggcgg agagcgaaaa gtttccgctg 120
gaccagtggt tcgagcgtat ggaatctcac ctgaaaaccg tggaggcata tgacgttact 180
ttttctggtt accgtggcca gcgtatcaaa ggctggctgc tggttccgaa actggaggaa 240
gaaaaactgc cgtgcgtagt tcagtacatc ggttacaacg gtggccgtgg ctttccgcac 300
gattggctgt tctggccgtc tatgggctac atttgcttcg tcatggatac tcgtggctcag 360
ggttccggct ggctgaaagg cgatactccg gattatccgg agggcccggg agaccgcgag 420
taccctggct tcatgacgcg tggattctg gatccgcgta cctattacta tcgccgcggt 480
ttaccgatg cagttcgtgc cgtagaggcc gcgcttctt tacctcaggt tgaccaggag 540
cgtattgtta tcgctggtgg ctcccagggt ggccgcatcg ccctggcggt atctgcgctg 600
agcaagaaag ctaaggcact gctgtgtgac gtcccgttcc tgtgtcactt ccgtcgcgct 660
gttcagctgg tagataccca tccgtacgcg gagattacta acttctgaa aactcaccgc 720
gacaaagaag aaatcgtttt ccgcaccctg tcctatttcg acggcgtaa cttcgcggct 780
cgtgcaaaaa ttccggcact gttctctggt ggtctgatgg acaacatcac ccctccttct 840
accgttttcg cggcatataa ctattatgcg ggtccgaaag aaatccgtat ctatccgtac 900
aacaaccacg aaggcgggtg tagctttcag gctgttgaac aagtgaatt cctgaagaaa 960
ctgtttgaga agggctaa 978

```

```

<210> SEQ ID NO 41
<211> LENGTH: 978
<212> TYPE: DNA
<213> ORGANISM: ARTIFICIAL SEQUENCE
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

```

```

<400> SEQUENCE: 41

```

```

atggcgttct tcgacctgcc tctggaagaa ctgaagaaat accgtccaga gcgttacgaa 60
gagaaggact tcgacgagtt ctgggaggaa actctggcgg agagcgaaaa gtttccgctg 120
gaccagtggt tcgagcgtat ggaatctcac ctgaaaaccg tggaggcata tgacgttact 180
ttttctggtt accgtggcca gcgtatcaaa ggctggctgc tggttccgaa actggaggaa 240
gaaaaactgc cgtgcgtagt tcagtacatc ggttacaacg gtggccgtgg ctttccgcac 300

```

-continued

```

gattggctgt tctggccgtc tatgggctac atttgettgc tcatggatac tctgggtcag 360
ggttccggct ggctgaaagg cgatactccg gattatccgg agggcccggg agaccgcag 420
taccctggct tcatgacgcg tggattctg gatccgcgta cctattacta tgcgccggtt 480
tttaccgatg cagttcgtgc cgtagaggcc gcggttctt tccctcaggt tgaccaggag 540
cgtattgta tgcctgggtg ctcccagggt ggccgcatcg ccctggcggg atctgcgctg 600
agcaagaaag ctaaggcact gctgtgtgac gtcccgttcc tgtgtcactt ccgtcgcgct 660
gttcagctgg tagataccca tccgtacgcg gagattacta acttccctgaa aactcaccgc 720
gacaaagaag aaatcgtttt ccgcaccctg tcctatttgc acggcgttga cttcgcggct 780
cgtgcaaaaa ttccggcact gttctctggt ggtctgatgg acaacatcac ccctccttct 840
accgttttgc cggcatataa ctattatgcg ggtccgaaag aaatccgtat ctatccgtac 900
aacaaccacg aaggcgggtg tagctttcag gctgttgaac aagtgaatt cctgaagaaa 960
ctgtttgaga agggctaa 978

```

```

<210> SEQ ID NO 42
<211> LENGTH: 978
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

```

```

<400> SEQUENCE: 42

```

```

atggcgttct tgcacctgcc tctggaagaa ctgaagaaat accgtccaga gcgttacgaa 60
gagaaggact tgcacgagtt ctgggaggaa actctggcgg agagcgaaaa gtttccgctg 120
gaccagtggt tgcagcgtat ggaatctcac ctgaaaaccg tggaggcata tgacgttact 180
ttttctgggt accgtggccg gcgtatcaaa ggtctgctgc tggttccgaa actggaggaa 240
gaaaaactgc cgtgcgtagt tcagtacatc ggttacaacg gtggccgtgg ctttccgcac 300
gattggctgt tctggccgtc tatgggctac atttgettgc tcatggatac tctgggtcag 360
ggttccggct ggctgaaagg cgatactccg gattatccgg agggcccggg agaccgcag 420
taccctggct tcatgacgcg tggattctg gatccgcgta cctattacta tgcgccggtt 480
tttaccgatg cagttcgtgc cgtagaggcc gcggttctt tccctcaggt tgaccaggag 540
cgtattgta tgcctgggtg ctcccagggt ggccgcatcg ccctggcggg atctgcgctg 600
agcaagaaag ctaaggcact gctgtgtgac gtcccgttcc tgtgtcactt ccgtcgcgct 660
gttcagctgg tagataccca tccgtacgcg gagattacta acttccctgaa aactcaccgc 720
gacaaagaag aaatcgtttt ccgcaccctg tcctatttgc acggcgttaa cttcgcggct 780
cgtgcaaaaa ttccggcact gttctctggt ggtctgatgg acaacatcac ccctccttct 840
accgttttgc cggcatataa ctattatgcg ggtccgaaag aaatccgtat ctatccgtac 900
aacaaccacg aaggcgggtg tagctttcag gctgttgaac aagtgaatt cctgaagaaa 960
ctgtttgaga agggctaa 978

```

```

<210> SEQ ID NO 43
<211> LENGTH: 978
<212> TYPE: DNA
<213> ORGANISM: ARTIFICIAL SEQUENCE
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

```

```

<400> SEQUENCE: 43

```

```

atggcgttct tgcacctgcc tctggaagaa ctgaagaaat accatccaga gcgttacgaa 60

```

-continued

```

gagaaggact tgcagcagtt ctgggaggaa actctggcgg agagcga aaa gtttccgctg 120
gaccagtggt tgcagcagtt ggaatctcac ctgaaaaccg tggaggcata tgacgttact 180
ttttctgggt accgtggcca gcgtatcaaa ggctggctgc tggttccgaa actggaggaa 240
gaaaaactgc cgtgcgtagt tcagtacatc ggttacaacg gtggccgtgg ctttccgcac 300
gattggctgt tctggccgtc tatgggctac atttgcttcg tcatggatac tctgggtcag 360
ggttccggct ggctgaaagg cgatactccg gattatccgg agggcccggg agaccgcag 420
taccctggct tcatgacgcg tggattctg gatccgcgta cctattacta tgcgccgctt 480
tttaccgatg cagttcgtgc cgtagaggcc gcggttctt tccctcatgt tgaccaggag 540
cgtattgta tgcctgggtg ctcccagggt ggcggcatcg ccctggcggg atctgcgctg 600
agcaagaaaag ctaaggcact gcagtgtagt gtcccgttc tgtgtcactt ccgtcgcgct 660
gttcagctgg tagatacca tccgtacgcg gagattacta acttcctgaa aactcaccgc 720
gacaaagaag aaatcgttt cgcaccctg tccattctg acggcgtaa cttcgcggct 780
cgtgcaaaaa ttccggcact gttctctggt ggtctgatgg acaacatcac cctccttct 840
accgttttcg cggcatataa ctattatgcg ggtccgaaag aaatccgtat ctatccgtac 900
aacaaccacg aaggcgggtg tagctttcag gctgttgaac aagtgaatt cctgaagaaa 960
ctgtttgaga agggctaa 978

```

```

<210> SEQ ID NO 44
<211> LENGTH: 978
<212> TYPE: DNA
<213> ORGANISM: ARTIFICIAL SEQUENCE
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

```

```

<400> SEQUENCE: 44

```

```

atggcgttct tgcacctgcc tctggaagaa ctgaagaaat accgtccaga gcgttacgaa 60
gagaaggact tgcagcagtt ctgggaggaa actctggcgg agagcga aaa gtttccgctg 120
gaccagtggt tgcagcagtt ggaatctcac ctgaaaaccg tggaggcata tgacgttact 180
ttttctgggt accgtggcca gcgtatcaaa ggctggctgc tggttccgaa actggaggaa 240
gaaaaactgc cgtgcgtagt tcagtacatc ggttacaacg gtggccgtgg ctttccgcac 300
gattggctgt tctggccgtc tatgggctac atttgcttcg tcatggatac tctgggtcag 360
ggttccggct ggctgaaagg cgatactccg gattatccgg agggcccggg agaccgcag 420
taccctggct tcatgacgcg tggattctg gatccgcgta cctattacta tgcgccgctt 480
tttaccgatg cagttcgtgc cgtagaggcc gcggttctt tccctcaggt tgaccaggag 540
cgtattgta tgcctgggtg ctcccagggt ggcggcatcg ccctggcggg atctgcgctg 600
agcaagaaaag ctaaggcact gctgtgtgac gtcccgttc tgtgtcactt ccgtcgcgct 660
gttcagctgg tagatacca tccgtacgcg gagattacta acttcctgaa aactcaccgc 720
gacaaagaag aaatcgttt cgcaccctg tccattctg acggcgtaa cttcgcggct 780
cgtgcaaaaa ttccgggact gttctctggt ggtctgatgg acaacatcac cctccttct 840
accgttttcg cggcatataa ctattatgcg ggtccgaaag aaatccgtat ctatccgtac 900
aacaaccacg aaggcgggtg tagctttcag gctgttgaac aagtgaatt cctgaagaaa 960
ctgtttgaga agggctaa 978

```

```

<210> SEQ ID NO 45
<211> LENGTH: 978

```

-continued

<212> TYPE: DNA
 <213> ORGANISM: ARTIFICIAL SEQUENCE
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 45

atggcgttct tcgacctgcc tctggaagaa ctgaagaaat accgtccaga gcgttacgaa	60
gagaaggact tcgacgagtt ctgggaggaa actctggcgg agagcgaaaa gtttccgctg	120
gaccagtgt tcgagcgtat ggaatctcac ctgaaaaccg tggaggcata tgacgttact	180
ttttctgggt accgtggcca gcgatcaaaa ggctggctgc tggttccgaa actggaggaa	240
gaaaaactgc cgtgcgtagt tcagtacatc ggttacaacg gtggccgtgg ctttccgcac	300
gattggctgt tctggccgtc tatgggctac atttgcctcg tcatggatac tctgggtcag	360
ggttccggct ggctgaaagg cgatactccg gattatccgg agggcccggg agaccgcag	420
tacctggct tcatgacgcg tggattctg gatccgcgta cctattacta tgcgccgctt	480
tttaccgatg cagttcgtgc cgtagaggcc gcgcttctt tccctcaggt tgaccaggag	540
cgtattgta tctctgggtg ctcccagggt ggcggcatcg ccctggcggg atctgcgctg	600
agcaagaaag ctaaggcact gctgtgtgac gtcccgttcc tgtgtcactt ccgtcgcgct	660
gttcagctgg tagataccca tccgtacgcg gagattacta acttctgaa aactcaccgc	720
gacaaagaag aaatcgtttt ccgcaccctg tcctattctg acggcgtaa cttcgcggct	780
cgtgcaaaaa ttccggcact gttctctgct ggtctgatgg acaacatcac ccctccttct	840
accgtttctg cggcatataa ctattatgcg ggtccgaaag aaatccgtat ctatccgtac	900
aacaaccacg aaggcgggtg tagctttcag gctgttgaac aagtgaaatt cctgaagaaa	960
ctgtttgaga agggctaa	978

<210> SEQ ID NO 46
 <211> LENGTH: 978
 <212> TYPE: DNA
 <213> ORGANISM: ARTIFICIAL SEQUENCE
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 46

atggcgttct tcgacctgcc tctggaagaa ctgaagaaat accgtccaga gcgttacgaa	60
gagaaggact tcgacgagtt ctgggaggaa actctggcgg agagcgaaaa gtttccgctg	120
gaccagtgt tcgagcgtat ggaatctcac ctgaaaaccg tggaggcata tgacgttact	180
ttttctgggt accgtggcca gcgatcaaaa ggctggctgc tggttccgaa actggaggaa	240
gaaaaactgc cgtgcgtagt tcagtacatc ggttacaacg gtggccgtgg ctttccgcac	300
gattggctgt tctggccgtc tatgggctac atttgcctcg tcatggatac tctgggtcag	360
ggttccggct ggctgaaagg cgatactccg gattatccgg agggcccggg agaccgcag	420
tacctggct tcatgacgcg tggattctg gatccgcgta cctattacta tgcgccgctt	480
tttaccgatg cagttcgtgc cgtagaggcc gcgcttctt tccctcaggt tgaccaggag	540
cgtattgta tctctgggtg ctcccagggt ggcggcatcg ccctggcggg atctgcgctg	600
agcaagaaag ctaaggcact gctgtgtgac gtcccgttcc tgtgtcactt ccgtcgcgct	660
gttcagctgg tagataccca tccgtacgcg gagattacta acttctgaa aactcaccgc	720
gacaaagaag aaatcgtttt ccgcaccctg tcctattctg acggcgtaa cttcgcggct	780
cgtgcaaaaa ttccggcact gttctctgct ggtctgatgg acaacatcac ccctccttct	840
accgtttctg cggcatataa ctattatgcg ggtccgaaag aaatccctat ctatccgtac	900

-continued

aacaaccacg aaggcgggtg tagctttcag gctggtgaac aagtgaatt cctgaagaaa 960
ctgtttgaga agggctaa 978

<210> SEQ ID NO 47
<211> LENGTH: 978
<212> TYPE: DNA
<213> ORGANISM: ARTIFICIAL SEQUENCE
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 47

atggcgttct tcgacctgcc tctggaagaa ctgaagaaat accgtccaga gcgttacgaa 60
gagaaggact tcgacgagtt ctgggaggaa actctggcgg agagcgaaaa gtttccgctg 120
gaccacgtgt tcgagcgtat ggaatctcac ctgaaaaccg tggaggcata tgacgttact 180
ttttctgggt accgtggcca gcgtatcaaa ggctggctgc tggttccgaa actggaggaa 240
gaaaaactgc cgtgcgtagt tcagtacatc ggttacaacg gtggccgtgg ctttccgcac 300
gattggctgt tctggccgtc tatgggctac atttgcttcg tcatggatac tcgtgggtcag 360
ggttccggct ggctgaaagg cgatactccg gattatccgg agggcccggg agaccgcgag 420
tacctggct tcatgacgcg tggattctg gatccgcgta cctattacta tcgcccgtt 480
tttaccgatg cagttcgtgc cgtagaggcc gcggcttctt tccctcaggt tgaccaggag 540
cgtattgtta tcgctgggtg ctcccagggt ggccgcatcg ccctggcggg atctgcgctg 600
agcaagaaag ctaaggcact gctgtgtgac gtcccgttcc tgagtcactt ccgtccgct 660
gttcagctgg tagataccca tccgtacgcg gagattacta acttctctgaa aactcaccgc 720
gacaaagaag aaatcgtttt ccgcaccctg tcctattctg acggcgtaa cttcgcggct 780
cgtgcaaaaa ttccggcact gttctctggt ggtctgatgg acaacatcac cctccttct 840
accgttttcg cggcatataa ctattatgcg ggtccgaaag aaatccgtat ctatccgtac 900
aacaaccacg aaggcgggtg tagctttcag gctggtgaac aagtgaatt cctgaagaaa 960
ctgtttgaga agggctaa 978

What is claimed is:

1. An isolated polypeptide having perhydrolytic activity 45
comprising the amino acid sequence of SEQ ID NO: 18.

2. The polypeptide of claim 1, wherein said polypeptide is
characterized by a peracetic acid formation specific activity
that is higher than the peracetic acid formation specific activ- 50
ity of the *Thermotoga maritima* C277T acetyl xylan esterase
having amino acid sequence SEQ ID NO: 5.

3. A process for producing a peroxy-carboxylic acid com-
prising:

(a) providing a set of reaction components comprising:

(1) at least one substrate selected from the group con- 55
sisting of:

(i) one or more esters having the structure

$[X]_m R_5$

wherein


X=an ester group of the formula $R_6-C(O)O$;

$R_6=C1$ to $C7$ linear, branched or cyclic hydrocar- 60
byl moiety, optionally substituted with hydroxyl
groups or $C1$ to $C4$ alkoxy groups, wherein R_6 65
optionally comprises one or more ether linkages
for $R_6=C2$ to $C7$;

R_5 =a $C1$ to $C6$ linear, branched, or cyclic hydro-
carbyl moiety or a five-membered cyclic het-
eroaromatic moiety or six-membered cyclic aro-
matic or heteroaromatic moiety optionally
substituted with hydroxyl groups; wherein each
carbon atom in R_5 individually comprises no
more than one hydroxyl group or no more than
one ester group or carboxylic acid group;
wherein R_5 optionally comprises one or more
ether linkages;

m is an integer ranging from 1 to the number of
carbon atoms in R_5 ; and wherein said esters have
solubility in water of at least 5 ppm at 25° C.;


(ii) one or more glycerides having the structure



wherein $R_1=C1$ to $C21$ straight chain or branched
chain alkyl optionally substituted with an

109

- hydroxyl or a C1 to C4 alkoxy group and R_3 and R_4 are individually H or $R_1C(O)$;
 (iii) one or more esters of the formula:



wherein R_1 is a C1 to C7 straight chain or branched chain alkyl optionally substituted with an hydroxyl or a C1 to C4 alkoxy group and R_2 is a C1 to C10 straight chain or branched chain alkyl, alkenyl, alkynyl, aryl, alkylaryl, alkylheteroaryl, heteroaryl, $(CH_2CH_2O)_n$, or $(CH_2CH(CH_3)-O)_nH$ and n is 1 to 10;

- (iv) one or more acylated monosaccharides, acylated disaccharides, or acylated polysaccharides; and
 (v) any combination of (i) through (iv);
 (2) a source of peroxygen; and
 (3) an enzyme catalyst comprising the polypeptide of claim 1;
 (b) combining the set of reaction components under suitable reaction conditions whereby peroxycarboxylic acid is produced; and
 (c) optionally diluting the peroxycarboxylic acid produced in step (b).

4. The process of claim 3 further comprising the step of: d) contacting a hard surface or inanimate object with the peroxycarboxylic acid produced in step (b) or step (c); whereby said hard surface or said inanimate object is disinfected, bleached, destained or a combination thereof.

5. The process of claim 3 wherein the inanimate object is a medical instrument.

6. The process of claim 3 further comprising the step of: d) contacting an article of clothing or a textile with peroxycarboxylic acid produced in step (b) or step (c); whereby the article of clothing or textile receives a benefit.

7. The process of claim 6 wherein the benefit is selected from the group consisting of disinfecting, sanitizing, bleaching, destaining, deodorizing, and combinations thereof.

8. The process of claim 3 further comprising the step of: d) contacting wood pulp or paper pulp with peroxycarboxylic acid produced in step (b) or step (c); whereby the wood pulp or paper pulp is bleached.

9. The process of claim 3 wherein the substrate is selected from the group consisting of: monoacetin; diacetin; triacetin; monopropionin; dipropionin; tripropionin; monobutyryn; dibutyryn; tributyrin; glucose pentaacetate; β -D-galactose pentaacetate; sorbitol hexaacetate; sucrose octaacetate; xylose tetraacetate; acetylated xylan; acetylated xylan fragments; β -D-ribofuranose-1,2,3,5-tetraacetate; tri-O-acetyl-D-galactal; tri-O-acetyl-D-glucal; monoesters or diesters of 1,2-ethanediol; 1,2-propanediol; 1,3-propanediol; 1,2-butanediol; 1,3-butanediol; 2,3-butanediol; 1,4-butanediol; 1,2-pentanediol; 2,5-pentanediol; 1,6-pentanediol; 1,2-hexanediol; 2,5-hexanediol; 1,6-hexanediol; 4-acetoxybenzoic acid; and mixtures thereof.

10. The process of claim 9 wherein the substrate is triacetin.

11. The process of claim 3 wherein the peroxycarboxylic acid produced is peracetic acid, perpropionic acid, perbutyric acid, perlactic acid, perglycolic acid, permethoxyacetic acid, per- β -hydroxybutyric acid, or mixtures thereof.

12. The process of claim 3 wherein the enzyme catalyst is in the form of a microbial cell, a permeabilized microbial cell, a microbial cell extract, a partially purified enzyme, or a purified enzyme.


110

13. A composition comprising:

(a) a set of reaction components comprising:

(1) at least one substrate selected from the group consisting of:

(i) one or more esters having the structure



wherein

X=an ester group of the formula $R_6-C(O)O$;


R_6 =C1 to C7 linear, branched or cyclic hydrocarbyl moiety, optionally substituted with hydroxyl groups or C1 to C4 alkoxy groups, wherein R_6 optionally comprises one or more ether linkages for R_6 =C2 to C7;

R_5 =a C1 to C6 linear, branched, or cyclic hydrocarbyl moiety or a five-membered cyclic heteroaromatic moiety or six-membered cyclic aromatic or heteroaromatic moiety optionally substituted with hydroxyl groups; wherein each carbon atom in R_5 individually comprises no more than one hydroxyl group or no more than one ester group or carboxylic acid group; wherein R_5 optionally comprises one or more ether linkages;

m is an integer ranging from 1 to the number of carbon atoms in R_5 ; and


wherein said esters have solubility in water of at least 5 ppm at 25° C.;

(ii) one or more glycerides having the structure



wherein R_1 =C1 to C21 straight chain or branched chain alkyl optionally substituted with an hydroxyl or a C1 to C4 alkoxy group and R_3 and R_4 are individually H or $R_1C(O)$;

(iii) one or more esters of the formula:



wherein R_1 is a C1 to C7 straight chain or branched chain alkyl optionally substituted with an hydroxyl or a C1 to C4 alkoxy group and R_2 is a C1 to C10 straight chain or branched chain alkyl, alkenyl, alkynyl, aryl, alkylaryl, alkylheteroaryl, heteroaryl, $(CH_2CH_2O)_n$, or $(CH_2CH(CH_3)-O)_nH$ and n is 1 to 10;

(iv) one or more acylated monosaccharides, acylated disaccharides, or acylated polysaccharides; and

(v) any combination of (i) through (iv);

(2) a source of peroxygen; and

(3) an enzyme catalyst comprising the polypeptide of claim 1; and

(b) at least one peroxycarboxylic acid formed upon combining the set of reaction components of (a).

111

14. A peracid generation and delivery system comprising:

- (a) a first compartment comprising
- (1) an enzyme catalyst comprising the polypeptide of claim 1;
 - (2) at least one substrate selected from the group consisting of:
 - (i) one or more esters having the structure

$[X]_m R_5$

wherein

X=an ester group of the formula $R_6-C(O)O$;


$R_6=C1$ to $C7$ linear, branched or cyclic hydrocarbyl moiety, optionally substituted with hydroxyl groups or $C1$ to $C4$ alkoxy groups, wherein R_6 optionally comprises one or more ether linkages for $R_6=C2$ to $C7$;

$R_5=a$ $C1$ to $C6$ linear, branched, or cyclic hydrocarbyl moiety or a five-membered cyclic heteroaromatic moiety or six-membered cyclic aromatic or heteroaromatic moiety optionally substituted with hydroxyl groups; wherein each carbon atom in R_5 individually comprises no more than one hydroxyl group or no more than one ester group or carboxylic acid group; wherein R_5 optionally comprises one or more ether linkages;

m is an integer ranging from 1 to the number of carbon atoms in R_5 ; and

wherein said esters have solubility in water of at least 5 ppm at $25^\circ C$.;


- (ii) one or more glycerides having the structure



112

wherein $R_1=C1$ to $C21$ straight chain or branched chain alkyl optionally substituted with an hydroxyl or a $C1$ to $C4$ alkoxy group and R_3 and R_4 are individually H or $R_1C(O)$;

- (iii) one or more esters of the formula:



wherein R_1 is a $C1$ to $C7$ straight chain or branched chain alkyl optionally substituted with an hydroxyl or a $C1$ to $C4$ alkoxy group and R_2 is a $C1$ to $C10$ straight chain or branched chain alkyl, alkenyl, alkynyl, aryl, alkylaryl, alkylheteroaryl, heteroaryl, $(CH_2CH_2O)_n$, or $(CH_2CH(CH_3)-O)_nH$ and n is 1 to 10;

- (iv) one or more acylated monosaccharides, acylated disaccharides, or acylated polysaccharides; and
- (v) any combination of (i) through (iv); and
- (3) an optional buffer; and
- (b) a second compartment comprising
- (1) source of peroxygen;
 - (2) a peroxide stabilizer; and
 - (3) an optional buffer.

15. The peracid generation and delivery system of claim 14 wherein the substrate comprises triacetin.

16. A laundry care product comprising the polypeptide of claim 1.

17. A personal care product comprising the polypeptide of claim 1.

* * * * *