



US008748700B2

(12) **United States Patent**
Hanger et al.

(10) **Patent No.:** **US 8,748,700 B2**
(45) **Date of Patent:** ***Jun. 10, 2014**

(54) **CONTROL OF AAD-1 MONOCOT VOLUNTEERS IN FIELDS OF DICOT CROPS**

(75) Inventors: **Gregory A. Hanger**, Carmel, IN (US); **Andrew E. Robinson**, Brownsburg, IN (US); **Norbert M. Satchivi**, Westfield, IN (US); **Richard S. Chambers**, Warriewood (AU); **Terry R. Wright**, Carmel, IN (US)

(73) Assignee: **Dow AgroSciences, LLC.**, Indianapolis, IN (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **13/390,988**

(22) PCT Filed: **Aug. 18, 2010**

(86) PCT No.: **PCT/US2010/045870**

§ 371 (c)(1),
(2), (4) Date: **May 7, 2012**

(87) PCT Pub. No.: **WO2011/022470**

PCT Pub. Date: **Feb. 24, 2011**

(65) **Prior Publication Data**

US 2012/0220460 A1 Aug. 30, 2012

Related U.S. Application Data

(60) Provisional application No. 61/328,944, filed on Apr. 28, 2010, provisional application No. 61/235,248, filed on Aug. 19, 2009.

(51) **Int. Cl.**
C12N 15/82 (2006.01)

(52) **U.S. Cl.**
USPC **800/300; 800/288**

(58) **Field of Classification Search**
USPC **800/300**
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

7,838,733 B2 * 11/2010 Wright et al. 800/300
2007/0089201 A1 4/2007 Briggs et al.
2009/0093366 A1 4/2009 Wright et al.

FOREIGN PATENT DOCUMENTS

WO WO 99/45781 A1 9/1999
WO WO 2005/107437 A2 11/2005
WO WO 2007/053482 A2 5/2007
WO WO 2008/143993 A2 11/2008

OTHER PUBLICATIONS

Hohe et al, A tool for understanding homologous recombination in plants (2003) Plant Cell Rep. 21:1135-1142.*

Deen et al, Control of volunteer glyphosate-resistant corn (*Zea mays*) in glyphosate-resistant soybean (2006) Weed Tech. 20:261-266.*

Loux et al, Weed control guide for Ohio and Indiana (Jan. 23, 2009), published by Ohio State University.*

Chander, et al., Genetic dissection of tocopherol content and composition in maize grain using quantitative trait loci analysis and the candidate gene approach, Molecular Breeding, vol. 22, No. 3, Apr. 12, 2008, pp. 353-365.

Zhang, et al., Mapping quantitative trait loci for oil, starch, and protein concentrations in grain with high-oil maize by SSR markers, Euphytica, Kluwer Academic Publishers, vol., 162, No. 3, Aug. 3, 2007, pp. 335-344.

Antonius J.M. Matzke et al, Position effects and epigenetic silencing of plant transgenes, Current Opinion in Plant Biology, vol. 1, No. 2, Apr. 1, 1998, pp. 142-148.

Dow Chemical Japan Ltd., "Approved Type 1 Use Regulation: Maize tolerant to aryloxyalkanoate herbicide (Modified aad-1, *Zea mays* subsp.," Japan Biosafety Clearing House, Online Jul. 30, 2009, pdf also available at <url:http://www.bch.biodic.go.jp/english/lmo_2009.html>.

Tan et al., "Imidazolinone-tolerant crops: history, current status and future," Pest Manag Sci, Online Dec. 31, 2004, pp. 246-257, vol. 61, Issue 3.

* cited by examiner

Primary Examiner — David H Kruse

Assistant Examiner — Mykola Kovalenko

(74) *Attorney, Agent, or Firm* — James Daly; Kenneth B. Ludwig; Faegre Baker Daniels LLP

(57) **ABSTRACT**

The subject invention relates in part to the control of AAD-1 monocot volunteers in fields planted with dicot crops such as soybeans or cotton. According to some embodiments of the subject invention, cyclohexanedione herbicides are selected as being an effective tool for controlling AAD-1 volunteers, as AAD genes do not impart tolerance to this class of gramini-cide chemistry. In addition, imidazolinone-class herbicides can be used in some preferred embodiments for selective control of conventional or herbicide-tolerant varieties of volunteer corn. AAD-1 corn comprising Event DAS-40278-9 is used in some particularly preferred embodiments.

21 Claims, No Drawings

CONTROL OF AAD-1 MONOCOT VOLUNTEERS IN FIELDS OF DICOT CROPS

CROSS REFERENCE TO RELATED APPLICATIONS

This application is a 371 national stage application of International Application No. PCT/US2010/045870, with an International Filing Date of Aug. 18, 2010, which claims priority to U.S. Application No. 61/235,248, filed on Aug. 19, 2009, and to U.S. Application No. 61/328,944, filed on Aug. 28, 2010, all of which are incorporated herein by reference in their entirety.

BACKGROUND

Corn (monocot) and soybeans (dicot), for example, can be rotated in various crop rotation cycles in various geographies. Cotton is also a dicot.

“Volunteer” plants are unwanted plants from the prior growing season that emerge in a field planted with crops for the current growing season. Volunteers are basically weeds, and can, like weeds, reduce harvest and yield of the crop of interest for the current growing season. The volunteers divert fertilizer resources and the like from the desired crops.

Unlike plain weeds, volunteers are often specifically engineered to be resistant to some herbicides. Thus, controlling volunteers can be more difficult than controlling naturally occurring weeds.

AAD (aryloxy alkanoate dioxygenase) genes as described herein impart high levels of tolerance to 2,4-D herbicides in plants that are transformed with an AAD gene.

AAD-1 genes (encoding SEQ ID NO:1, SEQ ID NO:2, and SEQ ID NO:3, for example, of the attached sequence listing; see WO 2005 107437) also impart high levels of tolerance to phenoxy- and aryloxyphenoxypropionate herbicides (“fops” such as fluazifop and haloxyfop) to corn and other monocot species transformed with the gene. (Fops are primarily used to control monocots, as dicots tend to have some natural resistance to fops.) Thus, AAD-1 allows the use of some fops as either selection agents or as herbicides on crops where crop destruction would be expected without the AAD-1 gene.

AAD-12 and AAD-13 genes also impart high levels of tolerance to pyridyloxyacetate herbicides (such as triclopyr and fluroxypyr; “pyrs”). Thus, AAD-12 and AAD-13 each allow the use of pyrs as either selection agents or as herbicides on crops where crop destruction would be expected without the AAD-12 or AAD-13 gene.

There are very numerous types of monocot- or grass-only herbicides (that kill monocots).

ACCase inhibitor herbicides include fops and dims.

BRIEF SUMMARY

The subject invention relates in part to the control of AAD-1 monocot volunteers in fields planted with dicot crops such as soybeans or cotton.

The subject invention also relates in part to the recognition that one potential downside to the use of AAD genes is that volunteers can be resistant to 2,4-D, as well as to fops (in the case of AAD-1). In the case of AAD-1, fop herbicides will no longer be effective for control of volunteer AAD-1 corn in fields planted with dicot crops such as soybean or cotton. The subject invention relates in part to the recognition that when an AAD gene is stacked with other herbicide resistance traits (such as glyphosate, glufosinate, and the like) in corn, control of volunteer corn plants in the following year can be an issue.

According to some embodiments of the subject invention, cyclohexanedione (‘dim’) herbicides (such as clethodim, sethoxydim, and the like) are selected, from almost innumerable

other options, as being an effective tool for controlling AAD-1 ‘volunteers,’ as AAD genes do not impart tolerance to this class of graminicide chemistry. Dims tend to be monocot-specific.

In addition, the following imidazolinone-class herbicides can be used, in some preferred embodiments, according to the subject invention: imazethapyr, imazamox, and imazaquin. Soy is naturally tolerant to these herbicides, so these imidazolinones can be used for selective control of conventional or herbicide-tolerant varieties of volunteer corn. (Non-commercialized CLEARFIELD cotton and imidazolinone-resistant soy was developed by BASF; cotton is not naturally tolerant to imidazolinones.)

This is assuming that the corn from the previous season was susceptible to imidazolinones. (Soybeans are naturally tolerant to certain imidazolinones.) Corn is generally susceptible to imidazolinones except for CLEARFIELD corn, which is tolerant to imidazolinones.

Use of other types of herbicides, including ALS—(acetolactate synthase) and/or AHAS—(acetohydroxyacid synthase) inhibitors could also be used according to the subject invention for the control of volunteer AAD monocot plants as well.

A third option, according to the subject invention, is to use other herbicides, as described herein, for controlling AAD-1 volunteers. These options include herbicides that provide a relatively fast “burn down,” and non-selective herbicides, such as paraquat, used pre-plant.

AAD-1 corn comprising Event DAS-40278-9 is used in some particularly preferred embodiments.

BRIEF DESCRIPTION OF THE SEQUENCES

SEQ ID NO:1 is the native protein sequence encoded by the AAD-1 v1 gene from *Sphingobium herbicidivorans*.

SEQ ID NO:2 is the amino acid sequence encoded by the AAD-1 v2 gene, which removed an internal NotI restriction site.

SEQ ID NO:3 is the amino acid sequence encoded by the AAD-1 v3 gene, which has plant codon usage for expression in a plant.

SEQ ID NO:4 is a sequence for corn Event DAS-40278-9, which includes 5' and 3' corn genomic flanking sequences, and the AAD-1 cassette insert sequence.

DETAILED DESCRIPTION

As used herein and unless otherwise specified, preferred dicots are soybeans or cotton.

ACCase inhibitor herbicides include fops and dims.

One aspect of the subject invention includes the use of ‘dim’ herbicides to remove volunteer AAD-1 corn in fields of dicots. In some specific embodiments, the AAD-1 gene is present in the corn as the AAD-1 corn event designated DAS-40278-9 having seed deposited with American Type Culture Collection (ATCC) with Accession No. PTA-10244 (Yellow Dent maize hybrid seed (*Zea Mays* L.):DAS-40278-9; Deposited in accordance with the Budapest Treaty on behalf of Dow AgroSciences LLC; Date of receipt of seeds/strain(s) by the ATTC: Jul. 10, 2009; viability confirmed Aug. 17, 2009), and progeny derived thereof. See also SEQ ID NO:4. This event is disclosed in U.S. Ser. No. 61/235,248 (filed Aug. 19, 2009). Such AAD-1 monocot volunteers could be present in the following year’s soybeans, cotton, or other broadleaf crop. Thus, the subject invention includes the application of a dim (or other as disclosed herein) herbicide to volunteer AAD-1 corn plants, particularly where the corn plant comprises the 40278-9 event.

Because of the specific detoxifying aspects of AAD genes, ‘dim’ herbicides would not be subject to detoxification by AAD-1, and AAD-1 monocot volunteers would remain

highly susceptible to dim herbicides. Thus, the subject invention includes the use of one or more dim herbicides, where the AAD-1 gene was used alone or stacked with other herbicide tolerance traits.

In some other preferred embodiments, imidazolinones can be used according to the subject invention to control monocot/corn volunteers in dicot fields. If the corn was susceptible to an imidazolinone and the dicots are resistant to imidazolinones, then a “dim” and/or imidazolinone herbicide could be used in the dicot fields to control the AAD-1 corn volunteers. However, if CLEARFIELD corn (which is resistant to imidazolinones), for example, was planted in the previous season, this would tend to preclude the use of imidazolinones the following season to control corn volunteers in dicot fields.

As alluded to above, selection of herbicides to use on the current planted fields, according to the subject invention, depends in part on the herbicide-tolerant trait(s) that are present in both the volunteers (the monocot crop of the prior season), if any, and the herbicide resistance trait(s) that are present in the field of dicot crops of the current growing season. Thus, additional herbicide chemistries can be selected to provide control of AAD-1 corn depending on the stack, if any, with other herbicide tolerant traits in the corn, and on the tolerance or susceptibility of the dicot (soybeans or cotton) in the field to be treated.

For example, if the volunteer AAD-1 corn was also stacked with a glyphosate- or glufosinate-trait, then glufosinate or glyphosate, respectively, could be used in the soybean field of the current season, assuming that the soybeans also have a resistance trait against glufosinate or glyphosate, respectively.

If the volunteer AAD-1 corn was also stacked with a glyphosate- and glufosinate-traits, then dims and/or imidazolinones could still be used in the soybean fields to control the AAD-1 corn volunteers (again assuming that the corn does not have a resistance trait against an imidazolinone if an imidazolinone is used to control the volunteers).

When an AAD-1 gene was used alone in corn in the previous growing season, imidazolinone, “dims,” glyphosate, and/or glufosinate could be used the following season to control the volunteer AAD-1 corn in planted fields of soybean or cotton that, assuming that the dicots are tolerant/resistant to any one or more of these further herbicides (such as glyphosate and/or glufosinate). That is, glyphosate and/or glufosinate could be used if the soybeans or cotton are engineered or bred to be tolerant to glyphosate and/or glufosinate, respectively.

When an AAD-1 gene was stacked in corn with, for example, a glyphosate or glufosinate resistant trait (such as is found in Roundup Ready [or GAT or other glyphosate-tolerant crops] or Liberty Link [or other glufosinate-tolerant] corn), then imidazolinone and/or a “dim” herbicide could be used to control volunteer AAD-1 corn in the dicot crops (assuming the dicot crop is resistant to imidazolinone if an imidazolinone herbicide is to be used).

Glufosinate or glyphosate, respectively, could also be used, assuming that the dicot crops are resistant to that herbicide and the corn is susceptible. That is, if the soybeans have a trait for glyphosate tolerance but not glufosinate tolerance, for example, then glyphosate could also be used (assuming ROUNDUP READY corn is not the volunteers). If the soybeans have a glufosinate tolerance trait (such as PAT) but not a glyphosate tolerance trait, then glufosinate could also be used (assuming that LIBERTY LINK corn is not the volunteers).

In any of those stacking scenarios (and even if AAD-1, glyphosate tolerance, and glufosinate tolerance, and imidazolinone tolerance traits were all present in the corn of the previous season), dims could be used in the soybean fields to control the AAD-1 corn volunteers.

In some embodiments, AAD-1 monocot volunteers can be controlled, using combinations of herbicides disclosed herein—see Examples 4 and 5, in fields of monocots. These embodiments include control of AAD-1 Event DAS-40278-9 corn volunteers in fields of monocots. Such embodiments include turf-in-turf (AAD-1 turf volunteers in a field of other (non-AAD1) turf).

EXAMPLES

Example 1

Control of Volunteer AAD Maize in a Field Planted with Soybean Using Alternative Herbicides

In one embodiment, volunteer transgenic maize lines containing an AAD expression cassette (AAD-1) are controlled within a field of soybean by the application of a herbicide or combination of herbicides. The specific herbicide used to control the volunteer transgenic AAD maize line is dependent upon the type of soybean seed being planted within the field (e.g., conventional soybean, Round-up Ready Soybean, Liberty Link Soybean, etc.).

Furthermore, the AAD trait may be stacked with other additional herbicide tolerant trait(s) via conventional breeding or a molecular stack. In such an example, the specific herbicide used to control the volunteer AAD maize stacked with another herbicide tolerant trait(s) will be dependent upon the additional herbicide tolerant trait(s) and the type of soybean being planted within the field.

The application of a given herbicide can be made before planting at pre-emergence/burndown or post-emergence after planting to control the volunteer transgenic AAD maize lines. Table 1 lists the herbicides to be used at the different stages of planting (pre-emergence or post-emergence) to control volunteer transgenic AAD maize. At or about a 1× Field Rate concentration of herbicide would be applied, as either a tank mix or alone, to the field for both pre-emergent and post-emergent volunteer control.

The control of transgenic maize plants containing the AAD expression cassette within a field of soybean would be applicable for the control of volunteer AAD transgenic monocot plants (including, but not limited too; corn, rice, sugar cane, switch grass, turf grass species, sorghum, barley, wheat, and oats, and durum) within a field being planted with a dicot crop (including, but not limited too; soybean, cotton, canola, flax, sunflower, legumes, alfalfa, peanut, and tomato). The example described above, in which volunteer transgenic AAD maize plants are controlled in a soybean field, is illustrative of the invention and not intended to restrict the scope of this embodiment.

Example 2

Control of Volunteer Corn (Conventional or Containing Non-AAD Herbicide Tolerance Traits) in a Field Planted with AAD Soybean Using Alternative Herbicides

In an embodiment, volunteer transgenic corn lines (containing the Clear Field trait, Roundup Ready or other Glyphosate Tolerant Trait, Liberty Link Trait, Imidazolinone tolerant trait, or any stacked combination thereof) or volunteer conventional corn lines are controlled within a field of transgenic AAD soybean (AAD-12) by the application of a herbicide or combination of herbicides. The specific herbicide used to control the volunteer corn plants is dependent upon the type of corn seed being planted within the field (e.g., conventional soybean, Round-up Ready Soybean, Liberty Link Soybean, etc.). Additionally, the specific herbicide used

to control the volunteer conventional or transgenic corn line is dependent upon the type of AAD transgenic soybean seed (i.e. stacked traits or alone) being planted within the field and the trait possessed by the volunteer corn line.

The application of a given herbicide can be made before planting at pre-emergence/burndown or post-emergence after planting to control the volunteer conventional or transgenic soybean lines. Table 2 and Table 3 list the herbicides to be used at the different stages of planting (pre-emergence or post-emergence) to control volunteer conventional or transgenic corn. At or about a 1x Field Rate concentration of herbicide would be applied, as either a tank mix or alone, to the field for both pre-emergent and post-emergent volunteer control.

The control of conventional or transgenic corn plants containing a herbicide tolerant expression cassette within a field of AAD transgenic soybean (either stacked with other herbicide tolerant traits or alone) would be applicable for the control of a conventional or herbicide tolerant transgenic monocot plant (including, but not limited too; corn, rice, sugar cane, switch grass, turf grass species, sorghum, barley, wheat, and oats, and durum) within a field being planted with an AAD transgenic dicot crop (including, but not limited too; soybean, cotton, canola, flax, sunflower, legumes, alfalfa, peanut, and tomato). The example described above, in which volunteer conventional or transgenic herbicide tolerant corn plants are controlled in a field planted with AAD transgenic soybean, is illustrative of the invention and not intended to restrict the scope of this embodiment.

Example 3

Control of Volunteer AAD-1 Maize in a Field Planted with AAD-12 Cotton Using Alternative Herbicides

In one embodiment, volunteer transgenic maize lines containing an AAD-1 expression cassette are controlled within a

field of AAD-12 cotton by the application of a herbicide or combination of herbicides. The specific herbicide used to control the volunteer transgenic AAD-1 maize line is dependent upon the type of AAD-12 cotton seed being planted within the field (e.g., AAD-12 cotton or AAD-12 cotton stacked with Round-up Ready, Liberty Link, or other herbicide tolerant traits).

Furthermore, the AAD-1 trait may be stacked with other additional herbicide tolerant trait(s) via conventional breeding or a molecular stack. In such an example, the specific herbicide used to control the volunteer AAD-1 maize stacked with another herbicide tolerant trait(s) will be dependent upon the additional herbicide tolerant trait(s) and the type of cotton being planted within the field.

The application of a given herbicide can be made before planting at pre-emergence/burndown or post-emergence after planting to control the volunteer transgenic AAD-1 maize lines. Table 4 lists the herbicides to be used at the different stages of planting (pre-emergence or post-emergence) to control volunteer transgenic AAD-1 maize. At or about a 1x Field Rate concentration of herbicide would be applied, as either a tank mix or alone, to the field for both pre-emergent and post-emergent volunteer control.

The control of transgenic maize plants containing the AAD-1 expression cassette within a field of AAD-12 cotton would be applicable for the control of volunteer AAD-1 transgenic monocot plants (including, but not limited too; corn, rice, sugar cane, switch grass, turf grass species, sorghum, barley, wheat, and oats, and durum) within a field being planted with an AAD-12 dicot crop (including, but not limited too; soybean, cotton, canola, flax, sunflower, legumes, alfalfa, peanut, and tomato). The example described above, in which volunteer transgenic AAD-1 maize plants are controlled in an AAD-12 cotton field, is illustrative of the invention and not intended to restrict the scope of this embodiment.

TABLE 1

Control of volunteer AAD-1 (trait alone or stacked with other HT traits) corn (or other monocot) in soybean (or other dicot crops)			
Previous year Corn hybrid	Current year Soybean variety being planted	Herbicides option - Burndown/PRE	Herbicides option - POST
Application: alone and tank mixes			
AAD-1	Soybean, conventional	Glyphosate, Dims, IMI,	Dims, IMI
	RR Soybean	glufosinate, paraquat,	Dims, Glyphosate, IMI
	LL Soybean	other ALS herbicides not	Dims, Glufosinate, IMI
	GAT + ALS	recommended for corn (need to be aware of planting restrictions) enabled by "ALS gene"	Dims, IMI, ALS inhibitors not selective for in-corn application
	AAD-12		Dims, IMI
	AAD-12 + GT		Dims, Glyphosate, IMI
	AAD-12 + PAT		Dims, Glufosinate, IMI
	AAD-12 + AHAS		Dims, IMI
	AAD-12 + PAT + GT		Dims, Glyphosate, glufosinate, IMI
	AAD-12 + PAT + AHAS		Dims, glufosinate, IMI
AAD-12 + GAT		Dims, IMI, glyphosate, ALS inhibitors not selective for in-corn application	
AAD-12 + PAT + GAT		Dims, IMI, glufosinate glyphosate, ALS inhibitors not selective for in-corn application	

TABLE 1-continued

Control of volunteer AAD-1 (trait alone or stacked with other HT traits) corn (or other monocot) in soybean (or other dicot crops)			
Previous year Corn hybrid	Current year Soybean variety being planted	Herbicides option - Burndown/PRE	Herbicides option - POST
AAD-1 + PAT	Soybean, conventional	Glyphosate, Dims, IMI, paraquat, other ALS herbicides not recommended for corn (need to be aware of planting restrictions) enabled by "ALS gene"	Dims, IMI
	RR Soybean		Dims, Glyphosate, IMI
	LL Soybean		Dims, IMI
	GAT		Dims, IMI, ALS inhibitors not selective for in-corn application
	AAD-12		Dims, IMI
	AAD-12 + GT		Dims, Glyphosate, IMI
	AAD-12 + PAT		Dims, IMI
	AAD-12 + AHAS		Dims, IMI
	AAD-12 + PAT + GT		Dims, Glyphosate, IMI
	AAD-12 + PAT + AHAS		Dims, IMI
AAD-12 + GAT	Dims, IMI, glyphosate, ALS inhibitors not selective for in- corn application		
AAD-12 + PAT + GAT	Dims, IMI, glyphosate, ALS inhibitors not selective for in- corn application		
AAD-1 + GT	Soybean, conventional	Dims, IMI, glufosinate, paraquat, other ALS herbicides not recommended for corn (need to be aware of planting restrictions) enabled by "ALS gene"	Dims, IMI
	RR Soybean		Dims, IMI
	LL Soybean		Dims, Glufosinate IMI
	GAT		Dims, IMI, ALS inhibitors not selective for in-corn application
	AAD-12		Dims, IMI
	AAD-12 + GT		Dims, IMI
	AAD-12 + PAT		Dims, Glufosinate, IMI
	AAD-12 + AHAS		Dims, IMI
	AAD-12 + PAT + GT		Dims, glufosinate, IMI
	AAD-12 + PAT + AHAS		Dims, glufosinate, IMI
AAD-12 + GAT	Dims, IMI, ALS inhibitors not selective for in-corn application		
AAD-12 + PAT + GAT	Dims, IMI, glufosinate, ALS inhibitors not selective for in- corn application		
AAD-1 + AHAS	Soybean, conventional	Glyphosate, Dims, glufosinate, paraquat, other ALS herbicides not recommended for corn (need to be aware of planting restrictions) enabled by "ALS gene"	Dims
	RR Soybean		Dims, Glyphosate
	LL Soybean		Dims, Glufosinate
	GAT		Dims, ALS inhibitors not selective for in-corn application
	AAD-12		Dims
	AAD-12 + GT		Dims, Glyphosate
	AAD-12 + PAT		Dims, Glufosinate
	AAD-12 + AHAS		Dims
	AAD-12 + PAT + GT		Dims, Glyphosate, glufosinate
	AAD-12 + PAT + AHAS		Dims, glufosinate
AAD-12 + GAT	Dims, glyphosate, ALS inhibitors not selective for in- corn application		
AAD-12 + PAT + GAT	Dims, glufosinate glyphosate, ALS inhibitors not selective for in-corn application		
AAD-1 + PAT + GT	Soybean, conventional	Dims, IMI, paraquat, other ALS herbicides not recommended for corn (need to be aware of planting restrictions) enabled by "ALS gene"	Dims, IMI
	RR Soybean		Dims, IMI
	LL Soybean		Dims, Glufosinate, IMI
	GAT		Dims, IMI, ALS inhibitors not selective for in-corn application
	AAD-12		Dims, IMI
	AAD-12 + GT		Dims, IMI
	AAD-12 + PAT		Dims, IMI
	AAD-12 + AHAS		Dims, IMI
	AAD-12 + PAT + GT		Dims, IMI
	AAD-12 + PAT + AHAS		Dims, IMI
AAD-12 + GAT	Dims, IMI, ALS inhibitors not selective for in-corn application		
AAD-12 + PAT + GAT	Dims, IMI, ALS inhibitors not selective for in-corn application		

TABLE 1-continued

Control of volunteer AAD-1 (trait alone or stacked with other HT traits) corn (or other monocot) in soybean (or other dicot crops)			
Previous year Corn hybrid	Current year Soybean variety being planted	Herbicides option - Burndown/PRE	Herbicides option - POST
AAD-1 + PAT + AHAS	Soybean, conventional	Glyphosate, Dims,	Dims
	RR Soybean	paraquat,	Dims, Glyphosate
	LL Soybean	other ALS herbicides not	Dims
	GAT	recommended for corn (need to be aware of planting restrictions)	Dims, glyphosate, ALS inhibitors not selective for in- corn application
	AAD-12	enabled by "ALS gene"	Dims
	AAD-12 + GT		Dims, Glyphosate
	AAD-12 + PAT		Dims
	AAD-12 + AHAS		Dims
	AAD-12 + PAT + GT		Dims, Glyphosate
	AAD-12 + PAT + AHAS		Dims
	AAD-12 + GAT		Dims
	AAD-12 + PAT + GAT		Dims, glyphosate, ALS inhibitors not selective for in- corn application
AAD-1 + GAT + ALS	Soybean, conventional	Dims, glufosinate,	Dims
	RR Soybean	paraquat,	Dims
	LL Soybean	other ALS herbicides not	Dims, Glufosinate
	GAT + ALS	recommended for corn (need to be aware of planting restrictions)	Dims
	AAD-12	enabled by "ALS gene"	Dims
	AAD-12 + GT		Dims
	AAD-12 + PAT		Dims, Glufosinate
	AAD-12 + AHAS		Dims
	AAD-12 + PAT + GT		Dims, Glufosinate
	AAD-12 + PAT + AHAS		Dims, Glufosinate
	AAD-12 + GAT		Dims
	AAD-12 + PAT + GAT		Dims, Glufosinate
AAD-1 + PAT + GAT + ALS	Soybean, conventional	Dims, paraquat,	Dims
	RR Soybean	other ALS herbicides not	Dims
	LL Soybean	recommended for corn (need to be aware of planting restrictions)	Dims
	GAT + ALS	enabled by "ALS gene"	Dims
	AAD-12		Dims
	AAD-12 + GT		Dims
	AAD-12 + PAT		Dims
	AAD-12 + AHAS		Dims
	AAD-12 + PAT + GT		Dims
	AAD-12 + PAT + AHAS		Dims
	AAD-12 + GAT		Dims
	AAD-12 + PAT + GAT		Dims

Gene and trait footnotes:

GT = any glyphosate specific tolerance trait including Roundup Ready (CP4), TIPS EPSPS (GA21, Glytol, DMMG), Athenix's EPSPS, GAT only without ALS, GOX, glyphosate decarboxylase, etc.

ALS = double mutant ALS gene insensitive to all ALS herbicide chemistries including IMI herbicides.

AAD-1 = Aryloxyalkanoate dioxygenase gene providing tolerance to all commercial phenoxy auxin and all aryloxyphenoxypropionate (fop) herbicides.

AAD-12 = Aryloxyalkanoate dioxygenase gene providing tolerance to phenoxyacetic auxin and pyridyloxyacetic auxin herbicides.

PAT = phosphinothricin acetyltransferase gene providing tolerance to glutamine synthetase inhibitors including, but not limited to, glufosinate. Similar phenotype is provided by genes such as BAR, DSM1, DSM2, et al.

AHAS = imidazolinone specific tolerance gene associated with point mutation at S623 of ALS gene (maize sequence) or equivalent amino acid in other spp (e.g., S653 in *Arabidopsis*).

RR = Roundup Ready trait, implies utility of CP4 gene as commercially deployed either alone or in combination with other genes but imparting glyphosate only tolerance.

CL = Clearfield crops, tolerant by nontransgenic means. Primary tolerance is to imidazolinone class of ALS-inhibiting chemistry with some partial tolerance to specific other herbicides with this mode of action. Use of CL designation is intended to distinguish from transgenic use of the AHAS gene.

LL = Liberty Link trait, implies utility of either PAT or BAR gene as commercially deployed either alone or in combination with other genes but imparting only tolerance to glutamine synthetase inhibitors such as glufosinate.

STS = designates resistance to sulfonylurea herbicide chemistry with use of ALS1 gene.

Herbicide footnotes:

IMI = any imidazolinone herbicide including, but not limited to, imazapyr, imazethapyr, imazamox, imazaquin.

DIMS = cyclohexanedione class of herbicides (dims) including, but not limited to, sethoxydim, clethodim, and for the purposes of this demonstration pinoxaden.

Fops = aryloxyphenoxypropionate herbicides (fops) including, but not limited to, quizalofop, haloxyfop, fenoxaprop, fluazifop, et al., their stereospecific isomers or racemic mixtures, and esters, acid, or salts thereof.

ALS inhibitors = any ALS inhibitor to the exclusion of IMI's for the sake of this demonstration (i.e., sulfonylureas, triazolopyrimidine sulfonamides, sulfonylaminocarbonyltriazolinone).

HPPD = p-Hydroxyphenyl pyruvate dioxygenase inhibitor class of chemistry including but not limited to mesotrione, sulcotrione, isoxaflutole, and pyrazolynate.

MSMA and DSMA = herbicides from the organoarsenicals chemistry family.

N/A = No suitable options available postemergence.

TABLE 2

Control of volunteer corn (or other monocot) [trait alone or stacked with other HT traits] in AAD-1 Soybean or other dicot crops			
Previous year Corn hybrid (or other monocot)	Current year Soybean variety (or other dicot) being planted	Herbicides option - Burndown/PRE	Herbicides option - POST
Appl: alone and tank mixes			
Conventional corn	AAD-1	Glufosinate, Glyphosate, Fops, Dims, IMI, Paraquat, 2,4-D, Dichlorprop, MCPA, MCP (Mecoprop), Dicamba, HPPD not recommended for corn	2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, IMI
	AAD-1 + GT	MCP (Mecoprop), Dicamba, HPPD not recommended for corn	2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, Glyphosate, IMI
	AAD-1 + PAT	other ALS herbicides not recommended for corn (need to be aware of planting restrictions)	2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, Glufosinate, IMI
	AAD-1 + AHAS	enabled by "ALS gene"	2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, IMI
	AAD-1 + PAT + GT		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, Glufosinate, Glyphosate, IMI
	AAD-1 + PAT + AHAS		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, Glufosinate, IMI
	AAD-1 + GAT + ALS		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, Glyphosate, IMI, ALS inhibitors not selective for in- corn application
	AAD-1 + PAT + GAT + ALS		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, Glufosinate, Glyphosate, IMI, ALS inhibitors not selective for in- corn application
CL Corn	AAD-1	Glufosinate, Glyphosate, Fops, Dims, Paraquat, 2,4- D, Dichlorprop, MCPA, MCP (Mecoprop), Dicamba, HPPD not recommended for corn	2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims
	AAD-1 + GT	MCP (Mecoprop), Dicamba, HPPD not recommended for corn	2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, Glyphosate,
	AAD-1 + PAT	other ALS herbicides not recommended for corn (need to be aware of planting restrictions)	2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, Glufosinate,
	AAD-1 + AHAS	enabled by "ALS gene"	2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims,
	AAD-1 + PAT + GT		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, Glufosinate, Glyphosate,
	AAD-1 + PAT + AHAS		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, Glufosinate,
	AAD-1 + GAT + ALS		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, Glyphosate, ALS inhibitors not selective for in- corn application
	AAD-1 + PAT + GAT + ALS		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, Glufosinate, Glyphosate, ALS inhibitors not selective for in-corn application

TABLE 2-continued

Control of volunteer corn (or other monocot) [trait alone or stacked with other HT traits] in AAD-1 Soybean or other dicot crops)			
Previous year Corn hybrid (or other monocot)	Current year Soybean variety (or other dicot) being planted	Herbicides option - Burndown/PRE	Herbicides option - POST
RR or GT Corn	AAD-1	Glufosinate, Fops, Dims, IMI, Paraquat, 2,4-D, Dichlorprop, MCPA, MCP (Mecoprop), Dicamba, HPPD not recommended for corn other ALS herbicides not recommended for corn (need to be aware of planting restrictions) enabled by "ALS gene"	2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, IMI
	AAD-1 + GT	MCP (Mecoprop), Dicamba, HPPD not recommended for corn other ALS herbicides not recommended for corn (need to be aware of planting restrictions) enabled by "ALS gene"	2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, IMI
	AAD-1 + PAT		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, Glufosinate, IMI
	AAD-1 + AHAS		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, IMI
	AAD-1 + PAT + GT		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, Glufosinate, IMI
	AAD-1 + PAT + AHAS		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, Glufosinate, IMI
	AAD-1 + GAT + ALS		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, IMI, ALS inhibitors not selective for in-corn application
	AAD-1 + PAT + GAT + ALS		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, Glufosinate, IMI, ALS inhibitors not selective for in- corn application
LL Corn	AAD-1	, Glyphosate, Fops, Dims, IMI, Paraquat, 2,4-D, Dichlorprop, MCPA, MCP (Mecoprop), Dicamba, HPPD not recommended for corn other ALS herbicides not recommended for corn (need to be aware of planting restrictions) enabled by "ALS gene"	2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, IMI
	AAD-1 + GT		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, Glyphosate, IMI
	AAD-1 + PAT		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims,, IMI
	AAD-1 + AHAS		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, IMI
	AAD-1 + PAT + GT		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims,, Glyphosate, IMI
	AAD-1 + PAT + AHAS		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims,, IMI
	AAD-1 + GAT + ALS		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, Glyphosate, IMI, ALS inhibitors not selective for in- corn application
	AAD-1 + PAT + GAT + ALS		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims,, Glyphosate, IMI, ALS inhibitors not selective for in- corn application
GAT + ALS	AAD-1	Glufosinate, Fops, Dims, Paraquat, 2,4-D, Dichlorprop, MCPA, MCP (Mecoprop), Dicamba, HPPD not recommended for corn	2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims
	AAD-1 + GT		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims,,
	AAD-1 + PAT		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, Glufosinate
	AAD-1 + AHAS		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims
	AAD-1 + PAT + GT		2,4-D, Dichlorprop, Fops, MCPA, MCP (Mecoprop), Dims, Glufosinate

TABLE 2-continued

Control of volunteer corn (or other monocot) [trait alone or stacked with other HT traits] in AAD-1 Soybean or other dicot crops)			
Previous year Corn hybrid (or other monocot)	Current year Soybean variety (or other dicot) being planted	Herbicides option - Burndown/PRE	Herbicides option - POST
	AAD-1 + PAT + AHAS		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, Glufosinate
	AAD-1 + GAT + ALS		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims
	AAD-1 + PAT + GAT + ALS		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, Glufosinate
CL (or AHAS) + PAT	AAD-1	Glyphosate, Fops, Dims,, Paraquat, 2,4-D, Dichlorprop, MCPA, MCPP (Mecoprop), Dicamba, HPPD not recommended for corn other ALS herbicides not recommended for corn (need to be aware of planting restrictions) enabled by "ALS gene"	2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims
	AAD-1 + GT		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, Glyphosate
	AAD-1 + PAT		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims,
	AAD-1 + AHAS		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims
	AAD-1 + PAT + GT		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, Glyphosate
	AAD-1 + PAT + AHAS		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims
	AAD-1 + GAT + ALS		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, Glyphosate, ALS inhibitors not selective for in- corn application
	AAD-1 + PAT + GAT + ALS		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, Glyphosate, ALS inhibitors not selective for in- corn application
CL (or AHAS) + GT	AAD-1	Glufosinate, Fops, Dims,, Paraquat, 2,4-D, Dichlorprop, MCPA, MCPP (Mecoprop), Dicamba, HPPD not recommended for corn other ALS herbicides not recommended for corn (need to be aware of planting restrictions) enabled by "ALS gene"	2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims
	AAD-1 + GT		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, Glufosinate
	AAD-1 + PAT		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims,
	AAD-1 + AHAS		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims
	AAD-1 + PAT + GT		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, Glufosinate
	AAD-1 + PAT + AHAS		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims
	AAD-1 + GAT + ALS		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, Glufosinate, ALS inhibitors not selective for in- corn application
	AAD-1 + PAT + GAT + ALS		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, Glufosinate, ALS inhibitors not selective for in- corn application

TABLE 2-continued

Control of volunteer corn (or other monocot) [trait alone or stacked with other HT traits] in AAD-1 Soybean or other dicot crops)			
Previous year Corn hybrid (or other monocot)	Current year Soybean variety (or other dicot) being planted	Herbicides option - Burndown/PRE	Herbicides option - POST
GT + PAT	AAD-1	Fops, Dims, IMI, Paraquat, 2,4-D, Dichlorprop, MCPA, MCP (Mecoprop),	2,4-D, Dichloprop, Fops, MCPA, MCP (Mecoprop), Dims, IMI
	AAD-1 + GT	Dicamba, HPPD not recommended for corn other ALS herbicides not recommended for corn	2,4-D, Dichloprop, Fops, MCPA, MCP (Mecoprop), Dims, IMI
	AAD-1 + PAT	(need to be aware of planting restrictions)	2,4-D, Dichloprop, Fops, MCPA, MCP (Mecoprop), Dims, IMI
	AAD-1 + AHAS	enabled by "ALS gene"	2,4-D, Dichloprop, Fops, MCPA, MCP (Mecoprop), Dims, IMI
	AAD-1 + PAT + GT		2,4-D, Dichloprop, Fops, MCPA, MCP (Mecoprop), Dims, IMI
	AAD-1 + PAT + AHAS		2,4-D, Dichloprop, Fops, MCPA, MCP (Mecoprop), Dims, IMI
	AAD-1 + GAT + ALS		2,4-D, Dichloprop, Fops, MCPA, MCP (Mecoprop), Dims, IMI, ALS inhibitors not selective for in-corn application
	AAD-1 + PAT + GAT + ALS		2,4-D, Dichloprop, Fops, MCPA, MCP (Mecoprop), Dims, IMI, ALS inhibitors not selective for in-corn application
GT + CL (or AHAS) + PAT	AAD-1	Fops, Dims, Paraquat, 2,4- D, Dichlorprop, MCPA, MCP (Mecoprop),	2,4-D, Dichloprop, Fops, MCPA, MCP (Mecoprop), Dims
	AAD-1 + GT	Dicamba, HPPD not recommended for corn other ALS herbicides not recommended for corn	2,4-D, Dichloprop, Fops, MCPA, MCP (Mecoprop), Dims
	AAD-1 + PAT	(need to be aware of planting restrictions)	2,4-D, Dichloprop, Fops, MCPA, MCP (Mecoprop), Dims
	AAD-1 + AHAS	enabled by "ALS gene"	2,4-D, Dichloprop, Fops, MCPA, MCP (Mecoprop), Dims
	AAD-1 + PAT + GT		2,4-D, Dichloprop, Fops, MCPA, MCP (Mecoprop), Dims
	AAD-1 + PAT + AHAS		2,4-D, Dichloprop, Fops, MCPA, MCP (Mecoprop), Dims
	AAD-1 + GAT + ALS		2,4-D, Dichloprop, Fops, MCPA, MCP (Mecoprop), Dims, ALS inhibitors not selective for in-corn application
	AAD-1 + PAT + GAT + ALS		2,4-D, Dichloprop, Fops, MCPA, MCP (Mecoprop), Dims, ALS inhibitors not selective for in-corn application
GAT + ALS + PAT	AAD-1	Fops, Dims, Paraquat, 2,4- D, Dichlorprop, MCPA, MCP (Mecoprop),	2,4-D, Dichloprop, Fops, MCPA, MCP (Mecoprop), Dims,
	AAD-1 + GT	Dicamba, HPPD not recommended for corn	2,4-D, Dichloprop, Fops, MCPA, MCP (Mecoprop), Dims,
	AAD-1 + PAT		2,4-D, Dichloprop, Fops, MCPA, MCP (Mecoprop), Dims,
	AAD-1 + AHAS		2,4-D, Dichloprop, Fops, MCPA, MCP (Mecoprop), Dims,

TABLE 2-continued

Control of volunteer corn (or other monocot) [trait alone or stacked with other HT traits] in AAD-1 Soybean or other dicot crops)			
Previous year Corn hybrid (or other monocot)	Current year Soybean variety (or other dicot) being planted	Herbicides option - Burndown/PRE	Herbicides option - POST
	AAD-1 + PAT + GT		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims,
	AAD-1 + PAT + AHAS		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims,
	AAD-1 + GAT + ALS		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims,
	AAD-1 + PAT + GAT + ALS		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims,
Dicamba	AAD-1	Glufosinate, Glyphosate, Fops, Dims, IMI, Paraquat, 2,4-D, Dichlorprop, MCPA, MCPP (Mecoprop)	2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, IMI
	AAD-1 + GT	other ALS herbicides not recommended for corn	2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, Glyphosate, IMI
	AAD-1 + PAT	(need to be aware of planting restrictions)	2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, Glufosinate, IMI
	AAD-1 + AHAS	enabled by "ALS gene"	2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, IMI
	AAD-1 + PAT + GT		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, Glufosinate, Glyphosate, IMI
	AAD-1 + PAT + AHAS		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, Glufosinate, IMI
	AAD-1 + GAT + ALS		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, Glyphosate, IMI, ALS inhibitors not selective for in- corn application
	AAD-1 + PAT + GAT + ALS		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, Glufosinate, Glyphosate, IMI, ALS inhibitors not selective for in- corn application
HPPD	AAD-1	Glufosinate, Glyphosate, Fops, Dims, IMI, Paraquat, 2,4-D, Dichlorprop, MCPA, MCPP (Mecoprop), Dicamba, HPPD not recommended for corn	2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, IMI
	AAD-1 + GT	other ALS herbicides not recommended for corn	2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, Glyphosate, IMI
	AAD-1 + PAT	(need to be aware of planting restrictions)	2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, Glufosinate, IMI
	AAD-1 + AHAS	enabled by "ALS gene"	2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, IMI
	AAD-1 + PAT + GT		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, Glufosinate, Glyphosate, IMI
	AAD-1 + PAT + AHAS		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, Glufosinate, IMI
	AAD-1 + GAT + ALS		2,4-D, Dichloprop, Fops, MCPA, MCPP (Mecoprop), Dims, Glyphosate, IMI, ALS inhibitors not selective for in- corn application

TABLE 2-continued

Control of volunteer corn (or other monocot) [trait alone or stacked with other HT traits] in AAD-1 Soybean or other dicot crops			
Previous year Corn hybrid (or other monocot)	Current year Soybean variety (or other dicot) being planted	Herbicides option - Burndown/PRE	Herbicides option - POST
	AAD-1 + PAT + GAT + ALS		2,4-D, Dichlorprop, Fops, MCPA, MCPP (Mecoprop), Dims, Glufosinate, Glyphosate, IMI, ALS inhibitors not selective for in- corn application

Gene and trait footnotes:

GT = any glyphosate specific tolerance trait including Roundup Ready (CP4), TIPS EPSPS (GA21, Glytol, DMMG), Athenix's EPSPS, GAT only without ALS, GOX, glyphosate decarboxylase, etc.

ALS = double mutant ALS gene insensitive to all ALS herbicide chemistries including IMI herbicides.

AAD-1 = Aryloxyalkanoate dioxygenase gene providing tolerance to all commercial phenoxy auxin and all aryloxyphenoxypropionate (fop) herbicides.

AAD-12 = Aryloxyalkanoate dioxygenase gene providing tolerance to phenoxyacetic auxin and pyridyloxyacetic auxin herbicides.

PAT = phosphinothricin acetyltransferase gene providing tolerance to glutamine synthetase inhibitors including, but not limited to, glufosinate. Similar phenotype is provided by genes such as BAR, DSM1, DSM2, et al.

AHAS = imidazolinone specific tolerance gene associated with point mutation at S623 of ALS gene (maize sequence) or equivalent amino acid in other spp (e.g., S653 in *Arabidopsis*).

RR = Roundup Ready trait, implies utility of CP4 gene as commercially deployed either alone or in combination with other genes but imparting glyphosate only tolerance.

CL = Clearfield crops, tolerant by nontransgenic means. Primary tolerance is to imidazolinone class of ALS-inhibiting chemistry with some partial tolerance to specific other herbicides with this mode of action. Use of CL designation is intended to distinguish from transgenic use of the AHAS gene.

LL = Liberty Link trait, implies utility of either PAT or BAR gene as commercially deployed either alone or in combination with other genes but imparting only tolerance to glutamine synthetase inhibitors such as glufosinate.

STS = designates resistance to sulfonyleurea herbicide chemistry with use of ALS1 gene.

Herbicide footnotes:

IMI = any imidazolinone herbicide including, but not limited to, imazapyr, imazethapyr, imazamox, imazaquin.

DIMS = cyclohexanedione class of herbicides (dims) including, but not limited to, sethoxydim, clethodim, and for the purposes of this demonstration poinoxaden.

Fops = aryloxyphenoxypropionate herbicides (fops) including, but not limited to, quizalofop, haloxyfop, fenoxaprop, fluazifop, et al., their stereospecific isomers or racemic mixtures, and esters, acid, or salts thereof.

ALS inhibitors = any ALS inhibitor to the exclusion of IMI's for the sake of this demonstration (i.e., sulfonyleureas, triazolopyrimidine sulfonanalides, sulfonyleurea carbonyl triazolinone).

HPPD = p-Hydroxyphenyl pyruvate dioxygenase inhibitor class of chemistry including but not limited to mesotrione, sulcotrione, isoxaflutole, and pyrazolynate.

MSMA and DSMA = herbicides from the organoarsenicals chemistry family.

N/A = No suitable options available postemergence.

TABLE 3

Control of volunteer corn (or other monocot) [trait alone or stacked with other HT traits] in AAD-12 Soybean (or other dicot crops)			
Previous year Corn hybrid (or other monocot)	Current year Soybean variety (or other dicot) being planted	Herbicides option - Burndown/PRE	Herbicides option - POST
Appl: alone and tank mixes			
Conventional corn	AAD-12	Glufosinate, Glyphosate,	2,4-D, Dims, IMI
	AAD-12 + GT	Fops, Dims, IMI, Paraquat,	2,4-D, Dims, Glyphosate, IMI
	AAD-12 + PAT	2,4-D, Dichlorprop, MCPA,	2,4-D, Dims, Glufosinate, IMI
	AAD-12 + AHAS	MCPP (Mecoprop),	2,4-D, Dims, IMI
	AAD-12 + PAT + GT	Dicamba, HPPD not recommended for corn	2,4-D, Dims, Glufosinate, Glyphosate, IMI
	AAD-12 + PAT + AHAS	other ALS herbicides not recommended for corn	2,4-D, Dims, Glufosinate, IMI
	AAD-12 + GAT + ALS	(need to be aware of planting restrictions) enabled by "ALS gene"	2,4-D, Dims, Glyphosate, IMI, ALS inhibitors not selective for in-corn application
CL Corn	AAD-12 + PAT + GAT + ALS		2,4-D, Dims, Glufosinate, Glyphosate, IMI, ALS inhibitors not selective for in- corn application
	AAD-12	Glufosinate, Glyphosate,	2,4-D, Dims
	AAD-12 + GT	Fops, Dims, Paraquat, 2,4- D, Dichlorprop, MCPA,	2,4-D, Dims, Glyphosate,
	AAD-12 + PAT	MCPP (Mecoprop),	2,4-D, Dims, Glufosinate,
	AAD-12 + AHAS	Dicamba, HPPD not recommended for corn	2,4-D, Dims, 2,4-D, Dims, Glufosinate, Glyphosate,

TABLE 3-continued

Control of volunteer corn (or other monocot) [trait alone or stacked with other HT traits] in AAD-12 Soybean (or other dicot crops)			
Previous year Corn hybrid (or other monocot)	Current year Soybean variety (or other dicot) being planted	Herbicides option - Burndown/PRE	Herbicides option - POST
	AAD-12 + PAT + AHAS AAD-12 + GAT + ALS	other ALS herbicides not recommended for corn (need to be aware of planting restrictions)	2,4-D, Dims, Glufosinate, 2,4-D, Dims, Glyphosate, ALS inhibitors not selective for in-corn application
	AAD-12 + PAT + GAT + ALS	enabled by "ALS gene"	2,4-D, Dims, Glufosinate, Glyphosate, ALS inhibitors not selective for in-corn application
RR or GT Corn	AAD-12 AAD-12 + GT AAD-12 + PAT AAD-12 + AHAS AAD-12 + PAT + GT AAD-12 + PAT + AHAS AAD-12 + GAT + ALS	Glufosinate, Fops, Dims, IMI, Paraquat, 2,4-D, Dichlorprop, MCPA, MCPP (Mecoprop), Dicamba, HPPD not recommended for corn other ALS herbicides not recommended for corn (need to be aware of planting restrictions)	2,4-D, Dims, IMI 2,4-D, Dims, IMI 2,4-D, Dims, Glufosinate, IMI 2,4-D, Dims, IMI 2,4-D, Dims, Glufosinate, IMI 2,4-D, Dims, Glufosinate, IMI 2,4-D, Dims, IMI, ALS inhibitors not selective for in-corn application
	AAD-12 + PAT + GAT + ALS	enabled by "ALS gene"	2,4-D, Dims, Glufosinate, IMI, ALS inhibitors not selective for in-corn application
LL Corn	AAD-12 AAD-12 + GT AAD-12 + PAT AAD-12 + AHAS AAD-12 + PAT + GT AAD-12 + PAT + AHAS AAD-12 + GAT + ALS	, Glyphosate, Fops, Dims, IMI, Paraquat, 2,4-D, Dichlorprop, MCPA, MCPP (Mecoprop), Dicamba, HPPD not recommended for corn other ALS herbicides not recommended for corn (need to be aware of planting restrictions)	2,4-D, Dims, IMI 2,4-D, Dims, Glyphosate, IMI 2,4-D, Dims,, IMI 2,4-D, Dims, IMI 2,4-D, Dims,, Glyphosate, IMI 2,4-D, Dims,, IMI 2,4-D, Dims, Glyphosate, IMI, ALS inhibitors not selective for in-corn application
	AAD-12 + PAT + GAT + ALS	enabled by "ALS gene"	2,4-D, Dims,, Glyphosate, IMI, ALS inhibitors not selective for in-corn application
GAT + ALS	AAD-12 AAD-12 + GT AAD-12 + PAT AAD-12 + AHAS AAD-12 + PAT + GT AAD-12 + PAT + AHAS AAD-12 + GAT + ALS AAD-12 + PAT + FGAT + ALS	Glufosinate, Fops, Dims, Paraquat, 2,4-D, Dichlorprop, MCPA, MCPP (Mecoprop), Dicamba, HPPD not recommended for corn	2,4-D, Dims 2,4-D, Dims,, 2,4-D, Dims, Glufosinate 2,4-D, Dims 2,4-D, Dims, Glufosinate 2,4-D, Dims, Glufosinate 2,4-D, Dims 2,4-D, Dims, Glufosinate
CL (or AHAS) + PAT	AAD-12 AAD-12 + GT AAD-12 + PAT AAD-12 + AHAS AAD-12 + PAT + GT AAD-12 + PAT + AHAS AAD-12 + GAT + ALS	Glyphosate, Fops, Dims,, Paraquat, 2,4-D, Dichlorprop, MCPA, MCPP (Mecoprop), Dicamba, HPPD not recommended for corn other ALS herbicides not recommended for corn (need to be aware of planting restrictions)	2,4-D, Dims 2,4-D, Dims, Glyphosate 2,4-D, Dims, 2,4-D, Dims 2,4-D, Dims, Glyphosate 2,4-D, Dims 2,4-D, Dims, Glyphosate, ALS inhibitors not selective for in-corn application
	AAD-12 + PAT + GAT + ALS	enabled by "ALS gene"	2,4-D, Dims, Glyphosate, ALS inhibitors not selective for in-corn application
CL (or AHAS) + GT	AAD-12 AAD-12 + GT AAD-12 + PAT AAD-12 + AHAS AAD-12 + PAT + GT AAD-12 + PAT + AHAS AAD-12 + GAT + ALS	Glufosinate, Fops, Dims,, Paraquat, 2,4-D, Dichlorprop, MCPA, MCPP (Mecoprop), Dicamba, HPPD not recommended for corn other ALS herbicides not recommended for corn (need to be aware of planting restrictions)	2,4-D, Dims 2,4-D, Dims, Glufosinate 2,4-D, Dims, 2,4-D, Dims 2,4-D, Dims, Glufosinate 2,4-D, Dims 2,4-D, Dims, Glufosinate, ALS inhibitors not selective for in-corn application
	AAD-12 + PAT + GAT + ALS	enabled by "ALS gene"	2,4-D, Dims, Glufosinate, ALS inhibitors not selective for in-corn application

TABLE 3-continued

Control of volunteer corn (or other monocot) [trait alone or stacked with other HT traits] in AAD-12 Soybean (or other dicot crops)			
Previous year Corn hybrid (or other monocot)	Current year Soybean variety (or other dicot) being planted	Herbicides option - Burndown/PRE	Herbicides option - POST
GT + PAT	AAD-12	Fops, Dims, IMI, Paraquat,	2,4-D, Dims, IMI
	AAD-12 + GT	2,4-D, Dichlorprop, MCPA,	2,4-D, Dims, IMI
	AAD-12 + PAT	MCPP (Mecoprop),	2,4-D, Dims, IMI
	AAD-12 + AHAS	Dicamba, HPPD not	2,4-D, Dims, IMI
	AAD-12 + PAT + GT	recommended for corn	2,4-D, Dims, IMI
	AAD-12 + PAT + AHAS	other ALS herbicides not	2,4-D, Dims, IMI
	AAD-12 + GAT + ALS	recommended for corn (need to be aware of planting restrictions)	2,4-D, Dims, IMI, ALS inhibitors not selective for in- corn application
GT + CL (or AHAS) + PAT	AAD-12	Fops, Dims, Paraquat, 2,4-	2,4-D, Dims
	AAD-12 + GT	D, Dichlorprop, MCPA,	2,4-D, Dims
	AAD-12 + PAT	MCPP (Mecoprop),	2,4-D, Dims
	AAD-12 + AHAS	Dicamba, HPPD not	2,4-D, Dims
	AAD-12 + PAT + GT	recommended for corn	2,4-D, Dims
	AAD-12 + PAT + AHAS	other ALS herbicides not	2,4-D, Dims
	AAD-12 + GAT + ALS	recommended for corn (need to be aware of planting restrictions)	2,4-D, Dims, ALS inhibitors not selective for in-corn application
GAT + ALS + PAT	AAD-12	Fops, Dims, Paraquat, 2,4-	2,4-D, Dims,
	AAD-12 + GT	D, Dichlorprop, MCPA,	2,4-D, Dims,
	AAD-12 + PAT	MCPP (Mecoprop),	2,4-D, Dims,
	AAD-12 + AHAS	Dicamba, HPPD not	2,4-D, Dims,
	AAD-12 + PAT + GT	recommended for corn	2,4-D, Dims,
	AAD-12 + PAT + AHAS		2,4-D, Dims,
	AAD-12 + GAT + ALS		2,4-D, Dims,
Dicamba	AAD-12	Glufosinate, Glyphosate,	2,4-D, Dims, IMI
	AAD-12 + GT	Fops, Dims, IMI, Paraquat,	2,4-D, Dims, Glyphosate, IMI
	AAD-12 + PAT	2,4-D, Dichlorprop, MCPA,	2,4-D, Dims, Glufosinate, IMI
	AAD-12 + AHAS	MCPP (Mecoprop)	2,4-D, Dims, IMI
	AAD-12 + PAT + GT	other ALS herbicides not	2,4-D, Dims, Glufosinate,
	AAD-12 + PAT + AHAS	recommended for corn	Glyphosate, IMI
	AAD-12 + GAT + ALS	(need to be aware of planting restrictions)	2,4-D, Dims, Glufosinate, IMI
HPPD	AAD-12	Glufosinate, Glyphosate,	2,4-D, Dims, IMI
	AAD-12 + GT	Fops, Dims, IMI, Paraquat,	2,4-D, Dims, Glyphosate, IMI
	AAD-12 + PAT	2,4-D, Dichlorprop, MCPA,	2,4-D, Dims, Glufosinate, IMI
	AAD-12 + AHAS	MCPP (Mecoprop),	2,4-D, Dims, IMI
	AAD-12 + PAT + GT	Dicamba, HPPD not	2,4-D, Dims, Glufosinate,
	AAD-12 + PAT + AHAS	recommended for corn	Glyphosate, IMI
	AAD-12 + GAT + ALS	other ALS herbicides not	2,4-D, Dims, Glufosinate, IMI
	recommended for corn (need to be aware of planting restrictions)	2,4-D, Dims, Glyphosate, IMI, ALS inhibitors not selective for in-corn application	

TABLE 3-continued

Control of volunteer corn (or other monocot) [trait alone or stacked with other HT traits] in AAD-12 Soybean (or other dicot crops)			
Previous year Corn hybrid (or other monocot)	Current year Soybean variety (or other dicot) being planted	Herbicides option - Burndown/PRE	Herbicides option - POST
	AAD-12 + PAT + GAT + ALS	enabled by "ALS gene"	2,4-D, Dims, Glufosinate, Glyphosate, IMI, ALS inhibitors not selective for in- corn application

Gene and trait footnotes:

GT = any glyphosate specific tolerance trait including Roundup Ready (CP4), TIPS EPSPS (GA21, Glyto1, DMMG), Athenix's EPSPS, GAT only without ALS, GOX, glyphosate decarboxylase, etc.

ALS = double mutant ALS gene insensitive to all ALS herbicide chemistries including IMI herbicides.

AAD-1 = Aryloxyalkanoate dioxygenase gene providing tolerance to all commercial phenoxy auxin and all aryloxyphenoxypropionate (fop) herbicides.

AAD-12 = Aryloxyalkanoate dioxygenase gene providing tolerance to phenoxyacetic auxin and pyridyloxyacetic auxin herbicides.

PAT = phosphinothricin acetyltransferase gene providing tolerance to glutamine synthetase inhibitors including, but not limited to, glufosinate. Similar phenotype is provided by genes such as BAR, DSM1, DSM2, et al.

AHAS = imidazolinone specific tolerance gene associated with point mutation at S623 of ALS gene (maize sequence) or equivalent amino acid in other spp (e.g., S653 in *Arabidopsis*).

RR = Roundup Ready trait, implies utility of CP4 gene as commercially deployed either alone or in combination with other genes but imparting glyphosate only tolerance.

CL = Clearfield crops, tolerant by nontransgenic means. Primary tolerance is to imidazolinone class of ALS-inhibiting chemistry with some partial tolerance to specific other herbicides with this mode of action. Use of CL designation is intended to distinguish from transgenic use of the AHAS gene.

LL = Liberty Link trait, implies utility of either PAT or BAR gene as commercially deployed either alone or in combination with other genes but imparting only tolerance to glutamine synthetase inhibitors such as glufosinate.

STS = designates resistance to sulfonyleurea herbicide chemistry with use of ALS1 gene.

Herbicide footnotes:

IMI = any imidazolinone herbicide including, but not limited to, imazapyr, imazethapyr, imazamox, imazaquin.

DIMS = cyclohexanedione class of herbicides (dims) including, but not limited to, sethoxydim, clethodim, and for the purposes of this demonstration pinoxaden.

Fops = aryloxyphenoxypropionate herbicides (fops) including, but not limited to, quizalofop, haloxyfop, fenoxaprop, fluzifop, et al., their stereospecific isomers or racemic mixtures, and esters, acid, or salts thereof.

ALS inhibitors = any ALS inhibitor to the exclusion of IMI's for the sake of this demonstration (i.e., sulfonyleureas, triazolopyrimidine sulfonamides, sulfonylaminocarbonyltriazolinone).

HPPD = p-Hydroxyphenyl pyruvate dioxygenase inhibitor class of chemistry including but not limited to mesotrione, sulcotrione, isoxaflutole, and pyrazolynate.

MSMA and DSMA = herbicides from the organoarsenicals chemistry family.

N/A = No suitable options available postemergence.

TABLE 4

Control of volunteer AAD-1 corn (or other monocot crop) (alone or stacked with other HT traits) in AAD-12-based Cotton (or other dicot crop)				
Previous year Corn hybrid	Current year Cotton hybrid being planted	Herbicides option - Burndown/PRE/PPI	Herbicides option - POST DIRECTED	Herbicides option - POST BROADCAST
Appl: alone and tank mixes				
AAD-1	Cotton, CL Cotton, STS Cotton	Glyphosate, Dims, Tribenuron, Trifloxysulfuron, Paraquat, Glufosinate, MSMA, Flumioxazin, Pendimethalin, Trifluralin, Prometryn, Clomazone,	Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin, Dims, Imazapyr + Imazethapyr, imazamox, imazaquin	Pyrithiobac, Dims, Imazapyr + Imazethapyr, imazamox, imazaquin
	RR Cotton	Fluometuron, Diuron, Fomesafen, Pyrithiobac	Dims, Glyphosate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Glyphosate, Pyrithiobac
	LL Cotton		Dims, Glufosinate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Glufosinate, Pyrithiobac

TABLE 4-continued

Control of volunteer AAD-1 corn (or other monocot crop) (alone or stacked with other HT traits) in AAD-12-based Cotton (or other dicot crop)				
Previous year Corn hybrid	Current year Cotton hybrid being planted	Herbicides option - Burndown/PRE/PPI	Herbicides option - POST DIRECTED	Herbicides option - POST BROADCAST
	GAT		Dims, Glyphosate, ALS herbicides (imidazolinones, sulfonyleureas, triazolopyrimidine sulfonanalides, sulfonylamino carbonyl triazolinones), Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Glyphosate, ALS herbicides (imidazolinones, sulfonyleureas, triazolopyrimidine sulfonanalides, sulfonylamino carbonyl triazolinones), Pyriithiobac
	AAD-12		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriithiobac
	AAD-12 + TIPS		Dims, Glyphosate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Glyphosate, Pyriithiobac
	AAD-12 + PAT		Dims, Glufosinate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Glufosinate, Pyriithiobac
	AAD-12 + AHAS		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Sulfonyleureas, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriithiobac Sulfonyleureas,
	AAD-12 + PAT + TIPS		Dims, Glyphosate, glufosinate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Glyphosate, glufosinate, Pyriithiobac
	AAD-12 + PAT + AHAS		Dims, glufosinate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, glufosinate, Sulfonyleureas, Pyriithiobac

TABLE 4-continued

Control of volunteer AAD-1 corn (or other monocot crop) (alone or stacked with other HT traits) in AAD-12-based Cotton (or other dicot crop)				
Previous year Corn hybrid	Current year Cotton hybrid being planted	Herbicides option - Burndown/PRE/PPI	Herbicides option - POST DIRECTED	Herbicides option - POST BROADCAST
	AAD-12 + GAT		Dims, Glyphosate, ALS herbicides (imidazolinones, sulfonyleureas, triazolopyrimidine sulfonanalides, sulfobylamino carbonyl triazolinones), Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Glyphosate, ALS herbicides (imidazolinones, sulfonyleureas, triazolopyrimidine sulfonanalides, sulfobylamino carbonyl triazolinones), Pyriithiobac
AAD-1 + PAT	CL Cotton	Glyphosate, Dims, Tribenuron, Trifloxysulfuron, Paraquat, MSMA, Flumioxazin, Pendimethalin, Trifluralin, Prometryn, Clomazone, Fluometuron, Diuron, Fomesafen, Pyriithiobac	Dims, Imazapyr + Imazethapyr, imazamox, imazaquin, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Pyriithiobac, Dims, Imazapyr + Imazethapyr, imazamox, imazaquin
	RR Cotton		Dims, Glyphosate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Glyphosate, Pyriithiobac
	LL Cotton		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims,, Pyriithiobac
	GAT		Dims, Glyphosate, ALS herbicides (imidazolinones, sulfonyleureas, triazolopyrimidine sulfonanalides, sulfobylamino carbonyl triazolinones)	Dims, Glyphosate, ALS herbicides (imidazolinones, sulfonyleureas, triazolopyrimidine sulfonanalides, sulfobylamino carbonyl triazolinones), Pyriithiobac
	AAD-12		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriithiobac
	AAD-12 + TIPS		Glyphosate, Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Glyphosate, Pyriithiobac
	AAD-12 + PAT		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims,, Pyriithiobac

TABLE 4-continued

Control of volunteer AAD-1 corn (or other monocot crop) (alone or stacked with other HT traits) in AAD-12-based Cotton (or other dicot crop)				
Previous year Corn hybrid	Current year Cotton hybrid being planted	Herbicides option - Burndown/PRE/PPI	Herbicides option - POST DIRECTED	Herbicides option - POST BROADCAST
	AAD-12 + AHAS		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Sulfonylureas, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Pyriithiobac Sulfonylureas,
	AAD-12 + PAT + TIPS		Dims, Glyphosate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Glyphosate,, Pyriithiobac
	AAD-12 + PAT + AHAS		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Sulfonylureas, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims,, Sulfonylureas, Pyriithiobac
	AAD-12 + GAT		Dims, Glyphosate, ALS herbicides (imidazolinones, sulfonylureas, triazolopyrimidine sulfonanalides, sulfobylamino carbonyl triazolinones), Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Glyphosate, ALS herbicides (imidazolinones, sulfonylureas, triazolopyrimidine sulfonanalides, sulfobylamino carbonyl triazolinones), Pyriithiobac
AAD-1 + TIPS	CL Cotton	Dims, Tribenuron, Trifloxysulfuron, Paraquat, Glufosinate, MSMA, Flumioxazin, Pendimethalin, Trifluralin, Prometryn, Clomazone, Fluometuron, Diuron,	Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin, Dims, Imazapyr + Imazethapyr, imazamox, imazaquin	Pyriithiobac, Dims, Imazapyr + Imazethapyr, imazamox, imazaquin
	RR Cotton	Fomesafen, Pyriithiobac	Dims, Glyphosate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriithiobac
	LL Cotton		Dims, Glufosinate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Glufosinate, Pyriithiobac

TABLE 4-continued

Control of volunteer AAD-1 corn (or other monocot crop) (alone or stacked with other HT traits) in AAD-12-based Cotton (or other dicot crop)				
Previous year Corn hybrid	Current year Cotton hybrid being planted	Herbicides option - Burndown/PRE/PPI	Herbicides option - POST DIRECTED	Herbicides option - POST BROADCAST
	GAT		Dims, ALS herbicides (imidazolinones, sulfonamide, sulfonamide, sulfobylamino carbonyl triazolinones), Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, ALS herbicides (imidazolinones, sulfonamide, sulfonamide, sulfobylamino carbonyl triazolinones), Pyriithiobac
	AAD-12		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Pyriithiobac
	AAD-12 + TIPS		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Pyriithiobac
	AAD-12 + PAT		Dims, Glufosinate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Glufosinate, Pyriithiobac
	AAD-12 + AHAS		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Sulfonamide, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Pyriithiobac Sulfonamide,
	AAD-12 + PAT + TIPS		Dims, glufosinate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, glufosinate, Pyriithiobac
	AAD-12 + PAT + AHAS		Dims, Glufosinate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Sulfonamide, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, glufosinate, Sulfonamide, Pyriithiobac
	AAD-12 + GAT		Dims, ALS herbicides (imidazolinones, sulfonamide, sulfonamide, sulfobylamino carbonyl triazolinones), Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, ALS herbicides (imidazolinones, sulfonamide, sulfonamide, sulfobylamino carbonyl triazolinones), Pyriithiobac

TABLE 4-continued

Control of volunteer AAD-1 corn (or other monocot crop) (alone or stacked with other HT traits) in AAD-12-based Cotton (or other dicot crop)				
Previous year Corn hybrid	Current year Cotton hybrid being planted	Herbicides option - Burndown/PRE/PPI	Herbicides option - POST DIRECTED	Herbicides option - POST BROADCAST
AAD-1 + AHAS	CL Cotton	Glyphosate, Dims, Tribenuron, Trifloxysulfuron, Paraquat, Glufosinate, MSMA, Flumioxazin, Pendimethalin, Trifluralin, Prometryn, Clomazone, Fluometuron, Diuron, Fomesafen, Pyriithiobac	Dims, Imazapyr + Imazethapyr, imazamox, imazaquin, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Pyriithiobac, Dims, Imazapyr + Imazethapyr, imazamox, imazaquin
	RR Cotton	Fomesafen, Pyriithiobac	Dims, Glyphosate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Glyphosate, Pyriithiobac
	LL Cotton		Dims, Prometryn, Lactofen, Fluometuron, Glufosinate, DSMA, MSMA, Oxyfluorfen, Paraquat Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Glufosinate, Pyriithiobac
	GAT		Dims, ALS herbicides (imidazolinones, triazolopyrimidine sulfonanalides, sulfobylamino carbonyl triazolinones), Prometryn, Lactofen, Fluometuron, Glufosinate, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Glyphosate, ALS herbicides (imidazolinones, triazolopyrimidine sulfonanalides, sulfobylamino carbonyl triazolinones), Pyriithiobac
	AAD-12		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriithiobac
	AAD-12 + TIPS		Glyphosate, Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Glyphosate, Pyriithiobac
	AAD-12 + PAT		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Glufosinate, Pyriithiobac
	AAD-12 + AHAS		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Pyriithiobac

TABLE 4-continued

Control of volunteer AAD-1 corn (or other monocot crop) (alone or stacked with other HT traits) in AAD-12-based Cotton (or other dicot crop)				
Previous year Corn hybrid	Current year Cotton hybrid being planted	Herbicides option - Burndown/PRE/PPI	Herbicides option - POST DIRECTED	Herbicides option - POST BROADCAST
	AAD-12 + PAT + TIPS		Dims, Glyphosate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Glyphosate, glufosinate, Pyriithiobac
	AAD-12 + PAT + AHAS		Dims, Glufosinate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, glufosinate, Pyriithiobac
	AAD-12 + GAT		Dims, Glyphosate, ALS herbicides (imidazolinones, triazolopyrimidine sulfonanalides, sulfobylamino carbonyl triazolinones), Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Glyphosate, ALS herbicides (imidazolinones, triazolopyrimidine sulfonanalides, sulfobylamino carbonyl triazolinones), Pyriithiobac
AAD-1 + PAT + TIPS	CL Cotton	Dims, Tribenuron, Trifloxysulfuron, Paraquat, MSMA, Flumioxazin, Pendimethalin, Trifluralin, Prometryn, Clomazone, Fluometuron, Diuron, Fomesafen, Pyriithiobac	Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin, Dims, Imazapyr + Imazethapyr, imazamox, imazaquin	Pyriithiobac, Dims, Imazapyr + Imazethapyr, imazamox, imazaquin
	RR Cotton		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Pyriithiobac
	LL Cotton		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Pyriithiobac
	GAT		Dims, ALS herbicides (imidazolinones, sulfonamide, triazolopyrimidine sulfonanalides, sulfobylamino carbonyl triazolinones), Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, ALS herbicides (imidazolinones, sulfonamide, triazolopyrimidine sulfonanalides, sulfobylamino carbonyl triazolinones), Pyriithiobac
	AAD-12		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Pyriithiobac

TABLE 4-continued

Control of volunteer AAD-1 corn (or other monocot crop) (alone or stacked with other HT traits) in AAD-12-based Cotton (or other dicot crop)				
Previous year Corn hybrid	Current year Cotton hybrid being planted	Herbicides option - Burndown/PRE/PPI	Herbicides option - POST DIRECTED	Herbicides option - POST BROADCAST
	AAD-12 + TIPS		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriothiac
	AAD-12 + PAT		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriothiac
	AAD-12 + AHAS		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriothiac Sulfonylureas,
	AAD-12 + PAT + TIPS		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriothiac
	AAD-12 + PAT + AHAS		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Sulfonylureas, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Sulfonylureas, Pyriothiac
	AAD-12 + GAT		Dims, ALS herbicides (imidazolinones, sulfonylureas, triazolopyrimidine sulfonanalides, sulfobylamino carbonyl triazolinones), Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, ALS herbicides (imidazolinones, sulfonylureas, triazolopyrimidine sulfonanalides, sulfobylamino carbonyl triazolinones), Pyriothiac
AAD-1 + PAT + AHAS	CL Cotton	Glyphosate, Dims, Tribenuron, Trifloxysulfuron, Paraquat, MSMA, Flumioxazin, Pendimethalin, Trifluralin, Prometryn, Clomazone, Fluometuron, Diuron,	Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Pyriothiac, Dims, Imazapyr + Imazethapyr, imazamox, imazaquin
	RR Cotton	Fomesafen, Pyriothiac	Dims, Glyphosate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Glyphosate, Pyriothiac

TABLE 4-continued

Control of volunteer AAD-1 corn (or other monocot crop) (alone or stacked with other HT traits) in AAD-12-based Cotton (or other dicot crop)				
Previous year Corn hybrid	Current year Cotton hybrid being planted	Herbicides option - Burndown/PRE/PPI	Herbicides option - POST DIRECTED	Herbicides option - POST BROADCAST
	LL Cotton		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriithiobac
	GAT		Dims, ALS herbicides (imidazolinones, triazolopyrimidine sulfonanalides, sulfobylamino carbonyl triazolinones), Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Glyphosate, ALS herbicides (imidazolinones, triazolopyrimidine sulfonanalides, sulfobylamino carbonyl triazolinones), Pyriithiobac
	AAD-12		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriithiobac
	AAD-12 + TIPS		Dims, Glyphosate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Glyphosate, Pyriithiobac
	AAD-12 + PAT		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriithiobac
	AAD-12 + AHAS		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriithiobac
	AAD-12 + PAT + TIPS		Dims, Glyphosate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Trifloxysulfuron, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Glyphosate, Pyriithiobac
	AAD-12 + PAT + AHAS		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriithiobac

TABLE 4-continued

Control of volunteer AAD-1 corn (or other monocot crop) (alone or stacked with other HT traits) in AAD-12-based Cotton (or other dicot crop)				
Previous year Corn hybrid	Current year Cotton hybrid being planted	Herbicides option - Burndown/PRE/PPI	Herbicides option - POST DIRECTED	Herbicides option - POST BROADCAST
	AAD-12 + GAT		Dims, ALS herbicides (imidazolinones, triazolopyrimidine sulfonanalides, sulfobylamino carbonyl triazolinones), Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Glyphosate, ALS herbicides (imidazolinones, triazolopyrimidine sulfonanalides, sulfobylamino carbonyl triazolinones), Pyriithiobac
AAD-1 + GAT	CL Cotton	Dims, Tribenuron, Trifloxysulfuron, Paraquat, Glufosinate, MSMA, Flumioxazin, Pendimethalin, Trifluralin, Prometryn,	Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Pyriithiobac, Dims, Imazapyr + Imazethapyr, imazamox, imazaquin
	RR Cotton	Clomazone, Fluometuron, Diuron, Fomesafen, Pyriithiobac	Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriithiobac
	LL Cotton		Dims, Glufosinate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Glufosinate, Pyriithiobac
	GAT		Fomesafen, Flumioxazin, Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Pyriithiobac
	AAD-12		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriithiobac
	AAD-12 + TIPS		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriithiobac
	AAD-12 + PAT		Dims, Glufosinate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Glufosinate, Pyriithiobac
	AAD-12 + AHAS		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriithiobac

TABLE 4-continued

Control of volunteer AAD-1 corn (or other monocot crop) (alone or stacked with other HT traits) in AAD-12-based Cotton (or other dicot crop)				
Previous year Corn hybrid	Current year Cotton hybrid being planted	Herbicides option - Burndown/PRE/PPI	Herbicides option - POST DIRECTED	Herbicides option - POST BROADCAST
	AAD-12 + PAT + TIPS		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, glufosinate, Pyriithiobac
	AAD-12 + PAT + AHAS		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, glufosinate, Pyriithiobac
	AAD-12 + GAT		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Pyriithiobac
AAD-1 + PAT + GAT	CL Cotton	Dims, Tribenuron, Trifloxysulfuron, Paraquat, Glufosinate, MSMA, Flumioxazin, Pendimethalin, Trifluralin, Prometryn,	Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Pyriithiobac, Dims, Imazapyr + Imazethapyr, imazamox, imazaquin
	RR Cotton	Clomazone, Fluometuron, Diuron, Fomesafen, Pyriithiobac	Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriithiobac
	LL Cotton		Dims, Glufosinate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Glufosinate, Pyriithiobac
	GAT		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Pyriithiobac
	AAD-12		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriithiobac
	AAD-12 + TIPS		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriithiobac
	AAD-12 + PAT		Dims, Glufosinate, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriithiobac

TABLE 4-continued

Control of volunteer AAD-1 corn (or other monocot crop) (alone or stacked with other HT traits) in AAD-12-based Cotton (or other dicot crop)				
Previous year Corn hybrid	Current year Cotton hybrid being planted	Herbicides option - Burndown/PRE/PPI	Herbicides option - POST DIRECTED	Herbicides option - POST BROADCAST
	AAD-12 + AHAS		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriithiobac
	AAD-12 + PAT + TIPS		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriithiobac
	AAD-12 + PAT + AHAS		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin,	Dims, Pyriithiobac
	AAD-12 + GAT		Dims, Prometryn, Lactofen, Fluometuron, DSMA, MSMA, Oxyfluorfen, Paraquat, Dimethipin, Diuron, Linuron, Fomesafen, Flumioxazin	Dims, Pyriithiobac

Gene and trait footnotes:

GT = any glyphosate specific tolerance trait including Roundup Ready (CP4), TIPS EPSPS (GA21, Glytol, DMMG), Athenix's EPSPS, GAT only without ALS, GOX, glyphosate decarboxylase, etc.

ALS = double mutant ALS gene insensitive to all ALS herbicide chemistries including IMI herbicides.

AAD-1 = Aryloxyalkanoate dioxygenase gene providing tolerance to all commercial phenoxy auxin and all aryloxyphenoxypropionate (fop) herbicides.

AAD-12 = Aryloxyalkanoate dioxygenase gene providing tolerance to phenoxyacetic auxin and pyridyloxyacetic auxin herbicides.

PAT = phosphinothricin acetyltransferase gene providing tolerance to glutamine synthetase inhibitors including, but not limited to, glufosinate. Similar phenotype is provided by genes such as BAR, DSM1, DSM2, et al.

AHAS = imidazolinone specific tolerance gene associated with point mutation at S623 of ALS gene (maize sequence) or equivalent amino acid in other spp (e.g., S653 in *Arabidopsis*).

RR = Roundup Ready trait, implies utility of CP4 gene as commercially deployed either alone or in combination with other genes but imparting glyphosate only tolerance.

CL = Clearfield crops, tolerant by nontransgenic means. Primary tolerance is to imidazolinone class of ALS-inhibiting chemistry with some partial tolerance to specific other herbicides with this mode of action. Use of CL designation is intended to distinguish from transgenic use of the AHAS gene.

LL = Liberty Link trait, implies utility of either PAT or BAR gene as commercially deployed either alone or in combination with other genes but imparting only tolerance to glutamine synthetase inhibitors such as glufosinate.

STS = designates resistance to sulfonyleurea herbicide chemistry with use of ALS1 gene.

Herbicide footnotes:

IMI = any imidazolinone herbicide including, but not limited to, imazapyr, imazethapyr, imazamox, imazaquin.

DIMS = cyclohexanedione class of herbicides (dims) including, but not limited to, sethoxydim, clethodim, and for the purposes of this demonstration pinoxaden.

Fops = aryloxyphenoxypropionate herbicides (fops) including, but not limited to, quizalofop, haloxyfop, fenoxaprop, fluazifop, et al., their stereospecific isomers or racemic mixtures, and esters, acid, or salts thereof.

ALS inhibitors = any ALS inhibitor to the exclusion of IMI's for the sake of this demonstration (i.e., sulfonyleureas, triazolopyrimidine sulfonamides, sulfonamide-carbonyl triazolinone).

HPPD = p-Hydroxyphenyl pyruvate dioxygenase inhibitor class of chemistry including but not limited to mesotrione, sulcotrione, isoxaflutole, and pyrazolynate.

MSMA and DSMA = herbicides from the organoarsenicals chemistry family.

N/A = No suitable options available postemergence.

Example 4

Control of Volunteer AAD Maize in a Field Planted with Maize Using Alternative Herbicides

In another embodiment, volunteer transgenic maize lines containing the AAD expression cassette (AAD-1) are controlled within a field of maize by the application of a herbicide or combination of herbicides. The specific herbicide used to control the volunteer transgenic AAD maize line is dependent upon the type of maize seed being planted within the field (e.g., Clear Field Maize, Round-up Ready Maize, Liberty Link Maize, etc.).

Furthermore, the AAD trait may be stacked with other additional herbicide tolerant trait(s) via conventional breeding or a molecular stack. In such an example, the specific

herbicide used to control the volunteer AAD maize stacked with another herbicide tolerant trait(s) will be dependent upon the additional herbicide tolerant trait(s) and the type of maize being planted within the field.

The application of a given herbicide can be made before planting at pre-emergence/burndown or post-emergence after planting to control the volunteer transgenic AAD maize lines. Table 5 lists the herbicides to be used at the different stages of planting (pre-emergence or post-emergence) to control volunteer transgenic AAD maize. At or about a 1× Field Rate concentration of herbicide would be applied, as either a tank mix or alone, to the field for both pre-emergent and post-emergent volunteer control.

The control of transgenic maize plants containing the AAD expression cassette within a field of maize would be appli-

cable for the control of an AAD transgenic monocot plant (including, but not limited too; corn, rice, sugar cane, switch grass, turf grass species, sorghum, barley, wheat, and oats, and durum) within a field being planted with a monocot crop (including, but not limited too; corn, rice, sugar cane, switch

grass, turf grass species, sorghum, barley, wheat, and oats, and durum). The example described above, in which volunteer transgenic AAD maize plants are controlled in a maize field, is illustrative of the invention and not intended to restrict the scope of this embodiment.

TABLE 5

Control of volunteer AAD-1 (trait alone or stacked with other HT traits) corn (or other monocot) in corn (or other monocot)			
Previous year Corn hybrid	2008 Corn hybrid being planted	Herbicides option - Burndown/PRE	Herbicides option - POST
Application: alone and tank mixes			
AAD-1	CL Corn	Glyphosate, Dims, glufosinate, paraquat IMI or other ALS herbicides not recommended for corn (need to be aware of planting restrictions)	IMI
	RR Corn		glyphosate
	LL Corn		glufosinate
	GAT + ALS		glyphosate, IMI, ALS inhibitors not usually recommended for in-corn application
	AAD-1 + PAT		glufosinate
	AAD-1 + GT		glyphosate
	AAD-1 + AHAS		IMI
	AAD-1 + PAT + GT		glufosinate, glyphosate
	AAD-1 + PAT + AHAS		glufosinate, glyphosate, IMI
	AAD-1 + PAT + GAT		Glyphosate, glufosinate, IMI, ALS inhibitors not usually recommended for in-corn application
	AAD-1 + GAT	glyphosate, ALS inhibitors not usually recommended for in-corn application	
AAD-1 + PAT	CL Corn	Glyphosate, Dims, paraquat IMI or other ALS herbicides not recommended for corn (need to be aware of planting restrictions)	IMI
	RR Corn		glyphosate
	LL Corn		N/A
	GAT		glyphosate, Imazapyr + imazethapyr, imazamox, imazaquin, ALS inhibitors not usually recommended for in-corn application
	AAD-1 Corn		N/A
	AAD-1 + GT		glyphosate
	AAD-1 + AHAS		IMI
	AAD-1 + PAT + GT		glyphosate
	AAD-1 + PAT + AHAS		IMI
	AAD-1 + PAT + GAT		glufosinate, IMI, ALS inhibitors not usually recommended for in-corn application
	AAD-1 + GAT	glyphosate, IMI, ALS inhibitors not usually recommended for in-corn application	
AAD-1 + GT	CL Corn	Dims, glufosinate, paraquat IMI or other ALS herbicides not recommended for corn (need to be aware of planting restrictions)	IMI
	RR Corn		N/A
	LL Corn		glufosinate
	GAT		IMI, ALS inhibitors not usually recommended for in-corn application
	AAD-1 Corn		N/A
	AAD-1 + PAT		glufosinate
	AAD-1 + AHAS		IMI
	AAD-1 + PAT + GT		glufosinate
	AAD-1 + PAT + AHAS		Glufosinate, IMI
	AAD-1 + GAT		IMI, ALS inhibitors not usually recommended for in-corn application
	AAD-1 + PAT + GAT	glufosinate, IMI, ALS inhibitors not usually recommended for in-corn application	

TABLE 5-continued

Control of volunteer AAD-1 (trait alone or stacked with other HT traits) corn (or other monocot) in corn (or other monocot)				
Previous year Corn hybrid	2008 Corn hybrid being planted	Herbicides option - Burndown/PRE	Herbicides option - POST	
AAD-1 + AHAS	CL Corn	Glyphosate, Dims,	N/A	
	RR Corn	glufosinate, paraquat	glyphosate	
	LL Corn	ALS herbicides not	glufosinate	
	GAT	recommended for corn (need to be aware of planting restrictions)	IMI, ALS inhibitors not usually recommended for in-corn application	
	AAD-1 Corn		N/A	
	AAD-1 + PAT		glufosinate	
	AAD-1 + GT		glyphosate	
	AAD-1 + PAT + GT		glufosinate, glyphosate	
	AAD-1 + PAT + AHAS		glufosinate	
	AAD-1 + GAT		Glyphosate, ALS inhibitors not usually recommended for in-corn application	
	AAD-1 + PAT + GAT		Glyphosate, glufosinate, ALS inhibitors not usually recommended for in-corn application	
	AAD-1 + PAT + GT	CL Corn	Dims, paraquat	IMI
		RR Corn	IMI or other ALS	N/A
LL Corn		herbicides not	N/A	
GAT		recommended for corn (need to be aware of planting restrictions)	IMI, ALS inhibitors not usually recommended for in-corn application	
AAD-1			N/A	
AAD-1 + PAT			N/A	
AAD-1 + GT			N/A	
AAD-1 + AHAS			IMI	
AAD-1 + PAT + AHAS			IMI	
AAD-1 + GAT			ALS inhibitors not usually recommended for in-corn application	
AAD-1 + PAT + GAT			ALS inhibitors not usually recommended for in-corn application	
AAD-1 + PAT + AHAS		CL Corn	Glyphosate, Dims,	N/A
		RR Corn	paraquat	glyphosate
	LL Corn	IMI or other ALS	N/A	
	GAT	herbicides not recommended for corn (need to be aware of planting restrictions)	Glyphosate, ALS inhibitors not usually recommended for in-corn application	
	AAD-1		N/A	
	AAD-1 + PAT		N/A	
	AAD-1 + GT		glyphosate	
	AAD-1 + AHAS		N/A	
	AAD-1 + PAT + GT		glyphosate	
	AAD-1 + GAT		glyphosate, ALS inhibitors not usually recommended for in-corn application	
	AAD-1 + PAT + GAT + ALS		glyphosate, ALS inhibitors not usually recommended for in-corn application	
	AAD-1 + GAT + ALS	CL Corn	Dims, glufosinate,	N/A
		RR Corn	paraquat	N/A
LL Corn		IMI or other ALS	glufosinate	
GAT + ALS		herbicides not recommended for corn (need to be aware of planting restrictions)	N/A	
AAD-1			N/A	
AAD-1 + PAT			glufosinate	
AAD-1 + GT			N/A	
AAD-1 + AHAS			N/A	
AAD-1 + PAT + GT			glufosinate	
AAD-1 + PAT + AHAS			glufosinate	
AAD-1 + GAT + ALS			glufosinate	

TABLE 5-continued

Control of volunteer AAD-1 (trait alone or stacked with other HT traits) corn (or other monocot) in corn (or other monocot)			
Previous year Corn hybrid	2008 Corn hybrid being planted	Herbicides option - Burndown/PRE	Herbicides option - POST
AAD-1 + PAT + GAT + ALS	CL Corn	Dims, paraquat	N/A
	RR Corn	IMI or other ALS	N/A
	LL Corn	herbicides not	N/A
	GAT + ALS	recommended for corn	N/A
	AAD-1	(need to be aware of	N/A
	AAD-1 + PAT	planting restrictions)	N/A
	AAD-1 + GT		N/A
	AAD-1 + AHAS		N/A
	AAD-1 + PAT + GT		N/A
AAD-1 + PAT + AHAS		N/A	

Gene and trait footnotes:

GT = any glyphosate specific tolerance trait including Roundup Ready (CP4), TIPS EPSPS (GA21, Glytol, DMMG), Athenix's EPSPS, GAT only without ALS, GOX, glyphosate decarboxylase, etc.

ALS = double mutant ALS gene insensitive to all ALS herbicide chemistries including IMI herbicides.

AAD-1 = Aryloxyalkanoate dioxygenase gene providing tolerance to all commercial phenoxy auxin and all aryloxyphenoxypionate (fop) herbicides.

AAD-12 = Aryloxyalkanoate dioxygenase gene providing tolerance to phenoxyacetic auxin and pyridyloxyacetic auxin herbicides.

PAT = phosphinothricin acetyltransferase gene providing tolerance to glutamine synthetase inhibitors including, but not limited to, glufosinate. Similar phenotype is provided by genes such as BAR, DSM1, DSM2, et al.

AHAS = imidazolinone specific tolerance gene associated with point mutation at S623 of ALS gene (maize sequence) or equivalent amino acid in other spp (e.g., S653 in *Arabidopsis*).

RR = Roundup Ready trait, implies utility of CP4 gene as commercially deployed either alone or in combination with other genes but imparting glyphosate only tolerance.

CL = Clearfield crops, tolerant by nontransgenic means. Primary tolerance is to imidazolinone class of ALS-inhibiting chemistry with some partial tolerance to specific other herbicides with this mode of action. Use of CL designation is intended to distinguish from transgenic use of the AHAS gene.

LL = Liberty Link trait, implies utility of either PAT or BAR gene as commercially deployed either alone or in combination with other genes but imparting only tolerance to glutamine synthetase inhibitors such as glufosinate.

STS = designates resistance to sulfonylurea herbicide chemistry with use of ALS1 gene.

Herbicide footnotes:

IMI = any imidazolinone herbicide including, but not limited to, imazapyr, imazethapyr, imazamox, imazaquin.

DIMS = cyclohexanedione class of herbicides (dims) including, but not limited to, sethoxydim, clethodim, and for the purposes of this demonstration pinoxaden.

Fops = aryloxyphenoxypionate herbicides (fops) including, but not limited to, quizalofop, haloxyfop, fenoxaprop, fluazifop, et al., their stereospecific isomers or racemic mixtures, and esters, acid, or salts thereof.

ALS inhibitors = any ALS inhibitor to the exclusion of IMI's for the sake of this demonstration (i.e., sulfonylureas, triazolopyrimidine sulfonamides, sulfonylaminocarbonyl triazolinone).

HPPD = p-Hydroxyphenyl pyruvate dioxygenase inhibitor class of chemistry including but not limited to mesotrione, sulcotrione, isoxaflutole, and pyrazolynate.

MSMA and DSMA = herbicides from the organoarsenicals chemistry family.

N/A = No suitable options available postemergence.

Example 5

Control of Volunteer Maize in a Field Planted with
AAD-Maize Using Alternative Herbicides

In an embodiment, volunteer transgenic maize lines (containing the Clear Field trait, Roundup Ready or other Glyphosate Tolerant Trait, Liberty Link Trait, Imidazolinone tolerant trait, or any stacked combination thereof) or volunteer conventional maize lines are controlled within a field of transgenic AAD maize (AAD-1) by the application of aryloxyphenoxypionate herbicide (Fops) including, but not limited to; quizalofop, haloxyfop, fenoxaprop, fluazifop, et al., their stereospecific isomers or racemic mixtures, and esters, acid, or salts thereof. In addition, the Fops herbicide may be applied with another herbicide(s) to control the volunteer maize lines described above.

The specific herbicide used to control the volunteer conventional or transgenic maize line is dependent upon the type of AAD transgenic maize seed (i.e. stacked traits or alone) being planted within the field and the trait possessed by the volunteer maize line. For example an AAD-1 transgenic maize line that has been stacked with another herbicide tolerant trait such as PAT could be sprayed with a herbicide

mixture containing a Fop and glufosinate, but only where the preceding volunteer plants do not contain PAT (or other glufosinate tolerant trait) and AAD-1.

The application of a given herbicide can be made before planting at pre-emergence/burndown or post-emergence after planting to control the volunteer conventional or transgenic maize lines. Table 6 lists the herbicides to be used at the different stages of planting (pre-emergence or post-emergence) to control volunteer conventional or transgenic maize. At or about a 1x Field Rate concentration of herbicide would be applied, as either a tank mix or alone, to the field for both pre-emergent and post-emergent volunteer control.

The control of conventional maize plants or transgenic maize plants containing a herbicide tolerant expression cassette within a field of AAD transgenic maize (either stacked with other herbicide tolerant traits or alone) would be applicable for the control of a herbicide tolerant transgenic monocot plant (including, but not limited too; corn, rice, sugar cane, switch grass, turf grass species, sorghum, barley, wheat, and oats, and durum) within a field being planted with an AAD monocot crop (including, but not limited too; corn, rice, sugar cane, switch grass, turf grass species, sorghum, barley, wheat, and oats, and durum). The example described above, in which volunteer transgenic herbicide tolerant maize plants are controlled in a field planted with AAD transgenic maize, is illustrative of the invention and not intended to restrict the scope of this embodiment.

TABLE 6

Control of volunteer corn (or other monocots) in AAD-1 (alone or stacked with other HT traits) corn (or other monocots)			
Previous year Corn hybrid	2008 Corn hybrid being planted	Herbicides option - Burndown/PRE	Herbicides option - POST
Application: alone and tank mixes			
Conventional corn	AAD-1	Glyphosate, Dims, Fops, glufosinate, paraquat	Fops
	AAD-1 + PAT	IMI or other ALS	Fops, glufosinate
	AAD-1 + GT	herbicides not recommended for corn (need to be aware of planting restrictions)	Fops, glyphosate
	AAD-1 + GAT + ALS		Fops, glyphosate, ALS inhibitors not selective for in-corn application
	AAD-1 + AHAS		Fops, glyphosate, IMI
	AAD-1 + CL		Fops, IMI
	AAD-1 + PAT + GT		Fops, glufosinate, glyphosate
	AAD-1 + PAT + GAT + ALS		Fops, glufosinate, glyphosate, IMI, ALS inhibitors not selective for in-corn application
	AAD-1 + PAT + AHAS		Fops, glufosinate, glyphosate, IMI
	CL Corn	AAD-1	Glyphosate, Dims, Fops, glufosinate, paraquat
AAD-1 + PAT		other ALS herbicides not recommended for corn (need to be aware of planting restrictions)	Fops, glufosinate
AAD-1 + GT			Fops, glyphosate
AAD-1 + GAT + ALS			Fops, glyphosate, ALS inhibitors not selective for in-corn application
AAD-1 + AHAS		Dims, Fops, glufosinate, paraquat	Fops, glyphosate
AAD-1 + CL			Fops
AAD-1 + PAT + GT		IMI or other ALS herbicides not recommended for corn (need to be aware of planting restrictions)	Fops, glufosinate, glyphosate
AAD-1 + PAT + GAT + ALS			Fops, glufosinate, glyphosate, ALS inhibitors not selective for in-corn application
AAD-1 + PAT + AHAS			Fops, glufosinate, glyphosate
RR or GT Corn		AAD-1	
	AAD-1 + PAT		Fops, glufosinate
	AAD-1 + GT		Fops
	AAD-1 + GAT + ALS		Fops, ALS inhibitors not selective for in-corn application
	AAD-1 + AHAS		Fops, IMI
	AAD-1 + CL		Fops, IMI
	AAD-1 + PAT + GT		Fops, glufosinate
	AAD-1 + PAT + GAT + ALS		Fops, glufosinate, IMI, ALS inhibitors not selective for in-corn application
	AAD-1 + PAT + AHAS		Fops, glufosinate, IMI
	LL Corn	AAD-1 + PAT + AHAS	
AAD-1		Glyphosate, Dims, Fops, paraquat	Fops
AAD-1 + PAT		IMI or other ALS	Fops
AAD-1 + GT		herbicides not recommended for corn (need to be aware of planting restrictions)	Fops, glyphosate
AAD-1 + GAT + ALS			Fops, glyphosate, ALS inhibitors not selective for in-corn application
AAD-1 + AHAS			Fops, glyphosate, IMI
AAD-1 + CL			Fops, IMI
AAD-1 + PAT + GT			Fops, glyphosate
AAD-1 + PAT + GAT + ALS			Fops, glyphosate, IMI, ALS inhibitors not selective for in-corn application
AAD-1 + PAT + AHAS			Fops, glyphosate, IMI

TABLE 6-continued

Control of volunteer corn (or other monocots) in AAD-1 (alone or stacked with other HT traits) corn (or other monocots)			
Previous year Corn hybrid	2008 Corn hybrid being planted	Herbicides option - Burndown/PRE	Herbicides option - POST
GAT + ALS	AAD-1	Glyphosate, Dims, Fops, glufosinate, paraquat	Fops
	AAD-1 + PAT		Fops, glufosinate
	AAD-1 + GT		Fops
	AAD-1 + GAT + ALS		Fops, ALS inhibitors not selective for in-corn application
	AAD-1 + AHAS		Fops
	AAD-1 + CL		Fops
	AAD-1 + PAT + GT		Fops, glufosinate
	AAD-1 + PAT + GAT + ALS		Fops, glufosinate
	AAD-1 + PAT + AHAS		Fops, glufosinate
	CL (or AHAS) + PAT		AAD-1
AAD-1 + PAT		Fops	
AAD-1 + GT		Fops, glyphosate	
AAD-1 + GAT + ALS		Fops, glyphosate, ALS inhibitors not selective for in-corn application	
AAD-1 + AHAS		Fops, glyphosate	
AAD-1 + CL		Fops, IMI	
AAD-1 + PAT + GT		Fops, glyphosate	
AAD-1 + PAT + GAT + ALS		Fops, glyphosate, ALS inhibitors not selective for in-corn application	
AAD-1 + PAT + AHAS		Fops, glyphosate	
CL (or AHAS) + GT		AAD-1	Dims, Fops, glufosinate, paraquat other ALS herbicides not recommended for corn (need to be aware of planting restrictions)
	AAD-1 + PAT	Fops, glufosinate	
	AAD-1 + GT	Fops	
	AAD-1 + GAT + ALS	Fops, ALS inhibitors not selective for in-corn application	
	AAD-1 + AHAS	Fops	
	AAD-1 + CL	Fops	
	AAD-1 + PAT + GT	Fops, glufosinate	
	AAD-1 + PAT + GAT + ALS	Fops, glufosinate, ALS inhibitors not selective for in-corn application	
	AAD-1 + PAT + AHAS	Fops, glufosinate	
	GT + PAT	AAD-1 + PAT + AHAS	
AAD-1		Fops	
AAD-1 + PAT		Fops	
AAD-1 + GT		Fops	
AAD-1 + GAT + ALS		Fops, ALS inhibitors not selective for in-corn application	
AAD-1 + AHAS		Fops, IMI	
AAD-1 + CL		Fops, IMI	
AAD-1 + PAT + GT		Fops	
AAD-1 + PAT + GAT + ALS		Fops, IMI, ALS inhibitors not selective for in-corn application	
AAD-1 + PAT + AHAS		Fops, IMI	
GT + CL (or AHAS) + PAT	AAD-1	Dims, Fops, paraquat other ALS herbicides not recommended for corn (need to be aware of planting restrictions)	Fops
	AAD-1 + PAT		Fops
	AAD-1 + GT		Fops
	AAD-1 + GAT + ALS		Fops, ALS inhibitors not selective for in-corn application
	AAD-1 + AHAS		Fops
	AAD-1 + CL		Fops
	AAD-1 + PAT + GT		Fops
	AAD-1 + PAT + GAT + ALS		Fops, ALS inhibitors not selective for in-corn application
	AAD-1 + PAT + AHAS		Fops
	AAD-1 + PAT + AHAS		Fops

TABLE 6-continued

Control of volunteer corn (or other monocots) in AAD-1 (alone or stacked with other HT traits) corn (or other monocots)			
Previous year Corn hybrid	2008 Corn hybrid being planted	Herbicides option - Burndown/PRE	Herbicides option - POST
GAT + ALS + PAT	AAD-1	Dims, Fops, paraquat	Fops
	AAD-1 + PAT		Fops
	AAD-1 + GT		Fops
	AAD-1 + GAT + ALS		Fops
	AAD-1 + AHAS		Fops
	AAD-1 + CL		Fops
	AAD-1 + PAT + GT		Fops
	AAD-1 + PAT + GAT + ALS		Fops
	AAD-1 + PAT + AHAS		Fops

Gene and trait footnotes:

GT = any glyphosate specific tolerance trait including Roundup Ready (CP4), TIPS EPSPS (GA21, Glytol, DMMG), Athenix's EPSPS, GAT only without ALS, GOX, glyphosate decarboxylase, etc.

ALS = double mutant ALS gene insensitive to all ALS herbicide chemistries including IMI herbicides.

AAD-1 = Aryloxyalkanoate dioxygenase gene providing tolerance to all commercial phenoxy auxin and all aryloxyphenoxypropionate (fop) herbicides.

AAD-12 = Aryloxyalkanoate dioxygenase gene providing tolerance to phenoxyacetic auxin and pyridyloxyacetic auxin herbicides.

PAT = phosphinothricin acetyltransferase gene providing tolerance to glutamine synthetase inhibitors including, but not limited to, glufosinate. Similar phenotype is provided by genes such as BAR, DSM1, DSM2, et al.

AHAS = imidazolinone specific tolerance gene associated with point mutation at S623 of ALS gene (maize sequence) or equivalent amino acid in other spp (e.g., S653 in *Arabidopsis*).

RR = Roundup Ready trait, implies utility of CP4 gene as commercially deployed either alone or in combination with other genes but imparting glyphosate only tolerance.

CL = Clearfield crops, tolerant by nontransgenic means. Primary tolerance is to imidazolinone class of ALS-inhibiting chemistry with some partial tolerance to specific other herbicides with this mode of action. Use of CL designation is intended to distinguish from transgenic use of the AHAS gene.

LL = Liberty Link trait, implies utility of either PAT or BAR gene as commercially deployed either alone or in combination with other genes but imparting only tolerance to glutamine synthetase inhibitors such as glufosinate.

STS = designates resistance to sulfonylurea herbicide chemistry with use of ALS1 gene.

Herbicide footnotes:

IMI = any imidazolinone herbicide including, but not limited to, imazapyr, imazethapyr, imazamox, imazaquin.

DIMS = cyclohexanedione class of herbicides (dims) including, but not limited to, sethoxydim, clethodim, and for the purposes of this demonstration pinoxaden.

Fops = aryloxyphenoxypropionate herbicides (fops) including, but not limited to, quizalofop, haloxyfop, fenoxaprop, fluazifop, et al., their stereospecific isomers or racemic mixtures, and esters, acid, or salts thereof.

ALS inhibitors = any ALS inhibitor to the exclusion of IMI's for the sake of this demonstration (i.e., sulfonylureas, triazolopyrimidine sulfonamides, sulfonylaminocarbonyltriazolinone).

HPPD = p-Hydroxyphenyl pyruvate dioxygenase inhibitor class of chemistry including but not limited to mesotrione, sulcotrione, isoxaflutole, and pyrazolynate.

MSMA and DSMA = herbicides from the organoarsenicals chemistry family.

N/A = No suitable options available postemergence.

Example 5

Testing of Post-Emergent Herbicide Application on Transgenic Maize Containing AAD-1 and PAT

Transgenic Hi-II maize plants containing the AAD-1 and PAT expression cassette (RB7 MAR v3::*Zea Maize* Ubiquitin Promoter v2::AAD-1 v3::*Zea Maize* Per5 3'UTR v2::Rice Actin1 Promoter v2::PAT v3::*Zea Maize* Lipase 3'UTR v1) from pDAB3404 and Hi-II non-transgenic control plants were propagated for testing. Seed were planted into Metro Mix Media (Sun Gro Horticulture Inc., Bellevue, Wash.) and

subirrigated with Hoagland's solution until wet, then allowed to gravity drain. Pots were placed in a glasshouse environment set to 27° C. until the germinated plants reached the V1-V3 growth stage. At or about, the V1-V3 growth stage, the maize plants were sprayed using a track sprayer and a drop down nozzle calibrated to deliver a 187 L/ha solution of herbicide at the 1× Field Rate (Table 7). After treatment the plants were placed in the greenhouse and scored via visual injury assessment after 14 days. Table 7 also lists the tolerance of the plants to the applied herbicides as determined by the visual injury assessment.

TABLE 7

Graminical active ingredients on AAD-1 + PAT + glyphosate tol Tolerance (-/+)				
Trt	Chemical Name	WT	3404	Rate (1x)/Comment
Select	clethodim	-	-	26.25-52.5 g ai/ha
Poast	sethoxydim	-	-	105-210 g ai/ha
Achieve Liquid	tralkoxydim	-	-	100 g ai/ha
Pursuit	imazethapyr	-/+	-/+	Need higher than 70 g ai/ha rate (50% injury @ 70 g ai/ha)
Raptor	imazamox	-	-	44 g ai/ha
Scepter	imazaquin	-/+	-/+	Need higher than 50 g ai/ha rate (50% injury @ 50 g ai/ha)
Assure II	quizalofop	-	+	35-70 g ai/ha
Discover	clodinafop	-	+	28-56 g ai/ha
Hoelon	diclofop-methyl	-	-	Not active on grasses
Fusilade DX	fluazifop-P-butyl	-	+	105 g ai/ha
Gallant Super	haloxyfop-methyl R	-	+	17.5-35 g ai/ha

TABLE 7-continued

Graminical active ingredients on AAD-1 + PAT + glyphosate tol				
Tolerance (-/+)				
Trt	Chemical Name	WT	3404	Rate (1x)/Comment
Puma Super	fenoxaprop-P-ethyl	-	+	23-46 g ai/ha
Clincher SF	cyhalofop-butyl	-	+	280 g ai/ha
Axial	pinoxaden	+/-	+/-	Not active but small amt of control on both WT and AAD-1
Glyphomax XRT	glyphosate-isopropylammonium	-	+	420-840 g ai/ha
Liberty	glufosinate-ammonium	-	+	240-480 g ai/ha
Hoelon	diclofop-methyl	-	-	Not active on grasses
Fusilade DX	fluazifop-P-butyl	-	+	105 g ai/ha
Gallant Super	haloxyfop-methyl R	-	+	17.5-35 g ai/ha
Puma Super	fenoxaprop-P-ethyl	-	+	23-46 g ai/ha
Clincher SF	cyhalofop-butyl	-	+	280 g ai/ha
Axial	pinoxaden	+/-	+/-	Not active but small amt of control on both WT and AAD-1
Glyphomax XRT	glyphosate-isopropylammonium	-	+	420-840 g ai/ha
Liberty	glufosinate-ammonium	-	+	240-480 g ai/ha

Note:

3404 column contains molecular stack of AAD-1, and PAT crossed conventionally with CP4 for glyphosate tolerance

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 4

<210> SEQ ID NO 1

<211> LENGTH: 295

<212> TYPE: PRT

<213> ORGANISM: Sphingobium herbicidovorans

<400> SEQUENCE: 1

```

Met His Ala Ala Leu Ser Pro Leu Ser Gln Arg Phe Glu Arg Ile Ala
1          5          10          15

Val Gln Pro Leu Thr Gly Val Leu Gly Ala Glu Ile Thr Gly Val Asp
          20          25          30

Leu Arg Glu Pro Leu Asp Asp Ser Thr Trp Asn Glu Ile Leu Asp Ala
          35          40          45

Phe His Thr Tyr Gln Val Ile Tyr Phe Pro Gly Gln Ala Ile Thr Asn
50          55          60

Glu Gln His Ile Ala Phe Ser Arg Arg Phe Gly Pro Val Asp Pro Val
65          70          75          80

Pro Leu Leu Lys Ser Ile Glu Gly Tyr Pro Glu Val Gln Met Ile Arg
          85          90          95

Arg Glu Ala Asn Glu Ser Gly Arg Val Ile Gly Asp Asp Trp His Thr
          100          105          110

Asp Ser Thr Phe Leu Asp Ala Pro Pro Ala Ala Val Val Met Arg Ala
          115          120          125

Ile Asp Val Pro Glu His Gly Gly Asp Thr Gly Phe Leu Ser Met Tyr
130          135          140

Thr Ala Trp Glu Thr Leu Ser Pro Thr Met Gln Ala Thr Ile Glu Gly
145          150          155          160

Leu Asn Val Val His Ser Ala Thr Arg Val Phe Gly Ser Leu Tyr Gln
          165          170          175

Ala Gln Asn Arg Arg Phe Ser Asn Thr Ser Val Lys Val Met Asp Val
          180          185          190

Asp Ala Gly Asp Arg Glu Thr Val His Pro Leu Val Val Thr His Pro
195          200          205

```

-continued

Gly Ser Gly Arg Lys Gly Leu Tyr Val Asn Gln Val Tyr Cys Gln Arg
 210 215 220

Ile Glu Gly Met Thr Asp Ala Glu Ser Lys Pro Leu Leu Gln Phe Leu
 225 230 235 240

Tyr Glu His Ala Thr Arg Phe Asp Phe Thr Cys Arg Val Arg Trp Lys
 245 250 255

Lys Asp Gln Val Leu Val Trp Asp Asn Leu Cys Thr Met His Arg Ala
 260 265 270

Val Pro Asp Tyr Ala Gly Lys Phe Arg Tyr Leu Thr Arg Thr Thr Val
 275 280 285

Gly Gly Val Arg Pro Ala Arg
 290 295

<210> SEQ ID NO 2
 <211> LENGTH: 295
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: AAD-1 v2 Translation
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (2)..(2)
 <223> OTHER INFORMATION: Different from v1
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (212)..(212)
 <223> OTHER INFORMATION: Different from v1

<400> SEQUENCE: 2

Met Ala Ala Ala Leu Ser Pro Leu Ser Gln Arg Phe Glu Arg Ile Ala
 1 5 10 15

Val Gln Pro Leu Thr Gly Val Leu Gly Ala Glu Ile Thr Gly Val Asp
 20 25 30

Leu Arg Glu Pro Leu Asp Asp Ser Thr Trp Asn Glu Ile Leu Asp Ala
 35 40 45

Phe His Thr Tyr Gln Val Ile Tyr Phe Pro Gly Gln Ala Ile Thr Asn
 50 55 60

Glu Gln His Ile Ala Phe Ser Arg Arg Phe Gly Pro Val Asp Pro Val
 65 70 75 80

Pro Leu Leu Lys Ser Ile Glu Gly Tyr Pro Glu Val Gln Met Ile Arg
 85 90 95

Arg Glu Ala Asn Glu Ser Gly Arg Val Ile Gly Asp Asp Trp His Thr
 100 105 110

Asp Ser Thr Phe Leu Asp Ala Pro Pro Ala Ala Val Val Met Arg Ala
 115 120 125

Ile Asp Val Pro Glu His Gly Gly Asp Thr Gly Phe Leu Ser Met Tyr
 130 135 140

Thr Ala Trp Glu Thr Leu Ser Pro Thr Met Gln Ala Thr Ile Glu Gly
 145 150 155 160

Leu Asn Val Val His Ser Ala Thr Arg Val Phe Gly Ser Leu Tyr Gln
 165 170 175

Ala Gln Asn Arg Arg Phe Ser Asn Thr Ser Val Lys Val Met Asp Val
 180 185 190

Asp Ala Gly Asp Arg Glu Thr Val His Pro Leu Val Val Thr His Pro
 195 200 205

Gly Ser Gly Cys Lys Gly Leu Tyr Val Asn Gln Val Tyr Cys Gln Arg
 210 215 220

Ile Glu Gly Met Thr Asp Ala Glu Ser Lys Pro Leu Leu Gln Phe Leu
 225 230 235 240

-continued

Tyr Glu His Ala Thr Arg Phe Asp Phe Thr Cys Arg Val Arg Trp Lys
 245 250 255

Lys Asp Gln Val Leu Val Trp Asp Asn Leu Cys Thr Met His Arg Ala
 260 265 270

Val Pro Asp Tyr Ala Gly Lys Phe Arg Tyr Leu Thr Arg Thr Thr Val
 275 280 285

Gly Gly Val Arg Pro Ala Arg
 290 295

<210> SEQ ID NO 3
 <211> LENGTH: 296
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: AAD-1 v3 Translation
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (2)..(3)
 <223> OTHER INFORMATION: Different from V1

<400> SEQUENCE: 3

Met Ala His Ala Ala Leu Ser Pro Leu Ser Gln Arg Phe Glu Arg Ile
 1 5 10 15

Ala Val Gln Pro Leu Thr Gly Val Leu Gly Ala Glu Ile Thr Gly Val
 20 25 30

Asp Leu Arg Glu Pro Leu Asp Asp Ser Thr Trp Asn Glu Ile Leu Asp
 35 40 45

Ala Phe His Thr Tyr Gln Val Ile Tyr Phe Pro Gly Gln Ala Ile Thr
 50 55 60

Asn Glu Gln His Ile Ala Phe Ser Arg Arg Phe Gly Pro Val Asp Pro
 65 70 75 80

Val Pro Leu Leu Lys Ser Ile Glu Gly Tyr Pro Glu Val Gln Met Ile
 85 90 95

Arg Arg Glu Ala Asn Glu Ser Gly Arg Val Ile Gly Asp Asp Trp His
 100 105 110

Thr Asp Ser Thr Phe Leu Asp Ala Pro Pro Ala Ala Val Val Met Arg
 115 120 125

Ala Ile Asp Val Pro Glu His Gly Gly Asp Thr Gly Phe Leu Ser Met
 130 135 140

Tyr Thr Ala Trp Glu Thr Leu Ser Pro Thr Met Gln Ala Thr Ile Glu
 145 150 155 160

Gly Leu Asn Val Val His Ser Ala Thr Arg Val Phe Gly Ser Leu Tyr
 165 170 175

Gln Ala Gln Asn Arg Arg Phe Ser Asn Thr Ser Val Lys Val Met Asp
 180 185 190

Val Asp Ala Gly Asp Arg Glu Thr Val His Pro Leu Val Val Thr His
 195 200 205

Pro Gly Ser Gly Arg Lys Gly Leu Tyr Val Asn Gln Val Tyr Cys Gln
 210 215 220

Arg Ile Glu Gly Met Thr Asp Ala Glu Ser Lys Pro Leu Leu Gln Phe
 225 230 235 240

Leu Tyr Glu His Ala Thr Arg Phe Asp Phe Thr Cys Arg Val Arg Trp
 245 250 255

Lys Lys Asp Gln Val Leu Val Trp Asp Asn Leu Cys Thr Met His Arg
 260 265 270

-continued

Ala Val Pro Asp Tyr Ala Gly Lys Phe Arg Tyr Leu Thr Arg Thr Thr
 275 280 285

Val Gly Gly Val Arg Pro Ala Arg
 290 295

<210> SEQ ID NO 4

<211> LENGTH: 8557

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: AAD-1 insert and flanking sequences

<400> SEQUENCE: 4

actggtatTT aatatactTT aataaatatt attagattcc tcgtcaaaga actTTTTtaca 60
 atatattctat ttagaatcat atatgtcata gTTTTTTTTc taagagtcta gTTTactagt 120
 aaaatccgac tcacattTTT cgaacttggg atgcaacact taaatagtac aaaaccttgg 180
 tatgcagtat tttacattgt aagattcaaa atttctaaag cagtatatat atgTTTccag 240
 aaacttatag atatagaaaa aacagagaga cgtatgCGaa aattCGataa aggtgtacat 300
 tggattCGca aggctaaata catattTatc gtggatccat gcagagTTTg ggtaataaaa 360
 ttagatactt ccaatcatgt gccacataat cacgtaacat tagtaattta aatgacatta 420
 ccatgtccaa ctgattTaaa acacaaactc ttcttgaacc atatagTTTg acaaaccaaa 480
 tatatataac tggagctact agttatgaat caattaaaaa ttactTTgaa gattcaacgt 540
 agtGCCagtt tggctctagc acatctaacc agaaggGcta aggctggctt caacaggaac 600
 agccaaatcc gagatCGagc catttgccat tTTTgggtag ttagTTTaaC tttcatatat 660
 cttcccatcc tTTTTTgcct agcctaaatg gctTTgatgt tgaagaccat attaattTgc 720
 ttcagtggca ctaggacaac catattggct ttggctgacc cgTTtagagtt agcctaatgg 780
 gtggaagggg agggaagggg aggatCGatg gtggcatgag agaggggTTg acgatcacga 840
 tgatgatgCG agtgaggagg agagggTggc gacgacacag gggagaaagg agagggacgc 900
 taggagcgTc aagggcgTgg gggagggggag ggtCGgaggg atgaaggatg acctaaatat 960
 tattgttgag tgatagaggg ttattcaact atccgaccCG tcgattTTga tggTatgtta 1020
 aattTgtgtg tcattTgttt gatggattta gtaaaggTTa tgggtctaga ggtgattTTt 1080
 gtgggTggg tTTTcacagag tTTaaactag cggattatat agtggTatag aagatatagt 1140
 tttattagaa catctccaaa atgtgactcg aaataatacc cccaaaattt aaaatactac 1200
 atcattTTga taaaaaggT aaagtagagc actgtTgGaa cagTTTTTaa aagTtTgTgc 1260
 ctatattTTa aaatagggtA ctgattTaaa atattgtTgt gggggataga tatccccggg 1320
 tccactagaa ggcgagaagg cctCGcgtgt ggccacgggc cagTTacccc gcaaggccat 1380
 cccttCGTgg gtcgagctag aattactggT agaatgggct gaccgaagaa ggcaacagac 1440
 tcgagcccaa acaatccatc ggctcgtgCG ctatccacag aaactacccg actTTcggc 1500
 gcatggcatc ctagaatatc ggggcgtatt agggatgagT cagcgagatt ttcggaagat 1560
 tagTtcagTt tGttcGctat tatttaggag acatatgatc ctcatgtacg tatggagTgc 1620
 cccacggTcg tGtatataag gtccagaggg taccatca tttctatCGa ccatctacct 1680
 atctcatcag cTTTTctca ttcaggagac ctCGcttGta acccaccaca tatagatcca 1740
 tcccaagaag tagTgtatta cgctctcta agcgGCCaa actTgcagaa aaccgctat 1800
 ccctctctCG tgcgtccagc acgaaccatt gagTtacaat caacagcacc gtacctTgaa 1860
 gcggaataca atgaaggTta gctacgattt acagcaaagc cagaatacaa tgaaccataa 1920

-continued

agtgattgaa	gctcgaata	tacgaaggaa	caaatat	taaaaaata	cgcaatgact	1980	
tggaacaaaa	gaaagtgata	tat	tcttaacaa	gcatccctc	taaagaatgg	2040	
cagtttctct	ttgcatgtaa	ctattatgct	cccttcgta	caaaaat	ggactactat	2100	
tggaacttc	ttctgaaaat	agtggccacc	gcttaattaa	ggcgcccat	gcccgggcaa	2160	
gcggccgctt	aattaaatt	aatggttaa	actaggaaat	ccaagcttg	atgcctgcag	2220	
atccccggg	atcctctaga	gtcgacctgc	agtgcagcgt	gaccggctg	tgccccctc	2280	
tagagataat	gagcattgca	tgtctaagtt	ataaaaaatt	accacatatt	ttttttgtca	2340	
cacttgtttg	aagtgcagtt	tatctatctt	tatacatata	tttaaacttt	actctacgaa	2400	
taatataatc	tatagtacta	caataatata	agtgttttag	agaatcatat	aatgaacag	2460	
ttagacatgg	tctaaaggac	aattgagtat	ttgacaaca	ggactctaca	gttttatctt	2520	
tttagtgctc	atgtgttctc	cttttttttt	gcaaatagct	tcacctatat	aatacttcat	2580	
ccat	tttatt	agtacatcca	tttagggttt	agggttaatg	gtttttatag	actaat	2640
tttagtacatc	tattttatc	tatttttagcc	tctaaattaa	gaaaactaaa	actctat	2700	
agttttttta	tttaatagtt	tagatataaa	atagaataaa	ataaagtgc	taaaaattaa	2760	
acaaatacc	tttaagaaat	taaaaaaact	aaggaaacat	tttcttgtt	tcgagtagat	2820	
aatgccagcc	tgtaaacgc	cgtcgacgag	tctaacggac	accaaccagc	gaaccagcag	2880	
cgtcgctgc	ggccaagcga	agcagacggc	acggcatctc	tgctcctg	tctggacccc	2940	
tctcgagagt	tccgctccac	cgttgactt	gctccgctgt	cggcatccag	aaattgcgtg	3000	
gcggagcggc	agacgtgagc	cggcacggca	ggcgccctcc	tcctcctc	acggcaccgg	3060	
cagctacggg	ggattcctt	cccaccgctc	cttcgctt	ccttcctc	ccgctgta	3120	
aaatagacac	cccctccaca	ccctcttcc	ccaacctcgt	gttggtcga	gcgcacacac	3180	
acacaaccag	atctccccca	aatccaccg	tcggcacctc	cgcttcaagg	taacccgctc	3240	
gtcctcccc	cccccccc	tctctacctt	ctctagatcg	gcgttcggt	ccatgcatgg	3300	
ttaggcccg	gtagttctac	ttctgttcat	gtttgtgta	gatccgtgtt	tgtgttagat	3360	
ccgtgctgct	agcgttcgta	cacggatg	acctgtacgt	cagacacgtt	ctgattgcta	3420	
acttgccagt	gtttctctt	ggggaatcct	gggatggctc	tagccgttcc	gcagacggga	3480	
tcgatttcat	gattttttt	gtttcgttgc	atagggtttg	gtttgccctt	ttcctttatt	3540	
tcaatatatg	ccgtgcactt	gtttgtcggg	tcactctt	atgctttttt	ttgtcttgg	3600	
tgtgatgatg	tggtctggtt	ggcggtcgt	tctagatcgg	agtagaattc	tgtttcaaac	3660	
tacctggtgg	atttattaat	tttggatctg	tatgtgtgtg	ccatacatat	tcatagttac	3720	
gaattgaaga	tgatggatgg	aatatcgat	ctaggatagg	tatacatggt	gatgcccgtt	3780	
ttactgatgc	atatacagag	atgctttttg	ttcgttgg	tgtgatgatg	tggtgtggtt	3840	
ggcggtcgt	tcattcgttc	tagatcggag	tagaatactg	tttcaacta	cctgggtgat	3900	
ttattaattt	tggaactgta	tgtgtgtgtc	atacatctc	atagttacga	gtttaagatg	3960	
gatgaaata	tcgatctagg	ataggtatac	atgttgatgt	gggttttact	gatgcatata	4020	
catgatggca	tatgcagcat	ctattcatat	gctctaacct	tgagtaccta	tctattataa	4080	
taaacaagta	tgttttataa	ttatttcgat	cttgatatac	ttggatgatg	gcatatgcag	4140	
cagctatatg	tggattttt	tagccctgcc	ttcatacgt	at	ttggtactgt	4200	
ttctttgtc	gatgctcacc	ctgttgtttg	gtgttacttc	tgcaggtac	ccccggggtc	4260	
gaccatggct	catgctgcc	tcagccctct	ctccaacgc	ttgagagaa	tagctgtcca	4320	

-continued

gccactcact	ggtgtccttg	gtgctgagat	cactggagtg	gacttgaggg	aaccacttga	4380
tgacagcacc	tggaatgaga	tattggatgc	cttccacact	taccaagtca	tctactttcc	4440
tggccaagca	atcaccaatg	agcagcacat	tgcatctctca	agaaggtttg	gaccagttga	4500
tccagtgcct	cttctcaaga	gcattgaagg	ctatccagag	gttcagatga	tccgcagaga	4560
agccaatgag	tctggaaggg	tgattggtga	tgactggcac	acagactcca	ctttccttga	4620
tgacactcca	gctgctggtg	tgatgagggc	catagatggt	cctgagcatg	gcggagacac	4680
tggttccctt	tcaatgtaca	cagcttggga	gaccttgtct	ccaaccatgc	aagccacat	4740
cgaagggctc	aacgttgtgc	actctgccac	acgtgtgttc	ggttccctct	accaagcaca	4800
gaaccgtcgc	ttcagcaaca	cctcagtcaa	ggtgatggat	gttgatgctg	gtgacagaga	4860
gacagtccat	cccttggttg	tgactcatcc	tggtctgga	aggaaaggcc	tttatgtgaa	4920
tcaagtctac	tgtcagagaa	ttgagggcat	gacagatgca	gaatcaaagc	cattgcttca	4980
gttccctctat	gagcatgcca	ccagatttga	cttcaactgc	cgtgtgaggt	ggaagaaaga	5040
ccaagtccct	gtctgggaca	acttgtgcac	catgcaccgt	gctgttccctg	actatgctgg	5100
caagttcaga	tacttgactc	gcaccacagt	tggtggagtt	aggcctgcc	gctgagtagt	5160
tagcttaatc	acctagagct	cgtttaaact	gagggcactg	aagtcgcttg	acgtgctgaa	5220
ttgtttgtga	tgttggtggc	gtattttgtt	taaataagta	agcatggctg	tgattttatc	5280
atatgatcga	tctttggggt	tttatttaac	acattgtaa	atgtgtatct	attaataact	5340
caatgtataa	gatgtgttca	ttcttcgggt	gccatagatc	tgcttatttg	acctgtgatg	5400
ttttgactcc	aaaaaccaa	atcacaactc	aataaactca	tggaatatgt	ccacctgttt	5460
cttgaagagt	tcactacca	ttccagttgg	catttatcag	tgttgcagcg	gcgctgtgct	5520
ttgtaacata	acaattgtta	cggcatatat	ccaatagcgg	cggcctcct	gcagggttta	5580
aacttgccgt	ggcctatttt	cagaagaagt	tccaatagt	agtccaaaat	ttttgtaacg	5640
aaggagcat	aatagttaca	tgcaaaggaa	aactgccatt	ctttagaggg	gatgcttggt	5700
taagaacaaa	aaatatatca	ctttcttttg	ttccaagtca	ttgcgtattt	ttttaaaaat	5760
atgtgtcct	tcgtatattt	cgagcttcaa	tcactttatg	gttctttgta	ttctggcttt	5820
gctgtaaatc	gtagctaacc	ttcttcctag	cagaaattat	taatacttgg	gatatttttt	5880
tagaatcaag	taaattacat	attaccacca	catcgagctg	cttttaaatt	catattacag	5940
ccatataggg	ttgattcatt	ttgcaaaatt	tccaggatat	tgacaacggt	aacttaataa	6000
tatcttgaaa	tattaaagct	attatgatta	gggtgcaaa	tgaccagagt	tggttcgggt	6060
tatatcaaaa	tcaaaccaaa	ccaactatat	cggtttgat	tggttcgggt	ttgccgggtt	6120
ttcagcattt	tctggttttt	tttttgtag	atgaatatta	ttttaatctt	actttgtcaa	6180
atttttgata	agtaaatata	tgtgtagta	aaaattaatt	ttttttaca	acatatgatc	6240
tattaaaata	ttcttatagg	agaattttct	taataacaca	tgatatttat	ttattttagt	6300
cgtttgacta	attttctgtt	gatgtacact	ttcaaagtta	accaaattta	gtaattaagt	6360
ataaaaatca	atatgatacc	taaataatga	tatgttctat	ttaattttaa	attatcgaaa	6420
tttcaactca	aattcgaaaa	agatatataa	gaattttgat	agattttgac	atatgaatat	6480
ggaagaacaa	agagattgac	gcatttttagt	aacacttgat	aagaaagtga	tcgtacaacc	6540
aattatttaa	agttaataaa	aatggagcac	ttcatattta	acgaaatatt	acatgccaga	6600
agagtcgcaa	atatttctag	atatttttta	aagaaaattc	tataaaaagt	cttaaaggca	6660
tatatataaa	aactatatat	ttatattttt	tacccaaaag	caccgcaagg	ggtagccctg	6720

-continued

```

gggtgtgcgga cggactctaa acaccgacag ctggcgcgcc aggtaggggg tgtgtctttg 6780
atctgagcta gctcaatgac cttacctcc aaatgcaaga tcgcccttcg ccccgggact 6840
atgttttgct ttggaacct ctcatccata gcagatgaag agggaaactct gcaccgcata 6900
gcagatctat tggagaagaa gctttcctca gaaatctcga ggggagccag ggcagaacag 6960
cgggtggcac catcacccgc acctcaagcg aagatgacct cttacaaacc gaaagtccggg 7020
agctcaccta cccgaaaaac tccgctgtcc acttcgcccc caaaggagtg gacacggatt 7080
actcgaagaa aggaagcgag tgtcccagat caggggacgg gaacacgcca agccatcttt 7140
ccgacgcctt cgcctcaaaa tgaggatgga aagaagagcg ccatcgcgct ggctcctttc 7200
taccocgacg tctcttcat cagggggaga ttggagttag caccctctt caacgatgag 7260
ccaacctgac aaggggaaga gcctccccag cgtgaggcgc gacgacggag gaatagaagc 7320
cagaacgtgc ggcgacatca cgaggctggg gaacgggatc cggcgcaacc cgtatcccgg 7380
gacgaagctt tagaagtagg aaaaactccc gacgagtggg tacaccgaga aaggcggaac 7440
tctcgcgcc gtgatcgccg acaagcttag gaccgagaac gagagcaagc cgagcaaggt 7500
gcaaggctgc gccgagagaa tgctctctt gctcggaaacc tgtaccccg cttcgtctcg 7560
gcaatgaaca cgccgagtga agtcggaggg gtactggccc agatagctga cggcctcccg 7620
cgaaccctag acacggaagg ctaccggcgg ctgcttactc gagcagttaa tcaccttcta 7680
cccatcacta atcctccaag cgacctacgc catgcatca acagccggcg agacacgcg 7740
agctccatca acgcttcgcg cgaccgatga cacgaaagt agataggga ccgagaggag 7800
tatgtccgag atcatgccat cctggcatga agtcatgcca cccgagctga gtcggttgcg 7860
gcctcgacca gtgtcccgtt ccagggacga tcaagatgac acacaactgg ctcccctcct 7920
tgggaccgac ctacgaacg ccgacatgaa gacacgtgcg gagtcttcgc acttactccg 7980
tgtctccggg ccatccagt gcccctaact tcaaggtctc caacgtcagc aagtatgagc 8040
gcaagcagga cctgggtggc tggttagcca tctacacgat tgtcacatgg gccgccggag 8100
cgacggagga cgtgatgaca gtgtatcttc ccattgtcct agggcaagac gcaatgcagt 8160
ggctccgaca tctaccccaa cattgcatag acaattggag cgacttcagt tgggtgcttca 8220
tcgccaactt ccagtcctc tttgacaagc cggcgagcc atgggaccta aaatccattg 8280
ggcatcaggg cgatgaaacg ctccggttgt acctcaagag gttttagacc atgaggaacc 8340
acacccccga agtcgcccag gcggggggtga ttgaagactt ctaccgagga tccaatgact 8400
cggctttcgt ccgagccata ctccagaaaa gcgtcggcca cctccgaaca cttgttccgg 8460
gaggcagacc tctacatcac cacggattaa cgggccagg acctcatcg aggcacgaaa 8520
gccgcgccac acgcgccag gtgtgacacg aaccagc 8557

```

The invention claimed is:

1. A method of controlling Aryloxy Alkanoate Dioxygenase (AAD-1) volunteer corn plants, comprising Event DAS-40278-9 as available in seed deposited under ATCC deposit number PTA-10244, in a field comprising dicot plants, said volunteer corn plants comprising an AAD-1 gene, wherein said method comprises applying a herbicide to said volunteer corn plants.

2. The method of claim 1, wherein said dicot plants are selected from the group consisting of soybeans and cotton.

3. The method of claim 1, wherein said AAD-1 gene encodes a protein selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, and SEQ ID NO:3.

4. The method of claim 1, wherein said herbicide is selected from the group consisting of a cyclohexanedione and an imidazolinone.

5. The method of claim 4, wherein said herbicide is a cyclohexanedione selected from the group consisting of clethodim and sethosydin.

6. The method of claim 4, wherein said herbicide is an imidazolinone selected from the group consisting of imazethapyr, imazamox, and imazaquin.

7. The method of claim 6, wherein said dicot plants are soybeans, and said volunteer corn plants are other than imidazolinone-tolerant corn.

8. The method of claim 1, wherein said volunteer corn plants comprise a glyphosate- and/or a glufosinate-tolerance gene.

9. A method of controlling AAD-1 volunteer corn plants in a field comprising dicot plants, said volunteer corn plants comprising SEQ ID NO:4 of corn Event DAS-40278-9, wherein said method comprises applying a herbicide to said volunteer corn plants, wherein said volunteer corn plants are susceptible to said herbicide, and said dicot plants are tolerant to said herbicide.

10. The method of claim 9, wherein said herbicide is selected from the group consisting of glyphosate and glufosinate.

11. The method of claim 9, wherein said dicot plants are selected from the group consisting of soybeans and cotton.

12. The method of claim 9, wherein said herbicide is selected from the group consisting of a cyclohexanedione and an imidazolinone.

13. The method of claim 12, wherein said volunteer corn plants comprise a glyphosate- and/or a glufosinate-tolerance gene.

14. The method of claim 13, wherein said herbicide is selected from the group consisting of a cyclohexanedione and an imidazolinone.

15. The method of claim 9, wherein said herbicide is selected from the group consisting of acetolactate synthase inhibitors and acetohydroxyacid synthase inhibitors.

16. The method of claim 9, wherein said AAD-1 gene encodes a protein selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, and SEQ ID NO:3.

17. The method of claim 9, wherein said herbicide is a cyclohexanedione selected from the group consisting of clethodim and sethoxydim.

18. The method of claim 9, wherein said herbicide is an imidazolinone selected from the group consisting of imazethapyr, imazamox, and imazaquin.

19. The method of claim 18, wherein said dicot plants are soybeans, and said corn is other than imidazolinone-tolerant corn.

20. The method of claim 1, wherein said herbicide is selected from the group consisting of glyphosate and glufosinate.

21. The method of claim 1, wherein said herbicide is selected from the group consisting of acetolactate synthase inhibitors and acetohydroxyacid synthase inhibitors.

* * * * *