



(10) **Patent No.:** US 9,009,087 B1  
(45) **Date of Patent:** Apr. 14, 2015

## OTHER PUBLICATIONS

Abawi, G. Y., R. J. Smith, and D. K. Brady. "Assessment of the value of long range weather forecasts in wheat harvest management." *Journal of agricultural engineering research* 62.1 (1995): 39-48.\*

Smith, G. P., and M. J. Gooding. "Models of wheat grain quality considering climate, cultivar and nitrogen effects." *Agricultural and forest meteorology* 94.3 (1999): 159-170.\*

de Toro, Alfredo, et al. "Cereal harvesting-strategies and costs under variable weather conditions." *Biosystems Engineering* 111.4 (2012): 429-439.\*

Strahan, Rod, and Jim Page. "Economics of on-farm grain storage and drying." *Stored Grain in Australia 2003*, Proceedings of the Australian Postharvest Technical Conference, Canberra. 2003, pp. 15-18.\*

\* cited by examiner

*Primary Examiner* — David Vincent

(74) *Attorney, Agent, or Firm* — Lazaris IP

(57) **ABSTRACT**

A modeling framework for evaluating the impact of weather conditions on farming and harvest operations applies real-time, field-level weather data and forecasts of meteorological and climatological conditions together with user-provided and/or observed feedback of a present state of a harvest-related condition to agronomic models and to generate a plurality of harvest advisory outputs for precision agriculture. A harvest advisory model simulates and predicts the impacts of this weather information and user-provided and/or observed feedback in one or more physical, empirical, or artificial intelligence models of precision agriculture to analyze crops, plants, soils, and resulting agricultural commodities, and provides harvest advisory outputs to a diagnostic support tool for users to enhance farming and harvest decision-making, whether by providing pre-, post-, or in situ-harvest operations and crop analyzes.

(57) **ABSTRACT**


A modeling framework for evaluating the impact of weather conditions on farming and harvest operations applies real-time, field-level weather data and forecasts of meteorological and climatological conditions together with user-provided and/or observed feedback of a present state of a harvest-related condition to agronomic models and to generate a plurality of harvest advisory outputs for precision agriculture. A harvest advisory model simulates and predicts the impacts of this weather information and user-provided and/or observed feedback in one or more physical, empirical, or artificial intelligence models of precision agriculture to analyze crops, plants, soils, and resulting agricultural commodities, and provides harvest advisory outputs to a diagnostic support tool for users to enhance farming and harvest decision-making, whether by providing pre-, post-, or in situ-harvest operations and crop analyzes.

(57) **ABSTRACT**

A modeling framework for evaluating the impact of weather conditions on farming and harvest operations applies real-time, field-level weather data and forecasts of meteorological and climatological conditions together with user-provided and/or observed feedback of a present state of a harvest-related condition to agronomic models and to generate a plurality of harvest advisory outputs for precision agriculture. A harvest advisory model simulates and predicts the impacts of this weather information and user-provided and/or observed feedback in one or more physical, empirical, or artificial intelligence models of precision agriculture to analyze crops, plants, soils, and resulting agricultural commodities, and provides harvest advisory outputs to a diagnostic support tool for users to enhance farming and harvest decision-making, whether by providing pre-, post-, or in situ-harvest operations and crop analyzes.

(57) **ABSTRACT**

**20 Claims, 2 Drawing Sheets**







FIG. 1





## 1

**MODELING THE IMPACT OF  
TIME-VARYING WEATHER CONDITIONS  
ON UNIT COSTS OF POST-HARVEST CROP  
DRYING TECHNIQUES USING  
FIELD-LEVEL ANALYSIS AND FORECASTS  
OF WEATHER CONDITIONS, FACILITY  
METADATA, AND OBSERVATIONS AND USER  
INPUT OF GRAIN DRYING DATA**

FIELD OF THE INVENTION

The present invention relates to harvest operations in precision agriculture. Specifically, the present invention relates to a system and method of applying real-time, field-level weather simulation and prediction to one or more agricultural models to generate a series of harvest advisory outputs in a tool for supporting farm operations management.

BACKGROUND OF THE INVENTION

Harvest operations for a variety of agricultural commodities are substantially influenced by environmental factors, such as the weather. While some weather conditions, such as precipitation, may create an obvious deterrent to harvest operations, more benign daily weather characteristics also play subtle yet significant roles. For example, many commodities require that the harvested product be at or below a product-dependent moisture threshold before they can be stably stored at ambient temperatures (at least without taking specific steps to keep the product stable, such as the maintenance of a constant airflow through the product). On the other extreme, delaying harvest for too long can result in the crop becoming overly dry, potentially exposing seeds to damage during the threshing process, or removing permissible water weight from the product. Such an occurrence of delayed harvest may result in lower crop revenue, since payments are often based on mass. Similarly, crop temperature thresholds may be a major consideration for long-term storage of some crops, for example tuberous crops such as potatoes and sugar beets, where cold conditions are advantageous.

The harvest operation itself is also often sensitive to plant, product, and soil moisture and temperature levels. For instance, green plants, or even deceased plants with a heightened moisture level, often create difficulty for harvest operations that are based on the use of a threshing action to separate the seed (or other product) from the parent plant. This can result in both yield loss due to un-threshed seeds passing through and out of the harvester, and seed damage due to the repeated or harsh threshing action that may be required. Frozen or excessively wet soils can also inhibit harvest operations for various crops, depending upon the harvest mechanism for the particular crop. Each of the product, plant, and soil moisture and temperature therefore impact both the timing and viability of harvest operations, and all of these qualities are highly influenced by complex interactions between plant and soil characteristics and environmental conditions.

As global agricultural operations continue to grow in size, the practicality of in-situ monitoring of field conditions on a regular basis is becoming increasingly diminished. Further, the often substantial equipment and labor resources involved in harvest operations are not easily moved across significant distances in an effort to find fields with viable or more favorable harvest conditions. The ability to both diagnose and predict the viability of harvest operations in a potentially remote field is therefore of increasing importance to the management of modern farm operations. Also, production agriculture is often a capital-intensive business with very thin

## 2

relative profit margins. The ability to more effectively manage the logistics associated with deployment of a farm operation's equipment and human resources is therefore becoming increasingly critical to profitability and long-term viability of the farm itself.

In part because agricultural research globally is largely carried out by institutions with local or regional focus, agronomic models are often based on sample datasets that are limited in size and/or geographic representativeness, with less than ideal documentation of (or accommodation for) associated weather and environmental conditions. Because of this, there are very few models that can be picked up and applied to other locations and timeframes without a potentially substantial loss in model accuracy. Models for the same processes can often lead to diametrically opposed conclusions when applied at differing locations because of model shortcomings that are due to a lack of understanding of the extent to which localized influences impact the associated processes during the development of the model.

Existing solutions do not provide a sufficient framework for utilizing weather analysis and prediction to accurately diagnose field-level weather conditions for precision agriculture to overcome the challenges above. Accordingly, there is a strong need not found in the existing art for a system and method that provides an improved process for application of weather information in agronomic modeling to produce a better understanding of farm and harvest operations. There is also a need not found in the existing art for support tools designed to provide real-time assessments of weather conditions and the impact on crops, plants, soils, and resulting agricultural products to enable improvements in farm and harvest operations.

BRIEF SUMMARY OF THE INVENTION

It is therefore one objective of the present invention to provide a system and method of applying real-time assessments of weather conditions to precision agriculture models for enhancing harvest operations. It is another objective to combine real-time and location-tagged data communication in farm operations with such real-time assessments of weather conditions for generating harvest advisory information in precision agriculture. It is still another objective of the present invention to combine user-provided and/or observed feedback of a present state of a harvest-related condition with real-time assessments of weather conditions for generating harvest advisory information in precision agriculture. It is yet another objective of the present invention to provide a diagnostic support tool for enhanced decision-making in harvest operations.

There are many other specific objectives of the present invention. One such objective is to provide a system and method of predicting dry-down of a crop over time, such as a mature small grain, coarse grain, or oilseed crop, for planning the timing of harvest operations. It is another such objective of the present invention to provide a system and method of predicting the time-varying unit costs, per percent moisture per unit of mass or volume, associated with fuel-based or forced-air mechanical drying of crops resulting from changing weather conditions and the characteristics of the drying facility. Yet another objective of the present invention is to provide a system and method of predicting the time-varying unit costs, per unit of mass or volume, associated with the combined impacts of time-varying grain moisture levels of a crop to be harvested and the time-varying unit costs per percent moisture per unit of mass or volume of forced-air or fuel-based mechanical drying.



It is another objective to provide tools for assessing a risk of delayed harvest operations to achieve favorable crop moisture levels, for evaluating soil temperature and soil moisture in a given field to provide guidance as to when soil conditions are likely to inhibit harvest and post-harvest tillage operations due to soil being exceedingly wet or frozen, and for evaluating the impact of time-varying soil temperature in a given field for the timing of harvest operations as it pertains to the internal temperature of root-based crops and their long-term storage stability at those temperatures.

Other objectives of the present invention include providing systems and methods for predicting the day-by-day and intra-day windows of opportunity for harvest operations owing to time-varying threshability as a result of interactions between a standing crop and atmospheric and soil conditions, to aid in more immediate planning of harvest operations; predicting the dry-down of hay (alfalfa, etc.) and other cuttings over time for planning the timing or windows of opportunity of cutting, windrowing, silageing, or bailing operations; and assisting in decision-making for application of a desiccant to affect a desired harvest window.

The present invention is a system and method of evaluating, diagnosing, and predicting various agronomic conditions attendant to farming activities in an advisory model configured to simulate and predict the impact of weather conditions on harvest operations. The present invention applies real-time, field-level data representative of assessments of localized weather conditions, together with real-time and location-tagged communication of data of various types and content, and long-range climatological and/or meteorological forecasting, to one or more physical, empirical, or artificial intelligence models of precision agriculture to analyze crops, plants, soils, and resulting agricultural commodities, and generate a plurality of harvest advisory outputs. Such outputs are provided either directly to farmers, to third parties, or in a harvest advisory tool based on the advisory model for supporting precision agricultural operations.

Through application of the harvest advisory modeling paradigms discussed herein, the prospects for providing improved guidance relating to farm and harvest operations has the potential to be very substantial in the field of precision agriculture. For example, diagnosing and predicting the in-field dry-down of agricultural commodities, the efficiency of fuel-based and/or forced-air grain drying activities, the associations between plant moisture and atmospheric conditions, and the soil temperature and moisture profiles all provide significant value-added benefit. Recent parallel advances in weather analysis and prediction, and in the availability of mechanisms for facilitating real-time and location-tagged data communication in farm operations, have created an enticing set of possible new applications for addressing the problems described above, and others. The application of both in-situ (though not necessarily in or near a given field) and remotely-sensed weather information, in combination with scientific and computational advances in the integration of data collected by these disparate weather observing systems, permit the diagnosis of field-level weather conditions with accuracy that greatly improve upon that obtained with the deployment of a basic weather station to each and every field.

Further, advances in the understanding of the interactions between the land surface and the overlying atmosphere, combined with other improvements to the physics of meteorological weather models, and the ever-increasing computational power available to operate these models, enable a level of both short-term and long-term accuracy and locality to weather forecasts that has not been previously attainable.

The present invention also applies time- and location-tagged feedback and observations on current and recent characteristics representing measurements of various crop states, such as moisture and temperature, to the one or more physical, empirical, or artificial intelligence models of precision agriculture. Such feedback and observations provide an enhanced collection of data along with the associated weather conditions and other relevant information to build more comprehensive datasets that are used to make far-reaching improvements to associated physical models for diagnosing and predicting underlying harvest-related conditions. Artificial intelligence is also incorporated to this more comprehensive dataset to draw automatic associations between available external data and the harvest-related condition to yield further models for simulating harvest conditions. Artificial intelligence in the present invention is also retrained as more and more data are accumulated, and the results may be tested against independent data in an effort to find the most reliable model. Such a model framework implicitly yields information as to the importance of related factors through the resulting weighting systems between inputs, subcomponents within the artificial intelligence layer, and the model output(s). Together, this feedback and observations, and physical, empirical, and artificial intelligence modeling paradigms discussed herein significantly enhance the analysis of crops, plants, soils, and resulting agricultural commodities within the present invention to generate harvest advisory outputs and improved guidance for farm and harvest operations.

Other objects, embodiments, features and advantages of the present invention will become apparent from the following description of the embodiments, taken together with the accompanying drawings, which illustrate, by way of example, the principles of the invention.

#### BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWINGS

The accompanying drawings, which are incorporated in and constitute a part of this specification, illustrate several embodiments of the invention and together with the description, serve to explain the principles of the invention.

FIG. 1 is a systemic diagram of a harvest advisory model and support tool for according to the present invention; and

FIG. 2 is a block diagram of information flow and steps performed within a harvest advisory model and support tool according to the present invention.

#### DETAILED DESCRIPTION OF THE INVENTION

In the following description of the present invention reference is made to the exemplary embodiments illustrating the principles of the present invention and how it is practiced. Other embodiments will be utilized to practice the present invention and structural and functional changes will be made thereto without departing from the scope of the present invention.

FIG. 1 is a system diagram of a harvest advisory model 100 for evaluating, diagnosing, and predicting various agronomic conditions that have an impact on farm and harvest operations, according to one embodiment of the present invention. The harvest advisory model 100 is performed within one or more systems and/or methods that includes several components, each of which define distinct activities required to apply real-time, field-level data representative of assessments of localized weather conditions, together with real-time and location-tagged communication of data of various types and



content, and long-range climatological and/or meteorological forecasting to analyze crops, plants, soils, and agricultural products, to generate a plurality of harvest condition advisory outputs, and enable a harvest advisory tool **200** configured to provide diagnostic support to farm and harvest operations.

The harvest advisory model **100** ingests, or receives, input data **102** that includes weather and location specifications **110**, such as meteorological data **111**, real-time, field-level weather data **112**, observed weather data **113** that has occurred, for example, during a current growing season, and extended-range weather and climatological forecast information **114**. The weather and location specifications **110** may also include other types of real-time and location-tagged data **115**, such as for example GPS data **116**, and data captured from orbiting satellites or remotely powered vehicles that when processed provide details at a field-level resolution, such as remotely-sensed satellite imagery data **117** and remotely-captured drone imagery data **118**. The input data **102** may further include crop and plant specifications **120** such as for example the type, variety and relative maturity of the crop **121**, and other planting, chemical application, and harvest data **122**, such as for example the date that a field was planted with seed, the population of planted seeds per specified area, when a field was sprayed for example with a pesticide, desiccant, other chemical, the type and amount of chemical applied, and anticipated temporal harvest data such as an expected harvest date or harvest window. The crop and plant specifications **120** may also include the desired crop temperature **123**, crop moisture **124**, seed moisture **125**, facility storage dimensions and specifications **126**, and airflow characteristics for a stored crop **127**. Additional information may include planting information such as plant depth **128** and row width **129**.

The input data **102** may also include field and soil specifications **140**, which may further include data such as for example the soil type **141** in which a crop was planted, and attributes such as soil temperature **142**, soil moisture **143**, current and/or forecasted soil conditions **144**, overall soil profile **145**, levels of vegetative debris **146** existing at the time of planting, and nutrient levels **147** present.

The input data **102** may also include physical, empirical, or observed agricultural data **130**, such as for example sampled crop moisture content **131**. This may include data such as samples and/or observations of actual grain moisture taken from a planted field at one or more times, and the associated dry-down of the grain in a particular field over time. Other examples of such data **130** include data regarding temporal harvestability windows **132** for a particular crop at a particular location over some recent period of time, or for various crops at various locations, over time. This may include, for example, temporal windows of threshability of a seed-based crop over some recent period of time, temporal windows of cut-windrow-bail operations for crops such as hay at a particular location over some recent period of time, and temporal windows of mature growth stages, desiccant efficacy, and harvestability for a particular desiccant for particular crop at a particular location over some recent period of time.

The physical, empirical, or observed agricultural data **130** may further include descriptive metadata reflecting actual performance, realization, or implementation of one or more of crops, plants, fields, soils, and facilities. This metadata may include crop and seed metadata **133** such as the relative maturity and dry-down characteristics of the variety, pest and disease susceptibility or resistance, whether a crop is irrigated or non-irrigated, and type and origin of seeds (for example, genetically modified or non-genetically modified, and company source). Such metadata may also include soils metadata

**134** such as information relating to previous crop grown on or in the soil, tillage practice, soil composition, presence of surface and/or subsurface drainage, nutrient composition, rate of degradation, rate of water movement in both a lateral and vertical direction, presence of materials such as salts in soil strata, and facility metadata **135** such as metadata relating to type characteristics, for example whether the facility is a forced-air drying facility or a fuel-based mechanical grain drying facility, and their associated resources and costs.

All of this input data is ingested into the harvest advisory model **100** via a data ingest module **171** as shown in the system diagram of FIG. 1. The harvest advisory model **100** ingests this input data **102** and applies it to one or more precision agriculture (agronomic) models **160** which may include artificial intelligence-derived precision agriculture models, to produce output data **180**. The output data **180** of the harvest advisory model **100** is represented a harvest condition output profile **181** that is provided to a harvest advisory tool **200** that can be used to predict, simulate, and forecast harvest conditions **270** as discussed further herein.

The input data **102** is applied to the one or more precision agriculture models **160** in a plurality of data processing modules **170** within a computing environment **150** that also includes one or more processors **152** and a plurality of software and hardware components. The one or more processors **152** and plurality of software and hardware components are configured to execute program instructions to perform the functions of the harvest advisory model **100** described herein, and embodied in the one or more data processing modules **170**.

The plurality of data processing modules **170** include, in addition to the data ingest module **171**, a physical and empirical models module **172**, and an artificial intelligence module **173** configured to apply a layer of artificial intelligence to the harvest advisory model **100** to develop relationships between input data **102** and outputs from other precision agriculture models to perform and improve the assessments, diagnoses, simulations, forecasts, and predictions that form the harvest condition output profile **182**. The plurality of data processing modules **170** also includes a harvest condition prediction module **177** configured to generate the harvest condition outputs **270**.

The physical and empirical models module **172** enables application of the input data **102** to the one or more precision agriculture models **160**. Such models **160** include crop, soil, plant, and other modeling paradigms, such as for example phenological models that include general crop-specific and crop variety-specific models, a common example being growing degree day (GDD) models. These models **160** also include soil models such as the EPIC, APEX, and ICBM soil models, and land surface models such as the NOAH, Mosaic, and VIC models. Other models contemplated within the scope of the present invention include crop-specific, site-specific, and attribute-specific physical models, such as models for simulating temperature, moisture, wetness, in-field crop dry-down, plant-atmosphere vapor diffusion, plant rewetting associated with precipitation, and deterministic and stochastic grain drying models that parameterize the unique drying characteristics of a specific crop to be dried based on characteristics of the drying process and statistical models, and models for analyzing storage characteristics of crops, seeds, soils, and storage facilities. It is contemplated that the input data **102** may be applied to existing precision agriculture models, as well as customized models for specific harvest conditions **270**, for example simulation of expected dry-down of grain in the particular field, grain and plant moisture levels,



and temperature and moisture content of a root-based crop over time, and other simulations, predictions, and forecasts as noted herein.

The harvest condition output profile **182**, as noted above, is output data **180** of the harvest advisory model **100**. Once the harvest advisory model **100** has generated the harvest condition output profile **182**, the present invention applies the harvest condition output profile **182** to develop the harvest advisory tool **200** for analyzing many types of information in harvest operations. The harvest condition output profile **182** provides estimates, for example, of standing crop dry-down rates, anticipated harvest dates and suitability, fuel consumption optimizers for forced-air drying, indicators of plant 'toughness' for anticipating harvest windows, and possible loss of field workability due to the formation of frost in the soils prior to post-harvest tillage.

FIG. 2 is a block diagram of information flow and steps performed within the harvest advisory model **100** and harvest advisory tool **200** for supporting users **300**. In the present invention, the model **100** is initialized **210** by receiving various input data **102**, such as weather and location specifications **110**. Crop and plant specifications **120** may also be applied at this phase. The harvest advisory model **100** uses this information to develop predictions of expected weather conditions **230** using one or more weather modeling paradigms. Additional input data **102** related to specific harvest conditions **270** to be modeled may then be applied, such as seed, soil, field and facility data, and any type of metadata. The present invention may utilize one or more predictive data models **175** for generating predictions or forecasts **220** relating to such additional input data **102**, depending on the type of harvest condition **270**.

The expected weather conditions **230**, and additional input data **102**, are then applied in step **240** to one or more precision agriculture models **160**. Output from such models **160** may also be further applied to develop **242** artificial intelligence models for further analysis of the harvest condition **270**, as discussed further herein. The present invention performs an initial evaluation **250** of the harvest condition **270**, and may ingest input data **102** representing further observations to perform modifications **260** of the initial evaluation. The present invention then produces the harvest condition output profile **182** and provides this to the harvest advisory tool **200**, for use by one or more users **300** or for further distribution or processing in an API module **176**.

As noted above, the harvest advisory model **100** includes a layer of artificial intelligence that may be applied to develop relationships between input data **102** and outputs from others of the precision agriculture models to perform the simulations, forecasts, diagnoses, and predictions represented in the harvest condition output profile **182**. The present invention contemplates that these relationships may be developed in a variety of different ways. For example, the artificial intelligence module **173** is configured to adjust to input data **102** that includes observations provided from fields that are local or even distributed across a region. Other data applied to the artificial intelligence module **173** includes, as noted above, output from other precision agriculture models, and it is further contemplated that the original datasets for such models may be collected from institutions around the world and analyzed within the present invention. Regardless, observations may be either quantitative (e.g. grain moisture level as tested) or qualitative (e.g. the plant material became harvestable or unharvestable at a given time and location, whether or not because of environmental factors). Such information can be analyzed with respect to both recent and current weather and soil conditions, and the corresponding agronomic model

outputs, to make appropriate adjustments to the unaltered agronomic model output that would otherwise be provided to the user. This information can also be used to provide updated guidance to other users in similar circumstances (with respect to locale, crop, etc.).

The artificial intelligence module **173** uses associated weather conditions, agronomic and other agricultural model outputs, environmental considerations, crop characteristics, and other relevant data, together with empirical observations where applicable, to build a more comprehensive dataset that can be used to make far-reaching improvements to (or even replacements for) the associated physical models for diagnosing and predicting the underlying harvest-related condition. For instance, the artificial intelligence layer can be applied to an adequately-sized dataset to draw automatic associations between the available external data and the harvest-related condition, effectively yielding a 'black box' model for simulating the condition. As more and more data are accumulated, the data can be sub-sampled, the artificial intelligence layer retrained, and the results tested against independent data in an effort to find the most reliable model. Further, such a model implicitly yields information as to the importance of related factors through the resulting weighting systems between inputs, subcomponents within the artificial intelligence layer, and the model output(s). This information may be used to identify which factors are particularly important or unimportant in the associated process, and thus help to target ways of improving the model over time.

The present invention contemplates that many different types of artificial intelligence may be employed within the scope thereof, and therefore, the artificial intelligence module **173** may include one or more of such types of artificial intelligence. The artificial intelligence module **173** may apply techniques including, but not limited to, k-nearest neighbor (KNN), logistic regression, support vector machines or networks (SVM), and one or more neural networks.

The use of artificial intelligence in the harvest advisory model **100** of the present invention enhances the utility of physical and empirical models by automatically and heuristically constructing appropriate relationships, mathematical or otherwise, relative to the complex interactions between growing and maturing plants, the environment in which they reside, the underlying processes and characteristics, and the observational input data made available to the system. For example, where predictive factors known to be related to a particular outcome are known and measured along with the actual outcomes in real-world situations, artificial intelligence techniques are used to 'train' or construct a model that relates the more readily-available predictors to the ultimate outcomes, without any specific a priori knowledge as to the form of those relationships.

The present invention adopts a combined approach for simulating the relationships between predictive data and eventual outcomes. In the harvest advisory model **100**, the general nature of the relationships can be quantified with one or more physical models and/or with artificial intelligence models, which are then applied to a combination of the predictive data, historical data, metadata, and other physical model outputs to better simulate the ultimate outcomes. This approach permits the better-understood portions of the problem at hand to be modeled using a physical or empirical model, while permitting the less well understood portions of the problem to be automatically modeled based on the relationships implicit in the data provided to the system. In addition, with sufficient input data and output reliability and accuracy, the physical models may be entirely supplanted by the use of artificial intelligence model(s).



Alternatively, the artificial intelligence layer need not be employed in the system to produce the desired output information.

The proliferation of devices for communicating data both collected and required during harvests provides the opportunity to address many issues with obtaining reliable, centralized datasets for modeling harvest activities. For example, issues arise because, although key predictive factors or ground truths are routinely measured in production agriculture, they are rarely reported into a centralized repository of data that could be used to develop models that simulate the relevant relationships. Further, the mere act of reporting and collecting this data does not in itself provide the ability to develop models based on the data. In one embodiment of the present invention, past, present, and predicted weather data for any location globally are constructed through combined application of one or more of in-situ weather observations, remotely-sensed weather observations, and meteorological analysis and prediction models. These data are made available via an API-based system for obtaining weather data for any given location and timeframe. Likewise, global, national, and regional databases of soil and land-cover characteristics are also made available via an API system, making these potentially predictive parameters readily available for association with any harvest-related observations that might become available. These data, along with user-provided crop metadata, can be used to diagnose and predict the growth and maturation of a crop to be harvested, with or without the aforementioned systems and methods including whether or not the processes are performed on a mobile device. Using a field data collection device, such as a smartphone or tablet, observations of conditions impacting harvest operations can be provided in near real-time. These observations may include grain moisture samples (whether from a trial harvest operation or manual sampling), crop harvestability metrics, field accessibility metrics, or any number of other characteristics of the crop or field that may impact the crop's behaviors or its harvest operations. Additionally, through combined application of locational information available from the device, and crop or field metadata relating to the aforementioned observations that might be collected through an application accessible on the device, data on additional predictive factors can be collected and associated with each observation. When combined with automatically-collected weather, soil, and other environmental data, these data can be used as the basis for automatically building artificial intelligence models that either simulate the future expected states of the harvest-impacting conditions directly, or that act to provide corrective adjustments to the outputs of one or more physical models for simulating the underlying processes.

The systems and methods of the present invention therefore provide the opportunity to substantially advance the state of the art in terms of planning and managing harvest operations with the use of combinations of physical and empirical models with artificial intelligence. It is further notable however that many types and sources of input data **100** allow for user-based and/or locality-based biases in the observations and predictive data that are available to develop the artificial intelligence modeling paradigms contemplated herein. Some of these biases may represent differences in perception (where subjective feedback is accepted), while others may be due to biases in the instrument(s) used to collect more quantitative observations, and even others may be due to variability in factors associated with the crop or farm operation that are outside of the realm of what is being collected in terms of metadata (for instance, the design of the particular harvesting equipment used can impact both harvestability and field-

accessibility for a given operation). Because of this, the present invention contemplates development of both generalized artificial intelligence models, using all available data and metadata, and localized or user-specific artificial intelligence models tailored to analyze, for example: a particular location, user, crop, operation, soil type, and seed variety. These highly-localized or -personalized models may yield information that is more closely-tailored to the particular location or user than that provided from original data.

As noted above, the harvest advisory model **100** ingests many different types of weather information for application to the modeling approaches discussed above. This includes real-time, historical, and forecasted field-level data representative of assessments of localized weather conditions, and may include long-range climatological and/or meteorological forecasting, in one or more predictive weather models **174**.

Such real-time, field-level data representative of assessments of localized weather conditions may be produced by many different sources of meteorological data to provide one or more of observed weather data and current field-level weather data, and predicted weather data, for example as data that is complementary to the data assimilation systems and forecasting systems noted below. Such additional sources of weather data may include data from both in-situ and remotely-sensed observation platforms. For example, numerical weather prediction models (NWP) and/or surface networks may be combined with data from weather radars and satellites to reconstruct the current weather conditions on any particular area to be analyzed. There are numerous industry NWP models available, and any such models may be used as sources of meteorological data in the present invention. Examples of NWP models at least include RUC (Rapid Update Cycle), WRF (Weather Research and Forecasting Model), GFS (Global Forecast System) (as noted above), and GEM (Global Environmental Model). Meteorological data is received in real-time, and may come from several different NWP sources, such as from the European Centre for Medium-Range Weather Forecasting (ECMWF), Meteorological Services of Canada's (MSC) Canadian Meteorological Centre (CMC), as well as the National Oceanic and Atmospheric Administration's (NOAA) Environmental Modeling Center (EMC), and many others. Additionally, internally or privately-generated "mesoscale" NWP models developed from data collected from real-time feeds to global and localized observation resources may also be utilized. Such mesoscale numerical weather prediction models may be specialized in forecasting weather with more local detail than the models operated at government centers, and therefore contain smaller-scale data collections than other NWP models used. These mesoscale models are very useful in characterizing how weather conditions may vary over small distances and over small increments of time. The present invention may be configured to ingest data from all types of NWP models, regardless of whether publicly, privately, or internally provided or developed.

Long-range climatological and/or meteorological forecasting may be generated by, in one aspect of the present invention, dynamical extended range weather forecast models generated from known methods for inter-seasonal to inter-annual climate prediction, which have evolved into a combination of deterministic and statistical modeling schemes that permit the capture of long-term low-frequency features. Such forecasting often requires global analyses for the specification of observed initial and boundary conditions, the use of sophisticated numerical weather prediction models, a statis-



## 11

tical treatment of the model results, and a verification of the forecast results as a feedback into forecast refinement.

In one embodiment of the present invention, providing long-range climatological and/or meteorological forecasting may involve the use of two data assimilation systems and two forecasting systems. The two data assimilation systems may be used to provide historic and current atmospheric and land surface initial conditions and also global ocean temperatures. For example, the two forecast systems may incorporate the U.S. National Centers for Environmental Predictions (NCEP) Global Forecast System (GFS) for atmospheric predictions and the Geophysical Fluid Dynamics Laboratory Modular Ocean Model to provide sea-surface temperature predictions. Software and data supporting the above are publicly available from the NCEP.

The present invention contemplates that one or more predictive weather models **174** may be incorporated to develop expected weather conditions **230** as in FIG. 2. Such models may be executed at any desirable time interval, such as daily or hourly, to provide a complete global dataset for use in initializing the subsequent model run, and to be used to supply localized values for harvest projections for individual crop locations.

Another type of input data **102** contemplated within the scope of the present invention is image data, such as field-level processed remotely-sensed imagery in the form of remotely-sensed satellite imagery data **117** and remotely-captured drone imagery data **118**, which may be ingested to provide additional support for an assessment of crop and soil states and yield metrics in the harvest advisory model **100**. One source of image data representing this remotely-sensed imagery is satellite systems, such as fine temporal resolution low-earth orbit satellites that provide a minimum of three spectral bands. Another source is unmanned or remotely-piloted vehicles such as those commonly referred to as "drones". Other sources are also contemplated, such as for example manned aerial reconnaissance, lower temporal frequency earth resources satellite such as LANDSAT and MODIS, ground-based robots, and sensors mounted on field and farm equipment. Regardless of the source, this image information is field-navigated to provide users with the most recent high-resolution depiction of a field to be or being harvested. Image data may be delivered on a web or application-based device configured for use with the harvest advisory tool **200**, and additional tools may be provided for spatially navigating the image data and overlaying a variety of weather data elements.

Regardless of the source, the field-level remotely-sensed raw or image data may be used by the harvest advisory model **100** to map the crop field and generate a time-series profile of harvest activity. It is contemplated that remotely-sensed satellite imagery data **117** and remotely-captured drone imagery data **118** may be analyzed prior to or in concert with application to the one or more modeling paradigms discussed herein. The remotely-sensed satellite imagery data **117** and remotely-captured drone imagery data **118** may be analyzed using a normalized difference vegetative index (NDVI) that provides the user with an evaluation of plant health, biomass, nutrient content and moisture or wetness content. Other approaches may also be employed for such analysis, such as: Modified Chlorophyll Absorption Ratio Index/Optimized Soil-Adjusted Vegetation Index (OSAVI), Triangular Chlorophyll Index/OSAVI, Moderate Resolution Imaging Spectrometer Terrestrial Chlorophyll Index/Improved Soil-Adjusted Vegetation Index (MSAVI), and Red-Edge Model/MSAVI.

## 12

The harvest advisory model **100** may further include additional sources of data, such as image-based data from systems such as video cameras, and data generated from one or more vehicle-based sensing systems, including those systems coupled to computing systems configured on harvesting equipment, or those systems configured to gather weather data from mobile devices present within vehicles, such as the mobile telephony devices and tablet computers as noted above. Crowd-sourced observational data may also be provided from farmers using mobile telephony devices or tablet computers using software tools such as mobile applications, and from other sources such as social media feeds. Meteorologist input may be still a further source of data. The data collected or used need not be in image format.

Visible adaptation of the output of any of the modeling paradigms employed within the harvest advisory model **100** and harvest advisory tool **200** to user-provided feedback data helps to foster a productive and ongoing feedback loop between the user **300** and the harvest advisory model **100** over time. While adding an artificial intelligence layer to the overall modeling system leads to output that is generally responsive to user feedback on the whole, it is possible that even the output of the artificial intelligence model (or artificial intelligence layer functioning together with a physical model) may differ from real-world observations. The present invention therefore contemplates, in another embodiment, an extra layer of forced-adaptation software for adjusting output to ensure responsiveness to feedback data and enforce the belief that data being provided is actually being used to automatically improve the corresponding model over time. Since even a model based on artificial intelligence cannot be reasonably forced to return an outcome that is consistent with what is observed, even after that observation has been included in the pool of data that was used to develop the model in the first place, a forced-adaptation layer that varies according to the nature of the problem and model(s) to which it resides is applied so that information provided to the user going forward is well-matched to the information the user has provided. For example, it is entirely possible that feedback and observational data provided by two different users, for seemingly identical situations from the perspective of the data the system has available, will conflict, even to the extent of potentially being diametrically opposed. Applied to the same adaptive learning model, the present invention may produce an indeterminate outcome because of the conflicting feedback, or even a conflicting outcome to what the user has reported. The forced adaptation layer is therefore an additional module atop the artificial intelligence layer that provides an appropriate albeit brute-force adjustment to system output to ensure that it is consistent with the feedback and observational data the user has provided for the appropriate timeframe (if one exists), so as to provide the perception of responsiveness to the user and encourage the provision of further feedback and observational data.

The present invention contemplates that many applications of the harvest advisory model **100** are used to provide specific output information for supporting farming and harvesting operations. In one such application, the harvest condition output profile **182** provides a pre-harvest prediction of in-field dry-down of a mature small grain, coarse grain, or oilseed crop over time, but crop type need not be limited to this. For example, root-based crops may be considered for either in-field crop dry-down or crop moisture acquisition. This prediction is provided via the harvest advisory tool **200** and is used for planning the timing of pre-harvest, harvest, and post-harvest operations for such crops.



The harvest advisory tool **200** provides this prediction in one or more approaches. In one such approach, the harvest advisory model **100** leverages weather prediction models to predict weather conditions that impact the rate of drying or wetting of a root-based crop or the grain associated with a mature small grain, coarse grain, or oilseed crop in a particular field. A physical model is also applied to simulate the expected dry-down of the grain in the particular field, based on the combination of crop and field characteristics and the expected weather conditions. This is accomplished by receiving one or more samples of actual grain moisture from the field at one or more times, and modifying, based on the differences between simulated and sampled moisture content at the corresponding times, the prediction of the expected dry-down of the grain in the particular field over time.

In another approach, a prediction of weather conditions that impact the rate of drying or wetting of a root-based crop or the grain associated with a mature small grain, coarse grain, or oilseed crop in a particular field is formed using crop metadata, and one or more samples of actual grain moisture from one or more fields at one or more times. The weather conditions, crop metadata, and root or grain moisture samples from one or more fields and one or more crops are applied to automatically develop an in-field root or grain dry-down model based on artificial intelligence. The harvest advisory model **100** then applies the artificial intelligence-based model to diagnose or predict the in-field dry-down of grain with similar characteristics in any field at any chosen time.

In another application, the harvest condition output profile **182** provides a prediction of the time-varying unit costs (per percent moisture per unit of mass or volume) associated with fuel-based or forced-air mechanical drying of grain resulting from changing weather conditions and the characteristics of a grain drying facility. This prediction is provided via the harvest advisory tool **200** for evaluating the impact of time-varying weather conditions on the unit costs of forced-air or fuel-based drying.

In this application, the harvest advisory model **100** leverages weather prediction models to predict weather conditions that impact the energy requirements associated with at least one of forced-air and fuel-based mechanical grain drying. This is accomplished by receiving metadata (whether past, present, or future) regarding the characteristics of at least one of forced-air and fuel-based mechanical grain drying facility and their associated resources, and applying the predicted weather data and facility metadata in at least one of an empirical, physical, and artificial intelligence model to predict the time-varying cost of at least one of forced-air and fuel-based grain drying. Factors that may impact the cost of forced-air or fuel-based drying include the atmospheric temperature and moisture content, the temperature of the grain itself (often reflective of the ambient temperature at the time of harvest), the moisture of the grain, the amount of airflow with respect to a measurement of area or volume, cost of the energy used to drive the airflow, and the energy efficiency of the facility itself. In the process of drying heat is necessary to evaporate moisture from the grain and a flow of air is needed to carry away the evaporated moisture. There are two basic mechanisms involved in the drying process; the migration of moisture from the interior of an individual grain to the surface, and the evaporation of moisture from the surface to the surrounding air. The rate of drying is determined by the moisture content and the temperature of the grain and the temperature, the (relative) humidity and the velocity of the air in contact with the grain. Likewise, facility design, such as continuous-flow dryers, batch dryers, tower dryers, and the like can impact the ultimate drying costs by 50% or more.

In yet another application, the harvest condition output profile **182** provides a prediction of the time-varying unit costs (per unit of mass or volume) associated with the combined impacts of time-varying grain moisture levels of the crop to be harvested, and the time-varying unit costs (per percent moisture per unit of mass or volume) of forced-air or fuel-based mechanical drying. This prediction is provided via harvest advisory tool **200** for evaluating the costs associated with in-field vs. facility-based drying of an agricultural commodity requiring sufficiently low moisture levels for stable long-term storage.

In this application, the harvest advisory model **100** predicts the in-field dry-down of the grain associated with a crop in a particular field using one or more physically-based and/or artificial intelligence models to simulate the combined impacts of weather conditions, crop characteristics, and field characteristics on grain moisture levels. The harvest advisory model **100** then uses predicted weather data to predict metadata associated with one or more of forced-air and fuel-based mechanical grain drying facilities, and analyzes relationships that quantify the process of grain drying, the per-unit, time-varying cost of drying the grain (cost, per percentage point of moisture, per unit of grain mass or volume). Then, using the predicted in-field grain dry-down and the predicted cost of one or more of forced-air and fuel-based grain drying, the effective time-varying overall cost of forced-air and/or fuel-based grain drying (cost per unit of grain mass or volume) is predicted.

Another application of the present invention includes aiding in the assessment of the expected costs of forced-air or fuel-based mechanical drying of a crop owing to early harvest, relative to the potential costs of crop losses due to adverse weather conditions and the susceptibility of the in-field crop to weather and other causes of yield loss. In this application, the harvest condition output profile **182** provides an assessment of the trade-offs associated with delaying harvest operations to achieve a more appealing crop moisture level verses the risks associated with leaving the crop exposed in a field. The present invention performs this assessment by applying predictions of the in-field grain dry-down and the time-varying cost of at least one of forced-air and fuel-based grain drying to predict the effective overall time-varying cost of mechanical forced-air or fuel-based grain drying (cost, per unit of grain mass or volume). It then predicts, using crop characteristics and predicted weather data, the time-varying potential for crop yield losses owing to adverse weather conditions, and provides time-varying comparative data on the costs of at least one of forced-air and fuel-based drying associated with earlier harvest of the crop, versus the risks associated with yield losses associated with adverse weather conditions that may be experienced prior to harvest of the crop.

The present invention may also be configured to diagnose and predict the soil temperature and moisture profiles in a given field to provide guidance as to when soils in that field are likely to become exceedingly wet or frozen, thereby inhibiting harvest and post-harvest tillage operations. In this application, the harvest condition output profile **182** provides an assessment of the trade-offs associated with delaying harvest operations to achieve a more appealing crop moisture level verses the risks associated with leaving the crop exposed in a field. The present invention performs this assessment by applying predictions of the in-field grain dry-down and the time-varying cost of at least one of forced-air and fuel-based grain drying to predict the effective overall time-varying cost of mechanical forced-air or fuel-based grain drying (cost, per unit of grain mass or volume). In parallel it also predicts, using crop characteristics, soil data, and predicted weather



15

data, the time-varying likelihood of at least one of excess moisture or frost developing within the soil profile, and provides time-varying comparative data on the costs of forced-air or fuel-based drying associated with earlier harvest of the crop, versus the risks associated with excessive moisture or frost developing within the soil profile prior to the time when at least one of harvest operations and post-harvest tillage operations can be completed.

In another application, the harvest condition output profile **182** enables a diagnosis and prediction of the impacts of time-varying soil temperature profile in a given field as it pertains to the internal temperature of root-based crops and their long-term storage stability at those temperatures. This is provided via harvest advisory tool **200** for aiding in the timing of harvest operations so as to optimize storage conditions.

The present invention provides this diagnosis and prediction in one or more approaches. In one approach, the harvest advisory model **100** leverages at least one of weather prediction models, land surface, and soil models to predict the time varying temperature and moisture content of a root-based crop intended to be harvested. Utilizing predicted weather conditions, the present invention then predicts, using the crop temperature and moisture, and airflow characteristics of the storage pile or facility, the stability of the crop for long-term storage.

In another approach, the present invention collects data on crop temperature and moisture, storage dimensions, and airflow characteristics for a stored crop, over time. It then compares this crop storage data against at least one of time-varying weather data and outputs of models for simulating crop moisture and temperature characteristics in the corresponding times and locations, for the corresponding crops, to develop artificial intelligence models for relating these crop storage characteristics to the more readily-available weather data and outputs of models for simulating crop moisture and temperature characteristics. Using the airflow characteristics of the storage pile or facility, and the artificial intelligence model and at least one of forecast weather data and forecast outputs of models for simulating crop temperature and moisture characteristics, the present invention then provides a forecast of the likely windows of harvestability on the current and future days and the potential for crop stability for long-term storage.

In another application, the harvest condition output profile **182** enables prediction of the day-by-day windows of opportunity for harvest operations owing to time-varying threshability as a result of interactions between a standing or wind-rowed crop and atmospheric and soil conditions. The present invention provides this prediction to aid in more immediate planning of harvest operations in one or more approaches. In one approach, the harvest advisory model **100** predicts the daily window of harvest opportunity due to the impact of plant wetness on crop threshability by collecting data on temporal windows of harvestability for a particular crop at a particular location over some recent period of time, and comparing this harvestability data against at least one of time-varying weather data and outputs of models for simulating plant wetness characteristics in the crop in order to identify relationships between these external data and the reported harvestability data. The harvest advisory model **100** then provides, based on these identified relationships and the at least one of time-varying weather forecast data and forecast outputs of models for simulating plant wetness characteristics, a forecast of the likely windows of harvestability on the current and future days.

In another approach, the present invention predicts the daily window of harvest opportunity due to the impact of

16

plant wetness on crop threshability by collecting data on windows of harvestability for various crops at various locations, over time, and comparing this harvestability data against at least one of time-varying weather data and outputs of models for simulating plant wetness characteristics in the corresponding times and locations, for the corresponding crops, in order to develop artificial intelligence models for relating these windows of harvestability to the more readily-available weather data and outputs of models for simulating plant wetness characteristics. The harvest advisory model **100** then provides, using the artificial intelligence model and at least one of forecast weather data and forecast outputs of models for simulating plant wetness characteristics, a forecast of the likely windows of harvestability on the current and future days.

The present invention may also be configured for an application in which the harvest condition output profile **182** enables prediction of the dry-down of cutting crops such as hay (and others, for example alfalfa) over time. The harvest advisory tool **200** uses this output of the harvest advisory model **100** for planning the timing of, or predicting windows of opportunity for, cutting, windrowing, silageing, or bailing operations for hay fields.

The present invention performs this collecting of data on temporal windows of cut-windrow-bail during consecutive and disjoint operations for hay at a particular location over some recent period of time, and compares this operational data against at least one of time-varying weather data and outputs of models for simulating plant wetness characteristics in the hay in order to identify relationships between these external data and the reported operational data. The harvest advisory model **100** then provides, based on these identified relationships and the at least one of time-varying weather forecast data and forecast outputs of models for simulating plant wetness characteristics, a forecast of the likely windows of, for example, cut-windrow-bail operations on the current and future days.

Additionally, the present invention may also be configured for an application in which the harvest condition output profile **182** provides assistance with the decision to apply a desiccant to affect a desired harvest window. The harvest advisory tool **200** uses this output of the harvest advisory model **100** in one or more approaches. In one such approach, the present invention collects data on temporal windows of mature growth stages, and desiccant efficacy and harvestability for a particular desiccant for particular crop at a particular location over some recent period of time. It then compares crop growth stage, desiccant, and harvestability data against at least one of time-varying weather data and outputs of models for simulating plant wetness and seed moisture characteristics in the crop in order to identify relationships between these external data and the reported crop growth stage, desiccant, and harvestability data. The present invention then provides, based on these identified relationships and the at least one of time-varying weather forecast data and forecast outputs of models for simulating plant wetness and seed moisture characteristics, a forecast of the likely windows of desiccant efficacy, and subsequently the crop moisture, on the current and future days.

In another approach, the harvest advisory model **100** collects data on temporal windows of mature growth stages, desiccant efficacy, and harvestability for a particular desiccant for a particular crop at a particular location over some recent period of time, and compares crop growth stage, desiccant, and harvestability data against at least one of time-varying weather data and outputs of models for simulating plant wetness and seed moisture characteristics in the crop in



17

order to develop artificial intelligence models relating these external data and the reported crop growth stage, desiccant, and harvestability data. The harvest advisory tool **200** then provides, using the artificial intelligence model and at least one of time-varying weather forecast data and forecast out-

puts of models for simulating plant wetness and seed moisture characteristics, a forecast of the likely windows of desiccant efficacy, and subsequently the crop moisture, on the current and future days.

The execution of the harvest advisory model **100** prior to, during, or following harvest activities, addresses multiple factors that can be used by farmers to improve and enhance management of harvest operations. The harvest advisory tool **200** provides farmers with many different types of harvest-related information, such as evaluations, predictions, forecast, and diagnoses of harvest conditions **270**. As noted above, this harvest-related information may include estimates of standing crop dry-down rates, anticipated harvest dates and suitability, fuel consumption optimizers for forced-air drying, indicators of plant ‘toughness’ for anticipating harvest windows, and possible loss of field workability due to the formation of frost in the soils prior to post-harvest tillage.

One specific example of such harvest-related information include predictions of expected in-field, pre-harvest dry-down **183** of a mature small grain, coarse grain or oilseed crop over time, so that users **300** can plan the timing of a harvest operation. Another example is a prediction of time-varying unit costs associated with at least one of fuel-based or forced-air mechanical drying **184** of grain resulting from changing weather conditions and characteristics of the grain drying facility, for post-harvest crop storage. Yet another example is an assessment of expected costs of forced-air or fuel-based mechanical drying of a crop owing to early harvest relative to the potential costs of crop losses due to adverse weather conditions and the susceptibility of the in-field crop to weather and other causes of yield loss. Still another example is a diagnosis and prediction of soil temperature and moisture profiles in a given field, to provide guidance as to when soils in that field are likely to become exceedingly wet or frozen, thereby inhibiting harvest and post-harvest tillage operations. A further example is a diagnosis and prediction of the impacts of a time-varying soil temperature profile in a given field as it pertains to the internal temperature of root-based crops and their long-term storage stability at those temperatures, thereby aiding in the timing of harvest operations so as to optimize storage conditions. Still a further example is prediction of harvest opportunity windows **184**, such as the daily window of harvest opportunity due to the impact of plant wetness on crop threshability, and the timing or windows of opportunity of cutting, windrowing, silageing, or bailing operations with regard to dry-down of hay (alfalfa et al.) cutting over time. Additional examples involve a determination of whether to apply a desiccant **186** to affect a desired harvest window, and a determination of field workability loss **187**.

The harvest advisory tool **200** contemplates that output data **180** may be generated for visual representation of the information contained, for example on a graphical user interface. With the harvest advisory tool **200**, users may be able to configure settings for, and view various aspects of, the harvest advisory model **100** using a display on such graphical user interfaces, and/or via web-based or application-based modules. Tools and pull-down menus on such a display (or in web-based or application-based modules) may also be provided to customize the type and nature of the input data **102** applied to the various modeling paradigms of the harvest advisory model **100**, as well as the output data **180** provided

18

in the harvest condition output profile **182**. Examples of this include the tactile and/or haptic notifications of changes in the harvest conditions discussed herein. Other types of notifications may include those provided via applications resident on mobile telephony, tablet, or wearable devices are also possible, such as for example a voice-based output generated to verbally notify farmers of possible disease or pest risk.

As noted above, the output data **180** of the harvest advisory model **100** may be used to generate a plurality of advisory services **190** in one or more application programming interface (API) modules **176**. These advisory services **190** provide enhanced decision-making support to harvest operations in precision agricultural production.

These advisory services **190** are driven by the generation of the harvest condition output profile **182** that results in the harvest action reporting and harvest summary reporting aspects of the output data **180**. Each of these advisory services **190** enriches the utilization and application of the augmented crop growth model **100**.

One such advisory service **190** is a current and historical weather reporting service **192**. Site-specific weather information is an important element of field recordkeeping in agricultural production. Using the field geo-positional information provided for the harvest advisory modeling, current and historical weather for the field location may be provided as an advisory service for specific geographical locations.

A soil modeling service **194** is another advisory service **190** contemplated within the scope of the present invention. One API module **176** applies the harvest condition output profile **182** and output data **180** to a sophisticated soil model to generate information that provides a better understanding of current and future soil conditions relative to historical soil conditions at harvest times. Such a soil model supports the use of existing soil properties e.g., organic matter, soil type, etc., tillage practices, presence of tile drainage, and irrigation history, along with the advanced short-, medium-, or long-range weather forecasting. This coupling of modeled soil characteristics with advanced weather forecasting information at harvest times provides a valuable tool for indicating crop potential. The output of an API module **176** generating this soil modeling service **194** may be further configured to provide a detailed past-through-future analysis of soil characteristics impacting harvest conditions, such as soil temperature and moisture assessments, including freeze and thaw information, the amount of water ‘throughput’, runoff or ponding of soils, and an important interpretation of workability of the soils.

Another advisory service **190** is soil conditions forecasting service **196**. The soil modeling service **194** can be used with prediction of expected weather conditions to provide an estimate of future soil conditions at harvest times in this advisory service **190**. Using the weather and location specifications **110**, a forecast of soil temperature and moisture conditions for the specified field location can be generated. Such a forecast in this service **196** enables anticipation of periods of suitable soil conditions for field operations, including workability, particularly those after periods of rainfall.

Another advisory service **190** as an output **180** of the harvest advisory model **100** is a harvest alerts service **198**. The importance of information provided to production agriculture in the present invention dictates a necessity for an effective and direct method of conveying information. The harvest alerts service **198** may utilize a ‘push’ technology for immediate and direct dissemination of information provided as output data **180** from the harvest advisory model **100**. The API module **176** that generates this service may be configured to so that as the time for conducting a harvest operation nears,



users may receive alerts of important combinations of weather conditions, risk factors, and decision-support aides for the purposes of managing and timing harvest operations. This is supported by user-specified elements including alert parameters, location(s), lead time(s), time(s) of day, and the particular device(s) to receive the alerts. For example, a custom harvesting operation may integrate their work orders with such an alerting system in order to optimize the logistics of their operation by avoiding costly weather-related delays to harvest operations, through either spatial or temporal reassignment of equipment and related resources.

As noted above, the present invention may in one embodiment is configured to diagnosis and prediction of the impacts of a time-varying soil temperature profile in a given field as it pertains to the internal temperature of root-based crops and their long-term storage stability at those temperatures. Such a model aids in the timing of harvest operations so as to optimize crop storage conditions.

Such an embodiment is performed in a system and method of evaluating a timing of harvest operations and crop storage conditions, comprising in which a profile of long-term storability of an agricultural commodity is developed with a harvest advisory model. The harvest advisory model in this embodiment predicts expected weather conditions relative to a timing of harvest operations and crop storage conditions by applying weather information including recent and current field-level weather data and extended-range weather forecast data, and field-specific information including location data representing a field where an agricultural commodity is planted to one or more predictive numerical weather models. The harvest advisory model forecasts harvestability windows over a specified period of time and crop stability for long-term storage, by simulating the expected weather conditions, crop-specific information for an agricultural commodity to be harvested, and the field-specific information in an agricultural model of one or more physical and empirical characteristics of crop moisture and crop temperature for harvestability of crops at corresponding times and locations. The agricultural model comprises at least one of a crop-specific growing degree day model, a soil model, a land surface model. Also, an artificial intelligence model may be developed to simulate crop storage characteristics from the expected weather conditions, one or more sampled observations from a planted field, and crop storage facility data by building a comprehensive dataset for the agricultural model of one or more physical and empirical characteristics of crop moisture and crop temperature for harvestability of crops at corresponding times and locations.

The systems and methods of the present invention may be implemented in many different computing environments 150. For example, they may be implemented in conjunction with a special purpose computer, a programmed microprocessor or microcontroller and peripheral integrated circuit element(s), an ASIC or other integrated circuit, a digital signal processor, electronic or logic circuitry such as discrete element circuit, a programmable logic device or gate array such as a PLD, PLA, FPGA, PAL, and any comparable means. In general, any means of implementing the methodology illustrated herein can be used to implement the various aspects of the present invention. Exemplary hardware that can be used for the present invention includes computers, handheld devices, telephones (e.g., cellular, Internet enabled, digital, analog, hybrids, and others), and other such hardware. Some of these devices include processors (e.g., a single or multiple microprocessors), memory, nonvolatile storage, input devices, and output devices. Furthermore, alternative software implementations including, but not limited to, distributed processing,

parallel processing, or virtual machine processing can also be configured to perform the methods described herein.

The systems and methods of the present invention may also be partially implemented in software that can be stored on a storage medium, executed on programmed general-purpose computer with the cooperation of a controller and memory, a special purpose computer, a microprocessor, or the like. In these instances, the systems and methods of this invention can be implemented as a program embedded on a mobile device or personal computer through such mediums as an applet, JAVA™ or CGI script, as a resource residing on one or more servers or computer workstations, as a routine embedded in a dedicated measurement system, system component, or the like. The system can also be implemented by physically incorporating the system and/or method into a software and/or hardware system.

Additionally, the data processing functions disclosed herein may be performed by one or more program instructions stored in or executed by such memory, and further may be performed by one or more modules configured to carry out those program instructions. Modules are intended to refer to any known or later developed hardware, software, firmware, artificial intelligence, fuzzy logic, expert system or combination of hardware and software that is capable of performing the data processing functionality described herein.

The foregoing descriptions of embodiments of the present invention have been presented for the purposes of illustration and description. It is not intended to be exhaustive or to limit the invention to the precise forms disclosed. Accordingly, many alterations, modifications and variations are possible in light of the above teachings, may be made by those having ordinary skill in the art without departing from the spirit and scope of the invention. It is therefore intended that the scope of the invention be limited not by this detailed description. For example, notwithstanding the fact that the elements of a claim are set forth below in a certain combination, it must be expressly understood that the invention includes other combinations of fewer, more or different elements, which are disclosed in above even when not initially claimed in such combinations.

The words used in this specification to describe the invention and its various embodiments are to be understood not only in the sense of their commonly defined meanings, but to include by special definition in this specification structure, material or acts beyond the scope of the commonly defined meanings. Thus if an element can be understood in the context of this specification as including more than one meaning, then its use in a claim must be understood as being generic to all possible meanings supported by the specification and by the word itself.

The definitions of the words or elements of the following claims are, therefore, defined in this specification to include not only the combination of elements which are literally set forth, but all equivalent structure, material or acts for performing substantially the same function in substantially the same way to obtain substantially the same result. In this sense it is therefore contemplated that an equivalent substitution of two or more elements may be made for any one of the elements in the claims below or that a single element may be substituted for two or more elements in a claim. Although elements may be described above as acting in certain combinations and even initially claimed as such, it is to be expressly understood that one or more elements from a claimed combination can in some cases be excised from the combination and that the claimed combination may be directed to a sub-combination or variation of a sub-combination.



## 21

Insubstantial changes from the claimed subject matter as viewed by a person with ordinary skill in the art, now known or later devised, are expressly contemplated as being equivalently within the scope of the claims. Therefore, obvious substitutions now or later known to one with ordinary skill in the art are defined to be within the scope of the defined elements.

The claims are thus to be understood to include what is specifically illustrated and described above, what is conceptually equivalent, what can be obviously substituted and also what essentially incorporates the essential idea of the invention.

The invention claimed is:

**1. A method comprising:**

ingesting, as input data, weather information and crop-specific information for a grain crop to be harvested, the weather information including recent and current field-level weather data and extended-range forecast weather data, and the crop-specific information including at least one of crop type data and harvest data;

modeling the input data in a plurality of data processing modules within a computing environment in which the plurality of data processing modules are executed in conjunction with at least one processor, the data processing modules configured to profile the impacts of time-varying weather conditions and characteristics of a grain drying facility, using at least one of fuel-based drying or forced-air mechanical drying of harvested grain, on unit costs of drying of harvested grain, by

1) predicting expected weather conditions relative to the energy requirements associated with migration of moisture from an interior of an individual grain to a surface, and evaporation of moisture from the surface to surrounding air associated with a grain drying facility, using at least one of fuel-based drying or forced-air mechanical drying of harvested grain,

2) applying the expected weather conditions, the crop-specific information, and facility metadata representing actual and/or realized performance characteristics of a grain drying facility using at least one of fuel-based drying or forced-air mechanical drying of harvested grain to at least one of an agricultural model of one or more physical and empirical characteristics impacting post-harvest storage of the harvested grain crop, and an artificial intelligence model of drying harvested grain, and

3) simulating time-varying unit costs per percentage of moisture per unit of mass or volume associated with at least one of fuel-based drying or forced-air mechanical drying; and

generating, as output data, one or more advisories representing a prediction of the time-varying unit costs per percentage of moisture per unit of mass or volume associated with at least one of fuel-based drying or forced-air mechanical drying in a harvest output condition profile.

**2. The method of claim 1, further comprising applying the harvest output condition profile to a harvest advisory tool configured to provide the one or more advisories representing the predicted time-varying unit costs to a user.**

**3. The method of claim 1, wherein the crop-specific information further includes one or more of crop relative maturity data, crop planting depth and row spacing data, crop post-maturity dry-down characteristics, targeted crop moisture or temperature thresholds, facility storage dimensions data, crop storage airflow data, and fuel-based grain dryer facility characteristics.**

## 22

**4. The method of claim 1, wherein the modeling further comprises applying the weather information to one or more predictive numerical weather models to generate the prediction of expected weather conditions.**

**5. The method of claim 1, wherein the agricultural model of one or more physical and empirical characteristics impacting post-harvest storage of the harvested grain crop includes at least one of deterministic and stochastic grain drying models based on at least one of characteristics of the drying process and statistical models that parameterize the unique drying characteristics of a specific crop to be dried.**

**6. The method of claim 1, wherein the applying the expected weather conditions, the crop-specific information, and facility metadata representing actual and/or realized performance characteristics of a grain drying facility using at least one of fuel-based drying or forced-air mechanical drying of harvested grain to an artificial intelligence model of drying harvested grain further comprises automatically developing a specific unit cost of grain drying model by building a comprehensive dataset of one or more physical and empirical characteristics impacting drying operations of the grain crop.**

**7. The method of claim 6, further comprising applying additional data representing sampled observations at or near the specification location at corresponding times in at least one of the agricultural model of one or more physical and empirical characteristics impacting post-harvest storage of the harvested grain crop and the artificial intelligence model of drying harvested grain, wherein the sampled observations include actual drying costs at a specific facility over time for one or more of different crops, different moisture levels, and different weather conditions.**

**8. The method of claim 6, further comprising:**

applying the expected weather conditions and the crop-specific information to an additional agricultural model of one or more physical and empirical characteristics impacting harvest operations of a grains crop to simulate an expected rate of drying of the grains crop in a particular field;

identifying differences in the simulation of expected rate of drying with one or more sampled observations at or near the specific location at corresponding; and

modifying the simulation based on the differences between the otherwise simulated rate of drying and the one or more sampled observations at or near the specification at the corresponding times.

**9. A system comprising:**

a computing environment including at least one computer-readable storage medium having program instructions stored therein and a computer processor operable to execute the program instructions to profile impacts of time-varying weather conditions and characteristics of a grain drying facility using at least one of fuel-based drying or forced-air mechanical drying of harvested grain on unit costs of drying of harvested grain by performing a harvest advisory model within a plurality of data processing modules, the plurality of data processing modules including:

a weather modeling module configured to diagnose and predict expected weather conditions relative to energy requirements associated with migration of moisture from an interior of an individual grain to a surface, and evaporation of moisture from the surface to surrounding air associated with a grain drying facility using at least one of fuel-based drying or forced-air mechanical drying of harvested grain by applying weather information including recent and current field-level weather data and



23

extended-range weather forecast data to one or more predictive numerical weather models;  
 one or more modules configured to 1) model the expected weather conditions, the crop-specific information, and facility metadata representing actual and/or realized performance characteristics of a grain drying facility using at least one of fuel-based drying or forced-air mechanical drying of harvested grain in an agricultural model of one or more physical and empirical characteristics impacting post-harvest storage of the harvested grain crop, 2) automatically develop an artificial intelligence, crop-specific information, and facility metadata by building a comprehensive dataset for the agricultural model of one or more physical and empirical characteristics impacting drying of the harvested grain crop, and 3) evaluate time-varying unit costs per percentage of moisture per unit of mass or volume associated with at least one of fuel-based drying or forced-air mechanical drying; and  
 a harvest condition prediction module configured to predict the time-varying unit costs per percentage of moisture per unit of mass or volume associated with at least one of fuel-based drying or forced-air mechanical drying and generate one or more advisories representing the prediction in a harvest output condition profile.

10. The system of claim 9, wherein the harvest output condition profile is applied to a diagnostic support tool configured to provide the one or more advisories representing the predicted time-varying unit costs to a user.

11. The system of claim 9, wherein the crop-specific information further includes one or more of crop relative maturity data, crop planting depth and row spacing data, crop post-maturity dry-down characteristics, targeted crop moisture or temperature thresholds, facility storage dimensions data, crop storage airflow data, and fuel-based grain dryer facility characteristics.

12. The system of claim 9, wherein the agricultural model of one or more physical and empirical characteristics impacting post-harvest storage of the harvested grain crop includes at least one of deterministic and stochastic grain drying models based on at least one of characteristics of the drying process and statistical models that parameterize the unique drying characteristics of a specific crop to be dried.

13. The system of claim 9, wherein the one or modules are further configured to model additional data representing sampled observations at or near the specification location at corresponding times in at least one of the agricultural model of one or more physical and empirical characteristics impacting post-harvest storage of the harvested grain crop and the artificial intelligence model of drying harvested grain, the sampled observations including actual drying costs at a specific facility over time for one or more of different crops, different moisture levels, and different weather conditions.

14. The system of claim 9, wherein the one or more modules are further configured to:

apply the expected weather conditions and the crop-specific information to an additional agricultural model of one or more physical and empirical characteristics impacting harvest operations of a grains crop to simulate an expected rate of drying of the grains crop in a particular field;

identify differences in the simulation of expected rate of drying with one or more sampled observations at or near the specific location at corresponding; and

24

modify the simulation based on the differences between the otherwise simulated rate of drying and the one or more sampled observations at or near the specification at the corresponding times.

15. A method of evaluating an impact of time-varying weather conditions on unit costs of drying a harvested grain crop, comprising:

within a computing environment comprised of a computer processor and at least one computer-readable storage medium operably coupled to the computer processor and having program instructions stored therein, the computer processor being operable to execute the program instructions to profile impact of time-varying weather conditions and characteristics of a grain drying facility using at least one of fuel-based drying or forced-air mechanical drying of harvested grain on unit costs of drying of harvested grain in a harvest advisory model configured to perform the steps of:

predicting expected weather conditions relative to energy requirements associated with migration of moisture from an interior of an individual grain to a surface, and evaporation of moisture from the surface to surrounding air associated with a grain drying facility using at least one of fuel-based drying or forced-air mechanical drying of harvested grain by applying weather information comprised of recent and current field-level weather data and extended-range weather forecast data to one or more predictive numerical weather models;

analyzing the expected weather conditions, the crop-specific information, and facility metadata representing actual and/or realized performance characteristics of a grain drying facility using at least one of fuel-based drying or forced-air mechanical drying of harvested grain in an agricultural model of one or more physical and empirical characteristics impacting post-harvest storage of the harvested grain crop, the agricultural model comprising at least one of deterministic and stochastic grain drying models based on at least one of characteristics of the drying process and statistical models that parameterize the unique drying characteristics of a specific crop to be dried;

analyzing the expected weather conditions, the crop-specific information, and the facility metadata to automatically develop an artificial intelligence model of drying harvested grain and their associated resources by building a comprehensive dataset of one or more physical and empirical characteristics impacting drying of the harvested grain crop; and

predicting time-varying unit costs per percentage of moisture per unit of mass or volume associated with at least one of fuel-based drying or forced-air mechanical drying.

16. The method of claim 15, further comprising generating one or more advisories of the predicted time-varying unit costs per percentage of moisture per unit of mass or volume associated with at least one of fuel-based drying or forced-air mechanical drying in a harvest output condition profile.

17. The method of claim 15, further comprising applying the harvest output condition profile to a diagnostic support tool configured to provide the one or more advisories representing the predicted time-varying unit costs to users performing harvest operations.

18. The method of claim 15, wherein the crop-specific information further includes one or more of crop relative maturity data, crop planting depth and row spacing data, crop post-maturity dry-down characteristics, targeted crop mois-



ture or temperature thresholds, facility storage dimensions data, crop storage airflow data, and fuel-based grain dryer facility characteristics.

19. The method of claim 15, further comprising analyzing additional data representing sampled observations at or near 5 the specification location at corresponding times in at least one of the agricultural model of one or more physical and empirical characteristics impacting post-harvest storage of the harvested grain crop and the artificial intelligence model of drying harvested grain, wherein the sampled observations 10 include actual drying costs at a specific facility over time for one or more of different crops, different moisture levels, and different weather conditions.

20. The method of claim 15, further comprising:  
applying the expected weather conditions and the crop- 15 specific information to an additional agricultural model of one or more physical and empirical characteristics impacting harvest operations of a grains crop to simulate an expected rate of drying of the grains crop in a particular field; and 20  
identifying differences in the simulation of expected rate of drying with one or more sampled observations at or near the specific location at corresponding to modify the simulation based on the differences between the other- 25 wise simulated rate of drying and the one or more sampled observations at or near the specification at the corresponding times.

\* \* \* \* \*