



US009234233B2

(12) **United States Patent**  
**Rose et al.**

(10) **Patent No.:** **US 9,234,233 B2**  
(45) **Date of Patent:** **Jan. 12, 2016**

(54) **METHODS FOR ENHANCING NUCLEIC ACID HYBRIDIZATION**

(71) Applicant: **Integrated DNA Technologies, Inc.**, Skokie, IL (US)

(72) Inventors: **Scott Rose**, Coralville, IA (US); **Mark A. Behlke**, Coralville, IA (US); **Richard Owczarzy**, Coralville, IA (US); **Joseph A. Walder**, Chicago, IL (US); **Derek M. Thomas**, Cedar Rapids, IA (US); **Michael R. Marvin**, Coralville, IA (US)

(73) Assignee: **Integrated DNA Technologies, Inc.**, Skokie, IL (US)

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **14/578,399**

(22) Filed: **Dec. 20, 2014**

(65) **Prior Publication Data**

US 2015/0141635 A1 May 21, 2015

**Related U.S. Application Data**

(62) Division of application No. 13/073,866, filed on Mar. 28, 2011, now Pat. No. 8,916,345.

(60) Provisional application No. 61/318,043, filed on Mar. 26, 2010.

(51) **Int. Cl.**  
**C07H 21/02** (2006.01)  
**C07H 19/04** (2006.01)

(Continued)

(52) **U.S. Cl.**  
CPC ..... **C12Q 1/6832** (2013.01); **C07H 21/04** (2013.01); **C12Q 1/6818** (2013.01); **G01N 33/5308** (2013.01); **G01N 33/542** (2013.01); **C12Q 1/6851** (2013.01)

(58) **Field of Classification Search**  
CPC ..... C12Q 1/6832; C07H 21/04  
USPC ..... 536/23.1, 26.6; 435/6.1  
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

2,264,303 A 12/1941 Dickey  
3,218,309 A 11/1965 Elslager et al.

(Continued)

FOREIGN PATENT DOCUMENTS

DE 2546535 4/1977  
EP 0070685 1/1983

(Continued)

OTHER PUBLICATIONS

Shi et al., "Design of Phosphoramidite Monomer for Optimal Incorporation of Functional Intercalator to Main Chain of Oligonucleotide," *Bioconjugate Chemistry*, vol. 16, No. 2, Mar. 1, 2005, pp. 306-311.


(Continued)

*Primary Examiner* — Jezia Riley

(74) *Attorney, Agent, or Firm* — Michael Best & Friedrich LLP





(57) **ABSTRACT**

A composition comprising an oligonucleotide having the structure 5'-Y<sub>1</sub>-L<sub>1</sub>-X-L<sub>2</sub>-Y<sub>2</sub>-3'. Y<sub>1</sub> comprises a sequence of one or more DNA or RNA nucleotides, including a first nucleotide N<sub>1</sub> having a 3' phosphate covalently linked to L<sub>1</sub>. Y<sub>2</sub> comprises a sequence of one or more DNA or RNA nucleotides, including a second nucleotide N<sub>2</sub> having a 5' phosphate covalently linked to L<sub>2</sub>. L<sub>1</sub> and L<sub>2</sub> each independently are a direct bond or a C<sub>1</sub>-C<sub>7</sub> alkyl, alkynyl, alkenyl, heteroalkyl, substituted alkyl, aryl, heteroaryl, substituted aryl, cycloalkyl, alkylaryl, or alkoxy group. X is



R<sub>1</sub> is a hydrogen or a C<sub>1</sub>-C<sub>8</sub> alkyl. M is a label or ligand comprising a fused polycyclic aromatic moiety.

**9 Claims, 23 Drawing Sheets**



(51)	<b>Int. Cl.</b>		6,531,589 B1	3/2003	Iyer et al.
	<i>C12Q 1/68</i>	(2006.01)	6,531,591 B1	3/2003	Fensholdt
	<i>C07H 21/04</i>	(2006.01)	6,653,473 B2	11/2003	Reed et al.
	<i>G01N 33/53</i>	(2006.01)	6,699,975 B2	3/2004	Reed et al.
	<i>G01N 33/542</i>	(2006.01)	6,727,356 B1	4/2004	Reed et al.
			6,790,945 B2	9/2004	Reed et al.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,407,189	A	10/1968	Merian
3,970,617	A	7/1976	Bruno
4,358,535	A	11/1982	Falkow et al.
4,439,356	A	3/1984	Khanna et al.
4,683,194	A	7/1987	Saiki et al.
4,683,202	A	7/1987	Mullis
4,711,955	A	12/1987	Ward et al.
4,820,630	A	4/1989	Taub
4,835,263	A	5/1989	Nguyen et al.
4,868,103	A	9/1989	Stavrianopoulos et al.
4,868,105	A	9/1989	Urdea et al.
4,889,818	A	12/1989	Gelfand et al.
4,914,210	A	4/1990	Levenson et al.
4,954,630	A	9/1990	Klein et al.
4,996,143	A	2/1991	Heller et al.
5,011,769	A	4/1991	Duck et al.
5,108,892	A	4/1992	Burke et al.
5,124,246	A	6/1992	Urdea et al.
5,188,934	A	2/1993	Menchen et al.
5,210,015	A	5/1993	Gelfand et al.
5,214,136	A	5/1993	Lin et al.
5,272,259	A	12/1993	Claussen et al.
5,304,645	A	4/1994	Klein et al.
5,312,728	A	5/1994	Lizardi et al.
5,326,679	A	7/1994	Yanagisawa et al.
5,328,824	A	7/1994	Ward et al.
5,451,463	A	9/1995	Nelson et al.
5,455,157	A	10/1995	Hinzpeter et al.
5,484,908	A	1/1996	Froehler
5,487,972	A	1/1996	Gelfand et al.
5,492,806	A	2/1996	Drmanac et al.
5,494,810	A	2/1996	Barany et al.
5,502,177	A	3/1996	Matteucci et al.
5,512,667	A	4/1996	Reed et al.
5,514,785	A	5/1996	Van Ness et al.
5,525,464	A	6/1996	Drmanac et al.
5,538,848	A	7/1996	Livak et al.
5,556,752	A	9/1996	Lockhart et al.
5,696,251	A	12/1997	Arnold, Jr. et al.
5,801,155	A	9/1998	Kutyavin et al.
5,804,375	A	9/1998	Gelfand et al.
5,824,796	A	10/1998	Petrie et al.
5,830,653	A	11/1998	Froehler et al.
5,846,726	A	12/1998	Nadeau et al.
5,849,482	A	12/1998	Meyer, Jr. et al.
5,876,930	A	3/1999	Livak et al.
5,925,517	A	7/1999	Tyagi et al.
5,942,610	A	8/1999	Nelson et al.
5,952,202	A	9/1999	Aoyagi et al.
6,007,992	A	12/1999	Lin et al.
6,028,183	A	2/2000	Lin et al.
6,084,102	A	7/2000	Kutyavin et al.
6,114,518	A	9/2000	Pitner et al.
6,117,973	A	9/2000	Batz et al.
6,117,986	A	9/2000	Nardone et al.
6,214,979	B1	4/2001	Gelfand et al.
6,312,894	B1	11/2001	Hedgpeth et al.
6,316,610	B2	11/2001	Lee et al.
6,323,337	B1	11/2001	Singer et al.
6,399,392	B1	6/2002	Haugland et al.
6,416,953	B1	7/2002	Heller
6,441,159	B1	8/2002	Lukhtanov et al.
6,448,407	B1	9/2002	Lee et al.
6,451,535	B1	9/2002	Jenne et al.
6,465,175	B2	10/2002	Horn et al.
6,465,644	B1	10/2002	Yan et al.
6,485,901	B1	11/2002	Gildea et al.
6,531,581	B1	3/2003	Nardone et al.

6,800,728	B2	10/2004	Schwartz
6,825,331	B2	11/2004	Manoharan et al.
6,875,858	B1	4/2005	DeFrancq et al.
7,060,809	B2	6/2006	Wengel et al.
7,122,383	B2	10/2006	Jones et al.
7,173,125	B2	2/2007	Schwartz et al.
7,439,341	B2	10/2008	Laikhter et al.
7,476,735	B2	1/2009	Laikhter et al.
7,605,243	B2	10/2009	Laikhter et al.
7,645,872	B2	1/2010	Laikhter et al.
7,803,536	B2	9/2010	Behlke et al.
7,803,936	B2	9/2010	Laikhter et al.
8,916,345	B2	12/2014	Rose
2002/0034754	A1	3/2002	Reed et al.
2002/0137070	A1	9/2002	Elghanian et al.
2003/0082547	A1	5/2003	Ewing et al.
2003/0096254	A1	5/2003	Reed et al.
2003/0144499	A1	7/2003	McGall et al.
2004/0180343	A1	9/2004	Weber
2005/0142598	A1	6/2005	Heindl et al.
2007/0218490	A1	9/2007	Laikhter et al.
2008/0085837	A1	4/2008	Coull et al.
2009/0053821	A1	2/2009	Laikhter et al.
2009/0259030	A1	10/2009	Cook et al.
2010/0298554	A1	11/2010	Laikhter et al.
2011/0060150	A1	3/2011	Laikhter et al.

FOREIGN PATENT DOCUMENTS

EP	0070686	1/1983
EP	0070687	1/1983
EP	0152886	8/1985
EP	0185494	6/1986
EP	0246864	11/1987
EP	0272007	11/1987
EP	0320308	6/1989
EP	0357011	3/1990
EP	0439182	7/1991
GB	1394368	5/1975
GB	1533121	11/1978
JP	52-88681	7/1977
JP	52-91031	8/1977
WO	89/09284	10/1989
WO	89/10979	11/1989
WO	90/14353	11/1990
WO	91/05060	4/1991
WO	92/10588	6/1992
WO	96/17957	6/1996
WO	96/28460	9/1996
WO	96/34983	11/1996
WO	96/40662	12/1996
WO	97/29154	8/1997
WO	97/39008	10/1997
WO	99/14226	3/1999
WO	99/37717	7/1999
WO	99/40226	8/1999
WO	99/51621	10/1999
WO	99/51775	10/1999
WO	99/64431	12/1999
WO	00/06771	2/2000
WO	00/70685	11/2000
WO	01/04129	1/2001
WO	01/42505	6/2001
WO	01/86001	11/2001
WO	03/019145	3/2003
WO	2004/026804	4/2004
WO	2004/113562	12/2004
WO	2005/049849	6/2005
WO	2006/081035	8/2006
WO	2006/127507	4/2007

(56)

## References Cited

## FOREIGN PATENT DOCUMENTS

WO	2010/147673	12/2010
WO	2011/120049	9/2011

## OTHER PUBLICATIONS

- Australian Patent Examination Report No. 2 for Application No. 2011230496 dated Jun. 22, 2015 (3 pages).
- 7th International Symposium, held 2001, pp. 1-14, 7th International Symposium & Exhibition Solid Phase Synthesis & Combinatorial Chemical Libraries, Sep. 18-22, 2001—University of Southampton, England, UK.
- Agrawal, S. et al., "Efficient methods for attaching non-radioactive labels to the 5' ends of synthetic oligodeoxyribonucleotides," *Nuc. Acids Res.* (1986) 14:6227-6245.
- Austermann, S. et al., "Inhibition of human immunodeficiency virus type 1 reverse transcriptase by 3'-blocked oligonucleotide primers," *Biochem. Pharm.* (1992) 43(12):2581-2589.
- Bailly, C. et al., "DNA recognition by intercalator-minor-groove binder hybrid molecules," *Bioconjugate Chem.* (1991) 2(6):379-393.
- Bollum, F.J., "Oligodeoxyribonucleotide-primed reactions catalyzed by calf thymus polymerase," *J. Bio. Chem.* (1962) 237(6):1945-1949.
- Boturnyn, D. et al., "Synthesis of Fluorescent probes for the Detection of Abasic Sites in DNA," *Tetrahedron* (1997) 53(15):5485-5492.
- Bustin, S.A., "Quantification of mRNA using real-time reverse transcription PCR (RT-PCR): trends and problems," *Mol. Endocrin.* (2002) 29:23-29.
- Cardullo, R.A et al., "Detection of nucleic acid hybridization by nonradiative fluorescence resonance energy transfer," *Proc. Natl. Acad. Sci. USA* (1988) 85:8790-8794.
- Caruthers, M.H. et al., "Chemical synthesis of deoxyoligonucleotides and deoxyoligonucleotide analogs," *Methods Enzymol.* (1992) 211:3-20.
- Conference Proceedings, retrieved 2010, two pages, Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries: Peptides, Proteins and Nucleic Acids—Small Molecule Organic Chemistry Diversity, Collected Papers, International Symposium, 7th, Southampton, United Kingdom, Sep. 18-22, 2001, Roger Epton, Editor.
- Dey, S. and Sheppard, T.L., "Ketone-DNA: a versatile postsynthetic DNA decoration platform," *Org. Lett.* (2001) 3(25):3983-3986.
- Dirks, R.W. et al., "Advances in fluorescent tracking of nucleic acids in living cells," *Biotechniques* (2006) 40:489-496.
- Dogan, Z. et al., "5'-tethered stilbene derivatives as fidelity- and affinity-enhancing modulators of DNA duplex stability," *J. Am. Chem. Soc.* (2004) 126:4762-4763.
- Doktycz, M.J. et al., "Studies of DNA dumbbells. I. Melting curves of 17 DNA dumbbells with different duplex stem sequences linked by T4 endloops: evaluation of the nearest-neighbor stacking interactions in DNA," *Biopolymers* (1992) 32:849-864.
- Easy Q&Q PCR, [http://www.eurogentec.be/code/en/product\\_easy\\_qq.htm](http://www.eurogentec.be/code/en/product_easy_qq.htm), visited May 9, 2002, 4 pages.
- Encyclopedia Britannica (2008) "Nucleic Acid," Encyclopedia Britannica Online (Jul. 11, 2008) <http://www.search.eb.com/eb/article-256731>.
- Gelfand, D.H., "Taq DNA Polymerase," *PCR Technology Principles and Applications for DNA Amplification*, Stockton Press, NY, Ehrlich ed. (1989) Ch. 2, 17-22.
- Genbank Accession No. AF298116, Rose et al., "*Mus musculus* bHLH protein Ptf1-p48 gene," (2001) 6 pages.
- Gibson, V. et al., "Molecular modelling of anthraquinone-oligodeoxynucleotide conjugates," *Pharm. Sci.* (1996) 2:545-548.
- Gorelick, M.V. et al., "Effect of benzannulation on the coloration of p-aminoazo compounds," *Zhurnal Organicheskoi Khimii* (1980) 16(9):1927-1933—Abstract only—Accession No. 1981:193676.
- Heesemann, J., "Studies on monolayers. I. Surface tension and absorption spectroscopic measurements of monolayers of surface-active azo and stilbene dyes," *J. Am. Chem. Soc.* (1980) 102(7):2167-2176.
- Heller, M.J. et al., "Chemiluminescent and fluorescent probes for DNA hybridization systems," *Rapid Detection and Identification of Infectious Agents* (1985) Academic Press, Inc., Orlando, Kingsbury et al. eds. 245-256.
- Ho, M.S. et al., "Azo polymers for reversible optical storage. 7. The effect of the size of the photochromic groups," *Macromolecules* (1995) 28(18):6124-6127 (Abstract only—Accession No. 1995:746851).
- Hugentobler, M. et al., "Compounds with a metal-arene  $\sigma$ -bond. Part 2. Cyclometalation of arylazo compounds. Part 2. Regioselectivity of the cyclopalladation of some substituted 1-aryazonaphthalenes," *Helvetica Chimica Acta* (1982) 65(4):1202-1211 (Abstract only—Accession No. 1982:616430).
- Ito, T. et al., "Reductive Electron Injection into Duplex DNA by Aromatic Amines," *J. Am. Chem. Soc.* (2004) 126:15552-15559.
- Iyer, R.P. et al., "3H-1,2-benzodithiole-3-one 1,1-dioxide as an improved sulfurizing reagent in the solid-phase synthesis of oligodeoxyribonucleoside phosphorothioates," *J. Am. Chem. Soc.* (1990) 112:1253-1254.
- Jenkins, J.H., Senior Librarian at the United States Patent and Trademark Office, email dated Mar. 18, 2010 and previously related email correspondence, pp. 1-2.
- Jiao, G.-S. et al., "Oligonucleotides with strongly fluorescent groups  $\Pi$ -conjugated to a nucleobase: syntheses, melting temperatures, and conformation," *Bioorg. Med. Chem. Lett.* (2003) 13(16):2785-2788.
- Ju, J. et al., "Fluorescence energy transfer dye-labeled primers for DNA sequencing and analysis," *Proc. Natl. Acad. Sci. USA* (1995) 92:4347-4351.
- Kerzhner, B.K. et al., "Photoisomerization of aromatic azo compounds adsorbed on a hydroxylated surface," *Zhurnal Obschei Khimii* (1983) 53(10):2303-2306 (Abstract only—Accession No. 1984:50840).
- Kuball, H-G. et al., "Helical twisting power of chiral mono- and bis-aminoanthraquinones," *J. Mater. Chem.* (1995) 5 (12):2167-2174.
- Lawyer, F.C. et al., "Isolation, characterization, and expression in *Escherichia coli* of the DNA polymerase gene from thermus aquaticus," *J. Biol. Chem.* (1989) 264(11):6427-6437.
- Lehman, I.R. et al., "Persistence of deoxyribonucleic acid polymerase I and its 5'  $\rightarrow$ 3' exonuclease activity in PoIA mutants of *Escherichia coli* K12," *J. Biol. Chem.* (1973) 248(22):7717-7723.
- Lewis, F.D. et al., "Hybrid oligonucleotides containing stilbene units. Excimer fluorescence and photodimerization," *J. Am. Chem. Soc.* (1995) 117:8785-8792.
- Mackay, I.M., "Real-time PCR in the microbiology laboratory," *Clin. Microbiol. Infect.* (2004) 10:190-212.
- Marshall, "Rules for the visible absorption spectra of halogenated fluorescein dyes," *Histochemical J.* (1975) 7:299-303.
- Matthews, J.A. et al., "Analytical strategies for the use of DNA probes," *Analy. Biochem.* (1988) 169:1-25.
- May, J.P. et al., "A new dark quencher for use in genetic analysis," *Chem. Commun.* (2003) 970-971.
- May, J.P. et al., "A novel dark quencher for oligonucleotide probes: synthesis and applications," poster presentation from TIDES 2002 IBC Oligonucleotide and Peptide Technology Conference, May 6-8, 2002, Las Vegas, Nevada (2 pages).
- May, J.P. et al., "Synthesis of a novel dark quencher for use with long wavelength dyes in oligonucleotide probes," *Innovation and Perspectives in Solid Phase Synthesis & Combinatorial Libraries: Peptides, Proteins and Nucleic Acids Small Molecule Organic Chemical Diversity, Collected Papers, 7th International Symposium, Southampton, United Kingdom, Sep. 18-22, 2002, 231-233.*
- Mineno, J. et al., "Fluorescent labeling of a DNA sequencing primer," *DNA Sequence—J. DNA Sequence and Mapping* (1993) 4:135-141.
- Misiura, K. et al., "Biotinyl and phosphotyrosinyl phosphoramidite derivatives useful in the incorporation of multiple reporter groups on synthetic oligonucleotides," *Nucleic Acids Research* (1990) 18(15):4345-4354.
- Miyashita, T. et al., "Novel dinucleoside phosphotriester unit conjugated with an intercalative moiety in a stereospecific manner enhances thermal stability of an alternate-stranded triple helix," *Tetrahedron Letters* (2003) 44:7399-7402.

(56)

## References Cited

## OTHER PUBLICATIONS

- Mori, K. et al., "Oligodeoxynucleotide analogs with 5'-linked anthraquinone," *FEBS Letters* (1989) 249(2):213-218.
- Morier-Teissier, E. et al., "Free radical production and DNA cleavage by copper chelating peptide-anthraquinones," *Anti-Cancer Drug Design* (1990) 5:291-305.
- Morrison, L.E. et al., "Solution-phase detection of polynucleotides using interacting fluorescent labels and competitive hybridization," *Anal. Biochem.* (1989) 183:231-244.
- Mullis, K. et al., "Specific Enzymatic Amplification of DNA In Vitro: The Polymerase Chain Reaction," *Cold Spring Harbor Symp. Quant. Biol.* (1986) 51:263-273.
- Narayanan, S. et al., "Clamping down on weak terminal base pairs: oligonucleotides with molecular caps as fidelity-enhancing elements at the 5'- and 3'-terminal residues," *Nucl. Acids. Res.* (2004) 32:2901-2911.
- Noble, S.A. et al., "Methylphosphonates as probes of protein-nucleic acid interactions," *Nuc. Acids Res.* (1984) 12(7):3387-3404.
- Owczarzy, R. et al., "Predicting stability of DNA duplexes in solutions containing magnesium and monovalent cations," *Biochem.* (2008) 47:5336-5353.
- Owczarzy, R. et al., "Predicting sequence-dependent melting stability of short duplex DNA oligomers," *Biopolymers* (1997) 44:217-239.
- Owczarzy, R., "Melting temperatures of nucleic acids: discrepancies in analysis," *Biophys. Chem.* (2005) 117:207-215.
- Patra, A. et al., "High fidelity base pairing at the 3'-terminus," *J. Am. Chem. Soc.* (2009) 131(35):12671-12681.
- Petersheim, M. et al., "Base-stacking and base-pairing contributions to helix stability: thermodynamics of double-helix formation with CCGG, CCGGp, CCGGAp, ACCGGp, CCGGUp, and ACCGGUp," *Biochem.* (1983) 22:256-263.
- PR Newswire, Announcing IBC's TIDES 2002: Oligonucleotide and Peptide Technology Conferences May 6-8, 2002, Las Vegas, NV, Copyright 2002, PR Newswire Association LLC, pp. 1-2, website (retrieved Mar. 18, 2010): [http://www.thefreelibrary.com/\\_/print/PrintArticle.aspx?id=82059345](http://www.thefreelibrary.com/_/print/PrintArticle.aspx?id=82059345).
- Proudnikov, D. et al., "Chemical methods of DNA and RNA fluorescent labeling," *Nucleic Acids Res.* (1996) 24(22):4535-4542.
- Puskas, L.G. et al., "Diamino-Antraquinone: A New Intercalating Agent. Synthesis and Linking to Oligodeoxynucleotide," *Nucleosides & Nucleotides* (1995) 14(3-5):967-968.
- Rose, S.D. et al., "The Role of PTF1-P48 in pancreatic acinar gene expression," *J. Biol. Chem.* (2001) 276 (47):44018-44026.
- Rychlik, W. et al., "Optimization of the annealing temperature for DNA amplification in vitro," *Nucl. Acids. Res.* (1990) 18:6409-6412.
- Santalucia, J, Jr., "A unified view of polymer, dumbbell, and oligonucleotide DNA nearest-neighbor thermodynamics," *Proc. Natl. Acad. Sci. USA* (1998) 95:1460-1465.
- Schuster, G.B., "Long-range charge transfer in DNA: transient structural distortions control the distance dependence," *Acc. Chem. Res.* (2000) 33:253-260.
- Setlow, P. et al., "Deoxyribonucleic acid polymerase: two distinct enzymes in one polypeptide," *J. Biol. Chem.* (1972) 247(1):224-231.
- Sijm, D.T.H.M. et al., "Aqueous solubility, octanol solubility, and octanol/water partition coefficient of nine hydrophobic dyes," *Envir. Toxic. Chem.* (1999) 18(6):1109-1117.
- Telser, J. et al., "Synthesis and characterization of DNA oligomers and duplexes containing covalently attached molecular labels: comparison of biotin, fluorescein, and pyrene labels by thermodynamic and optical spectroscopic measurements," *J. Am. Chem. Soc.* (1989) 111:6966-6967.
- Tu, C-P.D. et al., "3'-end labeling of DNA with [ $\alpha$ -<sup>32</sup>P]cordycepin-5'-triphosphate," *Gene* (1980) 10:177-183.
- Warshaw, M.M. et al., "Optical properties of sixteen dinucleoside phosphates," *J. Mol. Biol.* (1966) 20:29-38.
- Wetmur, J.G., "DNA probes: applications of the principles of nucleic acid hybridization," *Critical Review in Biochem. and Mol. Biol.* (1991) 26(3/4):227-259.
- Schnipper et al., "Synthesis and electrochemical properties of TTF modified oligodeoxynucleotides," *Chemical Communications*, No. 37, Jan. 1, 2009, p. 5552.
- Ansanuma et al., "Enantioselective Incorporation of Azobenzenes into Oligodeoxyribonucleotide for Effective Photoregulation of Duplex Formation\*\*," *Angew. Chem. Int. Ed.*, vol. 40, No. 14, Jan. 1, 2001, pp. 2671-2673.
- "Prokhorenko et al., "Incorporation of a pyrene nucleoside analogue into synthetic oligodeoxynucleotides using a nucleoside-like synthon", *Bioorganic & Medicinal Chemistry Letters*, Pergamon, vol. 5, No. 18, Sep. 21, 1995, pp. 2081-2084."
- Ossipov et al., "Dipyrido[3,2-a:2', 3'-c]phenazine-Tethered Oligo-DNA: Synthesis and Thermal Stability of Their DNA-DNA and DNA-RNA Duplexes and DNA-DNA-DNA Triplexes", *Helvetica Chimica Acta*, vol. 82, Jan. 1, 1999, pp. 2186-2200.
- Liang et al., "An interstrand-wedged duplex composed of alternating DNA base pairs and covalently attached intercalators," *Journal of Material Chemistry*, vol. 20, No. 3, Jan. 1, 2009, p. 575.
- Wang et al *Chinese j. of Chem.*, vol. 27, No. 8, pp. 1582-1588 (2009).
- Zielske, A.G., "(Tosyloxy)anthraquinones: Versatile Synthons for the Preparation of Various Aminoanthraquinones," *J. Org. Chem.* (1987) 52:1305-1309.
- International Search Report and Written Opinion for Application No. PCT/US2011/30215 dated May 23, 2011 (9 pages).
- International Preliminary Report on Patentability for Application No. PCT/US2011/30215 dated Oct. 2, 2012 (8 pages).
- European Patent Office Search Report for Application No. 11760380 dated Aug. 20, 2013 (10 pages).
- United States Patent Office Action for U.S. Appl. No. 13/073,866 dated Jun. 17, 2013 (6 pages).
- European Patent Office Action for Application No. 11760380.3 dated Apr. 14, 2014 (5 pages).
- Australian Patent Examination Report No. 1 for Application No. 2011230496 dated Jun. 16, 2014 (5 pages).
- Australian Patent Examination Report No. 1 for Application No. 2013202227 dated May 28, 2014 (3 pages).
- United States Patent Office Notice of Allowance for U.S. Appl. No. 13/073,866 dated Aug. 15, 2014 (5 pages).
- European Patent Office Action for Application No. 11760380.3 dated Feb. 18, 2015 (4 pages).
- Japanese Patent Office Action for Application No. 2013-502702 dated Mar. 2, 2015 (6 pages, English translation included).
- US 6,255,052, 05/2001, Batz et al. (withdrawn)




FIG. 1A






FIG. 1A (cont'd)



IB 1.1.1:



FQ:

FIG. 1B






FIG. 1B (cont'd)



Epoch Eclipse:



Black Hole 2:




FIG. 1C




FIG. 2A




FIG. 2B




FIG. 3A




FIG. 3B




FIG. 4A




FIG. 4B




FIG. 5A




FIG. 5B




FIG. 6A




FIG. 6B




FIG. 7A




FIG. 7B




FIG. 8A




FIG. 8B






FIG. 9A




FIG. 9B




FIG. 10A




FIG. 10B




FIG. 11A




FIG. 11B




FIG. 12




FIG. 13A




FIG. 13B




FIG. 13C




FIG. 13D






FIG. 13E




FIG. 14A




FIG. 14B

## METHODS FOR ENHANCING NUCLEIC ACID HYBRIDIZATION

### CROSS-REFERENCE TO RELATED APPLICATIONS

This patent application is a divisional of U.S. patent application Ser. No. 13/073,866 filed Mar. 28, 2011, which claims priority to U.S. Provisional Patent Application No. 61/318,043 filed Mar. 26, 2010. These applications are incorporated herein by reference in their entirety.

### SEQUENCE LISTING

The sequence listing is filed with the application in electronic format only and is incorporated by reference herein. The sequence listing text file "ASFILED\_Sequence\_US00" was created on Mar. 25, 2011, and is 49,585 bytes in size.

### FIELD OF THE DISCLOSURE

This disclosure pertains to novel oligonucleotide compounds with improved hybridization properties. Methods and reagents are provided which allow internal labeling of oligonucleotides by insertion of labels between adjacent residues without destabilizing the duplex. Because the nucleotide bases are not modified, such labeling groups can be introduced into any sequence. In some embodiments, such modifications increase duplex stability. In some embodiments, the labeling group is a fluorescence quencher. The disclosure further relates to the design of fluorescently labeled oligonucleotide probes with multiple quenching dyes capable of very efficient fluorescence quenching over a broad spectral range.

### BACKGROUND

Fluorescent energy transfer probes are an important tool in genetic analysis. These probes, also known as dual-labeled probes (DLPs) or self-quenching probes, are generally comprised of a fluorescent donor (a fluorophore) and a quencher linked to an oligonucleotide. This basic design, wherein a signal change is detected once the probe hybridizes to its intended target, is used in a variety of biological applications.

One method for detecting hybridization using fluorophores and quenchers is to link fluorescent donors and quenchers to a single oligonucleotide such that there is a detectable difference in fluorescence when the oligonucleotide is unhybridized as compared to when it is hybridized to its complementary sequence. In so-called molecular beacons, a partially self-complementary oligonucleotide is designed to form a hairpin and is labeled with a fluorescent donor at one end of the molecule and a quencher at the other end (U.S. Pat. No. 5,925,517). Intramolecular annealing to form the hairpin brings the donor and quencher into close proximity for fluorescent quenching to occur. Intermolecular annealing of such an oligonucleotide to a target sequence disrupts the hairpin, which increases the distance between the donor and quencher and results in a detectable increase in the fluorescent signal of the donor.

Oligonucleotides are not required to form a hairpin structure for this method to work efficiently. The fluorophore and quencher can be placed on an oligonucleotide such that when it is unhybridized and in a random coil conformation, the quencher is able to quench fluorescence from the fluorophore (U.S. Pat. No. 5,538,848). Once the oligonucleotide hybridizes to a complementary nucleotide sequence it becomes

more extended and the distance between the fluorophore and quencher is increased, resulting in reduced quenching and increased fluorescence.

Oligonucleotides labeled in a similar manner can also be used to monitor the kinetics of PCR amplification. In one version of this method, commonly known as a 5'-nuclease cleavage or hydrolysis assay, an oligonucleotide probe is designed to hybridize to the target sequence on the 3' side ("downstream") of one of the amplification primers. During PCR, the 5'-3' exonuclease activity of the DNA polymerase digests the 5' end of the probe thereby separating the fluorophore from the quencher. The fluorescence intensity of the sample increases as an increasing number of probe molecules are digested during the course of amplification (U.S. Pat. No. 5,210,015).

DLPs find use in other molecular/cellular biology and diagnostic assays, such as in end-point PCR, in situ hybridizations, in vivo DNA and RNA species detection, single nucleotide polymorphism (SNPs) analysis, enzyme assays, and in vivo and in vitro whole cell assays (see Dirks and Tanke, *Biotechniques* 2006, 40:489-486; Bustin, *Journal of Molecular Endocrinology* 2002, 29:23-39; Mackay, *Clin Microbiol Infect*, 2004, 10:190-212).

In one mechanism of fluorescence quenching termed ground state quenching, the fluorophore and the quencher associate to form a ground state complex which is not fluorescent. For ground state quenching to occur there need not be spectral overlap between the fluorophore and the quencher.

The most common mechanism of fluorescent quenching is fluorescence resonance energy transfer (FRET). In FRET, energy transfer occurs through space by dipolar coupling between the fluorophore and quencher and requires that there be overlap between the emission spectrum of the fluorescent donor and the absorbance spectrum of the quencher. This requirement complicates the design of probes that utilize FRET because quenchers are limited in their effective wavelength range. For example, the quencher known as BHQ-1, which absorbs light in the wavelength range of about 500-550 nm, quenches fluorescent light emitted by fluorescein, which fluoresces maximally at about 520 nm, but is of limited utility for Texas Red (emission maximum=615) or Cy5 (emission maximum=670). In contrast, the quencher BHQ-3, which absorbs light in the wavelength range of about 650-700 nm is almost completely ineffective at quenching fluorescein but is effective at quenching Cy5. In general, the number of quenchers that are known to be capable of quenching the fluorescence of any given fluorophore is limited.

Although fluorescent dyes themselves can be employed to quench fluorescence from other dyes, preferred quenchers will not fluoresce (or minimally fluoresce) so that background fluorescence is minimized. These quenchers are commonly referred to as dark quenchers. Dark quenchers allow for an increased signal to noise ratio in assays that employ DLPs, resulting in increased sensitivity. In addition, the lack of secondary fluorescence facilitates the use of additional fluorophores in multiplexed assay formats which utilize multiple distinct probes each containing a different fluorophore. If a quencher emitted light in a certain region, then additional probes could not bear fluorophores that emit light in that same region.

A number of factors are considered in designing a self-quenching probe. These include the ease of synthesis, the compatibility of the fluorophore and quencher, duplex stability, and the specificity of the probe in hybridizing to the intended target.

Duplex stability between complementary nucleic acid molecules is frequently expressed as the "melting tempera-

ture",  $T_m$ , of the duplex. Roughly speaking, the  $T_m$  indicates the temperature at which a duplex nucleic acid dissociates into two single strands. Nucleic acid hybridization is generally performed at a temperature slightly below the  $T_m$ , so that hybridization between a probe or primer and its target nucleic acid is optimized, while minimizing non-specific hybridization of the probe or primer to other, non-target nucleic acids. Duplex stability and  $T_m$  are also important in applications, such as PCR, where thermocycling may be involved. During such thermocycling melting steps, it is important that the sample temperature be raised sufficiently above the  $T_m$  so that duplexes of the target nucleic acid and its complement are dissociated. In subsequent steps of reannealing, however, the temperature must be brought sufficiently below the  $T_m$  that duplexes of the target nucleic acid and primer are able to form, while still remaining high enough to avoid non-specific hybridization events. For a general discussion, see Rychlik et al., *Nucleic Acids Research* 1990, 18:6409-6412.

Shorter oligonucleotides can help increase the specificity of a primer or probe, allowing for the discrimination of even a single mismatch between the probe and a potential complementary target. The shorter the oligonucleotide, the greater the effect of a single-base mismatch on duplex stability. However, the disadvantage of using such short oligonucleotides is that they hybridize weakly, even to a perfectly complementary sequence, and thus must be used at lower temperatures, which are unfavorable for reactions that use thermal stable enzymes, such as PCR. Certain modified nucleosides such as locked nucleic acids (LNAs) (U.S. Pat. No. 7,060,809) and C5-propynyl pyrimidines (U.S. Pat. No. 5,484,908) can be incorporated into oligonucleotides to increase duplex stability. Many nucleoside analogs, however, especially those having bulkier substituents attached to the base, are destabilizing. For example, fluorescein-dT can destabilize a duplex by up to 4° C. (*Bioorganic & Medicinal Chemistry Letters*, 13:2785-2788 2003).

Modified nucleosides employed to increase  $T_m$  are typically placed internally within an oligonucleotide sequence replacing a natural base. In contrast, non-nucleoside substituents when introduced internally within an oligonucleotide, either as a replacement for a base or as an insertion between bases, generally interfere with hybridization. For example, insertion of an abasic fluorescein group into an oligonucleotide has been observed to destabilize a duplex by 2-4° C. (*DNA Seq.* 4:135-141, 1993).

There are several classes of compounds that are known to increase binding affinity between complementary nucleic acid strands. One class is major groove binders, which includes proteins or ligands that bind to the major groove (the wider groove around a DNA helix). A second class, minor groove binders (MGBs), include non-covalently bound and covalently bound compounds. Because the minor groove of a helix is narrower in A-T rich regions, some noncovalently bound MGBs recognize the shape of the helix and preferably bind to specific sequence regions. For example, netropsin and distamycin preferably bind to A-T regions (see Bailly and Henichart, *ACS*, vol. 2, 379-393 (1991)). Covalently bound MGBs (U.S. Pat. No. 6,084,102) are typically linked to the 5' or 3' end of oligonucleotides (U.S. Pat. App. 2009/0259030) and are known to increase binding affinity and allow for shorter length probes.

A third class, intercalators, are generally flat polycyclic compounds, examples being acridine or lipoic acid derivatives (see U.S. Pat. No. 4,835,263). Intercalating compounds stabilize a duplex by fitting in between the bases of the nucleic acid monomers. They can be covalently or noncovalently

bound. Some minor groove compounds, such as 4',6-diamidino-2-phenylindole (DAPI), also intercalate.

Another group of compounds, capping reagents, are terminally attached compounds that favor Watson-Crick duplexes by stacking on the terminal base pair (Dogan, Z. et al., *J. Am. Chem. Soc.* 2004, 126, 4762-4763). Such groups include stilbene derivatives (Wu, T, et al., *J. Am. Chem. Soc.* 1995, 117, 8785-8792) and pyrenylmethylpyrrolindol (Narayanan, S. et al., *Nucleic Acids Res.* 2004, 32, 2901-2911).

The efficiency of quenching through FRET is extremely sensitive to the distance between the fluorophore and quencher ( $R_{FQ}$ ), varying with the reciprocal of  $R_{FQ}$  to the sixth power. Maximally efficient quenching minimizes background fluorescence and improves the sensitivity of the 5'-nuclease assay and other hybridization assays in which DLPs are used. Generally, for ease of synthesis and to avoid disruption of hybridization of the probe to the target sequence, the dye and quencher are attached to the ends of the oligonucleotide. For the 5'-nuclease assay, the most common configuration is to attach the dye at the 5'-end of the oligonucleotide and the quencher at the 3'-end. DLPs used in the 5'-nuclease assay are typically 25 to 30 bases in length. Even with the use of  $T_m$  enhancing modifications, such as LNA bases or a minor groove binder, probe length is still usually 14 to 18 bases. Any method that permits placement of the fluorophore and quencher in closer proximity within a probe without destabilizing the duplex formed between the probe and its target nucleic acid will improve quencher efficiency and enhance the performance of the probe.

#### BRIEF SUMMARY

This disclosure provides various compositions comprising oligonucleotides having modifying compounds placed internally within the oligonucleotide sequence between nucleotides. Even though these modifying compounds are inserted between adjacent nucleotides (as opposed to being substituted for one of the nucleotides), some of the modified oligonucleotides surprisingly form equally stable, or even more stable, duplexes with their complementary oligonucleotide sequences as compared to the stability of the duplex formed between the unmodified oligonucleotide and the complementary oligonucleotide. The modifying groups may include a variety of labels, including but not limited to fluorescence quenchers that enable the design of DLPs with very high quenching efficiency. Because the labeling group is not a modified base, the same modifying compound can be inserted into any position within any oligonucleotide sequence. This disclosure also provides methods for using and making the oligonucleotide compositions.


The compositions of the present disclosure each comprise an oligonucleotide having the general structure 5'-Y<sub>1</sub>-L<sub>1</sub>-X-L<sub>2</sub>-Y<sub>2</sub>-3', where:

Y<sub>1</sub> comprises a sequence of one or more DNA and/or RNA nucleotides, including a first nucleotide N<sub>1</sub> having a 3' phosphate covalently linked to L<sub>1</sub>;

Y<sub>2</sub> comprises a sequence of one or more DNA and/or RNA nucleotides, including a second nucleotide N<sub>2</sub> having a 5' phosphate covalently linked to L<sub>2</sub>;

L<sub>1</sub> and L<sub>2</sub> each independently are a direct bond or a C<sub>1</sub>-C<sub>7</sub> alkyl, alkynyl, alkenyl, heteroalkyl, substituted alkyl, aryl, heteroaryl, substituted aryl, cycloalkyl, alkylaryl, or alkoxy group;

5



X is or;

R<sub>1</sub> is a hydrogen or a C<sub>1</sub>-C<sub>8</sub> alkyl; and

M is a label.


These oligonucleotides may include any desired number of nucleotides, but preferably include 10-50 nucleotides, and even more preferably include 15-35 nucleotides. In some embodiments, L<sub>1</sub> and L<sub>2</sub> each may be a C<sub>1</sub>-C<sub>7</sub> alkyl, and preferably a C<sub>2</sub> alkyl. The 3' phosphate that is covalently linked to L<sub>1</sub>, and the 5' phosphate that is covalently linked to L<sub>2</sub>, each independently may be a phosphodiester, a phosphothioate, a phosphodithioate, a methyl phosphonate, a phosphoramidate, a phosphoramidite or a phosphotriester.

In some embodiments, Y<sub>1</sub> comprises a sequence of four or more DNA and/or RNA nucleotides, Y<sub>2</sub> comprises a sequence of four or more DNA and/or RNA nucleotides, M is a first quencher, and the oligonucleotide is adapted to hybridize to a second oligonucleotide having the structure 3'-Y<sub>3</sub>-Y<sub>4</sub>-5', where Y<sub>3</sub> comprises a sequence of four or more DNA and/or RNA nucleotides, including a third nucleotide N<sub>3</sub>, Y<sub>4</sub> comprises a sequence of four or more DNA and/or RNA nucleotides, including a fourth nucleotide N<sub>4</sub> that is directly attached to nucleotide N<sub>3</sub>. If the first oligonucleotide hybridizes to the second oligonucleotide, then N<sub>1</sub> base pairs with N<sub>3</sub> and N<sub>2</sub> base pairs with N<sub>4</sub> to form a duplex having a T<sub>m</sub> that is greater than the T<sub>m</sub> of a duplex formed between the second oligonucleotide and a third oligonucleotide having the structure 5'-Y<sub>1</sub>-Y<sub>2</sub>-3'. In such embodiments, the first oligonucleotide may be labeled with a fluorophore. For example, the fluorophore may be attached to the last nucleotide on the 5' end of the oligonucleotide, and in preferred embodiments, Y<sub>1</sub> may comprise a sequence of 8-12 DNA or RNA nucleotides. Compositions comprising the first oligonucleotide also may comprise the second oligonucleotide.

In some embodiments, Y<sub>1</sub> comprises a sequence of 8-12 DNA and/or RNA nucleotides, such as a sequence of 10 DNA and/or RNA nucleotides, where the nucleotide on the 5' end of Y<sub>1</sub> is labeled with a fluorophore, and M is a quencher.

In some embodiments, M comprises a fused polycyclic aromatic moiety.

In some embodiments, M is:




where L<sub>3</sub> is a direct bond or a C<sub>1</sub>-C<sub>8</sub> alkyl, alkenyl, alkenyl, heteroalkyl, substituted alkyl, cycloalkyl, or alkoxy, where R<sub>2</sub>-R<sub>6</sub> each independently are a hydrogen, an alkyl, an alkenyl, a heteroalkyl, a substituted alkyl, an aryl, a heteroaryl, a substituted aryl, a cycloalkyl, an alkylaryl, an alkoxy, an electron withdrawing group, or an electron donating group, and where one of R<sub>2</sub>-R<sub>6</sub> is —N=N—P, and where P is a fused polycyclic aromatic moiety. Electron withdrawing groups may be selected from the group consisting of —NO<sub>2</sub>,

6


—SO<sub>3</sub><sup>-</sup>, —SO<sub>2</sub><sup>-</sup>, —CN, —NCS, a ketone, an alkoxy, an ether, a carboxylic acid and a sulfonyl. Electron donating group is selected from the group consisting of an alkoxy, a heteroalkoxy, an alkyl, a cycloalkyl, a heteroalkyl, an amino, an alkylamino, or an arylamino.

In some embodiments, M is —L<sub>4</sub>-P, where L<sub>4</sub> is an alkyl, an alkynyl, an alkenyl, a heteroalkyl, a substituted alkyl, or an alkoxy group, and P is a fused polycyclic aromatic moiety. For example, L<sub>4</sub> may be —CH<sub>2</sub>-O—CH<sub>2</sub>-CH<sub>2</sub>-NH— or any other suitable linker.


Various of the oligonucleotides described herein include fused polycyclic aromatic moieties P. In some embodiments, P is



where R<sub>7</sub>-R<sub>9</sub> each independently are a hydrogen, an alkoxy, an alkyl, an alkylamino, an arylamino, a cycloalkyl, a heteroalkoxy, a heteroalkyl, or an amino, and R<sub>10</sub>-R<sub>13</sub> each independently are a hydrogen, a nitro, a cyano, a carboxylate, a sulfonyl, a sulfamoyl, an alkenyl, an alkynyl, an amino, an aryl, a heteroaryl, a biaryl, a bialkenyl, a bialkynyl, an alkoxy-carbonyl or a carbamoyl. In preferred embodiments, R<sub>9</sub> is



In some embodiments, P is



where R<sub>14</sub>-R<sub>19</sub> each independently are a hydrogen, an alkyl, a heteroalkyl, an aryl, a heteroaryl, an electron withdrawing group, or a five or six membered ring structure formed from the R<sub>1</sub>, R<sub>2</sub> pair, the R<sub>3</sub>, R<sub>4</sub> pair, the R<sub>4</sub>, R<sub>5</sub> pair, or the R<sub>5</sub>, R<sub>6</sub> pair.

Some of the oligonucleotides disclosed herein include a fluorophore, which may include, but is not limited to, 6-carboxyfluorescein (FAM), 2'7'-dimethoxy-4'5'-dichloro-6-carboxyfluorescein (JOE), tetrachlorofluorescein (TET), 6-carboxyrhodamine (R6G), N,N,N,N'-tetramethyl-6-carboxyrhodamine (TAMRA), 6-carboxy-X-rhodamine (ROX); 1-dimethylaminonaphthyl-5-sulfonate, 1-anilino-8-naphthalene sulfonate, 2-p-toluidinyl-6-naphthalene sul-

fonate, 5-(2'-aminoethyl)aminonaphthalene-1-sulfonic acid (EDANS), a coumarin dye, an acridine dye, indodicarbocyanine 3 (Cy3), indodicarbocyanine 5 (Cy5), indodicarbocyanine 5.5 (Cy5.5), 3-1-carboxy-pentyl)-3'-ethyl-5,5'-dimethylloxacarbo-cyanine (CyA); 1H,5H,11H,15H-Xantheno[2,3,4-ij:5,6,7-i'j']diquinolizin-18-ium, 9-[2(or 4)-[[[6-[2,5-dioxo-1-pyrrolidinyl)oxy]-6-oxohexyl]amino]sulfonyl]-4(or 2)-sulfo-phenyl]-2,3,6,7,12,13,16,17-octahydro-inner salt (TR or Texas Red), a BODIPY™ dye, benzoxazole, stilbene and pyrene. In some embodiments, the fluorophore may be attached to the 5' end, such as to the phosphate at the 5' end of the oligonucleotide.

Some of the oligonucleotides disclosed herein include more than one quencher, such as a first internal quencher (described above) and a second quencher. The second quencher may include, but is not limited to dabcy1, Eclipse® quencher, BHQ1, BHQ2 and BHQ3, Iowa Black® FQ, Iowa Black® RQ-n1 or Iowa Black® RQ-n2.

This disclosure also provides methods for using and making the oligonucleotide compositions. Methods for use may include methods for detecting target nucleic acids within a sample. For example, such methods may include contacting the sample with an oligonucleotide adapted to hybridize to the target nucleic acid, where the oligonucleotide includes an internal quencher and a fluorophore, and where the fluorescence of the fluorophore is reduced by fluorescence resonance energy transfer to the quencher or by ground state quenching by the quencher when the oligonucleotide is not hybridized to the second oligonucleotide, and detecting an increase in fluorescence indicating the presence of the second oligonucleotide in the sample.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A, 1B and 1C are structures of modified oligonucleotides tested for their effects on the stability of the duplex formed between the modified oligonucleotide and its complementary oligonucleotide. The modified oligonucleotides contain various modification compounds inserted between, and attached to the 3' and 5' phosphates of adjacent nucleotides.

FIG. 2A is an amplification plot that illustrates the relative fluorescence intensity ( $R_n$ ) of an HPRT q-PCR assay of the substitution data set for the FQ quencher located 6, 8, 10 and 12 positions from the 5'-fluorophore. FIG. 2B is an amplification plot that illustrates the baseline adjusted fluorescence of results in FIG. 2A ( $\Delta R_n$ ).

FIG. 3A is an amplification plot that illustrates the relative fluorescence intensity ( $R_n$ ) of the HPRT q-PCR assay of the insertion data set for FQ quencher located 6, 8, 10 and 12 positions from the 5'-fluorophore. FIG. 3B is an amplification plot that illustrates baseline adjusted fluorescence of the data in FIG. 3A ( $\Delta R_n$ ).

FIG. 4A is an amplification plot that shows the results of the baseline adjusted substitution analogs s10FQ and s6FQ and FIG. 4B is an amplification plot that shows the results for the baseline adjusted insertion analogs s10FQ and s6FQ where various plasmid copy target amounts ( $2 \times 10^2$ ,  $2 \times 10^3$ ,  $2 \times 10^4$ ,  $2 \times 10^5$ ,  $2 \times 10^6$  and  $2 \times 10^7$ ) were tested.

FIGS. 5A and 5B are amplification plots (baseline comparison ( $R_n$ ) and baseline adjusted ( $\Delta R_n$ ) comparison respectively) that demonstrate the improved performance of internal FQ (10) by itself or as a dual quenched probe compared to an internal BHQ1 (10) dual quenched probe.

FIGS. 6A and 6B are amplification plots (baseline comparison ( $R_n$ ) and baseline adjusted ( $\Delta R_n$ ) respectively) that demonstrate the increased performance of internal FQ (iFQ)

by itself or as a dual quenched containing probes compared to internal BHQ1 (iBHQ1) dual quenched probe using a different probe sequence. FIGS. 7A and 7B are amplification plots (baseline comparison ( $R_n$ ) and baseline adjusted comparison ( $\Delta R_n$ ) respectively) using an AB 7900HT instrument comparing a 5' FAM labeled dual quencher probes iFQ-3'FQ and iFQ-3'RQ-n1, illustrating the enhanced performance of an internally quenched probe in conjunction with a different 3' end quencher.

FIGS. 7A and 7B are amplification plots for the fluorescein (emission 520 nm) reporter dye probes, wherein the baseline plots are shown in FIG. 7A and baseline normalized plots are shown in FIG. 7B.

FIGS. 8A and 8B are amplification plots (baseline comparison ( $R_n$ ) and baseline adjusted comparison ( $\Delta R_n$ ) respectively) using an AB 7900HT instrument comparing a 5' MAX labeled dual quencher probes iFQ-3'FQ and iFQ-3'RQ-n1, illustrating the enhanced performance of an internally quenched probe in conjunction with a different 3' end quencher.

FIGS. 9A and 9B are amplification plots (baseline comparison ( $R_n$ ) and baseline adjusted comparison ( $\Delta R_n$ ) respectively) using an iQ5 BioRad instrument comparing a CY3 labeled dual quencher probes iFQ-3'FQ and iFQ-3'RQ-n1, illustrating the enhanced performance of an internally quenched probe in conjunction with a different 3' end quencher.

FIGS. 10A and 10B are amplification plots (baseline comparison ( $R_n$ ) and baseline adjusted comparison ( $\Delta R_n$ ) respectively) using a LC480 Roche instrument comparing a 5'TEX615 labeled dual quencher probes iFQ-3'FQ and iFQ-3'RQ-n1, illustrating the enhanced performance of an internally quenched probe in conjunction with a different 3' end quencher.

FIGS. 11A and 11B are amplification plots (baseline comparison ( $R_n$ ) and baseline adjusted comparison ( $\Delta R_n$ ) respectively) using a LC480 Roche instrument comparing a 5' CY5 labeled dual quencher probes iFQ-3'FQ and iFQ-3'RQ-n1, illustrating the enhanced performance of an internally quenched probe in conjunction with a different 3' end quencher.

FIG. 12 is a side by side baseline adjusted amplification plot ( $\Delta R_n$ ) where various plasmid copy target amounts ( $2 \times 10^2$ ,  $2 \times 10^3$ ,  $2 \times 10^4$ ,  $2 \times 10^5$ ,  $2 \times 10^6$ ,  $2 \times 10^7$ ) were tested for CY5 labeled dual quenched probes with the internal placement of the FQ quencher is between bases 9 and 10, versus 11-12, illustrating the ability to vary the positioning with the internal quencher.

FIGS. 13A, 13B, 13C, 13D and 13E are bar charts showing the  $\Delta T_m$  caused by modifying an oligonucleotide by inserting a modification compound between various nucleotides.

FIGS. 14A and 14B are plots showing the dependence of  $\Delta T_m$  on the position of the insertion within an oligonucleotide for various modification compounds.

#### DETAILED DESCRIPTION

This disclosure provides various compositions comprising oligonucleotides having modifying compounds placed internally within the oligonucleotide sequence between nucleotides. Some of the modified oligonucleotides surprisingly form equally stable, or even more stable, duplexes with their complementary oligonucleotide sequences as compared to the stability of the duplex formed between the unmodified oligonucleotide and the complementary oligonucleotide. The mechanism by which some modifying compounds confer stability to a duplex is unknown. It is particularly surprising

and unexpected that modifications of adjacent residues within an oligonucleotide with these compounds should increase duplex stability given the close proximity of the two phosphate groups on either side of reagent when in the double helix. The modifying groups may include a variety of labels, including but not limited to fluorescence quenchers that enable the design of DLPs with very high quenching efficiency. Because the labeling group is not a modified base, the same modifying compound can be inserted into any position within any oligonucleotide sequence. This disclosure also provides methods for using and making the oligonucleotide compositions.


The compositions of the present disclosure each comprise an oligonucleotide having the general structure 5'-Y<sub>1</sub>-L<sub>1</sub>-X-L<sub>2</sub>-Y<sub>2</sub>-3', where:

Y<sub>1</sub> comprises a sequence of one or more DNA and/or RNA nucleotides, including a first nucleotide N<sub>1</sub> having a 3' phosphate covalently linked to L<sub>1</sub>;

Y<sub>2</sub> comprises a sequence of one or more DNA and/or RNA nucleotides, including a second nucleotide N<sub>2</sub> having a 5' phosphate covalently linked to L<sub>2</sub>;

L<sub>1</sub> and L<sub>2</sub> each independently are a direct bond or a C<sub>1</sub>-C<sub>7</sub> alkyl, alkynyl, alkenyl, heteroalkyl, substituted alkyl, aryl, heteroaryl, substituted aryl, cycloalkyl, alkylaryl, or alkoxy group;

X is



R<sub>1</sub> is a hydrogen or a C<sub>1</sub>-C<sub>8</sub> alkyl; and M is a label.

FIG. 1 provides a non-exclusive exemplary list of modified oligonucleotides having a modifying compound inserted between adjacent nucleotides. These modified oligonucleotides may include any desired number of nucleotides, but preferably include 10-50 nucleotides, and even more preferably include 15-35 nucleotides. Moreover, depending on the application, the labeled oligonucleotide can be DNA, RNA or a chimeric oligonucleotide containing both DNA and RNA residues. Modified nucleosides such as LNA bases, 2'-O-methyl RNA and purine and pyrimidines analogs also may be included within the sequence. For use as a probe or primer, the length of the oligonucleotide is typically between 15 and 35 residues. Because the label is inserted between adjacent residues (as opposed to being a label attached to a particular nucleotide) the same modifying compound may be used to label essentially any sequence.

In some embodiments, L<sub>1</sub> and L<sub>2</sub> each may be a C<sub>1</sub>-C<sub>7</sub> alkyl, and preferably a C<sub>2</sub> alkyl. An increase in the stability of the duplex formed upon hybridization of the modified oligonucleotides to its target sequence can be achieved (see Example 1).

The 3' phosphate that is covalently linked to L<sub>1</sub>, and the 5' phosphate that is covalently linked to L<sub>2</sub>, each independently may be a phosphodiester, a phosphothioate, a phosphodithioate, a methyl phosphonate, a phosphoramidate, a phosphoramidite or a phosphotriester. A modified, neutrally charged phosphorous group could be used that would confer even greater stability.


In some embodiments, Y<sub>1</sub> comprises a sequence of four or more DNA and/or RNA nucleotides, Y<sub>2</sub> comprises a sequence of four or more DNA and/or RNA nucleotides, M is a first

quencher, and the oligonucleotide is adapted to hybridize to a second oligonucleotide having the structure 3'-Y<sub>3</sub>-Y<sub>4</sub>-5', where Y<sub>3</sub> comprises a sequence of four or more DNA and/or RNA nucleotides, including a third nucleotide N<sub>3</sub>, Y<sub>4</sub> comprises a sequence of four or more DNA and/or RNA nucleotides, including a fourth nucleotide N<sub>4</sub> that is directly attached to nucleotide N<sub>3</sub>. If the first oligonucleotide hybridizes to the second oligonucleotide, then N<sub>1</sub> base pairs with N<sub>3</sub> and N<sub>2</sub> base pairs with N<sub>4</sub> to form a duplex having a T<sub>m</sub> that is greater than the T<sub>m</sub> of a duplex formed between the second oligonucleotide and a third oligonucleotide having the structure 5'-Y<sub>1</sub>-Y<sub>2</sub>-3'. In such embodiments, the first oligonucleotide may be labeled with a fluorophore. For example, the fluorophore may be attached to the last nucleotide on the 5' end of the oligonucleotide, and in preferred embodiments, Y<sub>1</sub> may comprise a sequence of 8-12 DNA or RNA nucleotides for reasons discussed below. Compositions comprising the first oligonucleotide also may comprise the second oligonucleotide.

This disclosure also provides optimized positioning of a quencher relative to a fluorophore in a DLP. In some embodiments, the fluorophore may be attached to the nucleotide at the 5'-end of the oligonucleotide (e.g., to the 5' phosphate) and the quencher may be placed internally within the sequence between about 8 and 12 bases from the fluorophore, such as about 10 bases from the fluorophore (see Example 3). As such, in some embodiments, Y<sub>1</sub> comprises a sequence of 8-12 DNA and/or RNA nucleotides, such as a sequence of 10 DNA and/or RNA nucleotides, where the nucleotide on the 5' end of Y<sub>1</sub> is labeled with a fluorophore, and M is a quencher.

In some embodiments, M comprises a fused polycyclic aromatic moiety.

In some embodiments, M is:




where L<sub>3</sub> may be a direct bond or a C<sub>1</sub>-C<sub>8</sub> alkyl, alkenyl, alkenyl, heteroalkyl, substituted alkyl, cycloalkyl, or alkoxy, where R<sub>2</sub>-R<sub>6</sub> each independently may be a hydrogen, an alkyl, an alkenyl, a heteroalkyl, a substituted alkyl, an aryl, a heteroaryl, a substituted aryl, a cycloalkyl, an alkylaryl, an alkoxy, a ligand (e.g., amino acids, peptides, antibodies, fluorophores, biotin, enzyme conjugates, vitamins, steroids and other lipids, carbohydrates, digoxigenin and other haptens, etc.), an electron withdrawing group, or an electron donating group, and where one of R<sub>2</sub>-R<sub>6</sub> is —N=N—P, where P is a fused polycyclic aromatic moiety. Electron withdrawing groups may be selected from the group consisting of —NO<sub>2</sub>, —SO<sub>3</sub>, —SO<sub>2</sub><sup>-</sup>, —CN, —NCS, a ketone, an alkoxy, an ether, a carboxylic acid and a sulfonyl. Electron donating groups may be selected from the group consisting of an alkoxy, a heteroalkoxy, an alkyl, a cycloalkyl, a heteroalkyl, an amino, an alkylamino, or an arylamino.

In some embodiments, M is —L<sub>4</sub>-P, where L<sub>4</sub> may be an alkyl, an alkynyl, an alkenyl, a heteroalkyl, a substituted alkyl, or an alkoxy group, and P is a fused polycyclic aromatic moiety. For example, L<sub>4</sub> may be —CH<sub>2</sub>-O—CH<sub>2</sub>-CH<sub>2</sub>-NH— or any other suitable linker.




## 11


Various of the oligonucleotides described herein include fused polycyclic aromatic moieties P. In some embodiments, P may be an anthraquinone quencher having the following formula



where  $R_7$ - $R_9$  each independently are a hydrogen, an alkoxy, an alkyl, an alkylamino, an arylamino, a cycloalkyl, a heteroalkoxy, a heteroalkyl, or an amino, and  $R_{10}$ - $R_{13}$  each independently are a hydrogen, a nitro, a cyano, a carboxylate, a sulfonyl, a sulfamoyl, an alkenyl, an alkynyl, an amino, an aryl, a heteroaryl, a biaryl, a bialkenyl, a bialkynyl, an alkoxy-carbonyl or a carbamoyl. In preferred embodiments,  $R_9$  is



In some embodiments, P may be an azo quencher having the following formula



where  $R_{14}$ - $R_{19}$  each independently may be a hydrogen, an alkyl, a heteroalkyl, an aryl, a heteroaryl, an electron withdrawing group, an electron donating group, or a five or six membered ring structure formed from the  $R_1$ ,  $R_2$  pair, the  $R_3$ ,  $R_4$  pair, the  $R_4$ ,  $R_5$  pair, or the  $R_5$ ,  $R_6$  pair, and where  $R_{20}$  preferably is an electron withdrawing group, and most preferably  $-\text{NO}_2$ .

Some of the oligonucleotides disclosed herein include a fluorophore, which may include, but is not limited to, 6-carboxyfluorescein (FAM), 2'7'-dimethoxy-4'5'-dichloro-6-carboxyfluorescein (JOE), tetrachlorofluorescein (TET), 6-carboxyrhodamine (R6G), N,N,N',N'-tetramethyl-6-carboxyrhodamine (TAMRA), 6-carboxy-X-rhodamine (ROX); 1-dimethylaminonaphthyl-5-sulfonate, 1-anilino-8-naphthalene sulfonate, 2-p-toluidinyl-6-naphthalene sulfonate, 5-(2'-aminoethyl)aminonaphthalene-1-sulfonic acid (EDANS), a coumarin dye, an acridine dye, indodicarbocyanine 3 (Cy3), indodicarbocyanine 5 (Cy5), indodicarbocyanine 5.5 (Cy5.5), 3-1-carboxy-pentyl-3'-ethyl-5,5'-dimethylloxacarboxyanine (CyA); 1H,5H,11H,15H-Xantheno[2,3,4-ij:5,6,7-i'j']diquinolizin-18-ium, 9-[2(or 4)-[[[6-[2,5-dioxo-

## 12

1-pyrrolidinyloxy]-6-oxohexyl]amino]sulfonyl]-4(or 2)-sulfophenyl]-2,3,6,7,12,13,16,17-octahydro-inner salt (TR or Texas Red), a BODIPY™ dye, benzoxazole, stilbene and pyrene. In some embodiments, the fluorophore may be attached to the 5' end, such as to the phosphate at the 5' end of the oligonucleotide.

Some of the oligonucleotides disclosed herein may include more than one quencher, such as a first internal quencher (described above) and a second quencher. The second quencher may be placed internally or placed at a terminal end of the oligonucleotide. The second quencher may include, but is not limited to, an azo quencher and an anthraquinone quencher, although any quencher may be used. Examples of azo quenchers include, but are not limited to, the azo quencher shown above, dabcy1, Eclipse® quencher, BHQ1, BHQ2 and BHQ3. Examples of anthraquinone quenchers include, but are not limited to, the anthraquinone quencher shown above, Iowa Black® FQ, Iowa Black® RQ-n1 or Iowa Black® RQ-n2 (see, e.g., Laikhter et al., U.S. Patent App. 2004/0110308). Attachment of multiple quenchers to a probe not only can enhance quenching efficiency but also can provide effective quenching of various fluorophores that fluoresce over a broad spectral range (see Example 7).

The compositions of the present disclosure may be used in various assays for detecting target nucleic acids within a sample. Such methods may include contacting the sample with an oligonucleotide adapted to hybridize to the target nucleic acid, where the oligonucleotide includes an internal quencher and a fluorophore, and where the fluorescence of the fluorophore is reduced by fluorescence resonance energy transfer to the quencher and/or by ground state quenching by the quencher when the oligonucleotide is not hybridized to the second oligonucleotide, and detecting an increase in fluorescence indicating the presence of the second oligonucleotide in the sample. In some assays, such as the 5'-nuclease hydrolysis assay, the increase in fluorescence arises from cleavage of the labeled oligonucleotide. In some assays, the oligonucleotide forms a random-coil conformation when the oligonucleotide is unhybridized, such that the fluorescence of the fluorophore is reduced. In some assays, the oligonucleotide comprises a self-complementary sequence, and the quencher and fluorophore are attached to the oligonucleotide such that the fluorescence of the fluorophore is quenched by the quencher when the nucleic acid polymer undergoes intramolecular base pairing. These assays have many applications, including, but not limited to, monitoring PCR reactions, where synthesis of the PCR product results in an increase in fluorescence.

Function of dual-labeled probes in the 5'-nuclease hydrolysis assay requires that the fluorophore be effectively quenched by the quencher and also requires that the chemical modifiers employed (dye and quencher) do not interfere with nuclease cleavage. If cleavage is prevented or rendered inefficient by the presence of the chemical modifications, then fluorophore and quencher remain linked during PCR cycles and no detectable signal is generated. The chemical compositions disclosed herein function to efficiently quench the fluorophore and are compatible with probe hydrolysis using 5'-nuclease positive DNA polymerases, like Taq DNA polymerase. The best results are obtained when the internal quencher is positioned as close to the fluorophore as possible (to maximize quenching) yet still permits efficient cleavage of the nucleic acid bases between the dye and quencher (maximize the resulting fluorescent signal). Many fluorescent quenching groups can be placed internally and achieve efficient quenching, however many if not most of these chemical groups interfere with probe cleavage, especially when the

distance between fluorophore and quencher is less than 12 nucleotides. This principle is demonstrated in Example 3, where certain quenchers, such as BHQ-1, can achieve very efficient quenching of a 5'-fluorophore, such as Fluorescein (6-FAM), yet the actual magnitude of the fluorescent signal generated during real-time PCR is small, compromising the actual performance and sensitivity of the assay. In contrast, similar internal placement of the quenchers provided herein, are fully compatible with probe hydrolysis and final functional signal generation is large.

Traditional DLPs having a 5'-fluorophore and 3'-quencher perform poorly when probe length nears or exceeds 30 nucleotides (nt), because quenching efficiency drops to the point that the probe remains relatively bright even in the quenched state. The present disclosure provides probes of 30 nt length, 35 nt length, or longer as needed for the precise application, having internal quenchers substantially closer to the fluorophore. Quenching in the compositions of this disclosure remains highly efficient as the internal quencher can be inserted the same distance from the fluorophore regardless of probe length. Thus high quality, efficiently quenched probes of an expanded potential length range are possible, which may be of particular importance when working with nucleic acids which are very AT rich. Sequences that are AT-rich have lower melting temperatures and longer probes must be utilized to function in the temperature ranges typically needed for PCR.


The compounds of this disclosure can also be utilized in molecular beacon assays. Molecular beacon assays contain probes that contain terminal 3' and 5' ends that self-hybridize to form a stem-loop structure. One end typically contains a terminal fluorophore group and the other end contains a terminal quencher group. When the probe hybridizes to the target the quencher is no longer near the fluorophore and the signal increases. Typically the hybridizing portions of the probes are 4-7 base pairs long. Beacon probes that contain insertions as described herein, preferably within the hybridized portion of the beacon, and more preferably within 1-2 bases from the terminal end, the stability is increased. Therefore shorter hybridizing regions can be used to generate the same performance as a conventional beacon.

A wide variety of reactive fluorescent reporter dyes are known in the literature and can be used in the compositions of this disclosure, so long as they are quenched by the precise quencher group or combination of quencher groups employed. The precise fluorophore/quencher pair employed in a dual-labeled probe is usually carefully chosen from a relatively small set of pairings that work well together based upon the emission wavelength of the fluorophore and the absorption wavelength of the quencher. Placement of the quencher in an internal position, as provided here, closer to the fluorophore than is achievable with an end-labeled probe permits efficient quenching even of fluorophore/quencher pairs that usually do not work well together, expanding the utility of the quencher by enabling its use with a wider range of fluorophores. Attachment of a second quencher to the probe can even further expand the useful spectral range.

The oligonucleotide probes provided herein may incorporate one or more fluorophores. The fluorophores can be attached internally or at the 5'- or 3'-end. Typically, the fluorophore is an aromatic or heteroaromatic compound and can be a pyrene, anthracene, naphthalene, acridine, stilbene, indole, benzindole, oxazole, thiazole, benzothiazole, cyanine, carbocyanine, salicylate, anthranilate, coumarin, fluorescein, rhodamine or other like compound. Suitable fluorescent reporters include xanthene dyes, such as fluorescein or rhodamine dyes, including 6-carboxyfluorescein (FAM),

2'7'-dimethoxy-4'5'-dichloro-6-carboxyfluorescein (JOE), tetrachlorofluorescein (TET), 6-carboxyrhodamine (R6G), N,N,N,N'-tetramethyl-6-carboxyrhodamine (TAMRA), 6-carboxy-X-rhodamine (ROX). Suitable fluorescent reporters also include the naphthylamine dyes that have an amino group in the alpha or beta position. For example, naphthylamino compounds include 1-dimethylaminonaphthyl-5-sulfonate, 1-anilino-8-naphthalene sulfonate and 2-p-toluidinyl-6-naphthalene sulfonate, 5-(2'-aminoethyl)aminonaphthalene-1-sulfonic acid (EDANS). Other fluorescent reporter dyes include coumarins, such as 3-phenyl-7-isocyanatocoumarin; acridines, such as 9-isothiocyanatoacridine and acridine orange; N-(p-(2-benzoxazolyl)phenyl)maleimide; cyanines, such as indodicarbocyanine 3 (Cy3), indodicarbocyanine 5 (Cy5), indodicarbocyanine 5.5 (Cy5.5), 3-1-carboxy-pentyl-3'-ethyl-5,5'-dimethyloxycarbocyanine (CyA); 1H,5H,11H,15H-Xantheno[2,3,4-ij:5,6,7-i'j']diquinolizin-18-ium, 9-[2(or 4)-[[[6-[2,5-dioxo-1-pyrrolidinyl]oxy]-6-oxohexyl]amino]sulfonyl]-4(or 2)-sulfophenyl]-2,3,6,7,12,13,16,17-octahydro-inner salt (TR or Texas Red); BODIPY™ dyes; benzoxaazoles; stilbenes; pyrenes; and the like. See Haugland, "Molecular Probes Handbook of Fluorescent Probes and Research Chemicals" for further fluorophore examples.

Reagents for incorporation the modification compounds of the present disclosure into oligonucleotides may have the following general structure:



where  $R_{1,3}$  is a protecting group on the oxygen atom, most commonly a trityl group, and preferably a dimethoxytrityl group, and  $R_{2,2}$  is a phosphoramidite, a phosphate group, or a hydrogen phosphate used to couple the reagent to the growing oligonucleotide chain during synthesis. A phosphoramidite is preferred. A N,N-diisopropyl- $\beta$ -cyanoethyl phosphoramidite is the most preferred reactive group.

As used herein, the terms "nucleic acid" and "oligonucleotide," as used herein, refer to polydeoxyribonucleotides (containing 2-deoxy-D-ribose), polyribonucleotides (containing D-ribose), and to any other type of polynucleotide which is an N glycoside of a purine or pyrimidine base. There is no intended distinction in length between the terms "nucleic acid", "oligonucleotide", "oligomer" or "oligo", and these terms will be used interchangeably. These terms refer only to the primary structure of the molecule. Thus, these terms include double- and single-stranded DNA, as well as double- and single-stranded RNA. An oligonucleotide also can comprise nucleotide analogs in which the base, sugar, or phosphate backbone is modified as well as non-purine or non-pyrimidine nucleotide analogs oligonucleotides, which may comprise naturally occurring nucleosides or chemically modified nucleosides. In some embodiments, the compounds comprise modified sugar moieties, modified internucleoside linkages, or modified nucleobase moieties.

The term "base" as used herein includes purines, pyrimidines and non-natural bases and modifications well-known in the art. Purines include adenine, guanine and xanthine and modified purines such as 8-oxo-N<sup>6</sup>-methyladenine and 7-deazaxanthine. Pyrimidines include thymine, uracil and

cytosine and their analogs such as 5-methylcytosine and 4,4-ethanocytosine. Non-natural bases include 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N<sup>6</sup>-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N<sup>6</sup>-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N<sup>6</sup>-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methyl ester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, nitroindole, and 2,6-diaminopurine.

The term “base” is sometimes used interchangeably with “monomer”, and in this context it refers to a single nucleic acid or oligomer unit in a nucleic acid chain.

The term “probe” as used herein refers to nucleic acid oligonucleotides that produce a detectable response upon interaction with a target. The probes include at least one detectable moiety, a pair of moieties that form an energy transfer pair detectable upon some change of state of the probe in response to its interaction with a binding partner, or more than two moieties such as a fluorophore and more than one quencher.

The term “primer,” as used herein, refers to an oligonucleotide capable of acting as a point of initiation of DNA synthesis under suitable conditions. Such conditions include those in which synthesis of a primer extension product complementary to a nucleic acid strand is induced in the presence of four different nucleoside triphosphates and an agent for extension (e.g., a DNA polymerase or reverse transcriptase) in an appropriate buffer and at a suitable temperature. A primer is preferably a single-stranded DNA. The appropriate length of a primer depends on the intended use of the primer but typically ranges from 6 to 50 nucleotides, preferably from 15-35 nucleotides. Short primer molecules generally require cooler temperatures to form sufficiently stable hybrid complexes with the template. A primer need not reflect the exact sequence of the template nucleic acid, but must be sufficiently complementary to hybridize with the template. The design of suitable primers for the amplification of a given target sequence is well known in the art and described in the literature cited herein. Primers can incorporate additional features which allow for the detection or immobilization of the primer but do not alter the basic property of the primer, that of acting as a point of initiation of DNA synthesis. For example, primers may contain an additional nucleic acid sequence at the 5' end which does not hybridize to the target nucleic acid, but which facilitates cloning or detection of the amplified product. The region of the primer which is sufficiently complementary to the template to hybridize is referred to herein as the hybridizing region.

The term “hybridization,” as used herein, refers to the formation of a duplex structure by two single-stranded nucleic acids due to complementary base pairing. Hybridization can occur between fully complementary nucleic acid strands or between “substantially complementary” nucleic acid strands that contain minor regions of mismatch. Conditions under which hybridization of fully complementary nucleic acid strands is strongly preferred are referred to as “stringent hybridization conditions” or “sequence-specific

hybridization conditions”. Stable duplexes of substantially complementary sequences can be achieved under less stringent hybridization conditions; the degree of mismatch tolerated can be controlled by suitable adjustment of the hybridization conditions. Those skilled in the art of nucleic acid technology can determine duplex stability empirically considering a number of variables including, for example, the length and base pair composition of the oligonucleotides, ionic strength, and incidence of mismatched base pairs, following the guidance provided by the art (see, e.g., Sambrook et al., 1989, *Molecular Cloning—A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.; Wetmur, 1991, *Critical Review in Biochem. and Mol. Biol.* 26(3/4):227-259; and Owczarzy et al., 2008, *Biochemistry*, 47: 5336-5353, which are incorporated herein by reference).

The term “amplification reaction” refers to any chemical reaction, including an enzymatic reaction, which results in increased copies of a template nucleic acid sequence or results in transcription of a template nucleic acid. Amplification reactions include reverse transcription, the polymerase chain reaction (PCR), including Real Time PCR (see U.S. Pat. Nos. 4,683,195 and 4,683,202; PCR Protocols: A Guide to Methods and Applications (Innis et al., eds, 1990)), and the ligase chain reaction (LCR) (see Barany et al., U.S. Pat. No. 5,494,810). Exemplary “amplification reactions conditions” or “amplification conditions” typically comprise either two or three step cycles. Two step cycles have a high temperature denaturation step followed by a hybridization/elongation (or ligation) step. Three step cycles comprise a denaturation step followed by a hybridization step followed by a separate elongation or ligation step.

The following examples further illustrate the invention but, of course, should not be construed as in any way limiting its scope.

#### Example 1

This example demonstrates the improved stability of probes containing modifications of the present disclosure compared to a selection of other compounds.

##### Oligonucleotide Synthesis and Purification.

DNA oligonucleotides were synthesized using solid phase phosphoramidite chemistry, deprotected and desalted on NAP-5 columns (Amersham Pharmacia Biotech, Piscataway, N.J.) according to routine techniques (Caruthers et al., *Methods Enzymol* 1992, 211:3-20). The oligomers were purified using reversed-phase high performance liquid chromatography (RP-HPLC). The purity of each oligomer was determined by capillary electrophoresis (CE) carried out on a Beckman P/ACE MDQ system (Beckman Coulter, Inc., Fullerton, Calif.). All single strand oligomers were at least 90% pure. Electrospray-ionization liquid chromatography mass spectroscopy (ESI-LCMS) of the oligonucleotides was conducted using an Oligo HTCS system (Novatia, Princeton, N.J.), which consisted of ThermoFinnigan TSQ7000, Xcalibur data system, ProMass data processing software and Paradigm MS4™ HPLC (Michrom BioResources, Auburn, Calif.). Protocols recommended by manufacturers were followed. Experimental molar masses for all single strand oligomers were within 1.5 g/mol of expected molar mass. These results confirm identity of the oligomers.

##### Preparation of DNA Samples.

Melting experiments were carried out in buffer containing 3.87 mM NaH<sub>2</sub>PO<sub>4</sub>, 6.13 mM Na<sub>2</sub>HPO<sub>4</sub>, 1 mM Na<sub>2</sub>EDTA and 1000 mM NaCl. 1 M NaOH was used to titrate each solution to pH 7.0. Total sodium concentrations were estimated to be 1020 mM. The DNA samples were thoroughly

dialyzed against melting buffer in a 28-Well Microdialysis System (Life Technologies, Carlsbad, Calif.) following the manufacturer's recommended protocol. Concentrations of DNA oligomers were estimated from the samples' UV absorbance at 260 nm in a spectrophotometer (Beckman Coulter, Inc., Fullerton, Calif.), using extinction coefficients for each oligonucleotide that were estimated using the nearest neighbor model for calculating extinction coefficients. (See, Warshaw et al., J. Mol. Biol. 1966, 20:29-38).

#### Internal Modifications Studied.

FIGS. 1A-1C show the structures of modified portions of the various modified oligonucleotides studied in this example and in Example 2. The FQ (Integrated DNA Technologies, Inc., sometimes referred to as "iFQ" in this application), was introduced into oligonucleotides using phosphoramidite reagents at the time of synthesis. See Example 10 for synthesis of the phosphoramidite. In the first series of duplexes, the iFQ group was placed as an insertion between bases in the duplex so that a 10 base top strand annealed to a 10 base bottom strand and the iFQ group was not aligned to a base. Additionally, 10-mer oligonucleotides with C3 spacer insertions were also synthesized and studied. The C3 spacer represents the control wherein a linear insertion of a phosphate group plus propanediol is placed between bases, which is similar to the iFQ insertions without having the nathylene-azo ring structures present. Extinction coefficients at 260 nm of iFQ was estimated to be 13340; the C3 spacer does not contribute to UV absorbance. Two 20 base and two 25 base duplexes of similar design were also studied. A second set of 10 base duplexes was studied where the iFQ group was placed as a substitution such that a 10 base top strand (5 bases—iFQ—5 bases) was annealed to an 11 base bottom strand so that the iFQ group functioned as a substitution or replacement for a base. Four duplexes of this design were tested, one comprising each of the 4 bases (AGCT) to pair with the iFQ group.

#### Measurement of Melting Curves.

Oligomer concentrations were measured at least twice for each sample. If the estimated concentrations for any sample differed more than 4%, the results were discarded and new

absorbance measurements were performed. To prepare oligonucleotide duplexes, complementary DNA oligomers were mixed in 1:1 molar ratio, heated to 367 K (i.e., 94° C.) and slowly cooled to an ambient temperature. Each solution of duplex DNA was diluted with melting buffer to a total DNA concentration ( $C_T$ ) of 2  $\mu$ M.

Melting experiments were conducted on a single beam Beckman DU 650 spectrophotometer (Beckman-Coulter) with a Micro  $T_m$  Analysis accessory, a Beckman High Performance Peltier Controller (to regulate the temperature), and 1 cm path-length cuvettes. Melt data were recorded using a PC interfaced to the spectrophotometer. UV-absorbance values at 268 nm wavelength were measured at 0.1 degree increments in the temperature range from 383 to 368 K (i.e., 10-95° C.). Both heating (i.e., denaturation) and cooling (i.e., renaturation) transition curves were recorded in each sample at a controlled rate of temperature change (24.9 $\pm$ 0.3° C. per hour). Sample temperatures were collected from the internal probe located inside the Peltier holder, and recorded with each sample's UV-absorbance data. Melting profiles were also recorded for samples of buffer alone (no oligonucleotide), and these blank profiles were digitally subtracted from melting curves of the DNA samples. To minimize systematic errors, at least two melting curves were collected for each sample in different cuvettes and in different positions within the Peltier holder.

#### Determination of Melting Temperatures.

To determine each sample's melting temperature, the melting profiles were analyzed using methods that have been previously described (see, Doktycz et al., *Biopolymers* 1992, 32:849-864; Owczarzy et al., *Biopolymers* 1997, 44:217-239; Owczarzy R., *Biophys. Chem.* 2005, 117: 207-215.). Briefly, the experimental data for each sample was smoothed, using a digital filter, to obtain a plot of the sample's UV-absorbance as a function of its temperature. The fraction of single-stranded oligonucleotide molecules,  $\theta$ , was then calculated from that plot. The melting temperature or  $T_m$  of a sample was defined as the temperature where  $\theta=0.5$ .

Table 1 lists the sequences tested, the internal quenchers used, and the resulting melting temperatures.

TABLE 1

Melting Temperatures for nucleic acids containing internal quencher moieties, where iFQ = internal FQ azo quencher, and iSpC3 = internal C3 spacer.				
SEQ ID NO.	Duplex Sequence	N	$T_m$ (C.)	$\Delta T_m$ (C.)
1	5' -ATCGTTGCTA-3'	10	43.85	0.0
2	3' -TAGCAACGAT-5'	10		
3	5' -ATC/iFQ/GTTGCTA-3'	10	48.05	4.2
2	3' -TAGCAACGAT-5'	10		
4	5' -ATCG/iFQ/TTGCTA-3'	10	48.55	4.7
2	3' -TAGCAACGAT-5'	10		
5	5' -ATCGT/iFQ/TGCTA-3'	10	46.35	2.5
2	3' -TAGCAACGAT-5'	10		
6	5' -CTTGGATCGTTGCTAGTAGG-3'	20	69.55	0.0
7	3' -GAACCTAGCAACGATCATCC-5'	20		
8	5' -CTTGGATCGT/iFQ/TGCTAGTAGG-3'	20	71.35	1.8
7	3' -GAACCTAGCAACGATCATCC-5'	20		
9	5' -CACTTGGATCGTTGCTAGTAGGGTC-3'	25	76.15	0.0
10	3' -GTGAACCTAGCAACGATCATCCAG-5'	25		

TABLE 1-continued

Melting Temperatures for nucleic acids containing internal quencher moieties, where iFQ = internal FQ azo quencher, and iSpC3 = internal C3 spacer.				
SEQ ID NO.	Duplex Sequence	N	T <sub>m</sub> (C.)	ΔT <sub>m</sub> (C.)
11	5' -CACTTGGATC/iFQ/GTTGCTAGTAGGGTC-3'	25	77.05	0.9
10	3' -GTGAACCTAGCAACGATCATCCCAG-5'	25		
12	5' -ATC/iSpC3/GTTGCTA-3'	10	36.35	-7.5
2	3' -TAGCAACGAT-5'	10		
13	5' -ATCG/iSpC3/TTGCTA-3'	10	36.55	-7.3
2	3' -TAGCAACGAT-5'	10		
14	5' -ATCGT/iSpC3/TGCTA-3'	10	32.55	-11.3
2	3' -TAGCAACGAT-5'	10		
5	5' -ATCGT/iFQ/TGCTA-3'	10	47.35	3.5
15	3' -TAGCA/iSpC3/ACGAT-5'	10		
5	5' -ATCGT/iFQ/TGCTA-3'	10	42.24	-1.6
16	3' -TAGCAAACGAT-5'	11		
5	5' -ATCGT/iFQ/TGCTA-3'	10	45.27	1.4
17	3' -TAGCACACGAT-5'	11		
5	5' -ATCGT/iFQ/TGCTA-3'	10	40.44	-3.4
18	3' -TAGCAGACGAT-5'	11		
5	5' -ATCGT/iFQ/TGCTA-3'	10	45.27	1.4
19	3' -TAGCATACGAT-5'	11		

Three different insertion placement sites were studied using a 10-mer oligonucleotide scaffold. Use of the shorter sequences most clearly demonstrates the potential effects on T<sub>m</sub> and testing different placement sites illustrate that the T<sub>m</sub> effects can be sequence context dependent. The relative ΔT<sub>m</sub> shifts for the modified vs. unmodified 10mer sequences were averaged and are summarized in Table 2 below.

TABLE 2

Average ΔT <sub>m</sub> shifts for three 10mer sequences with internal modifiers			
Modifier	iFQ (insertion)	iFQ (substitution)	iSpC3
ΔT <sub>m</sub>	+3.8° C.	-0.6° C.	-8.7° C.

As Tables 1 and 2 illustrate, disrupting a DNA sequence with a modification that is small and offers no steric hindrance (or stabilization) like a propanediol group (C3 spacer) has a significant negative impact on the T<sub>m</sub> of a duplex (ΔT<sub>m</sub> of -8.7° C.). In contrast, the naphthylene-azo-class quencher studied (iFQ) significantly stabilized the duplex compared with the iC3 control. A greater degree of stabilization was seen when the iFQ was placed as an insertion (ΔT<sub>m</sub> of +12.5° C. relative to the iSpC3) than when the iFQ was placed as a base substitution (ΔT<sub>m</sub> of +8.1° C. relative to the iSpC3). Unexpectedly, use of the iFQ group as an insertion between bases stabilized the duplex compared to the unmodified parent duplex (ΔT<sub>m</sub> of +3.8° C. relative to the unmodified duplex), while base substitution resulted in slight destabilization (ΔT<sub>m</sub> of -0.6° C. relative to the unmodified duplex).

Therefore internal incorporation of the naphthylene-azo group within a DNA duplex stabilizes the duplex when placed as an insertion between bases.

Certain anthraquinone groups can stabilize a duplex when placed on the ends (J. Am. Chem. Soc., 131:12671-12681,

2009); however this effect has not been described for internal placement or using naphthylene-azo compounds. Therefore the use of internal naphthylene-azo-class quenchers would be preferred to maintain duplex stability.

## Example 2

The following example compares 10 base pair sets of duplexes with varying modification insertions placed at varying location along the 10-mer oligonucleotide. The structures of the modified portions of the various modified oligonucleotides are illustrated in FIGS. 1A-1C. As noted earlier, the synthesis of FQ phosphoramidite is described in Example 10. The structure "IB 1.1" is synthesized in the same manner as FQ except an aminoanthraquinone reagent is used instead of 4-nitro-1-naphthylamine. The IB RQ quenchers are anthraquinone-based compounds (U.S. Pat. Application 2004/0110308) which are commonly used with red wavelength fluorescent dyes.

The preparation of the DNA samples, the measurement of melting curves and determination of melting temperatures were performed as in Example 1. Table 3 lists the resulting T<sub>m</sub> data for each duplex studied, as well as the ΔT<sub>m</sub> relative to the duplex formed with the unmodified oligonucleotide.

TABLE 3

SEQ ID No.	Sequence	T <sub>m</sub>	ΔT <sub>m</sub>
1	5' ATCGTTGCTA	43.9	-
2	3' TAGCAACGAT		
20	5' ATC/GTTGCTA N-MDA "/"	31.0	-12.9
2	3' TAG CAACGAT		

21

TABLE 3-continued

SEQ ID No.	Sequence	$T_m$	$\Delta T_m$	
21 2	5' ATC/GTTGCTA C2 "/" 3' TAG CAACGAT	37.8	-6.1	5
22 2	5' ATC/GTTGCTA 2,2 DMP "/" 3' TAG CAACGAT	34.9	-9.0	
12 2	5' ATC/GTTGCTA iSpC3 "/" 3' TAG CAACGAT	36.3	-7.6	10
23 2	5' ATC/GTTGCTA C4 "/" 3' TAG CAACGAT	30.1	-13.8	
24 2	5' ATC/GTTGCTA C5 "/" 3' TAG CAACGAT	27.3	-16.6	15
25 2	5' ATC/GTTGCTA C6 "/" 3' TAG CAACGAT	26.1	-17.8	
26 2	5' ATC/GTTGCTA C7 "/" 3' TAG CAACGAT	24.7	-19.1	20
27 2	5' ATC/GTTGCTA iSpS9 "/" 3' TAG CAACGAT	28.5	-15.4	
28 2	5' ATC/GTTGCTA idSp "/" 3' TAG CAACGAT	33.5	-10.4	25
3 2	5' ATC/GTTGCTA iFQ "/" 3' TAG CAACGAT	48.0	+4.1	
29 2	5' ATC/GTTGCTA iBHQ2 "/" 3' TAG CAACGAT	45.0	+1.1	30
30 2	5' ATC/GTTGCTA iRQ-n1 "/" 3' TAG CAACGAT	38.7	-5.2	
31 2	5' ATC/GTTGCTA iRQ-n2 "/" 3' TAG CAACGAT	46.0	+2.1	35
32 2	5' ATC/GTTGCTA iEc "/" 3' TAG CAACGAT	40.1	-3.8	
33 2	5' ATC/GTTGCTA IB 1.1 "/" 3' TAG CAACGAT	47.6	+3.7	40
34 2	5' ATC/GTTGCTA NPDA "/" 3' TAG CAACGAT	29.3	-14.6	
35 2	5' ATCG/TTGCTA N-MDA "/" 3' TAGC AACGAT	34.1	-9.8	45
36 2	5' ATCG/TTGCTA C2 "/" 3' TAGC AACGAT	38.6	-5.3	
37 2	5' ATCG/TTGCTA 2,2 DMP "/" 3' TAGC AACGAT	35.5	-8.4	50
13 2	5' ATCG/TTGCTA iSpC3 "/" 3' TAGC AACGAT	36.6	-7.3	
38 2	5' ATCG/TTGCTA C4 "/" 3' TAGC AACGAT	32.0	-11.9	55
39 2	5' ATCG/TTGCTA C5 "/" 3' TAGC AACGAT	28.1	-15.8	
40 2	5' ATCG/TTGCTA C6 "/" 3' TAGC AACGAT	26.5	-17.4	
41 2	5' ATCG/TTGCTA C7 "/" 3' TAGC AACGAT	24.7	-19.2	60
42 2	5' ATCG/TTGCTA iSpS9 "/" 3' TAGC AACGAT	29.1	-14.8	
43 2	5' ATCG/TTGCTA idSp "/" 3' TAGC AACGAT	34.1	-9.8	65

22

TABLE 3-continued

SEQ ID No.	Sequence	$T_m$	$\Delta T_m$
4 2	5' ATCG/TTGCTA iFQ "/" 3' TAGC AACGAT	48.6	+4.7
44 2	5' ATCG/TTGCTA iBHQ2 "/" 3' TAGC AACGAT	43.3	-0.6
45 2	5' ATCG/TTGCTA iRQ-n1 "/" 3' TAGC AACGAT	34.7	-9.2
46 2	5' ATCG/TTGCTA iRQ-n2 "/" 3' TAGC AACGAT	44.7	+0.8
47 2	5' ATCG/TTGCTA iEc "/" 3' TAGC AACGAT	35.7	-8.2
48 2	5' ATCG/TTGCTA IB 1.1 "/" 3' TAGC AACGAT	45.9	+2.1
49 2	5' ATCG/TTGCTA NPDA "/" 3' TAGC AACGAT	30.0	-13.9
50 2	5' ATCGT/TGCTA C2 "/" 3' TAGCA ACGAT	34.0	-9.9
51 2	5' ATCGT/TGCTA 2,2 DMP "/" 3' TAGCA ACGAT	33.1	-10.8
14 2	5' ATCGT/TGCTA iSpC3 "/" 3' TAGCA ACGAT	32.6	-11.3
52 2	5' ATCGT/TGCTA C4 "/" 3' TAGCA ACGAT	26.4	-17.5
53 2	5' ATCGT/TGCTA C5 "/" 3' TAGCA ACGAT	23.3	-20.6
54 2	5' ATCGT/TGCTA C6 "/" 3' TAGCA ACGAT	21.9	-22.0
55 2	5' ATCGT/TGCTA C7 "/" 3' TAGCA ACGAT	20.4	-23.5
56 2	5' ATCGT/TGCTA iSpS9 "/" 3' TAGCA ACGAT	24.4	-19.5
57 2	5' ATCGT/TGCTA idSp "/" 3' TAGCA ACGAT	31.8	-12.1
5 2	5' ATCGT/TGCTA iFQ "/" 3' TAGCS ACGAT	46.3	+2.4
58 2	5' ATCGT/TGCTA iBHQ2 "/" 3' TAGCA ACGAT	41.5	-2.4
59 2	5' ATCGT/TGCTA iRQ-n1 "/" 3' TAGCA ACGAT	32.1	-11.8
60 2	5' ATCGT/TGCTA iRQ-n2 "/" 3' TAGCA ACGAT	42.6	-1.3
61 2	5' ATCGT/TGCTA iEc "/" 3' TAGCA ACGAT	33.1	-10.8
62 2	5' ATCGT/TGCTA IB 1.1 "/" 3' TAGCA ACGAT	44.7	+0.8
63 2	5' ATCGT/TGCTA NPDA "/" 3' TAGCA ACGAT	26.8	-17.1
64 2	5' A/TCGTTGCTA N-MDA "/" 3' T AGCAACGAT	40.9	-3.0
65 2	5' A/TCGTTGCTA C2 "/" 3' T AGCAACGAT	43.5	-0.4

TABLE 3-continued

SEQ ID No.	Sequence	$T_m$	$\Delta T_m$
66 2	5' A/TCGTTGCTA 2,2 DMP "/" 3' T AGCAACGAT	43.4	-0.4
67 2	5' A/TCGTTGCTA iSpC3 "/" 3' T AGCAACGAT	44.6	+0.7
68 2	5' A/TCGTTGCTA C4 "/" 3' T AGCAACGAT	43.3	-0.6
69 2	5' A/TCGTTGCTA C5 "/" 3' T AGCAACGAT	41.5	-2.4
70 2	5' A/TCGTTGCTA C6 "/" 3' T AGCAACGAT	41.6	-2.3
71 2	5' A/TCGTTGCTA C7 "/" 3' T AGCAACGAT	43.2	-0.7
72 2	5' A/TCGTTGCTA iSpS9 "/" 3' T AGCAACGAT	41.5	-2.4
73 2	5' A/TCGTTGCTA idSp "/" 3' T AGCAACGAT	43.0	-0.9
74 2	5' A/TCGTTGCTA iFQ "/" 3' T AGCAACGAT	51.8	+7.9
75 2	5' A/TCGTTGCTA iBHQ2 "/" 3' T AGCAACGAT	48.9	+5.0
76 2	5' A/TCGTTGCTA iRQ-n1 "/" 3' T AGCAACGAT	48.2	+4.3
77 2	5' A/TCGTTGCTA iRQ-n2 "/" 3' T AGCAACGAT	49.1	+5.2
78 2	5' A/TCGTTGCTA iEc "/" 3' T AGCAACGAT	44.5	+0.6
79 2	5' A/TCGTTGCTA IB 1.1 "/" 3' T AGCAACGAT	51.2	+7.3
80 2	5' A/TCGTTGCTA NPDA "/" 3' T AGCAACGAT	41.0	-2.9
81 2	5' ATCGTTGCT/A C2 "/" 3' TAGCAACGA T	44.2	+0.3
82 2	5' ATCGTTGCT/A 2,2 DMP "/" 3' TAGCAACGA T	43.8	-0.1
83 2	5' ATCGTTGCT/A iSpC3 "/" 3' TAGCAACGA T	44.5	+0.6
84 2	5' ATCGTTGCT/A C4 "/" 3' TAGCAACGA T	43.4	-0.5
85 2	5' ATCGTTGCT/A C5 "/" 3' TAGCAACGA T	42.8	-1.0
86 2	5' ATCGTTGCT/A C6 "/" 3' TAGCAACGA T	43.3	-0.6
87 2	5' ATCGTTGCT/A C7 "/" 3' TAGCAACGA T	40.9	-3.0
88 2	5' ATCGTTGCT/A iSpS9 "/" 3' TAGCAACGA T	44.2	+0.3
89 2	5' ATCGTTGCT/A idSp "/" 3' TAGCAACGA T	44.7	+0.8
90 2	5' ATCGTTGCT/A iFQ "/" 3' TAGCAACGA T	50.3	+6.4
91 2	5' ATCGTTGCT/A iBHQ2 "/" 3' TAGCAACGA T	50.7	+6.8

TABLE 3-continued

SEQ ID No.	Sequence	$T_m$	$\Delta T_m$
92 2	5' ATCGTTGCT/A iRQ-n1 "/" 3' TAGCAACGA T	48.6	+4.7
93 2	5' ATCGTTGCT/A iRQ-n2 "/" 3' TAGCAACGA T	47.5	+3.6
94 2	5' ATCGTTGCT/A iEc "/" 3' TAGCAACGA T	45.8	+1.9
95 2	5' ATCGTTGCT/A IB 1.1 "/" 3' TAGCAACGA T	50.9	+7.0
96 2	5' ATCGTTGCT/A NPDA "/" 3' TAGCAACGA T	43.9	+0.0

FIGS. 13A, 13B, 13C, 13D and 13E are bar charts each showing the  $\Delta T_m$  caused by each of the modifications at a particular location within the nucleotide. FIGS. 14A and 14B are plots showing the dependence of  $\Delta T_m$  on the position of the insertion within an oligonucleotide for each of the various modification compounds. In general, the insertion of a modification nearer the end of an oligonucleotide is less destabilizing than when it is inserted towards the middle of the oligonucleotide. The spacer modifications were all destabilizing, with the degree of destabilization increasing as the size of the spacer increased.

The FQ modification had a positive effect on stability whether it was inserted near an end or in the middle of the duplex. The IB 1.1 had nearly the same stability profile as FQ. The iRQ-n2 modification was also positive or negligible in its effect on duplex stability.

### Example 3

This example details the predictive modeling for oligonucleotides containing an internal quencher.

The thermodynamic impact of internal FQ azo quencher (iFQ) modification was determined from the difference between modified and native duplex DNAs. Melting experiments were conducted at 2  $\mu$ M DNA concentration ( $C_1$ ) and in 1M Na<sup>+</sup> buffer. Transition enthalpies ( $\Delta H^\circ$ ) and entropies ( $\Delta S^\circ$ ) were obtained from fits to individual melting profiles (Petersheim, M.& Turner, D. H. (1983) *Biochemistry*, 22, 256-263). The equilibrium constant  $K_a$  was calculated from the fraction of broken base pairs ( $\theta$ ) at each temperature,

$$K_a = \frac{2(1-\theta)}{\theta^2 C_1} \quad (1)$$

The graphs of  $-\ln K_a$  vs.  $1/T$  were least-square fit to straight lines and thermodynamic parameters were calculated from the slope and the intercept,

$$-\ln K_a = \frac{\Delta H^\circ}{RT} - \frac{\Delta S^\circ}{R} \quad (2)$$

The fits were limited to the range of  $\theta$  from 0.15 to 0.85, where  $\theta$  and  $K_a$  are the most accurate. The symbol R is the

## 25

ideal gas constant (1.9865 cal/(mol\*K)). Reported values of  $\Delta H^\circ$ , and  $\Delta S^\circ$  are averages from at least four heating and cooling melting profiles. This thermodynamic analysis assumes that the transition enthalpies and entropies are temperature-independent and melting transitions proceed in two-state manner. The results are summarized in Table 4. Average thermodynamic effect of inserted FQ azo quencher could be estimated from the following relationships,

$$\Delta H^\circ (\text{modified oligo}) = \Delta H^\circ (\text{native oligo}) - 1.272 \text{ cal/mol}$$

$$\Delta S^\circ (\text{modified oligo}) = \Delta S^\circ (\text{native oligo}) - 1.44 \text{ cal/(mol}\cdot\text{K)}$$

When these equations were employed to predict melting temperatures of 20 and 25 base pair duplexes (Table 5), the average error of predictions was 0.7° C.

TABLE 4

Thermodynamic effects of internal FQ modifications					
SEQ ID No.	DNA sequence (5' to 3') <sup>a</sup>	$\Delta H^\circ$ cal/mol	$\Delta\Delta H^\circ$ cal/mol	$\Delta S^\circ$ cal/(mol · K)	$\Delta\Delta S^\circ$ cal/(mol · K)
1	ATCGTTGCTA	-68012		-185.62	
3	ATC/iFQ/GTTGCTA	-71827	-3815	-194.73	-9.12
4	ATCG/iFQ/TTGCTA	-69783	-1771	-188.00	-2.38
5	ATCGT/iFQ/TGCTA	-66241	1771	-178.45	7.17
	Average effect		-1272		-1.44

<sup>a</sup>Complementary DNA strand was 5'-TAGCAACGAT-3'. Calculations were done using non-rounded values.

TABLE 5

Accuracy of $T_m$ predictions for two duplex DNAs containing internal FQ quencher that were not used to derive equations (1) and (2).					
SEQ ID No.	DNA sequence (5' to 3') <sup>a</sup>	$N_{bp}$	Exper. $T_m$ (° C.)	Predicted $T_m$ (° C.)	Error of $T_m$ prediction (° C.)
8	CTTGGATCGT/iFQ/TGCTAGTAGG	20	71.3	71.6	0.3
11	CACTTGGATC/iFQ/GTTGCTAGTAGGGTC	25	77.1	78.2	1.1

<sup>a</sup>Melting temperatures were calculated using the nearest-neighbor model (SantaLucia, J., Jr. (1998) *Proc. Natl. Acad. Sci. USA*, 95, 1460-1465) and equations (1) and (2).

The low error of prediction demonstrates the consistency and predictability of the effect of the addition of an internal quencher into an oligonucleotide.

## Example 4

This example demonstrates the functional performance through quantitative real time PCR (q-PCR) using probes containing the novel internal quenchers disclosed herein.

Validated qPCR assays were used to assess the performance of the different designs of fluorescence quenched probes. The primer sequences employed were specific for the human HPRT gene (NM 00194); probe and primer sequences and are listed below in Table 6.

## 26

TABLE 6

Sequences used in q-PCR assays		
SEQ ID No.	Sequence Name	Sequence
97	HPRT Forward	GACTTTGCTTTCCTTGGTGAG
98	HPRT Reverse	GGCTTATATCCAACACTTCGTG
99	HPRT Probe	ATGGTCAAGGTCGCAAGCTTGCTGGT

All oligonucleotides were synthesized by IDT (Integrated DNA Technologies, Coralville, Iowa). Probe oligonucleotides were HPLC purified. Mass identity of all oligonucleotides was verified by mass spectrometry. All target amplicons were cloned and sequence verified. The target plasmids

were linearized by restriction endonuclease digestion and 10 fold serial dilutions were performed to create standard curves.

HPRT q-PCR reactions were comprised of 0.4 U Immolase DNA Polymerase (Bioline, Taunton, Mass.), 0.8 mM dNTP mix, 3 mM MgCl<sub>2</sub>, and primer/probe concentrations at 200 nM each. Q-PCR reactions were performed on a Roche Lightcycler® 480 platform. Plasmid copy number standards were run in triplicate starting with 2×10<sup>7</sup> down to 2×10<sup>2</sup> (10 fold increments). The thermocycling profile used was 95<sup>10:00</sup>–(95<sup>0:15</sup>–60<sup>1:00</sup>)×40. Data analysis was performed using software supplied by the manufacturer.

Effects of Placement of Internal FQ (iFQ).

A series of oligonucleotide probes having a 5'-FAM reporter dye and an internal iFQ quencher were synthesized using the HPRT probe sequence varying the relative place-



ment of the iFQ group. In one set the iFQ group was placed as a base substitution (replacing a base within the sequence). In another set the iFQ group was placed as a base insertion between residues. The distance of the iFQ from the 5'-dye was varied including 6, 8, 10, or 12 positions from the 5'-end. Henceforth, a modification placed as an insertion at position 6 is indicated as an "i6" and as a substitution at position 8 is indicated as a "s8", etc. The probes were used in qPCR as outlined above using the HPRT primers (SEQ ID Nos. 97 and 98) and  $2 \times 10^6$  copies of a cloned HPRT amplicon plasmid target. Amplification plots for the substitution series probes (SEQ ID Nos. 100-103) are shown in FIG. 2A and baseline normalized plots are shown in FIG. 2B. Amplification plots for the insertion series (SEQ ID Nos. 104-107) are shown in FIG. 3A and baseline normalized plots are shown in FIG. 3B. It is clear that precise placement of the iFQ group within the probe affected probe characteristics, with changes seen in both the baseline fluorescence, magnitude of signal generation, and quantification cycle number (Cq, the cycle number where amplification signal is first detected).

Relative metrics for assessing probe quality from the amplification plots are reported in Table 7, including the baseline fluorescence measured in the qPCR device (as a measure of quenching efficiency) and the  $\Delta R_n$ , the difference in the fluorescence intensity between the start and end of the qPCR run (as a measure of the magnitude of signal generated). Having low baseline fluorescence coupled with high relative  $\Delta R_n$  signal generation leads to improved probe performance.

TABLE 7

Internal placement of iFQ in FAM labeled probes (HPRT assay)			
SEQ ID No.	Sequence (Substitution)	Baseline fluorescence	$\Delta R_n$
100	FAM-ATGGTCAAGGT/iFQ/GCAAGCTTGCTGGT-SpC3 (s12)	3.2	8.0
101	FAM-ATGGTCAAG/iFQ/TCGCAAGCTTGCTGGT-SpC3 (s10)	2.2	7.8
102	FAM-ATGGTCA/iFQ/GGTCGCAAGCTTGCTGGT-SpC3 (s8)	1.7	7.0
103	FAM-ATGGT/iFQ/AAGGTCGCAAGCTTGCTGGT-SpC3 (s6)	1.1	3.3
Sequence (Insertion)			
104	FAM-ATGGTCAAGGT/iFQ/CGCAAGCTTGCTGGT-SpC3 (i12)	6.0	11.6
105	FAM-ATGGTCAAG/iFQ/GTCGCAAGCTTGCTGGT-SpC3 (i10)	4.0	13.7
106	FAM-ATGGTCA/iFQ/AGGTCGCAAGCTTGCTGGT-SpC3 (i8)	2.0	12.5
107	FAM-ATGGT/iFQ/CAAGGTCGCAAGCTTGCTGGT-SpC3 (i6)	1.7	6.8

All probe designs functioned in the above qPCR assay, thus the iFQ group can be used in the methods of this disclosure as either a base substitution or as an insertion between bases. Placing the iFQ group as a base substitution resulted in slightly better quenching than when used as an insertion between bases, however as a group the substitution probes showed lower signal generation. Further, iFQ placed as a base substitution is slightly destabilizing (lowers  $T_m$ ) whereas iFQ placed as a base insertion is stabilizing (increases  $T_m$ ) (see Example 1 above). Given the better signal generation and improved duplex stabilization properties, placement of the iFQ as an insertion is considered to be a more preferred embodiment and all subsequent examples will be done using only iFQ insertion probes.

The relative distance of the iFQ group from the 5'-FAM reporter dye had a significant impact on baseline fluorescence and on signal generation. In FIG. 3A, background fluorescence levels were i6<i8<i10<i12. Placing the quencher group closer to the reporter dye reduced background fluorescence. This effect was expected as FRET based quenching improves with the proximity of the reporter and quencher. Unexpectedly, the relative placement of the iFQ group also affected the magnitude of signal generation during an amplification run and final functional fluorescence (positive signal) was i10>i8>i12>>i6 (FIG. 3B). Interestingly, the i6 probes showed poor fluorescence signal for both the insertion and substitution series. This reduced signal compromised assay quality and delayed the Cq point for these probes, indicating lower assay sensitivity. We hypothesize that placing the iFQ group too close to the fluorophore results in probes that are not fully cleaved by the DNA polymerase during amplification (5'-nuclease assay format), resulting in decreased release of reporter dye from quencher. Peak probe performance is realized at a point which is a compromise between quenching, which improves as quencher and fluorophore are placed more closely, and cleavage, which improves as quencher and fluorophore are separated by a greater number of nucleic acid bases. The precise range where this relationship is optimal is non-obvious and may be different for different reporter dye/quencher combinations. For this reporter dye/quencher combination, optimum performance of the assay was seen with i8 and i10 placement. For these probes, background fluorescence was low and signal generation was high.

## Example 5

The following example demonstrates the efficacy of the internal quencher probe design at varying target concentrations.

Example 4 examined the performance of 8 different probe designs using a single concentration of target nucleic acid. In the present example, six concentrations of target were tested in HPRT qPCR assays comparing performance of four of the probes from Example 4. HPRT specific probes using internal FQ (iFQ) quencher with FAM reporter were employed including substitution design s6 and s10 (SEQ ID Nos. 103 and 101) and insertion design i6 and i10 (SEQ ID Nos. 107 and 105). Amplification reactions were run as outlined in


Example 3 using input target plasmid copy numbers of  $2 \times 10^2$ ,  $2 \times 10^3$ ,  $2 \times 10^4$ ,  $2 \times 10^5$ ,  $2 \times 10^6$  and  $2 \times 10^7$ . Each reaction was run in triplicate.

FIG. 4A shows the results of the baseline adjusted substitution set and FIG. 4B shows the results for the baseline adjusted insertion set. There is a clear progression of the amplification plot curves corresponding to the expected difference of  $\sim 3.3$  cycles for every 10-fold change of input target nucleic acid, wherein the curves align from left to right the highest concentration of template to the lowest concentration. The insertion probe set outperformed the substitution probe set at all concentrations tested as evidenced by the improved magnitude of signal generated for all comparable data points. Further, the quencher i10 placement outperformed the quencher i6 placement series. Subsequent examples will therefore focus on use of the substitution i10 probe design.

#### Example 6

The following example demonstrates the use of an inserted internal quencher coupled with a second quencher positioned at the 3'-end of the probe.

A new series of probes were synthesized targeting the HPRT gene all having a 5'-fluorescein reporter dye (6-FAM) and having quenchers located at varying positions, including compositions having two quencher groups in the same probe molecule. Internal quenchers were added as insertions between bases. Single quencher probes (FQ quencher on the 3'-terminus, FQ quencher at the i10 position) were compared with dual quencher version of the same sequences. Dual quencher probes were made using the FQ chemical group or the Black Hole Quencher<sup>TM</sup>-1 (BHQ1, shown below), a different commercially available dark quencher (Biosearch Technologies, Novato, Calif.). Table 8 lists the probe sequences tested. HPRT qPCR assays were performed as described in Example 4, using  $2 \times 10^6$  copies of an HPRT amplicon-containing plasmid as target.



BHQ1

TABLE 8

Single and Dual Quenched Probes				
SEQ ID No.	Probe Name	Sequence	Baseline fluorescence	$\Delta R_n$
105	i10FQ	FAM-ATGGTCAAG/iFQ/GTCGCAAGCTTGCTGGT-SpC3	4.0	12.7
108	3' FQ	FAM-ATGGTCAAGGTCGCAAGCTTGCTGGT-FQ	10.5	11.0
109	i10FQ +3' FQ	FAM-ATGGTCAAG/iFQ/GTCGCAAGCTTGCTGGT-FQ	2.6	14.0
110	i10BHQ +3' BHQ	FAM-ATGGTCAAG/iBHQ1/GTCGCAAGCTTGCTGGT-BHQ1	2.5	7.5

These four probes were used in qPCR as outlined in Example 4 above using the HPRT primers (SEQ ID Nos. 97 and 98) and  $2 \times 10^6$  copies of a cloned HPRT amplicon plasmid target. Amplification plots are shown in FIG. 5A and baseline normalized plots are shown in FIG. 5B. Relative metrics for assessing probe quality from the amplification plots are also reported in Table 8, including the baseline fluorescence measured in the qPCR device (as a measure of quenching efficiency) and the  $\Delta R_n$  seen between the start and end of the qPCR run (as a measure of the magnitude of signal generated). Having low baseline fluorescence coupled with high relative  $\Delta R_n$  signal generation leads to improved probe performance.

The traditional probe design with a 5'-reporter dye and 3'-quencher (SEQ ID No. 108) showed the highest baseline fluorescence (i.e., worst quenching). The single quencher internal i10 placement probe (SEQ ID No. 105) showed significantly lower baseline fluorescence and the dual-quencher probes (SEQ ID Nos. 109, 110) showed the lowest baseline fluorescence (i.e., best quenching). Thus the dual quencher probes in FIGS. 5A and 5B showed slightly better performance than the i10FQ quencher alone, demonstrating that the presence of two quenchers not only does not negatively affect their quenching properties but rather improves the overall quenching properties.

The relative fluorescence signal generated using the different probe designs also varied with quencher type and placement. The three FQ probes generated a  $\Delta R_n$  of 11-14 with the highest relative fluorescent signal produced by the dual-quencher i10FQ-3'-FQ probe (SEQ ID No. 109). Interestingly, the same design using the alternative commercial dark quencher BHQ-1 (SEQ ID No. 110) performed significantly worse, showing a  $\Delta R_n$  of only 7.5. This probe also showed a delayed  $C_q$  value (the cycle number where amplification signal is first detected), indicating worse functional sensitivity. This clearly demonstrates that all dark quencher chemical

## 31

compositions are not functionally interchangeable and that the naphthylene-azo quencher (FQ) performs in a superior fashion using the methods of this disclosure.

## Example 7

Examples 4, 5, and 6 were performed using a single probe sequence specific for the human HPRT gene. The following example illustrates that the functional qPCR results detailed in Examples 4-6 are consistent when run using a different

## 32

plots are also reported in Table 10, including the baseline fluorescence measured in the qPCR device (as a measure of quenching efficiency) and the  $\Delta R_n$  seen between the start and end of the qPCR run (as a measure of the magnitude of signal generated). Note that the numbers reported are “relative fluorescence units” and that this example was performed on a different machine than the plots shown in Examples 4-6 above. The absolute numbers are different between machines; however the relative performance of the various probe designs is directly comparable.

TABLE 10

Single and dual quenched influenza SW H1 assays				
SEQ ID No.	Name	Sequence	Baseline	$\Delta R_n$
114	3'FQ	FAM-CAGAATATACATCCAGTCACAATTGGAAAA-FQ	2.6	1.5
115	i10FQ	FAM-CAGAATATA/iFQ/CATCCAGTCACAATTGGAAAA-SpC3	0.7	2.4
116	iF10Q +3'FQ	FAM-CAGAATATA/iFQ/CATCCAGTCACAATTGGAAAA-FQ	0.6	2.9
117	i10BHQ1 +3'BHQ1	FAM-CAGAATATA/iBHQ1/CATCCAGTCACAATTGGAAAA-BHQ1	0.45	0.8

probe sequence, for a different target gene, using a different thermal cycler platform and different reagents.

A new qPCR assay was employed specific for a strain of the H1N1 Influenza virus (SW H1, also known as the “swine flu”). Primer and probe sequences and are listed below in Table 9.

TABLE 9

Sequences used in an Influenza qPCR assay			
SEQ ID No.	Sequence Name	Sequence	
111	SW H1 Forward	GTGCTATAAACACCAGCCTYCCA	
112	SW H1 Reverse	CGGGATATTCCTTAATCCTGTRGC	
113	SW H1 Probe	CAGAATATACATCCAGTCACAATTGGAAAA	

A set of Influenza virus H1N1 specific (SW H1) probes were synthesized and qPCR assays were performed using the same methods as described in Example 4 except that the assays were run on an Applied Biosystems 7900HT Sequence Detection System (Applied Biosystems, Foster City, Calif.) according to manufacturer’s instructions, using 1x TaqMan Gene Expression Master Mix (Life Technologies, Carlsbad, Calif.), 250 nM probe and 1000 nM primers per the Center for Disease Control (CDC) document (CEC REF#1-007-005 Protocol for Detection and Characterization of Swine Influenza, 2009) recommendations. Assays run using different probe sequences and targets in the present example demonstrate that the performance of the probes of this disclosure are not sequence dependent, instrument dependent or polymerase formulation dependent.

The assays were run in duplicate with  $2 \times 10^6$  copies of plasmid target used. Table 10 lists the probe sequences tested. All internal quencher groups were placed as insertions. Relative metrics for assessing probe quality from the amplification

Amplification plots for the influenza qPCR assays are shown in FIG. 6A and baseline normalized amplification plots are shown in FIG. 6B. Similar to the results described in Example 6 for the HPRT assays, the probe containing the 3'FQ quencher alone did not perform as well as any of the internal FQ containing probes. In particular, the internal-quencher probes showed markedly lower baseline (background) fluorescence, with the dual-quencher probes having the lowest background. As before, the dual quencher probes outperformed the single quencher probes, with the iFQ+3'FQ combination working the best. The SWH1 probe is 30 bases long and represents a relatively long probe sequence for use as a dual-quenched probe in hydrolysis assays, which contributes to the greater difference seen between 3'-quencher probes and internal quencher probes in this example compared with the HPRT probes in Example 6 (the HPRT probe is 26 bases long). This illustrates another benefit of the methods of this disclosure. Long probes (as are frequently required in AT-rich target sequences) perform poorly using 3'-quencher design; however probe length does not affect performance of internally quenched probes.

Similar to the results observed in Example 6 above, the dual-quencher i10FQ+3'FQ probe performed the best of the set tested, showing both low baseline fluorescence and high positive signal strength. Of note, again in the influenza probe sequence context the dual-quencher i10BHQ1+3'BHQ1 probe showed low baseline fluorescence with very low signal strength and functioned poorly in the assay, showing a delayed Cq value relative to the other probes.

## Example 8

The following example demonstrates the efficacy of the internal quencher probes of the present disclosure when used with various fluorophores.

In previous examples, all probes contained a 5'-fluorescein reporter dye (6-FAM). Typically, quencher molecules perform well with a limited subset of reporter dyes that are matched such that the fluorophore fluorescence emission

wavelength overlaps well with the absorbance wavelengths of the quencher. A dye with emission in the red region, such as Cy5, typically requires use of a different quencher than one that is useful with a dye that has a shorter wavelength emission spectra, such as fluorescein. The probes in this example comprise different fluorophores having a wide range of emission wavelengths (Table 10) and demonstrate that the use of internal quencher probes in the dual-quencher format function well across a wide spectral range. All probes in the present example place the iFQ quencher at position i10 as an insertion. The dual-quencher probes of the present example are made using the i10FQ combined with either a 3'-FQ or with a 3'-IB RQ-n1 (also referred to as "RQ"). In traditional single 3'-quencher probe format, the FQ quencher (which has a peak absorbance around 534 nm) is typically employed with reporter dyes having emission in the 500-580 nm wavelength range. The RQ quencher (which has a bimodal peak absorbance around 610 and 640 nm) is typically employed with reporter dyes having emission in the 550-700 nm wavelength range.

TABLE 10

Fluorescent reporter dyes and their excitation and emission wavelengths.		
Dye	Excitation Max (nm)	Emission Max (nm)
6-FAM	495	520
MAX	524	557
Cy3	550	564
TEX-615	596	615
Cy5	648	668

A series of dual quencher probes specific for the human HPRT gene were synthesized having an insertion of the FQ quencher at the i10 position and either a 3'-FQ or a 3'-RQ quencher. Probe sequences are shown in Table 11 below. The probes were used in qPCR as described in Example 4 above using the HPRT primers (SEQ ID Nos. 97 and 98) and  $2 \times 10^6$  copies of a cloned HPRT amplicon plasmid target. The FAM and MAX probes were run using the Applied Biosystems AB7900HT Sequence Detection platform. The Cy3 probes were run using the BIO-RAD iQ5 platform. The TEX615 and Cy5 probes were run using the Roche LightCycler 480 platform.

Examples 6 and 7 demonstrated that the dual-quencher iFQ-3'FQ combination performed slightly better for FAM reporter dye probes than iFQ alone, although the single

quencher iFQ probes also performed well. Amplification plots for the fluorescein (emission 520 nm) reporter dye probes (SEQ ID Nos. 109 and 118) are shown in FIG. 7A and baseline normalized plots are shown in FIG. 7B. In this comparison, performance is nearly identical for the i10FQ-3'FQ vs. i10FQ-3'RQ probes, even though the RQ quencher is best suited for red wavelength dyes. The i10FQ-3'FQ probe did, however, perform slightly better.

Amplification plots for the MAX (emission 557 nm) reporter dye probes (SEQ ID Nos. 119 and 120) are shown in FIG. 8A and baseline normalized plots are shown in FIG. 8B. Baseline quenching was nearly identical for the i10FQ-3'FQ vs. i10FQ-3'RQ probes, however peak signal intensity was superior with the i10FQ-3'FQ probe. Both designs worked well but the i10FQ-3'FQ design is preferred.

Amplification plots for the Cy3 (emission 564 nm) reporter dye probes (SEQ ID Nos. 121 and 122) are shown in FIG. 9A and baseline normalized plots are shown in FIG. 9B. Baseline quenching was slightly lower for the i10FQ-3'FQ vs. i10FQ-3'RQ probes, and peak signal intensity was also slightly superior with the i10FQ-3'FQ probe. Both designs worked well but the i10FQ-3'FQ design is preferred.

Amplification plots for the TEX615 (emission 615 nm) reporter dye probes (SEQ ID Nos. 123 and 124) are shown in FIG. 10A and baseline normalized plots are shown in FIG. 10B. Baseline quenching was slightly lower for the i10FQ-3'RQ vs. i10FQ-3'FQ probes, however peak signal intensity was superior with the i10FQ-3'FQ probe. Both designs worked well but the i10FQ-3'FQ design is preferred.

Amplification plots for the Cy5 (emission 668 nm) reporter dye probes (SEQ ID Nos. 125 and 126) are shown in FIG. 11A and baseline normalized plots are shown in FIG. 11B. Baseline quenching was lower for the i10FQ-3'RQ vs. i10FQ-3'FQ probes; however peak signal intensity was identical for both designs. Both designs worked well but the i10FQ-3'RQ design showed lower baseline fluorescence.

The relative metrics for assessing probe quality from the amplification plots are also reported in Table 12, including the baseline fluorescence measured in the qPCR device (as a measure of quenching efficiency) and the  $\Delta R_n$  seen between the start and end of the qPCR run (as a measure of the magnitude of signal generated). Having low baseline fluorescence coupled with high relative  $\Delta R_n$  signal generation leads to improved probe performance. Note that different real-time PCR machines were employed for different probe pairs (see above) and that the arbitrary fluorescence units reporting fluorescence intensity vary between platforms.

TABLE 12

Dual-quencher probes with various reporter dyes and an internal i10FQ with either 3'-FQ or 3'-RQ				
SEQ ID No.	Name	Sequence	Baseline fluorescence	$\Delta R_n$
109	FAM i10FQ-3'FQ	FAM-ATGGTCAAG/iFQ/GTCGCAAGCTTGCTGGT-FQ	2.9	14
118	FAM i10FQ-3'RQ	FAM-ATGGTCAAG/iFQ/GTCGCAAGCTTGCTGGT-RQ	3.1	13.8
119	MAX i10FQ-3'FQ	MAX-ATGGTCAAG/iFQ/GTCGCAAGCTTGCTGGT-FQ	0.2	2.1
120	MAX i10FQ-3'RQ	MAX-ATGGTCAAG/iFQ/GTCGCAAGCTTGCTGGT-RQ	0.2	1.9
121	Cy3 i10FQ-3'FQ	Cy3-ATGGTCAAG/iFQ/GTCGCAAGCTTGCTGGT-FQ	350	1500

TABLE 12-continued

Dual-quencher probes with various reporter dyes and an internal i10FQ with either 3'-FQ or 3'-RQ				
SEQ ID No.	Name	Sequence	Baseline fluorescence	$\Delta R_a$
122	Cy3 i10FQ-3'RQ	Cy3-ATGGTCAAG/iFQ/GTCGCAAGCTTGCTGGT-RQ	250	1300
123	TEX i10FQ-3'FQ	TEX-ATGGTCAAG/iFQ/GTCGCAAGCTTGCTGGT-FQ	0.8	3.4
124	TEX i10FQ-3'RQ	TEX-ATGGTCAAG/iFQ/GTCGCAAGCTTGCTGGT-RQ	0.7	2.4
125	Cy5 i10FQ-3'FQ	Cy5-ATGGTCAAG/iFQ/GTCGCAAGCTTGCTGGT-FQ	0.4	1.5
126	Cy5 i10FQ-3'RQ	Cy5-ATGGTCAAG/iFQ/GTCGCAAGCTTGCTGGT-RQ	0.2	1.5

The use of an internal quencher paired with a 3'-quencher in the above dual-quencher probes enables use of a single quencher compound (FQ) in probes with reporter dyes ranging from FAM to Cy5. This novel format thus permits use of a quencher with dyes that it is not suitable to work with when used in the traditional 3'-quencher alone probe design format.

#### Example 9

The following example demonstrates that placement of the internal iFQ insertion can vary with the Cy5 reporter dye.

The probes studied in Example 8 all employed the FQ quencher placed as an insertion at position i10. This example compares function of a dual-quenched Cy5 probes having i10FQ vs. i12FQ.

Probe sequences are shown in Table 13 below. The probes were used in qPCR as described in Example 4 above using the HPRT primers (SEQ ID Nos. 97 and 98) and  $2 \times 10^7$  to  $2 \times 10^2$  copies of a cloned HPRT amplicon plasmid target. The Cy5 probes were run using the Roche LightCycler 480 platform.

TABLE 13

Dual-quencher probes with Cy5 reporter dye comparing i10FQ vs. i12FQ		
SEQ ID No.	Name	Sequence
125	Cy5 i10FQ-3'FQ	Cy5-ATGGTCAAG/iFQ/GTCGCAAGCTTGCTGGT-FQ
127	Cy5 i12FQ-3'FQ	Cy5-ATGGTCAAGGT/iFQ/CGCAAGCTTGCTGGT-RQ

The baseline adjusted amplification plot showing superimposed traces of all 6 dilution curves for both probes is shown in FIG. 12. Peak signal intensity was slightly superior for the i10FQ-3'FQ probe than for the i12FQ-3'FQ probe, however both probes performed equally well in quantitative detection of the target standard curve dilution set. Precise Cq values measured for each probe/target dilution are shown in Table 14 below, where it can be seen that the Cq values are nearly identical between the two probes.

TABLE 14

Cq values comparing sensitivity of i10FQ-3'FQ vs. i12FQ-3'FQ for an HPRT assay using Cy5 reporter dye		
Copy Number	CY5 i10FQ-3'FQ	CY5 i12FQ-3'FQ
2E7	14.2	14.4
2E6	17.5	17.4
2E5	20.9	21.0
2E4	24.8	24.6
2E3	28.0	27.9
2E2	31.6	32.4

Although there may be range in location for placement of the iFQ insertion for different probes, the preferred location may vary with different reporter dyes. More importantly, there exists some flexibility in placement such that iFQ group place anywhere in the i8-i12 range should work well.

#### Example 10


This example demonstrates the synthesis of a phosphoramidite useful for oligonucleotide synthesis that is derivatized with a quencher of the present disclosure. The synthetic method is shown below in Schemes 1 and 2.

#### Mono-DMT-phenyl diethanolamine (2)

A solution of 10 g of phenyl diethanolamine in 100 mL of pyridine was mixed for 3-4 h at room temperature with a solution of 6 g dimethoxytrityl-chloride (DMT-C1) in 150 mL of a 98:2 dichloromethane/pyridine solution. The reaction mixture was concentrated to dryness under vacuum. The residue was dissolved in 200 mL of ethyl acetate, washed with two portions of 100 mL of deionized water, and the organic layer dried over  $\text{Na}_2\text{SO}_4$ . The organic solution was concentrated and purified by column chromatography using a 300 g of silica gel column developed with 30/65/5 ethyl acetate/hexanes/triethylamine to yield 5.25 g (20% yield) of mono-DMT-phenyl diethanolamine. TLC:  $R_f$  0.55 (EtAc/hexanes/ $\text{Et}_3\text{N}$ —40/55/5).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  7.38 (d, J=8 Hz, 2H), 7.27 (d, J=8 Hz, 4H), 7.38 (d, J=8 Hz, 2H), 7.24-7.12 (m, 6H), 6.76 (d, J=8 Hz, 4H), 6.66 (d, J=8 Hz, 2H), 3.74 (s, 6H), 3.74 (t, J=7.5 Hz, 2H), 3.54 (t, J=7.5 Hz, 2H), 3.51 (t, J=7.5 Hz, 2H), 3.33 (t, J=7.5 Hz, 2H), 2.23 (br. s, 1H).

37

Scheme 1




Mono-DMT-4-(1-nitro-4-naphthylazo)-N,N-diethanolamine (3)

Cold concentrated HCl (17 mL) was added dropwise at 0° C. over 15 min to a suspension of 4-nitro-1-naphthylamine (2 g) in cold water (6 mL) at 0° C. Then NaNO<sub>2</sub> (1.6 g) in cold water (4 mL) was added dropwise at 0° C. over 15 min and the 4-nitro-1-naphthylamine dissolved upon stirring. LiBF<sub>4</sub> (1.38 g) in H<sub>2</sub>O (3 mL) was added dropwise at 0° C. The reaction mixture was stirred at 0° C. for 30 min. A brownish yellow powder (3.08 g) of naphthyl-1-nitro-4-tetrafluoroborate azonium salt (1) was obtained after filtering and rinsing the solution with cold water, methanol, and ether. A solution of 4 g of mono-DMT-phenyl diethanolamine (2) in 50 mL of dimethylsulfoxide (DMSO) was added with stirring at 10-15° C. over 10-15 min to a chilled solution of 2.8 g of azonium salt (1) in 50 mL of DMSO at 10-15° C. in a water bath. After an additional 15 min of stirring, 3 mL of triethylamine was added to the reaction mixture followed by 100 mL of ethyl acetate. The reaction mixture was washed with 3×30 mL of deionized water and the organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>. The solvent was removed and product was purified by column chromatography with 300 g of silica gel to provide 1.8 g of

38

mono-DMT-4-(1-nitro-4-naphthylazo)-NN-diethanolamine (3). TLC: R<sub>f</sub> 0.65 (DCM/Et<sub>3</sub>N—80/20). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 9.04 (d, J=8.4 Hz, 1H), 8.68 (d, J=8.4 Hz, 1H), 8.34 (d, J=8.4 Hz, 1H), 7.96 (d, J=9.2 Hz, 2H), 7.81-7.71 (m, 3H), 7.39 (d, J=8 Hz, 2H), 7.27 (d, J=8 Hz, 4H), 7.24-7.19 (m, 3H), 6.78 (d, J=8 Hz, 4H), 6.77 (d, J=8 Hz, 2H), 3.88 (t, J=7.5 Hz, 2H), 3.75 (s, 6H), 3.78-3.68 (m, 4H), 3.47 (t, J=7.5 Hz, 2H), 1.57 (br. s, 1H).

10



Mono-DMT-4-(1-nitro-4-naphthylazo)-N,N-diethanolamine phosphoramidite (4)

A solution of 0.2 ml of N,N-diisopropylamino-cyanoethylphosphoramidolchloride was stirred into a solution of 0.3 g of alcohol (3) in 20 mL of anhydrous THF and 1 mL of triethylamine for 5 min at 0-5° C. After 15 min of additional stirring the reaction mixture was warmed to room temperature. The solvent was evaporated under a vacuum and the residue purified by column chromatography through 50 g of silica gel (EtOAc/PE/TEA: 10/85/5-40/55/5). TLC: R<sub>f</sub> 0.65 (DCM/Et<sub>3</sub>N—80/20). <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 9.05 (d, J=8.4 Hz, 1H),

8.68 (d, J=8.4 Hz, 1H), 8.34 (d, J=8.4 Hz, 1H), 7.96 (d, J=9.2 Hz, 2H), 7.81-7.71 (m, 3H), 7.39 (d, J=8 Hz, 2H), 7.27 (d, J=8 Hz, 4H), 7.24-7.19 (m, 3H), 6.78 (d, J=8 Hz, 4H), 6.76 (d, J=8 Hz, 2H), 3.85-3.75 (m, J=7.5 Hz, 4H), 3.76 (s, 6H), 3.70 (t, J=7.5 Hz, 2H), 3.41 (t, J=7.5 Hz, 2H), 2.58 (t, J=8.0 Hz, 2H), 1.20 (s, 3H), 1.18 (s, 3H), 1.17 (s, 3H), 1.15 (s, 3H). <sup>31</sup>P NMR  $\delta$  148.39.

analysis of the fluorescence curves yielded the melting temperature of the beacon alone (stem  $T_m$ ) and the melting temperature of the probe plus complement (loop  $T_m$ ). Each sample independently was measured a minimum of three times.

Table 15 lists the sequences that were synthesized and tested. The "RQ" is IBRQn-1.

TABLE 15

Molecular beacon sequences		
SEQ ID No.	Name	Sequence
128	beacon	CGCGATCAGACAAGGAGTGGGCTTCATGGATCGCG
129	complement	TTACATGAAGCCCACTCCTTGTCTATC
130	3' Dabcyl	/56-FAM/CGCGATCAGACAAGGAGTGGGCTTCATGGATCGCG/3Dabcyl/
131	3' BHQ-1	/56-FAM/CGCGATCAGACAAGGAGTGGGCTTCATGGATCGCG/3BHQ-1/
132	3' FQ	/56-FAM/CGCGATCAGACAAGGAGTGGGCTTCATGGATCGCG/3FQ/
133	FQ0-FQ	/56-FAM/CGCGATCAGACAAGGAGTGGGCTTCATGGATCGCG/FQ/FQ/
134	FQ1-FQ	/56-FAM/CGCGATCAGACAAGGAGTGGGCTTCATGGATCGC/FQ/G/3FQ/
135	FQ2-FQ	/56-FAM/CGCGATCAGACAAGGAGTGGGCTTCATGGATCG/FQ/CG/3FQ/
136	FQ3-FQ	/56-FAM/CGCGATCAGACAAGGAGTGGGCTTCATGGATC/FQ/GCG/3FQ/
137	FQ4-FQ	/56-FAM/CGCGATCAGACAAGGAGTGGGCTTCATGGAT/FQ/CGCG/3FQ/
138	FQ5-FQ	/56-FAM/CGCGATCAGACAAGGAGTGGGCTTCATGGA/FQ/TCGCG/3FQ/
139	FQ6-FQ	/56-FAM/CGCGATCAGACAAGGAGTGGGCTTCATGG/FQ/ATCGCG/3FQ/
140	3' RQSp	/56-FAM/CGCGATCAGACAAGGAGTGGGCTTCATGGATCGCG/3RQSp/
141	FQ0-RQ	/56-FAM/CGCGATCAGACAAGGAGTGGGCTTCATGGATCGCG/FQ//3RQSp/
142	FQ1-RQ	/56-FAM/CGCGATCAGACAAGGAGTGGGCTTCATGGATCGC/FQ/G/3RQSp/
143	FQ2-RQ	/56-FAM/CGCGATCAGACAAGGAGTGGGCTTCATGGATCG/FQ/CG/3RQSp/
144	FQ3-RQ	/56-FAM/CGCGATCAGACAAGGAGTGGGCTTCATGGATC/FQ/GCG/3RQSp/
145	FQ4-RQ	/56-FAM/CGCGATCAGACAAGGAGTGGGCTTCATGGAT/FQ/CGCG/3RQSp/
146	FQ5-RQ	/56-FAM/CGCGATCAGACAAGGAGTGGGCTTCATGGA/FQ/TCGCG/3RQSp/
147	FQ6-RQ	/56-FAM/CGCGATCAGACAAGGAGTGGGCTTCATGG/FQ/ATCGCG/3RQSp/

The phosphoramidite (4) can be added to an oligonucleotide during synthesis using standard phosphoramidite oligonucleotide synthetic techniques.

#### Example 11

The following example demonstrates the utility of the disclosed insertions in a molecular beacon probe.

All Molecular Beacon oligonucleotides were synthesized and purified by Integrated DNA Technologies, Inc. Molecular beacon oligonucleotides at 200 nM were incubated in a buffer consisting of 16.0 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 67.0 mM Tris-HCl pH 8.3, 0.01% Tween-20 in the presence and absence of a 1000 nM complementary oligonucleotide. The samples were incubated at 30° C. in a CFX884 Real Time System (BioRad, Hercules, Calif.), and subjected to an increase in temperature from 30-95° C. at a ramp rate of 1° C./min, with fluorescence measurements taken at each degree interval. First derivative

Table 16 lists the results of the assay for each beacon.

TABLE 16

Quencher	Stem $T_m$	Stem $\Delta T_m$	Loop $T_m$	Loop $\Delta T_m$
3' Dabcyl	66		63	
3' BHQ-1	70		62.7	
3' FQ	70	—	63	—
FQ0-FQ	72	2	62	-1
FQ1-FQ	68	-2	63	0
FQ2-FQ	70	0	62	-1
FQ3-FQ	71	1	62	-1
FQ4-FQ	72	2	62	-1
FQ5-FQ	74	4	63	0
FQ6-FQ	75.3	5.3	64	1
3' RQSp	74	—	61	—
FQ0-RQ	74	0	63	2
FQ1-RQ	70	-4	62	1
FQ2-RQ	71	-3	61	0
FQ3-RQ	71	-3	61	0

TABLE 16-continued

Quencher	Stem $T_m$	Stem $\Delta T_m$	Loop $T_m$	Loop $\Delta T_m$
FQ4-RQ	73	-1	62	1
FQ5-RQ	76	2	62	1
FQ6-RQ	77	3	61.3	0.3

The stability is improved when the FQ quencher is placed internally close to the 3' quencher, particularly with a 3' RQ quencher.

All references, including publications, patent applications, and patents, cited herein are hereby incorporated by reference to the same extent as if each reference were individually and specifically indicated to be incorporated by reference and were set forth in its entirety herein.

The use of the terms "a" and "an" and "the" and similar referents in the context of this disclosure (especially in the context of the following claims) are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context. The terms "comprising," "having," "including," and "containing" are to be construed as open-ended terms (i.e., meaning "including, but not limited to,") unless otherwise noted. Recitation of ranges of values herein are merely intended to serve as a shorthand method of referring individually to each separate value falling within the range, unless otherwise indicated herein, and each

separate value is incorporated into the specification as if it were individually recited herein. All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g., "such as") provided herein, is intended merely to better illuminate the various inventions disclosed herein and does not pose a limitation on the scope of any inventions unless otherwise claimed. No language in the specification should be construed as indicating any non-claimed element as essential to the practice of the invention.

Preferred embodiments of various inventions are described herein, including the best mode known to the inventors for carrying out the various inventions. Variations of those preferred embodiments may become apparent to those of ordinary skill in the art upon reading the foregoing description. The inventors expect skilled artisans to employ such variations as appropriate, and the inventors intend for the various inventions to be practiced otherwise than as specifically described herein. Accordingly, this disclosure includes all modifications and equivalents of the subject matter recited in the claims appended hereto as permitted by applicable law. Moreover, any combination of the above-described elements in all possible variations thereof is encompassed by the disclosure unless otherwise indicated herein or otherwise clearly contradicted by context.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 147

<210> SEQ ID NO 1  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 1

atcgttgcta

10

<210> SEQ ID NO 2  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 2

tagcaacgat

10

<210> SEQ ID NO 3  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (3)..(4)  
 <223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 3 and 4.

<400> SEQUENCE: 3

atcgttgcta

10

<210> SEQ ID NO 4



-continued

---

<211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (4)..(5)  
 <223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 4  
 and 5.

<400> SEQUENCE: 4

atcggtgcta 10

<210> SEQ ID NO 5  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (5)..(6)  
 <223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 5  
 and 6.

<400> SEQUENCE: 5

atcggtgcta 10

<210> SEQ ID NO 6  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 6

cttggatcgt tgctagtagg 20

<210> SEQ ID NO 7  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 7

cctactagca acgatccaag 20

<210> SEQ ID NO 8  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (10)..(11)  
 <223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 10  
 and 11.

<400> SEQUENCE: 8

cttggatcgt tgctagtagg 20

<210> SEQ ID NO 9  
 <211> LENGTH: 25  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:

-continued

---

<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 9

cacttggatc gttgctagta gggtc 25

<210> SEQ ID NO 10  
 <211> LENGTH: 25  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 10

gaccctacta gcaacgatcc aagtg 25

<210> SEQ ID NO 11  
 <211> LENGTH: 25  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (10)..(11)  
 <223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 10  
 and 11.

<400> SEQUENCE: 11

cacttggatc gttgctagta gggtc 25

<210> SEQ ID NO 12  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (3)..(4)  
 <223> OTHER INFORMATION: iSpC3 is inserted between nucleotide positions  
 3 and 4.

<400> SEQUENCE: 12

atcgttgcta 10

<210> SEQ ID NO 13  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (4)..(5)  
 <223> OTHER INFORMATION: iSpC3 is inserted between nucleotide positions  
 4 and 5.

<400> SEQUENCE: 13

atcgttgcta 10

<210> SEQ ID NO 14  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (5)..(6)

-continued

---

<223> OTHER INFORMATION: iSpC3 is inserted between nucleotide positions  
5 and 6.

<400> SEQUENCE: 14

atcgttgcta 10

<210> SEQ ID NO 15

<211> LENGTH: 10

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (5)..(6)

<223> OTHER INFORMATION: iSpC3 is inserted between nucleotide positions  
5 and 6.

<400> SEQUENCE: 15

tagcaacgat 10

<210> SEQ ID NO 16

<211> LENGTH: 11

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 16

tagcaaacga t 11

<210> SEQ ID NO 17

<211> LENGTH: 11

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 17

tagcacacga t 11

<210> SEQ ID NO 18

<211> LENGTH: 11

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 18

tagcagacga t 11

<210> SEQ ID NO 19

<211> LENGTH: 11

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 19

tagcatacga t 11

<210> SEQ ID NO 20

<211> LENGTH: 10

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

-continued

---

<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (3)..(4)  
<223> OTHER INFORMATION: N-MDA is inserted between nucleotide positions  
3 and 4.

<400> SEQUENCE: 20

atcgttgcta 10

<210> SEQ ID NO 21  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (3)..(4)  
<223> OTHER INFORMATION: C2 is inserted between nucleotide positions 3  
and 4.

<400> SEQUENCE: 21

atcgttgcta 10

<210> SEQ ID NO 22  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (3)..(4)  
<223> OTHER INFORMATION: 2,2 DMP is inserted between nucleotide  
positions 3 and 4.

<400> SEQUENCE: 22

atcgttgcta 10

<210> SEQ ID NO 23  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (3)..(4)  
<223> OTHER INFORMATION: C4 is inserted between nucleotide positions 3  
and 4.

<400> SEQUENCE: 23

atcgttgcta 10

<210> SEQ ID NO 24  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (3)..(4)  
<223> OTHER INFORMATION: C5 is inserted between nucleotide positions 3  
and 4.

<400> SEQUENCE: 24

atcgttgcta 10

-continued

---

<210> SEQ ID NO 25  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (3)..(4)  
<223> OTHER INFORMATION: C6 is inserted between nucleotide positions 3  
and 4.

<400> SEQUENCE: 25

atcgttgcta 10

<210> SEQ ID NO 26  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (3)..(4)  
<223> OTHER INFORMATION: C7 is inserted between nucleotide positions 3  
and 4.

<400> SEQUENCE: 26

atcgttgcta 10

<210> SEQ ID NO 27  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (3)..(4)  
<223> OTHER INFORMATION: iSpS9 is inserted between nucleotide positions  
3 and 4.

<400> SEQUENCE: 27

atcgttgcta 10

<210> SEQ ID NO 28  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (3)..(4)  
<223> OTHER INFORMATION: idSp is inserted between nucleotide positions 3  
and 4.

<400> SEQUENCE: 28

atcgttgcta 10

<210> SEQ ID NO 29  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (3)..(4)  
<223> OTHER INFORMATION: iBHQ2 is inserted between nucleotide positions  
3 and 4.

-continued

---

<400> SEQUENCE: 29

atcgttgcta 10

<210> SEQ ID NO 30

<211> LENGTH: 10

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (3)..(4)

<223> OTHER INFORMATION: iRQ-n1 is inserted between nucleotide positions 3 and 4.

<400> SEQUENCE: 30

atcgttgcta 10

<210> SEQ ID NO 31

<211> LENGTH: 10

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (3)..(4)

<223> OTHER INFORMATION: iRQ-n2 is inserted between nucleotide positions 3 and 4.

<400> SEQUENCE: 31

atcgttgcta 10

<210> SEQ ID NO 32

<211> LENGTH: 10

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (3)..(4)

<223> OTHER INFORMATION: iEc is inserted between nucleotide positions 3 and 4.

<400> SEQUENCE: 32

atcgttgcta 10

<210> SEQ ID NO 33

<211> LENGTH: 10

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (3)..(4)

<223> OTHER INFORMATION: IB 1.1 is inserted between nucleotide positions 3 and 4.

<400> SEQUENCE: 33

atcgttgcta 10

<210> SEQ ID NO 34

<211> LENGTH: 10

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

---

<223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (3)..(4)  
 <223> OTHER INFORMATION: NPDA is inserted between nucleotide positions 3  
 and 4.

<400> SEQUENCE: 34

atcgttgcta 10

<210> SEQ ID NO 35  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (4)..(5)  
 <223> OTHER INFORMATION: N-MDA is inserted between nucleotide positions  
 4 and 5.

<400> SEQUENCE: 35

atcgttgcta 10

<210> SEQ ID NO 36  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (4)..(5)  
 <223> OTHER INFORMATION: C2 is inserted between nucleotide positions 4  
 and 5.

<400> SEQUENCE: 36

atcgttgcta 10

<210> SEQ ID NO 37  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (4)..(5)  
 <223> OTHER INFORMATION: 2,2 DMP is inserted between nucleotide  
 positions 4 and 5.

<400> SEQUENCE: 37

atcgttgcta 10

<210> SEQ ID NO 38  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (4)..(5)  
 <223> OTHER INFORMATION: C4 is inserted between nucleotide positions 4  
 and 5.

<400> SEQUENCE: 38

atcgttgcta 10

-continued

---

<210> SEQ ID NO 39  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (4)..(5)  
 <223> OTHER INFORMATION: C5 is inserted between nucleotide positions 4  
 and 5.  
  
 <400> SEQUENCE: 39  
  
 atcgttgcta 10

<210> SEQ ID NO 40  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (4)..(5)  
 <223> OTHER INFORMATION: C6 is inserted between nucleotide positions 4  
 and 5.  
  
 <400> SEQUENCE: 40  
  
 atcgttgcta 10

<210> SEQ ID NO 41  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (4)..(5)  
 <223> OTHER INFORMATION: C7 is inserted between nucleotide positions 4  
 and 5.  
  
 <400> SEQUENCE: 41  
  
 atcgttgcta 10

<210> SEQ ID NO 42  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (4)..(5)  
 <223> OTHER INFORMATION: iSpS9 is inserted between nucleotide positions  
 4 and 5.  
  
 <400> SEQUENCE: 42  
  
 atcgttgcta 10

<210> SEQ ID NO 43  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (4)..(5)  
 <223> OTHER INFORMATION: idSp is inserted between nucleotide positions



-continued

---

4 and 5.

<400> SEQUENCE: 43

atcgttgcta 10

<210> SEQ ID NO 44  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (4)..(5)  
<223> OTHER INFORMATION: iBHQ2 is inserted between nucleotide positions  
4 and 5.

<400> SEQUENCE: 44

atcgttgcta 10

<210> SEQ ID NO 45  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (4)..(5)  
<223> OTHER INFORMATION: iRQ-n1 is inserted between nucleotide positions  
4 and 5.

<400> SEQUENCE: 45

atcgttgcta 10

<210> SEQ ID NO 46  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (4)..(5)  
<223> OTHER INFORMATION: iRQ-n2 is inserted between nucleotide positions  
4 and 5.

<400> SEQUENCE: 46

atcgttgcta 10

<210> SEQ ID NO 47  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (4)..(5)  
<223> OTHER INFORMATION: iEc is inserted between nucleotide positions 4  
and 5.

<400> SEQUENCE: 47

atcgttgcta 10

<210> SEQ ID NO 48  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence

-continued

---

<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (4)..(5)  
<223> OTHER INFORMATION: IB 1.1 is inserted between nucleotide positions  
4 and 5.

<400> SEQUENCE: 48

atcgttgcta 10

<210> SEQ ID NO 49  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (4)..(5)  
<223> OTHER INFORMATION: NPDA is inserted between nucleotide positions 4  
and 5.

<400> SEQUENCE: 49

atcgttgcta 10

<210> SEQ ID NO 50  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (5)..(6)  
<223> OTHER INFORMATION: C2 is inserted between nucleotide positions 5  
and 6.

<400> SEQUENCE: 50

atcgttgcta 10

<210> SEQ ID NO 51  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (5)..(6)  
<223> OTHER INFORMATION: 2,2 DMP is inserted between nucleotide  
positions 5 and 6.

<400> SEQUENCE: 51

atcgttgcta 10

<210> SEQ ID NO 52  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (5)..(6)  
<223> OTHER INFORMATION: C4 is inserted between nucleotide positions 5  
and 6.

<400> SEQUENCE: 52

atcgttgcta 10

-continued

---

<210> SEQ ID NO 53  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (5)..(6)  
<223> OTHER INFORMATION: C5 is inserted between nucleotide positions 5  
and 6.

<400> SEQUENCE: 53

atcgttgcta 10

<210> SEQ ID NO 54  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (5)..(6)  
<223> OTHER INFORMATION: C6 is inserted between nucleotide positions 5  
and 6.

<400> SEQUENCE: 54

atcgttgcta 10

<210> SEQ ID NO 55  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (5)..(6)  
<223> OTHER INFORMATION: C7 is inserted between nucleotide positions 5  
and 6.

<400> SEQUENCE: 55

atcgttgcta 10

<210> SEQ ID NO 56  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (5)..(6)  
<223> OTHER INFORMATION: iSpS9 is inserted between nucleotide positions  
5 and 6.

<400> SEQUENCE: 56

atcgttgcta 10

<210> SEQ ID NO 57  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (5)..(6)

-continued

---

<223> OTHER INFORMATION: idSp is inserted between nucleotide positions 5 and 6.

<400> SEQUENCE: 57

atcgttgcta 10

<210> SEQ ID NO 58

<211> LENGTH: 10

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (5)..(6)

<223> OTHER INFORMATION: iBHQ2 is inserted between nucleotide positions 5 and 6.

<400> SEQUENCE: 58

atcgttgcta 10

<210> SEQ ID NO 59

<211> LENGTH: 10

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (5)..(6)

<223> OTHER INFORMATION: iRQ-n1 is inserted between nucleotide positions 5 and 6.

<400> SEQUENCE: 59

atcgttgcta 10

<210> SEQ ID NO 60

<211> LENGTH: 10

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (5)..(6)

<223> OTHER INFORMATION: iRQ-n2 is inserted between nucleotide positions 5 and 6.

<400> SEQUENCE: 60

atcgttgcta 10

<210> SEQ ID NO 61

<211> LENGTH: 10

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (5)..(6)

<223> OTHER INFORMATION: iEc is inserted between nucleotide positions 5 and 6.

<400> SEQUENCE: 61

atcgttgcta 10

<210> SEQ ID NO 62

<211> LENGTH: 10

<212> TYPE: DNA

-continued

---

<213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (5)..(6)  
 <223> OTHER INFORMATION: IB 1.1 is inserted between nucleotide positions  
 5 and 6.

<400> SEQUENCE: 62

atcgttgcta 10

<210> SEQ ID NO 63  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (5)..(6)  
 <223> OTHER INFORMATION: NPDA is inserted between nucleotide positions 5  
 and 6.

<400> SEQUENCE: 63

atcgttgcta 10

<210> SEQ ID NO 64  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(2)  
 <223> OTHER INFORMATION: N-MDA is inserted between nucleotide positions  
 1 and 2.

<400> SEQUENCE: 64

atcgttgcta 10

<210> SEQ ID NO 65  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(2)  
 <223> OTHER INFORMATION: C2 is inserted between nucleotide positions 1  
 and 2.

<400> SEQUENCE: 65

atcgttgcta 10

<210> SEQ ID NO 66  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(2)  
 <223> OTHER INFORMATION: 2,2 DMP is inserted between nucleotide  
 positions 1 and 2.

<400> SEQUENCE: 66

-continued

---

 atcgttgcta 10

<210> SEQ ID NO 67  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(2)  
 <223> OTHER INFORMATION: iSpC3 is inserted between nucleotide positions  
 1 and 2.  
  
 <400> SEQUENCE: 67

atcgttgcta 10

<210> SEQ ID NO 68  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(2)  
 <223> OTHER INFORMATION: C4 is inserted between nucleotide positions 1  
 and 2.  
  
 <400> SEQUENCE: 68

atcgttgcta 10

<210> SEQ ID NO 69  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(2)  
 <223> OTHER INFORMATION: C5 is inserted between nucleotide positions 1  
 and 2.  
  
 <400> SEQUENCE: 69

atcgttgcta 10

<210> SEQ ID NO 70  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(2)  
 <223> OTHER INFORMATION: C6 is inserted between nucleotide positions 1  
 and 2.  
  
 <400> SEQUENCE: 70

atcgttgcta 10

<210> SEQ ID NO 71  
 <211> LENGTH: 10  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature

-continued

---

<222> LOCATION: (1)..(2)  
<223> OTHER INFORMATION: C7 is inserted between nucleotide positions 1 and 2.

<400> SEQUENCE: 71

atcgttgcta 10

<210> SEQ ID NO 72  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(2)  
<223> OTHER INFORMATION: iSpS9 is inserted between nucleotide positions 1 and 2.

<400> SEQUENCE: 72

atcgttgcta 10

<210> SEQ ID NO 73  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(2)  
<223> OTHER INFORMATION: idSp is inserted between nucleotide positions 1 and 2.

<400> SEQUENCE: 73

atcgttgcta 10

<210> SEQ ID NO 74  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(2)  
<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 1 and 2.

<400> SEQUENCE: 74

atcgttgcta 10

<210> SEQ ID NO 75  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(2)  
<223> OTHER INFORMATION: iBHQ2 is inserted between nucleotide positions 1 and 2.

<400> SEQUENCE: 75

atcgttgcta 10

<210> SEQ ID NO 76  
<211> LENGTH: 10

-continued

---

<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(2)  
<223> OTHER INFORMATION: iRQ-n1 is inserted between nucleotide positions  
1 and 2.

<400> SEQUENCE: 76

atcgttgcta 10

<210> SEQ ID NO 77  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(2)  
<223> OTHER INFORMATION: iRQ-n2 is inserted between nucleotide positions  
1 and 2.

<400> SEQUENCE: 77

atcgttgcta 10

<210> SEQ ID NO 78  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(2)  
<223> OTHER INFORMATION: iEc is inserted between nucleotide positions 1  
and 2.

<400> SEQUENCE: 78

atcgttgcta 10

<210> SEQ ID NO 79  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(2)  
<223> OTHER INFORMATION: IB 1.1 is inserted between nucleotide positions  
1 and 2.

<400> SEQUENCE: 79

atcgttgcta 10

<210> SEQ ID NO 80  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(2)  
<223> OTHER INFORMATION: NPDA is inserted between nucleotide positions 1  
and 2.

<400> SEQUENCE: 80



-continued

---

atcgttgcta 10

<210> SEQ ID NO 81  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (9)..(10)  
<223> OTHER INFORMATION: C2 is inserted between nucleotide positions 9  
and 10.

<400> SEQUENCE: 81

atcgttgcta 10

<210> SEQ ID NO 82  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (9)..(10)  
<223> OTHER INFORMATION: 2,2 DMP is inserted between nucleotide  
positions 9 and 10.

<400> SEQUENCE: 82

atcgttgcta 10

<210> SEQ ID NO 83  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (9)..(10)  
<223> OTHER INFORMATION: iSpC3 is inserted between nucleotide positions  
9 and 10.

<400> SEQUENCE: 83

atcgttgcta 10

<210> SEQ ID NO 84  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (9)..(10)  
<223> OTHER INFORMATION: C4 is inserted between nucleotide positions 9  
and 10.

<400> SEQUENCE: 84

atcgttgcta 10

<210> SEQ ID NO 85  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:

-continued

---

<221> NAME/KEY: misc\_feature  
<222> LOCATION: (9)..(10)  
<223> OTHER INFORMATION: C5 is inserted between nucleotide positions 9 and 10.

<400> SEQUENCE: 85

atcgttgcta 10

<210> SEQ ID NO 86  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (9)..(10)  
<223> OTHER INFORMATION: C6 is inserted between nucleotide positions 9 and 10.

<400> SEQUENCE: 86

atcgttgcta 10

<210> SEQ ID NO 87  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (9)..(10)  
<223> OTHER INFORMATION: C7 is inserted between nucleotide positions 9 and 10.

<400> SEQUENCE: 87

atcgttgcta 10

<210> SEQ ID NO 88  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (9)..(10)  
<223> OTHER INFORMATION: iSpS9 is inserted between nucleotide positions 9 and 10.

<400> SEQUENCE: 88

atcgttgcta 10

<210> SEQ ID NO 89  
<211> LENGTH: 10  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (9)..(10)  
<223> OTHER INFORMATION: idSp is inserted between nucleotide positions 9 and 10.

<400> SEQUENCE: 89

atcgttgcta 10

<210> SEQ ID NO 90

-continued

---

```

<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (9)..(10)
<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 9
and 10.

<400> SEQUENCE: 90

atcgttgcta                                     10

<210> SEQ ID NO 91
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (9)..(10)
<223> OTHER INFORMATION: iBHQ2 is inserted between nucleotide positions
9 and 10.

<400> SEQUENCE: 91

atcgttgcta                                     10

<210> SEQ ID NO 92
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (9)..(10)
<223> OTHER INFORMATION: iRQ-n1 is inserted between nucleotide positions
9 and 10.

<400> SEQUENCE: 92

atcgttgcta                                     10

<210> SEQ ID NO 93
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (9)..(10)
<223> OTHER INFORMATION: iRQ-n2 is inserted between nucleotide positions
9 and 10.

<400> SEQUENCE: 93

atcgttgcta                                     10

<210> SEQ ID NO 94
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (9)..(10)
<223> OTHER INFORMATION: iEc is inserted between nucleotide positions 9
and 10.

```

-continued

---

<400> SEQUENCE: 94

atcgttgcta	10
------------	----

<210> SEQ ID NO 95

<211> LENGTH: 10

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (9)..(10)

<223> OTHER INFORMATION: IB 1.1 is inserted between nucleotide positions 9 and 10.

<400> SEQUENCE: 95

atcgttgcta	10
------------	----

<210> SEQ ID NO 96

<211> LENGTH: 10

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (9)..(10)

<223> OTHER INFORMATION: NPDA is inserted between nucleotide positions 9 and 10.

<400> SEQUENCE: 96

atcgttgcta	10
------------	----

<210> SEQ ID NO 97

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 97

gactttgctt tccttggtca g	21
-------------------------	----

<210> SEQ ID NO 98

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 98

ggcttatatc caacacttcg tg	22
--------------------------	----

<210> SEQ ID NO 99

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 99

atggtcaagg tcgcaagctt gctggt	26
------------------------------	----

<210> SEQ ID NO 100

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (1)..(1)

<223> OTHER INFORMATION: n is dA modified with FAM.

-continued

---

```

<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (11)..(12)
<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 11
and 12.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (25)..(25)
<223> OTHER INFORMATION: n is dT modified with SpC3.

<400> SEQUENCE: 100

```

```

ntgggtcaagg tgcaagcttg ctggn 25

```

```

<210> SEQ ID NO 101
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: n is dA modified with FAM.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (9)..(10)
<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 9
and 10.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (25)..(25)
<223> OTHER INFORMATION: n is dT modified with SpC3.

<400> SEQUENCE: 101

```

```

ntgggtcaagt cgcaagcttg ctggn 25

```

```

<210> SEQ ID NO 102
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: n is dA modified with FAM.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (7)..(8)
<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 7
and 8.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (25)..(25)
<223> OTHER INFORMATION: n is dT modified with SpC3.

<400> SEQUENCE: 102

```

```

ntgggtcaggt cgcaagcttg ctggn 25

```

```

<210> SEQ ID NO 103
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: n is dA modified with FAM.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (5)..(6)
<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 5

```

-continued

---

and 6.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (25)..(25)  
 <223> OTHER INFORMATION: n is dT modified with SpC3.  
 <400> SEQUENCE: 103  
 ntggtaaggt cgcaagcttg ctggn 25

<210> SEQ ID NO 104  
 <211> LENGTH: 26  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(1)  
 <223> OTHER INFORMATION: n is dA modified with FAM.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (11)..(12)  
 <223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 11  
 and 12.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (26)..(26)  
 <223> OTHER INFORMATION: n is dT modified with SpC3.  
 <400> SEQUENCE: 104  
 ntgggtcaagg tcgcaagctt gctggn 26

<210> SEQ ID NO 105  
 <211> LENGTH: 26  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(1)  
 <223> OTHER INFORMATION: n is dA modified with FAM.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (9)..(10)  
 <223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 9  
 and 10.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (26)..(26)  
 <223> OTHER INFORMATION: n is dT modified with SpC3.  
 <400> SEQUENCE: 105  
 ntgggtcaagg tcgcaagctt gctggn 26

<210> SEQ ID NO 106  
 <211> LENGTH: 26  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(1)  
 <223> OTHER INFORMATION: n is dA modified with FAM.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (7)..(8)  
 <223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 7  
 and 8.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (26)..(26)

-continued

---

<223> OTHER INFORMATION: n is dT modified with SpC3.

<400> SEQUENCE: 106

ntggtcaagg tcgcaagctt gctggn

26

<210> SEQ ID NO 107

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (1)..(1)

<223> OTHER INFORMATION: n is dA modified with FAM.

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (5)..(6)

<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 5 and 6.

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (26)..(26)

<223> OTHER INFORMATION: n is dT modified with SpC3.

<400> SEQUENCE: 107

ntggtcaagg tcgcaagctt gctggn

26

<210> SEQ ID NO 108

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (1)..(1)

<223> OTHER INFORMATION: n is dA modified with FAM.

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (26)..(26)

<223> OTHER INFORMATION: n is dT modified with FQ.

<400> SEQUENCE: 108

ntggtcaagg tcgcaagctt gctggn

26

<210> SEQ ID NO 109

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (1)..(1)

<223> OTHER INFORMATION: n is dA modified with FAM.

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (9)..(10)

<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 9 and 10.

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (26)..(26)

<223> OTHER INFORMATION: n is dT modified with FQ.

<400> SEQUENCE: 109

ntggtcaagg tcgcaagctt gctggn

26

<210> SEQ ID NO 110

<211> LENGTH: 26

-continued

---

<212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(1)  
 <223> OTHER INFORMATION: n is dA modified with FAM.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (9)..(10)  
 <223> OTHER INFORMATION: iBHQ1 is inserted between nucleotide positions  
 9 and 10.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (26)..(26)  
 <223> OTHER INFORMATION: n is dT modified with BHQ1.

<400> SEQUENCE: 110  
  
 ntggtcaagg tcgcaagctt gctggg 26

<210> SEQ ID NO 111  
 <211> LENGTH: 23  
 <212> TYPE: DNA  
 <213> ORGANISM: H1N1 influenza A virus

<400> SEQUENCE: 111  
  
 gtgctataaa caccagccty cca 23

<210> SEQ ID NO 112  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: H1N1 influenza A virus

<400> SEQUENCE: 112  
  
 cgggatattc cttaatcctg trgc 24

<210> SEQ ID NO 113  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: H1N1 influenza A virus

<400> SEQUENCE: 113  
  
 cagaatatac atccagtcac aattggaaaa 30

<210> SEQ ID NO 114  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: H1N1 influenza A virus  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(1)  
 <223> OTHER INFORMATION: n is dC modified with FAM.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (30)..(30)  
 <223> OTHER INFORMATION: n is dA modified with FQ.

<400> SEQUENCE: 114  
  
 nagaatatac atccagtcac aattggaaan 30

<210> SEQ ID NO 115  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: H1N1 influenza A virus  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(1)  
 <223> OTHER INFORMATION: n is dC modified with FAM.



-continued

---

```

<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (9)..(10)
<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 9
and 10.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (30)..(30)
<223> OTHER INFORMATION: n is dA modified with SpC3.

<400> SEQUENCE: 115

```

```

nagaatatac atccagtcac aattggaaan 30

```

```

<210> SEQ ID NO 116
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: H1N1 influenza A virus
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: n is dC modified with FAM.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (9)..(10)
<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 9
and 10.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (30)..(30)
<223> OTHER INFORMATION: n is dA modified with FQ.

<400> SEQUENCE: 116

```

```

nagaatatac atccagtcac aattggaaan 30

```

```

<210> SEQ ID NO 117
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: H1N1 influenza A virus
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: n is dC modified with FAM.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (9)..(10)
<223> OTHER INFORMATION: iBHQ1 is inserted between nucleotide positions
9 and 10.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (30)..(30)
<223> OTHER INFORMATION: n is dA modified with BHQ1.

<400> SEQUENCE: 117

```

```

nagaatatac atccagtcac aattggaaan 30

```

```

<210> SEQ ID NO 118
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: n is dA modified with FAM.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (9)..(10)
<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 9
and 10.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (26)..(26)
<223> OTHER INFORMATION: n is dT modified with RQ.

```

-continued

---

<400> SEQUENCE: 118

ntgggtcaagg tcgcaagctt gctggn

26

<210> SEQ ID NO 119

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (1)..(1)

<223> OTHER INFORMATION: n is dA modified with MAX.

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (9)..(10)

<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 9 and 10.

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (26)..(26)

<223> OTHER INFORMATION: n is dT modified with FQ.

<400> SEQUENCE: 119

ntgggtcaagg tcgcaagctt gctggn

26

<210> SEQ ID NO 120

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (1)..(1)

<223> OTHER INFORMATION: n is dA modified with MAX.

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (9)..(10)

<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 9 and 10.

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (26)..(26)

<223> OTHER INFORMATION: n is dt modified with RQ.

<400> SEQUENCE: 120

ntgggtcaagg tcgcaagctt gctggn

26

<210> SEQ ID NO 121

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (1)..(1)

<223> OTHER INFORMATION: n is dA modified with Cy3.

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (9)..(10)

<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 9 and 10.

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (26)..(26)

<223> OTHER INFORMATION: n is dT modified with FQ.

<400> SEQUENCE: 121

ntgggtcaagg tcgcaagctt gctggn

26

<210> SEQ ID NO 122

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

-continued

---

```

<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: n is dA modified with Cy3.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (9)..(10)
<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 9
and 10.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (26)..(26)
<223> OTHER INFORMATION: n is dT modified with RQ.

```

```

<400> SEQUENCE: 122

```

```

ntggtcaagg tcgcaagctt gctggn 26

```

```

<210> SEQ ID NO 123
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: n is dA modified with TEX.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (9)..(10)
<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 9
and 10.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (26)..(26)
<223> OTHER INFORMATION: n is dT modified with FQ.

```

```

<400> SEQUENCE: 123

```

```

ntggtcaagg tcgcaagctt gctggn 26

```

```

<210> SEQ ID NO 124
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: n is dA modified with TEX.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (9)..(10)
<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 9
and 10.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (26)..(26)
<223> OTHER INFORMATION: n is dT modified with RQ.

```

```

<400> SEQUENCE: 124

```

```

ntggtcaagg tcgcaagctt gctggn 26

```

```

<210> SEQ ID NO 125
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: n is dA modified with Cy5.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (9)..(10)
<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 9
and 10.
<220> FEATURE:
<221> NAME/KEY: misc_feature

```

-continued

---

<222> LOCATION: (26)..(26)  
 <223> OTHER INFORMATION: n is dT modified with FQ.  
  
 <400> SEQUENCE: 125  
  
 ntggtcaagg tcgcaagctt gctggn 26  
  
  
 <210> SEQ ID NO 126  
 <211> LENGTH: 26  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(1)  
 <223> OTHER INFORMATION: n is dA modified with Cy5.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (9)..(10)  
 <223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 9  
 and 10.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (26)..(26)  
 <223> OTHER INFORMATION: n is dT modified with RQ.  
  
 <400> SEQUENCE: 126  
  
 ntggtcaagg tcgcaagctt gctggn 26  
  
  
 <210> SEQ ID NO 127  
 <211> LENGTH: 26  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(1)  
 <223> OTHER INFORMATION: n is dA modified with Cy5.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (11)..(12)  
 <223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 11  
 and 12.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (26)..(26)  
 <223> OTHER INFORMATION: n is dT modified with RQ.  
  
 <400> SEQUENCE: 127  
  
 ntggtcaagg tcgcaagctt gctggn 26  
  
  
 <210> SEQ ID NO 128  
 <211> LENGTH: 35  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
  
 <400> SEQUENCE: 128  
  
 cgcgatcaga caaggagtgg gcttcatgga tcgcg 35  
  
  
 <210> SEQ ID NO 129  
 <211> LENGTH: 27  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
  
 <400> SEQUENCE: 129  
  
 ttacatgaag cccactcett gtctatc 27  
  
  
 <210> SEQ ID NO 130

-continued

---

```

<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: n is dC modified with FAM.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (35)..(35)
<223> OTHER INFORMATION: n is dG modified with Dabcyl.

<400> SEQUENCE: 130

ngcgatcaga caaggagtgg gcttcatgga tcgcn                                     35

<210> SEQ ID NO 131
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: n is dC modified with FAM.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (35)..(35)
<223> OTHER INFORMATION: n is dG modified with BHQ-1.

<400> SEQUENCE: 131

ngcgatcaga caaggagtgg gcttcatgga tcgcn                                     35

<210> SEQ ID NO 132
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: n is dC modified with FAM.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: n is dG modified with FQ.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (35)..(35)
<223> OTHER INFORMATION: n is a, c, g, or t

<400> SEQUENCE: 132

ngcgatcaga caaggagtgg gcttcatgga tcgcn                                     35

<210> SEQ ID NO 133
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: n is dC modified with FAM.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (35)..(35)
<223> OTHER INFORMATION: n is dG modified with FQ.
<220> FEATURE:
<221> NAME/KEY: misc_feature

```

-continued

---

<222> LOCATION: (35)..(35)  
<223> OTHER INFORMATION: n is dG modified with FQ/FQ.  
  
<400> SEQUENCE: 133  
  
ngcgatcaga caaggagtgg gcttcatgga tcgcn 35

<210> SEQ ID NO 134  
<211> LENGTH: 35  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(1)  
<223> OTHER INFORMATION: n is dC modified with FAM.  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (34)..(35)  
<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 34  
and 35.  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (35)..(35)  
<223> OTHER INFORMATION: n is dG modified with FQ.  
  
<400> SEQUENCE: 134  
  
ngcgatcaga caaggagtgg gcttcatgga tcgcn 35

<210> SEQ ID NO 135  
<211> LENGTH: 35  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(1)  
<223> OTHER INFORMATION: n is dC modified with FAM.  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (33)..(34)  
<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 33  
and 34.  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (35)..(35)  
<223> OTHER INFORMATION: n is dG modified with FQ.  
  
<400> SEQUENCE: 135  
  
ngcgatcaga caaggagtgg gcttcatgga tcgcn 35

<210> SEQ ID NO 136  
<211> LENGTH: 35  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(1)  
<223> OTHER INFORMATION: n is dC modified with FAM.  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (32)..(33)  
<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 32  
and 33.  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (35)..(35)  
<223> OTHER INFORMATION: n is dG modified with FQ.  
  
<400> SEQUENCE: 136

-continued

---

ngcgatcaga caaggagtgg gcttcatgga tcgcn

35

<210> SEQ ID NO 137  
<211> LENGTH: 35  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(1)  
<223> OTHER INFORMATION: n is dC modified with FAM.  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (31)..(32)  
<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 31 and 32.  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (35)..(35)  
<223> OTHER INFORMATION: n is dG modified with FQ.  
  
<400> SEQUENCE: 137

ngcgatcaga caaggagtgg gcttcatgga tcgcn

35

<210> SEQ ID NO 138  
<211> LENGTH: 35  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(1)  
<223> OTHER INFORMATION: n is dC modified with FAM.  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (30)..(31)  
<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 30 and 31.  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (35)..(35)  
<223> OTHER INFORMATION: n is dG modified with FQ.  
  
<400> SEQUENCE: 138

ngcgatcaga caaggagtgg gcttcatgga tcgcn

35

<210> SEQ ID NO 139  
<211> LENGTH: 35  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (1)..(1)  
<223> OTHER INFORMATION: n is dC modified with FAM.  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (29)..(30)  
<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 29 and 30.  
<220> FEATURE:  
<221> NAME/KEY: misc\_feature  
<222> LOCATION: (35)..(35)  
<223> OTHER INFORMATION: n is dG modified with FQ.  
  
<400> SEQUENCE: 139

ngcgatcaga caaggagtgg gcttcatgga tcgcn

35

-continued

---

<210> SEQ ID NO 140  
 <211> LENGTH: 35  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(1)  
 <223> OTHER INFORMATION: n is dC modified with FAM.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (35)..(35)  
 <223> OTHER INFORMATION: n is dG modified with RQ.

<400> SEQUENCE: 140

ngcgatcaga caaggagtgg gcttcatgga tcgcn 35

<210> SEQ ID NO 141  
 <211> LENGTH: 35  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(1)  
 <223> OTHER INFORMATION: n is dC modified with FAM.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (35)..(35)  
 <223> OTHER INFORMATION: n is dG modified with FQ/RQ.

<400> SEQUENCE: 141

ngcgatcaga caaggagtgg gcttcatgga tcgcn 35

<210> SEQ ID NO 142  
 <211> LENGTH: 35  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(1)  
 <223> OTHER INFORMATION: n is dC modified with FAM.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (34)..(35)  
 <223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 34  
 and 35.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (35)..(35)  
 <223> OTHER INFORMATION: n is dG modified with RQ.

<400> SEQUENCE: 142

ngcgatcaga caaggagtgg gcttcatgga tcgcn 35

<210> SEQ ID NO 143  
 <211> LENGTH: 35  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(1)  
 <223> OTHER INFORMATION: n is dC modified with FAM.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (33)..(34)  
 <223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 33



-continued

---

and 34.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (35)..(35)  
 <223> OTHER INFORMATION: n is dG modified with RQ.  
 <400> SEQUENCE: 143  
 ngcgatcaga caaggagtgg gcttcatgga tcgcn 35

<210> SEQ ID NO 144  
 <211> LENGTH: 35  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(1)  
 <223> OTHER INFORMATION: n is dC modified with FAM.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (32)..(33)  
 <223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 32  
 and 33.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (35)..(35)  
 <223> OTHER INFORMATION: n is dG modified with RQ.  
 <400> SEQUENCE: 144  
 ngcgatcaga caaggagtgg gcttcatgga tcgcn 35

<210> SEQ ID NO 145  
 <211> LENGTH: 35  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(1)  
 <223> OTHER INFORMATION: n is dC modified with FAM.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (31)..(32)  
 <223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 31  
 and 32.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (35)..(35)  
 <223> OTHER INFORMATION: n is dG modified with RQ.  
 <400> SEQUENCE: 145  
 ngcgatcaga caaggagtgg gcttcatgga tcgcn 35

<210> SEQ ID NO 146  
 <211> LENGTH: 35  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (1)..(1)  
 <223> OTHER INFORMATION: n is dC modified with FAM.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (30)..(31)  
 <223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 30  
 and 31.  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (35)..(35)

<223> OTHER INFORMATION: n is dG modified with RQ.

<400> SEQUENCE: 146

ngcgatcaga caaggagtgg gcttcatgga tcgcn

35

<210> SEQ ID NO 147

<211> LENGTH: 35

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (1)..(1)

<223> OTHER INFORMATION: n is dC modified with FAM.

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (29)..(30)

<223> OTHER INFORMATION: iFQ is inserted between nucleotide positions 29 and 30.

<220> FEATURE:

<221> NAME/KEY: misc\_feature

<222> LOCATION: (35)..(35)

<223> OTHER INFORMATION: n is dG modified with RQ.

<400> SEQUENCE: 147

ngcgatcaga caaggagtgg gcttcatgga tcgcn

35

What is claimed is:

1. A composition, comprising:


a first oligonucleotide having the structure 5'-Y<sub>1</sub>-L<sub>1</sub>-X-L<sub>2</sub>-Y<sub>2</sub>-3', wherein:

Y<sub>1</sub> comprises a sequence of DNA or RNA nucleotides, including a first nucleotide N<sub>1</sub> having a 3' phosphate covalently linked to L<sub>1</sub>;


Y<sub>2</sub> comprises a sequence of DNA or RNA nucleotides, including a second nucleotide N<sub>2</sub> having a 5' phosphate covalently linked to L<sub>2</sub>;

L<sub>1</sub> and L<sub>2</sub> each independently are a direct bond or a C<sub>1</sub>-C<sub>7</sub> alkyl, alkynyl, alkenyl, heteroalkyl, substituted alkyl, aryl, heteroaryl, substituted aryl, cycloalkyl, alkylaryl, or alkoxy group;

X is




R<sub>1</sub> is a hydrogen or a C<sub>1</sub>-C<sub>8</sub> alkyl; and M is




L<sub>3</sub> is a direct bond or a C<sub>1</sub>-C<sub>8</sub> alkyl, alkenyl, heteroalkyl, substituted alkyl cycloalkyl, or alkoxy;

R<sub>2</sub>-R<sub>6</sub> each independently are a hydrogen, an alkyl, an alkenyl, a heteroalkyl, a substituted alkyl, an aryl, a heteroaryl, a substituted aryl, a cycloalkyl, an alkylaryl, an alkoxy, an electron withdrawing group, or an electron donating group, and one of

R<sub>2</sub>-R<sub>6</sub> is



wherein P is




and

R<sub>7</sub>-R<sub>9</sub> each independently are a hydrogen, an alkoxy, an alkyl, an alkylamino, an arylamino, a cycloalkyl, a heteroalkoxy, a heteroalkyl, or an amino; and

R<sub>10</sub>-R<sub>13</sub> each independently are a hydrogen, a nitro, a cyano, a carboxylate, a sulfonyl, a sulfamoyl, an alkenyl, an alkynyl, an amino, an aryl, a heteroaryl, a biaryl, a bialkenyl, a bialkynyl, an alkoxy carbonyl or a carbamoyl, or

111

wherein P is



and

$R_{14}$ - $R_{19}$ , each independently are a hydrogen, an alkyl, a heteroalkyl, an aryl, a heteroaryl, an electron withdrawing group, or a five or six membered ring structure formed from the  $R_1$ ,  $R_2$  pair, the  $R_3$ ,  $R_4$  pair, the  $R_4$ ,  $R_5$  pair or the  $R_5$ ,  $R_6$  pair;

wherein the first oligonucleotide is hybridized to a second oligonucleotide having the structure 3'- $Y_3$ - $Y_4$ -5', wherein:

$Y_3$  comprises a sequence of DNA or RNA nucleotides, including a third nucleotide  $N_3$ ; and

$Y_4$  comprises a sequence of DNA or RNA nucleotides, including a fourth nucleotide  $N_4$  that is directly attached to nucleotide  $N_3$ ;

wherein  $N_1$  base pairs with  $N_3$  and  $N_2$  base pairs with  $N_4$  such that the duplex has a  $T_m$  that is greater than the  $T_m$  of a duplex formed between the second oligonucleotide and a third oligonucleotide having the structure 5'- $Y_1$ - $Y_2$ -3'.

2. The composition of claim 1, wherein  $L_1$  and  $L_2$  each is a  $C_1$ - $C_7$  alkyl.


3. The composition of claim 2, wherein  $L_1$  and  $L_2$  are  $C_2$  alkyls.

4. The composition of claim 1, wherein the 3' phosphate that is covalently linked to  $L_1$ , and the 5' phosphate that is

112

covalently linked to  $L_2$ , each independently are a phosphodiester, a phosphothioate, a phosphodithioate, a methyl phosphonate, a phosphoramidate, a phosphoramidite or a phosphotriester.

5. The composition of claim 1, wherein:  
 $R_9$  is



15

6. The composition of claim 1, wherein the first oligonucleotide is labeled with a fluorophore.

7. The composition of claim 6, wherein the fluorophore is 6-carboxyfluorescein (FAM), 2'7'-dimethoxy-4'5'-dichloro-6-carboxyfluorescein (JOE), tetrachlorofluorescein (TET), 6-carboxyrhodamine (R6G), N,N,N,N'-tetramethyl-6-carboxyrhodamine (TAMRA), 6-carboxy-X-rhodamine (ROX); 1-dimethylaminonaphthyl-5-sulfonate, 1-anilino-8-naphthalene sulfonate, 2-p-toluidinyl-6-naphthalene sulfonate, 5-(2'-aminoethyl)aminonaphthalene-1-sulfonic acid (EDANS), a coumarin dye, an acridine dye, indodicarbocyanine 3 (Cy3), indodicarbocyanine 5 (Cy5), indodicarbocyanine 5.5 (Cy5.5), 3-(1-carboxy-pentyl)-3'-ethyl-5,5'-dimethyloxycarbocyanine (CyA); 1H,5H, 11H,15H-Xantheno[2,3,4-ij:5,6,7-i'j']diquinolizin-18-ium, 9-[2(or 4)-[[[6-[2,5-dioxo-1-pyrrolidinyl)oxy]-6-oxohexyl]amino]sulfonyl]-4(or 2)-sulfophenyl]-2,3,6,7,12,13,16,17-octahydro-inner salt (TR or Texas Red), a BODIPY™ dye, benzoxazole, stilbene or pyrene.

8. The composition of claim 6, wherein the fluorophore is attached to the 5' phosphate at the end of the oligonucleotide.

9. The composition of claim 6, wherein the first oligonucleotide is labeled with a second quencher.

\* \* \* \* \*