

US009267144B2

(12) **United States Patent**  
**Sidorov et al.**

(10) **Patent No.:** **US 9,267,144 B2**  
(45) **Date of Patent:** **Feb. 23, 2016**

(54) **PLASTID TRANSFORMATION OF MAIZE**

(75) Inventors: **Vladimir Sidorov**, Chesterfield, MO (US); **Jeffrey M. Staub**, Wildwood, MO (US); **Yuechun Wan**, Madison, WI (US); **Guangning Ye**, Chesterfield, MO (US)

(73) Assignee: **Monsanto Technology LLC**, St. Louis, MO (US)

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 1151 days.

(21) Appl. No.: **12/106,073**

(22) Filed: **Apr. 18, 2008**

(65) **Prior Publication Data**

US 2008/0289063 A1 Nov. 20, 2008

**Related U.S. Application Data**

(60) Continuation of application No. 11/244,263, filed on Oct. 5, 2005, now abandoned, which is a division of application No. 10/248,492, filed on Jan. 23, 2003, now abandoned.

(60) Provisional application No. 60/319,095, filed on Jan. 23, 2002.

(51) **Int. Cl.**  
**C12N 15/82** (2006.01)

(52) **U.S. Cl.**  
CPC ..... **C12N 15/8214** (2013.01)

(58) **Field of Classification Search**  
None  
See application file for complete search history.

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

5,627,061	A *	5/1997	Barry et al.	800/288
5,693,507	A	12/1997	Daniell et al.	
5,767,368	A	6/1998	Zhong et al.	
6,603,061	B1 *	8/2003	Armstrong et al.	800/294
7,129,391	B1 *	10/2006	Daniell	800/278
7,803,991	B2 *	9/2010	Daniell	800/300
2002/0042934	A1 *	4/2002	Staub et al.	800/300
2005/0198704	A1	9/2005	Sticklen	

**FOREIGN PATENT DOCUMENTS**

WO WO 99/10513 \* 3/1999

**OTHER PUBLICATIONS**

Zhong et al (1996, Plant Physiol. 110:1097-1107).  
Khan and Maliga, "Fluorescent antibiotic resistance marker for tracking plastid transformation in higher plants." Nat Biotechnol. Sep. 1999;17(9):910-5.  
Maier et al., "Complete sequence of the maize chloroplast genome: gene content, hotspots of divergence and fine tuning of genetic information by transcript editing." J Mol Biol. Sep. 1, 1995;251(5):614-28.

Sidorov et al., "Technical Advance: Stable chloroplast transformation in potato: use of green fluorescent protein as a plastid marker." Plant J. Jul. 1999;19(2):209-216.

Sikdar et al., "Plastid transformation in Arabidopsis thaliana." vol. 18, Nos. 1-2 / Nov. 1998 Plant Cell Reports 1 8:20-24, 1998;

Staub and Maliga, "Long regions of homologous DNA are incorporated into the tobacco plastid genome by transformation." Plant Cell. Jan. 1992;4(1):39-45.

Staub et al., "High-yield production of a human therapeutic protein in tobacco chloroplasts." Nat Biotechnol. Mar. 2000;18(3):333-8.

Svab and Maliga, "High-frequency plastid transformation in tobacco by selection for a chimeric aadA gene." Proc Natl Acad Sci U S A. Feb. 1, 1993;90(3):913-7.

Svab et al., "Stable transformation of plastids in higher plants." Proc Natl Acad Sci U S A. Nov. 1990;87(21):8526-30.

Zhong et al., "The Competence of Maize Shoot Meristems for Integrative Transformation and Inherited Expression of Transgenes." Plant Physiol. Apr. 1996;110(4):1097-1107.

Zoubenko et al., "Efficient targeting of foreign genes into the tobacco plastid genome." Nucleic Acids Res. Sep. 25, 1994;22(19):3819-24.

Ahmadabadi, Mohammad et al., A leaf-based regeneration and transformation system for maize (*Zea mays* L.), Transgenic Res (2007) 16:437-448.

Bogorad, Lawrence, Engineering chloroplasts: an alternative site for foreign genes, proteins, reactions and products, Tibtech Jun. 2000, vol. 18, pp. 257-263.

Daniell, Henry et al., Breakthrough in chloroplast genetic engineering of agronomically important crops, Trends in Biotechnology, vol. 23, No. 5, May 2005, pp. 238-245.

Lee, Sa Mi et al., Plastid Transformation in the Monocotyledonous Cereal Crop, Rice (*Oryza sativa*) and Transmission of Transgenes to Their Progeny, Mol. Cells, vol. 21, No. 3, pp. 401-410 (2006).

Verma, Dheeraj et al., Chloroplast Vector Systems for Biotechnology Applications, Plant Physiology, Dec. 2007, vol. 145, pp. 1129-1143.

Clarke, Jihong Liu et al., Chloroplast Biotechnology, Genomics and Evolution: Current Status, Challenges and Future Directions, Plant Molecular Biology, May 18, 2011, pp. 207-209, vol. 76.

Dhingra, Amit et al., Chloroplast Genetic Engineering Via Organogenesis or Somatic Embryogenesis, Methods in Molecular Biology, 2006, pp. 245-262, vol. 323.

Daniell, Henry et al., Expression of the Native Cholera Toxin B Subunit Gene and Assembly as Functional Oligomers in Transgenic Tobacco Chloroplasts, Journal of Molecular Biology, 2001, pp. 1001-1009, vol. 311.

Svab, Zora et al., High-Frequency Plastid Transformation in Tobacco by Selection for a Chimeric aadA Gene, Proceedings of the National Academy of Sciences, Feb. 1993, pp. 913-917, vol. 90.

\* cited by examiner

*Primary Examiner* — Anne Kubelik

(74) *Attorney, Agent, or Firm* — Thompson Coburn LLP; J. Wendy Davis

(57) **ABSTRACT**

A method is provided for transforming maize plants to express DNA sequences of interest from plant cell plastids. The method allows the transformation of maize plant tissue with heterologous DNA constructs by microprojectile bombardment of green light grown cultures and dark grown embryogenic cultures. The invention also provides for maize cells in which the plastids contain heterologous DNA constructs.

**14 Claims, 14 Drawing Sheets**




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7






Figure 8




Figure 9




Figure 10




Figure 11




Figure 12





Figure L3

Figure 14



**PLASTID TRANSFORMATION OF MAIZE****CROSS REFERENCE TO RELATED APPLICATIONS**

This application is a continuation of U.S. application Ser. No. 11/244,263 filed Oct. 5, 2005, which is a divisional of U.S. application Ser. No. 10/248,492 filed Jan. 23, 2003, which claims priority to U.S. Application No. 60/319,095 filed Jan. 23, 2002, the contents of all of which are hereby incorporated in their entireties by reference into this application.

**BACKGROUND OF INVENTION**

This invention relates to the application of genetic engineering techniques to plants. Specifically, the invention relates to compositions and methods for transformation of nucleic acid sequences into plant cell plastids.

Molecular biological techniques have enabled researchers to introduce pieces of DNA from one organism to another organism. Such techniques, referred to as recombinant DNA technology, have positively impacted the areas of medicine and agriculture. Conventional cloning methods have enabled the introduction of new pharmaceuticals and improved crops of agricultural importance. As the need for the introduction of multiple pieces of DNA and larger fragments of DNA into numerous target hosts increases, the need for novel cloning strategies increases accordingly.

The plastids of higher plants are an attractive target for genetic engineering. Plant plastids (chloroplasts, amyloplasts, elaioplasts, chromoplasts, etc.) are the major biosynthetic centers that in addition to photosynthesis are responsible for production of industrially important compounds such as amino acids, complex carbohydrates, fatty acids, and pigments. Plastids are derived from a common precursor known as a proplastid and thus the plastids present in a given plant species all have the same genetic content. Plant cells contain 500-10,000 copies of a small 120-160 kilobase circular genome, each molecule of which has a large (approximately 25 kb) inverted repeat. Thus, it is possible to engineer plant cells to contain up to 20,000 copies of a particular gene of interest, which potentially can result in very high levels of foreign gene expression.

Plastid transformation has been restricted to a few dicot species. In all successful experiments (Svab et al., Proc. Natl. Acad. Sci. USA 87, 8526-8530, 1990; Svab and Maliga, Proc. Natl. Acad. Sci. USA 90:913-917, 1993; Sikdar et al., Plant Cell Reports 18:20-24, 1998; Sidorov et al., Plant Journal 19(2):209-216, 1999), green leaves that contain developed chloroplasts were the target material used for particle bombardment. The most efficient selectable marker for isolation of plastid transformants is *aadA*, which confers resistance to the antibiotics spectinomycin and streptomycin (Svab et al., Proc. Natl. Acad. Sci. USA 87, 8526-8530, 1990; Svab and Maliga, Proc. Natl. Acad. Sci. USA 90:913-917, 1993; Sikdar et al., Plant Cell Reports 18:20-24, 1998; Sidorov et al., Plant Journal 19(2):209-216, 1999). NPTII, conferring resistance to kanamycin, has been reported for tobacco (Carrer et al., Mol Gen Genet 241:49-56, 1993) and glyphosate (U.S. Patent Application 200242934) were also used for selecting plastid transformants.

Recently, plastid transformation in rice was reported using a nongreen embryogenic culture as initial material for transformation (Khan and Maliga, Nature Biotechnology 17(9): 910-915, 1999). Selection was for streptomycin resistance conferred by *aadA*, as rice is not sensitive to spectinomycin.

Selection was applied for only two weeks in liquid suspension and during the subsequent plant regeneration phase. Regenerated plants were chimeric, with some leaves containing sectors with transformed plastids. However, the transformed cells were also apparently only heteroplasmic, having both transformed and non-transformed plastids. This pattern of heteroplasmy is most likely due to the short period of selection that did not allow enough time for amplification of transformed plastids prior to plant regeneration.

However, a need still exists for a method of producing homoplasmic monocotyledonous plants, which are most useful. The present invention should be applicable to all monocots.

**SUMMARY OF INVENTION**

Novel methods for producing transplastomic monocotyledonous plants are provided. Such methods employ novel combinations of starting explant material with an appropriate monocotyledonous optimized plastid transformation vector and selection/tissue culture protocols. The invention described herein provides a method for obtaining homoplasmic monocotyledonous plants. In preferred embodiments of the present invention, the explant is chosen from among green multiple bud cultures, greening embryogenic callus, dark-grown embryogenic callus, or pre-cultured immature embryos. Plant cells containing transformed plastids are identified and selected and the cells amplified. Transplastomic plants are regenerated therefrom. In a preferred embodiment, explants from maize tissue are used. The invention described herein provides transplastomic monocotyledonous plant cells and sectors. Also provided herein are monocotyledonous plants, including maize, rice and wheat, whose plastids are transformed to contain a heterologous nucleic acid sequence conferring a desired trait or outcome to the resulting transgenic plant.

**BRIEF DESCRIPTION OF DRAWINGS**

FIG. 1. Plasmid map of construct pMON49219.  
 FIG. 2. Plasmid map of construct pMON38722.  
 FIG. 3. Plasmid map of construct pMON56102.  
 FIG. 4. Plasmid map of construct pMON49296.  
 FIG. 5. Plasmid map of construct pMON56108.  
 FIG. 6. Plasmid map of construct pMON56110.  
 FIG. 7. Plasmid map of construct pMON56113.  
 FIG. 8. Plasmid map of construct pMON56114.  
 FIG. 9. Plasmid map of construct pMON38762.  
 FIG. 10. Plasmid map of construct pMON30098.  
 FIG. 11. Plasmid map of construct pMON25496.  
 FIG. 12. Plasmid map of construct pMON38762.  
 FIG. 13. Plasmid map of construct pMON45294.  
 FIG. 14. Map of expected PCR products.

**BRIEF DESCRIPTION OF SEQUENCES**

SEQ ID NO:1 PCR primer Xu10  
 SEQ ID NO:2 PCR primer Xu11  
 SEQ ID NO:3 PCR primer Xu12  
 SEQ ID NO:4 PCR primer Xu13  
 SEQ ID NO:5 PCR primer Xu7  
 SEQ ID NO:6 PCR primer Xu8  
 SEQ ID NO:7 maize *rbcL* flanking region  
 SEQ ID NO:8 maize inverted repeat region  
 SEQ ID NO:9 PCR primer  
 SEQ ID NO:10 PCR primer  
 SEQ ID NO:11 PCR primer



SEQ ID NO:12 PCR primer  
 SEQ ID NO:13 maize plastid promoter from rrn16  
 SEQ ID NO:14 maize Prn with G10L  
 SEQ ID NO:15 termination region from tobacco rps16  
 gene  
 SEQ ID NO:16 tobacco Prn with atpB gene leader plus  
 DB  
 SEQ ID NO:17 termination region from maize petD gene  
 SEQ ID NO:18 termination region from maize rbcL gene  
 SEQ ID NO:19 maize Prn plus G10L plus 14 aa of GFP  
 SEQ ID NO:20 PCR primer 10  
 SEQ ID NO:21 PCR primer 60  
 SEQ ID NO:22 PCR primer 61  
 SEQ ID NO:23 PCR primer 62  
 SEQ ID NO:24 PCR primer 63  
 SEQ ID NO:25 PCR primer 64  
 SEQ ID NO:26 PCR primer 65  
 SEQ ID NO:27-PCR primer 66  
 SEQ ID NO:28 PCR primer GFP 5':1  
 SEQ ID NO:29 PCR primer GFP 3':2  
 SEQ ID NO:30 PCR primer 202  
 SEQ ID NO:31 PCR primer 201  
 SEQ ID NO:32 PCR primer 198  
 SEQ ID NO:33 PCR primer 199  
 SEQ ID NO:34 PCR primer 200  
 SEQ ID NO:35 PCR primer 194

#### DETAILED DESCRIPTION

##### Definitions

In order to provide a clear and consistent understanding of the specification and the claims, including the scope given to such terms, the following definitions are provided.

A first nucleic acid sequence is “operably linked” with a second nucleic acid sequence when the sequences are arranged so that the first nucleic acid sequence affects the function of the second nucleic acid sequence. Preferably, the two sequences are part of a single contiguous nucleic acid molecule and more preferably are adjacent. For example, a promoter is operably linked to a gene if the promoter regulates or mediates transcription of the gene in a cell.

A “recombinant” nucleic acid is made by an artificial combination of two otherwise separated segments of sequence, e.g., by chemical synthesis or by the manipulation of isolated segments of nucleic acids by genetic engineering techniques. Techniques for nucleic-acid manipulation are well known (see for example, Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press, 1989; Maliga et al., *Methods in Plant Molecular Biology*, Cold Spring Harbor Press, 1995; Birren et al., *Genome Analysis: volume 1, Analyzing DNA*, (1997), volume 2, *Detecting Genes*, (1998), volume 3, *Cloning Systems*, (1999) volume 4, *Mapping Genomes*, (1999), Cold Spring Harbor, N.Y.).

Methods for chemical synthesis of nucleic acids are discussed, for example, in Beaucage and Carruthers (*Tetra. Letts.* 22:1859-1862, 1981) and Matteucci et al. (*J. Am. Chem. Soc.* 103:3185, 1981). Chemical synthesis of nucleic acids can be performed, for example, on commercial automated oligonucleotide synthesizers.

A “synthetic nucleic acid sequence” can be designed and chemically synthesized for enhanced expression in particular host cells and for the purposes of cloning into appropriate vectors. Host cells often display a preferred pattern of codon usage (Campbell et al., *Plant Physiol.* 92:1-11, 1990). Synthetic DNAs designed to enhance expression in a particular host should therefore reflect the pattern of codon usage in the

host cell. Computer programs are available for these purposes including but not limited to the “BestFit” or “Gap” programs of the Sequence Analysis Software Package, Genetics Computer Group, Inc., University of Wisconsin Biotechnology Center, Madison, Wis. 53711.

“Amplification” of nucleic acids or “nucleic acid reproduction” refers to the production of additional copies of a nucleic acid sequence and is carried out using polymerase chain reaction (PCR) technologies. A variety of amplification methods are known in the art and are described, inter alia, in U.S. Pat. Nos. 4,683,195 and 4,683,202 and in *PCR Protocols: A Guide to Methods and Applications*, ed. Innis et al., Academic Press, San Diego, 1990. In PCR, a primer refers to a short oligonucleotide of defined sequence that is annealed to a DNA template to initiate the polymerase chain reaction.

“Transformed”, “transfected”, or “transgenic” refers to a cell, tissue, organ, or organism into which has been introduced a foreign nucleic acid, such as a recombinant vector. Preferably, the introduced nucleic acid is integrated into the genomic DNA of the recipient cell, tissue, organ or organism such that the introduced nucleic acid is inherited by subsequent progeny. A “transgenic” or “transformed” cell or organism also includes progeny of the cell or organism and progeny produced from a breeding program employing such a “transgenic” plant as a parent in a cross and exhibiting an altered phenotype resulting from the presence of a recombinant construct or vector.

The term “gene” refers to chromosomal DNA, plasmid DNA, cDNA, synthetic DNA, or other DNA that encodes a peptide, polypeptide, protein, or RNA molecule, and regions flanking the coding sequence involved in the regulation of expression. Some genes can be transcribed into mRNA and translated into polypeptides (structural genes); other genes can be transcribed into RNA (e.g., rRNA, tRNA); and other types of genes function as regulators of expression (regulator genes).

“Expression” of a gene refers to the transcription of a gene to produce the corresponding mRNA and translation of this mRNA to produce the corresponding gene product, i.e., a peptide, polypeptide, or protein. Gene expression is controlled or modulated by regulatory elements including 5' regulatory elements such as promoters.

“Genetic component” refers to any nucleic acid sequence or genetic element that may also be a component or part of an expression vector. Examples of genetic components include, but are not limited to, promoter regions, 5' untranslated leaders, introns, genes, 3' untranslated regions, and other regulatory sequences or sequences that affect transcription or translation of one or more nucleic acid sequences.

The terms “recombinant DNA construct,” “recombinant vector,” “expression vector” or “expression cassette” refer to any agent such as a plasmid, cosmid, virus, BAC (bacterial artificial chromosome), autonomously replicating sequence, phage, or linear or circular single-stranded or double-stranded DNA or RNA nucleotide sequence, derived from any source, capable of genomic integration or autonomous replication, comprising a DNA molecule in which one or more DNA sequences have been linked in a functionally operative manner using well-known recombinant DNA techniques.

As used herein, “heterologous” in reference to a nucleic acid is a nucleic acid that originates from a foreign species, or, if from the same species, is substantially modified from its native form in composition and/or genomic locus by deliberate human intervention. For example, a promoter operably linked to a heterologous structural gene is from a species different from that from which the structural gene was

derived, or, if from the same species, one or both are substantially modified from their original form. A heterologous protein may originate from a foreign species, or, if from the same species, is substantially modified from its original form by deliberate human intervention.

As used herein, "recombinant" includes reference to a cell or vector that has been modified by the introduction of a heterologous nucleic acid sequence or that the cell is derived from a cell so modified. Thus, for example, recombinant cells express genes that are not found in identical form within the native (non-recombinant) form of the cell or express native genes that are otherwise abnormally expressed, under expressed or not expressed at all as a result of deliberate human intervention.

By "host cell" is meant a cell that contains a vector and supports the replication, transcription, or transcription and translation (expression) of the expression construct. Host cells for use in the present invention can be prokaryotic cells, such as *E. coli*, or eukaryotic cells such as yeast, plant, insect, amphibian, or mammalian cells. Preferably, host cells are monocotyledenous plant cells.

As used herein, the term "plant" includes reference to whole plants, plant organs (for example, leaves, stems, roots, etc.), seeds, and plant cells and progeny of same. Plant cell, as used herein, includes, without limitation, seed suspension cultures, embryos, meristematic regions, callus tissue, leaves, roots, shoots, gametophytes, sporophytes, pollen, and microspores. The class of plants that can be used in the methods of the present invention is generally as broad as the class of higher plants amenable to transformation techniques, including monocotyledenous plants. Particularly preferred plants include barley, corn, oat, rice, rye, sorghum, sugarcane, triticale, and wheat.

As used herein, "transgenic plant" includes reference to a plant that comprises within its nuclear genome a heterologous polynucleotide. Generally, the heterologous polynucleotide is stably integrated within the nuclear genome such that the polynucleotide is passed on to successive generations. The heterologous polynucleotide may be integrated into the genome alone or as part of a recombinant expression cassette. "Transgenic" is used herein to include any cell, cell line, callus, tissue, plant part or plant, the genotype of which has been altered by the presence of heterologous nucleic acid including those transgenics initially so altered as well as those created by sexual crosses or asexual propagation from the initial transgenic. The term "transgenic" as used herein does not encompass the alteration of the genome (chromosomal or extra-chromosomal) by conventional plant breeding methods or by naturally occurring events such as random cross-fertilization, non-recombinant viral infection, non-recombinant bacterial transformation, non-recombinant transposition, or spontaneous mutation.

As used herein, "sector" refers to a proliferating lineage of cells in which exogenous DNA is stably integrated and inherited through organelle replication in daughter cells.

As used herein, "transplastomic" refers to a plant cell having a heterologous nucleic acid introduced into the plant cell plastid. The introduced nucleic acid may be integrated into the plastid genome or may be contained in an autonomously replicating plasmid. Preferably, the nucleic acid is integrated into the plant cell plastid's genome. A plant cell can be both transgenic and transplastomic.

The term "introduced" in the context of inserting a nucleic acid sequence into a cell, means "transfection" or "transformation" or "transduction" and includes reference to the incorporation of a nucleic acid sequence into a eukaryotic or prokaryotic cell where the nucleic acid sequence may be

incorporated into the genome of the cell (for example, chromosome, plasmid, plastid, or mitochondrial DNA), converted into an autonomous replicon, or transiently expressed (for example, transfected mRNA).

The present invention describes a method for the production of a plastid transformant of maize using microprojectile bombardment of green, light-grown cultures. As target tissues, use of greening embryogenic cultures, dark-grown embryogenic cultures and pre-cultured immature embryos is also described. Successful plastid transformation includes several steps: a) engineering of plastid targeted constructs with an efficient selectable marker; b) establishment of regenerable cell cultures containing plastids suitable for transformation; c) development of a proper selection scheme favorable for sorting out of transformed plastids; and d) recovery of stable plastid transformants.

In developing the constructs of the invention, the various fragments comprising the regulatory regions and open reading frame may be subjected to different processing conditions, such as ligation, restriction enzyme digestion, PCR, in vitro mutagenesis, linkers and adapters addition, and the like. Thus, nucleotide transitions, transversions, insertions, deletions, or the like, may be performed on the DNA that is employed in the regulatory regions or the nucleic acid sequences of interest for expression in the plastids. Methods for restriction digests, Klenow blunt end treatments, ligations, and the like are well known to those in the art and are described, for example, by Maniatis et al. (in *Molecular cloning: a laboratory manual* (1982) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.).

During the preparation of the constructs, the various fragments of DNA will often be cloned in an appropriate cloning vector, which allows for amplification of the DNA, modification of the DNA or manipulation by joining or removing of sequences, linkers, or the like. Normally, the vectors will be capable of replication in at least a relatively high copy number in *E. coli*. A number of vectors are readily available for cloning, including such vectors as pBR322, pUC series, M13 series, and pBluescript (Stratagene; La Jolla, Calif.).

The constructs for use in the methods of the present invention are prepared to direct the expression of the nucleic acid sequences directly from the host plant cell plastid. Examples of such constructs and methods are known in the art and are generally described, for example, in Svab et al. (1990) *Proc. Natl. Acad. Sci. USA* 87:8526-8530 and Svab and Maliga (1993) *Proc. Natl. Acad. Sci. USA* 90:913-917 and in U.S. Pat. No. 5,693,507.

The skilled artisan will recognize that any convenient element that is capable of initiating transcription in a plant cell plastid, also referred to as "plastid functional promoters," can be employed in the constructs of the present invention. A number of plastid functional promoters are available in the art for use in the constructs and methods of the present invention. Such promoters include, but are not limited to, the promoter of the D1 thylakoid membrane protein, *psbA* (Staub et al. *EMBO Journal*, 12(2):601-606, 1993), and the 16S rRNA promoter region, *Prn* (Svab and Maliga, 1993, *Proc. Natl. Acad. Sci. USA* 90:913-917). The expression cassettes can include additional elements for expression of the protein, such as transcriptional and translational enhancers, ribosome binding sites, and the like.

As translation is a limiting step for plastid transgene expression, a variety of translational control elements need to be tested for efficacy. Efficient transgene translation will ensure that the markers used for selection of plastid transformed cells will function. Examples of such translational enhancing sequences include the heterologous bacteriophage

gene 10 leader (G10L) (U.S. Pat. No. 6,271,444), the G10L including the downstream box (DB) (U.S. patent application Ser. No. 09/668,188), or the 14 amino acids from the green fluorescent protein (GFP) (U.S. patent application Ser. No. 09/351,124) enhancer elements.

Regulatory transcript termination regions may be provided in the expression constructs of this invention as well. Transcript termination regions may be provided by any convenient transcription termination region derived from a gene source, for example, the transcript termination region that is naturally associated with the transcript initiation region. The skilled artisan will recognize that any convenient transcript termination region that is capable of terminating transcription in a plant cell may be employed in the constructs of the present invention.

The expression cassettes for use in the methods of the present invention also preferably contain additional nucleic acid sequences providing for the integration into the host plant cell plastid genome or for autonomous replication of the construct in the host plant cell plastid. Preferably, the plastid expression constructs contain regions of homology for integration into the host plant cell plastid. The regions of homology employed can target the constructs for integration into any region of the plastid genome; preferably the regions of homology employed target the construct to either the inverted repeat region of the plastid genome or the large single copy region. Where more than one construct is to be used in the methods, the constructs can employ the use of the regions of homology to target the insertion of the construct into the same or a different position of the plastid genome.

Additional expression cassettes can comprise any nucleic acid to be introduced into a host cell plastid by the methods encompassed by the present invention including, for example, DNA sequences or genes from another species, or even genes or sequences that originate with or are present in the same species, but are incorporated into recipient cells by genetic engineering methods rather than classical reproduction or breeding techniques. An introduced piece of DNA can be referred to as exogenous DNA. Exogenous as used herein is intended to refer to any gene or DNA segment that is introduced into a recipient cell, regardless of whether a similar gene may already be present in such a cell. The type of DNA included in the exogenous DNA can include DNA that is already present in the plant cell, DNA from another plant, DNA from a different organism, or a DNA generated externally, such as a DNA sequence containing an antisense message of a gene, or a DNA sequence encoding a synthetic or modified version of a gene.

A number of factors affect successful plastid transformation. Among them the most critical are 1) the target material used for delivering plastid-expressing genes into and regenerating transformants from and 2) the use of proper selectable markers and effective selection regimes. Green leaves, which contain large numbers of well-developed chloroplasts, are the ideal target material for introducing plastid-expressing foreign genes into plastids. In all successful stable plastid transformation reports to-date, green leaves are the only target material used because of the availability of well-established leaf-based plant regeneration systems in those species (tobacco, *Arabidopsis*, and potato). However, in corn, like most monocotyledonous plant species, plant regeneration from leaf tissue is impossible or very inefficient, which severely limits the possibility of using leaf tissue for corn plastid transformation. Instead, we have been using maize light- and dark-grown embryogenic cultures or immature embryos and light-grown green multiple bud cultures for plastid transformation. Immature embryos and embryogenic cultures

derived from immature embryos are typically highly transformable using nuclear transformation approaches, but they contain immature plastids that may not be optimally suited for plastid transformation. Light-grown cultures are established by culturing embryogenic callus or excised meristem region from in vitro germinating seedlings on media containing different levels and different combinations of plant growth regulators under light and selecting for green and highly regenerable (either embryogenic or multiple bud) callus cultures. The light-grown cultures have been used for maize plastid transformation because of the presence of developing or developed chloroplasts and good plant regeneration capacity. However, these green cultures are more heterogeneous than embryogenic cultures and contain some developed tissues that are not highly transformable. The additional novel strategy of using greening embryogenic callus as the target material was developed because such cultures contain developing plastids that may be a more amenable target for plastid transformation, but also maintain the high transformation capacity normally attributed to embryogenic cultures. To select for plastid transformants following microprojectile bombardment, different selection regimes have been developed for the different target materials.

In addition, streptomycin or glyphosate selection with green multiple bud or greening embryogenic cultures could be more effective because identification of plant cells resistant to streptomycin relies on the green phenotype of resistant cells, and glyphosate selection could be most effective with the meristematic tissues, the natural sink for glyphosate. Green multiple bud corn cultures can be grown in many ways (U.S. Pat. No. 5,767,368; Zhong et al., 1996 *Plant Physiol.* 110:1097-1107), but any method that produces cells containing chloroplasts or their precursors can be utilized in this invention. The use of "greening embryogenic" cultures has not been reported. Several corn genotypes, including a triple hybrid (Pa91×H99)×A188, H99, Honey and Pearl, HiII and LH198×HiII were used in the following experiments. However, any corn genotype could be used by one of skill in the art.

Because plant cells contain a large copy number of plastid genomes, an effective selectable marker and selection regime are very important for selecting homoplasmic transformants. Selectable markers that have been used for plastid transformation include *aadA*, which confers resistance to the antibiotics spectinomycin and streptomycin, and the CP4 gene that confers resistance to the herbicide glyphosate. In the case of streptomycin, the antibiotic is not lethal to plant cells. Selection relies on green phenotype of resistant cells. In contrast, glyphosate is a plastid-lethal marker and destroys plastid membrane structure at the lethal concentration. Effective selection therefore may be facilitated by a sub-lethal selection scheme, as outlined in US Patent Application 200242934.

## EXAMPLES

The following examples further illustrate the present invention. They are in no way to be construed as a limitation in scope and meaning of the claims.

### Example 1

#### Vector Constructions

Plastid vectors optimally contain several critical elements: a highly expressed selectable marker, a reporter gene for easy screening and identification of transgenic plastids, and a

transgene insertion site that does not negatively influence expression or function of the transgene or the neighboring endogenous plastid genes.

#### Insertion Sites:

Plastid vectors are designed to target transgene integration into the plastid genome via homologous recombination. The location of transgene insertion must be chosen such that the insertion does not cause any disruption of normal plastid function. This is achieved by cloning the transgenes into a plastid intergenic region where no open reading frames exist, preferably such that readthrough transcription from the transgenes into neighboring resident plastid genes is avoided. Two plastid genomic locations are targeted for insertion of transgenes: the Large Single Copy region and the Inverted Repeat region. Because the Inverted Repeat region is present in two copies per genome, the transgenes will also be present in two copies in transformed lines.

#### a) Large Single Copy Region

In tobacco, the site between the *rbcL* and *accD* genes in the tobacco Large Single Copy region was shown to be a successful insertion site (Svab and Maliga, 1992, *ibid.*). Based on success in tobacco, we also chose the maize plastid genomic region downstream of *rbcL* as the site of insertion for maize plastid transgenes. However, the gene order between tobacco and maize in this genomic region is different. Complete nucleotide sequencing of the maize plastid genome (Maier et al., 1995 *J. Mol. Biol.* 251:614-628; Genbank accession X86563) revealed that a large inversion occurred during evolutionary separation of the monocot and dicot plastid lineages such that the region downstream of *rbcL* is completely different between tobacco and maize. Therefore, insertion of transgenes in this region requires thoughtful identification of a suitable non-coding, intergenic region.

An ~3.4 kb region surrounding the maize plastid *rbcL* gene was cloned for use as a homologous targeting sequence to direct transgene insertion. First, an ~2.8 kb PstI/SpeI DNA fragment from plasmid pZmc301 (gift of Dr. Ralph Bock, Freiburg, Germany) that carries a portion of the *rbcL* coding region and downstream *psaI* and *orf185* genes was sub-cloned to create plasmid pMON49204. Then, a further sub-clone of pMON49204, an ~1 kb BamHI/PstI fragment, was generated and used as template for PCR mutagenesis to create an engineered NotI site used for subsequent insertion of plastid transgenes. The NotI site was created by overlap PCR mutagenesis using PCR primers (SEQ ID NO:1-4). The newly created NotI site and surrounding region was verified by sequencing and then reinserted back into pMON49204 as a BamHI/PstI fragment, to create plasmid pMON49206. Lastly, an additional ~560 bp of homologous flanking sequence, carrying a further portion of the maize *rbcL* gene, was PCR amplified from total maize genomic DNA using PCR primers (SEQ ID NO:5-6). The amplified fragment was cloned next to pMON49206 to create the Single Copy Region transformation vector pMON49219 (FIG. 1). The nucleotide sequence of the cloned ~3.4 kb region is shown in SEQ ID NO:7.

Translation of the ~3.4 kb sequence in all six possible reading frames identified the known *rbcL*, *psaI* and *orf185* genes. In addition, several small open reading frames of less than 50 amino acids were identified. To facilitate cloning of transgenes into the homologous flanking region, the unique NotI restriction site was engineered ~580 bp downstream of *rbcL* in the *rbcL*-*psaI* intergenic region, away from any small open reading frames. The NotI insertion site is flanked by ~1.5 kb and ~1.8 kb of homologous DNA on either side of the transgenes to direct integration into the plastid genome by

homologous recombination. Plasmid pMON49219 shown in FIG. 1 carries this ~3.4 kb plastid fragment with the engineered NotI site.

#### b) Inverted Repeat Region

The second targeting location for insertion of maize transgenes is upstream of *trnV*, in the intergenic region between the *trnV/rn16* operon and the divergently transcribed *rps12/3'-rps7* operon. This region is located in the Large Inverted Repeat region and is therefore present in two copies per plastid genome. The analogous region is routinely used for transgene insertion in tobacco (Staub and Maliga, 1993; Zoubenko et al., 1994 *Nucleic Acids Res.* 22:3819-3824). However, the sequence and gene content at the site of insertion differs between tobacco and maize.

An ~4.8 kb plastid DNA fragment was chosen for PCR amplification and cloning as the maize homologous flanking region. The sequence of the fragment is shown in SEQ ID NO:8. For amplification of the *rn16/trnV* gene flanking region, one PCR primer (5'-gggtaccgtcgacttacagggcgcgctctaccactgagctaataagccc-3') (SEQ ID NO:9) was designed after the *trnI* gene coding region that lies downstream of the *rn16* gene. Note that sites for the restriction enzymes KpnI and Sall were incorporated to the end of the PCR primer for subsequent cloning purposes. The second PCR primer (5'-taagcttgccggccgccggtgacaattgaaatccaattttccattatttgac-3') (SEQ ID NO:10) was designed to anneal in the *rn16/trnV-rps12/3'-rps7* intergenic region and also encodes a unique engineered NotI restriction site (underlined). For amplification of the *rps12/3'-rps7* gene flanking region, one primer was designed after a part of the *rps7* gene coding region (5'-aactagtcggccaccatgttaactaatcgattacgaaaattggatcgg-3') (SEQ ID NO:11). Note that SpeI and SmaI sites were incorporated onto the end of the PCR primer for subsequent cloning purposes. The second primer was designed to also anneal in the same location in the *rn16/trnV-rps12/3'-rps7* intergenic region, and also encode the unique NotI site (5'-ggaattcgccggcgcctctatcgaaaataggattgactatggattcgagcc-3') (SEQ ID NO:12). As a result of the PCR amplifications, the unique NotI insertion site was created approximately mid-way between the ~4.8 kb homologous flanking region, with ~2.3 kb and ~2.5 kb of homologous DNA on either side of the transgene insertion site. The pMON38722 clone with this maize flanking region and unique NotI site is shown in FIG. 2.

In tobacco, the transgene insertion site in the analogous region of the plastid genome disrupts an open reading frame of no known function that is not found in other plant species. Insertion into this location in tobacco has no deleterious effect (Staub and Maliga, 1992, *ibid.*). In maize, two different unidentified open reading frames, each with no known function, are present in the analogous position. Therefore, it is unknown whether insertion into these open reading frames would affect plastid gene functions. To avoid any possible effect on plastid gene function, the insertion site was chosen in the intergenic region between the *trnV* gene and the nearby ORF85 gene, away from any putative plastid gene regulatory elements.

#### Selectable Markers:

For selection of plastid transformed cells, the *aadA* gene that gives resistance to spectinomycin and streptomycin or the CP4 gene that confers tolerance to glyphosate was chosen.

For plastid transformation vectors carrying *aadA*, streptomycin was used as the selective agent because maize plastids are naturally resistant to spectinomycin. Selection using this antibiotic is based on inhibition of plastid protein synthesis, which prevents accumulation of photosynthetic proteins and chlorophyll, thus resulting in bleaching. Resistant cells are

identified by their green color on selective media. Therefore, this antibiotic can only be used with light-grown, chlorophyll-containing green tissue culture systems.

Glyphosate is also used as a selective agent. Glyphosate inhibits aromatic amino acid biosynthesis but also has more pleiotropic effects including bleaching and inhibition of growth. Resistant cells are differentiated by growth and by greening if grown in the light. Therefore, this selectable marker is useful for both dark-grown or light-grown tissue culture systems.

Based on observations with glyphosate selection in tobacco, most selection regimes have focused on transitions from sub-lethal to lethal concentrations of the selective agent. Selection using this approach has been described in US Patent Application 200242934. It is also possible to use a steady-state concentration of glyphosate throughout the selection period, enough to inhibit growth of cells in culture but only gradually become lethal.

#### Transgene Expression Elements:

For expression of the *aadA* or CP4 transgenes, expression signals that provide for constitutive transcription and translation of the transgene are most desirable. The expression signals need to be derived from resident maize plastid genes to ensure that the appropriate trans-acting factors are present for faithful gene expression. In most cases, we have used the maize 16S ribosomal RNA (*rrn16*) operon promoter (*ZmPrn*) (SEQ ID NO:13) to drive transgene expression. This promoter region includes the mapped transcriptional start site of the *rrn16* operon (Strittmatter et al., 1985, EMBO J, 4 (3): 599-604).

For translation, several different elements are being tested. The plastid *atpB* gene leader and first 14 amino acids of its coding region (gift of Dr. Pal Maliga), fused to the *ZmPrn*, are being tested. The encoded b-subunit of plastid ATPase accumulates in all plastid types, therefore it was predicted that the *atpB* gene leader would provide for constitutive translation. Furthermore, the 14-amino acid translational fusion was shown to enhance translation in some cases (Pal Maliga, personal communication).

The heterologous bacteriophage T7 gene 10 leader (G10L) is also being tested. This element has been shown to enhance translation in plastids (Staub et al., 2000 Nature Biotech. 18:333-338; U.S. patent application Ser. No. 09/113,690). In some cases, an additional translational fusion of 14 amino acids of the green fluorescent protein (14aaGFP), that has also been shown to enhance translation (U.S. patent application Ser. No. 09/351,124), is used in addition to the G10L. In other cases, the "downstream box" sequence from the bacteriophage T7 gene 10 coding region (EC DB), which also enhances translation (U.S. patent application Ser. No. 09/668,188), is used in addition to the G10L.

#### Screenable Marker:

To facilitate identification of maize plastid transformants, a GFP transgene was included in the transformation vectors. GFP can be monitored in live tissue and used to follow the growth of transplastomic sectors (Sidorov et al. 1999 Plant Journal 19:209-216; U.S. patent application Ser. No. 09/351,124). The GFP transgene was also driven by the *ZmPrn* promoter. The translational enhancer sequence, G10L (Staub et al., 2000, *ibid.*) was included to ensure constitutive high level translation.

#### Plastid Vectors:

pMON49219 (FIG. 1): maize genes *rbcL*, *psaI* and ORF185 Large Single Copy flanking region clone, including the unique NotI site used for transgene insertion.

pMON38722 (FIG. 2): maize genes *rrn16* (16S rDNA), *trnV* (tRNA-Val), -ORF85 ORF58, and *rps12* Exons I and II

in the Inverted Repeat flanking region clone, including unique NotI site used for transgene insertion.

pMON56102 (FIG. 3): *ZmPrn*:G10L (SEQ ID NO:14): GFP:NtTrps16 (SEQ ID NO:15) plus NtPrn.atpB+DB (SEQ ID NO:16): *aadA*:*ZmTpetD* (SEQ ID NO:17) in the same direction as the *rbcL* gene in the pMON49219 flanking region.

pMON49296 (FIG. 4): *ZmPrn*:GFP:*ZmTrbcL* (SEQ ID NO:18) plus *ZmPrn*:G10L:14aaGFP (SEQ ID NO:19) -*aadA*:*ZmTpetD* in the same direction as the *rbcL* gene in the pMON49219 flanking region.

pMON56108 (FIG. 5): *ZmPrn*:G10L:GFP:NtTrps16 plus *ZmPrn*:G10L:EC DB-*aadA*:*ZmTpetD* in the same direction as the *rbcL* gene in the pMON49219 flanking region.

pMON56110 (FIG. 6): *ZmPrn*:GFP:*ZmTrbcL* plus *ZmPrn*:G10L:EC DB-CP4-:*ZmTpetD* in the same direction as the *rbcL* gene in the pMON49219 flanking region.

pMON56113 (FIG. 7): *ZmPrn*:G10L:GFP:NtTrps16 plus *ZmPrn*:G10L:EC DB-*aadA*:*ZmTpetD* upstream and in the same direction as the *rrn16* gene in the pMON38722 flanking region.

pMON56114 (FIG. 8): *ZmPrn*:G10L:GFP:NtTrps16 plus *ZmPrn*:G10L:EC DB-CP4:*ZmTpetD* upstream and in the same direction as the *rrn16* gene in the pMON38722 flanking region.

#### Nuclear Transformation Vectors

Control nuclear transformations are valuable to determine ideal selection conditions and frequency of transformation using new maize explant types. Furthermore, transient nuclear expression of reporter genes can assist in developing optimal bombardment parameters. For these reasons, nuclear transformation vectors are routinely employed: pMON38762 (FIG. 9): GFP (green fluorescent protein) gene containing the *hsp70* intron, driven by the 35S promoter and the nos3' terminator plus the streptomycin resistance gene (*aadA*) containing the *rac* intron, driven by the 35S promoter and the *hsp17* 3' terminator.

pMON30098 (FIG. 10): GFP gene driven by e35S promoter and HSP70 intron and nos 3"-terminator.

pMON25496 (FIG. 11): CP4 gene with *Arabidopsis* EPSPS chloroplast transit peptide driven by rice actin promoter and rice actin intron plus CP4 gene with *Arabidopsis* EPSPS chloroplast transit peptide driven by 35S promoter with maize HSP70 intron.

pMON38762 (FIG. 12): *aadA* gene driven by e35S promoter with *cab* gene leader and rice actin intron plus GFP gene driven by e35S promoter and HSP70intron.

pMON45294 (FIG. 13): *aadA* gene driven by e35S promoter with *cab* gene leader and rice actin intron plus GFP gene with *Arabidopsis* EPSPS chloroplast transit peptide driven by e35S promoter and HSP70intron.

#### Example 2

##### Initiation and Establishment of Green Multiple Bud Cultures as Target Materials

#### Plant Materials:

For the present invention, plants of hybrids Pa91×H99 were grown in a greenhouse or in the field with normal growth conditions for maize plants. Plants of Pa91×H99 were pollinated with pollen from line A188 plants to form embryos with genotype of (Pa91×H99)×A188 (also known as triple hybrid, or PHA). The hybrid ears were harvested approximately 10 days after pollination. Immature embryos were isolated the same day when the ears were harvested or after the ears were stored in a refrigerator (4° C.) for one to several days.

Establishment of Light-Grown Green Multiple Bud Cultures:

Green multiple bud cultures (also sometimes called green meristematic cultures) can be established from a variety of tissues. In some cases, green multiple bud cultures are established from dark-grown embryogenic callus that is transferred to light and maintained for several subcultures on MS2 or MS3 medium (Table 1), transferring green meristematic zones at each subculture until a homogeneous callus is established. For the present invention, the meristematic regions of young seedlings derived from immature embryos were used to establish the green multiple bud callus cultures as described below.

Immature embryos of H99 or (Pa91xH99)xA188 were isolated from sterilized ears and placed on CM4C medium (Table 2) with scutellum side touched to medium. The embryos were kept in the light culture room or Percival incubator (16-h light and 8-h dark, 27° C.). In 5-7 days the growing seedlings were approximately 1-3 cm long. A section (approximately 5 mm long) containing the nodal region (with apical meristem and leaf primordia) was dissected and laid horizontally on MS2 or MS3 medium. Initially the primary shoots continued to elongate and were trimmed weekly. Eventually, the meristematic regions enlarged and formed clusters of shoots/shoot primordia in approximately 6 weeks. The cultures were green, highly regenerable and considered meristematic. They were proliferated and maintained under the same growth conditions by breaking into smaller pieces and transferring onto fresh medium every 3-4 weeks. The cultures on MS2 and MS3 media were similar morphologically, except that the cultures were more compact on MS2 and more differentiated on MS3.

TABLE 1

Media used for inducing organogenic cultures from embryogenic callus cultures and meristem region.	
Medium	Reference and description of the medium
MS2	MS medium (Murashige and Skoog, 1962 <i>Physiol. Plant</i> 15:473-479) supplemented with 40 g/l maltose, 500 mg/l casein hydrolysate, 1.95 g/l MES, 2 mg/l BA, 0.5 mg/l 2,4-D, 100 mg/l ascorbic acid, pH 5.8.
MS3	The same as MS2, except that BA is 1 mg/l and 2,4-D is replaced by 2.2 mg/l picloram.

TABLE 2

Callus induction media used for inducing callus from immature embryos.		
Embryo genotype	Callus induction medium	Reference and description of the medium
H99 (Pa91xH99)xA188	CM4C	Cheng et al., 1997 <i>Plant Physiol.</i> 115:971-980.
LH198xHiII	N6 1-100-12	N6 medium supplemented with 1.38 g/L proline, 100 mg/L casein hydrolysate, 20 μM silver nitrate, 1 mg/L 2,4-D, 0.125 mg/L Cu, 0.125 mg/L Co and 1.25 mg/L Mo.

In some cases, the genotype Honey and Pearl was used. In this case, mature seeds were purchased from a commercial source, surface sterilized and then placed on MSOD medium (MS medium with 40 g/L maltose, 20 g/L glucose, 150 mg/L asparagine, 100 mg/L inositol, 0.65 mg/L niotinic acid, 125

mg/L pyridoxine, 125 mg/L thiamine, 125 mg/L pantothenate, and 7 g/L Phytagar) and germinated in the dark for 4 days. After 4 days, the meristem region is excised about 3 mm below and 2 mm above the nodal area. The explants are then placed standing up in MS3 media containing 1 mg/L BAP and 2.2 mg/L picloram in the light at 26° C. for 18 hr of light. Primary shoots are then trimmed every 3-4 days until multiple buds are visible (about 3-4 weeks). At this time, the leaves on top of the buds are trimmed off, the buds are separated and then subcultured every three weeks.

#### DNA Delivery by Particle Gun Bombardment

##### Preparation of Target Materials for Bombardment

Actively proliferating meristematic areas from the green callus were selected, and leaves or leaf primordia were removed from one to a few days prior to particle gun bombardment. The selected calli were then transferred into the middle of the plates with solid MS3 or MS2 medium. The prepared plates were incubated at light conditions.

##### Precipitation of Plasmid DNA with Particles and Bombardment:

The plasmid DNA was precipitated with gold (0.4 or 0.6 μm) or tungsten particles according to standard operation procedure for helium gun. Particles (0.5 mg) are sterilized with 100% ethanol, washed 2-3 times each with 1 mL aliquots of sterile water. They are then resuspended in 500 μL of sterile 50% glycerol. Twenty-five μL aliquots of particle suspension are added into 1.5 mL sterile microfuge tubes, followed by 5 μL of DNA of interest (at 1 mg/mL) and mixing by finger-vortexing. A fresh CaCl<sub>2</sub> and spermidine premix is then prepared by mixing 2.5M CaCl<sub>2</sub> and 0.1M spermidine at a ratio of 5 to 2. Thirty-five μL of the freshly prepared "premix" is added to the particle-DNA mixture, and mixed quickly by finger-vortexing. The mixture is incubated at room temperature for 20 min. The supernatant is removed after a pulse spin in the microfuge. The DNA-particles mixture is washed twice, first by resuspending in 200 μL of 70% ethanol, followed by pulse spin and removing of the supernatant, repeated with 200 μL of 100% ethanol. The DNA-particles mixture is finally resuspended in 40 μL of 100% ethanol. After thorough mixing, an aliquot of 5 μL is loaded onto the center of the macrocarrier already installed in the macrocarrier holder, and allowed to dry in a low humidity environment, preferably with desiccant. Each tube can be used for 5-6 bombardments.

In comparison to standard protocol, double the concentration of DNA was also used for particle preparation. The plate with the target tissue was bombarded twice using the helium gun (Bio Rad, Richmond, Calif.) using the protocol described by the manufacturer, at target shelf levels L3. The gap distance was set at 1.0 cm and the rupture pressures at 1100-1550 psi.

##### Selection of Plastid Transformants in Green Multiple Bud Cultures Using *aadA* as a Selectable Marker

DNA vector pMON56102 (FIG. 3) containing *aadA* and *gfp* genes (see Example 1) was used to bombard the (Pa91xH99)xA188 target tissues. After bombardment, the material remained on the same medium overnight (16-20 h) in the light culture room or in a Percival incubator. They were then transferred onto selection medium of MS3 with 100 mg/L streptomycin. The cultures were transferred onto fresh selection medium every 3-4 weeks until highly resistant callus colonies or sectors were observed or all the cultures were dead. During selection, the cultures were examined periodically under a Leica MZ80 GFP plus fluorescence microscope for GFP expression.

MS3 medium with 100 mg/L streptomycin was used for selection. Twenty-two plates with light-grown (Pa91xH99)x

A188 multiple bud culture were shot with pMON56102. After 4 weeks of selection, all plates were analyzed. In one plate a leaf arising directly from callus tissue with a positive GFP sector was observed. This leaf was sectored, having a linear green (streptomycin resistant) sector surrounded by white (streptomycin sensitive) sectors oriented from the base to the tip of the leaf, although the green sector made up the bulk of this leaf. Using fluorescence microscopy, GFP positive cells were identified from the base to the tip of the leaf located around the midvein of the leaf, coincident with the location of the green sector. The remainder of the leaf, coincident with the location of the white sectors, had red fluorescence due to the chlorophyll autofluorescence in untransformed cells.

Samples were also analyzed by laser scanning confocal microscopy. In control nuclear transformants expressing a non-targeted GFP gene, the GFP fluorescence is distributed all over the cytoplasm. The chloroplasts emit only red autofluorescence. The analysis of the putative transformed leaf confirmed that GFP fluorescence is localized in chloroplasts within the green sector. Confocal microscopy also confirmed that the recovered transformant is chimeric, carrying mostly homoplasmic cells in the green sector, but also some heteroplasmic cells at the junction between the green and white sectors.

The multiple bud callus with a small GFP positive area was transferred to selective medium with 50 mg/L streptomycin in an effort to amplify the transformed tissues. Although these cells and sectors did not regenerate into whole plants, it is expected that a plant could be regenerated from the protocol. More cells and sectors need to be obtained and evaluated.

#### Molecular Analysis of the Transformation Event

A small portion of the transformed leaf sector was sacrificed for DNA extraction and used for PCR analysis. The corn leaf sample was ground in 150  $\mu$ L of CTAB and incubated at 65° C. for 30 min. The mixture was then cooled to room temperature and extracted with 150  $\mu$ L of chloroform: isoamyl alcohol (24:1). The mixture was spun at 14,000 rpm for 10 min. The supernatant was collected to the new tube and two volumes of 100% ethanol were added. The solution was then kept at -20° C. for 30 min to precipitate the nucleic acids. DNA was spun down at 14,000 rpm for 10 min. The DNA pellet was then washed with 75% ethanol, air dried, and dissolved in 24  $\mu$ L of water.

PCR reactions were performed using Roche Expand Long Template PCR System. Two microliters of the above DNA solution was used as the template. The other components were used according to the manufacturer's recommendations. The PCR mixture was first denatured at 94° C. for 2 min, then repeated 35 cycles of 94° C. for 10 sec, 53° C. for 30 sec and 68° C. for 2 min, and at last elongated at 68° C. for 10 min. The PCR products were then separated on 1% agarose gel.

FIG. 14 shows the PCR primers used and the expected length of DNA products. The vertical dashed line indicates the junction of the foreign DNA insert on the transforming DNA and the flanking wild-type plastid DNA (ptDNA). The results show that the transforming DNA has integrated into the maize plastid genome. Specifically, the PCR 61/PCR62 (SEQ ID NO:22 and SEQ ID NO:23), PCR60/PCR65 (SEQ ID NO:21 and SEQ ID NO:26), and PCR60/PCR66 (SEQ ID NO:21 and SEQ ID NO:27) reactions show the predicted bands of 1941, 2280, and 2201 bp, respectively, that are only possible if the foreign DNA has integrated into the correct genomic location. The other PCR products show that the various portions of the transforming DNA are also intact, as expected for a homologous integration event.

A further experiment using a green multiple bud explant has produced at least one plastid transformed callus. The experiment utilized light-grown (Pa91xH99)' A188 multiple bud culture maintained on MS3 medium.

In this experiment, a cobombardment protocol was used, with both a nuclear transformation vector encoding a GFP gene (pMON30098; FIG. 10) and a plastid transformation vector encoding both aadA and GFP genes (pMON49296; FIG. 4). The GFP gene on the pMON30098 vector is driven by the e35S promoter and has the hsp70 intron and therefore can only be expressed in the nucleus. This vector was included to monitor "transient" GFP expression useful to isolate those callus pieces that were productively hit during the transformation bombardment. After bombardment, selection for plastid transformants on streptomycin medium is achieved via the pMON49296 plastid-encoded aadA gene.

For bombardments, the standard DNA precipitation protocol was used, except that twice the amount of pMON30098 (2  $\mu$ g/ $\mu$ L) was used relative to the pMON49296 (1  $\mu$ g/ $\mu$ L) vector. Gold particles of 0.4  $\mu$ m were used, a rupture disc pressure of 1350 psi, plate level 3 and 2 shots per plate. Tissues were placed on MS3 plus osmoticum prior to bombardment. After a one-day delay, bombarded tissues were transferred to MS3 plus streptomycin. A total of 24 plates of green multiple bud tissues were bombarded.

After the one-day delay period, all callus tissue was transferred to MS3 plus 100  $\mu$ M streptomycin for selection. At ~4-week intervals, the callus tissue was subcultured to fresh MS3 medium plus streptomycin. Prior to the first subculture, the callus was observed under the fluorescence microscope for GFP activity. Callus pieces were divided into two types: good GFP expressors (treatment #1) and poor or no GFP expressors (treatment #2). It should be noted that GFP expression in this case was expected to be from the nuclear-encoded GFP gene, as transient expression from the plastid-encoded GFP gene is not expected to be visible at this early timepoint.

Callus from treatment #1 was first subcultured to 200, 500 or 1000  $\mu$ M streptomycin. At the second subculture ~4 weeks later, all calli were subcultured to 1000  $\mu$ M streptomycin. For treatment #2, at the second subculture, all callus was subcultured directly to 1000  $\mu$ M streptomycin and maintained at that concentration throughout subsequent subcultures.

After ~5 months in culture, essentially all of the callus had turned yellow and appeared to be nonregenerable or was brown and dying. However, small GFP positive sectors of ~2 mm to ~4 mm were observed on 17 independent callus pieces. GFP fluorescence was observed in 11 callus pieces from treatment #1 and 6 callus pieces from treatment #2.

GFP fluorescence could arise from nuclear transformation of the pMON30098 construct and expression of its encoded GFP gene, from the plastid GFP gene on pMON49296 but from a nuclear insertion and fortuitous expression from a nuclear promoter, or from the plastid GFP gene after predicted insertion into the plastid genome and plastid expression. To prove plastid-localized expression of GFP in the callus lines, confocal microscopy was performed on several lines as shown in Table 3. Of 9 samples tested, confocal sample #6 (callus number 110-9) had exclusively a plastid localization of green GFP fluorescence. In addition to the predominantly green GFP fluorescence of plastids in sample 6, a few plastids showing only red chlorophyll autofluorescence were observed. These results indicate that callus 110-9 is a plastid transformant but still contains some wild-type plastids and is therefore heteroplasmic. Confocal microscopy sample #7 (callus number 110-5) showed some plastid-localized GFP expression, but also carried GFP localized to cell

walls. This may indicate that this sample was transformed in both the plastid and the nucleus, but no further analysis of this sample was attempted.

TABLE 3

Calli Number	Treatment #	GFP +/-	confocal sample #	GFP localization
110-1aa	1	+	3	cell wall
110-2	1	+	1	cell wall
110-3	1	+	4	cell wall
110-5	1	+	7	cell wall/plastid
110-8	1	+	5	cell wall
110-9	1	+	6	Plastid
110-13b	2	+	8	cell wall
110-16b	2	+	9	cell wall
110-17	2	+	2	cell wall

To confirm plastid transformation in callus 110-9 by molecular means, PCR analysis was performed. First, PCR primers were designed that distinguished between the nuclear GFP gene on plasmid pMON30098 (PCR pair GFP5":1/GFP3":2) (SEQ ID NO:28 and SEQ ID NO:29) and the plastid GFP gene on plasmid pMON49296 (PCR pair 202/201) (SEQ ID NO:30 and SEQ ID NO:31). Amplification of the 536 bp plastid GFP gene fragment was successful, whereas the nuclear GFP gene fragment did not amplify in callus 110-9. This result indicates that the plastid-localized GFP arises from expression of the gene from pMON49296, and therefore verifies plastid transformation in this sample. Additional PCR confirmation of plastid transformation in callus line 110-9 was obtained using primer pairs 198/199 (SEQ ID NO:32 and SEQ ID NO:33), 200/194 (SEQ ID NO:34 and SEQ ID NO:35) and 202/194 (SEQ ID NO:30 and SEQ ID NO:35) to successfully amplify the 492 bp *aadA* coding region fragment, the 545 bp fragment overlapping the GFP and *aadA* genes, and the 1065 bp larger fragment overlapping the GFP and *aadA* genes, respectively. Taken together, these results show that both the GFP and *aadA* transgenes reside in the plastid genome of the 110-9 line (FIG. 14).

#### Selection of Plastid Transformants in Green Multiple Bud Cultures Using CP4 as a Selectable Marker

Plastid transformation vectors pMON56110 (FIG. 6) or pMON56114 (FIG. 8) (containing CP4 and GFP genes), pMON53161 (containing CP4 gene only) and pMON56101 (containing CP4 and *aadA* genes) were used for transformation of PHA green multiple bud cultures and glyphosate selection. Actively growing green callus pieces that are one to two weeks post subculture on fresh MS2 or MS3 medium were used as target material. Prior to bombardment, the leaves and leaf primordia were carefully removed. Callus pieces were arranged in the center of a plate (about 1.5 inches in diameter) containing MS2 or MS3 medium with or without osmoticum (0.2 M mannitol). In the case where osmoticum was used, the culture was incubated on osmoticum for 3-4 hr prior to bombardment. Post bombardment, the materials were removed from osmoticum to MS3 or MS2 medium the next day if osmoticum was used, or they were left on the same medium. After a delay time of 1-5 days, they were transferred to the same medium containing 10-25  $\mu$ M of glyphosate for 2 weeks. They were then moved to 100-250  $\mu$ M of glyphosate for the remaining selection period. The cultures were transferred to fresh selection medium at 3-4 week intervals. After 3-4 rounds of selection at the higher levels of glyphosate, most of the tissues turned yellow to brownish and eventually died. The tolerant, green pieces were moved to hormone-free medium without glyphosate for regeneration. It is also pos-

sible to use a steady-state concentration (100  $\mu$ M) of glyphosate throughout the selection period, which is enough to inhibit growth of cells in culture but only gradually becomes lethal.

5 Samples of the tolerant lines are analyzed by PCR or Southern blotting at callus or plant stage to confirm the nature of tolerance.

#### Recovery of Plants from Putatively Transformed Sectors

Putatively transformed sectors are normally kept on selection to allow them to continue growth so that they can be regenerated. Under such circumstances, a couple of approaches will be taken to rescue the putative sectors. First, because the *gfp* gene is included in most of the vectors, GFP can be used as a "screenable" marker. The GFP positive sectors can be isolated under a dissecting microscope and placed on medium without any selective agent to allow the sectors to recover and to continue to grow by monitoring GFP expression; or the dissected sectors can be placed on media alternating with and without the selection to allow continued growth of the sectors without running the risk of losing the transformed copies of the plastid genomes; or the dissected sectors could be put on medium with low levels of selection to balance the growth and maintenance of the transformed copies. Second, the dissected sectors could also be placed on top of nurse cultures, which are resistant to glyphosate, in the presence of glyphosate for a period of time to help the putative sectors to grow and to amplify its transgenic copies of the plastid genome.

#### Example 3

##### Greening Embryogenic Callus as Target Material

Light-grown multiple bud cultures have been shown above to be successful for plastid transformation. However, the frequency of transformation in the light-grown cultures is very low. In contrast, dark-grown embryogenic callus is typically highly transformable when nuclear transformation is performed, but these cultures contain only undeveloped plastids and so may not be ideal for plastid transformation. "Greening" embryogenic cultures that contain more developed plastids but still maintain the high transformability of dark-grown cultures would be advantageous and may be useful as target materials for plastid transformation.

To establish greening embryogenic cultures, dark-grown embryogenic callus must first be initiated. Young ears of inbred or hybrid maize lines (e.g., H99, PHA for Type I embryogenic callus and HiII or LH198 $\times$ HiII for Type II embryogenic callus) are sterilized in 50% bleach (bleach contains 5.25% sodium hypochlorite) for 20 min and washed with sterile water 3 times. Immature embryos are isolated and cultured with scutellum side facing up on callus induction media, which differs depending on the plant genotype as shown in Table 1. The embryo cultures are kept in the dark culture room at 26-27° C. Embryogenic callus that develops from the embryos is maintained on the same medium through routine subculture every 2 to 3 weeks.

For greening embryogenic cultures, the dark-grown embryogenic calli are spread out on MS2 or MS3 media and cultured at 26° C. with 16 hr light/8 hr darkness. The somatic embryos start to green up from the embryo side within 3-10 days. The greening somatic embryos are then picked up and used as the target material for bombardment using the helium gun. In some cases, it may be desirable to bombard the dark-grown embryogenic callus prior to incubation in the light and subsequently perform the selection on MS2 or MS3 greening medium in the light.



For streptomycin selection experiments, tissues are maintained on the same medium for a one-day delay period before selection. The tissues are then placed on MS2 or MS3 selection media with 50 or 100 mg/L streptomycin. This selection level allows for shoot development and callus growth at a slightly inhibited level but also allows bleaching of nontransformed tissue. At approximately monthly intervals, the growing tissue is separated from the dead and non-regenerable tissue and the leaves are peeled away from the callus. The callus pieces are then placed on higher concentrations of streptomycin at levels of 100 or 200 mg/L. This level decreases the regenerability of nontransformed cells, greatly reducing the growth of the callus. After further selection for one month, the living green tissue is placed on hormone-free regeneration media. Shoots that emerge as green or GFP positive are further analyzed. In some cases, it may be desirable to increase the streptomycin concentration to 500 or 1000 mg/L for one subculture prior to plant regeneration to further eliminate any possibility of non-transformed cells.

For glyphosate selection experiments, greening type II embryogenic calli (e.g., LH198×HiII or HiII) are selected on MS2 or MS3 media initially with 10-25  $\mu$ M of glyphosate for about 2 weeks, then stepping up to 100-250  $\mu$ M glyphosate. At 100-250  $\mu$ M glyphosate, growth is dramatically inhibited and the non-transformed tissues eventually die.

#### Example 4

##### Dark-Grown Embryogenic Callus Cultures as Target Material

Dark-grown embryogenic callus cultures of various genotypes (e.g., LH198×HiII or HiII) were established from immature embryos as described previously and maintained in continuous dark at 26° C. Callus is subcultured at intervals of about every 2 weeks.

For bombardment of dark grown callus, embryogenic pieces of about 1-2 mm are placed on a filter paper on top of callus induction media. For glyphosate selection, the calli are incubated on osmoticum-containing medium for 3-4 hr prior to bombardment, followed by 1-4-day delay on maintenance medium without glyphosate post bombardment. They are then placed on the first selection of 100-500  $\mu$ M glyphosate for one week. The selection level is subsequently increased to 500  $\mu$ M then 1 mM, until it reaches 3 mM. After several rounds of subculture on 3 mM, distinct, glyphosate-tolerant callus will be obtained. They will then be placed on hormone-free medium for regeneration.

#### Example 5

##### Pre-Cultured Immature Embryos as Target Material Using Glyphosate Selection

Genotypes of LH198×HiII or HiII (derivative of A188×B73) were used as the source for immature embryos. Ten days after pollination, the ears were harvested and within 5 days of the harvest date the immature embryos were excised. The corn ears were sterilized with 50% bleach (v/v) with 2 drops of Tween-20 for 20 min. They were then rinsed three times with sterile distilled water. Using a scalpel, the tops of the kernels were removed. Using a small spatula, the embryo was excised and placed onto N6 media with AgNO<sub>3</sub> and 3% sucrose, with scutellum side away from media. The immature embryos were pre-cultured in the dark at 28° C. for 3 days.

Prior to bombardment, about 50 embryos were arranged into a 2 cm diameter circle onto the same N6 medium plus

osmoticum. Bombardment used M5 tungsten particles and the standard DNA precipitation procedures. Ten  $\mu$ L of DNA/tungsten mix was added to each macrocarrier. Each plate was bombarded twice at sample level L3 (9 cm) and rupture pressure 1350 psi. Approximately two hours after bombardment, the embryos were removed from osmoticum media and placed back onto N6 medium for 5-6 days in the dark at 28° C. After this delay period, the coleoptiles of the embryos were removed and the embryos were transferred to N6 medium with 50  $\mu$ M glyphosate. After two weeks the explants were transferred to N6 plus 500  $\mu$ M glyphosate, and callus was spread with a spatula if needed. Every two weeks the tissue was transferred to fresh selection medium, and cultured at 28° C. in the dark.

#### Example 6

##### Nuclear Transformation Using aadA or CP4 Selection

Light-grown multiple bud culture of PHA was used for the nuclear transformation. The vectors pMON38762 and pMON45294 (see the section describing the vectors) were bombarded using similar parameters. For streptomycin selection, MS3 medium supplemented with 500 and 1000 mg/L of selective agent was used. GFP positive shoots were selected in both cases. Of a total of 224 pieces of the light-grown green multiple bud tissue bombarded with pMON38762 (cytoplasmic aadA), over 10 independently transformed callus lines were obtained by selection on medium containing a combination of streptomycin and spectinomycin each at 1000 mg/L. Transformation frequency was slightly lower with nuclear CTP:aadA construct pMON45294.

Light-grown multiple bud cultures of PHA were also used for glyphosate selection according to the same transformation protocol as and following similar selection regimes to plastid transformation. A CP4 vector and a gfp vector were co-bombarded into the green multiple bud tissue. The former allows for selection of the transformed cells, whereas the later allows for identification and monitoring of the transformed sectors. Following a few rounds of selection at 100-250  $\mu$ M of glyphosate, some green tolerant clones were identified. These clones will be analyzed molecularly to confirm the insertion of the transgenes into the nuclear genome.

For selection of the dark-grown embryogenic calli with glyphosate, LH198×HiII calli were placed on filter paper on osmoticum-containing medium and incubated for 4 hr before bombardment. They were bombarded with 0.6 $\mu$  gold particles coated with equal amounts of pMON30098 (containing GFP) and pMON25496 (containing CP4) and bombarded at 1100 psi according to the Standard Operation Protocol for the Biolistic PDS-1000/He particle gun. After a delay of 4 days on the maintenance medium, the calli were selected on 1 mM glyphosate for 2 weeks, followed by 3 mM for several rounds, each for 2 weeks. At the end of 4<sup>th</sup> round of 3 mM glyphosate, glyphosate-tolerance and GFP expression were recorded. Forty-six percent of the initially bombarded calli were tolerant to glyphosate. Among the glyphosate-tolerant clones, over 40% also showed various levels of GFP expression, ranging from spotty to uniform, from very weak to very strong GFP expression.

All publications and patent applications mentioned in this specification are indicative of the level of skill of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent appli-

cation was specifically and individually indicated to be incorporated by reference.

Although the invention has been described in detail for the purpose of illustration, it is understood that such detail is

solely for that purpose, and variations can be made therein by those skilled in the art without departing from the spirit and scope of the invention which is defined by the following claims.

---

 SEQUENCE LISTING
 

---

<160> NUMBER OF SEQ ID NOS: 35

<210> SEQ ID NO 1

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 1

tgaaatagat cttgatacga tc

22

<210> SEQ ID NO 2

<211> LENGTH: 27

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 2

gcggccgctc aaaagaaagc tattgtg

27

<210> SEQ ID NO 3

<211> LENGTH: 38

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 3

agctttcttt tgagcggccg catccaattt caagttag

38

<210> SEQ ID NO 4

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 4

agtggatcca cttaaaaaaa aaaac

25

<210> SEQ ID NO 5

<211> LENGTH: 27

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 5

ggctcgagct gcggaatctt ctactgg

27

<210> SEQ ID NO 6

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 6

-continued

---

cgcatgtacc tgcagtcgca ttc 23

<210> SEQ ID NO 7  
 <211> LENGTH: 3438  
 <212> TYPE: DNA  
 <213> ORGANISM: Zea mays

<400> SEQUENCE: 7

ggtaccgggc cccccctcga gctgcggaat cttctactgg tacatggaca actgtttggg 60  
 ctgatggact taccagtctt gatcgttaca aggacgatg ctatcacatc gagcccgttc 120  
 ctggggaccc agatcaatat atctgttatg tagcttatcc attagaccta tttgaagagg 180  
 gttctgttac taacatgttt acttccattg tgggtaacgt atttggttc aaagccttac 240  
 gcgctctacg tttggaggat ctacgaattc ccctgctta ttcaaaaact ttccaaggtc 300  
 cgcctcacgg tatccaagtt gaaagggata agttgaacaa gtacggtcgt cctttattgg 360  
 gatgtactat taaacccaaa ttgggattat ccgcaaaaaa ttacggtaga gcgtgttatg 420  
 agtgtctacg cgggtgactt gattttacca aagatgatga aaacgtaaac tcacaacat 480  
 ttatgcgctg gagagaccgt ttcgtctttt gtgccgaagc aatttataaa gcacaagccg 540  
 aaactggtga aatcaagggg cactacttga atgcgactgc aggtacatgc gaagaaatga 600  
 ttaagagagc tgtatttgca aggaattag gggttcctat tgtaatgcat gactacttaa 660  
 caggaggatt caccgcaaat actactttgt ctcatattg ccgcgacaac ggccacttc 720  
 ttcacattca ccgagcaatg catgcagtta ttgatagaca gaaaaatcat ggtatgcatt 780  
 tccgtgtatt agctaaagca ttgcgtatgt cggggggaga tcatatccac tccggtacag 840  
 tagtaggtaa gttagaaggg gaacgcgaaa taactttagg ttttgttgat ttattgcgcg 900  
 atgattttat tgaaaaagat cgttctcgcg gtatctttt cactcaggac tgggtatcca 960  
 tgccagggtg tataccggtg gcttctgggg gtattcatgt ttggcatatg ccagctctga 1020  
 ccgaaatctt tggagatgat tccgtattac aatttgggtg aggaacttta ggacatcctt 1080  
 ggggaaatgc acctggtgca gcagctaadc gtgtggcttt agaagcctgt gtacaagctc 1140  
 gtaacgaagg gcgcatctt gctcgtgaag gtaatgaaat tatcaaagca gcttgcaaat 1200  
 ggagtgctga actagccgca gcttgtgaaa tatggaagga gatcaaattt gatggtttca 1260  
 aagcgatgga taccatataa aataaaaaaa aaagcaaat agaaagagaa aaaataagtt 1320  
 aagaaatgca gtaattcttc tttattcttc taattgattg caattcaatt cggctcaatc 1380  
 ttttctaaaa aaaaaaaga ctgagccgaa ttgaaataga tcttgatagc atcatgagac 1440  
 ttgacaaatc gagattcttc tattctatat atctagaata tagaaaggta taatacaata 1500  
 aacaaatata aatcaaaata tagtattatc gtacgataat ggaatcaaat acgcagtatt 1560  
 tacagaaaag agtcttcggt tattgggaaa gaatcaatat acttttaatg tcgaatcggg 1620  
 attcactaag acagaaataa agcattgggt cgaactcttc tttggtgtta aggtagtagc 1680  
 tgtgaatagc catcgactac ccggaaaggt agaagaatgg gacctattct gggacataca 1740  
 atgcattaca gacgatgat cattaccctt caactgggta ttctattcca cttctacttt 1800  
 ttaaattaaa gggattctg aaagaggtat ttcttttatt ttcacaatag ctttcttttg 1860  
 agcggccgca tccaatttca agttcgatta gaaggataga aaggcgatga gaatgggaaa 1920  
 agaaaaataa aatattttat tttaatgagt tccccttttg agttcccctt ttttgcaatt 1980  
 ttcttattat caattccatt catttatttt atagaatact tttttattct atatctataa 2040  
 aaaatcttta ttttgtaaac taaaaaac aatagtcaat attccttata atagatatac 2100

-continued

---

```

ttaattatat cataagaatc ttaagatata tttcgaatag atagaaatag taaatttgaa 2160
ttgagacaca tattctatga cagattttaa cttaccctct attttegtac ctttagtagg 2220
cttagtattt ccggcaattg caatgacttc cttatttctt tatgtgcaga aaaataagat 2280
tgtctagaaa cgacggggcc aaattttctc aatgtatttc caggatcata atacagatat 2340
ttttttgtgt aagtactata atatgatagg gtatgtagct ctttctacac acaaatgaaa 2400
aacgtctatg gatgcagata taagctacga gcataaatgc atgcatatgc gtagccgact 2460
atagcgagtt ttttttttta agtggatcca ctaatttttt ttgaaatagaa agtcaatgta 2520
tctaaccaat tatttcacag gagtactagc tactgaaggc tatttcagaa tcaaaaaaag 2580
taaagtgaaa aatcatttag cttattctct caatttcaat tgaccgctac tggatttagt 2640
atatctaata tgaattggcg atcagaacac atatggatag aacttctaaa aggttctcga 2700
aaaagaggta attttttctg ggctgtatt ctttttctag gttcattagg attcttagcg 2760
gttggggctt ccagttatct tggtagaat atgatatctg tacttccatc tcaacaaatt 2820
cttttttttc cacagggggc cgtgatgtct ttctacggaa tcgcgggcct attcattagc 2880
tcctatttgt ggtgcactat tttatggaat gtaggcagtg gttatgaccg attcgataga 2940
aaagagggaa tagtgtgcat ttttcgttgg ggattccctg gaataaaacg tcgcatcttc 3000
cttcaattcc ttgtgcggga tatccaatca attagaattc aggttaaaga gggcttttat 3060
cctcgtcgta tcctttatat ggaaatccgg ggccaggggg tcattccctt gactcgtact 3120
gatgagaagt tttttactcc acgagaaatt gaacaaaaag ctgccgaatt ggctatttc 3180
ttgcgtgtac caattgaagt attttgagta ccaattgcaa ttttttgcaa ttgtaagggg 3240
attttcgagt atttatctaa agtaaaggaa ggaacaaacg aggataagag aaaattgctt 3300
cgaatttggt ttgtccaagt gagatatatg gcatactatt cttccatttt tatacgaag 3360
gccttttttt atttctctat tccactccat ttagatctaa gaaagaacct aatacaatga 3420
aattccacta gttctaga 3438

```

```

<210> SEQ ID NO 8
<211> LENGTH: 4852
<212> TYPE: DNA
<213> ORGANISM: Zea mays

```

```

<400> SEQUENCE: 8

```

```

ggtagcgtcg acttatcagg ggcgcgctct accactgagc taatagcccg tcgcgcggcc 60
tcccaaaggg aggccgctac gccaaaagcg agaaaaactc catcccttcc cttttgacat 120
ccccatgccg ccacacgggg ggacatgggg acgtcaaaaa ggggatccta tcaactatcaa 180
ctaatttggt ccgacctagg ataataagct catgagcttg gtcttacttc accctaaacg 240
aaagaagact tccatatcca agtttagctc agacgtagct gccttctttt tgggcgtgaa 300
gcagtgtcaa accaaaatac ccaataagca taagcattag ctctccctga aaaggaggtg 360
atccagccgc acctccagt acggctacct tgtaacgact tcaactccagt cgcaagccta 420
gccttaggca tccccctcct tacggttaag ggtaatgact tcaaacctgg ccagctccta 480
tagtgtgacg ggcggtgtgt acaaggcccg ggaacggatt cgccgcccga tggctgaccg 540
gcgattacta gcgattcctg cttcatgcag gcgagttgca gcctgcaatc cgaactgagg 600
acgggttttt ggagttagct caccctcgcg agatcgcgac cttttgtccc gccattgta 660
gcacgtgtgt cgcccagggc ataaggggca tgatgacttg gcctcatcct ctcttctctc 720

```

-continued

---

cggcttaaca	ccggcggtct	gttcaggggt	ccaaactcat	agtggcaact	aaacacgagg	780
gttgcgctcg	ttgcgagact	taaccaca	ccttacggca	cgagctgacg	acagccatgc	840
accacctgtg	tccgcgttcc	cgagggcacc	cctctctttc	aagaggattc	gcggcatgtc	900
aagccctggg	aaggttcttc	gccttgcatc	gaattaaacc	acatgctcca	ccgcttgtgc	960
gggccccgt	caattccttt	gagtttcatt	cttgcaacg	tactccccag	gcgggatact	1020
taacgcgtta	gctacagcac	tgacgggtc	gagtcgcaca	gcacctagta	tccatcgttt	1080
acggctagga	ctactggggg	ctctaatccc	atttgetccc	ctagctttcg	tctctcagtg	1140
tcagtgtcgg	cccagcagag	tgctttcgcc	gttgggttcc	ttccgatct	caatgcattt	1200
caccgctcca	ccggaaatc	cctctgcccc	taccgtactc	cagcttggtg	gtttccaccg	1260
cctgtccagg	gttgagccct	gggatttgac	ggcggacttg	aaaagccacc	tacagacgct	1320
ttacgcccc	tcattccgga	taacgcttgc	atcctctgtc	ttaccgggc	tgctggcaca	1380
gagttagccg	atgcttattc	ctcagatacc	gtcattgttt	cttctccgag	aaaagaagtt	1440
gacgaccggt	aggccttcca	cctccacgcg	gcattgctcc	gtcaggcttt	cgcccattgc	1500
ggaaaattcc	ccactgctgc	ctcccgtagg	agtctgggcc	gtgtctcagt	cccagtgtgg	1560
ctgatcatcc	tctcggacca	gctactgatc	atcgccctgg	taagctattg	cctcaccaac	1620
tagctaata	gacgcgagcc	cctccttggg	cggattttct	cttttgetcc	tcagcctacg	1680
gggtattagc	aaccgtttcc	agttgttgtt	cccctcccaa	gggcaggttc	ttacgcgtta	1740
ctcaccggtt	cgccactgga	aacaccactt	cccgttcgac	ttgcatgtgt	taagcatgcc	1800
gccagcgttc	atcctgagcc	aggatcgaac	tctccatgag	attcatagtt	gcattactta	1860
tagcttcctt	attegtagac	aaagcggatt	cggaaattgtc	tttcttcca	aggataactt	1920
gtatccatgc	gcttcagatt	attagcctgg	agttcgccac	cagcagtata	gccaacctta	1980
ccctatcacg	tcaatcccac	aagcctccta	tccattcccg	ttcgatcgtg	gtgggggagt	2040
aagtcaaaat	agaaaaaact	cacattaggt	ttagggataa	tcaggctcga	actgatgact	2100
tccaccacgt	caaggtgaca	ctctaccgct	gagttatata	ccttccccgt	cccacgaga	2160
aagagaatta	acgaatccta	aggcaaaggg	gcgagaaacg	caaggccact	cttctccggg	2220
gctttctttc	cgcaactatta	tggacagtca	aataatggga	aaaattggat	tcaattgtca	2280
acccggcggc	cgctcctatc	gaaaatagga	ttgactatgg	attcgagcca	tagcacatgg	2340
tttcataaaa	tctgtacgat	tttcccgatc	taaactgagt	gggtttccat	gaagaagatc	2400
ttgttcagca	tgttctatc	gatactggta	ggagaagaac	ccgactcggg	attcttaaaa	2460
aaagagggga	agcagaacca	agtcaagatg	atatggatcg	ccccttcttc	ttgcgcaaaa	2520
gatcttacca	tttccgaag	aactggggct	acatttctct	tccatttcca	ttccagagtt	2580
tctatctggt	tccacgcctt	ttttttgaga	cctcgaaaca	tgaaatggac	aaattccttc	2640
tcttaggaac	acatacaaga	aaaaggataa	tggtagcccc	cccattaact	aatttatgaa	2700
tttcatagta	atagaaatcc	atgtcctacc	gagacagaat	ttcgaacttg	ctatcctctt	2760
gcctaatagg	caaagattga	cctctgtaga	aagactgatt	cattcggatc	gatatgagga	2820
cccaactccg	ttgcattgcc	agaatccatg	ttccatattt	gaagcgggtt	gacctctgtg	2880
cttctctcat	ggtacaatcc	tcttctgct	gagccccctt	tctcctcggg	ccacagagaa	2940
aaaatgtagg	actggtgccc	acagttcatc	acggaagaaa	gaactcacag	agccggggat	3000
cgctaactaa	tagaatagta	ctactaacta	ataactaat	atagataact	aataactaata	3060
tatagatatt	aataactaata	tatagatata	tagaaataga	tatctagaaa	tagaaacgaa	3120

-continued

---

```

ctaataatata gataatcgaa atcgaaaaga actgtctttt ctgtataactt tccccgttct 3180
atgctaccgc gggctcttatg caatcgatcg gatcatatag atatcccttc aacacaacat 3240
aggatcatcga aaggatctcg gacgactcac caaagcacga aagccagtta gaaaatggat 3300
tcctatattga agagtgccta accgcatgga taagctcaca ttaaccctgc aatattggat 3360
ccaattcggg atttttcttg ggaagtttcg ggaagaaatt ggaataatat cgattcatac 3420
agaggaaaaa gttctctatt gatgcaaacg ctgtacctca gaggataggg atagaggaag 3480
agggaaaaat cgaaatgaaa taaataaaga ataaagccaa aaaaataagt cgaagataga 3540
agagcccaga ttcaaatga agaatggaa actcgaaaag gatccttctg attctcaaag 3600
aatgaggggc aaggggattg ataccgagaa agatttcttc ttattataag acgtgatttg 3660
atccgcatat gtttggtaaa agaacaatct tctccttaa tcataaatgg aaagtgttca 3720
attagaacat gaaaacgtga ctcaattggc cttagttagt cttcgggacg gagtgggaaga 3780
aggcgggaga ctctcgaacg aggaaaagga tcccttcgaa agaattgaac gaggagccgt 3840
atgaggtgaa aatctcatgt acggttctgt agagggacag taaggatgac ttatctgtcg 3900
acttttccac tatcaacccc aaaaaacca actctgcctt acgtaaagt gccagagtac 3960
gattaacctc tggatttgaa atcactgctt atatacctgg tattggccat aatttacaag 4020
aacattctgt agtattagta agaggaggaa gggtaagga tttaccctgt gtgagatatac 4080
gcattattcg aggaacccta gatgctgtcg cagtaaagaa tcgtcaacaa gggcgttcta 4140
gtgcgttgta gattcttata caagacttgt atcatttgat gatgccatgt gaatcgctag 4200
aaacatgtga agtgtatggc taaccacaata acgaaagttt cgtaagggga ctggagcagg 4260
ctaccatgag acaaaagatc ttctttctaa agagattcga ttcggaactc ttatatgtcc 4320
aaggtcaata tggaaattct ttcagagggt ttccttact ttgtccgtgt caacaaacaa 4380
ttcgaataac ctcgactttt tcagaacagg tccgagtcaa atagcaatga ttcgaagcac 4440
ttctttttcc attacactat ttcggaaacc taaggactcg atcgtatgga tatggaaaat 4500
acaggatttc cggctcctagc gggaaaagga gggaaaacgga tactcaattt aaagtgagta 4560
aacagaattc catactgat ctcatagatc cctatagaat tctgtgaaa gccgtattcg 4620
atgaaagtcg tatgtacggc ttggagggag atctttccta tctttcgaga tccaccctac 4680
aatatggggc caaaaagcca aaaaaataag tgattcgttt ttagccctta taaaaagaaa 4740
acggattctt gaacctctt cacgctcatg tcacgtcgag gtactgcaga aaaaagaacc 4800
gcaaaatccg atccaatttt tcgtaatcga ttagttaaca tgggtggccc gg 4852

```

```

<210> SEQ ID NO 9
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

```

```

<400> SEQUENCE: 9

```

```

gggtaccgtc gacttatcag gggcgcgctc taccactgag ctaatagccc 50

```

```

<210> SEQ ID NO 10
<211> LENGTH: 54
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

```

-continued

---

<400> SEQUENCE: 10

taagcttgcg gccgcccgtg tgacaattga atccaatttt tcccattatt tgac 54

<210> SEQ ID NO 11  
 <211> LENGTH: 52  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 11

aactagtccc gggccaccat gttaactaat cgattacgaa aaattggatc gg 52

<210> SEQ ID NO 12  
 <211> LENGTH: 51  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 12

ggaattcgcg gccgctccta tcgaaaatag gattgactat ggattcgagc c 51

<210> SEQ ID NO 13  
 <211> LENGTH: 156  
 <212> TYPE: DNA  
 <213> ORGANISM: Zea mays

<400> SEQUENCE: 13

ggaattcact cccccaccac gatcgaacgg gaatggatag gaggcttggtg ggattgacgt 60  
 gatagggtag ggttggtat actgctgggtg gcgaactcca ggctaataat ctgaagcgt 120  
 tggatacaag ttatccttg aaggaaagac aattcc 156

<210> SEQ ID NO 14  
 <211> LENGTH: 211  
 <212> TYPE: DNA  
 <213> ORGANISM: Zea mays  
 <220> FEATURE:  
 <221> NAME/KEY: promoter  
 <222> LOCATION: (1)..(160)  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (161)..(208)  
 <223> OTHER INFORMATION: G10L sequence

<400> SEQUENCE: 14

gcggccaatt cactccccca ccacgatcga acgggaatgg ataggaggct tgtgggattg 60  
 acgtgatagg gtagggttg ctatactgct ggtggcgaac tccaggctaa taatctgaag 120  
 cgcattggata caagttatcc ttggaaggaa agacaattcc ggatcctgta gaaataattt 180  
 tgtttaactt taagaaggag atatacccat g 211

<210> SEQ ID NO 15  
 <211> LENGTH: 162  
 <212> TYPE: DNA  
 <213> ORGANISM: Nicotiana tabacum

<400> SEQUENCE: 15

gggtgcagaa attcaattaa ggaaataaat taaggaaata caaaaagggg ggtagtcatt 60  
 tgtatataac tttgatgac ttttctcttc ttttttttg ttttctctcc ctttctttt 120  
 ctatttgat tttttatca ttgcttccat tgaattaagc tt 162

-continued

---

```

<210> SEQ ID NO 16
<211> LENGTH: 236
<212> TYPE: DNA
<213> ORGANISM: Nicotiana tabacum
<220> FEATURE:
<221> NAME/KEY: promoter
<222> LOCATION: (1)..(95)
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (96)..(185)
<223> OTHER INFORMATION: plastid atpB gene leader
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (186)..(236)
<223> OTHER INFORMATION: plastid atpB gene (first 15 aa)

<400> SEQUENCE: 16

aagcttgatc tcgctcccc gccgctcgttc aatgagaatg gataagaggc tcgtgggatt      60
gacgtgaggg ggcagggatg gctatatttc tgggagaatt aaccgatcga cgtgcaagcg      120
gacatttatt ttaaattcga taatttttgc aaaaacattt cgacatattt atttatttta      180
ttattatgag aatcaatcct actacttctg gttctggggg ttccaeggct agcatg         236

<210> SEQ ID NO 17
<211> LENGTH: 158
<212> TYPE: DNA
<213> ORGANISM: Zea mays

<400> SEQUENCE: 17

tctagattta gggatttttc gagttgattc attccaccgc gaagtcccct gcatgggtat      60
ctaggaaata gttacttcca agtgaatctt cctagatac ctaaaatcta ttttattatg      120
atccatttgc cgaaaatata gattaagcta gcggccgc                               158

<210> SEQ ID NO 18
<211> LENGTH: 223
<212> TYPE: DNA
<213> ORGANISM: Zea mays

<400> SEQUENCE: 18

ggctctagaa ataaaaaaaa aagcaaaata gaaagagaaa aaataagtta agaaatgcag      60
taattcttct ttattctctt aattgattgc aattcaattc ggctcaatct tttctaaaaa      120
aaaaaaaaagac tgagccgaat tgaaatagat cttgatacga tcatgagact tgacaaatcg      180
agattcttct attctatata tcaagaatat agaaaggtaa gct                          223

<210> SEQ ID NO 19
<211> LENGTH: 249
<212> TYPE: DNA
<213> ORGANISM: Zea mays
<220> FEATURE:
<221> NAME/KEY: promoter
<222> LOCATION: (1)..(156)
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (157)..(204)
<223> OTHER INFORMATION: G10L
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (205)..(249)
<223> OTHER INFORMATION: 14 aa of GFP

<400> SEQUENCE: 19

ggaattcact cccccaccac gatcgaacgg gaatggatag gaggcttgtg ggattgacgt      60
gatagggtag ggttggtat actgctggtg gcgaactcca ggctaataat ctgaagcgca      120

```



-continued

---

tggatacaag ttatccttgg aaggaaagac aattccggat cctgtagaaa taatdddttgtt 180  
 taactttaaag aaggagatat acccatgggt aaaggagaag aactdddttcac tggagttgtc 240  
 ccaagcatg 249

<210> SEQ ID NO 20  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 20

tgaaatagat cttgatacga tc 22

<210> SEQ ID NO 21  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 21

gaattgggac aactccagtg 20

<210> SEQ ID NO 22  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 22

tgaaaggcga gatcaccaag 20

<210> SEQ ID NO 23  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 23

aagagaaaga gactccccag 20

<210> SEQ ID NO 24  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 24

gcggaatcct ctactgtac 20

<210> SEQ ID NO 25  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 25

gtacacgcaa gaaataggcc 20

-continued

---

<210> SEQ ID NO 26  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 26

gttgtaggga gggacttatg 20

<210> SEQ ID NO 27  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 27

attgacttac tacaccccg 20

<210> SEQ ID NO 28  
 <211> LENGTH: 26  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 28

gttcactggc gtggtcccaa tcctgg 26

<210> SEQ ID NO 29  
 <211> LENGTH: 25  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 29

tcatccatgc catgcgtgat cccag 25

<210> SEQ ID NO 30  
 <211> LENGTH: 26  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 30

cctgttcctt ggccaacact tgtcac 26

<210> SEQ ID NO 31  
 <211> LENGTH: 26  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 31

ccatgtgtaa tcccagcagc agttac 26

<210> SEQ ID NO 32  
 <211> LENGTH: 23  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: synthetic construct

-continued

&lt;400&gt; SEQUENCE: 32

ggcagaagcg gtgatcgccg aag

23

&lt;210&gt; SEQ ID NO 33

&lt;211&gt; LENGTH: 28

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: synthetic construct

&lt;400&gt; SEQUENCE: 33

gcctcaaata gatcctgttc aagaaccg

28

&lt;210&gt; SEQ ID NO 34

&lt;211&gt; LENGTH: 27

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: synthetic construct

&lt;400&gt; SEQUENCE: 34

ctgggattac acatggcatg gatgaac

27

&lt;210&gt; SEQ ID NO 35

&lt;211&gt; LENGTH: 28

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: synthetic construct

&lt;400&gt; SEQUENCE: 35

cgatacttcg gcgatcaccg cttctgcc

28

The invention claimed is:

**1.** A method for transforming a plastid of a maize cell, said method comprising:

- (a) culturing an immature embryo of a maize plant in the presence of light to obtain a seedling;
- (b) culturing a nodal section from said seedling in a medium comprising cytokinin in the presence of light to obtain a green multiple bud culture;
- (c) introducing an exogenous DNA molecule of interest comprising an expression cassette into at least one cell of said green multiple bud culture, wherein said expression cassette is expressible in a plastid and becomes integrated into a plastid genome of said at least one cell; and
- (d) selecting for at least one plastid transformed maize cell from said green multiple bud culture comprising said expression cassette.

**2.** The method of claim 1, wherein said nodal section comprises meristematic tissue.

**3.** The method of claim 1, wherein said medium further comprises an auxin.

**4.** The method of claim 1, wherein said cytokinin is benzyl adenine (BA) or picloram.

**5.** The method of claim 1, wherein said exogenous DNA molecule is introduced into at least one cell of said green multiple bud culture through particle bombardment.

**6.** The method of claim 1, wherein said expression cassette comprises a selectable marker gene conferring tolerance to a selective agent, and wherein said selecting step comprises selection with said selective agent.

**7.** The method of claim 6, wherein said selectable marker gene is an aadA gene and said selective agent is spectinomycin or streptomycin.

**8.** The method of claim 6, wherein said selectable marker gene is a CP4 gene and said selective agent is glyphosate.

**9.** The method of claim 6, wherein said selectable marker gene is under the control of a plastid functional promoter.

**10.** The method of claim 6, wherein said expression cassette further comprises a screenable marker gene.

**11.** The method of claim 10, wherein said screenable marker gene is a GFP gene.

**12.** The method of claim 1, wherein said maize plant is selected from the group consisting of H99, (PA91×H99)×A188, Honey and Pearl, HiII and LH198×HiII.

**13.** The method of claim 1, wherein said expression cassette is integrated into the rbcL-psaI intergenic region of the Large Single Copy region of the maize plastid genome or the rrn16/trnV-rps12/3'-rps7 intergenic region of the Inverted Repeat region of the maize plastid genome.

**14.** A method of producing a transplastomic maize plant, said method comprising:

- (a) transforming a plastid of at least one maize plant cell according to the method of claim 1; and
- (b) regenerating a maize plant from the at least one plastid transformed maize plant cell to produce said transplastomic maize plant.

\* \* \* \* \*