



US009283230B2

(12) **United States Patent**
Clunas et al.

(10) **Patent No.:** **US 9,283,230 B2**
(45) **Date of Patent:** **Mar. 15, 2016**

(54) **PHENOTHIAZINE DIAMINIUM SALTS AND THEIR USE**

(30) **Foreign Application Priority Data**

Feb. 11, 2011 (SG) 201101060-0

(75) Inventors: **Scott Clunas**, Old Aberdeen (GB); **John Mervyn David Storey**, Old Aberdeen (GB); **James Peter Sinclair**, Old Aberdeen (GB); **Thomas Craven Baddeley**, Old Aberdeen (GB); **Ahtsham Ishaq**, Old Aberdeen (GB); **Michael Simpson**, Old Aberdeen (GB); **Craig Williamson**, Surrey (GB); **Barry Alan Wood**, Old Aberdeen (GB); **Claude Michel Wischik**, Old Aberdeen (GB); **Charles Robert Harrington**, Old Aberdeen (GB); **Janet Elizabeth Rickard**, Old Aberdeen (GB); **David Horsley**, Old Aberdeen (GB); **Yin Sze Loh**, Singapore (SG); **Colin Marshall**, Old Aberdeen (GB); **Karrar Ahmad Khan**, Old Aberdeen (GB)

(51) **Int. Cl.**
A61K 31/5415 (2006.01)
C07D 279/20 (2006.01)
C07D 309/04 (2006.01)
C07C 309/05 (2006.01)
A61K 9/20 (2006.01)
C07C 309/04 (2006.01)
A61K 9/28 (2006.01)
C07C 309/30 (2006.01)
C07C 309/35 (2006.01)

(73) Assignee: **WisTa Laboratories Ltd.**, Singapore (SG)

(52) **U.S. Cl.**
CPC **A61K 31/5415** (2013.01); **A61K 9/2095** (2013.01); **C07C 309/04** (2013.01); **C07C309/05** (2013.01); **C07D 279/20** (2013.01); **A61K 9/2893** (2013.01); **C07C 309/30** (2013.01); **C07C 309/35** (2013.01)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(58) **Field of Classification Search**
None
See application file for complete search history.

(21) Appl. No.: **13/984,841**

(56) **References Cited**

(22) PCT Filed: **Aug. 15, 2011**

FOREIGN PATENT DOCUMENTS

(86) PCT No.: **PCT/GB2011/001221**

WO WO 2007/110627 A2 10/2007
WO WO 2008/155533 A2 12/2008
WO WO 2009/044127 * 4/2009 A61K 31/542

§ 371 (c)(1),
(2), (4) Date: **Aug. 9, 2013**

OTHER PUBLICATIONS

(87) PCT Pub. No.: **WO2012/107706**

International Search Report and Written Opinion issued in related International Patent Application No. PCT/GB2011/001221, mailed Nov. 3, 2011.
Chemical Abstracts Service, XP002661598, Database Accession No. 1236208-20-0, 1 page (2010).

PCT Pub. Date: **Aug. 16, 2012**

(65) **Prior Publication Data**

US 2013/0315992 A1 Nov. 28, 2013

* cited by examiner


Related U.S. Application Data

(60) Provisional application No. 61/485,880, filed on May 13, 2011.

Primary Examiner — Bethany Barham

Assistant Examiner — Barbara Frazier

(74) *Attorney, Agent, or Firm* — Foley & Lardner LLP



(57)

ABSTRACT

The invention relates to compounds of general formula (I): wherein: each of R^1 and R^9 is independently selected from: —H, C_{1-4} alkyl, C_{2-4} alkenyl, and halogenated C_{1-4} alkyl; each of R^{3NA} and R^{3NB} is independently selected from: —H, C_{1-4} alkyl, C_{2-4} alkenyl, and halogenated C_{1-4} alkyl; each of R^{7NA} and R^{7NB} is independently selected from: —H, C_{1-4} alkyl, C_{2-4} alkenyl, and halogenated C_{1-4} alkyl; and wherein: each of R^A and R^B is independently selected from: C_{1-4} alkyl, halogenated C_{1-4} alkyl, and C_{6-10} aryl; or R^A and R^B are linked to form a group selected from: C_{1-6} alkylene and C_{6-10} arylene; and pharmaceutically acceptable salts thereof, which are useful in the treatment of, for example, Alzheimer's disease. In other aspects the invention also relates to novel formulations of 3,7-diamino-10H-phenothiazinium salts.


**5 Claims, 22 Drawing Sheets**

Figure 1:




Figure 2:




Figure 3:




Figure 4:




Figure 5:





Figure 6:



Mon May 03 00:42:11 2010

FIND PEAKS:

Spectrum: *6421JPS85/a

Region: 4000.00 400.00

Absolute threshold: 84.166

Sensitivity: 50

Peak list:

Position:	Intensity:
526.09	59.505
536.32	54.578
562.62	51.500
593.54	67.112
648.19	73.812
677.40	77.342
690.57	78.882

Handwritten:
JPS in air
20th Jun 2010
[NICOLET IR100]

Figure 7:




Figure 8:




Figure 9:




Figure 10:




Figure 11:




Figure 12:




Figure 13:




Figure 14:




Figure 15a:




Figure 15b – expanded view:




Figure 15c:




Figure 16:






Figure 17A:




X-ray structure (unit cell) of LMTEs

Figure 17B:



X-ray structure (unit cell) of LMT.EDSA

Figure 17C:



X-ray structure (unit cell) of LMTM

Figure 18:





Figure 19:



PHENOTHIAZINE DIAMINIUM SALTS AND THEIR USE

TECHNICAL FIELD

This invention pertains generally to the field of phenothiazine compounds, in particular certain phenothiazine diaminium salts, including uses and formulations thereof. In some embodiments the invention relates to bis(sulfonic acid) salts of diaminophenothiazine compounds such as N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine. The compounds of the invention are useful, for example, in the treatment of tauopathies such as Alzheimer's disease (AD).

BACKGROUND

A number of patents and publications are cited herein in order to more fully describe and disclose the invention and the state of the art to which the invention pertains. Each of these references is incorporated herein by reference in its entirety into the present disclosure, to the same extent as if each individual reference was specifically and individually indicated to be incorporated by reference.

Throughout this specification, including the claims which follow, unless the context requires otherwise, the word "comprise," and variations such as "comprises" and "comprising," will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

It must be noted that, as used in the specification and the appended claims, the singular forms "a," "an," and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a pharmaceutical carrier" includes mixtures of two or more such carriers, and the like.

Ranges are often expressed herein as from "about" one particular value, and/or to "about" another particular value. When such a range is expressed, another embodiment includes from the one particular value and/or to the other particular value. Similarly, when values are expressed as approximations, by the use of the antecedent "about," it will be understood that the particular value forms another embodiment.

Any sub-titles herein are included for convenience only, and are not to be construed as limiting the disclosure in any way.

Conditions of dementia are frequently characterised by a progressive accumulation of intracellular and/or extracellular deposits of proteinaceous structures such as β -amyloid plaques and neurofibrillary tangles (NFTs) in the brains of affected patients. The appearance of these lesions largely correlates with pathological neurofibrillary degeneration and brain atrophy, as well as with cognitive impairment (see, e.g., Mukaetova-Ladinska, E. B. et al., 2000, *Am. J. Pathol.*, Vol. 157, No. 2, pp. 623-636).

In Alzheimer's disease, both neuritic plaques and NFTs contain paired helical filaments (PHFs), of which a major constituent is the microtubule-associated protein tau (see, e.g., Wischik et al., 1988, *PNAS USA*, Vol. 85, pp. 4506-4510). Plaques also contain extracellular β -amyloid fibrils derived from the abnormal processing of amyloid precursor protein (APP) (see, e.g., Kang et al., 1987, *Nature*, Vol. 325, p. 733). An article by Wischik et al. (in 'Neurobiology of Alzheimer's Disease', 2nd Edition, 2000, Eds. Dawbarn, D. and Allen, S. J., The Molecular and Cellular Neurobiology Series, Bios Scientific Publishers, Oxford) discusses in detail the putative role of tau protein in the pathogenesis of neuro-

degenerative dementias. Loss of the normal form of tau, accumulation of pathological PHFs, and loss of synapses in the mid-frontal cortex all correlate with associated cognitive impairment. Furthermore, loss of synapses and loss of pyramidal cells both correlate with morphometric measures of tau-reactive neurofibrillary pathology, which parallels, at a molecular level, an almost total redistribution of the tau protein pool from a soluble to a polymerised form (i.e., PHFs) in Alzheimer's disease.

Tau exists in alternatively-spliced isoforms, which contain three or four copies of a repeat sequence corresponding to the microtubule-binding domain (see, e.g., Goedert, M., et al., 1989, *EMBO J.*, Vol. 8, pp. 393-399; Goedert, M., et al., 1989, *Neuron*, Vol. 3, pp. 519-526). Tau in PHFs is proteolytically processed to a core domain (see, e.g., Wischik, C. M., et al., 1988, *PNAS USA*, Vol. 85, pp. 4884-4888; Wischik et al., 1988, *PNAS USA*, Vol. 85, pp. 4506-4510; Novak, M., et al., 1993, *EMBO J.*, Vol. 12, pp. 365-370) which is composed of a phase-shifted version of the repeat domain; only three repeats are involved in the stable tau-tau interaction (see, e.g., Jakes, R., et al., 1991, *EMBO J.*, Vol. 10, pp. 2725-2729). Once formed, PHF-like tau aggregates act as seeds for the further capture and provide a template for proteolytic processing of full-length tau protein (see, e.g., Wischik et al., 1996, *PNAS USA*, Vol. 93, pp. 11213-11218).

The phase shift which is observed in the repeat domain of tau incorporated into PHFs suggests that the repeat domain undergoes an induced conformational change during incorporation into the filament. During the onset of AD, it is envisaged that this conformational change could be initiated by the binding of tau to a pathological substrate, such as damaged or mutated membrane proteins (see, e.g., Wischik, C. M., et al., 1997, in "Microtubule-associated proteins: modifications in disease", Eds. Avila, J., Brandt, R. and Kosik, K. S. (Harwood Academic Publishers, Amsterdam) pp. 185-241).

In the course of their formation and accumulation, PHFs first assemble to form amorphous aggregates within the cytoplasm, probably from early tau oligomers which become truncated prior to, or in the course of, PHF assembly (see, e.g., Mena, R., et al., 1995, *Acta Neuropathol.*, Vol. 89, pp. 50-56; Mena, R., et al., 1996, *Acta Neuropathol.*, Vol. 91, pp. 633-641). These filaments then go on to form classical intracellular NFTs. In this state, the PHFs consist of a core of truncated tau and a fuzzy outer coat containing full-length tau (see, e.g., Wischik et al., 1996, *PNAS USA*, Vol. 93, pp. 11213-11218).


The assembly process is exponential, consuming the cellular pool of normal functional tau and inducing new tau synthesis to make up the deficit (see, e.g., Lai, R. Y. K., et al., 1995, *Neurobiology of Ageing*, Vol. 16, No. 3, pp. 433-445). Eventually, functional impairment of the neurone progresses to the point of cell death, leaving behind an extracellular NFT. Cell death is highly correlated with the number of extracellular NFTs (see, e.g., Wischik et al., in 'Neurobiology of Alzheimer's Disease', 2nd Edition, 2000, Eds. Dawbarn, D. and Allen, S. J., The Molecular and Cellular Neurobiology Series, Bios Scientific Publishers, Oxford). As tangles are extruded into the extracellular space, there is progressive loss of the fuzzy outer coat of the neurone with corresponding loss of N-terminal tau immunoreactivity, but preservation of tau immunoreactivity associated with the PHF core (see, e.g., Bondareff, W. et al., 1994, *J. Neuropath. Exper. Neurol.*, Vol. 53, No. 2, pp. 158-164).

Diaminophenothiazine Compounds

Methythyonium Chloride (MTC) (also known as Methylene blue (MB); methylthionine chloride; tetramethylthionine chloride; 3,7-bis(dimethylamino) phenothiazin-5-ium chloride; C.I. Basic Blue 9; tetramethylthionine chloride;

3

3,7-bis(dimethylamino) phenazathionium chloride; Swiss blue; C.I. 52015; C.I. Solvent Blue 8; aniline violet; and Urolene Blue®) is a low molecular weight (319.86), water soluble, tricyclic organic compound of the following formula:



Methylthioninium Chloride (MTC) is a well known phenothiazine dye and redox indicator and has also been used as an optical probe of biophysical systems, as an intercalator in nanoporous materials, as a redox mediator, and in photoelectrochromic imaging. Methylthioninium chloride (MTC) and other diaminophenothiazines have been described as inhibitors of protein aggregation in diseases in which proteins aggregate pathologically.

In particular, diaminopenothiazines including MTC have been shown to inhibit tau protein aggregation and to disrupt the structure of PHFs, and reverse the proteolytic stability of the PHF core (see, e.g., WO 96/30766, Hofmann-La Roche). Such compounds were disclosed for use in the treatment or prophylaxis of various diseases, including Alzheimer's disease.

WO2007/110630 (WisTa Laboratories Ltd) also discloses certain specific diaminophenothiazine compounds related to MTC, including ETC, DEMTC, DMETC, DEETC, MTZ, ETZ, MTI, MTLHI, ETI, ETLHI, MTN, and ETN, which are useful as drugs, for example in the treatment of Alzheimer's disease.

Additionally, WO 2005/030676 (The University Court of the University of Aberdeen) discusses radiolabelled phenothiazines, and their use in diagnosis and therapy, for example, of tauopathies.

Methylthioninium chloride (MTC) has also been disclosed for other medical uses. For example it is currently used to treat methemoglobinemia (a condition that occurs when the blood cannot deliver oxygen where it is needed in the body). MTC is also used as a medical dye (for example, to stain certain parts of the body before or during surgery); a diagnostic (for example, as an indicator dye to detect certain compounds present in urine); a mild urinary antiseptic; a stimulant to mucous surfaces; a treatment and preventative for kidney stones; and in the diagnosis and treatment of melanoma.

MTC has been used to treat malaria, either singly (see, e.g., Guttman, P. and Ehrlich, P., 1891, "Uber die wirkung des methylenblau bei malaria," Berl. Klin. Wochenschr., Vol. 28, pp. 953-956) or in combination with chloroquine (see, e.g., Schirmer, H., et al., 2003, "Methylene blue as an antimalarial agent," Redox Report, Vol. 8, pp. 272-275; Rengelshausen, J., et al., 2004, "Pharmacokinetic interaction of chloroquine and methylene blue combination against malaria," European Journal of Clinical Pharmacology, Vol. 60, pp. 709-715).

MTC (under the name Virostat®, from Bioenvision Inc., New York) has also shown potent viricidal activity in vitro. Specifically Virostat® is effective against viruses such as HIV and West Nile Virus in laboratory tests. Virostat® is also currently in clinical trials for the treatment of chronic Hepa-


4

titis C, a viral infection of the liver. The virus, HCV, is a major cause of acute hepatitis and chronic liver disease, including cirrhosis and liver cancer. MTC, when combined with light, can also prevent the replication of nucleic acid (DNA or RNA). Plasma, platelets and red blood cells do not contain nuclear DNA or RNA. When MTC is introduced into the blood components, it crosses bacterial cell walls or viral membrane then moves into the interior of the nucleic acid structure. When activated with light, the compound then binds to the nucleic acid of the viral or bacterial pathogen, preventing replication of the DNA or RNA. Because MTC can inactivate pathogens, it has the potential to reduce the risk of transmission of pathogens that would remain undetected by testing.

Oral and parenteral formulations of MTC have been commercially available in the United States, usually under the name Urolene Blue®.

Reduced ('Leuco') Forms

MTC, a phenothiazin-5-ium salt, may be considered to be an "oxidized form" in relation to the corresponding 10H-phenothiazine compound, N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine, which may be considered to be a "reduced form":



The "reduced form" (or "leuco form") is known to be unstable and can be readily and rapidly oxidized to give the corresponding "oxidized" form.

May et al. (Am J Physiol Cell Physiol, 2004, Vol. 286, pp. C1390-C1398) have shown that human erythrocytes sequentially reduce and take up MTC; that MTC itself is not taken up by the cells; that it is the reduced form of MTC that crosses the cell membrane; that the rate of uptake is enzyme dependent; and that both MTC and reduced MTC are concentrated in cells (reduced MTC re-equilibrates once inside the cell to form MTC).

MTC and similar drugs are taken up in the gut and enter the bloodstream. Unabsorbed drug percolates down the alimentary canal, to the distal gut. One important undesired side-effect is the effect of the unabsorbed drug in the distal gut, for example, sensitisation of the distal gut and/or antimicrobial effects of the unabsorbed drug on flora in the distal gut, both


5

leading to diarrhoea. Therefore, it is desirable to minimize the amount of drug that percolates to the distal gut. By increasing the drug's uptake in the gut (i.e., by increasing the drug's bioavailability), dosage may be reduced, and the undesired side-effects, such as diarrhoea, may be ameliorated.

Since it is the reduced form of MTC that is taken up by cells, it may be desirable to administer the reduced form to patients. This may also reduce reliance on the rate limiting step of enzymatic reduction.

WO 02/055720 (The University Court of the University of Aberdeen) discloses the use of reduced forms of certain diaminophenothiazines for the treatment of protein aggregating diseases, primarily tauopathies.

WO02007/110627 (WisTa Laboratories Ltd) disclosed certain 3,7-diamino-10H-phenothiazinium salts, effective as drugs or pro-drugs for the treatment of diseases including Alzheimer's disease. These compounds are also in the "reduced" or "leuco" form when considered in respect of MTC. These included the following salts:



Although providing certain advantages over the use of MTC, the synthesis of LMT.2HCl under certain conditions may result in CH₃Cl being trapped within the crystal. This then needs to be removed since CH₃Cl is toxic and levels need to be kept below safety levels.

Furthermore LMT.2HBr contains bromide ions. This is in principle less desirable since bromide is toxic either at high levels or with chronic dosing and, at lower levels, can cause side effects such as confusion in patients.

Therefore it can be seen the provision of further salts of methylthionium compounds, having one or more desirable properties over those already known, would be a contribution to the art.

Furthermore the provision of novel formulations of methylthionium compounds which enhance stability, absorption, and/or otherwise improve their effectiveness as therapeutics would be a contribution to the art.

6

SUMMARY OF THE INVENTION


The present inventors have now identified a new class of stable phenothiazine diaminium compounds which have improved properties as compared to previously disclosed diaminophenothiazine compounds and salts.

The properties of the compounds are described hereinafter, whereby it can be seen that in preferred embodiments the invention can provide one or more of improved physical, pharmacokinetic, biochemical or other beneficial properties.

In other aspects the present inventors also provide novel formulations of 3,7-diamino-10H-phenothiazinium salts.

In one aspect the present invention provides certain compounds, specifically, certain phenothiazine diaminium compounds, as described herein.

The compound may be selected from compounds of general formula (I):



wherein:

each of R¹ and R⁹ is independently selected from: —H, C₁₋₄alkyl, C₂₋₄alkenyl, and halogenated C₁₋₄alkyl;

each of R^{3NA} and R^{3NB} is independently selected from: —H, C₁₋₄alkyl, C₂₋₄alkenyl, and halogenated C₁₋₄alkyl;

each of R^{7NA} and R^{7NB} is independently selected from:
—H, C_{1-4} alkyl, C_{2-4} alkenyl, and halogenated C_{1-4} alkyl;
and wherein:

each of R^A and R^B is independently selected from:
 C_{1-4} alkyl, halogenated C_{1-4} alkyl, and C_{6-10} aryl;
or
 R^A and R^B are linked to form a group R^{AB} , wherein R^{AB} is
selected from:
 C_{1-6} alkylene and C_{6-10} arylene;
and pharmaceutically acceptable salts thereof.

Another aspect of the invention pertains to processes for synthesizing a compound as described above.

Another aspect of the invention pertains to a pharmaceutical composition comprising a compound as described herein and a pharmaceutically acceptable carrier or diluent.

Another aspect of the invention pertains to a method of preparing a pharmaceutical composition comprising admixing a compound as described herein and a pharmaceutically acceptable carrier or diluent.

Another aspect of the invention pertains to a pharmaceutical composition in solid dosage form, comprising a compound as described herein and further comprising at least one diluent suitable for dry compression, and optionally one or more other excipients.

Another aspect of the invention pertains to a process for the manufacture of a pharmaceutical composition by a dry compression method, said composition being a solid dosage form comprising a compound as described herein, at least one diluent suitable for dry compression, and optionally one or more other excipients.

Another aspect of the invention pertains to a free-flowing, cohesive powder, comprising a compound as described herein and at least one diluent suitable for dry compression, and optionally one or more other excipients, said powder being capable of being compressed into a solid dosage form.

Another aspect of the present invention pertains to a method of reversing and/or inhibiting the aggregation of a protein (e.g., a tau protein, a synuclein, etc.), for example, aggregation of a protein associated with a neurodegenerative disease and/or clinical dementia, comprising contacting the protein with an effective amount of a compound or composition as described herein. Such a method may be performed in vitro, or in vivo.

Another aspect of the present invention pertains to a method of treatment or prophylaxis of a disease condition in a subject comprising administering to said subject a prophylactically or therapeutically effective amount of a compound as described herein, preferably in the form of a pharmaceutical composition, preferably a pharmaceutical composition in solid dosage form, as further described herein.

Another aspect of the present invention pertains to a compound or composition as described herein for use in a method of treatment or prophylaxis (e.g., of a disease condition) of the human or animal body by therapy.

Another aspect of the present invention pertains to use of a compound or composition as described herein, in the manufacture of a medicament for use in the treatment or prophylaxis of a disease condition.

In some embodiments, the disease condition is a disease of protein aggregation.

In some embodiments, the disease condition is a tauopathy, e.g., a neurodegenerative tauopathy, e.g., Alzheimer's disease or other disease described hereinafter.

In some embodiments, the disease condition is skin cancer, e.g., melanoma.

In some embodiments, the disease condition is a viral, bacterial or protozoal disease condition, e.g., Hepatitis C, HIV, West Nile Virus (WNV), or malaria.

Another aspect of the present invention pertains to a method of inactivating a pathogen in a sample (for example a blood or plasma sample), comprising the steps of introducing a compound or composition as described herein, into the sample, and then exposing the sample to light.

Another aspect of the present invention pertains to a kit comprising (a) a compound as described herein, preferably provided as a pharmaceutical composition and in a suitable container and/or with suitable packaging; and (b) instructions for use, for example, written instructions on how to administer the compound or composition.

As will be appreciated by one of skill in the art, features and preferred embodiments of one aspect of the invention will also pertain to other aspects of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows the ^1H NMR spectrum of an exemplary compound of the invention (LMT.2MsOH) in deuterated methanol (CD_3OD) at 600 MHz.

FIG. 2 shows the ^{13}C NMR spectrum of LMT.2MsOH in CD_3OD at a frequency of 100.56 MHz.

FIG. 3 shows the DEPT-135 spectrum of LMT.2MsOH in CD_3OD at a frequency of 100.56 MHz.

FIG. 4 shows the HSQC spectrum of LMT.2MsOH in CD_3OD at a frequency of 100.56 MHz.

FIG. 5 shows an expanded section of the HSQC spectrum of LMT.2MsOH in CD_3OD at a frequency of 100.56 MHz.

FIG. 6 shows the infrared (FT-IR) spectrum of LMT.2MsOH (KBr).

FIG. 7 shows the electron impact (EI) mass spectrum of LMT.2MsOH.

FIG. 8 shows the electrospray ionisation (ESI) mass spectrum of LMT.2MsOH.

FIG. 9 shows the UV/Vis spectrum of LMT.2MsOH in de-ionised water.

FIG. 10 shows the HPLC trace for LMT.2MsOH.

FIG. 11 shows a powder X-ray diffractogram for LMT.2MsOH, measured with $\text{Cu K}\alpha$ radiation.

FIG. 12 shows the FT-Raman spectrum for crystalline LMT.2MsOH. The most intense signals are found at 1615 cm^{-1} , 1588 cm^{-1} , 1258 cm^{-1} , and 1042 cm^{-1} .

FIG. 13 shows the thermogravimetric profile for crystalline LMT.2MsOH. A constant weight was detected by TG and TG-FTIR up to the beginning of decomposition at $240\text{--}270^\circ\text{C}$.

FIG. 14 shows the differential scanning calorimetry analysis for crystalline LMT.2MsOH. A sharp m.p. at 271°C . ($\Delta H=87\text{ J/g}$) was immediately followed by decomposition.

FIGS. 15a and 15b show the dynamic vapour sorption (DVS) curve for crystalline LMT.2MsOH measured at 25°C . with $5\%/h$ scanning rate. The horizontal dashed lines indicate steps of water uptake of one equivalent. A stable weight of the sample (less than 0.5% weight change) was observed in the relative humidity (r.h.) range between 0% and 70% . Above this r.h., the water uptake increased rapidly, and the sample ultimately deliquesced. Upon drying, the water content decreased again to approximately 4 equiv. at 50% r.h. The DVS curve of the crystalline dihydrochloride salt (LMT.2HCl) is shown for comparison as a dashed line, the DVS curve of the dihydrobromide salt (LMT.2HBr) as a dotted line.

9

FIG. 15c shows the dynamic vapour sorption (DVS) curve for crystalline LMT.2MsOH as a function of time. The relative humidity is also indicated (right axis). The horizontal dashed lines indicate steps of one equivalent water uptake.

FIG. 16 shows polarizing microscopy pictures of the LMT.2MsOH (left) and recrystallized LMT.2MsOH (right). Crystals of up to 100 μm in size were obtained by recrystallization from 2-PrOH/water. Crystals are irregularly shaped.

FIGS. 17a-c shows the X-ray crystal structures of LMTEsOH, LMT.EDSA, and LMT.2MsOH

FIG. 18 shows a comparison of the plasma concentration in pig of the MT moiety over time following dosing of LMT.2HBr, LMT.2HCl and LMT.2MsOH.

FIG. 19 is a diagram of the apparatus used in the dissolution studies (see Formulation Example 12).

DETAILED DESCRIPTION OF THE INVENTION


The present inventors have identified a new class of phenothiazine diaminium compounds which have desirable physical or other properties and/or surprisingly improved activity as compared to previously disclosed diamino-phenothiazine compounds and salts.

In other aspects they have additionally provided novel formulations of phenothiazine diaminium compounds, including (but not limited to) the class above.

The Compounds

In general terms, unless context demands otherwise, the compounds of the invention may be described as bis(sulfonate) salts (or bis(sulfonic acid) salts) of 3,7-diamino-10H-phenothiazine compounds. In other words, the compounds are salts of the corresponding 3,7-diamino-10H-phenothiazine compounds with organic sulfonic acids.

More specifically, a compound of the invention is a bis(sulfonate) salt of a compound of general formula:



wherein R^1 , R^9 , R^{3NA} , R^{3NB} , R^{7NA} and R^{7NB} are as defined above.

In some embodiments, the salt is a bis(alkylsulfonate) salt or a bis(arylsulfonate) salt.

In some embodiments, the salt is selected from a bis(methanesulfonate) salt, a bis(ethanesulfonate) salt, a bis(p-toluenesulfonate) salt, a bis(benzenesulfonate) salt, an ethanedisulfonate salt, a propanedisulfonate salt, or a naphthalenedisulfonate salt.

In some embodiments, the salt is a bis(methanesulfonate) salt (which may also be called a bis(mesylate) salt).

In some embodiments, the salt is a bis(ethanesulfonate) salt (which may also be called a bis(esylate) salt).

In some embodiments, the salt is a bis(p-toluenesulfonate) salt (which may also be called a bis(tosylate) salt).

In some embodiments, the salt is a bis(benzenesulfonate) salt.

In some embodiments, the salt is an ethanedisulfonate salt.


In some embodiments, the salt is a propanedisulfonate salt.

In some embodiments, the salt is a naphthalenedisulfonate salt, preferably a naphthalene-1,5-disulfonate salt.

10

In other words, the compounds of the invention can be considered to be products obtainable from the reaction of a 3,7-diamino-10H-phenothiazine compound, for example as set out above, with two organic sulfonic acid moieties ($R^A\text{SO}_3\text{H}$ and $R^B\text{SO}_3\text{H}$). The two organic sulfonic acid moieties may optionally be present on the same molecule, i.e. where R^A and R^B are linked.

In some embodiments, compounds of the invention are selected from compounds of general formula (I):



wherein:

each of R^1 and R^9 is independently selected from: —H, C_{1-4} alkyl, C_{2-4} alkenyl, and halogenated C_{1-4} alkyl;

each of R^{3NA} and R^{3NB} is independently selected from: —H, C_{1-4} alkyl, C_{2-4} alkenyl, and halogenated C_{1-4} alkyl;

each of R^{7NA} and R^{7NB} is independently selected from: —H, C_{1-4} alkyl, C_{2-4} alkenyl, and halogenated C_{1-4} alkyl;

and wherein:

each of R^A and R^B is independently selected from:

C_{1-4} alkyl, halogenated C_{1-4} alkyl, and C_{6-10} aryl;

or


R^A and R^B are linked to form a group R^{AB} , wherein R^{AB} is selected from:

C_{1-6} alkylene and C_{6-10} arylene;

and pharmaceutically acceptable salts, solvates, and hydrates thereof.


Compounds of the invention are represented herein by a general formula showing the structure of the 3,7-diamino-10H-phenothiazine compound, with the 3,7-diamino groups being in protonated form.

The resultant doubly positively-charged species is associated with two sulfonate counterion moieties (which may optionally be present on the same molecule, i.e. where R^A and R^B are linked):



However, as will be understood by one skilled in the art, the same salt could equally be represented in other ways, such as, for example:

11



etc.

Further Definitions And Preferences

The term “C₁₋₄ alkyl”, as used herein, pertains to a monovalent moiety obtained by removing a hydrogen atom from a hydrocarbon compound having from 1 to 4 carbon atoms, which may be aliphatic or alicyclic, or a combination thereof.

Similarly, the term “C₂₋₄ alkenyl” pertains to a monovalent moiety obtained by removing a hydrogen atom from a C₂₋₄ alkene compound (i.e. a hydrocarbon compound containing at least one double bond and from 2 to 4 carbon atoms).

The term “C₁₋₆ alkylene”, as used herein, pertains to a bidentate moiety obtained by removing two hydrogen atoms, either both from the same carbon atom, or one from each of two different carbon atoms, of an aliphatic linear hydrocarbon compound having from 1 to 6 carbon atoms.

In some embodiments, C₁₋₄alkyl groups may be selected from: linear C₁₋₄alkyl groups, such as -Me, -Et, -nPr, -iPr, and -nBu; branched C₃₋₄alkyl groups, such as -iPr, -iBu, -sBu, and -tBu; and cyclic C₃₋₄alkyl groups, such as -cPr and -cBu.

In some embodiments, C₂₋₄alkenyl groups may be selected from linear C₁₋₄alkenyl groups, such as —CH=CH₂ (vinyl) and —CH₂—CH=CH₂ (allyl).

In some embodiments, halogenated C₁₋₄alkyl groups may be selected from: —CF₃, —CH₂CF₃, and —CF₂CF₃.

The term “C₆₋₁₀ aryl”, as used herein, pertains to a monovalent moiety obtained by removing a hydrogen atom from an aromatic ring atom of a C₆₋₁₀ aromatic compound, said compound having one ring, or two or more rings (e.g., fused), and having from 6 to 10 ring atoms, and wherein at least one of said ring(s) is an aromatic ring.

The term “C₆₋₁₀ arylene”, as used herein, pertains to a bidentate moiety obtained by removing two hydrogen atoms from an aromatic compound having from 6 to 10 carbon atoms.

In some embodiments, C₆₋₁₀ aryl groups may be selected from C₆₋₁₀ carboaryl groups such as phenyl, and naphthyl.

12

In some embodiments, C₆₋₁₀ arylene groups may be selected from phenylene and naphthylene.

Said C₁₋₄ alkyl and C₁₋₆ alkylene groups may be unsubstituted or may optionally be substituted, for example with one or more groups selected from halo (e.g. F, Cl, Br, or I), amino (e.g. —NH₂, —NHR, or —NR₂, wherein each R is independently C₁₋₄alkyl), hydroxy (—OH), alkoxy (—OR, wherein R is independently C₁₋₄alkyl), nitro (—NO₂), etc.

Said C₆₋₁₀ aryl and C₆₋₁₀ arylene groups may be unsubstituted or may optionally be substituted, for example with one or more groups selected from C₁₋₄ alkyl, for example—Me, halogenated C₁₋₄alkyl, for example—CF₃, halo (e.g. F, Cl, Br, or I), amino (e.g. —NH₂, —NHR, or —NR₂, wherein each R is independently C₁₋₄alkyl), hydroxy (—OH), alkoxy (—OR, wherein R is independently C₁₋₄alkyl), nitro (—NO₂), etc.

Groups R^A and R^B

Each of R^A and R^B is independently selected from:

C₁₋₄alkyl, halogenated C₁₋₄alkyl, and C₆₋₁₀ aryl;

or

R^A and R^B are linked to form a group R^{AB}, wherein R^{AB} is selected from:

C₁₋₆ alkylene and C₆₋₁₀ arylene;

In some embodiments, each of R^A and R^B is independently selected from:

C₁₋₄alkyl, halogenated C₁₋₄alkyl, and C₆₋₁₀ aryl.

In some embodiments, each of R^A and R^B is independently C₁₋₄ alkyl.

In some embodiments, each of R^A and R^B is independently selected from Me, Et, nPr, iPr, nBu, iBu, tBu.

In some embodiments, each of R^A and R^B is independently selected from Me and Et.

In some embodiments, each of R^A and R^B is independently C₆₋₁₀ aryl.


In some embodiments, each of R^A and R^B is independently selected from benzene, 1-naphthalene, 2-naphthalene and p-toluene.

In some embodiments, each of R^A and R^B is independently selected from Me, Et, benzene and p-toluene.

In some embodiments, R^A and R^B are the same.

In some embodiments, R^A and R^B are different.


In some embodiments, R^A and R^B are the same and are independently Me. The compound may then be referred to as a diaminophenothiazine bis(methanesulfonate) salt which is of general formula (Ia):



In some embodiments, R^A and R^B are linked to form a group R^{AB}.

In these embodiments, the compounds of the invention may alternately be represented by general formula Ib:

13



wherein R^{AB} is selected from C_{1-6} alkylene and C_{6-10} arylene.

In some embodiments, R^{AB} is a C_{1-6} alkylene group.

In some embodiments, R^{AB} is a C_{1-6} alkylene group selected from $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$ and $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$.

In some embodiments, R^{AB} is a C_{1-6} alkylene group selected from methylene ($-\text{CH}_2-$), ethylene ($-\text{CH}_2\text{CH}_2-$) and propylene ($-\text{CH}_2\text{CH}_2\text{CH}_2-$).

In some embodiments, R^{AB} is ethylene.

In some embodiments, R^{AB} is a C_{6-10} arylene group.

In some embodiments, R^{AB} is a C_{6-10} arylene group selected from phenylene and naphthylene.

In some embodiments, R^{AB} is phenylene.

In some embodiments, R^{AB} is selected from 1,2-phenylene, 1,3-phenylene, and 1,4-phenylene.


In some embodiments, R^{AB} is phenylene optionally substituted with one or more substituents, for example selected from C_{1-4} alkyl, halogenated C_{1-4} alkyl, and halo.

In some embodiments, R^{AB} is naphthylene.


In some embodiments, R^{AB} is selected from 1,2-naphthylene, 1,3-naphthylene, 1,4-naphthylene, 1,5-naphthylene, 1,6-naphthylene, 1,7-naphthylene and 1,8-naphthylene.

In some embodiments, R^{AB} is selected from:

1,5-naphthylene i.e.



and 1,8-naphthylene. i.e.




In some embodiments, R^{AB} is naphthylene optionally substituted with one or more substituents, for example selected from C_{1-4} alkyl, halogenated C_{1-4} alkyl, and halo.

Groups R^1 and R^9

In some embodiments, each of R^1 and R^9 is independently $-\text{H}$, $-\text{Me}$, $-\text{Et}$, or $-\text{CF}_3$.

14

(Ib)



In some embodiments, each of R^1 and R^9 is independently $-\text{H}$, $-\text{Me}$, or $-\text{Et}$.

In some embodiments, R^1 and R^9 are the same.

In some embodiments, R^1 and R^9 are different.

In some embodiments, each of R^1 and R^9 is independently $-\text{H}$.

In some embodiments, each of R^1 and R^9 is independently $-\text{Me}$.

In some embodiments, each of R^1 and R^9 is independently $-\text{Et}$.

Groups R^{3NA} and R^{3NB}

Each of R^{3NA} and R^{3NB} is independently selected from: $-\text{H}$, C_{1-4} alkyl, C_{2-4} alkenyl, and halogenated C_{1-4} alkyl.

In some embodiments, each of R^{3NA} and R^{3NB} is independently selected from: C_{1-4} alkyl, C_{2-4} alkenyl, and halogenated C_{1-4} alkyl.

In some embodiments, each of R^{3NA} and R^{3NB} is independently $-\text{Me}$, $-\text{Et}$, $-\text{nPr}$, $-\text{nBu}$, $-\text{CH}_2-\text{CH}=\text{CH}_2$, or $-\text{CF}_3$.

In some embodiments, each of R^{3NA} and R^{3NB} is independently $-\text{Me}$, $-\text{nPr}$, $-\text{nBu}$, $-\text{CH}_2-\text{CH}=\text{CH}_2$, or $-\text{CF}_3$.

In some embodiments, each of R^{3NA} and R^{3NB} is independently $-\text{Me}$ or $-\text{Et}$.

In some embodiments, R^{3NA} and R^{3NB} are the same.

In some embodiments, R^{3NA} and R^{3NB} are different.

In some embodiments, each of R^{3NA} and R^{3NB} is independently $-\text{Me}$.

Groups R^{7NA} and R^{7NB}

Each of R^{7NA} and R^{7NB} is independently selected from: $-\text{H}$, C_{1-4} alkyl, C_{2-4} alkenyl, and halogenated C_{1-4} alkyl.

In some embodiments, each of R^{7NA} and R^{7NB} is independently selected from: C_{1-4} alkyl, C_{2-4} alkenyl, and halogenated C_{1-4} alkyl.

In some embodiments, each of R^{7NA} and R^{7NB} is independently $-\text{Me}$, $-\text{Et}$, $-\text{nPr}$, $-\text{nBu}$, $-\text{CH}_2-\text{CH}=\text{CH}_2$, or $-\text{CF}_3$.

In some embodiments, each of R^{7NA} and R^{7NB} is independently $-\text{Me}$, $-\text{nPr}$, $-\text{nBu}$, $-\text{CH}_2-\text{CH}=\text{CH}_2$, or $-\text{CF}_3$.

In some embodiments, each of R^{7NA} and R^{7NB} is independently $-\text{Me}$ or $-\text{Et}$.

In some embodiments, R^{7NA} and R^{7NB} are the same.

In some embodiments, R^{7NA} and R^{7NB} are different.

In some embodiments, each of R^{7NA} and R^{7NB} is independently $-\text{Me}$.

Groups R^{3NA} , R^{3NB} , R^{7NA} and R^{7NB}

In some embodiments:

each of R^{3NA} and R^{3NB} is independently C_{1-4} alkyl, C_{2-4} alkenyl, or halogenated C_{1-4} alkyl;

each of R^{7NA} and R^{7NB} is independently C_{1-4} alkyl, C_{2-4} alkenyl, or halogenated C_{1-4} alkyl.

In some embodiments:

each of R^{3NA} and R^{3NB} is independently $-\text{Me}$, $-\text{Et}$, $-\text{nPr}$, $-\text{nBu}$, $-\text{CH}_2-\text{CH}=\text{CH}_2$, or $-\text{CF}_3$;

each of R^{7NA} and R^{7NB} is independently $-\text{Me}$, $-\text{Et}$, $-\text{nPr}$, $-\text{nBu}$, $-\text{CH}_2-\text{CH}=\text{CH}_2$, or $-\text{CF}_3$.

In some embodiments:

each of R^{3NA} and R^{3NB} is independently $-\text{Me}$ or $-\text{Et}$;

each of R^{7NA} and R^{7NB} is independently $-\text{Me}$ or $-\text{Et}$.

In some embodiments, R^{3NA} and R^{3NB} and R^{7NA} and R^{7NB} are all the same.

In some embodiments, R^{3NA} and R^{3NB} and R^{7NA} and R^{7NB} are the same and are all $-\text{Me}$ or all $-\text{Et}$.

In some embodiments, R^{3NA} and R^{3NB} and R^{7NA} and R^{7NB} are the same and are all $-\text{Me}$.

Salts and Solvates

Although the compounds described herein are themselves salts, they may also be provided in the form of a mixed salt (i.e., the compound of the invention in combination with another salt). Such mixed salts are intended to be encom-

15

passed by the term “and pharmaceutically acceptable salts thereof”. Unless otherwise specified, a reference to a particular compound also includes salts thereof.

The compounds of the invention may also be provided in the form of a solvate or hydrate. The term “solvate” is used herein in the conventional sense to refer to a complex of solute (e.g., compound, salt of compound) and solvent. If the solvent is water, the solvate may be conveniently referred to as a hydrate, for example, a mono-hydrate, a di-hydrate, a tri-hydrate, etc. Unless otherwise specified, any reference to a compound also includes solvate and hydrate forms thereof.

Naturally, solvates or hydrates of salts of the compounds are also encompassed by the present invention.

Isotopic Variation

In some embodiments, one or more carbon atoms of the compound is ^{11}C , ^{13}C or ^{14}C .

In some embodiments, one or more carbon atoms of the compound is ^{11}C .

In some embodiments, one or more carbon atoms of the compound is ^{13}C .

In some embodiments, one or more carbon atoms of the compound is ^{14}C .

In some embodiments, one or more nitrogen atoms of the compound is ^{15}N .

16

In some embodiments, one or more or all of the carbon atoms of one or more or all of the groups $\text{R}^{3\text{NA}}$, $\text{R}^{3\text{NB}}$, $\text{R}^{7\text{NA}}$, $\text{R}^{7\text{NB}}$, R^1 , R^9 , R^A and R^B is ^{11}C , ^{13}C , or ^{14}C .

In some embodiments, one or more or all of the carbon atoms of one or more or all of the groups $\text{R}^{3\text{NA}}$, $\text{R}^{3\text{NB}}$, $\text{R}^{7\text{NA}}$ and $\text{R}^{7\text{NB}}$ is ^{11}C , ^{13}C , or ^{14}C .

Combinations


All compatible combinations of the embodiments described above are explicitly disclosed herein as if each combination was specifically and individually recited.

In particular, in the compounds of the invention, the groups $\text{R}^{3\text{NA}}$, $\text{R}^{3\text{NB}}$, $\text{R}^{7\text{NA}}$, $\text{R}^{7\text{NB}}$, R^1 , R^9 , R^A and R^B (and R^{AB}) are defined as independent variables and it will be recognised by those skilled in the art that any compatible combination of these groups and substituents may be utilised in the compounds and methods of the present invention.


All compatible combinations of these and other defined variables are therefore specifically embraced by the present invention, and are disclosed herein as if each and every combination were individually and explicitly recited.

Some Preferred Embodiments


In some embodiments, the compound of the invention may be selected from the following compounds and pharmaceutically acceptable salts, solvates, and hydrates thereof:



-continued



One particular compound of the invention is compound 1: 40 compounds oxidize (e.g., autoxidize) to give the corresponding oxidized forms. Thus, it is likely, if not inevitable, that compositions comprising the compounds of the present invention will contain, as an impurity, at least some of the corresponding oxidized compound.



N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diaminium bis(methanesulfonate).

This compound may also be referred to as:

N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine bis (hydromethanesulfonate)

Leuco methylthioninium bis(hydromethanesulfonate)

Leuco methylthioninium bis(mesylate)

LMTM

LMT.2MsOH

Purity

The compounds of the present invention may conveniently be described as being in a "stabilized reduced form". The

45 Thus, another aspect of the present invention pertains to compounds as described herein, in substantially purified form and/or in a form substantially free from contaminants (e.g., the corresponding oxidized compound, other contaminants).

In some embodiments, the substantially purified form is at least 50% by weight pure, e.g., at least 60% by weight pure, e.g., at least 70% by weight pure, e.g., at least 80% by weight pure, e.g., at least 90% by weight pure, e.g., at least 95% by weight pure, e.g., at least 97% by weight pure, e.g., at least 98% by weight pure, e.g., at least 99% by weight pure.

55 In some embodiments, the contaminants represent no more than 50% by weight, e.g., no more than 40% by weight, e.g., no more than 30% by weight, e.g., no more than 20% by weight, e.g., no more than 10% by weight, e.g., no more than 5% by weight, e.g., no more than 3% by weight, e.g., no more than 2% by weight, e.g., no more than 1% by weight.

60 Product-by-Process

In some embodiments, the compound is one which is obtained by, or is obtainable by, a method as described herein.


CHEMICAL SYNTHESIS

65 Methods for the chemical synthesis of the compounds of the present invention are described herein. These and/or other


19

well known methods may be modified and/or adapted in known ways in order to facilitate the synthesis of additional compounds within the scope of the present invention.

Compounds of formula (I):




may be prepared from compounds of formula (II):



wherein R^1 , R^9 , R^{3NA} , R^{3NB} , R^{7NA} , and R^{7NB} are as defined previously.

Compounds of formula (II) may, for example, be prepared from compounds of formula (III):



wherein R^{Prot} is an amine protecting group and R^1 , R^9 , R^{3NA} , R^{3NB} , R^{7NA} , R^{7NB} , R^A and R^B are as defined previously.

By way of non-limiting example, R^{Prot} may be an acyl group, for example an acetyl ($-\text{C}(=\text{O})\text{Me}$) or a benzoyl ($-\text{C}(=\text{O})\text{Ph}$) group.

The compounds of formula (II) may be prepared e.g. by deprotection of the compounds of formula (III), or by other known methods. Conversely, compounds of formula (II) may be produced by protection of compounds of formula (III).

Compounds of formulae (II) and (III) are known, and may be prepared from known and/or commercially available starting materials, e.g. from corresponding phenothiazine compounds, using known methods.

For example, intermediates of formula (II) and (III) were used in the methods for the synthesis of 3,7-diamino-10H-phenothiazine hydrochloride, hydrobromide, and hydroiodide salts disclosed in WO2007/110627.

As disclosed in that document, a suitable phenothiazine can be converted to the corresponding 3,7-dinitro-phenothiazine, for example using sodium nitrite with acetic acid and chloroform.

The ring amino group may then be protected, for example as the acetate, for example using acetic anhydride and pyridine.


20

The nitro groups may then be reduced to amino groups, for example using tin (II) chloride with ethanol.


The amino groups may then be substituted, for example disubstituted, for example methyl disubstituted, for example using methyl iodide, sodium hydroxide, DMSO, and tetra-n-butyl ammonium bromide, to provide a N-acetyl protected 3,7-dialkylamino-10H-phenothiazine.

Examples of such a method are illustrated in Schemes 1a and 1b. The use of any one or more of the reagents described herein in the process is of course encompassed by the present invention:

Scheme 1a






Scheme 1b



21

-continued




The amino group of this N-acetyl intermediate can then be deprotected, i.e. the N-acetyl group may be removed, for example using aqueous acid.

Compounds of formulae (II) and (III) may also be made using the methods disclosed in WO2008/007074. This document discloses compounds of formula (III) and compounds of formula (II) wherein R^{Prot} is an acyl group, for example an acetyl group.


In one approach, an appropriate thioninium chloride (e.g., methyl thioninium chloride, ethyl thioninium chloride, etc) may first be reduced and acetylated to give the corresponding 1-(3,7-bis-dimethylamino-phenothiazin-10-yl)-ethanone, for example, by reaction with hydrazine (NH_2NH_2), methyl hydrazine ($MeNHNH_2$), or sodium borohydride ($NaBH_4$); and acetic anhydride ($(H_3CCO)_2O$); for example, in the presence of a suitable base, for example, pyridine (C_5H_5N) or Hünig's base (diisopropylethylamine, $C_8H_{19}N$), for example, in a suitable solvent, for example, ethanol or acetonitrile. The reduced and acetylated compound (of formula (III)) may then be deprotected (by removing the acetyl group), for example by reaction with a suitable acid, to give a compound of formula (II) or may be used directly. Advantageously, this reaction may produce a product with a high degree of purity.

An example is shown in the following scheme.




22

-continued




In another approach, an appropriate thioninium salt, for example, ethyl thioninium semi zinc chloride, may be simultaneously reduced and the ring amino group protected, for example, by reaction with a reducing agent phenylhydrazine, ethanol, acetic anhydride, and pyridine.


An example is shown in the following scheme:



In one aspect, the present invention therefore provides a method of preparing a 3,7-diamino-10H-phenothiazine compound of formula (I):



from a compound of formula (II):



wherein R^A , R^B , R^1 , R^9 , R^{3NA} , R^{3NB} , R^{7NA} , and R^{7NB} are as previously defined.


In some embodiments, the method comprises the step of: salt formation (SF).

23


In some embodiments, salt formation (SF) comprises treatment of a compound of formula (II) with an appropriate sulfonic acid.

In some embodiments, salt formation comprises treatment of a solution of a compound of formula (II) with an appropriate sulfonic acid, in an organic solvent.

In a further aspect, the present invention provides a method of preparing a 3,7-diamino-10H-phenothiazine compound of formula (I):



from a compound of formula (III):



wherein R^A , R^B , R^1 , R^9 , R^{3NA} , R^{3NB} , R^{7NA} , and R^{7NB} are as previously defined and wherein R^{Prot} is an amine protecting group.

A wide variety of amine protecting groups are widely used and well known in organic synthesis. See, for example, *Protective Groups in Organic Synthesis* (T. Green and P. Wuts; 4th Edition; John Wiley and Sons, 2006).

In some embodiments, the amine protecting group is an acid-cleavable protecting group.

In some embodiments, the amine protecting group is an acyl group, such as an acetyl group.


In some embodiments, the method comprises the steps of: ring amino deprotection (DP); and salt formation (SF).

Ring amino deprotection (DP) comprises removal of the protecting group to convert the N-protected ring amine group ($-\text{NR}^{Prot}-$) to a free ring amine group ($-\text{NH}-$). Deprotection of a compound of formula (III) produces the corresponding compound of formula (II).

Methods for the removal of amine protecting groups are known in the art. See, for example, *Protective Groups in Organic Synthesis* (T. Green and P. Wuts; 4th Edition; John Wiley and Sons, 2006).

In some embodiments, the step of ring amino deprotection (DP) and the step of salt formation (SF) are performed simultaneously (i.e., as one step). For example:

24



In some embodiments, simultaneous ring amino deprotection (DP) and salt formation (SF) comprises treatment of the compound of formula (III) with an appropriate sulfonic acid, to produce a bis(sulfonate) salt of formula (I).

In some embodiments, simultaneous ring amine deprotection and salt formation may comprise treatment of a solution of a compound of formula (III) in an organic solvent with the sulfonic acid and water.

In some embodiments, the organic solvent is toluene.

In the methods of the invention, the sulfonic acid may be selected from alkylsulfonic acids and arylsulfonic acids. It may be a sulfonic acid of formula $R^A\text{SO}_3\text{H}$ or $R^B\text{SO}_3\text{H}$, wherein R^A and R^B are as defined herein.

In some embodiments, the sulfonic acid may be a disulfonic acid, i.e. a compound containing two sulfonic acid moieties per molecule. These sulfonic acid moieties may be linked by e.g. an alkylene or arylene group.

In some embodiments the sulfonic acid may be selected from: methanesulfonic acid (MsOH), ethanesulfonic acid (EsOH), benzenesulfonic acid (BSA), naphthalenesulfonic acid (NSA), p-toluenesulfonic acid (TsOH), ethanedisulfonic acid (EDSA), propanedisulfonic acid (PDSA) and naphthalene-1,5-disulfonic acid (NDSA).

In some embodiments, the phenothiazine starting material (i.e. the compound of formula (III)) is first heated in said organic solvent until completely dissolved and the resultant solution is filtered before addition of the reagents (i.e. the sulfonic acid and water).

In some embodiments, the compound is heated in said organic solvent at a temperature of about 60-80° C., for example at a temperature of about 70° C.

In some embodiments, the sulfonic acid is added in an amount of at least 2 molar equivalents, for example about 2.2 molar equivalents, relative to the phenothiazine starting material. If a disulfonic acid is used, it will be understood the molar amount of the acid will be at least 1 molar equivalent, for example about 1.1 molar equivalents, so as to achieve the same number of sulfonic acid moieties per molecule of phenothiazine starting material.

It may be desirable to add the sulfonic acid slowly to prevent a temperature increase (exotherm). Therefore, in some embodiments, the sulfonic acid is added gradually.

In some embodiments, the sulfonic acid is added at a temperature of about 15-25° C.

In some embodiments, after addition of the sulfonic acid and water, the reaction is heated to a temperature of about 80-90° C.

In some embodiments, the reaction is maintained at this temperature until judged complete by e.g. chromatographic analysis.

25

In some embodiments, after reaction, the solution is treated with a counter solvent to precipitate the product. In some embodiments, the counter solvent is an alcohol, for example ethanol.

It may be desirable to 'seed' the reaction mixture with a small amount, for example, about 1 mg per gram of starting material (compound of formula (II), of the desired bis(sulfonate) product. Without wishing to be bound by theory, it is thought that addition of the seed ensures early and efficient precipitation of the desired product, reducing the opportunity for possible side-reactions and by-product formation. The seed is also thought to be of use in controlling the particle size of the precipitated product.


Hence in some embodiments, after reaction, the resultant mixture is seeded with a small amount of the desired bis(sulfonate) salt.

In some embodiments, the seed comprises particles of the desired bis(sulfonate) salt which have been ground.


In some embodiments, the seed comprises particles of the desired bis(sulfonate) salt which have been ground to a size of less than about 100 μm .

In some embodiments, the precipitated product is isolated by filtration.


In some embodiments, after filtration, the product is washed with an organic solvent, for example ethanol or acetonitrile. Salt formation (SF) produces the bis(sulfonate) salt of formula (I) from the compound of formula (II):




As explained above, the bis(sulfonate) salt may also be prepared directly from a corresponding amino-protected (e.g. N-acetyl) compound of formula (III).




26



In this case, salt formation may be performed at the same time as deprotection, for example by using the appropriate sulfonic acid, e.g. methanesulfonic acid, for the deprotection step. An example is illustrated in the following scheme:



In a further aspect, the present invention provides a method of preparing a compound of formula (I):



wherein R^A , R^B , R^1 , R^9 , R^{3NA} , R^{3NB} , R^{7NA} , and R^{7NB} are as previously defined the method comprising:

- 55 preparing a compound of formula (II) or (III) as defined herein, followed by salt formation (SF) and/or ring amine deprotection (DP).

60 The steps of salt formation (SF) and ring amine deprotection (DP) are as described above.

In some embodiments, preparing said compound of formula (II) or (III) comprises a method as disclosed in WO2007/110627.

65 In some embodiments, preparing said compound of formula (II) or (III) comprises a method as disclosed in WO2008/007074.

27

In some embodiments, preparing a compound of formula (II) comprises ring amino deprotection (DP) of a compound of formula (III), as set out above.


In some embodiments, preparing a compound of formula (III) comprises one or more steps selected from:

- nitration (NO),
- ring amino protection (AP),
- nitro reduction (NR),
- amine substitution (AS).

In some embodiments, preparing a compound of formula (III) comprises the steps of reduction (RED), and ring amino protection (AP).

The steps may be performed in any logical order. In some embodiments, the steps are performed in the order listed (i.e., any step in the list is performed at the same time as, or subsequent to, the preceding step in the list).


In some embodiments, nitration (NO) comprises: nitration (NO), wherein a 10H-phenothiazine is converted to a 3,7-dinitro-10H-phenothiazine, for example:



In some embodiments, nitration is performed using a nitrite, for example, sodium nitrite, for example, sodium nitrite with acetic acid, and a solvent such as dimethyl sulfoxide, dimethyl formamide, acetonitrile, tetrahydrofuran, dimethoxyethane, acetone, dichloromethane or chloroform.

In some embodiments, ring amino protection (AP) comprises:

ring amino protection (AP), wherein the ring amino group ($-\text{NH}-$) of a 3,7-dinitro-10H-phenothiazine is converted to a protected ring amino group ($-\text{NR}^{\text{Prot}}$), for example:




In some embodiments, ring amino protection is achieved as an acetate, for example, using acetic anhydride, for example, using acetic anhydride and a base such as an amine base, for example triethylamine or pyridine.

In some embodiments, nitro reduction (NR) step comprises:

28

nitro reduction (NR), wherein each of the nitro ($-\text{NO}_2$) groups of a protected 3,7-dinitro-10H-phenothiazine is converted to an amino ($-\text{NH}_2$) group, for example:




In some embodiments, nitro reduction may be performed using, for example, tin (II) chloride, for example, tin (II) chloride with ethanol.

In some embodiments, nitro reduction may be performed using, for example, palladium on charcoal (Pd/C) and hydrogen in, for example, 2-methyl-tetrahydrofuran.

In some embodiments, nitro reduction may be performed using, for example, zinc and aqueous ammonium chloride in methanol and THF

In some embodiments, amine substitution (AS) step comprises:

amine substitution (AS), wherein each of the amino ($-\text{NH}_2$) groups of a protected 3,7-diamino-10H-phenothiazine is converted to disubstituted amino group, for example:



In some embodiments, amine substitution is performed using an alkyl halide, for example, an alkyl iodide, for example, methyl iodide, for example, methyl iodide with sodium hydroxide, DMSO, toluene and tetra-n-butyl ammonium bromide.

In some embodiments, amine substitution comprises treatment with formaldehyde (e.g. paraformaldehyde, formalin) under reducing conditions. For example, treatment with formalin and hydrogen gas, in the presence of a Pd/C catalyst; or treatment with paraformaldehyde in the presence of a reducing agent such as sodium cyanoborohydride and acetic acid.

In some embodiments, the reduction (RED) step is: reduction (RED), wherein a 3,7-di(disubstituted amino)-thioninium salt is reduced to give the corresponding 3,7-di(disubstituted amino)-10H-phenothiazine, for example by treatment with a reducing agent, such as hydrazine (NH_2NH_2), methyl hydrazine (MeNHNH_2),

or sodium borohydride and a base, such as pyridine, triethylamine, or Hünig's base (diisopropylethylamine). In some embodiments, the ring amino protection (AP) step is:


ring amino protection (AP), wherein a 3,7-di(disubstituted amino)-10H-phenothiazine is protected, for example by treatment with acetic anhydride, to give the corresponding protected 3,7-di(disubstituted amino)-10H-phenothiazine, for example the corresponding N-acetyl 3,7-di(disubstituted amino)-10H-phenothiazine.

In some embodiments, the steps are performed in the order listed (i.e., any step in the list is performed at the same time as, or subsequent to, the preceding step in the list).

In some embodiments, the step of reduction (RED) and the step of ring amino protection (AP) are performed simultaneously (i.e., as one step).

For example, in some embodiments, the combined reduction (RED) step and ring amino protection (AP) step is:

reduction (RED) and ring amino protection (AP), wherein a 3,7-di(disubstituted amino)-thioninium salt is reduced to give the corresponding 3,7-di(disubstituted amino)-10H-phenothiazine, and the ring amino group (—NH—) of the 3,7-di(disubstituted amino)-10H-phenothiazine is converted to a protected ring amino group (—R^{Prot}) to give the corresponding protected 3,7-di(disubstituted amino)-10H-phenothiazine, for example:



wherein Y is a counterion. In some embodiments, Y represents Cl⁻.

In some embodiments, the 3,7-di(disubstituted amino)-thioninium salt is methylthioninium chloride (MTC).

In some embodiments, the combined reduction (RED) step and ring amino protection (AP) step is achieved using a hydrazine, such as phenylhydrazine, MeNHNH₂, or NH₂NH₂·H₂O and acetic anhydride.

In some embodiments, the step is performed under a nitrogen atmosphere.

In some embodiments the combined reduction (RED) step and ring amino protection (AP) step is performed using, for example, phenylhydrazine, ethanol, acetic anhydride, and pyridine.

In some embodiments, the combined reduction (RED) step and ring amino protection (AP) step is performed using, for example, hydrazine hydrate, acetonitrile, acetic anhydride, and triethylamine, under a nitrogen atmosphere.

In some embodiments, the protected 3,7-di(disubstituted amino)-10H-phenothiazine, for example the N-acetyl 3,7-di(disubstituted amino)-10H-phenothiazine, undergoes a purification step.

In some embodiments, purification comprises addition of an organic solvent, for example toluene, and an acid, for example acetic acid, to dissolve the compound, followed by a washing step.

In some embodiments, washing comprises addition of water and/or aqueous acetic acid to the solution of the compound; agitation and/or heating; and separation of the organic layer.

In some embodiments, washing is repeated, for example up to three times.

In some embodiments, washing is followed by isolation of the purified product.

In some embodiments, isolation of the purified product comprises cooling, precipitation and filtration of the product.

Crystalline Forms

In some embodiments, the compound of the invention is provided in crystalline form.

In some embodiments, the crystalline form is 'Form A' as described herein.

In some embodiments, the crystalline form has the structure depicted in FIG. 17 and/or is characterised by the crystal data shown in an Annex Table 1 and/or the atomic co-ordinates shown in an Annex Table 2 and/or the bond lengths and angles shown in an Annex Table 3 and/or the anisotropic displacement parameters shown in an Annex Table 4 and/or the hydrogen coordinates and isotropic displacement parameters shown in an Annex Table 5.

Reversing and/or Inhibiting the Aggregation of a Protein

One aspect of the invention is the use of a compound or composition as described herein, to regulate (e.g., to reverse and/or inhibit) the aggregation of a protein, for example, aggregation of a protein associated with a neurodegenerative disease and/or clinical dementia. The aggregation may be in vitro, or in vivo, and may be associated with a disease state as discussed below.

Thus, one aspect of the invention pertains to a method of regulating (e.g., reversing and/or inhibiting) the aggregation of a protein, for example, aggregation of a protein associated with a neurodegenerative disease and/or clinical dementia, comprising contacting the protein with an effective amount of a compound or composition as described herein. The method may be performed in vitro, or in vivo.

Similarly, one aspect of the invention pertains to a method of regulating (e.g., reversing and/or inhibiting) the aggregation of a protein in the brain of a mammal, which aggregation is associated with a disease state as described herein, the treatment comprising the step of administering to said mammal in need of said treatment, a prophylactically or therapeutically effective amount of a compound or composition as described herein, that is an inhibitor of said aggregation.

Methods of Treatment

Another aspect of the present invention pertains to a method of treatment comprising administering to a patient in need of treatment a prophylactically or therapeutically effective amount of a compound as described herein, preferably in the form of a pharmaceutical composition.

Use in Methods of Therapy

Another aspect of the present invention pertains to a compound or composition as described herein, for use in a method of treatment (e.g., of a disease condition) of the human or animal body by therapy.

Use in the Manufacture of Medicaments

Another aspect of the present invention pertains to use of a compound or composition as described herein, in the manufacture of a medicament for use in treatment (e.g., of a disease condition).

In some embodiments, the medicament comprises a compound of the invention.

In some embodiments, the medicament is a composition as described hereinbelow.

Disease Conditions Treated—Diseases of Protein Aggregation

The compounds and compositions of the present invention are useful in the treatment or prophylaxis of diseases of protein aggregation.

Thus, in some embodiments, the disease condition is a disease of protein aggregation, and, for example, the treatment is with an amount of a compound or composition as described herein, sufficient to inhibit the aggregation of the protein associated with said disease condition.

In general, the protein aggregation is that which arises from an induced conformational polymerisation interaction, i.e.,

one in which a conformational change of the protein, or in a fragment thereof, gives rise to templated binding and aggregation of further (precursor) protein molecules in a self-propagating manner. Once nucleation is initiated, an aggregation cascade may ensue which involves the induced conformational polymerisation of further protein molecules, leading to the formation of toxic product fragments in aggregates which are substantially resistant to further proteolysis. The protein aggregates thus formed are thought to be a proximal cause of disease states manifested as neurodegeneration, clinical dementia, and other pathological symptoms.

The following Table lists various disease-associated aggregating proteins and the corresponding diseases of protein aggregation. The use of the compounds and compositions of the invention in respect of these proteins or diseases is encompassed by the present invention.

Diseases of protein aggregation				
Protein	Disease	Aggregating domain and/or mutations	Fibril subunit size (kDa)	Reference
Neurodegenerative disorders				
Prion protein	Prion diseases (CJD, nvCJD, Fatal familial insomnia, Gerstmann-Straussler-Scheinker syndrome, Kuru)	Inherited and sporadic forms PrP-27-30; many mutations.	27	Prusiner (1998)
			27	Prusiner (1998)
		Fibrillogenic domains: 113-120, 178-191, 202-218.		Gasset et al. (1992)
Tau protein	Alzheimer's disease, Down's syndrome, FTDP-17, CBD, post-encephalitic parkinsonism, Pick's disease, parkinsonism with dementia complex of Guam	Inherited and sporadic forms	10-12	Wischnik et al. (1988)
			10-12	Wischnik et al. (1988)
		Truncated tau (tubulin-binding domain) 297-391. Mutations in tau in FTDP-17. Many mutations in presenilin proteins.		Hutton et al. (1998) Czech et al. (2000)
Amyloid β -protein	Alzheimer's disease, Down's syndrome	Inherited and sporadic forms Amyloid β -protein; 1-42(3). Mutations in APP in rare families.	4	Glenner & Wong, (1984)
			4	Glenner & Wong, (1984) Goate et al. (1991)
Huntingtin	Huntington's disease	N-termini of protein with expanded glutamine repeats.	40	DiFiglia et al. (1997)
Ataxins (1, 2, 3, 7)	Spinocerebellar ataxias (SCA1, 2, 3, 7)	Proteins with expanded glutamine repeats.		Paulson et al. (1999)
Atrophin	Dentatorubropallidoluysian atrophy (DRPLA)	Proteins with expanded glutamine repeats.		Paulson et al. (1999)
Androgen receptor	Spinal and bulbar muscular atrophy	Proteins with expanded glutamine repeats.		Paulson et al. (1999)

-continued

Diseases of protein aggregation				
Protein	Disease	Aggregating domain and/or mutations	Fibril subunit size (kDa)	Reference
Neuroserpin	Familial encephalopathy with neuronal inclusion bodies (FENIB)	Neuroserpin; S49P, S52R.	57	Davis et al. (1999)
α -Synuclein	Parkinson's disease, dementia with Lewy bodies, multiple system atrophy	Inherited and sporadic forms	19	Spillantini et al. (1998) also PCT/GB2007/001105
		A53T, A30P in rare autosomal-dominant PD families.		Polymeropoulos et al. (1997)
TDP-43	FTLD-TDP	Several TDP-43 mutations	10-43	Mackenzie et al. (2010)
	Amyotrophic lateral sclerosis	Several TDP-43 mutations	10-43	Mackenzie et al. (2010)
Cystatin C	Hereditary cerebral angiopathy (Icelandic)	Cystatin C less 10 residues; L68Q.	12-13	Abrahamson et al. (1992)
Superoxide dismutase 1	Amyotrophic lateral sclerosis	SOD1 mutations.	16	Shibata et al. (1996)
Non-neurodegenerative disorders				
Haemoglobin	Sickle cell anaemia	Haemoglobin beta chain (S).		Carrell & Gooptu (1998)
	Inclusion body haemolysis	Many mutations.		
Serpins	α 1-Antitrypsin deficiency (emphysema, cirrhosis)	Mutations		Lomas et al. (1992)
	Antithrombin deficiency (thromboembolic disease)	Mutations		Carrell & Gooptu (1998)
	C1-inhibitor deficiency (angioedema)	Mutations		Carrell & Gooptu (1998)
Immunoglobulin light chain	Plasma cell dyscrasias (primary systemic AL amyloidosis)	Light chain or fragments.	0.5-25	Westermarck et al. (1985)
Serum amyloid A	Reactive, secondary systemic AA amyloidosis	76-residue fragment (critical residues 2-12).	4.5-7.5	Westermarck et al. (1985)
	Chronic inflammatory disease			
Transthyretin	Familial amyloid polyneuropathy (systemic; FAP I)	Tetramer dissociated to conformational monomer variant. Many mutations (some not associated with amyloid; several different types of disease).	10-14	Gustavsson et al. (1991)
	Senile cardiac amyloidosis	Normal transthyretin	10-14	Gustavsson et al. (1991)
Gelsolin	Familial amyloidosis - Finnish type (FAP IV)	D187Q leads to truncated 173-225/243 (critical residues 182-192).	9.5	Maury & Baumann (1990)
β 2-Microglobulin	Haemodialysis amyloidosis	β 2-Microglobulin	12-25	Gorevic et al. (1985)
	Prostatic amyloid			
Apolipoprotein AI	Familial amyloid polyneuropathy (systemic; FAP III)	N-terminal 83-93 residues; G26R, W50R, L60R	9	Booth et al. (1997)
Lysozyme	Familial visceral amyloidosis	Lysozyme or fragments (with or without I56T, D67H)	14	Pepys et al. (1993)
Amylin (Islet amyloid polypeptide)	Type II diabetes (NIDDM)	Fragments (critical core of 20-29); no mutations	3.9	Westermarck (1990)
Fibrinogen α -chain	Hereditary renal amyloidosis	Fibrinogen fragments	7-10	Uemichi et al. (1992)

Diseases of protein aggregation				
Protein	Disease	Aggregating domain and/or mutations	Fibril subunit size (kDa)	Reference
Procalcitonin	Medullary carcinoma of thyroid	Calcitonin fragments	3.4	Sletten et al. (1976)
Atrial natriuretic factor	Cardiac amyloidosis	ANF, no mutants	3.5	Johansson et al. (1987)
Insulin	Injection localised amyloidosis	Insulin		Dische et al. (1988)
Multiple proteins	Inclusion body myositis	β -amyloid, tau, ubiquitin, ApoE, and presenilin-1		Askenas et al (2009)
Other proteins forming amyloid	(in vitro)	Other proteins		Chiti et al. (1999)

As described in WO 02/055720, WO2007/110630, and WO2007/110627, diaminothiazines have utility in the inhibition of such protein aggregating diseases.

Thus it will be appreciated that, except where context requires otherwise, description of embodiments with respect to tau protein or tau-like proteins (e.g., MAP2; see below), should be taken as applying equally to the other proteins discussed herein (e.g., β -amyloid, synuclein, prion, etc.) or other proteins which may initiate or undergo a similar pathological aggregation by virtue of conformational change in a domain critical for propagation of the aggregation, or which imparts proteolytic stability to the aggregate thus formed (see, e.g., the article by Wischik et al. in "Neurobiology of Alzheimer's Disease", 2nd Edition, 2000, Eds. Dawbarn, D. and Allen, S. J., The Molecular and Cellular Neurobiology Series, Bios Scientific Publishers, Oxford). All such proteins may be referred to herein as "aggregating disease proteins."

Likewise, where mention is made herein of "tau-tau aggregation", or the like, this may also be taken to be applicable to other "aggregating-protein aggregation", such as β -amyloid aggregation, prion aggregation, synuclein aggregation, etc. The same applies for "tau proteolytic degradation" etc.

Preferred Aggregating Disease Proteins

Preferred embodiments of the invention are based on tau protein. The term "tau protein," as used herein, refers generally to any protein of the tau protein family. Tau proteins are characterised as being one among a larger number of protein families which co-purify with microtubules during repeated cycles of assembly and disassembly (see, e.g., Shelanski et al., 1973, Proc. Natl. Acad. Sci. USA, Vol. 70, pp. 765-768), and are known as microtubule-associated-proteins (MAPs). Members of the tau family share the common features of having a characteristic N-terminal segment, sequences of approximately 50 amino acids inserted in the N-terminal segment, which are developmentally regulated in the brain, a characteristic tandem repeat region consisting of 3 or 4 tandem repeats of 31-32 amino acids, and a C-terminal tail.

MAP2 is the predominant microtubule-associated protein in the somatodendritic compartment (see, e.g., Matus, A., in "Microtubules" [Hyams and Lloyd, Eds.] pp. 155-166, John Wiley and Sons, New York, USA). MAP2 isoforms are almost identical to tau protein in the tandem repeat region, but differ substantially both in the sequence and extent of the N-terminal domain (see, e.g., Kindler and Garner, 1994, Mol. Brain Res., Vol. 26, pp. 218-224). Nevertheless, aggregation in the tandem-repeat region is not selective for the tau repeat domain. Thus it will be appreciated that any discussion herein in relation to tau protein or tau-tau aggregation should be taken as relating also to tau-MAP2 aggregation, MAP2-MAP2 aggregation, and so on.

20

In some embodiments, the protein is tau protein.

In some embodiments, the protein is a synuclein, e.g., α - or β -synuclein.

In some embodiments, the protein is TDP-43.

25 TAR DNA-Binding Protein 43 (TDP-43) is a 414 amino acid protein encoded by TARDBP on chromosome 1p36.2. The protein is highly conserved, widely expressed, and predominantly localised to the nucleus but can shuttle between the nucleus and cytoplasm (Mackenzie et al 2010). It is involved in transcription and splicing regulation and may have roles in other processes, such as: microRNA processing, apoptosis, cell division, stabilisation of messenger RNA, regulation of neuronal plasticity and maintenance of dendritic integrity. Furthermore, since 2006 a substantial body of evidence has accumulated in support of the TDP-43 toxic gain of function hypothesis in amyotrophic lateral sclerosis (ALS). TDP-43 is an inherently aggregation-prone protein and aggregates formed in vitro are ultrastructurally similar to the TDP-43 deposits seen in degenerating neurones in ALS patients (Johnson et al 2009). Johnson et al (2008) showed that when TDP-43 is overexpressed in a yeast model only the aggregated form is toxic. Several in vitro studies have also shown that C-terminal fragments of TDP-43 are more likely than full-length TDP-43 to form insoluble cytoplasmic aggregates that become ubiquitinated, and toxic to cells (Arai et al 2010; Igaz et al 2009; Nonaka et al 2009; Zhang et al 2009). Though Nonaka et al (2009) suggested that these cytoplasmic aggregates bind the endogenous full-length protein depleting it from the nucleus, Zhang et al (2009) found retention of normal nuclear expression, suggesting a purely toxic effect for the aggregates. Yang et al (2010) have described the capture of full-length TDP-43 within aggregates of C- and N-terminal fragments of TDP-43 in NSC34 motor neurons in culture. Neurite outgrowth, impaired as a result of the presence of such truncated fragments, could be rescued by overexpression of the full-length protein. Although the role of neurite outgrowth in vivo has not been established, this model would support the suggestion made by Nonaka and colleagues for a role of TDP-43 aggregation in ALS pathogenesis.

60 Mutant TDP-43 expression in cell cultures has repeatedly been reported to result in increased generation of C-terminal fragments, with even greater cytoplasmic aggregation and toxic effects than the wild-type protein (Kabashi et al 2008; Sreedharan et al 2008; Johnson et al 2009; Nonaka et al 2009; Arai et al 2010; Barmarda et al 2010; Kabashi et al 2010).

Where the protein is tau protein, in some embodiments of the present invention, there is provided a method of inhibiting

production of protein aggregates (e.g. in the form of paired helical filaments (PHFs), optionally in neurofibrillary tangles (NFTs) in the brain of a mammal, the treatment being as described above.

Preferred Indications—Diseases of Protein Aggregation

Notably it is not only Alzheimer's disease (AD) in which tau protein (and aberrant function or processing thereof) may play a role. The pathogenesis of neurodegenerative disorders such as Pick's disease and progressive supranuclear palsy (PSP) appears to correlate with an accumulation of pathological truncated tau aggregates in the dentate gyrus and stellate pyramidal cells of the neocortex, respectively. Other dementias include fronto-temporal dementia (FTD); FTD with parkinsonism linked to chromosome 17 (FTDP-17); disinhibition-dementia-parkinsonism-amyotrophy complex (DDPAC); pallido-ponto-nigral degeneration (PPND); Guam-ALS syndrome; pallido-nigro-luysian degeneration (PNLD); cortico-basal degeneration (CBD) and others (see, e.g., the article by Wischik et al. in "Neurobiology of Alzheimer's Disease", 2nd Edition, 2000, Eds. Dawbarn, D. and Allen, S. J., The Molecular and Cellular Neurobiology Series, Bios Scientific Publishers, Oxford; especially Table 5.1). All of these diseases, which are characterized primarily or partially by abnormal tau aggregation, are referred to herein as "tauopathies".

Thus, in some embodiments, the disease condition is a tauopathy.

In some embodiments, the disease condition is a neurodegenerative tauopathy.

In some embodiments, the disease condition is selected from Alzheimer's disease (AD), Pick's disease, progressive supranuclear palsy (PSP), fronto temporal dementia (FTD), FTD with parkinsonism linked to chromosome 17 (FTDP 17), frontotemporal lobar degeneration (FTLD) syndromes; disinhibition-dementia-parkinsonism-amyotrophy complex (DDPAC), pallido-ponto-nigral degeneration (PPND), Guam-ALS syndrome, pallido nigro luysian degeneration (PNLD), cortico-basal degeneration (CBD), dementia with argyrophilic grains (AgD), dementia pugilistica (DP) or chronic traumatic encephalopathy (CTE), Down's syndrome (DS), dementia with Lewy bodies (DLB), subacute sclerosing panencephalitis (SSPE), MCI, Niemann-Pick disease, type C (NPC), Sanfilippo syndrome type B (mucopolysaccharidosis III B), or myotonic dystrophies (DM), DM1 or DM2, or chronic traumatic encephalopathy (CTE).

In some embodiments, the disease condition is a lysosomal storage disorder with tau pathology. NPC is caused by mutations in the gene NPC1, which affects cholesterol metabolism (Love et al 1995) and Sanfilippo syndrome type B is caused by a mutation in the gene NAGLU, in which there is lysosomal accumulation of heparin sulphate (Ohmi et al. 2009). In these lysosomal storage disorders, tau pathology is observed and its treatment may decrease the progression of the disease. Other lysosomal storage disorders may also be characterised by accumulation of tau.

Use of phenothiazine diaminium salts in the treatment of Parkinson's Disease and MCI is described in more detail in PCT/GB2007/001105 and PCT/GB2008/002066.

In some embodiments, the disease condition is Parkinson's Disease, MCI, or Alzheimer's disease.

In some embodiments, the disease condition is Huntington's Disease or other polyglutamine disorder such as spinal bulbar muscular atrophy (or Kennedy disease), and dentatorubropallidoluysian atrophy and various spinocerebellar ataxias.

In some embodiments, the disease condition is an FTLD syndrome (which may for example be a tauopathy or TDP-43 proteinopathy, see below).

In some embodiments, the disease condition is PSP or ALS.

In some embodiments, treatment (e.g., treatment of a neurodegenerative tauopathy, e.g., Alzheimer's disease) may optionally be in combination with one or more other agents, for example, one or more cholinesterase inhibitors (such as Donepezil (also known as Aricept™), Rivastigmine (also known as Exelon™), Galantamine (also known as Reminyl™), NMDA receptor antagonists (such as Memantine (also known as Ebixa™, Namenda™), muscarinic receptor agonists, and/or inhibitors of amyloid precursor protein processing that leads to enhanced generation of beta-amyloid.

TDP-43 proteinopathies include amyotrophic lateral sclerosis (ALS; ALS-TDP) and frontotemporal lobar degeneration (FTLD-TDP).

The role of TDP-43 in neurodegeneration in ALS and other neurodegenerative disorders has been reviewed in several recent publications (Chen-Plotkin et al 2010; Gendron et al 2010; Geser et al 2010; Mackenzie et al 2010).

ALS is a neurodegenerative disease, characterised by progressive paralysis and muscle wasting, consequent on the degeneration of both upper and lower motor neurones in the primary motor cortex, brainstem and spinal cord. It is sometimes referred to as motor neuron disease (MND) but there are diseases other than ALS which affect either upper or lower motor neurons. A definite diagnosis requires both upper and lower motor neurone signs in the bulbar, arm and leg musculature with clear evidence of clinical progression that can not be explained by any other disease process (Wijesekera and Leigh 2009).

Although the majority of cases are ALS-TDP, there are other cases where the pathological protein differs from TDP-43. Misfolded SOD1 is the pathological protein in ubiquitin-positive inclusions in ALS with SOD1 mutations (Seetharaman et al 2009) and in a very small subset (approximately 3-4%) of familial ALS, due to mutations in FUS (fused in sarcoma protein), the ubiquitinated pathological protein is FUS (Vance et al 2009; Blair et al 2010). FUS, like TDP-43, appears to be important in nuclear-cytoplasmic shuttling although the ways in which impaired nuclear import of FUS remains unclear. A new molecular classification of ALS, adapted from Mackenzie et al (2010), reflects the distinct underlying pathological mechanisms in the different subtypes (see Table below).

New Molecular Classification of ALS (modified from Mackenzie et al 2010). In the majority of cases, TDP-43 is the pathological ubiquitinated protein found in ALS.

Ubiquitin-positive inclusions in ALS

Ubiquitinated disease protein	TDP-43	FUS	SOD1
Clinico-pathologic subtype	ALS-TDP	ALS-FUS	ALS-SOD1
Associated genotype	TARDBP	FUS	SOD1
Frequency of ALS cases	Common	Rare	Rare

Amyotrophic lateral sclerosis has been recognised as a nosological entity for almost a century and a half and it is recognised in ICD-10 is classified as a subtype of MND in ICD 10 (G12.2). Reliable clinical diagnostic are available for

ALS, which differ little from Charcot's original description, and neuropathological criteria, reflecting the underlying molecular pathology, have also been agreed.

While ALS is classified pathologically into three subgroups, ALS-TDP, ALS-SOD1 and ALS-FUS, both latter conditions are rare. The largest study to date showed all sporadic ALS cases to have TDP-43 pathology (Mackenzie et al 2007). Only around 5% of ALS is familial (Byrne et al 2010) and mutations in SOD1, the commonest mutations found in FALS, account for between 12-23% of cases (Andersen et al 2006). SOD1 may also be implicated in 2-7% of SALS. Mutations in FUS appear to be far less common, accounting for only around 3-4% of FALS (Blair et al 2010). So it can be reliably predicted that a clinical case of SALS will have TDP-43 based pathology. Similarly this can be reliably predicted in FALS due to mutations in TDP-43, which

middle age. There is often a positive family history of similar disorders in a first degree relative.

Behavioural variant FTD is characterised by early prominent change in social and interpersonal function, often accompanied by repetitive behaviours and changes in eating pattern. In semantic dementia there are prominent word finding problems, despite otherwise fluent speech, with degraded object knowledge and impaired single word comprehension on cognitive assessment. Progressive non-fluent aphasia presents with a combination of motor speech problems and grammatical deficits. The core clinical diagnostic features for these three FTLD syndromes are shown in the Table below and the full criteria in Neary et al (1998).

Clinical Profile and Core Diagnostic Features of FTLD Syndromes

FTLD Syndrome-Clinical Profile	Core Diagnostic Features
<p>Frontotemporal Dementia Character change and disordered social conduct are the dominant features initially and throughout the disease course. Instrumental functions of perception, spatial skills, praxis and memory are intact or relatively well preserved.</p>	<ol style="list-style-type: none"> 1. Insidious onset and gradual progression 2. Early decline in social interpersonal conduct 3. Early impairment in regulation of personal conduct 4. Early emotional blunting 5. Early loss of insight
<p>Semantic Dementia Semantic disorder (impaired understanding of word meaning and/or object identity) is the dominant feature initially and throughout the disease course. Other aspects of cognition, including autobiographic memory, are intact or relatively well preserved.</p>	<ol style="list-style-type: none"> A) Insidious onset and gradual progression B) Language disorder characterised by <ol style="list-style-type: none"> 1. Progressive, fluent empty speech 2. Loss of word meaning manifest by impaired naming and comprehension 3. Semantic paraphasias and/or 4. Perceptual disorder characterised by <ol style="list-style-type: none"> 1. Prosopagnosia: impaired recognition of identity of familiar faces and/or 2. Associative agnosia: impaired recognition of object identity C) Preserved perceptual matching and drawing reproduction D) Preserved single word repetition E) Preserved ability to read aloud and write to dictation orthographically regular words
<p>Progressive Non-fluent Aphasia Disorder of expressive language is the dominant feature initially and throughout the disease course. Other aspects of cognition are intact or relatively well preserved.</p>	<ol style="list-style-type: none"> A) Insidious onset and gradual progression B) Non-fluent spontaneous speech with at least one of the following: agrammatism, phonemic paraphasias or anomia

account for around 4% of cases (Mackenzie et al 2010). ALS with mutations in: VCP, accounting for 1-2% of FALS (Johnson et al 2010), ANG (Seilhean et al 2009), and CHMP2B (Cox et al 2010) have also been reported to be associated with TDP-43 positive pathology. Although SOD1, FUS and ATXN2 mutations have not been found to be associated with TDP-43 positive aggregates, it has however been reported that TDP-43 is implicated in the pathological processes putatively arising from these mutations (Higashi et al 2010; Ling et al 2010; Elden et al 2010).

It is therefore established that TDP-43 has an important, and potentially central role, in the pathogenesis of the vast majority of SALS cases and may be implicated in the pathogenesis of a significant proportion of FALS. ALS is now widely considered to be a TDP-43 proteinopathy (Neumann et al 2009) and numerous in vitro, and in vivo studies provide support to the hypothesis that toxic gain of function, due to TDP-43 aggregation is responsible for at least some of the neurotoxicity in the disease.

FTLD syndromes are insidious onset, inexorably progressive, neurodegenerative conditions, with peak onset in late

The discovery that TDP-43-positive inclusions characterize ALS and FTLD-TDP (Neumann et al 2006) was quickly followed by the identification of missense mutations in the TARDBP gene in both familial and sporadic cases of ALS (Gitcho et al 2008; Sreedharan et al., 2008). So far, 38 different TARDBP mutations have been reported in 79 genealogically unrelated families worldwide (Mackenzie et al 2010). TARDBP mutations account for approximately 4% of all familial and around 1.5% of sporadic ALS cases.

As of December 2010, mutations in thirteen genes which are associated with familial and sporadic ALS have been identified. Linkage of ALS to five other chromosome loci has been demonstrated but thus far specific mutations have not been identified.

Methylthionium (MT) in TDP-43 Proteinopathies

MT has a mode of action which targets and can reduce TDP-43 protein aggregation in cells, which is a pathological feature of the vast majority of both familial and sporadic ALS and is also characteristic of FTLD-P.

In addition laboratory data shows that methylthionium inhibits the formation of TDP-43 aggregates in SH-SY5Y

cells. Following treatment with 0.05 μ M MT, the number of TDP-43 aggregates was reduced by 50%. These findings were confirmed by immunoblot analysis (Yamashita et al 2009).

The compounds and compositions of the invention may therefore be useful for the treatment of amyotrophic lateral sclerosis (ALS) and frontotemporal lobar degeneration (FTLD).

Methylthioninium (MT) in Huntington's Disease and Polyglutamine Disorders

MT can reduce polyglutamine protein aggregation in cells, which is a pathological feature of Huntington's disease. Huntington's disease is caused by expansion of a translated CAG repeat located in the N-terminus of huntingtin. Wild-type chromosomes contain 6-34 repeats whereas, in Huntington's disease, chromosomes contain 36-121 repeats. The age of onset of disease correlates inversely with the length of the CAG tracts that code for polyglutamine repeats within the protein.

Laboratory data shows that methylthioninium inhibits the formation of aggregates of a huntingtin derivative containing a polyglutamine stretch of 102 residues in zebrafish (van Bebber et al. 2010). MT, when tested at 0, 10 and 100 μ M, prevented the formation of such aggregates in zebrafish in a dose dependent manner.

The compounds and compositions of the invention may therefore be useful for the treatment of Huntington's disease and other polyglutamine disorders such as spinal bulbar muscular atrophy (or Kennedy disease), and dentatorubropallidoluysian atrophy and various spinocerebellar ataxias (Orr & Zoghbi, 2007).

Mitochondrial Diseases and Lafora Disease

The organ most frequently affected in mitochondrial disorders, particularly respiratory chain diseases (RCDs), in addition to the skeletal muscle, is the central nervous system (CNS). CNS manifestations of RCDs comprise stroke-like episodes, epilepsy, migraine, ataxia, spasticity, movement disorders, psychiatric disorders, cognitive decline, or even dementia (mitochondrial dementia). So far mitochondrial dementia has been reported in MELAS, MERRE, LHON, CPEO, KSS, MNGIE, NARP, Leigh syndrome, and Alpers-Huttenlocher disease (Finsterer, 2009). There are four complexes in the mitochondrial respiration chain, involving a series of electron transfers. Abnormal function of any of these complexes can result in mitochondrial diseases secondary to an abnormal electron transport chain and subsequent abnormal mitochondrial respiration. Complex III of the mitochondrial respiration chain acts to transfer electrons to cytochrome c.

Compounds and compositions of the invention may also be used to treat mitochondrial diseases which are associated with a deficient and/or impaired complex III function of the respiration chain. The compounds have the ability to act as effective electron carrier and/or transfer, as the thioninium moiety has a low redox potential converting between the oxidised and reduced form. In the event of an impaired and/or deficient function of Complex III leading to mitochondrial diseases, compounds of the invention are also able to perform the electron transportation and transfer role of complex III because of the ability of the thioninium moiety to shuttle between the oxidised and reduced form, thus acting as an electron carrier in place of sub-optimally functioning complex III, transferring electrons to cytochrome c.

Compounds and compositions of the invention also have the ability to generate an active thioninium moiety that has the ability to divert misfolded protein/amino acid monomers/oligomers away from the Hsp70 ADP-associated protein

accumulation and/or refolding pathways, and instead rechannel these abnormal folded protein monomers/oligomers to the pathway that leads directly to the Hsp70 ATP-dependent ubiquitin-proteasome system (UPS), a pathway which removes these misfolded proteins/amino acid monomers/oligomers via the direct route (Jinwal et al. 2009).

Lafora disease (LD) is an autosomal recessive teenage-onset fatal epilepsy associated with a gradual accumulation of poorly branched and insoluble glycogen, termed polyglucosan, in many tissues. In the brain, polyglucosan bodies, or Lafora bodies, form in neurons. Inhibition of Hsp70 ATPase by MT (Jinwal et al. 2009) may upregulate the removal of misfolded proteins. Lafora disease is primarily due to a lysosomal ubiquitin-proteasomal system (UPS) defect because of a mutation in either the Laforin or Malin genes, both located on Chromosome 6, which result in inclusions that may accelerate the aggregation of misfolded tau protein. Secondary mitochondrial damage from the impaired UPS may further result in a suppressed mitochondrial activity and impaired electron transport chain leading to further lipofuscin and initiating the seizures that are characteristic of Lafora disease.

The MT moiety may disaggregate existing tau aggregates, reduce more tau accumulating and enhance lysosomal efficiency by inhibiting Hsp70 ATPase. MT may lead to a reduction in tau tangles by enhancing the ubiquitin proteasomal system removal of tau monomers/oligomers, through its inhibitory action on Hsp70 ATPase.

Thus compounds and compositions of the present invention may have utility in the treatment of Lafora disease.

Disease Conditions Treated—Other Disease Conditions

In some embodiments, the disease condition is skin cancer.

In some embodiments, the disease condition is melanoma.

In some embodiments, the disease condition is a viral, bacterial or protozoal disease condition.

In some embodiments, the (protozoal) disease condition is malaria. Treatment may be in combination with one or more antimicrobial agents, for example, chloroquine and/or atovaquone.

In some embodiments, the (viral) disease condition is caused by Hepatitis C, HIV, or West Nile Virus (WNV).

Other Uses

Another aspect of the present invention pertains to use of a compound as described herein, in a method of inactivating a pathogen in a sample (for example a blood or plasma sample), comprising the steps of introducing the compound into the sample, and exposing the sample to light.

For example, in some embodiments, the method comprises the steps of introducing the compound into the sample, and then exposing the sample to light.

Use as Ligands

The compounds described herein that are capable of inhibiting the aggregation of tau protein will also be capable of acting as ligands or labels of tau protein (or aggregated tau protein). Thus, in some embodiments, the compound of the invention is a ligand of tau protein (or aggregated tau protein).

Such compounds (ligands) may incorporate, be conjugated to, be chelated with, or otherwise be associated with, other chemical groups, such as stable and unstable detectable isotopes, radioisotopes, positron-emitting atoms, magnetic resonance labels, dyes, fluorescent markers, antigenic groups, therapeutic moieties, or any other moiety that may aid in a prognostic, diagnostic, or therapeutic application.

For example, in some embodiments, the compound is as defined herein, but with the additional limitation that the compound incorporates, is conjugated to, is chelated with, or is otherwise associated with, one or more (e.g., 1, 2, 3, 4, etc.) detectable labels, for example, isotopes, radioisotopes,

positron-emitting atoms, magnetic resonance labels, dyes, fluorescent markers, antigenic groups, or therapeutic moieties.

In some embodiments, the compound is a ligand as well as a label, e.g., a label for tau protein (or aggregated tau protein), and incorporates, is conjugated to, is chelated with, or is otherwise associated with, one or more (e.g., 1, 2, 3, 4, etc.) detectable labels.

For example, in some embodiments, the compound is as defined above, but with the additional limitation that the compound incorporates, is conjugated to, is chelated with, or is otherwise associated with, one or more (e.g., 1, 2, 3, 4, etc.) detectable labels.

Labelled compounds (e.g., when ligated to tau protein or aggregated tau protein) may be visualised or detected by any suitable means, and the skilled person will appreciate that any suitable detection means as is known in the art may be used.

For example, the compound (ligand-label) may be suitably detected by incorporating a positron-emitting atom (e.g., ^{11}C) (e.g., as a carbon atom of one or more alkyl group substituents, e.g., methyl group substituents) and detecting the compound using positron emission tomography (PET) as is known in the art.

Such ^{11}C . labelled compounds may be prepared by adapting the methods described herein in known ways, for example, in analogy to the methods described in WO 02/075318 (see FIGS. 11a, 11b, 12 therein) and WO 2005/030676.

Thus, another aspect of the present invention pertains to a method of labelling tau protein (or aggregated tau protein) comprising the step of: (i) contacting the tau protein (or aggregated tau protein) with a compound that incorporates, is conjugated to, is chelated with, or is otherwise associated with, one or more (e.g., 1, 2, 3, 4, etc.) detectable labels. The compound may be provided as a composition as described herein.

Another aspect of the present invention pertains to a method of detecting tau protein (or aggregated tau protein) comprising the steps of: (i) contacting the tau protein (or aggregated tau protein) with a compound that incorporates, is conjugated to, is chelated with, or is otherwise associated with, one or more (e.g., 1, 2, 3, 4, etc.) detectable labels, and (ii) detecting the presence and/or amount of said compound bound to tau protein (or aggregated tau protein). The compound may be provided as a composition as described herein.

Another aspect of the present invention pertains to a method of diagnosis or prognosis of a tau proteinopathy in a subject believed to suffer from the disease, comprising the steps of: (i) introducing into the subject a compound capable of labelling tau protein or aggregated tau protein, particularly tau protein (e.g., a compound that incorporates, is conjugated to, is chelated with, or is otherwise associated with, one or more (e.g., 1, 2, 3, 4, etc.) detectable labels); (ii) determining the presence and/or amount of said compound bound to tau protein or aggregated tau protein in the brain of the subject; and (iii) correlating the result of the determination made in (ii) with the disease state of the subject. The compound may be provided as a composition as described herein.

Another aspect of the present invention pertains to a compound capable of labelling tau protein or aggregated tau protein (e.g., a compound that incorporates, is conjugated to, is chelated with, or is otherwise associated with, one or more (e.g., 1, 2, 3, 4, etc.) detectable labels), for use in a method of diagnosis or prognosis of a tau proteinopathy. The compound may be provided as a composition as described herein.

Another aspect of the present invention pertains to use of a compound of the invention capable of labelling tau protein or

aggregated tau protein, particularly tau protein (e.g., a compound that incorporates, is conjugated to, is chelated with, or is otherwise associated with, one or more (e.g., 1, 2, 3, 4, etc.) detectable labels), in a method of manufacture of a diagnostic or prognostic reagent for use in the diagnosis or prognosis of a tau proteinopathy. The compound may be provided as a composition as described herein.

Those skilled in the art will appreciate that instead of administering ligands/labels directly, they could be administered in a precursor form, for conversion to the active form (e.g., ligating form, labelling form) by an activating agent present in, or administered to, the same subject.

The ligands disclosed herein may be used as part of a method of diagnosis or prognosis. It may be used to select a patient for treatment, or to assess the effectiveness of a treatment or a therapeutic (e.g., an inhibitor of tau protein aggregation) administered to the subject.

Treatment

The term “treatment,” as used herein in the context of treating a condition, pertains generally to treatment and therapy, whether of a human or an animal (e.g., in veterinary applications), in which some desired therapeutic effect is achieved, for example, the inhibition of the progress of the condition, and includes a reduction in the rate of progress, a halt in the rate of progress, regression of the condition, amelioration of the condition, and cure of the condition. Treatment as a prophylactic measure (i.e., prophylaxis, prevention) is also included.

The term “therapeutically-effective amount,” as used herein, pertains to that amount of a compound of the invention, or a material, composition or dosage from comprising said compound, which is effective for producing some desired therapeutic effect, commensurate with a reasonable benefit/risk ratio, when administered in accordance with a desired treatment regimen.

Similarly, the term “prophylactically effective amount,” as used herein, pertains to that amount of a compound of the invention, or a material, composition or dosage from comprising said compound, which is effective for producing some desired prophylactic effect, commensurate with a reasonable benefit/risk ratio, when administered in accordance with a desired treatment regimen.

“Prophylaxis” in the context of the present specification should not be understood to circumscribe complete success i.e. complete protection or complete prevention. Rather prophylaxis in the present context refers to a measure which is administered in advance of detection of a symptomatic condition with the aim of preserving health by helping to delay, mitigate or avoid that particular condition.

The term “treatment” includes combination treatments and therapies, in which two or more treatments or therapies are combined, for example, sequentially or simultaneously. Examples of treatments and therapies include, but are not limited to, chemotherapy (the administration of active agents, including, e.g., drugs, antibodies (e.g., as in immunotherapy), prodrugs (e.g., as in photodynamic therapy, GDEPT, ADEPT, etc.); surgery; radiation therapy; and gene therapy.

For example, it may be beneficial to combine treatment with a compound as described herein with one or more other (e.g., 1, 2, 3, 4) agents or therapies.

The particular combination would be at the discretion of the physician who would select dosages using his/her common general knowledge and dosing regimens known to a skilled practitioner.

The agents (i.e., a compound as described herein, plus one or more other agents) may be administered simultaneously or sequentially, and may be administered in individually varying

dose schedules and via different routes. For example, when administered sequentially, the agents can be administered at closely spaced intervals (e.g., over a period of 5-10 minutes) or at longer intervals (e.g., 1, 2, 3, 4 or more hours apart, or even longer periods apart where required), the precise dosage regimen being commensurate with the properties of the therapeutic agent(s).

The agents (i.e., a compound as described here, plus one or more other agents) may be formulated together in a single dosage form, or alternatively, the individual agents may be formulated separately and presented together in the form of a kit, optionally with instructions for their use.

Routes of Administration

The compound of the invention, or pharmaceutical composition comprising it, may be administered to a subject/patient by any convenient route of administration, whether systemically/peripherally or topically (i.e., at the site of desired action).

Routes of administration include, but are not limited to, oral (e.g., by ingestion); buccal; sublingual; transdermal (including, e.g., by a patch, plaster, etc.); transmucosal (including, e.g., by a patch, plaster, etc.); intranasal (e.g., by nasal spray); ocular (e.g., by eyedrops); pulmonary (e.g., by inhalation or insufflation therapy using, e.g., via an aerosol, e.g., through the mouth or nose); rectal (e.g., by suppository or enema); vaginal (e.g., by pessary); parenteral, for example, by injection, including subcutaneous, intradermal, intramuscular, intravenous, intraarterial, intracardiac, intrathecal, intraspinal, intracapsular, subcapsular, intraorbital, intraperitoneal, intratracheal, subcuticular, intraarticular, subarachnoid, and intrasternal (including, e.g., intracatheter injection into the brain); by implant of a depot or reservoir, for example, subcutaneously or intramuscularly.

Preferred compositions are oral compositions, formulated as described in more detail hereinafter.

The Subject/Patient

The subject/patient may be an animal, a mammal, a placental mammal, a rodent (e.g., a guinea pig, a hamster, a rat, a mouse), murine (e.g., a mouse), a lagomorph (e.g., a rabbit), avian (e.g., a bird), canine (e.g., a dog), feline (e.g., a cat), equine (e.g., a horse), porcine (e.g., a pig), ovine (e.g., a sheep), bovine (e.g., a cow), a primate, simian (e.g., a monkey or ape), a monkey (e.g., marmoset, baboon), a monotreme (e.g. platypus), an ape (e.g., gorilla, chimpanzee, orangutang, gibbon), or a human.

Furthermore, the subject/patient may be any of its forms of development, for example, a foetus.

In some embodiments, the subject/patient is a human.

Compositions/Formulations

While it is possible for the compound of the invention to be used (e.g., administered) alone, it is often preferable to present it as a composition or formulation.

Another aspect of the invention therefore provides a composition comprising a compound as described herein, and a pharmaceutically acceptable carrier or diluent.

In some embodiments, the composition is a pharmaceutical composition (e.g., formulation, preparation, medicament) comprising a compound as described herein, and a pharmaceutically acceptable carrier, diluent, or excipient.

In some embodiments, the composition is a pharmaceutical composition comprising at least one compound, as described herein, together with one or more other pharmaceutically acceptable ingredients well known to those skilled in the art, including, but not limited to, pharmaceutically acceptable carriers, diluents, excipients, adjuvants, fillers, buffers, preservatives, anti-oxidants, lubricants, stabilisers,

solubilisers, surfactants (e.g., wetting agents), masking agents, colouring agents, flavouring agents, and sweetening agents.

In some embodiments, the composition further comprises other active agents, for example, other therapeutic or prophylactic agents.

Suitable carriers, diluents, excipients, etc. can be found in standard pharmaceutical texts. See, for example, *Handbook of Pharmaceutical Additives*, 2nd Edition (eds. M. Ash and I. Ash), 2001 (Synapse Information Resources, Inc., Endicott, N.Y., USA), *Remington's Pharmaceutical Sciences*, 20th edition, pub. Lippincott, Williams & Wilkins, 2000; and *Handbook of Pharmaceutical Excipients*, 2nd edition, 1994.

Another aspect of the present invention pertains to methods of making a pharmaceutical composition comprising admixing at least one [¹¹C]-radiolabelled compound, as defined herein, together with one or more other pharmaceutically acceptable ingredients well known to those skilled in the art, e.g., carriers, diluents, excipients, etc. If formulated as discrete units (e.g., tablets, etc.), each unit contains a predetermined amount (dosage) of the compound.

The term "pharmaceutically acceptable," as used herein, pertains to compounds, ingredients, materials, compositions, dosage forms, etc., which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of the subject in question (e.g., human) without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio. Each carrier, diluent, excipient, etc. must also be "acceptable" in the sense of being compatible with the other ingredients of the formulation.

The formulations may be prepared by any methods well known in the art of pharmacy. Such methods include the step of bringing into association the compound with a carrier which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association the compound with carriers (e.g., liquid carriers, finely divided solid carrier, etc.), and then shaping the product, if necessary.

The formulation may be prepared to provide for rapid or slow release; immediate, delayed, timed, or sustained release; or a combination thereof.


Formulations suitable for parenteral administration (e.g., by injection), include aqueous or non-aqueous, isotonic, pyrogen-free, sterile liquids (e.g., solutions, suspensions), in which the compound is dissolved, suspended, or otherwise provided (e.g., in a liposome or other microparticulate). Such liquids may additionally contain other pharmaceutically acceptable ingredients, such as anti-oxidants, buffers, preservatives, stabilisers, bacteriostats, suspending agents, thickening agents, and solutes which render the formulation isotonic with the blood (or other relevant bodily fluid) of the intended recipient. Examples of excipients include, for example, water, alcohols, polyols, glycerol, vegetable oils, and the like. Examples of suitable isotonic carriers for use in such formulations include Sodium Chloride Injection, Ringer's Solution, or Lactated Ringer's Injection. Typically, the concentration of the compound in the liquid is from about 1 ng/ml to about 10 µg/ml, for example from about 10 ng/ml to about 1 µg/ml. The formulations may be presented in unit-dose or multi-dose sealed containers, for example, ampoules and vials, and may be stored in a freeze-dried (lyophilised) condition requiring only the addition of the sterile liquid carrier, for example water for injections, immediately prior to use.

47

Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules, and tablets.

EXAMPLES OF SOME PREFERRED FORMULATIONS

One aspect of the present invention pertains to a dosage unit (e.g., a pharmaceutical tablet or capsule) comprising 20



48

Processes generally used for tablet formulation and film coating often require the use of heat accompanied by low humidity during the drying process.

5 LMTM and the other leuco-methylthionium salts are potentially prone to oxidation to methylthionium moiety (MT) and to degradation e.g. to L Azure B (LAB) (see Scheme, below):

to 300 mg of a compound as described herein (e.g., obtained by, or obtainable by, a method as described herein; having a purity as described herein; etc.), and a pharmaceutically acceptable carrier, diluent, or excipient.

In some embodiments, the dosage unit is a tablet.

In some embodiments, the dosage unit is a capsule.

In some embodiments, said capsules are gelatine capsules.

In some embodiments, said capsules are HPMC (hydroxypropylmethylcellulose) capsules.

In some embodiments, the amount is 30 to 200 mg.

In some embodiments, the amount is about 30 mg.

In some embodiments, the amount is about 60 mg.

In some embodiments, the amount is about 100 mg.

In some embodiments, the amount is about 150 mg.

In some embodiments, the amount is about 200 mg.

Throughout the present specification dosage amounts, e.g. as set out above, may refer to the amount of the compound itself or may refer to the amount of free base equivalent (i.e. the amount of LMT moiety) contained in the dosage unit. Both these alternatives are expressly disclosed by the present invention.

In some embodiments, the pharmaceutically acceptable carrier, diluent, or excipient is or comprises one or both of a glyceride (e.g., Gelucire 44/14®; lauroyl macrogol-32 glycerides PhEur, USP) and colloidal silicon dioxide (e.g., 2% Aerosil 200®; Colloidal Silicon Dioxide PhEur, USP).

Novel Formulations—Solid Dosage Forms

40 For a material such as e.g. LMTM, which is prone to oxidation (as explained above), conventional formulation processes may therefore lead to degradation and hence, potentially, to instability in the performance of the product.

45 The principle behind the formulations of the present invention is therefore the provision of a method of manufacture of compressed pharmaceutical formulations and capsules containing leuco-methylthionium salts e.g. bis(methanesulfonate) (LMTM) as the active substance, by direct tablet compression technology or by other unique tableting techniques, and by encapsulation, in which the active substance exists substantially in a stable form.

50 The most commonly used method for the preparation of solid dosage forms is wet granulation (also called moist granulation). This involves adding a granulating fluid to a powder. The granulating fluid may be water or some other solvent that is sufficiently volatile that can subsequently be removed by drying. The granulating fluid may also include a binder. Once the solvent has been removed, the resulting mass is milled.

55 60 Wet granulation is often preferred over direct compression because wet granulation is more likely to overcome any problems associated with the physical characteristics of various ingredients in the formulation. Wet granulation provides material which has the required flow and cohesive properties necessary to obtain an acceptable solid dosage form. The content uniformity of the solid dosage form is generally improved with wet granulation because all of the granules

49

usually contain the same amount of drug. Segregation of the drug from excipients is also avoided.

In direct compression, the individual constituents of the composition to be compressed are mixed without previous granulation and then directly compressed. Whilst this appears to be an elegant and simple process, it may be difficult to obtain with it commercially usable tablets which have sufficient strength yet which also disintegrate sufficiently rapidly after administration. Also, many active substances cannot be processed by direct compression since they cannot be compressed without a granulation step.

It has now, surprisingly, been found that compounds of the present invention are stable in a dry compressed solid dosage form such as a tablet, during manufacture and storage, and that the amount of degradation products such as L Azure B (LAB) and methylthioninium (MT) formed can be controlled within the specifications (for example, LAB less than 2% and MT less than 12%).


This is in contrast to the behaviour of e.g. LMTM when processed by conventional wet granulation processes. Without wishing to be bound by theory, in conventional wet granulation processes LMTM, for instance, may be very unstable and a substantial amount of LAB and MT may be formed.

Accordingly, one aspect of the present invention provides a pharmaceutical composition comprising a compound of the invention, in solid dosage form. The composition preferably further comprises at least one diluent suitable for dry compression. The pharmaceutical composition is characterised in that the compound exists in a substantially stable form.

Another aspect of the invention provides a free-flowing, cohesive powder, comprising a compound of the invention and at least one diluent suitable for dry compression, and optionally one or more other excipients, said powder being capable of being compressed into a solid dosage form.

These compositions and formulations are initially described herein with respect to the bis(sulfonate) salts of the present invention, in particular LMTM. However, the advantages of the present formulation methods are equally applicable to other members of the leuco-methylthioninium family of salts

For example, the formulations described herein are applicable also to the 3,7-diamino-10H-phenothiazinium salts disclosed in WO2007/110627 (WisTa Laboratories Ltd), which were briefly discussed above. These include leuco-methylthioninium bis(hydrobromide) (LMT.2HBr, LMTB) and leuco-methylthioninium bis(hydrochloride) (LMT.2HCl, LMTC). Therefore, in a broader aspect, the present invention provides a pharmaceutical composition comprising a compound of the following formula I:



wherein:

R^1 , R^9 , R^{3NA} , R^{3NB} , R^{7NA} and R^{7NB} are as previously defined;


and wherein each of HX^1 and HX^2 is independently a protic acid;

50

or a pharmaceutically acceptable salt, solvate, or hydrate thereof;

in a solid dosage form as described herein.

For completeness, it is noted that, as would be understood by one skilled in the art, the above formula could equally be written as:



wherein X^1 and X^2 are the corresponding counterions.

Preferably X^1 and X^2 are independently sulfonate (such as alkylsulfonate or arylsulfonate, for example R^ASO_3^- or R^BSO_3^- as defined above) or halide (Cl^- , Br^- , I^-). In other words, HX^1 and HX^2 are preferably independently sulfonic acids ($\text{R}^A\text{SO}_3\text{H}$, $\text{R}^B\text{SO}_3\text{H}$) or hydrohalides (HCl , HBr , HI).

As used hereafter, the term ‘active ingredient’ refers to the relevant leuco(methylthioninium) salt. In other words it refers to a compound of formula (I), such as a compound of the invention, for example LMTM.

Another aspect of the invention provides a process for the manufacture of said pharmaceutical compositions, by a dry compression method. The process preferably comprises dry compression of an intimate powder mixture of the active compound with at least one diluent suitable for dry compression, and optionally one or more other excipients.

In some embodiments, the process comprises direct compression.

In some embodiments, the process comprises simple direct compression.

In some embodiments, the process comprises dry granulation.

In some embodiments, the process comprises moist granulation of excipients, followed by addition of the active ingredient extra-granularly.

Solid dosage forms according to the invention advantageously exhibit long-term chemical and physical stability of the active ingredient (compound of the invention—e.g. LMTM). The pharmaceutical compositions according to the invention also have fast dissolution rates, even after long-term storage.

A “substantially stable” form of the active ingredient means, in the present context, a form which does not react to form impurities such as oxidative impurities or other degradation products to any significant extent during the formulation process, or on storage of the formulated product.

Therefore, in the present context, it may refer to a material which contains, for example, less than 20% w/w, less than 15% w/w, or less than 10% w/w of oxidative impurities or other degradation products. In other words the material contains at least 80% w/w, at least 85% w/w, or at least 90% w/w of the pure active ingredient, in its original (unreacted) form.

In some embodiments, the material containing the active ingredient may contain, for example, less than 20% w/w, less than 15% w/w, less than 12% w/w, or less than 10% w/w of MT. In some embodiments, the material may contain, for example, less than 5% w/w, less than 3% w/w, or less than 2% w/w of LAB.

A "stable" tablet is, in the context of the present invention, a tablet that remains substantially stable after prolonged storage under controlled conditions of temperature and humidity. Stability testing may be carried out with the solid dosage forms directly exposed to the chosen environmental conditions, or with the solid dosage forms contained within packaging.

Content of Active Ingredient

The amount of the active ingredient in the uncoated composition is generally more than about 10% w/w, but can be more than 20%, or more than 30% w/w. The amount of the active ingredient is generally less than about 70% w/w, and usually less than 60% or less than 50% w/w in a tablet formulation. Typically, the amount of the active ingredient in the uncoated tablet core composition is thus from about 10% w/w (or 20% or 30%) to about 70% w/w (or 60% or 50%). Where a coating is applied to the composition, as described below, the overall weight of the composition is increased and thus the percentage of the active ingredient in the overall composition is somewhat reduced.

Diluents

The active ingredient may not be inherently compressible and thus may require addition of suitable diluents to aid compression.

The pharmaceutical compositions of the invention therefore commonly comprise at least 15% w/w, more commonly at least 20%, at least 30%, at least 40% or at least 50% w/w of diluent(s).

Diluents that may be used include one or more of microcrystalline cellulose, lactose, mannitol, calcium salts such as calcium phosphate dibasic, calcium sulphate and calcium carbonate, and sugars such as lactose, sucrose, dextrose and maltodextrin.

Preferred diluents are microcrystalline cellulose, lactose and mannitol. Spray-dried forms of lactose and mannitol are particularly suitable forms of those compounds for direct compression or dry granulation techniques.

It has unexpectedly been found that when an active ingredient described herein, for example a compound of the present invention, such as LMTM, is formulated with dry compression diluents such as one or more of microcrystalline cellulose, spray dried lactose, anhydrous lactose and mannitol, the resulting solid dosage forms are stable in the sense that the active ingredient remains chemically stable, even after extended storage.

The invention thus provides a method of preparing low, medium- or high-dose tablets, for example low, medium-, or high-dose LMTM tablets, that are stable and have good dissolution profiles, acceptable degrees of hardness and resistance to chipping, as well as a short disintegration time.

Dissolution of Compositions of the Invention

The present inventors have also surprisingly found that the unique solid dosage forms described herein provide a very fast dissolution rate.

As explained hereinbefore, and without wishing to be bound by theory, it is thought that the active methylthioninium (MT) moiety may preferably be absorbed from the stomach and/or the upper GI tract. A fast-disintegrating and fast-dissolving formulation of the leuco(methylthioninium) salts would therefore be advantageous, since this would deliver the maximum possible amount of drug to the intended point of absorption.

The fast dissolution rate of the solid dosage forms described herein means that they are capable of dissolving rapidly in the stomach and/or upper GI tract and hence presenting the active ingredient there effectively, for rapid absorption.

In some embodiments, the formulations of the invention, when evaluated using a standard pharmacopeial method, provide at least 80% dissolution within 30 minutes, preferably at least 80% dissolution within 15 minutes, more preferably at least 80% dissolution within 10 minutes.

In some embodiments, the formulations of the invention, when evaluated using a standard pharmacopeial method, provide at least 90% dissolution within 30 minutes, preferably at least 90% dissolution within 15 minutes, more preferably at least 90% dissolution within 10 minutes.

In some embodiments, the formulations of the invention, when evaluated using a standard pharmacopeial method provide at least 95% dissolution within 30 minutes, preferably at least 95% dissolution within 15 minutes, more preferably at least 95% dissolution within 10 minutes.

Dissolution rates may be measured by standard pharmacopeial methods as described in United States Pharmacopeia (USP) General Chapter <711>. The current USP is USP 34 (2011). For example, dissolution rates for the formulations of the invention may be measured using apparatus according to USP Dissolution Apparatus 2 (Paddle).

In some embodiments, the dissolution rates above are evaluated in 0.1M hydrochloric acid at a working concentration of ~5 µg/ml LMT, with stirring at 50 rpm paddle speed. In some embodiments, the dissolution rates are evaluated by spectrophotometric analysis. In some embodiments, analysis comprises UV/vis spectrophotometry ($\lambda_{max\ LMT}=255\text{ nm}$).

As a consequence of their surprisingly high dissolution rate, the formulation methods described herein can provide the active compound with a high degree of bioavailability.

The fast dissolution rate is maintained after prolonged storage, even if storage is under 'stressed' conditions (i.e. increased temperature and humidity). The fast dissolution rate, and hence the good bioavailability, of compositions formulated according to the processes of the present invention is also highly tolerant of variations in the formulation itself.

Other Ingredients

The pharmaceutical composition will generally also include a lubricant. Examples of lubricants include magnesium stearate, calcium stearate, sodium stearyl fumarate, stearic acid, glyceryl behaptate, polyethylene glycol, ethylene oxide polymers (for example, those available under the registered trademark Carbowax from Union Carbide, Inc., Danbury, Conn.), sodium lauryl sulphate, magnesium lauryl stearate, mixtures of magnesium stearate with sodium lauryl sulphate, and hydrogenated vegetable oil. Preferred lubricants include calcium stearate, magnesium stearate and sodium stearyl fumarate. Most preferred as the lubricant is magnesium stearate. Lubricants generally comprise from about 0.5 to about 5.0% of the total (uncoated) tablet weight. The amount of lubricant employed is generally from about 1.0 to about 2.0%, preferably 0.5 to 2.0% w/w.

In addition to the diluent(s) and lubricant(s), other conventional excipients may also be present in the pharmaceutical compositions of the invention. Such additional excipients include disintegrants, binders, flavouring agents, colours and glidants. Some excipients can serve multiple functions, for example as both binder and tablet disintegrant.

A tablet disintegrant may be present in an amount necessary to achieve rapid dissolution. Disintegrants are excipients which oppose the physical forces of particle bonding in a tablet or capsule when the dosage form is placed in an aqueous environment. Examples of disintegrants include crosslinked polyvinylpyrrolidone (crospovidone), sodium starch glycolate, crosslinked sodium carboxymethyl cellulose (sodium croscarmellose), and pregelatinized starch. Generally the amount of disintegrant can be from 0 to about

25% w/w, more commonly from about 1% to about 15% w/w, and usually less than 10% or less than 5% w/w, of the composition.

Binders are excipients which contribute to particle adhesion in a solid formulation. Examples of binders include cellulose derivatives (carboxymethylcellulose, hydroxypropyl methylcellulose, hydroxypropyl cellulose, hydroxyethylcellulose, ethylcellulose, microcrystalline cellulose) and sugars such as lactose, sucrose, dextrose, glucose, maltodextrin, and mannitol, xylitol, polymethacrylates, polyvinylpyrrolidone, sorbitol, pregelatinized starch, alginic acids, and salts thereof such as sodium alginate, magnesium aluminum silicate, polyethylene glycol, carrageenan and the like. Generally, the amount of binder can vary widely, eg from 0% to 95% w/w of the composition. As noted above, excipients may serve multiple functions. For instance, the tableting diluent may also serve as a binder.

Glidants are substances added to a powder to improve its flowability. Examples of glidants include magnesium stearate, colloidal silicon dioxide (such as the grades sold as Aerosil), starch and talc. Glidants may be present in the pharmaceutical composition at a level of from 0 to about 5% w/w. Again, however, it should be noted that excipients may serve multiple functions. The lubricant, for example magnesium stearate, may also function as a glidant.

Examples of colours that may be incorporated into the pharmaceutical compositions of the invention include titanium dioxide and/or dyes suitable for food such as those known as FD&C dyes and natural colouring agents. A colouring agent is unlikely to be used in the powder mixture that is compressed in accordance with the aspects of the invention discussed above, but may form part of a coating applied to the composition, as described below, in which case the colouring agent may be present in the film coat in an amount up to about 2.0% w/w.

The tablet is desirably coated with a conventional film coating which imparts toughness, ease of swallowing, and an elegant appearance to the final product. Many polymeric film-coating materials are known in the art. A preferred film-coating material is hydroxypropylmethylcellulose (HPMC) or polyvinyl alcohol-part hydrolysed (PVA). HPMC and PVA may be obtained commercially, for example from Colorcon, in coating formulations containing excipients which serve as coating aids, under the registered trademark Opadry. Opadry formulations may also contain talc, polydextrose, triacetin, polyethyleneglycol, polysorbate 80, titanium dioxide, and one or more dyes or lakes. Other suitable film-forming polymers may also be used, including hydroxypropylcellulose, vinyl copolymers such as polyvinyl pyrrolidone and polyvinyl acetate, and acrylate-methacrylate copolymers. Use of a film coating is beneficial for ease of handling and because a blue coloured uncoated core may stain the inside of the mouth during swallowing. Coating also improves light stability of the dosage form.

Coating of the tablets may conveniently be carried out using a conventional coating pan. In preferred embodiments of the process, the coating pan is pre-heated using heated inlet air until the exhaust temperature reaches 35°-55° C., more preferably 40-50° C. This may typically require application of heated inlet air at an inlet temperature of 45-75° C., preferably 50-65° C., for 10-15 minutes. The tablet cores containing the active ingredient (e.g. LMTM) are then added to the coating pan and the aqueous film coat applied. The spray rate is controlled such that the bed temperature is maintained at 38-48° C., more preferably 42-44° C., until the desired weight gain (coating weight) has been achieved.

Dry Compression Methods

'Dry compression', as used herein, refers to compression techniques which do not involve the use of heat or moisture. Dry compression may comprise direct compression of the active ingredient with suitable diluents or it may comprise dry granulation (for example slugging/double compression method or roller compaction).

Direct compression may comprise simple direct compression of the active ingredient with diluents suitable for direct compression. Alternatively it may comprise granulation, for example moist granulation, of the excipients to produce a dry granular excipient mixture which can then be directly compressed with the dry active ingredient (and optionally further dry excipients). This may be referred to as 'extra-granular incorporation' of the active ingredient.

Accordingly, in some embodiments the solid dosage forms of the invention may be produced in a manufacturing process which comprises simple direct compression. In this embodiment, the tablet ingredients, i.e. the active ingredient (e.g. LMTM), diluent(s) and other optional excipients, are blended together in solid particulate form to create an intimate mixture, e.g. in a tumbling blender, and are then compressed using a tablet machine.

In other embodiments, the composition is prepared by a dry granulation process. Dry granulation refers to the process of granulating without the use of granulating fluids. In order for a material to be dry-granulated, at least one of its constituents, either the active ingredient or a diluent, must have cohesive properties. Dry granulation may be performed by a process known as "slugging". In "slugging", the material to be granulated is first made into a large compressed mass or "slug", typically using a tablet press with large flat-faced tooling (an example of a linear press is illustrated in U.S. Pat. No. 4,880, 373). A fairly dense slug may be formed by allowing sufficient time for the air to escape from the material to be compacted. Compressed slugs are then milled through a desired mesh screen manually or automatically as, for example, by way of a comminuting mill. Formation of granules by "slugging" is also known as precompression. When tablets are made from the granulated slugged material, the process is referred to as the "double compression method".

Dry granulation may also be performed using a "roller compactor". In a roller compactor, material particles are consolidated and densified by passing the material between two high-pressure rollers. The densified material from a roller compactor is then reduced to a uniform granule size by milling. The uniform granules may then be mixed with other substances, such as a lubricant, to tablet the material (as, for example, by way of a rotary tableting machine). In addition to pharmaceutical use, roller compaction is used in other industries, such as the food industry, animal feed industry and fertilizer industry.

Dry granulation is nowadays generally understood to mean roller compaction or slugging, and is well known to those skilled in the art (see, for instance, *Pharmaceutical Dosage Forms: Tablets* (Lieberman, Lachman, and Schwartz (Eds); Marcel Dekker, Inc, 2nd Edition, 1989) and *Remington's Pharmaceutical Sciences* (A. R. Gennaro (Ed); Mack Publishing Co, Easton, Pa., 18th edition, 1990)).

In further embodiments of the invention, tablets are prepared by moist granulation of excipients and incorporation of the active ingredient (e.g. LMTM) extra-granularly. Typically, such a process involves wet massing diluents such as lactose and/or microcrystalline cellulose with water, optionally with the addition of a binder such as polyvinyl pyrrolidone. The wet mass is dried, then passed through a mesh, to form granules. The active ingredient and any remaining

excipients, such as a lubricant, are then blended with the dry granules and compressed to form tablets.

Use of Acids in the Compositions of the Invention

Compositions containing leuco(methylthionium) compounds, including compounds of the invention such as LMTM may, in some embodiments, be stabilised by addition of an appropriate amount of certain acids to the bulk substance prior to formulation. These acids may be used to prevent the formation of further MT, both during formulation and throughout the life of the product, thereby providing a stable pharmaceutical composition for the purposes of obtaining regulatory approval with associated cost savings in packaging.

According to the present invention, therefore, there is also provided a pharmaceutical composition comprising an active ingredient as described herein and a pharmaceutically acceptable carrier, characterised in that said formulation additionally comprises an acid in an amount sufficient to prevent the formation of MT. Without wishing to be bound by theory, it is thought that acids having a pK1 of greater than 1.5 are preferred. In some embodiments, the acid is present in an amount of from 5% to 25% w/w.

Preferably the composition is prepared by a dry compression method as described above.

Preferred acids for the purposes of the invention are maleic acid (pK1 1.9), phosphoric acid (pK1 2.12), ascorbic acid (pK1 4.17), sorbic acid (pK1 4.76), aspartic acid, and sialic acid. The stabilising effect of the added acid may be enhanced by the selection of an appropriate carrier. The carrier is preferably mannitol, a cellulosic material, or a starch, or mixtures thereof. The carrier is typically present in an amount of at least 40% w/w of the formulation.

Particle Size

It has also been found that a significant reduction in the formation of MT can also be achieved by the selection of an appropriate particle size range for the dry powder blend, typically wherein more than 10% of the particles have a size greater than 10 microns. Therefore according to another aspect of the invention, there is provided a pharmaceutical composition comprising an active ingredient as described herein and a pharmaceutically acceptable carrier, additionally characterised in that said composition comprises particles of which more than 10% have a size greater than 10 microns.

Carriers

It has been found that a significant reduction in the formation of MT can also be achieved by the choice of an appropriate carrier, particularly one having a particle shape which is averse to the entry of water. Elcema TM, for example, which has long, lamellar particles which are smooth and flat in shape with a non-porous surface, appears to reduce MT formation by limiting the access of water. Ethylcellulose, mannitol and Starch 1500 TM and Microcrystalline cellulose are also particularly suitable for this purpose.

Therefore according to another aspect of the invention, there is provided a pharmaceutical composition comprising a leuco(methylthionium) compound, for example a compound of the invention such as LMTM, and a pharmaceutically acceptable carrier, characterised in that said carrier is Elcema TM, ethylcellulose, mannitol, or Starch 1500 TM.

Encapsulation

Stabilised dry powder blends in accordance with the invention may be formulated, for example, by compressing into tablets or filling into capsules (with or without prior conversion to a granulated powder by means such as described in Formulation Examples 1 to 4) to give pharmaceutical compositions having excellent shelf life.

Capsules according to the invention are typically of gelatin or preferably HPMC. Preferred excipients include lactose, starch, a cellulose, milk sugar and high MW polyethylene glycols.

Conclusions

Pharmaceutical compositions and formulations prepared according to the methods described above are more stable, immediately after completion of manufacture, than formulations produced using conventional aqueous granulation. Furthermore they can demonstrate enhanced stability on storage.

For example, a pharmaceutical formulation thus prepared, with a content of 10 to 50% by weight of LMTM, preferably 15% to 40% by weight of LMTM, makes it possible that in standard stability tests, for example in long term accelerated stability testing, at a temperature of 25° C. and a relative humidity 60±5% the content of L Azure B does not increase by more than 2%, relative to LMTM peak area, within a period of 24 months.

During processing and on storage leuco(methylthionium) compounds, such as LMTM may also oxidise to produce a small amount of MT (see Scheme, above).

The presence of relatively small concentrations (e.g. less than 12%) of MT in the leuco-formulations of the present invention, although undesirable, is not considered to have adverse clinical significance per se as even if the body is presented with MT in its charged or oxidized form from LMTM and the various other leuco salts, this can then be reduced to the uncharged (reduced) MT form prior to absorption. In addition to the small amount of MT formed during processing such as blending and tableting, leuco-methylthionium salts of the present invention may react with oxygen absorbed on the excipients and present within the tablet to give more of the MT particularly in the presence of moisture.

One advantage of the formulations of this invention is to minimise the amount of MT formed in the tablets, e.g. to less than 12% over 2 years when stored at 25° C. at a relative humidity of 60%. This refers to the cumulative amount of MT formed during both processing and storage of the tablets: generally, the formulation methods of the invention result in less than 5% MT formation during processing; a maximum of around 5-7% MT is then formed during storage of the finished pack. This provides a shelf-life of at least 24 months.

This is demonstrated in the Formulation Examples, below.

Dosage

It will be appreciated by one of skill in the art that appropriate dosages of the compound, and compositions comprising the compound, can vary from patient to patient. Determining the optimal dosage will generally involve the balancing of the level of therapeutic benefit against any risk or deleterious side effects. The selected dosage level will depend on a variety of factors including, but not limited to, the activity of the particular compound, the route of administration, the time of administration, the rate of excretion of the compound, the duration of the treatment, other drugs, compounds, and/or materials used in combination, the severity of the condition, and the species, sex, age, weight, condition, general health, and prior medical history of the patient. The amount of compound and route of administration will ultimately be at the discretion of the physician, veterinarian, or clinician, although generally the dosage will be selected to achieve local concentrations at the site of action which achieve the desired effect without causing substantial harmful or deleterious side-effects.

Administration can be effected in one dose, continuously or intermittently (e.g., in divided doses at appropriate intervals) throughout the course of treatment. Methods of determining the most effective means and dosage of administra-

57

tion are well known to those of skill in the art and will vary with the formulation used for therapy, the purpose of the therapy, the target cell(s) being treated, and the subject being treated. Single or multiple administrations can be carried out with the dose level and pattern being selected by the treating physician, veterinarian, or clinician.

In general, a suitable dose of the compound is in the range of about 100 ng to about 25 mg (more typically about 1 μ g to about 10 mg) per kilogram body weight of the subject per day.

In some embodiments, the compound is administered to a human patient according to the following dosage regime: about 100 mg, 3 times daily.

In some embodiments, the compound is administered to a human patient according to the following dosage regime: about 150 mg, 2 times daily.

In some embodiments, the compound is administered to a human patient according to the following dosage regime: about 200 mg, 2 times daily.


EXAMPLES

The following examples are provided solely to illustrate the present invention and are not intended to limit its scope.

Example 1

Synthesis and Characterisation

Laboratory synthesis of 10-Acetyl-N,N,N',N'-tetramethylphenothiazine-3,7-diamine



step i; NaNO₂, DMSO, CH₃COOH, step ii; (H₃CCO)₂O, Et₃N, DMF, step iii; Pd/C, 2-MeTHF, H₂ step iv; H₂CO, H₂ or step iii; Zn, (aq) NH₄Cl, MeOH, THF step iv; H₂CO, NaCNBH₃, CH₃COOH

Synthesis of 3,7-dinitro-10H-phenothiazine (2)

To a 3 necked 1 litre round bottom flask (RBF) fitted with a thermometer, dropping funnel and a condenser was added phenothiazine (MW 199.28 g/mol, 25.00 g, 125.5 mmol) and dimethylsulfoxide (250 ml) the mixture was stirred for 2 minutes or until the phenothiazine dissolved. The condenser was then connected to a Dreschel bottle half filled with water. Sodium nitrite (MW 69.00 g/mol, 51.94 g, 752.7 mmol) was then added to the RBF and acetic acid (150 ml) was added to the dropping funnel. The acetic acid was then added to the

58

RBF in a drop-wise fashion over a 20 minute period. The light yellow slurry becomes red in colour and a solid precipitated out of solution. Upon completion of the acetic acid addition the mixture was stirred for 2 hours at ambient temperature (36-20° C.) before increasing the temperature to 95° C. and stirring for 17 hours. After this time the mixture was cooled to 50° C. and methanol (100 ml) was added and the mixture cooled further to 22° C. The cooled mixture was then filtered and the cake washed with methanol (3x25 ml). The washed cake was left on the filter with the vacuum applied for 30 minutes before being dried for 15 hours at 50° C. to give the product as a brown solid (MW 289.27 g/mol, 29.45 g, 81%).

Notes

1. The addition of acetic acid produced NO_x gases, which was converted to nitric acid by allowing the gas to bubble into a Dreschel bottle half filled with water.
2. The addition of acetic acid is exothermic and the mixture rises from 22° C. to 36° C.
3. Methanol was added to help dissolve any sodium acetate and as an anti-solvent to maximise the product yield.
4. The synthesis was also successful using dimethylformamide (DMF), acetonitrile (MeCN), tetrahydrofuran (THF), acetone or dimethoxyethane (DME) as the reaction solvent.

NMR: The product (5 mg) was dissolved in DMSO-d₆ (1.5 ml) and may require to be warmed to completely dissolve the solid.

δ_H (400 MHz; DMSO-d₆): 6.72 (2H, d, J 8.8, ArH), 7.77 (2H, d, J 2.8, ArH), 7.87 (2H, dd, J 2.8, 8.8, ArH)

Synthesis of 3,7-dinitro-10-acetylphenothiazine (3)

To a 3 necked 500 ml round bottom flask fitted with a thermometer and a condenser was added 3,7-dinitro-10H-phenothiazine (MW 289.27 g/mol, 29.00 g, 100 mmol), dimethylformamide (58 ml), acetic anhydride (MW 102.09 g/mol, 102.09 g, 1000 mmol) and triethylamine (MW 101.19 g/mol, 40.88 g, 401 mmol). The mixture was heated to 105° C. and stirred at this temperature for 3 hours. The mixture was cooled to ambient temperature (21° C.) before being cooled to 5° C. whereby it was stirred for 1 hour. The product was

59

isolated by filtration and washed with methanol (3×30 ml) to give a light yellow crystalline solid, which was dried at 50° C. for 15 hours (MW 331.31 g/mol, 26.94 g, 81%).

Notes

1. Crystals of the product form during the reaction, after ~1 hour at 105° C.
2. Upon cooling the bulk of the product precipitates at ~70° C.
3. Product was orange in colour before it was washed with methanol.

NMR: The product (10 mg) was dissolved in DMSO-d₆ (1.5 ml).

δ_H (400 MHz; DMSO-d₆): 2.25 (3H, s, CH₃), 7.92 (2H, d, J 8.8, ArH), 8.28 (2H, dd, J 8.8, 2, ArH), 8.47 (2H, d, J 2, ArH)

Synthesis of 10-Acetyl-N,N,N',N'-tetramethylphenothiazine-3,7-diamine (5)

To a 3 necked 100 ml round bottom flask fitted with a thermometer and a condenser was added 3,7-dinitro-10-acetylphenothiazine (MW 331.31 g/mol, 5 g, 15.09 mmol), palladium on carbon (10%, dry, 0.5 g) and 2-methyltetrahydrofuran (25 ml). The flask was evacuated and purged with hydrogen 5 times before the mixture was heated to 56° C. After 17 hours the reduction was judged to have reached completion (see tlc conditions) giving compound 4, and formalin was added (MW 30.03 g/mol, 14.7 g, 181.1 mmol). The flask was once again evacuated and purged 5 times with hydrogen. After 71 hours from the addition of formalin (total time 88 hours) at 56° C. the tetra-methylation was judged to be complete by tlc. The mixture was filtered at 50° C., the grey catalyst was washed with 2-methyltetrahydrofuran (3×5 ml) and the filtrate and washings were combined. To this solution was added methanol (5 ml) to homogenise the mixture. Cooling to 5° C. resulted in a colourless solid precipitating from solution. A further two volumes of methanol (10 ml) were added and the slurry was stirred for 50 minutes at 5° C. The crude product was isolated by filtration to give a colourless solid, which was washed with methanol (3×5 ml) and dried at 50° C. for 16 hours (MW 327.45 g/mol, 2.26 g, 46%). The filtrate from the isolation process had water added (50 ml), which gave further solid. The suspension was stirred at 5° C. for 2 hours before being collected by filtration, washed with methanol (3×5 ml) and dried at 50° C. for 13 hours. (MW 327.45 g/mol, 0.83 g, 17%). The total yield of product was (3.09 g, 63%).

Notes

1. Normal phase tlc conditions, eluent 75% ethyl acetate, 25% petroleum spirit (40-60° C.), and UV lamp at 254 nm.
2. The retention factor of the dinitro starting material is 0.68 as a yellow spot, the retention factor of the hydrogenation product is 0.25 as a blue spot and the retention factor of the methylation product is 0.67 as a light blue spot.
3. The method for the tlc analysis of the hydrogenation step was direct spotting whereas the analysis of the methylation product had water added to a reaction aliquot which was extracted with ethyl acetate and then spotted.
4. After 17 hours the tlc shows two spots, the major spot was the reduction product the minor spot is unknown.
5. After 88 hours the tlc shows mainly the tetra-methylated product as the major spot.
6. Typically the reduction and methylation would be complete within 72 hours.
7. 1H NMR spectroscopy of the two samples gave identical spectra, traces of 2-methyltetrahydrofuran were detected along with an unknown signal at 5 ppm.

NMR: The product (10 mg) was dissolved in CDCl₃ (1.5 ml).

60

δ_H (400 MHz; CDCl₃): 2.09 (3H, s, CH₃), 2.86 (12H, s, NCH₃), 6.54 (2H, d, J 8, ArH), 6.64 (2H, s, ArH), 7.19 (2H, brd s, ArH)

Alternative Synthesis of 10-Acetyl-N,N,N',N'-tetramethylphenothiazine-3,7-diamine (5)

To a 50 ml round bottom flask was added 3,7-dinitro-10-acetylphenothiazine (MW 331.31 g/mol, 1 g, 3.02 mmol), zinc dust (MW 65.39 g/mol, 1.38 g, 21.13 mmol), methanol (6 ml) and tetrahydrofuran (2 ml). The mixture was heated to 50° C. after which a warm solution (45-50° C.) of aqueous ammonium chloride (MW 53.49 g/mol, 2.26 g, 42.26 mmol dissolved in 6 ml of water) was added slowly to maintain a gentle reflux. The mixture was then heated to 70° C. and stirred at this temperature for two hours after which it was cooled to ambient temperature (23° C.). The cooled mixture was filtered to remove the zinc salts and the filtrate containing compound 4 was treated with paraformaldehyde (MW 30.03 g/mol, 1.09 g, 36.22 mmol), sodium cyanoborohydride (MW 62.84 g/mol, 1.14 g, 18.11 mmol) and acetic acid (2 ml). The mixture was heated to 50° C. and stirred at this temperature for 3 hours. After cooling to ambient temperature (23° C.), water (2×10 ml) was added and the colourless slurry was stirred for 16 hours. The solid was then collected by filtration and washed with methanol (3×2 ml) to give the title compound (MW 327.45 g/mol, 0.91 g, 92%) as an off-white solid.

Notes


1. Reduction reaction using zinc and aqueous ammonium chloride was fast and clean taking only 2 hours to reach completion with no other spots recorded by tlc analysis.
2. The reductive methylation using sodium cyanoborohydride, paraformaldehyde and acetic acid was fast and clean, taking only 3 hours to reach completion.

NMR: The product (10 mg) was dissolved in CDCl₃ (1.5 ml).

δ_H (400 MHz; CDCl₃): 2.17 (3H, s, CH₃), 2.94 (12H, s, NCH₃), 6.61 (2H, d, J 8, ArH), 6.71 (2H, s, ArH), 7.26 (2H, brd s, ArH)

Synthesis 1

Synthesis of N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diaminium bis(methanesulphonate) (LMT.2MsOH)



i. MSA, H₂O, toluene, 85° C. ii. EtOH

61

10-Acetyl-N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine (AcMT) (150 g) was added to a 3-neck round bottomed flask. Toluene (1.8 l) was added and the mixture heated to reflux for 30 min. The solution was allowed to cool to 70° C. before being passed through an in-line 5 μ filter to a jacketed vessel fitted with distillation apparatus.¹ Toluene (150 ml) was added to the round bottom flask. This was used to rinse the transfer line and filter. Approximately 1.4 l. of toluene was distilled off under reduced pressure.² The temperature was lowered to 18° C. before water (42 ml) was added.³ This was followed by the addition of methanesulphonic acid (MSA) (65.5 ml, 99%, 2.2 equiv.) over a 5 min. period.⁴ A second portion of water (18 ml) was added. The mixture was heated to 85° C. for 3 h by which time the reaction was judged complete by tlc analysis. The biphasic solution was allowed to cool to 50° C. before absolute EtOH (150 ml) was added over 20 min.⁵ The mixture was seeded using 150 mg of N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diaminium bis(methanesulphonate).⁶⁻⁸ A second portion of EtOH (600 ml) was added over 90 min.⁹ and the reaction allowed to cool to 20° C. over 1 h.¹⁰ It was stirred at this temperature for 1 h. before the solid was collected by filtration. The cake was washed with 3 \times 300 ml of MeCN,¹¹ sucked dry for 5 min. and placed under vacuum overnight to give the product as a yellow crystalline solid (85-90% yield).

μ_{max} (KBr)/cm⁻¹; 3430 (NH), 3014 (=CH), 2649 (C—H), 1614 (C=C), 1487 (C—C), 1318 (S=O), 1199 (SO₂-O), 1059 (S=O), 823 (ArC—H)

δ_H (600 MHz; CD₃OD); 2.71 (6H, s, SCH₃), 3.21 (12H, s, NCH₃), 6.75 (2H, d, J=8.8 Hz, ArH), 7.22 (4H, d J 2.9 Hz, ArH), 7.24 (4H, dd J 2.9, 8.8 Hz, ArH),

δ_C (100 MHz; CD₃OD); 38.2 (SCH₃), 45.9 (NCH₃), 115.0 (CH), 118.2 (CH), 118.7 (QC), 119.9 (QH), 137.1 (CH), 142.8 (QC)

MP: 271° C.

m/z (EI+): Calculated 285.129970; Observed 285.131292 (100%, [M-2MSA]⁺).

m/z (ES-): Calculated 95; Observed 95 (100%, [M-LMT]⁻).

Elemental analysis % (C₁₈H₂₇N₃O₆S₃): Calculated C (45.26), N (8.80), S (20.14), H (5.70); Observed C (45.19), N (8.76), S (19.84), H (5.53)

Notes

1. Heating to reflux ensures complete dissolution of AcMT for transfer through 5 μ filter. Toluene is a good solvent and a 70° C. target is a compromise between ensuring the material stays in solution and minimising potential damage to plastic transfer hoses and filter.
2. 500 ml of remaining toluene ensures reaction volume meets minimum stir depth of reactor.
3. Volume of water is controlled to ensure product crystallises out as a free flowing precipitate. Seeding the reaction reduces the impact of small variations in water volume.
4. 2.2 equivalents of MSA are used to effect the hydrolysis and form the salt whilst leaving a sufficient quantity of excess acid (0.2 equiv.) to ensure the stability of the product in solution. Addition of MSA causes a slight exotherm, hence the 5 minute addition time.
5. EtOH is used as counter solvent to precipitate the product. A portion is added before the seed to ensure the seed does not dissolve. An extended addition time ensures controlled crystallisation of the product (see notes 7 and 8).
6. It is possible to carry out the reaction without the use of a seed, however its incorporation ensures the early precipitation of LMT.2MsOH which in turn prevents formation of by-products (such as the alcohol ester EMS a potential

62

genotoxic by-product—not detected in the synthetic process) and encapsulation of EtOH.

7. The seed is also useful as a means of controlling the particle size of the product. When the seed material was used which has been ground in a mortar and pestle to <100 μ m a significant reduction in the average particle size of the product is observed. When <100 μ seed which had not been ground was used no such effect was observed. Therefore, without wishing to be bound by theory, it appears that the ability of the seed to control the particle size is not a function of the seed particle size, it is linked to the proportion of internal or 'new' crystal faces that the crushing of the seed has exposed.
8. Finally, when the seed material was relatively large and uncrushed a considerable amount of product (skin) may adhere to the side of the reactor vessel during the EtOH addition. This may be reduced by introducing a heat/cool cycle into the process after the EtOH addition. However, an unexpected bonus of the utilisation of the crushed seed was that the level of skinned material present after EtOH addition was reduced by ~90%. Therefore it was no longer necessary to carry out the heat/cool cycle. It seems that this is linked to the small seed size rather than new faces because when the reaction was carried out using uncrushed <100 μ seed the same reduction in skinning was observed.
9. The rate of EtOH addition has an effect on particle size and EtOH inclusion. Fast addition (<1 h) reduces particle size however EtOH inclusion increases. A slow addition (2 h) has the opposite effect hence a balance must be struck.
10. Rate of cooling has a similar although reduced effect. Fast cool down (<1 h) leads to reduction in particle size with a concomitant increase in EtOH levels. A slow cool has the opposite effect.
11. EtOH is equally effective as MeCN at removing the related substances, however its use is accompanied by a slight increase in the level of retained EtOH.

Characterisation of N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diaminium bis(methanesulphonate) (LMT.2MsOH)

Elemental Analysis (Microanalysis)

The analysis has good correlation between the theoretical values and the analysis values for carbon, nitrogen, hydrogen and sulphur.


Results of the elemental analysis:

Molecular Formula C ₁₈ H ₂₇ N ₃ O ₆ S ₃		
Element	% Theoretical	% Found
C	45.26	45.19
H	5.70	5.53
N	8.80	8.76
S	20.14	19.84

¹H—Nuclear Magnetic Resonance (NMR) Spectroscopy

The ¹H NMR spectrum was obtained in deuterated methanol CD₃OD, on a Varian 600 MHz instrument and is shown in FIG. 1.

63

Assignment of the ^1H NMR spectrum is below:


Assignment	Chemical Shift (ppm)	Protons	Group
15/16	2.71	6H, s	2 x SCH ₃
11/12/13/14	3.21	12H, s	2 x N(CH ₃) ₂
1/9	6.75	2H, d, 8.8 Hz	2 x C—H (Aromatic)
4/6	7.22	2H, d, 2.9 Hz	2 x C—H (Aromatic)
2/8	7.24	2H, dd, 8.8 and 2.9 Hz	2 x C—H (Aromatic)

 ^{13}C —Nuclear Magnetic Resonance (NMR) Spectroscopy

The ^{13}C NMR spectrum was obtained on a Varian 400 MHz NMR instrument at a frequency of 100.56 MHz in deuterated methanol CD₃OD and is shown in FIG. 2.

The initial assignment of the ^{13}C -NMR spectrum was based on correlation with charts of known chemical shifts, (Literature Reference: Structure Determination of Organic Compounds: Tables of Spectral Data, Pretsch E., et al., Springer, London, p 122).

Further assignments utilised DEPT-135, HSQC and HMBC experiments to unambiguously confirm the assignments. DEPT-135 (Distortionless Enhancement by Polarisation Transfer), HSQC (Heteronuclear Single Quantum Coherence) and HMBC (Heteronuclear Multiple Bond Correlation) spectra were obtained on a Varian 400 MHz NMR instrument at a frequency of 100.56 MHz (see FIGS. 3-5).

Assignment	Chemical shift (PPM)	NMR Region	DEPT-135
15/16	38.2	Alkyl	CH ₃
11/12/13/14	45.9	Alkyl	CH ₃
1/9	115.0	Aromatic - C	CH
4/6	118.2	Aromatic - C	CH
4a/5a	118.7	Aromatic - C	C
2/8	119.9	Aromatic - C	C
3/7	137.1	Aromatic - C	CH
9a/10a	142.8	Aromatic - C	C

Infrared Spectroscopy (IR)

A sample was thoroughly mixed and ground in a mortar and pestle with 200 mg of anhydrous KBr. This mixture was then pressed into a disc, using a die at a pressure of 1500 psi. The IR spectrum was then obtained on a Nicolet Avatar 320 FT-IR spectrometer. The spectrum is shown in FIG. 6.

64

Assignment of the infrared spectrum:

Peak Wavenumber (cm ⁻¹)	Peak Type	Assignment
~3430	broad	N—H stretch
3014	medium	=C—H stretch
2649	medium	C—H stretch
1614	medium	C=C stretch
1487	strong	C—C stretch
1318	strong	S=O stretch
1199	strong	SO ₂ —O stretch
1059	strong	S=O stretch
823	strong	Aromatic C—H stretch

Mass Spectrometry (MS)

Mass spectroscopic analysis was carried out using a Waters, LCT Premier XE mass spectrometer. A flow rate of 1 ml/hr was adopted. The source used for the analysis of the active component was electron impact ionisation in the positive mode. The source used for the analysis of the methanesulphonate counter ion was electrospray ionisation in the positive mode.

Using electron impact ionisation a major peak is observed at 285 (see FIG. 7). This corresponds to the molecular ion C₁₆H₁₉N₃S. A comparison of the exact mass measured and the theoretical value is provided below:

Theoretical	Peak m/z	Abundance (%)	Assignment
285.129970	285.131292	100	C ₁₆ H ₁₉ N ₃ S

The measured accurate mass is in good agreement with the calculated mass for C₁₆H₁₉N₃S.

Using electrospray ionisation a major peak is observed at 95 (see FIG. 8). This corresponds to the molecular ion of the counter ion CH₃O₃S:

Peak m/z	Abundance (%)	Assignment
95	100	CH ₃ O ₃ S

Ultraviolet-Visible Spectroscopy (UV-Vis)

A 5 mg sample was dissolved in de-ionised water, and made up to 100 ml in a volumetric flask. The analysis was carried out using quartz curvets in a Perkin Elmer Lambda 25 UV/Vis spectrometer. The UV-Vis spectrum is shown in FIG. 9.

Assignment of the UV-Vis spectrum:

λ_{max} (nm)	Absorbance
226	1.7615
255	3.5860
332	0.7527
664	0.1845

The extinction coefficient ϵ for the lambda max at 255 nm was 34254.64. This was calculated according to the Beer-Lambert Law:

$$\epsilon = \frac{A}{C \times l}$$

65

where A=Absorbance Log (I_0/I) 3.5860; C=Concentration Mol/L; l=path length 1 cm

High Performance Liquid Chromatography (HPLC)

A 100 mg sample was submitted for HPLC analysis. The analysis was carried out on an Agilent 1200 series with VWD 5 Detector or PDA for identity, according to the method summarised in the table below.

HPLC method:

Parameters	Conditions
Column	Zorbax SB-CN, 50 × 4.6 mm, 3.5 μm.
Column temperature	283 K
Mobile phase	A: 0.1% v/v Formic acid in water B: 100% acetonitrile
Flow rate	1 ml/min
Injection volume	5 μl
Stop time	22 min.
Wavelength	UV at 255 nm Bandwidth at 4 nm. Reference wavelength set at off. PDA scan 190 nm to 800 nm (Identity only)
Auto sampler temperature	278 K Protected from light.

	Time (min.)	Mobile Phase A (%)	Mobile Phase B (%)
Mobile Phase Gradient	0.0	100	0
	10.0	90	10
	17.0	50	50
	18.0	50	50
	18.1	100	0
	22.0	100	0

The HPLC trace is shown in FIG. 10. The organic purity was found to be 99.45% w/w.

HPLC Analysis (% Purity) including retention times		
LMT	MT ⁺	Leuco Azure B
6.39 min. 99.45	14.38 min. 0.55	5.77 min. <0.05

Crystalline Form

In the above-described method, LMT.2MsOH is produced in crystalline form. The crystalline form of LMT.2MsOH is illustrated by the X-ray powder diffraction spectrum shown in FIG. 11. The XRPD exhibits sharp signals, indicative of a high degree of crystalline order. Variations in relative peak intensity may be observed, which are attributable to orientation effects in combination with differences in particle size. Only slight variations in relative peak intensity (less than 50%) are observed as a function of sample thickness (0.1 mm vs. 1.0 mm).

The crystal form is further characterised by FT-Raman, thermogravimetric (TG), differential scanning calorimetric (DSC), dynamic vapour sorption (DVS) analysis, and microscopy (FIGS. 12-16). This form may conveniently be referred to as 'Form A'.

Crystals for single crystal X-ray analysis were obtained from ethanol, methanesulfonic acid and water. See FIG. 17c. Instrumental Details

X-Ray Powder Diffraction: Bruker 08 Advance, Cu Ka radiation ($\lambda=1.54180 \text{ \AA}$), 40 kV/40 mA, LynxEye detector, 0.02° step size in 2θ , 37 s per step, 2.5°-50° 2θ scanning range. The samples were prepared on silicon single crystal sample holders with 0.1 or 1.0 mm depth without any special treatment other than the application of slight pressure to get a flat surface. All samples were rotated during the measurement.

66

Differential Scanning Calorimetry: Perkin Elmer DSC 7. Gold crucibles closed under N₂, heating rate 20° C./min, scan from -50° C. to 280° C.

Dynamic Vapor Sorption: Projekt Messtechnik SPS 11-100n water vapor sorption analyzer. The samples were placed in aluminum crucibles on top of a microbalance and were equilibrated at 25° C. and 50% r.h. before starting a pre-defined humidity program at 25° C. (50-0-95-50% r.h., scanning with $\Delta \text{ r.h.}=5\% \text{ h}^{-1}$ and with 'isohumid' equilibration periods at the extreme values).

FT-Raman Spectroscopy: Bruker RFS100. Nd:YAG 1064 nm excitation, 50 mW laser power, Ge-detector, 128 scans, range 50-3500 cm^{-1} , 2 cm^{-1} resolution. Aluminum sample holder.

Polarizing Light Microscopy: Leitz Orthoplan microscope with Leica OFC280 CCO camera.


TG: TA Instruments TGA Q5000. Open aluminum crucible, N₂ atmosphere, heating rate 10° C. min^{-1} , range 25-300° C.

TG-FTIR: Netzsch Thermo-Microbalance TG 209 with Bruker FT-IR Spectrometer Vector 22. Aluminum crucible with micro-hole, N₂ atmosphere, heating rate 10° C. min^{-1} , range 25-250° C.

Without wishing to be bound by theory, it is suggested that this form represents the only stable polymorphic form of LMT.2MsOH. Polymorphism studies have shown that Form A is reproduced in nearly all crystallisation systems (studies were performed using de-gassed solvents, under an inert atmosphere).

Amorphous LMT.2MsOH can be prepared by evaporation of an aqueous solution of LMT.2MsOH, however the amorphous material recrystallises to Form A upon further drying.

Industrial Scale Synthesis of AcMT and LMT.2MsOH



i; N₂H₄·H₂O, Et₃N, MeCN, N₂, 65° C., 1 h; ii; Ac₂O, N₂, 95° C., 2 h; iii; MSA, H₂O, Toluene, N₂, 85° C., iv; EtOH.

67

Large scale synthesis of 10-acetyl-N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine (AcMT)

Acetonitrile (MeCN) (300 l) was added to reactor 1 (R1) and cooled to $-5-0^{\circ}\text{C}$. Methylthionium chloride rehydrate (MTC.3H₂O) (150 kg) was added and the temperature increased to $15-25^{\circ}\text{C}$. Triethylamine (Et₃N) (100 l) was added followed by MeCN rinse (20 l). Hydrazine hydrate (N₂H₄.H₂O) (12 l) was added over 30 min. The reaction temperature was increased to $60-70^{\circ}\text{C}$. over 1 h and then maintained at this temperature for 1 h before being reduced to $40-50^{\circ}\text{C}$. Acetic anhydride (Ac₂O) (240 l) was added over 1 h followed by MeCN (20 l) rinse. Batch temperature was increased to $90-100^{\circ}\text{C}$. for 2 h. Temperature was reduced to $55-65^{\circ}\text{C}$. and water (340 l) was added over 2 h whilst maintaining the temperature. The batch temperature was then reduced to $-5-5^{\circ}\text{C}$. over 2 h. and held there for 6 h. The solid was collected by filtration. The cake was fully de-liquored before water (400 l) was added to R1. The temperature in R1 was allowed to rise to $15-25^{\circ}\text{C}$. before the water was used in portions to wash the filter cake. The product was dried under a stream of nitrogen for 6 h. before being offloaded (Yield: 90-110 kg).

Large scale purification of 10-acetyl-N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine (AcMT)

Water (300 l) was added to R1, followed by 10-Acetyl-N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine (AcMT) (100 kg). Toluene (400 l) and 80% aqueous acetic acid (40 l) were added, followed by a water rinse (50 l). The batch temperature was increased to $75-85^{\circ}\text{C}$. for 1 h. The agitator was stopped and the layers allowed to settle for 30 min. The lower aqueous layer is removed and fresh water (300 l), 80% aqueous acetic acid (40 l) followed by water rinse (50 l) were then added. The mixture was stirred at $75-85^{\circ}\text{C}$. for 1 h before the agitator was stopped and the layers allowed to settle for 30 min. The lower aqueous layer was removed and fresh water (300 l), 80% aqueous acetic acid (40 l) followed by water rinse (50 l) were then added. The mixture was stirred at $75-85^{\circ}\text{C}$. for 1 h before the agitator was stopped. The layers were allowed to settle for 30 min before the lower layer was removed and water (390 l) was added and the mixture stirred for 1 h. The agitator was stopped and the layers allowed to settle for 30 min. The lower aqueous layer was removed and the temperature reduced to $-5-5^{\circ}\text{C}$. The jacket temperature was increased to 80°C . and then when it reached 60°C . the temperature was reduced to $-10-0^{\circ}\text{C}$. over 2 h. The mixture was stirred for 4 h before it was transferred to the filter. The cake was fully de-liquored before toluene (150 l) was added to R1. The toluene was stirred in R1 for 30 min. before it was used in portions to wash the filter cake. The product was dried on the filter under a stream of nitrogen for 48 h until loss on drying $<1\%$ before being offloaded (Yield: 75-90 kg).

Large scale synthesis of N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diaminium bis(methanesulphonate) (LMT.2MsOH)


AcMT (18-22 kg) was added to R1. Toluene (volume (l)=16×AcMT weight) was added and the mixture heated to $90-100^{\circ}\text{C}$. for 30 min. The solution was allowed to cool to $60-80^{\circ}\text{C}$. before being passed through an in-line 5μ filter to reactor 2 (R2). Toluene (50 l) was added to reactor 1 (still at $\sim 70^{\circ}\text{C}$.) and stirred for 30 min. This was used to rinse the transfer line and filter. The above process was repeated once more. The process of removing the excess toluene from R2 by

68

distillation under reduced pressure was then started. The capacity of R2 permitting, two more portions of AcMT (18-22 kg each) were transferred from R1 to R2 following the method described above. The distillation was complete when the batch volume in R2 was reduced to ~ 340 l. The temperature was increased to $95-105^{\circ}\text{C}$. for 15-30 min. before being cooled to $15-25^{\circ}\text{C}$. Water (20 l) was added to R2. This was followed by the addition of methanesulphonic acid (MSA) (33 l, 99%, 2.2 equiv.) whilst keeping the batch temperature at $15-30^{\circ}\text{C}$. A second portion of water (10 l) was added and the mixture stirred at this temperature for 2 h. The mixture was heated to $80-90^{\circ}\text{C}$. for 3-4 h. The biphasic solution was allowed to cool to $48-58^{\circ}\text{C}$. before absolute EtOH (75 l) was added over 15-30 min. The stirrer was stopped and the mixture was seeded using 150 g of crushed ($<100\mu$) N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diaminium bis(methanesulphonate). A second portion of EtOH (300 l) was added over 80-110 min. Jacket temperature was set to 10°C . and when the temperature reached 25°C . the jacket temperature was reset to 20°C . It was stirred at $15-25^{\circ}\text{C}$. for 2 h. before the solid was collected by filtration. The cake was thoroughly de-liquored. MeCN (300 l) was added to R2 and stirred for 15 min before being used portion-wise to wash the filter cake. A second 300 l of MeCN was added to R2 and the wash process repeated. The product was dried on the filter until loss on drying $<0.2\%$ before being offloaded (80-90% yield).

Synthesis 2

Synthesis and analysis of N,N,N',N'-Tetramethyl-10H-phenothiazine-3,7-diaminium bis(ethanesulfonate) (LMT.2EsOH)

step i; H₂O, MeOH, EsOH, IPA, Acetone

Synthetic Method for LMT.2EsOH

The synthesis of LMT.2EsOH was carried out by acid hydrolysis of 10-Acetyl-N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine. The acid used was ethanesulfonic acid and the solvent combination was aqueous methanol.

Experimental Details

To a 100 ml round bottom flask was added 10-Acetyl-N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine (5 g, 15.27 mmol, MW 327.45 g/mol), (70%, aq) ethanesulfonic

69

acid (7.21 g, 45.81 mmol, MW 110.13 g/mol) and methanol (25 ml). The mixture was heated to 75° C. and stirred at this temperature 4 hours before the mixture was cooled over ice water. No solid formed and the methanol was removed under vacuum to give a viscous green oil. To this oil was added isopropanol (25 ml) and the mixture was heated to reflux to ensure a homogenous solution. Once cooled acetone was added until a precipitate formed. The suspension was cooled over ice water for 1 hour before being filtered to give the crude product as a yellow solid, which turned green upon exposure to air. The crude was washed with acetone (3×5 ml) and air dried for 3 days to give the crude product (3.35 g, 43%, MW 505.68 g/mol) as a light green solid.

μ_{max} (KBr)/ cm^{-1} : 3448 (NH), 3263 (=CH), 3030 (=CH), 2987 (CH), 2938 (CH), 2582 (SO₃H), 2452 (SO₃H), 1487 (C—C), 1211 (O=S=O), 1188 (O=S=O), 1145 (O=S=O), 1026.

δ_H (400 MHz; D₂O): 1.07 (6H, t, J 7.6, CH₃), 2.72 (4H, q, J 7.6, SCH₂), 3.02 (12H, s, NCH₃), 6.54 (2H, d, J 9.2, ArH), 7.02 (4H, brd s, ArH);

δ_C (100 MHz; D₂O): 142.3 (QC), 136.6 (QC), 119.9 (CH), 118.4 (QC), 118.2 (CH), 115.2 (CH), 46.2 (NCH₃), 45.3 (SCH₂), 8.3 (CH₃).

MP: 208-210° C. (IPA/Acetone)

m/z (EI+): Calculated mass 285.129970; Observed 285.129761 (100%, [M-2EsOH]⁺).


m/z (ES-): Calculated mass 109; Observed 109 (100%, [M-LMT]⁻).

Crystallography

A 1 g sample of LMT.2EsOH was dissolved in acetic acid (~0.1 g) and ethyl acetate was layered on top and allowed to slowly diffuse over 3 days in the dark. Crystals developed and were collected and analysed by X-ray diffraction and confirmed the product as the bis(ethanesulfonate). See FIG. 17a.

Synthesis 3

Synthesis and analysis of N,N,N',N'-Tetramethyl-10H-phenothiazine-3,7-diaminium bis(p-toluenesulfonate) (LMT.2TsOH)



Synthetic Method for LMT.2TsOH

The synthesis of LMT.2TsOH was carried out by neutralising N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-di-

70

aminium dichloride with sodium carbonate and extracting the neutral species into organic solvent. The extract was treated with p-toluenesulphonic acid and the mixture concentrated to dryness.

Experimental Details

To a 50 ml beaker was added sodium carbonate (0.59 g, 5.58 mmol, MW 105.99 g/mol) and water (10 ml), the mixture was stirred until the solid had dissolved. To a 100 ml separating funnel was added N,N,N',N'-Tetramethyl-10H-phenothiazine-3,7-diaminium dichloride (1 g, 2.79 mmol, MW 358.33 g/mol), tetrahydrofuran (35 ml) and diethylether (5 ml) then the aqueous solution of sodium carbonate. The neutral species was extracted into the organic solvent layer and separated from the aqueous layer. To the organic extract was added p-toluenesulphonic acid monohydrate (1.06 g, 5.58 mmol, MW 190.20 g/mol) pre-dissolved in tetrahydrofuran (5 ml) and the mixture was concentrated to dryness to give the product (MW 629.8216 g/mol) as a crunchy green amorphous foam.

μ_{max} (KBr)/ cm^{-1} : 3440 (NH), 3270 (=CH), 3032 (=CH), 2628 (SO₃, H), 1484 (C—C), 1194 (O=S=O), 1122 (O=S=O), 1032.

δ_H (400 MHz; D₂O): 2.24 (6H, s, CH₃), 3.09 (12H, s, NCH₃), 6.62 (2H, d, J 8.4, ArH), 7.10 (4H, s, ArH), 7.13 (4H, d, J 8.4, Ts-H), 7.61 (4H, d, J 8.4, Ts-H)

δ_C (100 MHz; D₂O): 19.9 (CH₃), 45.9 (NCH₃), 115.0 (CH), 118.2 (CH), 118.6 (QC), 119.9 (CH), 125.5 (CH), 128.5 (CH), 137.0 (QC), 140.5 (QC), 141.9 (QC), 142.8 (QC).


Mp: 108° C. (THF/Et₂O)

m/z (EI+): Calculated mass 285.129970; Observed 285.129398 (100%, [M-2TsOH]⁺).

m/z (ES-): Calculated mass 171.0116; Observed 171.0121 (100%, [M-LMT]⁻).

Synthesis 4

Synthesis and analysis of N,N,N',N'-Tetramethyl-10H-phenothiazine-3,7-diaminium ethanesulfonate (LMT.EDSA)



The synthesis of LMT.EDSA was carried out by acid hydrolysis of 10-acetyl-N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine. The acid used was 1,2-ethanesulfonic acid and the solvent combination was aqueous ethanol.

71

Experimental Details

To a 25 ml round bottom flask was added 10-acetyl-N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine (1 g, 3.05 mmol, MW 327.45 g/mol), 1,2-ethanedithiolic acid monohydrate (0.95 g, 4.58 mmol, MW 208.21 g/mol), water (1 ml) and ethanol (5 ml). The mixture was heated to 85° C. and stirred at this temperature for 2.5 hours where a yellow green solid precipitated from solution. The slurry was cooled over ice water for 30 min before filtering to give the crude product as a green yellow solid. The crude was washed with ethanol (3×3 ml) and air dried for 15 min before being oven dried for 3.5 hours at 70° C. to give the crude product (1.33 g, 91%, MW 475.61 g/mol) as a yellow solid.

Purification of LMT.EDSA

To a 50 ml conical flask was added crude LMT.EDSA (1 g, 2.10 mmol, MW 475.61 g/mol) and water (10 ml). The slurry was heated to 95° C. and stirred at this temperature until the solid dissolved. The solution was then allowed to cool to 25° C. where a light green crystalline solid formed. The slurry was then cooled over ice water for 30 min before filtering. The solid collected was washed with methanol (3×3 ml) and air dried for 18 hours to give the purified product (0.88 g, 88%, MW 475.61 g/mol) as a crystalline light green solid.

μ_{max} (KBr)/cm⁻¹; 3408 (NH), 3280 (=CH), 3221 (C—H), 3036 (=CH), 2574 (SO₃H), 2480 (SO₃H), 1484 (C—C), 1226 (O=S=O)

δ_H (400 MHz; D₂O); 2.98 (12H, s, NCH₃), 3.06 (4H, s, SCH₂), 6.45 (2H, d, J 6, ArH), 6.95 (4H, d J 4, ArH)

δ_C (100 MHz; D₂O); 46.2 (NCH₃), 46.4 (SCH₂), 115.1 (CH), 118.1 (CH), 118.4 (QC), 119.8 (CH), 136.5 (QC), 142.1 (QC)

MP: decomposes at 268° C. (H₂O)

m/z (EI+): Calculated 285.129970; Observed 285.130948 (100%, [M-EDSA]⁺).


m/z (ES-): Calculated 188.9528; Observed 188.9535 (100%, [M-LMT]⁻).

Crystallography

A 40 mg sample of LMT.EDSA was dissolved in hot deuterated water (~1 ml) and allowed to slowly cool in the dark. Crystals developed which were collected and analysed by X-ray diffraction and confirmed the product as the monohydrate of the 1:1 LMT to EDSA adduct. See FIG. 17b.


Synthesis 5

Synthesis and analysis of N,N,N',N'-Tetramethyl-10H-phenothiazine-3,7-diaminium naphthalenedisulfonate (LMT.NDSA)



72

-continued

15 step i; H₂O, EtOH, NDSA

Synthetic Method for LMT.NDSA

The synthesis of LMT.NDSA was carried out by acid hydrolysis of 10-acetyl-N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine. The acid used was 1,5-naphthalenedisulfonic acid and the solvent combination was aqueous ethanol.

Experimental Details

To a 25 ml round bottom flask was added 10-Acetyl-N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diamine (1 g, 3.05 mmol, MW 327.45 g/mol), 1,5-naphthalenedisulfonic acid tetrahydrate (1.65 g, 4.58 mmol, MW 360.36 g/mol), water (1 ml) and ethanol (5 ml). The mixture was heated to 85° C. and stirred at this temperature for 30 minutes where the mixture was still insoluble. To the hot mixture was added water (4 ml) and the reaction heated to 95° C. and stirred at this temperature for 8 hours. The suspension was cooled over ice water for 10 minutes before being filtered to give the crude product as a light green solid. The crude was washed with ethanol (3×5 ml) and air dried for 3 days to give the crude product (1.75 g, 100%, MW 573.71 g/mol) as a light green blue solid.

μ_{max} (KBr)/cm⁻¹; 3382 (NH), 3302 (=CH), 3040 (=CH), 2525 (SO₃H), 1478 (C—C), 1238 (O=S=O), 1219 (O=S=O), 1179, 1158, 1030.

δ_H (400 MHz; D₂O); 3.06 (12H, s, NCH₃), 6.70 (2H, brd, ArH), 7.14 (4H, brd, ArH), 7.43 (2H, t, J 8.0, 7.6, Naph-H), 7.94 (2H, d, J 7.2, Naph-H), 8.87 (2H, d, J 78.4, Naph-H), 9.10 (1H, s, NH)

δ_C (100 MHz; D₂O); 46.0 (NCH₃), 115.3 (CH), 117.5 (QC), 118.7 (CH), 120.4 (CH), 124.6 (CH), 124.7 (CH), 129.6 (CH), 129.9 (QC), 138.3 (QC), 141.7 (QC), 143.8 (QC).

MP; decomposes at 256° C. (MeCN)

m/z (EI+): Calculated mass 285.129970; Observed 285.130367 (100%, [M-NDSA]⁺).

m/z (ES-): Calculated mass 286.9684; Observed 286.9697 (100%, [M-LMT]⁻).

Example 2

Solubility Studies

i) Solubility of N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diaminium dibromide, dichloride and bis(methanesulphonate) (LMT.2HBr, LMT.2HCl and LMT.2MsOH) salts

Two aqueous solutions (pH 2.00 and 3.01 at 21.4° C.) were prepared by carefully adding HCl (5 M) to deionised water.

In each experiment a 5 ml aliquot of one of the aforementioned solutions was heated to 37° C. A portion of the appro-

priate salt (LMT.2MsOH, LMT.2HCl or LMT.2HBr) was added and the mixture stirred for a few moments to allow for complete dissolution of the solid. This step was repeated until no further dissolution took place.

The results are shown in the Table:

Salt	pH (21.4° C.)	g/5 ml* (37° C.)
LMT·2HBr	3.01	4.726-5.236
LMT·2HBr	2.00	4.822-5.096
LMT·2HCl	3.01	4.978-6.029
LMT·2HCl	2.00	4.404-4.961
LMT·2MsOH	2.00	8.825-9.943

*Lower limit of range corresponds to total weight at which complete dissolution was observed. Upper limit is total weight added before saturation was achieved

As can be seen LMT.2MsOH has a good aqueous solubility.

ii) pH Dependence of LMT. 2MsOH Salt

In related experiments three buffered stock solutions were prepared (pH 2, pH 3, and pH 7) as follows:

pH 2 Buffered Aqueous Solution

A solution of (0.2M) potassium chloride (KCl) (0.745 g in 50 mL of deionised water) was initially prepared. From this solution 50 mL was taken and diluted with approximately 80 mL of deionised water. A (0.2 M) hydrochloric acid (HCl) solution was then used to adjust the pH to 2, before further dilution with deionised water to make up to 200 mL. A final pH of 2.00 at 21.6° C. was recorded.

pH 3 Buffered Aqueous Solution

A solution of (0.1M) potassium hydrogen phthalate (2.042 g in 100 mL of deionised water) was initially prepared. From this solution 100 mL was taken and diluted with approximately 50 mL of deionised water. A 0.2 M HCl solution was then used to adjust the pH to 3, before further dilution with deionised water to make up to 200 mL. A final pH of 2.99 at 21.7° C. was recorded.

pH 7 Buffered Aqueous Solution

A solution of (0.1M) potassium phosphate monobasic (KH₂PO₄) (1.370 g in 100 mL of deionised water) was initially prepared. From this solution 100 mL was taken and diluted with approximately 80 mL of deionised water. A 0.5 M sodium hydroxide (NaOH) solution was then used to adjust the pH to 7, before further dilution with deionised water to make up to 200 mL. A final pH of 7.07 at 22° C. was recorded.

Method

A 5 mL aliquot of an aqueous buffered solution was added to a vial which contained a micro-flea. This vial was placed into a water bath set at 25° C. To the solution was added LMT.2MsOH in 1-1.5 g portions. After each addition, a 10 minutes stir time was allowed to ensure maximum opportunity for dissolution. The homogeneity of the mixture was judged by eye. If solid was still present, after the stir time, as judged by visual inspection, the saturation point was judged to have been reached.

Results

The viscosity of the resulting mixtures precluded adequate isolation of the excess solid, therefore it was not possible to determine exact solubility values. Consequently each of the results are reported as a range in which the total mass of LMT.2MsOH added prior to the saturation point constitutes the lower limit and the total mass of LMT.2MsOH added, post saturation point, provides the upper limit.

The results from each of the three experiments are shown below:

pH	Solubility (g/mL)
2.00	1.600-1.773
2.99	1.981-2.092
7.07	2.033-2.114

As can be seen, the solubility tailed off slightly as the pH was reduced, however LMT.2MsOH performed well in each of the three aqueous systems.

In conclusion LMT.2MsOH has better aqueous solubility than MTC (not shown) and enhanced solubility compared to the corresponding chloride and bromide salts. This suggests an increased utility in respect of the treatment and uses described herein.

Example 3

Inhibition of Aggregation and Toxicity

Methods: Solid Phase Assay for Tau Aggregation

The tau-tau aggregation assay uses purified recombinant tau fragments in a solid-phase immunoassay. Methods are described in detail in e.g. WO 96/30766. Briefly, the assay measures the binding of truncated tau (amino acids 297-391) in solution to solid-phase bound truncated tau (residues 297-390). Binding of the former is detected with the antibody mAb 423, which specifically recognises peptides containing a C-terminal Glu-391 residue. The Tau complex formed in vitro is similar to the aggregated complex that forms in Alzheimer's disease as a consequence of the stability of the pathological Tau-Tau binding interaction through a 94/95-amino acid repeat domain (residues 297-390), found in the proteolytically stable core of the paired helical filament.

The B₅₀ value (expressed as mean±SE) is determined as the concentration of compound at which tau-tau binding is decreased by 50%.

Methods: Cell-Based Tau Aggregation Assay

The assay is based on 3T6 mouse cells that have been engineered to express both full-length human tau protein (htau40) under the control of an inducible promoter (pO-PRSVI), and to express low levels of truncated tau (295-390, dGA) under the control of a constitutive promoter (pcDNA3.1). Expression of large quantities of htau40 is induced by the addition of IPTG (10-50 μM), which in turn leads to the production of additional truncated tau by a process in which aggregation and processing of the full length-tau occurs in the presence of dGA tau which acts as template. Addition of tau-tau aggregation inhibitors to the assay blocks this process. Methods are described in more detail in WO 02/055720.

Results are expressed as the concentration at which there is a 50% inhibition of generation of the 12 kD fragment. This is referred to as the EC₅₀ value.

Cells (4A and clones thereof) are grown to ~80% confluency in a 10-cm dish, before splitting to two 24-well plates and allowed to grow for 24 hrs. Test item is added at various concentrations and, after 24 hrs, IPTG is added. After overnight incubation the medium is removed, the wells are washed with PBS and cells are collected by the addition of Laemmli buffer. Samples were stored at -20° C. for subsequent gel electrophoresis, Western blotting and antibody labelling. Samples are separated by SDS PAGE, transferred to PVDF membrane and the tau labelled with 7/51 antibody detected by ECL on a Kodak Image Station. Compound was typically tested at four concentrations in triplicate over a range of concentrations with all the samples being run on one

75

gel. The ratio of the intensities of the dGA to tau40 bands, normalised to control samples in which there had been no drug, was plotted against drug concentration and the EC₅₀ value was determined graphically from the concentration at which the ratio falls to 0.5.

The method is summarised in Table 1 immediately below. MTC (TR×0014.047) was run as a control in all experiments and the EC₅₀ value was normalised to MTC having an EC₅₀=0.59 μM.

TABLE 1

summary of assay procedure for measuring EC ₅₀	
Timing	Action
Day 1	Split cells to 24-well plates
Day 2	Add drug at various concentrations
Day 3	Late afternoon, add IPTG
Day 4	Morning, collect in Laemmli buffer, store -20° C. before further processing
Processing Day	Run samples on SDS Page gels, transfer to PVDF membrane, labeled with 7/51 anti tau antibody. Blots are quantified using the Kodak 1D software and data

TABLE 1-continued

summary of assay procedure for measuring EC ₅₀	
Timing	Action
	is transferred to the Systat statistics package for graphing.

Methods: Cellular Toxicity Assay

Cells (3T6 mouse fibroblast) are grown to ~80% confluency in a 10-cm dish, before splitting to 96 well plates, 10% of the 10-cm dish per 96-well plate, 50 μl per well. One column of 8 wells is left empty (to be a reagent blank in the assay). The cells were allowed to grow overnight before drug was added to four wells at the starting concentration (typically 200 μM for MTC or LMT.2HBr) and in subsequent wells using a 1:2 dilution series with the final four wells of cells being used as a control without drug. This allows two drugs to be tested per 96-well plate. The cells were left in the presence of drug for 48 hrs, after which medium was removed and cells washed with PBS. Cell number was determined using a Cytotox 96 well kit (Promega) which is based on the lactate dehydrogenase (LDH) assay. The assay quantitatively measures LDH, a stable cytosolic enzyme released on cell lysis. Released LDH is measured with an enzymatic assay

76

which results in the conversion of a tetrazolium salt into a red formazan product. The amount of colour formed is proportional to the number of cells lysed.

Briefly, cells are lysed with 50 μl/well 1× lysis buffer for 45-60 minutes, followed by 50 μl/well LDH assay reagent for 30 minutes and the reaction stopped with 50 μl/well stop buffer. The absorbance is read at 490 nm. The absorbance relative to untreated wells (untreated cells=1.0) was plotted against drug concentration. The LD₅₀ was determined graphically from the concentration at which the absorbance is decreased by 50%. MTC (TR×0014.047) was run as a control in all experiments when testing LMT.2HBr and the LD₅₀ value was corrected to MTC with an LD₅₀=65 μM.

Results:

Various bis(sulfonate) salts according to the invention were tested and compared with bis(halide) salts N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diaminium dichloride (LMTc, LMT.2HCl) and N,N,N',N'-tetramethyl-10H-phenothiazine-3,7-diaminium di(bromide) LMT.2HBr and with methylthioninium chloride (MTC).

In vitro data for the different methylthioninium salt forms are summarised in Table 2 immediately below:

TABLE 2

Summary of the in vitro data.				
Compound	LD ₅₀ (μM)	EC ₅₀ (μM)	THx	B ₅₀ (μM)
MTC	65 ± 5	0.59 ± 0.04	110	195.6 ± 16.1 (n = 10)
LMT.2HBr	61 ± 4 (n = 20)	0.66 ± 0.15 (n = 8)	92	472.4 ± 27.6 (n = 3)
LMT.2MsOH	34 ± 4 (n = 8)	0.19 ± 0.04 (n = 8)	179	238.2 ± 74.2 (n = 3)
LMT.2HCl	64 ± 8 (n = 10)	0.63 ± 0.10 (n = 7)	102	360.8 ± 38.2 (n = 3)
LMT.2TsOH	87 ± 10 (n = 8)	0.62 ± 0.34 (n = 2)	140	296.0 ± 37.9 (n = 3)
LMT.NDSA	77 ± 15 (n = 8)	0.71 ± 0.34 (n = 4)	108	333.7 ± 63.2 (n = 2)
LMT.EDSA	78 ± 6 (n = 8)	0.68 ± 0.32 (n = 4)	115	399.9 ± 17.6 (n = 2)
LMT.2EsOH	52 ± 3 (n = 8)	0.52 ± 0.13 (n = 3)	100	297.0 ± 75.1 (n = 3)
MSA*	—	NE (20)	—	>500
EDSA*	—	NE (20)	—	>500

THx, therapeutic index (THx = LD₅₀/EC₅₀)

Values expressed as mean ± SE.

NE = not effective (at max dose tested)

*MSA = methansulfonic acid; EDSA = ethanedisulfonic acid

Comments

The EC₅₀ values (mean±SE) for LMT.2MsOH and LMT.2HCl are 0.19±0.04 μM and 0.63±0.10 μM, respectively, with corresponding therapeutic indices of 179 and 102.

The relative potency of compounds in the cell-based model of tau-tau aggregation is LMT.2MsOH>MTC, LMT.2HBr, LMT.2HCl. The therapeutic index is 63% greater for LMT.2MsOH compared with MTC.

The order of potency in the cell-based assay is MTC, LMT.2MsOH>LMT.2HCl>LMT.2HBr. The B₅₀ values for LMT.2MsOH and LMT.2HCl are 238.2±74.2 μM and 360.8±38.2 μM, respectively. The order of relative potency in the cell-free assay is LMT.2MsOH>MTC, LMT.2HCl, LMT.2HBr.

Example 4

Toxicology, Impurities and Effect on the Hemopoietic System

LMT.2HBr, LMT.2HCl, LMT.2MsOH or MTC were administered daily for 14 days to female Wistar rats; the doses were 95 mg MT/kg/day from Days 1 to 10 and 60 mg MT/kg/day from Days 11 to 14. Clinical signs of raised body posture, subdued behaviour and general weakness were seen in all

treated groups. Treatment-related deaths occurred in the LMT.2HBr— and MTC-treated groups.

Changes in red blood cell parameters were seen in the blood and bone marrow of all treated groups that were indicative of a regenerative anaemia. These included: decreased numbers of red blood cells, low haemoglobin concentration and increased numbers of reticulocytes in blood and an increase in the numbers of red cell precursors in bone marrow. This was corroborated histologically by increased levels of erythropoiesis in the spleen.

A decrease in the numbers of neutrophilic granulocytes was seen in the bone marrow of all treated animals though the magnitude of this effect was considerably greater in the LMT.2HBr-treated group than in the other groups. This difference was also noted in the severity of the neutropaenia observed in prepared blood smears where there was a marked depletion of mature neutrophils in LMT.2HBr-treated animals, a modest decrease with MTC and no decrease in the LMT.2HCl or LMT.2MsOH groups. The results of this study suggest that, in rats at least, LMT.2HBr has a higher propensity to cause neutrophil depletion than LMT.2HCl, LMT.2MsOH or MTC. Decreased numbers of mature neutrophils and granulocytes were also observed in the bone marrow at the high dose (45 mg MT/kg/day) in a 6-month study of LMT.2HBr in the rat. The decreased neutrophils or neutropaenia observed after LMT.2HBr although reversible would make patients more susceptible to bacterial infections as their primary role is in destruction of bacteria. Thus LMT.2MsOH shows improved properties compared with LMT.2HBr in rats in terms of both tolerability (dose-related deaths) and in neutrophil response.

Table: Neutrophil response in rats following 14 day oral administration of different salt forms of LMT. Total neutrophils are recorded as a percentage of total white cells (approximately 100 white blood cells (range 100 to 107) were examined from each slide; frequency in the presence of immature neutrophils was recorded per animal group; dose-related deaths recorded as animal numbers per group of 8 rats.

Compound	Neutrophils	Early neutrophils	Dose-related deaths
Vehicle control	15.50%	0/8	0/8
LMT.2HBr	3.00%*	8/8	2/8
LMT.2HCl	19.90%	2/8	0/8
LMT.2MsOH	18.30%	1/8	0/8

*P < 0.001 compared with control

Although LMT.2HCl and LMT.2MsOH are comparable in the above analysis, there is a distinction in the impurities found in the two salt forms. For LMT.2HCl, the presence of methyl chloride was detected during synthesis and trapped within the product in such a way that it was difficult to remove entirely. By contrast impurities such as ethyl and methyl methanesulfonate (EMS, MMS) could be controlled to much lower levels in the LMT.2MsOH synthetic process

Studies on the hemopoietic system were performed in rat, monkey and minipig.

The lowest doses at which methemoglobinemia was observed were 15 mg MT/kg/day in rats (MTC and LMT.2HBr) or 30 mg MT/kg/day (LMT.2MsOH), 5.3 mg MT/kg/day in primates (MTC), and 10 mg MT/kg/day (LMT.2MsOH and LMT.2HBr) in minipigs.

After the first 28-days of dosing in the 9-month LMT.2MsOH study in minipigs, there are no indications of methemoglobinemia at 3 mg MT/kg/day.

However, as would be expected, as dose levels of MTC, LMT.2HBr or LMT.2MsOH increased, signs of oxidative stress to RBCs emerged in a dose-related fashion, evidenced by increasing levels of methemoglobin and ultimately at doses that were not tolerated, Heinz body (aggregates of denatured, precipitated hemoglobin within red cells) formation.

Example 5

Pharmacokinetics

FIG. 18 shows a comparison of the plasma concentration in pig of the MT moiety over time following dosing of LMT.2HBr, LMT.2HCl and LMT.2MsOH at (two oral doses, 2 and 15 mg/kg).

As can be seen the C_{max} (at T_{max} of 1 hour) for LMT.2MsOH was more than 2-fold greater than that for LMT.2HCl or LMT.2HBr. Thus LMT.2MsOH can provide a more effective exposure to MT than LMT.2HCl or LMT.2HBr.

Example 6

Gastric Irritation Studies

Study (28-day rat with MTC or LMT.2HBr): Incidence and severity of selected microscopic findings in stomach from terminal animals

Incidence and severity of selected findings in sternum, femur, liver and spleen: terminal kill															
Tissue and finding	Level (mg/kg/day)	Male							Female						
		1M	2M	3M	4M	5M	6M	7M	1F	2F	3F	4F	5F	6F	7F
		MTC			LMT.2HBr				MTC			LMT.2HBr			
	No. examined:	5	0	0	5	0	0	5	5	0	0	5	0	0	5
Stomach (non glandular)	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Gastritis	2	—	—	1	—	—	—	—	—	—	—	—	—	—	1
	3	—	—	—	—	—	—	1	—	—	1	—	—	—	1
Inflammatory cell infiltration	1	—	—	1	—	—	—	—	—	—	—	—	—	—	—

Key: “—” = finding not present, 1 = minimal, 2 = slight, 3 = moderate, 4 = moderately severe, 5 = severe

From the above the following can be predicted with 10 per group

Incidence and severity of selected findings in sternum, femur, liver and spleen: terminal kill															
Tissue and finding	Level (mg/kg/day)	Male						Female							
		1M	2M	3M	4M	5M	6M	7M	1F	2F	3F	4F	5F	6F	7F
		MTC			LMT·2HBr			MTC			LMT·2HBr				
	No. examined:	10	0	0	10	0	0	10	10	0	0	10	0	0	10
Stomach (non glandular) Gastritis	1	—	—	—	—	—	—	2	—	—	—	—	—	—	
	2	—	—	—	2	—	—	—	—	—	—	—	—	2	
	3	—	—	—	—	—	—	2	—	—	2	—	—	2	
total					2			4			2			4	

Study (28-day rat study with LMT.2MsOH): Incidence and severity of selected microscopic findings in sternum, liver, spleen and stomach from terminal animals

Incidence and severity of selected findings in sternum, liver and spleen: terminal kill									
Tissue and finding	Level (mg/kg/day)	Males				Females			
		1M	2M	3M	4M	1F	2F	3F	4F
		Grade *				—			
	No. examined:	10	0	0	10	10	0	0	10
Stomach (non glandular) Gastritis	1	—	—	—	—	—	—	—	—
	2	—	—	—	2	—	—	—	2
	3	—	—	—	—	—	—	—	1
total					2				3
Inflammatory cell infiltration	1	—	—	—	4	—	—	—	4
	2	—	—	—	—	—	—	—	1

* Key: “—” = finding not present, 1 = minimal, 2 = slight, 3 = moderate, 4 = moderately severe, 5 = severe

These results show that LMT.2MsOH causes less gastric irritation than LMT.2HBr.

Example 7

Formulations

Formulation Example 1

Preparation of LMTM Tablets Using Direct Compression

Tablets having the following compositions were prepared by a direct compression method:

Ingredient	Tablet strength (LMT mg/tablet)				
	50	75	100	125	150
LMTM	84.43	126.65	168.86	211.08	253.29
Spray-dried mannitol	344.57	302.35	290.14	392.92	425.71
Microcrystalline cellulose (Avicel PH102 or PH112)	50.00	75.00	95.00	125.00	150.00
Crospovidone (crosslinked polyvinylpyrrolidone)	15.00	15.00	15.00	15.00	15.00

-continued

Ingredient	Tablet strength (LMT mg/tablet)				
	50	75	100	125	150
Magnesium stearate	6.00	6.00	6.00	6.00	6.00
Total tablet core weight	500.00	525.00	575.00	750.00	850.00

The LMTM, spray-dried mannitol, microcrystalline cellulose, crospovidone and magnesium stearate were blended in a tumbling blender, and then compressed using a tableting machine.

The tablet cores were then film coated with an aqueous suspension of Opadry* blue (*registered trademark of Colcon for a range of film coating materials).

Formulation Example 2

Preparation of LMTM Tablets Using Dry Granulation (Roller Compaction)

Tablets having the following compositions were prepared by a dry granulation method:

81

Ingredient	Tablet strength (LMT mg/tablet)				
	50	75	100	125	150
	(mg/tablet)				
LMTM	84.43	126.65	168.86	211.08	253.29
Spray-dried mannitol	344.57	302.35	290.14	392.92	425.71
Microcrystalline cellulose (Avicel PH102 or PH112)	50.00	75.00	95.00	125.00	150.00
Crospovidone (crosslinked polyvinylpyrrolidone)	15.00	15.00	15.00	15.00	15.00
Magnesium stearate	6.00	6.00	6.00	6.00	6.00
Total tablet core weight	500.00	525.00	575.00	750.00	850.00

The LMTM, spray-dried mannitol, microcrystalline cellulose, crospovidone and magnesium stearate were blended in a tumbling blender. The mix was then dry granulated using a roller compactor and then milled with an oscillating granulator using a suitable screen. In this case, half of magnesium stearate was used prior to roller compaction and half of the magnesium stearate was then added to the granulation and blended prior to compression on a conventional tableting machine.

The tablet cores were then film coated with an aqueous suspension of Opadry* blue (*registered trademark of Colorcon for a range of film coating materials).

Formulation Example 3

Preparation of LMTM Tablets by Dry Granulation (Slugging)

Tablets having the following compositions were prepared by a further dry granulation method.

Ingredient	Tablet strength (LMT mg/tablet)				
	50	75	100	125	150
	(mg/tablet)				
LMTM	84.43	126.65	168.86	211.08	253.29
Spray-dried mannitol	344.57	302.35	290.14	392.92	425.71
Microcrystalline cellulose (Avicel PH102 or PH112)	50.00	75.00	95.00	125.00	150.00
Crospovidone (crosslinked polyvinylpyrrolidone)	15.00	15.00	15.00	15.00	15.00
Magnesium stearate	6.00	6.00	6.00	6.00	6.00
Total tablet core weight	500.00	525.00	575.00	750.00	850.00

The LMTM and excipients were blended in a tumbling blender, and then compressed to produce slugs (plain, flat faced tablets) using a tableting machine.

82

The slugs were then milled using an oscillating granulator fitted with a 20 mesh screen.

In this example, half of magnesium stearate was used prior to slugging and then half of the magnesium stearate added to the granulation and blended prior to compression on a conventional tableting machine.

The tablet cores were then film coated with an aqueous suspension of Opadry* blue (*registered trademark of Colorcon for a range of film coating materials).

Formulation Example 4

Preparation of LMTM Tablets by Wet Granulation of Excipients and Incorporation of LMTM Extra-Granularly

Tablets having the following compositions were prepared by a wet granulation method:

Ingredient	Tablet strength (LMT mg/tablet)				
	50	75	100	125	150
	(mg/tablet)				
LMTM	84.43	126.65	168.86	211.08	253.29
Mannitol	334.57	292.35	280.14	380.92	413.71
Microcrystalline cellulose (Avicel PH102)	50.00	75.00	95.00	125.00	150.00
Crospovidone (crosslinked polyvinylpyrrolidone)	15.00	15.00	15.00	15.00	15.00
Polyvinylpyrrolidone	10.00	10.00	10.00	12.00	12.00
Magnesium stearate	6.00	6.00	6.00	6.00	6.00
Total tablet core weight	500.00	525.00	575.00	750.00	850.00

The mannitol, crospovidone (a third of the total) and microcrystalline cellulose were blended in a tumbling blender. The blended material was then granulated using a solution of PVP in water. The wet mass was dried in a fluid bed dryer and then milled using an oscillating granulator fitted with a suitable screen.

The milled material was then blended with the remainder of the crospovidone and magnesium stearate, and the LMTM, prior to compression on a conventional tablet machine. The tablet cores were then film coated with an aqueous suspension of Opadry* blue (*registered trademark of Colorcon for a range of film coating materials).

Formulation Example 5

Preparation of LMTM Capsules

Ingredient	Capsule strength (LMT mg/capsule)					
	50	75	100	125	150	200
	mg/capsule					
LMTM	84.43	126.65	168.86	211.08	253.29	337.72
Spray-dried mannitol	191.07	148.85	116.64	79.42	42.21	37.78
Crospovidone (crosslinked polyvinylpyrrolidone)	3.00	3.00	3.00	3.00	3.00	3.00
Magnesium stearate	1.50	1.50	1.50	1.50	1.50	1.50
Total capsule fill weight	280.00	280.00	290.00	295.00	300.00	380.00

83

The LMTM and the excipients were blended in a tumbling blender. The resulting drug blends were filled into capsules (50, 75, 100, 125 and 150 mg formulations into size 1 capsules and the 200 mg formulation into size 0 capsules) using a capsule filling machine. Both gelatine capsules and HPMC capsules were prepared.

Formulation Example 6

Results of Stability Testing LMTM 75 mg Film Coated Tablets

Test	Time Point (months)	Storage Location	
		25° C./60% RH	40° C./75% RH
Assay as % LMT free base	0	102.2	102.2
	1	101.5	94.8
	3	100.0	94.2
	6	96.4	not done
	9	95.6	not done
	12	96.0	not done

Formulation Example 7

Results of Stability Testing LMTM 100 mg Film Coated Tablets

Test	Time Point (months)	Storage Location	
		25° C./60% RH	40° C./75% RH
Assay as % LMT free base	0	101.0	101.0
	1	96.7	93.7
	3	95.9	92.8
	6	96.0	94.2
	9	97.1	not done
	12	96.8	not done

Formulation Example 8

Results of Stability Testing LMTM 75 mg Film Coated Tablets

Test	Time Point (months)	Storage Location	
		25° C./60% RH	40° C./75% RH
% MT formed	0	2.16	2.06
	1	2.05	3.79
	3	2.19	4.51
	6	2.83	5.71
	9	3.53	not done
	12	3.28	not done

84

Formulation Example 9

Results of Stability Testing LMTM 100 mg Film Coated Tablets

Test	Time Point (months)	Storage Location	
		25° C./60% RH	40° C./75% RH
% MT formed	0	2.07	2.07
	1	1.78	3.27
	3	1.81	4.92
	6	2.51	5.07
	9	2.72	
	12	2.88	

Formulation Example 10

Material	mg/tablet (as LMT)	mg/tablet (as LMTM)	% (core only)
Tablet Core			
LMTB (batch number 0802100070)	100.00	163.03	32.61
Spray Dried Mannitol (Pearlitol 200 SD)	329.00	265.97	53.19
Microcrystalline cellulose	50.00	50.00	10.00
Crospovidone	15.00	15.00	3.00
Magnesium Stearate	6.00	6.00	1.20
Tablet Core Total	500.00	500.00	100.00
Film Coat			
Polyvinyl Alcohol (part hydrolysed)	8.80	8.80	
Talc	4.00	4.00	
Titanium Dioxide	3.10	3.10	
Macrogol PEG 3350	2.47	2.47	
Lecithin (soya)	0.70	0.70	
Iron Oxide Yellow	0.47	0.47	
Indigo Carmine	0.45	0.45	
Aluminium Lake			
Total Film Coated Tablet	520.00	520.00	
Manufacturer	Piramal, Morpeth, UK		
Tablet Core Batch Number	A02581		
Date of Manufacture	15 TH October 2009		

Tablets having the above formulation were prepared by a direct compression method as described above and then film coated (see Formulation Example 1).

Formulation Example 11

LMTM 75 mg Film-Coated Tablets

Material	mg/tablet (theoretical)	mg/tablet (actual)	% (core only)
Tablet Core			
LMTM (batch numbers)	75.00	126.80	24.15

85

-continued

Material	mg/tablet (theoretical)	mg/tablet (actual)	% (core only)
800225510 & 80224450)			
Spray Dried Mannitol (Pearlitol 200 SD)	354.00	302.20	57.56
Microcrystalline cellulose	75.00	75.00	14.29
Crospovidone	15.00	15.00	2.86
Magnesium stearate	6.00	6.00	1.14
Tablet core total	525.00	525.00	100.00
Film Coat			
Polyvinyl Alcohol (part hydrolysed)	13.86	13.86	
Talc	6.30	6.30	
Titanium Dioxide	4.89	4.89	
Macrogol PEG 3350	3.89	3.89	
Lecithin (soya)	1.10	1.10	
Iron Oxide Yellow	0.75	0.75	
Indigo Carmine	0.71	0.71	
Aluminium Lake			
Total Film Coated Tablet	556.5	556.5	
Manufacturer	Piramal, Morpeth, UK		
Tablet Core Batch Number	A04827		
Date of Manufacture	5 th August 2010		

Tablets having the above formulation were prepared by a direct compression method as described above and then film coated (see Formulation Example 1).

Formulation Example 12

Dissolution Studies

LMTB film-coated tablets (3×100 mg) and LMTM tablets (4×75 mg), prepared as in Formulation Examples 10 and 11, were stirred (see FIG. 19) at a paddle speed of 50 rpm and the dissolution rate was assessed, using a standard pharmacopeial method (USP 34) and the conditions specified below.

Instrumental Conditions

Parameter	Condition
Media	0.1M HCl (Degassed with He purging)
Media Volume	1000 ml, 6 vessels
Dissolved Oxygen	<3.00 ppm
Bath Temperature	37° C. ± 0.5° C.
Paddles	Teflon Coated
Paddle Speed	50 rpm
Pull Volume	10 ml - no media replacement
Filter	HDPE 10 µm
Time points	10, 15, 30 and 45 min
Vessels	6 (protected from light)
λ_{max} LMT	255 nm
Sample Working Concentration (µg/ml)	ca 5 µg/ml (as free base) LMT
Standard Working Concentration (µg/ml)	ca 5 µg/ml (as free base) LMT

(Q=75% at 45 mins. For S1, 6 of 6 tablets not less than 80% dissolution at 45 minutes).

86

Results are shown in the following tables.

LMTM (4×75 mg; Batch No: A04827)

Dissolution (% dissolved):

Vessel	T = 10 min	T = 15 min	T = 30 min	T = 45 min
1	94	95	97	99
2	90	91	94	95
3	94	94	97	97
4	95	94	97	97
5	92	92	94	94
6	93	92	96	97
Mean	93	93	96	97

LMTB (3×100 mg; Batch No: A02581)

Dissolution (% dissolved):

Vessel	T = 10 min	T = 15 min	T = 30 min	T = 45 min
1	91	95	96	96
2	96	100	99	99
3	95	98	98	99
4	93	95	96	96
5	96	98	99	100
6	98	102	102	102
Mean	95	98	98	99

Tablets which had been stored for varying periods of time, under normal (25° C./60% RH) or 'stressed' conditions (40° C./75% RH), were also tested using the same method.

Results are shown in the tables below.

LMTM (4×75 mg; Batch No: A04827)—stored at 25° C./60% RH

Dissolution (% dissolved):

Storage Time	Vessel	T = 10 min	T = 15 min	T = 30 min	T = 45 min
1 month	1	97	97	97	99
	2	96	98	101	101
	3	98	99	102	102
	4	95	97	98	100
	5	97	98	101	101
	6	98	98	100	101
3 months	Mean	97	98	100	101
	1	91	93	95	97
	2	92	95	96	96
	3	93	94	95	97
	4	92	93	96	96
	5	93	94	95	96
6 months	6	90	91	94	95
	Mean	92	93	95	96
	1	89	89	90	91
	2	91	90	93	94
	3	98	97	98	98
	4	97	97	99	99
9 months	5	94	94	96	96
	6	88	90	93	93
	Mean	93	93	95	95
	1	92	93	92	94
	2	90	94	95	97
	3	86	91	90	93
60	4	85	91	96	94
	5	90	85	94	94
	6	92	96	94	96
	Mean	89	92	93	94

LMTM (4×75 mg; Batch No: A04827)—stored at 40° C./75% RH

Dissolution (% dissolved):

Storage Time	Vessel	T = 10 min	T = 15 min	T = 30 min	T = 45 min
1 month	1	94	95	97	98
	2	94	96	96	97
	3	94	96	94	96
	4	94	95	95	95
	5	100	102	103	101
	6	93	94	96	97
	Mean	95	96	97	97
3 months	1	92	93	95	96
	2	93	94	95	97
	3	89	91	92	92
	4	89	89	89	91
	5	93	95	96	97
	6	93	95	98	97
	Mean	91	93	94	95
6 months	1	69	84	92	94
	2	93	94	97	91
	3	64	85	92	94
	4	74	89	92	94
	5	91	95	95	96
	6	73	90	93	94
	Mean	77	89	94	94

LMTB (3×100 mg; Batch No: A02581)—stored at 25° C./60% RH

Dissolution (% dissolved):

Storage Time	Vessel	T = 10 min	T = 15 min	T = 30 min	T = 45 min
3 weeks	1	96	98	98	98
	2	94	97	97	98
	3	94	97	97	97
	4	98	100	101	101
	5	92	94	95	95
	6	92	95	97	97
	Mean	94	97	98	98
3 months	1	89	92	92	92
	2	89	92	93	92
	3	93	96	96	96
	4	95	98	99	98
	5	95	96	96	96
	6	96	99	98	97
	Mean	93	96	96	95
6 months	1	96	97	96	97
	2	95	101	100	101
	3	95	97	96	97
	4	95	95	95	96
	5	96	98	99	99
	6	95	94	94	96
	Mean	95	97	97	98
9 months	1	87	91	93	91
	2	88	92	94	92
	3	90	93	91	92
	4	91	95	93	94
	5	91	93	93	92
	6	94	95	95	93
	Mean	90	93	93	92
12 months	1	92	97	98	97
	2	91	92	92	92
	3	95	96	95	96
	4	94	95	95	95
	5	89	89	89	89
	6	97	98	98	98
	Mean	93	94	95	94

LMTB (3×100 mg; Batch No: A02581)—stored at 40° C./75% RH

Dissolution (% dissolved):

Storage Time	Vessel	T = 10 min	T = 15 min	T = 30 min	T = 45 min
3 weeks	1	94	98	99	98
	2	96	100	100	101
	3	94	97	96	97
	4	94	98	98	98
	5	95	97	98	98
	6	95	97	98	97
	Mean	95	98	98	98
3 months	1	92	93	94	93
	2	93	97	97	97
	3	90	92	92	92
	4	84	89	94	94
	5	84	97	97	97
	6	93	94	93	94
	Mean	89	94	95	95
6 months	1	8	72	96	96
	2	48	82	95	96
	3	91	93	94	94
	4	94	98	98	99
	5	13	71	93	93
	6	74	87	92	93
	Mean	55	84	95	95

25 Annex—Crystallographic Data

Crystallographic Data for LMT·EDSA (FIG. 17a):

TABLE 1

Crystal data and structure refinement for LMT·EDSA.	
Identification code	6408CM136
Empirical formula	C ₁₈ H ₂₇ N ₃ O ₇ S ₃
Formula weight	493.62
Temperature	100(2) K
Wavelength	0.71073 Å
Crystal system	Monoclinic
Space group	C2/c
Unit cell dimensions	a = 18.2832(3) Å α = 90°. b = 11.8667(3) Å β = 114.1990(10)°. c = 10.9539(2) Å γ = 90°.
Volume	2167.74(8) Å ³
Z	4
Density (calculated)	1.519 Mg/m ³
Absorption coefficient	0.389 mm ⁻¹
F(000)	1048
Crystal size	0.28 × 0.21 × 0.18 mm ³
Theta range for data collection	2.11 to 27.51°.
Index ranges	-23 ≤ h ≤ 23, -15 ≤ k ≤ 15, -14 ≤ l ≤ 14
Reflections collected	25214
Independent reflections	2487 [R(int) = 0.0486]
Completeness to theta = 25.00°	99.9%
Absorption correction	Semi-empirical from equivalents
Max. and min. transmission	0.9333 and 0.8989
Refinement method	Full-matrix least-squares on F ²
Data/restraints/parameters	2487/0/144
Goodness-of-fit on F ²	1.080
Final R indices [I > 2σ(I)]	R1 = 0.0315, wR2 = 0.0906
R indices (all data)	R1 = 0.0336, wR2 = 0.0925
Largest diff. peak and hole	0.333 and -0.654 e · Å ⁻³

TABLE 2

Atomic coordinates (×10 ⁴) and equivalent isotropic displacement parameters (Å ² × 10 ³) for LMT·EDSA. U(eq) is defined as one third of the trace of the orthogonalized U ^{ij} tensor.				
	x	y	z	U(eq)
S(1)	10000	2270(1)	12500	19(1)
S(2)	3802(1)	204(1)	9934(1)	11(1)
N(1)	10000	-370(1)	12500	18(1)

TABLE 2-continued

Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for LMT·EDSA. U(eq) is defined as one third of the trace of the orthogonalized U^{ij} tensor.				
	x	y	z	U(eq)
N(2)	7750(1)	1619(1)	7782(1)	12(1)
O(1)	3943(1)	-435(1)	11137(1)	18(1)
O(2)	3830(1)	1425(1)	10131(1)	17(1)
O(3)	3063(1)	-144(1)	8793(1)	15(1)
C(1)	9411(1)	1332(1)	11218(1)	13(1)
C(2)	9493(1)	154(1)	11332(1)	14(1)
C(3)	9040(1)	-512(1)	10228(1)	16(1)
C(4)	8481(1)	-30(1)	9061(1)	16(1)
C(5)	8383(1)	1126(1)	8996(1)	13(1)
C(6)	8852(1)	1814(1)	10051(1)	13(1)
C(8)	7127(1)	2225(1)	8087(1)	16(1)
C(9)	8070(1)	2352(1)	7003(1)	17(1)
C(10)	4593(1)	-141(1)	9448(1)	13(1)
O(1S)	5000	2293(1)	2500	24(1)

TABLE 3

Bond lengths [\AA] and angles [$^\circ$] for LMT·EDSA.	
S(1)—C(1)	1.7696(13)
S(1)—C(1)#1	1.7696(13)
S(2)—O(1)	1.4488(10)
S(2)—O(2)	1.4629(10)
S(2)—O(3)	1.4747(10)
S(2)—C(10)	1.7802(13)
N(1)—C(2)	1.3826(15)
N(1)—C(2)#1	1.3826(15)
N(1)—H(1)	0.8800
N(2)—C(5)	1.4785(16)
N(2)—C(9)	1.4959(17)
N(2)—C(8)	1.4970(17)
N(2)—H(2)	0.9300
C(1)—C(6)	1.3913(17)
C(1)—C(2)	1.4049(18)
C(2)—C(3)	1.3972(19)
C(3)—C(4)	1.3908(19)
C(3)—H(3)	0.9500
C(4)—C(5)	1.3804(19)
C(4)—H(4)	0.9500
C(5)—C(6)	1.3881(18)
C(6)—H(6)	0.9500
C(8)—H(8A)	0.9800
C(8)—H(8B)	0.9800
C(8)—H(8C)	0.9800
C(9)—H(9A)	0.9800
C(9)—H(9B)	0.9800
C(9)—H(9C)	0.9800
C(10)—C(10)#2	1.522(2)
C(10)—H(10A)	0.9900
C(10)—H(10B)	0.9900
O(1S)—H(1O1)	0.7486
O(1S)—H(2O1)	0.9717
C(1)—S(1)—C(1)#1	102.05(9)
O(1)—S(2)—O(2)	113.66(6)
O(1)—S(2)—O(3)	112.60(6)
O(2)—S(2)—O(3)	111.52(6)
O(1)—S(2)—C(10)	106.77(6)
O(2)—S(2)—C(10)	106.71(6)
O(3)—S(2)—C(10)	104.89(6)
C(2)—N(1)—C(2)#1	126.50(17)
C(2)—N(1)—H(1)	116.7
C(2)#1—N(1)—H(1)	116.7
C(5)—N(2)—C(9)	113.51(10)
C(5)—N(2)—C(8)	112.13(10)
C(9)—N(2)—C(8)	111.06(11)
C(5)—N(2)—H(2)	106.5
C(9)—N(2)—H(2)	106.5
C(8)—N(2)—H(2)	106.5
C(6)—C(1)—C(2)	120.17(12)
C(6)—C(1)—S(1)	116.69(10)
C(2)—C(1)—S(1)	123.14(10)

TABLE 3-continued

Bond lengths [\AA] and angles [$^\circ$] for LMT·EDSA.		
5	N(1)—C(2)—C(3)	118.76(13)
	N(1)—C(2)—C(1)	122.44(13)
	C(3)—C(2)—C(1)	118.79(12)
	C(4)—C(3)—C(2)	120.95(13)
	C(4)—C(3)—H(3)	119.5
10	C(2)—C(3)—H(3)	119.5
	C(5)—C(4)—C(3)	119.15(13)
	C(5)—C(4)—H(4)	120.4
	C(3)—C(4)—H(4)	120.4
	C(4)—C(5)—C(6)	121.24(12)
15	C(4)—C(5)—N(2)	118.50(12)
	C(6)—C(5)—N(2)	120.25(12)
	C(5)—C(6)—C(1)	119.54(12)
	C(5)—C(6)—H(6)	120.2
	C(1)—C(6)—H(6)	120.2
20	N(2)—C(8)—H(8A)	109.5
	N(2)—C(8)—H(8B)	109.5
	H(8A)—C(8)—H(8B)	109.5
	N(2)—C(8)—H(8C)	109.5
	H(8A)—C(8)—H(8C)	109.5
25	H(8B)—C(8)—H(8C)	109.5
	N(2)—C(9)—H(9A)	109.5
	N(2)—C(9)—H(9B)	109.5
	H(9A)—C(9)—H(9B)	109.5
	N(2)—C(9)—H(9C)	109.5
30	H(9A)—C(9)—H(9C)	109.5
	H(9B)—C(9)—H(9C)	109.5
	C(10)#2—C(10)—S(2)	111.21(12)
	C(10)#2—C(10)—H(10A)	109.4
	S(2)—C(10)—H(10A)	109.4
35	C(10)#2—C(10)—H(10B)	109.4
	S(2)—C(10)—H(10B)	109.4
	H(10A)—C(10)—H(10B)	108.0
	H(1O1)—O(1S)—H(2O1)	100.8

Symmetry transformations used to generate equivalent atoms:

$$\#1 -x+2, y, -z+5/2 \quad \#2 -x+1, -y, -z+2$$

TABLE 4

Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for LMT·EDSA. The anisotropic displacement factor exponent takes the form: $-2p^2[h^2 a^*2U^{11} + \dots + 2hk a^* b^* U^{12}]$							
	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}	
50	S(1)	20(1)	10(1)	14(1)	0	-6(1)	0
	S(2)	10(1)	12(1)	9(1)	0(1)	2(1)	0(1)
	N(1)	22(1)	9(1)	14(1)	0	-3(1)	0
	N(2)	12(1)	13(1)	9(1)	0(1)	2(1)	-1(1)
55	O(1)	16(1)	22(1)	13(1)	5(1)	5(1)	0(1)
	O(2)	17(1)	13(1)	17(1)	-2(1)	4(1)	0(1)
	O(3)	11(1)	16(1)	13(1)	-1(1)	0(1)	-1(1)
	C(1)	12(1)	13(1)	11(1)	-1(1)	2(1)	-2(1)
	C(2)	12(1)	13(1)	13(1)	0(1)	2(1)	0(1)
	C(3)	19(1)	11(1)	14(1)	-2(1)	3(1)	-1(1)
60	C(4)	16(1)	15(1)	12(1)	-2(1)	2(1)	-1(1)
	C(5)	12(1)	15(1)	10(1)	1(1)	2(1)	0(1)
	C(6)	13(1)	12(1)	12(1)	0(1)	4(1)	0(1)
	C(8)	13(1)	18(1)	15(1)	1(1)	5(1)	1(1)
	C(9)	18(1)	22(1)	12(1)	2(1)	6(1)	-2(1)
	C(10)	11(1)	17(1)	10(1)	-2(1)	3(1)	-1(1)
65	O(1S)	25(1)	14(1)	18(1)	0	-7(1)	0

91

TABLE 5

Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for LMT·EDSA.				
	x	y	z	U(eq)
H(1)	10000	-1111	12500	22
H(2)	7492	1019	7227	15
H(3)	9114	-1305	10275	19
H(4)	8171	-489	8319	19
H(6)	8791	2610	9978	15
H(8A)	6937	1734	8616	23
H(8B)	6675	2426	7248	23
H(8C)	7359	2911	8596	23
H(9A)	8321	3023	7531	26
H(9B)	7629	2582	6164	26
H(9C)	8469	1933	6802	26
H(10A)	4571	-955	9239	16
H(10B)	4521	284	8628	16
H(10I)	5146	2050	3190	29
H(20I)	4556	1790	2015	29

Crystallographic Data for LMT·2EsOH (FIG. 17b)

TABLE 1

Crystal data and structure refinement for LMT·2EsOH.				
Identification code	6408cm173c_0m			
Empirical formula	$\text{C}_{20}\text{H}_{31}\text{N}_3\text{O}_6\text{S}_3$			
Formula weight	505.66			
Temperature	100(2) K			
Wavelength	0.71073 \AA			
Crystal system	Monoclinic			
Space group	C2/c			
Unit cell dimensions	a = 40.8384(12) \AA	$\alpha = 90^\circ$.		
	b = 25.2658(7) \AA	$\beta = 115.4540(10)^\circ$.		
	c = 20.3833(6) \AA	$\gamma = 90^\circ$.		
Volume	18990.2(9) \AA ³			
Z	32			
Density (calculated)	1.415 Mg/m ³			
Absorption coefficient	0.354 mm ⁻¹			
F(000)	8576			
Crystal size	0.32 × 0.24 × 0.18 mm ³			
Theta range for data collection	0.98 to 25.00°.			
Index ranges	-48 ≤ h ≤ 48, -29 ≤ k ≤ 30, -24 ≤ l ≤ 23			
Reflections collected	108984			
Independent reflections	16707 [R(int) = 0.0912]			
Completeness to theta = 25.00°	99.9%			
Absorption correction	Semi-empirical from equivalents			
Max. and min. transmission	0.9391 and 0.8952			
Refinement method	Full-matrix least-squares on F ²			
Data/restraints/parameters	16707/25/1205			
Goodness-of-fit on F ²	1.085			
Final R indices	R1 = 0.0628, wR2 = 0.1638			
[I > 2sigma(I)]				
R indices (all data)	R1 = 0.0986, wR2 = 0.1918			
Largest diff. peak and hole	2.683 and -0.811 e · \AA ⁻³			

TABLE 2

Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for LMT·2EsOH. U(eq) is defined as one third of the trace of the orthogonalized U^{ij} tensor.				
	x	y	z	U(eq)
S(1A)	256(1)	3950(1)	1866(1)	69(1)
N(1A)	-401(1)	3176(1)	1220(2)	29(1)
N(2A)	980(1)	2230(1)	2700(2)	25(1)
N(3A)	-743(1)	5348(1)	744(2)	25(1)
C(1A)	-207(1)	4102(2)	1408(3)	33(1)
C(2A)	-297(1)	4633(2)	1295(3)	32(1)
C(3A)	-652(1)	4786(2)	916(2)	24(1)
C(4A)	-926(1)	4413(2)	664(2)	27(1)

92

TABLE 2-continued

Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for LMT·2EsOH. U(eq) is defined as one third of the trace of the orthogonalized U^{ij} tensor.				
	x	y	z	U(eq)
C(5A)	-836(1)	3880(2)	784(3)	28(1)
C(6A)	-479(1)	3714(2)	1142(2)	25(1)
C(7A)	-60(1)	2948(2)	1584(2)	25(1)
C(8A)	-27(1)	2402(2)	1631(2)	24(1)
C(9A)	308(1)	2154(2)	1988(2)	25(1)
C(10A)	614(1)	2468(2)	2318(2)	24(1)
C(11A)	588(1)	3011(2)	2281(3)	29(1)
C(12A)	254(1)	3255(2)	1917(3)	32(1)
C(13A)	1017(1)	1896(2)	3334(3)	36(1)
C(14A)	1092(1)	1933(2)	2196(3)	41(1)
C(15A)	-662(2)	5526(2)	130(3)	50(2)
C(16A)	-570(1)	5712(2)	1375(3)	33(1)
S(1B)	2816(1)	1091(1)	1548(1)	26(1)
N(1B)	2213(1)	258(1)	1144(2)	21(1)
N(2B)	3639(1)	-517(1)	2667(2)	27(1)
N(3B)	1743(1)	2378(1)	956(2)	22(1)
C(1B)	2355(1)	1198(2)	1329(2)	19(1)
C(2B)	2237(1)	1719(2)	1291(2)	22(1)
C(3B)	1873(1)	1825(2)	1032(2)	21(1)
C(4B)	1620(1)	1417(2)	832(2)	22(1)
C(5B)	1738(1)	896(2)	890(2)	21(1)
C(6B)	2104(1)	778(2)	1128(2)	21(1)
C(7B)	2566(1)	73(2)	1537(2)	22(1)
C(8B)	2628(1)	-468(2)	1684(2)	23(1)
C(9B)	2974(1)	-666(2)	2055(2)	24(1)
C(10B)	3263(1)	-320(2)	2304(2)	24(1)
C(11B)	3213(1)	220(2)	2178(2)	23(1)
C(12B)	2866(1)	417(2)	1788(2)	22(1)
C(13B)	3693(2)	-973(2)	3183(3)	38(1)
C(14B)	3785(1)	-651(2)	2126(2)	29(1)
C(15B)	1835(1)	2684(2)	426(3)	34(1)
C(16B)	1872(1)	2660(2)	1668(3)	35(1)
S(1C)	5390(1)	3672(1)	1826(1)	25(1)
N(1C)	4792(1)	2826(1)	1436(2)	25(1)
N(2C)	6224(1)	2099(1)	3029(2)	20(1)
N(3C)	4310(1)	4945(1)	1101(2)	26(1)
C(1C)	4925(1)	3774(2)	1581(2)	20(1)
C(2C)	4803(1)	4289(2)	1510(2)	21(1)
C(3C)	4436(1)	4388(2)	1235(2)	21(1)
C(4C)	4184(1)	3984(2)	1037(2)	25(1)
C(5C)	4307(1)	3464(2)	1126(2)	23(1)
C(6C)	4676(1)	3350(2)	1388(2)	21(1)
C(7C)	5145(1)	2651(2)	1833(2)	22(1)
C(8C)	5213(1)	2113(2)	1990(2)	22(1)
C(9C)	5559(1)	1925(2)	2384(2)	21(1)
C(10C)	5847(1)	2278(2)	2639(2)	20(1)
C(11C)	5788(1)	2813(2)	2493(2)	21(1)
C(12C)	5443(1)	3002(2)	2087(2)	20(1)
C(13C)	6374(1)	1945(2)	2499(2)	26(1)
C(14C)	6284(1)	1670(2)	3576(2)	24(1)
C(15C)	4375(2)	5182(2)	496(3)	47(1)
C(16C)	4468(2)	5280(2)	1771(3)	47(2)
S(1D)	7907(1)	1349(1)	2060(1)	32(1)
N(1D)	7269(1)	547(1)	1633(2)	29(1)
N(2D)	8670(1)	-331(2)	2894(2)	28(1)
N(3D)	6848(1)	2694(1)	1136(2)	27(1)
C(1D)	7440(1)	1484(2)	1723(2)	25(1)
C(2D)	7333(1)	2011(2)	1602(2)	27(1)
C(3D)	6969(1)	2136(2)	1304(2)	26(1)
C(4D)	6709(1)	1744(2)	1117(3)	32(1)
C(5D)	6818(1)	1220(2)	1238(3)	32(1)
C(6D)	7179(1)	1080(2)	1536(2)	24(1)
C(7D)	7614(1)	338(2)	1971(2)	22(1)
C(8D)	7660(1)	-209(2)	2063(2)	24(1)
C(9D)	8001(1)	-439(2)	2370(2)	25(1)
C(10D)	8302(1)	-111(2)	2608(2)	24(1)
C(11D)	8268(1)	437(2)	2537(2)	26(1)
C(12D)	7925(1)	662(2)	2212(2)	23(1)
C(13D)	8743(1)	-762(2)	3435(3)	43(1)
C(14D)	8764(1)	-504(2)	2294(3)	31(1)
C(15D)	6807(2)	2848(2)	397(3)	41(1)
C(16D)	7078(1)	3086(2)	1700(3)	36(1)
S(2)	525(1)	500(1)	1331(1)	35(1)
S(3)	827(1)	2850(1)	152(1)	25(1)

TABLE 2-continued

Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for LMT·2EsOH. U(eq) is defined as one third of the trace of the orthogonalized U^i tensor.				
	x	y	z	U(eq)
S(4)	1743(1)	4426(1)	389(1)	29(1)
S(5)	3048(1)	3051(1)	1414(1)	30(1)
S(6)	5663(1)	497(1)	1723(1)	33(1)
S(7)	5874(1)	2562(1)	455(1)	34(1)
S(8)	8160(1)	2896(1)	1391(1)	23(1)
S(9)	8481(1)	615(1)	291(1)	26(1)
O(1)	619(4)	884(5)	1948(7)	56(2)
O(2)	392(10)	52(10)	1452(19)	70(6)
O(3)	886(4)	594(6)	1706(8)	55(2)
O(1')	342(2)	769(3)	1689(4)	61(2)
O(2')	419(6)	-77(6)	1226(11)	70(5)
O(3')	893(2)	480(4)	1357(5)	58(2)
O(4)	1009(1)	2335(1)	358(2)	35(1)
O(5)	1031(1)	3269(1)	635(2)	41(1)
O(6)	718(1)	2970(1)	-613(2)	27(1)
O(7)	1911(1)	3966(1)	826(2)	45(1)
O(8)	1422(1)	4299(1)	-282(2)	45(1)
O(9)	1667(1)	4851(1)	790(2)	30(1)
O(10)	3321(1)	2861(1)	1185(2)	41(1)
O(11)	2835(1)	2637(2)	1525(3)	68(1)
O(12)	3217(1)	3392(1)	2047(2)	46(1)
O(13)	5318(1)	542(2)	1772(2)	56(1)
O(14)	5728(1)	-26(1)	1506(2)	60(1)
O(15)	5964(1)	678(1)	2388(2)	42(1)
O(16)	5932(1)	2520(1)	-212(2)	32(1)
O(17)	5774(1)	2071(2)	681(2)	57(1)
O(18)	6182(1)	2836(1)	1038(2)	39(1)
O(19)	8126(1)	2360(1)	1113(2)	31(1)
O(20)	8540(1)	3032(1)	1876(2)	35(1)
O(21)	7918(1)	3019(1)	1720(2)	31(1)
O(22)	8264(1)	640(1)	-494(2)	26(1)
O(23)	8590(1)	62(1)	534(2)	31(1)
O(24)	8786(1)	983(1)	553(2)	33(1)
C(1S)	298(2)	782(2)	463(4)	62(2)
C(2S)	387(2)	1337(2)	396(3)	45(1)
C(3S)	2064(1)	4682(2)	95(3)	30(1)
C(4S)	1928(1)	5196(2)	-337(3)	39(1)
C(5S)	2748(1)	3452(2)	689(3)	37(1)
C(6S)	2934(1)	3925(2)	551(3)	32(1)
C(7S)	5504(1)	2984(2)	239(3)	39(1)
C(8S)	5570(2)	3516(2)	-63(4)	54(2)
C(9S)	5626(2)	921(2)	1016(3)	38(1)
C(10S)	5990(3)	881(4)	989(5)	100(3)
C(11S)	8047(1)	3340(2)	650(3)	28(1)
C(12S)	7648(1)	3320(2)	118(3)	38(1)
C(13S)	8200(1)	808(2)	708(3)	27(1)
C(14S)	7876(1)	454(2)	490(3)	34(1)
C(15S)	414(1)	2769(2)	254(3)	36(1)
C(16S)	186(2)	3261(2)	66(3)	49(2)

TABLE 3

Bond lengths [\AA] and angles [$^\circ$] for LMT·2EsOH.	
S(1A)—C(1A)	1.756(5)
S(1A)—C(12A)	1.759(5)
N(1A)—C(7A)	1.390(6)
N(1A)—C(6A)	1.389(6)
N(1A)—H(1AA)	0.8800
N(2A)—C(10A)	1.485(5)
N(2A)—C(14A)	1.492(6)
N(2A)—C(13A)	1.495(6)
N(2A)—H(2AA)	0.9300
N(3A)—C(3A)	1.474(5)
N(3A)—C(16A)	1.489(6)
N(3A)—C(15A)	1.494(6)
N(3A)—H(3A)	0.9300
C(1A)—C(2A)	1.384(6)
C(1A)—C(6A)	1.404(6)
C(2A)—C(3A)	1.371(6)
C(2A)—H(2A)	0.9500

TABLE 3-continued

Bond lengths [\AA] and angles [$^\circ$] for LMT·2EsOH.	
C(3A)—C(4A)	1.384(6)
C(4A)—C(5A)	1.389(6)
C(4A)—H(4A)	0.9500
C(5A)—C(6A)	1.384(6)
C(5A)—H(5A)	0.9500
C(7A)—C(8A)	1.387(6)
C(7A)—C(12A)	1.399(6)
C(8A)—C(9A)	1.391(6)
C(8A)—H(8A)	0.9500
C(9A)—C(10A)	1.384(6)
C(9A)—H(9A)	0.9500
C(10A)—C(11A)	1.376(6)
C(11A)—C(12A)	1.388(6)
C(11A)—H(11A)	0.9500
C(13A)—H(13A)	0.9800
C(13A)—H(13B)	0.9800
C(13A)—H(13C)	0.9800
C(14A)—H(14A)	0.9800
C(14A)—H(14B)	0.9800
C(14A)—H(14C)	0.9800
C(15A)—H(15A)	0.9800
C(15A)—H(15B)	0.9800
C(15A)—H(15C)	0.9800
C(16A)—H(16A)	0.9800
C(16A)—H(16B)	0.9800
C(16A)—H(16C)	0.9800
S(1B)—C(12B)	1.761(4)
S(1B)—C(1B)	1.762(4)
N(1B)—C(6B)	1.385(5)
N(1B)—C(7B)	1.392(6)
N(1B)—H(1B)	0.8800
N(2B)—C(10B)	1.474(6)
N(2B)—C(14B)	1.502(6)
N(2B)—C(13B)	1.512(6)
N(2B)—H(2BB)	0.9300
N(3B)—C(3B)	1.479(5)
N(3B)—C(16B)	1.496(6)
N(3B)—C(15B)	1.501(6)
N(3B)—H(3B)	0.9300
C(1B)—C(2B)	1.392(6)
C(1B)—C(6B)	1.410(6)
C(2B)—C(3B)	1.372(6)
C(2B)—H(2B)	0.9500
C(3B)—C(4B)	1.391(6)
C(4B)—C(5B)	1.391(6)
C(4B)—H(4B)	0.9500
C(5B)—C(6B)	1.389(6)
C(5B)—H(5B)	0.9500
C(7B)—C(8B)	1.399(6)
C(7B)—C(12B)	1.406(6)
C(8B)—C(9B)	1.377(6)
C(8B)—H(8B)	0.9500
C(9B)—C(10B)	1.381(6)
C(9B)—H(9B)	0.9500
C(10B)—C(11B)	1.384(6)
C(11B)—C(12B)	1.388(6)
C(11B)—H(11B)	0.9500
C(13B)—H(13D)	0.9800
C(13B)—H(13E)	0.9800
C(13B)—H(13F)	0.9800
C(14B)—H(14D)	0.9800
C(14B)—H(14E)	0.9800
C(14B)—H(14F)	0.9800
C(15B)—H(15D)	0.9800
C(15B)—H(15E)	0.9800
C(15B)—H(15F)	0.9800
C(16B)—H(16D)	0.9800
C(16B)—H(16E)	0.9800
C(16B)—H(16F)	0.9800
S(1C)—C(12C)	1.760(4)
S(1C)—C(1C)	1.765(4)
N(1C)—C(7C)	1.387(5)
N(1C)—C(6C)	1.395(5)
N(1C)—H(1C)	0.8800
N(2C)—C(10C)	1.469(5)
N(2C)—C(14C)	1.498(5)
N(2C)—C(13C)	1.503(5)
N(2C)—H(2CC)	0.9300

TABLE 3-continued

Bond lengths [Å] and angles [°] for LMT·2EsOH.	
N(3C)—C(3C)	1.482(5)
N(3C)—C(15C)	1.494(6)
N(3C)—C(16C)	1.497(6)
N(3C)—H(3C)	0.9300
C(1C)—C(2C)	1.377(6)
C(1C)—C(6C)	1.411(6)
C(2C)—C(3C)	1.380(6)
C(2C)—H(2C)	0.9500
C(3C)—C(4C)	1.381(6)
C(4C)—C(5C)	1.390(6)
C(4C)—H(4C)	0.9500
C(5C)—C(6C)	1.397(6)
C(5C)—H(5C)	0.9500
C(7C)—C(8C)	1.397(6)
C(7C)—C(12C)	1.413(6)
C(8C)—C(9C)	1.378(6)
C(8C)—H(8C)	0.9500
C(9C)—C(10C)	1.386(6)
C(9C)—H(9C)	0.9500
C(10C)—C(11C)	1.384(6)
C(11C)—C(12C)	1.377(6)
C(11C)—H(11C)	0.9500
C(13C)—H(13G)	0.9800
C(13C)—H(13H)	0.9800
C(13C)—H(13I)	0.9800
C(14C)—H(14G)	0.9800
C(14C)—H(14H)	0.9800
C(14C)—H(14I)	0.9800
C(15C)—H(15G)	0.9800
C(15C)—H(15H)	0.9800
C(15C)—H(15I)	0.9800
C(16C)—H(16G)	0.9800
C(16C)—H(16H)	0.9800
C(16C)—H(16I)	0.9800
S(1D)—C(12D)	1.760(4)
S(1D)—C(1D)	1.761(5)
N(1D)—C(7D)	1.381(6)
N(1D)—C(6D)	1.389(6)
N(1D)—H(1D)	0.8800
N(2D)—C(10D)	1.471(6)
N(2D)—C(13D)	1.486(6)
N(2D)—C(14D)	1.495(6)
N(2D)—H(2D)	0.9300
N(3D)—C(3D)	1.483(6)
N(3D)—C(15D)	1.495(6)
N(3D)—C(16D)	1.504(6)
N(3D)—H(3D)	0.9300
C(1D)—C(2D)	1.389(6)
C(1D)—C(6D)	1.405(6)
C(2D)—C(3D)	1.379(6)
C(2D)—H(2DD)	0.9500
C(3D)—C(4D)	1.381(6)
C(4D)—C(5D)	1.385(7)
C(4D)—H(4D)	0.9500
C(5D)—C(6D)	1.377(6)
C(5D)—H(5D)	0.9500
C(7D)—C(8D)	1.394(6)
C(7D)—C(12D)	1.409(6)
C(8D)—C(9D)	1.385(6)
C(8D)—H(8D)	0.9500
C(9D)—C(10D)	1.384(6)
C(9D)—H(9D)	0.9500
C(10D)—C(11D)	1.391(6)
C(11D)—C(12D)	1.388(6)
C(11D)—H(11D)	0.9500
C(13D)—H(13J)	0.9800
C(13D)—H(13K)	0.9800
C(13D)—H(13L)	0.9800
C(14D)—H(14J)	0.9800
C(14D)—H(14K)	0.9800
C(14D)—H(14L)	0.9800
C(15D)—H(15J)	0.9800
C(15D)—H(15K)	0.9800
C(15D)—H(15L)	0.9800
C(16D)—H(16J)	0.9800
C(16D)—H(16K)	0.9800
C(16D)—H(16L)	0.9800
S(2)—O(2)	1.32(3)

TABLE 3-continued

Bond lengths [Å] and angles [°] for LMT·2EsOH.	
S(2)—O(3)	1.359(16)
S(2)—O(1')	1.423(7)
S(2)—O(3')	1.481(8)
S(2)—O(2')	1.510(16)
S(2)—O(1)	1.503(13)
S(2)—C(1S)	1.756(6)
S(3)—O(5)	1.442(3)
S(3)—O(6)	1.459(3)
S(3)—O(4)	1.469(3)
S(3)—C(15S)	1.795(5)
S(4)—O(7)	1.446(4)
S(4)—O(9)	1.461(3)
S(4)—O(8)	1.466(3)
S(4)—C(3S)	1.779(5)
S(5)—O(11)	1.438(4)
S(5)—O(12)	1.456(4)
S(5)—O(10)	1.463(3)
S(5)—C(5S)	1.775(5)
S(6)—O(14)	1.453(4)
S(6)—O(15)	1.456(4)
S(6)—O(13)	1.458(4)
S(6)—C(9S)	1.749(5)
S(7)—O(17)	1.442(4)
S(7)—O(18)	1.479(4)
S(7)—O(16)	1.482(3)
S(7)—C(7S)	1.744(5)
S(8)—O(21)	1.447(3)
S(8)—O(19)	1.452(3)
S(8)—O(20)	1.479(3)
S(8)—C(11S)	1.776(4)
S(9)—O(24)	1.458(3)
S(9)—O(22)	1.458(3)
S(9)—O(23)	1.486(3)
S(9)—C(13S)	1.765(5)
O(1)—O(3)	1.56(2)
C(1S)—C(2S)	1.470(8)
C(1S)—H(1S1)	0.9900
C(1S)—H(1S2)	0.9900
C(2S)—H(2S1)	0.9800
C(2S)—H(2S2)	0.9800
C(2S)—H(2S3)	0.9800
C(3S)—C(4S)	1.534(7)
C(3S)—H(3S1)	0.9900
C(3S)—H(3S2)	0.9900
C(4S)—H(4S1)	0.9800
C(4S)—H(4S2)	0.9800
C(4S)—H(4S3)	0.9800
C(5S)—C(6S)	1.506(7)
C(5S)—H(5S1)	0.9900
C(5S)—H(5S2)	0.9900
C(6S)—H(6S1)	0.9800
C(6S)—H(6S2)	0.9800
C(6S)—H(6S3)	0.9800
C(7S)—C(8S)	1.550(7)
C(7S)—H(7S1)	0.9900
C(7S)—H(7S2)	0.9900
C(8S)—H(8S1)	0.9800
C(8S)—H(8S2)	0.9800
C(8S)—H(8S3)	0.9800
C(9S)—C(10S)	1.511(10)
C(9S)—H(9S1)	0.9900
C(9S)—H(9S2)	0.9900
C(10S)—H(10A)	0.9800
C(10S)—H(10B)	0.9800
C(10S)—H(10C)	0.9800
C(11S)—C(12S)	1.522(7)
C(11S)—H(11E)	0.9900
C(11S)—H(11F)	0.9900
C(12S)—H(12A)	0.9800
C(12S)—H(12B)	0.9800
C(12S)—H(12C)	0.9800
C(13S)—C(14S)	1.498(6)
C(13S)—H(13M)	0.9900
C(13S)—H(13N)	0.9900
C(14S)—H(14M)	0.9800
C(14S)—H(14N)	0.9800
C(14S)—H(14O)	0.9800
C(15S)—C(16S)	1.500(7)

TABLE 3-continued

Bond lengths [Å] and angles [°] for LMT·2EsOH.	
C(15S)—H(15M)	0.9900
C(15S)—H(15N)	0.9900
C(16S)—H(16M)	0.9800
C(16S)—H(16N)	0.9800
C(16S)—H(16O)	0.9800
C(1A)—S(1A)—C(12A)	102.6(2)
C(7A)—N(1A)—C(6A)	126.5(4)
C(7A)—N(1A)—H(1AA)	116.8
C(6A)—N(1A)—H(1AA)	116.8
C(10A)—N(2A)—C(14A)	112.3(4)
C(10A)—N(2A)—C(13A)	112.8(4)
C(14A)—N(2A)—C(13A)	111.3(4)
C(10A)—N(2A)—H(2AA)	106.7
C(14A)—N(2A)—H(2AA)	106.7
C(13A)—N(2A)—H(2AA)	106.7
C(3A)—N(3A)—C(16A)	114.4(3)
C(3A)—N(3A)—C(15A)	111.3(4)
C(16A)—N(3A)—C(15A)	110.1(4)
C(3A)—N(3A)—H(3A)	106.8
C(16A)—N(3A)—H(3A)	106.8
C(15A)—N(3A)—H(3A)	106.8
C(2A)—C(1A)—C(6A)	120.3(4)
C(2A)—C(1A)—S(1A)	116.7(3)
C(6A)—C(1A)—S(1A)	123.0(3)
C(3A)—C(2A)—C(1A)	120.4(4)
C(3A)—C(2A)—H(2A)	119.8
C(1A)—C(2A)—H(2A)	119.8
C(2A)—C(3A)—C(4A)	120.6(4)
C(2A)—C(3A)—N(3A)	120.2(4)
C(4A)—C(3A)—N(3A)	119.1(4)
C(3A)—C(4A)—C(5A)	118.8(4)
C(3A)—C(4A)—H(4A)	120.6
C(5A)—C(4A)—H(4A)	120.6
C(6A)—C(5A)—C(4A)	121.8(4)
C(6A)—C(5A)—H(5A)	119.1
C(4A)—C(5A)—H(5A)	119.1
C(5A)—C(6A)—N(1A)	119.7(4)
C(5A)—C(6A)—C(1A)	118.0(4)
N(1A)—C(6A)—C(1A)	122.3(4)
C(8A)—C(7A)—N(1A)	119.7(4)
C(8A)—C(7A)—C(12A)	118.4(4)
N(1A)—C(7A)—C(12A)	121.9(4)
C(7A)—C(8A)—C(9A)	121.9(4)
C(7A)—C(8A)—H(8A)	119.0
C(9A)—C(8A)—H(8A)	119.0
C(10A)—C(9A)—C(8A)	118.4(4)
C(10A)—C(9A)—H(9A)	120.8
C(8A)—C(9A)—H(9A)	120.8
C(11A)—C(10A)—C(9A)	120.9(4)
C(11A)—C(10A)—N(2A)	117.9(4)
C(9A)—C(10A)—N(2A)	121.2(4)
C(10A)—C(11A)—C(12A)	120.4(4)
C(10A)—C(11A)—H(11A)	119.8
C(12A)—C(11A)—H(11A)	119.8
C(11A)—C(12A)—C(7A)	119.9(4)
C(11A)—C(12A)—S(1A)	116.5(3)
C(7A)—C(12A)—S(1A)	123.5(3)
N(2A)—C(13A)—H(13A)	109.5
N(2A)—C(13A)—H(13B)	109.5
H(13A)—C(13A)—H(13B)	109.5
N(2A)—C(13A)—H(13C)	109.5
H(13A)—C(13A)—H(13C)	109.5
H(13B)—C(13A)—H(13C)	109.5
N(2A)—C(14A)—H(14A)	109.5
N(2A)—C(14A)—H(14B)	109.5
H(14A)—C(14A)—H(14B)	109.5
N(2A)—C(14A)—H(14C)	109.5
H(14A)—C(14A)—H(14C)	109.5
H(14B)—C(14A)—H(14C)	109.5
N(3A)—C(15A)—H(15A)	109.5
N(3A)—C(15A)—H(15B)	109.5
H(15A)—C(15A)—H(15B)	109.5
N(3A)—C(15A)—H(15C)	109.5
H(15A)—C(15A)—H(15C)	109.5
H(15B)—C(15A)—H(15C)	109.5
N(3A)—C(16A)—H(16A)	109.5
N(3A)—C(16A)—H(16B)	109.5
H(16A)—C(16A)—H(16B)	109.5

TABLE 3-continued

Bond lengths [Å] and angles [°] for LMT·2EsOH.	
N(3A)—C(16A)—H(16C)	109.5
H(16A)—C(16A)—H(16C)	109.5
H(16B)—C(16A)—H(16C)	109.5
C(12B)—S(1B)—C(1B)	101.6(2)
C(6B)—N(1B)—C(7B)	125.1(4)
C(6B)—N(1B)—H(1B)	117.4
C(7B)—N(1B)—H(1B)	117.4
C(10B)—N(2B)—C(14B)	111.3(3)
C(10B)—N(2B)—C(13B)	114.6(4)
C(14B)—N(2B)—C(13B)	110.7(4)
C(10B)—N(2B)—H(2BB)	106.5
C(14B)—N(2B)—H(2BB)	106.5
C(13B)—N(2B)—H(2BB)	106.5
C(3B)—N(3B)—C(16B)	113.0(3)
C(3B)—N(3B)—C(15B)	111.8(3)
C(16B)—N(3B)—C(15B)	111.1(4)
C(3B)—N(3B)—H(3B)	106.9
C(16B)—N(3B)—H(3B)	106.9
C(15B)—N(3B)—H(3B)	106.9
C(2B)—C(1B)—C(6B)	120.4(4)
C(2B)—C(1B)—S(1B)	117.8(3)
C(6B)—C(1B)—S(1B)	121.5(3)
C(3B)—C(2B)—C(1B)	119.7(4)
C(3B)—C(2B)—H(2B)	120.2
C(1B)—C(2B)—H(2B)	120.2
C(2B)—C(3B)—C(4B)	121.1(4)
C(2B)—C(3B)—N(3B)	120.3(4)
C(4B)—C(3B)—N(3B)	118.7(4)
C(5B)—C(4B)—C(3B)	119.2(4)
C(5B)—C(4B)—H(4B)	120.4
C(3B)—C(4B)—H(4B)	120.4
C(4B)—C(5B)—C(6B)	121.1(4)
C(4B)—C(5B)—H(5B)	119.5
C(6B)—C(5B)—H(5B)	119.5
N(1B)—C(6B)—C(5B)	120.1(4)
N(1B)—C(6B)—C(1B)	121.4(4)
C(5B)—C(6B)—C(1B)	118.5(4)
N(1B)—C(7B)—C(8B)	119.9(4)
N(1B)—C(7B)—C(12B)	121.8(4)
C(8B)—C(7B)—C(12B)	118.3(4)
C(9B)—C(8B)—C(7B)	121.6(4)
C(9B)—C(8B)—H(8B)	119.2
C(7B)—C(8B)—H(8B)	119.2
C(8B)—C(9B)—C(10B)	119.0(4)
C(8B)—C(9B)—H(9B)	120.5
C(10B)—C(9B)—H(9B)	120.5
C(9B)—C(10B)—C(11B)	121.4(4)
C(9B)—C(10B)—N(2B)	120.8(4)
C(11B)—C(10B)—N(2B)	117.6(4)
C(10B)—C(11B)—C(12B)	119.5(4)
C(10B)—C(11B)—H(11B)	120.3
C(12B)—C(11B)—H(11B)	120.3
C(11B)—C(12B)—C(7B)	120.3(4)
C(11B)—C(12B)—S(1B)	118.2(3)
C(7B)—C(12B)—S(1B)	121.2(3)
N(2B)—C(13B)—H(13D)	109.5
N(2B)—C(13B)—H(13E)	109.5
H(13D)—C(13B)—H(13E)	109.5
N(2B)—C(13B)—H(13F)	109.5
H(13D)—C(13B)—H(13F)	109.5
H(13E)—C(13B)—H(13F)	109.5
N(2B)—C(14B)—H(14D)	109.5
N(2B)—C(14B)—H(14E)	109.5
H(14D)—C(14B)—H(14E)	109.5
N(2B)—C(14B)—H(14F)	109.5
H(14D)—C(14B)—H(14F)	109.5
H(14E)—C(14B)—H(14F)	109.5
N(3B)—C(15B)—H(15D)	109.5
N(3B)—C(15B)—H(15E)	109.5
H(15D)—C(15B)—H(15E)	109.5
N(3B)—C(15B)—H(15F)	109.5
H(15D)—C(15B)—H(15F)	109.5
H(15E)—C(15B)—H(15F)	109.5
N(3B)—C(16B)—H(16D)	109.5
N(3B)—C(16B)—H(16E)	109.5
H(16D)—C(16B)—H(16E)	109.5
N(3B)—C(16B)—H(16F)	109.5
H(16D)—C(16B)—H(16F)	109.5

TABLE 3-continued

Bond lengths [Å] and angles [°] for LMT·2EsOH.	
H(16E)—C(16B)—H(16F)	109.5
C(12C)—S(1C)—C(1C)	101.7(2)
C(7C)—N(1C)—C(6C)	125.4(4)
C(7C)—N(1C)—H(1C)	117.3
C(6C)—N(1C)—H(1C)	117.3
C(10C)—N(2C)—C(14C)	114.9(3)
C(10C)—N(2C)—C(13C)	110.2(3)
C(14C)—N(2C)—C(13C)	111.2(3)
C(10C)—N(2C)—H(2CC)	106.7
C(14C)—N(2C)—H(2CC)	106.7
C(13C)—N(2C)—H(2CC)	106.7
C(3C)—N(3C)—C(15C)	111.2(4)
C(3C)—N(3C)—C(16C)	112.9(4)
C(15C)—N(3C)—C(16C)	111.5(4)
C(3C)—N(3C)—H(3C)	106.9
C(15C)—N(3C)—H(3C)	106.9
C(16C)—N(3C)—H(3C)	106.9
C(2C)—C(1C)—C(6C)	120.3(4)
C(2C)—C(1C)—S(1C)	117.7(3)
C(6C)—C(1C)—S(1C)	121.8(3)
C(1C)—C(2C)—C(3C)	119.7(4)
C(1C)—C(2C)—H(2C)	120.2
C(3C)—C(2C)—H(2C)	120.2
C(2C)—C(3C)—C(4C)	121.8(4)
C(2C)—C(3C)—N(3C)	118.6(4)
C(4C)—C(3C)—N(3C)	119.5(4)
C(3C)—C(4C)—C(5C)	118.7(4)
C(3C)—C(4C)—H(4C)	120.7
C(5C)—C(4C)—H(4C)	120.7
C(4C)—C(5C)—C(6C)	120.9(4)
C(4C)—C(5C)—H(5C)	119.5
C(6C)—C(5C)—H(5C)	119.5
N(1C)—C(6C)—C(5C)	119.9(4)
N(1C)—C(6C)—C(1C)	121.3(4)
C(5C)—C(6C)—C(1C)	118.7(4)
N(1C)—C(7C)—C(8C)	119.9(4)
N(1C)—C(7C)—C(12C)	122.0(4)
C(8C)—C(7C)—C(12C)	118.1(4)
C(9C)—C(8C)—C(7C)	121.5(4)
C(9C)—C(8C)—H(8C)	119.2
C(7C)—C(8C)—H(8C)	119.2
C(8C)—C(9C)—C(10C)	119.3(4)
C(8C)—C(9C)—H(9C)	120.3
C(10C)—C(9C)—H(9C)	120.3
C(11C)—C(10C)—C(9C)	120.5(4)
C(11C)—C(10C)—N(2C)	117.5(4)
C(9C)—C(10C)—N(2C)	121.9(4)
C(12C)—C(11C)—C(10C)	120.4(4)
C(12C)—C(11C)—H(11C)	119.8
C(10C)—C(11C)—H(11C)	119.8
C(11C)—C(12C)—C(7C)	120.1(4)
C(11C)—C(12C)—S(1C)	118.4(3)
C(7C)—C(12C)—S(1C)	121.3(3)
N(2C)—C(13C)—H(13G)	109.5
N(2C)—C(13C)—H(13H)	109.5
H(13G)—C(13C)—H(13H)	109.5
N(2C)—C(13C)—H(13I)	109.5
H(13G)—C(13C)—H(13I)	109.5
H(13H)—C(13C)—H(13I)	109.5
N(2C)—C(14C)—H(14G)	109.5
N(2C)—C(14C)—H(14H)	109.5
H(14G)—C(14C)—H(14H)	109.5
N(2C)—C(14C)—H(14I)	109.5
H(14G)—C(14C)—H(14I)	109.5
H(14H)—C(14C)—H(14I)	109.5
N(3C)—C(15C)—H(15G)	109.5
N(3C)—C(15C)—H(15H)	109.5
H(15G)—C(15C)—H(15H)	109.5
N(3C)—C(15C)—H(15I)	109.5
H(15G)—C(15C)—H(15I)	109.5
H(15H)—C(15C)—H(15I)	109.5
N(3C)—C(16C)—H(16G)	109.5
N(3C)—C(16C)—H(16H)	109.5
H(16G)—C(16C)—H(16H)	109.5
N(3C)—C(16C)—H(16I)	109.5
H(16G)—C(16C)—H(16I)	109.5
H(16H)—C(16C)—H(16I)	109.5
C(12D)—S(1D)—C(1D)	102.4(2)

TABLE 3-continued

Bond lengths [Å] and angles [°] for LMT·2EsOH.	
C(7D)—N(1D)—C(6D)	126.5(4)
C(7D)—N(1D)—H(1D)	116.7
C(6D)—N(1D)—H(1D)	116.7
C(10D)—N(2D)—C(13D)	114.5(4)
C(10D)—N(2D)—C(14D)	111.3(3)
C(13D)—N(2D)—C(14D)	110.7(4)
C(10D)—N(2D)—H(2D)	106.6
C(13D)—N(2D)—H(2D)	106.6
C(14D)—N(2D)—H(2D)	106.6
C(3D)—N(3D)—C(15D)	111.2(4)
C(3D)—N(3D)—C(16D)	114.3(4)
C(15D)—N(3D)—C(16D)	111.2(4)
C(3D)—N(3D)—H(3D)	106.5
C(15D)—N(3D)—H(3D)	106.5
C(16D)—N(3D)—H(3D)	106.5
C(2D)—C(1D)—C(6D)	120.2(4)
C(2D)—C(1D)—S(1D)	117.4(3)
C(6D)—C(1D)—S(1D)	122.4(3)
C(3D)—C(2D)—C(1D)	119.7(4)
C(3D)—C(2D)—H(2DD)	120.1
C(1D)—C(2D)—H(2DD)	120.1
C(2D)—C(3D)—C(4D)	120.7(4)
C(2D)—C(3D)—N(3D)	120.7(4)
C(4D)—C(3D)—N(3D)	118.4(4)
C(3D)—C(4D)—C(5D)	119.2(5)
C(3D)—C(4D)—H(4D)	120.4
C(5D)—C(4D)—H(4D)	120.4
C(6D)—C(5D)—C(4D)	121.5(4)
C(6D)—C(5D)—H(5D)	119.2
C(4D)—C(5D)—H(5D)	119.2
C(5D)—C(6D)—N(1D)	118.7(4)
C(5D)—C(6D)—C(1D)	118.6(4)
N(1D)—C(6D)—C(1D)	122.7(4)
N(1D)—C(7D)—C(8D)	119.7(4)
N(1D)—C(7D)—C(12D)	121.5(4)
C(8D)—C(7D)—C(12D)	118.7(4)
C(9D)—C(8D)—C(7D)	121.8(4)
C(9D)—C(8D)—H(8D)	119.1
C(7D)—C(8D)—H(8D)	119.1
C(10D)—C(9D)—C(8D)	118.3(4)
C(10D)—C(9D)—H(9D)	120.9
C(8D)—C(9D)—H(9D)	120.9
C(9D)—C(10D)—C(11D)	121.7(4)
C(9D)—C(10D)—N(2D)	121.0(4)
C(11D)—C(10D)—N(2D)	117.1(4)
C(12D)—C(11D)—C(10D)	119.5(4)
C(12D)—C(11D)—H(11D)	120.3
C(10D)—C(11D)—H(11D)	120.3
C(11D)—C(12D)—C(7D)	120.0(4)
C(11D)—C(12D)—S(1D)	116.4(3)
C(7D)—C(12D)—S(1D)	123.4(3)
N(2D)—C(13D)—H(13J)	109.5
N(2D)—C(13D)—H(13K)	109.5
H(13J)—C(13D)—H(13K)	109.5
N(2D)—C(13D)—H(13L)	109.5
H(13J)—C(13D)—H(13L)	109.5
H(13K)—C(13D)—H(13L)	109.5
N(2D)—C(14D)—H(14J)	109.5
N(2D)—C(14D)—H(14K)	109.5
H(14J)—C(14D)—H(14K)	109.5
N(2D)—C(14D)—H(14L)	109.5
H(14J)—C(14D)—H(14L)	109.5
H(14K)—C(14D)—H(14L)	109.5
N(3D)—C(15D)—H(15J)	109.5
N(3D)—C(15D)—H(15K)	109.5
H(15J)—C(15D)—H(15K)	109.5
N(3D)—C(15D)—H(15L)	109.5
H(15J)—C(15D)—H(15L)	109.5
H(15K)—C(15D)—H(15L)	109.5
N(3D)—C(16D)—H(16J)	109.5
N(3D)—C(16D)—H(16K)	109.5
H(16J)—C(16D)—H(16K)	109.5
N(3D)—C(16D)—H(16L)	109.5
H(16J)—C(16D)—H(16L)	109.5
H(16K)—C(16D)—H(16L)	109.5
O(2)—S(2)—O(3)	118.5(17)
O(2)—S(2)—O(1')	87.9(12)
O(3)—S(2)—O(1')	108.0(8)

TABLE 3-continued

Bond lengths [Å] and angles [°] for LMT·2EsOH.	
O(2)—S(2)—O(3')	117.2(14)
O(3)—S(2)—O(3')	31.5(6)
O(1')—S(2)—O(3')	138.3(6)
O(2)—S(2)—O(2')	23.8(18)
O(3)—S(2)—O(2')	115.1(10)
O(1')—S(2)—O(2')	110.6(8)
O(3')—S(2)—O(2')	101.4(8)
O(2)—S(2)—O(1)	111.9(14)
O(3)—S(2)—O(1)	65.7(9)
O(1')—S(2)—O(1)	42.6(5)
O(3')—S(2)—O(1)	95.8(7)
O(2')—S(2)—O(1)	134.1(9)
O(2)—S(2)—C(1S)	117.0(16)
O(3)—S(2)—C(1S)	118.1(6)
O(1')—S(2)—C(1S)	99.1(4)
O(3')—S(2)—C(1S)	97.8(4)
O(2')—S(2)—C(1S)	104.8(8)
O(1)—S(2)—C(1S)	114.6(5)
O(5)—S(3)—O(6)	113.5(2)
O(5)—S(3)—O(4)	112.5(2)
O(6)—S(3)—O(4)	111.9(2)
O(5)—S(3)—C(15S)	107.5(2)
O(6)—S(3)—C(15S)	106.2(2)
O(4)—S(3)—C(15S)	104.6(2)
O(7)—S(4)—O(9)	113.8(2)
O(7)—S(4)—O(8)	113.4(2)
O(9)—S(4)—O(8)	111.3(2)
O(7)—S(4)—C(3S)	106.0(2)
O(9)—S(4)—C(3S)	106.6(2)
O(8)—S(4)—C(3S)	104.9(2)
O(11)—S(5)—O(12)	112.3(3)
O(11)—S(5)—O(10)	114.1(3)
O(12)—S(5)—O(10)	109.9(2)
O(11)—S(5)—C(5S)	107.4(2)
O(12)—S(5)—C(5S)	107.2(2)
O(10)—S(5)—C(5S)	105.5(2)
O(14)—S(6)—O(15)	112.2(2)
O(14)—S(6)—O(13)	113.4(3)
O(15)—S(6)—O(13)	111.4(2)
O(14)—S(6)—C(9S)	105.6(2)
O(15)—S(6)—C(9S)	108.3(2)
O(13)—S(6)—C(9S)	105.4(3)
O(17)—S(7)—O(18)	114.0(2)
O(17)—S(7)—O(16)	114.2(2)
O(18)—S(7)—O(16)	110.7(2)
O(17)—S(7)—C(7S)	105.8(3)
O(18)—S(7)—C(7S)	105.3(2)
O(16)—S(7)—C(7S)	106.0(2)
O(21)—S(8)—O(19)	114.3(2)
O(21)—S(8)—O(20)	111.7(2)
O(19)—S(8)—O(20)	111.95(19)
O(21)—S(8)—C(11S)	106.3(2)
O(19)—S(8)—C(11S)	108.1(2)
O(20)—S(8)—C(11S)	103.7(2)
O(24)—S(9)—O(22)	113.23(19)
O(24)—S(9)—O(23)	112.94(19)
O(22)—S(9)—O(23)	111.07(18)
O(24)—S(9)—C(13S)	106.1(2)
O(22)—S(9)—C(13S)	107.7(2)
O(23)—S(9)—C(13S)	105.3(2)
S(2)—O(1)—O(3)	52.7(7)
S(2)—O(3)—O(1)	61.6(8)
C(2S)—C(1S)—S(2)	115.9(5)
C(2S)—C(1S)—H(1S1)	108.3
S(2)—C(1S)—H(1S1)	108.3
C(2S)—C(1S)—H(1S2)	108.3
S(2)—C(1S)—H(1S2)	108.3
H(1S1)—C(1S)—H(1S2)	107.4
C(1S)—C(2S)—H(2S1)	109.5
C(1S)—C(2S)—H(2S2)	109.5
H(2S1)—C(2S)—H(2S2)	109.5
C(1S)—C(2S)—H(2S3)	109.5
H(2S1)—C(2S)—H(2S3)	109.5
H(2S2)—C(2S)—H(2S3)	109.5
C(4S)—C(3S)—S(4)	111.3(3)
C(4S)—C(3S)—H(3S1)	109.4
S(4)—C(3S)—H(3S1)	109.4
C(4S)—C(3S)—H(3S2)	109.4

TABLE 3-continued

Bond lengths [Å] and angles [°] for LMT·2EsOH.	
S(4)—C(3S)—H(3S2)	109.4
H(3S1)—C(3S)—H(3S2)	108.0
C(3S)—C(4S)—H(4S1)	109.5
C(3S)—C(4S)—H(4S2)	109.5
H(4S1)—C(4S)—H(4S2)	109.5
C(3S)—C(4S)—H(4S3)	109.5
H(4S1)—C(4S)—H(4S3)	109.5
H(4S2)—C(4S)—H(4S3)	109.5
C(6S)—C(5S)—S(5)	112.7(3)
C(6S)—C(5S)—H(5S1)	109.0
S(5)—C(5S)—H(5S1)	109.0
C(6S)—C(5S)—H(5S2)	109.0
S(5)—C(5S)—H(5S2)	109.0
H(5S1)—C(5S)—H(5S2)	107.8
C(5S)—C(6S)—H(6S1)	109.5
C(5S)—C(6S)—H(6S2)	109.5
H(6S1)—C(6S)—H(6S2)	109.5
C(5S)—C(6S)—H(6S3)	109.5
H(6S1)—C(6S)—H(6S3)	109.5
H(6S2)—C(6S)—H(6S3)	109.5
C(8S)—C(7S)—S(7)	110.6(4)
C(8S)—C(7S)—H(7S1)	109.5
S(7)—C(7S)—H(7S1)	109.5
C(8S)—C(7S)—H(7S2)	109.5
S(7)—C(7S)—H(7S2)	109.5
H(7S1)—C(7S)—H(7S2)	108.1
C(7S)—C(8S)—H(8S1)	109.5
C(7S)—C(8S)—H(8S2)	109.5
H(8S1)—C(8S)—H(8S2)	109.5
C(7S)—C(8S)—H(8S3)	109.5
H(8S1)—C(8S)—H(8S3)	109.5
H(8S2)—C(8S)—H(8S3)	109.5
C(10S)—C(9S)—S(6)	104.5(5)
C(10S)—C(9S)—H(9S1)	110.8
S(6)—C(9S)—H(9S1)	110.8
C(10S)—C(9S)—H(9S2)	110.8
S(6)—C(9S)—H(9S2)	110.8
H(9S1)—C(9S)—H(9S2)	108.9
C(9S)—C(10S)—H(10A)	109.5
C(9S)—C(10S)—H(10B)	109.5
H(10A)—C(10S)—H(10B)	109.5
C(9S)—C(10S)—H(10C)	109.5
H(10A)—C(10S)—H(10C)	109.5
H(10B)—C(10S)—H(10C)	109.5
C(12S)—C(11S)—S(8)	113.0(3)
C(12S)—C(11S)—H(11E)	109.0
S(8)—C(11S)—H(11E)	109.0
C(12S)—C(11S)—H(11F)	109.0
S(8)—C(11S)—H(11F)	109.0
H(11E)—C(11S)—H(11F)	107.8
C(11S)—C(12S)—H(12A)	109.5
C(11S)—C(12S)—H(12B)	109.5
H(12A)—C(12S)—H(12B)	109.5
C(11S)—C(12S)—H(12C)	109.5
H(12A)—C(12S)—H(12C)	109.5
H(12B)—C(12S)—H(12C)	109.5
C(14S)—C(13S)—S(9)	111.6(3)
C(14S)—C(13S)—H(13M)	109.3
S(9)—C(13S)—H(13M)	109.3
C(14S)—C(13S)—H(13N)	109.3
S(9)—C(13S)—H(13N)	109.3
H(13M)—C(13S)—H(13N)	108.0
C(13S)—C(14S)—H(14M)	109.5
C(13S)—C(14S)—H(14N)	109.5
H(14M)—C(14S)—H(14N)	109.5
C(13S)—C(14S)—H(14O)	109.5
H(14M)—C(14S)—H(14O)	109.5
H(14N)—C(14S)—H(14O)	109.5
C(16S)—C(15S)—S(3)	112.7(4)
C(16S)—C(15S)—H(15M)	109.0
S(3)—C(15S)—H(15M)	109.0
C(16S)—C(15S)—H(15N)	109.0
S(3)—C(15S)—H(15N)	109.0
H(15M)—C(15S)—H(15N)	107.8
C(15S)—C(16S)—H(16M)	109.5
C(15S)—C(16S)—H(16N)	109.5
H(16M)—C(16S)—H(16N)	109.5
C(15S)—C(16S)—H(16O)	109.5

103

TABLE 3-continued

Bond lengths [Å] and angles [°] for LMT·2EsOH.	
H(16M)—C(16S)—H(16O)	109.5
H(16N)—C(16S)—H(16O)	109.5

Symmetry transformations used to generate equivalent atoms:

TABLE 4

Anisotropic displacement parameters ($\text{Å}^2 \times 10^3$) for LMT·2EsOH. The anisotropic displacement factor exponent takes the form: $-2h^2 a^* U^{11} + \dots + 2 h k a^* b^* U^{12}$						
	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
S(1A)	18(1)	21(1)	121(2)	12(1)	-15(1)	-2(1)
N(1A)	15(2)	24(2)	36(2)	1(2)	0(2)	-3(2)
N(2A)	19(2)	20(2)	27(2)	5(2)	1(2)	0(2)
N(3A)	17(2)	23(2)	27(2)	-1(2)	3(2)	1(2)
C(1A)	20(3)	22(2)	39(3)	3(2)	-4(2)	-1(2)
C(2A)	17(2)	24(2)	39(3)	2(2)	-4(2)	-1(2)
C(3A)	22(2)	22(2)	25(2)	-1(2)	7(2)	-2(2)
C(4A)	17(2)	29(2)	31(3)	-5(2)	7(2)	2(2)
C(5A)	15(2)	28(2)	37(3)	-6(2)	9(2)	-6(2)
C(6A)	18(2)	23(2)	26(2)	-2(2)	3(2)	-3(2)
C(7A)	22(2)	23(2)	25(2)	0(2)	5(2)	-2(2)
C(8A)	21(2)	25(2)	21(2)	-3(2)	6(2)	-7(2)
C(9A)	25(2)	22(2)	24(2)	2(2)	7(2)	-3(2)
C(10A)	19(2)	24(2)	22(2)	3(2)	4(2)	2(2)
C(11A)	19(2)	23(2)	33(3)	2(2)	1(2)	-8(2)
C(12A)	20(3)	22(2)	42(3)	6(2)	2(2)	-1(2)
C(13A)	28(3)	39(3)	30(3)	15(2)	0(2)	-2(2)
C(14A)	34(3)	44(3)	37(3)	1(2)	7(2)	13(2)
C(15A)	96(5)	26(3)	36(3)	3(2)	37(3)	2(3)
C(16A)	32(3)	26(2)	30(3)	-9(2)	4(2)	6(2)
S(1B)	17(1)	18(1)	39(1)	3(1)	7(1)	-1(1)
N(1B)	21(2)	16(2)	25(2)	-2(2)	8(2)	-3(2)
N(2B)	29(2)	24(2)	20(2)	4(2)	1(2)	6(2)
N(3B)	21(2)	19(2)	21(2)	-2(2)	3(2)	2(2)
C(1B)	17(2)	20(2)	18(2)	0(2)	5(2)	0(2)
C(2B)	21(2)	23(2)	16(2)	-3(2)	3(2)	-4(2)
C(3B)	22(2)	18(2)	18(2)	-2(2)	6(2)	1(2)
C(4B)	16(2)	23(2)	20(2)	-1(2)	3(2)	0(2)
C(5B)	16(2)	21(2)	21(2)	-1(2)	5(2)	-4(2)
C(6B)	25(2)	20(2)	17(2)	0(2)	9(2)	1(2)
C(7B)	22(2)	24(2)	20(2)	-3(2)	9(2)	0(2)
C(8B)	29(3)	19(2)	27(2)	0(2)	17(2)	-2(2)
C(9B)	32(3)	20(2)	25(2)	3(2)	16(2)	4(2)
C(10B)	31(3)	24(2)	14(2)	1(2)	7(2)	9(2)
C(11B)	24(2)	23(2)	19(2)	-2(2)	6(2)	-1(2)
C(12B)	28(3)	18(2)	20(2)	0(2)	9(2)	2(2)
C(13B)	53(3)	30(3)	26(3)	14(2)	11(2)	17(2)
C(14B)	22(2)	30(2)	27(3)	1(2)	4(2)	6(2)
C(15B)	42(3)	22(2)	35(3)	3(2)	15(2)	3(2)
C(16B)	39(3)	30(3)	25(3)	-8(2)	2(2)	10(2)
S(1C)	16(1)	21(1)	34(1)	2(1)	6(1)	0(1)
N(1C)	15(2)	22(2)	31(2)	-8(2)	4(2)	-5(2)
N(2C)	16(2)	19(2)	23(2)	0(2)	6(2)	0(1)
N(3C)	21(2)	28(2)	27(2)	1(2)	7(2)	5(2)
C(1C)	16(2)	25(2)	15(2)	-1(2)	2(2)	0(2)
C(2C)	20(2)	24(2)	17(2)	-2(2)	6(2)	-1(2)
C(3C)	21(2)	20(2)	19(2)	1(2)	6(2)	5(2)
C(4C)	20(2)	34(3)	20(2)	-2(2)	7(2)	4(2)
C(5C)	20(2)	26(2)	20(2)	-1(2)	6(2)	-1(2)
C(6C)	22(2)	24(2)	15(2)	-4(2)	7(2)	-1(2)
C(7C)	18(2)	29(2)	17(2)	-4(2)	6(2)	2(2)
C(8C)	17(2)	22(2)	27(2)	-8(2)	9(2)	-6(2)
C(9C)	21(2)	20(2)	22(2)	-3(2)	10(2)	-2(2)
C(10C)	14(2)	25(2)	18(2)	-3(2)	5(2)	1(2)
C(11C)	15(2)	24(2)	21(2)	-5(2)	6(2)	-7(2)
C(12C)	18(2)	19(2)	21(2)	-4(2)	7(2)	2(2)
C(13C)	17(2)	30(2)	29(3)	2(2)	9(2)	3(2)
C(14C)	23(2)	23(2)	23(2)	2(2)	7(2)	-2(2)
C(15C)	46(3)	45(3)	56(4)	22(3)	29(3)	11(3)
C(16C)	38(3)	33(3)	46(3)	-18(2)	-4(3)	8(2)
S(1D)	25(1)	23(1)	46(1)	-3(1)	13(1)	-4(1)
N(1D)	26(2)	23(2)	36(2)	-4(2)	11(2)	-4(2)
N(2D)	22(2)	35(2)	23(2)	2(2)	7(2)	2(2)

104

TABLE 4-continued

Anisotropic displacement parameters ($\text{Å}^2 \times 10^3$) for LMT·2EsOH. The anisotropic displacement factor exponent takes the form: $-2h^2 a^* U^{11} + \dots + 2 h k a^* b^* U^{12}$						
	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
N(3D)	27(2)	24(2)	34(2)	0(2)	16(2)	2(2)
C(1D)	28(3)	28(2)	20(2)	-3(2)	10(2)	0(2)
C(2D)	30(3)	29(2)	23(2)	-5(2)	13(2)	-4(2)
C(3D)	33(3)	24(2)	24(2)	-1(2)	13(2)	1(2)
C(4D)	27(3)	31(3)	33(3)	0(2)	10(2)	-1(2)
C(5D)	26(3)	29(3)	36(3)	-6(2)	8(2)	-6(2)
C(6D)	29(3)	24(2)	19(2)	-3(2)	10(2)	0(2)
C(7D)	23(2)	27(2)	16(2)	0(2)	9(2)	-1(2)
C(8D)	30(3)	27(2)	19(2)	0(2)	15(2)	-3(2)
C(9D)	29(3)	29(2)	16(2)	3(2)	9(2)	-2(2)
C(10D)	24(2)	30(2)	17(2)	-2(2)	8(2)	0(2)
C(11D)	26(3)	32(2)	17(2)	-6(2)	7(2)	-6(2)
C(12D)	28(3)	24(2)	18(2)	-5(2)	11(2)	-1(2)
C(13D)	33(3)	60(4)	33(3)	19(3)	11(2)	10(3)
C(14D)	30(3)	33(3)	32(3)	3(2)	16(2)	5(2)
C(15D)	51(3)	34(3)	41(3)	6(2)	23(3)	10(2)
C(16D)	35(3)	26(2)	47(3)	-12(2)	19(3)	-3(2)
S(2)	36(1)	27(1)	45(1)	9(1)	18(1)	2(1)
S(3)	23(1)	22(1)	24(1)	-3(1)	4(1)	-1(1)
S(4)	16(1)	23(1)	40(1)	-4(1)	4(1)	-1(1)
S(5)	23(1)	24(1)	41(1)	8(1)	12(1)	4(1)
S(6)	42(1)	19(1)	29(1)	0(1)	9(1)	-2(1)
S(7)	31(1)	28(1)	39(1)	4(1)	12(1)	1(1)
S(8)	20(1)	19(1)	24(1)	0(1)	4(1)	-1(1)
S(9)	23(1)	23(1)	27(1)	-1(1)	7(1)	4(1)
O(1)	67(4)	53(4)	52(5)	19(4)	29(4)	10(4)
O(2)	56(6)	25(9)	107(14)	17(8)	13(8)	-20(7)
O(3)	55(4)	54(4)	57(5)	23(4)	25(4)	8(3)
O(1')	79(5)	68(4)	48(4)	25(3)	39(4)	31(4)
O(2')	53(4)	20(7)	109(13)	14(6)	9(7)	-4(5)
O(3')	35(3)	71(5)	68(5)	41(4)	21(4)	9(3)
O(4)	26(2)	26(2)	44(2)	3(2)	7(2)	2(1)
O(5)	40(2)	33(2)	35(2)	-10(2)	0(2)	-8(2)
O(6)	23(2)	31(2)	26(2)	-2(1)	9(1)	-6(1)
O(7)	27(2)	27(2)	72(3)	11(2)	11(2)	0(2)
O(8)	16(2)	42(2)	57(2)	-18(2)	-3(2)	-2(2)
O(9)	24(2)	29(2)	38(2)	-2(1)	14(2)	-3(1)
O(10)	43(2)	49(2)	33(2)	12(2)	18(2)	30(2)
O(11)	37(2)	48(2)	95(4)	34(2)	4(2)	-11(2)
O(12)	78(3)	30(2)	34(2)	-2(2)	28(2)	3(2)
O(13)	52(3)	72(3)	56(3)	-16(2)	35(2)	-33(2)
O(14)	94(3)	25(2)	35(2)	-2(2)	3(2)	13(2)
O(15)	37(2)	40(2)	35(2)	-9(2)	1(2)	5(2)
O(16)	23(2)	37(2)	36(2)	-16(2)	13(2)	-8(1)
O(17)	59(3)	36(2)	68(3)	23(2)	21(2)	-5(2)
O(18)	28(2)	52(2)	32(2)	-9(2)	9(2)	0(2)
O(19)	33(2)	19(2)	34(2)	-1(1)	7(2)	-2(1)
O(20)	20(2)	24(2)	45(2)	-4(2)	-1(2)	-2(1)
O(21)	32(2)	34(2)	28(2)	3(1)	14(2)	2(1)
O(22)	26(2)	27(2)	24(2)	-2(1)	9(1)	-2(1)
O(23)	26(2)	23(2)	39(2)	3(1)	9(2)	8(1)
O(24)	26(2)	25(2)	42(2)	-5(2)	8(2)	-4(1)
C(1S)	60(4)	50(4)	56(4)	14(3)	5(3)	6(3)
C(2S)	48(3)	52(3)	36(3)	10(3)	19(3)	-2(3)
C(3S)	20(2)	37(3)	30(3)	-4(2)	6(2)	5(2)
C(4S)	32(3)	57(3)	28(3)	14(2)	12(2)	8(2)
C(5S)	23(3)	31(3)	50(3)	11(2)	9(2)	8(2)
C(6S)	37(3)	32(3)	26(3)	6(2)	13(2)	3(2)
C(7S)	32(3)	47(3)	38(3)	-2(2)	14(2)	1(2)
C(8S)	48(4)	22(3)	73(4)	9(3)	8(3)	-3(2)
C(9S)	45(3)	31(3)	43(3)	11(2)	24(3)	7(2)
C(10S)	118(8)	88(6)	104(7)	-11(5)	58(6)	-22(6)
C(11S)	30(3)	25(2)	30(3)	3(2)	14(2)	-2(2)
C(12S)	40(3)	40(3)	30(3)	3(2)	11(2)	2(2)
C(13S)	29(3)	25(2)	26(2)	0(2)	11(2)	1(2)
C(14S)	35(3)	32(3)	37(3)	-2(2)	18(2)	-2(2)
C(15S)	38(3)	39(3)	35(3)	2(2)	19(2)	-2(2)
C(16S)	51(4)	50(3)	63(4)	5(3)	40(3)	4(3)

105

TABLE 5

Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for LMT·2EsOH.				
	x	y	z	U(eq)
H(1AA)	-585	2958	1019	35
H(2AA)	1142	2510	2886	30
H(3A)	-993	5379	587	30
H(2A)	-112	4894	1482	38
H(4A)	-1173	4519	414	32
H(5A)	-1024	3623	616	33
H(8A)	-239	2190	1412	29
H(9A)	327	1779	2005	30
H(11A)	801	3220	2506	34
H(13A)	956	2107	3668	55
H(13B)	1268	1770	3586	55
H(13C)	853	1592	3162	55
H(14A)	950	1606	2041	62
H(14B)	1350	1846	2447	62
H(14C)	1048	2153	1770	62
H(15A)	-399	5529	288	75
H(15B)	-759	5883	-21	75
H(15C)	-775	5281	-280	75
H(16A)	-638	5607	1762	49
H(16B)	-652	6075	1222	49
H(16C)	-306	5694	1555	49
H(1B)	2048	25	885	25
H(2BB)	3779	-238	2945	33
H(3B)	1492	2364	765	27
H(2B)	2408	2001	1444	26
H(4B)	1368	1494	658	26
H(5B)	1566	616	765	25
H(8B)	2428	-704	1523	28
H(9B)	3012	-1036	2140	29
H(11B)	3416	453	2358	28
H(13D)	3579	-889	3507	57
H(13E)	3953	-1033	3472	57
H(13F)	3582	-1293	2905	57
H(14D)	3650	-952	1829	43
H(14E)	4042	-744	2383	43
H(14F)	3758	-345	1811	43
H(15D)	2099	2713	612	50
H(15E)	1729	3039	363	50
H(15F)	1737	2500	-42	50
H(16D)	1800	2460	1997	53
H(16E)	1763	3014	1591	53
H(16F)	2137	2692	1883	53
H(1C)	4627	2587	1195	30
H(2CC)	6358	2390	3285	24
H(3C)	4060	4939	950	32
H(2C)	4971	4574	1649	25
H(4C)	3932	4060	845	30
H(5C)	4137	3182	1006	27
H(8C)	5015	1871	1821	26
H(9C)	5601	1557	2480	25
H(11C)	5987	3052	2673	25
H(13G)	6256	1619	2248	39
H(13H)	6636	1887	2761	39
H(13I)	6327	2230	2142	39
H(14G)	6168	1769	3891	36
H(14H)	6544	1621	3871	36
H(14I)	6177	1339	3323	36
H(15G)	4636	5189	631	70
H(15H)	4280	5544	406	70
H(15I)	4252	4970	55	70
H(16G)	4450	5092	2174	70
H(16H)	4333	5614	1683	70
H(16I)	4723	5354	1894	70
H(1D)	7088	320	1462	34
H(2D)	8828	-56	3133	33
H(3D)	6618	2711	1123	32
H(2DD)	7510	2283	1724	32
H(4D)	6459	1833	909	38
H(5D)	6639	950	1112	38
H(8D)	7452	-429	1911	28
H(9D)	8028	-812	2416	30
H(11D)	8478	655	2709	31
H(13J)	8654	-656	3792	65
H(13K)	9004	-828	3683	65
H(13L)	8618	-1085	3187	65

106

TABLE 5-continued

Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for LMT·2EsOH.				
	x	y	z	U(eq)
H(14J)	8622	-820	2060	46
H(14K)	9023	-587	2493	46
H(14L)	8707	-220	1935	46
H(15J)	7042	2819	377	61
H(15K)	6720	3214	295	61
H(15L)	6632	2612	33	61
H(16J)	7101	2976	2179	54
H(16K)	6965	3437	1582	54
H(16L)	7320	3101	1708	54
H(1S1)	354	568	117	74
H(1S2)	34	756	315	74
H(2S1)	348	1552	757	67
H(2S2)	230	1466	-93	67
H(2S3)	641	1363	479	67
H(3S1)	2296	4750	523	36
H(3S2)	2109	4415	-213	36
H(4S1)	1692	5133	-749	59
H(4S2)	2102	5315	-517	59
H(4S3)	1901	5469	-21	59
H(5S1)	2641	3235	241	44
H(5S2)	2548	3575	803	44
H(6S1)	3029	4152	984	48
H(6S2)	2760	4126	137	48
H(6S3)	3134	3806	444	48
H(7S1)	5283	2814	-129	47
H(7S2)	5465	3051	679	47
H(8S1)	5796	3675	290	81
H(8S2)	5368	3758	-148	81
H(8S3)	5587	3453	-521	81
H(9S1)	5428	806	551	46
H(9S2)	5579	1289	1119	46
H(10A)	6182	981	1462	150
H(10B)	5993	1119	613	150
H(10C)	6028	516	875	150
H(11E)	8109	3705	840	33
H(11F)	8195	3253	387	33
H(12A)	7581	2956	-55	57
H(12B)	7605	3553	-295	57
H(12C)	7500	3438	363	57
H(13M)	8341	797	1242	33
H(13N)	8118	1177	569	33
H(14M)	7744	447	-41	52
H(14N)	7716	589	698	52
H(14O)	7955	95	671	52
H(15M)	272	2476	-63	44
H(15N)	475	2669	763	44
H(16M)	317	3544	407	74
H(16N)	-43	3188	97	74
H(16O)	135	3371	-430	74

Crystallographic Data for LMT·2MsOH (FIG. 17c)

TABLE 1

Crystal data and structure refinement for LMT·2MsOH.	
Identification code	64412SC171
Empirical formula	C18H27N3O6S3
Formula weight	477.61
Temperature	150(2) K
Wavelength	0.71073 \text{\AA}
Crystal system	Triclinic
Space group	P-1
Unit cell dimensions	a = 11.6401(6) \text{\AA} α = 104.682(2)° b = 12.0744(6) \text{\AA} β = 92.386(2)° c = 18.4846(9) \text{\AA} γ = 116.151(2)°
Volume	2220.42(19) \text{\AA}^3
Z	4
Density (calculated)	1.429 Mg/m^3
Absorption coefficient	0.374 mm^-1
F(000)	1008
Crystal size	0.30 x 0.18 x 0.04 mm^3
Theta range for data collection	1.16 to 27.57°

107

TABLE 1-continued

Crystal data and structure refinement for LMT·2MsOH.	
Index ranges	-15 ≤ h ≤ 15, -15 ≤ k ≤ 15, -24 ≤ l ≤ 24
Reflections collected	42564
Independent reflections	10184 [R(int) = 0.0662]
Completeness to theta = 25.00°	99.6%
Absorption correction	Semi-empirical from equivalents
Max. and min. transmission	0.9852 and 0.8962
Refinement method	Full-matrix least-squares on F ²
Data/restraints/parameters	10184/198/552
Goodness-of-fit on F ²	1.071
Final R indices [I > 2sigma(I)]	R1 = 0.0593, wR2 = 0.1399
R indices (all data)	R1 = 0.0909, wR2 = 0.1566
Largest diff. peak and hole	1.192 and -0.905 e · Å ⁻³

TABLE 2

Atomic coordinates (×10 ⁴) and equivalent isotropic displacement parameters (Å ² × 10 ³) for eul1_0m. U(eq) is defined as one third of the trace of the orthogonalized U ^{ij} tensor.				
	x	y	z	U(eq)
C(1A)	2847(3)	7453(4)	3069(2)	28(1)
C(2A)	2545(3)	8117(3)	2643(2)	27(1)
C(3A)	3528(3)	9173(4)	2496(2)	29(1)
C(4A)	4823(3)	9566(4)	2760(2)	37(1)
C(5A)	5121(3)	8863(4)	3154(2)	39(1)
C(6A)	4156(3)	7809(4)	3317(2)	34(1)
C(7A)	3630(5)	5911(5)	3768(2)	48(1)
C(8A)	4139(5)	5179(5)	4015(2)	55(1)
C(9A)	3314(5)	4011(5)	4119(2)	58(1)
C(10A)	1978(5)	3531(5)	3953(2)	52(1)
C(11A)	1451(5)	4217(4)	3678(2)	45(1)
C(12A)	2292(4)	5408(4)	3601(2)	42(1)
C(13A)	3947(4)	10426(4)	1555(2)	39(1)
C(14A)	3035(4)	10981(4)	2698(2)	36(1)
C(15A)	479(5)	2617(4)	4788(2)	56(1)
C(16A)	422(9)	1431(7)	3338(3)	21(2)
C(16')	-175(7)	1295(6)	3509(4)	38(2)
C(1B)	1734(3)	3733(3)	1573(2)	20(1)
C(2B)	467(3)	2802(3)	1532(2)	20(1)
C(3B)	-556(3)	2949(3)	1228(2)	20(1)
C(4B)	-328(3)	4011(3)	986(2)	21(1)
C(5B)	938(3)	4959(3)	1054(2)	22(1)
C(6B)	1992(3)	4819(3)	1335(2)	21(1)
C(7B)	4382(3)	5861(3)	1707(2)	21(1)
C(8B)	5559(3)	6955(3)	1766(2)	22(1)
C(9B)	6724(3)	7111(3)	2126(2)	22(1)
C(10B)	6691(3)	6167(3)	2435(2)	20(1)
C(11B)	5535(3)	5073(3)	2382(2)	20(1)
C(12B)	4385(3)	4907(3)	2011(2)	20(1)
C(13B)	-2276(3)	1245(3)	1673(2)	27(1)
C(14B)	-2081(3)	892(3)	317(2)	27(1)
C(15B)	8932(3)	7573(3)	3209(2)	26(1)
C(16B)	8431(3)	5579(3)	2180(2)	29(1)
C(1S)	3536(4)	59(4)	4695(2)	38(1)
C(2S)	8797(4)	7948(3)	434(2)	33(1)
C(3S)	5403(4)	3853(4)	327(2)	40(1)
C(4S)	6718(6)	3037(5)	4356(2)	67(2)
N(1A)	4480(3)	7134(4)	3726(2)	46(1)
N(2A)	3126(3)	9915(3)	2118(2)	29(1)
N(3A)	1056(5)	2317(4)	4117(2)	68(1)
N(1B)	3245(2)	5734(3)	1338(2)	24(1)
N(2B)	-1895(2)	1871(3)	1060(1)	21(1)
N(3B)	7903(2)	6237(2)	2774(2)	21(1)
O(1S)	1876(5)	153(5)	5602(3)	31(1)
O(2S)	1457(5)	367(5)	4352(3)	27(1)
O(3S)	3166(4)	2156(4)	5329(3)	28(1)
O(2S')	3198(4)	1901(4)	4683(3)	26(1)
O(1S')	2515(5)	924(5)	5683(3)	25(1)
O(3S')	1291(5)	-191(6)	4380(3)	27(1)
O(4S)	10898(2)	10147(2)	1042(1)	34(1)
O(5S)	9224(3)	9462(3)	1796(2)	40(1)
O(6S)	10527(2)	8353(2)	1521(2)	39(1)
O(7S)	6954(2)	2997(3)	-257(1)	34(1)

108

TABLE 2-continued

Atomic coordinates (×10 ⁴) and equivalent isotropic displacement parameters (Å ² × 10 ³) for eul1_0m. U(eq) is defined as one third of the trace of the orthogonalized U ^{ij} tensor.				
	x	y	z	U(eq)
O(8S)	6130(2)	2435(2)	845(1)	31(1)
O(9S)	4703(3)	1484(2)	-383(2)	44(1)
O(10S)	7552(3)	4905(3)	3786(2)	53(1)
O(11S)	8403(4)	3384(4)	3483(2)	73(1)
O(12S)	6252(3)	2791(3)	2924(2)	57(1)
S(1)	2478(1)	713(1)	4948(1)	30(1)
S(2)	9944(1)	9074(1)	1257(1)	23(1)
S(3)	5818(1)	2588(1)	111(1)	22(1)
S(4)	7286(1)	3562(1)	3574(1)	26(1)
S(1A)	1567(1)	6322(1)	3376(1)	32(1)
S(1B)	2994(1)	3393(1)	1838(1)	26(1)

TABLE 3

Bond lengths [Å] and angles [°] for eul1_0m.	
C(1A)—C(2A)	1.390(5)
C(1A)—C(6A)	1.408(4)
C(1A)—S(1A)	1.753(4)
C(2A)—C(3A)	1.388(5)
C(2A)—H(2A)	0.9500
C(3A)—C(4A)	1.388(5)
C(3A)—N(2A)	1.472(5)
C(4A)—C(5A)	1.387(6)
C(4A)—H(4A)	0.9500
C(5A)—C(6A)	1.390(6)
C(5A)—H(5A)	0.9500
C(6A)—N(1A)	1.393(5)
C(7A)—C(12A)	1.386(6)
C(7A)—N(1A)	1.394(6)
C(7A)—C(8A)	1.409(5)
C(8A)—C(9A)	1.380(7)
C(8A)—H(8A)	0.9500
C(9A)—C(10A)	1.387(7)
C(9A)—H(9A)	0.9500
C(10A)—C(11A)	1.399(5)
C(10A)—N(3A)	1.502(7)
C(11A)—C(12A)	1.386(7)
C(11A)—H(11A)	0.9500
C(12A)—S(1A)	1.768(3)
C(13A)—N(2A)	1.506(4)
C(13A)—H(13A)	0.9800
C(13A)—H(13B)	0.9800
C(13A)—H(13C)	0.9800
C(14A)—N(2A)	1.499(5)
C(14A)—H(14A)	0.9800
C(14A)—H(14B)	0.9800
C(14A)—H(14C)	0.9800
C(15A)—N(3A)	1.477(6)
C(15A)—H(15A)	0.9800
C(15A)—H(15B)	0.9800
C(15A)—H(15C)	0.9800
C(16A)—N(3A)	1.482(6)
C(16A)—H(16H)	0.9800
C(16A)—H(16I)	0.9800
C(16A)—H(16J)	0.9800
C(16')—N(3A)	1.563(6)
C(16')—H(16K)	0.9800
C(16')—H(16L)	0.9800
C(16')—H(16M)	0.9800
C(1B)—C(2B)	1.390(4)
C(1B)—C(6B)	1.398(4)
C(1B)—S(1B)	1.770(3)
C(2B)—C(3B)	1.394(4)
C(2B)—H(2B)	0.9500
C(3B)—C(4B)	1.384(4)
C(3B)—N(2B)	1.479(4)
C(4B)—C(5B)	1.386(4)
C(4B)—H(4B)	0.9500
C(5B)—C(6B)	1.407(4)
C(5B)—H(5B)	0.9500

TABLE 3-continued

Bond lengths [Å] and angles [°] for eul1_0m.	
C(6B)—N(1B)	1.387(4)
C(7B)—N(1B)	1.391(4)
C(7B)—C(8B)	1.397(4)
C(7B)—C(12B)	1.405(4)
C(8B)—C(9B)	1.398(4)
C(8B)—H(8B)	0.9500
C(9B)—C(10B)	1.385(4)
C(9B)—H(9B)	0.9500
C(10B)—C(11B)	1.387(4)
C(10B)—N(3B)	1.478(4)
C(11B)—C(12B)	1.386(4)
C(11B)—H(11B)	0.9500
C(12B)—S(1B)	1.766(3)
C(13B)—N(2B)	1.494(4)
C(13B)—H(13D)	0.9800
C(13B)—H(13E)	0.9800
C(13B)—H(13F)	0.9800
C(14B)—N(2B)	1.503(4)
C(14B)—H(14D)	0.9800
C(14B)—H(14E)	0.9800
C(14B)—H(14F)	0.9800
C(15B)—N(3B)	1.497(4)
C(15B)—H(15D)	0.9800
C(15B)—H(15E)	0.9800
C(15B)—H(15F)	0.9800
C(16B)—N(3B)	1.503(4)
C(16B)—H(16A)	0.9800
C(16B)—H(16B)	0.9800
C(16B)—H(16C)	0.9800
C(1S)—S(1)	1.755(4)
C(1S)—H(1S1)	0.9800
C(1S)—H(1S2)	0.9800
C(1S)—H(1S3)	0.9800
C(2S)—S(2)	1.768(3)
C(2S)—H(2S1)	0.9800
C(2S)—H(2S2)	0.9800
C(2S)—H(2S3)	0.9800
C(3S)—S(3)	1.755(4)
C(3S)—H(3S1)	0.9800
C(3S)—H(3S2)	0.9800
C(3S)—H(3S3)	0.9800
C(4S)—S(4)	1.762(4)
C(4S)—H(4S1)	0.9800
C(4S)—H(4S2)	0.9800
C(4S)—H(4S3)	0.9800
N(1A)—H(1A)	0.8800
N(2A)—H(2A1)	0.9300
N(3A)—H(3A)	0.9300
N(1B)—H(1B)	0.8800
N(2B)—H(2B1)	0.9300
N(3B)—H(3B)	0.9300
O(1S)—S(1)	1.573(5)
O(2S)—S(1)	1.422(5)
O(3S)—S(1)	1.508(5)
O(2S')—S(1)	1.528(5)
O(1S')—S(1)	1.312(5)
O(3S')—S(1)	1.473(5)
O(4S)—S(2)	1.449(2)
O(5S)—S(2)	1.447(3)
O(6S)—S(2)	1.472(2)
O(7S)—S(3)	1.458(2)
O(8S)—S(3)	1.467(2)
O(9S)—S(3)	1.433(3)
O(10S)—S(4)	1.451(3)
O(11S)—S(4)	1.417(3)
O(12S)—S(4)	1.443(3)
C(2A)—C(1A)—C(6A)	119.8(3)
C(2A)—C(1A)—S(1A)	117.7(2)
C(6A)—C(1A)—S(1A)	122.0(3)
C(3A)—C(2A)—C(1A)	120.3(3)
C(3A)—C(2A)—H(2A)	119.9
C(1A)—C(2A)—H(2A)	119.9
C(2A)—C(3A)—C(4A)	120.5(4)
C(2A)—C(3A)—N(2A)	116.9(3)
C(4A)—C(3A)—N(2A)	122.3(3)
C(5A)—C(4A)—C(3A)	118.9(4)
C(5A)—C(4A)—H(4A)	120.5
C(3A)—C(4A)—H(4A)	120.5

TABLE 3-continued

Bond lengths [Å] and angles [°] for eul1_0m.	
C(4A)—C(5A)—C(6A)	121.7(3)
C(4A)—C(5A)—H(5A)	119.1
C(6A)—C(5A)—H(5A)	119.1
C(5A)—C(6A)—N(1A)	120.6(3)
C(5A)—C(6A)—C(1A)	118.6(4)
N(1A)—C(6A)—C(1A)	120.8(4)
C(12A)—C(7A)—N(1A)	122.0(3)
C(12A)—C(7A)—C(8A)	118.7(5)
C(1A)—C(7A)—C(8A)	119.3(4)
C(9A)—C(8A)—C(7A)	120.2(5)
C(9A)—C(8A)—H(8A)	119.9
C(7A)—C(8A)—H(8A)	119.9
C(8A)—C(9A)—C(10A)	120.1(4)
C(8A)—C(9A)—H(9A)	120.0
C(10A)—C(9A)—H(9A)	120.0
C(9A)—C(10A)—C(11A)	120.6(5)
C(9A)—C(10A)—N(3A)	121.4(4)
C(11A)—C(10A)—N(3A)	117.8(4)
C(12A)—C(11A)—C(10A)	118.5(4)
C(12A)—C(11A)—H(11A)	120.7
C(10A)—C(11A)—H(11A)	120.7
C(7A)—C(12A)—C(11A)	121.8(4)
C(7A)—C(12A)—S(1A)	121.7(4)
C(11A)—C(12A)—S(1A)	116.2(3)
N(2A)—C(13A)—H(13A)	109.5
N(2A)—C(13A)—H(13B)	109.5
H(13A)—C(13A)—H(13B)	109.5
N(2A)—C(13A)—H(13C)	109.5
H(13A)—C(13A)—H(13C)	109.5
H(13B)—C(13A)—H(13C)	109.5
N(2A)—C(14A)—H(14A)	109.5
N(2A)—C(14A)—H(14B)	109.5
H(14A)—C(14A)—H(14B)	109.5
N(2A)—C(14A)—H(14C)	109.5
H(14A)—C(14A)—H(14C)	109.5
H(14B)—C(14A)—H(14C)	109.5
N(3A)—C(15A)—H(15A)	109.5
N(3A)—C(15A)—H(15B)	109.5
H(15A)—C(15A)—H(15B)	109.5
N(3A)—C(15A)—H(15C)	109.5
H(15A)—C(15A)—H(15C)	109.5
H(15B)—C(15A)—H(15C)	109.5
N(3A)—C(16A)—H(16H)	109.5
N(3A)—C(16A)—H(16I)	109.5
N(3A)—C(16A)—H(16J)	109.5
N(3A)—C(16')—H(16K)	109.5
N(3A)—C(16')—H(16L)	109.5
H(16K)—C(16')—H(16L)	109.5
N(3A)—C(16')—H(16M)	109.5
H(16K)—C(16')—H(16M)	109.5
H(16L)—C(16')—H(16M)	109.5
C(2B)—C(1B)—C(6B)	121.3(3)
C(2B)—C(1B)—S(1B)	116.9(2)
C(6B)—C(1B)—S(1B)	121.5(2)
C(1B)—C(2B)—C(3B)	118.7(3)
C(1B)—C(2B)—H(2B)	120.7
C(3B)—C(2B)—H(2B)	120.7
C(4B)—C(3B)—C(2B)	121.2(3)
C(4B)—C(3B)—N(2B)	118.7(3)
C(2B)—C(3B)—N(2B)	119.5(3)
C(3B)—C(4B)—C(5B)	119.7(3)
C(3B)—C(4B)—H(4B)	120.1
C(5B)—C(4B)—H(4B)	120.1
C(4B)—C(5B)—C(6B)	120.4(3)
C(4B)—C(5B)—H(5B)	119.8
C(6B)—C(5B)—H(5B)	119.8
N(1B)—C(6B)—C(1B)	122.5(3)
N(1B)—C(6B)—C(5B)	118.7(3)
C(1B)—C(6B)—C(5B)	118.7(3)
N(1B)—C(7B)—C(8B)	119.3(3)
N(1B)—C(7B)—C(12B)	121.7(3)
C(8B)—C(7B)—C(12B)	119.0(3)
C(7B)—C(8B)—C(9B)	120.9(3)
C(7B)—C(8B)—H(8B)	119.5
C(9B)—C(8B)—H(8B)	119.5
C(10B)—C(9B)—C(8B)	118.7(3)
C(10B)—C(9B)—H(9B)	120.6
C(8B)—C(9B)—H(9B)	120.6

111

TABLE 3-continued

Bond lengths [Å] and angles [°] for eul1_0m.	
C(9B)—C(10B)—C(11B)	121.4(3)
C(9B)—C(10B)—N(3B)	121.0(3)
C(11B)—C(10B)—N(3B)	117.4(3)
C(12B)—C(11B)—C(10B)	119.7(3)
C(12B)—C(11B)—H(11B)	120.1
C(10B)—C(11B)—H(11B)	120.1
C(11B)—C(12B)—C(7B)	120.2(3)
C(11B)—C(12B)—S(1B)	117.6(2)
C(7B)—C(12B)—S(1B)	121.8(2)
N(2B)—C(13B)—H(13D)	109.5
N(2B)—C(13B)—H(13E)	109.5
H(13D)—C(13B)—H(13E)	109.5
N(2B)—C(13B)—H(13F)	109.5
H(13D)—C(13B)—H(13F)	109.5
H(13E)—C(13B)—H(13F)	109.5
N(2B)—C(14B)—H(14D)	109.5
N(2B)—C(14B)—H(14E)	109.5
H(14D)—C(14B)—H(14E)	109.5
N(2B)—C(14B)—H(14F)	109.5
H(14D)—C(14B)—H(14F)	109.5
H(14E)—C(14B)—H(14F)	109.5
N(3B)—C(15B)—H(15D)	109.5
N(3B)—C(15B)—H(15E)	109.5
H(15D)—C(15B)—H(15E)	109.5
N(3B)—C(15B)—H(15F)	109.5
H(15D)—C(15B)—H(15F)	109.5
H(15E)—C(15B)—H(15F)	109.5
N(3B)—C(16B)—H(16A)	109.5
N(3B)—C(16B)—H(16B)	109.5
H(16A)—C(16B)—H(16B)	109.5
N(3B)—C(16B)—H(16C)	109.5
H(16A)—C(16B)—H(16C)	109.5
H(16B)—C(16B)—H(16C)	109.5
S(1)—C(1S)—H(1S1)	109.5
S(1)—C(1S)—H(1S2)	109.5
H(1S1)—C(1S)—H(1S2)	109.5
S(1)—C(1S)—H(1S3)	109.5
H(1S1)—C(1S)—H(1S3)	109.5
H(1S2)—C(1S)—H(1S3)	109.5
S(2)—C(2S)—H(2S1)	109.5
S(2)—C(2S)—H(2S2)	109.5
H(2S1)—C(2S)—H(2S2)	109.5
S(2)—C(2S)—H(2S3)	109.5
H(2S1)—C(2S)—H(2S3)	109.5
H(2S2)—C(2S)—H(2S3)	109.5
S(3)—C(3S)—H(3S1)	109.5
S(3)—C(3S)—H(3S2)	109.5
H(3S1)—C(3S)—H(3S2)	109.5
S(3)—C(3S)—H(3S3)	109.5
H(3S1)—C(3S)—H(3S3)	109.5
H(3S2)—C(3S)—H(3S3)	109.5
S(4)—C(4S)—H(4S1)	109.5
S(4)—C(4S)—H(4S2)	109.5
H(4S1)—C(4S)—H(4S2)	109.5
S(4)—C(4S)—H(4S3)	109.5
H(4S1)—C(4S)—H(4S3)	109.5
H(4S2)—C(4S)—H(4S3)	109.5
C(6A)—N(1A)—C(7A)	124.8(3)
C(6A)—N(1A)—H(1A)	117.6
C(7A)—N(1A)—H(1A)	117.6
C(3A)—N(2A)—C(14A)	110.2(3)
C(3A)—N(2A)—C(13A)	114.9(3)
C(14A)—N(2A)—C(13A)	111.1(3)
C(3A)—N(2A)—H(2A1)	106.7
C(14A)—N(2A)—H(2A1)	106.7
C(13A)—N(2A)—H(2A1)	106.7
C(15A)—N(3A)—C(10A)	111.1(3)
C(15A)—N(3A)—C(16A)	130.2(6)
C(10A)—N(3A)—C(16A)	101.3(4)
C(15A)—N(3A)—C(16')	102.0(5)
C(10A)—N(3A)—C(16')	119.1(4)
C(16A)—N(3A)—C(16')	28.2(3)
C(15A)—N(3A)—H(3A)	103.9
C(10A)—N(3A)—H(3A)	103.9
C(16A)—N(3A)—H(3A)	103.9
C(16')	116.0
C(16')—N(3A)—H(3A)	116.0
C(6B)—N(1B)—C(7B)	125.8(3)
C(6B)—N(1B)—H(1B)	117.1

112

TABLE 3-continued

Bond lengths [Å] and angles [°] for eul1_0m.	
C(7B)—N(1B)—H(1B)	117.1
C(3B)—N(2B)—C(13B)	114.9(2)
C(3B)—N(2B)—C(14B)	109.1(2)
C(13B)—N(2B)—C(14B)	111.1(3)
C(3B)—N(2B)—H(2B1)	107.1
C(13B)—N(2B)—H(2B1)	107.1
C(14B)—N(2B)—H(2B1)	107.1
C(10B)—N(3B)—C(15B)	114.9(2)
C(10B)—N(3B)—C(16B)	110.6(2)
C(15B)—N(3B)—C(16B)	110.8(2)
C(10B)—N(3B)—H(3B)	106.7
C(15B)—N(3B)—H(3B)	106.7
C(16B)—N(3B)—H(3B)	106.7
O(1S')—S(1)—O(2S)	131.2(3)
O(1S')—S(1)—O(3S')	123.5(3)
O(2S)—S(1)—O(3S)	25.1(2)
O(1S')—S(1)—O(3S)	71.7(3)
O(2S)—S(1)—O(3S)	110.8(3)
O(3S')—S(1)—O(3S)	134.9(3)
O(1S')—S(1)—O(2S')	116.5(3)
O(2S)—S(1)—O(2S')	84.3(3)
O(3S')—S(1)—O(2S')	107.5(3)
O(3S)—S(1)—O(2S')	45.0(2)
O(1S')—S(1)—O(1S)	33.5(2)
O(2S)—S(1)—O(1S)	109.1(3)
O(3S')—S(1)—O(1S)	93.0(3)
O(3S)—S(1)—O(1S)	103.3(3)
O(2S')—S(1)—O(1S)	148.0(3)
O(1S')—S(1)—C(1S)	107.0(2)
O(2S)—S(1)—C(1S)	114.7(2)
O(3S')—S(1)—C(1S)	102.3(2)
O(3S)—S(1)—C(1S)	113.6(2)
O(2S')—S(1)—C(1S)	95.2(2)
O(1S)—S(1)—C(1S)	104.4(2)
O(4S)—S(2)—O(5S)	112.76(15)
O(4S)—S(2)—O(6S)	111.68(16)
O(5S)—S(2)—O(6S)	112.07(17)
O(4S)—S(2)—C(2S)	108.24(17)
O(5S)—S(2)—C(2S)	106.71(17)
O(6S)—S(2)—C(2S)	104.86(16)
O(9S)—S(3)—O(7S)	112.41(17)
O(9S)—S(3)—O(8S)	113.91(16)
O(7S)—S(3)—O(8S)	111.12(14)
O(9S)—S(3)—C(3S)	105.99(19)
O(7S)—S(3)—C(3S)	107.38(18)
O(8S)—S(3)—C(3S)	105.42(16)
O(11S)—S(4)—O(12S)	112.8(2)
O(11S)—S(4)—O(10S)	114.1(2)
O(12S)—S(4)—O(10S)	110.8(2)
O(11S)—S(4)—C(4S)	106.2(3)
O(12S)—S(4)—C(4S)	107.6(2)
O(10S)—S(4)—C(4S)	104.6(2)
C(1A)—S(1A)—C(12A)	100.76(19)
C(12B)—S(1B)—C(1B)	101.93(14)

Symmetry transformations used to generate equivalent atoms:

TABLE 4

Anisotropic displacement parameters ($\text{Å}^2 \times 10^3$) for eul1_0m.
The anisotropic displacement factor exponent takes the form: $-2\pi^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
C(1A)	22(1)	45(2)	19(1)	2(1)	1(1)	21(1)
C(2A)	17(1)	41(2)	20(1)	4(1)	0(1)	14(1)
C(3A)	19(1)	46(2)	15(1)	2(1)	1(1)	12(1)
C(4A)	19(1)	57(2)	19(1)	-4(1)	2(1)	12(1)
C(5A)	19(1)	66(2)	20(1)	-6(1)	-2(1)	21(1)
C(6A)	23(1)	59(2)	17(1)	-4(1)	-2(1)	26(1)
C(7A)	75(2)	90(2)	15(1)	8(2)	5(1)	73(2)
C(8A)	86(2)	98(2)	19(1)	6(2)	0(1)	83(2)
C(9A)	101(2)	92(2)	20(2)	6(2)	-2(2)	86(2)
C(10A)	100(2)	76(2)	18(1)	9(1)	3(2)	78(2)
C(11A)	86(2)	71(2)	16(1)	14(1)	7(1)	69(2)

TABLE 4-continued

Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for eul1_0m. The anisotropic displacement factor exponent takes the form: $-2\pi^2[h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$						
	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
C(12A)	75(2)	75(2)	15(1)	15(1)	8(1)	67(2)
C(13A)	27(2)	44(2)	20(2)	8(2)	4(2)	-3(2)
C(14A)	39(2)	30(2)	25(2)	7(2)	10(2)	5(2)
C(15A)	104(4)	34(2)	28(2)	4(2)	-10(2)	34(3)
C(1B)	17(1)	22(1)	18(1)	4(1)	3(1)	10(1)
C(2B)	21(1)	22(1)	18(1)	6(1)	4(1)	10(1)
C(3B)	19(1)	23(1)	19(1)	5(1)	4(1)	10(1)
C(4B)	20(1)	23(1)	22(1)	6(1)	3(1)	12(1)
C(5B)	22(1)	21(1)	24(1)	7(1)	4(1)	11(1)
C(6B)	18(1)	22(1)	23(1)	5(1)	6(1)	10(1)
C(7B)	21(1)	20(1)	23(1)	6(1)	6(1)	11(1)
C(8B)	23(1)	19(1)	26(1)	6(1)	6(1)	11(1)
C(9B)	21(1)	19(1)	26(1)	5(1)	7(0)	9(1)
C(10B)	19(1)	21(1)	19(1)	3(1)	4(1)	10(1)
C(11B)	20(1)	21(1)	20(1)	5(1)	5(1)	10(1)
C(12B)	18(1)	20(1)	20(1)	6(1)	5(1)	8(1)
C(13B)	21(2)	30(2)	24(2)	10(2)	4(1)	7(1)
C(14B)	24(2)	26(2)	21(2)	0(1)	0(1)	6(1)
C(15B)	24(2)	22(2)	25(2)	0(1)	0(1)	8(1)
C(16B)	27(2)	31(2)	27(2)	0(2)	2(1)	18(2)
C(1S)	33(2)	49(2)	28(2)	7(2)	0(2)	20(2)
C(2S)	35(2)	29(2)	26(2)	2(2)	-6(2)	12(2)
C(3S)	62(3)	44(2)	31(2)	12(2)	9(2)	40(2)
C(4S)	95(4)	51(3)	30(2)	16(2)	22(2)	11(3)
N(1A)	42(2)	88(3)	23(2)	2(2)	-6(1)	52(2)
N(2A)	20(1)	32(2)	19(1)	5(1)	2(1)	0(1)
N(3A)	159(4)	58(2)	17(2)	1(2)	-12(2)	84(3)
N(1B)	18(1)	25(1)	34(2)	17(1)	7(1)	9(1)
N(2B)	17(1)	25(1)	21(1)	6(1)	2(1)	10(1)
N(3B)	18(1)	20(1)	21(1)	2(1)	1(1)	8(1)
O(4S)	39(2)	25(1)	33(1)	10(1)	12(1)	9(1)
O(5S)	37(2)	35(2)	34(2)	0(1)	14(1)	10(1)
O(6S)	30(1)	27(1)	54(2)	14(1)	-11(1)	8(1)
O(7S)	27(1)	48(2)	42(2)	28(1)	16(1)	22(1)
O(8S)	33(1)	42(1)	34(1)	23(1)	13(1)	25(1)
O(9S)	33(2)	28(1)	53(2)	7(1)	-5(1)	3(1)
O(10S)	96(2)	27(1)	26(1)	4(1)	-11(2)	24(2)
O(11S)	84(3)	128(3)	45(2)	25(2)	15(2)	82(3)
O(12S)	43(2)	54(2)	28(2)	2(1)	-5(1)	-11(1)
S(1)	21(1)	19(1)	45(1)	13(1)	-12(1)	5(1)
S(2)	23(1)	22(1)	20(1)	5(1)	3(1)	9(1)
S(3)	19(1)	22(1)	28(1)	10(1)	6(1)	11(1)
S(4)	29(1)	22(1)	18(1)	6(1)	2(1)	4(1)
S(1A)	29(1)	43(1)	39(1)	19(1)	5(1)	26(1)
S(1B)	17(1)	22(1)	39(1)	15(1)	1(1)	7(1)

TABLE 5

Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for eul1_0m.				
	x	y	z	U(eq)
H(2A)	1662	7847	2452	32
H(4A)	5495	10304	2671	45
H(5A)	6007	9108	3318	47
H(8A)	5053	5489	4110	66
H(9A)	3662	3536	4304	69
H(11A)	536	3874	3547	54
H(13A)	4800	11131	1827	58
H(13B)	3517	10752	1260	58
H(13C)	4057	9728	1211	58
H(14A)	2513	10627	3064	54
H(14B)	2623	11370	2445	54
H(14C)	3910	11643	2966	54
H(15A)	1173	3266	5213	84
H(15B)	-13	1828	4928	84
H(15C)	-105	2958	4669	84
H(16H)	-60	541	3357	32
H(16I)	1086	1487	3019	32
H(16J)	-179	1679	3122	32

TABLE 5-continued

Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for eul1_0m.				
	x	y	z	U(eq)
H(16K)	-683	569	3701	56
H(16L)	94	977	3039	56
H(16M)	-708	1698	3407	56
H(2B)	301	2081	1708	24
H(4B)	-1035	4090	775	25
H(5B)	1095	5709	911	26
H(8B)	5567	7602	1559	27
H(9B)	7524	7850	2158	27
H(11B)	5531	4439	2601	24
H(13D)	-1727	843	1744	41
H(13E)	-3189	582	1529	41
H(13F)	-2161	1901	2148	41
H(14D)	-1922	1307	-87	41
H(14E)	-2973	184	197	41
H(14F)	-1468	547	355	41
H(15D)	9282	8071	2855	39
H(15E)	9634	7519	3485	39
H(15F)	8555	8004	3570	39
H(16A)	7739	4713	1898	43
H(16B)	9145	5506	2426	43
H(16C)	8752	6089	1829	43
H(1S1)	4324	513	5083	56
H(1S2)	3770	163	4204	56
H(1S3)	3106	-863	4655	56
H(2S1)	8396	8376	210	50
H(2S2)	8124	7229	571	50
H(2S3)	9237	7616	64	50
H(3S1)	5170	3997	-144	60
H(3S2)	6146	4646	652	60
H(3S3)	4661	3623	594	60
H(4S1)	6520	2129	4255	100
H(4S2)	5930	3121	4436	100
H(4S3)	7389	3569	4811	100
H(1A)	5282	7506	3975	55
H(2A1)	2289	9350	1848	35
H(3A)	1596	2011	4279	82
H(1B)	3327	6282	1083	29
H(2B1)	-2456	2212	994	26
H(3B)	7679	5766	3119	25

References

- 40 Abrahamson, M., Jonsdottir, S., Olafsson, I. & Grubb, A. (1992) Hereditary cystatin C amyloid angiopathy identification of the disease-causing mutation and specific diagnosis by polymerase chain reaction based analysis. *Human Genetics* 89, 377-380.
- 45 Andersen, P. (2006) Amyotrophic lateral sclerosis associated with mutations in the CuZn superoxide dismutase gene. *Current Neurology and Neuroscience Reports* 6, 37-46.
- Arai, T., Hasegawa, M., Nonoka, T., Kametani, F., Yamashita, M., Hosokawa, M., Niizato, K., Tsuchiya, K., Kobayashi, Z., Ikeda, K., Yoshida, M., Onaya, M., Fujishiro, H. & Akiyama, H. (2010) Phosphorylated and cleaved TDP-43 in ALS, FTLN and other neurodegenerative disorders and in cellular models of TDP-43 proteinopathy. *Neuropathology* 30, 170-181.
- 55 Askanas, V., Engel, W. K. & Nogalska, A. (2009) Inclusion body myositis: a degenerative muscle disease associated with intra-muscle fiber multi-protein aggregates, proteasome inhibition, endoplasmic reticulum stress and decreased lysosomal degradation. *Brain Pathology* 19, 493-506.
- 60 Barmada, S. J., Skibinski, G., Korb, E., Rao, E. J., Wu, J. Y. & Finkbeiner, S. (2010) Cytoplasmic mislocalization of TDP-43 is toxic to neurons and enhanced by a mutation associated with familial amyotrophic lateral sclerosis. *Journal of Neuroscience* 30, 639-649.
- 65 Blair, I. P., Williams, K. L., Warraich, S. T., Durnall, J. C., Thoeng, A. D., Manavis, J., Blumbergs, P. C., Vucic, S.,

- Kiernan, M. C. & Nicholson, G. A. (2010) FUS mutations in amyotrophic lateral sclerosis: clinical, pathological, neurophysiological and genetic analysis. *Journal of Neurology Neurosurgery and Psychiatry* 81, 639-645.
- Booth, D. R., Sunde, M., Bellotti, V., Robinson, C. V., Hutchinson, W. L., Fraser, P. E., Hawkins, P. N., Dobson, C. M., Radford, S. E., Blake, C. C. F. & Pepys, M. B. (1997) Instability, unfolding and aggregation of human lysozyme variants underlying amyloid fibrillogenesis. *Nature* 385, 787-793.
- Byrne, S., Walsh, C., Lynch, C., Bede, P., Elamin, M., Kenna, K., McLaughlin, R. & Hardiman, O. (2011) Rate of familial amyotrophic lateral sclerosis: a systematic review and meta-analysis. *Journal of Neurology, Neurosurgery & Psychiatry* 82, 623-627.
- Carrell, R. W. & Gooptu, B. (1998) Conformational changes and disease—serpins, prions and Alzheimer's. *Current Opinion in Structural Biology* 8, 799-809.
- Chen-Plotkin, A. S., Lee, V. M. Y. & Trojanowski, J. Q. (2010) TAR DNA-binding protein 43 in neurodegenerative disease. *Nature Reviews Neurology* 6, 211-220.
- Chiti, F., Webster, P., Taddei, N., Clark, A., Stefani, M., Ramponi, G. & Dobson, C. (1999) Designing conditions for in vitro formation of amyloid protofilaments and fibrils. *Proceedings of the National Academy of Sciences, USA* 96, 3590-3594.
- Cox, L. E., Ferraiuolo, L., Goodall, E. F., Heath, P. R., Higginbottom, A., Mortiboys, H., Hollinger, H. C., Hartley, J. A., Brockington, A., Burness, C. E., Morrison, K. E., Wharton, S. B., Grierson, A. J., Ince, P. G., Kirby, J. & Shaw, P. J. (2010) Mutations in CHMP2B in lower motor neuron predominant amyotrophic lateral sclerosis (ALS). *PLOS One* 5, e9872.
- Czech, C., Tremp, G. & Pradier, L. (2000) Presenilins and Alzheimer's disease: biological functions and pathogenic mechanisms. *Progress in Neurobiology* 60, 363-384.
- Davis, R. L., Shrimpton, A. E., Holohan, P. D., Bradshaw, C., Feiglin, D., Collins, G. H., Sonderegger, P., Kinter, J., Becker, L. M., Lachawan, F., Krasnewich, D., Muenke, M., Lawrence, D. A., Yerby, M. S., Shaw, C.-M., Gooptu, B., Elliott, P. R., Finch, J. T., Carrell, R. W. & Lomas, D. A. (1999) Familial dementia caused by polymerization of mutant neuroserpin. *Nature* 401, 376-379.
- DiFiglia, M., Sapp, E., Chase, K. O., Davies, S. W., Bates, G. P., Vonsattel, J. P. & Aronin, N. (1997) Aggregation of huntingtin in neuronal intranuclear inclusions and dystrophic neurites in brain. *Science* 277, 1990-1993.
- Dische, F. E., Wernstedt, C., Westermark, G. T., Westermark, P., Pepys, M. B., Rennie, J. A., Gilbey, S. G. & Watkins, P. J. (1988) Insulin as an amyloid-fibril protein at sites of repeated insulin injections in a diabetic patient. *Diabetologia* 31, 158-161.
- Elden, A. C., Kim, H.-J., Hart, M. P., Chen-Plotkin, A. S., Johnson, B. S., Fang, X., Arakola, M., Geser, F., Greene, R., Lu, M. M., Padmanabhan, A., Clay-Falcone, D., McCluskey, L., Elman, L., Juhr, D., Gruber, P. J., Rub, U., Auburger, G., Trojanowski, J. Q., Lee, V. M. Y., Van Deerlin, V. M., Bonini, N. M. & Gitler, A. D. (2010) Ataxin-2 intermediate-length polyglutamine expansions are associated with increased risk for ALS. *Nature* 466, 1069-1075.
- Finsterer, J. (2009) Mitochondrial disorders, cognitive impairment and dementia. *J. Neurol. Sci* 283:143-148
- Gasset, M., Bladwin, M. A., Lloyd, D. H., abriel, J.-M., Holtzman, D. M., Cohen, F. E., Fletterick, R. & Prusiner, S. B. (1992) Predicted a-helical region of the prion protein

- when synthesized as peptides form amyloid. *Proceedings of the National Academy of Sciences, USA* 89, 10940-10944.
- Gendron, T. F., Josephs, K. A. & Petrucelli, L. (2010) Review: Transactive response DNA-binding protein 43 (TDP-43): mechanisms of neurodegeneration. *Neuropathology and Applied Neurobiology* 36, 97-112.
- Geser, F., Lee, V. M.-Y. & Trojanowski, J. Q. (2010) Amyotrophic lateral sclerosis and frontotemporal lobar degeneration: A spectrum of TDP-43 proteinopathies. *Neuropathology* 30, 103-112.
- Gitcho, M. A., Baloh, R. H., Chakraverty, S., Mayo, K., Norton, J. B., Levitch, D., Hatanpaa, K. J., White, C. L., Ill, Bigio, E. H., Caselli, R., Baker, M., Al-Lozi, M. T., Morris, J. C., Pestronk, A., Rademakers, R., Goate, A. M. & Cairns, N. J. (2008) TDP-43 A315T mutation in familial motor neuron disease. *Annals of Neurology* 63, 535-538.
- Glennner, G. G. & Wong, C. W. (1984) Alzheimer's disease: initial report of the purification and characterisation of a novel cerebrovascular amyloid protein. *Biochemical and Biophysical Research Communications* 120, 885-890.
- Goate, A., Chartier-Harlin, M.-C., Mullan, M., Brown, J., Crawford, F., Fidani, L., Giuffra, L., Haynes, A., Irving, N., James, L., Mant, R., Newton, P., Rooke, K., Roques, P., Talbot, C., Pericak-Vance, M., Roses, A., Williamson, R., Rossor, M., Owen, M. & Hardy, J. (1991) Segregation of a missense mutation in the amyloid precursor protein gene with familial Alzheimer's disease. *Nature* 349, 704-706.
- Gorevic, P. D., Casey, T. T., Stone, W. J., DiRaimondo, C. R., Prelli, F. C. & Frangione, B. (1985) b-2 Microglobulin is an amyloidogenic protein in man. *Journal of Clinical Investigation* 76, 2425-2429.
- Gustavsson, A., Engstrim, U. & Westermark, P. (1991) Normal transthyretin and synthetic transthyretin fragments form amyloid-like fibrils in vitro. *Biochemical and Biophysical Research Communications* 175, 1159-1164.
- Higashi, S., Tsuchiya, Y., Araki, T., Wada, K. & Kabuta, T. (2010) TDP-43 physically interacts with amyotrophic lateral sclerosis-linked mutant CuZn superoxide dismutase. *Neurochemistry International* 57, 906-913.
- Hutton, M., Lendon, C., Rizzu, P., Baker, M., Froelich, S., Houlden, H., Pickering-Brown, S., Chakraverty, S., Isaacs, A., Grover, A., Hackett, J., Adamson, J., Lincoln, S., Dickson, D., Davies, P., Petersen, R. C., Stevens, M., de Graaf, E., Wauters, E., van Baren, J., Hillebrand, M., Joosse, M., Kwon, J. M., Nowotny, P., Che, L. K., Norton, J., Morris, J. C., Reed, L. A., Trojanowski, J. Q., Basun, H., Lannfelt, L., Neystat, M., Fahn, S., Dark, F., Tannenberg, T., Dodd, P. R., Hayward, N., Kwok, J. B. J., Schofield, P. R., Andreadis, A., Snowden, J., Craufurd, D., Neary, D., Owen, F., Oostra, B. A., Hardy, J., Goate, A., van Swieten, J., Mann, D., Lynch, T. & Heutink, P. (1998) Association of missense and 5'-splice-site mutations in tau with the inherited dementia FTDP-17. *Nature* 393, 702-705.
- Igaz, L. M., Kwong, L. K., Chen-Plotkin, A., Winton, M. J., Unger, T. L., Xu, Y., Neumann, M., Trojanowski, J. Q. & Lee, V. M. Y. (2009) Expression of TDP-43 C-terminal fragments in vitro recapitulates pathological features of TDP-43 proteinopathies. *Journal of Biological Chemistry* 284, 8516-8524.
- Jinwal, U K, Miyata, Y, Koren, J, III, Jones, J R, Trotter, J H et al. (2009) Chemical manipulation of Hsp70 ATPase activity regulates tau stability. *J. Neurosci.* 29:12079-12088

- Johansson, B., Wernstedt, C. & Westermark, P. (1987) Atrial natriuretic peptide deposited as atrial amyloid fibrils. *Biochemical and Biophysical Research Communications* 148, 1087-1092.
- Johnson, B. S., McCaffery, J. M., Lindquist, S. & Gitler, A. D. (2008) A yeast TDP-43 proteinopathy model: Exploring the molecular determinants of TDP-43 aggregation and cellular toxicity. *Proceedings of the National Academy of Sciences* 105, 6439-6444.
- Johnson, B. S., Snead, D., Lee, J. J., McCaffery, J. M., Shorter, J. & Gitler, A. D. (2009) TDP-43 is intrinsically aggregation-prone, and amyotrophic lateral sclerosis-linked mutations accelerate aggregation and increase toxicity. *Journal of Biological Chemistry* 284, 20329-20339.
- Johnson, J. O., Mandrioli, J., Benatar, M., Abramzon, Y., Van Deerlin, V. M., Trojanowski, J. Q., Gibbs, J. R., Brunetti, M., Gronka, S., Wu, J., Ding, J., McCluskey, L., Martinez-Lage, M., Falcone, D., Hernandez, D. G., Arepalli, S., Chong, S., Schymick, J. C., Rothstein, J., Landi, F., Wang, Y.-D., Calvo, A., Mora, G., Sabatelli, M., Monsurro, M. R., Battistini, S., Salvi, F., Spataro, R., Sola, P., Borghero, G., Galassi, G., Scholz, S. W., Taylor, J. P., Restagno, G., Chiò, A. & Traynor, B. J. (2010) Exome sequencing reveals VCP mutations as a cause of familial ALS. *Neuron* 68, 857-864.
- Kabashi, E., Lin, L., Tradewell, M. L., Dion, P. A., Bercier, V., Bourguoin, P., Rochefort, D., Bel Hadj, S., Durham, H. D., Velde, C. V., Rouleau, G. A. & Drapeau, P. (2010) Gain and loss of function of ALS-related mutations of TARDBP (TDP-43) cause motor deficits in vivo. *Human Molecular Genetics* 19, 671-683.
- Kabashi, E., Valdmanis, P. N., Dion, P., Spiegelman, D., McConkey, B. J., Velde, C. V., Bouchard, J.-P., Lacomblez, L., Pochigaeva, K., Salachas, F., Pradat, P.-F., Camu, W., Meininger, V., Dupre, N. & Rouleau, G. A. (2008) TARDBP mutations in individuals with sporadic and familial amyotrophic lateral sclerosis. *Nature Genetics* 40, 572-574.
- Ling, S.-C., Albuquerque, C. P., Han, J. S., Lagier-Tourenne, C., Tokunaga, S., Zhou, H. & Cleveland, D. W. (2010) ALS-associated mutations in TDP-43 increase its stability and promote TDP-43 complexes with FUS/TLS. *Proceedings of the National Academy of Sciences* 107, 13318-13323.
- Lomas, D. A., Evans, D. L., Finch, J. T. & Carrell, R. W. (1992) The mechanism of Z α 1-antitrypsin accumulation in the liver. *Nature* 357, 605-607.
- Love, S., Bridges, L. R. & Case, C. P. (1995) Neurofibrillary tangles in Niemann-Pick disease type C. *Brain* 118, 119-129.
- Mackenzie, I. R. A., Bigio, E. H., Ince, P. G., Geser, F., Neumann, M., Cairns, N. J., Kwong, L. K., Forman, M. S., Ravits, J., Stewart, H., Eisen, A., McClusky, L., Kretzschmar, H. A., Monoranu, C. M., Highley, J. R., Kirby, J., Siddique, T., Shaw, P. J., Lee, V. M. Y. & Trojanowski, J. Q. (2007) Pathological TDP-43 distinguishes sporadic amyotrophic lateral sclerosis from amyotrophic lateral sclerosis with SOD1 mutations. *Annals of Neurology* 61, 427-434.
- Mackenzie, I. R. A., Rademakers, R. & Neumann, M. (2010) TDP-43 and FUS in amyotrophic lateral sclerosis and frontotemporal dementia. *The Lancet Neurology* 9, 995-1007.
- Maury, C. P. & Baumann, M. (1990) Isolation and characterization of cardiac amyloid in familial amyloid polyneuropathy type IV (Finnish): relation of the amyloid protein to variant gelsolin. *Biochimica et Biophysica Acta* 1096, 84-86.


- Neary, D., Snowden, J. S., Gustafson, L., Passant, U., Stuss, D., Black, S., Freedman, M., Kertesz, A., Robert, P. H., Albert, M., Boone, K., Miller, B. L., Cummings, J. & Benson, D. F. (1998) Frontotemporal lobar degeneration: a consensus on clinical diagnostic criteria. *Neurology* 51, 1546-1554.
- Neumann, M. (2009) Molecular neuropathology of TDP-43 proteinopathies. *International Journal of Molecular Sciences* 10, 232-246.
- Neumann, M., Sampathu, D. M., Kwong, L. K., Truax, A. C., Micsenyi, M. C., Chou, T. T., Bruce, J., Schuck, T., Grossman, M., Clark, C. M., McCluskey, L. F., Miller, B. L., Masliah, E., Mackenzie, I. R., Feldman, H., Feiden, W., Kretzschmar, H. A., Trojanowski, J. Q. & Lee, V. M. Y. (2006) Ubiquitinated TDP-43 in frontotemporal lobar degeneration and amyotrophic lateral sclerosis. *Science* 314, 130-133.
- Nonaka, T., Kametani, F., Arai, T., Akiyama, H. & Hasegawa, M. (2009) Truncation and pathogenic mutations facilitate the formation of intracellular aggregates of TDP-43. *Human Molecular Genetics* 18, 3353-3364.
- Ohmi, K., Kudo, L. C., Ryazantsev, S., Zhao, H.-Z., Karsten, S. L. & Neufeld, E. F. (2009) Sanfilippo syndrome type B, a lysosomal storage disease, is also a tauopathy. *Proceedings of the National Academy of Sciences* 106, 8332-8337.
- Orr, H. T. & Zoghbi, H. Y. (2007) Trinucleotide repeat disorders. *Annual Review of Neuroscience* 30, 575-621.
- Paulson, H. L. (1999) Human genetics '99: trinucleotide repeats. *American Journal of Human Genetics* 64, 339-345.
- Pepys, M. B., Hawkins, P. N., Booth, D. R., Vigushin, D. M., Tennent, G. A., Soutar, A. K., Totty, N., Nguyen, O., Blake, C. C. F., Terry, C. J., Feest, T. G., Zalin, A. M. & Hsuan, J. J. (1993) Human lysozyme gene mutations cause hereditary systemic amyloidosis. *Nature* 362, 553-557.
- Polymeropoulos, M. H., Lavedan, C., Leroy, E., Ide, S. E., Dehejia, A., Dutra, A., Pike, B., Root, H., Rubenstein, J., Boyer, R., Stenroos, E. S., Chandrasekharappa, S., Athanassiadou, A., Papaetropoulos, T., Johnson, W. G., Lazarini, A. M., Duvoisin, R. C., Di Iorio, G., Golbe, L. I. & Nussbaum, R. L. (1997) Mutation in the α -synuclein gene identified in families with Parkinson's disease. *Science* 276, 2045-2047.
- Prusiner, S. B., Scott, M. R., DeArmond, S. J. & Cohen, F. E. (1998) Prion protein biology. *Cell* 93, 337-348.
- Seetharaman, S. V., Prudencio, M., Karch, C., Holloway, S. P., Borchelt, D. R. & Hart, P. J. (2009) Immature copper-zinc superoxide dismutase and familial amyotrophic lateral sclerosis. *Experimental Biology and Medicine* 234, 1140-1154.
- Seilhean, D., Cazeneuve, C., Thuries, V., Russaouen, O., Millecamps, S., Salachas, F., Meininger, V., LeGuern, E. & Duyckaerts, C. (2009) Accumulation of TDP-43 and α -actin in an amyotrophic lateral sclerosis patient with the K171 ANG mutation. *Acta Neuropathologica* 118, 561-573.
- Shibata, N., Hirano, A., Kobayashi, M., Siddique, T., Deng, H. X., Hung, W. Y., Kato, T. & Asayama, K. (1996) Intense superoxide dismutase-1 immunoreactivity in intracytoplasmic hyaline inclusions of familial amyotrophic lateral sclerosis with posterior column involvement. *Journal of Neuropathology and Experimental Neurology* 55, 481-490.
- Sletten, K., Westermark, P. & Natvig, J. B. (1976) Characterization of amyloid fibril proteins from medullary carcinoma of the thyroid. *Journal of Experimental Medicine* 143, 993-998.

- Spillantini, M. G., Crowther, R. A., Jakes, R., Hasegawa, M. & Goedert, M. (1998) α -Synuclein in filamentous inclusions of Lewy bodies from Parkinson's disease and dementia with Lewy bodies. *Proceedings of the National Academy of Sciences, USA* 95, 6469-6473.
- Sreedharan, J., Blair, I. P., Tripathi, V. B., Hu, X., Vance, C., Rogelj, B., Ackerley, S., Durnall, J. C., Williams, K. L., Buratti, E., Baralle, F., de Belleruche, J., Mitchell, J. D., Leigh, P. N., Al-Chalabi, A., Miller, C. C., Nicholson, G. & Shaw, C. E. (2008) TDP-43 mutations in familial and sporadic amyotrophic lateral sclerosis. *Science* 319, 1668-1672.
- Uemichi, T., Liuepnicks, J. j. & Benson, M. D. (1994) Hereditary renal amyloidosis with a novel variant fibrinogen. *Journal of Clinical Investigation* 93, 731-736.
- van Bebber, F., Paquet, D., Hruscha, A., Schmid, B. & Haass, C. (2010) Methylene blue fails to inhibit Tau and polyglutamine protein dependent toxicity in zebrafish. *Neurobiology of Disease* 39, 265-271.
- Vance, C., Rogelj, B., Hortobagyi, T., De Vos, K. J., Nishimura, A. L., Sreedharan, J., Hu, X., Smith, B., Ruddy, D., Wright, P., Ganesalingam, J., Williams, K. L., Tripathi, V., Al-Saraj, S., Al-Chalabi, A., Leigh, P. N., Blair, I. P., Nicholson, G., de Belleruche, J., Gallo, J.-M., Miller, C. C. & Shaw, C. E. (2009) Mutations in FUS, an RNA processing protein, cause familial amyotrophic lateral sclerosis type 6. *Science* 323, 1208-1211.
- Westermarck, P., Engstrom, U., Johnson, K. H., Westermarck, G. T. & Betsholtz, C. (1990) Islet amyloid polypeptide: pinpointing amino acid residues linked to amyloid fibril formation. *Proceedings of the National Academy of Sciences, USA* 87, 5036-5040.
- Westermarck, P., Johnson, K. H. & Pitkanen, P. (1985) Systemic amyloidosis: A review with emphasis on pathogenesis. *Applied Physiology* 3, 55-68.
- Westermarck, P., Johnson, K. H., O'Brien, T. D. & Betsholtz, C. (1992) Islet amyloid polypeptide—a novel controversy in diabetes research. *Diabetologia* 35, 297-303.
- Wijesekera, L. & Leigh, P. N. (2009) Amyotrophic lateral sclerosis. *Orphanet Journal of Rare Diseases* 4, 3.
- Wischik, C. M., Novak, M., Thogersen, H. C., Edwards, P. C., Runswick, M. J., Jakes, R., Walker, J. E., Milstein, C., M., R. & Klug, A. (1988) Isolation of a fragment of tau derived from the core of the paired helical filament of Alzheimer's disease. *Proceedings of the National Academy of Sciences, USA* 85, 4506-4510.
- Yamashita, M., Nonaka, T., Arai, T., Kametani, F., Buchman, V. L., Ninkina, N., Bachurin, S. O., Akiyama, H., Goedert, M. & Hasegawa, M. (2009) Methylene blue and dimebon inhibit aggregation of TDP-43 in cellular models. *FEBS Letters* 583, 2419-2424.

- Zhang, Y.-J., Xu, Y.-F., Cook, C., Gendron, T. F., Roettges, P., Link, C. D., Lin, W.-L., Tong, J., Castanedes-Casey, M., Ash, P., Gass, J., Rangachari, V., Buratti, E., Baralle, F., Golde, T. E., Dickson, D. W. & Petrucelli, L. (2009) Aberrant cleavage of TDP-43 enhances aggregation and cellular toxicity. *Proceedings of the National Academy of Sciences* 106, 7607-7612.

The invention claimed is:

1. A method of treating a tauopathy comprising administering to a subject in need thereof the compound of the following formula:



or a pharmaceutically acceptable salt, solvate, or hydrate thereof.

2. The method of claim 1, wherein the tauopathy is Alzheimer's disease (AD), Pick's disease, progressive supranuclear palsy (PSP), frontotemporal dementia (FTD), FTD with parkinsonism linked to chromosome 17 (FTDP 17), frontotemporal lobar degeneration (FTLD) syndromes; disinhibition-dementia-parkinsonism-amyotrophy complex (DDPAC), pallido-ponto-nigral degeneration (PPND), Guam-ALS syndrome, pallido nigro luisian degeneration (PNLD), cortico-basal degeneration (CBD), dementia with argyrophilic grains (AgD), dementia pugilistica (DP) or chronic traumatic encephalopathy (CTE), Down's syndrome (DS), dementia with Lewy bodies (DLB), subacute sclerosing panencephalitis (SSPE), Niemann-Pick disease, type C (NPC), Sanfilippo syndrome type B, mucopolysaccharidosis III B (MPS III B)), or myotonic dystrophies (DM), DM1 or DM2.

3. The method of claim 2, wherein the tauopathy is Alzheimer's disease.

4. The method of claim 2, wherein the tauopathy is PSP.

5. The method of claim 2, wherein the tauopathy is FTD, FTDP-17, or FTLD.

* * * * *