



US009289400B2

(12) **United States Patent**
Kortagere et al.

(10) **Patent No.:** **US 9,289,400 B2**
(45) **Date of Patent:** **Mar. 22, 2016**

(54) **D₃ DOPAMINE RECEPTOR AGONISTS TO TREAT DYSKINESIA IN PARKINSON'S DISEASE**

215/54 (2013.01); *C07C 217/62* (2013.01);
C07D 209/14 (2013.01); *C07D 235/16*
(2013.01); *C07D 471/04* (2013.01)

(75) Inventors: **Sandhya Kortagere**, Newtown, PA (US); **Eldo V. Kuzhikandahil**, Chester, NJ (US)

(58) **Field of Classification Search**

CPC A61K 31/137
USPC 514/646, 649, 678, 741
See application file for complete search history.

(73) Assignees: **Drexel University**, Philadelphia, PA (US); **Rutgers, The State University of New Jersey**, New Brunswick, NJ (US)

(56) **References Cited**

U.S. PATENT DOCUMENTS

2,407,167 A 9/1946 Kulz
5,352,688 A * 10/1994 Kaminski 514/357
6,031,003 A 2/2000 Nemeth et al.
2008/0058356 A1 3/2008 Crespo Crespo et al.
2009/0281149 A1 11/2009 Scott et al.
2010/0143322 A1 6/2010 Wolf et al.

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **13/764,623**

FOREIGN PATENT DOCUMENTS

DE 102011015842 * 11/2012
EP 1138666 B1 3/2004
WO 9323035 A2 11/1993
WO 99/43678 9/1999
WO 2004113391 A2 12/2004
WO 2005020882 A2 3/2005
WO 2006078239 A1 7/2006
WO 2006083692 A2 8/2006
WO 2006085149 A2 8/2006
WO 2006097744 A2 9/2006
WO 2007/002516 1/2007
WO 2007022415 A2 2/2007

(22) PCT Filed: **Aug. 10, 2011**

(86) PCT No.: **PCT/US2011/047263**

§ 371 (c)(1),
(2), (4) Date: **Feb. 11, 2013**

(87) PCT Pub. No.: **WO2012/021629**

PCT Pub. Date: **Feb. 16, 2012**

(65) **Prior Publication Data**

US 2014/0228355 A1 Aug. 14, 2014

Related U.S. Application Data

(60) Provisional application No. 61/372,733, filed on Aug. 11, 2010.

(51) **Int. Cl.**

A61K 31/137 (2006.01)
A61K 31/4045 (2006.01)
A61K 31/4184 (2006.01)
A61K 31/428 (2006.01)
A61K 31/473 (2006.01)
A61K 31/48 (2006.01)
A61K 31/506 (2006.01)
A61K 31/519 (2006.01)
C07C 211/29 (2006.01)
C07C 215/08 (2006.01)
C07C 215/54 (2006.01)
C07C 217/62 (2006.01)
C07D 209/14 (2006.01)
C07D 235/16 (2006.01)
C07D 471/04 (2006.01)
A61K 45/06 (2006.01)
A61K 31/198 (2006.01)
A61K 31/5513 (2006.01)

(52) **U.S. Cl.**

CPC **A61K 31/137** (2013.01); **A61K 31/198** (2013.01); **A61K 31/4045** (2013.01); **A61K 31/4184** (2013.01); **A61K 31/428** (2013.01); **A61K 31/473** (2013.01); **A61K 31/48** (2013.01); **A61K 31/506** (2013.01); **A61K 31/519** (2013.01); **A61K 31/5513** (2013.01); **A61K 45/06** (2013.01); **C07C 211/29** (2013.01); **C07C 215/08** (2013.01); **C07C**

(Continued)

OTHER PUBLICATIONS

PubChemSubstance Direct Submission SID 84973441, 2-[(2-chlorobenzyl)amino]butan-1-ol-1-ol-6 Substance Summary (Jul. 22, 2009), [Retrieved from the Internet Mar. 28, 2013: <<http://pubchem.ncbi.nlm.nih.gov/summary/summary.cgi?sid=84973441>>.

(Continued)

Primary Examiner — Sreeni Padmanabhan

Assistant Examiner — Sahar Javanmard

(74) *Attorney, Agent, or Firm* — Saul Ewing LLP; Kathryn Doyle; Domingos J. Silva

(57) **ABSTRACT**

The present invention provides a method of inhibiting, suppressing or preventing levodopa-induced dyskinesia in a patient suffering from Parkinson's Disease, comprising the step of administering to the patient a pharmaceutical composition comprising at least one compound of the invention. The present invention further provides a method of inhibiting, suppressing or preventing Parkinson's Disease in a patient, comprising the step of administering to the patient a pharmaceutical composition comprising at least one compound of the invention.

13 Claims, 35 Drawing Sheets

(56)

References Cited

FOREIGN PATENT DOCUMENTS

WO	2007133772	A2	11/2007
WO	2008052953	A1	5/2008
WO	2008107335	A1	9/2008
WO	2008113364	A2	9/2008
WO	2009136175	A1	11/2009
WO	2009140204	A2	11/2009

OTHER PUBLICATIONS

Clarke CE, Deane K. Cabergoline versus bromocriptine for levodopa-induced complications in Parkinson's disease. Cochrane Database of Systematic Reviews 2001, Issue 1. DOI: 10.1002/14651858.CD001519.

Klein, et al. "Synthesis of chiral 1,4-disubstituted-1,2,3-triazole derivatives from amino acids." 2009, *Molecules* 14:5124-5143.

Newman, et al., "N-(4-(4-(2,3-dichloro- or 2-methoxyphenyl)piperazin-1-yl)butyl)heterobiarylcarboxamides with functionalized linking chains as high affinity and enantioselective D3 receptor antagonists." 2009, *J Med Chem* 52(8):2559-2570.

Grundt, et al., "Heterocyclic analogues of N-(4-(4-(2,3-dichlorophenyl)piperazin-1-yl)butyl)arylcarboxamides with

functionalized linking chains as novel dopamine D3 receptor ligands: potential substance abuse therapeutic agents." 2007, *J Med Chem* 50:4135-4146.

Boeckler et al., "CoMFA and CoMSIA investigations revealing novel insights into the binding modes of dopamine D3 receptor agonists." 2005, *J Med Chem* 48:2493-2508.

Westrich et al., "Development of tolerance in D3 dopamine receptor signaling is accompanied by distinct changes in receptor conformation." 2010, *Biochemical Pharmacology* 79:897-907.

Ask, et al., "Selective Inhibition of Monoamine Oxidase by p-Aminostyryl Phenylalkylamines in Catecholaminergic Neurons", *Neuropharmacology*, vol. 25, No. 1, pp. 33-40, 1986.

Patent Examination Report No. 1, dated Feb. 5, 2014, Australian Government.

Costall, et al., "Dyskinetic phenomena caused by the intrastriatal injection of phenylethylamine, phenylpiperazine, tetrahydroisoquinoline and tetrahydronaphthalene derivatives in the guinea pig", *Eur J Pharmacol.* 31(1), Mar. 1975, 94-109.

References cited in U.S. Appl. No. 13/957,861 (which is a divisional of U.S. Appl. No. 13/764,623) and various non-US counterpart applications.

* cited by examiner

Fig. 1A




Fig. 1B




Fig. 1C




Fig. 1D




Fig. 2A




Fig. 2B




Fig. 2C




Fig. 3




Fig. 4

DRD3 ----MASLSQLSSHLNYTCGAENSTG----ASQARPHAYYA---LSYCALILAIIVFGNGL
 2RH1 DYKDDDAMGQPGNGSAFLLAPNRSHPDHDVTQQRDEVWVVGMGIVMSLIVLAIIVFGNVL
 : : * _ : : * _ : * * _ : _ : _ : : * * * * * *

DRD3 VCMAVLKERALQTTTNYLVVSLAVADLLVATLVMPWVVYLEVTGGVWNFSRICCDVFVTL
 2RH1 VITAIKFERLQTVTNYFITSLACADLMGLAVVPFGAAHILMK-MWTFGNFWCEFWTSI
 * * : * _ * * * * * : : * * * * * : _ * : * : _ : : * _ _ : * : _ : : :

DRD3 **D**VMMCTASILNLCAISIDRYTAVVMPVHYQHGTGQSSCRRVALMITAVWVLAFAVSCPLL
 2RH1 DVLCVTASIE TLCVIAVDRYFAITSPFKYQSLLTKNKARVILMVWIVSGLTSEFLPIQMH
 * * : * * * * * _ * * * * * : : * * * * * : _ * : * * : * * : : _ :

DRD3 FG-----FNTTGDPTVCS-ISNPDFVIYSSVVSEYLPFGVTVLVYARIYVVLK-----
 2RH1 WYRATHQEAINCYAEETCCDFFTNQAYAIASSIVSFYVPLVIMVFVYSRVFQEAKRQLNI
 : : * . : * * _ : : * : _ * * * * * : * : * * * * * : * :

DRD3 -----QRRRKRI LTRQNS
 2RH1 FEMLRIDEGLRLKIYKDTEGYTIGIGHLLTKSPSLNAAKSELDKAIGRNTNGVITKDEA
 * _ : : * * * * * :


DRD3 Q-----CNSVRPGFQPQTLSPDPAHLELKR----YYSICQDTALGGPGFQERGGELKR
 2RH1 EKLFNQDVDAAVRGILRNAKLKPVYDSLDAVRRALINMVFQMGETGVAGFTNSLRMLQ-
 : : * * : : _ * _ * : * : * * _ * * : * :

DRD3 EEKTRNSLSPTIAPKLSLEVRKLSNGRLSTSLKLGPLQPRGVPLREKKATQMVAVLGF
 2RH1 -QKRWDEAAVNLAKSRYNQTPNRAKRVITTFRTGTWDAYKFCLKEHKALKTLGIIMGTF
 : * : _ : _ : * _ : : * : * : : * _ : _ _ * : * * * * : : _ * : * * * *

DRD3 IVCWLP**FF**LTHVLNTHCQTCHVSPELYSATTWLGYUNSA LNPVIYTTFNIEFRKAF LKIL
 2RH1 TLCWL PFFIVNIVH-VIQDNLIRKEVYILLNWIGYVNSGFNPLIYCRS-PDFRIAFQELL
 : * * * * * : _ : : : * : * * * _ * : * * * * * : * * * * * : * * * * * :

DRD3 SC-----
 2RH1 CLRSSLKAYGNGYSSNGNTGEQSG

Fig. 5



color key

- | | |
|--------------|----------------------------------|
| ○ - nonpolar | ● - ligand interaction site |
| ○ - polar | - main chain H-bond |
| ○ - acidic | - side chain H-bond |
| ○ - basic | → - direction of proton donation |

Fig. 6A




Fig. 6B




Fig. 6C




Fig. 6D




Fig. 7A




Fig. 7B




Fig. 8




Fig. 9A




Fig. 9B




Fig. 9C




Fig. 10A & Fig. 10B






Fig. 11A & Fig. 11B

(A)



(B)




Fig. 12B




Fig. 12A




Fig. 13A




Fig. 13B




Fig. 14




Fig. 15




Fig. 16A




Fig. 16B




Fig. 17A




Fig. 17B




Fig. 17C




Fig. 17D




Fig. 18




Fig. 19A

Comparing Effect of Typical vs Atypical D3R agonists on LID




Fig. 19B




Fig. 20




Fig. 21A




Fig. 21B




Fig. 22A







Fig. 22B





	hSERT	hNET	hDAT
LogIC50	1.138	-0.06543	0.2069
IC50	13.75	0.8601	1.610

Fig. 23B



	hSERT	hNET	hDAT
LogIC50	1.555	-0.5326	0.9306
IC50	35.85	0.2933	8.523

**D₃ DOPAMINE RECEPTOR AGONISTS TO
TREAT DYSKINESIA IN PARKINSON'S
DISEASE**

CROSS-REFERENCE TO RELATED
APPLICATIONS

This application is a continuation of International Application No. PCT/US2011/047263, filed Aug. 10, 2011, which claims the benefit of U.S. Provisional Application No. 61/372,733, filed Aug. 11, 2010, all of which applications are hereby incorporated by reference in their entireties herein.

BACKGROUND OF THE INVENTION

The family of G-protein coupled receptors (GPCRs) is one of the most important classes of proteins from both functional and structural standpoints. The human genome contains nearly 950 genes coding for GPCRs, of which nearly 450 genes have been implicated as therapeutic targets. Ligand binding to GPCRs induces multiple receptor conformations and different ligands may stabilize different receptor conformations (Kenakin & Miller, 2010, *Pharmacol. Rev.* 62(2): 265-304). The concept of functional selectivity is based on the hypothesis that different receptor conformations recruit different signaling proteins which leads to preferential activation of one signaling pathway over another (Mailman, 2007, *Trends Pharmacol. Sci.* 28(8):390-396). In addition to selecting the signaling pathways, agonist-induced receptor conformations can also potentially affect receptor signaling properties.

Among the GPCRs, the subfamily of dopamine receptors has attracted attention from biologists and pharmacologists. In the central nervous system, dopamine receptors are widely expressed and involved in the control of locomotion, cognition, emotion and neuroendocrine secretion. In the peripheral system, dopamine receptors are present more prominently in kidney, vasculature and pituitary, where they affect mainly sodium homeostasis, vascular tone, and hormone secretion. While there are numerous examples of functionally-selective ligands, which preferentially activates one signaling cascade but not others, functionally-selective ligands that alter receptor signaling properties are rare and have not been described for dopamine receptors.

The neurotransmitter dopamine controls a wide variety of physiological and behavioral functions in mammals via five major subtypes of dopamine receptors. They are broadly classified into the "D₁ like" and "D₂ like" dopamine receptors based on pharmacology and function. The D₁-like consists of D₁ and D₅ receptors, while the D₂-like consists of D₂, D₃, and D₄ receptors. The D₃ receptor primarily couples to the pertussis toxin-sensitive G α -proteins (Gi/Go) (Ahlgren-Beckendorf & Levant, 2004, *J. Recept. Signal Transduct. Res.* 24(3):117-130). When transfected into different cell lines, the D₃ receptor couples to adenylyl cyclase V isoform (Robinson & Caron, 1997, *Mol. Pharmacol.* 52:508-514) and initiates signaling events including phosphorylation of mitogen-activated protein (MAP) kinases (Cussac et al., 1999, *Mol. Pharmacol.* 56(5):1025-103). D₂ and D₃ dopamine receptors also modulate potassium and calcium channel function (Seabrook et al., 1994, *Br. J. Pharmacol.* 111:391-393; Werner et al., 1996, *Mol. Pharmacol.* 49:656-661). Transfected D₃ receptors couple robustly to natively expressed G-protein coupled inward rectifier potassium (GIRK) and voltage-gated P/Q type calcium channels, and inhibit firing of spontaneous action potentials and secretory activity in the AtT-20 neuroendocrine cell line (Kuzhikandathil & Oxford,

1999, *J. Neurosci.* 19(5):1698-1707; Kuzhikandathil & Oxford, 2000, *J. Gen. Physiol.* 115:697-706; Kuzhikandathil et al., 1998, *Mol. Cell. Neurosci.* 12:390-402). The D₃ receptor further couples to natively expressed adenylyl cyclase V (Kuzhikandathil & Bartoszyk, 2006, *Neuropharm.* 51:873-884), MAP kinases (Westrich & Kuzhikandathil, 2007, *Biochim. Biophys. Acta-MCR* 1773:1747-1758) and ion channels (Kuzhikandathil & Oxford, 1999, *J. Neurosci.* 19(5): 1698-1707; Kuzhikandathil & Oxford, 2000, *J. Gen. Physiol.* 115:697-706; Kuzhikandathil et al., 1998, *Mol. Cell. Neurosci.* 12:390-402; Kuzhikandathil et al., 2004, *Mol. Cell. Neurosci.* 26:144-155) in AtT-20 cells.

The expression of D₃ dopamine receptor is altered under many pathological conditions and following chronic treatment. In Parkinson's disease, levodopa-induced dyskinesias are associated with a specific up regulation of D_{3R} expression in putamen and globus pallidus internal segment, regions that normally express the D₂ receptor (Bezard et al., 2003, *Nat. Med.* 9(6):762-767; Guigoni et al., 2005, *Parkinsonism Related Disorders* 11 Suppl 1, S25-29). In rodent models, the behavioral sensitization associated with levodopa treatment is mediated by upregulated D₃ receptors (Guillin et al., 2001, *Nature* 411(6833):86-89). In schizophrenia, D₃ receptor expression levels are increased two fold in the basal ganglia. Antipsychotic treatment has also been reported to change the expression of D₃ receptor. The density of D₃ receptor is increased in chronic cocaine users in striatum and substantia nigra, as well as in the nucleus accumbens. Stress and depression-induced down regulation of D₃ receptor expression is reversed following chronic antidepressant treatment.

Dopamine receptors are targets for the treatment of various neurological and psychiatric disorders, such as Parkinson's disease, schizophrenia, drug addiction, depression, bipolar disorder, attention deficit hyperactivity syndrome, Tourette's syndrome, Huntington's disease and migraine.

Parkinson's disease (also known as Parkinson disease) is a degenerative disorder of the central nervous system that often impairs the sufferer's motor skills, speech, and other functions (Jankovic, 2008, *J. Neurol. Neurosurg. Psychiatr.* 79(4): 368-76). Parkinson's Disease is characterized by muscle rigidity, tremor, a slowing of physical movement (bradykinesia) and a loss of physical movement (akinesia) in extreme cases. The primary symptoms of Parkinson's Disease are the results of decreased stimulation of the motor cortex by the basal ganglia, normally caused by insufficient formation and action of dopamine, which is produced in the dopaminergic neurons of the brain (specifically the substantia nigra). Secondary symptoms may include high level cognitive dysfunction and subtle language problems. Parkinson's Disease is both chronic and progressive. At present, there is no cure for Parkinson's Disease, but medications may provide relief from the symptoms.

The most widely used form of treatment is L-dopa (levodopa). Levodopa is transformed into dopamine by L-aromatic amino acid decarboxylase (also known as dopa-decarboxylase) in the dopaminergic neurons. However, only 1-5% of levodopa enters the dopaminergic neurons. The remaining levodopa is often metabolized to dopamine elsewhere, causing a wide variety of side effects ("Symptomatic pharmacological therapy in Parkinson's disease". Parkinson's Disease. London: Royal College of Physicians, 2006, pp. 59-100). Due to feedback inhibition, levodopa administration results in a reduction of the endogenous formation of levodopa, and so eventually becomes counterproductive. Levodopa may also be co-administered with carbidopa ((2S)-3-(3,4-dihydroxyphenyl)-2-hydrazino-2-methylpropanoic acid), which prevents levodopa metabolism elsewhere in the

3

body. The dopamine agonists bromocriptine, pergolide, pramipexole, ropinirole, piribedil, cabergoline, apomorphine, and lisuride have been found to be moderately effective.

Levodopa-induced dyskinesia (LID) is a particularly serious side effect of the long-term use of levodopa. These motor fluctuations occur in more than half of Parkinson's Disease patients after 5-10 years of levodopa treatment, with the percentage of affected patients increasing over time, and LID is thought to be potentially irreversible. Dyskinesia most commonly occurs at the time of peak levodopa plasma concentrations and is thus referred to as peak-dose dyskinesia. As patients advance, they may evidence diphasic dyskinesia, which occurs when the drug concentration rises or falls. Attempts to moderate dyskinesia by the use of other treatments, such as bromocriptine (Parlodel™), appear to be ineffective. In order to avoid dyskinesia, patients with the young-onset form of the disease are often hesitant to commence levodopa therapy until absolutely necessary for fear of suffering severe dyskinesia later on. Currently, there is no pharmacotherapeutic means of treating LID in patients suffering from Parkinson's Disease.


Interestingly, there is an alteration of dopamine receptor expression in most disorders associated with the dopaminergic system, such as Parkinson's Disease. Changes in dopamine receptor expression are also observed following chronic treatment of these neurological and psychiatric disorders. In the case of D₃ dopamine receptor, changes in expression have been reported in Parkinson's disease, schizophrenia, depression, and drug addiction. Following chronic drug treatment, studies have reported an upregulation of D₃ receptor in LID in Parkinson's disease and antipsychotic-induced tardive dyskinesia in schizophrenia.

The ability of using levodopa as a therapeutic agent in the treatment of Parkinson's Disease is severely hampered by the likelihood that levodopa-induced dyskinesia (LID) will eventually develop. There is a need in the art for novel therapeutic agents that treat, ameliorate or prevent levodopa-induced dyskinesia in patients suffering from Parkinson's Disease. The present invention fulfills this need.

BRIEF SUMMARY OF THE INVENTION

The invention includes a pharmaceutical composition comprising at least one pharmaceutically acceptable carrier and at least one compound selected from the group consisting of:

a compound of formula (I):



wherein in formula (I):


R¹, R² and R³ are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl-(C₁₋₃)alkyl, substituted aryl-(C₁₋₃)alkyl, carboxy, alkylcarboxy, formyl, alkyl-carbonyl, aryl-carbonyl, and heteroaryl-carbonyl;

4


R⁴ and R⁵ are independently selected from the group consisting of H, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl-(C₁₋₃)alkyl, and substituted aryl-(C₁₋₃)alkyl; and,

n is 2, 3, 4 or 5;

a compound of formula (IIa) or (IIb):



(IIa)



(IIb)

wherein in formula (IIa) or (IIb):

R¹ and R² are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl-(C₁₋₃)alkyl, substituted aryl-(C₁₋₃)alkyl, carboxy, alkylcarboxy, formyl, alkyl-carbonyl, aryl-carbonyl, and heteroaryl-carbonyl;

R³ and R⁴ are independently selected from the group consisting of H, C₁₋₆ alkyl, aryl, heteroaryl, and substituted C₁₋₆ alkyl;

R⁵ is selected from the group consisting of H, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl-(C₁₋₃)alkyl, and substituted aryl-(C₁₋₃)alkyl; and,

m is 1, 2, 3 or 4;

2,7-diamino-5-(4-fluorophenyl)-4-oxo-3,4,5,6-tetrahydropyrido[2,3-d]pyrimidine-6-carbonitrile;

(Z)-2-(1H-benzof[d]imidazol-2-yl)-N'-hydroxy-3-(4-methoxyphenyl)propanimidamide;

mixtures thereof, or a pharmaceutically acceptable salt thereof.


In one embodiment, in formula (I) R¹, R² and R³ are independently selected from the group consisting of H, cyano, halo, alkoxy, nitro, C₁₋₆ alkyl, and carboxy. In another embodiment, in formula (I) R⁴ and R⁵ are independently selected from the group consisting of H, C₁₋₆ alkyl, and substituted C₁₋₆ alkyl. In yet another embodiment, in formula (I) n is 2. In yet another embodiment, in formula (IIa) or (IIb) m is 1 or 2.

In one embodiment, the at least one compound is selected from the group consisting of: 2-amino-4-(2-chlorophenyl)butan-1-ol; 2-(3-aminoethyl)phenol; 4-(2-chlorophenyl)-2-methylamino-butane (also known as 4-(2-chlorophenyl)-N-methylbutan-2-amine); 4-(2-chlorophenyl)-butan-2-amine; 4-(2-fluorophenyl)butan-2-amine; 4-(2-bromophenyl)butan-2-amine; 4-(2-iodophenyl)butan-2-amine; 4-(2-methoxyphenyl)butan-2-amine; 2-(3-aminobutyl)phenol; 3-(3,4-diethoxyphenyl)propan-1-amine; 4-(4-chlorophenyl)butan-2-

5

amine; 4-(4-methoxyphenyl)butan-2-amine; 2-(5-chloro-1-methyl-1H-indol-3-yl)ethanamine; 1-(5-fluoro-1-methyl-1H-indol-3-yl)propan-2-amine; 1-(5-methoxy-1-methyl-1H-indol-3-yl)propan-2-amine; 2,7-diamino-5-(4-fluorophenyl)-4-oxo-3,4,5,6-tetrahydropyrido[2,3-d]pyrimidine-6-carbonitrile; (Z)-2-(1H-benzo[d]imidazol-2-yl)-N'-hydroxy-3-(4-methoxyphenyl)propanimidamide; mixtures thereof, or a pharmaceutically acceptable salt thereof.

The invention also includes a method of treating, ameliorating or preventing levodopa-induced dyskinesia in a patient suffering from Parkinson's Disease. The method comprises administering to the patient a pharmaceutical composition comprising a therapeutically effective amount of at least one compound selected from the group consisting of: a compound of formula (I):





wherein in formula (I):

R¹, R² and R³ are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl-(C₁₋₃)alkyl, substituted aryl-(C₁₋₃)alkyl, carboxy, alkylcarboxy, formyl, alkyl-carbonyl, aryl-carbonyl, and heteroaryl-carbonyl;

R⁴ and R⁵ are independently selected from the group consisting of H, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl-(C₁₋₃)alkyl, and substituted aryl-(C₁₋₃)alkyl; and,

n is 2, 3, 4 or 5;

a compound of formula (IIa) or (IIb):



wherein in formula (IIa) or (IIb):

R¹ and R² are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl-(C₁₋₃)alkyl, substituted aryl-(C₁₋₃)alkyl,

6

alkyl, carboxy, alkylcarboxy, formyl, alkyl-carbonyl, aryl-carbonyl, and heteroaryl-carbonyl;

R³ and R⁴ are independently selected from the group consisting of H, C₁₋₆ alkyl, aryl, heteroaryl, and substituted C₁₋₆ alkyl;

R⁵ is selected from the group consisting of H, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl-(C₁₋₃)alkyl, and substituted aryl-(C₁₋₃)alkyl; and,

m is 1, 2, 3 or 4;

2,7-diamino-5-(4-fluorophenyl)-4-oxo-3,4,5,6-tetrahydropyrido[2,3-d]pyrimidine-6-carbonitrile;

(Z)-2-(1H-benzo[d]imidazol-2-yl)-N'-hydroxy-3-(4-methoxyphenyl)propanimidamide;

mixtures thereof, or a pharmaceutically acceptable salt thereof.

(I)

In one embodiment, in formula (I) R¹, R² and R³ are independently selected from the group consisting of H, cyano, halo, alkoxy, nitro, C₁₋₆ alkyl, and carboxy. In another embodiment, in formula (I) R⁴ and R⁵ are independently selected from the group consisting of H, C₁₋₆ alkyl, and substituted C₁₋₆ alkyl. In yet another embodiment, in formula (I) n is 2. In yet another embodiment, in formula (IIa) or (IIb) m is 1 or 2.

In one embodiment, the at least one compound is selected from the group consisting of: 2-amino-4-(2-chlorophenyl)butan-1-ol; 2-(3-aminohexyl)phenol; 4-(2-chlorophenyl)butan-2-amine; 4-(2-chlorophenyl)-2-methylamino-butane; 4-(2-fluorophenyl)butan-2-amine; 4-(2-bromophenyl)butan-2-amine; 4-(2-iodophenyl)butan-2-amine; 4-(2-methoxyphenyl)butan-2-amine; 2-(3-aminobutyl)phenol; 3-(3,4-diethoxyphenyl)propan-1-amine; 4-(4-chlorophenyl)butan-2-amine; 4-(4-methoxyphenyl)butan-2-amine; 2-(5-chloro-1-methyl-1H-indol-3-yl)ethanamine; 1-(5-fluoro-1-methyl-1H-indol-3-yl)propan-2-amine; 1-(5-methoxy-1-methyl-1H-indol-3-yl)propan-2-amine; 2,7-diamino-5-(4-fluorophenyl)-4-oxo-3,4,5,6-tetrahydropyrido[2,3-d]pyrimidine-6-carbonitrile; (Z)-2-(1H-benzo[d]imidazol-2-yl)-N'-hydroxy-3-(4-methoxyphenyl)propanimidamide; mixtures thereof, or a pharmaceutically acceptable salt thereof.

(IIa)

In one embodiment, the composition further comprises a drug selected from the group consisting of levodopa, clozapine, bromocriptine, pergolide, pramipexole, ropinirole, piribedil, cabergoline, apomorphine, and lisuride, a salt thereof and mixtures thereof.

(IIb)

In one embodiment, the pharmaceutical composition is co-administered to the patient with a second pharmaceutical composition comprising levodopa.

In one embodiment, the pharmaceutical composition is administered to the patient a given period of time before a second pharmaceutical composition comprising levodopa is administered to the patient.


In one embodiment, the given period of time varies from about 2 minutes to about 24 hours.

In one embodiment, the patient is human.

The invention further includes a method of treating, ameliorating or preventing Parkinson's Disease in a patient. The method comprises administering to the patient a pharmaceutical composition comprising a therapeutically effective amount of at least one compound selected from the group consisting of:

7

a compound of formula (I):





wherein in formula (I):

R^1 , R^2 and R^3 are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C_{1-6} alkyl, substituted C_{1-6} alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl- (C_{1-3}) alkyl, substituted aryl- (C_{1-3}) alkyl, carboxy, alkylcarboxy, formyl, alkyl-carbonyl, aryl-carbonyl, and heteroaryl-carbonyl;

R^4 and R^5 are independently selected from the group consisting of H, C_{1-6} alkyl, substituted C_{1-6} alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl- (C_{1-3}) alkyl, and substituted aryl- (C_{1-3}) alkyl; and,

n is 2, 3, 4 or 5;

a compound of formula (IIa) or (IIb):



wherein in formula (IIa) or (IIb):

R^1 and R^2 are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C_{1-6} alkyl, substituted C_{1-6} alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl- (C_{1-3}) alkyl, substituted aryl- (C_{1-3}) alkyl, carboxy, alkylcarboxy, formyl, alkyl-carbonyl, aryl-carbonyl, and heteroaryl-carbonyl;

R^3 and R^4 are independently selected from the group consisting of H, C_{1-6} alkyl, aryl, heteroaryl, and substituted C_{1-6} alkyl;

R^5 is selected from the group consisting of H, C_{1-6} alkyl, substituted C_{1-6} alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl- (C_{1-3}) alkyl, and substituted aryl- (C_{1-3}) alkyl; and,

m is 1, 2, 3 or 4;

2,7-diamino-5-(4-fluorophenyl)-4-oxo-3,4,5,6-tetrahydropyrido[2,3-d]pyrimidine-6-carbonitrile;

(Z)-2-(1H-benzo[d]imidazol-2-yl)-N'-hydroxy-3-(4-methoxyphenyl)propanimidamide;

mixtures thereof, or a pharmaceutically acceptable salt thereof.

8

In one embodiment, the at least one compound is selected from the group consisting of: 2-amino-4-(2-chlorophenyl)butan-1-ol; 2-(3-aminohexyl)phenol; 4-(2-chlorophenyl)-2-methylamino-butane; 4-(2-chlorophenyl)-butan-2-amine; 4-(2-fluorophenyl)butan-2-amine; 4-(2-bromophenyl)butan-2-amine; 4-(2-iodophenyl)butan-2-amine; 4-(2-methoxyphenyl)butan-2-amine; 2-(3-aminobutyl)phenol; 3-(3,4-diethoxyphenyl)propan-1-amine; 4-(4-chlorophenyl)butan-2-amine; 4-(4-methoxyphenyl)butan-2-amine; 2-(5-chloro-1-methyl-1H-indol-3-yl)ethanamine; 1-(5-fluoro-1-methyl-1H-indol-3-yl)propan-2-amine; 1-(5-methoxy-1-methyl-1H-indol-3-yl)propan-2-amine; 2,7-diamino-5-(4-fluorophenyl)-4-oxo-3,4,5,6-tetrahydropyrido[2,3-d]pyrimidine-6-carbonitrile; (Z)-2-(1H-benzo[d]imidazol-2-yl)-N'-hydroxy-3-(4-methoxyphenyl)propanimidamide; mixtures thereof, or a pharmaceutically acceptable salt thereof.

In one embodiment, the composition further comprises at least one drug selected from the group consisting of levodopa, clozapine, bromocriptine, pergolide, pramipexole, ropinirole, piribedil, cabergoline, apomorphine, and lisuride, a salt thereof and mixtures thereof.

In one embodiment, the patient is further administered a second pharmaceutical composition comprising at least one drug selected from the group consisting of levodopa, clozapine, bromocriptine, pergolide, pramipexole, ropinirole, piribedil, cabergoline, apomorphine, and lisuride, a salt thereof and mixtures thereof.

BRIEF DESCRIPTION OF THE DRAWINGS

For the purpose of illustrating the invention, there are depicted in the drawings certain embodiments of the invention. However, the invention is not limited to the precise arrangements and instrumentalities of the embodiments depicted in the drawings.

FIG. 1, comprising FIGS. 1A-D, illustrates representative voltage (FIGS. 1A and 1C) and current (FIGS. 1B and 1D) clamp recording in AtT-20 cells stably expressing either human $D_{2.5}$ (FIGS. 1A and 1B) or D_3 (FIGS. 1C and 1D). For voltage clamp recording (FIGS. 1A and 1C), the cells were held at -65 mV and 100 nM dopamine (black bar) applied for 1 minute. For current clamp recording (FIGS. 1B and 1D), the spontaneous action potentials were hyperpolarized by one minute application of 100 nM quinpirole (QP), which is an agonist for D_2 -like receptors.

FIG. 2, comprising FIGS. 2A-C, illustrates agonist-induced modulation of D_3 receptor tolerance and SRT properties. Representative voltage clamp (FIG. 2A) and current clamp recording (FIG. 2B) from AtT- D_3 cells treated with 300 nM PBZI ((3aS,9bR)-3-propyl-1,2,3a,4,5,9b-hexahydrobenzo[e]indol-8-ol) or 100 nM quinpirole ((4aR,8aR)-5-propyl-4,4-a,5,6,7,8,8a,9-octahydro-1H-pyrazolo[3,4-g]quinoline) are illustrated. PBZI elicited 3 responses that did not show either tolerance or SRT properties; on the same cell, quinpirole elicited GIRK response with tolerance and SRT. Similarly, representative voltage clamp recordings illustrated in FIG. 2C show that 300 nM FAUC73 (square crossed bar) induces GIRK currents that do not show D_3 receptor tolerance and SRT. In contrast, in the same cell, 300 nM PD128907 (black bar) elicited tolerance and SRT. The cells were held at -65 mV and the duration of agonist application was ~ 60 seconds.

FIG. 3 is a flowchart illustration of the hybrid structure based (HSB) protocol that was used to design these novel D_3 agonists.

FIG. 4 illustrates the primary amino acid sequence alignment of D₃ receptor (SEQ ID NO:1) with β 2AR (pdb code 2RH1—coded as BADRE-2RH1; SEQ ID NO:2). Residues that form the TM helices are shown underlined. Identical residues are indicated by “*” and similarity is indicated by “:” and “.”.

FIG. 5 is a schematic representation of PBZI binding to the D₃ receptor with the pharmacophore elements superimposed. Unfilled ovals labeled “gr” represent hydrophobic elements, unfilled ovals labeled “pur” represent hydrogen bond donor-acceptor pairs, and distance constraints are represented by dotted black lines. Hydrogen bonded interactions are shown by arrows, ionic interactions in lines and pi-pi interactions in lines extending across the two six membered rings. Gray spheres and contours indicate matching regions between ligand and receptors.

FIG. 6, comprising FIGS. 6A-D, is a series of schemes illustrating the mode of interaction of ES609 (FIG. 6A), PD128907 (FIG. 6B), PBZI (FIG. 6C), and Dopamine (FIG. 6D). The binding site residues are colored by their nature, with hydrophobic residues in green (labeled gr), polar residues in purple (labeled pur), and charged residues highlighted with bold contours. Gray spheres and contours indicate matching regions between ligand and receptors. Hydrogen bonded interactions are shown by arrows, ionic interactions in lines and pi-pi interactions in lines extending across the two six membered rings. The figures were generated using the LIGX module of MOE program. The 3D pharmacophore used to screen the molecules is overlaid on the PBZI structure (FIG. 6C) with open red circles representing hydrophilic interactions (labeled red), blue open circles representing hydrophobic and aromatic interactions (labeled bl) and black dotted lines representing distance between the pharmacophore elements.

FIG. 7, comprising FIGS. 7A-B, is a set of molecular models illustrating that the HSB method allows for the identification of residues and conformations involved in D₃ receptor tolerance and SRT properties. FIG. 7A illustrates the structural super positioning of D₃ receptor bound to PD128907, PBZI and ES609, with the receptor represented in cartoon format. The transmembrane helices are numbered 1 through 7 and the extra and intracellular loops are labeled EC1-EC3 and IC1-IC3 respectively. FIG. 7B illustrates a molecular model of dopamine D₃ receptor with 4-(2-chlorophenyl)-butan-2-amine (ES609) docked. The receptor is depicted as ribbons, and the ligand is rendered in a space filled model.

FIG. 8 illustrates a flowchart for the experimental development plan.

FIG. 9, comprising FIGS. 9A-C, illustrates voltage clamp experiments. FIG. 9A illustrates a representative voltage clamp recording of D₃ receptor-induced GIRK response upon two sequential 100 nM dopamine (DA) treatment. The DA was dissolved in extracellular solution with 30 mM potassium (to enhance GIRK currents). Duration of agonist application is ~60 seconds. The ratio of second to first GIRK response is a quantitative measure of tolerance. FIG. 9B is a graph bar that indicates that D₃ receptor tolerance is elicited by 100 nM dopamine (DA), quinpirole (QP), and PD 128907; but not by PBZI or 4-(2-chlorophenyl)-butan-2-amine (ES609). FIG. 9C is a graph bar illustrating that tolerance property of D₃ receptor is agonist-dependent. Cumulative data showed the ratio of 2nd to 1st agonist-induced GIRK response in AtT-20 cells stably expressing the human D₃ dopamine receptor. GIRK responses were elicited using saturating concentration of dopamine (DA, 100 nM, N=5), quinpirole (100 nM, N=5), PD128907 (100 nM, N=5), 7OH-DPAT (100 nM, N=4), 7OH-

PIPAT (100 nM, N=4), sarizotan (100 nM, N=6), pramipexole (300 nM, N=4), rotigotine (100 nM, N=4), PBZI (300 nM, N=10), and FAUC 73 (300 nM, N=4). Error bars represent \pm SEM. *, **, P<0.05, ANOVA, post-hoc Holm-Sidak test.

FIG. 10, comprising FIGS. 10A-B, illustrates representative voltage clamp recording showing that PBZI (PEUN1) may compete with dopamine (DA) and prevent the development of tolerance and SRT at D₃ receptors expressed stably in AtT-20 cells. In FIG. 10A, 100 nM DA elicited D₃ receptor tolerance and SRT. In FIG. 10B, 1000 nM PBZI applied simultaneously with DA blocked the development of tolerance and SRT.

FIG. 11, comprising FIGS. 11A-B, illustrates dose response curves for PBZI (FIG. 11A) and 4-(2-chlorophenyl)-butan-2-amine (ES609; FIG. 11B) in two different functional assays. As illustrated in FIG. 11A, GIRK current response to PBZI in AtT-20 cells expressing D₃ receptors was measured using whole cell voltage clamp recording. The current response was normalized for cell size (using membrane capacitance). FIG. 11B illustrates inhibition of 10 μ M forskolin induced cAMP levels by activation of D₃ receptors in AtT-20 cells with various doses of 4-(2-chlorophenyl)-butan-2-amine. The cAMP levels were measured by using an ELISA kit from GE Healthcare. The triangle in FIG. 11B illustrates the inhibition elicited by 300 nM quinpirole.

FIG. 12, comprising FIGS. 12A-B, illustrates the effects of PBZI and PD128907 on locomotion in Balb/c mice. As illustrated in FIG. 12A, PBZI induced a dose-dependent and monophasic inhibition of locomotion. The horizontal arrows indicate the hypoactivity induced by 10 mg/kg, sc PBZI (circles) plotted in 10 minute bins. The 1 mg/kg dose (triangles) had no significant effect. The locomotor activity induced by PBZI was normalized to locomotor activity in control vehicle (saline) injected mice. The data points indicated by the horizontal arrows shows statistically significant reductions in locomotor activity (p=0.002, Newman-Keuls multiple (α =0.05), n=4 mice per treatment group) that recovers to control levels by 80 minutes. As illustrated in FIG. 12B, in contrast to PBZI, Balb/c mice administered 0.4 mg/kg, sc PD128907 (circles, n=4) induces an initial hypoactivity (horizontal arrows) that is followed by hyperactivity (area indicated by hatched rectangle), which returns to levels exhibited by saline injected mice (n=4) by 120 minutes. A lower dose of PD128907 (0.05 mg/kg) elicited hypoactivity for a shorter duration (first 20 minutes) but elicited the same robust hyperactivity as the higher dose of PD128907. The PD128907-induced locomotor activity was normalized to locomotor activity in control vehicle (saline) injected mice.

FIG. 13, comprising FIGS. 13A-B, illustrates the comparison of PBZI, PD128907 (tolerance-inducing D₃ agonist) and control saline on 6 mg/kg levodopa-induced dyskinesia. The drugs were administered 10 minutes prior to levodopa injection. PD128907 increased AIMs score even before the levodopa was administered. PBZI delayed and reduced the dyskinesia induced by levodopa (FIG. 13A). As illustrated in FIG. 13B, integration of the area under the curves yields the total integrated AIMs score which is significantly reduced by the administration of PBZI (*P<0.05, ANOVA, post-hoc Holm's test).

FIG. 14 is a bar graph illustrating the dose dependent improvement in forelimb bradykinesia following levodopa (L-DOPA) injection, measured using the forelimb stepping test. (*, P<0.01, ANOVA, post hoc Holm's test).

FIG. 15 illustrates the time course of the in vivo experiments.

11

FIG. 16, comprising FIGS. 16A-B, illustrates the finding that activation of D₃ receptor by dopamine and PBZI modulates neuronal firing differently.

FIG. 16A illustrates representative current clamp recording in an AtT-20 neuroendocrine cell stably expressing the human D₃ dopamine receptor. Activation of the D₃ receptor by 100 nM dopamine (black bar), dissolved in standard external solution with 5 mM potassium, hyperpolarized the cell and inhibited the spontaneous action potentials during the first application but not to second or third application. FIG. 16B illustrates that activation of the D₃ receptor by 300 nM PBZI (gray hatched bar), dissolved in standard external solution with 5 mM potassium, hyperpolarized the cell and inhibited the spontaneous action potentials during the first and second treatment.

FIG. 17, comprising FIGS. 17A-D, is a set of graphs illustrating that the novel D₃ receptor agonist, ES609, abolishes tolerance and SRT properties. In FIG. 17A, representative voltage clamp recording showed that 100 nM ES609 (cross hatched bar) induced GIRK currents that did not show tolerance and SRT in an AtT-20 cell stably expressing the human D₃ receptor. In contrast, in the same cell, 100 nM quinpirole (QP, black bar) elicited tolerance and SRT. In FIG. 17B, representative voltage clamp recording showed that neither 300 nM PBZI (hatched bar) nor 300 nM ES609 (cross hatched bar) induced GIRK currents in parental AtT-20 cells in the absence of exogenous D₃ receptor expression. In FIG. 17C, representative voltage clamp recording showed that 100 nM ES609 (cross hatched bar) and 100 nM quinpirole (black bar) induced GIRK currents in AtT-20 cell stably expressing the human D_{2S} receptor. The ES609-induced GIRK current was significantly less than the quinpirole-induced current in D_{2S} receptor expressing cells. The cells were held at -65 mV and the duration of agonist application was ~60 seconds. FIG. 17D illustrates cumulative dose response of ES609-induced GIRK response in AtT-20 cells stably expressing the human D₃ receptor. The black filled circle is the GIRK response elicited by a saturating dose of quinpirole (QP, 300 nM) and showed that ES609 is a full agonist at the D₃ receptor. Error bars represent ±SEM. The GIRK currents were divided by cell capacitance to normalize for cell size. The data points were fit with a four parameter Hill equation.

FIG. 18 is a bar graph illustrating the finding that PBZI and ES609 improve motor deficits in a rat PD model.

FIG. 19, comprising FIGS. 19A-B, is a pair of graphs illustrating the finding that PBZI & ES609 improve levodopa-induced dyskinesia in a rat PD model.

FIG. 20 is a bar graph illustrating the finding that atypical (but not typical) D₃ receptor agonists prevent L-DOPA induced dyskinesia.

FIG. 21, comprising FIGS. 21A-B, is a set of graphs illustrating the effect of PBZI and ES609 on neuron viability in hippocampal cultures (from Day 2 to Day 5 in culture).

FIG. 22, comprising FIGS. 22A-B, is a set of graphs illustrating the effect of PBZI & ES609 on neuroprotection of hippocampal cells treated with 10 mM hydrogen peroxide for 4 hours.

FIG. 23, comprising FIGS. 23A-B, is a set of graphs illustrating the finding that ES609 inhibits monoamine transporter uptake activity.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to the discovery that signal transduction pathways for the D₃ dopamine receptor (amino acid sequence of SEQ ID NO:1) have specific tolerance and slow response termination (SRT) properties. In a neurological

12

disorder that exhibits altered D₃ dopamine receptor expression, the tolerance and slow response termination (SRT) properties are aberrantly expressed and might contribute to the pathology.

In a non-limiting aspect, agonist-induced tolerance in D₃ receptors is associated with a unique conformational state of the receptor. This association suggested that the tolerance and SRT properties of the D₃ receptor are ligand-dependent and that functionally-selective agonists that altered the tolerance-specific conformation would abolish the tolerance and SRT properties. Screening known D₃ receptor agonists and determining their ability to abolish the tolerance and SRT properties allowed the identification of two agonists, cis-8-OH-PBZI (PBZI) and FAUC 73, which while being full agonists at D₃ receptors, completely abolished the receptors' tolerance and SRT properties. In order to distinguish these new D₃ receptor agonists from classical tolerance- and SRT-inducing agonists, they were collectively named atypical D₃ receptor agonists. A pharmacophore model based on the interactions of PBZI with the D₃ receptor was designed as an input to the Hybrid Structure Based (HSB) *in silico* screening method, allowing the identification of an additional novel agonist, ES 609, which also abolished D₃ receptor tolerance and SRT properties.

The present invention thus further relates to the discovery of novel functionally-selective D₃ dopamine receptor agonists that do not elicit the tolerance and slow response termination properties of D₃ receptors. In one aspect, these agonists, rather than preferentially activating signaling pathways, modify the signaling properties of the receptor. In another aspect, this new class of atypical D₃ receptor agonists pharmacologically converts the D₃ receptor to the functional equivalent of a D₂ receptor.

The functional properties of this novel class of atypical D₃ receptor agonists was characterized by studying their ability to activate the D₃ receptor-adenylyl cyclase and G-protein coupled inward rectifier potassium channel signaling pathways. This novel family of D₃ agonists may be used to treat, ameliorate or prevent levodopa-induced dyskinesia (LID) in patients suffering from Parkinson's Disease, or neurological disorders in which there is ectopic overexpression of D₃ receptors.

DEFINITIONS

As used herein, each of the following terms has the meaning associated with it in this section.

Unless defined otherwise, all technical and scientific terms used herein generally have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Generally, the nomenclature used herein and the laboratory procedures in cell culture, molecular genetics, organic chemistry, and peptide chemistry are those well-known and commonly employed in the art.

As used herein, the articles "a" and "an" refer to one or to more than one (i.e. to at least one) of the grammatical object of the article. By way of example, "an element" means one element or more than one element.

As used herein, the term "about" will be understood by persons of ordinary skill in the art and will vary to some extent on the context in which it is used.

As used herein, the term "L-DOPA" refers to levodopa, also known as L-3,4-dihydroxyphenylalanine or a salt thereof.

As used herein, the term "quinpirole" or "QP" refers to (4aR,8aR)-5-propyl-4,4-a,5,6,7,8,8a,9-octahydro-1H-pyrazolo[3,4-g]quinolone or a salt thereof.

As used herein, the term “GR218231” refers to (+)-(2R)-1,2,3,4-tetrahydro-6-[[4-methoxyphenyl)sulfonyl]methyl]-N,N-dipropyl-2-naphthalenamine or a salt thereof.

As used herein, the term “PBZI” refers to (3aS,9bR)-3-propyl-1,2,3a,4,5,9b-hexahydrobenzo[e]indol-8-ol or a salt thereof.

As used herein, the term “clozapine” refers to 8-chloro-11-(4-methylpiperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine or a salt thereof.

As used herein, the term “WST-1” refers to 2-(4-iodophenyl)-3-(4-nitrophenyl)-5-(2,4-disulfophenyl)-2H-tetrazolium, monosodium salt or a salt thereof.

As used herein, the term “PD128907” refers to (4aR,10bR)-3,4-a,4,10b-tetrahydro-4-propyl-2H,5H-[1]benzopyrano-[4,3-b]-1,4-oxazin-9-ol hydrochloride or a salt thereof.

As used herein, the term “70H-DPAT” refers to 7-hydroxy-N,N-dipropyl-2-aminotetralin or a salt thereof.

As used herein, the term “6-OHDA” refers to 6-hydroxydopamine or a salt thereof.

As used herein, the term “pramipexole” refers to (S)-N⁶-propyl-4,5,6,7-tetrahydro-1,3-benzothiazole-2,6-diamine or a salt thereof.

As used herein, the term “rotigotine” refers to (S)-6-[propyl(2-thiophen-2-ylethyl)amino]-5,6,7,8-tetrahydronaphthalen-1-ol or a salt thereof.

As used herein, the term “ES609” or “ES0609” refers to 4-(2-chlorophenyl)-butan-2-amine or a salt thereof.

As used herein, the term “sarizotan” refers to 1-[(2R)-3,4-dihydro-2H-chromen-2-yl]-N-([5-(4-fluorophenyl)pyridin-3-yl]methyl)methanamine or a salt thereof.

As used herein, the term “FAUC73” refers to (4-ethynylcyclohex-3-enyl) dipropylamine or a salt thereof.

As used herein, the term “eticlopride” refers to 3-chloro-5-ethyl-N-[(2S)-1-ethylpyrrolidin-2-yl]methyl}-6-hydroxy-2-methoxybenzamide or a salt thereof.

As used herein, the term “LID” refers to levodopa-induced dyskinesia.

As used herein, the term “tolerance property” as applying to the D₃ receptor refers to the progressive decrease in receptor signaling function upon repeated stimulation by classical agonists, including dopamine.

As used herein, the term “SRT property” or “slow response termination” as applying to the D₃ receptor refers to the increase in time taken to terminate the signaling function of the D₃ receptor, after removal of the agonist.

As used herein, the term “polypeptide” refers to a polymer composed of amino acid residues, related naturally occurring structural variants, and synthetic non-naturally occurring analogs thereof linked via peptide bonds. Synthetic polypeptides may be synthesized, for example, using an automated polypeptide synthesizer. As used herein, the term “protein” typically refers to large polypeptides. As used herein, the term “peptide” typically refers to short polypeptides. Conventional notation is used herein to represent polypeptide sequences: the left-hand end of a polypeptide sequence is the amino-terminus, and the right-hand end of a polypeptide sequence is the carboxyl-terminus.

As used herein, amino acids are represented by the full name thereof, by the three letter code corresponding thereto, or by the one-letter code corresponding thereto, as indicated below:

Full Name	Three-Letter Code	One-Letter Code
Aspartic Acid	Asp	D
Glutamic Acid	Glu	E
Lysine	Lys	K
Arginine	Arg	R
Histidine	His	H

-continued

Full Name	Three-Letter Code	One-Letter Code
Tyrosine	Tyr	Y
Cysteine	Cys	C
Asparagine	Asn	N
Glutamine	Gln	Q
Serine	Ser	S
Threonine	Thr	T
Glycine	Gly	G
Alanine	Ala	A
Valine	Val	V
Leucine	Leu	L
Isoleucine	Ile	I
Methionine	Met	M
Proline	Pro	P
Phenylalanine	Phe	F
Tryptophan	Trp	W

As used herein, the term “treatment” or “treating,” is defined as the application or administration of a therapeutic agent, i.e., a compound of the invention (alone or in combination with another pharmaceutical agent), to a patient, or application or administration of a therapeutic agent to an isolated tissue or cell line from a patient (e.g., for diagnosis or ex vivo applications), who has LID, a symptom of LID or the potential to develop LID, with the purpose to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve or affect LID, the symptoms of LID or the potential to develop LID. The term “treatment” or “treating” is also defined as the application or administration of a therapeutic agent, i.e., a compound of the invention (alone or in combination with another pharmaceutical agent), to a patient, or application or administration of a therapeutic agent to an isolated tissue or cell line from a patient (e.g., for diagnosis or ex vivo applications), who has Parkinson’s Disease, a symptom of Parkinson’s Disease or the potential to develop Parkinson’s Disease, with the purpose to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve or affect Parkinson’s Disease, the symptoms of Parkinson’s Disease or the potential to develop Parkinson’s Disease. Such treatments may be specifically tailored or modified, based on knowledge obtained from the field of pharmacogenomics.

As used herein, the term “prevent” or “prevention” means no disorder or disease development if none had occurred, or no further disorder or disease development if there had already been development of the disorder or disease. Also considered is the ability of one to prevent some or all of the symptoms associated with the disorder or disease.

As used herein, the term “patient” or “subject” refers to a human or a non-human mammal. Non-human mammals include, for example, livestock and pets, such as ovine, bovine, porcine, canine, feline and murine mammals. Preferably, the patient or subject is human.

As used herein, the terms “effective amount,” “pharmaceutically effective amount” and “therapeutically effective amount” refer to a nontoxic but sufficient amount of an agent to provide the desired biological result. That result may be reduction and/or alleviation of the signs, symptoms, or causes of a disease, or any other desired alteration of a biological system. An appropriate therapeutic amount in any individual case may be determined by one of ordinary skill in the art using routine experimentation.

As used herein, the term “pharmaceutically acceptable” refers to a material, such as a carrier or diluent, which does not abrogate the biological activity or properties of the compound, and is relatively non-toxic, i.e., the material may be administered to an individual without causing undesirable

biological effects or interacting in a deleterious manner with any of the components of the composition in which it is contained.

As used herein, the language “pharmaceutically acceptable salt” refers to a salt of the administered compounds prepared from pharmaceutically acceptable non-toxic acids, including inorganic acids, organic acids, solvates, hydrates, or clathrates thereof. Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, sulfuric, and phosphoric. Appropriate organic acids may be selected, for example, from aliphatic, aromatic, carboxylic and sulfonic classes of organic acids, examples of which are formic, acetic, propionic, succinic, camphorsulfonic, citric, fumaric, gluconic, isethionic, lactic, malic, mucic, tartaric, para-toluenesulfonic, glycolic, glucuronic, maleic, furoic, glutamic, benzoic, anthranilic, salicylic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, pantothenic, benzenesulfonic (besylate), stearic, sulfanilic, alginic, galacturonic, and the like.

As used herein, the term “composition” or “pharmaceutical composition” refers to a mixture of at least one compound useful within the invention with a pharmaceutically acceptable carrier. The pharmaceutical composition facilitates administration of the compound to a patient. Multiple techniques of administering a compound exist in the art including, but not limited to, intravenous, oral, aerosol, parenteral, ophthalmic, pulmonary and topical administration.

As used herein, the term “pharmaceutically acceptable carrier” means a pharmaceutically acceptable material, composition or carrier, such as a liquid or solid filler, stabilizer, dispersing agent, suspending agent, diluent, excipient, thickening agent, solvent or encapsulating material, involved in carrying or transporting a compound useful within the invention within or to the patient such that it may perform its intended function. Typically, such constructs are carried or transported from one organ, or portion of the body, to another organ, or portion of the body. Each carrier must be “acceptable” in the sense of being compatible with the other ingredients of the formulation, including the compound useful within the invention, and not injurious to the patient. Some examples of materials that may serve as pharmaceutically acceptable carriers include: sugars, such as lactose, glucose and sucrose; starches, such as corn starch and potato starch; cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients, such as cocoa butter and suppository waxes; oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; glycols, such as propylene glycol; polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; esters, such as ethyl oleate and ethyl laurate; agar; buffering agents, such as magnesium hydroxide and aluminum hydroxide; surface active agents; alginic acid; pyrogen-free water; isotonic saline; Ringer’s solution; ethyl alcohol; phosphate buffer solutions; and other non-toxic compatible substances employed in pharmaceutical formulations. As used herein, “pharmaceutically acceptable carrier” also includes any and all coatings, antibacterial and antifungal agents, and absorption delaying agents, and the like that are compatible with the activity of the compound useful within the invention, and are physiologically acceptable to the patient. Supplementary active compounds may also be incorporated into the compositions. The “pharmaceutically acceptable carrier” may further include a pharmaceutically acceptable salt of the compound useful within the invention. Other additional ingredients that may be included in the pharmaceutical compositions used in the practice of the invention are known in the

art and described, for example in Remington’s Pharmaceutical Sciences (Genaro, Ed., Mack Publishing Co., 1985, Easton, Pa.), which is incorporated herein by reference.

As used herein, the term “alkyl,” by itself or as part of another substituent means, unless otherwise stated, a straight or branched chain hydrocarbon having the number of carbon atoms designated (i.e. C₁₋₆ means one to six carbon atoms) and includes straight, branched chain, or cyclic substituent groups. Examples include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl, neopentyl, hexyl, and cyclopropylmethyl. Most preferred is (C₁-C₆)alkyl, particularly ethyl, methyl, isopropyl, isobutyl, n-pentyl, n-hexyl and cyclopropylmethyl.

As used herein, the term “substituted alkyl” means alkyl, as defined above, substituted by one, two or three substituents selected from the group consisting of halogen, —OH, alkoxy, —NH₂, —N(CH₃)₂, —C(=O)OH, trifluoromethyl, —CN, —C(=O)O(C₁-C₄)alkyl, —C(=O)NH₂, —SO₂NH₂, —C(=NH)NH₂, and —NO₂, preferably containing one or two substituents selected from halogen, —OH, alkoxy, —NH₂, trifluoromethyl, —N(CH₃)₂, and —C(=O)OH, more preferably selected from halogen, alkoxy and —OH. Examples of substituted alkyls include, but are not limited to, 2,2-difluoropropyl, 2-carboxycyclopentyl and 3-chloropropyl.

As used herein, the term “alkoxy” employed alone or in combination with other terms means, unless otherwise stated, an alkyl group having the designated number of carbon atoms, as defined above, connected to the rest of the molecule via an oxygen atom, such as, for example, methoxy, ethoxy, 1-propoxy, 2-propoxy (isopropoxy) and the higher homologs and isomers. Preferred are (C₁-C₃) alkoxy, particularly ethoxy and methoxy.

As used herein, the term “halo” or “halogen” alone or as part of another substituent means, unless otherwise stated, a fluorine, chlorine, bromine, or iodine atom, preferably, fluorine, chlorine, or bromine, more preferably, fluorine or chlorine.

As used herein, the term “heteroalkyl” by itself or in combination with another term means, unless otherwise stated, a stable straight or branched chain alkyl group consisting of the stated number of carbon atoms and one or two heteroatoms selected from the group consisting of O, N, and S, and wherein the nitrogen and sulfur atoms may be optionally oxidized and the nitrogen heteroatom may be optionally quaternized. The heteroatom(s) may be placed at any position of the heteroalkyl group, including between the rest of the heteroalkyl group and the fragment to which it is attached, as well as attached to the most distal carbon atom in the heteroalkyl group. Examples include: —O—CH₂—CH₂—CH₃, —CH₂—CH₂—CH₂—OH, —CH₂—CH₂—NH—CH₃, —CH₂—S—CH₂—CH₃, and —CH₂CH₂—S(=O)—CH₃. Up to two heteroatoms may be consecutive, such as, for example, —CH₂—NH—OCH₃, or —CH₂—CH₂—S—CH₃.

As used herein, the term “aromatic” refers to a carbocycle or heterocycle with one or more polyunsaturated rings and having aromatic character, i.e. having (4n+2) delocalized π (pi) electrons, where n is an integer.

As used herein, the term “aryl,” employed alone or in combination with other terms, means, unless otherwise stated, a carbocyclic aromatic system containing one or more rings (typically one, two or three rings) wherein such rings may be attached together in a pendent manner, such as a biphenyl, or may be fused, such as naphthalene. Examples include phenyl, anthracyl, and naphthyl. Preferred are phenyl and naphthyl, most preferred is phenyl.

As used herein, the term “aryl-(C₁-C₃)alkyl” means a functional group wherein a one to three carbon alkylene chain is attached to an aryl group, e.g., —CH₂CH₂-phenyl. Preferred is aryl-CH₂— and aryl-CH(CH₃)—. The term “substituted aryl-(C₁-C₃)alkyl” means an aryl-(C₁-C₃)alkyl functional group in which the aryl group is substituted. Preferred is substituted aryl(CH₂)—. Similarly, the term “heteroaryl-(C₁-C₃)alkyl” means a functional group wherein a one to three carbon alkylene chain is attached to a heteroaryl group, e.g., —CH₂CH₂-pyridyl. Preferred is heteroaryl-(CH₂)—. The term “substituted heteroaryl-(C₁-C₃)alkyl” means a heteroaryl-(C₁-C₃)alkyl functional group in which the heteroaryl group is substituted. Preferred is substituted heteroaryl-(CH₂)—.

As used herein, the term “heterocycle” or “heterocyclyl” or “heterocyclic” by itself or as part of another substituent means, unless otherwise stated, an unsubstituted or substituted, stable, mono- or multi-cyclic heterocyclic ring system that consists of carbon atoms and at least one heteroatom selected from the group consisting of N, O, and S, and wherein the nitrogen and sulfur heteroatoms may be optionally oxidized, and the nitrogen atom may be optionally quaternized. The heterocyclic system may be attached, unless otherwise stated, at any heteroatom or carbon atom that affords a stable structure. A heterocycle may be aromatic or non-aromatic in nature. In one embodiment, the heterocycle is a heteroaryl.

As used herein, the term “heteroaryl” or “hetero aromatic” refers to a heterocycle having aromatic character. A polycyclic heteroaryl may include one or more rings that are partially saturated. Examples include tetrahydroquinoline and 2,3-dihydrobenzofuryl.

Examples of non-aromatic heterocycles include monocyclic groups such as aziridine, oxirane, thiirane, azetidine, oxetane, thietane, pyrrolidine, pyrroline, imidazoline, pyrazolidine, dioxolane, sulfolane, 2,3-dihydrofuran, 2,5-dihydrofuran, tetrahydrofuran, thiophane, piperidine, 1,2,3,6-tetrahydropyridine, 1,4-dihydropyridine, piperazine, morpholine, thiomorpholine, pyran, 2,3-dihydropyran, tetrahydropyran, 1,4-dioxane, 1,3-dioxane, homopiperazine, homopiperidine, 1,3-dioxepane, 4,7-dihydro-1,3-dioxepin and hexamethyleneoxide.

Examples of heteroaryl groups include pyridyl, pyrazinyl, pyrimidinyl (particularly 2- and 4-pyrimidinyl), pyridazinyl, thienyl, furyl, pyrrolyl (particularly 2-pyrrolyl), imidazolyl, thiazolyl, oxazolyl, pyrazolyl (particularly 3- and 5-pyrazolyl), isothiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,3,4-triazolyl, tetrazolyl, 1,2,3-thiadiazolyl, 1,2,3-oxadiazolyl, 1,3,4-thiadiazolyl and 1,3,4-oxadiazolyl.

Examples of polycyclic heterocycles include indolyl (particularly 3-, 4-, 5-, 6- and 7-indolyl), indolinyl, quinolyl, tetrahydroquinolyl, isoquinolyl (particularly 1- and 5-isoquinolyl), 1,2,3,4-tetrahydroisoquinolyl, cinnolinyl, quinoxaliny (particularly 2- and 5-quinoxaliny), quinazoliny, phthalazinyl, 1,8-naphthyridinyl, 1,4-benzodioxanyl, coumarin, dihydrocoumarin, 1,5-naphthyridinyl, benzofuryl (particularly 3-, 4-, 5-, 6- and 7-benzofuryl), 2,3-dihydrobenzofuryl, 1,2-benzisoxazolyl, benzothienyl (particularly 3-, 4-, 5-, 6-, and 7-benzothienyl), benzoxazolyl, benzothiazolyl (particularly 2-benzothiazolyl and 5-benzothiazolyl), purinyl, benzimidazolyl (particularly 2-benzimidazolyl), benzotriazolyl, thioxanthinyl, carbazolyl, carbolinyl, acridinyl, pyrrolizidinyl, and quinolizidinyl.

The aforementioned listing of heterocyclyl and heteroaryl moieties is intended to be representative and not limiting.

As used herein, the term “substituted” means that an atom or group of atoms has replaced hydrogen as the substituent attached to another group.


For aryl, aryl-(C₁-C₃)alkyl and heterocyclyl groups, the term “substituted” as applied to the rings of these groups refers to any level of substitution, namely mono-, di-, tri-, tetra-, or penta-substitution, where such substitution is permitted. The substituents are independently selected, and substitution may be at any chemically accessible position. In one embodiment, the substituents vary in number between one and four. In another embodiment, the substituents vary in number between one and three. In yet another embodiment, the substituents vary in number between one and two. In yet another embodiment, the substituents are independently selected from the group consisting of C₁₋₆ alkyl, —OH, C₁₋₆ alkoxy, halo, amino, acetamido and nitro. In yet another embodiment, the substituents are independently selected from the group consisting of C₁₋₆ alkyl, C₁₋₆ alkoxy, halo, acetamido, and nitro. As used herein, where a substituent is an alkyl or alkoxy group, the carbon chain may be branched, straight or cyclic, with straight being preferred.

“Instructional material,” as that term is used herein, includes a publication, a recording, a diagram, or any other medium of expression that may be used to communicate the usefulness of the compounds of the invention. In some instances, the instructional material may be part of a kit useful for effecting alleviating or treating the various diseases or disorders recited herein. Optionally, or alternately, the instructional material may describe one or more methods of alleviating the diseases or disorders in a cell or a tissue of a mammal. The instructional material of the kit may, for example, be affixed to a container that contains the compounds of the invention or be shipped together with a container that contains the compounds. Alternatively, the instructional material may be shipped separately from the container with the intention that the recipient uses the instructional material and the compound cooperatively. For example, the instructional material is for use of a kit; instructions for use of the compound; or instructions for use of a formulation of the compound.

Compounds of the Invention

The compounds useful within the invention may be synthesized using techniques well-known in the art of organic synthesis.

In one aspect, the compound useful within the invention has the formula (I):



wherein in formula (I):

R¹, R² and R³ are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl-(C₁₋₃)alkyl, substituted aryl-(C₁₋₃)alkyl, carboxy, alkylcarboxy, formyl, alkyl-carbonyl, aryl-carbonyl, and heteroaryl-carbonyl;

19

R^4 and R^5 are independently selected from the group consisting of H, C_{1-6} alkyl, substituted C_{1-6} alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl- (C_{1-3}) alkyl, and substituted aryl- (C_{1-3}) alkyl; and,

n is 2, 3, 4 or 5;


or a pharmaceutically acceptable salt thereof.

In one embodiment, R^1 , R^2 and R^3 are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C_{1-6} alkyl, substituted C_{1-6} alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, carboxy, alkylcarboxy, formyl, alkyl-carbonyl, aryl-carbonyl, and heteroaryl-carbonyl. In another embodiment, R^1 , R^2 and R^3 are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C_{1-6} alkyl, substituted C_{1-6} alkyl, heteroalkyl, carboxy, alkylcarboxy, formyl, and alkyl-carbonyl. In yet another embodiment, R^1 , R^2 and R^3 are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C_{1-6} alkyl, substituted C_{1-6} alkyl, heteroalkyl, and carboxy. In yet another embodiment, R^1 , R^2 and R^3 are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C_{1-6} alkyl, and carboxy. In yet another embodiment, R^1 , R^2 and R^3 are independently selected from the group consisting of H, cyano, halo, alkoxy, nitro, C_{1-6} alkyl, and carboxy. In yet another embodiment, R^1 and R^2 are H, and R^3 is chloro. In yet another embodiment, R^1 , R^2 and R^3 are independently selected from the group consisting of H, fluoro, chloro, bromo, iodo, methoxy, ethoxy, hydroxyl, methyl, ethyl or other C_{1-6} alkyl.

In one embodiment, n is 2, 3 or 4. In another embodiment, n is 2 or 3. In yet another embodiment, n is 2.


In one embodiment, R^4 and R^5 are independently selected from the group consisting of H, C_{1-6} alkyl, substituted C_{1-6} alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, and substituted aryl. In another embodiment, R^4 and R^5 are independently selected from the group consisting of H, C_{1-6} alkyl, substituted C_{1-6} alkyl, heteroalkyl, heterocyclyl, and substituted heterocyclyl. In yet another embodiment, R^4 and R^5 are independently selected from the group consisting of H, C_{1-6} alkyl, substituted C_{1-6} alkyl, and heteroalkyl. In yet another embodiment, R^4 and R^5 are independently selected from the group consisting of H, C_{1-6} alkyl, and substituted C_{1-6} alkyl. In yet another embodiment, R^4 and R^5 are methyl. In yet another embodiment, R^5 is H, methyl, ethyl, prop-1-yl, prop-2-yl, hydroxymethyl, 1-hydroxy-ethyl, 2-hydroxy-ethyl, 1-hydroxy-prop-1-yl, 2-hydroxy-prop-1-yl, 3-hydroxy-prop-1-yl, 1-hydroxy-prop-2-yl or 2-hydroxy-prop-2-yl.

In one embodiment, the compound useful within the invention is selected from the group consisting of:
2-amino-4-(2-chlorophenyl)butan-1-ol




20


2-(3-aminohexyl)phenol




4-(2-chlorophenyl)-butan-2-amine




4-(2-chlorophenyl)-2-methylamino-butane (also known as 4-(2-chlorophenyl)-N-methylbutan-2-amine)



4-(2-fluorophenyl)butan-2-amine




4-(2-bromophenyl)butan-2-amine




21


4-(2-iodophenyl)butan-2-amine




4-(2-methoxyphenyl)butan-2-amine




2-(3-aminobutyl)phenol



3-(3,4-diethoxyphenyl)propan-1-amine




4-(4-chlorophenyl)butan-2-amine



22



4-(4-methoxyphenyl)butan-2-amine



15 mixtures thereof, or a pharmaceutically acceptable salt thereof.

In another aspect, the compound useful within the invention has the formula (IIa) or (IIb):

20



40 wherein in formula (IIa) or (IIb):

R^1 and R^2 are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C_{1-6} alkyl, substituted C_{1-6} alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl- (C_{1-3}) alkyl, substituted aryl- (C_{1-3}) alkyl, carboxy, alkylcarboxy, formyl, alkyl-carbonyl, aryl-carbonyl, and heteroaryl-carbonyl;

45 R^3 and R^4 are independently selected from the group consisting of H, C_{1-6} alkyl, aryl, heteroaryl, and substituted C_{1-6} alkyl;

50 R^5 is selected from the group consisting of H, C_{1-6} alkyl, substituted C_{1-6} alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl- (C_{1-3}) alkyl, and substituted aryl- (C_{1-3}) alkyl; and,

55 m is 1, 2, 3 or 4;

or a pharmaceutically acceptable salt thereof.

In one embodiment, R^1 and R^2 are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C_{1-6} alkyl, substituted C_{1-6} alkyl, carboxy, alkylcarboxy, formyl, alkyl-carbonyl, aryl-carbonyl, and heteroaryl-carbonyl. In another embodiment, R^1 and R^2 are independently selected from the group consisting of H, cyano, hydroxyl, halo, and alkoxy, C_{1-6} alkyl, and substituted C_{1-6} alkyl.


65 In one embodiment, R^3 and R^4 are independently selected from the group consisting of H, and C_{1-6} alkyl.

23


In one embodiment, R^5 is selected from the group consisting of H, C_{1-6} alkyl, and substituted C_{1-6} alkyl.

In one embodiment, m is 1, 2 or 3.


In one embodiment, the compound useful within the invention is selected from the group consisting of
2-(5-chloro-1-methyl-1H-indol-3-yl)ethanamine



1-(5-fluoro-1-methyl-1H-indol-3-yl)propan-2-amine




1-(5-methoxy-1-methyl-1H-indol-3-yl)propan-2-amine



mixtures thereof, or a pharmaceutically acceptable salt thereof.


In yet another aspect, the compound useful within the invention is selected from the group consisting of:

2,7-diamino-5-(4-fluorophenyl)-4-oxo-3,4,5,6-tetrahydropyrido[2,3-d]pyrimidine-6-carbonitrile



24


(Z)-2-(1H-benzo[d]imidazol-2-yl)-N'-hydroxy-3-(4-methoxyphenyl)propanimidamide



mixtures thereof, or a pharmaceutically acceptable salt thereof.

Methods of the Invention

In one aspect, the invention includes a method of treating, ameliorating or preventing levodopa-induced dyskinesia in a patient suffering from Parkinson's Disease. The method comprises administering to the patient a pharmaceutical composition comprising a therapeutically effective amount of at least one compound selected from the group consisting of:
a compound of formula (I):





wherein in formula (I):

R^1 , R^2 and R^3 are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C_{1-6} alkyl, substituted C_{1-6} alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl-(C_{1-3})alkyl, substituted aryl-(C_{1-3})alkyl, carboxy, alkylcarboxy, formyl, alkyl-carbonyl, aryl-carbonyl, and heteroaryl-carbonyl;

R^4 and R^5 are independently selected from the group consisting of H, C_{1-6} alkyl, substituted C_{1-6} alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl-(C_{1-3})alkyl, and substituted aryl-(C_{1-3})alkyl; and,
n is 2, 3, 4 or 5;

a compound of formula (IIa) or (IIb):



25

wherein in formula (IIa) or (IIb):

R¹ and R² are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl-(C₁₋₃)alkyl, substituted aryl-(C₁₋₃)alkyl, carboxy, alkylcarboxy, formyl, alkyl-carbonyl, aryl-carbonyl, and heteroaryl-carbonyl;

R³ and R⁴ are independently selected from the group consisting of H, C₁₋₆ alkyl, aryl, heteroaryl, and substituted C₁₋₆ alkyl;

R⁵ is selected from the group consisting of H, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl-(C₁₋₃)alkyl, and substituted aryl-(C₁₋₃)alkyl; and,

m is 1, 2, 3 or 4;

2,7-diamino-5-(4-fluorophenyl)-4-oxo-3,4,5,6-tetrahydropyrido[2,3-d]pyrimidine-6-carbonitrile;

(Z)-2-(1H-benzo[d]imidazol-2-yl)-N¹-hydroxy-3-(4-methoxyphenyl)propanimidamide;

mixtures thereof, or a pharmaceutically acceptable salt thereof.

In one embodiment, R¹, R² and R³ are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, carboxy, alkylcarboxy, formyl, alkyl-carbonyl, aryl-carbonyl, and heteroaryl-carbonyl. In another embodiment, R¹, R² and R³ are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, heteroalkyl, carboxy, alkylcarboxy, formyl, and alkyl-carbonyl. In yet another embodiment, R¹, R² and R³ are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, heteroalkyl, and carboxy. In yet another embodiment, R¹, R² and R³ are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, heteroalkyl, and carboxy. In yet another embodiment, R¹, R² and R³ are independently selected from the group consisting of H, cyano, halo, alkoxy, nitro, C₁₋₆ alkyl, and carboxy. In yet another embodiment, R¹ and R² are H, and R³ is chloro. In yet another embodiment, R¹, R² and R³ are independently selected from the group consisting of H, fluoro, chloro, bromo, iodo, methoxy, and ethoxy.

In one embodiment, R¹ and R² are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, carboxy, alkylcarboxy, formyl, alkyl-carbonyl, aryl-carbonyl, and heteroaryl-carbonyl. In another embodiment, R¹ and R² are independently selected from the group consisting of H, cyano, hydroxyl, halo, and alkoxy, C₁₋₆ alkyl, and substituted C₁₋₆ alkyl.

In one embodiment, n is 2, 3 or 4. In another embodiment, n is 2 or 3. In yet another embodiment, n is 2.

In one embodiment, R⁴ and R⁵ are independently selected from the group consisting of H, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, and substituted aryl. In another embodiment, R⁴ and R⁵ are independently selected from the group consisting of H, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, heteroalkyl, heterocyclyl, and substituted heterocyclyl. In yet another embodiment, R⁴ and R⁵ are independently selected from the group consisting of H, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, and heteroalkyl. In yet another embodiment, R⁴ and R⁵ are independently selected from the group consisting of H, C₁₋₆ alkyl, and substituted C₁₋₆ alkyl. In yet another embodiment, R⁴ and R⁵ are methyl.

26

In yet another embodiment, R⁵ is H, methyl, ethyl, prop-1-yl, prop-2-yl, hydroxymethyl, 1-hydroxy-ethyl, 2-hydroxy-ethyl, 1-hydroxy-prop-1-yl, 2-hydroxy-prop-1-yl, 3-hydroxy-prop-1-yl, 1-hydroxy-prop-2-yl or 2-hydroxy-prop-2-yl.

In one embodiment, R³ and R⁴ are independently selected from the group consisting of H, and C₁₋₆ alkyl.

In one embodiment, R⁵ is selected from the group consisting of H, C₁₋₆ alkyl, and substituted C₁₋₆ alkyl.

In one embodiment, m is 1, 2 or 3.


In one embodiment, the compound useful within the invention is selected from the group consisting of: 2-amino-4-(2-chlorophenyl)butan-1-ol; 2-(3-aminoethyl)phenol; 4-(2-chlorophenyl)-butan-2-amine; 4-(2-chlorophenyl)-2-methylamino-butane; 4-(2-fluorophenyl)butan-2-amine; 4-(2-bromophenyl)butan-2-amine; 4-(2-iodophenyl)butan-2-amine; 4-(2-methoxyphenyl)butan-2-amine; 2-(3-aminobutyl)phenol; 3-(3,4-diethoxyphenyl)propan-1-amine; 4-(4-chlorophenyl)butan-2-amine; 4-(4-methoxyphenyl)butan-2-amine; 2-(5-chloro-1-methyl-1H-indol-3-yl)ethanamine; 1-(5-fluoro-1-methyl-1H-indol-3-yl)propan-2-amine; 1-(5-methoxy-1-methyl-1H-indol-3-yl)propan-2-amine; 2,7-diamino-5-(4-fluorophenyl)-4-oxo-3,4,5,6-tetrahydropyrido[2,3-d]pyrimidine-6-carbonitrile; (Z)-2-(1H-benzo[d]imidazol-2-yl)-N¹-hydroxy-3-(4-methoxyphenyl)propanimidamide;

mixtures thereof, or a pharmaceutically acceptable salt thereof.

In one embodiment, the composition further comprises a drug selected from the group consisting of levodopa, clozapine, bromocriptine, pergolide, pramipexole, ropinirole, piribedil, cabergoline, apomorphine, and lisuride, a salt thereof and mixtures thereof. In another embodiment, the pharmaceutical composition is co-administered to the patient with a second pharmaceutical composition comprising levodopa. In yet another embodiment, the pharmaceutical composition is administered to the patient a given period of time before a second pharmaceutical composition comprising levodopa is administered to the patient. In yet another embodiment, the given period of time varies from about 2 minutes to about 24 hours. In yet another embodiment, the patient is human.

In another aspect, the invention includes a method of treating, ameliorating or preventing Parkinson's Disease in a patient. The method comprises administering to the patient a pharmaceutical composition comprising a therapeutically effective amount of at least one compound selected from the group consisting of:

a compound of formula (I):



(I)



wherein in formula (I):

R¹, R² and R³ are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C₁₋₆ alkyl, substituted C₁₋₆ alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl-(C₁₋₃)alkyl, substituted aryl-(C₁₋₃)alkyl, carboxy, alkylcarboxy, formyl, alkyl-carbonyl, aryl-carbonyl, and heteroaryl-carbonyl;

27

R^4 and R^5 are independently selected from the group consisting of H, C_{1-6} alkyl, substituted C_{1-6} alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl- (C_{1-3}) alkyl, and substituted aryl- (C_{1-3}) alkyl; and,
n is 2, 3, 4 or 5;

a compound of formula (IIa) or (IIb):



wherein in formula (IIa) or (IIb):

R^1 and R^2 are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C_{1-6} alkyl, substituted C_{1-6} alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl- (C_{1-3}) alkyl, substituted aryl- (C_{1-3}) alkyl, carboxy, alkylcarboxy, formyl, alkyl-carbonyl, aryl-carbonyl, and heteroaryl-carbonyl;

R^3 and R^4 are independently selected from the group consisting of H, C_{1-6} alkyl, aryl, heteroaryl, and substituted C_{1-6} alkyl;

R^5 is selected from the group consisting of H, C_{1-6} alkyl, substituted C_{1-6} alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, substituted aryl, aryl- (C_{1-3}) alkyl, and substituted aryl- (C_{1-3}) alkyl; and,

m is 1, 2, 3 or 4;

2,7-diamino-5-(4-fluorophenyl)-4-oxo-3,4,5,6-tetrahydropyrido[2,3-d]pyrimidine-6-carbonitrile;

(Z)-2-(1H-benzo[d]imidazol-2-yl)-N'-hydroxy-3-(4-methoxyphenyl)propanimidamide;

mixtures thereof, or a pharmaceutically acceptable salt thereof.

In one embodiment, R^1 , R^2 and R^3 are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C_{1-6} alkyl, substituted C_{1-6} alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, carboxy, alkylcarboxy, formyl, alkyl-carbonyl, aryl-carbonyl, and heteroaryl-carbonyl. In another embodiment, R^1 , R^2 and R^3 are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C_{1-6} alkyl, substituted C_{1-6} alkyl, heteroalkyl, carboxy, alkylcarboxy, formyl, and alkyl-carbonyl. In yet another embodiment, R^1 , R^2 and R^3 are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C_{1-6} alkyl, and carboxy. In yet another embodiment, R^1 , R^2

28

and R^3 are independently selected from the group consisting of H, cyano, halo, alkoxy, nitro, C_{1-6} alkyl, and carboxy. In yet another embodiment, R^1 and R^2 are H, and R^3 is chloro. In yet another embodiment, R^1 , R^2 and R^3 are independently selected from the group consisting of H, fluoro, chloro, bromo, iodo, methoxy, and ethoxy.

In one embodiment, R^1 and R^2 are independently selected from the group consisting of H, cyano, hydroxyl, amino, acetamido, halo, alkoxy, nitro, C_{1-6} alkyl, substituted C_{1-6} alkyl, carboxy, alkylcarboxy, formyl, alkyl-carbonyl, aryl-carbonyl, and heteroaryl-carbonyl. In another embodiment, R^1 and R^2 are independently selected from the group consisting of H, cyano, hydroxyl, halo, and alkoxy, C_{1-6} alkyl, and substituted C_{1-6} alkyl.

In one embodiment, n is 2, 3 or 4. In another embodiment, n is 2 or 3. In yet another embodiment, n is 2.

In one embodiment, R^4 and R^5 are independently selected from the group consisting of H, C_{1-6} alkyl, substituted C_{1-6} alkyl, heteroalkyl, heterocyclyl, substituted heterocyclyl, aryl, and substituted aryl. In another embodiment, R^4 and R^5 are independently selected from the group consisting of H, C_{1-6} alkyl, substituted C_{1-6} alkyl, heteroalkyl, heterocyclyl, and substituted heterocyclyl. In yet another embodiment, R^4 and R^5 are independently selected from the group consisting of H, C_{1-6} alkyl, substituted C_{1-6} alkyl, and heteroalkyl. In yet another embodiment, R^4 and R^5 are independently selected from the group consisting of H, C_{1-6} alkyl, and substituted C_{1-6} alkyl. In yet another embodiment, R^4 and R^5 are methyl.

In yet another embodiment, R^5 is H, methyl, ethyl, prop-1-yl, prop-2-yl, hydroxymethyl, 1-hydroxy-ethyl, 2-hydroxy-ethyl, 1-hydroxy-prop-1-yl, 2-hydroxy-prop-1-yl, 3-hydroxy-prop-1-yl, 1-hydroxy-prop-2-yl or 2-hydroxy-prop-2-yl.

In one embodiment, R^3 and R^4 are independently selected from the group consisting of H, and C_{1-6} alkyl.

In one embodiment, R^5 is selected from the group consisting of H, C_{1-6} alkyl, and substituted C_{1-6} alkyl.

In one embodiment, m is 1, 2 or 3.

In one embodiment, the compound useful within the invention is selected from the group consisting of: 2-amino-4-(2-chlorophenyl)butan-1-ol; 2-(3-aminoethyl)phenol; 4-(2-chlorophenyl)-butan-2-amine; 4-(2-chlorophenyl)-2-methylamino-butane; 4-(2-fluorophenyl)butan-2-amine; 4-(2-bromophenyl)butan-2-amine; 4-(2-iodophenyl)butan-2-amine; 4-(2-methoxyphenyl)butan-2-amine; 2-(3-aminobutyl)phenol; 3-(3,4-diethoxyphenyl)propan-1-amine; 4-(4-chlorophenyl)butan-2-amine; 4-(4-methoxyphenyl)butan-2-amine; 2-(5-chloro-1-methyl-1H-indol-3-yl)ethanamine; 1-(5-fluoro-1-methyl-1H-indol-3-yl)propan-2-amine; 1-(5-methoxy-1-methyl-1H-indol-3-yl)propan-2-amine; 2,7-diamino-5-(4-fluorophenyl)-4-oxo-3,4,5,6-tetrahydropyrido[2,3-d]pyrimidine-6-carbonitrile; (Z)-2-(1H-benzo[d]imidazol-2-yl)-N'-hydroxy-3-(4-methoxyphenyl)propanimidamide; mixtures thereof, or a pharmaceutically acceptable salt thereof.

In one embodiment, the composition further comprises a drug selected from the group consisting of levodopa, clozapine, bromocriptine, pergolide, pramipexole, ropinirole, piribedil, cabergoline, apomorphine, and lisuride, a salt thereof and mixtures thereof. In another embodiment, the patient is further administered a composition comprising a drug selected from the group consisting of levodopa, clozapine, bromocriptine, pergolide, pramipexole, ropinirole, piribedil, cabergoline, apomorphine, and lisuride, a salt thereof and mixtures thereof.

Molecular Basis of LID in Patients Suffering from Parkinson's Disease

The molecular mechanisms underlying the development of LID in patients suffering from Parkinson's Disease are still not well understood. Studies have shown that expression of a number of genes is altered in dyskinetic animals. In particular, in both rodents and primates, studies have reported a specific increase in the D₃ dopamine receptor expression in the basal ganglia of dyskinetic animals. The functional consequence of the increased D₃ receptor expression, in areas that normally express the D₂ dopamine receptor, is not known.

According to the concept of functional selectivity, ligands may differentially activate signaling pathways coupled to a single GPCR (Kenakin, 2003, Trends Pharmacol. Sci. 24(7): 346-354; Urban et al., 2007, J. Pharmacol. Exp. Ther. 320(1): 1-13). Functional selectivity has been previously reported for D₂ receptors (Gay et al., 2004, Mol. Pharmacol. 66(1):97-105). Sarizotan, a ligand with affinity at D₂-like dopamine receptors, exhibits functional selectivity at D_{2L} and D_{4.2} dopamine receptors (Kuzhikandathil & Bartoszyk, 2006, Neuropharmacology 51:873-884). Sarizotan is a partial agonist for the D_{2L}- and D_{4.2}-GIRK channel signaling pathway but a full agonist at the D_{2L}- and D_{4.2}-adenylyl cyclase pathway (Kuzhikandathil & Bartoszyk, 2006, Neuropharmacology 51:873-884). Subsequent studies revealed that sarizotan induces tolerance and slow response termination (SRT) properties on D_{2S} dopamine receptors. This suggests that, in addition to differentially activating signaling pathways, certain agonists could modulate intrinsic receptor properties such as the tolerance and slow response termination properties.

The D₃ dopamine receptor exhibits tolerance and slow response termination (SRT) properties that distinguish it from the D₂ dopamine receptors (FIG. 1). The tolerance property of D₃ receptor describes the progressive decrease in receptor signaling function upon repeated stimulation by classical agonists, including dopamine. The SRT property describes the prolongation of time taken to terminate the signaling function of the D₃ receptor, after removal of the agonist.

D₃ receptor tolerance and SRT has been observed in cultured substantia nigra neurons, and with human and mouse D₃ receptors heterologously expressed in *Xenopus oocytes*, CHO-K1 cells and AtT-20 cells. Tolerance and SRT properties are elicited by the native agonist dopamine and by various synthetic agonists tested to date. The tolerance and SRT properties of D₃ receptors are elicited over a broad range of agonist concentrations, being observed at 10, 30, 100 and 1000 nM. The properties are manifest in D₃-GIRK, D₃-ACV and D₃-MAP kinase signaling pathways.

Differences in the properties of D₂ and D₃ receptors give rise to a differential modulation of neuronal firing (FIG. 1). These results suggest a model in which aberrant expression of D₃ receptor tolerance and SRT properties could result in aberrant modulation of neuronal firing in the basal ganglia of the dyskinetic animals and contribute to the development of LID in Parkinson's Disease. According to this model, if D₃ receptor tolerance and SRT properties could be abolished, the modulation of neuronal firing by the over-expressed D₃ receptor in the basal ganglia of dyskinetic animals would be similar to the natively expressed D₂ receptors and potentially prevent the expression of dyskinesia.

Initial Studies on Dopamine D₃ Agonists

To identify agonists that could abolish the D₃ dopamine receptor tolerance and SRT, synthetic D₃ receptor-preferring agonists were functionally screened. The functional activity of the compounds were tested by stimulating human D₂, D₃ and D₄ dopamine receptors individually expressed in the AtT-

20 cell line. These dopamine receptors couple to and activate the G-protein coupled inward rectifier potassium (GIRK) channels and also inhibit the endogenous adenylyl cyclase in AtT-20 cells (Kuzhikandathil et al., 2004, Mol. Cell. Neurosci. 26:144-155; Westrich & Kuzhikandathil, 2007, Biochim. Biophys. Acta-Mol. Cell. Res. 1773:1747-1758).

The activation of the GIRK channels was measured electrophysiologically using whole cell voltage clamp recording (Kuzhikandathil et al., 2004, Mol. Cell. Neurosci. 26:144-155). The GIRK channel responses elicited with new ligands were compared to quinpirole, a classical full agonist of D₂, D₃ and D₄ dopamine receptors. The adenylyl cyclase inhibition was determined using a commercially available ELISA assay (Westrich & Kuzhikandathil, 2007, Biochim. Biophys. Acta-Mol. Cell. Res. 1773:1747-1758).

An initial functional screen identified cis-8-hydroxy-3-(n-propyl)-1,2,3a,4,5,9b-hexahydro-1H-benz[e]indole hydrobromide (PBZI). This compound has been reported as a D₃ dopamine receptor agonist; nevertheless, the present studies indicate that, unlike traditional D₃ receptor agonists, PBZI does not elicit tolerance and SRT properties (FIG. 2).

Hybrid Structure Based (HSB) Protocol

The Hybrid Structure Based (HSB) protocol (Koratgere & Welsh, 2006, J. Comp. Aided Mol. Des. 20:789-802; FIG. 3) was used to identify additional compounds belonging to the novel class of D₃ receptor agonists that do not elicit tolerance and SRT properties. The HSB protocol consists of 5 sub-phases (FIG. 3). Phase I corresponds to building a comprehensive electronic database of vendor available small molecules. Phase II corresponds to developing and screening compounds using a hybrid pharmacophore starting from the 3D structural snapshots from the MD simulation. Phase III corresponds to subjecting compounds that pass the hybrid pharmacophore screening to clustering, filtering, chemical space analysis and classification models to develop the enriched database of small molecules. Phase IV corresponds to docking molecules from the enriched database to the DRD3 receptor, with scores derived from consensus scoring schemes. Phase V corresponds to testing best ranking compounds for their activity against all the dopamine receptors. Develop Comprehensive Electronic Database of Small Molecules:

A subset of the Zinc database (Irwin & Soichet, 2005, J. Chem. Inf. Models 45:177-82) consisting of compounds from commercial vendors, along with other compounds including natural products, ligands from PDB and FDA approved drugs, form the entire database of nearly 3 million compounds. All compounds were acquired as sdf formatted files, converted into Mol2 format and energy minimized using Tripos force field. Further, all molecules in the database were filtered for redundancy and renamed according to their corresponding vendor listing.

Develop a Combined Ligand-Protein Pharmacophore:

Generating the combined pharmacophore (also called the hybrid pharmacophore) is an important step of the method. Hence, customizing the pharmacophore to capture the essential features of interactions between PBZI and D₃ receptor is important.

Such information was obtained from the 3D structural complex of PBZI-D₃ receptor complex. The homology model of D₃ receptor was built using the crystal structure of beta2 adrenergic receptor (B2AR) with a partial inverse agonist bound to it (2RH1) (Cherezov et al., 2007, Science 318(5854):1258-65). The transmembrane (TM) regions and the loop regions bore significant homology to the corresponding residues from the beta adrenergic crystal structure. The third intracellular loop in the B2AR was longer than the

corresponding loop in the D₃ receptor and hence a large deletion shown in the alignment (FIG. 4) was not modeled. However, the rest of the third intracellular loop, shown to be very significant for the SRT and receptor tolerance, was modeled using the aligned coordinates from the B2AR structure. Further the entire structure was refined using energy minimization and molecular dynamics simulation with a 2 ns long production run. All simulations were performed using NAMD (Kale et al., 1999, J. Comput. Physics 151(1):283-312) with CHARMM force field (Brooks et al., 1983, J. Comput. Chem. 4:187-217).

The refined structure was used for further docking experiments. Dopamine, PD128907 and PBZI were docked using GOLD program (ver4.0) (Jones et al., 1997, J. Mol. Biol. 267:727-48). The docked complexes were energy minimized using NAMD program. Key interactions between PBZI and the D₃ receptor were used to build the hybrid pharmacophore (FIG. 5). Various pharmacophore models were generated involving a combination of the four pharmacophore elements that are key to designing D₃ receptor agonists, such as the salt bridge with the aspartic acid in TM3 and hydrophobic interactions with the aromatic amino acid cluster from TM5 and TM6. Further, a small interchange of residues between TM2 and TM3 was shown to be instrumental in designing highly selective D₄ agonists and antagonist (Kortagere et al., 2004, Mol. Pharmacol. 66:1491-99). These biochemical and functional evidence were integrated into the design of the hybrid pharmacophore, which was then used to screen small molecule databases (which were described above).

A preliminary screening using the hydrophobic core and the two hydrogen bond acceptor-donor pair resulted in 85 compounds that satisfy the pharmacophore and dock to the D₃ receptor with high scores. One representative compound 4-(2-chlorophenyl)-butan-2-amine (FIG. 6) was used for in vitro and in vivo functional characterization. 4-(2-Chlorophenyl)-butan-2-amine docked very similar to the parent PBZI compound in the D₃ receptor, making the required salt bridge interaction with Asp110 and hydrophobic aromatic interactions with aromatic residues such as Phe345 and His349 (FIG. 6).

Filtering, Chemical Space Analysis and Clustering Modules:

Filtering Schemes:

Compounds that result from pharmacophore based screening are clustered based on their physicochemical properties, such as shape, log P, volume, TPSA and molecular weight, derived from CHEMAXON program. Lipinski's rule of five is applied as a first filter, and a regression based blood-brain-barrier (BBB) penetration model that may filter out compounds for BBB penetration is the second level of filter (Kortagere et al., 2008, Pharm. Res. 8:1836-45). The BBB regression model is a generalized model, described as:

$$\log \text{BB}(\text{pred}) = 0.3408 * \log P - 0.0192 * \text{TPSA} + 0.2503 * a_{\text{N}} + 0.1467 * a_{\text{O}} + 0.1069 * \log s - 0.0011 * \text{mass} - 0.0001 * \text{volume} + 0.0602 * \# \text{rot. bonds.}$$

where: a_N is number for nitrogen atoms, a_O is number of oxygen atoms, TPSA is topological polar surface area, log S is solubility and log P is a water/octanol partition coefficient and measure of hydrophobicity, and # rot. bonds is the number of rotatable bonds.

Chemical Space Analysis and Clustering Techniques:

Chemical space analysis of the compounds is performed to analyze if the compounds belong to one or more chemical classes. Outliers that are chemically divergent against the query structure are removed; however a broad cutoff is used to allow for scaffold hopping. Clustering and QSAR based classification of the compounds is performed using support vec-

tor machine (SVM) techniques. Methods and application of the SVM based models to classify compounds that may penetrate BBB and to classify PXR activators have been done with good accuracy.

Hence adopting this method to classify the compounds based on the target and to generate an enrichment of the database is useful to avoid screening undesired molecules. The molecules in the database developed herein do not contain stereoisomers. During the enrichment process, compounds that are chemically identical are removed for redundancy by retaining only a single copy of the molecule.

Develop Customized Scoring Schemes:

The GOLD program is used for preliminary docking. The active site for established ligands is defined as all residues that may encompass the ligand within an 8 Å radius sphere. The "library screening mode" option in GOLD docking program is used for fast docking. Further, given the non-deterministic nature of genetic algorithms, 50 independent docking runs are performed for each ligand. The full set of docked structures is energy minimized using the molecular modeling package SYBYL (Tripos Inc., St Louis, Mo., USA). The docked receptor-ligand complexes is scored using a customizable knowledge based scoring function based on the nature of interaction of every ligand atom with the protein atom. Details of the method and the normalization scheme have been discussed in Kortagere et al. (Pharm. Res. 2009, 4:1001-11) Further, a consensus scoring scheme that involves the Goldscore, Chemscore, contact score and shape weighted scoring scheme is used to rank the compounds. A recent application on classifying PXR compounds using the consensus scheme has been detailed in Kortagere et al. Similar schemes are implemented to derive the best ranking compounds that bind to D₃ receptor. The docking program used provides some level receptor flexibility at the binding site. However, a complete induced fit model cannot be achieved using this level of screening as it is computationally expensive and may not well be required. As an alternative, another docking program called Glide, that has been well demonstrated to use induce fit concept on the final 25 best docked molecules instead of at the screening level, may be used. This ensures that (a) the best docked complexes are redocked and rescored; and (b) since no one docking or scoring program may efficiently capture the intricacies of the docking process, by using more than one docking program, it is ensured that the best ranked molecules that are short listed for experimental validation are screened efficiently.

Furthermore, to assess the selectivity of these best ranking compounds to the dopamine D₃ receptor, they are docked and scored against the other four dopamine receptors. Only those compounds that score the best against the dopamine D₃ receptor are used as lead molecules in the iterative HSB method. Preliminary results have validated the HSB protocol, as a novel compound, 4-(2-chlorophenyl)-butan-2-amine (FIG. 7), was identified as a D₃ receptor agonist that does NOT elicit the tolerance and SRT properties.

Evaluation of the Function, Selectivity and Cytotoxicity Profile of Novel D₃ Receptor Agonists in Cell-Based Assays

The lead compounds identified above are evaluated in cell based assays to determine their ability to activate D₃ receptors without inducing tolerance and SRT properties. The selectivity at various dopamine receptor subtypes are assessed by generating functional dose-response curves from AtT-20 neuroendocrine cell lines stably expressing the various dopamine receptor subtypes. The cytotoxicity is assessed by measuring the effect of lead compounds on proliferation and cell death of three different cell lines representing an endocrine, neuronal

and hepatic cell type. Each criterion that determines if a lead compound progresses through the preclinical development protocol is shown in FIG. 8.

Agonism of D₃ Dopamine Receptor:

To determine whether a compound is an agonist of the D₃ dopamine receptor the AtT-20 neuroendocrine cell line that stably expresses the human D₃ receptors is used. In the initial screen, the ability of 300 nM lead compound to activate G-protein coupled inward rectifier potassium (GIRK) channels (FIG. 1) is measured using whole cell voltage clamp recording. The protocol has been previously used for performing whole cell voltage clamp and measuring agonist-induced GIRK response. The acute ligand-induced GIRK currents are compared to that elicited by equivalent concentration of native agonist dopamine or full agonist quinpirole. The specificity of the response is determined by measuring GIRK response elicited by the lead compound in non-transfected AtT-20 cells and also in the presence of D₂-like dopamine receptor antagonist eticlopride and D₃ receptor selective antagonist GR218231 ((+)-(2R)-1,2,3,4-tetrahydro-6-[[4-methoxyphenyl)sulfonyl]methyl]-N,N-dipropyl-2-naphthalenamine). To be considered for further development, the lead compound should generate a GIRK response that is equal to or greater than that elicited by equivalent concentration (300 nM) of dopamine or quinpirole.

Abolishment of the D₃ Receptor Tolerance and SRT:

To determine whether a compound abolishes the D₃ receptor tolerance and SRT, the D₃ receptor-activated GIRK response to two consecutive treatments of the lead compound is measured. As illustrated by FIG. 9, the ratio of the second response to first response is close to 1 for agonists that abolish tolerance but is approximately 0.4 for agonists that elicit tolerance (for e.g. dopamine and quinpirole). In one embodiment, for further consideration, the lead compound should have a ratio of second to first GIRK response between 0.7 and 1.0.

In a second series of experiments, using the coupling to GIRK channels as the assay, the ability of the lead compound to compete with 100 nM, 300 nM and 1000 nM dopamine to block the development of tolerance and SRT at D₃ receptors is determined. As illustrated in FIG. 10, PBZI (which reduces AIM scores in the LID model) was able to compete effectively with dopamine in this assay.

Determination of the EC₅₀ of Lead Compound at Various D₂-Like Dopamine Receptors:

AtT-20 neuroendocrine cell lines stably expressing the human isoforms of D_{2S}, D_{2L}, D₃, D_{4.2} and D_{4.4} dopamine receptors are available. In addition, AtT-20 cells stably expressing D₁ dopamine receptors and 5HT-1A serotonin receptors are also available. Furthermore, non-transfected AtT-20 cells express endogenous somatostatin and muscarinic receptors.

As the initial step, membranes isolated from the cell lines stably expressing the various dopamine receptors are used to perform competitive radioligand binding assays. This procedure yields the K_i of the lead compound for the various dopamine receptor subtypes. As the next step, the various cell lines are used to obtain dose response curves in two functional assays.

In the first assay, whole cell voltage clamp recording is used to measure GIRK response to increasing concentrations of the lead compound (0.01 nM to 3,000 nM dose range). The EC₅₀ values for each receptor subtype is determined by fitting the data points using the Hill equation.

In the second functional assay, the ability of D₂-like dopamine receptors to inhibit adenylyl cyclase activity is assessed. The ability of the lead compound to dose-dependently decrease forskolin-induced cAMP levels is measured.

The IC₅₀ values for each receptor subtype is determined by fitting the data points using the Hill equation. Previously, the EC₅₀ and IC₅₀ values for dopamine and quinpirole has been determined in these cell lines in both functional assays. In one embodiment, to be advanced, the lead compound has to have an EC₅₀ or IC₅₀ value that is less than or equal to that of dopamine or quinpirole in either one of these assays. FIG. 11 illustrates examples of dose response curves for PBZI and 4-(2-chlorophenyl)-butan-2-amine at D₃ dopamine receptor.

Assessment of Cytotoxicity of the Lead Compound:

To determine the cytotoxicity of the lead compound, the cell proliferation labeling reagent WST-1 (2-(4-iodophenyl)-3-(4-nitrophenyl)-5-(2,4-disulfohenyl)-2H-tetrazolium, monosodium salt; Roche Applied Science) is used. The effect of the lead compound on the proliferation of three different cell lines (AtT-20 cells as a model for neuroendocrine cells, CAD cells as a model for catecholaminergic neuronal cells and HepG2 as a model for hepatic cells) is determined. Together, the three cell lines provide a preliminary screen for cytotoxicity.

The cells are treated with a wide concentration range of the lead compound (0.01 nM to 100 μM) for 24 hours and then incubated with WST-1 for an additional 4 hours and the absorbance values are determined using a spectrophotometer. The criterion for being selected in the cytotoxicity screen is that the lead compound should exhibit an EC₅₀ greater than 10 μM. As the lead compound development advances, the compound is also submitted to in vivo ADMET, toxicity and pharmacokinetic analysis.

Assessment of the Effect of Novel D₃ Receptor Agonists in Locomotor Behavior Assays

The compounds useful within the invention are evaluated for their ability to reduce AIMs scores in the rat intrastriatal 6-OHDA lesion model of LID.

Assessment of the Effect of the Lead Compound on Locomotor Activity in Balb/c Mice:

Tolerance- and non-tolerance inducing agonists that target dopamine D₃ receptors differentially alter locomotor activity. Specifically, tolerance-inducing compounds (e.g., PD128907, also known as (4aR,10bR)-3,4-a,4,10b-tetrahydro-4-propyl-2H,5H-[1]benzopyrano-[4,3-b]-1,4-oxazin-9-ol hydrochloride; quinpirole; and PBZI) induce biphasic locomotor effects, characterized by an initial inhibition and a subsequent stimulation of locomotion.

The preliminary results (Example 5) are consistent with studies that have demonstrated that administration of classical D₂/D₃ receptor agonists such as quinpirole, PD128907 or 7-hydroxy-N,N-dipropyl-2-aminotetralin (7OH-DPAT) elicited a biphasic locomotor response characterized by an initial inhibition and a subsequent stimulation of locomotion. The studies reported herein indicate that administration of D₃ receptor agonists such as PBZI that abolish tolerance and SRT to Balb/c mice only elicit hypoactivity. In contrast, PD128907 elicited tolerance and SRT at D₃ dopamine receptors and induces both hypoactivity and hyperactivity (FIG. 12). Since PBZI significantly reduced the AIM score in the rat LID model (FIG. 13), the agonist-induced locomotor activity assay may be used as a predictor of compounds that successfully reduce dyskinesia in the LID model.

In one embodiment, a goal of the study is to test this hypothesis as it may lead to development of a simple behavioral assay for screening lead compounds that might be beneficial for treating LID. Therefore, three different doses of the lead compound are administered and their effects on locomotor activity are determined. The dosage is determined based on the functional EC₅₀ values and binding affinity of the lead

compound from the cell-based assays described above. The compound is administered to Balb/c mice and locomotor activity is monitored using the TruScan Automated Behavioral Monitoring system (Coulbourn Instruments). In a non-limiting aspect, the lead compound that abolishes D₃ receptor tolerance and SRT only elicits hypoactivity in the locomotor assay, consistent with the PBZI results (FIG. 12).

Assessment of the Effect of the Lead Compound in the Rat Intrastratial 6-OHDA Lesion Model:

If the tolerance and SRT properties of the D₃ receptor contributes to the expression of LID in the intrastratial 6-hydroxydopamine (6-OHDA) lesioned rats, administration of non-tolerance-inducing D₃ receptor agonists, such as PBZI, prior to levodopa administration should reduce or abolish the symptoms associated with dyskinesia.

The unilateral intrastratial 6-OHDA lesioned rat model of Parkinson's Disease has been extensively characterized and widely used to test neuroprotective and transplantation strategies to treat Parkinson's Disease. The partial and slowly progressing degeneration of the nigral dopamine neurons and the concomitant development of motor deficits in this rat Parkinson's Disease model appear to mimic the progressive deterioration observed in patients suffering from Parkinson's Disease. Of particular interest is the development of LID in this animal model, which are similar to the abnormal involuntary movements (AIMs) seen clinically.

The intrastratial 6-OHDA rat Parkinson's Disease model described by Winkler et al., 2002 is herein used. In particular, rats in which unilateral intrastratial lesion is induced by injection of 6-OHDA (7 µg) at three different sites in the ventrolateral striatum are used. The coordinates for the three injection sites are, in mm: injection site 1: AP: +1.0; ML: -3.0; DV: -5.0; injection site 2: AP: -0.1; ML: -3.7; DV: -5.0; injection site 3: AP: -1.2; ML: -4.5; DV: -5.0. Anterior-posterior (AP) and mediolateral (ML) coordinates are from bregma. The dorso-ventral (DV) coordinates are from dura. The toothbar location is 0.

The intrastratial 6-OHDA lesioned and sham lesioned rats are obtained from the Custom Surgical Services division of Charles River Laboratories, Wilmington, Mass., USA. The commercial vendor tests the lesioned animals for amphetamine-induced rotation one week post lesion prior to shipping. A previous study has demonstrated that animals lesioned by the three-site injection exhibit a significant increase in amphetamine-induced rotation after one week. The lesioned animals exhibit bradykinesia/akinesia in the limb contralateral to the lesion. In the experiments, the ability of levodopa to improve bradykinesia/akinesia in the contralateral paw is first determined by performing the forelimb stepping test. The stepping test, subsequently, also provides a way to test if the lead compound interferes with the ability of levodopa to improve bradykinesia/akinesia. The effect of lesion on forelimb bradykinesia/akinesia is measured three weeks post-lesion using the stepping test. Once a significant deficit in stepping ability is established in the paw contralateral to the lesion, increasing doses of levodopa (2, 4 and 6 mg/kg) are administered along with 15 mg/kg benserazide (peripheral DOPA-decarboxylase inhibitor) and stepping ability assessed using the Stepping test, 60 minutes post injection. The dose of levodopa that is selected for the chronic administration phase is the dose that significantly improves the score in the stepping test (FIG. 14).

Seven and half weeks after the lesion, chronic levodopa treatment (6 mg/kg, single dose per day given with 15 mg/kg benserazide) is initiated. Stepping test and AIMs scores are determined on day 1, 8, 10, 17 and 24 after the initiation of chronic levodopa treatment. Upon chronic levodopa admin-

istration, the lesioned animals exhibit abnormal involuntary movements (AIMs) that include locomotor, axial, limb, and orolingual components. AIMs are scored using a rating scale based on frequency and severity of the dyskinetic symptoms.

It should be noted that the rank order of the various AIMs components in terms of frequency and severity in the rat model is: limb>orolingual=axial>>locomotor. This lesion model has been used to test the effect of PBZI on LID (Example 3).

To determine AIMs score, the rats are placed in a transparent glass cylinder and recorded using a digital video recorder. An experimentally blinded scorer observe for one minute at every 20th minute, from 20 minutes before to 180 minutes after the injection of levodopa. Levodopa-induced AIMs score is determined according to the rat dyskinesia scale. 6-OHDA lesioned animals develop LID approximately 10 days after initiation of the chronic levodopa treatment. The ability of PBZI to reduce the levodopa-induced AIMs score was determined by administering PBZI 10 minutes prior to levodopa administration on day 10, 17 and 24 after initiation of the chronic levodopa administration. The stepping test was also performed on these days to assess the effect of PBZI on the ability of levodopa to improve the stepping deficit in the lesioned animals. Two doses of the lead compound are tested. The dosage is determined by the affinity of the compound for the receptor and the results of the locomotor behavior assay described above.

The time course of the experiment is shown in FIG. 15. One week after the 3-site unilateral intrastratial 6-OHDA injection, the rats are subjected to the amphetamine-induced rotation test to identify lesioned animals. Three weeks post lesion, the animals are subjected to the stepping test to determine the pre-levodopa stepping deficit. Five weeks post lesion, increasing doses (2 to 6 mg/kg) of levodopa are administered once daily only on indicated days and stepping test performed. Seven and half weeks post lesion, levodopa (6 mg/kg) is administered once daily every day for 3.5 weeks. On day 1 and 8 after initiation of the chronic administration, stepping test and AIMs scores are determined following levodopa injection. On days 10, 17 and 24 stepping test and AIMs scores are determined following co-injection of levodopa and compound. At the end of 11-weeks, the animals are decapitated and brain tissue used for histological analysis: brain sections of animals are stained for tyrosine hydroxylase to assess loss of dopaminergic neurons in the lesioned hemisphere. Brain sections from some animals are assessed for D₂ and D₃ receptor binding and mRNA expression.

Combination Therapies

The compounds of the present invention are intended to be useful in the methods of present invention in combination with one or more additional compounds useful for treating Parkinson's Disease. These additional compounds may comprise compounds of the present invention or compounds, e.g., commercially available compounds, known to treat, prevent, or reduce the symptoms of Parkinson's Disease.

In non-limiting examples, the compounds of the invention may be used in combination with one or more of the following drugs: levodopa, clozapine, bromocriptine, pergolide, pramipexole, ropinirole, piribedil, cabergoline, apomorphine, and lisuride, a salt thereof and mixtures thereof.

A synergistic effect may be calculated, for example, using suitable methods such as, for example, the Sigmoid-E_{max} equation (Holford & Scheiner, 19981, Clin. Pharmacokinet 6: 429-453), the equation of Loewe additivity (Loewe & Muischnek, 1926, Arch. Exp. Pathol Pharmacol. 114: 313-326) and the median-effect equation (Chou & Talalay, 1984, Adv. Enzyme Regul. 22: 27-55). Each equation referred to

above may be applied to experimental data to generate a corresponding graph to aid in assessing the effects of the drug combination. The corresponding graphs associated with the equations referred to above are the concentration-effect curve, isobologram curve and combination index curve, respectively.

Administration/Dosage/Formulations

The regimen of administration may affect what constitutes an effective amount. The therapeutic formulations may be administered to the patient either prior to or after the onset of a LID. Further, several divided dosages, as well as staggered dosages may be administered daily or sequentially, or the dose may be continuously infused, or may be a bolus injection. Further, the dosages of the therapeutic formulations may be proportionally increased or decreased as indicated by the exigencies of the therapeutic or prophylactic situation.

Administration of the compositions of the present invention to a patient, preferably a mammal, more preferably a human, may be carried out using known procedures, at dosages and for periods of time effective to treat LID in the patient. An effective amount of the therapeutic compound necessary to achieve a therapeutic effect may vary according to factors such as the state of the disease or disorder in the patient; the age, sex, and weight of the patient; and the ability of the therapeutic compound to treat LID in the patient. Dosage regimens may be adjusted to provide the optimum therapeutic response. For example, several divided doses may be administered daily or the dose may be proportionally reduced as indicated by the exigencies of the therapeutic situation. A non-limiting example of an effective dose range for a therapeutic compound of the invention is from about 1 and 5,000 mg/kg of body weight/per day. One of ordinary skill in the art would be able to study the relevant factors and make the determination regarding the effective amount of the therapeutic compound without undue experimentation.

Actual dosage levels of the active ingredients in the pharmaceutical compositions of this invention may be varied so as to obtain an amount of the active ingredient that is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient.

In particular, the selected dosage level will depend upon a variety of factors including the activity of the particular compound employed, the time of administration, the rate of excretion of the compound, the duration of the treatment, other drugs, compounds or materials used in combination with the compound, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

A medical doctor, e.g., physician or veterinarian, having ordinary skill in the art may readily determine and prescribe the effective amount of the pharmaceutical composition required. For example, the physician or veterinarian could start doses of the compounds of the invention employed in the pharmaceutical composition at levels lower than that required in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved.

In particular embodiments, it is especially advantageous to formulate the compound in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the patients to be treated; each unit containing a predetermined quantity of therapeutic compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical vehicle. The dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the therapeutic compound and

the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding/formulating such a therapeutic compound for the treatment of LID in a patient.

In one embodiment, the compositions of the invention are formulated using one or more pharmaceutically acceptable excipients or carriers. In one embodiment, the pharmaceutical compositions of the invention comprise a therapeutically effective amount of a compound of the invention and a pharmaceutically acceptable carrier.

The carrier may be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils.

The proper fluidity may be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms may be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, sodium chloride, or polyalcohols such as mannitol and sorbitol, in the composition. Prolonged absorption of the injectable compositions may be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate or gelatin. In one embodiment, the pharmaceutically acceptable carrier is not DMSO alone.

In one embodiment, the compositions of the invention are administered to the patient in dosages that range from one to five times per day or more. In another embodiment, the compositions of the invention are administered to the patient in range of dosages that include, but are not limited to, once every day, every two, days, every three days to once a week, and once every two weeks. It will be readily apparent to one skilled in the art that the frequency of administration of the various combination compositions of the invention will vary from individual to individual depending on many factors including, but not limited to, age, disease or disorder to be treated, gender, overall health, and other factors. Thus, the invention should not be construed to be limited to any particular dosage regime and the precise dosage and composition to be administered to any patient will be determined by the attending physical taking all other factors about the patient into account.

Compounds of the invention for administration may be in the range of from about 1 μ g to about 10,000 mg, about 20 μ g to about 9,500 mg, about 40 μ g to about 9,000 mg, about 75 μ g to about 8,500 mg, about 150 μ g to about 7,500 mg, about 200 μ g to about 7,000 mg, about 3050 μ g to about 6,000 mg, about 500 μ g to about 5,000 mg, about 750 μ g to about 4,000 mg, about 1 mg to about 3,000 mg, about 10 mg to about 2,500 mg, about 20 mg to about 2,000 mg, about 25 mg to about 1,500 mg, about 50 mg to about 1,000 mg, about 75 mg to about 900 mg, about 100 mg to about 800 mg, about 250 mg to about 750 mg, about 300 mg to about 600 mg, about 400 mg to about 500 mg, and any and all whole or partial increments therebetween.

In some embodiments, the dose of a compound of the invention is from about 1 mg and about 2,500 mg. In some embodiments, a dose of a compound of the invention used in compositions described herein is less than about 10,000 mg, or less than about 8,000 mg, or less than about 6,000 mg, or less than about 5,000 mg, or less than about 3,000 mg, or less than about 2,000 mg, or less than about 1,000 mg, or less than about 500 mg, or less than about 200 mg, or less than about 50 mg. Similarly, in some embodiments, a dose of a second

compound (i.e., a drug used for treating Parkinson's Disease) as described herein is less than about 1,000 mg, or less than about 800 mg, or less than about 600 mg, or less than about 500 mg, or less than about 400 mg, or less than about 300 mg, or less than about 200 mg, or less than about 100 mg, or less than about 50 mg, or less than about 40 mg, or less than about 30 mg, or less than about 25 mg, or less than about 20 mg, or less than about 15 mg, or less than about 10 mg, or less than about 5 mg, or less than about 2 mg, or less than about 1 mg, or less than about 0.5 mg, and any and all whole or partial increments thereof.

In one embodiment, the present invention is directed to a packaged pharmaceutical composition comprising a container holding a therapeutically effective amount of a compound of the invention, alone or in combination with a second pharmaceutical agent; and instructions for using the compound to treat, prevent, or reduce one or more symptoms of LID in a patient.

Formulations may be employed in admixtures with conventional excipients, i.e., pharmaceutically acceptable organic or inorganic carrier substances suitable for oral, parenteral, nasal, intravenous, subcutaneous, enteral, or any other suitable mode of administration, known to the art. The pharmaceutical preparations may be sterilized and if desired mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure buffers, coloring, flavoring and/or aromatic substances and the like. They may also be combined where desired with other active agents, e.g., other analgesic agents.

The term "container" includes any receptacle for holding the pharmaceutical composition. For example, in one embodiment, the container is the packaging that contains the pharmaceutical composition. In other embodiments, the container is not the packaging that contains the pharmaceutical composition, i.e., the container is a receptacle, such as a box or vial that contains the packaged pharmaceutical composition or unpackaged pharmaceutical composition and the instructions for use of the pharmaceutical composition. Moreover, packaging techniques are well known in the art. It should be understood that the instructions for use of the pharmaceutical composition may be contained on the packaging containing the pharmaceutical composition, and as such the instructions form an increased functional relationship to the packaged product. However, it should be understood that the instructions may contain information pertaining to the compound's ability to perform its intended function, e.g., treating, preventing, or reducing a LID in a patient.

Routes of administration of any of the compositions of the invention include oral, nasal, rectal, intravaginal, parenteral, buccal, sublingual or topical. The compounds for use in the invention may be formulated for administration by any suitable route, such as for oral or parenteral, for example, transdermal, transmucosal (e.g., sublingual, lingual, (trans)buccal, (trans)urethral, vaginal (e.g., trans- and perivaginally), (intra) nasal and (trans)rectal), intravesical, intrapulmonary, intraduodenal, intragastrical, intrathecal, subcutaneous, intramuscular, intradermal, intra-arterial, intravenous, intra-bronchial, inhalation, and topical administration.

Suitable compositions and dosage forms include, for example, tablets, capsules, caplets, pills, gel caps, troches, dispersions, suspensions, solutions, syrups, granules, beads, transdermal patches, gels, powders, pellets, magmas, lozenges, creams, pastes, plasters, lotions, discs, suppositories, liquid sprays for nasal or oral administration, dry powder or aerosolized formulations for inhalation, compositions and formulations for intravesical administration and the like. It

should be understood that the formulations and compositions that would be useful in the present invention are not limited to the particular formulations and compositions that are described herein.

5 Oral Administration

For oral application, particularly suitable are tablets, dragees, liquids, drops, suppositories, or capsules, caplets and gelcaps. The compositions intended for oral use may be prepared according to any method known in the art and such compositions may contain one or more agents selected from the group consisting of inert, non-toxic pharmaceutically excipients which are suitable for the manufacture of tablets. Such excipients include, for example an inert diluent such as lactose; granulating and disintegrating agents such as cornstarch; binding agents such as starch; and lubricating agents such as magnesium stearate. The tablets may be uncoated or they may be coated by known techniques for elegance or to delay the release of the active ingredients. Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert diluent.

For oral administration, the compounds of the invention may be in the form of tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g., polyvinylpyrrolidone, hydroxypropylcellulose or hydroxypropylmethylcellulose); fillers (e.g., cornstarch, lactose, microcrystalline cellulose or calcium phosphate); lubricants (e.g., magnesium stearate, talc, or silica); disintegrates (e.g., sodium starch glycollate); or wetting agents (e.g., sodium lauryl sulphate). If desired, the tablets may be coated using suitable methods and coating materials such as OPADRY™ film coating systems available from Colorcon, West Point, Pa. (e.g., OPADRY™ OY Type, OYC Type, Organic Enteric OY-P Type, Aqueous Enteric OY-A Type, OY-PM Type and OPADRY™ White, 32K18400). Liquid preparation for oral administration may be in the form of solutions, syrups or suspensions. The liquid preparations may be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol syrup, methyl cellulose or hydrogenated edible fats); emulsifying agent (e.g., lecithin or acacia); non-aqueous vehicles (e.g., almond oil, oily esters or ethyl alcohol); and preservatives (e.g., methyl or propyl p-hydroxy benzoates or sorbic acid).

Granulating techniques are well known in the pharmaceutical art for modifying starting powders or other particulate materials of an active ingredient. The powders are typically mixed with a binder material into larger permanent free-flowing agglomerates or granules referred to as a "granulation." For example, solvent-using "wet" granulation processes are generally characterized in that the powders are combined with a binder material and moistened with water or an organic solvent under conditions resulting in the formation of a wet granulated mass from which the solvent must then be evaporated.

Melt granulation generally consists in the use of materials that are solid or semi-solid at room temperature (i.e. having a relatively low softening or melting point range) to promote granulation of powdered or other materials, essentially in the absence of added water or other liquid solvents. The low melting solids, when heated to a temperature in the melting point range, liquefy to act as a binder or granulating medium. The liquefied solid spreads itself over the surface of powdered materials with which it is contacted, and on cooling, forms a solid granulated mass in which the initial materials are bound together. The resulting melt granulation may then be provided to a tablet press or be encapsulated for preparing the oral

dosage form. Melt granulation improves the dissolution rate and bioavailability of an active (i.e. drug) by forming a solid dispersion or solid solution.

U.S. Pat. No. 5,169,645 discloses directly compressible wax-containing granules having improved flow properties. The granules are obtained when waxes are admixed in the melt with certain flow improving additives, followed by cooling and granulation of the admixture. In certain embodiments, only the wax itself melts in the melt combination of the wax(es) and additives(s), and in other cases both the wax(es) and the additives(s) will melt.

The present invention also includes a multi-layer tablet comprising a layer providing for the delayed release of one or more compounds of the invention, and a further layer providing for the immediate release of a medication for treatment of Parkinson's Disease. Using a wax/pH-sensitive polymer mix, a gastric insoluble composition may be obtained in which the active ingredient is entrapped, ensuring its delayed release.

Parenteral Administration

For parenteral administration, the compounds of the invention may be formulated for injection or infusion, for example, intravenous, intramuscular or subcutaneous injection or infusion, or for administration in a bolus dose and/or continuous infusion. Suspensions, solutions or emulsions in an oily or aqueous vehicle, optionally containing other formulatory agents such as suspending, stabilizing and/or dispersing agents may be used.

Additional Administration Forms

Additional dosage forms of this invention include dosage forms as described in U.S. Pat. Nos. 6,340,475, 6,488,962, 6,451,808, 5,972,389, 5,582,837, and 5,007,790. Additional dosage forms of this invention also include dosage forms as described in U.S. Patent Applications Nos. 20030147952, 20030104062, 20030104053, 20030044466, 20030039688, and 20020051820. Additional dosage forms of this invention also include dosage forms as described in PCT Applications Nos. WO 03/35041, WO 03/35040, WO 03/35029, WO 03/35177, WO 03/35039, WO 02/96404, WO 02/32416, WO 01/97783, WO 01/56544, WO 01/32217, WO 98/55107, WO 98/11879, WO 97/47285, WO 93/18755, and WO 90/11757.

Controlled Release Formulations and Drug Delivery Systems
In certain embodiments, the formulations of the present invention may be, but are not limited to, short-term, rapid-offset, as well as controlled, for example, sustained release, delayed release and pulsatile release formulations.

The term sustained release is used in its conventional sense to refer to a drug formulation that provides for gradual release of a drug over an extended period of time, and that may, although not necessarily, result in substantially constant blood levels of a drug over an extended time period. The period of time may be as long as a month or more and should be a release which is longer than the same amount of agent administered in bolus form.

For sustained release, the compounds may be formulated with a suitable polymer or hydrophobic material which provides sustained release properties to the compounds. As such, the compounds for use the method of the invention may be administered in the form of microparticles, for example, by injection or in the form of wafers or discs by implantation.

In a preferred embodiment of the invention, the compounds of the invention are administered to a patient, alone or in combination with another pharmaceutical agent, using a sustained release formulation.

The term delayed release is used herein in its conventional sense to refer to a drug formulation that provides for an initial release of the drug after some delay following drug adminis-

tration and that may, although not necessarily, include a delay of from about 10 minutes up to about 12 hours.

The term pulsatile release is used herein in its conventional sense to refer to a drug formulation that provides release of the drug in such a way as to produce pulsed plasma profiles of the drug after drug administration.

The term immediate release is used in its conventional sense to refer to a drug formulation that provides for release of the drug immediately after drug administration.

As used herein, short-term refers to any period of time up to and including about 8 hours, about 7 hours, about 6 hours, about 5 hours, about 4 hours, about 3 hours, about 2 hours, about 1 hour, about 40 minutes, about 20 minutes, or about 10 minutes and any or all whole or partial increments thereof after drug administration after drug administration.

As used herein, rapid-offset refers to any period of time up to and including about 8 hours, about 7 hours, about 6 hours, about 5 hours, about 4 hours, about 3 hours, about 2 hours, about 1 hour, about 40 minutes, about 20 minutes, or about 10 minutes, and any and all whole or partial increments thereof after drug administration.

Dosing

The therapeutically effective amount or dose of a compound of the present invention will depend on the age, sex and weight of the patient, the current medical condition of the patient and the progression of Parkinson's Disease in the patient being treated. The skilled artisan will be able to determine appropriate dosages depending on these and other factors.

A suitable dose of a compound of the present invention may be in the range of from about 0.01 mg to about 5,000 mg per day, such as from about 0.1 mg to about 1,000 mg, for example, from about 1 mg to about 500 mg, such as about 5 mg to about 250 mg per day. The dose may be administered in a single dosage or in multiple dosages, for example from 1 to 4 or more times per day. When multiple dosages are used, the amount of each dosage may be the same or different. For example, a dose of 1 mg per day may be administered as two 0.5 mg doses, with about a 12-hour interval between doses.

It is understood that the amount of compound dosed per day may be administered, in non-limiting examples, every day, every other day, every 2 days, every 3 days, every 4 days, or every 5 days. For example, with every other day administration, a 5 mg per day dose may be initiated on Monday with a first subsequent 5 mg per day dose administered on Wednesday, a second subsequent 5 mg per day dose administered on Friday, and so on.

The compounds for use in the method of the invention may be formulated in unit dosage form. The term "unit dosage form" refers to physically discrete units suitable as unitary dosage for patients undergoing treatment, with each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, optionally in association with a suitable pharmaceutical carrier. The unit dosage form may be for a single daily dose or one of multiple daily doses (e.g., about 1 to 4 or more times per day). When multiple daily doses are used, the unit dosage form may be the same or different for each dose.

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, numerous equivalents to the specific procedures, embodiments, claims, and examples described herein. Such equivalents were considered to be within the scope of this invention and covered by the claims appended hereto. For example, it should be understood, that modifications in reaction conditions, including but not limited to reaction times, reaction size/volume, and experimental reagents, such as solvents, catalysts, pressures,

atmospheric conditions, e.g., nitrogen atmosphere, and reducing/oxidizing agents, with art-recognized alternatives and using no more than routine experimentation, are within the scope of the present application. It is to be understood that wherever values and ranges are provided herein, all values and ranges encompassed by these values and ranges, are meant to be encompassed within the scope of the present invention. Moreover, all values that fall within these ranges, as well as the upper or lower limits of a range of values, are also contemplated by the present application.

The following examples further illustrate aspects of the present invention. However, they are in no way a limitation of the teachings or disclosure of the present invention as set forth herein.

EXAMPLES

The invention is now described with reference to the following Examples. These Examples are provided for the purpose of illustration only, and the invention is not limited to these Examples, but rather encompasses all variations that are evident as a result of the teachings provided herein.

Materials:

Unless otherwise noted, all starting materials and resins were obtained from commercial suppliers and used without purification.

Quinpirole (Sigma-Aldrich, St. Louis, Mo.), PD128907 (Tocris, Ellisville, Mo.), 70H-DPAT (Tocris), pramipexole (Tocris), rotigotine (Tocris), 4-(2-chlorophenyl)-butan-2-amine (ES609; Asinex, Moscow, Russia) and cis-8-Hydroxy-3-(n-propyl)-1,2,3a,4,5,9b-hexahydro-1H-benz[e]indole hydrobromide (PBZI; Sigma-Aldrich) were dissolved in water and used at indicated concentrations. Sarizotan (Merck KGaA, Gibbstown, N.J.), FAUC73 (Sigma-Aldrich) and 70H-PIPAT (Tocris) were dissolved in DMSO. A 10 mM stock of dopamine (Sigma-Aldrich) was freshly dissolved in 100 mM ascorbic acid and used at a final concentration of 100 nM.

Cell Culture:

AtT-20 mouse pituitary cells were grown in Ham's F10 medium with 5% FBS, 10% heat-inactivated horse serum, 2 mM glutamine and 50 µg/ml gentamicin (Invitrogen, Carlsbad, Calif.). AtT-20 cells stably expressing the human D_{2S}, D_{2L}, D₃, and D_{4.2} and receptor were maintained in the above F10 culture media supplemented with 500 µg/ml G418 (Invitrogen). For electrophysiological characterization, cells were plated onto glass coverslips coated with 40 µg/ml poly L-lysine (Sigma-Aldrich). The generation and characterization of the AtT-20 cells stably expressing various human dopamine receptors have been previously reported (Kuzhikandathil & Oxford, 2000, *J. Gen. Physiol.* 115:697-706; Kuzhikandathil et al., 1998, *Mol. Cell. Neurosci.* 12:390-402; Kuzhikandathil & Bartoszyk, 2006, *Neuropharm.* 51:873-884; Westrich & Kuzhikandathil, 2007, *Biochim. Biophys. Acta-MCR* 1773:1747-1758; Kuzhikandathil et al., 2004, *Mol. Cell. Neurosci.* 26:144-155; Westrich et al., 2010, *Biochem. Pharmacol.* 79:897-907).

Measurement of cAMP:

Cyclic AMP (cAMP) levels were assessed using the cAMP Biotrak Enzymeimmunoassay (EIA) kit (GE Healthcare, Piscataway, N.J., USA) as described previously (Kuzhikandathil & Bartoszyk, 2006, *Neuropharm.* 51:873-884). The cAMP levels in each treated sample were assayed in triplicate and the entire experiment repeated three independent times.

Electrophysiology:

Agonist-activated currents were measured by the whole-cell patch clamp technique in voltage clamp and current clamp mode as described previously (Kuzhikandathil & Oxford, 2000, *J. Gen. Physiol.* 115:697-706; Kuzhikandathil et al., 1998, *Mol. Cell. Neurosci.* 12:390-402; Kuzhikandathil & Bartoszyk, 2006, *Neuropharm.* 51:873-884; Westrich & Kuzhikandathil, 2007, *Biochim. Biophys. Acta-MCR* 1773:1747-1758; Kuzhikandathil et al., 2004, *Mol. Cell. Neurosci.* 26:144-155; Westrich et al., 2010, *Biochem. Pharmacol.* 79:897-907). Drug solutions were delivered to cells via a multi-barreled micropipette array. The current responses were normalized to the cell capacitance, to account for variation in cell size.

Statistics:

Analysis of variance (ANOVA) and the Holm-Sidak multiple pair-wise comparison tests and Student's t-test was performed with the SigmaPlot® 11 software (SPSS Inc.). Data were considered statistically significant when the probability value (P) was less than 0.05.

Computational Modeling (HSB method):

The HSB protocol for designing small molecule inhibitors to G-protein coupled receptors has been described (Kortagere & Welsh, 2006, *J. Comput. Aided Mol. Des.* 20(12):789-8026). Briefly, the method involved creating a focused library of small molecules derived from commercial vendors. All molecules in the database were converted to the UNITY format and screened using UNITY module integrated in SYBYL (SYBYL 8.0, Tripos International).

To create a 3D pharmacophore based on the interactions of PBZI in D₃ receptor binding site, a homology model of D₃ receptor was created using the crystal structure of the beta adrenergic (β₂-AR) receptor in complex with a partial inverse agonist (PDB code: 2RH1) (Cherezov et al., 2007, *Science* 318(5854):1258-1265) as a template with the homology modeling program Modeller (ver 9.4) (Sali et al., 1993, *J. Mol. Biol.* 234(3):779-815; Westrich et al., 2010, *Biochem. Pharmacol.* 79:897-907; Kortagere et al., 2011, *Biochem. Pharmacol.* 81(1):157-163). PBZI has a fairly rigid conformation owing to its tricyclic structure. A conformational analysis using stochastic search method adopted in MOE was utilized to obtain the best conformation of the alkyl tail region extending from the amine group. All the conformations obtained were clustered based on energy and a representative member from the most populated cluster was chosen for further optimization using the AM1 semi-empirical quantum chemical method adopted in MOE. A similar procedure was used for other ligands in this study. The optimized conformations were used for further docking experiments. PBZI was docked to the 3D structure of the refined D₃ structure using docking program GOLD (ver 4.1) (Jones et al., 1995, *J. Mol. Biol.* 245(1):43-53). Twenty independent runs were performed and the docked complexes were scored using Goldscore (Jones et al., 1995, *J. Mol. Biol.* 245(1):43-53), chemscore (Eldridge et al., 1997, *J. Comput. Aided Mol. Des.* 11(5):425-445) and customized scoring scheme (Kortagere & Welsh, 2006, *J. Comput. Aided Mol. Des.* 20(12):789-802).

To obtain a more realistic conformation of the agonist bound model, the model was immersed into an explicit water-POPC lipid bilayer-water model membrane system using the Desmond module (Kevin et al., 2006, Proceedings of the ACM/IEEE Conference on Supercomputing (SC06), New York, N.Y., IEEE) of the Schrodinger suite program. The model membrane was pre-aligned to the β₂-AR crystal structure to adopt its orientation to the D₃ receptor-PBZI model. Default conditions for bilayer composition including those of

Na⁺ and Cl⁻ ions were selected and the entire simulation was performed using the default all atom OPLS force field. A four step protocol consisting of routines for pre-relaxation, minimization, heating, equilibration and production run of 3 ns was followed to completely refine the model (Chien et al., 2010, Science 330(6007):1091-1115). Throughout the simulations, the interactions of PBZI with residues from TM3 and TM5 were maintained using low levels of constraints. However all constraints were removed during the production run to completely relax the ligand in the protein environment. Key interactions between PBZI and D₃ receptor, namely salt bridge with Asp110, hydrogen bonded interaction with Ser192, aromatic ring interactions with His349, hydrophobic interactions with Val111 and other aromatic residues from TM6 were used to build a four point hybrid pharmacophore as shown in FIG. 6C. Electronic libraries of vendor-available small molecules were screened to identify hits that correspond to the pharmacophore.

Finally, the hits were filtered using Lipinski's Rule of 5 (Lipinski et al., 2001, Adv. Drug Deliv. Rev. 46(1-3):3-26), blood brain barrier (BBB) penetration (Kortagere et al., 2008, Pharm. Res. 25(8):1836-1845) and off-target screening against Pregnane xenobiotic receptors (Kortagere et al., 2009, Pharm. Res. 26(4):1001-1011) and hERG channels (Chekmarev et al., 2008, Chem. Res. Toxicol. 21(6):1304-1314). The resulting 290 hits were docked to the binding site of D₃ receptor using GOLD program and scored using a variety of scoring functions as described previously for docking of PBZI. Fifteen best ranking hits were obtained from the vendors and functionally evaluated.

Example 1

In Vitro Characterization of Compounds

PBZI and 4-(2-chlorophenyl)-butan-2-amine were characterized for their interactions with dopamine receptors. Both compounds activated G-protein coupled inward rectifier potassium (GIRK) channels via D₃ dopamine receptors (FIG. 2) and inhibited adenylyl cyclase activity (FIG. 11). The in vitro functional data for both compounds are shown in Table 1. Of most interest was the observation that unlike traditional D₂/D₃ receptor agonists, PBZI and 4-(2-chlorophenyl)-butan-2-amine abolished the tolerance and SRT properties of the D₃R (FIG. 2). These two compounds represent a new class of D₃ dopamine receptor agonists that abolishes the tolerance and SRT properties of the D₃ receptor while being a full agonist.

TABLE 1

Characterization of PBZI & 4-(2-chlorophenyl)-butan-2-amine effect on D ₂ -like receptor signaling function in AtT-20 stable cell lines				
	D _{2S} dopamine receptor	D _{2L} dopamine receptor	D ₃ dopamine receptor	D _{4.2} dopamine receptor
PBZI				
Adenylyl cyclase inhibition	partial agonist (EC ₅₀ = 140 nM)	full agonist (EC ₅₀ = 67 nM)	full agonist (EC ₅₀ = 35 nM)	no response

TABLE 1-continued

Characterization of PBZI & 4-(2-chlorophenyl)-butan-2-amine effect on D ₂ -like receptor signaling function in AtT-20 stable cell lines				
	D _{2S} dopamine receptor	D _{2L} dopamine receptor	D ₃ dopamine receptor	D _{4.2} dopamine receptor
GIRK channel activation	partial agonist (EC ₅₀ = 53 nM)	full agonist (EC ₅₀ ND)	full agonist (EC ₅₀ = 30 nM)	partial agonist (EC ₅₀ ND)
4-(2-chloro-phenyl)-butan-2-amine (ES0609)				
Adenylyl cyclase inhibition	Partial agonist (EC ₅₀ = 2.4 nM)	No response	Full agonist (EC ₅₀ = 0.15 nM)	Partial agonist (EC ₅₀ = 0.82 nM)
GIRK channel activation	Partial agonist (EC ₅₀ -ND)	Partial agonist (EC ₅₀ -ND)	Full agonist (EC ₅₀ = 30 nM)	no response

ND: not determined

Example 2

Ligand Dependent D₃ Receptor Tolerance and SRT Properties

FIG. 9A illustrates the D₃ dopamine receptor-induced activation of native GIRK channels when the receptor is stimulated with the endogenous agonist, dopamine. The tolerance property of D₃ receptor is quantified as the ratio of 2nd to 1st agonist-induced response. The D₃ receptor tolerance property is also observed in the D₃ receptor-adenylyl cyclase and -mitogen activated protein kinase pathways (Kuzhikandathil & Bartoszyk, 2006, Neuropharm. 51:873-884; Westrich & Kuzhikandathil, 2007, Biochim. Biophys. Acta-MCR 1773: 1747-1758). In addition, FIG. 9A shows the D₃ receptor SRT property, which is the delayed termination of the agonist-induced response after agonist removal. Tolerant D₃ receptor adopts a distinct conformation (Westrich et al., 2010, Biochem. Pharmacol. 79:897-907), suggesting that tolerance and SRT properties might be modulated by functionally-selective agonists that alter this distinct conformational state. To identify agonists that might modulate D₃ receptor tolerance and SRT properties, ten distinct agonists were screened for their ability to induce tolerance and SRT, using the D₃ receptor-GIRK channel signaling pathway as an assay. The selected agonists included the endogenous ligand dopamine, ligands that exhibited selectivity for D₃, and compounds used clinically to treat Parkinson's disease. The results in FIG. 9C suggest that, while most agonists induced tolerance, two agonists, cis-80H-PBZI and FAUC73, abolished the tolerance property. Interestingly, PD128907 induced enhanced tolerance that was significantly different than the other agonists.

FIGS. 2A and 2C illustrate representative voltage clamp recordings of AtT-20 cells stably expressing the human D₃ dopamine receptor and treated with agonists that either induce tolerance or do not induce tolerance. PBZI (FIG. 2A) and FAUC73 (FIG. 2C) abolished both tolerance and SRT properties; however in the same cell, quinpirole (FIG. 2A) or PD128907 (FIG. 2C) induced severe tolerance and SRT. Control experiments in parental AtT-20 cells as well as pretreatment with D₂/D₃ antagonist, eticlopride (100 nM), showed that the agonistic effect of PBZI and FAUC73 are specific for

47

D₃ receptors. Furthermore, the ability to abolish tolerance and SRT was not concentration dependent-PBZI tested at doses from 100 nM to 10 μM did not induce tolerance. Together, these results suggest that the D₃ receptor tolerance and SRT properties are ligand-dependent. Of the two tolerance- and SRT-abolishing compounds, PBZI is water soluble and more extensively characterized, in vitro and in vivo.

Example 3

In Vivo Characterization of Compounds: In Vivo Behavioral Response

To determine the in vivo behavioral effect of the novel class of D₃ dopamine receptor agonists, the effects of PBZI and a classical tolerance and SRT inducing D₃ dopamine receptor agonist, PD128907 ((4aR,10bR)-3,4-a,4,10b-tetrahydro-4-propyl-2H,5H-[1]benzopyrano-[4,3-b]-1,4-oxazin-9-ol hydrochloride), were compared. FIG. 12 illustrates that PD128907 elicited a biphasic locomotor response in a novel open field activity test. In contrast, PBZI (which does not elicit tolerance and SRT) induced only a monophasic locomotor response. This novel result thus identified a potential behavioral screen for the new class of D₃ dopamine receptor agonist described herein.

Example 4

In Vivo Characterization of Compounds: Intrastratial 6-OHDA Rat Parkinson's Disease Model

In order to further evaluate the therapeutic potential of the novel class of D₃ receptor agonists, the hypothesis that up regulation of D₃ receptor in the striatum contributes to the development of levodopa-induced dyskinesia (LID) in Parkinson's disease was tested. The unilateral intrastratial 6-OHDA lesioned rat model of Parkinson's Disease has been extensively characterized and widely used to test neuroprotective and transplantation strategies to treat Parkinson's Disease (Winkler et al., 2002, Neurobiol. Dis. 10(2):165-86). The partial and slowly progressing degeneration of the nigral dopamine neurons and the concomitant development of motor deficits in this rat Parkinson's Disease model appeared to mimic the progressive deterioration observed in patients suffering from Parkinson's Disease (Winkler et al., 2002, Neurobiol. Dis. 10(2):165-86). Of particular interest is the development of LID in this animal model, wherein the movements are similar to the abnormal involuntary movements (AIMs) seen clinically.

The intrastratial 6-OHDA rat Parkinson's Disease model described by Winkler and co-workers was used to test the effect of PBZI and ES0609 on LID (Winkler et al., 2002, Neurobiol. Dis. 10(2):165-86). To determine AIMs score, the rats were placed in transparent glass cylinder and recorded using a digital video recorder. 6-OHDA lesioned animals develop LID approximately 10 days after initiation of the chronic levodopa treatment. The ability of PBZI or 4-(2-chlorophenyl)-butan-2-amine to reduce the levodopa-induced AIMs score was determined by administering the novel D₃ receptor agonists 10 minutes prior to levodopa administration on day 10, 17 and 24 after initiation of the chronic levodopa administration. The stepping test was also performed on these days to assess the effect of these new agonists on the ability of levodopa to improve the stepping deficit in the lesioned animals.

48

Example 5

In Vivo Characterization of Compounds: Reduction of Dyskinesia

The ability of a tolerance inducing (PD128907) and the non-tolerance inducing D₃ receptor agonists (PBZI or 4-(2-chlorophenyl)-butan-2-amine) to reduce dyskinesia associated with chronic levodopa treatment was tested in the rat model. Control (saline), PD128907 (0.1 mg/kg, sc), PBZI (17 mg/kg, sc) or 4-(2-chlorophenyl)-butan-2-amine (20 mg/kg) were administered ten minutes before levodopa treatment. FIG. 13A illustrates the effect of control, PD128907 (a tolerance inducing D₂/D₃ agonist) and PBZI (a non tolerance inducing D₃/D₃ agonist) on the total AIMs score (includes locomotor, axial, limb and orolingual) over time of a day 17 (chronic L-DOPA phase) lesioned animal. Interestingly, the administration of PD128907 to the lesioned animals in of itself induced dyskinesia, even before levodopa administration. This is consistent with clinical reports that LID may also be elicited by dopamine receptor agonists used in the treatment of Parkinson's Disease.

The results in FIG. 13A suggest that PBZI itself did not induce dyskinesia, and delayed the onset of levodopa induced dyskinesia. The cumulative data obtained by calculating the area under the curve in FIG. 13B illustrates that PBZI significantly reduces (by ~300%) the total integrated AIMs score. Stepping test data obtained during co-administration of PBZI and levodopa suggest that PBZI did not attenuate the beneficial effect of levodopa on bradykinesia/akinesia. These results suggest that PBZI and its analogs are promising candidates for preclinical development as new therapeutics for the treatment of LID in Parkinson's disease.

Example 6

In Vivo Characterization of Compounds: Locomotor Behavior in Mice

The effects of PBZI on locomotor behavior in Balb/c mice were evaluated. The specific objectives of this experiment were: (a) to determine whether PBZI would induce a decrease in locomotion in Balb/c mice; (b) to evaluate PBZI's effects over a longer test session; and (c) to determine if PD128907 elicited a biphasic locomotor effect in Balb/c mice.

A total of 12 young adult, male Balb/c mice were obtained from Charles River, Inc. (Wilmington, Mass.), and housed in groups of four in standard cages. The animals were maintained on a 12 hour light/12 hour dark cycle and permitted ad lib access to food and water. Experiments were conducted during the animal's light phase. The strain of mouse was selected based on the previous work evaluating the relationship between dopaminergic compounds and motor activity in Balb/c mice. The mice received a single injection of PBZI (0, 1, or 10 mg/kg, sc), and immediately thereafter were individually placed into a test arena (TruScan arena, Coulbourn Instruments Inc.) for 2-hr in normal illumination. For PD128907, separate groups of Balb/c mice received subcutaneous injections of saline (n=4) or 0.05 or 0.4 mg/kg PD128907 (n=4, for each). Locomotion ('distance traveled') scores of drug-injected animals were collapsed across 10-min intervals and for comparison purposes were normalized to saline injected animals (FIG. 12A). The experiment was conducted on three separate occasions. Locomotion varied significantly as a function of a Dose X Time interaction (F(2,4)=5.40, p=0.002). Newman-Keuls multiple (α=0.05) of the main effects comprising this interaction revealed that

locomotion was significantly reduced (by approximately 90%) within 20-min in mice receiving 10 mg/kg of PBZI compared with mice in the other groups. Of further importance, there were no between-group differences during the other time intervals. This effect persisted for 40-min post-injection in Balb/c mice.

On the contrary, PD128907, as reported previously for other strains of mice, elicited a biphasic effect in Balb/c mice. A hypoactivity was observed in the first 40-min post-injection followed by hyperactivity during the 70 to 110 minute post-injection period (FIG. 12B). The biphasic effect of PD128907 was observed at a dose of 0.05 mg/kg or higher. While the lower dose of PD128907 (0.05 mg/kg) elicited hypoactivity of a shorter duration (20 minutes), it elicited the same duration and magnitude of hyperactivity. The difference in dose required to elicit hypoactivity is likely due to the differences in D_3 dopamine receptor binding affinity for PBZI ($K_i=22$ nM) and PD128907 ($K_i=2.3$ nM). With both PBZI and PD128907, the locomotor activity returned to control saline levels by 120 minutes post-injection. These results clearly demonstrate that agonists that induce D_3 receptor tolerance (PD128907) elicit a biphasic locomotor response. However, agonists that do not induce D_3 receptor tolerance (PBZI) elicit a monophasic hypoactive locomotor response.

Example 7

Effect of D_3 Receptor Signaling Properties on Cellular Function and Signaling Pathways

The stably-expressed D_3 receptor couples to endogenous GIRK channels in AtT-20 neuroendocrine cells and modulates spontaneous action potentials and secretion (Kuzhikandathil & Oxford, 1999, *J. Neurosci.* 19(5):1698-1707; Kuzhikandathil & Oxford, 2000, *J. Gen. Physiol.* 115:697-706; Kuzhikandathil et al., 1998, *Mol. Cell. Neurosci.* 12:390-402). Activation of the stably expressed human D_3 dopamine receptor by dopamine inhibited spontaneous action potentials during the first application, but not upon subsequent applications (FIG. 16A). In contrast, the activation of the D_3 receptor by the tolerance and SRT abolishing agonist, PBZI, inhibited spontaneous action potentials during the first application and subsequent applications (FIG. 16B). This result suggests that the modulation of neuronal firing by D_3 receptor agonists that induce tolerance is very different from those that do not induce tolerance. The latter class of agonists converts the D_3 receptor to the functional equivalent of a D_2 receptor.

Example 8

Identification of a Novel Agonist that does not Induce Tolerance and SRT By Using a D_3 Receptor-PBZI Pharmacophore Model

The results in FIGS. 9C, 2A and 2C showed that PD128907 induces severe tolerance and SRT; interestingly, PBZI, which does not induce tolerance and SRT shares a few core structural elements with PD128907 (FIG. 6). The stark difference in the ability of PD128907 and PBZI to induce tolerance and SRT suggested that comparative modeling studies of these compounds docked in the D_3 receptor homology model might yield information to develop a pharmacophore model to screen for additional compounds that abolish tolerance and SRT properties.

To further understand the difference in signaling properties induced by PD128907 and PBZI, they were docked to the binding site of the D_3 receptor. The docking was defined by

salt bridge interactions of the protonatable amine with Asp110, hydrogen bond interactions with conserved serine residues in transmembrane (TM) 5 and aromatic interactions with residues from TM6 and TM7. The ligand bound complexes were minimized and further refined using molecular dynamics (MD) simulations. A structural super positioning of the refined complexes yielded a root mean square deviation (rmsd) of 3.5 Å with well-marked differences in the TM bundle as well as the loop regions, leading to the hypothesis that each of these agonists elicit a unique conformational change in the receptor as previously observed for quinpirole. PD128907 forms a conserved salt bridge with Asp110, hydrogen bonds with Ser193 on TM5 and aromatic interactions with His349, Trp342 and Tyr373. Since the tetrahydropyran ring of PD128907 is more electronegative than the equivalent hexahydrobenzo group of PBZI, the latter probably has fewer interactions with the hydrophobic groups in TM3 as well as other aromatic residues in TM6. PBZI forms the conserved salt bridge with Asp110, hydrogen bonds with Ser192 in TM5 and has favorable pi-pi interactions with His349 and Phe345. In addition, due to the hydrophobic nature of hexahydrobenzo group, it forms favorable interactions with Val111 in TM3, Phe197 in TM5, Trp342 in TM6 and Tyr373 in TM7.

In order to verify if these interactions of PBZI with the D_3 receptor were significant, a three dimensional pharmacophore incorporating the hydrophobic elements, salt bridge interaction and aromatic-pi interactions was designed. Using this three dimensional pharmacophore, a three million compound library was screened using the HSB method for small molecules that could mimic the pharmacophore features of PBZI. The hits from the screen were then subjected to filtering schemes that include Lipinski's drug-like properties, Pregnane xenobiotic receptor activation and more importantly blood brain barrier (BBB) penetration. The 290 hits that resulted from the filtering schemes were docked to the binding site of D_3 receptor and scored using a variety of scoring schemes. The scoring schemes were customized to rank only those molecules that formed a salt bridge interaction with Asp110 and had favorable interactions with aromatic cluster formed by TM5, TM6 and TM7.

The fifteen compounds identified by the HSB in silico screen were evaluated for their ability to activate D_3 receptors and induce GIRK response, tolerance and SRT. The functional studies identified a novel D_3 receptor agonist, ES609 (4-(2-chlorophenyl)-butan-2-amine), which did not induce tolerance and SRT. FIG. 17 shows representative traces and cumulative data for ES609 and suggests that, as in the case of PBZI and FAUC73, ES609 also abolishes D_3 receptor tolerance and SRT properties (FIG. 17A). Control experiments showed that ES 609 did not elicit GIRK currents in parental AtT-20 cells (FIG. 17B) and the currents induced in AtT-20 cells stably expressing the D_3 receptor were blocked by pretreatment with the antagonist, eticlopride. Docking experiments confirmed that ES609 follows the interaction pattern of PBZI, with favorable pi-stacking interactions with His349 and salt bridge with Asp110 (FIG. 6A). Further refinement of the D_3 receptor-ES609 complex using MD simulation studies showed that ES609 elicits a similar conformation as that of PBZI. ES609-bound D_3 receptor structure super positioned on to the PBZI-bound D_3 receptor structure with a rmsd of 1.2 Å (FIG. 7A). The results show that while ES609 shares the same core structure as that of dopamine (FIG. 6D), its interaction pattern as well as the conformational change it elicits in the D_3 receptor are similar to those of PBZI.

Functional Characterization of PBZI and ES609

To compare the functional effects of the new class of D₃ receptor agonists, PBZI (Table 2) and ES609 (Table 3) were tested on AtT-20 cells stably expressing human D₃, D_{2S}, D_{2L} or D_{4.2}. Functional efficacy was determined by assessing the ability of PBZI and ES609 to inhibit adenylyl cyclase or activate GIRK channels coupled to these “D₂-like” dopamine receptors. The EC₅₀ values of PBZI and ES609 for inhibiting adenylyl cyclase and activating GIRK channels were in the range of 0.2 nM to 30 nM for D₃ receptors. By comparing to responses elicited by a saturating concentration (300 nM) of the full agonist, quinpirole, the results showed that PBZI and ES609 are full agonists at D₃ receptors in both the adenylyl cyclase and GIRK channel assays. In contrast, both compounds were partial agonists at the D_{2S} dopamine receptor. At D_{2L} dopamine receptors, PBZI was a full agonist in both assays; in contrast, ES609 did not elicit any response in the adenylyl cyclase assay and was a partial agonist in the GIRK channel assay. At D_{4.2} dopamine receptors, PBZI elicited no response and ES609 was a partial agonist in the adenylyl cyclase assay. In the GIRK channel assay, PBZI was a partial agonist at D_{4.2} receptors, while ES609 elicited no response. Together these results suggested that both PBZI and ES609, are full agonists at D₃ receptor, and were either partial agonists or elicited no responses at other D₂-like dopamine receptors.

TABLE 2

Cis-8OH-PBZI induced inhibition of adenylyl cyclase and activation of GIRK channels in AtT-20 cells stably expressing the individual dopamine receptor subtypes				
D ₂ -like dopamine receptors	Adenylyl cyclase inhibition EC ₅₀ (nM)	Adenylyl cyclase inhibition Ratio of PBZI to quinpirole response (at 300 nM)	GIRK channel activation EC ₅₀ (nM)	GIRK channel activation Ratio of PBZI to quinpirole response (at 300 nM)
D _{2S}	>150	0.29*	53 ± 19.8	0.48*
D _{2L}	67 ± 15	0.83	ND	0.75
D ₃	35 ± 2.8	0.97	29 ± 16.1	1.3
D _{4.2}	No response	0 [†]	ND	0.35*

*P < 0.05, statistically significant, Student's t-test.

[†]no adenylyl cyclase inhibition in the presence of 300 nM PBZI; but 300 nM quinpirole elicits ~70% inhibition.

ND—not determined; a full dose-response experiment to determine EC₅₀ was not performed; data was obtained for a single concentration (300 nM) of PBZI.

± Standard Error of Mean.

TABLE 3

ES609 induced inhibition of adenylyl cyclase and activation of GIRK channels in AtT-20 cells stably expressing the individual dopamine receptor subtypes				
D ₂ -like dopamine receptors	Adenylyl cyclase inhibition EC ₅₀ (nM)	Adenylyl cyclase inhibition Ratio of ES609 to quinpirole response (at 300 nM)	GIRK channel activation EC ₅₀ (nM)	GIRK channel activation Ratio of ES609 to quinpirole response (at 300 nM)
D _{2S}	2.4 ± 0.2	0.29*	ND	0.30*
D _{2L}	No response	0 [†]	ND	0.32*

TABLE 3-continued

ES609 induced inhibition of adenylyl cyclase and activation of GIRK channels in AtT-20 cells stably expressing the individual dopamine receptor subtypes				
D ₂ -like dopamine receptors	Adenylyl cyclase inhibition EC ₅₀ (nM)	Adenylyl cyclase inhibition Ratio of ES609 to quinpirole response (at 300 nM)	GIRK channel activation EC ₅₀ (nM)	GIRK channel activation Ratio of ES609 to quinpirole response (at 300 nM)
D ₃	0.15 ± 0.06	0.71	30 ± 7.7	0.86
D _{4.2}	0.82 ± 0.3	0.60*	No response	0 ^{††}

*P < 0.05, statistically significant, Student's t-test.

[†]no adenylyl cyclase inhibition in the presence of 300 nM ES609; but 300 nM quinpirole elicits ~80% inhibition.

^{††}300 nM ES609 elicits no GIRK response; but, 300 nM quinpirole elicits a full GIRK response.

ND—not determined; a full dose-response experiment to determine EC₅₀ was not performed; data was obtained for a single concentration (300 nM) of ES609.

± Standard Error of Mean.

The results described herein demonstrate that the D₃ receptor tolerance and SRT properties are ligand-dependent and help identify a new class of atypical D₃ receptor agonists that do not induce these properties. PBZI, a water soluble compound that is structurally similar to PD128907 but does not induce tolerance and SRT, was identified. Previous binding studies have shown that for D₂-like receptors, PBZI has a Ki of 27 nM for D₃, 1800 nM for D_{2S} and 280 nM for D_{4.2} (Scheideler et al., 1997, Eur. J. Pharmacol. 339(2-3):261-270). Binding at D₁-like receptors and a panel of other neurotransmitter receptors, ion channels and transporters were negligible. Thus PBZI exhibited D₃ receptor selectivity in receptor binding assay. This is consistent with the result from the functional studies, which shows that PBZI is a full agonist at D₃ receptors and a partial agonist at the D_{2S} dopamine receptors (Table 2). The partial agonistic effect at D_{2S} receptors is also consistent with previous in vitro results. In vivo, animals administered PBZI show specific increases in c-fos expression in medial prefrontal cortex and the shell region of nucleus accumbens, regions with high D₃ receptor expression. Effect of PBZI on D_{2L} receptors had not been previously determined. The results described herein show that PBZI is a full agonist at the D_{2L} dopamine receptor (Table 2). Based on the pre- and post-synaptic location of D_{2S} and D_{2L} receptors, respectively, PBZI would be predicted to primarily have postsynaptic effects. This is supported by previous in vivo studies as well (Fink-Jensen et al., 1998, Eur. J. Pharmacol. 342(2-3):153-161).

The classical D₂-like receptor agonist, PD128907, is selective for D₃ receptor with a Ki of 0.4 nM for D₃, 202 nM for D_{2S} and 114 nM for D_{4.2} (Scheideler et al., 1997, Eur. J. Pharmacol. 339(2-3):261-270); however the studies described herein showed that it induces severe tolerance and SRT. Interestingly, the chemical structure of PD128907 is similar to PBZI. Although PD128907 and PBZI share a similar core structure and function as full agonists at D₃ receptor, they have dramatically different effects on the tolerance and SRT properties. The results described herein suggest that the functional differences between PBZI and PD128907 are due to the different conformation these agonists induce in the D₃ receptor. The majority of these conformational changes are confined to the regions closest to the binding site and extracellular loop (EC) 2 loop region. Comparison of the PBZI- and PD128907-bound structures suggests that the maximum shift occurs in TM4, which is coupled with large movements of EC2. In addition, the conformational changes are also observed in intracellular loop (IC) 2 which has previously shown to be

important for mediating the tolerance property (Westrich et al., 2010, *Biochem. Pharmacol.* 79:897-907). Other significant conformational changes include a downward shift along the length of the TM6 helix and an unwinding of the first turn of the TM3 helix in the PD128907-bound form.

In this study, by carefully monitoring the nature of interactions and the conformational effects produced by the binding of PBZI and PD 128907 in computational models, a novel atypical D₃ receptor agonist, ES609, which exhibits selectivity and abolishes the tolerance and SRT properties, was designed. The HSB method may be used to identify additional atypical D₃ receptor agonists. ES609 was designed based on the nature of interactions of PBZI with D₃ receptor. The hypothesis was to increase the strength of the pi-pi interactions with the aromatic core by introducing electron withdrawing groups on the ligand. Thus the halogen group at ortho position in ES609 was highly suitable to the design, along with the added hydrophobicity proximal to the protonatable amine (FIG. 6D). The latter strengthened the interactions with hydrophobic groups on TM3 similar to the interactions of FAUC73 with residues in TM3. His349 may play a major role in promoting ligand biased signaling in D_{2L} receptor. These results are consistent with the findings that strengthening the pi-pi interactions with His349 and other members of the aromatic core may contribute favorably to the atypical properties of PBZI, FAUC73 and ES609. Studies are geared towards understanding the specific role of EC2 and TM4 in promoting functional selectivity and to further understand the complete structure-activity-relationship of these atypical agonists in complex with D₃ receptor. Functionally, ES609

and PBZI have similar efficacy for activating D₃ receptor coupled signal transduction pathways. However in contrast to PBZI, ES609 is either a partial agonist or elicits no response at other D₂-like receptors (D_{2S}, D_{2L} and D_{4.2}) when tested in two different signaling pathways. The selectivity exhibited by ES609 suggests that its structure could serve as a template for designing future selective D₃ receptor agonists. The functional affinity of ES609, defined by its EC₅₀, is similar to other D₂-like dopamine receptor agonists in the two signaling pathways that we studied. Furthermore, its small molecular weight and water solubility makes it ideally suited for in vivo studies. Given that PBZI and ES609 have similar signaling properties in vitro, they are expected to have a similar effect in vivo.

The new class of atypical D₃ receptor agonists described here represents a novel variation to the concept of functional selectivity. Traditionally, functional selectivity is ascribed to the ability of different ligands to elicit varying responses in different signal transduction pathways coupled to the same receptor. The ability of ligands to activate different pathways coupled to the same receptor to different degree is due to the different conformations that the ligand engenders in the receptor. Given that the D₃ receptor tolerance and SRT properties are determined by distinct conformational states, the identification of ligands that alters these conformational states is predicted to modulate the two properties. Consistent with this prediction, in this study a group of ligands that alter the two D₃ receptor properties was identified. The results described herein indicate that the functional selectivity concept may not be limited to selective

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 2

<210> SEQ ID NO 1

<211> LENGTH: 400

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

Met Ala Ser Leu Ser Gln Leu Ser Ser His Leu Asn Tyr Thr Cys Gly
1 5 10 15

Ala Glu Asn Ser Thr Gly Ala Ser Gln Ala Arg Pro His Ala Tyr Tyr
20 25 30

Ala Leu Ser Tyr Cys Ala Leu Ile Leu Ala Ile Val Phe Gly Asn Gly
35 40 45

Leu Val Cys Met Ala Val Leu Lys Glu Arg Ala Leu Gln Thr Thr Thr
50 55 60

Asn Tyr Leu Val Val Ser Leu Ala Val Ala Asp Leu Leu Val Ala Thr
65 70 75 80

Leu Val Met Pro Trp Val Val Tyr Leu Glu Val Thr Gly Gly Val Trp
85 90 95

Asn Phe Ser Arg Ile Cys Cys Asp Val Phe Val Thr Leu Asp Val Met
100 105 110

Met Cys Thr Ala Ser Ile Leu Asn Leu Cys Ala Ile Ser Ile Asp Arg
115 120 125

Tyr Thr Ala Val Val Met Pro Val His Tyr Gln His Gly Thr Gly Gln
130 135 140

Ser Ser Cys Arg Arg Val Ala Leu Met Ile Thr Ala Val Trp Val Leu
145 150 155 160

Ala Phe Ala Val Ser Cys Pro Leu Leu Phe Gly Phe Asn Thr Thr Gly

-continued

165					170					175					
Asp	Pro	Thr	Val	Cys	Ser	Ile	Ser	Asn	Pro	Asp	Phe	Val	Ile	Tyr	Ser
			180					185					190		
Ser	Val	Val	Ser	Phe	Tyr	Leu	Pro	Phe	Gly	Val	Thr	Val	Leu	Val	Tyr
			195				200						205		
Ala	Arg	Ile	Tyr	Val	Val	Leu	Lys	Gln	Arg	Arg	Arg	Lys	Arg	Ile	Leu
			210				215						220		
Thr	Arg	Gln	Asn	Ser	Gln	Cys	Asn	Ser	Val	Arg	Pro	Gly	Phe	Pro	Gln
			225				230						235		240
Gln	Thr	Leu	Ser	Pro	Asp	Pro	Ala	His	Leu	Glu	Leu	Lys	Arg	Tyr	Tyr
				245					250					255	
Ser	Ile	Cys	Gln	Asp	Thr	Ala	Leu	Gly	Gly	Pro	Gly	Phe	Gln	Glu	Arg
			260					265					270		
Gly	Gly	Glu	Leu	Lys	Arg	Glu	Glu	Lys	Thr	Arg	Asn	Ser	Leu	Ser	Pro
			275					280					285		
Thr	Ile	Ala	Pro	Lys	Leu	Ser	Leu	Glu	Val	Arg	Lys	Leu	Ser	Asn	Gly
			290				295						300		
Arg	Leu	Ser	Thr	Ser	Leu	Lys	Leu	Gly	Pro	Leu	Gln	Pro	Arg	Gly	Val
				305			310						315		320
Pro	Leu	Arg	Glu	Lys	Lys	Ala	Thr	Gln	Met	Val	Ala	Ile	Val	Leu	Gly
				325					330					335	
Ala	Phe	Ile	Val	Cys	Trp	Leu	Pro	Phe	Phe	Leu	Thr	His	Val	Leu	Asn
			340					345						350	
Thr	His	Cys	Gln	Thr	Cys	His	Val	Ser	Pro	Glu	Leu	Tyr	Ser	Ala	Thr
			355				360						365		
Thr	Trp	Leu	Gly	Tyr	Val	Asn	Ser	Ala	Leu	Asn	Pro	Val	Ile	Tyr	Thr
			370				375						380		
Thr	Phe	Asn	Ile	Glu	Phe	Arg	Lys	Ala	Phe	Leu	Lys	Ile	Leu	Ser	Cys
			385				390						395		400

<210> SEQ ID NO 2

<211> LENGTH: 500

<212> TYPE: PRT


<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

Asp	Tyr	Lys	Asp	Asp	Asp	Ala	Met	Gly	Gln	Pro	Gly	Asn	Gly	Ser	Ala
				5					10					15	
Phe	Leu	Leu	Ala	Pro	Asn	Arg	Ser	His	Ala	Pro	Asp	His	Asp	Val	Thr
			20					25					30		
Gln	Gln	Arg	Asp	Glu	Val	Trp	Val	Val	Gly	Met	Gly	Ile	Val	Met	Ser
			35				40						45		
Leu	Ile	Val	Leu	Ala	Ile	Val	Phe	Gly	Asn	Val	Leu	Val	Ile	Thr	Ala
			50				55						60		
Ile	Ala	Lys	Phe	Glu	Arg	Leu	Gln	Thr	Val	Thr	Asn	Tyr	Phe	Ile	Thr
				65			70						75		80
Ser	Leu	Ala	Cys	Ala	Asp	Leu	Val	Met	Gly	Leu	Ala	Val	Val	Pro	Phe
				85					90					95	
Gly	Ala	Ala	His	Ile	Leu	Met	Lys	Met	Trp	Thr	Phe	Gly	Asn	Phe	Trp
			100					105					110		
Cys	Glu	Phe	Trp	Thr	Ser	Ile	Asp	Val	Leu	Cys	Val	Thr	Ala	Ser	Ile
			115				120						125		
Glu	Thr	Leu	Cys	Val	Ile	Ala	Val	Asp	Arg	Tyr	Phe	Ala	Ile	Thr	Ser
				130			135						140		

What is claimed is:

1. A method of treating or ameliorating levodopa-induced dyskinesia in a patient suffering from Parkinson's Disease, said method comprising administering to said patient a therapeutically effective amount of at least one compound of formula (I):



wherein in formula (I):

R^1 , R^2 and R^3 are independently selected from the group consisting of H, cyano, halo, hydroxyl, alkoxy, nitro, C1-6 alkyl, and carboxy;

R^4 and R^5 are independently selected from the group consisting of H, C1-6 alkyl, and substituted C1-6 alkyl; and, n is 2,

mixtures thereof, or a pharmaceutically acceptable salt thereof.

2. The method of claim 1, wherein said at least one compound is selected from the group consisting of:

2-amino-4-(2-chlorophenyl)butan-1-ol;

2-(3-aminohexyl)phenol;

4-(2-chlorophenyl)-2-methylamino-butane;

4-(2-fluorophenyl)butan-2-amine;

4-(2-bromophenyl)butan-2-amine;

4-(2-iodophenyl)butan-2-amine;

4-(2-methoxyphenyl)butan-2-amine;

2-(3-aminobutyl)phenol;

3-(3,4-diethoxyphenyl)propan-1-amine;

4-(4-chlorophenyl)butan-2-amine;

4-(4-methoxyphenyl)butan-2-amine;

mixtures thereof, or a pharmaceutically acceptable salt thereof.

3. The method of claim 1, wherein said patient is further administered a drug selected from the group consisting of levodopa, clozapine, bromocriptine, pergolide, pramipexole, ropinirole, piribedil, cabergoline, apomorphine, and lisuride, a salt thereof and mixtures thereof.


4. The method of claim 1, wherein said compound is co-administered to said patient with a second pharmaceutical composition comprising levodopa.

5. The method of claim 1, wherein said compound is administered to said patient a given period of time before a second pharmaceutical composition comprising levodopa is administered to said patient.

6. The method of claim 5, wherein said given period of time varies from about 2 minutes to about 24 hours.

7. The method of claim 1, wherein said patient is human.

8. A method of treating or ameliorating Parkinson's Disease in a patient in need thereof, said method comprising administering to said patient a therapeutically effective amount of at least one compound of formula (I):



(I) wherein in formula (I):

R^1 , R^2 and R^3 are independently selected from the group consisting of H, cyano, halo, hydroxyl, alkoxy, nitro, C1-6 alkyl, and carboxy;

R^4 and R^5 are independently selected from the group consisting of H, C1-6 alkyl, and substituted C1-6 alkyl; and, n is 2,

mixtures thereof, or a pharmaceutically acceptable salt thereof.

9. The method of claim 8, wherein said at least one compound is selected from the group consisting of:

2-amino-4-(2-chlorophenyl)butan-1-ol;

2-(3-aminohexyl)phenol;

4-(2-chlorophenyl)-2-methylamino-butane;

4-(2-fluorophenyl)butan-2-amine;

4-(2-bromophenyl)butan-2-amine;

4-(2-iodophenyl)butan-2-amine;

4-(2-methoxyphenyl)butan-2-amine;

2-(3-aminobutyl)phenol;

3-(3,4-diethoxyphenyl)propan-1-amine;

4-(4-chlorophenyl)butan-2-amine;

4-(4-methoxyphenyl)butan-2-amine;

mixtures thereof, or a pharmaceutically acceptable salt thereof.

10. The method of claim 8, wherein said compound is administered to said patient as a composition that further comprises at least one drug selected from the group consisting of levodopa, clozapine, bromocriptine, pergolide, pramipexole, ropinirole, piribedil, cabergoline, apomorphine, and lisuride, a salt thereof and mixtures thereof.

11. The method of claim 8, wherein said patient is further administered a second pharmaceutical composition comprising at least one drug selected from the group consisting of levodopa, clozapine, bromocriptine, pergolide, pramipexole, ropinirole, piribedil, cabergoline, apomorphine, and lisuride, a salt thereof and mixtures thereof.

12. A method of treating or ameliorating levodopa-induced dyskinesia in a patient suffering from Parkinson's Disease, said method comprising administering to said patient a therapeutically effective amount of at least one compound selected from the group consisting of:

2-amino-4-(2-chlorophenyl)butan-1-ol;

2-(3-aminohexyl)phenol;

4-(2-chlorophenyl)-2-methylamino-butane;

4-(2-fluorophenyl)butan-2-amine;

4-(2-bromophenyl)butan-2-amine;

4-(2-iodophenyl)butan-2-amine;

4-(2-methoxyphenyl)butan-2-amine;

2-(3-aminobutyl)phenol;

3-(3,4-diethoxyphenyl)propan-1-amine;

4-(4-chlorophenyl)butan-2-amine;

4-(4-methoxyphenyl)butan-2-amine;

mixtures thereof, or a pharmaceutically acceptable salt thereof.

13. A method of treating or ameliorating Parkinson's Disease in a patient, said method comprising administering to said patient a therapeutically effective amount of at least one compound selected from the group consisting of:

61

2-amino-4-(2-chlorophenyl)butan-1-ol;
2-(3-aminohexyl)phenol;
4-(2-chlorophenyl)-2-methylamino-butane;
4-(2-fluorophenyl)butan-2-amine;
4-(2-bromophenyl)butan-2-amine; 5
4-(2-iodophenyl)butan-2-amine;
4-(2-methoxyphenyl)butan-2-amine;
2-(3-aminobutyl)phenol;
3-(3,4-diethoxyphenyl)propan-1-amine;
4-(4-chlorophenyl)butan-2-amine; 10
4-(4-methoxyphenyl)butan-2-amine;
mixtures thereof, or a pharmaceutically acceptable salt
thereof.

* * * * *

62