



US009312505B2

(12) **United States Patent**
Brooks et al.

(10) **Patent No.:** **US 9,312,505 B2**
(45) **Date of Patent:** **Apr. 12, 2016**

(54) **ORGANIC ELECTROLUMINESCENT MATERIALS AND DEVICES**

(71) Applicant: **Universal Display Corporation**, Ewing, NJ (US)

(72) Inventors: **Jason Brooks**, Philadelphia, PA (US);
Glenn Morello, Pittsburgh, PA (US);
Chuanjun Xia, Lawrenceville, NJ (US);
Jun Deng, Murrysville, PA (US)

(73) Assignee: **Universal Display Corporation**, Ewing, NJ (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 455 days.

(21) Appl. No.: **13/787,169**

(22) Filed: **Mar. 6, 2013**

(65) **Prior Publication Data**

US 2014/0084261 A1 Mar. 27, 2014

Related U.S. Application Data

(60) Provisional application No. 61/705,400, filed on Sep. 25, 2012.

(51) **Int. Cl.**

H01L 51/54 (2006.01)
C09K 11/06 (2006.01)
H01L 51/50 (2006.01)
C07F 15/00 (2006.01)
H01L 51/00 (2006.01)

(52) **U.S. Cl.**

CPC **H01L 51/50** (2013.01); **C07F 15/0086** (2013.01); **C09K 11/06** (2013.01); **H01L 51/0087** (2013.01); **C09K 2211/1029** (2013.01); **C09K 2211/1033** (2013.01); **C09K 2211/1044** (2013.01); **C09K 2211/185** (2013.01); **H01L 51/0085** (2013.01); **H01L 51/5016** (2013.01)

(58) **Field of Classification Search**

None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,769,292 A 9/1988 Tang et al.
5,061,569 A 10/1991 VanSlyke et al.
5,247,190 A 9/1993 Friend et al.
5,703,436 A 12/1997 Forrest et al.
5,707,745 A 1/1998 Forrest et al.
5,834,893 A 11/1998 Bulovic et al.
5,844,363 A 12/1998 Gu et al.
6,013,982 A 1/2000 Thompson et al.
6,087,196 A 7/2000 Sturm et al.
6,091,195 A 7/2000 Forrest et al.
6,097,147 A 8/2000 Baldo et al.
6,294,398 B1 9/2001 Kim et al.
6,303,238 B1 10/2001 Thompson et al.
6,337,102 B1 1/2002 Forrest et al.
6,468,819 B1 10/2002 Kim et al.
6,528,187 B1 3/2003 Okada

6,687,266 B1 2/2004 Ma et al.
6,835,469 B2 12/2004 Kwong et al.
6,893,743 B2* 5/2005 Sato et al. H01L 51/0059
257/102

6,921,915 B2 7/2005 Takiguchi et al.
7,087,321 B2 8/2006 Kwong et al.
7,090,928 B2 8/2006 Thompson et al.
7,154,114 B2 12/2006 Brooks et al.

(Continued)

FOREIGN PATENT DOCUMENTS

EP 0650955 5/1995
EP 1725079 11/2006
EP 2034538 3/2009
JP 200511610 1/2005
JP 2007123392 5/2007
JP 2007254297 10/2007
JP 2008074939 4/2008
WO 0139234 5/2001
WO 0202714 1/2002
WO 0215645 2/2002

(Continued)

OTHER PUBLICATIONS

U.S. Appl. No. 61/490,111, filed May 26, 2011.*


(Continued)

Primary Examiner — Marie R. Yamnitzky

(74) *Attorney, Agent, or Firm* — Duane Morris LLP

(57) **ABSTRACT**

In certain embodiments, the invention provides metal complexes having Formula (I):



wherein each L^x is independently a monodentate ligand, and any two adjacent L^x may optionally combine to form a bidentate ligand; wherein M^1 is cobalt(I), rhodium(I), iridium(I), nickel(II), platinum(II), palladium(II), silver(III), gold(III), or copper(III); wherein m is a value from 1 to the maximum number of ligands that may be attached to M^1 ; wherein $m+n$ is the maximum number of ligands that may be attached to M^1 ; wherein G^1 is O or CR^4R^5 ; and R^1 to R^5 are various substituents, which can optionally combine with each other, among themselves, or with any L^x . In certain embodiments, the invention provides devices, such as organic light emitting devices, that comprise such metal complexes.

15 Claims, 7 Drawing Sheets

(56)

References Cited

U.S. PATENT DOCUMENTS

7,250,226 B2 7/2007 Tokito et al.
 7,279,704 B2 10/2007 Walters et al.
 7,332,232 B2 2/2008 Ma et al.
 7,338,722 B2 3/2008 Thompson et al.
 7,393,599 B2 7/2008 Thompson et al.
 7,396,598 B2 7/2008 Takeuchi et al.
 7,431,968 B1 10/2008 Shtein et al.
 7,445,855 B2 11/2008 Mackenzie et al.
 7,534,505 B2 5/2009 Lin et al.
 2002/0034656 A1 3/2002 Thompson et al.
 2002/0134984 A1 9/2002 Igarashi
 2002/0158242 A1 10/2002 Son et al.
 2003/0138657 A1 7/2003 Li et al.
 2003/0152802 A1 8/2003 Tsuboyama et al.
 2003/0162053 A1 8/2003 Marks et al.
 2003/0175553 A1 9/2003 Thompson et al.
 2003/0230980 A1 12/2003 Forrest et al.
 2004/0036077 A1 2/2004 Ise
 2004/0137267 A1 7/2004 Igarashi et al.
 2004/0137268 A1 7/2004 Igarashi et al.
 2004/0174116 A1 9/2004 Lu et al.
 2005/0006625 A1* 1/2005 Seo et al. C07D 213/74
 252/301.16
 2005/0025993 A1 2/2005 Thompson et al.
 2005/0112407 A1 5/2005 Ogasawara et al.
 2005/0238919 A1 10/2005 Ogasawara
 2005/0244673 A1 11/2005 Satoh et al.
 2005/0260441 A1 11/2005 Thompson et al.
 2005/0260449 A1 11/2005 Walters et al.
 2006/0008670 A1 1/2006 Lin et al.
 2006/0202194 A1 9/2006 Jeong et al.
 2006/0240279 A1 10/2006 Adamovich et al.
 2006/0251923 A1 11/2006 Lin et al.
 2006/0263635 A1 11/2006 Ise
 2006/0280965 A1 12/2006 Kwong et al.
 2007/0190359 A1 8/2007 Knowles et al.
 2007/0278938 A1 12/2007 Yabunouchi et al.
 2008/0015355 A1 1/2008 Schafer et al.
 2008/0018221 A1 1/2008 Egen et al.
 2008/0106190 A1 5/2008 Yabunouchi et al.
 2008/0124572 A1 5/2008 Mizuki et al.
 2008/0220265 A1 9/2008 Xia et al.
 2008/0297033 A1 12/2008 Knowles et al.
 2009/0008605 A1 1/2009 Kawamura et al.
 2009/0009065 A1 1/2009 Nishimura et al.
 2009/0017330 A1 1/2009 Iwakuma et al.
 2009/0030202 A1 1/2009 Iwakuma et al.
 2009/0039776 A1 2/2009 Yamada et al.
 2009/0045730 A1 2/2009 Nishimura et al.
 2009/0045731 A1 2/2009 Nishimura et al.
 2009/0101870 A1 4/2009 Prakash et al.
 2009/0108737 A1 4/2009 Kwong et al.
 2009/0115316 A1 5/2009 Zheng et al.
 2009/0165846 A1 7/2009 Johannes et al.
 2009/0167162 A1 7/2009 Lin et al.
 2009/0179554 A1 7/2009 Kuma et al.
 2012/0302753 A1* 11/2012 Li et al. C07F 15/0086
 546/4
 2014/0091265 A1* 4/2014 Stoessel et al. H01L 51/0087
 252/519.21

FOREIGN PATENT DOCUMENTS

WO 03040257 5/2003
 WO 03060956 7/2003
 WO 2004093207 10/2004
 WO 2004107822 12/2004
 WO 2005014551 2/2005
 WO 2005019373 3/2005
 WO 2005030900 4/2005
 WO 2005089025 9/2005
 WO 2005123873 12/2005
 WO 2006009024 1/2006
 WO 2006056418 6/2006

WO 2006072002 7/2006
 WO 2006082742 8/2006
 WO 2006098120 9/2006
 WO 2006100298 9/2006
 WO 2006103874 10/2006
 WO 2006114966 11/2006
 WO 2006132173 12/2006
 WO 2007002683 1/2007
 WO 2007004380 1/2007
 WO 2007063754 6/2007
 WO 2007063796 6/2007
 WO 2008056746 5/2008
 WO 2008101842 8/2008
 WO 2008132085 11/2008
 WO 2009000673 12/2008
 WO 2009003898 1/2009
 WO 2009008311 1/2009
 WO 2009018009 2/2009
 WO 2009021126 2/2009
 WO 2009050290 4/2009
 WO 2009062578 5/2009
 WO 2009063833 5/2009
 WO 2009066778 5/2009
 WO 2009066779 5/2009
 WO 2009086028 7/2009
 WO 2009100991 8/2009
 WO WO 2012/163471 A1 * 12/2012

OTHER PUBLICATIONS

Adachi, Chihaya et al., "Organic Electroluminescent Device Having a Hole Conductor as an Emitting Layer," Appl. Phys. Lett., 55(15): 1489-1491 (1989).
 Adachi, Chihaya et al., "Nearly 100% Internal Phosphorescence Efficiency in an Organic Light Emitting Device," J. Appl. Phys., 90(10): 5048-5051 (2001).
 Adachi, Chihaya et al., "High-Efficiency Red Electrophosphorescence Devices," Appl. Phys. Lett., 78(11):1622-1624 (2001).
 Aonuma, Masaki et al., "Material Design of Hole Transport Materials Capable of Thick-Film Formation in Organic Light Emitting Diodes," Appl. Phys. Lett., 90:183503-1-183503-3, (2007).
 Baldo et al., "Highly Efficient Phosphorescent Emission from Organic Electroluminescent Devices," Nature, vol. 395, 151-154, (1998).
 Baldo et al., "Very high-efficiency green organic light-emitting devices based on electrophosphorescence," Appl. Phys. Lett., vol. 75, No. 1, 4-6 (1999).
 Gao, Zhiqiang et al., "Bright-Blue Electroluminescence From a Silyl-Substituted ter-(phenylene-vinylene) derivative," Appl. Phys. Lett., 74(6): 865-867 (1999).
 Hamada, Yuji et al., "High Luminance in Organic Electroluminescent Devices with Bis(10-hydroxybenzo[h]quinolinato) beryllium as an Emitter," Chem. Lett., 905-906 (1993).
 Holmes, R.J. et al., "Blue Organic Electrophosphorescence Using Exothermic Host-Guest Energy Transfer," Appl. Phys. Lett., 82(15):2422-2424 (2003).
 Hu, Nan-Xing et al., "Novel High Tg Hole-Transport Molecules Based on Indolo[3,2-b]carbazoles for Organic Light-Emitting Devices," Synthetic Metals, 111-112:421-424 (2000).
 Huang, Jinsong et al., "Highly Efficient Red-Emission Polymer Phosphorescent Light-Emitting Diodes Based on Two Novel Tris(1-phenylisoquinolinato-C2,N)iridium(III)Derivatives," Adv. Mater., 19:739-743 (2007).
 Huang, Wei-Sheng et al., "Highly Phosphorescent Bis-Cyclometalated Iridium Complexes Containing Benzoimidazole-Based Ligands," Chem. Mater, 16(12):2480-2488 (2004).
 Hung, L.S. et al., "Anode Modification in Organic Light-Emitting Diodes by Low-Frequency Plasma Polymerization of CHF3," Appl. Phys. Lett., 78(5):673-675 (2001).
 Ikeda, Hisao et al., "P-185 Low-Drive-Voltage OLEDs with a Buffer Layer Having Molybdenum Oxide," SID Symposium Digest, 37:923-926 (2006).

(56)

References Cited

OTHER PUBLICATIONS

- Inada, Hiroshi and Shirota, Yasuhiko, "1,3,5-Tris[4-(diphenylamino)phenyl]benzene and its Methylsubstituted Derivatives as a Novel Class of Amorphous Molecular Materials," *J. Mater. Chem.*, 3(3):319-320 (1993).
- Kanno, Hiroshi et al., "Highly Efficient and Stable Red Phosphorescent Organic Light-Emitting Device Using bis[2-(2-benzothiazoyl)phenolato]zinc(II) as host material," *Appl. Phys. Lett.*, 90:123509-1-123509-3 (2007).
- Kido, Junji et al., "1,2,4-Triazole Derivative as an Electron Transport Layer in Organic Electroluminescent Devices," *Jpn. J. Appl. Phys.*, 32:L917-L920 (1993).
- Kuwabara, Yoshiyuki et al., "Thermally Stable Multilayered Organic Electroluminescent Devices Using Novel Starburst Molecules, 4,4',4''-Tri(N-carbazolyl)triphenylamine (TCTA) and 4,4',4''-Tris(3-methylphenylphenyl-amino) triphenylamine (m-MTDATA), as Hole-Transport Materials," *Adv. Mater.*, 6(9):677-679 (1994).
- Kwong, Raymond C. et al., "High Operational Stability of Electrophosphorescent Devices," *Appl. Phys. Lett.*, 81(1) 162-164 (2002).
- Lamansky, Sergey et al., "Synthesis and Characterization of Phosphorescent Cyclometalated Iridium Complexes," *Inorg. Chem.*, 40(7):1704-1711 (2001).
- Lee, Chang-Lyoul et al., "Polymer Phosphorescent Light-Emitting Devices Doped with Tris(2-phenylpyridine) Iridium as a Triplet Emitter," *Appl. Phys. Lett.*, 77(15)2280-2282 (2000).
- Lo, Shih-Chun et al., "Blue Phosphorescence from Iridium(III) Complexes at Room Temperature," *Chem. Mater.*, 18 (21)5119-5129 (2006).
- Ma, Yuguang et al., "Triplet Luminescent Dinuclear-Gold(I) Complex-Based Light-Emitting Diodes with Low Turn-On voltage," *Appl. Phys. Lett.*, 74(10):1361-1363 (1999).
- Mi, Bao-Xiu et al., "Thermally Stable Hole-Transporting Material for Organic Light-Emitting Diode an Isoindole Derivative," *Chem. Mater.*, 15(16):3148-3151 (2003).
- Nishida, Jun-ichi et al., "Preparation, Characterization, and Electroluminescence Characteristics of α -Diimine-type Platinum(II) Complexes with Perfluorinated Phenyl Groups as Ligands," *Chem. Lett.*, 34(4): 592-593 (2005).
- Niu, Yu-Hua et al., "Highly Efficient Electrophosphorescent Devices with Saturated Red Emission from a Neutral Osmium Complex," *Chem. Mater.*, 17(13):3532-3536 (2005).
- Noda, Tetsuya and Shirota, Yasuhiko, "5,5'-Bis(dimesitylboryl)-2,2'-bithiophene and 5,5''-Bis(dimesitylboryl)-2,2',2''-terthiophene as a Novel Family of Electron-Transporting Amorphous Molecular Materials," *J. Am. Chem. Soc.*, 120 (37):9714-9715 (1998).
- Okumoto, Kenji et al., "Green Fluorescent Organic Light-Emitting Device with External Quantum Efficiency of Nearly 10%," *Appl. Phys. Lett.*, 89:063504-1-063504-3 (2006).
- Palilis, Leonidas C., "High Efficiency Molecular Organic Light-Emitting Diodes Based on Silole Derivatives and Their Exciplexes," *Organic Electronics*, 4:113-121 (2003).
- Paulose, Betty Marie Jennifer S. et al., "First Examples of Alkenyl Pyridines as Organic Ligands for Phosphorescent Iridium Complexes," *Adv. Mater.*, 16(22):2003-2007 (2004).
- Ranjan, Sudhir et al., "Realizing Green Phosphorescent Light-Emitting Materials from Rhenium(I) Pyrazolato Diimine Complexes," *Inorg. Chem.*, 42(4):1248-1255 (2003).
- Sakamoto, Youichi et al., "Synthesis, Characterization, and Electron-Transport Property of Perfluorinated Phenylene Dendrimers," *J. Am. Chem. Soc.*, 122(8):1832-1833 (2000).
- Sotoyama, Wataru et al., "Efficient Organic Light-Emitting Diodes with Phosphorescent Platinum Complexes Containing NCN-Coordinating Tridentate Ligand," *Appl. Phys. Lett.*, 86:153505-1-153505-3 (2005).
- Sun, Yiru and Forrest, Stephen R., "High-Efficiency White Organic Light Emitting Devices with Three Separate Phosphorescent Emission Layers," *Appl. Phys. Lett.*, 91:263503-1-263503-3 (2007).
- T. Östergård et al., "Langmuir-Blodgett Light-Emitting Diodes of Poly(3-Hexylthiophene) Electro-Optical Characteristics Related to Structure," *Synthetic Metals*, 88:171-177 (1997).
- Takizawa, Shin-ya et al., "Phosphorescent Iridium Complexes Based on 2-Phenylimidazo[1,2- α]pyridine Ligands Tuning of Emission Color toward the Blue Region and Application to Polymer Light-Emitting Devices," *Inorg. Chem.*, 46(10):4308-4319 (2007).
- Tang, C.W. and VanSlyke, S.A., "Organic Electroluminescent Diodes," *Appl. Phys. Lett.*, 51(12):913-915 (1987).
- Tung, Yung-Liang et al., "Organic Light-Emitting Diodes Based on Charge-Neutral Ru II Phosphorescent Emitters," *Adv. Mater.*, 17(8)1059-1064 (2005).
- Van Slyke, S. A. et al., "Organic Electroluminescent Devices with Improved Stability," *Appl. Phys. Lett.*, 69 (15):2160-2162 (1996).
- Wang, Y. et al., "Highly Efficient Electroluminescent Materials Based on Fluorinated Organometallic Iridium Compounds," *Appl. Phys. Lett.*, 79(4):449-451 (2001).
- Wong, Keith Man-Chung et al., "A Novel Class of Phosphorescent Gold(III) Alkynyl-Based Organic Light-Emitting Devices with Tunable Colour," *Chem. Commun.*, 2906-2908 (2005).
- Wong, Wai-Yeung, "Multifunctional Iridium Complexes Based on Carbazole Modules as Highly Efficient Electrophosphors," *Angew. Chem. Int. Ed.*, 45:7800-7803 (2006).
- Baldo et al., "Very high-efficiency green organic light-emitting devices based on electrophosphorescence," *Appl. Phys. Lett.*, vol. 75, No. 1, 4-6 (1999).
- Guo, Tzung-Fang et al., "Highly Efficient Electrophosphorescent Polymer Light-Emitting Devices," *Organic Electronics*, 1: 15-20 (2000).
- Huang, Jinsong et al., "Highly Efficient Red-Emission Polymer Phosphorescent Light-Emitting Diodes Based on Two Novel Tris(1-phenylisoquinolinato-C2,N)iridium(III) Derivatives," *Adv. Mater.*, 19:739-743 (2007).
- Ikai, Masamichi et al., "Highly Efficient Phosphorescence From Organic Light-Emitting Devices with an Exciton-Block Layer," *Appl. Phys. Lett.*, 79(2):156-158 (2001).
- Salbeck, J. et al., "Low Molecular Organic Glasses for Blue Electroluminescence," *Synthetic Metals*, 91: 209-215 (1997).
- Shirota, Yasuhiko et al., "Starburst Molecules Based on pi-Electron Systems as Materials for Organic Electroluminescent Devices," *Journal of Luminescence*, 72-74:985-991 (1997).
- T. Östergård et al., "Langmuir-Blodgett Light-Emitting Diodes of Poly(3-Hexylthiophene) Electro-Optical Characteristics Related to Structure," *Synthetic Metals*, 88: 171-177 (1997).

* cited by examiner




FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7

1
**ORGANIC ELECTROLUMINESCENT
 MATERIALS AND DEVICES**

CROSS-REFERENCE TO RELATED
 APPLICATIONS

This application claims priority to U.S. Provisional Application Ser. No. 61/705,400, filed Sep. 25, 2012, the entire contents of which are incorporated herein by reference.

JOINT RESEARCH AGREEMENT

The claimed invention was made by, on behalf of, and/or in connection with one or more of the following parties to a joint university corporation research agreement: Regents of the University of Michigan, Princeton University, University of Southern California, and the Universal Display Corporation. The agreement was in effect on and before the date the claimed invention was made, and the claimed invention was made as a result of activities undertaken within the scope of the agreement.

FIELD OF THE INVENTION

The present invention relates to organic light emitting devices (OLEDs). In particular, the invention relates to metal complexes having a bridged N,N-diaryl ligand and their use in various devices, including OLEDs.

BACKGROUND


Opto-electronic devices that make use of organic materials are becoming increasingly desirable for a number of reasons. Many of the materials used to make such devices are relatively inexpensive, so organic opto-electronic devices have the potential for cost advantages over inorganic devices. In addition, the inherent properties of organic materials, such as their flexibility, may make them well suited for particular applications such as fabrication on a flexible substrate. Examples of organic opto-electronic devices include organic light emitting devices (OLEDs), organic phototransistors, organic photovoltaic cells, and organic photodetectors. For OLEDs, the organic materials may have performance advantages over conventional materials. For example, the wavelength at which an organic emissive layer emits light may generally be readily tuned with appropriate dopants.

OLEDs make use of thin organic films that emit light when voltage is applied across the device. OLEDs are becoming an increasingly interesting technology for use in applications such as flat panel displays, illumination, and backlighting. Several OLED materials and configurations are described in U.S. Pat. Nos. 5,844,363, 6,303,238, and 5,707,745, which are incorporated herein by reference in their entirety.

One application for phosphorescent emissive molecules is a full color display. Industry standards for such a display call for pixels adapted to emit particular colors, referred to as "saturated" colors. In particular, these standards call for saturated red, green, and blue pixels. Color may be measured using CIE coordinates, which are well known to the art.

One example of a green emissive molecule is tris(2-phenylpyridine) iridium, denoted Ir(ppy)₃, which has the following structure:

2



In this, and later figures herein, we depict the dative bond from nitrogen to metal (here, Ir) as a straight line.

As used herein, the term "organic" includes polymeric materials as well as small molecule organic materials that may be used to fabricate organic opto-electronic devices. "Small molecule" refers to any organic material that is not a polymer, and "small molecules" may actually be quite large. Small molecules may include repeat units in some circumstances. For example, using a long chain alkyl group as a substituent does not remove a molecule from the "small molecule" class. Small molecules may also be incorporated into polymers, for example as a pendent group on a polymer backbone or as a part of the backbone. Small molecules may also serve as the core moiety of a dendrimer, which consists of a series of chemical shells built on the core moiety. The core moiety of a dendrimer may be a fluorescent or phosphorescent small molecule emitter. A dendrimer may be a "small molecule," and it is believed that all dendrimers currently used in the field of OLEDs are small molecules.

As used herein, "top" means furthest away from the substrate, while "bottom" means closest to the substrate. Where a first layer is described as "disposed over" a second layer, the first layer is disposed further away from substrate. There may be other layers between the first and second layer, unless it is specified that the first layer is "in contact with" the second layer. For example, a cathode may be described as "disposed over" an anode, even though there are various organic layers in between.

As used herein, "solution processable" means capable of being dissolved, dispersed, or transported in and/or deposited from a liquid medium, either in solution or suspension form.

A ligand may be referred to as "photoactive" when it is believed that the ligand directly contributes to the photoactive properties of an emissive material. A ligand may be referred to as "ancillary" when it is believed that the ligand does not contribute to the photoactive properties of an emissive material, although an ancillary ligand may alter the properties of a photoactive ligand.

As used herein, and as would be generally understood by one skilled in the art, a first "Highest Occupied Molecular Orbital" (HOMO) or "Lowest Unoccupied Molecular Orbital" (LUMO) energy level is "greater than" or "higher than" a second HOMO or LUMO energy level if the first energy level is closer to the vacuum energy level. Since ionization potentials (IP) are measured as a negative energy relative to a vacuum level, a higher HOMO energy level corresponds to an IP having a smaller absolute value (an IP that is less negative). Similarly, a higher LUMO energy level corresponds to an electron affinity (EA) having a smaller absolute value (an EA that is less negative). On a conventional energy level diagram, with the vacuum level at the top, the LUMO energy level of a material is higher than the HOMO energy level of the same material. A "higher" HOMO or LUMO energy level appears closer to the top of such a diagram than a "lower" HOMO or LUMO energy level.


3

As used herein, and as would be generally understood by one skilled in the art, a first work function is “greater than” or “higher than” a second work function if the first work function has a higher absolute value. Because work functions are generally measured as negative numbers relative to vacuum level, this means that a “higher” work function is more negative. On a conventional energy level diagram, with the vacuum level at the top, a “higher” work function is illustrated as further away from the vacuum level in the downward direction. Thus, the definitions of HOMO and LUMO energy levels follow a different convention than work functions.

More details on OLEDs, and the definitions described above, can be found in U.S. Pat. No. 7,279,704, which is incorporated herein by reference in its entirety.

SUMMARY OF THE INVENTION

In at least one aspect, the invention provides metal complexes having an N,N-diaryl ligand, such as compounds having Formula (I):



wherein each L^x is independently a monodentate ligand, and any two adjacent L^x may optionally combine to form a bidentate ligand; wherein M^1 is cobalt(I), rhodium(I), iridium(I), nickel(II), platinum(II), palladium(II), silver(III), gold(III), or copper(III); wherein m is a value from 1 to the maximum number of ligands that may be attached to M^1 ; wherein $m+n$ is the maximum number of ligands that may be attached to M^1 ; wherein G^1 is O or CR^4R^5 ; wherein R^1 is selected from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof; wherein R^2 represents mono-, di-, or tri-substitution, wherein each R^2 is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^2 may optionally combine to form a ring, which can be further substituted; wherein R^3 represents mono-, di-, tri-, or tetra-substitution, wherein each R^3 is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^3 may optionally combine to form a ring, which can be further substituted; wherein R^4 and R^5 are independently selected


4

from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein R^4 and R^5 may optionally combine to form a ring, which can be further substituted; wherein R^1 may optionally combine with any R^3 to form a ring system, which can be further substituted; wherein any of R^4 or R^5 may optionally combine with any R^2 or R^3 to form a ring system, which can be further substituted; and wherein any of R^1 , R^2 , or R^3 may optionally combine with one or more L^x to form a bidentate, tridentate, or tetradentate ligand.

In some embodiments, M^1 is platinum(II), palladium(II), or gold(III). In some other embodiments, M^1 is platinum (II).

In some embodiments, G^1 is O, $C(CH_3)_2$, or $C(C_6H_5)_2$.

In some embodiments, the metal complex is a compound having Formula (II):




wherein rings A^1 , A^2 , and A^3 are independently five- or six-membered carbocyclic or heterocyclic aromatic rings having 0-3 nitrogen atoms and 0-1 additional heteroatoms selected from the group consisting of oxygen, sulfur, and selenium; Z^1 and Z^2 are independently carbon or nitrogen; wherein R^2 represents mono- or di-substitution, wherein each R^2 is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^2 may optionally combine to form a ring, which can be further substituted; wherein R^3 represents mono-, di-, tri-, or tetra-substitution, wherein each R^3 is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^3 may optionally combine to form a ring, which can be further substituted; wherein R^6 represents mono-, di-, or tri-substitution, wherein each R^6 is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^6 may optionally combine to form a ring, which can be further substituted; wherein R^7 and R^8 are independently selected

5


phino, and combinations thereof, and wherein any two adjacent R^6 may optionally combine to form a ring, which can be further substituted; wherein R^7 and R^8 represent mono-, di-, tri-, or tetra-substitution, wherein each R^7 or R^8 is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^7 or any two adjacent R^8 may optionally combine to form a ring, which can be further substituted; wherein any R^3 may optionally combine with any R^6 to form a ring system, which can be further substituted; wherein any R^2 may optionally combine with any R^8 to form a ring system, which can be further substituted; and wherein any R^6 may optionally combine with any R^7 to form a ring system, which can be further substituted; wherein any R^7 may optionally combine with any R^8 to form a ring system, which can be further substituted. The other variables have the definitions provided in the above embodiments.

In some such embodiments, the metal complex is a compound having Formula (IIa):




In some other such embodiments, the metal complex is a compound selected from the group consisting of:

Compound 1


**6**

-continued


Compound 2



Compound 3







Compound 4



7

-continued



8


-continued

Compound 5

5

10

15




Compound 6

20

25

30



Compound 9

Compound 10


Compound 11

Compound 7

35

40

45



and


Compound 8

50

55

60


65



Compound 12

9

In some embodiments, the metal complex is a compound having Formula (IV):




wherein rings A⁴, A⁵, and A⁶ are independently five- or six-membered carbocyclic or heterocyclic aromatic rings having 0-3 nitrogen atoms and 0-1 additional heteroatoms selected from the group consisting of oxygen, sulfur, and selenium; wherein Z³, Z⁴, and Z⁵ are independently carbon or nitrogen; wherein G² is oxygen, sulfur, CR¹²R^{12a}, SiR¹²R^{12a}, or NR¹²; wherein each R¹² and R^{12a} is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein R¹² and R^{12a} may optionally combine with each other or with any one R¹⁰ or R¹¹ to form a ring system, which can be further substituted; wherein R⁹ and R¹⁰ represent mono-, di-, tri-, or tetra-substitution, wherein each R⁹ or R¹⁰ is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R⁹ or any two adjacent R¹⁰ may optionally combine to form a ring, which can be further substituted; wherein R¹¹ represents mono-, di-, or tri-substitution, wherein each R¹¹ is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R¹¹ may optionally combine to form a ring, which can be further substituted; wherein any R³ may optionally combine with any R¹¹ to form a ring system, which can be further substituted; and wherein any R² may optionally combine with any R⁹ to form a ring system, which can be further substituted. The other variables have the definitions provided in the above embodiments.

In some such embodiments, the metal complex is a compound having Formula (IVa):


10

(IVa)




In some embodiments where the metal complex has a Formula (IV) or Formula (IVa), G² is NR¹², and R¹² is phenyl. In some other embodiments, the metal complex is a compound selected from the group consisting of:


Compound 13



Compound 14




Compound 15



11

-continued

Compound 16



5


10

15

12

-continued

Compound 20




20

25

30

35

Compound 17




40

45

50

Compound 18




55


60

65


Compound 19




Compound 21



Compound 22




Compound 23




13


-continued



Compound 24



Compound 25




Compound 26

In some embodiments, the metal complex is a compound having Formula (V):

14


(V)



wherein rings A^7 , A^8 , and A^9 are independently five- or six-membered carbocyclic or heterocyclic aromatic rings having 0-3 nitrogen atoms and 0-1 additional heteroatoms selected from the group consisting of oxygen, sulfur, and selenium; wherein Z^6 and Z^7 independently are carbon or nitrogen; G^3 is oxygen, sulfur, $CR^{16}R^{16a}$, $SiR^{16}R^{16a}$, or NR^{16} ; wherein each R^{16} and R^{16a} is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein R^{16} and R^{16a} may optionally combine with each other or with any one R^{13} or R^{14} to form a ring system, which can be further substituted; wherein R^{13} and R^{14} represent mono-, di-, or tri-substitution, wherein each R^{13} or R^{14} is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^{13} or any two adjacent R^{14} may optionally combine to form a ring, which can be further substituted; wherein R^{15} represents mono-, di-, tri- or tetra-substitution, wherein each R^{15} is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^{15} may optionally combine to form a ring, which can be further substituted; wherein any R^2 may optionally combine with any R^{13} to form a ring system, which can be further substituted; and wherein any R^{14} may optionally combine with any R^{15} to form a ring system, which can be further substituted. The other variables have the definitions provided in the above embodiments.


In some such embodiments, the metal complex is a compound having Formula (Va):

15




In some embodiments where the metal complex has a Formula (V) or Formula (Va), G^3 is NR^{16} , and R^{16} is phenyl.

In some other embodiments, the metal complex is a compound selected from the group consisting of:



Compound 27



Compound 28

16

-continued

(Va)

5

10

15

20

25

30

35

40


45

50


55

60


65




Compound 29



Compound 30




Compound 31



Compound 32


17

-continued



Compound 33

In some embodiments, the metal complex is a compound having Formula (VI):



(VI)


wherein rings A⁷, A⁸, and A⁹ are independently five- or six-membered carbocyclic or heterocyclic aromatic rings having 0-3 nitrogen atoms and 0-1 additional heteroatoms selected from the group consisting of oxygen, sulfur, and selenium; wherein Z⁸ and Z⁹ are independently carbon or nitrogen; wherein G⁴ is oxygen, sulfur, CR²⁰R^{20a}, SiR²⁰R^{20a}, or NR²⁰; wherein each R²⁰ and R^{20a} is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein R²⁰ and R^{20a} may optionally combine with each other or with any one R¹⁸ or R¹⁹ to form a ring system, which can be further substituted; wherein G⁵ is oxygen, sulfur, CR²¹R^{21a}, SiR²¹R^{21a}, or NR²¹; wherein each R²¹ and R^{21a} is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein R²¹ and R^{21a} may optionally combine with each other or with any one R² or R¹⁷ to form a ring system, which can be further substituted; wherein R¹⁷ and R¹⁸ represent mono-, di-, tri-, or tetra-substitution, wherein each R¹⁷ or R¹⁸ is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl,

18


carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R¹⁷ or any two adjacent R¹⁸ may optionally combine to form a ring, which can be further substituted; wherein R¹⁹ represents mono-, di-, or tri-substitution, wherein each R¹⁹ is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R¹⁹ may optionally combine to form a ring, which can be further substituted; wherein any R³ may optionally combine with any R¹⁹ to form a ring system, which can be further substituted. The other variables have the definitions provided in the above embodiments.

In some other embodiments, the metal complex is a compound selected from the group consisting of:


Compound 34



Compound 35




Compound 36




19

-continued


Compound 37




Compound 38



Compound 39




Compound 40



20

In some embodiments, the metal complex is a compound having Formula (VII):



20 wherein each L^y is independently a monodentate ligand, and any two adjacent L^y may optionally combine to form a bidentate ligand; wherein M^2 is cobalt(I), rhodium(I), iridium(I), nickel(II), platinum(II), palladium(II), silver(III), gold(III), or copper(III); wherein p is a value from 1 to the maximum

25 number of ligands that may be attached to M^2 ; wherein $p+q$ is the maximum number of ligands that may be attached to M^2 ; wherein G^6 is O or $CR^{24}R^{25}$; wherein R^{21} is selected from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof; wherein R^{22} represents mono-, di-, or tri-

30 substitution, wherein each R^{22} is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^2 may optionally combine to form a ring, which can be further substituted; wherein R^{23} represents mono-, di-, tri-, or tetra-

35 substitution, wherein each R^{23} is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^3 may

40 optionally combine to form a ring, which can be further substituted; wherein R^{24} and R^{25} are independently selected from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein R^{24} and R^{25} may optionally combine to form a ring, which can be further substituted;

45 wherein R^{21} may optionally combine with any R^{23} to form a ring system, which can be further substituted; wherein any of R^{24} or R^{25} may optionally combine with any R^{22} or R^{23} to form a ring system, which can be further substituted; and wherein any of R^{21} , R^{22} , or R^{23} may optionally combine with one or more L^y to form a bidentate, tridentate, or tetradentate

50 ligand.

55

60


65

21


In some embodiments, M^2 is platinum(II), palladium(II), or gold(III). In some other embodiments, M^2 is platinum (II).

In some embodiments, G^6 is O, $C(CH_3)_2$, or $C(C_6H_5)_2$.


In some other embodiments, the metal complex is a compound selected from the group consisting of:




Compound 41



Compound 42




Compound 43




Compound 44

22


-continued




Compound 45



Compound 46




Compound 47



Compound 48

23

-continued



Compound 49


5

10

15

Compound 50

20




25

30

35

Compound 51




40

45

50

Compound 52




55

60


65

24


-continued




Compound 53



Compound 54



Compound 55




Compound 56

and

25


-continued



Compound 57

26

-continued








In another aspect, the invention provides devices that include metal complexes having a bridged N,N-diaryl ligand, such as those described in the foregoing paragraphs. In some embodiments, the invention provides a first device comprising a first organic light emitting device, which further comprises: an anode; a cathode; and an organic layer disposed between the anode and the cathode, which comprises a boron-nitrogen heterocycle according to any of the above embodiments. In some embodiments, the first device is a consumer product. In some embodiments, the first device is an organic light emitting device (OLED). In some embodiments, the first device comprises a lighting panel.


In some embodiments, the organic layer of the first device is an emissive layer. In some such embodiments, the emissive layer comprises an emissive dopant. In some embodiments, a metal complex having a bridged N,N-diaryl ligand (as described above) is an emissive dopant. In some other embodiments, a metal complex having a bridged N,N-diaryl ligand (as described above) is a non-emissive dopant.

In some embodiments, the organic layer of the first device comprises a host. In some such embodiments, the host comprises a triphenylene containing benzo-fused thiophene or benzo-fused furan; wherein any substituent in the host is an unfused substituent independently selected from the group consisting of C_nH_{2n+1} , OC_nH_{2n+1} , OAr_1 , $N(C_nH_{2n+1})_2$, $N(Ar_1)(Ar_2)$, $CH=CH-C_nH_{2n+1}$, $C\equiv C-C_nH_{2n+1}$, Ar_1 , Ar_1-Ar_2 , and $C_nH_{2n}-Ar_1$, or the host has no substitution; wherein n is from 1 to 10; and wherein Ar_1 and Ar_2 are independently selected from the group consisting of benzene, biphenyl, naphthalene, triphenylene, carbazole, and heteroaromatic analogs thereof. In some other embodiments, the host comprises a compound selected from the group consisting of: carbazole, dibenzothiophene, dibenzofuran, dibenzoselenophene, azacarbazole, aza-dibenzothiophene, aza-dibenzofuran, and aza-dibenzoselenophene.

In some embodiments, the host is a compound selected from the group consisting of:



27



and combinations thereof.

In some embodiments, the host is a metal complex.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows an organic light emitting device.

FIG. 2 shows an inverted organic light emitting device that does not have a separate electron transport layer.

FIG. 3 shows an embodiment of a metal complex having a bridged N,N-diaryl ligand.

FIG. 4 shows a positive-mode MALDI-TOF mass spectrum for Host 1 and Comparative Example 3. The arrow shows the major fragment, which is the loss of N-phenyl. The inset shows the structure and mass of the bond-broken photoproduct.

FIG. 5 shows a positive-mode MALDI-TOF mass spectrum for Host 1 and Comparative Example 2. The arrow

28

shows the major fragment, which is the loss of N-phenyl. The inset shows the structure and mass of the bond-broken photoproduct.

FIG. 6 shows a positive-mode MALDI-TOF mass spectrum for Host 1 and Comparative Example 1. The arrow shows the major fragment, which is the loss of N-phenyl. The inset shows the structure and mass of the bond-broken photoproduct.

FIG. 7 shows the geometry for Compound 3, which shows the N-phenyl bond intact. The left structure shows the optimized ground state triplet. The right structure shows an optimized N-phenyl bond-breaking triplet. Slight differences in the geometry of the phenyl and geminal dimethyl groups can be seen, which may account for the small calculated energy difference of 5.22 kcal/mol.

DETAILED DESCRIPTION

Generally, an OLED comprises at least one organic layer disposed between and electrically connected to an anode and a cathode. When a current is applied, the anode injects holes and the cathode injects electrons into the organic layer(s). The injected holes and electrons each migrate toward the oppositely charged electrode. When an electron and hole localize on the same molecule, an “exciton,” which is a localized electron-hole pair having an excited energy state, is formed. Light is emitted when the exciton relaxes via a photoemissive mechanism. In some cases, the exciton may be localized on an excimer or an exciplex. Non-radiative mechanisms, such as thermal relaxation, may also occur, but are generally considered undesirable.

The initial OLEDs used emissive molecules that emitted light from their singlet states (“fluorescence”) as disclosed, for example, in U.S. Pat. No. 4,769,292, which is incorporated by reference in its entirety. Fluorescent emission generally occurs in a time frame of less than 10 nanoseconds.

More recently, OLEDs having emissive materials that emit light from triplet states (“phosphorescence”) have been demonstrated. Baldo et al., “Highly Efficient Phosphorescent Emission from Organic Electroluminescent Devices,” *Nature*, vol. 395, 151-154, 1998; (“Baldo-I”) and Baldo et al., “Very high-efficiency green organic light-emitting devices based on electrophosphorescence,” *Appl. Phys. Lett.*, vol. 75, No. 3, 4-6 (1999) (“Baldo-II”), which are incorporated by reference in their entireties. Phosphorescence is described in more detail in U.S. Pat. No. 7,279,704 at cols. 5-6, which are incorporated by reference.

FIG. 1 shows an organic light emitting device 100. The figures are not necessarily drawn to scale. Device 100 may include a substrate 110, an anode 115, a hole injection layer 120, a hole transport layer 125, an electron blocking layer 130, an emissive layer 135, a hole blocking layer 140, an electron transport layer 145, an electron injection layer 150, a protective layer 155, a cathode 160, and a barrier layer 170. Cathode 160 is a compound cathode having a first conductive layer 162 and a second conductive layer 164. Device 100 may be fabricated by depositing the layers described, in order. The properties and functions of these various layers, as well as example materials, are described in more detail in U.S. Pat. No. 7,279,704 at cols. 6-10, which are incorporated by reference.

More examples for each of these layers are available. For example, a flexible and transparent substrate-anode combination is disclosed in U.S. Pat. No. 5,844,363, which is incorporated by reference in its entirety. An example of a p-doped hole transport layer is m-MTDATA doped with F.sub.4-TCNQ at a molar ratio of 50:1, as disclosed in U.S. Patent

Application Publication No. 2003/0230980, which is incorporated by reference in its entirety. Examples of emissive and host materials are disclosed in U.S. Pat. No. 6,303,238 to Thompson et al., which is incorporated by reference in its entirety. An example of an n-doped electron transport layer is BPhen doped with Li at a molar ratio of 1:1, as disclosed in U.S. Patent Application Publication No. 2003/0230980, which is incorporated by reference in its entirety. U.S. Pat. Nos. 5,703,436 and 5,707,745, which are incorporated by reference in their entireties, disclose examples of cathodes including compound cathodes having a thin layer of metal such as Mg:Ag with an overlying transparent, electrically-conductive, sputter-deposited ITO layer. The theory and use of blocking layers is described in more detail in U.S. Pat. No. 6,097,147 and U.S. Patent Application Publication No. 2003/0230980, which are incorporated by reference in their entireties. Examples of injection layers are provided in U.S. Patent Application Publication No. 2004/0174116, which is incorporated by reference in its entirety. A description of protective layers may be found in U.S. Patent Application Publication No. 2004/0174116, which is incorporated by reference in its entirety.

FIG. 2 shows an inverted OLED **200**. The device includes a substrate **210**, a cathode **215**, an emissive layer **220**, a hole transport layer **225**, and an anode **230**. Device **200** may be fabricated by depositing the layers described, in order. Because the most common OLED configuration has a cathode disposed over the anode, and device **200** has cathode **215** disposed under anode **230**, device **200** may be referred to as an "inverted" OLED. Materials similar to those described with respect to device **100** may be used in the corresponding layers of device **200**. FIG. 2 provides one example of how some layers may be omitted from the structure of device **100**.

The simple layered structure illustrated in FIGS. 1 and 2 is provided by way of non-limiting example, and it is understood that embodiments of the invention may be used in connection with a wide variety of other structures. The specific materials and structures described are exemplary in nature, and other materials and structures may be used. Functional OLEDs may be achieved by combining the various layers described in different ways, or layers may be omitted entirely, based on design, performance, and cost factors. Other layers not specifically described may also be included. Materials other than those specifically described may be used. Although many of the examples provided herein describe various layers as comprising a single material, it is understood that combinations of materials, such as a mixture of host and dopant, or more generally a mixture, may be used. Also, the layers may have various sublayers. The names given to the various layers herein are not intended to be strictly limiting. For example, in device **200**, hole transport layer **225** transports holes and injects holes into emissive layer **220**, and may be described as a hole transport layer or a hole injection layer. In one embodiment, an OLED may be described as having an "organic layer" disposed between a cathode and an anode. This organic layer may comprise a single layer, or may further comprise multiple layers of different organic materials as described, for example, with respect to FIGS. 1 and 2.

Structures and materials not specifically described may also be used, such as OLEDs comprised of polymeric materials (PLEDs) such as disclosed in U.S. Pat. No. 5,247,190 to Friend et al., which is incorporated by reference in its entirety. By way of further example, OLEDs having a single organic layer may be used. OLEDs may be stacked, for example as described in U.S. Pat. No. 5,707,745 to Forrest et al, which is incorporated by reference in its entirety. The OLED structure may deviate from the simple layered structure illustrated in

FIGS. 1 and 2. For example, the substrate may include an angled reflective surface to improve out-coupling, such as a mesa structure as described in U.S. Pat. No. 6,091,195 to Forrest et al., and/or a pit structure as described in U.S. Pat. No. 5,834,893 to Bulovic et al., which are incorporated by reference in their entireties.

Unless otherwise specified, any of the layers of the various embodiments may be deposited by any suitable method. For the organic layers, preferred methods include thermal evaporation, ink-jet, such as described in U.S. Pat. Nos. 6,013,982 and 6,087,196, which are incorporated by reference in their entireties, organic vapor phase deposition (OVPD), such as described in U.S. Pat. No. 6,337,102 to Forrest et al., which is incorporated by reference in its entirety, and deposition by organic vapor jet printing (OVJP), such as described in U.S. patent application Ser. No. 10/233,470, now U.S. Pat No 7,431,968, which is incorporated by reference in its entirety. Other suitable deposition methods include spin coating and other solution based processes. Solution based processes are preferably carried out in nitrogen or an inert atmosphere. For the other layers, preferred methods include thermal evaporation. Preferred patterning methods include deposition through a mask, cold welding such as described in U.S. Pat. Nos. 6,294,398 and 6,468,819, which are incorporated by reference in their entireties, and patterning associated with some of the deposition methods such as ink jet and OVJD. Other methods may also be used. The materials to be deposited may be modified to make them compatible with a particular deposition method. For example, substituents such as alkyl and aryl groups, branched or unbranched, and preferably containing at least 3 carbons, may be used in small molecules to enhance their ability to undergo solution processing. Substituents having 20 carbons or more may be used, and 3-20 carbons is a preferred range. Materials with asymmetric structures may have better solution processability than those having symmetric structures, because asymmetric materials may have a lower tendency to recrystallize. Dendrimer substituents may be used to enhance the ability of small molecules to undergo solution processing.

Devices fabricated in accordance with embodiments of the present invention may further optionally comprise a barrier layer. One purpose of the barrier layer is to protect the electrodes and organic layers from damaging exposure to harmful species in the environment including moisture, vapor and/or gases, etc. The barrier layer may be deposited over, under or next to a substrate, an electrode, or over any other parts of a device including an edge. The barrier layer may comprise a single layer, or multiple layers. The barrier layer may be formed by various known chemical vapor deposition techniques and may include compositions having a single phase as well as compositions having multiple phases. Any suitable material or combination of materials may be used for the barrier layer. The barrier layer may incorporate an inorganic or an organic compound or both. The preferred barrier layer comprises a mixture of a polymeric material and a non-polymeric material as described in U.S. Pat. No. 7,968,146, PCT Pat. Application Nos. PCT/US2007/023098 and PCT/US2009/042829, which are herein incorporated by reference in their entireties. To be considered a "mixture", the aforesaid polymeric and non-polymeric materials comprising the barrier layer should be deposited under the same reaction conditions and/or at the same time. The weight ratio of polymeric to non-polymeric material may be in the range of 95:5 to 5:95. The polymeric material and the non-polymeric material may be created from the same precursor material. In one example,

31

the mixture of a polymeric material and a non-polymeric material consists essentially of polymeric silicon and inorganic silicon.


Devices fabricated in accordance with embodiments of the invention may be incorporated into a wide variety of consumer products, including flat panel displays, computer monitors, medical monitors, televisions, billboards, lights for interior or exterior illumination and/or signaling, heads up displays, fully transparent displays, flexible displays, laser printers, telephones, cell phones, personal digital assistants (PDAs), laptop computers, digital cameras, camcorders, viewfinders, micro-displays, vehicles, a large area wall, theater or stadium screen, or a sign. Various control mechanisms may be used to control devices fabricated in accordance with the present invention, including passive matrix and active matrix. Many of the devices are intended for use in a temperature range comfortable to humans, such as 18 degrees C. to 30 degrees C., and more preferably at room temperature (20-25 degrees C.).

The materials and structures described herein may have applications in devices other than OLEDs. For example, other optoelectronic devices such as organic solar cells and organic photodetectors may employ the materials and structures. More generally, organic devices, such as organic transistors, may employ the materials and structures.

The terms halo, halogen, alkyl, cycloalkyl, alkenyl, alkynyl, arylkyl, heterocyclic group, aryl, aromatic group, and heteroaryl are known to the art, and are defined in U.S. Pat. No. 7,279,704 at cols. 31-32, which are incorporated herein by reference.

Metal complexes having ligands with N,N-diaryl groups can function as emissive agents, e.g., phosphorescent emitters. In some instances, however, such structures may exhibit instability due to rupture of the N-aryl bond, thereby limiting the utility of such compounds as emissive agents. In some embodiments of the invention, a bridging group is incorporated into the N,N-diaryl ligand structure to stabilize the ligand and strengthen the N-aryl bond. The disclosed bridging structures can also improve the steric properties of the ligand and make it more suitable for certain types of coordination, such as tetradentate coordination in a square planar configuration (e.g., with platinum (II)).

In at least one aspect, the invention provides metal complexes having an N,N-diaryl ligand, such as compounds having Formula (I):



wherein each L^x is independently a monodentate ligand, and any two adjacent L^x may optionally combine to form a bidentate ligand; wherein M^1 is cobalt(I), rhodium(I), iridium(I), nickel(II), platinum(II), palladium(II), silver(III), gold(III), or copper(III); wherein m is a value from 1 to the maximum number of ligands that may be attached to M^1 ; wherein $m+n$ is the maximum number of ligands that may be attached to


32

M^1 ; wherein G^1 is O or CR^4R^5 ; wherein R^1 is selected from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof; wherein R^2 represents mono-, di-, or tri-substitution, wherein each R^2 is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^2 may optionally combine to form a ring, which can be further substituted; wherein R^3 represents mono-, di-, tri-, or tetra-substitution, wherein each R^3 is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^3 may optionally combine to form a ring, which can be further substituted; wherein R^4 and R^5 are independently selected from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein R^4 and R^5 may optionally combine to form a ring, which can be further substituted; wherein R^1 may optionally combine with any R^3 to form a ring system, which can be further substituted; wherein any of R^4 or R^5 may optionally combine with any R^2 or R^3 to form a ring system, which can be further substituted; and wherein any of R^1 , R^2 , or R^3 may optionally combine with one or more L^x to form a bidentate, tridentate, or tetradentate ligand.

In some embodiments, M^1 is platinum(II), palladium(II), or gold(III). In some other embodiments, M^1 is platinum (II).

In some embodiments, G^1 is O, $C(CH_3)_2$, or $C(C_6H_5)_2$.

In some embodiments, the metal complex is a compound having Formula (II):




wherein rings A^1 , A^2 , and A^3 are independently five- or six-membered carbocyclic or heterocyclic aromatic rings having 0-3 nitrogen atoms and 0-1 additional heteroatoms selected from the group consisting of oxygen, sulfur, and selenium; Z^1 and Z^2 are independently carbon or nitrogen; wherein R^2 represents mono- or di-substitution, wherein each R^2 is

33

selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^2 may optionally combine to form a ring, which can be further substituted; wherein R^3 represents mono-, di-, tri-, or tetra-substitution, wherein each R^3 is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^3 may optionally combine to form a ring, which can be further substituted; wherein R^6 represents mono-, di-, or tri-substitution, wherein each R^6 is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^6 may optionally combine to form a ring, which can be further substituted; wherein R^7 and R^8 represent mono-, di-, tri-, or tetra-substitution, wherein each R^7 or R^8 is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^7 or any two adjacent R^8 may optionally combine to form a ring, which can be further substituted; wherein any R^3 may optionally combine with any R^6 to form a ring system, which can be further substituted; wherein any R^2 may optionally combine with any R^8 to form a ring system, which can be further substituted; and wherein any R^6 may optionally combine with any R^7 to form a ring system, which can be further substituted; wherein any R^7 may optionally combine with any R^8 to form a ring system, which can be further substituted. The other variables have the definitions provided in the above embodiments.


In some such embodiments, the metal complex is a compound having Formula (IIa):




In some other such embodiments, the metal complex is a compound selected from the group consisting of:

34

Compound 1




Compound 2







Compound 3

(IIa)



35

-continued



36


-continued

Compound 4

5

10

15




Compound 8

Compound 5

25

30

35




Compound 9

Compound 6

40

45

50




Compound 10

Compound 7

55

60

65




and


37

-continued

Compound 12



In some embodiments, the metal complex is a compound having Formula (IV):




wherein rings A^4 , A^5 , and A^6 are independently five- or six-membered carbocyclic or heterocyclic aromatic rings having 0-3 nitrogen atoms and 0-1 additional heteroatoms selected from the group consisting of oxygen, sulfur, and selenium; wherein Z^3 , Z^4 , and Z^5 are independently carbon or nitrogen; wherein G^2 is oxygen, sulfur, $CR^{12}R^{12a}$, $SiR^{12}R^{12a}$, or NR^{12} ; wherein each R^{12} and R^{12a} is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein R^{12} and R^{12a} may optionally combine with each other or with any one R^{10} or R^{11} to form a ring system, which can be further substituted; wherein R^9 and R^{10} represent mono-, di-, tri-, or tetra-substitution, wherein each R^9 or R^{10} is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^9 or any two adjacent R^{10} may optionally combine to form a ring, which can be further substituted; wherein R^{11} represents mono-, di-, or tri-substitution, wherein each R^{11} is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino,

38

silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^{11} may optionally combine to form a ring, which can be further substituted; wherein any R^3 may optionally combine with any R^{11} to form a ring system, which can be further substituted; and wherein any R^2 may optionally combine with any R^9 to form a ring system, which can be further substituted. The other variables have the definitions provided in the above embodiments.


In some such embodiments, the metal complex is a compound having Formula (IVa):




In some embodiments where the metal complex has a Formula (IV) or Formula (IVa), G^2 is NR^{12} , and R^{12} is phenyl.

In some other embodiments, the metal complex is a compound selected from the group consisting of:

Compound 13




Compound 14



39

-continued

Compound 15




5

10

15

Compound 16




20

25

30

Compound 17 35




40

45

50

Compound 18



55


60

65


40

-continued


Compound 19




Compound 20



Compound 21




Compound 22




41


-continued




Compound 23



Compound 24




Compound 25



Compound 26

42


In some embodiments, the metal complex is a compound having Formula (V):



wherein rings A^7 , A^8 , and A^9 are independently five- or six-membered carbocyclic or heterocyclic aromatic rings having 0-3 nitrogen atoms and 0-1 additional heteroatoms selected from the group consisting of oxygen, sulfur, and selenium; wherein Z^6 and Z^7 independently are carbon or nitrogen; G^3 is oxygen, sulfur, $CR^{16}R^{16a}$, $SiR^{16}R^{16a}$, or NR^{16} ; wherein each R^{16} and R^{16a} is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein R^{16} and R^{16a} may optionally combine with each other or with any one R^{13} or R^{14} to form a ring system, which can be further substituted; wherein R^{13} and R^{14} represent mono-, di-, or tri-substitution, wherein each R^{13} or R^{14} is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^{13} or any two adjacent R^{14} may optionally combine to form a ring, which can be further substituted; wherein R^{15} represents mono-, di-, tri- or tetra-substitution, wherein each R^{15} is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^{15} may optionally combine to form a ring, which can be further substituted; wherein any R^2 may optionally combine with any R^{13} to form a ring system, which can be further substituted; and wherein any R^{14} may optionally combine with any R^{15} to form a ring system, which can be further substituted. The other variables have the definitions provided in the above embodiments.

In some such embodiments, the metal complex is a compound having Formula (Va):


43



In some embodiments where the metal complex has a Formula (V) or Formula (Va), G^3 is NR^{16} , and R^{16} is phenyl.

In some other embodiments, the metal complex is a compound selected from the group consisting of:


Compound 27



25

30


Compound 28



40

45

Compound 29



55

60

65

44

-continued


(Va)

Compound 30

5

10

15




Compound 31

20

25

30




Compound 32

35

40

45

Compound 28



, and


Compound 33

50

55


60

65



In some embodiments, the metal complex is a compound having Formula (VI):

45




wherein rings A⁷, A⁸, and A⁹ are independently five- or six-membered carbocyclic or heterocyclic aromatic rings having 0-3 nitrogen atoms and 0-1 additional heteroatoms selected from the group consisting of oxygen, sulfur, and selenium; wherein Z⁸ and Z⁹ are independently carbon or nitrogen; wherein G⁴ is oxygen, sulfur, CR²⁰R^{20a}, SiR²⁰R^{20a}, or NR²⁰; wherein each R²⁰ and R^{20a} is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein R²⁰ and R^{20a} may optionally combine with each other or with any one R¹⁸ or R¹⁹ to form a ring system, which can be further substituted; wherein G⁵ is oxygen, sulfur, CR²¹R^{21a}, SiR²¹R^{21a}, or NR²¹; wherein each R²¹ and R^{21a} is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein R²¹ and R^{21a} may optionally combine with each other or with any one R² or R¹⁷ to form a ring system, which can be further substituted; wherein R¹⁷ and R¹⁸ represent mono-, di-, tri-, or tetra-substitution, wherein each R¹⁷ or R¹⁸ is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R¹⁷ or any two adjacent R¹⁸ may optionally combine to form a ring, which can be further substituted; wherein R¹⁹ represents mono-, di-, or tri-substitution, wherein each R¹⁹ is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R¹⁹ may optionally combine to form a ring, which can be further substituted; wherein any R³ may optionally combine with any R¹⁹ to form a ring system, which can be further substituted. The other variables have the definitions provided in the above embodiments.


In some other embodiments, the metal complex is a compound selected from the group consisting of:

46


(VI)




Compound 34



Compound 35



Compound 36




Compound 37


47

-continued


Compound 38




Compound 39



Compound 40



In some embodiments, the metal complex is a compound having Formula (VII):



(VII)

48


wherein each L^y is independently a monodentate ligand, and any two adjacent L^y may optionally combine to form a bidentate ligand; wherein M^2 is cobalt(I), rhodium(I), iridium(I), nickel(II), platinum(II), palladium(II), silver(III), gold(III), or copper(III); wherein p is a value from 1 to the maximum number of ligands that may be attached to M^2 ; wherein $p+q$ is the maximum number of ligands that may be attached to M^2 ; wherein G^6 is O or $CR^{24}R^{25}$; wherein R^{21} is selected from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof; wherein R^{22} represents mono-, di-, or tri-substitution, wherein each R^{22} is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^2 may optionally combine to form a ring, which can be further substituted; wherein R^{23} represents mono-, di-, tri-, or tetra-substitution, wherein each R^{23} is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^3 may optionally combine to form a ring, which can be further substituted; wherein R^{24} and R^{25} are independently selected from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein R^{24} and R^{25} may optionally combine to form a ring, which can be further substituted; wherein R^{21} may optionally combine with any R^{23} to form a ring system, which can be further substituted; wherein any of R^{24} or R^{25} may optionally combine with any R^{22} or R^{23} to form a ring system, which can be further substituted; and wherein any of R^{21} , R^{22} , or R^{23} may optionally combine with one or more L^y to form a bidentate, tridentate, or tetradentate ligand.

In some embodiments, M^2 is platinum(II), palladium(II), or gold(III). In some other embodiments, M^2 is platinum(II).

In some embodiments, G^6 is O, $C(CH_3)_2$, or $C(C_6H_5)_2$.


In some other embodiments, the metal complex is a compound selected from the group consisting of:

Compound 41



49

-continued



Compound 42


5

10

15

50

-continued




Compound 46

Compound 43


20

25

30



Compound 47



Compound 44


35

40

45

50

Compound 48





Compound 45

55

60


65

Compound 49



51

-continued



Compound 50

52


-continued

5

10

15

20




Compound 53

25

Compound 51

30


35



40

45

50



Compound 54



Compound 55

55

Compound 52

60

65




Compound 56

and

53

-continued




In another aspect, the invention provides devices that include metal complexes having a bridged N,N-diaryl ligand, such as those described in the foregoing paragraphs. In some embodiments, the invention provides a first device comprising a first organic light emitting device, which further comprises: an anode; a cathode; and an organic layer disposed between the anode and the cathode, which comprises a boron-nitrogen heterocycle according to any of the above embodiments. In some embodiments, the first device is a consumer product. In some embodiments, the first device is an organic light emitting device (OLED). In some embodiments, the first device comprises a lighting panel.

In some embodiments, the organic layer of the first device is an emissive layer. In some such embodiments, the emissive layer comprises an emissive dopant. In some embodiments, a metal complex having a bridged N,N-diaryl ligand (as described above) is an emissive dopant. In some other embodiments, a metal complex having a bridged N,N-diaryl ligand (as described above) is a non-emissive dopant.

In some embodiments, the organic layer of the first device comprises a host. In some such embodiments, the host comprises a triphenylene containing benzo-fused thiophene or benzo-fused furan; wherein any substituent in the host is an unfused substituent independently selected from the group consisting of C_nH_{2n+1} , OC_nH_{2n+1} , OAr_1 , $N(C_nH_{2n+1})_2$, $N(Ar_1)(Ar_2)$, $CH=CH-C_nH_{2n+1}$, $C\equiv C-C_nH_{2n+1}$, Ar_1 , Ar_1-Ar_2 , and $C_nH_{2n}-Ar_1$, or the host has no substitution; wherein n is from 1 to 10; and wherein Ar_1 and Ar_2 are independently selected from the group consisting of benzene, biphenyl, naphthalene, triphenylene, carbazole, and heteroaromatic analogs thereof. In some other embodiments, the host comprises a compound selected from the group consisting of: carbazole, dibenzothiophene, dibenzofuran, dibenzoselenophene, azacarbazole, aza-dibenzothiophene, aza-dibenzofuran, and aza-dibenzoselenophene.

In some embodiments, the host is a compound selected from the group consisting of:



Compound 57

5

10

15

20

25

30

35

40

45

50






55

60


65

54

-continued



55



and combinations thereof.

In some embodiments, the host is a metal complex.

Combination with Other Materials


The materials described herein as useful for a particular layer in an organic light emitting device may be used in combination with a wide variety of other materials present in the device. For example, emissive dopants disclosed herein may be used in conjunction with a wide variety of hosts, transport layers, blocking layers, injection layers, electrodes and other layers that may be present. The materials described or referred to below are non-limiting examples of materials that may be useful in combination with the compounds disclosed herein, and one of skill in the art can readily consult the literature to identify other materials that may be useful in combination.

56

HIL/HTL:

A hole injecting/transporting material to be used in the present invention is not particularly limited, and any compound may be used as long as the compound is typically used as a hole injecting/transporting material. Examples of the material include, but not limit to: a phthalocyanine or porphyrin derivative; an aromatic amine derivative; an indolocarbazole derivative; a polymer containing fluorohydrocarbon; a polymer with conductivity dopants; a conducting polymer, such as PEDOT/PSS; a self-assembly monomer derived from compounds such as phosphonic acid and silane derivatives; a metal oxide derivative, such as MoO_x ; a p-type semiconducting organic compound, such as 1,4,5,8,9,12-Hexaazatriphenylenehexacarbonitrile; a metal complex, and a cross-linkable compounds.

Examples of aromatic amine derivatives used in HIL or HTL include, but not limit to the following general structures:




Each of Ar^1 to Ar^9 is selected from the group consisting aromatic hydrocarbon cyclic compounds such as benzene, biphenyl, triphenyl, triphenylene, naphthalene, anthracene, phenalene, phenanthrene, fluorene, pyrene, chrysene, perylene, azulene; group consisting aromatic heterocyclic compounds such as dibenzothiophene, dibenzofuran, dibenzoselenophene, furan, thiophene, benzofuran, benzothiophene, benzoselenophene, carbazole, indolocarbazole, pyridylindole, pyrrolodipyridine, pyrazole, imidazole, triazole, oxazole, thiazole, oxadiazole, oxatriazole, dioxazole, thiadiazole, pyridine, pyridazine, pyrimidine, pyrazine, triazine, oxazine, oxathiazine, oxadiazine, indole, benzimidazole, indazole, indoxazine, benzoxazole, benzisoxazole, benzothiazole, quinoline, isoquinoline, cinnoline, quinazoline, quinoxaline, naphthyridine, phthalazine, pteridine, xanthene, acridine, phenazine, phenothiazine, phenoxazine, benzofuro-pyridine, furodipyridine, benzothienopyridine, thienodipyridine, benzoselenophenopyridine, and selenophenodipyridine; and group consisting 2 to 10 cyclic structural units which are groups of the same type or different types selected from the aromatic hydrocarbon cyclic group and the aromatic heterocyclic group and are bonded to each other directly or via at least one of oxygen atom, nitrogen atom, sulfur atom, silicon atom, phosphorus atom, boron atom, chain structural unit and the aliphatic cyclic group. Wherein each Ar is further substituted by a substituent selected from the group consist-

57


ing of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof.

In one aspect, Ar¹ to Ar⁹ is independently selected from the group consisting of:



k is an integer from 1 to 20; X¹⁰¹ to X¹⁰⁸ is C (including CH) or N; Z¹⁰¹ is NAr¹, O, or S; Ar¹ has the same group defined above.

Examples of metal complexes used in HIL or HTL include, but not limit to the following general formula:



Met is a metal; (Y¹⁰¹-Y¹⁰²) is a bidentate ligand, Y¹⁰¹ and Y¹⁰² are independently selected from C, N, O, P, and S; L¹⁰¹ is another ligand; k' is an integer value from 1 to the maximum number of ligands that may be attached to the metal; and k'+k'' is the maximum number of ligands that may be attached to the metal.

In one aspect, (Y¹⁰¹-Y¹⁰²) is a 2-phenylpyridine derivative.

In another aspect, (Y¹⁰¹-Y¹⁰²) is a carbene ligand.

In another aspect, Met is selected from Ir, Pt, Os, and Zn.


In a further aspect, the metal complex has a smallest oxidation potential in solution vs. Fc⁺/Fc couple less than about 0.6 V.

Host:

The light emitting layer of the organic EL device of the present invention preferably contains at least a metal complex as light emitting material, and may contain a host material using the metal complex as a dopant material. Examples of the host material are not particularly limited, and any metal complexes or organic compounds may be used as long as the triplet energy of the host is larger than that of the dopant. While the Table below categorizes host materials as preferred for devices that emit various colors, any host material may be used with any dopant so long as the triplet criteria is satisfied.


58

Examples of metal complexes used as host are preferred to have the following general formula:



Met is a metal; (Y¹⁰³-Y¹⁰⁴) is a bidentate ligand, Y¹⁰³ and Y¹⁰⁴ are independently selected from C, N, O, P, and S; L¹⁰¹ is another ligand; k' is an integer value from 1 to the maximum number of ligands that may be attached to the metal; and k'+k'' is the maximum number of ligands that may be attached to the metal.

In one aspect, the metal complexes are:



(O—N) is a bidentate ligand, having metal coordinated to atoms O and N.





In another aspect, Met is selected from Ir and Pt.

In a further aspect, (Y¹⁰³-Y¹⁰⁴) is a carbene ligand.

Examples of organic compounds used as host are selected from the group consisting aromatic hydrocarbon cyclic compounds such as benzene, biphenyl, triphenyl, triphenylene, naphthalene, anthracene, phenalene, phenanthrene, fluorene, pyrene, chrysene, perylene, azulene; group consisting aromatic heterocyclic compounds such as dibenzothiophene, dibenzofuran, dibenzoselenophene, furan, thiophene, benzofuran, benzothiophene, benzoselenophene, carbazole, indolocarbazole, pyridylindole, pyrrolodipyridine, pyrazole, imidazole, triazole, oxazole, thiazole, oxadiazole, oxatriazole, dioxazole, thiadiazole, pyridine, pyridazine, pyrimidine, pyrazine, triazine, oxazine, oxathiazine, oxadiazine, indole, benzimidazole, indazole, indoxazine, benzoxazole, benzisoxazole, benzothiazole, quinoline, isoquinoline, cinnoline, quinazoline, quinoxaline, naphthyridine, phthalazine, pteridine, xanthene, acridine, phenazine, phenothiazine, phenoxazine, benzofuropyridine, furodipyridine, benzothienopyridine, thienodipyridine, benzoselenophenopyridine, and selenophenodipyridine; and group consisting 2 to 10 cyclic structural units which are groups of the same type or different types selected from the aromatic hydrocarbon cyclic group and the aromatic heterocyclic group and are bonded to each other directly or via at least one of oxygen atom, nitrogen atom, sulfur atom, silicon atom, phosphorus atom, boron atom, chain structural unit and the aliphatic cyclic group. Wherein each group is further substituted by a substituent selected from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof.

In one aspect, host compound contains at least one of the following groups in the molecule:


59




60

-continued

5




10



15

20



25

30 R^{101} to R^{107} is independently selected from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, when it is aryl or heteroaryl, it has the similar definition as Ar's mentioned above.

35 k is an integer from 1 to 20; k''' is an integer from 0 to 20.

40 X^{101} to X^{108} is selected from C (including CH) or N.

Z^{101} and Z^{102} is selected from NR^{101} , O, or S.


HBL:

45 A hole blocking layer (HBL) may be used to reduce the number of holes and/or excitons that leave the emissive layer. The presence of such a blocking layer in a device may result in substantially higher efficiencies as compared to a similar device lacking a blocking layer. Also, a blocking layer may be used to confine emission to a desired region of an OLED.

50 In one aspect, compound used in HBL contains the same molecule or the same functional groups used as host described above.

55 In another aspect, compound used in HBL contains at least one of the following groups in the molecule:


60



65

61

-continued




k is an integer from 1 to 20; L^{101} is another ligand, k' is an integer from 1 to 3.

ETL:


Electron transport layer (ETL) may include a material capable of transporting electrons. Electron transport layer may be intrinsic (undoped), or doped. Doping may be used to enhance conductivity. Examples of the ETL material are not particularly limited, and any metal complexes or organic compounds may be used as long as they are typically used to transport electrons.

In one aspect, compound used in ETL contains at least one of the following groups in the molecule:



62

-continued




R^{101} is selected from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, when it is aryl or heteroaryl, it has the similar definition as Ar's mentioned above.

Ar^1 to Ar^3 has the similar definition as Ar's mentioned above.

k is an integer from 1 to 20.

X^{101} to X^{108} is selected from C (including CH) or N.

In another aspect, the metal complexes used in ETL contains, but not limit to the following general formula:



(O—N) or (N—N) is a bidentate ligand, having metal coordinated to atoms O, N or N,N; L^{101} is another ligand; k' is an integer value from 1 to the maximum number of ligands that may be attached to the metal.

In any above-mentioned compounds used in each layer of the OLED device, the hydrogen atoms can be partially or fully deuterated. Thus, any specifically listed substituent, such as, without limitation, methyl, phenyl, pyridyl, etc. encompasses undeuterated, partially deuterated, and fully deuterated versions thereof. Similarly, classes of substituents such as, without limitation, alkyl, aryl, cycloalkyl, heteroaryl, etc. also encompass undeuterated, partially deuterated, and fully deuterated versions thereof.

In addition to and/or in combination with the materials disclosed herein, many hole injection materials, hole transporting materials, host materials, dopant materials, exciton/hole blocking layer materials, electron transporting and electron injecting materials may be used in an OLED. Non-limiting examples of the materials that may be used in an OLED in combination with materials disclosed herein are listed in Table 1 below. Table 1 lists non-limiting classes of materials, non-limiting examples of compounds for each class, and references that disclose the materials.

TABLE 1








MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Hole injection materials		
Phthalocyanine and porphyrin compounds		Appl. Phys. Lett. 69, 2160 (1996)
Starburst triarylaminines		J. Lumin. 72-74, 985 (1997)
CF _x Fluorohydrocarbon polymer	$\text{---}[\text{CH}_x\text{F}_y]_n\text{---}$	Appl. Phys. Lett. 78, 673 (2001)
Conducting polymers (e.g., PEDOT: PSS, polyaniline, polythiophene)		Synth. Met. 87, 171 (1997) WO2007002683
Phosphonic acid and silane SAMs		US20030162053
Triarylamine or polythiophene polymers with conductivity dopants		EP1725079A1

TABLE 1-continued


MATERIAL

EXAMPLES OF MATERIAL

PUBLICATIONS




Organic compounds with
conductive inorganic
compounds, such as
molybdenum and tungsten
oxides



US20050123751
SID Symposium
Digest, 37, 923 (2006)
WO2009018009

n-type semiconducting
organic complexes



US20020158242

TABLE 1-continued





MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Metal organometallic complexes		US20060240279
Cross-linkable compounds		US20080220265
Polythiophene based polymers and copolymers		WO 2011075644 EP2350216
Hole transporting materials		
Triarylamines (e.g., TPD, α -NPD)		Appl. Phys. Lett. 51, 913 (1987)

TABLE 1-continued




MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
		U.S. Pat. No. 5,061,569
		EP650955
		J. Mater. Chem. 3, 319 (1993)

TABLE 1-continued





MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
		Appl. Phys. Lett. 90, 183503 (2007)
		Appl. Phys. Lett. 90, 183503 (2007)
Triarylamine on spirofluorene core		Synth. Met. 91, 209 (1997)
Arylamine carbazole compounds		Adv. Mater. 6, 677 (1994), US20080124572

TABLE 1-continued





MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Triarylamine with (di)benzothiophene/ (di)benzofuran		US20070278938, US20080106190 US20110163302
Indolocarbazoles		Synth. Met. 111, 421 (2000)
Isoindole compounds		Chem. Mater. 15, 3148 (2003)
Metal carbene complexes		US20080018221

TABLE 1-continued








MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Arylcarbazoles		Appl. Phys. Lett. 78, 1622 (2001)
Metal 8-hydroxyquinolates (e.g., Alq ₃ , BALq)		Nature 395, 151 (1998)
		US20060202194
		WO2005014551
		WO2006072002
Metal phenoxybenzothiazole compounds		Appl. Phys. Lett. 90, 123509 (2007)
Conjugated oligomers and polymers (e.g., polyfluorene)		Org. Electron. 1, 15 (2000)

TABLE 1-continued

MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Aromatic fused rings		WO2009066779, WO2009066778, WO2009063833, US20090045731, US20090045730, WO2009008311, US20090008605, US20090009065
Zinc complexes		WO2010056066
Chrysene based compounds		WO2011086863
Green hosts		
Arylcarbazoles		Appl. Phys. Lett. 78, 1622 (2001)
		US20030175553
		WO2001039234

TABLE 1-continued






MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Aryltriphenylene compounds		US20060280965
		US20060280965
		WO2009021126
Poly-fused heteroaryl compounds		US20090309488 US20090302743 US20100012931
Donor acceptor type molecules		WO2008056746

TABLE 1-continued






MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
		WO2010107244
Aza-carbazole/DBT/DBF		JP2008074939
		US20100187984
Polymers (e.g., PVK)		Appl. Phys. Lett. 77, 2280 (2000)
Spirofluorene compounds		WO2004093207

TABLE 1-continued






MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Metal phenoxybenzoxazole compounds		WO2005089025
		WO2006132173
		JP200511610
Spirofluorene-carbazole compounds		JP2007254297
		JP2007254297

TABLE 1-continued






MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Indolocabazoles		WO2007063796
		WO2007063754
5-member ring electron deficient heterocycles (e.g., triazole, oxadiazole)		J. Appl. Phys. 90, 5048 (2001)
		WO2004107822
Tetraphenylene complexes		US20050112407

TABLE 1-continued







MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Metal phenoxypyridine compounds		WO2005030900
Metal coordination complexes (e.g., Zn, Al with N^N ligands)		US20040137268, US20040137267
Blue hosts		
Arylcarbazoles		Appl. Phys. Lett, 82, 2422 (2003)
		US20070190359
Dibenzothiophene/ Dibenzofuran-carbazole compounds		WO2006114966, US20090167162
		US20090167162

TABLE 1-continued







MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
		WO2009086028
		US20090030202, US20090017330
		US20100084966
Silicon aryl compounds		US20050238919
		WO2009003898
Silicon/Germanium aryl compounds		EP2034538A

TABLE 1-continued







MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Aryl benzoyl ester		WO2006100298
Carbazole linked by non-conjugated groups		US20040115476
Aza-carbazoles		US20060121308
High triplet metal organometallic complex		U.S. Pat. No. 7,154,114
Phosphorescent dopants Red dopants		
Heavy metal porphyrins (e.g., PtOEP)		Nature 395, 151 (1998)
Iridium(III) organometallic complexes		Appl. Phys. Lett. 78, 1622 (2001)

TABLE 1-continued

MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
		US2006835469
		US2006835469
		US20060202194
		US20060202194
		US20070087321

TABLE 1-continued

MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
		US20080261076 US20100090591
		US20070087321
	<p data-bbox="1070 1606 1141 1634">H₁₇C₈</p>	Adv. Mater. 19, 739 (2007)
	<p data-bbox="1336 1877 1417 1906">Ir(acac)</p>	WO2009100991
		WO2008101842

TABLE 1-continued







MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Platinum(II) organometallic complexes		U.S. Pat. No. 7,232,618
		WO2003040257
		US20070103060
Osmium(III) complexes		Chem. Mater. 17, 3532 (2005)
Ruthenium(II) complexes		Adv. Mater. 17, 1059 (2005)
Rhenium (I), (II), and (III) complexes		US20050244673

TABLE 1-continued




MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Iridium(III) organometallic complexes	Green dopants  and its derivatives	Inorg. Chem. 40, 1704 (2001)
		US20020034656
		U.S. Pat. No. 7,332,232

TABLE 1-continued






MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
	 <p>The structure shows an iridium (Ir) center coordinated to two bipyridine ligands. The first bipyridine ligand has an isopropyl group at the 2-position. The second bipyridine ligand has a phenyl group at the 6-position. The third coordination site is occupied by a ligand consisting of a benzene ring fused to a five-membered ring containing a nitrogen atom, with a phenyl group attached to the benzene ring.</p>	US20090108737
	 <p>The structure shows an iridium (Ir) center coordinated to two bipyridine ligands. The first bipyridine ligand is unsubstituted. The second bipyridine ligand has a 4-methylphenyl group at the 6-position. The third coordination site is occupied by a ligand consisting of a benzene ring fused to a five-membered ring containing a nitrogen atom.</p>	WO2010028151
	 <p>The structure shows an iridium (Ir) center coordinated to two bipyridine ligands. The first bipyridine ligand is unsubstituted. The second bipyridine ligand has a phenyl group at the 6-position. The third coordination site is occupied by a ligand consisting of a benzene ring fused to an imidazole ring.</p>	EP1841834B
	 <p>The structure shows an iridium (Ir) center coordinated to three bipyridine ligands. Each bipyridine ligand has a phenyl group at the 6-position.</p>	US20060127696
	 <p>The structure shows an iridium (Ir) center coordinated to three bipyridine ligands. Each bipyridine ligand is unsubstituted.</p>	US20090039776

TABLE 1-continued

MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
		U.S. Pat. No. 6,921,915
		US20100244004
		U.S. Pat. No. 6,687,266
		Chem. Mater. 16, 2480 (2004)
		US20070190359

TABLE 1-continued







MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
		US20060008670 JP2007123392
		WO2010086089, WO2011044988
		Adv. Mater. 16, 2003 (2004)
		Angew. Chem. Int. Ed. 2006, 45, 7800
		WO2009050290
		US20090165846

TABLE 1-continued

MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
		US20080015355
		US20010015432
		US20100295032
Monomer for polymer metal organometallic compounds		U.S. Pat. No. 7,250,226, U.S. Pat. No. 7,396,598
Pt(II) organometallic complexes, including polydentated ligands		Appl. Phys. Lett. 86, 153505 (2005)

TABLE 1-continued






MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
		Appl. Phys. Lett. 86, 153505 (2005)
		Chem. Lett. 34, 592 (2005)
		WO2002015645
		US20060263635
		US20060182992 US20070103060

TABLE 1-continued





MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Cu complexes		WO2009000673
		US20070111026
Gold complexes		Chem. Commun. 2906 (2005)
Rhenium(III) complexes		Inorg. Chem. 42, 1248 (2003)

TABLE 1-continued





MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Osmium(II) complexes		U.S. Pat. No. 7,279,704
Deuterated organometallic complexes		US20030138657
Organometallic complexes with two or more metal centers		US20030152802
		U.S. Pat. No. 7,090,928

TABLE 1-continued

MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Iridium(III) organometallic complexes		WO2002002714
		WO2006009024
		US20060251923 US20110057559 US20110204333
		U.S. Pat. No. 7,393,599, WO2006056418, US20050260441, WO2005019373
		U.S. Pat. No. 7,534,505
		WO2011051404

TABLE 1-continued

MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
		U.S. Pat. No. 7,445,855
		US20070190359, US20080297033 US20100148663
		U.S. Pat. No. 7,338,722
		US20020134984
		Angew. Chem. Int. Ed. 47, 1 (2008)

TABLE 1-continued

MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
		Chem. Mater. 18, 5119 (2006)
		Inorg. Chem. 46, 4308 (2007)
		WO2005123873
		WO2005123873
		WO2007004380

TABLE 1-continued

MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Osmium(II) complexes		WO2006082742
Osmium(II) complexes		U.S. Pat. No. 7,279,704
Gold complexes		Appl. Phys. Lett. 74,1361 (1999)
Platinum(II) complexes		WO2006098120, WO2006103874

TABLE 1-continued






MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Pt tetradentate complexes with at least one metal-carbene bond		U.S. Pat. No. 7,655,323
Exciton/hole blocking layer materials		
Bathocuprine compounds (e.g., BCP, BPhen)		Appl. Phys. Lett. 75, 4 (1999)
		Appl. Phys. Lett. 79, 449 (2001)
Metal 8-hydroxyquinolates (e.g., BALq)		Appl. Phys. Lett. 81, 162 (2002)
5-member ring electron deficient heterocycles such as triazole, oxadiazole, imidazole, benzoimidazole		Appl. Phys. Lett. 81, 162 (2002)

TABLE 1-continued





MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Triphenylene compounds		US20050025993
Fluorinated aromatic compounds		Appl. Phys. Lett. 79, 156 (2001)
Phenothiazine-S-oxide		WO2008132085
Silylated five-membered nitrogen, oxygen, sulfur or phosphorus dibenzoheterocycles		WO2010079051

TABLE 1-continued






MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Aza-carbazoles		US20060121308
Electron transporting materials		
Anthracene-benzimidazole compounds		WO2003060956
		US20090179554
Aza triphenylene derivatives		US20090115316
Anthracene-benzothiazole compounds		Appl. Phys. Lett. 89, 063504 (2006)

TABLE 1-continued






MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Metal 8-hydroxyquinolates (e.g., Alq ₃ , Zrq ₄)		Appl. Phys. Lett. 51, 913 (1987) U.S. Pat. No. 7,230,107
Metal hydroxybenoquinolates		Chem. Lett. 5, 905 (1993)
Bathocuprine compounds such as BCP, BPhen, etc		Appl. Phys. Lett. 91, 263503 (2007)
5-member ring electron deficient heterocycles (e.g., triazole, oxadiazole, imidazole, benzimidazole)		Appl. Phys. Lett. 79, 449 (2001)
5-member ring electron deficient heterocycles (e.g., triazole, oxadiazole, imidazole, benzimidazole)		Appl. Phys. Lett. 55, 1489 (1989)

TABLE 1-continued







MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Silole compounds		Jpn. J. Apply. Phys. 32, L917 (1993)
Arylborane compounds		J. Am. Chem. Soc. 120, 9714 (1998)
Fluorinated aromatic compounds		J. Am. Chem. Soc. 122, 1832 (2000)
Fullerene (e.g., C60)		US20090101870
Triazine complexes		US20040036077

TABLE 1-continued

MATERIAL	EXAMPLES OF MATERIAL	PUBLICATIONS
Zn (N [^] N) complexes		U.S. Pat. No. 6,528,187

EXAMPLES

Example 1


Computational Examples

DFT calculations were performed for certain example compounds and comparative compounds. The example compounds and comparative compounds are shown on the following page. In certain models, a hot, Host 1, was used, which is also shown.

-continued

Comparative example 3


20



25

30

Comparative example 1




35

40

Comparative Example 4

45




50

55

60

65


Comparative example 2



135

-continued

Comparative Example 5



5


10

15

136

-continued


Comparative Example 8



20

25

Comparative Example 9




30

35


40

45

Comparative Example 6



Compound 1



50



55

60

65


Compound 2

Comparative Example 7



137

-continued



Compound 3

5

10

15

20

25


Compound 4

30

35

40

45




50


Compound 8

55

60


**138**

-continued




Compound 56


Compound 57



Compound 58



Host 1



65

70


75

80

85

Geometry optimization calculations were performed within the Gaussian 09 software package using the B3LYP hybrid functional and CEP-31g effective core potential basis set. Complexes were optimized as both ground state singlets and triplets. Three bonds strengths of interest were examined: N-aryl, N-phenyl, and N-Phenyl2, the identities of which are shown in Structure A atop the following page.

139



Structure A

Bond strength calculations were executed at the same level of theory by two approaches which were dependent on either (a) creating two fragments or (b) breaking a bond that does not cause fragmentation.

Fragmentation of the phenyl group from the bridging nitrogen (N_{phenyl}) results in the transition metal fragment in a doublet spin state and the phenyl group also in a doublet spin state. The complete fragmentation is only observed for Comparative Examples 4-6, where the N_{phenyl} bond is not tethered. For these complexes, the N_{phenyl} fragment can be completely removed whereas for Compound 1-4, the phenyl fragment remains bound to the complex on one or both sides. For complexes that do not fragment the phenyl bonded to the bridging nitrogen both singlet→singlet and triplet→triplet spin states bond strength energies were calculated. The results are shown below in TABLE 2.

140

phenylimidazole ligand increases the N-phenyl bond strength. In the triplet state, Comparative Example 1 has an N-phenyl bond strength of 8.02 kcal/mole. Comparative Examples 4 and 6, without tethering substitutions are found to have similar N-phenyl bond strength compared to Comparative Example 1. Compound 2 is shown to have a stronger N-phenyl bond of 10.99 kcal/mol, however, the triplet energy of 475 nm is lower than Comparative Example 1. Tethered examples, Compounds 1-4 are found to have stronger N-phenyl bonds while in several cases retaining the desirable high triplet energy. A special case is found for Compound 3, which is tethered by geminal dimethyl groups on both sides. As shown in the table, designated by “*”, these substitutions are calculated to have a large impact on the N-phenyl bond strength. Linking the phenyl group to the complex on both sides prevents the phenyl from twisting or moving out of the bonding position. Optimization results in a distorted geometry of Compound 3, such that the phenyl is not directly lined up with the nitrogen and the N-phenyl bond. The distortion results in a relatively small calculated energy difference of 5.22 kcal/mol for the N-phenyl bond strength of Compound 3. However, the N-phenyl bond is reformed, so the small difference in energy is related to the slight geometry difference, not a broken bond. The calculated geometric distortion for Compound 3 is shown in FIG. 7. Thus, tethering (or bridging) the phenyl group on both sides retains the N-phenyl bond and shifts the lowest bond strength to the N-aryl bond. In addition the other structures, Compound 1, 2 and 4, are found to have greater N-phenyl bond strengths compared to Comparative Example 1.

Calculating the N_{aryl} bond strengths was done by removing the dimethylphenyl group and separately calculating the energies of the doublet complex and dimethylphenyl. There are no major differences in the N-aryl bond strengths besides the differing ground state triplet energies between the complexes.

TABLE 2

Compound	Calc T1 (nm)	N-Phenyl Bond Strength, singlet/triplet (kcal/mol)	N-Aryl Bond Strength, singlet/triplet (kcal/mol)	N-Phenyl-2 Bond Strength, singlet/triplet (kcal/mol)	Weakest bond (kcal/mol)	Device Lifetime (relative to comparative example 1)
Comparative Example 1	467	66.82/8.02	74.08/15.28	85.89/27.09	8.02	1
Comparative Example 2	502	67.69/13.35	77.15/22.81		13.35	10
Comparative Example 3	508	68.79/15.32	n/a		15.32	100
Comparative Example 4	467	66.92/8.17	74.15/15.40	84.98/26.23	8.17	
Comparative Example 5	475	66.35/10.99	75.29/19.93	85.41/30.06	10.99	
Comparative Example 6	467	66.40/7.82	74.21/15.63	86.00/27.42	7.82	
Comparative Example 7		67.86/25.67			25.67	
Comparative Example 8		69.19/15.01			15.01	
Comparative Example 9		67.33/14.26			14.26	
Compound 1	475	70.31/11.52	75.84/17.05	75.73/16.95	11.52	
Compound 2	464	73.10/12.36	75.57/14.83	81.82/21.08	12.36	
Compound 3	466	62.84/5.22*	71.73/14.10	83.04/25.42	14.10	
Compound 4	464	78.26/17.70	76.56/16.00	83.92/23.36	16.00	
Compound 8		73.59/29.07			29.07	
Compound 56		75.16/21.18			21.18	
Compound 57		76.07/22.91			22.91	
Compound 58		77.67/24.97			24.97	

As shown in TABLE 2, calculations indicate that linking the phenyl in the N-phenyl position to the phenyl of the

Finally, calculations were performed on the N-phenyl-2 bond in which the bridging N atom is detached from the

phenyl of the phenylimidazole ligand. This bond is calculated to be the strongest bond of the three investigated except for complex Compound 1. The N-phenyl-2 bond strength is calculated to be a mere 0.1 kcal/mol weaker than the N-aryl bond for complex Compound 1. In all cases, the N-phenyl-2 bond is never calculated to be the weakest bond of the complex. This suggests that the breaking the N-phenyl-2 bond is not a significant source of decomposition. In addition, Compound 8, 56, 57, and 58 further demonstrate the versatility of this approach. In each case the N-aryl bond strength is calculated to be greater than the corresponding untethered comparative example, Comparative Example 7, 8 and 9.

Example 2

MALDI-TOF Fragmentation


The relative instability of the N—C ring bond is demonstrated experimentally by excited state fragmentation patterns observed in a MALDI mass spectrometer. Host 1 and Comparative Examples 1-3 at an approximate 20 wt % ratio were dissolved in dichloromethane and spot cast on an Applied Biosystems Voyager DE-STR MALDI sample target plate. The samples were analyzed in the positive mode. FIGS. 4-6 show that there is a major fragment that results from the loss of the N-phenyl ring, i.e. N—C ring bond breakage. In each case, it is found that the broken N—C bond is an abundant fragment associated with the dopant.

Without being bound to any theory, it is believed that linking or tethering (or bridging) the C-Ring back to the complex will result in an improvement in stability particularly for phosphorescent metal complexes with bridging aryl amine substitutions.

It is understood that the various embodiments described herein are by way of example only, and are not intended to limit the scope of the invention. For example, many of the materials and structures described herein may be substituted with other materials and structures without deviating from the spirit of the invention. The present invention as claimed may therefore include variations from the particular examples and preferred embodiments described herein, as will be apparent to one of skill in the art. It is understood that various theories as to why the invention works are not intended to be limiting.

We claim:

1. A compound having Formula (II):



wherein rings A¹, A², and A³ are independently five- or six-membered carbocyclic or heterocyclic aromatic

rings having 0-3 nitrogen atoms and 0-1 additional heteroatoms selected from the group consisting of oxygen, sulfur, and selenium;

Z¹ and Z² are independently carbon or nitrogen;

wherein R² represents mono- or di-substitution, wherein each R² is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R² may optionally combine to form a ring, which can be further substituted;

wherein R³ represents mono-, di-, tri-, or tetra-substitution, wherein each R³ is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R³ may optionally combine to form a ring, which can be further substituted;

wherein G¹ is O or CR⁴R⁵;

wherein R⁴ and R⁵ are independently selected from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein R⁴ and R⁵ may optionally combine to form a ring, which can be further substituted;

wherein any of R⁴ or R⁵ may optionally combine with any R² or R³ to form a ring system, which can be further substituted;

wherein R⁶ represents mono-, di-, or tri-substitution, wherein each R⁶ is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R⁶ may optionally combine to form a ring, which can be further substituted;

wherein R⁷ and R⁸ represent mono-, di-, tri-, or tetra-substitution, wherein each R⁷ or R⁸ is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R⁷ or any two adjacent R⁸ may optionally combine to form a ring, which can be further substituted;

wherein any R³ may optionally combine with any R⁶ to form a ring system, which can be further substituted;

wherein any R² may optionally combine with any R⁸ to form a ring system, which can be further substituted; and


wherein any R⁶ may optionally combine with any R⁷ to form a ring system, which can be further substituted;

wherein any R⁷ may optionally combine with any R⁸ to form a ring system, which can be further substituted.

143

2. The compound of claim 1, wherein G¹ is O, C(CH₃)₂, or C(C₆H₅)₂.

3. The compound of claim 1, which is a compound having Formula (IIa):



(IIa)

144

-continued


Compound 3

5

10

15

20




4. The compound of claim 1, which is selected from the group consisting of:

25

Compound 1

Compound 4




30

35


40

45



Compound 2

Compound 5




50

55


60

65



145

-continued



Compound 6


5

10

15

146

-continued




Compound 10

Compound 7


20

Compound 11



Compound 8

35



Compound 9


40

45

50

55

Compound 12




60

65

5. A first device comprising a first organic light emitting device which comprises:
 an anode;
 a cathode; and
 an organic layer disposed between the anode and the cathode,
 the organic layer comprising a compound having Formula (II):

147



wherein rings A^1 , A^2 , and A^3 are independently five- or six-membered carbocyclic or heterocyclic aromatic rings having 0-3 nitrogen atoms and 0-1 additional heteroatoms selected from the group consisting of oxygen, sulfur, and selenium;

Z^1 and Z^2 are independently carbon or nitrogen;

wherein R^2 represents mono- or di-substitution, wherein each R^2 is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^2 may optionally combine to form a ring, which can be further substituted;

wherein R^3 represents mono-, di-, tri-, or tetra-substitution, wherein each R^3 is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^3 may optionally combine to form a ring, which can be further substituted;

wherein G^1 is O or CR^4R^5 ;

wherein R^4 and R^5 are independently selected from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein R^4 and R^5 may optionally combine to form a ring, which can be further substituted;

wherein any of R^4 or R^5 may optionally combine with any R^2 or R^3 to form a ring system, which can be further substituted;

wherein R^6 represents mono-, di-, or tri-substitution, wherein each R^6 is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^6 may optionally combine to form a ring, which can be further substituted;

148

wherein R^7 and R^8 represent mono-, di-, tri-, or tetra-substitution, wherein each R^7 or R^8 is selected independently from the group consisting of hydrogen, deuterium, halide, alkyl, cycloalkyl, heteroalkyl, arylalkyl, alkoxy, aryloxy, amino, silyl, alkenyl, cycloalkenyl, heteroalkenyl, alkynyl, aryl, heteroaryl, acyl, carbonyl, carboxylic acids, ester, nitrile, isonitrile, sulfanyl, sulfinyl, sulfonyl, phosphino, and combinations thereof, and wherein any two adjacent R^7 or any two adjacent R^8 may optionally combine to form a ring, which can be further substituted;

wherein any R^3 may optionally combine with any R^6 to form a ring system, which can be further substituted; wherein any R^2 may optionally combine with any R^8 to form a ring system, which can be further substituted; and wherein any R^6 may optionally combine with any R^7 to form a ring system, which can be further substituted; wherein any R^7 may optionally combine with any R^8 to form a ring system, which can be further substituted.

6. The first device of claim 5, wherein the first device is a consumer product.

7. The first device of claim 5, wherein the first device is an organic light-emitting device.

8. The first device of claim 5, wherein the first device comprises a lighting panel.

9. The first device of claim 5, wherein the organic layer is an emissive layer and the compound is an emissive dopant.

10. The first device of claim 5, wherein the organic layer is an emissive layer and the compound is a non-emissive dopant.

11. The first device of claim 5, wherein the organic layer further comprises a host.

12. The first device of claim 11, wherein the host comprises a triphenylene containing benzo-fused thiophene or benzo-fused furan;


wherein any substituent in the host is an unfused substituent independently selected from the group consisting of C_nH_{2n+1} , OC_nH_{2n+1} , OAr_1 , $N(C_nH_{2n+1})_2$, $N(Ar_1)(Ar_2)$, $CH=CH-C_nH_{2n+1}$, $C\equiv C-C_nH_{2n+1}$, Ar_1 , Ar_1-Ar_2 , and $C_nH_{2n}-Ar_1$, or the host has no substitution;

wherein n is from 1 to 10; and

wherein Ar_1 and Ar_2 are independently selected from the group consisting of benzene, biphenyl, naphthalene, triphenylene, carbazole, and heteroaromatic analogs thereof.


13. The first device of claim 11, wherein the host comprises a compound selected from the group consisting of: carbazole, dibenzothiophene, dibenzofuran, dibenzoselenophene, azacarbazole, aza-dibenzothiophene, aza-dibenzofuran, and aza-dibenzoselenophene.

14. The first device of claim 11, wherein the host is selected from the group consisting of:




149

-continued



150

-continued



and combinations thereof.

55 **15.** The first device of claim 11, wherein the host comprises a metal complex.

* * * * *