



US009551708B2

(12) **United States Patent**
Snider et al.

(10) **Patent No.:** **US 9,551,708 B2**
(45) **Date of Patent:** ***Jan. 24, 2017**

(54) **METHODS OF TREATING A SUBJECT
HAVING HEART FAILURE**

(71) Applicant: **Critical Care Diagnostics, Inc.**, San
Diego, CA (US)

(72) Inventors: **James V. Snider**, San Diego, CA (US);
Robert W. Gerwien, Newington, CT
(US)

(73) Assignee: **Critical Care Diagnostics, Inc.**, San
Diego, CA (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

This patent is subject to a terminal dis-
claimer.

(21) Appl. No.: **14/299,851**

(22) Filed: **Jun. 9, 2014**

(65) **Prior Publication Data**

US 2014/0286944 A1 Sep. 25, 2014

Related U.S. Application Data

(63) Continuation of application No. 13/552,533, filed on
Jul. 18, 2012, now Pat. No. 8,748,110.

(60) Provisional application No. 61/508,923, filed on Jul.
18, 2011, provisional application No. 61/509,359,
filed on Jul. 19, 2011.

(51) **Int. Cl.**

G01N 33/53 (2006.01)

G01N 33/68 (2006.01)

A61P 9/04 (2006.01)

G01N 33/566 (2006.01)

C07K 14/705 (2006.01)

C07K 14/715 (2006.01)

(52) **U.S. Cl.**

CPC **G01N 33/566** (2013.01); **G01N 33/53**
(2013.01); **G01N 33/6869** (2013.01); **G01N**
33/6893 (2013.01); **C07K 14/7155** (2013.01);
G01N 2333/7155 (2013.01); **G01N 2800/32**
(2013.01); **G01N 2800/52** (2013.01)

(58) **Field of Classification Search**

None

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

7,432,060 B2 10/2008 Lee
7,655,415 B2 2/2010 Lee
7,670,769 B2 3/2010 Lee
7,985,558 B2 7/2011 Lee
7,989,210 B2 8/2011 Lee
7,998,683 B2 8/2011 Snider et al.
8,090,562 B2 1/2012 Snider et al.
8,420,785 B2 4/2013 Snider
8,530,173 B2 9/2013 Lee
8,597,958 B2 12/2013 Lee

8,617,825 B2 12/2013 Snider et al.
8,728,742 B2 5/2014 Snider
8,734,769 B2 5/2014 Lee
8,748,110 B2 6/2014 Snider et al.
8,748,116 B2 6/2014 Lee
2007/0248981 A1 10/2007 Snider et al.
2009/0264779 A1 10/2009 Snider et al.
2009/0305265 A1 12/2009 Snider et al.
2010/0009356 A1 1/2010 Snider et al.
2010/0055683 A1 3/2010 Snider et al.
2011/0053170 A1 3/2011 Snider et al.
2011/0250703 A1 10/2011 Lee
2011/0262941 A1 10/2011 Snider
2011/0280887 A1 11/2011 Lee
2012/0040381 A1 2/2012 Snider
2012/0065897 A1 3/2012 Snider
2012/0276551 A1 11/2012 Snider
2013/0177931 A1 7/2013 Snider
2013/0244236 A1 9/2013 Snider et al.
2013/0251664 A1 9/2013 Lee
2013/0273562 A1 10/2013 Lee
2013/0317030 A1 11/2013 Lee
2013/0345805 A1 12/2013 Snider et al.
2014/0045200 A1 2/2014 Snider
2014/0051773 A1 2/2014 Snider
2014/0058743 A1 2/2014 Snider et al.

FOREIGN PATENT DOCUMENTS

RU 2355295 5/2009

OTHER PUBLICATIONS

U.S. Appl. No. 14/290,465, filed May 29, 2014, Lee.

U.S. Appl. No. 14/244,526, filed Apr. 3, 2014, Snider et al.

Aaronson, et al., "Development and prospective validation of a
clinical index to predict survival in ambulatory patients referred for
cardiac transplant evaluation," *Circulation* 95(12):2660-2667
(1997).

Bittner et al., "Prediction of mortality and morbidity with a 6-minute
walk test in patients with left ventricular dysfunction. SOLVD
Investigators," *JAMA* 270(14):1702-1707 (1993).

International Search Report issued in international application PCT/
US2012/047238, mailed Nov. 1, 2012.

Korhonen et al., "Smoking cessation program with exercise
improves cardiovascular disease biomarkers in sedentary women,"
Journal of Women's Health 20:1051-1064 (2011).

(Continued)

Primary Examiner — Zachary Howard

(74) *Attorney, Agent, or Firm* — Fish & Richardson P.C.

(57) **ABSTRACT**

Methods of treating a subject having a cardiovascular dis-
ease, selecting a therapy for a subject having a cardiovas-
cular disease, identifying a subject having a cardiovascular
disease that will benefit or not benefit from exercise therapy,
determining whether a subject having a cardiovascular dis-
ease should begin, continue, not begin, discontinue, or avoid
exercise therapy, determining whether a subject having a
cardiovascular disease should continue, discontinue, or
avoid exercise therapy, reducing the risk of an adverse
outcome (e.g., death) in a subject having a cardiovascular
disease, and predicting the efficacy of exercise therapy in a
subject having a cardiovascular disease. These methods
include determining a level of soluble ST2 in a subject.

30 Claims, 8 Drawing Sheets

(56)

References Cited

OTHER PUBLICATIONS

Ky et al., “High-sensitivity ST2 for prediction of adverse outcomes in chronic heart failure,” *Cir. Heart Fail.* 4:180-187 (2011).
Lu et al., “Establishment of reference intervals for soluble ST2 from a United States population,” *Clin. Chim. Acta* 411:21-22 (2010) (Abstract only).

Mancini et al., “Value of peak exercise oxygen consumption for optimal timing of cardiac transplantation in ambulatory patients with heart failure,” *Circulation* 83:778-786 (1991).
O’Connor et al., “Efficacy and safety of exercise training in patients with chronic heart failure, HF-ACTION randomized Controlled Trial,” *JAMA* 301:1439-1450 (2009).
Written Opinion of the International Searching Authority issued in international application PCT/US2012/047238, mailed Nov. 1, 2012.
International Preliminary Report on Patentability for PCT/US2012/047238, issued Jan. 21, 2014.




Fig. 1




Fig. 2




Fig. 3




Fig. 4

Met	Gly	Phe	Trp	Ile	Leu	Ala	Ile	Leu	Thr	Ile	Leu	Met	Tyr	Ser	Thr
1				5					10					15	
Ala	Ala	Lys	Phe	Ser	Lys	Gln	Ser	Trp	Gly	Leu	Glu	Asn	Glu	Ala	Leu
			20					25					30		
Ile	Val	Arg	Cys	Pro	Arg	Gln	Gly	Lys	Pro	Ser	Tyr	Thr	Val	Asp	Trp
		35				40						45			
Tyr	Tyr	Ser	Gln	Thr	Asn	Lys	Ser	Ile	Pro	Thr	Gln	Glu	Arg	Asn	Arg
	50				55						60				
Val	Phe	Ala	Ser	Gly	Gln	Leu	Leu	Lys	Phe	Leu	Pro	Ala	Ala	Val	Ala
65				70					75					80	
Asp	Ser	Gly	Ile	Tyr	Thr	Cys	Ile	Val	Arg	Ser	Pro	Thr	Phe	Asn	Arg
			85					90					95		
Thr	Gly	Tyr	Ala	Asn	Val	Thr	Ile	Tyr	Lys	Lys	Gln	Ser	Asp	Cys	Asn
			100					105					110		
Val	Pro	Asp	Tyr	Leu	Met	Tyr	Ser	Thr	Val	Ser	Gly	Ser	Glu	Lys	Asn
		115				120						125			
Ser	Lys	Ile	Tyr	Cys	Pro	Thr	Ile	Asp	Leu	Tyr	Asn	Trp	Thr	Ala	Pro
	130					135					140				
Leu	Glu	Trp	Phe	Lys	Asn	Cys	Gln	Ala	Leu	Gln	Gly	Ser	Arg	Tyr	Arg
145				150					155					160	
Ala	His	Lys	Ser	Phe	Leu	Val	Ile	Asp	Asn	Val	Met	Thr	Glu	Asp	Ala
			165					170					175		
Gly	Asp	Tyr	Thr	Cys	Lys	Phe	Ile	His	Asn	Glu	Asn	Gly	Ala	Asn	Tyr
			180					185					190		
Ser	Val	Thr	Ala	Thr	Arg	Ser	Phe	Thr	Val	Lys	Asp	Glu	Gln	Gly	Phe
		195				200						205			
Ser	Leu	Phe	Pro	Val	Ile	Gly	Ala	Pro	Ala	Gln	Asn	Glu	Ile	Lys	Glu
	210					215					220				
Val	Glu	Ile	Gly	Lys	Asn	Ala	Asn	Leu	Thr	Cys	Ser	Ala	Cys	Phe	Gly
225				230						235				240	
Lys	Gly	Thr	Gln	Phe	Leu	Ala	Ala	Val	Leu	Trp	Gln	Leu	Asn	Gly	Thr
			245					250					255		
Lys	Ile	Thr	Asp	Phe	Gly	Glu	Pro	Arg	Ile	Gln	Gln	Glu	Glu	Gly	Gln
			260					265					270		
Asn	Gln	Ser	Phe	Ser	Asn	Gly	Leu	Ala	Cys	Leu	Asp	Met	Val	Leu	Arg
		275				280						285			
Ile	Ala	Asp	Val	Lys	Glu	Glu	Asp	Leu	Leu	Leu	Gln	Tyr	Asp	Cys	Leu
	290					295					300				
Ala	Leu	Asn	Leu	His	Gly	Leu	Arg	Arg	His	Thr	Val	Arg	Leu	Ser	Arg
305				310						315					320
Lys	Asn	Pro	Ser	Lys	Glu	Cys	Phe								
				325											

(SEQ ID NO:1)

Fig. 5

gaggagggac ctacaaagac tggaaactat tcttagctcc gtcactgact ccaagttcat 60
cccctctgtc ttccagtttg gttgagatat aggctactct tcccaactca gtcttgaaga 120
gtatcaccaa ctgcctcatg tgtggtgacc ttccactgtcg tatgccagtg actcatcttg 180
agtaatctca acaacgagtt accaatactt gctcttgatt gataaacaga atgggggtttt 240
ggatcttagc aattctcaca attctcatgt attccacagc agcaaagttt agtaaacaat 300
catggggcct ggaaaatgag gctttaattg taagatgtcc tagacaagga aaacctagtt 360
acaccgtgga ttggtattac tcacaaacaa acaaaagtat tcccactcag gaaagaaatc 420
gtgtgtttgc ctccaggcca cttctgaagt ttctaccagc tgcagttgct gattctggta 480
tttataacctg tattgtcaga agtcccacat tcaataggac tggatatgcg aatgtcacca 540
tatataaaaa acaatcagat tgcaatgttc cagattattt gatgtattca acagtatctg 600
gatcagaaaa aaattccaaa atttattgtc ctaccattga cctctacaac tggacagcac 660
ctcttgagtg gtttaagaat tgtcaggctc ttcaaggatc aaggtaacag ggcacacaag 720
catttttggc cattgataat gtgatgactg aggacgcagg tgattacacc tgtaaattta 780
tacacaatga aaatggagcc aattatagtg tgacggcgac caggctcctc acgggtcaagg 840
atgagcaagg cttttctctg ttccagtaa tgggagcccc tgcacaaaaat gaaataaagg 900
aagtggaaat tggaaaaaac gcaaacctaa cttgctctgc ttgttttggg aaaggcactc 960
agttcttggc tgccgtcctg tggcagctta atggaacaaa aattacagac tttggtgaac 1020
caagaattca acaagaggaa gggcaaaatc aaagtttcag caatgggctg gcttgtctag 1080
acatgggttt aagaatagct gacgtgaagg aagaggattt attgctgcag tacgactgtc 1140
tggccctgaa ttgcatggc ttgagaaggc acaccgtaag actaagtagg aaaaatccaa 1200
gtaaggagtg tttctgagac tttgatcacc tgaactttct ctagcaagtg taagcagaat 1260
ggagtgtggc tccaagagat ccatacaagc aatgggaatg gcctgtgcca taaaatgtgc 1320
ttctcttctt cgggatgttg tttgctgtct gatctttgta gactgttctt gtttgctggg 1380
agcttctctg ctgcttaaat tgttcgtcct ccccaactcc ctctatcgt tggtttgtct 1440
agaacactca gctgcttctt tggtcacctt tgttttctaa ctttatgaac tccctctgtg 1500
tcaactgtatg tgaaaggaaa tgcaccaaca accgtaaaact gaacgtgttc ttttgtgctc 1560
ttttataact tgcattacat gttgtaagca tgggtccgttc tatacctttt totggtcata 1620
atgaacactc attttgttag cgaggggtgg aaagtgaaca aaaaggggaa gtatcaaact 1680
actgccattt cagtgaagaa atcctaggtg ctactttata ataagacatt tgttaggcca 1740
ttcttgcatg gatataaaga aatacctgag actgggtgat ttatatgaaa agaggtttaa 1800
ttggctcaca gttctgcagg ctgtatggga agcatggcgg catctgcttc tggggacacc 1860
tcaggagctt tactcatggc agaaggcaaa gcaaaggcag gcacttcaca cagtaaaagc 1920
aggagcgaga gagaggtgcc aactgaaac agccagatct catgagaagt cactcactat 1980
tgcaaggaca gcatcaaaga gatggtgcta aaccattcat gatgaactca ccccatgat 2040
ccaatcacct cccaccaggc tccacctcga atactgggga ttaccattca gcatgagatt 2100
tgggcaggaa cacagaccca aaccatacca cacacattat cattgttaaa ctttgtaaag 2160
tatttaaggc acatggaaca cacgggaagt ctggtagctc agccatttc tttattgcat 2220
ctgttattca ccatgtaatt caggtaccac gtattccagg gagcctttct tggccctcag 2280
tttgagatg acacactttc caagtactct tgtagcatcc tgtttgtatc atagcactgg 2340
tcacattgcc ttacctaaat ctgtttgaca gtctgtcaca cagactgca agctccatga 2400
gggcagggac atcatctctt ccatcttttg gtccttagtg caatacctgg cagctagcca 2460
gtgctcagct aaatatattg tgaactgaata aatgaatgca caacaaaaa aaaaaaaaaa 2520
aaaaaaaaaa aaaaaaaaaa aa 2542
(SEQ ID NO:2)

Fig. 6

Met	Gly	Phe	Trp	Ile	Leu	Ala	Ile	Leu	Thr	Ile	Leu	Met	Tyr	Ser	Thr
1				5					10					15	
Ala	Ala	Lys	Phe	Ser	Lys	Gln	Ser	Trp	Gly	Leu	Glu	Asn	Glu	Ala	Leu
			20					25					30		
Ile	Val	Arg	Cys	Pro	Arg	Gln	Gly	Lys	Pro	Ser	Tyr	Thr	Val	Asp	Trp
		35					40					45			
Tyr	Tyr	Ser	Gln	Thr	Asn	Lys	Ser	Ile	Pro	Thr	Gln	Glu	Arg	Asn	Arg
	50					55					60				
Val	Phe	Ala	Ser	Gly	Gln	Leu	Leu	Lys	Phe	Leu	Pro	Ala	Ala	Val	Ala
65				70						75					80
Asp	Ser	Gly	Ile	Tyr	Thr	Cys	Ile	Val	Arg	Ser	Pro	Thr	Phe	Asn	Arg
			85					90					95		
Thr	Gly	Tyr	Ala	Asn	Val	Thr	Ile	Tyr	Lys	Lys	Gln	Ser	Asp	Cys	Asn
			100					105					110		
Val	Pro	Asp	Tyr	Leu	Met	Tyr	Ser	Thr	Val	Ser	Gly	Ser	Glu	Lys	Asn
		115					120					125			
Ser	Lys	Ile	Tyr	Cys	Pro	Thr	Ile	Asp	Leu	Tyr	Asn	Trp	Thr	Ala	Pro
	130					135					140				
Leu	Glu	Trp	Phe	Lys	Asn	Cys	Gln	Ala	Leu	Gln	Gly	Ser	Arg	Tyr	Arg
145				150						155					160
Ala	His	Lys	Ser	Phe	Leu	Val	Ile	Asp	Asn	Val	Met	Thr	Glu	Asp	Ala
			165					170					175		
Gly	Asp	Tyr	Thr	Cys	Lys	Phe	Ile	His	Asn	Glu	Asn	Gly	Ala	Asn	Tyr
			180					185					190		
Ser	Val	Thr	Ala	Thr	Arg	Ser	Phe	Thr	Val	Lys	Asp	Glu	Gln	Gly	Phe
		195					200					205			
Ser	Leu	Phe	Pro	Val	Ile	Gly	Ala	Pro	Ala	Gln	Asn	Glu	Ile	Lys	Glu
	210					215					220				
Val	Glu	Ile	Gly	Lys	Asn	Ala	Asn	Leu	Thr	Cys	Ser	Ala	Cys	Phe	Gly
225				230						235					240
Lys	Gly	Thr	Gln	Phe	Leu	Ala	Ala	Val	Leu	Trp	Gln	Leu	Asn	Gly	Thr
			245					250					255		
Lys	Ile	Thr	Asp	Phe	Gly	Glu	Pro	Arg	Ile	Gln	Gln	Glu	Glu	Gly	Gln
			260					265					270		
Asn	Gln	Ser	Phe	Ser	Asn	Gly	Leu	Ala	Cys	Leu	Asp	Met	Val	Leu	Arg
		275				280						285			
Ile	Ala	Asp	Val	Lys	Glu	Glu	Asp	Leu	Leu	Leu	Gln	Tyr	Asp	Cys	Leu
	290					295					300				
Ala	Leu	Asn	Leu	His	Gly	Leu	Arg	Arg	His	Thr	Val	Arg	Leu	Ser	Arg
305				310						315					320
Lys	Asn	Pro	Ile	Asp	His	His	Ser	Ile	Tyr	Cys	Ile	Ile	Ala	Val	Cys
			325						330				335		
Ser	Val	Phe	Leu	Met	Leu	Ile	Asn	Val	Leu	Val	Ile	Ile	Leu	Lys	Met
			340					345					350		
Phe	Trp	Ile	Glu	Ala	Thr	Leu	Leu	Trp	Arg	Asp	Ile	Ala	Lys	Pro	Tyr
		355					360					365			
Lys	Thr	Arg	Asn	Asp	Gly	Lys	Leu	Tyr	Asp	Ala	Tyr	Val	Val	Tyr	Pro
	370					375					380				
Arg	Asn	Tyr	Lys	Ser	Ser	Thr	Asp	Gly	Ala	Ser	Arg	Val	Glu	His	Phe
385					390					395					400

Fig. 7A

Val	His	Gln	Ile	Leu	Pro	Asp	Val	Leu	Glu	Asn	Lys	Cys	Gly	Tyr	Thr
				405					410					415	
Leu	Cys	Ile	Tyr	Gly	Arg	Asp	Met	Leu	Pro	Gly	Glu	Asp	Val	Val	Thr
			420					425					430		
Ala	Val	Glu	Thr	Asn	Ile	Arg	Lys	Ser	Arg	Arg	His	Ile	Phe	Ile	Leu
		435					440					445			
Thr	Pro	Gln	Ile	Thr	His	Asn	Lys	Glu	Phe	Ala	Tyr	Glu	Gln	Glu	Val
	450					455					460				
Ala	Leu	His	Cys	Ala	Leu	Ile	Gln	Asn	Asp	Ala	Lys	Val	Ile	Leu	Ile
465					470					475					480
Glu	Met	Glu	Ala	Leu	Ser	Glu	Leu	Asp	Met	Leu	Gln	Ala	Glu	Ala	Leu
			485					490					495		
Gln	Asp	Ser	Leu	Gln	His	Leu	Met	Lys	Val	Gln	Gly	Thr	Ile	Lys	Trp
			500					505					510		
Arg	Glu	Asp	His	Ile	Ala	Asn	Lys	Arg	Ser	Leu	Asn	Ser	Lys	Phe	Trp
		515					520					525			
Lys	His	Val	Arg	Tyr	Gln	Met	Pro	Val	Pro	Ser	Lys	Ile	Pro	Arg	Lys
	530					535					540				
Ala	Ser	Ser	Leu	Thr	Pro	Leu	Ala	Ala	Gln	Lys	Gln				
545					550				555						

(SEQ ID NO:3)

Fig. 7B

```
aaagagagggc tggctgttgt atttagtaaa gctataaagc tgtaagagaa attggccttc 60
tgagttgtga aactgtgggc agaaagttga ggaagaaaga actcaagtac aacccaatga 120
ggttgagata taggctactc ttcccaactc agtcttgaag agtatcacca actgcctcat 180
gtgtggtgac cttcactgtc gtatgccagt gactcatctg gagtaatctc aacaacgagt 240
taccaatact tgctcttgat tgataaacag aatgggggtt tggatcttag caattctcac 300
aattctcatg tattccacag cagcaaagtt tagtaaacia tcatgggggc tggaaaatga 360
ggctttaatt gtaagatgtc ctagacaagg aaaacctagt tacaccgtgg attggtatta 420
ctcacaaaca aacaaaagta ttcccactca ggaaagaaat cgtgtgtttg cctcaggcca 480
acttctgaag tttctaccag ctgcagttgc tgattctggt atttatacct gtattgtcag 540
aagtcccaca ttcaatagga ctggatatgc gaatgtcacc atatataaaa aacaatcaga 600
ttgcaatggt ccagattatt tgatgtattc aacagtatct ggatcagaaa aaaattccaa 660
aatttattgt cctaccattg acctctacaa ctggacagca cctcttgagt ggtttaagaa 720
ttgtcaggct cttcaaggat caaggtagag ggccgacaag tcatttttgg tcattgataa 780
tgtgatgact gaggacgcag gtgattacac ctgtaaattt atacacaatg aaaatggagc 840
caattatagt gtgacggcga ccaggctcct caccgtcaag gatgagcaag gcttttctct 900
gtttccagta atcggagccc ctgcacaaaa tgaaataaag gaagtggaaa ttggaaaaaa 960
cgcaaaccta acttgctctg cttgttttgg aaaaggcact cagttcttgg ctgccgtcct 1020
gtggcagctt aatggaacaa aaattacaga ctttggtgaa ccaagaattc aacaagagga 1080
agggcaaaat caaagtttca gcaatgggct ggcttgtcta gacatgggtt taagaatagc 1140
tgacgtgaag gaagaggatt tattgtcgca gtacgactgt ctggccctga atttgcattg 1200
cttgagaagg cacaccgtaa gactaagtag gaaaaatcca attgatcatc atagcatcta 1260
ctgcataatt gcagtatgta gtgtattttt aatgctaate aatgtcctgg ttatcatcct 1320
aaaaatgttc tggattgagg ccactctgct ctggagagac atagctaaac cttacaagac 1380
taggaatgat ggaaagctct atgatgctta tgttgtctac ccacggaact acaaateccag 1440
tacagatggg gccagtcgtg tagagcactt tgttcaccag attctgcctg atgttcttga 1500
aaataaatgt ggctatacct tatgcattta tgggagagat atgctacctg gagaagatgt 1560
agtcactgca gtggaaacca acatacgaaa gagcaggcgg cacattttca tcctgacccc 1620
tcagatcact cacaataagg agtttgccta cgagcaggag gttgccctgc actgtgccct 1680
catccagaac gacgccaagg tgatacttat tgagatggag gctctgagcg agctggacat 1740
gctgcaggct gaggcgcttc aggactccct ccagcatctt atgaaagtac aggggaccat 1800
caagtggagg gaggaccaca ttgccaataa aaggctcctg aattctaaat tctggaagca 1860
cgtgaggtac caaatgcctg tgccaagcaa aattcccaga aaggcctcta gtttgactcc 1920
cttggctgcc cagaagcaat agtgctgct gtgatgtgca aaggcatctg agtttgaagc 1980
tttctgact tctcttagct ggcttatgcc cctgcactga agtgtgagga gcaggaatat 2040
taaagggatt caggcctc 2058
(SEQ ID NO:4)
```

Fig. 8

METHODS OF TREATING A SUBJECT HAVING HEART FAILURE

CLAIM OF PRIORITY

This application is a continuation of U.S. patent application Ser. No. 13/552,553, filed on Jul. 18, 2012 (issued as U.S. Pat. No. 8,748,110), which claims the benefit of U.S. Provisional Patent Application Ser. No. 61/508,923, filed on Jul. 18, 2011, and U.S. Provisional Patent Application Ser. No. 61/509,359, filed on Jul. 19, 2011, the entire contents of which are herein incorporated by reference.

TECHNICAL FIELD

This invention relates to methods of treating cardiovascular diseases using exercise therapy, and predicting the efficacy of exercise therapy.

BACKGROUND

Circulating biomarkers play a critical role in the diagnosis and management of patients with chronic heart failure (Braunwald, *N. Engl. J. Med.* 358:2148-2159, 2008). Natriuretic peptides, such as brain natriuretic peptide (BNP) and N-terminal pro-BNP (NT-proBNP), have been demonstrated to be powerful tools for the diagnosis, risk stratification, and management of patients with heart failure (Felker et al., *Canadian Med. Assoc. J.* 175:611-617, 2006). In addition to being useful for clinical management, biomarkers can provide insights into the mechanisms underlying important physiologic relationships. Exercise intolerance, typically manifested as exertional dyspnea, is a major morbidity of chronic heart failure. Both maximal (e.g., as measured by peak oxygen uptake [peak VO_2]) (Aaronson et al., *Circulation* 95:2660-2667, 1997; Mancini et al., *Circulation* 83:778-786, 1991) and submaximal exercise capacity (e.g., as measured by distance in the 6-minute walk test) (Bittner et al., *JAMA* 270:1702-1707, 1993) have been demonstrated to be of substantial prognostic importance in chronic heart failure.

A variety of therapies can be used to treat patients diagnosed with a cardiovascular disease. For example, exercise therapy is commonly used to treat patients diagnosed with a cardiovascular disease (see, for example, Korhonen et al., *J. Womens Health* 20:1051-1064, 2011).

SUMMARY

Applicants have discovered a correlation between the level of soluble ST2 and the efficacy of exercise therapy in subjects having a cardiovascular disease, and a correlation between soluble ST2 level, exercise therapy, and the risk of an adverse outcome (e.g., death) in a subject having a cardiovascular disease.

Provided herein are methods of treating a subject having a cardiovascular disease that, in some embodiments, include determining a level of soluble ST2 in a biological sample from the subject, identifying a subject that has a decreased level of soluble ST2 in the biological sample compared to a risk reference level of soluble ST2, and selecting the identified subject for exercise therapy. Also provided are methods of treating a subject having a cardiovascular disease that include determining a level of soluble ST2 in a biological sample from the subject, identifying a subject that has an elevated level of soluble ST2 in the biological sample

compared to a risk reference level of soluble ST2; and instructing the identified subject not to begin, to discontinue, or to avoid exercise therapy.

Also provided are methods of selecting a therapy for a subject having a cardiovascular disease that include determining a level of soluble ST2 in a biological sample from the subject, and comparing the level of soluble ST2 in the biological sample to a risk reference level of soluble ST2, where a decreased level of soluble ST2 in the biological sample compared to the risk reference level indicates that the subject should begin or continue exercise therapy, and an elevated level of soluble ST2 in the biological sample compared to the risk reference level indicates that the subject should not begin or should discontinue exercise therapy.

Also provided are methods of identifying a subject having a cardiovascular disease that will benefit from exercise therapy that include determining a level of soluble ST2 in a biological sample from the subject, and selecting a subject that has a decreased level of soluble ST2 in the biological sample as compared to a risk reference level of soluble ST2, where the selected subject is identified as a subject that will benefit from exercise therapy. Also provided are methods of identifying a subject having a cardiovascular disease that will not benefit from exercise therapy that include determining a level of soluble ST2 in a biological sample from the subject, and selecting a subject that has an elevated level of soluble ST2 in the biological sample compared to a risk reference level of soluble ST2, where the selected subject is identified as a subject that will not benefit from exercise therapy.

Also provided are methods of determining whether a subject having a cardiovascular disease should begin, continue, not begin, or discontinue exercise therapy that include determining a level of soluble ST2 in a biological sample from the subject, where a decreased level of soluble ST2 in the biological sample compared to a risk reference level of soluble ST2 indicates that the subject should begin or continue exercise therapy, and an elevated level of soluble ST2 indicates that the subject should not begin or discontinue exercise therapy.

Also provided are methods of determining whether a subject having a cardiovascular disease should discontinue or continue exercise therapy that include determining a level of soluble ST2 in a biological sample from the subject at a first time point before or after the start of exercise therapy, and determining a level of soluble ST2 in a biological sample from the subject undergoing exercise therapy at a second time point after the start of exercise therapy and after the first time point, where an elevation in the level of soluble ST2 in the biological sample at the second time point compared to the level of soluble ST2 in the biological sample at the first time point indicates that the subject should discontinue exercise therapy, and a decrease in the level of soluble ST2 in the biological sample at the second time point compared to the level of soluble ST2 in the biological sample at the first time point indicates that the subject should continue exercise therapy.

In some embodiments of the methods described herein, the methods include determining that the subject has a level of soluble ST2 that is above a first reference level (e.g., a first level indicating that the subject has a cardiovascular disease, or is at risk of an adverse cardiovascular event; e.g., as described in U.S. Pat. No. 7,998,683; US2011/0262941; US2012/0040381; U.S. Pat. No. 8,090,562; US2012/0065897; U.S. Pat. Nos. 7,670,769; 7,655,415; 7,989,210; US2011/0250703; U.S. Pat. Nos. 7,432,060; 7,985,558;

US2011/028088, all of which are incorporated herein by reference) and below a second, risk reference level (e.g., below a level indicating that the subject is at risk of an adverse event if they engage in exercise, as described herein), and the subject is selected for exercise therapy. Thus in some embodiments, the methods include determining that the subject has a level of ST2 that falls within a range that is associated with the presence of a cardiovascular disease that would benefit from exercise therapy and the absence of high risk of an adverse event associated with exercise therapy.

Also provided are methods of reducing the risk of an adverse outcome in a subject having a cardiovascular disease that include determining a level of soluble ST2 in a biological sample from the subject, identifying a subject that has a decreased level of soluble ST2 in the biological sample compared to a risk reference level of soluble ST2, and selecting the identified subject for exercise therapy. Also provided are methods of reducing the risk of an adverse outcome in a subject having a cardiovascular disease that include determining a level of soluble ST2 in a biological sample from the subject, identifying a subject that has an elevated level of soluble ST2 in the biological sample compared to a risk reference level of soluble ST2, and instructing the subject to not begin or to discontinue exercise therapy. In some embodiments of these methods, the risk of adverse outcome is risk of death.

Also provided are methods of predicting the efficacy of exercise therapy in a subject having a cardiovascular disease that include determining a level of soluble ST2 in a biological sample from the subject, and comparing the level of soluble ST2 in the biological sample to an efficacy reference level of soluble ST2, where a decreased level of soluble ST2 in the biological sample compared to the efficacy reference level of soluble ST2 indicates that the exercise therapy will be effective in the subject, and an elevated level of soluble ST2 in the biological sample compared to the efficacy reference level of soluble ST2 indicates that the exercise therapy will not be effective in the subject.

In any of the methods described herein, the biological sample contains blood or serum. In any of the methods described herein, the determining is performed using an antibody or an antibody fragment that binds to soluble ST2. In any of the methods described herein, the reference level of soluble ST2 is a predetermined threshold value. In any of the methods described herein, the reference level of soluble ST2 is a level of soluble ST2 in a healthy subject. In any of the methods described herein, the subject is hypercholesterolemic, hypertriglyceridemic, hyperlipidemic, a smoker, hypertensive, or has a body mass index of greater than 30. Some embodiments of the methods described herein further include determining a level of cardiac troponin I, B-type natriuretic peptide, atrial natriuretic peptide, or C-reactive protein in the biological sample. Some embodiments of the methods described herein further include determining a level of level of cardiac troponin I, B-type natriuretic peptide, atrial natriuretic peptide, or C-reactive protein in the biological sample at the first time point or the biological sample at the second time point.

In some embodiments of any of the methods described herein, the cardiovascular disease is selected from the group of: cardiac hypertrophy, myocardial infarction, stroke, arteriosclerosis, and heart failure. In some embodiments of any of the methods described herein, the subject is administered at least one therapeutic agent selected from the group of: anti-inflammatory agents, anti-thrombotic agents, anti-coagulants, anti-platelet agents, lipid-reducing agents (e.g., a

statin), direct thrombin inhibitors, glycoprotein IIb/IIIb receptor inhibitors, calcium channel blockers, beta-adrenergic receptor blockers, cyclooxygenase-2 inhibitors, and renin-angiotensin-aldosterone system (RAAS) inhibitors. In some embodiments, the RAAS inhibitor is selected from the group of: an angiotensin-converting enzyme inhibitor, an angiotensin II receptor blocker, and an aldosterone antagonist.

As used herein, the term "cardiovascular disease" refers to a disorder of the heart and blood vessels, and includes disorders of the arteries, veins, arterioles, venules, and capillaries. Non-limiting examples of cardiovascular diseases include cardiac hypertrophy, myocardial infarction, stroke, arteriosclerosis, and heart failure. Additional examples of cardiovascular diseases are known in the art.

By the term "soluble ST2" is meant a soluble protein containing a sequence at least 90% identical (e.g., at least 95%, 96%, 97%, 98%, 99%, or 100% identical) to NCBI Accession No. NP_003847.2 (SEQ ID NO: 1) or containing a sequence at least 90% identical (e.g., at least 95%, 96%, 97%, 98%, 99%, or 100% identical) to amino acids 19-328 of SEQ ID NO: 1, or a nucleic acid containing a sequence at least 90% identical (e.g., at least 95%, 96%, 97%, 98%, 99%, or 100% identical) to NCBI Accession No. NM_003856.2 (SEQ ID NO: 2) or containing a sequence at least 90% identical (e.g., at least 95%, 96%, 97%, 98%, 99%, or 100% identical) to nucleotides 285 to 1214 of SEQ ID NO: 2.

By the term "elevated" or "elevation" is meant a difference, e.g., the presence of a statistically significant or detectable increase in a determined or measured level (e.g., a human soluble ST2 protein level) compared to a reference level (e.g., a level of human soluble ST2 in a subject not having a disease, a subject not presenting with two or more symptoms of a disease, or a subject not identified as being at risk of developing a disease, or a threshold level of human soluble ST2). In some embodiments, the reference is a threshold level, and any level above that is considered "elevated." Additional reference levels of human soluble ST2 are described herein and are known in the art.

By the term "reference level" is meant a threshold level or a level in a control subject or control patient population. A reference level will depend on the assay performed and can be determined by one of ordinary skill in the art. Non-limiting examples of reference levels are described herein and are known in the art. Reference levels of human soluble ST2 can be determined using methods known in the art.

In some embodiments, the reference level is a risk reference level, e.g., a risk reference level of soluble ST2 in a subject who experienced or was more likely to experience an adverse outcome and engaged in exercise, a level in a population of subjects who experienced or were more likely to experience an adverse outcome and engaged in exercise, or a threshold level of soluble ST2 above which the risk of an adverse outcome is increased in those who engage in exercise therapy.

In some embodiments, the reference level is an efficacy reference level, e.g., an efficacy reference level of soluble ST2 is a level in a subject who experienced a therapeutic benefit from exercise therapy, a level in a population of subjects who experienced a therapeutic benefit from exercise therapy, or a threshold level of soluble ST2 below which the subject is likely to experience a therapeutic benefit from exercise therapy.

By the term "additional marker" is meant a protein, nucleic acid, lipid, or carbohydrate, or a combination (e.g., two or more) thereof, that is diagnostic of the presence of a

particular disease. The methods described herein can include detecting the level of soluble human ST2 and at least one additional marker in a biological sample from a subject. Non-limiting examples of additional markers that can be detected include: proANP, NT-proANP, ANP, proBNP, NT-proBNP, BNP, troponin, CRP, creatinine, Blood Urea Nitrogen (BUN), liver function enzymes, albumin, and bacterial endotoxin; and those markers described in U.S. Patent Application Publication Nos.: 2007/0248981; 2011/0053170; 2010/0009356; 2010/0055683; and 2009/0264779 (each of which is hereby incorporated by reference).

By the term “hypertriglyceridemia” is meant a triglyceride level that is greater than or equal to 180 ng/mL (e.g., greater than or equal to 200 ng/mL).

By the term “hypercholesterolemia” is meant an increased level of at least one form of cholesterol or total cholesterol in a subject. For example, a subject with hypercholesterolemia can have a high density lipoprotein (HDL) level of ≥ 40 mg/dL (e.g., >50 mg/dL or >60 mg/dL), a low density lipoprotein (LDL) level of >130 mg/dL (e.g., >160 mg/dL or >200 mg/dL), and/or a total cholesterol level of >200 mg/dL (e.g., 240 mg/dL).

By the term “hypertension” is meant an increased level of systolic and/or diastolic blood pressure. For example, a subject with hypertension can have a systolic blood pressure that is >120 mmHg (e.g., >140 mmHg or >160 mmHg) and/or a diastolic blood pressure that is >80 mmHg (e.g., >90 mmHg or >100 mmHg).

By the term “healthy subject” is meant a subject that does not have a disease (e.g., cardiovascular disease). For example, a healthy subject has not been diagnosed as having a disease and is not presenting with two or more (e.g., two, three, four, or five) symptoms of a disease state.

By “risk of death” is meant the risk of death in a subject from a disease or complications associated with a disease (e.g., a cardiovascular disease) compared to a reference population. The term risk of death as used herein excludes intentional or accidental death, e.g., death by blunt or crushing trauma, such as a car accident.

As used herein, a “biological sample” includes one or more of blood, serum, plasma, urine, and body tissue. Generally, a biological sample is a sample containing serum, blood, or plasma.

By the term “statin” is meant a therapeutic molecule that inhibits the enzyme HMG-CoA reductase. Non-limiting examples of statins include: atorvastatin, fluvastatin, lovastatin, pitavastatin, pravastatin, rosuvastatin, and simvastatin. Additional examples of statins are known in the art.

By “adverse outcome” is meant any detrimental event that occurs in a subject as a result of a disease (e.g., a cardiovascular disease). Non-limiting examples of adverse outcomes in a subject having a cardiovascular disease include: organ failure, organ transplantation, hospitalization or rehospitalization, recurrence of one or more symptoms of a cardiovascular disease, development of one or more additional symptoms of a cardiovascular disease, an increase in the frequency, intensity, or duration of one or more symptoms of a cardiovascular disease experienced by the subject, a first or subsequent myocardial infarction, or death (mortality). In preferred embodiments, the adverse outcome is mortality.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Methods and materials are described herein for use in the present invention; other, suitable methods and materials known in the art can also be used.

The materials, methods, and examples are illustrative only and not intended to be limiting. All publications, patent applications, patents, sequences, database entries, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control.

The details of one or more embodiments of the invention are set forth in the accompanying drawings and the description below. Other features, objects, and advantages of the invention will be apparent from the description and drawings, and from the claims.

DESCRIPTION OF DRAWINGS

FIG. 1 is a Kaplan-Meier graph showing the surviving proportion of heart failure subjects over time within the identified groups. The survival data for subjects having a soluble ST2 level less than or equal to 20.4 ng/mL (top line); greater than 20.4 ng/mL, but less than or equal to 28.6 ng/mL (middle line); and greater than 28.6 ng/mL (bottom line) are shown (log-rank test, $p < 0.001$).

FIG. 2 is a graph and table showing the percentage of mortality in heart failure subjects over time. The percentage mortality data for heart failure subjects having a soluble ST2 level equal to or less than 35 ng/mL (left bar for each time point), or greater than 35 ng/mL (right bar for each time point) are shown. The table also shows the relative risk of mortality at each time point for subjects having a soluble ST2 level greater than 35 ng/mL compared to subjects having a soluble ST2 level less than or equal to 35 ng/mL.

FIG. 3 is a Kaplan-Meier graph showing the surviving proportion of heart failure subjects over time within the identified groups. The survival data for subjects having a soluble ST2 level equal to or less than 35 ng/mL and performing an exercise treatment regimen (top line), subjects having a level of soluble ST2 greater than 35 ng/mL and not performing an exercise treatment regimen (second line from top), subject having a level of soluble ST2 greater than 35 ng/mL and not performing an exercise treatment regime (second line from bottom), and subjects having a level of soluble ST2 equal to or less than 35 ng/mL and performing an exercise treatment regime (bottom line) are shown.

FIG. 4 is a graph of the hazard ratio for mortality within 1 year in heart failure subjects performing an exercise treatment regime compared to heart failure subjects not performing an exercise treatment regime having the different soluble ST2 levels shown.

FIG. 5 shows the protein sequence of Soluble Human ST2. Signal Peptide Amino acids 1-18 (underlined); Mature Peptide after removal of signal sequence amino acids (19-328).

FIG. 6 shows the mRNA sequence of soluble human ST2. Nucleotides 285 to 1214 encode the amino acid sequence (without the signal sequence) of soluble human ST2.

FIGS. 7A-B shows the protein sequence of the long form of human St2 (membrane-bound). Signal Peptide Amino acids 1-18 (underlined); Mature Peptide after removal of signal sequence amino acids (19-556).

FIG. 8 shows the mRNA sequence of human ST2. Nucleotides 326 to 1939 encode the amino acid sequence (without the signal sequence) of the long form of human ST2.

DETAILED DESCRIPTION

Provided herein are methods of treating a subject having a cardiovascular disease; selecting a therapy for a subject

having a cardiovascular disease; identifying a subject that will benefit or will not benefit from exercise therapy; determining whether a subject having a cardiovascular disease should begin, continue, or discontinue exercise therapy; and predicting the efficacy of exercise therapy in a subject having a cardiovascular disease. Also provided are methods of reducing the risk an adverse outcome (e.g., death) in a subject having a cardiovascular disease. These methods require determining a level of soluble ST2 in a biological sample from the subject.

Cardiovascular Diseases

A cardiovascular disease is a disorder of the heart and blood vessels (e.g., disorders of the arteries, veins, arterioles, venules, and capillaries). Cardiovascular diseases can be diagnosed using methods known in the art. Non-limiting examples of cardiovascular disease include congestive heart failure, stroke, acute coronary artery disease, arrhythmia, asymmetric septal hypertrophy (e.g., left ventricular hypertrophy with resultant diastolic dysfunction), cardiomyopathy, valvular dysfunction, pericarditis, atherosclerosis, and myocardial infarction. A subject can be diagnosed as having a cardiovascular disease by a medical professional (e.g., a physician, a physician's assistant, a nurse, a nurse's assistant, or a laboratory technician) using exemplary methods described herein. Additional methods for diagnosing a cardiovascular disease are known in the art.

Heart failure is a clinical syndrome of diverse etiologies linked by the common feature of impaired heart pumping and characterized by the failure of the heart to pump blood commensurate with the requirements of the metabolizing tissues. Heart failure can be diagnosed in a subject by the observation of one or more of the following non-limiting symptoms in a subject: dyspnea, fatigue and weakness, edema in the legs, ankles, and feet, rapid or irregular heartbeat, reduced ability to exercise, persistent cough or wheezing, white or pink blood-tinged phlegm, abdominal swelling (ascites), sudden weight gain from fluid retention, lack of appetite, nausea, difficulty concentrating or decreased alertness, and chest pain. Additional non-limiting methods for diagnosing heart failure in a subject include the use of commercially available diagnostic tests (e.g., enzyme-linked immunosorbent assays) known in the art.

A myocardial infarction is a focus of necrosis resulting from inadequate perfusion of the cardiac tissue. Myocardial infarction generally occurs from an abrupt decrease in coronary blood flow that follows a thrombotic occlusion of a coronary artery previously narrowed by atherosclerosis. Generally, a myocardial infarction occurs when an atherosclerotic plaque fissures, ruptures, or ulcerates, and a mural thrombus forms leading to coronary artery occlusion. Non-limiting methods of diagnosing a myocardial infarction include the use of a number of commercially available diagnostic tests known in the art. Generally, these diagnostic tests may be divided into four main categories: (1) nonspecific indexes of tissue necrosis and inflammation; (2) electrocardiograms; (3) serum enzyme changes (e.g., creatine phosphokinase levels); and (4) cardiac imaging. A myocardial infarction can also be diagnosed by the observation of one or more of the following symptoms in a subject: chest pain (typically on the left side of the body), neck or jaw pain, shoulder or arm pain, clammy skin, dyspnea, nausea, and vomiting. Additional methods of diagnosing a myocardial infarction are known in the art.

A stroke can be diagnosed in a subject by the observation of one or more symptoms and/or by a physical examination (e.g., interventional and non-interventional diagnostic tools, such as computed tomography and magnetic resonance

imaging). Non-limiting symptoms of a stroke include: paralysis, weakness, decreased sensation and/or vision, numbness, tingling, aphasia (e.g., inability to speak or slurred speech, or difficulty reading or writing), agnosia (i.e., inability to recognize or identify sensory stimuli), loss of memory, coordination difficulties, lethargy, sleepiness or unconsciousness, lack of bladder or bowel control, and cognitive decline (e.g., dementia, limited attention span, and inability to concentrate). In some examples, medical imaging techniques can be used identify a subject having an infarct or a hemorrhage in the brain.

Cardiac hypertrophy is typically characterized by left ventricular hypertrophy, usually of a non-dilated chamber, that occurs without an obvious antecedent cause. Cardiac hypertrophy can be diagnosed through the use of electrocardiography or echocardiography.

Arteriosclerosis is a cardiovascular disease characterized by a hardening or loss of elasticity. Arteriosclerosis can be diagnosed by the detection of one or more of the following physical symptoms: a weak or absent pulse below the narrowed area of an artery, decreased blood pressure in an affected limb, bruits caused by turbulent flow in an artery, an aneurysm in the abdomen or behind the knee, poor wound healing, and increased levels of cholesterol. Atherosclerosis can also be detected using imaging techniques including, but not limited to: Doppler ultrasound, electrocardiography, angiography, computed tomography, or magnetic resonance (e.g., magnetic resonance angiography).

A subject can be diagnosed as having a cardiovascular disease following admission to a hospital or following presentation to a health care clinic. Effective therapeutic treatment of a cardiovascular disease can be determined by observing a decrease in the number of symptoms of a cardiovascular disease in a subject or a decrease in the frequency, intensity, and/or duration of one or more symptoms of a cardiovascular disease (e.g., any of the symptoms described herein) in a subject. Effective therapeutic treatment of a cardiovascular disease can also be determined by detecting a decrease in the levels of one or more markers of a cardiovascular disease (e.g., any of the markers of cardiovascular disease known in the art or described herein) in a biological sample from a subject over time (e.g., a significant decrease in the level of at least one marker at a second time point compared to the level of the biomarker at a first time point prior to the start of treatment or at an earlier time point during the treatment period). Successful treatment of a cardiovascular disease can also be determined by a decreased risk of an adverse event (e.g., a decreased risk of death, hospitalization or rehospitalization, organ failure, organ transplantation, or a first or subsequent myocardial infarction) (e.g., compared to the risk of an adverse event in patient population diagnosed with the same cardiovascular disease but receiving no treatment or a different treatment).

Exercise Therapy

Subjects diagnosed as having a cardiovascular disease are often directed/instructed by a healthcare provider to perform an exercise therapy regime. Exercise therapy is a treatment regime that involves the periodic performance of physical activity by a subject (e.g., a subject diagnosed as having a cardiovascular disease). The physical activity performed during this type of therapy can be aerobic (e.g., walking, jogging/running, swimming, biking, or rowing) or anaerobic (e.g., weight lifting or resistance training) exercise. The physical activity can be performed in the presence of a health care professional (e.g., a physical therapist, a nurse, a nurse's assistant, a physician's assistant, or a physician). The physical activity can be performed at least once a week

(e.g., at least once a day, twice a day, two times a week, three times a week, four times a week, five times a week, or six times a week). In some embodiments, a single episode of physical activity within the exercise therapy regime can last between 5 minutes and 3 hours, between 5 minutes and 2 hours, between 5 minutes and 1 hour, or between 10 minutes and 1 hour. The intensity and/or type of the physical activity performed by the subject can vary with the physical condition of the subject (e.g., age, severity of cardiovascular disease, additional disease states, weight, and blood pressure). A subject can continue to perform an exercise therapy regime over an extended period of time (e.g., over a period of 1 month to 1 year, 1 month to 2 years, 1 year to 3 years, 2 years to 5 years, or 4 years to 10 years). In some embodiments, a subject can be monitored by a health care professional to adjust one or more parameters of the exercise therapy regime, including the frequency, intensity, length of individual episodes of physical activity, and the type of physical activity performed. The efficacy of exercise therapy in a subject having a cardiovascular disease can be determined using any of the methods for determining successful treatment of a cardiovascular disease (e.g., those methods described herein or known in the art). A subject can continue to perform an exercise therapy regime until instructed by a health care professional to discontinue the exercise therapy or until a specific therapeutic outcome has been achieved (e.g., a decrease in the number of symptoms of a cardiovascular disease has been achieved, a decrease in severity, intensity, or frequency of one or more symptoms of a cardiovascular disease has been achieved, or a decrease in the levels of one or more markers of a cardiovascular disease has been achieved in the subject).

ST2

The ST2 gene is a member of the interleukin-1 receptor family whose protein product exists both as a trans-membrane form as well as a soluble receptor that is detectable in serum (Kieser et al., *FEBS Lett.* 372(2-3):189-193, 1995; Kumar et al., *J. Biol. Chem.* 270(46):27905-27913, 1995; Yanagisawa et al., *FEBS Lett.* 302(1):51-53, 1992; Kuroiwa et al., *Hybridoma* 19(2):151-159, 2000). Soluble ST2 was described to be markedly up-regulated in an experimental model of heart failure (Weinberg et al., *Circulation* 106(23):2961-2966, 2002), and data suggest that human soluble ST2 concentrations are also elevated in those with chronic severe heart failure (Weinberg et al., *Circulation* 107(5):721-726, 2003), as well as in those with acute myocardial infarction (Shimpo et al., *Circulation* 109(18):2186-2190, 2004).

Without wishing to be bound by theory, the transmembrane form of ST2 is thought to play a role in modulating responses of T helper type 2 cells (Lohning et al., *Proc. Natl. Acad. Sci. U.S.A.* 95(12):6930-6935, 1998; Schmitz et al., *Immunity* 23(5):479-490, 2005), and may play a role in development of tolerance in states of severe or chronic inflammation (Brint et al., *Nat. Immunol.* 5(4):373-379, 2004), while the soluble form of ST2 is up-regulated in growth stimulated fibroblasts (Yanagisawa et al., 1992, supra). Experimental data suggest that the ST2 gene is markedly up-regulated in states of cardiomyocyte stretch (Weinberg et al., 2002, supra) in a manner analogous to the induction of the BNP gene (Bruneau et al., *Cardiovasc. Res.* 28(10):1519-1525, 1994).

Tominaga et al. (*FEBS Lett.* 258:301-304, 1989) isolated murine genes that were specifically expressed by growth stimulation in BALB/c-3T3 cells. Haga et al. (*Eur. J. Biochem.* 270:163-170, 2003) describes that the ST2 gene was named on the basis of its induction by growth stimulation. The ST2 gene encodes two protein products: ST2 or sST2,

which is a soluble secreted form, and ST2L, a transmembrane receptor form that is very similar to the interleukin-1 receptors. The HUGO Nomenclature Committee designated the human homolog of ST2, the cloning of which was described in Tominaga et al., *Biochim. Biophys. Acta.* 1171:215-218, 1992, as Interleukin 1 Receptor-Like 1 (IL1RL1). The two terms are used interchangeably in the art.

The mRNA sequence of the shorter, soluble isoform of human ST2 can be found at GenBank Acc. No. NM_003856.2 (SEQ ID NO: 2), and the polypeptide sequence is at GenBank Acc. No. NP_003847.2 (SEQ ID NO: 1). The mRNA sequence for the longer form of human ST2 is at GenBank Acc. No. NM_016232.4 (SEQ ID NO: 4), and the polypeptide sequence is at GenBank Acc. No. NP_057316.3 (SEQ ID NO: 3). Additional information is available in the public databases at GeneID: 9173, MIM ID #601203, and UniGene No. Hs.66.

Methods for detecting and measuring soluble ST2 are known in the art, e.g., as described in U.S. Patent Application Publication Nos. 2003/0124624, 2004/0048286, and 2005/0130136, and U.S. patent application Ser. No. 13/083,333 and PCT Application No. PCT/US2011/031801, the entire contents of which are incorporated herein by reference. These U.S. patent application publications describe methods of determining the level of soluble ST2 using an antibody or antibody fragment that binds to soluble ST2.

In some embodiments, the antibody is a monoclonal antibody produced by the hybridoma deposited at the ATCC and designated by Patent Deposit Designation PTA-10431 (the 7E4 antibody), or is an antigen-binding fragment of the antibody produced by the hybridoma deposited at the ATCC and designated by the Patent Deposit Designation PTA-10431 (fragments of the 7E4 antibody). In some embodiments, the antibody is a monoclonal antibody produced by the hybridoma deposited at the ATCC and designated by Patent Deposit Designation PTA-10432 (the 9F8 antibody), or is an antigen-binding fragment of the antibody produced by the hybridoma deposited at the ATCC and designated by the Patent Deposit Designation PTA-10432 (fragments of the 9F8 antibody). Combinations of two or more of the antibodies or fragments described herein (e.g., two or more of a 7E4 antibody, 7E4 antibody fragments, 9F8 antibody, and 9F8 antibody fragments) are useful in any of the methods described herein.

The human soluble ST2-binding monoclonal antibodies produced by the hybridomas designated by Patent Deposit Designation PTA-10431 and Patent Deposit Designation PTA-10432 were each generated by immunizing a non-human mammal with a recombinant human soluble ST2 isolated from human embryonic kidney (HEK)-293 cells.

Kits for measuring soluble ST2 are also commercially available, e.g., the ST2 ELISA Kit manufactured by Medical & Biological Laboratories Co., Ltd. (MBL International Corp., Woburn, Mass.), No. 7638. In addition, devices for measuring ST2 and other biomarkers are described in U.S. Patent Application Publication No. 2005/0250156 (incorporated herein by reference in its entirety).

As described in detail herein, soluble ST2 levels can be determined in any biological sample from a subject, including blood, serum, plasma, urine, and body tissue. Generally, the level of soluble ST2 is determined in a sample containing serum, blood, or plasma. The level of soluble ST2 can be determined in a biological sample that has been stored for a period of time (e.g., for at least 1 hour, 1 day, 1 week, or 1 month) at a temperature at or below 10° C. (e.g., below 0° C., below -20° C., or around -196° C.).

Elevated concentrations of soluble ST2 are markedly prognostic for death in patients with heart failure, with a dramatic divergence in survival curves for those with elevated soluble ST2 soon after presentation (Weinberg et al., *Circulation* 107:721-726, 2003; Mueller et al., *Clin Chem.* 54(4):752-756, 2008; Daniels et al., *Am. Heart J.* 160:721-728, 2010; Ky et al., *Circ. Heart Fail.* 4(2):180-187, 2011; and Manzano-Fernandez et al., *Am. J. Cardiol.* 107:259-267, 2011). The relationship between soluble ST2 and death in heart failure patients was shown to be independent of etiology, and superseded all other biomarker predictors of mortality in this setting, including other markers of inflammation, myonecrosis, renal dysfunction, and most notably NT-proBNP, a marker well known as having value for predicting death in heart failure patients.

Reference Levels of ST2

As described herein, the level of soluble ST2 in a subject indicates whether a subject having a cardiovascular disease should be selected for exercise therapy, whether a subject will benefit or will not benefit from exercise therapy, or whether a subject should begin, continue, or discontinue exercise therapy. In addition, the level of soluble ST2 in a subject can be used to select a therapy including exercise for a subject having a cardiovascular disease. Additional clinical and therapeutic uses of detecting a level of soluble ST2 are described herein and are known in the art. Reference levels of human soluble ST2 can be determined using methods known in the art (e.g., using the antibodies described in U.S. patent application Ser. No. 13/083,333 and PCT Application No. PCT/US2011/031801). In general, it will be desirable to use a reference level of soluble ST2 determined using the same method as is used to determine the level of soluble ST2 in the subject.

Risk Reference Levels

The methods described herein can include comparing the level of soluble ST2 in a biological sample to a reference level of soluble ST2. A reference level of soluble ST2 can be or represent a level of soluble ST2 found in a biological sample from a subject (e.g., a control subject who experienced or was more likely to experience an adverse outcome and engaged in exercise) or a population (e.g., a population of subjects who experienced or were more likely to experience an adverse outcome and engaged in exercise), or can be or represent a threshold level of soluble ST2 above which the risk of an adverse outcome is increased in those who engage in exercise therapy. These reference levels are referred to herein as the "risk reference level." In general, the presence of a level of ST2 below the risk reference level indicates that the subject does not have an elevated risk of an adverse outcome if they engage in exercise therapy, while the presence of a level of ST2 above the risk reference level indicates that the subject has an elevated risk of an adverse outcome if they do engage in exercise therapy.

In some embodiments, the risk reference level of soluble ST2 is a threshold level of soluble ST2 or a percentile (e.g., 75th, 80th, 85th, 90th, or 95th percentile) of soluble ST2 levels in a population of subjects who experienced an adverse outcome and engaged in exercise.

In some embodiments, efficacy reference level is about 55 ng/mL, determined using the Presage ST2 kit, or the equivalent thereof.

Efficacy Reference Levels

Alternatively or in addition, the methods described herein can include comparing the level of soluble ST2 in a biological sample to a reference level of soluble ST2 that is or represents a level of soluble ST2 found in a biological sample from a subject (e.g., a control subject who experienced a therapeutic benefit from exercise therapy, i.e., did not or was less likely to experience an adverse outcome and engaged in exercise) or a population (e.g., a population of subjects who experienced a therapeutic benefit from exercise therapy, i.e., did not experience or were less likely to experience an adverse outcome and engaged in exercise), or can be or represent a threshold level of soluble ST2 below which the subject is likely to experience a therapeutic benefit from exercise therapy, i.e., a threshold level of soluble ST2 below which the likelihood of an adverse outcome is decreased in those who engage in exercise therapy. These reference levels are referred to herein as the "efficacy reference level." In general, the presence of a level of ST2 below the efficacy reference level indicates that the subject has an increased likelihood of not experiencing an adverse outcome if they engage in exercise therapy, while the presence of a level of ST2 above the efficacy reference level indicates that the subject has an elevated risk of an adverse outcome if they do engage in exercise therapy.

In some embodiments, the efficacy reference level of soluble ST2 is a threshold level of soluble ST2 or a percentile (e.g., 75th, 80th, 85th, 90th, or 95th percentile) of soluble ST2 levels in a population of subjects who experienced an adverse outcome and engaged in exercise.

In some embodiments, efficacy reference level is about 35 ng/mL, determined using the Presage ST2 kit, or the equivalent thereof.

Healthy Controls and Other Reference Levels

In some embodiments, the methods described herein can also include comparing the level of soluble ST2 in a biological sample to a reference level of soluble ST2 that represent the average level of soluble ST2 present in a population of subjects: a population of subjects diagnosed as having a specific cardiovascular disease, a population of healthy subjects not diagnosed with a disease (e.g., a healthy male patient population or a healthy female patient population), a population of subjects not at risk of developing a cardiovascular disease, or a population of subjects not presenting with two or more symptoms of a cardiovascular disease. A reference level can also be a baseline level or a level in the same patient measured at an earlier or later point in time. Additional non-limiting examples of reference levels of human soluble ST2 include the level of human soluble ST2 in a subject or a patient population that: does not have high risk cardiovascular disease; does not have renal failure; does not have hypertriglyceridemia, hypercholesterolemia, hypertension, and/or a body mass index of <30 (e.g., a BMI under 25); and/or does not suffer from a pulmonary disease, sepsis, or Kawasaki disease.

In some embodiments, the reference level of soluble ST2 is a threshold level of soluble ST2. In some embodiments, the threshold level of soluble ST2 is a median level of soluble ST2 or a percentile (e.g., 75th, 80th, 85th, 90th, or 95th percentile) of soluble ST2 levels in a healthy patient population, e.g., a healthy male patient population or a healthy female patient population (e.g., any of the values or ranges listed in Table 1).

TABLE 1

sST2 Concentrations at Specific Thresholds - US Self-Reported Healthy Cohort ¹						
Percentiles	Entire Cohort		Male		Female	
	ST2 (ng/mL)	95% CI	ST2 (ng/mL)	95% CI	ST2 (ng/mL)	95% CI
2.5	8.0	7.1 to 8.6	8.6	7.7 to 11.8	7.3	5.5 to 8.4
5	9.3	8.4 to 10.2	11.8	8.6 to 12.7	8.5	7.3 to 9.4
10	11.5	10.3 to 11.9	13.7	12.2 to 14.8	10.2	9.0 to 11.2
25	14.5	13.7 to 15.2	17.6	16.8 to 18.7	12.4	11.9 to 13.5
median	18.8	18.2 to 19.9	23.6	21.3 to 25.1	16.2	15.4 to 17.4
75	25.3	23.8 to 26.9	30.6	28.7 to 33.3	19.9	18.8 to 20.8
90	34.3	32.4 to 35.6	37.2	35.5 to 40.9	23.7	22.2 to 25.8
95	37.9	35.9 to 41.3	45.4	39.4 to 48.6	29.0	24.6 to 33.2
97.5	45.6	40.1 to 48.7	48.5	45.8 to 58.5	33.1	29.6 to 39.9

¹These levels were determined using the antibodies described in U.S. patent application Ser. No. 13/083,333 and PCT Application No. PCT/US2011/031801.

In some embodiments, the threshold level is 28.6 ng/mL, about 28 ng/mL to about 35 ng/mL, 35 ng/mL, about 35 ng/mL to about 45 ng/mL, about 45 ng/mL to about 55 ng/mL, or the range of 55 to 60 ng/mL (e.g., the entire range or any level between 55 to 60 ng/mL) (e.g., using the antibodies described in U.S. patent application Ser. No. 13/083,333 and PCT Application No. PCT/US2011/031801). Reference levels of human soluble ST2 can be determined using methods known in the art (e.g., using the antibodies described in U.S. patent application Ser. No. 13/083,333 and PCT Application No. PCT/US2011/031801). Additional reference levels of soluble ST2 are known in the art. As is known in the art, the reference level of soluble ST2 can vary based on the assay used to determine soluble ST2 levels.

Subjects

The methods described herein can be performed on a variety of subjects having a cardiovascular disease (e.g., any of the subjects described herein). In some embodiments of any of the methods, the subject has been previously diagnosed as having a cardiovascular disease. In some embodiments of any of the methods, the subject is hypercholesterolemic, hypertriglyceridemic, hyperlipidemic, a smoker, hypertensive, or has a body mass index of greater than 25 (e.g., between 25 and 30, or greater than 30). In some embodiments of any of the methods, the subject can already be receiving a therapeutic agent (e.g., one or more of the additional therapeutic agents described herein or known in the art for treating a cardiovascular disease). In some embodiments, the subject has heart failure.

In some embodiments of any of the methods, the subject can have previously been admitted to a hospital or can be receiving treatment on an outpatient basis. In some embodiments of any of the methods, the patient can be 20 to 40 years old, 40 to 50 years old, 50 to 60 years old, 60 to 70 years old, 70 to 80 years old, 80 to 90 years old, or 90 to 100 years old.

Methods of Treating a Subject Having a Cardiovascular Disease

Provided herein are methods of treating a subject having a cardiovascular disease (e.g., any of the cardiovascular diseases described herein or known in the art, e.g., heart failure). These methods include determining a level of soluble ST2 in a biological sample from the subject, identifying a subject that has a decreased (e.g., a significant or detectable decrease) level of soluble ST2 in the biological sample compared to a risk or efficacy reference level of soluble ST2 (e.g., as described herein), and selecting the

identified subject for exercise therapy (e.g., any form of exercise therapy described herein or known in the art). Also provided are methods of treating a subject that include determining a level of soluble ST2 in a biological sample from the subject, identifying a subject that has an increased (e.g., a significant or detectable increase) level of soluble ST2 in the biological sample compared to a risk or efficacy reference level of soluble ST2 (e.g., as described herein), and instructing the identified subject not to begin, to discontinue, or to avoid exercise therapy. The level of soluble ST2 in the biological sample can be determined using any of the methods described herein (e.g., methods using an antibody or antibody fragment that binds specifically to soluble ST2). The biological sample can be any of the biological samples described herein. In some embodiments, the biological sample is collected from a subject within 2 or 4 years of diagnosis with a cardiovascular disease, a myocardial infarction, or heart failure.

The selected subject can begin performance of any of the exercise therapy regimes described herein. Some embodiments of these methods further include monitoring the subject to determine whether the exercise therapy regime should be continued or altered in one or more aspects (e.g., any of the aspects of exercise therapy regimes described herein). Methods for monitoring the subject to determine whether the exercise therapy should be continued or discontinued are described herein. These methods can be performed by any health care professional (e.g., a physician, a physical therapist, a nurse, a physician's assistant, a laboratory technician, or a nursing assistant).

In some embodiments, the treating results in a reduced (e.g., a significant decrease) in the risk of an adverse outcome (e.g., any of the adverse outcomes described herein) in the subject. In some embodiments, the treating results in a decrease in the number of symptoms of a cardiovascular disease, a decrease (e.g., a significant or detectable decrease) in the intensity, frequency, or duration of one or more symptoms of a cardiovascular disease, or a decrease in the level of at least one marker of a cardiovascular disease in a biological sample from the subject.

Methods of Selecting a Therapy for a Subject Having a Cardiovascular Disease

Also provided are methods of selecting a therapy for a subject having a cardiovascular disease. These methods include determining a level of soluble ST2 in a biological sample from the subject and comparing the level of soluble ST2 in the biological sample to a risk or efficacy reference level of soluble ST2 (e.g., any of the reference levels of

soluble ST2 described herein or known in the art), where a decreased (e.g., a significant or detectable decrease) level of soluble ST2 in the biological sample compared to the risk or efficacy reference level of soluble ST2 indicates that the subject should begin or continue exercise therapy (e.g., any of the exercise therapy regimes described herein), and an elevated (e.g., a significant or detectable increase) level of soluble ST2 in the biological sample compared to the risk or efficacy reference level of soluble ST2 indicates that the subject should not begin or should discontinue exercise therapy. The level of soluble ST2 in the biological sample can be determined using any of the methods described herein, and the biological sample can be any of the biological samples described herein. In some embodiments, the biological sample is collected from a subject within 2 or 4 years of diagnosis with a cardiovascular disease, a myocardial infarction, or heart failure. These methods can be performed by any health care professional (e.g., a physician, a physical therapist, a nurse, a physician's assistant, a laboratory technician, or a nursing assistant). In some embodiments, the subject can already be performing an exercise therapy regime.

Methods of Identifying a Subject that Will Benefit/not Benefit from Exercise Therapy

Also provided herein are methods for identifying a subject that will benefit from exercise therapy (benefit from the performance of any of the exercise therapy regimes described herein). The methods include determining a level of soluble ST2 in a biological sample from the subject, and selecting a subject that has a decreased (e.g., a significant or detectable decrease) level of soluble ST2 in the biological sample compared to an efficacy reference level of soluble ST2 (e.g., any of the reference levels of soluble ST2 described herein or known in the art), where the selected subject is identified as a subject that will benefit from exercise therapy (e.g., any of the exercise therapy regimes described herein). Also provided are methods of identifying a subject that will not benefit from exercise therapy (will not benefit from the performance of any of the exercise therapy regimes described herein) that include determining a level of soluble ST2 in a biological sample from the subject, and selecting a subject that has an elevated (e.g., a significant or detectable increase) level of soluble ST2 in the biological sample compared to an efficacy reference level of soluble ST2 (e.g., any of the reference levels of soluble ST2 described herein or known in the art), where the selected subject is identified as a subject that will not benefit from exercise therapy (e.g., any of the exercise therapy regimes described herein).

In some embodiments, the biological sample is collected from a subject within 2 or 4 years of diagnosis with a cardiovascular disease, a myocardial infarction, or heart failure. In some embodiments, the efficacy reference level of soluble ST2 is a threshold soluble ST2 level of less than or equal to 35 ng/mL or a range of 55 to 60 ng/mL (e.g., the entire range or any level between 55 to 60 ng/mL). The level of soluble ST2 in the biological sample can be determined using any of the methods described herein (e.g., methods using an antibody or antibody fragment that binds specifically to soluble ST2), and the biological sample can be any of the biological samples described herein.

In some embodiments of these methods, the subject has been previously diagnosed as having a cardiovascular disease (e.g., previously diagnosed as having heart failure). In some embodiments, the subject is hypercholesterolemic, hypertriglyceridemic, hyperlipidemic, a smoker, hypertensive, or has a body mass index of greater than 25 (e.g., between

25 and 30, or greater than 30). In some embodiments, the subject can already be receiving a therapeutic agent (e.g., one or more of the additional therapeutic agents described herein or known in the art for treating a cardiovascular disease). In some embodiments, the subject can have previously been admitted to a hospital or can be receiving treatment on an outpatient basis. In some embodiments, the patient can be 20 to 40 years old, 40 to 50 years old, 50 to 60 years old, 60 to 70 years old, 70 to 80 years old, 80 to 90 years old, or 90 to 100 years old.

In some embodiments, the benefit from exercise therapy can be one or more of the following: a reduction (e.g., a significant decrease) in the risk of an adverse outcome (e.g., any of the adverse outcomes described herein) in the subject, a reduction in the number of symptoms of a cardiovascular disease, a reduction (e.g., a detectable or observable decrease) in the intensity, frequency, or duration of one or more symptoms of a cardiovascular disease, or a reduction (e.g., detectable decrease) in the levels of at least one marker of a cardiovascular disease in a biological sample from the subject (e.g., as compared to a subject or population of subjects having the same cardiovascular disease but not receiving therapy or receiving a different therapy). The benefit from exercise therapy can be determined at various time points in a subject (e.g., after at least 6 months of exercise therapy, after 1 year of exercise therapy, or after 2 years of exercise therapy).

Methods of Determining Whether a Subject should Begin, Continue, or Discontinue Exercise Therapy

Also provided are methods of determining whether a subject having a cardiovascular disease should begin or continue exercise therapy (e.g., any of the exercise therapy regimes described herein) that include determining a level of soluble ST2 in a biological sample from the subject, wherein a decreased (e.g., a significant or detectable decrease) level of soluble ST2 in the biological sample compared to a risk or efficacy reference level of soluble ST2 (e.g., any of the reference levels described herein) indicates that the subject should begin or continue exercise therapy. Also provided are methods of determining whether a subject having a cardiovascular disease should not begin or discontinue exercise therapy (e.g., any of the exercise therapy regimes described herein) that include determining a level of soluble ST2 in a biological sample from the subject, where an elevated (e.g., a significant or detectable increase) level of soluble ST2 in the biological sample compared to a risk or efficacy reference level of soluble ST2 (e.g., any of the reference levels of soluble ST2 described herein) indicates that the subject should not begin or discontinue exercise therapy.

Also provided are methods of determining whether a subject having a cardiovascular disorder should discontinue or avoid exercise therapy (e.g., any of the exercise therapy regimes described herein) that include determining a level of soluble ST2 in a biological sample from the subject at a first time point before or after the start of exercise therapy, and determining a level of soluble ST2 in a biological sample from the subject undergoing (performing) exercise therapy at a second time point after the start of exercise therapy and after the first time point, where an increase (e.g., a significant or detectable increase) in the level of soluble ST2 in the biological sample of the second time point compared to the level of soluble ST2 in the biological sample at the first time point indicates that the subject should discontinue or avoid exercise therapy. Also provided are methods of determining whether a subject having a cardiovascular disorder should continue exercise therapy (e.g., any of the exercise therapy regimes described herein) that include determining a level of

soluble ST2 in a biological sample from the subject at a first time point before or after the start of exercise therapy, and determining a level of soluble ST2 in a biological sample from the subject undergoing (performing) exercise therapy at a second time point after the start of exercise therapy and after the first time point, where a decrease (e.g., a significant or detectable decrease) or no change (e.g., no significant change) in the level of soluble ST2 in the biological sample of the second time point compared to the level of soluble ST2 in the biological sample at the first time point indicates that the subject should continue exercise therapy.

Also provided are methods of determining whether a subject having a cardiovascular disorder should continue exercise therapy (e.g., any of the exercise therapy regimes described herein) that include determining a level of soluble ST2 in a biological sample from the subject at a first time point before or after the start of exercise therapy, and instructing the subject to begin or continue exercise therapy if the level of soluble ST2 is below a risk or efficacy reference level; and determining a level of soluble ST2 in a biological sample from the subject undergoing (performing) exercise therapy at a second time point after the start of exercise therapy and after the first time point, and instructing the subject to continue the exercise therapy if the level at the second time point is still below the risk or efficacy reference level, or to stop the exercise therapy if the level at the second time point is above the risk or efficacy reference level.

In some embodiments, one or more additional levels of soluble ST2 can be determined in the subject (e.g., determined in biological samples obtained at one or more additional time points after the second time point). In some embodiments, the level of soluble ST2 is determined in a biological sample obtained from the subject having cardiovascular disease at least every month (e.g., at least every two months, at least every three months, at least every four months, at least every five months, or at least every six months) during the performance of a exercise therapy regime. In such embodiments, an increase (e.g., a significant or detectable increase) in the level of soluble ST2 in a biological sample taken at a later time point compared to the level of soluble ST2 in a biological sample taken at an earlier time point (e.g., the immediately prior biological sample), or the presence of a level above a risk or efficacy reference level, indicates that the subject should discontinue exercise therapy, and a decrease (e.g., a significant or detectable decrease) or no change (e.g., no significant change) in the level of soluble ST2 in a biological sample taken at a later time point compared to the level of soluble ST2 in a biological sample taken at an earlier time point (e.g., the immediately prior biological sample), or the presence of a level below a risk or efficacy reference level, indicates that the subject should continue exercise therapy. In some embodiments of these methods, the subjects are monitored by a health care professional (e.g., a physician, a physical therapist, a nurse, a nurse's assistant, a physician's assistant, or a laboratory technician).

In some embodiments, the biological sample is collected from a subject within 2 years of diagnosis with a cardiovascular disease, a myocardial infarction, or heart failure. The level of soluble ST2 in the biological sample can be determined using any of the methods described herein, and the biological sample can be any of the biological samples described herein. These methods can reduce (e.g., a significant reduction) in the risk of death or an adverse outcome (e.g., any of the adverse outcomes described herein) in the subject, reduce the number of symptoms of a cardiovascular disease, reduce (e.g., a detectable or observable reduction) in

the intensity, frequency, or duration in one or more symptoms of a cardiovascular disease, or reduce (e.g., detectable reduction) in the level of at least one marker (e.g., any of the markers described herein) of a cardiovascular disease in a biological sample from the subject (e.g., as compared to a subject or population of subjects having the same cardiovascular disease but not receiving therapy or receiving a different therapy).

Methods of Reducing the Risk of an Adverse Outcome in a Subject

Also provided herein are methods of reducing (e.g., a significant reduction) the risk of an adverse outcome (e.g., risk of death) in a subject having a cardiovascular disease that include determining a level of soluble ST2 in a biological sample from the subject, identifying a subject that has a decreased (e.g., a significant or detectable decrease) level of soluble ST2 in the biological sample compared to a risk reference level of soluble ST2 (e.g., as described herein), and selecting the identified subject for exercise therapy (e.g., any of the exercise therapy regimes described herein). Also provided are methods of reducing (e.g., a significant reduction) the risk of an adverse outcome (e.g., risk of death) in a subject having a cardiovascular disease that include determining a level of soluble ST2 in a biological sample from the subject, identifying a subject that has an elevated (e.g., a significant or detectable increase) level of soluble ST2 in the biological sample compared to a risk reference level of soluble ST2 (e.g., as described herein), and instructing the identified subject to avoid or discontinue exercise therapy (e.g., any of the exercise therapy regimes described herein).

In some embodiments, the adverse outcome can be one or more of: death, organ failure, organ transplantation, hospitalization or rehospitalization, recurrence of one or more symptoms of a cardiovascular disease, development of one or more additional symptoms of a cardiovascular disease, an increase in the frequency, intensity, or duration of one or more symptoms of a cardiovascular disease experienced by the subject, or a first or subsequent myocardial infarction. In some embodiments, the biological sample is collected from a subject within 2 years of diagnosis with a cardiovascular disease, a myocardial infarction, or heart failure. Any of these methods can be performed by a health care professional (e.g., a physician, a physical therapist, a nurse, a nurse's assistant, a physician's assistant, or a laboratory technician).

Methods of Predicting the Efficacy of Exercise Therapy

Also provided are methods of predicting the efficacy of exercise therapy (e.g., any of the exercise therapy regimes described herein) in a subject having a cardiovascular disease. These methods include determining a level of soluble ST2 in a biological sample from the subject, and comparing the level of soluble ST2 in the biological sample to an efficacy reference level of soluble ST2 (e.g., any of the reference levels of soluble ST2 described herein), wherein a decreased (e.g., a significant or detectable decrease) level of soluble ST2 in the biological sample compared to the efficacy reference level of soluble ST2 (e.g., as described herein) indicates that exercise therapy will be effective in the subject, and an elevated (e.g., a significant or detectable increase) level of soluble ST2 in the biological sample compared to the efficacy reference level of soluble ST2 (e.g., as described herein) indicates that the exercise therapy will not be effective in the subject. In some embodiments, the efficacy reference level of soluble ST2 is a threshold soluble ST2 level of 28.6 ng/mL, less than or equal to 35 ng/mL, about 28 ng/mL to about 35 ng/mL (e.g., the entire range or any level between 28 to 35 ng/mL), 35 ng/mL, about 35

ng/mL to about 45 ng/mL (e.g., the entire range or any level between 35 to 45 ng/mL), about 45 ng/mL to about 55 ng/mL (e.g., the entire range or any level between 45 to 55 ng/mL), about 35 ng/mL to about 55 ng/mL (e.g., the entire range or any level between 35 to 55 ng/mL), or the range of 55 to 60 ng/mL (e.g., the entire range or any level between 55 to 60 ng/mL) (e.g., determined using the antibodies described in U.S. patent application Ser. No. 13/083,333 and PCT Application No. PCT/US2011/031801). In some embodiments, the biological sample is collected from a subject within 2 years of diagnosis with a cardiovascular disease, a myocardial infarction, or heart failure. The level of soluble ST2 in the biological sample can be determined using any of the methods described herein, and the biological sample can be any of the biological samples described herein.

In some embodiments, the efficacy of exercise therapy can be one or more of the following: a reduction (e.g., a significant decrease) in the risk of an adverse outcome (e.g., any of the adverse outcomes described herein) in the subject, a reduction in the number of symptoms of a cardiovascular disease, a reduction (e.g., a detectable or observable decrease) in the intensity, frequency, or duration of one or more symptoms of a cardiovascular disease, or a reduction (e.g., detectable decrease) in the levels of at least one marker of a cardiovascular disease (e.g., any of the markers described herein or known in the art) in a biological sample from the subject (e.g., as compared to a subject or population of subjects having the same cardiovascular disease but not receiving therapy or receiving a different therapy). The efficacy of exercise therapy can be determined at various time points in a subject (e.g., after at least 6 months of exercise therapy, after 1 year of exercise therapy, or after 2 years of exercise therapy).

Additional Therapeutic Markers

Any of the methods described herein can further include determining the level of at least one additional marker (e.g., at least one additional marker of a cardiovascular disease) in a biological sample from a subject. In some embodiments, the biological sample used to determine the level of the at least one additional marker may be the same sample(s) that is used to determine a level of soluble ST2 in a subject. In some embodiments, the biological sample used to determine the level of the at least one additional marker is a different sample than the sample(s) used to determine a level of soluble ST2 in a subject. The biological sample can be any of the biological samples described herein.

The additional marker can be any protein, nucleic acid, lipid, or carbohydrate, or a combination (e.g., two or more) thereof, that is diagnostic of the presence of a particular disease (e.g., diagnostic of a cardiovascular disease). Several additional markers useful for the diagnosis of a cardiovascular disease are known in the art, and include, without limitation, cardiac troponin I, B-type natriuretic peptide (e.g., proBNP, NT-proBNP, and BNP), atrial natriuretic peptide (e.g., proANP, NT-proANP, and ANP), troponin, C-reactive protein, creatinine, Blood Urea Nitrogen (BUN), liver function enzymes, albumin, and bacterial endotoxin. Additional non-limiting markers of a cardiovascular disease are described in U.S. Patent Application Publication Nos.: 2007/0248981; 2011/0053170; 2010/0009356; 2010/0055683; and 2009/0264779 (each of which is hereby incorporated by reference). Additional markers of a cardiovascular disease are known in the art.

Methods for determining the level of the above described markers of a cardiovascular disease are known in the art. Diagnostic tests for determining the level of several of these

markers are commercially available. For example, diagnostic tests for determining the level of C-reactive protein (e.g., Exocell), B-type natriuretic peptide (e.g., Alpcos Immunoassays), atrial natriuretic peptide (e.g., Cusabio Biotech Co., Ltd.), and cardiac troponin I (e.g., Calbiotech Inc.) are commercially available.

Additional Therapeutic Treatments

In any of the methods described herein, the subject can further be administered an additional therapeutic treatment (e.g., at least one therapeutic treatment in addition to exercise therapy). In some embodiments of the methods described herein, the subject can be receiving at least one therapeutic treatment at the time the exercise therapy begins. In some embodiments, a health care professional may adjust (e.g., increase or decrease) the dosage or frequency of administration of at least one therapeutic agent administered to the subject prior to the start of exercise therapy or at a time point during the exercise therapy. In some embodiments of the methods described herein, the efficacy of the exercise therapy allows for a decrease in the number of therapeutic agents or allows for a decrease in the dose or frequency of administration of one or more therapeutic agents to a subject having a cardiovascular disease.

Non-limiting examples of therapeutic treatment of a cardiovascular disease (in addition to exercise therapy) include the administration of one or more of the following agents: statins, anti-inflammatory agents, anti-thrombotic agents, anti-coagulants, anti-platelet agents, lipid-reducing agents, direct thrombin inhibitors, glycoprotein IIb/IIIa receptor inhibitors, calcium channel blockers, beta-adrenergic receptor blockers, cyclooxygenase-2 inhibitors, and renin-angiotensin-aldosterone system (RAAS) inhibitors.

Non-limiting examples of lipid-reducing agents that can be used to treat a cardiovascular disease in a subject (alone or in combination any other therapy, including exercise therapy) include: a statin, gemfibrozil, cholestyramine, colestipol, nicotinic acid, and probucol. Statins are molecules that are capable of inhibiting the activity of HMG-CoA reductase. Non-limiting examples of statins that can be administered to a subject having a cardiovascular disease (alone or in combination with any other therapy, including exercise therapy) include: atorvastatin, cerivastatin, fluvastatin, lovastatin, pitavastatin, pravastatin, rosuvastatin, and simvastatin. Additional examples of statins and other lipid-reducing agents are known in the art.

Non-limiting examples of anti-inflammatory agents that can be used to treat a cardiovascular disease in a subject (alone or in combination any other therapy, including exercise therapy) include: Alclufenac, Alclometasone Dipropionate, Algestone Acetonide, Alpha Amylase, Amcinafal, Amcinafide, Amfenac Sodium, Amiprilose Hydrochloride, Anakinra, Anilolac, Anitrazafen, Apazone, Balsalazide Disodium, Bendazac, Benoxaprofen, Benzydamine Hydrochloride, Bromelains, Broperamole, Budesonide, Carprofen, Cicloprofen, Cintazone, Cliprofen, Clobetasol Propionate, Clobetasone Butyrate, Clopirac, Cloticasone Propionate, Cormethasone Acetate, Cortodoxone, Deflazacort, Desonide, Desoximetasone, Dexamethasone Dipropionate, Diclofenac Potassium, Diclofenac Sodium, Diflorasone Diacetate, Diflumidone Sodium, Diflunisal, Difluprednate, Diftalone, Dimethyl Sulfoxide, Drocinnolide, Endrysone, Enlimomab, Enolicam Sodium, Epirizole, Etodolac, Etofenamate, Felbinac, Fenamole, Fenbufen, Fenclofenac, Fenclozoc, Fendosal, Fempipalone, Fentiazac, Flazalone, Fluazacort, Flufenamic Acid, Flumizole, Flunisolide Acetate, Flunixin, Flunixin Meglumine, Fluocortin Butyl, Fluorometholone Acetate, Fluquazone, Flurbiprofen, Fluretofen,

Fluticasone Propionate, Furaprofen, Furobufen, Halcinonide, Halobetasol Propionate, Halopredone Acetate, Ibufenac, Ibuprofen, Ibuprofen Aluminum, Ibuprofen Piconol, Ilo-nidap, Indomethacin, Indomethacin Sodium, Indoprofen, Indoxole, Intrazole, Isoflupredone Acetate, Isoxepac, Isoxi-
 5 cam, Ketoprofen, Lofemizole Hydrochloride, Lornoxicam, Loteprednol Etabonate, Meclofenamate Sodium, Meclofe-
 namic Acid, Meclorisone Dibutyrate, Mefenamic Acid, Mesalamine, Meseclazone, Methylprednisolone Suleptan-
 ate, Morniflumate, Nabumetone, Naproxen, Naproxen
 10 Sodium, Naproxol, Nimazone, Olsalazine Sodium, Orgot-
 ein, Orpanoxin, Oxaprozin, Oxyphenbutazone, Paranyline
 Hydrochloride, Pentosan Polysulfate Sodium, Phenbutazone
 Sodium Glycerate, Pirfenidone, Piroxicam, Piroxicam Cin-
 15 namate, Piroxicam Olamine, Pirprofen, Prednazate, Prifelone, Prodolic Acid, Proquazone, Proxazole, Proxazole
 Citrate, Rimexolone, Romazarit, Salcolex, Salnacedin, Sal-
 salate, Salycilates, Sanguinarium Chloride, Seclazone, Ser-
 metacin, Sudoxicam, Sulindac, Suprofen, Talmetacin, Tal-
 niflumate, Talosalate, Tebufelone, Tenidap, Tenidap Sodium,
 Tenoxicam, Tesicam, Tesimide, Tetrydamine, Tiopinac,
 Tixocortol Pivalate, Tolmetin, Tolmetin Sodium, Triclonide,
 Triflumidate, Zidometacin, Glucocorticoids, and Zomepirac
 20 Sodium. One preferred anti-inflammatory agent is aspirin. Additional examples of anti-inflammatory agents are known in the art.

Non-limiting examples of anti-thrombotic agents that can be used to treat a cardiovascular disease in a subject (alone or in combination with any other therapy, including exercise
 25 therapy) include: plasminogen proactivator, tissue plasmi-
 nogen activator, Anisoylated Plasminogen-Streptokinase
 Activator Complex, Pro-Urokinase, (Pro-UK), rTPA (re-
 combinant alteplase or activase), recombinant Pro-UK,
 Abbokinase, Eminase, Streptase Anagrelide Hydrochloride,
 30 Bivalirudin, Dalteparin Sodium, Danaparoid Sodium,
 Dazoxiben Hydrochloride, Efgatran Sulfate, Enoxaparin
 Sodium, Ifetroban, Ifetroban Sodium, Tinzaparin Sodium,
 Retaplast, Trifenagrel, Warfarin, and Dextrins. Additional
 35 examples of anti-thrombotic agents are known in the art.

Non-limiting examples of anti-coagulants that can be used to treat a cardiovascular disease in a subject (alone or in combination with any other therapy, including exercise
 40 therapy) include: Ancrod, Anticoagulant Citrate Dextrose
 Solution, Anticoagulant Citrate Phosphate Dextrose
 Adenine Solution, Anticoagulant Citrate Phosphate Dex-
 trose Solution, Anticoagulant Heparin Solution, Anticoagu-
 lant Sodium Citrate Solution, Ardeparin Sodium, Bivaliru-
 45 din, Bromindione, Dalteparin Sodium, Desirudin,
 Dicumarol, Heparin Calcium, Heparin Sodium, Lyapolate
 Sodium, Nafamostat Mesylate, Phenprocoumon, Tinzaparin
 Sodium, and Warfarin Sodium. Additional examples of
 50 anti-coagulants are known in the art.

Non-limiting examples of anti-platelet agents that can be used to treat a cardiovascular disease in a subject (alone or in combination with any other therapy, including exercise
 55 therapy) include: Clopidogrel, Sulfinpyrazone, Aspirin,
 Dipyridamole, Clofibrate, Pyridinol Carbamate, Prostaglan-
 din E, Glucagon, Antiserotonin drugs, Caffeine, Theophyllin
 Pentoxifyllin, Ticlopidine, and Anagrelide. Additional
 60 examples of anti-platelet agents are known in the art.

Non-limiting examples of direct thrombin inhibitors that can be used to treat a cardiovascular disease in a subject (alone or in combination with any other therapy, including
 65 exercise therapy) include: hirudin, hirugen, hirulog, aga-
 troban, PPACK, and thrombin aptamers. Additional
 examples of thrombin inhibitors are known in the art.

Non-limiting examples of glycoprotein IIb/IIIb receptor inhibitors that can be used to treat a cardiovascular disease in a subject (alone or in combination with any other therapy, including exercise therapy) include: ReoPro (abcixamab),
 5 lamifiban, and tirofiban. Additional examples of glycopro-
 tein IIb/IIIb receptor inhibitors are known in the art.

Non-limiting examples of calcium channel blockers that can be used to treat a cardiovascular disease in a subject (alone or in combination with any other therapy, including
 10 exercise therapy) include: dihydropyridines, such as nife-
 dipine; phenyl alkyl amines, such as verapamil; and benzo-
 thiazepines, such as diltiazem. Additional non-limiting
 examples of calcium channel blockers include amrinone,
 15 amlodipine, bencyclane, felodipine, fendiline, flunarizine,
 isradipine, nicardipine, nimodipine, perhexilene, gallopamil,
 tiapamil, tiapamil analogues (such as 1993RO-11-2933),
 phenytoin, barbiturates, and the peptides dynorphin, omega-
 conotoxin, and omega-agatoxin, and the like and/or phar-
 20 maceutically acceptable salts thereof. Additional examples
 of calcium channel blockers are known in the art.

Non-limiting examples of beta-adrenergic receptor block-
 25 ers that can be used to treat a cardiovascular disease in a
 subject (alone or in combination with any other therapy,
 including exercise therapy) include: atenolol, acebutolol,
 alprenolol, befunolol, betaxolol, bunitrolol, carteolol, celip-
 30 rolol, hedroxalol, indenolol, labetalol, levobunolol, mepin-
 dolol, methypranol, metindol, metoprolol, metrizoranolol,
 oxprenolol, pindolol, propranolol, practolol, practolol,
 sotalolnadolol, tiprenolol, tomalolol, timolol, bupranolol,
 35 penbutolol, trimepranol, 2-(3-(1,1-dimethylethyl)-amino-2-
 hydroxypropoxy)-3-pyridenecarbonitril HCl, 1-butylamino-
 3-(2,5-dichlorophenoxy)-2-propanol, 1-isopropylamino-3-
 (4-(2-cyclopropylmethoxyethyl)phenoxy)-2-propanol,
 3-isopropylamino-1-(7-methylindan-4-yloxy)-2-butanol,
 2-(3-t-butylamino-2-hydroxy-propylthio)-4-(5-carbamoyl-
 40 2-thienyl)thiazol, and 7-(2-hydroxy-3-t-butylaminopropoxy)
 phthalide. The above beta-adrenergic receptor blockers can
 be used as isomeric mixtures, or in their respective levoro-
 tating or dextrorotating form. Additional examples of beta-
 45 adrenergic receptor blockers are known in the art.

Non-limiting examples of cyclooxygenase-2 inhibitors that can be used to treat a cardiovascular disease in a subject (alone or in combination with any other therapy, including
 50 exercise therapy) include those described in U.S. Pat. Nos.
 5,474,995; 5,521,213; 5,536,752; 5,550,142; 5,552,422;
 5,604,253; 5,604,260; 5,639,780; 5,677,318; 5,691,374;
 5,698,584; 5,710,140; 5,733,909; 5,789,413; 5,817,700;
 5,849,943; 5,861,419; 5,922,742; 5,925,631; 5,643,933;
 5,474,995; and 5,543,297; WO 95/00501, and WO 95/18799
 55 (each of which is incorporated herein by reference). Addi-
 tional examples of cyclooxygenase-2 inhibitors are known
 in the art.

Renin-angiotensin-aldosterone system (RAAS) inhibitors can be used to treat a cardiovascular disease in a subject (alone or in combination with any other therapy, such as
 60 exercise therapy). RAAS inhibitors include agents that inter-
 fere with the function and synthesis or catabolism of angio-
 tensin II. RAAS agents include, but are not limited to,
 angiotensin-converting enzyme (ACE) inhibitors, angio-
 tensin II receptor blockers, agents that activate the catabo-
 65 lism of angiotensin II, agents that prevent the synthesis of
 angiotensin I (from which angiotensin II is ultimately
 derived), and aldosterone antagonists. The RAAS is
 involved in the regulation of hemodynamics and water and
 electrolyte balance. Factors that lower blood volume, renal
 perfusion pressure, or the concentration of Na⁺ in plasma
 tend to activate the system, while factors that increase these

parameters tend to suppress its function. RAAS inhibitors are compounds that act to interfere with the production of angiotensin II from angiotensinogen or angiotensin I or interfere with the activity of angiotensin II. Such inhibitors are well known in the art and include compounds that act to inhibit the enzymes involved in the ultimate production of angiotensin II, including renin and ACE. They also include compounds that interfere with the activity of angiotensin II, once produced.

Angiotensin II receptor blockers include angiotensin II antagonists which interfere with the activity of angiotensin II by binding to angiotensin II receptors and interfere with their activity. Angiotensin II receptor blockers are well known and include peptide compounds and non-peptide compounds. Most angiotensin II receptor blockers are slightly modified congeners in which agonist activity is attenuated by replacement of phenylalanine in position 8 with some other amino acid. Examples of angiotensin II receptor blockers include: peptidic compounds (e.g., saralasin, [(San1)(Val5)(Ala8)] angiotensin-(1-8) octapeptide, and related analogs); N-substituted imidazole-2-one (U.S. Pat. No. 5,087,634); imidazole acetate derivatives, including 2-N-butyl-4-chloro-1-(2-chlorobenzile), imidazole-5-acetic acid (see, Long et al., *J. Pharmacol. Exp. Ther.* 247:1-7, 1988); 4, 5, 6, 7-tetrahydro-1H-imidazo [4, 5-c] pyridine-6-carboxylic acid, and analog derivatives thereof (U.S. Pat. No. 4,816,463); N2-tetrazole beta-glucuronide analogs (U.S. Pat. No. 5,085,992); substituted pyrroles, pyrazoles, and tryazoles (U.S. Pat. No. 5,081,127); phenol and heterocyclic derivatives, such as 1, 3-imidazoles (U.S. Pat. No. 5,073,566); imidazo-fused 7-member ring heterocycles (U.S. Pat. No. 5,064,825); peptides (e.g., U.S. Pat. No. 4,772,684); antibodies to angiotensin II (e.g., U.S. Pat. No. 4,302,386); and aralkyl imidazole compounds, such as biphenyl-methyl substituted imidazoles (e.g., EP 253,310); N-morpholinoacetyl-(-1-naphthyl)-L-alanyl-(4, thiazolyl)-L-alanyl (35, 45)-4-amino-3-hydroxy-5-cyclo-hexapentanoyl-N-hexylamide; SKF108566 (E-alpha-2-[2-butyl-1-(carboxyphenyl)methyl] 1H-imidazole-5-yl[methylane]-2-thiophenepropanoic acid); Losartan; Remikirin; and A2 agonists.

Non-limiting examples of ACE inhibitors include acylmercapto and mercaptoalkanoyl prolines, such as captopril (U.S. Pat. No. 4,105,776) and zofenopril (U.S. Pat. No. 4,316,906); carboxyalkyl dipeptides, such as enalapril (U.S. Pat. No. 4,374,829), lisinopril (U.S. Pat. No. 4,374,829), quinapril (U.S. Pat. No. 4,344,949), ramipril (U.S. Pat. No. 4,587,258), and perindopril (U.S. Pat. No. 4,508,729); carboxyalkyl dipeptide mimics, such as cilazapril (U.S. Pat. No. 4,512,924) and benazapril (U.S. Pat. No. 4,410,520); and phosphinylalkanoyl prolines, such as fosinopril (U.S. Pat. No. 4,337,201) andtrandolopril.

Additional non-limiting examples of RAAS inhibitors include: derivatives of peptides (U.S. Pat. No. 5,116,835); amino acids connected by nonpeptide bonds (U.S. Pat. No. 5,114,937); di- and tri-peptide derivatives (U.S. Pat. No. 5,106,835); amino acids and derivatives thereof (U.S. Pat. Nos. 5,104,869 and 5,095,119); diol sulfonamides and sulfinyls (U.S. Pat. No. 5,098,924); modified peptides (U.S. Pat. No. 5,095,006); peptidyl beta-aminoacyl aminodiol carbamates (U.S. Pat. No. 5,089,471); pyroloimidazolones (U.S. Pat. No. 5,075,451); fluorine and chlorine statine, or statone containing peptides (U.S. Pat. No. 5,066,643); peptidyl amino diols (U.S. Pat. Nos. 5,063,208 and 4,845,079); N-morpholino derivatives (U.S. Pat. No. 5,055,466); pepstatin derivatives (U.S. Pat. No. 4,980,283); N-heterocyclic alcohols (U.S. Pat. No. 4,885,292); monoclonal antibodies

to renin (U.S. Pat. No. 4,780,401); and a variety of other peptides and analogs thereof (U.S. Pat. Nos. 5,071,837, 5,064,965, 5,063,207, 5,036,054, 5,036,053, 5,034,512, and 4,894,437) (each of which is incorporated by reference).

Additional examples of RAAS inhibitors include aldosterone antagonists. Non-limiting examples of aldosterone antagonists include: Spironolactone, Eplerenone, Canrenone (canrenoate potassium), Prorenone (prorenoate potassium), and Mexrenone (mexrenoate potassium).

EXAMPLES

The invention is further described in the following examples, which do not limit the scope of the invention described in the claims.

Example 1

Soluble ST2 Levels are Predictive of Mortality and Adverse Events in Heart Failure Patients

Analysis was performed on the samples and data collected from 2331 heart failure patients enrolled in the HF-ACTION study. A total of 2329 of the patients underwent baseline exercise testing and were randomized to exercise therapy or normal clinical therapy. Of these patients, blood samples were collected from 912 patients representing both arms of the study, 453 were in the exercise arm and 459 were in the normal treatment arm. Nine hundred and ten of these patients had a sufficient sample size for soluble ST2 measurement. The median soluble ST2 level in this cohort was 23.7 ng/ml, and ranged from 2.2 ng/mL to 344.2 ng/mL. As in the previously published heart failure cohorts, soluble ST2 levels were predictive of death for the full duration of the follow-up period (up to 1460 days). The prognostic value of soluble ST2 levels for risk of death in heart failure patients during the entire follow-up period (up to 1460 days) is illustrated in FIG. 1. Patients in the highest soluble ST2 level group (soluble ST2 levels of greater than 28.6 ng/mL) have the greatest risk of mortality, with this risk presenting early in the follow-up period. The increased risk of mortality for those patients in the highest soluble ST2 level group is maintained throughout the entire follow-up period.

The predictive strength of soluble ST2 levels was also assessed by Cox proportional hazards analysis, using soluble ST2 level as both a log (ln)-transformed continuous variable (HR 3.38, $p < 0.0001$), as well as a dichotomous variable at a concentration of 35 ng/ml (HR 2.59, $p < 0.0001$). FIG. 2 shows the surviving portion of heart failure patients over time for subjects having ST2 concentrations above or below a level of 35 ng/mL.

Example 2

Subjects with Elevated Levels of Soluble ST2 Performing an Exercise Therapy Regime have an Increased Risk of Death

As reported in O'Connor et al. (*JAMA* 301(14):1439-1450, 2009) there was a modest, but insignificant, decrease in mortality rate in heart failure patients who underwent exercise therapy relative to those who received standard care. An assessment of soluble ST2 levels in this cohort shows that the therapeutic efficacy of exercise therapy is different in patients having a high versus low level of soluble ST2, with more benefit, e.g., lower mortality rate, observed in heart failure patients with low soluble ST2 levels. This

correlation is illustrated in a Kaplan-Meier analysis of the data from the following heart failure patient groups: patients having a level of soluble ST2 less than or equal to 35 ng/mL and performing an exercise therapy regime (line 1, top line); patients having a level of soluble ST2 less than or equal to 35 ng/nL and not performing an exercise therapy regime (line 2, second line from the top); patients having a level of soluble ST2 greater than 35 ng/mL and not performing an exercise therapy regime (line 3, second line from the bottom); and patients having a level of soluble ST2 greater than 35 ng/mL and performing an exercise therapy regime (line 4, bottom line) (FIG. 3). In both treatment arms (i.e., those performing an exercise therapy regime or those receiving standard care (not performing an exercise therapy regime)), heart failure patients with low soluble ST2 levels have better survival over the 4 year follow-up period than patients with elevated soluble ST2 levels. By 1 year, the patients with low soluble ST2 levels who performed an exercise therapy regime had a significantly lower mortality rate (~3%) compared to the usual care treatment group (not performing an exercise therapy regime), and this benefit persisted for the full 4-year follow-up period. Conversely, the data from heart failure patients with elevated soluble ST2 levels show no survival benefit from the performance of an exercise therapy regime out to 2 years of followup. After 2 years, there was an apparent adverse effect (increased mortality) of performing an exercise treatment regime in heart failure patients with elevated soluble ST2 levels compared to the usual care (not performing an exercise therapy regime) patients.

The ST2 analysis cutpoint of 35 ng/mL was selected by choosing a Presage ST2 Assay concentration value above the 90th and below the 95th percentile of the group. These reference values were subsequently confirmed to be consistent with an additional 3,450 subjects measured in a large

observational population study, the Framingham Offspring Cohort (Wang et al. 2004) by showing that ST2 values are 32.9 ng/ml and 37.3 ng/ml at the 90th and 95th percentiles respectively, bracketing and confirming the selected 35 ng/ml value.

The relationship between the mortality risk of performing an exercise therapy regime and soluble ST2 levels in heart failure patients is also illustrated in FIG. 4. The data in FIG. 4 show the calculated hazard ratio for mortality within 1 year in heart failure subjects performing an exercise treatment regime to heart failure subjects not performing an exercise treatment regime having different soluble ST2 levels. These data show that, as soluble ST2 levels increase there is a steady increase in the hazard ratio, reflecting worse survival, and that a level of soluble ST2 between 55 and 60 ng/mL has a hazard ratio of 1.0. These data indicated that heart failure patients with soluble ST2 levels below 55-60 ng/mL are likely to benefit from performing an exercise therapy regime, with greater benefit achieved at lower soluble ST2 levels. Heart failure patients with soluble ST2 levels above 55-60 ng/mL are not likely to achieve a therapeutic benefit from performing an exercise therapy regime and are at increased risk of experiencing an adverse outcome (e.g., mortality).

OTHER EMBODIMENTS

It is to be understood that while the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 4

<210> SEQ ID NO 1
<211> LENGTH: 328
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

Met Gly Phe Trp Ile Leu Ala Ile Leu Thr Ile Leu Met Tyr Ser Thr
1 5 10 15

Ala Ala Lys Phe Ser Lys Gln Ser Trp Gly Leu Glu Asn Glu Ala Leu
20 25 30

Ile Val Arg Cys Pro Arg Gln Gly Lys Pro Ser Tyr Thr Val Asp Trp
35 40 45

Tyr Tyr Ser Gln Thr Asn Lys Ser Ile Pro Thr Gln Glu Arg Asn Arg
50 55 60

Val Phe Ala Ser Gly Gln Leu Leu Lys Phe Leu Pro Ala Ala Val Ala
65 70 75 80

Asp Ser Gly Ile Tyr Thr Cys Ile Val Arg Ser Pro Thr Phe Asn Arg
85 90 95

Thr Gly Tyr Ala Asn Val Thr Ile Tyr Lys Lys Gln Ser Asp Cys Asn
100 105 110

Val Pro Asp Tyr Leu Met Tyr Ser Thr Val Ser Gly Ser Glu Lys Asn
115 120 125

Ser Lys Ile Tyr Cys Pro Thr Ile Asp Leu Tyr Asn Trp Thr Ala Pro
130 135 140

-continued

Leu Glu Trp Phe Lys Asn Cys Gln Ala Leu Gln Gly Ser Arg Tyr Arg
145 150 155 160

Ala His Lys Ser Phe Leu Val Ile Asp Asn Val Met Thr Glu Asp Ala
165 170 175

Gly Asp Tyr Thr Cys Lys Phe Ile His Asn Glu Asn Gly Ala Asn Tyr
180 185 190

Ser Val Thr Ala Thr Arg Ser Phe Thr Val Lys Asp Glu Gln Gly Phe
195 200 205

Ser Leu Phe Pro Val Ile Gly Ala Pro Ala Gln Asn Glu Ile Lys Glu
210 215 220

[illegible]

Lys Gly Thr Gln Phe Leu Ala Ala Val Leu Trp Gln Leu Asn Gly Thr
245 250 255

Lys Ile Thr Asp Phe Gly Glu Pro Arg Ile Gln Gln Glu Glu Gly Gln
260 265 270

Asn Gln Ser Phe Ser Asn Gly Leu Ala Cys Leu Asp Met Val Leu Arg
275 280 285

Ile Ala Asp Val Lys Glu Glu Asp Leu Leu Leu Gln Tyr Asp Cys Leu
290 295 300

Ala Leu Asn Leu His Gly Leu Arg Arg His Thr Val Arg Leu Ser Arg
305 310 315 320

Lys Asn Pro Ser Lys Glu Cys Phe
325

<210> SEO ID NO 2

```
<211> LENGTH: 2542
```

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

gaaggagggac ctacaaagac tggaaactat tcttagctcc gtcactgact ccaagttcat 60

cccctctgtc tttcagtttg gttgagatat aggctactct tcccaactca gtcttgaaga 120

qtatcaccaa ctgcctcatg tqtggtgacc ttcactgtcg tatgccagtg actcatctgg 180

agtaatctca acaacgaatt accaatactt gctcttgatt gataaacaga atgggggtttt 240

qqatcttagc aattctcaca attctcatgt attccacagc agcaaagttt agtaaacaat 300

catggggcct ggaaaatgaq gctttaattg taagatgtcc tagacaagga aaacctaagt 360

acaccgtqqa ttggtattac tcacaaacaa acaaaaagtat tcccactcag qaaagaaatc 420

gtgtgttttgc ctcagqccaa cttctgaagt ttctaccagc tqcagttgct gattcttgta 480

tttatacctg tattgtcaga agtcccacat tcaataggac tqgatatgcg aatgtcacca 540

tatataaaaa acaatcaqat tqcaatgttc caqattatth gatqtattca acaqtatctg 600

gatcagaaaa aaattccaaa atttattgtc ctaccattga cctctacaac tggacagcac 660

ctcttgagtg gtttaagaat tgtcaaggctc ttcaaggatc aaqgtacagg gcgcacaaqt 720

cattttttggt cattgataat gtgatgactg aggcacgcagg tgattacacc tqttaaattta 780

tacacaatga aaatqgaacc aattataatg tgacqgcgcac caqgtccttc acqgtcaagg 840

atgagcaagg cttttctctg tttccagtaa tcggagcccc tgcacaaaat gaaataaagg 900

aaqtgqaaat tqgaaaaaac qcaaacctaa cttgctctgc ttgtttttgga aaagqcactc 960

agttcttggc tgccgtcctg tggcagctta atggaacaaa aattacagac tttggtgaac 1020

-continued

caagaattca	acaagaggaa	gggcaaaatc	aaagtttcag	caatgggctg	gcttgctctag	1080
acatgggtttt	aagaatagct	gacgtgaagg	aagaggatth	attgctgcag	tacgactgtc	1140
tggccctgaa	tttgcattggc	ttgagaaggc	acaccgtaag	actaagtagg	aaaaatccaa	1200
gtaaggagtg	tttctgagac	tttgatcacc	tgaactttct	ctagcaagtg	taagcagaat	1260
ggagtgtggt	tccaagagat	ccatcaagac	aatgggaatg	gcctgtgcc	taaaatgtgc	1320
ttctcttctt	cgggatgttg	tttgctgtct	gatctttgta	gactgttctt	gtttgctggg	1380
agcttctctg	ctgcttaa	tggtcgtctt	ccccactctt	ctcctatcgt	tggtttgtct	1440
agaacactca	gctgcttctt	tggtcctctt	tggtttctaa	ctttatgaac	tccctctgtg	1500
tcactgtatg	tgaaggaaa	tgcaccaaca	accgtaaa	gaacgtgttc	ttttgtgtct	1560
ttttataact	tgcattacat	gttgtaagca	tggtccgttc	tatacttttt	tctggtcata	1620
atgaacactc	atthttgttag	cgagggtggt	aaagtgaaca	aaaaggggaa	gtatcaaact	1680
actgccattt	cagtgaagaa	atcctaggtg	ctactttata	ataagacatt	tgtagggcca	1740
ttcttgcat	gatataaaga	aatacctgag	actgggtgat	ttatatgaaa	agaggtttaa	1800
ttggctcaca	gttctgcagg	ctgtatggga	agcatggcgg	catctgcttc	tggggacacc	1860
tcaggagctt	tactcatggc	agaaggcaaa	gcaaaggcag	gcacttcaca	cagtaaaagc	1920
aggagcgaga	gagaggtgcc	acactgaaac	agccagatct	catgagaagt	cactcactat	1980
tgcaaggaca	gcacaaaga	gatgggtgcta	aaccattcat	gatgaactca	ccccatgat	2040
ccaatcacct	cccaccaggc	tccacctcga	atactgggga	ttaccattca	gcattgagatt	2100
tgggcaggaa	cacagacca	aaccatacca	cacacattat	cattgttaaa	ctttgtaaag	2160
tatttaagg	acatggaaca	cacgggaagt	ctggtagctc	agccatttc	tttattgcat	2220
ctgttattca	ccatgtaatt	caggtaccac	gtattccagg	gagcctttct	tggccctcag	2280
tttgcatgat	acacactttc	caagtactct	tgtagcatcc	tggtttgtatc	atagcactgg	2340
tcacattgcc	ttacctaaat	ctgtttgaca	gtctgtctca	cacgactgca	agctccatga	2400
gggcagggac	atcatctctt	ccatctttgg	gtccttagtg	caatacctgg	cagctagcca	2460
gtgctcagct	aatattttgt	tgactgaata	aatgaatgca	caaccaaaaa	aaaaaaaaaa	2520
aaaaaaaaaa	aaaaaaaaaa	aa				2542
<210> SEQ ID NO 3						
<211> LENGTH: 556						
<212> TYPE: PRT						
<213> ORGANISM: Homo sapiens						
<400> SEQUENCE: 3						
Met Gly Phe Trp Ile Leu Ala Ile Leu Thr Ile Leu Met Tyr Ser Thr						
1 5 10 15						
Ala Ala Lys Phe Ser Lys Gln Ser Trp Gly Leu Glu Asn Glu Ala Leu						
20 25 30						
Ile Val Arg Cys Pro Arg Gln Gly Lys Pro Ser Tyr Thr Val Asp Trp						
35 40 45						
Tyr Tyr Ser Gln Thr Asn Lys Ser Ile Pro Thr Gln Glu Arg Asn Arg						
50 55 60						
Val Phe Ala Ser Gly Gln Leu Leu Lys Phe Leu Pro Ala Ala Val Ala						
65 70 75 80						
Asp Ser Gly Ile Tyr Thr Cys Ile Val Arg Ser Pro Thr Phe Asn Arg						
85 90 95						
Thr Gly Tyr Ala Asn Val Thr Ile Tyr Lys Lys Gln Ser Asp Cys Asn						

-continued

100								105				110					
Val	Pro	Asp	Tyr	Leu	Met	Tyr	Ser	Thr	Val	Ser	Gly	Ser	Glu	Lys	Asn		
		115					120					125					
Ser	Lys	Ile	Tyr	Cys	Pro	Thr	Ile	Asp	Leu	Tyr	Asn	Trp	Thr	Ala	Pro		
	130					135					140						
Leu	Glu	Trp	Phe	Lys	Asn	Cys	Gln	Ala	Leu	Gln	Gly	Ser	Arg	Tyr	Arg		
145					150					155					160		
Ala	His	Lys	Ser	Phe	Leu	Val	Ile	Asp	Asn	Val	Met	Thr	Glu	Asp	Ala		
				165					170					175			
Gly	Asp	Tyr	Thr	Cys	Lys	Phe	Ile	His	Asn	Glu	Asn	Gly	Ala	Asn	Tyr		
		180					185						190				
Ser	Val	Thr	Ala	Thr	Arg	Ser	Phe	Thr	Val	Lys	Asp	Glu	Gln	Gly	Phe		
	195					200						205					
Ser	Leu	Phe	Pro	Val	Ile	Gly	Ala	Pro	Ala	Gln	Asn	Glu	Ile	Lys	Glu		
	210					215					220						
Val	Glu	Ile	Gly	Lys	Asn	Ala	Asn	Leu	Thr	Cys	Ser	Ala	Cys	Phe	Gly		
225					230					235					240		
Lys	Gly	Thr	Gln	Phe	Leu	Ala	Ala	Val	Leu	Trp	Gln	Leu	Asn	Gly	Thr		
			245						250					255			
Lys	Ile	Thr	Asp	Phe	Gly	Glu	Pro	Arg	Ile	Gln	Gln	Glu	Glu	Gly	Gln		
		260						265					270				
Asn	Gln	Ser	Phe	Ser	Asn	Gly	Leu	Ala	Cys	Leu	Asp	Met	Val	Leu	Arg		
		275					280					285					
Ile	Ala	Asp	Val	Lys	Glu	Glu	Asp	Leu	Leu	Leu	Gln	Tyr	Asp	Cys	Leu		
	290					295					300						
Ala	Leu	Asn	Leu	His	Gly	Leu	Arg	Arg	His	Thr	Val	Arg	Leu	Ser	Arg		
305					310					315					320		
Lys	Asn	Pro	Ile	Asp	His	His	Ser	Ile	Tyr	Cys	Ile	Ile	Ala	Val	Cys		
			325						330				335				
Ser	Val	Phe	Leu	Met	Leu	Ile	Asn	Val	Leu	Val	Ile	Ile	Leu	Lys	Met		
		340						345					350				
Phe	Trp	Ile	Glu	Ala	Thr	Leu	Leu	Trp	Arg	Asp	Ile	Ala	Lys	Pro	Tyr		
		355					360					365					
Lys	Thr	Arg	Asn	Asp	Gly	Lys	Leu	Tyr	Asp	Ala	Tyr	Val	Val	Tyr	Pro		
	370					375					380						
Arg	Asn	Tyr	Lys	Ser	Ser	Thr	Asp	Gly	Ala	Ser	Arg	Val	Glu	His	Phe		
385					390					395					400		
Val	His	Gln	Ile	Leu	Pro	Asp	Val	Leu	Glu	Asn	Lys	Cys	Gly	Tyr	Thr		
			405						410					415			
Leu	Cys	Ile	Tyr	Gly	Arg	Asp	Met	Leu	Pro	Gly	Glu	Asp	Val	Val	Thr		
		420					425						430				
Ala	Val	Glu	Thr	Asn	Ile	Arg	Lys	Ser	Arg	Arg	His	Ile	Phe	Ile	Leu		
		435					440					445					
Thr	Pro	Gln	Ile	Thr	His	Asn	Lys	Glu	Phe	Ala	Tyr	Glu	Gln	Glu	Val		
	450					455					460						
Ala	Leu	His	Cys	Ala	Leu	Ile	Gln	Asn	Asp	Ala	Lys	Val	Ile	Leu	Ile		
465					470					475					480		
Glu	Met	Glu	Ala	Leu	Ser	Glu	Leu	Asp	Met	Leu	Gln	Ala	Glu	Ala	Leu		
			485						490				495				
Gln	Asp	Ser	Leu	Gln	His	Leu	Met	Lys	Val	Gln	Gly	Thr	Ile	Lys	Trp		
		500						505					510				
Arg	Glu	Asp	His	Ile	Ala	Asn	Lys	Arg	Ser	Leu	Asn	Ser	Lys	Phe	Trp		
	515						520					525					

-continued

Lys His Val Arg Tyr Gln Met Pro Val Pro Ser Lys Ile Pro Arg Lys
530 535 540

Ala Ser Ser Leu Thr Pro Leu Ala Ala Gln Lys Gln
545 550 555

<210> SEQ ID NO 4
<211> LENGTH: 2058
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

aaagagagggc	tggctgttgt	atttagtaaa	gctataaagc	tgtaagagaa	attggctttc	60
tgagttgtga	aactgtgggc	agaaagttga	ggaagaaaga	actcaagtac	aacccaatga	120
ggttgagata	taggctactc	ttcccaactc	agtcttgaag	agtatcacca	actgcctcat	180
gtgtggtgac	cttcactgtc	gtatgccagt	gactcatctg	gagtaatctc	aacaacgagt	240
taccaatact	tgctcttgat	tgataaacag	aatgggggtt	tggatcttag	caattctcac	300
aattctcatg	tattccacag	cagcaaagtt	tagtaaacia	tcatggggcc	tggaaaatga	360
ggctttaatt	gtaagatgtc	ctagacaagg	aaaacctagt	tacaccgtgg	attggtatta	420
ctcacaaaca	aacaaaagta	ttcccactca	ggaaagaaat	cgtgtgtttg	cctcaggcca	480
acttctgaag	tttctaccag	ctgcagttgc	tgattctggg	atttatacct	gtattgtcag	540
aagtcccaca	ttcaatagga	ctggatatgc	gaatgtcacc	atatataaaa	aacaatcaga	600
ttgcaatggt	ccagattatt	tgatgtattc	aacagtatct	ggatcagaaa	aaaattccaa	660
aatttattgt	cctaccattg	acctctacaa	ctggacagca	cctcttgagt	ggtttaagaa	720
ttgtcagggt	cttcaaggat	caagggtacag	ggcgcacaa	tcatttttgg	tcattgataa	780
tgtgatgact	gaggacgcag	gtgattacac	ctgtaaat	atacacatg	aaaatggagc	840
caattatagt	gtgacggcga	ccagggtcct	cacgggtca	gatgagcaag	gcttttctct	900
gtttccagta	atcgagagcc	ctgcacaaaa	tgaaataaag	gaagtggaaa	ttggaaaaaa	960
cgcaaaccta	acttgctctg	cttggtttgg	aaaaggcact	cagttcttgg	ctgccgtcct	1020
gtggcagcct	aatggaacaa	aaattacaga	ctttggtgaa	ccaagaattc	aacaagagga	1080
agggcaaaat	caaagtttca	gcaatgggct	ggcttgtcta	gacatgggtt	taagaatagc	1140
tgacgtgaag	gaagaggatt	tattgctgca	gtacgactgt	ctggccctga	atttgcattg	1200
cttgagaagg	cacaccgtaa	gactaagtag	gaaaaatcca	attgatcatc	atagcatcta	1260
ctgcataatt	gcagtatgta	gtgtattttt	aatgctaata	aatgtcctgg	ttatcatcct	1320
aaaaatgttc	tggattgagg	ccactctgct	ctggagagac	atagctaaac	cttacaagac	1380
taggaatgat	ggaaagctct	atgatgctta	tggtgtctac	ccacggaact	acaaatccag	1440
tacagatggg	gccagtcgtg	tagagcactt	tggtcaccag	attctgcctg	atgttcttga	1500
aaataaatgt	ggctatacct	tatgcattta	tgggagagat	atgctacctg	gagaagatgt	1560
agtcactgca	gtggaaacca	acatacgaaa	gagcaggcgg	cacattttca	tcctgacccc	1620
tcagatcact	cacaataagg	agtttgctta	cgagcaggag	gttgccctgc	actgtgcctt	1680
catccagaac	gacgccaagg	tgatacttat	tgagatggag	gctctgagcg	agctggacat	1740
gctgcagggt	gaggcgcttc	aggactccct	ccagcatctt	atgaaagtac	aggggacct	1800

-continued

caagtggagg gaggaccaca ttgccaataa aaggtccctg aattctaaat tctggaagca	1860
cgtgaggtac caaatgcctg tgccaagcaa aattcccaga aaggcctcta gtttgactcc	1920
cttggtgcc cagaagcaat agtgctgct gtgatgtgca aaggcatctg agtttgaagc	1980
tttctgact tctcctagct ggcttatgcc cctgcactga agtgtgagga gcaggaatat	2040
taaagggatt caggcctc	2058

- What is claimed is:
1. A method of treating a subject having heart failure, the method comprising:
- (a) performing an immunoassay to determine a level of soluble ST2 in a biological sample from a subject having heart failure at a first time point before the start of exercise therapy;
 - (b) beginning exercise therapy in a subject having a decreased level of soluble ST2 in the biological sample at the first time point as compared to an efficacy reference level;
 - (c) performing an immunoassay to determine a level of soluble ST2 in a biological sample from the subject undergoing exercise therapy at a second time point after the start of exercise therapy and after the first time point;
 - (d) identifying a subject having a decreased level of soluble ST2 in the biological sample at the second time point as compared to a risk reference level; and
 - (e) continuing to treat the identified subject with exercise therapy.
2. The method of claim 1, wherein the biological samples at the first and second time points comprise blood, plasma, or serum.
3. The method of claim 1, wherein the subject is hypercholesterolemic, hypertriglyceridemic, hyperlipidemic, a smoker, hypertensive, or has a body mass index of greater than 30.
4. The method of claim 1, further comprising determining a level of cardiac troponin, B type natriuretic peptide, atrial natriuretic peptide, or C-reactive protein in the biological sample at the first time point or in the biological sample at the second time point.
5. The method of claim 1, wherein the subject is previously diagnosed as having heart failure.
6. The method of claim 1, wherein the immunoassay is an enzyme-linked immunosorbent assay (ELISA).
7. The method of claim 1, wherein one or both of the efficacy reference level and the risk reference level is a threshold level of soluble ST2.
8. The method of claim 1, wherein the immunoassay in step (a) and the immunoassay in step (c) are performed using one or both of (i) an antibody produced by the hybridoma deposited at the American Type Culture Collection (ATCC) and designated by the Patent Deposit Designation PTA-10431, and (ii) an antibody produced by the hybridoma deposited at the ATCC and designated by the Patent Deposit Designation PTA-10432.
9. A method of treating a subject having heart failure, the method comprising:
- (a) performing an immunoassay to determine a level of soluble ST2 in a biological sample from a subject having heart failure at a first time point before the start of exercise therapy;
 - (b) beginning exercise therapy in a subject having a decreased level of soluble ST2 in the biological sample at the first time point as compared to an efficacy reference level;
 - (c) performing an immunoassay to determine a level of soluble ST2 in a biological sample from the subject undergoing exercise therapy at a second time point after the start of exercise therapy and after the first time point;
 - (d) identifying a subject having an elevated level of soluble ST2 in the biological sample at the second time point as compared to the risk reference level; and
 - (e) treating the identified subject with a treatment of heart failure that does not include exercise therapy, wherein the treatment comprises a therapeutically effective amount of a renin-angiotensin-aldosterone system inhibitor and a therapeutically effective amount of a beta-adrenergic receptor blocker.
10. The method of claim 9, wherein the biological samples at the first and second time points comprise blood, plasma, or serum.
11. The method of claim 9, wherein the subject is hypercholesterolemic, hypertriglyceridemic, hyperlipidemic, a smoker, hypertensive, or has a body mass index of greater than 30.
12. The method of claim 9, further comprising determining a level of cardiac troponin, B type natriuretic peptide, atrial natriuretic peptide, or C-reactive protein in the biological sample at the first time point or in the biological sample at the second time point.
13. The method of claim 9, wherein the subject is previously diagnosed as having heart failure.
14. The method of claim 9, wherein the immunoassay is an enzyme-linked immunosorbent assay (ELISA).
15. The method of claim 9, wherein one or both of the efficacy reference level and the risk reference level is a threshold level of soluble ST2.
16. A method of treating a subject having heart failure, the method comprising:
- (a) performing an immunoassay to determine a level of soluble ST2 in a biological sample from a subject having heart failure at a first time point after the start of exercise therapy;
 - (b) continuing exercise therapy in a subject having a decreased level of soluble ST2 in the biological sample at the first time point as compared to an efficacy reference level;
 - (c) performing an immunoassay to determine a level of soluble ST2 in a biological sample from the subject undergoing exercise therapy at a second time point after the start of exercise therapy and after the first time point;
 - (d) identifying a subject having a decreased level of soluble ST2 in the biological sample at the second time point as compared to a risk reference level; and

37

(e) continuing to treat the identified subject with exercise therapy.

17. The method of claim 16, wherein the biological samples at the first and second time points comprise blood, plasma, or serum.

18. The method of claim 16, wherein the subject is hypercholesterolemic, hypertriglyceridemic, hyperlipidemic, a smoker, hypertensive, or has a body mass index of greater than 30.

19. The method of claim 16, further comprising determining a level of cardiac troponin, B type natriuretic peptide, atrial natriuretic peptide, or C-reactive protein in the biological sample at the first time point or in the biological sample at the second time point.

20. The method of claim 16, wherein the subject is previously diagnosed as having heart failure.

21. The method of claim 16, wherein the immunoassay is an enzyme-linked immunosorbent assay (ELISA).

22. The method of claim 16, wherein one or both of the efficacy reference level and the risk reference level is a threshold level of soluble ST2.

23. The method of claim 16, wherein the immunoassay in step (a) and the immunoassay in step (c) are performed using one or both of (i) an antibody produced by the hybridoma deposited at the American Type Culture Collection (ATCC) and designated by the Patent Deposit Designation PTA-10431, and (ii) an antibody produced by the hybridoma deposited at the ATCC and designated by the Patent Deposit Designation PTA-10432.

24. A method of treating a subject having heart failure, the method comprising:

- (a) performing an immunoassay to determine a level of soluble ST2 in a biological sample from a subject having heart failure at a first time point after the start of exercise therapy;

38

(b) continuing exercise therapy in a subject having a decreased level of soluble ST2 in the biological sample at the first time point as compared to an efficacy reference level;

(c) performing an immunoassay to determine a level of soluble ST2 in a biological sample from the subject undergoing exercise therapy at a second time point after the start of exercise therapy and after the first time point; and

(d) identifying a subject having an elevated level of soluble ST2 in the biological sample at the second time point as compared to the risk reference level; and

(e) treating the identified subject with a treatment of heart failure that does not include exercise therapy, wherein the treatment comprises a therapeutically effective amount of renin-angiotensin-aldosterone system inhibitor and a therapeutically effective amount of a beta-adrenergic receptor blocker.

25. The method of claim 24, wherein the biological samples at the first and second time points comprise blood, plasma, or serum.

26. The method of claim 24, wherein the subject is hypercholesterolemic, hypertriglyceridemic, hyperlipidemic, a smoker, hypertensive, or has a body mass index of greater than 30.

27. The method of claim 24, further comprising determining a level of cardiac troponin, B type natriuretic peptide, atrial natriuretic peptide, or C-reactive protein in the biological sample at the first time point or in the biological sample at the second time point.

28. The method of claim 24, wherein the subject is previously diagnosed as having heart failure.

29. The method of claim 24, wherein the immunoassay is an enzyme-linked immunosorbent assay (ELISA).

30. The method of claim 24, wherein one or both of the efficacy reference level and the risk reference level is a threshold level of soluble ST2.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 9,551,708 B2
APPLICATION NO. : 14/299851
DATED : January 24, 2017
INVENTOR(S) : James V. Snider and Robert W. Gerwien

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

In the Specification

Column 1, Line 7, Delete “13/552,553,” and insert -- 13/552,533, --, therefor.

Column 1, Line 12, Delete “herein” and insert -- hereby --, therefor.

In the Claims

Column 35, Line 39, Claim 3, delete “hyperlidemic,” and insert -- hyperlipidemic, --, therefor.

Column 36, Line 35, Claim 11, delete “hyperlidemic,” and insert -- hyperlipidemic, --, therefor.

Column 37, Line 7, Claim 18, delete “hyperlidemic,” and insert -- hyperlipidemic, --, therefor.

Column 38, Line 22, Claim 26, delete “hyperlidemic,” and insert -- hyperlipidemic, --, therefor.

Signed and Sealed this
Twenty-sixth Day of September, 2017



Joseph Matal
*Performing the Functions and Duties of the
Under Secretary of Commerce for Intellectual Property and
Director of the United States Patent and Trademark Office*