

US009562098B2

(12) **United States Patent**
Mary et al.

(10) **Patent No.: US 9,562,098 B2**
(45) **Date of Patent: Feb. 7, 2017**

(54) **ANTI-CD28 HUMANIZED ANTIBODIES**
(71) Applicants: **Effimune**, Nantes (FR); **Institut National de la Sante et de la Recherche Medicale (INSERM)**, Paris (FR)
(72) Inventors: **Caroline Mary**, Sainte Pazanne (FR); **Nicolas Poirier**, Nantes (FR); **Bernard Vanhove**, Reze (FR)

(73) Assignees: **OSE Immunotherapeutics**, Nantes (FR); **Institut National de la Sante et de la Recherche Medicale (INSERM)**, Paris (FR)
(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 180 days.

(21) Appl. No.: **14/326,119**

(22) Filed: **Jul. 8, 2014**

(65) **Prior Publication Data**
US 2015/0071916 A1 Mar. 12, 2015

Related U.S. Application Data
(62) Division of application No. 13/577,103, filed as application No. PCT/IB2011/050646 on Feb. 16, 2011, now Pat. No. 8,785,604.

(30) **Foreign Application Priority Data**
Feb. 18, 2010 (EP) 10290080
Jul. 13, 2010 (EP) 10290389

(51) **Int. Cl.**
A61K 39/395 (2006.01)
C07K 16/28 (2006.01)
A61K 47/48 (2006.01)
A61K 39/00 (2006.01)

(52) **U.S. Cl.**
CPC **C07K 16/2818** (2013.01); **A61K 39/3955** (2013.01); **A61K 47/48215** (2013.01); **C07K 16/2896** (2013.01); **A61K 2039/505** (2013.01); **C07K 2317/24** (2013.01); **C07K 2317/524** (2013.01); **C07K 2317/53** (2013.01); **C07K 2317/55** (2013.01); **C07K 2317/56** (2013.01); **C07K 2317/64** (2013.01); **C07K 2317/76** (2013.01)

(58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**
U.S. PATENT DOCUMENTS
8,785,604 B2 * 7/2014 Mary C07K 16/2818
424/144.1
9,085,629 B2 * 7/2015 McKinnon C07K 16/2818
2005/0142642 A1 6/2005 Sun et al.
2008/0025994 A1 1/2008 Steward et al.
2011/0313135 A1 * 12/2011 Vanhove C07K 16/2818
530/387.3

FOREIGN PATENT DOCUMENTS
WO 02/051871 7/2002
WO 2004/058820 7/2004
WO 2010/009391 1/2010
WO 2010/070047 6/2010
WO 2010/082136 7/2010

OTHER PUBLICATIONS
Hunig et al., Med Microbiol Immunol (2010) 199:239-246.*
Kashmiri, SRD Grafting—A New Approach to Antibody Humanization, Methods, 36, pp. 25-34, 2005.
Tan, Humanization of an Anti-CD28 Antibody Using Germline Human Antibody Sequences, Blood, 96, p. 31A, 2000.
Poirier, Inducing CTLA-4-Dependent Immune Regulation by Selective CD28 Blockade Promotes Regulatory T Cells in Organ Transplantation, Science Translational Medicine, 2, pp. 1-11, 2010.
Saldanha, Molecular Engineering I: Humanization, Handbook of Therapeutic Antibodies, pp. 119-144, 2007.
Vanhove, Selective Blockade of CD28 and Not CTLA-4 with a Single-Chain Fv-alpha 1-Antitrypsin Fusion Antibody, Blood, 102(2): 564-70, 2003.
Nunes, CD28 mAbs With Distinct Binding Properties Differ in their Ability to Induce T Cell Activation: Analysis of Early and Late Activation Events, International Immunology, 5, pp. 311-315, 1993.

* cited by examiner

Primary Examiner — Ilia Ouspenski
(74) *Attorney, Agent, or Firm* — Morgan, Lewis & Bockius LLP

(57) **ABSTRACT**
The invention relates to humanized antibodies directed against the human lymphocyte receptor CD28. When used in a monovalent form these antibodies are antagonists, i.e. capable of blocking of the CD28/B7 interaction, without activating CD28. These antibodies can be used in particular as therapeutic agents for blocking T cell activation through the CD28 receptor.

12 Claims, 11 Drawing Sheets

1
 8
 9
 37
 38
 66
 67
 95
 96
 124
 125
 153
 154
 182
 183
 211
 212
 231

ATG GAA TGG TGC TGG GTC TTT CTC TTC CTC TCA GTA ACT GCA GGT GTC CAC TCC AAG GTC CAA CTG CAG CAG TCT GGA GCT
 M E W C W V F L F L L S V T A G V H S K V Q L Q Q S G A
 V12K R19K L20V
 GAG CTG AAG AAA CCC GGG GCG TCG GTG AAA GTC TCC TGC AAG GCG TCT GGT TAC ACC TTC ACT GAA TAT ATT ATA CAC TGG ATA AAG
 E L K F G A S V K S C K A S G Y T F T E Y I I H W I K
 CTG AGG TCT GGA CAG GGT CTT GAG TGG ATT GGG TGG TTT TAC CCT GGA AGT AAT GAT ATA CAG TAC AAT GCG CAA TTC AAG GGC AAG
 L R S G Q G L E W I G W F Y P G S N D I Q Y N A F K G K
 K62Q
 GCC ACA TTG ACT GCG GAC AAA TCC ACC ACC GTC TAT ATG GAA CTT ACT GGA TTG ACA CCC GAG GAC TCT GCG GTC TAT TTT TGT
 A T L T A D K S S S T V Y M E L T G L T E D S A V Y F C
 S84P
 GCA AGA CCG GAC GAT TTC TCT GGT TAC TAC GAC GCC CTT CCT TAC TGG GGC CAA GGG ACT CTG GTC ACT GTC TCT GCA GCT AGC ACC AAG
 A R R D D F S G Y D A L P Y W G Q G T L V T V S A A S T K
 G P S V F P L A P S S K S T S G G T A A L G C L V C L V K D Y F
 153
 GGC CCA TCG GTC TTC CCC CTG GCA CCC TCC TCC ACC ACC GTC TAT ATG GAA CTT ACT GGA TTG ACA CCC GAG GAC TCT GCG GTC TAT TTT TGT
 G P S V F P L A P S S K S T S G G T A A L G C L V C L V K D Y F
 182
 CCC GAA CCG GTG ACG GTG TCG TCG AAC TCA GGC GCC CTG ACC AGC GGC GTG CAC ACC TTC CCG GCT GTC CTA CAG TCC TCA GGA CTC
 P E P V T V S W N S G A L T S G V H T F F A V L Q S S G L
 211
 TAC TCC CTC AGC AGC GTG ACC GTG ACC GTG CCC TCC AGC AGC TTG GGC ACC CAG ACC TAC ATC TGC AAC GTG AAT CAC AAG CCC AGC AAC
 Y S L S S V V T V P S S S L G T Q T Y I C N V N H K P S N
 231
 ACC AAG GTG CAC AAG AAA Gtt GAG CCC AAA TCT TGT GAC AAA ACT CAC ACA TGC GCC GCA TAA
 T K V D K K V E P K S C D K T H T C A A Stop

Figure 1

1 atg agt gtg ccc act oag gtc ctg ggg ttg ctg ctg tgg ctt aca gat gcc aga tgt GAC ATC CAG ATG ACT CAG TCT CCA
 M S V P T Q V L G L L L L W L T D A R C D I Q M T Q S P
 8
 9 TCT TCC CTA TCT GCA ICT GTG GGA CAC AGG GTC ACC ATC ACC TGT AAA ACA AAT GAG AAT ATT TAC AGT AAT TTA GCA TGG TAT CAG
 S L S S V G D H V T I T C K F N E N I Y S N L A W Y Q
 37
 38 CAG AAA CAC GGA AAA TCT CCT CAG CTC CTG ATC TAT GCT GCA ACA CAC TTA GTA GAG GGT GTC CCA TCA AGG TTC AGT GGC AGT GSA
 Q K G K S P Q L L I Y A A T H L V E G P S R F S G S G
 66
 67 TCA GCC ACA CAG TAT TCC CTC ACA ATC ACC AGC CTG CAG CCA GAA GAT TTT GGG AAT TAT TAC TGT CAA CAC TTT TGG GST ACT CCG
 S G T Q Y S L I H S L Q E D F G N Y Y C Q H F W S T P
 95
 96 TGC ACC TTC GGA GGG ACC AAG CTG GAA ATA AAA CCG AGG GTG GCT GCA CCA TCT GTC TTC ATC TTC CCG CCA TCT GAT GAG CAG
 C F G G G T K L E I K R T V A A P S V F I F P P S D E Q
 124
 125 TTG AAA ICT GGA ACT GCC TCT GTT GTG TGC CTG CTG AAT AAC TTC TAT CCC ASA GAG GCC AAA GTA CAG TGG AAG GTG GAT AAC GCC
 L K S G T A S V V C L L N F Y P R E A K V Q W K V D N A
 153
 154 CTC CAA TCG GGT AAC TCC CAG GAG AGT GTC ACA GAG CAG GAC AGC AAG GAC AGC ACC TAC AGC CTC AGC AGC ACC CTG ACG CTG AGC
 I Q S G N S Q E S V T E Q D S K D S T Y S L S S T L F L S
 182
 183 AAA GCA GAC TAC GAG AAA CAC AAA GTC TAC GCC TGC ACC CAT CAG GGC CTG AGT TCG CCC GTC ACA AAG AGC TTC AAC AGG
 K A D Y E K H K V Y A C E V T H Q G L S S F V T K S F N R
 211
 212 GGA GAG TGT TAA
 G E C Stop
 214

Figure 2





Figure 3

1	ATC	GAA	TGG	TGC	TGG	GTG	TTC	CTG	TTC	CTG	CTG	TCC	GTG	ACC	GCT	45
1	Met	Glu	Trp	Cys	Trp	Val	Phe	Leu	Phe	Leu	Leu	Ser	Val	Thr	Ala	15
46	GGC	GTG	CAC	TCC	AAG	CAG	GTG	CAG	CTG	CAG	CAG	TCT	GCC	GCC	GAG	90
16	Gly	Val	His	Ser	Lys	Gln	Val	Gln	Leu	Gln	Gln	Ser	Gly	Ala	Glu	30
91	CTG	AAG	AAG	CCT	GGC	GCC	TCC	GTG	AAG	GTG	TCC	TGC	AAG	GCC	TCC	135
31	Leu	Lys	Lys	Pro	Gly	Ala	Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	45
136	GGC	TAC	ACC	TTC	ACC	GAG	TAC	ATC	ATC	CAC	TGG	ATC	AAG	CTG	AGA	180
46	Gly	Tyr	Thr	Phe	Thr	Glu	Tyr	Ile	Ile	His	Trp	Ile	Lys	Leu	Arg	60
181	TCC	GCC	CAG	GGC	CTG	GAA	TGG	ATC	GCC	TGG	TTC	TAC	CCT	GGC	TCC	225
61	Ser	Gly	Cln	Gly	Leu	Glu	Trp	Ile	Gly	Trp	Phe	Tyr	Pro	Gly	Ser	75
226	AAC	GAC	ATC	CAG	TAC	AAC	GCC	CAG	TTC	AAG	GCC	AAG	GCC	ACC	CTG	270
76	Asn	Asp	Ile	Gln	Tyr	Asn	Ala	Gln	Phe	Lys	Gly	Lys	Ala	Thr	Leu	90
271	ACC	GCC	GAC	AAG	TCC	TCC	TCC	ACC	GTG	TAC	ATG	GAA	CTG	ACC	GCC	315
91	Thr	Ala	Asp	Lys	Ser	Ser	Ser	Thr	Val	Tyr	Met	Glu	Leu	Thr	Gly	105
316	CTG	ACC	CCT	GAG	GAC	TCC	GCC	GTG	TAC	TTC	TGC	GCC	AGG	CGG	GAC	360
106	Leu	Thr	Pro	Glu	Asp	Ser	Ala	Val	Tyr	Phe	Cys	Ala	Arg	Arg	Asp	120
361	GAC	TTC	TCT	GGC	TAC	GAC	GCC	CTG	CCT	TAT	TGG	GGC	CAG	GCC	ACC	405
121	Asp	Phe	Ser	Gly	Tyr	Asp	Ala	Leu	Pro	Tyr	Trp	Gly	Gln	Gly	Thr	135
406	CTG	GTG	ACC	GTG	TCC	GCC	GAC	AAA	ACT	CAC	ACA	TGC	CCA	CCG	TGC	450
136	Leu	Val	Thr	Val	Ser	Ala	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	150
451	CCA	GCA	CCT	GAG	TTC	CTG	GGG	GGA	CCA	TCA	GTC	TTC	CTG	TTC	CCC	495
151	Pro	Ala	Pro	Glu	Phe	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	165
496	CCA	AAA	CCC	AAG	GAC	ACT	CTC	ATG	ATC	TCC	CGG	ACC	CCT	GAG	GTC	540
166	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	180
541	ACG	TGC	GTG	GTG	GTG	GAC	GTG	AGC	CAG	GAA	GAC	CCC	GAG	GTC	CAG	585
181	Thr	Cys	Val	Val	Val	Asp	Val	Ser	Gln	Glu	Asp	Pro	Glu	Val	Gln	195
588	TTC	AAC	TGG	TAC	GTG	GAT	GCC	GTG	GAG	GTG	CAT	AAT	GCC	AAG	ACA	630
196	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	210
631	AAG	CCG	CGG	GAG	GAG	CAG	TTC	AAC	AGC	ACG	TAC	CGT	GTG	GTC	AGC	675
211	Lys	Pro	Arg	Glu	Glu	Gln	Phe	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	225
676	GTC	CTC	ACC	GTC	CTG	CAC	CAG	GAC	TGG	CTG	AAC	GCC	AAG	GAG	TAC	720
226	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	240
721	AAG	TGC	AAG	GTC	TCC	AAC	AAA	GCC	CTC	CCG	TCC	TCC	ATC	GAG	AAA	765
241	Lys	Cys	Lys	Val	Ser	Asp	Lys	Clv	Leu	Pro	Ser	Ser	Ile	Glu	Lys	255
766	ACC	ATC	TCC	AAA	GCC	AAA	GGG	CAG	CCC	CGA	GAG	CCA	CAG	GTG	TAC	810
256	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	270
811	ACC	CTG	CCC	CCA	TCC	CAG	GAG	GAG	ATG	ACC	AAG	AAC	CAG	GTC	AGC	855
271	Thr	Leu	Pro	Pro	Ser	Gln	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	285
856	CTG	ACC	TGC	CTG	GTC	AAA	GCC	TTC	TAC	CCC	AGC	GAC	ATC	GCC	GTG	900
286	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	300
901	GAG	TGG	GAG	AGC	AAT	GGG	CAG	CCG	GAG	AAC	AAC	TAC	AAG	ACC	ACG	945
301	Gln	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	315
946	CCT	CCC	GTG	CTG	GAC	TCC	GAC	GCC	TCC	TTC	TTC	CTC	TAC	AGC	AGG	990
316	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Arg	330
991	CTC	ACC	GTG	CAC	AAG	AGC	AGG	TGG	CAG	GAG	GGG	AAT	GEC	TTC	TCA	1035
331	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Glu	Gly	Asn	Val	Phe	Ser	345
1036	TGC	TCC	GTG	ATG	CAT	GAG	GCT	CTG	CAC	AAC	CAC	TAC	ACA	CAG	AAG	1080
346	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	360
1081	AGC	CTC	TCC	CTG	TCT	CTG	GGT	AAA	TGA							1107
361	Ser	Leu	Ser	Leu	Ser	Leu	Gly	Lys	End							

Figure 4

1	ATG	TCC	GTG	CCT	ACC	CAG	GTG	CTG	GGA	CTG	CTG	CTG	CTG	TGG	CTG	45
1	Met	Ser	Val	Pro	Thr	Gln	Val	Leu	Gly	Leu	Leu	Leu	Leu	Trp	Leu	15
46	ACC	GAC	GCC	AGA	TGC	GAC	ATC	CAG	ATG	ACC	CAG	TCC	CCC	TCC	TCC	90
16	Thr	Asp	Ala	Arg	Cys	Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	30
91	CTG	TCT	GCC	TCC	CTG	GGC	GAC	CGG	GTG	ACC	ATC	ACC	TGT	AAG	ACC	135
31	Leu	Ser	Ala	Ser	Val	Gly	Asp	Arg	Val	Thr	Ile	Thr	Cys	Lys	Thr	45
136	AAC	GAG	AAC	ATC	TAC	TCC	AAC	CTG	GCC	TGG	TAT	CAC	CAG	AAG	GAC	180
46	Asn	Glu	Asn	Ile	Tyr	Ser	Asn	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Asp	60
181	GGC	AAG	TCC	CCT	CAG	CTC	CTG	ATC	TAC	GCC	GCC	ACC	CAT	CTG	GTC	225
61	Gly	Lys	Ser	Pro	Gln	Leu	Leu	Ile	Tyr	Ala	Ala	Thr	His	Leu	Val	75
226	GAG	GGC	GTG	CCC	TCT	AGA	TTC	TCC	GGC	TCC	GCC	TCT	GCC	ACC	CAG	270
76	Glu	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Gln	90
271	TAC	TCC	CTG	ACC	ATC	AGC	TCC	CTG	CAG	CCT	GAG	GAC	TTC	GGC	AAC	315
91	Tyr	Ser	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro	Gln	Asp	Phe	Gly	Asn	105
316	TAC	TAC	TGC	CAG	CAC	TTC	TGG	GGC	ACC	CCT	TGT	ACC	TTC	GGC	GGA	360
106	Tyr	Tyr	Cys	Gln	His	Phe	Trp	Gly	Thr	Pro	Cys	Thr	Phe	Gly	Gly	120
361	GGC	ACC	AAG	CTG	GAA	ATC	AAG	CGG	GAC	AAA	ACT	CAC	ACA	TGC	CCA	405
121	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Arg	Asp	Lys	Thr	His	Thr	Cys	Pro	135
406	CCG	TGC	CCA	GCA	CCT	GAG	TTC	CTG	GGG	GGA	CCA	TCA	GTC	TTC	CTG	450
136	Pro	Cys	Pro	Ala	Pro	Glu	Phe	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	150
451	TTC	CCC	CCA	AAA	CCC	AAG	GAC	ACT	CTC	ATG	ATC	TCC	CGG	ACC	CCT	495
151	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	165
496	GAG	GTC	ACG	TGC	CTG	GTG	GTG	GAC	GTG	AGC	CAG	GAA	GAC	CCC	GAG	540
166	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	Gln	Glu	Asp	Pro	Glu	180
541	GTC	CAG	TTC	AAC	TGG	TAC	GTG	GAT	GGC	GTG	GAC	GTG	CAT	AAT	GCC	585
181	Val	Gln	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	195
586	AAG	ACA	AAG	CCG	CGG	GAG	GAG	CAG	TTC	AAC	AGC	ACG	TAC	CGT	GTG	630
196	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Phe	Asn	Ser	Thr	Tyr	Arg	Val	210
631	GTC	ACC	GTC	CTC	ACC	GTC	CTG	CAC	CAG	GAC	TGG	CTG	AAC	GCC	AAG	675
211	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	225
676	GAG	TAC	AAG	TGC	AAG	GTC	TCC	AAC	AAA	GGC	CTC	CCG	TCC	TCC	ATC	720
226	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Gly	Leu	Pro	Ser	Ser	Ile	240
721	GAG	AAA	ACC	ATC	TCC	AAA	GCC	AAA	GGG	CAG	CCC	CGA	GAG	CCA	CAG	765
241	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	255
766	GTG	TAC	ACC	CTG	CCC	CCA	TCC	CAG	GAG	CAG	ATG	ACC	AAG	AAC	CAG	810
256	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Gln	Glu	Glu	Met	Thr	Lys	Asn	Gln	270
811	GTC	ACC	CTG	ACC	TGC	CTG	GTC	AAA	GGC	TTC	TAC	CCC	AGC	GAC	ATC	855
271	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	285
856	GCC	GTG	GAG	TGG	GAG	AGC	AAT	GGG	CAG	CCG	GAG	AAC	AAC	TAC	AAG	900
286	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	300
901	ACC	ACG	CCT	CCC	GTG	CTG	GAC	TCC	GAC	GGC	TCC	TTC	TTC	CTC	TAC	945
301	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	315
946	AGC	AGG	CTC	ACC	GTG	GAC	AAG	AGC	AGG	TGG	CAG	GAG	GGG	AAT	GTC	990
316	Ser	Arg	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Glu	Gly	Asn	Val	330
991	TTC	TCA	TGC	TCC	GTG	ATG	CAT	GAG	GCT	CTG	CAC	AAC	CAC	TAC	ACA	1035
331	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	345
1036	CAG	AAG	AGC	CTC	TCC	CTG	TCT	CTG	GGT	AAA	TGA					1068
346	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Leu	Gly	Lys	End					

Figure 5

1	ATG	GAA	TGG	TGC	TGG	GTG	TTC	CTG	TTC	CTG	CTG	TCC	GTG	ACC	GCT	45
1	Met	Glu	Trp	Cys	Trp	Val	Phe	Leu	Phe	Leu	Leu	Ser	Val	Thr	Ala	15
46	GGC	GTG	CAC	TCC	AAG	CAG	GTC	CAG	CTG	CAG	CAG	TCT	GGC	GCC	GAG	90
16	Gly	Val	His	Ser	Lys	Gln	Val	Gln	Leu	Gln	Gln	Ser	Gly	Ala	Glu	30
91	CTG	AAG	AAG	CCT	GGC	GCC	TCC	GTC	AAG	GTG	TCC	TGC	AAG	GCC	TCC	135
31	Leu	Lys	Lys	Pro	Gly	Ala	Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	45
136	GGC	TAC	ACC	TTC	ACC	GAG	TAC	ATC	ATC	CAC	TGG	ATC	AAG	CTG	AGA	180
46	Gly	Tyr	Thr	Phe	Thr	Glu	Tyr	Ile	Ile	His	Trp	Ile	Lys	Leu	Arg	60
181	TCC	GGC	CAG	GGC	CTG	GAA	TGG	ATC	GGC	TGG	TTC	TAC	CCT	GGC	TCC	225
61	Ser	Gly	Gln	Gly	Leu	Glu	Trp	Ile	Gly	Trp	Phe	Tyr	Pro	Gly	Ser	75
226	AAC	GAC	ATC	CAG	TAC	AAC	GCC	CAG	TTC	AAG	GGC	AAG	GCC	ACC	CTG	270
76	Asn	Asp	Ile	Gln	Tyr	Asn	Ala	Gln	Phe	Lys	Gly	Lys	Ala	Thr	Leu	90
271	ACC	GCC	GAC	AAG	TCC	TCC	TCC	ACC	GTG	TAC	ATG	GAA	CTG	ACC	GGC	315
91	Thr	Ala	Asp	Lys	Ser	Ser	Ser	Thr	Val	Tyr	Met	Glu	Leu	Thr	Gly	105
316	CTG	ACC	CCT	GAG	GAC	TCC	GCC	GTG	TAC	TTC	TGC	GCC	AGG	CGG	GAC	360
106	Leu	Thr	Pro	Glu	Asp	Ser	Ala	Val	Tyr	Phe	Cys	Ala	Arg	Arg	Asp	120
361	GAC	TTC	TCT	GGC	TAC	GAC	GCC	CTG	CCT	TAT	TGG	GGC	CAG	GCC	ACC	405
121	Asp	Phe	Ser	Gly	Tyr	Asp	Ala	Leu	Pro	Tyr	Trp	Gly	Gln	Gly	Thr	135
406	CTG	GTG	ACC	GTG	TCC	GCC	GAG	CCC	AAA	TCT	TGT	GAC	AAA	ACT	CAC	450
136	Leu	Val	Thr	Val	Ser	Ala	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	150
451	ACA	TGC	CCA	CCG	TGC	CCA	GCA	CCT	GAG	TTC	CTG	GGG	CGA	CCA	TCA	495
151	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Phe	Leu	Gly	Gly	Pro	Ser	165
496	GTC	TTC	CTG	TTC	CCC	CCA	AAA	CCC	AAG	GAC	ACT	CTC	ATG	ATC	TCC	540
166	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	180
541	CGG	ACC	CCT	GAG	GTC	ACG	TGC	GTG	GTG	GTG	GAC	GTG	AGC	CAG	GAA	585
181	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	Gln	Glu	195
586	GAC	CCC	GAG	GTC	CAG	TTC	AAC	TGG	TAC	GTG	CAT	GGC	GTG	GAG	GTG	630
196	Asp	Pro	Glu	Val	Gln	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	210
631	CAT	AAT	GCC	AAG	ACA	AAG	CCG	CGG	GAG	GAG	CAG	TTC	AAC	AGC	ACC	675
211	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Phe	Asn	Ser	Thr	225
676	TAC	CGT	GTG	GTC	AGC	GTC	CTC	ACC	GTC	CTG	CAC	CAG	GAC	TGG	CTG	720
226	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	240
721	AAC	GGC	AAG	GAG	TAC	AAG	TGC	AAG	GTC	TCC	AAC	AAA	GGC	CTC	CCG	765
241	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Gly	Leu	Pro	255
766	TCC	TCC	ATC	GAG	AAA	ACC	ATC	TCC	AAA	GCC	AAA	GGG	CAG	CCC	CGA	810
256	Ser	Ser	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	270
811	GAG	CCA	CAG	GTG	TAC	ACC	CTG	CCC	CCA	TCC	CAG	GAG	GAG	ATG	ACC	855
271	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Gln	Glu	Glu	Met	Thr	285
856	AAG	AAC	CAG	GTC	AGC	CTG	ACC	TGC	CTG	GTC	AAA	GGC	TTC	TAC	CCC	900
286	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	300
901	AGC	GAC	ATC	GCC	GTG	GAG	TGG	GAG	AGC	AAT	GGG	CAG	CCG	GAG	AAC	945
301	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	315
946	AAC	TAC	AAG	ACC	ACG	CCT	CCC	GTG	CTC	GAC	TCC	GAC	GGC	TCC	TTC	990
316	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	330
991	TTC	CTC	TAC	AGC	AGG	CTC	ACC	GTG	GAC	AAG	AGC	AGG	TGG	CAG	GAG	1035
331	Phe	Leu	Tyr	Ser	Arg	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Glu	345
1036	GGG	AAT	GTC	TTC	TCA	TGC	TCC	GTG	ATG	CAT	GAG	GCT	CTG	CAC	AAC	1080
346	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	360
1081	CAC	TAC	ACA	CAG	AAG	AGC	CTC	TCC	CTG	TCT	CTG	GGT	AAA	TGA	1122	
361	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Leu	Gly	Lys	End		

Figure 6

1	ATC	TCC	GTG	CCT	ACC	CAG	GTG	CTG	GGA	CTG	CTG	CTG	CTG	TGC	CTG	45
7	Met	Ser	Val	Pro	Thr	Gln	Val	Leu	Gly	Leu	Leu	Leu	Leu	Trp	Leu	15
46	ACC	GAC	GCC	AGA	TGC	GAC	ATC	CAG	ATG	ACC	CAG	TCC	CCC	TCC	TCC	90
16	Thr	Asp	Ala	Arg	Cys	Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	30
91	CTG	TCT	GCC	TCC	GTG	GGC	GAC	CGG	GTC	ACC	ATC	ACC	TGT	AAG	ACC	135
31	Leu	Ser	Ala	Ser	Val	Gly	Asp	Arg	Val	Thr	Ile	Thr	Cys	Lys	Thr	45
136	AAC	GAG	AAC	ATC	TAC	TCC	AAC	CTG	GCC	TGG	TAT	CAG	CAG	AAG	GAC	180
46	Asn	Glu	Asn	Ile	Tyr	Ser	Asn	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Asp	60
181	GGC	AAG	TCC	CCT	CAG	CTG	CTG	ATC	TAC	GCC	GCC	ACC	CAT	CTG	GTG	225
61	Gly	Lys	Ser	Pro	Gln	Leu	Leu	Ile	Tyr	Ala	Ala	Thr	His	Leu	Val	75
226	GAG	GGC	GTG	CCC	TCT	AGA	TTC	TCC	GGC	TCC	GGC	TCT	GGC	ACC	CAG	270
76	Glu	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Gln	90
271	TAC	TCC	CTG	ACC	ATC	AGC	TCC	CTG	CAG	CCT	GAG	GAC	TTC	GGC	AAC	315
91	Tyr	Ser	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro	Glu	Asp	Phe	Gly	Asn	105
316	TAC	TAC	TGC	CAG	CAC	TTC	TGG	GGC	ACC	CCT	TGT	ACC	TTC	GGC	GGA	360
106	Tyr	Tyr	Cys	Gln	His	Phe	Trp	Gly	Thr	Pro	Cys	Thr	Phe	Gly	Gly	120
361	GGC	ACC	AAG	CTG	GAA	ATC	AAG	CGG	GAG	CCC	AAA	TCT	TGT	GAC	AAA	405
121	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Arg	Gln	Pro	Lys	Ser	Cys	Asp	Lys	135
406	ACT	CAC	ACA	TGC	CCA	CCG	TGC	CCA	GCA	CCT	GAG	TTC	CTG	GGG	GGA	450
136	Thr	His	Thr	Cys	Phe	Pro	Cys	Phe	Ala	Pro	Gln	Phe	Leu	Gly	Gly	150
451	CCA	TCA	GTC	TTC	CTG	TTC	CCC	CCA	AAA	CCC	AAG	CAC	ACT	CTC	ATG	495
151	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	165
486	ATC	TCC	CGG	ACC	CCT	GAG	GTC	ACG	TGC	GTC	GTG	GTG	GAC	GTC	AGC	540
166	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	180
541	CAG	GAA	GAC	CCC	GAG	GTC	CAG	TTC	AAC	TGG	TAC	GTG	CAT	GGC	GTG	585
181	Gln	Glu	Asp	Pro	Glu	Val	Gln	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	195
586	GAG	GTG	CAT	AAT	GCC	AAG	ACA	AAG	CCG	CGG	GAG	GAG	CAG	TTC	AAC	630
196	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Phe	Asn	210
631	AGC	ACG	TAC	CGT	GTG	GTC	AGC	GTC	CTC	ACC	GTC	CTG	CAC	CAG	GAC	675
211	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	225
676	TGC	CTG	AAC	GGC	AAG	GAG	TAC	AAG	TGC	AAG	GTC	TCC	AAC	AAA	GGC	720
226	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Gly	240
721	CTC	CCG	TCC	TCC	ATC	GAG	AAA	ACC	ATC	TCC	AAA	GCC	AAA	GGG	CAG	765
241	Leu	Pro	Ser	Ser	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	255
766	CCC	CGA	GAG	CCA	CAG	GTG	TAC	ACC	CTG	CCC	CCA	TCC	CAG	GAG	GAG	810
256	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Gln	Glu	Glu	270
811	ATG	ACC	AAG	AAC	CAG	GTC	AGC	CTG	ACC	TGC	CTG	GTC	AAA	GGC	TTC	855
271	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	285
856	TAC	CCC	AGC	GAC	ATC	GCC	GTG	GAG	TGG	GAG	AGC	AAT	GGG	CAG	CCG	900
286	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	300
901	GAG	AAC	AAC	TAC	AAG	ACC	ACG	CCT	CCC	GTG	CTG	GAC	TCC	GAC	GGC	945
301	Glu	Asp	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	315
946	TCC	TTC	TTC	CTC	TAC	AGC	AGG	CTC	ACC	GTG	GAC	AAG	AGC	AGG	TCC	990
316	Ser	Phe	Phe	Leu	Tyr	Ser	Arg	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	330
991	CAG	GAG	GGG	AAT	GTC	TTC	TCA	TGC	TCC	GTG	ATG	CAT	GAG	GCT	CTG	1035
331	Gln	Glu	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	345
1036	CAC	AAC	CAC	TAC	ACA	CAG	AAG	AGC	CTC	TCC	CTG	TCT	CTG	GGT	AAA	1080
346	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Leu	Gly	Lys	360
1081	TGA															1083
361	End															

Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13

ANTI-CD28 HUMANIZED ANTIBODIES

SEQUENCE LISTING SUBMISSION VIA
EFS-WEB

A computer readable text file, entitled "045636-5202_01-SequenceListing.txt," created on or about Jul. 8, 2014, with a file size of about 36 kb contains the sequence listing for this application and is hereby incorporated by reference in its entirety.

The present invention relates to humanized antibodies binding CD28, to monovalent fragments thereof, and to their therapeutic uses, in particular in the context of regulating T cell activation.

Abnormal activation of T cells is involved in the pathogenesis of many autoimmune diseases, and also in transplant rejection phenomena, where they cause an immune response directed against the transplanted organ to develop.

One of the most important systems for regulating T lymphocyte activation is the molecular system B7/CD28/CTLA4. This system plays, for example, an essential role in the mechanisms of transplant rejection (WOODWARD et al., *Transplantation*, 66, 14-20, 1998). The molecules B7.1 (CD80) and B7.2 (CD86) borne by the APCs can activate the receptor CD28 and also the receptor CTLA4 of T lymphocytes. The activation of CD28 sends the T lymphocyte a positive signal which stimulates the cell; on the other hand, the activation of CTLA4 sends a negative signal which leads to a non-response (anergy) (FALLARINO et al., *J. Exp. Med.*, 188, 205-210, 1998).

Resting T lymphocytes express a large amount of CD28 and very little CTLA4. When there is a first cognitive contact between an APC and a T lymphocyte, the CD28/B7 interaction is favored, which activates the cell. It is only several hours after the initiation of activation that, due to the increase in membrane expression of CTLA4, the affinity of which for B7 is 5 to 10 times greater than that of CD28, the B7/CD28 interaction shifts in favor of a B7/CTLA4 interaction.

Regulatory T lymphocytes express a large amount of CD28 and of CTLA4 that prevent or allow, respectively, the suppressive activity of regulatory T lymphocytes. In the presence of APC expressing high level of B7, the CD28/B7 interaction prevents the suppressive activity of regulatory T lymphocytes (Sansom et al., *Trends Immunol.* 24, 314-319, 2003).

Selective inhibition of the agonist signal given to the T cell by CD28, leaving the antagonist system consisting of the pair CTLA4/B7 intact, via specific blocking of the CD28/B7 interaction, would make it possible to prevent T lymphocyte activation and to promote immune suppression by regulatory T lymphocytes. Such specific blocking of the CD28/B7 interaction can be obtained using some antibodies directed against CD28.

These antibodies are to be used in a monovalent form (for instance as Fab or scFv fragments), since when used in their divalent native form, their binding to CD28 brings about the dimerization and the activation of this receptor. Fab fragments each contain a light chain and the first half of a heavy chain; scFv fragments consist of the variable portions of the heavy and light chains of a parent antibody, connected to one another via a flexible linker (CLACKSON et al., *Nature*, 352, 624-628, 1991), thus forming a single-chain protein.

One such antibody is antibody CD28.3, produced by the hybridoma cell line CNCM I-2582, and disclosed in PCT application WO 02/051871. This antibody, when used in a monovalent form such as scFv fragments, is capable of

blocking in vitro the CD28 receptor without activating it (PCT WO 02/051871; VANHOVE et al., *Blood*, 102, 564-70, 2003), and has shown also its efficiency in vivo in models of organ transplantation in mice and in primates (POIRIER et al., *World Transplant Congress*, Sydney, Australia. Aug. 16-21, 2008; POIRIER et al., *Sci Trans Med*, 2:17, p17ra10, 2010).

A drawback of all monoclonal antibodies derived from murine sources, is their immunogenicity when administered to human subjects. They provoke anti-mouse immune response, which results in a lesser efficiency of the treatment, in particular when repeated administration is required.

This drawback can, in principle, be avoided by the use of humanized antibodies. The aim of humanization is to obtain a recombinant antibody which has similar antigen-binding properties as the mouse monoclonal antibody from which the complementarity-determining regions (CDRs) sequences were derived, and which is far less immunogenic in humans.

The CDRs are the portions of the variable domains of an antibody which directly contact the antigen and determine the antigen-binding specificity; the framework regions (FRs) which are located between the CDRs in the variable domains do not directly contact the antigen, but serves as a scaffold to maintain the global structure of the variable domains.

Several approaches to antibody humanization have been reported. The more widely used are based on "CDR grafting", which involves the transplantation of the CDRs of a murine antibody into appropriate human FRs. However, in many antibodies, some FR residues are important for antigen binding, because they influence the conformation of CDRs and thus their antigen binding properties, in particular the binding affinity. A loss in binding affinity is particularly detrimental in the case of an antibody intended to be used in a monovalent form which generally exhibit less affinity for the antigen than the native divalent antibody. Thus, in most cases, it is further necessary, in order to obtain a sufficient binding affinity, to reintroduce one or several framework residues from the mouse antibody in the human FRs, with the risk of simultaneously bringing back unwanted immunogenicity.

Another approach to antibody humanisation, called "de-immunization", involves the identification within the FRs regions of the antibody, of B-cell and T-cell epitopes recognized as "foreign" and therefore potentially immunogenic in humans, and to remove them by appropriate amino-acids substitutions. This approach however also entails the risk that FR residues important for antigen binding are deleted. Moreover, some immunogenic epitopes may lie in the CDRs and trying to remove them involves a very high risk of destroying not only the antigen-binding affinity but also the antigen-binding specificity of the antibody.

Therefore, a major issue in antibody humanisation is to determine which amino acid residues are critical for retaining the antigen-binding properties. Various methods have been proposed for predicting the more appropriate sites for substitution in the FRs regions. Although they provide general principles that may be of some help in the first steps of humanization, the final result greatly varies from an antibody to another. Thus, for a given antibody, it is very difficult to foretell which substitutions will provide the desired result. In the case wherein not only substitutions in the FRs, but also in the CDRs would be necessary to decrease satisfactorily the immunogenicity in humans, the final result becomes totally unpredictable.

The inventors have succeeded in producing humanized CD28.3 (hereinafter referred to as hCD28.3), with a low

3

immunogenicity, and which, although it has several amino-acids substitutions including a non-conservative K→Q substitution in the CDR2 of the heavy chain, retains the CD28 binding properties of the parent mouse CD28.3. When used in a monovalent form, the hCD28.3 of the invention also retains the CD28 binding properties of the parent mouse CD28.3.

The present invention provides an anti-CD28 antibody, characterised in that it is selected among:

- a) an antibody having a CD28-binding site consisting of: a first variable domain (also defined herein as the “heavy chain variable domain”) defined by the following sequence:

VQLQQSGAELKKPGASVKVSCKASGYTFTEYIIH-
WIKLRSGQGLEWI GWFYPGSNDIQYNAQFKGKATL-
TADKSSSTVYMEGLTLPEDSAVYFCARRDDFSG
YDALPYWGQGTLVTVSA (SEQ ID NO: 1), wherein said variable domain may optionally further comprise a Q residue at its N-terminal end;

- a second variable domain (also defined herein as the “light chain variable domain”) defined by the following sequence:

DIQMTQSPSSLSASVGDRVTITCKTNENIYSN-
LAWYQQKDGKSPQLL IYAATHLVEGVPSRFSGSGS-
GTQYSLTISSLQPEDFGNYCQHFHWGTPXTFGGGT-
KLEI KR (SEQ ID NO: 2), wherein X=C, A, or N.

- b) an antibody having a CD28-binding site consisting of: a first variable domain having the CDRs of the variable domain of SEQ ID NO: 1; a second variable domain having the CDRs of the variable domain of SEQ ID NO: 2.

The term “anti-CD28 antibody” herein refers to any antigen-binding protein having at least one antigen-binding site (consisting of the variable domains of the light chain and of the heavy chain) able to specifically bind human CD28. It encompasses antibodies in a divalent form (such as native immunoglobulin molecules or F(ab)₂ fragments) with two CD28-binding sites, as well as antibodies in a monovalent form which have a single CD28-binding site, (for instance Fab, Fab', Fv and scFv fragments). In most cases, antibodies in a monovalent form will be preferred.

It includes in particular recombinant antibodies comprising a CD28-binding site associated with one or more heterologous polypeptide(s).

By way of example, an antibody of the invention may be a recombinant Fab or Fab' fragment containing the constant domain CH1 of a human immunoglobulin fused at the C-terminal end of the variable domain of SEQ ID NO: 1, and the constant domain CL of a human immunoglobulin fused at the C-terminal end of the variable domain of SEQ ID NO: 2. An example of such a recombinant Fab fragment is a Fab fragment with a heavy chain having the sequence of amino-acids 21-251 of SEQ ID NO: 4 and a light chain having the sequence of amino-acids 21-234 of SEQ ID NO: 6.

Also, a hCD28.3 antibody of the invention may comprise, besides the variable domains of SEQ ID NO: 1 and SEQ ID NO: 2, defined above, one or more of the following components:

- a human constant region (Fc). This constant region can be selected among constant domains from any class of immunoglobulins, including IgM, IgG, IgD, IgA and IgE, and any isotype, including IgG1, IgG2, IgG3 and IgG4. Preferred constant regions are selected among constant domains of IgG, in particular IgG4.
a protein which makes it possible to prolong the plasma half-life when it is administered in vivo under monovalent form as disclosed for instance in PCT WO

4

02/051871; in a preferred embodiment, said protein is the CH2-CH3 domains of an IgG molecule, as disclosed in PCT/IB/2010/000196; according to said embodiment, a hCD28.3 monovalent antibody of the invention is an heterodimer of:

- a first protein chain consisting essentially of, from its N-terminus to its C-terminus:
a region A having the sequence SEQ ID NO: 1;
a region B consisting of a peptide linker and the CH2 and CH3 domains of an IgG immunoglobulin;
a second protein chain consisting essentially of, from its N-terminus to its C-terminus:
a region A' having the sequence SEQ ID NO: 2;
a region B identical to the region B of the first polypeptide.

Preferably, the peptide linker is the hinge region of human IgG1 immunoglobulins having the sequence EPKSCDKTH-TCPPCP (SEQ ID NO: 7), and the CH2 and CH3 domains are those of an immunoglobulin of the IgG4 subclass. One can also use a shortened version of said hinge region, having the sequence DKTHTCPPCP (SEQ ID NO: 8).

According to a preferred embodiment, the polypeptide sequence of the first protein chain is the sequence of amino-acids 21-368 of SEQ ID NO: 10, and the polypeptide sequence of the second protein chain is the sequence of amino-acids 21-355 of SEQ ID NO: 12. According to another preferred embodiment, the polypeptide sequence of the first protein chain is the sequence of amino-acids 21-373 of SEQ ID NO: 14, and the polypeptide sequence of the second protein chain is the sequence of amino-acids 21-360 of SEQ ID NO: 16.

Optionally, a hCD28.3 antibody of the invention may further comprise one or more of the following components:
a protein having pharmacological activity (for example a toxin);
one or more tag polypeptide(s).

Alternatively, to prolong their plasma half life, in particular when they are under the form of Fab fragments, the antibodies of the invention can be conjugated with water soluble polymers such as polyethylene glycol (PEGylation). PEGylation is a classical way to enhance the pharmacokinetic properties of therapeutic polypeptides, and can be achieved by techniques known in the art.

In this respect, the inventors found that the replacement of the original cysteine residue at position 96 of the variable domain of the native CD 28.3 by an alanine or an asparagine residue (resulting in an antibody having a light chain containing a variable domain of SEQ ID NO: 2 wherein X=A or N) allowed a better efficacy in pegylation of the antibody using maleimide-activated polyethylene glycol (targeting reactive cystein residues), without modifying substantially its binding activity, although cysteine-96 is comprised in the CDR3 of the antibody light chain. The benefit of the replacement of the original cysteine residue at position 96 of the variable domain mainly consist in a specific branching of the polyethylene glycol onto the C-terminal cysteine residue of the heavy chain. Without replacement of the original cysteine residue at position 96 of the variable domain of the native CD 28.3, maleimide-activated polyethylene glycol can bind to that cysteine residue and impair the binding activity of the Fab molecule.

The inventors also found that addition of a di-alanine extension after the C-terminal cysteine of the heavy chain also resulted in a better pegylation efficiency.

The invention also encompasses a polynucleotide selected among:

a) a polynucleotide encoding a polypeptide having the CDRs of SEQ ID NO: 1, in particular a polynucleotide encoding a polypeptide of SEQ ID NO: 1;

b) a polynucleotide encoding a polypeptide having the CDRs of SEQ ID NO: 2, in particular a polynucleotide encoding a polypeptide of SEQ ID NO: 2;

c) a polynucleotide encoding an hCD28.3 antibody of the invention, as defined above.

Polynucleotides of the invention generally also comprise additional sequences: for instance they may advantageously comprise a sequence encoding a leader sequence or signal peptide allowing secretion of said protein chain.

The present invention also encompasses recombinant vectors, in particular expression vectors, comprising a polynucleotide of the invention, associated with transcription- and translation-controlling elements which are active in the host cell chosen. Vectors which can be used to construct expression vectors in accordance with the invention are known in themselves, and will be chosen in particular as a function of the host cell intended to be used.

The present invention also encompasses host-cells transformed with a polynucleotide of the invention. Preferably, said host cell is transformed with a polynucleotide comprising a sequence encoding the heavy chain of a hCD28.3 antibody of the invention and a polynucleotide comprising a sequence encoding the light chain of a hCD28.3 antibody of the invention, and expresses said antibody. Said polynucleotides can be inserted in the same expression vector, or in two separate expression vectors.

Host cells which can be used in the context of the present invention can be prokaryotic or eukaryotic cells. Among the eukaryotic cells which can be used, mention will in particular be made of plant cells, cells from yeast, such as *Saccharomyces*, insect cells, such as *Drosophila* or *Spodoptera* cells, and mammalian cells such as HeLa, CHO, 3T3, C127, BHK, COS, etc., cells.

The construction of expression vectors of the invention and the transformation of the host cells can be carried out by the conventional techniques of molecular biology.

Still another object of the invention is a method for preparing a hCD28.3 antibody of the invention. Said method comprises culturing a host-cell transformed with a polynucleotide comprising a sequence encoding the heavy chain of a hCD28.3 antibody of the invention and a polynucleotide comprising a sequence encoding the light chain of a hCD28.3 antibody of the invention and recovering said antibody from said culture.

If the antibody is secreted by the host-cell, it can be recovered directly from the culture medium; if not, cell lysis will be carried out beforehand. The antibody can then be purified from the culture medium or from the cell lysate, by conventional procedures, known in themselves to those skilled in the art, for example by fractionated precipitation, in particular precipitation with ammonium sulfate, electrophoresis, gel filtration, affinity chromatography, etc.

The hCD28.3 antibodies of the invention can be used to obtain medicinal products. These medicinal products are also part of the object of the invention.

The present invention also comprises a therapeutic composition comprising a hCD28.3 antibody of the invention, together with a pharmaceutically acceptable excipient.

Preferably, said composition is a composition for parenteral administration, formulated to allow the administration of a dose of from 0.5 to 20 mg/Kg, advantageously of from

5 to 10 mg/Kg of an hCD28.3 antibody of the invention. The injection route of the composition can be preferably subcutaneous or intra-venous.

For instance, hCD28.3 antibodies of the invention can be used to obtain immunosuppressant medicinal products which selectively blocks T cell activation phenomena involving the CD28 receptor. Such immunosuppressant medicinal products which act by selective blocking of CD28 have applications in all T lymphocyte-dependent pathological conditions, including in particular transplant rejection, graft-versus-host disease, T lymphocyte-mediated autoimmune diseases, such as type I diabetes, rheumatoid arthritis or multiple sclerosis, and type IV hypersensitivity, which is involved in allergic phenomena and also in the pathogenesis of chronic inflammatory diseases, in particular following infection with a pathogenic agent (in particular leprosy, tuberculosis, leishmaniasis, listeriosis, etc.).

The present invention will be understood more clearly from the further description which follows, which refers to nonlimiting examples of the preparation and properties of a hCD28.3 antibody in accordance with the invention.

The construction of expression vectors of the invention and the transformation of host-cells can be made by the standard techniques of molecular biology.

A hCD28.3 antibody of the invention can be obtained by culturing a host cell containing an expression vector comprising a nucleic acid sequence encoding said antibody, under conditions suitable for the expression thereof, and recovering said antibody from the host cell culture.

The present invention will be further illustrated by the following additional description, which refers to examples illustrating the properties of hCD28.3 antibodies of the invention. It should be understood however that these examples are given only by way of illustration of the invention and do not constitute in any way a limitation thereof.

LEGENDS OF THE DRAWINGS

FIG. 1: Nucleotidic (SEQ ID NO: 3) and amino acid (SEQ ID NO: 4) sequences of the Signal-VH-hCH1 construction. Bold: leader sequence (Nucleotide, SEQ ID NO: 17; amino acid, SEQ ID NO: 18); Underlined: positions of the CDRs (SEQ ID NO: 19, 20 and 21) of the parent CD28.3 antibody. Italics: human CH1 region (Nucleotide, SEQ ID NO: 22; amino acid, SEQ ID NO: 23); Highlighted and double underlined: substitutions made in the CD28.3 antibody VH region.

FIG. 2: Nucleotidic (SEQ ID NO: 5) and amino acid (SEQ ID NO: 6) sequences of the Signal-VL-hC κ construction. Bold: leader sequence (Nucleotide, SEQ ID NO: 24; amino acid, SEQ ID NO: 25); Underlined: positions of the CDRs (SEQ ID NO: 26, 27, and 28) of the parent CD28.3 antibody. Italics: human c kappa region (Nucleotide, SEQ ID NO: 29; amino acid, SEQ ID NO: 30); Highlighted and double underlined: substitutions made in the CD28.3 antibody VL region.

FIG. 3: A) optical density at 405 nm for increasing concentrations of FR104, hCD28.3 Fab or CD28.3 Fab in the Binding ELISA; B) calculation of the regression curves, allowing for determining comparative AC50 values.

FIG. 4: Nucleotidic (SEQ ID NO: 9) and amino acid (SEQ ID NO: 10) sequences of the hVHCD28.3-short hinge γ 1-h γ 4CH2CH3 construction. Bold: leader sequence (SEQ ID NO: 18). Underlined: CDRs (SEQ ID NO: 19, 20, and 21).

Double underlined: hinge region (SEQ ID NO: 8). Dotted underlined: CH2-CH3 domains (SEQ ID NO: 31) of the human IgG4.

FIG. 5: Nucleotidic (SEQ ID NO: 11) and amino acid (SEQ ID NO: 12) sequences of the hVLCD28.3-short hinge γ 1-hy4CH2CH3 construction. Bold: leader sequence (SEQ ID NO: 25). Underlined: CDRs. Double underlined: hinge region (SEQ ID NO: 8). Dotted underlined: CH2-CH3 domains (SEQ ID NO: 31) of the human IgG4.

FIG. 6: Nucleotidic (SEQ ID NO: 13) and amino acid (SEQ ID NO: 14) sequences of the hVHCD28.3-full hinge γ 1-hy4CH2CH3 construction. Bold: leader sequence (SEQ ID NO: 18). Underlined: CDRs (SEQ ID NO: 19, 20, and 21). Double underlined: hinge region (SEQ ID NO: 7). Dotted underlined: CH2-CH3 domains (SEQ ID NO: 31) of the human IgG1.

FIG. 7: Nucleotidic (SEQ ID NO: 15) and amino acid (SEQ ID NO: 16) sequences of the hVLCD28.3-full hinge γ 1-hy4CH2CH3 construction. Bold: leader sequence (SEQ ID NO: 25). Underlined: CDRs (SEQ ID NO: 26, 27 and 28). Double underlined: hinge region (SEQ ID NO: 7). Dotted underlined: CH2-CH3 domains (SEQ ID NO: 31) of the human IgG1.

FIG. 8: Anti-CD28 binding properties of hVH/VL CD28.3 monovalent antibodies. COS cells were co-transfected with 2 μ g (each) pSignal-hVH-short hinge γ 1-hy4CH2-CH3 and pSignal-hVL-short hinge γ 1-hy4CH2-CH3, or co-transfected with 2 μ g (each) pSignal-hVH-full hinge γ 1-hy4CH2-CH3 and pSignal-hVL-full hinge γ 1-hy4CH2-CH3. After 6 days, supernatants were collected and monovalent antibodies were dosed using a first sandwich ELISA. Supernatants were also assessed with a binding ELISA on immobilized CD28 target molecules and bound monovalent anti-CD28 antibodies were revealed with anti-human Fc antibodies labeled with peroxidase. A: Optical density obtained with indicated molecules according to their concentration. B: table with regression curves and the calculation of ED50 (effective dose 50), the concentration needed to reach 50% binding activity in this assay.

FIG. 9: hVH/VL CD28.3 monovalent antibodies inhibit IL-2 secretion by activated T cells. Jurkat T cells were stimulated with SEE superantigen and Raji antigen-presenting-cells during 48 h, in the presence of indicated concentrations of purified hVH/VL-short hinge γ 1-hy4CH2-CH3 monovalent antibodies. Supernatant were collected and IL-2 measured by ELISA.

FIG. 10: SP sepharose HP-chromatography (left) and SDS-PAGE (right) under unreduced conditions after pegylation of C96-Fabs from humanised CD28.3 antibody. Lane 1: marker; lane 2: load; lane 3: Peak 1; lane 4: Peak 2; lane 5: Peak 3.

FIG. 11: Binding properties for CD28 of recombinant hCD28.3 Fabs with or without C96 mutations. The graph shows binding activity (Y axis) according to Fab concentration (X axis).

FIG. 12: SP sepharose HP-chromatography (left) and SDS-PAGE (right) under unreduced conditions after pegylation of C96A-Fabs from humanised CD28.3 antibody. Lane 1: MW markers; lane 2: Pegylated proteins pre-chromatography; lane 3: peak 1 containing the monopegylated Fab, representing 41% of the starting material.

FIG. 13: SP sepharose HP-chromatography (left) and SDS-PAGE (right) under unreduced conditions after pegylation of C96A-Fabs from humanised CD28.3 antibody with a CAAC-terminal sequence in the heavy chain. Lane 1: MW markers; lane 2: Pegylated proteins pre-chromatography; lane 3: peak.

EXAMPLE 1

Construction and Eucaryotic Expression of a hCD28.3 Monovalent Antibody (Fab Fragment)

Heavy Chain:

The sequence encoding the VH region of hCD28.3 (SEQ ID NO: 1) in fusion with the sequence encoding the human CH1 region (NCBI Accession number AAF03881) and with a sequence encoding the leader peptide of the heavy chain of the native murine CD28.3 antibody, was synthesized chemically, and introduced in the cloning vector pGA18 (Genart) for amplification. The sequence was then excised by digestion with KpnI/BamHI restriction enzymes and subcloned into the KpnI/BamHI sites of the plasmid pcDNA3.1-hygro (Invitrogen). Positive clones were amplified and purified by Midiprep-endotoxin free (Macherey-Nagel) for transfection step.

The resulting plasmid is designated pSignal-VH-hCH1. It comprises a construct containing the sequence encoding the VH region of hCD28.3 between the sequence encoding the CD28.3 heavy chain leader peptide and the sequence encoding the human CH1 region (NCBI Accession number AAF03881). The nucleotidic and amino acid sequences of this construct are shown on FIG. 1. They are also represented as SEQ ID NO: 3 and SEQ ID NO: 4 in the enclosed sequence listing.

Light Chain:

The sequence encoding the VL region of hCD28.3 (SEQ ID NO: 2) in fusion with the sequence encoding the human c kappa region (NCBI accession number BAC01725) and with a sequence encoding the leader peptide of the light chain of the native murine CD28.3 antibody, was synthesized chemically, and introduced in the cloning vector pGA18 (Genart) for amplification. The sequence was then excised by digestion with KpnI/BamHI restriction enzymes and subcloned into the KpnI/BamHI sites of the plasmid pcDNA3.A-hygro (Invitrogen). Positive clones were amplified and purified by Midiprep-endotoxin free (Macherey-Nagel) for transfection step.

The resulting plasmid is designated pSignal-VL-hC κ . It comprises a construct containing the sequence encoding the VL region of hCD28.3 between the sequence encoding the CD28.3 light chain signal peptide and the sequence encoding the human c kappa region (NCBI accession number BAC01725). The nucleotidic and amino acid sequences of this construct are shown on FIG. 2. They are also represented as SEQ ID NO: 5 and SEQ ID NO: 6 in the enclosed sequence listing.

Eucaryotic Expression

COS cells were co-transfected with 2 μ g (each) pSignal-VL-hCH1 and pSignal-VH-hCH1 using the Fugene lipofection kit (Roche Diagnostics, Basel, Switzerland) according to the manufacturer's instructions. Cultures were maintained for 3 days at 37° C., divided one third, and put back into culture for an additional 3 days, after which time the cell supernatants were collected.

The activity of the hCD28.3 monovalent antibody is evaluated directly in the supernatant by ELISA, as described in Example 2 below.

EXAMPLE 2

Detection of the hCD28.3 Fab Fragment Binding Activity by ELISA

The binding properties of the hCD28.3 Fab fragment have been compared with those obtained after transfection of Cos cells with plasmids coding for CD28.3 Fab (not humanized), using two ELISA assays

First (Sandwich ELISA), the concentrations of the hCD28.3 and CD28.3 Fab fragments in the culture supernatants of transfected COS cells have been determined using a sandwich ELISA. Briefly, the anti-CD28 Fab contained in the supernatants are first captured by a rabbit polyclonal antibody, specific for the heavy and light variable domains of CD28.3 (obtained after immunization of rabbits with a single-chain-FIT containing the heavy and light variable domains of the native CD28.3, and purified by immunoadsorption on CD28.3 Fab-Sepharose). The captured proteins are then revealed with a murine monoclonal antibody directed to the kappa chain of human IgG, followed by a polyclonal goat anti-mouse antibody labelled with peroxidase. Bound antibody was revealed by colorimetry using the TMB substrate, and read at 405 nm.

The OD corresponding to different dilutions of the supernatant are then compared to a standard curve obtained with known quantities of a CD28.3 Fab, called FR104, purified from culture supernatant of transformed CHO cells with standard techniques of chromatography, and dosed with a BCA (bisynchronous acid) assay. FR104 contains the native (not humanized), VH and VL regions of the CD28.3 antibody. Therefore, we can evaluate the amount of Fab proteins present in cell supernatants.

Second (Binding ELISA), for testing the binding activity of hCD28.3 Fab fragments compared to CD28.3 Fab, chimeric human CD28/Fc (R&D Systems, Abingdon, United Kingdom) was used at 2 µg/ml in carbonate buffer 0.05M pH 9.2 to coat the wells (50 µL/well) of microtiter plates (Nunc Immunoplates) overnight at 4° C. These immobilized CD28 target molecules will bind only immunoreactive molecules with anti-CD28 activity.

The wells were then washed 3 times successively with 200 µL PBS-0.05% Tween, and saturated with 100 µL PBS Tween 0.1% BSA 1% for 2 hours at 37° C.

Then, after 3 washings with 200 µL PBS-0.05% Tween, supernatants containing known concentrations of CD28.3 or hCD28.3 Fab fragments were added (50 µL/well) at different dilutions in PBS-0.1% Tween and incubated for 2 hours at 37° C. After 3 washings with 200 µL PBS-0.05% Tween, a murine monoclonal antibody directed to the kappa chain of human IgG, (1/10000 dilution) was added (1 hour, 37° C.), followed by peroxidase-conjugated goat anti-mouse antibodies (1/2000 dilution), followed by colorimetric revelation using the TMB substrate and reading at 405 nm.

Then the results are plotted as the absorbance (Y axis), measured with the binding ELISA, according to the Fab concentration (X axis), measured with the sandwich ELISA. An AC50 (Antibody Concentration 50) is determined after calculating the slope of the curve in its linear range as the concentration of the anti-CD28 Fab needed to reach 50% of the maximal optical density (OD) in the binding assay.

The results are shown on FIG. 3 and Table I.

FIG. 3A shows the optical density at 405 nm for increasing concentrations of FR104, hCD28.3 Fab or CD28.3 Fab in the Binding ELISA.

FIG. 3 B shows the calculation of the regression curves, allowing for determining comparative AC50 values.

Table I below summarises the OD50, the equation, and the AC50 for the standard FR104, and the Fab fragments VH-wild type+VL-wild type and Fab hCD28.3

TABLE I

	OD50	Equation	AC50
Std FR104	1.792	$y = 1.1424\text{Ln}(x) - 3.6351$	115
CD28.3 Fab	1.82	$y = 0.9776\text{Ln}(x) - 3.2483$	162
hCD28.3 Fab	1.804	$y = 1.0217\text{Ln}(x) - 3.2859$	151

These results show that 50% of the binding activity to CD28 could be reached at a concentration similar for Fab fragments VH-wild type+VL-wild type (CD28.3 Fab) and hCD28.3 Fab. The concentration is slightly lower for the standard, probably because it is purified before the assay. Thus hCD28.3 retains the CD28-binding properties of the wild type VH and VL sequences of CD28.

EXAMPLE 3

Construction and Eucaryotic Expression of a hCD28.3 Monovalent Antibody (FV-FC Fragment) with a Short $\gamma 1$ Hinge and a $\gamma 4$ CH2-CH3 Domain

Heavy Chain:

The sequence encoding the VH region of hCD28.3 (SEQ ID NO: 1) in C-terminal fusion with the sequence encoding a portion of the hinge region of the human IgG1 (SEQ ID NO: 8), with CH2-CH3 domains of the human IgG4 (nucleotides 787 to 1440 of the sequence NCBI Accession number BC025985) and in N-terminal position with a sequence encoding the leader peptide of the heavy chain of the native murine CD28.3 antibody, was synthesized chemically, and introduced in the cloning vector pMA (Geneart) for amplification. The sequence was then excised by digestion with NheI/EcoRI restriction enzymes and subcloned into the NheI/EcoRI sites of the plasmid pCIneo (Promega). After transformation of *E. coli* cells, positive clones were amplified and extracted plasmids were purified by Midiprep-endotoxin free columns (Macherey-Nagel).

The resulting plasmid is designated pSignal-hVH-shorthingey1-h $\gamma 4$ CH2-CH3. It comprises a construct containing the sequence encoding the VH region of hCD28.3 between the sequence encoding the CD28.3 heavy chain signal peptide and the sequence encoding a part of the human $\gamma 1$ hinge region and of the human $\gamma 4$ CH2-CH3 domains. The nucleotidic and amino acid sequences of this construct are shown on FIG. 4. They are also represented as SEQ ID NO: 9 and SEQ ID NO: 10 in the enclosed sequence listing.

Light Chain:

The sequence encoding the VL region of hCD28.3 (SEQ ID NO: 2) in fusion with the sequence encoding a portion of the hinge region of the human IgG1 (SEQ ID NO: 8), with CH2-CH3 domains of the human IgG4 (nucleotides 787 to 1440 of the sequence NCBI Accession number BC025985) and in N-terminal position with a sequence encoding the leader peptide of the heavy chain of the native murine CD28.3 antibody, was synthesized chemically, and introduced in the cloning vector pMA (Geneart) for amplification. The sequence was then excised by digestion with NheI/EcoRI restriction enzymes and subcloned into the NheI/EcoRI sites of the plasmid pCIneo (Promega). After transformation of *E. coli* cells, positive clones were amplified and extracted plasmids were purified by Midiprep-endotoxin free columns (Macherey-Nagel).

The resulting plasmid is designated pSignal-hVL-shorthingey1-h $\gamma 4$ CH2-CH3. It comprises a construct containing the sequence encoding the VL region of hCD28.3

11

between the sequence encoding the CD28.3 light chain signal peptide and the sequence encoding a part of the human $\gamma 1$ hinge region and of the human $\gamma 4$ CH2-CH3 domains. The nucleotidic and amino acid sequences of this construct are shown on FIG. 5. They are also represented as SEQ ID NO: 11 and SEQ ID NO: 12 in the enclosed sequence listing.

Eukaryotic Expression

COS cells were co-transfected with 1 μ g (each) pSignal-hVL-shorthinge $\gamma 1$ -h $\gamma 4$ CH2-CH3 and pSignal-hVH-shorthinge $\gamma 1$ -h $\gamma 4$ CH2-CH3, using the Lipofectamine lipofection kit (Invitrogen) according to the manufacturer's instructions. Cultures were maintained for 3 days at 37° C., after which time the cell supernatants were collected. The activity of the monovalent antibody is evaluated directly in the supernatant by ELISA, as described in Example 5 below.

EXAMPLE 4

Construction and Eucaryotic Expression of a hCD28.3 Monovalent Antibody (FV-FC Fragment) with a Full Length $\gamma 1$ Hinge and a $\gamma 4$ CH2-CH3 Domain

Heavy Chain:

The sequence encoding the VH region of hCD28.3 (SEQ ID NO: 1) in C-terminal fusion with the sequence encoding a full length hinge region of the human IgG1 (SEQ ID NO: 7), with CH2-CH3 domains of the human IgG4 (nucleotides 787 to 1440 of the sequence NCBI Accession number BC025985) and in N-terminal position with a sequence encoding the leader peptide of the heavy chain of the native murine CD28.3 antibody, was synthesized chemically, and introduced in the cloning vector pMA (Genent) for amplification. The sequence was then excised by digestion with NheI/EcoRI restriction enzymes and subcloned into the NheI/EcoRI sites of the plasmid pCIneo (Promega). After transformation of *E. coli* cells, positive clones were amplified and extracted plasmids were purified by Midiprep-endotoxin free columns (Macherey-Nagel).

The resulting plasmid is designated pSignal-hVH-fullhinge $\gamma 1$ -h $\gamma 4$ CH2-CH3. It comprises a construct containing the sequence encoding the VH region of hCD28.3 between the sequence encoding the CD28.3 heavy chain signal peptide and the sequence encoding the human $\gamma 1$ hinge region and the human $\gamma 4$ CH2-CH3 domains. The nucleotidic and amino acid sequences of this construct are shown on FIG. 6. They are also represented as SEQ ID NO: 13 and SEQ ID NO: 14 in the enclosed sequence listing.

Light Chain:

The sequence encoding the VL region of hCD28.3 (SEQ ID NO: 2) in fusion with the sequence encoding the full length hinge region of the human IgG1 (SEQ ID NO: 7), with CH2-CH3 domains of the human IgG4 (nucleotides 787 to 1440 of the sequence NCBI Accession number BC025985) and in N-terminal position with a sequence encoding the leader peptide of the heavy chain of the native murine CD28.3 antibody, was synthesized chemically, and introduced in the cloning vector pMA (Genent) for amplification. The sequence was then excised by digestion with NheI/EcoRI restriction enzymes and subcloned into the NheI/EcoRI sites of the plasmid pCIneo (Promega). After transformation of *E. coli* cells, positive clones were amplified and extracted plasmids were purified by Midiprep-endotoxin free columns (Macherey-Nagel).

The resulting plasmid is designated pSignal-hVL-fullhinge $\gamma 1$ -h $\gamma 4$ CH2-CH3. It comprises a construct contain-

12

ing the sequence encoding the VL region of hCD28.3 between the sequence encoding the CD28.3 light chain signal peptide and the sequence encoding the human $\gamma 1$ full length hinge region and of the human $\gamma 4$ CH2-CH3 domains.

The nucleotidic and amino acid sequences of this construct are shown on FIG. 7. They are also represented as SEQ ID NO: 15 and SEQ ID NO: 16 in the enclosed sequence listing.

Eukaryotic Expression

COS cells were co-transfected with 1 μ g (each) pSignal-hVH-fullhinge $\gamma 1$ -h $\gamma 4$ CH2-CH3 and pSignal-hVL-fullhinge $\gamma 1$ -h $\gamma 4$ CH2-CH3 plasmids using the Lipofectamine lipofection kit (Invitrogen) according to the manufacturer's instructions. Cultures were maintained for 3 days at 37° C., after which time the cell supernatants were collected.

The activity of the hCD28.3 monovalent antibody is evaluated directly in the supernatant by ELISA, as described in Example 5 below.

EXAMPLE 5

Evaluation of the HCD28.3-Full Length $\gamma 1$ Hinge- $\gamma 4$ CH2-CH3 Domains and HCD28.3-Short $\gamma 1$ Hinge- $\gamma 4$ CH2-CH3 Domains Monovalent Antibodies Binding Activity by ELISA

The binding properties of the hCD28.3 monovalent antibodies hCD28.3-full $\gamma 1$ hinge- $\gamma 4$ CH2-CH3 domains and hCD28.3-short $\gamma 1$ hinge- $\gamma 4$ CH2-CH3 domains produced by transfected COS cells have been analysed using two ELISA assays.

First (Sandwich ELISA), the concentrations of the hCD28.3 monovalent antibodies in the culture supernatants of transfected COS cells have been determined using a sandwich ELISA. Briefly, the monovalent antibodies contained in the supernatants are first captured by a goat polyclonal antibody directed to human IgG. The captured proteins are then revealed with a biotinylated goat polyclonal anti-human IgG, Fc specific, antibody, then, a Peroxidase-conjugated streptavidin. Bound antibody was revealed by colorimetry using the TMB substrate, and read at 405 nm.

The OD corresponding to different dilutions of the supernatant are then compared to a standard curve obtained with known quantities of hCD28.3 monovalent antibodies, purified from culture supernatant of transformed CHO cells with standard techniques of chromatography, and dosed with a BCA (bisynchronous acid) assay.

Second (Binding ELISA), for testing the binding activity of hCD28.3 monovalent antibodies, chimeric human CD28/Fc (R&D Systems, Abingdon, United Kingdom) was used at 2 μ g/ml in carbonate buffer 0.05M pH 9.2 to coat the wells (50 μ L/well) of microtiter plates (Nunc Immunoplates) overnight at 4° C. These immobilized CD28 target molecules will bind only immunoreactive molecules with anti-CD28 activity.

The wells were then washed 3 times successively with 200 μ L PBS-0.05% Tween, and saturated with 100 μ L PBS Tween 0.1% BSA 1% for 2 hours at 37° C.

Then, after 3 washings with 200 μ L PBS-0.05% Tween, supernatants containing known concentrations of the monovalent antibodies to be tested were added (50 μ L/well) at different dilutions in PBS-0.1% Tween and incubated for 2 hours at 37° C. After 3 washings with 200 μ L PBS-0.05% Tween, we added (1/500 dilution; 1 hour, 37° C.) a rabbit polyclonal antiserum, specific for the heavy and light variable domains of CD28.3 (obtained after immunization of rabbits with a single-chain-FIT containing the heavy and

13

light variable domains of the native CD28.3, and purified by immunoabsorption on CD28.3 Fab-Sepharose). This was followed by peroxidase-conjugated donkey anti-rabbit antibodies (1/2000 dilution), followed by colorimetric revelation using the TMB substrate and reading at 405 nm.

Then the results are plotted as the absorbance (Y axis), measured with the binding ELISA, according to the monovalent antibody concentration (X axis), measured with the sandwich ELISA. An AC50 (Antibody Concentration 50) is determined after calculating the slope of the curve in its linear range as the concentration of the monoclonal antibody needed to reach 50% of the maximal optical density (OD) in the binding assay.

FIG. 8 compares binding activities of hCD28.3-full IgG1 hinge-IgG4CH2-CH3 domains with hCD28.3-short IgG1 hinge-IgG4CH2-CH3 domains monovalent antibodies in the Binding ELISA (FIG. 8A).

FIG. 8B summarises the equation, the regression factor and the AC50 for the monovalent antibodies.

These results show that 50% of the binding activity to CD28 could be reached at a concentration similar for hCD28.3-full γ 1 hinge- γ 4CH2-CH3 domains or hCD28.3-short γ 1 hinge- γ 4CH2-CH3 domains monovalent antibodies.

EXAMPLE 6

HCD28.3 Monovalent Antibodies Prevents T Cell Activation

To verify that hCD28.3 monovalent antibody blocks CD28-dependent T cell activation, we stimulated human T cells (Jurkat cells) with SEE superantigen presented by a Raji B cell line. The endotoxin SEE, when presented to the class II-positive B cell lymphoblastoid line Raji, activates the V β 8-expressing T cell line Jurkat to secrete IL-2 (Herman et al, 1990, J. Exp. Med. 172:709). Since Jurkat cells express high level of CD28 and Raji cells express CD80/86, this reaction is partially dependant on CD28. We measured synthesis of interleukin-2 in this assay by ELISA (ELISA Max™ Set Deluxe Human IL-2 Kit; Biolegend #431805) after 48 h, in the presence of increasing concentrations of hVH/VL CD28.3-short γ 1 hinge- γ 4CH2-CH3 domains.

The results are shown on FIG. 9. They reveal that hCD28.3 monovalent antibodies reduce IL-2 synthesis by T cells in a dose-dependent manner.

EXAMPLE 7

Preparation of a Pegylated HCD28.3 Monovalent Antibody

A hCD28.3 Fab fragment prepared as described in Example 1 was pegylated with maleimide-activated 40 KDa PEG using standard conditions for reduction and PEGylation.

Briefly, Fab antibody fragments were concentrated to 1 mg/mL and then diafiltrated against 20 mM Sodium phosphate, 2 mM EDTA, and pH 7.0. Fab' antibody fragments were then reduced by addition of cysteamine chloride in a molar equivalent ratio=30:1 at room temperature. After 5 hours, solution was applied on a desalting column. Polyethylene glycol (PEG) (Sunbright GL2 400MA, NOF Corporation) was dissolved in 20 mM Phosphate, 2 mM EDTA, pH 7.0 to give 9% (w/w) solution. Desalted Fab solution and PEG were mixed in a molar equivalent ratio=1:1.5 and incubated at ambient temperature for 3 h. Following PEGy-

14

lation the Fab-peg was purified by chromatography using SP Sepharose HP medium. The target protein was eluted with a salt gradient from 0 to 1 M NaCl. Eluted peaks were analysed by SDS-Page. Peak 1 represented monopegylated material, peak 2 unpegylated material and peak 3 polypegylated material.

The results are shown on FIG. 10.

These results show that a significant part of the Fab proteins from the CD28.3 mAb presents a perturbed pegylation profile which results in a yield of monopegylated Fabs of about 5% only (peak 1).

The CD28.3 mAb contains a cystein residue (C96) that is not engaged in intra or inter-chain disulfide bridges, at position 96 of the variable domain light chain. Free cystein will possess a higher reactivity than cystein residues engaged into disulfide bridges and will therefore preferentially be targets of maleimide-activated pegs. Therefore it is likely that a second, unwanted pegylation occurs on this residue.

To solve that problem we performed a VL-C96 mutation study to determine whether it was possible to substitute the C96 residue by another amino acid without modifying the binding properties of the antibody.

Plasmid coding for humanized anti-CD28.3 Fabs with unmodified C96 in the light chain, or with C96 to A, G, S, V, T, N or R mutations were constructed and transfected into Cos cells by lipofection, as disclosed in Example 1. Cell supernatants were first analyzed by sandwich ELISA to determine total Fab concentration, as disclosed in Example 2. Then supernatants were analysed by ELISA to determine binding activity on immobilized recombinant CD28, as disclosed in Example 2.

The results are shown in FIG. 11. These results show that unlike all other substitutions tested, the C96A substitution resulted in a fully active antibody and that the C96N substitution resulted only in a moderate reduction of activity.

The C96A Fab fragment variant was pegylated and purified by chromatography as described above. Pegylated proteins pre-chromatography and elution peaks were analysed by SDS-Page. The results are shown on FIG. 12. Peak 1 represents monopegylated material.

These results show that the C96A Fab fragment can be pegylated with an efficacy reaching 41% (FIG. 12).

Advantage of the CAA C-Terminal End of the Heavy Chain.

The immediate molecular environment of a free cystein might modify its accessibility to maleimide-pegylation and therefore modify the yield of the pegylation reaction. One possible option for the C-terminal cystein is to be the last amino acid of the heavy chain. Another option is the addition of "stuff amino acids" at the C-terminal position, after the last cysteine. We therefore compared pegylation efficacy of a Fab' molecule from the C96A variant of the humanized CD28.3 Mab with the last C-terminal cystein being the last amino acid of the heavy chain (C variant; data shown in FIG. 12) with a similar molecule with the last C-terminal cystein being followed by two alanins (CAA variant). Our data clearly and reproducibly demonstrated that the CAA variant could be pegylated with a 20% higher efficacy (FIG. 13). Indeed pegylation yield that was of 41% for the C96A-C variant reached 52% for the C96A-CAA variant.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 31

<210> SEQ ID NO 1
 <211> LENGTH: 120
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: hCD28.3 VH

<400> SEQUENCE: 1

Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Lys Lys Pro Gly Ala Ser
 1 5 10 15
 Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Glu Tyr Ile
 20 25 30
 Ile His Trp Ile Lys Leu Arg Ser Gly Gln Gly Leu Glu Trp Ile Gly
 35 40 45
 Trp Phe Tyr Pro Gly Ser Asn Asp Ile Gln Tyr Asn Ala Gln Phe Lys
 50 55 60
 Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser Thr Val Tyr Met
 65 70 75 80
 Glu Leu Thr Gly Leu Thr Pro Glu Asp Ser Ala Val Tyr Phe Cys Ala
 85 90 95
 Arg Arg Asp Asp Phe Ser Gly Tyr Asp Ala Leu Pro Tyr Trp Gly Gln
 100 105 110
 Gly Thr Leu Val Thr Val Ser Ala
 115 120

<210> SEQ ID NO 2
 <211> LENGTH: 108
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: hCD28.3 VL
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (96)..(96)
 <223> OTHER INFORMATION: Xaa = Cys, Ala, or Asn

<400> SEQUENCE: 2

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Lys Thr Asn Glu Asn Ile Tyr Ser Asn
 20 25 30
 Leu Ala Trp Tyr Gln Gln Lys Asp Gly Lys Ser Pro Gln Leu Leu Ile
 35 40 45
 Tyr Ala Ala Thr His Leu Val Glu Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Gln Tyr Ser Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80
 Glu Asp Phe Gly Asn Tyr Tyr Cys Gln His Phe Trp Gly Thr Pro Xaa
 85 90 95
 Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg
 100 105

<210> SEQ ID NO 3
 <211> LENGTH: 756
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Signal-VH-hCH1

-continued

<400> SEQUENCE: 3

```

atggaatggt gctgggtctt tctcttcctc ctgtcagtaa ctgcaggtgt ccaactccaag    60
gtccaactgc agcagctctgg agctgagctg aagaaacccg gggcgctcgg gaaagtctcc    120
tgcaaggcgt ctggttacac cttcactgaa tatattatac actggataaa gctgaggtct    180
ggacagggtc ttgagtggat tgggtgggtt taccctggaa gtaatgatat acagtacaat    240
gcgcaattca agggcaaggc cacattgact gcggacaaat cctccagcac cgtctatatg    300
gaacttactg gattgacacc cgaggactct gcggtctatt tttgtgcaag acgcgacgat    360
ttctctgggt acgacgcctt tccttactgg ggccaaggga ctctggtcac tgtctctgca    420
gctagcacca agggcccatc ggtcttcccc ctggcacccct cctccaagag cacctctggg    480
ggcacagcgg ccctgggctg cctgggtcaag gactacttcc ccgaaccggt gacgggtgctg    540
tggaactcag gcgcctgac cagcggcgtg cacaccttcc cggtctcct acagtcctca    600
ggactctact ccctcagcag cgtgggtgacc gtgccctcca gcagcttggg caccagacc    660
tacatctgca acgtgaatca caagcccagc aacaccaagg tggacaagaa agttgagccc    720
aaatcttggt acaaaaactca cacatgcgcc gcataa                                756

```

<210> SEQ ID NO 4

<211> LENGTH: 251

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Signal-VH-hCH1

<400> SEQUENCE: 4

```

Met Glu Trp Cys Trp Val Phe Leu Phe Leu Leu Ser Val Thr Ala Gly
1           5           10           15
Val His Ser Lys Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Lys Lys
20           25           30
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe
35           40           45
Thr Glu Tyr Ile Ile His Trp Ile Lys Leu Arg Ser Gly Gln Gly Leu
50           55           60
Glu Trp Ile Gly Trp Phe Tyr Pro Gly Ser Asn Asp Ile Gln Tyr Asn
65           70           75           80
Ala Gln Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser
85           90           95
Thr Val Tyr Met Glu Leu Thr Gly Leu Thr Pro Glu Asp Ser Ala Val
100          105          110
Tyr Phe Cys Ala Arg Arg Asp Asp Phe Ser Gly Tyr Asp Ala Leu Pro
115          120          125
Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ala Ala Ser Thr Lys
130          135          140
Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly
145          150          155          160
Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro
165          170          175
Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr
180          185          190
Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val
195          200          205
Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn
210          215          220

```

-continued

Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys Lys Val Glu Pro
225 230 235 240

Lys Ser Cys Asp Lys Thr His Thr Cys Ala Ala
245 250

<210> SEQ ID NO 5
<211> LENGTH: 705
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Signal-VL-hCkappa

<400> SEQUENCE: 5

atgagtgtgc cactcaggt cctgggggtg ctgctgctgt ggcttacaga tgccagatgt 60
gacatccaga tgactcagtc tccatcttcc ctatctgcat ctgtgggaga cagggtcacc 120
atcacgtgta aaacaaatga gaatttttac agtaatttag catggtatca gcagaaagac 180
ggaaaatctc ctcagctcct gatctatgct gcaacacact tagtagaggg tgtgccatca 240
aggttcagtg gcagtgatc aggcacacag tattccctca caatcagcag cctgcagcca 300
gaagatthtg ggaattatta ctgtcaacac ttttggggta ctccgtgcac gttcggaggg 360
gggaccaagc tggaaataaa acggacgggtg gctgcacat ctgtcttcat cttcccgcc 420
tctgatgagc agttgaaatc tggaaactgcc tctgttgtgt gcctgctgaa taacttctat 480
cccagagagg ccaaagtaca gtggaagggtg gataacgccc tccaatcggg taactcccag 540
gagagtgtca cagagcagga cagcaaggac agcacctaca gcctcagcag caccctgacg 600
ctgagcaaag cagactacga gaaacacaaa gtctacgct gcgaagtcac ccatcagggc 660
ctgagttcgc ccgtcacaaa gagcttcaac aggggagagt gttaa 705

<210> SEQ ID NO 6
<211> LENGTH: 234
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Signal-VL-hCkappa
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (116)..(116)
<223> OTHER INFORMATION: Xaa = Cys or Ala

<400> SEQUENCE: 6

Met Ser Val Pro Thr Gln Val Leu Gly Leu Leu Leu Leu Trp Leu Thr
1 5 10 15
Asp Ala Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser
20 25 30
Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Lys Thr Asn Glu Asn
35 40 45
Ile Tyr Ser Asn Leu Ala Trp Tyr Gln Gln Lys Asp Gly Lys Ser Pro
50 55 60
Gln Leu Leu Ile Tyr Ala Ala Thr His Leu Val Glu Gly Val Pro Ser
65 70 75 80
Arg Phe Ser Gly Ser Gly Ser Gly Thr Gln Tyr Ser Leu Thr Ile Ser
85 90 95
Ser Leu Gln Pro Glu Asp Phe Gly Asn Tyr Tyr Cys Gln His Phe Trp
100 105 110
Gly Thr Pro Xaa Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg
115 120 125

-continued

Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln
 130 135 140

Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr
 145 150 155 160

Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser
 165 170 175

Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr
 180 185 190

Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys
 195 200 205

His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro
 210 215 220

Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 225 230

<210> SEQ ID NO 7
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro
 1 5 10 15

<210> SEQ ID NO 8
 <211> LENGTH: 10
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

Asp Lys Thr His Thr Cys Pro Pro Cys Pro
 1 5 10

<210> SEQ ID NO 9
 <211> LENGTH: 1107
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: hVHCD28.3-short hingegamma1-hgamma4CH2CH3

<400> SEQUENCE: 9

atggaatggt gctgggtgtt cctgttctctg ctgtccgtga ccgctggcgt gcaactccaag 60
 cagggtgcagc tgcagcagtc tggcgccgag ctgaagaagc ctggcgccctc cgtcaagggtg 120
 tcctgcaagg cctccggcta caccttcacc gagtacatca tccactggat caagctgaga 180
 tccggccagg gcctggaatg gatcggctgg ttctaccctg gctccaacga catccagtac 240
 aacgcccagt tcaagggcaa ggccaccctg accgcccaga agtcctcctc caccgtgtac 300
 atggaactga ccggcctgac ccctgaggac tccgccgtgt acttctgcgc caggcgggac 360
 gacttctctg gctacgacgc cctgccttat tggggccagg gcaccctggt gaccgtgtcc 420
 gccgacaaaa ctcacacatg cccaccgtgc ccagcacctg agttcctggg gggaccatca 480
 gtcttctctg tcccccaaaa acccaaggac actctcatga tctcccggac cctgagggtc 540
 acgtgcgtgg tgggtggacgt gagccaggaa gaccccagg tccagttcaa ctggtacgtg 600
 gatggcgtgg aggtgcataa tgccaagaca aagccgcggg aggagcagtt caacagcacg 660
 taccgtgtgg tcagcgtcct caccgtcctg caccaggact ggctgaacgg caaggagtac 720
 aagtgcaagg tctccaacaa aggcctcccg tcctccatcg agaaaacat ctccaaagcc 780

-continued

```

aaagggcagc cccgagagcc acaggtgtac accctgcccc catcccagga ggagatgacc 840
aagaaccagg tcagcctgac ctgcctggtc aaaggcttct accccagcga catcgccgtg 900
gagtgggaga gcaatgggca gccggagaac aactacaaga ccacgcctcc cgtgctggac 960
tccgacggct ccttcttct ctacagcagg ctcaccgtgg acaagagcag gtggcaggag 1020
gggaatgtct tctcatgctc cgtgatgcat gaggctctgc acaaccacta cacacagaag 1080
agcctctccc tgtctctggg taaatga 1107

```

```

<210> SEQ ID NO 10
<211> LENGTH: 368
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: hVHCD28.3-short hingegamma1-hgamma4CH2CH3

```

```

<400> SEQUENCE: 10

```

```

Met Glu Trp Cys Trp Val Phe Leu Phe Leu Leu Ser Val Thr Ala Gly
1           5           10           15
Val His Ser Lys Gln Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Lys
20           25           30
Lys Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr
35           40           45
Phe Thr Glu Tyr Ile Ile His Trp Ile Lys Leu Arg Ser Gly Gln Gly
50           55           60
Leu Glu Trp Ile Gly Trp Phe Tyr Pro Gly Ser Asn Asp Ile Gln Tyr
65           70           75           80
Asn Ala Gln Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser
85           90           95
Ser Thr Val Tyr Met Glu Leu Thr Gly Leu Thr Pro Glu Asp Ser Ala
100          105          110
Val Tyr Phe Cys Ala Arg Arg Asp Asp Phe Ser Gly Tyr Asp Ala Leu
115          120          125
Pro Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ala Asp Lys Thr
130          135          140
His Thr Cys Pro Pro Cys Pro Ala Pro Glu Phe Leu Gly Gly Pro Ser
145          150          155          160
Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg
165          170          175
Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp Pro
180          185          190
Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala
195          200          205
Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val Val
210          215          220
Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr
225          230          235          240
Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys Thr
245          250          255
Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu
260          265          270
Pro Pro Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys
275          280          285
Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser
290          295          300

```


-continued

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp
 305 310 315 320

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys Ser
 325 330 335

Arg Trp Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
 340 345 350

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Leu Gly Lys
 355 360 365

<210> SEQ ID NO 11
 <211> LENGTH: 1068
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: hVLCD28.3-short hingegamma1-hgamma4CH2CH3

<400> SEQUENCE: 11

atgtccgtgc ctaccaggt gctgggactg ctgctgctgt ggctgaccga cgccagatgc 60
 gacatccaga tgaccagtc cccctcctcc ctgtctgcct ccgtgggcca ccgggtgacc 120
 atcacctgta agaccaacga gaacatctac tccaacctgg cctggtatca gcagaaggac 180
 ggcaagtccc ctcagctgct gatctacgcc gccacccatc tgggtggaggc cgtgcctct 240
 agattctcgc gctccggctc tggcaccag tactccctga ccatcagctc cctgcagcct 300
 gaggacttcg gcaactacta ctgccagcac ttctggggca ccccttgtag cttcggcgga 360
 ggaccaagc tggaaatcaa gcgggacaaa actcacacat gcccaccgtg cccagcacct 420
 gaggctcctg ggggaccatc agtcttctg ttcccccaa aaccaagga cactctcatg 480
 atctcccga cccctgaggt cacgtgcgtg gtggtggagc tgagccagga agaccccgag 540
 gttccagttca actggtacgt ggatggcgtg gaggtgcata atgccaagac aaagccgcgg 600
 gaggagcagt tcaacagcac gtaccgtgtg gtcagcgtcc tcaccgtcct gcaccaggac 660
 tggctgaacg gcaaggagta caagtgcaag gtctccaaca aaggcctccc gtectccatc 720
 gagaaaacca tctccaaagc caaagggcag ccccgagagc cacaggtgta caccctgccc 780
 ccatcccagg aggagatgac caagaaccag gtcagcctga cctgcctggt caaaggcttc 840
 taccacagcg acatcgccgt ggagtgggag agcaatgggc agccggagaa caactacaag 900
 accacgcctc ccgtgctgga ctccgacggc tccttcttcc tctacagcag gctcaccgtg 960
 gacaagagca ggtggcagga ggggaatgtc ttctcatgct ccgtgatgca tgaggctctg 1020
 cacaaccact acacacagaa gagcctctcc ctgtctctgg gtaaatga 1068

<210> SEQ ID NO 12
 <211> LENGTH: 355
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: hVLCD28.3-short hingegamma1-hgamma4CH2CH3

<400> SEQUENCE: 12

Met Ser Val Pro Thr Gln Val Leu Gly Leu Leu Leu Leu Trp Leu Thr
 1 5 10 15

Asp Ala Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser
 20 25 30

Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Lys Thr Asn Glu Asn
 35 40 45

Ile Tyr Ser Asn Leu Ala Trp Tyr Gln Gln Lys Asp Gly Lys Ser Pro
 50 55 60

-continued

Gln Leu Leu Ile Tyr Ala Ala Thr His Leu Val Glu Gly Val Pro Ser
 65 70 75 80
 Arg Phe Ser Gly Ser Gly Ser Gly Thr Gln Tyr Ser Leu Thr Ile Ser
 85 90 95
 Ser Leu Gln Pro Glu Asp Phe Gly Asn Tyr Tyr Cys Gln His Phe Trp
 100 105 110
 Gly Thr Pro Cys Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg
 115 120 125
 Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Phe Leu Gly
 130 135 140
 Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met
 145 150 155 160
 Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln
 165 170 175
 Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val
 180 185 190
 His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr
 195 200 205
 Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly
 210 215 220
 Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile
 225 230 235 240
 Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val
 245 250 255
 Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser
 260 265 270
 Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu
 275 280 285
 Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro
 290 295 300
 Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val
 305 310 315 320
 Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met
 325 330 335
 His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser
 340 345 350
 Leu Gly Lys
 355

<210> SEQ ID NO 13
 <211> LENGTH: 1122
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: hVHCD28.3-full hingegamma1-hgamma4CH2CH3
 <400> SEQUENCE: 13

atggaatggt gctgggtggt cctgttctctg ctgtccgtga ccgctggcgt gcaactccaag 60
 cagggtgcagc tgcagcagtc tggcgccgag ctgaagaagc ctggcgcctc cgtcaagggtg 120
 tcctgcaagg cctccggcta caccttcacc gagtacatca tccactggat caagctgaga 180
 tccggccagg gcttggatg gatcggtctg ttctaccctg gctccaacga catccagtac 240
 aacgcccagt tcaagggcaa ggccaccctg accgcccaga agtcctcctc caccgtgtac 300
 atggaactga ccggcctgac cctgaggac tccgccgtgt acttctgcgc caggcgggac 360

-continued

```

gacttctctg gctacgacgc cctgccttat tggggccagg gcaccctggt gaccgtgtcc 420
gccgagccca aatcttgtga caaaactcac acatgcccac cgtgcccagc acctgagttc 480
ctgggggggac catcagtctt cctgttcccc ccaaaccaca aggacactct catgatctcc 540
cggacccttg aggtcacgtg cgtggtggtg gacgtgagcc aggaagaccg cgaggtccag 600
ttcaactggt acgtggatgg cgtggaggtg cataatgcca agacaaagcc gcgggaggag 660
cagttcaaca gcacgtaccg tgtggtcagc gtctcaccg tcctgcacca ggactggctg 720
aacggcaagg agtacaagtg caaggtctcc aacaaaggcc tcccgtctc catcgagaaa 780
accatctcca aagccaaagg gcagccccga gagccacagg tgtacaccct gccccatcc 840
caggaggaga tgaccaagaa ccaggtcagc ctgacctgcc tggtaaagg cttctacccc 900
agcgacatcg ccgtggagtg ggagagcaat gggcagccgg agaacaacta caagaccag 960
cctcccgtgc tggactccga cggctccttc ttctctaca gcaggctcac cgtggacaag 1020
agcaggtggc aggaggggaa tgtcttctca tgctccgtga tgcattgaggc tctgcacaac 1080
cactacacac agaagagcct ctccctgtct ctgggtaaat ga 1122

```

<210> SEQ ID NO 14

<211> LENGTH: 373

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: hVHCD28.3-full hingegamma1-hgamma4CH2CH3

<400> SEQUENCE: 14

```

Met Glu Trp Cys Trp Val Phe Leu Phe Leu Leu Ser Val Thr Ala Gly
1           5           10          15
Val His Ser Lys Gln Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Lys
20          25          30
Lys Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr
35          40          45
Phe Thr Glu Tyr Ile Ile His Trp Ile Lys Leu Arg Ser Gly Gln Gly
50          55          60
Leu Glu Trp Ile Gly Trp Phe Tyr Pro Gly Ser Asn Asp Ile Gln Tyr
65          70          75          80
Asn Ala Gln Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser
85          90          95
Ser Thr Val Tyr Met Glu Leu Thr Gly Leu Thr Pro Glu Asp Ser Ala
100         105         110
Val Tyr Phe Cys Ala Arg Arg Asp Asp Phe Ser Gly Tyr Asp Ala Leu
115        120        125
Pro Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ala Glu Pro Lys
130        135        140
Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Phe
145        150        155        160
Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr
165        170        175
Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val
180        185        190
Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val
195        200        205
Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser
210        215        220

```

-continued

Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu
 225 230 235 240

Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser
 245 250 255

Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro
 260 265 270

Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys Asn Gln
 275 280 285

Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala
 290 295 300

Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr
 305 310 315 320

Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Arg Leu
 325 330 335

Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser Cys Ser
 340 345 350

Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser
 355 360 365

Leu Ser Leu Gly Lys
 370

<210> SEQ ID NO 15
 <211> LENGTH: 1083
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: hVLCD28.3-full hingegamma1-hgamma4CH2CH3

<400> SEQUENCE: 15

atgtccgtgc ctaccaggt gctgggactg ctgctgctgt ggctgaccga cgccagatgc 60

gacatccaga tgaccagtc cccctcctcc ctgtctgect ccgtgggcca ccgggtgacc 120

atcacctgta agaccaacga gaacatctac tccaacctgg cctggtatca gcagaaggac 180

ggcaagtccc ctgagctgct gatctacgcc gccacccatc tgggtggaggc cgtgccctct 240

agattctcgc gctccggctc tggcaccag tactccctga ccatcagctc cctgcagcct 300

gaggacttcg gcaactacta ctgccagcac ttctggggca ccccttgtag cttcggcgga 360

ggcaccaagc tggaaatcaa gcgggagccc aaatcttgtag acaaaactca cacatgcccc 420

ccgtgcccag cacctgagtt cctgggggga ccatcagctc tcctgttccc cccaaaaccc 480

aaggacactc tcatgatctc ccggaccctc gaggtcacgt gcgtgggtgg ggacgtgagc 540

caggaagacc ccgaggtcca gttcaactgg tacgtggatg gcgtggaggt gcataatgcc 600

aagacaaagc cgcgggagga gcagttcaac agcacgtacc gtgtggtcag cgtcctcacc 660

gtcctgcacc aggactggct gaacggcaag gactacaagt gcaaggtctc caacaaaggc 720

ctcccgtcct ccatcgagaa aacctctcc aaagccaaag ggcagccccg agagccacag 780

gtgtacacc tgccccatc ccaggaggag atgaccaaga accaggtcag cctgacctgc 840

ctgggtcaaag gcttctacc cagcgacatc gccgtggagt gggagagcaa tgggcagccg 900

gagaacaact acaagaccac gcctcccgtg ctggactccg acggctcctt ctctctctac 960

agcaggctca ccgtggacaa gagcaggtgg caggagggga atgtcttctc atgctccgtg 1020

atgcatgagg ctctgcacaa ccactacaca cagaagagcc tctccctgct tctgggtaaa 1080

tga 1083

-continued

```

<210> SEQ ID NO 16
<211> LENGTH: 360
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: hVLCD28.3-full hinge $\gamma$ 1-h $\gamma$ 4CH2CH3

<400> SEQUENCE: 16

Met Ser Val Pro Thr Gln Val Leu Gly Leu Leu Leu Trp Leu Thr
1          5          10          15

Asp Ala Arg Cys Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser
          20          25          30

Ala Ser Val Gly Asp Arg Val Thr Ile Thr Cys Lys Thr Asn Glu Asn
          35          40          45

Ile Tyr Ser Asn Leu Ala Trp Tyr Gln Gln Lys Asp Gly Lys Ser Pro
          50          55          60

Gln Leu Leu Ile Tyr Ala Ala Thr His Leu Val Glu Gly Val Pro Ser
65          70          75          80

Arg Phe Ser Gly Ser Gly Ser Gly Thr Gln Tyr Ser Leu Thr Ile Ser
          85          90          95

Ser Leu Gln Pro Glu Asp Phe Gly Asn Tyr Tyr Cys Gln His Phe Trp
          100         105         110

Gly Thr Pro Cys Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys Arg
          115         120         125

Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala
130         135         140

Pro Glu Phe Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro
145         150         155         160

Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val
          165         170         175

Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val
          180         185         190

Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln
195         200         205

Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln
210         215         220

Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly
225         230         235         240

Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro
          245         250         255

Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr
          260         265         270

Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser
          275         280         285

Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr
290         295         300

Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr
305         310         315         320

Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe
          325         330         335

Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys
          340         345         350

Ser Leu Ser Leu Ser Leu Gly Lys
          355         360

```

-continued

<210> SEQ ID NO 17
 <211> LENGTH: 60
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Nucleotide for leader sequence

<400> SEQUENCE: 17

atggaatggt gctgggtctt tctcttctc ctgtcagtaa ctgcaggtgt ccaactccaag 60

<210> SEQ ID NO 18
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Amind acid for leader sequence

<400> SEQUENCE: 18

Met Glu Trp Cys Trp Val Phe Leu Phe Leu Leu Ser Val Thr Ala Gly
 1 5 10 15

Val His Ser Lys
 20

<210> SEQ ID NO 19
 <211> LENGTH: 4
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: AA - VH - CDR1

<400> SEQUENCE: 19

Thr Glu Tyr Ile
 1

<210> SEQ ID NO 20
 <211> LENGTH: 17
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: AA - VH - CDR2

<400> SEQUENCE: 20

Ile Gly Trp Phe Tyr Pro Gly Ser Asn Asp Ile Gln Tyr Asn Ala Gln
 1 5 10 15

Phe

<210> SEQ ID NO 21
 <211> LENGTH: 13
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: AA - VH - CDR3

<400> SEQUENCE: 21

Ala Arg Arg Asp Asp Phe Ser Gly Tyr Asp Ala Leu Pro
 1 5 10

<210> SEQ ID NO 22
 <211> LENGTH: 336
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: VT - hCH1

<400> SEQUENCE: 22

gctagcacca agggcccatc ggtcttcccc ctggcacct cctccaagag cacctctggg 60

-continued

```

ggcacagegg ccctgggctg cctgggtcaag gactacttcc ccgaaccggt gacgggtgctg 120
tggaactcag ggcacctgac cagcggcgctg cacaccttcc cggctgtcct acagtctctca 180
ggactctact ccctcagcag cgtgggtgacc gtgccctcca gcagcttggg caccagacc 240
tacatctgca acgtgaatca caagcccagc aacaccaagg tggacaagaa agttgagccc 300
aaatcttggtg acaaaactca cacatgctgcc gcataa 336

```

```

<210> SEQ ID NO 23
<211> LENGTH: 100
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: AA - hCH1

```

```

<400> SEQUENCE: 23

```

```

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
1           5           10           15
Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
          20           25           30
Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
          35           40           45
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
          50           55           60
Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
65           70           75           80
Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
          85           90           95
Lys Val Glu Pro
          100

```

```

<210> SEQ ID NO 24
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: NT - leader seq

```

```

<400> SEQUENCE: 24

```

```

atgagtgctgc ccactcaggt cctgggggttg ctgctgctgt ggcttacaga tgccagatgt 60

```

```

<210> SEQ ID NO 25
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: AA - leader seq

```

```

<400> SEQUENCE: 25

```

```

Met Ser Val Pro Thr Gln Val Leu Gly Leu Leu Leu Leu Trp Leu Thr
1           5           10           15
Asp Ala Arg Cys
          20

```

```

<210> SEQ ID NO 26
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: aa - vl - CDR1

```

```

<400> SEQUENCE: 26

```

-continued

Asn Leu Ala Trp Tyr Gln Gln Lys
1 5

<210> SEQ ID NO 27
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: aa - v1 - CDR2

<400> SEQUENCE: 27

Ala Ala Thr His Leu Val Glu Gly
1 5

<210> SEQ ID NO 28
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: aa - v1 - CDR3

<400> SEQUENCE: 28

His Phe Trp Gly Thr Pro Cys
1 5

<210> SEQ ID NO 29
<211> LENGTH: 318
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: NT - human c kappa region

<400> SEQUENCE: 29

gtggctgcac catctgtctt catcttcccg ccatctgatg agcagttgaa atctggaact 60
gcctctggtg tgtgcttctt gaataacttc tatcccagag aggccaaagt acagtggaag 120
gtggataacg cctccaatc gggtaactcc caggagagtg tcacagagca ggacagcaag 180
gacagcacct acagcctcag cagcaccctg acgtgagca aagcagacta cgagaaacac 240
aaagtctacg cctgcgaagt caccatcag ggctgagtt cgcccgtcac aaagagcttc 300
aacaggggag agtggttaa 318

<210> SEQ ID NO 30
<211> LENGTH: 105
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: aa - human c kappa region

<400> SEQUENCE: 30

Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu
1 5 10 15
Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro
20 25 30
Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly
35 40 45
Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr
50 55 60
Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His
65 70 75 80
Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val
85 90 95

-continued

Thr Lys Ser Phe Asn Arg Gly Glu Cys
100 105

<210> SEQ ID NO 31
<211> LENGTH: 217
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: CH2-CH3 domains

<400> SEQUENCE: 31

Ala Pro Glu Phe Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys
1 5 10 15
Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val
20 25 30
Val Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr
35 40 45
Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu
50 55 60
Gln Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His
65 70 75 80
Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys
85 90 95
Gly Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln
100 105 110
Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met
115 120 125
Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro
130 135 140
Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn
145 150 155 160
Tyr Lys Thr Thr Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu
165 170 175
Tyr Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val
180 185 190
Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln
195 200 205
Lys Ser Leu Ser Leu Ser Leu Gly Lys
210 215

The invention claimed is:

1. A method of treating a T-lymphocyte-mediated auto-immune disease in a subject in need thereof comprising administering an anti-CD28 monovalent antibody to the subject, wherein the anti-CD28 monovalent antibody is selected from the group consisting of:

- (a) an antibody having a CD28-binding site consisting of:
a heavy chain variable domain of SEQ ID NO: 1; and
a light chain variable domain of SEQ ID NO: 2; and
(b) an antibody having a CD28-binding site consisting of:
a heavy chain variable domain having all three complementarity determining regions (CDRs) of the variable domain of SEQ ID NO: 1; and
a light chain variable domain having all three CDRs of the variable domain of SEQ ID NO: 2.

2. The method of claim 1, wherein the monovalent antibody is a heterodimer of:

- a first protein chain having the sequence of amino-acids 21-251 of SEQ ID NO: 4; and

a second protein chain having the sequence of amino-acids 21-234 of SEQ ID NO: 6.

3. The method of claim 1, wherein the monovalent antibody is a heterodimer of:

(I) a first protein chain consisting essentially of, from its N-terminus to its C-terminus:

- i: a region A which is a heavy chain variable domain of SEQ ID NO: 1; and
ii: a region B consisting of a peptide linker followed by the CH2 and CH3 domains of an IgG immunoglobulin, and

(II) a second protein chain consisting essentially of, from its N-terminus to its C-terminus:

- i: a region A' which is a light chain variable domain of SEQ ID NO: 2; and
ii: a region B identical to the region B of the first protein chain.

4. The method of claim 3, wherein the peptide linker is selected from the group consisting of:

43

a peptide of SEQ ID NO: 7; and
a peptide of SEQ ID NO: 8.

5 5. The method of claim 3, wherein the CH2 and CH3 domains are those of an immunoglobulin of the IgG4 subclass.

6. The method of claim 5, wherein the monovalent antibody is selected from the group consisting of:

a monovalent antibody wherein the polypeptide sequence of the first protein chain is the sequence of amino-acids 21-368 of SEQ ID NO: 10, and the polypeptide sequence of the second protein chain is the sequence of amino-acids 21-355 of SEQ ID NO: 12; and

10 a monovalent antibody wherein the polypeptide sequence of the first protein chain is the sequence of amino-acids 21-373 of SEQ ID NO: 14, and the polypeptide sequence of the second protein chain is the sequence of amino-acids 21-360 of SEQ ID NO: 16.

44

7. The method of claim 2 wherein the second protein chain comprises a variable domain of SEQ ID NO: 2.

8. The method of claim 1, wherein the monovalent antibody is administered in a composition and the composition further comprises a pharmaceutically acceptable excipient.

9. The method of claim 2, wherein the second protein chain comprises a variable domain of SEQ ID NO: 2 where X represents an alanine or an asparagine residue.

10 10. The method of claim 2, wherein the monovalent antibody is pegylated.

11. The method of claim 1, wherein the heavy chain variable domain further comprises a Q residue at the N-terminal end.

15 12. The method of claim 1, wherein the autoimmune disease is selected from the group consisting of type I diabetes, rheumatoid arthritis, and multiple sclerosis.

* * * * *