

US009598375B2

(12) **United States Patent**
Jones et al.(10) **Patent No.:** **US 9,598,375 B2**
(45) **Date of Patent:** **Mar. 21, 2017**(54) **SUBSTITUTED IMIDAZOLE DERIVATIVES
AND METHODS OF USE THEREOF**

- (71) Applicant: **vTv Therapeutics LLC**, High Point, NC (US)
- (72) Inventors: **David Jones**, Milford, OH (US); **Raju Bore Gowda**, Oak Ridge, NC (US); **Rongyuan Xie**, Greensboro, NC (US)
- (73) Assignee: **vTv Therapeutics LLC**, High Point, NC (US)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 241 days.

(21) Appl. No.: **14/049,261**(22) Filed: **Oct. 9, 2013**(65) **Prior Publication Data**

US 2014/0039025 A1 Feb. 6, 2014

Related U.S. Application Data

(62) Division of application No. 12/888,660, filed on Sep. 23, 2010, now Pat. No. 8,580,833.

(60) Provisional application No. 61/247,206, filed on Sep. 30, 2009.

(51) **Int. Cl.**
C07D 233/64 (2006.01)
C07D 233/60 (2006.01)(52) **U.S. Cl.**
CPC **C07D 233/64** (2013.01); **C07D 233/60** (2013.01)(58) **Field of Classification Search**
CPC **C07D 233/64**
See application file for complete search history.(56) **References Cited**

U.S. PATENT DOCUMENTS

3,255,202 A	6/1966	Johnson
3,708,598 A	1/1973	Griot
3,951,968 A	4/1976	Fauran et al.
4,024,271 A	5/1977	Durant et al.
4,032,522 A	6/1977	Baldwin et al.
4,166,452 A	9/1979	Generales, Jr.
4,265,874 A	5/1981	Bonsen et al.
4,356,108 A	10/1982	Schwab et al.
4,873,313 A	10/1989	Crawford et al.
4,933,422 A	6/1990	Hammer
4,963,539 A	10/1990	Delaney
5,011,849 A	4/1991	Gassner et al.
5,153,226 A	10/1992	Chucholowski et al.
5,166,214 A	11/1992	Billheimer et al.
5,179,210 A	1/1993	Ebel
5,192,785 A	3/1993	Lo et al.
5,202,424 A	4/1993	Vlassara et al.
5,318,984 A	6/1994	Billheimer et al.
5,358,960 A	10/1994	Ulrich et al.
5,500,436 A	3/1996	Schoenafinger et al.
5,523,317 A	6/1996	Masaki et al.
5,585,344 A	12/1996	Vlassara et al.
5,589,496 A	12/1996	Hamanaka et al.

5,663,186 A	9/1997	Nelson et al.
5,688,653 A	11/1997	Ulrich et al.
5,703,092 A	12/1997	Xue et al.
5,795,907 A	8/1998	Kalindjian et al.
5,817,626 A	10/1998	Findeis et al.
5,817,823 A	10/1998	Hong et al.
5,840,294 A	11/1998	Kisilevsky et al.
5,864,018 A	1/1999	Morser et al.
5,922,770 A	7/1999	Peschke et al.
5,939,526 A	8/1999	Gaugler et al.
5,962,500 A	10/1999	Eide et al.
5,962,535 A	10/1999	Miyamoto et al.
6,034,250 A	3/2000	Goldstein et al.
6,100,098 A	8/2000	Newkirk
6,197,791 B1	3/2001	Venkatesan et al.
6,201,002 B1	3/2001	Beere et al.
6,221,667 B1	4/2001	Reiner et al.
6,265,351 B1	7/2001	Porta et al.
6,268,479 B1	7/2001	Stern et al.
6,274,615 B1	8/2001	Pappolla et al.
6,277,853 B1	8/2001	Perez et al.
6,300,356 B1	10/2001	Segal et al.
6,316,474 B1	11/2001	McCauley et al.

(Continued)

FOREIGN PATENT DOCUMENTS

EP 0584588 A1 3/1994
EP 0586806 A1 3/1994

(Continued)


OTHER PUBLICATIONS

Albericio et al., "Coupling Reagents and Activation," *Methods in Enzymology* 289:104-126 (1997).

(Continued)

Primary Examiner — Samira Jean-Louis(74) *Attorney, Agent, or Firm* — Samuel B. Rollins(57) **ABSTRACT**

The present invention provides imidazole derivatives of Formula (I) and pharmaceutically acceptable salts thereof.



and their use in the treatment of diseases such as Alzheimer's Disease.

9 Claims, No Drawings

(56)

References Cited

U.S. PATENT DOCUMENTS

6,323,218	B1	11/2001	Bush et al.
6,353,009	B1	3/2002	Fujiwara et al.
6,416,733	B1	7/2002	Barrett et al.
6,441,023	B1	8/2002	Venkatesan et al.
6,441,049	B2	8/2002	Reitz et al.
6,441,064	B1	8/2002	Shah et al.
6,472,145	B2	10/2002	Reiner et al.
6,538,013	B2	3/2003	Goebel et al.
6,541,639	B2	4/2003	Zhou et al.
6,613,801	B2	9/2003	Mjalli et al.
6,673,810	B2	1/2004	Lam et al.
6,673,927	B2	1/2004	Gordon et al.
6,677,299	B2	1/2004	Stern et al.
6,730,796	B2	5/2004	Cheng et al.
6,825,164	B1	11/2004	Stern et al.
6,919,338	B2	7/2005	Mortlock et al.
7,067,554	B2	6/2006	Mjalli et al.
7,087,632	B2	8/2006	Mjalli et al.
7,329,684	B2	2/2008	Mjalli et al.
7,361,678	B2	4/2008	Mjalli et al.
7,421,177	B2	9/2008	Schmidt et al.
7,423,177	B2	9/2008	Mjalli et al.
7,714,013	B2	5/2010	Mjalli et al.
7,737,285	B2	6/2010	Mjalli et al.
7,776,919	B2	8/2010	Mjalli et al.
7,884,219	B2	2/2011	Hari
8,372,988	B2	2/2013	Hari
8,580,833	B2	11/2013	Jones et al.
2001/0039256	A1	11/2001	Stern et al.
2002/0006957	A1	1/2002	Mjalli et al.
2002/0116725	A1	8/2002	Stern et al.
2002/0122799	A1	9/2002	Stern et al.
2002/0193432	A1	12/2002	Mjalli et al.
2003/0032663	A1	2/2003	Mjalli et al.
2003/0207896	A1	11/2003	Konno et al.
2003/0236282	A1	12/2003	Hurnaus et al.
2004/0063770	A1	4/2004	Ahn et al.
2004/0082542	A1	4/2004	Mjalli et al.
2004/0097407	A1	5/2004	Mjalli et al.
2004/0127692	A1	7/2004	David et al.
2005/0026811	A1	2/2005	Mjalli et al.
2006/0020042	A1	1/2006	McDonald et al.
2006/0247253	A1	11/2006	Leban et al.
2007/0021386	A1	1/2007	Mjalli et al.
2007/0135437	A1	6/2007	Benjamin et al.
2009/0035302	A1	2/2009	Mjalli et al.
2010/0048726	A1	2/2010	McDonald et al.
2010/0256119	A1	10/2010	Mjalli et al.
2012/0088778	A1	4/2012	Mjalli et al.
2014/0039025	A1	2/2014	Jones et al.

FOREIGN PATENT DOCUMENTS

EP	0633026	A1	1/1995
EP	0707000	A1	4/1996
EP	1139990		10/2001
FR	1476560		4/1967
FR	2773800	A1	7/1999
GB	2005674	A	4/1979
JP	60-080656	A	3/1994
JP	90-040651	A	2/1997
JP	2003-012690	A	1/2003
JP	2003-040888	A	2/2003
JP	2003-313170	A	11/2003
JP	2003-313172	A	11/2003
JP	2004-221557	A	8/2004
WF	WO 00/66102	A2	11/2000
WO	WO 93/09100	A1	5/1993
WO	WO 95/01340	A1	1/1995
WO	WO 95/02591	A1	1/1995
WO	WO 95/09838	A1	4/1995
WO	WO 95/30647	A1	11/1995
WO	WO 95/35279	A1	12/1995
WO	WO 96/32385	A1	10/1996
WO	WO 97/22618	A1	6/1997

WO	WO 97/26913	A1	7/1997
WO	WO 97/39121	A1	10/1997
WO	WO 97/39125	A1	10/1997
WO	WO 98/22138	A1	5/1998
WO	WO 98/27108	A2	6/1998
WO	WO 98/33492	A1	8/1998
WO	WO 98/35945	A1	8/1998
WO	WO 98/37877	A1	9/1998
WO	WO 99/07402	A1	2/1999
WO	WO 99/16755	A1	4/1999
WO	WO 99/18987	A1	4/1999
WO	WO 99/25690	A2	5/1999
WO	WO 99/50230	A1	10/1999
WO	WO 99/54485	A1	10/1999
WO	WO 00/19994	A1	4/2000
WO	WO 00/20458	A1	4/2000
WO	WO 00/20621	A1	4/2000
WO	WO 00/38635	A1	7/2000
WO	WO 01/12598	A2	2/2001
WO	WO 01/32604	A1	5/2001
WO	WO 01/92210	A1	12/2001
WO	WO 02/069965	A1	9/2002
WO	WO 02/070473	A2	9/2002
WO	WO 03/024937	A1	3/2003
WO	WO 03/053922	A2	7/2003
WO	WO 03/075921	A2	9/2003
WO	WO 03/086390	A1	10/2003
WO	WO 03/075921	A3	12/2003
WO	WO 2004/035061	A1	4/2004
WO	WO 2004/046141	A	6/2004
WO	WO 2004/087653	A2	10/2004
WO	WO 2004/110350	A2	12/2004
WO	WO 2005/000295	A1	1/2005
WO	WO 2005/019185	A1	3/2005
WO	WO 2006/124897	A2	11/2006
WO	WO 2008/067121	A2	6/2008
WO	WO 2008/153957	A1	12/2008
WO	WO 2010/126745	A1	11/2010
WO	WO 2011/103091	A1	8/2011

OTHER PUBLICATIONS

- Barton, "Protection of N—H Bonds and NR₃," Protective Groups in Organic Chemistry, McOmie, Ed., pp. 43-93 (1973).
- Behl et al., "Amyloid beta peptide induces necrosis rather than apoptosis," *Brain Research* 645:253-264 (1994).
- Behl, "Hydrogen Peroxide Mediates Amyloid beta Protein Toxicity," *Cell* 77:817-827 (1994).
- Berge et al., "Pharmaceutical Salts," *Journal of Pharmaceutical Sciences* 66(1):1-19 (1977).
- Bierhaus et al., "Advanced Glycation End Product (AGE)-Mediated Induction of Tissue Factor in Cultured Endothelial Cells Is Dependent on RAGE," *Circulation* 96:2262-2271 (1997).
- Blacker et al., "Reliability and Validity of NINCDS-ADRDA Criteria for Alzheimer's Disease," *Arch. Neurol.* 51:1198-1204 (1994).
- Bonnardel-Phu et al., "Acute Modulation of Albumin Microvascular Leakage by Advanced Glycation End Products in Microcirculation of Diabetic Rats In Vivo," *Diabetes* 48:2052-2058 (1999).
- Buttke et al., "Synthesis, Structure, and Photophysical Properties of Polyarylated Imidazoles and Oxazoles," *J. prakt. Chem.* 339:721-728 (1997).
- Chartier-Harlin et al., "Early-onset Alzheimer's disease caused by mutations at codon 717 of the beta-amyloid precursor protein gene," *Nature* 353:844-846 (1991).
- Checler, "Processing of the beta-Amyloid Precursor Protein and Its Regulation in Alzheimer's Disease," *Journal of Neurochemistry* 65(4):1431-1444 (1995).
- Chitale et al., "Antagonism of Rho-kinase stimulates rat penile erection via a nitric oxide-independent pathway," *Nature Medicine* 7(1):119-122 (2001).
- Coskun et al., "The first regio- and diastereoselective synthesis of homochiral perhydroimidazoisoxazoles via the 1,3-dipolar cycloaddition of imidazoline 3-oxides with (1S)-(-)-beta-pinene," *Tetrahedron: Asymmetry* 12:1463-1467 (2001).

(56)

References Cited

OTHER PUBLICATIONS

- Crall, Jr. et al., "The Extramural and Intramural Coronary Arteries in Juvenile Diabetes Mellitus," *The American Journal of Medicine* 64:221-230 (1978).
- Database CAPLUS Abstract of Kuz'Menko et al., "Unusual decomposition of benzimidazolium phenacyl salts. Synthesis of 1,4-diarylimidazoles," *Khimiya Geterotsiklicheskokh Soedinenii*, No. 3, pp. 388-392 (1982).
- Database CAPLUS on STN Abstract of Dobrev et al., "Addition of N,N-disubstituted amides to N-benzoyldiphenylketimine in the presence of lithium amide in liquid ammonia," *God. Sofii. Univ., Khim. Fak.* 70(1):201-207 (1978).
- Database CAPLUS on STN Abstract of Frappier et al., "Peptide Alkaloids. X. Approach for the synthesis of peptidic alkaloids. 1. Reactivity of N-tolylsulfonlaziridines towards reactive nucleophiles," *Tetrahedron* 34(19):2911-2916 (1976).
- Database CAPLUS on STN Abstract of Hamada et al., "Preparation of anilide derivatives for determination of enzymes," 1992.
- Database CAPLUS on STN Abstract of Pirkle et al., "Separation of the enantiomers of N-protected alpha-amino acids as anilide and 3,5-dimethylanilide derivatives," *Journal of Chromatography* 479(2):419-423 (1989).
- Database CAPLUS on STN Abstract of Sivanandaiah et al., "Synthesis of peptides mediated by KOBt," *International Journal of Peptide and Protein Research* 44(1):24-30 (1994).
- Database CAPLUS on STN Abstract of Tsukida et al., "Aminocarboxylic acids, selectin inhibitors containing them, and their uses" (1999).
- Database CAPLUS on STN Abstract of Vidugiriene et al., "Synthesis and study of derivatives of 3-(alkylamino)-2-(methylthio)carboxylic acids," *Chemija* 2:101-106 (1990).
- Database HCAPLUS on STN Abstract of Katzenellenbogen et al., "Preparation of non-steroidal estrogen receptor subtype-selective ligands," Accession No. 2000:240935, Reg. No. 234093-17-5 (2000).
- Deane et al., "RAGE mediated amyloid-beta peptide transport across the blood-brain barrier and accumulation in brain," *Nature Medicine* 9(7):907-913 (2003).
- Degenhardt et al., "Chemical Modification of Proteins by Methylglyoxal," *Cellular and Molecular Biology* 44(7):1139-1145 (1998).
- Denny et al., "Potential Antitumor Agents. 59. Structure-Activity Relationships for 2-Phenylbenzimidazole-4-carboxamides, a New Class of 'Minimal' DNA-Intercalating Agents Which May Not Act via Topoisomerase II," *Journal of Medicinal Chemistry* 33(2):814-819 (1990).
- Digenis et al., "Peptidyl Carbamates Incorporating Amino Acid Isoteres as Novel Elastase Inhibitors," *Journal of Medicinal Chemistry* 29(8):1468-1476 (1986).
- Dyer et al., "Accumulation of Maillard Reaction Products in Skin Collagen in Diabetes and Aging," *J. Clin. Invest.* 91:2463-2469 (1993).
- Dyer et al., "Formation of Pentosidine during Nonenzymatic Browning of Proteins by Glucose," *The Journal of Biological Chemistry* 266(18):11654-11660 (1991).
- Eriks et al., "Histamine H2-Receptor Agonists. Synthesis, in Vitro Pharmacology, and Qualitative Structure-Activity Relationships of Substituted 4- and 5-(2-Aminoethyl)thiazoles," *J. Med. Chem.* 35(17):3239-3246 (1992).
- Evans et al., "Synthesis of a group of 1H-benzimidazoles and their screening for antiinflammatory activity," *Eur. J. Med. Chem.* 31:635-642 (1996).
- Fang et al., "RAGE-dependent signaling in microglia contributes to neuroinflammation, A-beta accumulation, and impaired learning/memory in a mouse model of Alzheimer's disease," *The FASEB Journal* 24:1043-1055 (2010).
- Fink et al., "Novel structural templates for estrogen-receptor ligands and prospects for combinatorial synthesis of estrogens," *Chemistry & Biology* 6(4):205-219 (1999).
- Galasko et al., "Clinical-Neuropathological Correlations in Alzheimer's Disease and Related Dementias," *Arch. Neurol.* 51:888-895 (1994).
- Games et al., "Alzheimer-type neuropathology in transgenic mice overexpressing V717F beta-amyloid precursor protein," *Nature* 373:523-527 (1995).
- Girouard et al., "Neurovascular coupling in the normal brain and in hypertension, stroke, and Alzheimer disease," *J. Appl. Physiol.* 100:328-335 (2006).
- Golub et al., "Molecular Classification of Cancer: Class Discovery and Class Prediction by Gene Expression Monitoring," *Science* 286:531-537 (1999).
- Goova et al., "Blockade of Receptor for Advanced Glycation End-Products Restores Effective Wound Healing in Diabetic Mice," *The American Journal of Pathology* 159:513-525 (2001).
- Greene et al., *Protective Groups in Organic Synthesis*, 2nd Ed., John Wiley & Sons, Inc., New York, Chapter 7, pp. 222-287 (1991).
- Groutas et al., "Synthesis and Pharmacological Studies of N-[4[2-Hydroxy-3[[2-4-(1H-imidazol-1-yl)phenoxy]ethyl]amino]propoxy]phenyl]methanesulfonamide, a Novel Antiarrhythmic Agent with Class II and Class III Activities," *J. Med. Chem.* 33:1087-1090 (1990).
- Gualtieri et al., "Antiviral Agents. 2. Analogs of 2-(alpha-Hydroxybenzyl)benzimidazole," *Journal of Medicinal Chemistry* 15(4):420-422 (1972).
- Haass et al., "Cellular Processing of beta-Amyloid Precursor Protein and the Genesis of Amyloid beta-Peptide," *Cell* 75:1039-1042 (1993).
- Hambly et al., "Reappraisal of the role of the diabetic state in coronary artery disease," *Chest* 70(2):251-257 (1976).
- Hammes et al., "Diabetic retinopathy risk correlates with intracellular concentrations of the glycoxidation product N(epsilon)-(carboxymethyl) lysine independently of glycohaemoglobin concentrations," *Diabetologia* 42:603-607 (1999).
- Heinze et al., "Synthesis of tetraarylimidazoles and pentaarylimidazolium salts," *Chemische Berichte* 101(10):3504-3516 (1968).
- Hofmann et al., "RAGE Mediates a Novel Proinflammatory Axis: A Central Cell Surface Receptor for S100/Calgranulin Polypeptides," *Cell* 97:889-901 (1999).
- Hori et al., "The Receptor for Advanced Glycation End Products (RAGE) Is a Cellular Binding Site for Amphotericin," *The Journal of Biological Chemistry* 270(43):25752-25761 (1995).
- Huttunen et al., "Receptor for Advanced Glycation End Products (RAGE)-mediated Neurite Outgrowth and Activation of NF-kB Require the Cytoplasmic Domain of the Receptor but Different Downstream Signaling Pathways," *The Journal of Biological Chemistry* 274(28):19919-19924 (1999).
- International Preliminary Report on Patentability for related International Application No. PCT/2010/049934 mailed Apr. 12, 2012.
- International Search Report and Written Opinion for related International Application No. PCT/US2010/049934 mailed Nov. 10, 2010.
- Johnson et al., "MDL 29311: Antioxidant With Marked Lipid- and Glucose-Lowering Activity in Diabetic Rats and Mice," *Diabetes* 42:1179-1186 (1993).
- Kamboh, "Molecular Genetics of Late-Onset Alzheimer's Disease," *Annals of Human Genetics* 98:381-404 (2004).
- Kannel et al., "Diabetes and Cardiovascular Disease: The Framingham Study," *JAMA* 241(19):2035-2038 (1979).
- Kannel et al., "Diabetics and Glucose Tolerance as Risk Factors for Cardiovascular Disease: The Framingham Study," *Diabetics Care* 2(2):120-126 (1979).
- Kennedy et al., "Familial Alzheimer's disease," *Brain* 116:309-324 (1993).
- Kislinger et al., "Receptor for Advanced Glycation End Products Mediates Inflammation and Enhanced Expression of Tissue Factor in Vasculature of Diabetic Apolipoprotein E-Null Mice," *Arterioscler Thromb Vasc Biol.* 21:905-910 (2001).
- Kumar et al., "RAGE at the Blood-Brain Barrier Mediates Neurovascular Dysfunction Caused by Amyloid-beta1-40 Peptide," *Neurosci. Program*, p. 414-#275.19 (2000).

(56)

References Cited

OTHER PUBLICATIONS

- Lampe et al., "Cardiotonic Agents. 6. Histamine Analogues as Potential Cardiovascular Selective H₂ Agonists," *Journal of Medicinal Chemistry* 33(6):1688-1697 (1990).
- Lander et al., "Activation of the Receptor for Advanced Glycation End Products Triggers a p21(ras)-dependent Mitogen-activated Protein Kinase Pathway Regulated by Oxidant Stress," *The Journal of Biological Chemistry* 272(28):17810-17814 (1997).
- Leder et al., "v-Ha-ras transgene abrogates the initiation step in mouse skin tumorigenesis: Effects of phorbol esters and retinoic acid," *Proc. Natl. Acad. Sci. USA* 87:9178-9182 (1990).
- Levy-Lahad et al., "Candidate Gene for the Chromosome 1 Familial Alzheimer's Disease Locus," *Science, New Series* 269(5226):973-977 (1995).
- Li et al., "Characterization and Functional Analysis of the Promoter of RAGE, the Receptor for Advanced Glycation End Products," *The Journal of Biological Chemistry* 272(26):16498-16506 (1997).
- Li et al., "Sp1-binding Elements in the Promoter of RAGE Are Essential for Amphoterin-mediated Gene Expression in Cultured Neuroblastoma Cells," *The Journal of Biological Chemistry* 273:30870-30878 (1998).
- Lugering et al., "The myleloic related protein MRP8/14 (27E10 antigen)—usefulness as a potential marker for disease activity in ulcerative colitis and putative biological function," *European Journal of Clinical Investigation* 25:659-664 (1995).
- Mackic et al., "Human Blood-Brain Barrier Receptors for Alzheimer's Amyloid-beta 1-40: Asymmetrical Binding, Endocytosis, and Transcytosis at the Apical Side of Brain Microvascular Endothelial Cell Monolayer," *J. Clin. Invest.* 102(4):734-743 (1998).
- McKhann et al., "Clinical diagnosis of Alzheimer's disease: Report of the NINCDS-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease," *Neurology* 34:939-944 (1984).
- Miyata et al., "Beta2-Microglobulin Modified with Advanced Glycation End Products Is a Major Component of Hemodialysis-associated Amyloidosis," *J. Clin. Invest.* 92:1243-1252 (1993).
- Miyata et al., "The Receptor for Advanced Glycation End Products (RAGE) Is a Central Mediator of the Interaction of AGE-beta2Microglobulin with Human Mononuclear Phagocytes Via an Oxidant-sensitive Pathway," *J. Clin. Invest.* 98(5):1088-1094 (1996).
- Morcos et al., "Activation of Tubular Epithelial Cells in Diabetic Nephropathy," *Diabetes* 51:3532-3544 (2002).
- Morris et al., "Place navigation impaired in rats with hippocampal lesions," *Nature* 297:681-683 (1982).
- Needleman et al., "A General Method Applicable to the Search for Similarities in the Amino Acid Sequence of Two Proteins," *J. Mol. Biol.* 48:443-453 (1970).
- Neeper et al., "Cloning and Expression of a Cell Surface Receptor for Advanced Glycosylation End Products of Proteins," *The Journal of Biological Chemistry* 267(21):14998-15004 (1992).
- Ohkubo et al., "Studies on Cerebral Protective Agents. VII. Synthesis of Novel 4-Arylazole Derivatives with Anti-anoxic Activity," *Chem. Pharm. Bull.* 43(6):947-954 (1995).
- Oldfield et al., "Advanced glycation end products cause epithelial-myofibroblast transdifferentiation via the receptor for advanced glycation end products (RAGE)," *The Journal of Clinical Investigation* 108(12):1853-1863 (2001).
- Pappolla et al., "The Heat Shock/Oxidative Stress Connection: Relevance to Alzheimer Disease," *Molecular and Chemical Neuropathology* 28:21-24 (1996).
- Park et al., "Suppression of accelerated diabetic atherosclerosis by the soluble receptor for advanced glycation endproducts," *Nature Medicine* 4(9):1025-1031 (1998).
- Parkkinen et al., "Amphoterin, the 30-kDa Protein in a Family of HMG1-type Polypeptides, Enhanced Expression in Transformed Cells, Leading Edge Localization, and Interactions with Plasminogen Activation," *The Journal of Biological Chemistry* 268(26):19726-19738 (1993).
- Pastor et al., "Molecular Genetics of Alzheimer's Disease," *Current Psychiatry Reports* 6:125-133 (2004).
- Pearson et al., "Improved tools for biological sequence comparison," *Proc. Natl. Acad. Sci. USA* 85:2444-2448 (1988).
- Penning et al., "Structure-Activity Relationship Studies on 1-[2-(4-Phenylphenoxy)ethyl]pyrrolidine (SC-22716), a Potent Inhibitor of Leukotriene A₄ (LTA₄) Hydrolase," *J. Med. Chem.* 43:721-735 (2000).
- Pike et al., "Neurodegeneration Induced by beta-Amyloid Peptides in vitro: The Role of Peptide Assembly State," *The Journal of Neuroscience* 13(4):1676-1687 (1993).
- Porretta et al., "Chemotherapeutic agents with an imidazole moiety. III. Synthesis and microbiological activity of new 1,4-diaryl-imidazole and 1,4-pyrrolimidazolephenylene derivatives," *II Farmaco* 46(7,8):913-924 (1991).
- Pyorala et al., "Diabetes and Atherosclerosis: An Epidemiologic View," *Diabetes/Metabolism Reviews* 3(2):463-524 (1987).
- Rammes et al., "Myeloid-related Protein (MRP) 8 and MRP14, Calcium-binding Proteins of the S100 Family, Are Secreted by Activated Monocytes via a Novel, Tubulin-dependent Pathway," *The Journal of Biological Chemistry* 272(14):9496-9502 (1997).
- Ranginwala et al., "Clinical Criteria for the Diagnosis of Alzheimer Disease: Still Good After All These Years," *Am. J. Geriatr. Psychiatry* 16(5):384-388 (2008).
- Rauvala et al., "Isolation and Some Characteristics of an Adhesive Factor of Brain That Enhances Neurite Outgrowth in Central Neurons," *The Journal of Biological Chemistry* 262(34):16625-16635 (1987).
- Reddy et al., "N(epsilon)-(Carboxymethyl)lysine Is a Dominant Advanced Glycation End Product (AGE) Antigen in Tissue Proteins," *Biochemistry* 34:10872-10878 (1995).
- Ritthaler et al., "Expression of Receptors for Advanced Glycation End Products in Peripheral Occlusive Vascular Disease," *American Journal of Pathology* 146(3):688-694 (1995).
- Robertson et al., "Atherosclerosis in Persons with Hypertension and Diabetes Mellitus," *Laboratory Investigation* 18(5):538-551 (1968).
- Rogaev et al., "Familial Alzheimer's disease in kindreds with missense mutations in a gene on chromosome 1 related to the Alzheimer's disease type 3 gene," *Nature* 376:775-778 (1995).
- Sanfilippo et al., "Synthesis of (Aryloxy)alkylamines. 1. Novel Antisecretory Agents with H⁺K⁺-ATPase Inhibitory Activity," *J. Med. Chem.* 31:1778-1785 (1988).
- Sanfilippo et al., "Synthesis of (Aryloxy)alkylamines. 2. Novel Imidazo-fused Heterocycles with Calcium Channel Blocking and Local Anesthetic Activity," *J. Med. Chem.* 31(11):2221-2227 (1988).
- Schafer et al., "The S100 family of EF-hand calcium-binding proteins: functions and pathology," *TIBS* 21:134-140 (1996).
- Schleicher et al., "Increased Accumulation of the Glycoxidation Product N(epsilon)-(carboxymethyl)lysine in Human Tissues in Diabetes and Aging," *J. Clin. Invest.* 99(3):457-468 (1997).
- Schmidt et al., "Advanced Glycation Endproducts Interacting with Their Endothelial Receptor Induce Expression of Vascular Cell Adhesion Molecule-1(VCAM-1) in Cultured Human Endothelial Cells and in Mice," *J. Clin. Invest.* 96:1395-1403 (1995).
- Schmidt et al., "Isolation and Characterization of Two Binding Proteins for Advanced Glycosylation End Products from Bovine Lung Which Are Present on the Endothelial Cell Surface," *The Journal of Biological Chemistry* 267(21):14987-14977 (1992).
- Schmidt et al., "Receptor for advanced glycation end products (AGEs) has a central role in vessel wall interactions and gene activation in response to circulating AGE proteins," *Proc. Natl. Acad. Sci. USA* 91:8807-8811 (1994).
- Schmidt et al., "The dark side of glucose," *Nature Medicine* 1(10):1002-1004 (1995).
- Schmidt et al., "The role of RAGE in amyloid-beta peptide-mediated pathology in Alzheimer's disease," *Current Opinion in Investigational Drugs* 10(7):672-680 (2009).
- Schmidt et al., "The V-Domain of Receptor for Advanced Glycation Endproducts (RAGE) Mediates Binding of AGEs: A Novel Target for Therapy of Diabetic Complications," *Supplement to Circulation* 96(8):Abstract No. 194 (1997).

(56)

References Cited

OTHER PUBLICATIONS

- Scozzafava et al., "Carbonic anhydrase activators—Part 21. Novel activators of isozymes I, II and IV incorporating carboxamido and ureido histamines moieties," *Eur. J. Med. Chem.* 35:31-39 (2000).
- Selkoe, "Normal and Abnormal Biology of the beta-Amyloid Precursor Protein," *Annual Review of Neuroscience* 17:489-517 (1994).
- Selkoe, "The Molecular Pathology of Alzheimer's Disease," *Neuron* 6:487-498 (1991).
- Selkoe, "Translating cell biology into therapeutic advances in Alzheimer's disease," *Nature* 399:A23-31 (1999).
- Semprini et al., "Evidence for differential S100 gene over-expression in psoriatic patients from genetically heterogeneous pedigrees," *Hum. Genet.* 111:310-313 (2002).
- Sherrington et al., "Cloning of a gene bearing missense mutations in early-onset familial Alzheimer's disease," *Nature* 375:754-760 (1995).
- Smith et al., "Comparison of Biosequences," *Advances in Applied Mathematics* 2:482-489 (1981).
- Snowdon, "Healthy Aging and Dementia: Findings from the Nun Study," *Annals of Internal Medicine* 139(5):450-454 (2003).
- Sousa et al., "Interaction of the Receptor for Advanced Glycation End Products (RAGE) with Transthyretin Triggers Nuclear Transcription Factor κ B (NF- κ B) Activation," *Laboratory Investigation* 80(7):1101-1110 (2000).
- Strittmatter et al., "Apolipoprotein E: Highly-avidity binding to beta-amyloid and increased frequency of type 4 allele in late-onset familial Alzheimer disease," *Proc. Natl. Acad. Sci. USA* 90:1977-1981 (1993).
- Taguchi et al., "Blockade of RAGE-amphoterin signalling suppresses tumor growth and metastases," *Nature* 405:354-360 (2000).
- Tanaka et al., "The Receptor for Advanced Glycation End Products Is Induced by the Glycation Products Themselves and Tumor Necrosis Factor- α through Nuclear Factor- κ B, and by 17 β -Estradiol through Sp-1 in Human Vascular Endothelial Cells," *The Journal of Biological Chemistry* 275(33):25781-25790 (2000).
- Teillet et al., "Food Restriction Prevents Advanced Glycation End Product Accumulation and Retards Kidney Aging in Lean Rats," *J. Am. Soc. Nephrol.* 11:1488-1497 (2000).
- Varney et al., "Crystal-Structure-Based Design and Synthesis of Novel C-Terminal Inhibitors of HIV Protease," *Journal of Medicinal Chemistry* 37(15):2274-2284 (1994).
- Vlassara, "Advanced Glycation End-products and Atherosclerosis," *Annals of Medicine* 28:419-426 (1996).
- Waller et al., "Status of the Coronary Arteries at Necropsy in Diabetes Mellitus with Onset After Age 30 Years: Analysis of 229 Diabetic Patients With and Without Clinical Evidence of Coronary Heart Disease and Comparison to 183 Control Subjects," *The American Journal of Medicine* 69:498-506 (1980).
- Wang et al., "The Polypeptide of Soluble Amyloid beta Protein in Cultured Cell Media: Detection and Quantification of Amyloid beta Protein and Variants by Immunoprecipitation-Mass Spectrometry," *The Journal of Biological Chemistry* 271(50):31894-31902 (1996).
- Wautier et al., "Advanced glycation end products (AGEs) on the surface of diabetic erythrocytes bind to the vessel wall via a specific receptor inducing oxidant stress in the vasculature: A link between surface-associated AGEs and diabetic complications," *Proc. Natl. Acad. Sci. USA* 91:7742-7746 (1994).
- Wautier et al., "Receptor-mediated Endothelial Cell Dysfunction in Diabetic Vasculopathy: Soluble Receptor for Advanced Glycation End Products Blocks Hyperpermeability in Diabetic Rats," *J. Clin. Invest.* 97(1):238-243 (1996).
- Wisniewski et al., "Apolipoprotein E: a pathological chaperone protein in patients with cerebral and systemic amyloid," *Neuroscience Letters* 135:235-238 (1992).
- Yan et al., "Amyloid-beta peptide-Receptor for Advanced Glycation Endproduct interaction elicits neuronal expression of macrophage-colony stimulating factor: A proinflammatory pathway in Alzheimer disease," *Proc. Natl. Acad. Sci. USA* 94:5296-5301 (1997).
- Yan et al., "An intracellular protein that binds amyloid-beta peptide and mediates neurotoxicity in Alzheimer's disease," *Nature* 389:689-695 (1997).
- Yan et al., "Enhanced Cellular Oxidant Stress by the Interaction of Advanced Glycation End Products with Their Receptors/Binding Proteins," *The Journal of Biological Chemistry* 269(13):9889-9897 (1994).
- Yan et al., "RAGE and Alzheimer's Disease: A Progression Factor for Amyloid-beta-Induced Cellular Perturbation?" *Journal of Alzheimer's Disease* 16:833-843 (2009).
- Yan et al., "RAGE and amyloid-beta peptide neurotoxicity in Alzheimer's disease," *Nature* 382:685-691 (1996).
- Yan et al., "Receptor-dependent cell stress and amyloid accumulation in systemic amyloidosis," *Nature* 6(6):643-651 (2000).
- Yankner et al., "Neurotrophic and Neurotoxic Effects of Amyloid beta Protein: Reversal by Tachykinin Neuropeptides," *Science, New Series* 250(4978):279-282 (1990).
- Yeh et al., "Requirement for p38 and p44/p42 Mitogen-Activated Protein Kinases in RAGE-Mediated Nuclear Factor- κ B Transcriptional Activation and Cytokine Secretion," *Diabetes* 50:1495-1504 (2001).
- Zimmer et al., "The S100 Protein Family: History, Function, and Expression," *Brain Research Bulletin* 37(4):417-429 (1995).
- Davis, et al., "RAGE Deletion Increases Anti-Oxidant and Anti-Inflammatory Biochemical Profiles in Human APP Transgenic Mice," Poster presented at Alzheimer's Association International Conference, Washington, DC, Jul. 20, 2015.
- Sabbagh, M., "Evaluation of Phase 2b Safety of Azeliragon (TTP488)" presented at Clinical Trials on Alzheimer's Disease program, Barcelona, Spain, Nov. 6, 2015.
- Amendment No. 6 to Form S-1 Registration Statement for vTv Therapeutics Inc., Jul. 24, 2015. pp. 1-2, 83, 86-94.
- Investor Presentation—Jul. 2015. Slides 9-18.
- Aricept® package insert, Feb. 2012.
- Barile et al., "The RAGE Axis in Early Diabetic Retinopathy," *Investigative Ophthalmology & Visual Science* 46(8):2916-2924 (2005).
- Bishop et al., "Neural Mechanisms of ageing and cognitive decline," *Nature* 464:529-535 (2010).
- Bonetta, "Door Slams on RAGE," *Alzheimer Research Forum Print News*, Nov. 9, 2011.
- Burstein A, et al. "Azeliragon Phase 2b Survival Analysis Supports Beneficial Effects on Delaying Time to Cognitive Deterioration in Patients with Mild Alzheimer's Disease." Poster Presented at the Alzheimer's Association International Conference. Jul. 27, 2016. Toronto, Canada.
- Burstein et al. "Evaluation of the relationship between TTP488 plasma concentration and changes in ADAS-cog relative to placebo." Poster session presented at: the Alzheimer's Association International Conference, Jul. 13-18, 2013, Boston, Massachusetts.
- Burstein et al., "Effect of TTP488 in patients with mild to moderate Alzheimer's disease," *BMC Neurology* 14:12 (2014), 19 pages.
- Donahue et al., "RAGE, LRP-1, and amyloid-beta protein in Alzheimer's disease," *Acta Neuropathol.* 112:405-415 (2006).
- G. Basta et al., *63 Cardiovascular Research* 582-592 (2004).
- G.P. Sims et al., *28 Annual Review of Immunology*, 367-368 (2010).
- Galasko et al., "A clinical trial of an inhibitor of RAGE-A-beta interactions in Alzheimer's disease," *RI clinical trial manuscript*, Aug. 8, 2012.
- Galasko et al., "A Randomized Clinical Trial of an inhibitor of RAGE-A-beta interactions in patients with mild to moderate AD," DRAFT of presentation in Clinical Trials on Alzheimer's Disease program, San Diego, California, Nov. 3, 2011.
- Galasko et al., "Clinical trial of an inhibitor of RAGE-A-beta interactions in Alzheimer disease," *Neurology* 82:1537-1542 (2014).
- Galasko et al., Supplements 1-6 to "Clinical trial of an inhibitor of RAGE-A-beta interactions in Alzheimer disease," *Neurology* 82:1537-1542 (2014).
- International Search Report and Written Opinion for related International Application No. PCT/US2013/062964, mailed Nov. 19, 2013.

(56)

References Cited

OTHER PUBLICATIONS

Kostura et al. Efficacy of RAGE antagonist in murine model of Alzheimer's disease. Poster session presented at: the Alzheimer's Association International Congress; Jul. 13-18, 2014; Copenhagen, Denmark.

Namenda® package insert, 2007, Jan. 2011.

Perrone et al., "The Complexity of Sporadic Alzheimer's Disease Pathogenesis: The Role of RAGE as Therapeutic Target to Promote Neuroprotection by Inhibiting Neurovascular Dysfunction," International Journal of Alzheimer's Disease, vol. 2012, 13 pages.

R. Ramasamy (Yan) et al., 15 Glycobiology 16R-18R (2005).

Sabbagh et al., "PF-04494700, an Oral Inhibitor of Receptor for Advanced Glycation End Products (RAGE), in Alzheimer Disease," Alzheimer Disease & Associated Disorders 25(3):206-212 (2011).

Spite et al., "Novel Lipid Mediators Promote Resolution of Acute Inflammation: Impact of Aspirin and Statins," Circulation Research, 107:1170-1184 (2010).

T. Wendt et al., 185 Atherosclerosis 70-77 (2006).

Takuma et al., "RAGE-mediated signaling contributes to intraneuronal transport of amyloid-beta and neuronal dysfunction," PNAS 106(47):20021-20026 (2009).

Thompson, A. J. et al., "Protein Conformational Misfolding and Amyloid Formation: Characteristics of a New Class of Disorders that Include Alzheimer's and Prion Diseases," Current Medicinal Chemistry, 9:1751-1762 (2002).

Vellas, et al., "Long-term changes in ADAS-cog: What is clinically relevant for disease modifying trials in Alzheimer?" (vol. 11, No. 4, 2007; Journal of Nutrition, Health & Aging).

1

**SUBSTITUTED IMIDAZOLE DERIVATIVES
AND METHODS OF USE THEREOF**

FIELD OF THE INVENTION

This invention relates to compounds which are inhibitors of the interaction between the receptor for advanced glycation endproducts (RAGE) and its physiological ligands such as advanced glycated end products (AGEs), S100/calgranulin/EN-RAGE, β -amyloid, and amphoterin, for the treatment of RAGE mediated diseases.

BACKGROUND OF THE INVENTION

The Receptor for Advanced Glycated Endproducts (RAGE) is a member of the immunoglobulin super family of cell surface molecules. The extracellular (N-terminal) domain of RAGE includes three immunoglobulin-type regions, one V (variable) type domain followed by two C-type (constant) domains (Neeper et al., *J. Biol. Chem.* 267:14998-15004 (1992)). A single transmembrane spanning domain and a short, highly charged cytosolic tail follow the extracellular domain. The N-terminal, extracellular domain can be isolated by proteolysis of RAGE to generate soluble RAGE (sRAGE) comprised of the V and C domains.

RAGE is expressed in most tissues, and in particular, is found in cortical neurons during embryogenesis (Hori et al. (1995)). Increased levels of RAGE are also found in aging tissues (Schleicher et al., *J. Clin. Invest.* 99 (3): 457-468 (1997)), and the diabetic retina, vasculature and kidney (Schmidt et al., *Nature Med.* 1:1002-1004 (1995)). Activation of RAGE in different tissues and organs leads to a number of pathophysiological consequences. RAGE has been implicated in a variety of conditions including: acute and chronic inflammation (Hofmann et al., *Cell* 97:889-901 (1999)), the development of diabetic late complications such as increased vascular permeability (Wautier et al., *J. Clin. Invest.* 97:238-243 (1996)), nephropathy (Teillet et al., *J. Am. Soc. Nephrol.* 11:1488-1497 (2000)), atherosclerosis (Vlassara et al., *The Finnish Medical Society DUODECIM, Ann. Med.* 28:419-426 (1996)), and retinopathy (Hammes et al., *Diabetologia* 42:603-607 (1999)). RAGE has also been implicated in Alzheimer's disease (Yan et al., *Nature* 382: 685-691 (1996)), erectile dysfunction, and in tumor invasion and metastasis (Taguchi et al., *Nature* 405: 354-357 (2000)).

Advanced glycation endproducts (AGEs) have been implicated in a variety of disorders including complications associated with diabetes and normal aging. Incubation of proteins or lipids with aldose sugars results in nonenzymatic glycation and oxidation of amino groups on proteins to form Amadori adducts. Over time, the adducts undergo additional rearrangements, dehydrations, and cross-linking with other proteins to form complexes known as AGEs. Factors which promote formation of AGEs include delayed protein turnover (e.g. as in amyloidoses), accumulation of macromolecules having high lysine content, and high blood glucose levels (e.g. as in diabetes) (Hori et al., *J. Biol. Chem.* 270: 25752-761, (1995)).

AGEs display specific and saturable binding to cell surface receptors on endothelial cells of the microvasculature, monocytes and macrophages, smooth muscle cells, mesangial cells, and neurons.

In addition to AGEs, other compounds can bind to, and inhibit the interaction of physiological ligands with RAGE. In normal development, RAGE interacts with amphoterin, a polypeptide which mediates neurite outgrowth in cultured embryonic neurons (Hon et al., (1995)). RAGE has also

2

been shown to interact with EN-RAGE, a protein having substantial similarity to calgranulin (Hofmann et al. (1999)). RAGE has also been shown to interact with β -amyloid (Yan et al., *Nature* 389:689-695 (1997); Yan et al., *Nature* 382: 685-691 (1996); Yan et al., *Proc. Natl. Acad. Sci.*, 94:5296-5301 (1997)).


Binding of ligands such as AGEs, S100/calgranulin/EN-RAGE, β -amyloid, CML (N ϵ -Carboxymethyl lysine), and amphoterin to RAGE has been shown to modify expression of a variety of genes. For example, in many cell types interaction between RAGE and its ligands generates oxidative stress, which thereby results in activation of the free radical sensitive transcription factor NF- κ B, and the activation of NF- κ B regulated genes, such as the cytokines IL-1 β , TNF- α , and the like.

In addition, several other regulatory pathways, such as those involving p21 ras, MAP kinases, ERK1 and ERK2, have been shown to be activated by binding of AGEs and other ligands to RAGE. In fact, transcription of RAGE itself is regulated at least in part by NF- κ B. Thus, an ascending, and often detrimental, spiral is fueled by a positive feedback loop initiated by ligand binding. Inhibiting binding of physiological ligands to RAGE provides for the down-regulation of the pathophysiological changes brought about by excessive concentrations of AGEs and other ligands for RAGE as described above.

Thus, there is a need for the development of compounds that inhibit the binding of physiological ligands to RAGE.

SUMMARY OF THE INVENTION

The present invention relates to compounds of Formula (I):



or pharmaceutically acceptable salts thereof, wherein R¹ and R² are independently selected from the group consisting of —CH₃, —CH₂CH₃, —CH(CH₃)₂, and —CH₂CH₂CH₃; and Q¹ is selected from the group consisting of —CH₂OCH₂CH₃ and —CH₂CH₂CH₂CH₃.

This invention also provides for methods of preparation of compounds of Formula (I) or pharmaceutically acceptable salts thereof, pharmaceutical compositions comprising compounds of Formula (I) or pharmaceutically acceptable salts thereof; and methods for the use of compounds of Formula (I) or pharmaceutically acceptable salts thereof in treating diseases mediated by RAGE.

Compounds of Formula (I) or pharmaceutically acceptable salts thereof are useful as inhibitors of the interaction of the receptor for advanced glycation endproducts (RAGE) with ligands such as advanced glycated end products (AGEs), S100/calgranulin/EN-RAGE, β -amyloid, and amphoterin. The compounds are also useful in treating a variety of diseases or conditions in humans that may be responsive to the inhibition of RAGE. Such diseases or

3

conditions include, but are not limited to, acute and chronic inflammation, the development of diabetic late complications such as increased vascular permeability, nephropathy, atherosclerosis, and retinopathy, the development of Alzheimer's disease and related disorders, erectile dysfunction, tumor invasion and metastasis, and osteoporosis.

The scope of the present invention includes combinations of the various aspects, embodiments, and preferences as herein described.

BRIEF DESCRIPTION OF DRAWINGS


Not applicable

DETAILED DESCRIPTION OF THE INVENTION

The following definitions are meant to clarify, but not limit, the terms defined. If a particular term used herein is not specifically defined, such term should not be considered indefinite. Rather, such terms are used within their plain and ordinary meanings.

As used herein, the various functional groups represented will be understood to have a point of attachment at the functional group having the hyphen. In other words, in the case of $-\text{CH}_2\text{CH}_2\text{CH}_3$, it will be understood that the point of attachment is the CH_2 group at the far left.

In a first embodiment, the present invention includes a compound of Formula (I):



or a pharmaceutically acceptable salt thereof, wherein

R^1 and R^2 are independently selected from the group consisting of $-\text{CH}_3$, $-\text{CH}_2\text{CH}_3$, $-\text{CH}(\text{CH}_3)_2$, and $-\text{CH}_2\text{CH}_2\text{CH}_3$; and

Q^1 is selected from the group consisting of $-\text{CH}_2\text{OCH}_2\text{CH}_3$ and $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$.

In a second embodiment, the present invention provides a compound of Formula (I) or a pharmaceutically acceptable salt thereof wherein R^1 is $-\text{CH}_3$.

In a third embodiment, the present invention provides a compound of Formula (I) or a pharmaceutically acceptable salt thereof wherein R^1 is $-\text{CH}_2\text{CH}_3$.

In a fourth embodiment, the present invention provides a compound of Formula (I) or a pharmaceutically acceptable salt thereof according to any one of the previous embodiments wherein R^2 is $-\text{CH}_3$.

In a fifth embodiment, the present invention provides a compound of Formula (I) or a pharmaceutically acceptable salt thereof according to any one of the first to third embodiments wherein R^2 is $-\text{CH}_2\text{CH}_3$.

In a sixth embodiment, the present invention provides a compound of Formula (I) or a pharmaceutically acceptable salt thereof according to any one of the previous embodiments wherein Q^1 is $-\text{CH}_2\text{OCH}_2\text{CH}_3$.

4

In a seventh embodiment, the present invention provides a compound of Formula (I) or a pharmaceutically acceptable salt thereof according to any one of the first to fifth embodiments wherein Q^1 is $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$.

In an eighth embodiment, the present invention provides a compound of Formula (I) or a pharmaceutically acceptable salt thereof according to any one of the first to seventh embodiments wherein the compound is a free amine.

In a ninth embodiment, the present invention provides a compound of Formula (I) or a pharmaceutically acceptable salt thereof according to any one of the first to seventh embodiments wherein the compound is a pharmaceutically acceptable salt.

In a tenth embodiment, the present invention provides a compound of Formula (I) or a pharmaceutically acceptable salt thereof according to any one of the first to seventh embodiments wherein the compound is a hydrochloride salt.

In an eleventh embodiment, the present invention provides a compound of Formula (I) or a pharmaceutically acceptable salt thereof according to any one of the first to tenth embodiments wherein the group $-\text{CH}_2\text{CH}(\text{OH})\text{CH}_2\text{NR}^1\text{R}^2$ is in the S configuration.

In a twelfth embodiment, the present invention provides a compound of Formula (I) or a pharmaceutically acceptable salt thereof according to any one of the first to tenth embodiments wherein the group $-\text{CH}_2-\text{CH}(\text{OH})-\text{CH}_2-\text{NR}^1\text{R}^2$ is in the R configuration.

Specific embodiments of the compound of Formula (I) or a pharmaceutically acceptable salt thereof include:

- (R)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-dimethylamino-propan-2-ol;
- (R)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-diethylamino-propan-2-ol;
- (S)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-dimethylamino-propan-2-ol;
- (S)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-diethylamino-propan-2-ol;
- (R)-1-(4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-dimethylamino-propan-2-ol;
- (S)-1-(4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-dimethylamino-propan-2-ol;
- (R)-1-(4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-diethylamino-propan-2-ol; and
- (S)-1-(4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-diethylamino-propan-2-ol;

or a pharmaceutically acceptable salt thereof.

Another aspect of the present invention includes a pharmaceutical composition comprising a compound of Formula (I) or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.

One aspect of the present invention includes a method for treating a RAGE-mediated disease comprising administering to a subject a compound of Formula (I) or a pharmaceutically acceptable salt thereof. Another aspect includes use of a compound of Formula (I) or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for treating a RAGE-mediated disease. A still further aspect includes a compound of Formula (I) or a pharmaceutically acceptable salt thereof for use in the treatment of a RAGE-mediated disease. In one embodiment, the disease is Alzheimer's Disease. In one embodiment, such treatment modifies the presentation of Alzheimer's Disease. In another

embodiment, such treatment improves cognitive performance of a subject suffering from mild to moderate Alzheimer's Disease.

Pharmaceutically acceptable salts of the compounds of the present invention are also included within the scope of the invention. The term "pharmaceutically acceptable salt(s)" as used herein refers to non-toxic salts of a compound of Formula (I) which are generally prepared by reacting the free base (i.e. free amine) of the compound of Formula (I) with a suitable organic or inorganic acid such as, but not limited to, hydrochloride, hydrobromide, phosphate, sulfate, trifluoroacetate, trichloroacetate, acetate, oxalate, maleate, pyruvate, malonate, succinate, citrate, tartrate, fumarate, mandelate, benzoate, cinnamate, methiodide, methbromide, methchloride, methanesulfonate, ethanesulfonate, picrate and the like, and include acids related to the pharmaceutically-acceptable salts listed in the Journal of Pharmaceutical Science, 66, 2 (1977) p. 1-19. Other salts which are not pharmaceutically acceptable may be useful in the preparation of compounds of the invention and these form a further aspect of the invention.

Unless otherwise stated, structures depicted herein are also meant to include compounds which differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structure except for the replacement of a hydrogen atom by a deuterium or tritium, or the replacement of a carbon atom by a ^{13}C - or ^{14}C -enriched carbon are within the scope of the invention.

The compound of Formula (I) contains one chiral center. The scope of the present invention includes mixtures of stereoisomers as well as purified enantiomers or enantiomerically/diastereomerically enriched mixtures. Also included within the scope of the invention are the individual isomers of the compounds represented by the formulae of the present invention, as well as any wholly or partially equilibrated mixtures thereof. The present invention also includes any tautomers of the compounds represented by the formulas above.

Examples of compounds of Formula (I) or a pharmaceutically acceptable salt thereof having potentially useful biological activity are herein described. The ability of compounds of Formula (I) or pharmaceutically acceptable salts thereof to inhibit the interaction of RAGE with its physiological ligands was established with representative compounds of Formula (I) or a pharmaceutically acceptable salt thereof using the assay(s) described in the Examples section below.

The invention further provides pharmaceutical compositions comprising a compound of Formula (I) or a pharmaceutically acceptable salt thereof. The term "pharmaceutical composition" is used herein to denote a composition that may be administered to a mammalian host, e.g., orally, topically, parenterally, by inhalation spray, or rectally, in unit dosage formulations containing conventional non-toxic carriers, diluents, adjuvants, vehicles and the like. The term "parenteral" as used herein, includes subcutaneous injections, intravenous, intramuscular, intracisternal injection, or by infusion techniques.

The pharmaceutical compositions containing a compound of the invention may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous, or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any known method, and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents, and preserving

agents in order to provide pharmaceutically elegant and palatable preparations. Tablets may contain the active ingredient in admixture with non-toxic pharmaceutically-acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example corn starch or alginic acid; binding agents, for example, starch, gelatin or acacia; and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may also be coated by the techniques described in U.S. Pat. Nos. 4,356,108; 4,166,452; and 4,265,874, to form osmotic therapeutic tablets for controlled release.

Formulations for oral use may also be presented as hard gelatin capsules where the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or a soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions may contain the active compounds in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide such as lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example, heptadecaethyl-eneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as a liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active compound in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example, sweetening, flavoring, and coloring agents may also be present.

The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example, olive oil or arachis oil, or a mineral oil, for example a liquid paraffin, or a mixture

thereof. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavoring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents. The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to the known methods using suitable dispersing or wetting agents and suspending agents described above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conveniently employed as solvent or suspending medium. For this purpose, any bland fixed oil may be employed using synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

The compositions may also be in the form of suppositories for rectal administration of the compounds of the invention. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will thus melt in the rectum to release the drug. Such materials include cocoa butter and polyethylene glycols, for example.

For topical use, creams, ointments, jellies, solutions or suspensions, lotions, eye ointments and eye or ear drops, impregnated dressings and aerosols etc., containing the compounds of the invention are contemplated. These topical formulations may contain appropriate conventional additives such as preservatives, solvents to assist drug penetration and emollients in ointments and creams. The formulations may also contain compatible conventional carriers, such as cream or ointment bases and ethanol or oleyl alcohol for lotions. Such carriers may be present as from about 0.1% up to about 99% of the formulation. More usually they will form up to about 80% of the formulation. For the purpose of this application, topical applications shall include mouth washes and gargles.

The compounds of the present invention may also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes may be formed from a variety of phospholipids.

The compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol, polyhydroxyethylaspartamidephenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polypsilone caprolactone, polyhydroxy butyric acid, poly-

orthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and cross-linked or amphipathic block copolymers of hydrogels.

For administration by inhalation the compounds according to the invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, tetrafluoroethane, heptafluoropropane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of e.g. gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of a compound of the invention and a suitable powder base such as lactose or starch.

Compounds that antagonize the interaction of RAGE with its physiological ligands are potentially useful in treating diseases or conditions that may be responsive to the inhibiting of the RAGE receptor. The present invention provides a method of treatment comprising: administering to a subject a compound of Formula (I) or a pharmaceutically acceptable salt thereof. In an embodiment of this aspect, the present invention provides a method for the inhibition of the interaction of RAGE with its physiological ligands. In another embodiment of this aspect, the present invention provides a method for treating a disease state selected from the group consisting of acute and chronic inflammation including skin inflammation such as psoriasis, atopic dermatitis, inflammation associated with organ, tissue, or cell transplantation, and lung inflammation including, asthma and chronic obstructive pulmonary disease, sepsis, diabetes, diabetes related complications, renal failure, hyperlipidemic atherosclerosis associated with diabetes, neuronal cytotoxicity, restenosis, Down's syndrome, dementia associated with head trauma, amyotrophic lateral sclerosis, multiple sclerosis, amyloidosis, an autoimmune disease, wound healing, periodontal disease, neuropathy, neuronal degeneration, vascular permeability, nephropathy, atherosclerosis, retinopathy, Alzheimer's disease, erectile dysfunction, tumor invasion and/or metastasis, and osteoporosis which comprises administering to a subject a therapeutically effective amount of a compound of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

I. RAGE and the Complications of Diabetes

As noted above, the compounds of the present invention are useful in the treatment of the complications of diabetes. It has been shown that nonenzymatic glycoxidation of macromolecules ultimately resulting in the formation of advanced glycation endproducts (AGEs) is enhanced at sites of inflammation, in renal failure, in the presence of hyperglycemia and other conditions associated with systemic or local oxidant stress (Dyer, D., et al., *J. Clin. Invest.*, 91:2463-2469 (1993); Reddy, S., et al., *Biochem.*, 34:10872-10878 (1995); Dyer, D., et al., *J. Biol. Chem.*, 266:11654-11660 (1991); Degenhardt, T., et al., *Cell Mol. Biol.*, 44:1139-1145 (1998)). Accumulation of AGEs in the vasculature can occur focally, as in the joint amyloid composed of AGE- β 2-microglobulin found in patients with dialysis-related amyloidosis (Miyata, T., et al., *J. Clin. Invest.*, 92:1243-1252 (1993); Miyata, T., et al., *J. Clin. Invest.*, 98:1088-1094 (1996)), or generally, as exemplified by the vasculature and tissues of patients with diabetes (Schmidt, A-M., et al., *Nature Med.*, 1:1002-1004 (1995)). The progressive accumulation of AGEs over time in patients with diabetes suggests that endogenous clearance mechanisms are not able to function effectively at sites of AGE deposi-

tion. Such accumulated AGEs have the capacity to alter cellular properties by a number of mechanisms. Although RAGE is expressed at low levels in normal tissues and vasculature, in an environment where the receptor's ligands accumulate, it has been shown that RAGE becomes upregulated (Li, J. et al., *J. Biol. Chem.*, 272:16498-16506 (1997); Li, J., et al., *J. Biol. Chem.*, 273:30870-30878 (1998); Tanaka, N., et al., *J. Biol. Chem.*, 275:25781-25790 (2000)). RAGE expression is increased in endothelium, smooth muscle cells and infiltrating mononuclear phagocytes in diabetic vasculature. Also, studies in cell culture have demonstrated that AGE-RAGE interaction caused changes in cellular properties important in vascular homeostasis.

II. RAGE and Cellular Dysfunction in the Amyloidoses

Also as noted above, the compounds of the present invention are useful in treating amyloidoses and/or Alzheimer's Disease. RAGE appears to be a cell surface receptor which binds β -sheet fibrillar material regardless of the composition of the subunits (amyloid- β peptide, A β , amylin, serum amyloid A, prion-derived peptide) (Yan, S.-D., et al., *Nature*, 382:685-691 (1996); Yan, S.-D., et al., *Nat. Med.*, 6:643-651 (2000)). Deposition of amyloid has been shown to result in enhanced expression of RAGE. For example, in the brains of patients with Alzheimer's disease (AD), RAGE expression increases in neurons and glia (Yan, S.-D., et al., *Nature* 382:685-691 (1996)). The consequences of A β interaction with RAGE appear to be quite different on neurons versus microglia. Whereas microglia become activated as a consequence of A β -RAGE interaction, as reflected by increased motility and expression of cytokines, early RAGE-mediated neuronal activation is superceded by cytotoxicity at later times. Further evidence of a role for RAGE in cellular interactions of A β concerns inhibition of A β -induced cerebral vasoconstriction and transfer of the peptide across the blood-brain barrier to brain parenchyma when the receptor was blocked (Kumar, S., et al., *Neurosci. Program*, p141 (2000)). Inhibition of RAGE-amyloid interaction has been shown to decrease expression of cellular RAGE and cell stress markers (as well as NF- κ B activation), and diminish amyloid deposition (Yan, S.-D., et al., *Nat. Med.*, 6:643-651 (2000)) suggesting a role for RAGE-amyloid interaction in both perturbation of cellular properties in an environment enriched for amyloid (even at early stages) as well as in amyloid accumulation.

In other studies using a mouse model of Alzheimer's Disease, it has been shown that RAGE antagonists can reverse the formation of plaques and the loss of cognition. In U.S. Patent Publication No. US 2005/0026811, small molecule RAGE antagonists were used to inhibit the progression of A β deposition and reduced the volume of pre-existing plaques in Alzheimer's Disease mice (US 2005/0026811 at ¶¶581-586). Furthermore, treatment with such small molecule RAGE antagonists improved cognition in these Alzheimer's Disease mouse models (US 2005/0026811 at ¶¶587-590). Thus, in a mouse model of Alzheimer's Disease, those mice who had developed A β plaques and cognitive loss and were treated with small molecule RAGE antagonists exhibited a reduction in plaque volume and an improvement in cognitive performance as compared to those Alzheimer's Disease mice who were not treated with the small molecule RAGE antagonists, showing that the RAGE antagonist compounds may delay or slow loss of cognitive performance, or may improve cognitive performance of a subject suffering from dementia of Alzheimer's type.

Also, it had been shown in both cellular assays and in animal studies that RAGE mediates the transcytosis of circulating A β across the blood-brain barrier (BBB). Such

increased transcytosis of A β results in neuronal oxidant stress and sustained reductions in cerebral blood flow. The effects of RAGE can be inhibited by a RAGE modulator (e.g., anti-RAGE antibody or sRAGE) (see e.g., Mackic et al., *J. Clin. Invest.*, 102:734-743 (1998); see also Kumar et al., *Neurosci., Program*, p 141 (2000)). These findings were confirmed by additional studies (see e.g., U.S. Pat. No. 6,825,164 at col. 17, line 48 to col. 18, line 43; Deane et al., *Nature Medicine*, 9:907-913 (2003)). Reduced cerebral perfusion can promote ischemic lesions which can act synergistically with A β to exacerbate dementia. Also, insufficient cerebral blood flow may alter A β trafficking across the blood brain barrier thereby reducing A β clearance and promoting accumulation of A β in brain (see Girouard and Iadecola, *J. Appl. Physiol.*, 100, 328-335 (2006) at page 332). Thus, the increase in cerebral blood flow promoted by RAGE antagonists may reduce the symptoms or delay onset of development of Alzheimer's Disease, or both. For example, RAGE antagonists may delay or slow loss of cognitive performance, or may improve cognitive performance of a subject suffering from dementia of Alzheimer's type, or both.

III. RAGE and Propagation of the Immune/Inflammatory Response

As noted above, the compounds of the present invention are useful in treating inflammation. For example, S100/calgranulins have been shown to comprise a family of closely related calcium-binding polypeptides characterized by two EF-hand regions linked by a connecting peptide (Schafer, B. et al., *TIBS*, 21:134-140 (1996); Zimmer, D., et al., *Brain Res. Bull.*, 37:417-429 (1995); Rammes, A., et al., *J. Biol. Chem.*, 272:9496-9502 (1997); Luger, N., et al., *Eur. J. Clin. Invest.*, 25:659-664 (1995)). Although they lack signal peptides, it has long been known that S100/calgranulins gain access to the extracellular space, especially at sites of chronic immune/inflammatory responses, as in cystic fibrosis and rheumatoid arthritis. RAGE is a receptor for many members of the S100/calgranulin family, mediating their proinflammatory effects on cells such as lymphocytes and mononuclear phagocytes. Also, studies on delayed-type hypersensitivity response, colitis in IL-10 null mice, collagen-induced arthritis, and experimental autoimmune encephalitis models suggest that RAGE-ligand interaction (presumably with S100/calgranulins) has a proximal role in the inflammatory cascade as implicated in the inflammatory diseases such as but not limited to rheumatoid arthritis and multiple sclerosis.

RAGE is also implicated in inflammatory diseases of the skin such as but not limited to atopic dermatitis, eczema, and psoriasis. Psoriasis in particular is characterized by inflamed itchy lesions. Psoriasis may be accompanied by arthropathic symptoms that are similar to those seen in rheumatoid arthritis. There is considerable evidence that psoriasis is a polygenic autoimmune disorder. Psoriatic lesions are rich in cytokines, in particular IL-1 and IL-8, both potent proinflammatory mediators. IL-8 in particular is a chemotactic factor for neutrophils; neutrophils are also known to synthesize and secrete S100 proteins, one of the ligands for RAGE which is implicated in propagation of the immune and inflammatory response. Psoriasin, (S100A7) a new member of the S100 gene family, is a secreted protein isolated from psoriatic skin. Semprini et al. (*Hum. Genet.* 2002 October, 111(4-5), 310-3) have shown a linkage of psoriasis genetic susceptibility to distinct overexpression of S100 proteins in skin. Therefore, a modulator of RAGE would be expected to regulate the immune response in psoriasis.

IV. RAGE and Amphoterin

As noted above, the compounds of the present invention are useful in treating tumor and tumor metastasis. For example, amphoterin is a high mobility group I nonhistone chromosomal DNA binding protein (Rauvala, H., et al., *J. Biol. Chem.*, 262:16625-16635 (1987); Parkikinen, J., et al., *J. Biol. Chem.* 268:19726-19738 (1993)) which has been shown to interact with RAGE. It has been shown that amphoterin promotes neurite outgrowth, as well as serving as a surface for assembly of protease complexes in the fibrinolytic system (also known to contribute to cell mobility). In addition, a local tumor growth inhibitory effect of blocking RAGE has been observed in a primary tumor model (C6 glioma), the Lewis lung metastasis model (Taguchi, A., et al., *Nature* 405:354-360 (2000)), and spontaneously arising papillomas in mice expressing the v-Ha-ras transgene (Leder, A., et al., *Proc. Natl. Acad. Sci.*, 87:9178-9182 (1990)).

V. RAGE and Respiratory Diseases

Airway inflammation is important in the pathogenesis of asthma. Such inflammation may give rise to significant exacerbations and increases in asthma severity, as well as to be a major factor in a decline in asthmatic status. In severe exacerbations of asthma there is an intense, mechanistically heterogeneous inflammatory response involving neutrophil and eosinophil accumulation and activation. Neutrophils are a significant source of S100 proteins, key ligands for RAGE implicated in the propagation of the immune response and inflammation. Therefore, modulators of RAGE would be expected to possess therapeutic value in the treatment of asthma. Further, the propagation step in the immune response in the lung driven by S100—RAGE interaction would be expected to lead to the activation and/or recruitment of inflammatory cells, such as neutrophils, which in chronic obstructive pulmonary diseases such as emphysema, are significant sources of damaging proteases. Therefore, a RAGE modulator would be expected possess potential in the treatment of chronic obstructive pulmonary diseases.

As used herein, the phrase “therapeutically effective amount” shall mean that amount of a drug or pharmaceutical agent that will elicit the therapeutic response of an subject that is being sought.

In these methods, factors which may influence what constitutes a therapeutically effective amount include, but are not limited to, the size and weight of the subject, the biodegradability of the therapeutic agent, the activity of the therapeutic agent, the size of the effected area, as well as its bioavailability. The phrase includes amounts which, as compared to a corresponding subject who has not received such amount, results in improved treatment, healing, or amelioration of a side effect, or a decrease in the rate of advancement of a disease or disorder.

In an embodiment, the present invention provides a method for treating restenosis comprising: administering to a subject a therapeutically effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof. In an embodiment, the subject is suffering from diabetes.

In an embodiment, the present invention provides a method for treating acute or chronic inflammation comprising: administering to a subject a therapeutically effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof.

In an embodiment, the present invention provides a method for treating dementia associated with head trauma comprising: administering to a subject a therapeutically effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof. In an embodiment, the

cognitive performance of the subject is improved. In another embodiment, the cognitive performance of the subject is maintained. In another embodiment, the rate of loss of cognitive performance of the subject is slowed.

In an embodiment, the present invention provides a method for treating Alzheimer’s Disease comprising: administering to a subject a therapeutically effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof. With respect to Alzheimer’s Disease, the present invention is believed useful in alteration the course of the underlying dementing process. Alzheimer’s Disease may be diagnosed by NINCDS and DSM criteria, Mini-Mental State Examination, and Clinical Dementia Rating within particular limits. One aspect of the present invention includes improving cognitive performance comprising administering a compound of Formula (I) or a pharmaceutically acceptable salt thereof. Cognitive performance may be assessed with the cognitive subscale of the Alzheimer’s Disease Assessment Scale (ADAS-cog), as is known in the art, which scores cognitive function on a 0 to 70 scale, with higher scores indicating greater cognitive impairment. Thus, a reduction in score demonstrates cognitive improvement. One aspect of the present invention includes administering to a subject a compound of Formula (I) or a pharmaceutically acceptable salt thereof to reduce an ADAS-cog score of a subject in need of such reduction. Such a subject may be a human be suffering from dementia of Alzheimer’s type, mild to moderate Alzheimer’s Diseases, or severe Alzheimer’s Disease.

In addition, the progression of Alzheimer’s Disease may also be assessed in a human through examination of four areas of function: General, Cognitive, Behavioral, and Activities of Daily Living. Such an assessment may be performed using a Clinician’s Interview Based Impression of Change (CIBIC or CIBIC plus). One aspect of the present invention includes improvement in subject’s function comprising administering a compound of Formula (I) or a pharmaceutically acceptable salt thereof. In one embodiment, the subject’s function is one or more of general, cognitive, behavioral, and activities of daily living.

In an embodiment, the present invention provides a method for improving wound healing in a diabetic subject comprising: administering to the subject a therapeutically effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof, so as to improve the rate of wound healing in the subject relative to an untreated wound.

In an embodiment, the present invention provides a method for treating in a subject inflammation associated with transplantation of an organ, a tissue or a plurality of cells from a first site to a second site comprising: administering to the subject a therapeutically effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof, so as to reduce inflammation in the subject. In an embodiment, the first and second sites are in different subjects. In another embodiment, the first and second sites are in the same subject. In another embodiment, the transplanted organ, cells or tissue comprise a cell or tissue of a pancreas, skin, liver, kidney, heart, bone marrow, blood, bone, muscle, artery, vein, cartilage, thyroid, nervous system, or stem cells.

In another embodiment, at least one compound of Formula (I) or a pharmaceutically acceptable salt thereof is utilized, either alone or in combination with one or more known therapeutic agents

As used herein, the phrase "a subject" refers to mammalian subjects, preferably humans, who either suffer from one or more of the aforesaid diseases or disease states or are at risk for such.

In a further aspect of the present invention, the RAGE inhibitors of the invention may be used in adjuvant therapeutic or combination therapeutic treatments with other known therapeutic agents.

The following is a non-exhaustive listing of adjuvants and additional therapeutic agents which may be utilized in combination with the RAGE inhibitors of the present invention:

Pharmacologic classifications of anticancer agents:

1. Alkylating agents: Cyclophosphamide, nitrosoureas, carboplatin, cisplatin, procarbazine
2. Antibiotics: Bleomycin, Daunorubicin, Doxorubicin
3. Antimetabolites: Methotrexate, Cytarabine, Fluorouracil
4. Plant alkaloids: Vinblastine, Vincristine, Etoposide, Paclitaxel,
5. Hormones: Tamoxifen, Octreotide acetate, Finasteride, Flutamide
6. Biologic response modifiers: Interferons, Interleukins, Anti-tumor antibodies

Pharmacologic classifications of treatment for Rheumatoid Arthritis (Inflammation)

1. Analgesics: Aspirin
2. NSAIDs (Nonsteroidal anti-inflammatory drugs): Ibuprofen, Naproxen, Diclofenac
3. DMARDs (Disease-Modifying Antirheumatic drugs): Methotrexate, gold preparations, hydroxychloroquine, sulfasalazine
4. Biologic Response Modifiers, DMARDs: Etanercept, Infliximab Glucocorticoids

Pharmacologic classifications of treatment for Diabetes Mellitus

1. Sulfonylureas: Tolbutamide, Tolazamide, Glyburide, Glipizide
2. Biguanides: Metformin
3. Miscellaneous oral agents: Acarbose, Troglitazone
4. Insulin

Pharmacologic classifications of treatment for Alzheimer's Disease

1. Cholinesterase Inhibitor: Tacrine, Donepezil
2. Antipsychotics: Haloperidol, Thioridazine
3. Antidepressants: Desipramine, Fluoxetine, Trazodone, Paroxetine
4. Anticonvulsants: Carbamazepine, Valproic acid

In a further embodiment, the present invention provides a method of treating a RAGE mediated disease, the method comprising administering to a subject a therapeutically effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof in combination with a therapeutic agent selected from the group consisting of alkylating agents, antimetabolites, plant alkaloids, antibiotics, hormones, biologic response modifiers, analgesics, NSAIDs, DMARDs, glucocorticoids, sulfonylureas, biguanides, insulin, cholinesterase inhibitors, antipsychotics, antidepressants, and anticonvulsants.

In a further embodiment, the present invention provides the pharmaceutical composition of the invention as described above, further comprising one or more therapeutic agents selected from the group consisting of alkylating agents, antimetabolites, plant alkaloids, antibiotics, hormones, biologic response modifiers, analgesics, NSAIDs,

DMARDs, glucocorticoids, sulfonylureas, biguanides, insulin, cholinesterase inhibitors, antipsychotics, antidepressants, and anticonvulsants.

Such other therapeutic agents may be administered by a like route or different route that the compound of Formula (I) or a pharmaceutically acceptable salt thereof. Where a compound of Formula (I) or a pharmaceutically acceptable salt thereof is used in combination with another therapeutic agent, the composition may contain the compound of Formula (I) or a pharmaceutically acceptable salt thereof in combination with the other therapeutic agent(s). Alternatively, where separate dosage formulations are used, the compound of Formula (I) or a pharmaceutically acceptable salt thereof and one or more additional therapeutic agents may be administered at essentially the same time (e.g., concurrently) or at separately staggered times (e.g., sequentially).


Generally speaking, a compound of Formula (I) or a pharmaceutically acceptable salt thereof may be administered at a dosage level of from about 0.003 to 500 mg/kg of the body weight of the subject being treated. In an embodiment, a compound of Formula (I) or a pharmaceutically acceptable salt thereof may be administered at a dosage range between about 0.003 and 200 mg/kg of body weight per day. In an embodiment, a compound of Formula (I) or a pharmaceutically acceptable salt thereof may be administered at a dosage range between about 0.1 to 100 mg/kg of body weight per day. The amount of active ingredient that may be combined with the carrier materials to produce a single dosage may vary depending upon the host treated and the particular mode of administration. For example, a formulation intended for oral administration to humans may contain 1 mg to 2 grams of a compound of Formula (I) or a pharmaceutically acceptable salt thereof with an appropriate and convenient amount of carrier material which may vary from about 5 to 95 percent of the total composition. A dosage form intended for topical administration to the skin may be prepared at 0.1% to 99% compound to topical excipient ratio. A dosage form intended for inhaled administration of 0.01 to 200 mg of compound in a suitable carrier to deliver an inhaled dosage of compound. Dosage unit forms of systemically delivered compound may generally contain between from about 5 mg to about 500 mg of active ingredient. This dosage may be individualized by the clinician based on the specific clinical condition of the subject being treated. Thus, it will be understood that the specific dosage level for any particular subject will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination, size of effected area and the severity of the particular disease undergoing therapy.

The compounds of this invention may be made by a variety of methods well known to those of ordinary skill in the art including the methods are set out below in the Examples.


In another aspect, the present invention also provides a method for the synthesis of compounds useful as intermediates in the preparation of compounds of the present invention along with methods for their preparation.

In an embodiment, the present invention provides a method for synthesizing a compound of Formula (I) or a pharmaceutically acceptable salt thereof

15



comprising: mixing a compound of Formula (X)



and an amine having the formula R^1R^2NH ,
wherein

R^1 and R^2 are independently selected from the group consisting of $-CH_3$, $-CH_2CH_3$, $-CH(CH_3)_2$, and $-CH_2CH_2CH_3$; and

Q^1 is selected from the group consisting of $-CH_2OCH_2CH_3$ and $-CH_2CH_2CH_2CH_3$.

In an embodiment of the method of synthesis, R^1 and R^2 are the same.

In another embodiment of the method of synthesis, R^1 and R^2 are $-CH_3$.

In another embodiment of the method of synthesis, R^1 and R^2 are $-CH_2CH_3$.

In another embodiment of the method of synthesis, Q^1 is $-CH_2OCH_2CH_3$.

In another embodiment of the method of synthesis, Q^1 is $-CH_2CH_2CH_2CH_3$.

In another embodiment of the method of synthesis, the compound of Formula (X) is in the S configuration.

In another embodiment of the method of synthesis, the compound of Formula (X) is in the R configuration.

In another embodiment of the method of synthesis, mixture of the compound of Formula (X) and R^1R^2NH is heated above room temperature. In a further embodiment, the mixture may be heated with microwave radiation.

In another embodiment of the method of synthesis, the compound of Formula (X) and R^1R^2NH are mixed in a solvent. The solvent may be selected from an aprotic solvent. A suitable aprotic solvent includes THF.

EXAMPLES

LC-MS data were obtained using gradient elution on a parallel MUX™ system, running four Waters 1525 binary HPLC pumps, equipped with a Mux-UV 2488 multichannel UV-Vis detector (recording at 215 and 254 nm) and a Leap Technologies HTS PAL Auto sampler using a Sepax GP-C18 4.6x50 mm column. A three minute gradient may be run from 25% of solution B (97.5% acetonitrile, 2.5% water, 0.05% TFA) and 75% of solution A (97.5% water, 2.5% acetonitrile, 0.05% TFA) to 100% of solution B. The system is interfaced with a Waters Micromass ZQ mass

16

spectrometer using electrospray ionization. All MS data was obtained in the positive mode unless otherwise noted.


1H NMR data was obtained on a Varian 400 MHz spectrometer.

Abbreviations used in the Examples are as follows:

d =	day
DCM =	dichloromethane
DMF =	N,N-dimethylformamide
DMSO =	dimethylsulfoxide
ELISA =	enzyme - linked immunosorbent assay
ether =	diethyl ether
EtOAc =	ethyl acetate
g =	gram
h =	hour
Hz =	hertz
L =	liter
LC =	liquid chromatography
M =	molar
m/z =	mass to charge ratio
MeOH =	methanol
mg =	milligram
min =	minute
mL =	milliliter
mM =	millimolar
mmol =	millimole
mol =	mole
MS =	mass spectrometry
N =	normal
NMR =	nuclear magnetic resonance spectroscopy
ppm =	parts per million
rt or RT =	room temperature
TFA =	trifluoroacetic acid
THF =	tetrahydrofuran
TLC =	thin layer chromatography

Intermediate A1

4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenol



Pyridinium bromide perbromide (33.6 g, 0.105 mole) was added to a solution of 4-acetylphenyl acetate (17.8 g, 0.1 mole) in dioxane (100 mL). The heterogeneous mixture was stirred for 5 hours. During the course of the reaction the intensity of the red color decreased and a white solid was formed. The reaction mixture was diluted with ether (200 mL) and washed with water (3x100 mL), brine (75 mL), dried ($MgSO_4$) and removed in vacuo to give the desired product as an oil, which solidified upon standing at room temperature (26.0 g). This product was used in the next transformation without further purification.

A solution of acetic acid 4-(2-bromo-acetyl)-phenyl ester (8.6 g, 33.6 mmol) in DCM (20 mL) was added to a mixture of 4-(4-chlorophenoxy)aniline (6.4 g, 29.2 mmol) and $NaHCO_3$ (4.2 g, 50 mmol) in methanol (100 mL). The formation of a yellow precipitate occurred after 1 h, but the reaction still did not go completion as indicated by both TLC and HPLC. The reaction mixture was further stirred overnight. The solvents were removed in vacuo and the residue was added to ice-water (200 g). The flask was then rinsed

17

with more water (100 mL). After 1 hour, the yellow solid was collected by filtration and washed with water (200 mL). The filtrate (water) in the filtering flask was removed and vacuum kept going on for an hour to remove most of the water. To dry further, the solid was washed with isovaleryl ester, and the amide of the unreacted aniline.


A solution of acetic acid 4-{2-[4-(4-chloro-phenoxy)-phenylamino]-acetyl}-phenyl ester (79.17 g, 200 mmol, 1.0 eq.) in dichloromethane (800 mL) and triethylamine (56 mL, 400 mmol, 2.0 eq.) was cooled to -0° C. and treated with valeryl chloride (35.6 mL, 300 mmol, 1.5 eq.). The reaction mixture was stirred and warmed to room temperature over 24 h. This reaction mixture was then further treated with additional triethylamine (28 mL, 200 mmol, 1.0 eq.) and valeryl chloride (11.9 mL, 100 mmol, 0.5 eq.). Analysis of the reaction by TLC and LC/MS showed that some starting material remained, but the desired keto-amide was the major product. The reaction was evaporated in vacuo, recharged with ethyl acetate and filtered. The solvent was evaporated in vacuo, and the residue was then purified by flash column chromatography over silica gel (EtOAc/hexanes ~25%). The resultant oil was dissolved in ethyl acetate, washed with 1N HCl, dried and evaporated in vacuo. This material was then used as is in the next transformation.

A mixture of acetic acid 4-(2-{[4-(4-chloro-phenoxy)-phenyl]-pentanoyl-amino}-acetyl)-phenyl ester (from above, based on 200 mmol) with ammonium acetate (308 g, 4000 mmol, 20.0 eq) in acetic acid (300 mL) was stirred at $100-110^{\circ}$ C. overnight. After completion of the reaction (indicated by HPLC), the mixture was cooled below 60° C. and poured over ice. After stirring, the solid was filtered, washed with diethyl ether (twice), ethyl acetate (twice), ether (once) and air dried, yielding ~55.0 g (65.6%) of 4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenol as a finely divided off-white solid.

$^1\text{H-NMR}$ (400 MHz; CDCl_3): δ 7.65 (d, 2H), 7.37 (d, 2H), 7.30 (d, 2H), 7.13 (s, 1H), 7.09 (d, 2H), 7.03 (d, 2H), 6.84 (d, 2H), 2.70-2.66 (m, 2H), 1.69-1.61 (m, 2H), 1.33-1.28 (m, 2H), 0.86 (t, 3H).

Intermediate A2

2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-4-[4-((R)-1-oxiranylmethoxy)-phenyl]-1H-imidazole



A mixture of 4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenol (0.42 g, 1.0 mmol, 1.0 eq.) and Cs_2CO_3 (1.0 g, 3.0 mmol, 3.0 eq.) in DMF (3 mL) was stirred and preheated to 80° C. The reaction mixture was then treated with a solution of (2R)-(-)-glycidyl tosylate (0.27 g, 1.2 mmol, 1.2 eq.) in 1 mL of DMF dropwise, and further stirred at 80° C. for ~30-60 min following completion of the addition. Analysis of the reaction by TLC and LC/MS showed that the starting phenol had been consumed and the desired alkylated-phenol was the major product. The reaction was then cooled and diluted with EtOAc and washed with brine. The organic phase was dried with Na_2SO_4 and evaporated in vacuo. The crude alkylated-


18

phenol was then purified by flash column chromatography over silica gel (EtOAc/hexanes ~1:3).

$^1\text{H-NMR}$ (400 MHz; CDCl_3): δ 7.72 (d, 2H), 7.36 (d, 2H), 7.30 (d, 2H), 7.15 (s, 1H), 7.09 (d, 2H), 7.03 (d, 2H), 6.94 (d, 2H), 4.26-4.22 (m, 1H), 4.02-3.98 (m, 1H), 3.40-3.36 (m, 1H), 2.92-2.90 (m, 1H), 2.79-2.77 (m, 1H), 2.69-2.65 (m, 2H), 1.71-1.63 (m, 2H), 1.37-1.27 (m, 2H), 0.86 (t, 3H).

Intermediate A3

2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-4-[4-((S)-1-oxiranylmethoxy)-phenyl]-1H-imidazole




A mixture of 4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenol (0.42 g, 1.0 mmol, 1.0 eq.) and Cs_2CO_3 (1.0 g, 3.0 mmol, 3.0 eq.) in DMF (3 mL) was stirred and preheated to 80° C. The reaction mixture was then treated with a solution of (2S)-(+)-glycidyl tosylate (0.27 g, 1.2 mmol, 1.2 eq.) in 1 mL of DMF dropwise, and further stirred at 80° C. for ~30-60 min following completion of the addition. Analysis of the reaction by TLC and LC/MS showed that the starting phenol had been consumed and the desired alkylated-phenol was the major product. The reaction was then cooled and diluted with EtOAc and washed with brine. The organic phase was dried with Na_2SO_4 and evaporated in vacuo. The crude alkylated-phenol was then purified by flash column chromatography over silica gel (EtOAc/hexanes ~1:3).

$^1\text{H-NMR}$ (400 MHz; CDCl_3): δ 7.72 (d, 2H), 7.36 (d, 2H), 7.30 (d, 2H), 7.15 (s, 1H), 7.09 (d, 2H), 7.03 (d, 2H), 6.94 (d, 2H), 4.26-4.23 (m, 1H), 4.01-3.98 (m, 1H), 3.40-3.36 (m, 1H), 2.93-2.91 (m, 1H), 2.79-2.77 (m, 1H), 2.69-2.65 (m, 2H), 1.71-1.63 (m, 2H), 1.37-1.25 (m, 2H), 0.86 (t, 3H).

Intermediate B1

4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenol



Pyridinium bromide perbromide (33.6 g, 0.105 mole) was added to a solution of 4-acetylphenyl acetate (17.8 g, 0.1 mole) in dioxane (100 mL). The heterogeneous mixture was stirred for 5 hours. During the course of the reaction the intensity of the red color decreased and a white solid was formed. The reaction mixture was diluted with ether (200 mL) and washed with water (3x100 mL), brine (75 mL), dried (MgSO_4) and removed in vacuo to give the desired product as an oil, which solidified upon standing at room

19

temperature (26.0 g). This product was used in the next transformation without further purification.


A solution of acetic acid 4-(2-bromo-acetyl)-phenyl ester (8.6 g, 33.6 mmol) in DCM (20 mL) was added to a mixture of 4-(4-chlorophenoxy)aniline (6.4 g, 29.2 mmol) and NaHCO₃ (4.2 g, 50 mmol) in methanol (100 mL). The formation of a yellow precipitate occurred after 1 h, but the reaction still did not go completion as indicated by both TLC and HPLC. The reaction mixture was further stirred overnight. The solvents were removed in vacuo and the residue was added to ice-water (200 g). The flask was then rinsed with more water (100 mL). After 1 hour, the yellow solid was collected by filtration and washed with water (200 mL). The filtrate (water) in the filtering flask was removed and vacuum kept going on for an hour to remove most of the water. To dry further, the solid was washed with isovaleryl ester, and the amide of the unreacted aniline.

A solution of acetic acid 4-{2-[4-(4-chloro-phenoxy)-phenylamino]-acetyl}-phenyl ester (0.33 mmol, 1.0 eq.) in THF (3 mL) was cooled to -78° C., treated with ethoxyacetyl chloride (0.33 mmol, 1.0 eq.) and stirred for ~5 min. This cold reaction mixture was then treated with pyridine (0.33 mmol, 1.0 eq.) dropwise and allowed to stir for ~1 h. Analysis of the reaction by TLC and LC/MS showed that the starting material has been consumed and the desired ketoamide was the major product. The reaction was then diluted with Et₂O and washed with H₂O, the organic phase was dried with Na₂SO₄ and evaporated in vacuo, and the crude keto-aniline was used in the subsequent step without further purification.

A mixture of N-(4-chlorophenoxyphenyl)-N-(4-acetoxymethyl)-n-pentanamide (0.1011 mol, 1.0 eq) and ammonium acetate (175 g, 2.27 mol, 22.4 eq) in acetic acid (150 mL) was heated at 100-110° C. After completion of the reaction as indicated by HPLC or TLC, the mixture was cooled below 60° C. and is added to chilled water. The solid was filtered, washed with water and ethyl acetate and air dried to produce the desired 4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenol.

Intermediate B2

1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-4-[4-((R)-1-oxiranylmethoxy)-phenyl]-1H-imidazole



A mixture of 4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenol (0.21 g, 0.5 mmol, 1.0 eq.) and Cs₂CO₃ (0.49 g, 1.5 mmol, 3.0 eq.) in DMF (2 mL) was stirred and preheated to 80° C. The reaction mixture was then treated with a solution of (2R)-(-)-glycidyl tosylate (0.17 g, 0.75 mmol, 1.5 eq.) in 1 mL of DMF dropwise, and further stirred at 80° C. for ~30 min following completion of the addition. Analysis of the reaction by TLC and LC/MS showed that the starting phenol had been consumed and the desired alkylated-phenol was the major product. The reaction was then cooled and diluted with EtOAc and washed with brine. The organic phase was dried with Na₂SO₄ and evaporated in vacuo. The crude alkylated-


20

phenol was then purified by flash column chromatography over silica gel (EtOAc/hexanes ~1:3).

¹H-NMR (400 MHz; CDCl₃): δ 7.74 (d, 2H), 7.49 (d, 2H), 7.36 (d, 2H), 7.28 (s, 1H), 7.09 (d, 2H), 7.03 (d, 2H), 6.95 (d, 2H), 4.48 (s, 2H), 4.27-4.23 (m, 1H), 4.03-3.99 (m, 1H), 3.62-3.57 (m, 2H), 3.40-3.37 (m, 1H), 2.94-2.92 (m, 1H), 2.80-2.78 (m, 1H), 1.21 (t, 3H).

Intermediate B3

1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-4-[4-((S)-1-oxiranylmethoxy)-phenyl]-1H-imidazole




A mixture of 4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenol (0.21 g, 0.5 mmol, 1.0 eq.) and Cs₂CO₃ (0.49 g, 1.5 mmol, 3.0 eq.) in DMF (2 mL) was stirred and preheated to 80° C. The reaction mixture was then treated with a solution of (2S)-(+)-glycidyl tosylate (0.17 g, 0.75 mmol, 1.5 eq.) in 1 mL of DMF dropwise, and further stirred at 80° C. for ~30 min following completion of the addition. Analysis of the reaction by TLC and LC/MS showed that the starting phenol had been consumed and the desired alkylated-phenol was the major product. The reaction was then cooled and diluted with EtOAc and washed with brine. The organic phase was dried with Na₂SO₄ and evaporated in vacuo. The crude alkylated-phenol was then purified by flash column chromatography over silica gel (EtOAc/hexanes ~1:3).

¹H-NMR (400 MHz; CDCl₃): δ 7.74 (d, 2H), 7.49 (d, 2H), 7.36 (d, 2H), 7.28 (s, 1H), 7.09 (d, 2H), 7.03 (d, 2H), 6.95 (d, 2H), 4.48 (s, 2H), 4.27-4.24 (m, 1H), 4.03-3.99 (m, 1H), 3.62-3.57 (m, 2H), 3.40-3.37 (m, 1H), 2.94-2.92 (m, 1H), 2.80-2.78 (m, 1H), 1.20 (t, 3H).

Example 1

(R)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-dimethylamino-propan-2-ol dihydrochloride



A solution of 2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-4-[4-((R)-1-oxiranylmethoxy)-phenyl]-1H-imidazole (100 mg, 2.1 mmol, from intermediate A2) in 3 mL of dimethylamine in THF (2M) was stirred at 76° C. for 1 h in a microwave reactor. Upon completion (determined by LC/MS), the reaction was evaporated in vacuo and purified by silica gel flash column chromatography using a gradient of EtOAc to 96% EtOAc/(2M NH₃/MeOH) as an eluent to

21


afford (R)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-dimethylamino-propan-2-ol.

The resultant free base was converted to the corresponding dihydrochloride salt by dissolution in 1 mL of DCM and 3 mL of HCl/dioxane (4.0 M) and removal of solvent in vacuo.

¹H-NMR (400 MHz; CD₃OD): δ 7.90 (s, 1H), 7.73 (d, 2H), 7.62 (d, 2H), 7.44 (d, 2H), 7.23 (d, 2H), 7.16-7.10 (m, 4H), 4.39 (m, 1H), 4.09 (d, 2H), 3.37 (d, 2H), 2.98-2.96 (m, 8H), 1.69-1.66 (m, 2H), 1.37-1.31 (m, 2H), 0.88 (t, 3H).

Example 2

(R)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-diethylamino-propan-2-ol dihydrochloride




A solution of 2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-4-[4-((R)-1-oxiranylmethoxy)-phenyl]-1H-imidazole (100 mg, 2.1 mmol, from intermediate A2) in 1 mL of diethylamine and 2 mL of THF was stirred at 76° C. for 1 h in a microwave reactor. Upon completion (determined by LC/MS), the reaction was evaporated in vacuo and purified by silica gel flash column chromatography using a gradient of EtOAc to 96% EtOAc/(2M NH₃/MeOH) as an eluent to afford (R)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-diethylamino-propan-2-ol.

The resultant free base was converted to the corresponding dihydrochloride salt by dissolution in 1 mL of DCM and 3 mL of HCl/dioxane (4.0 M) and removal of solvent in vacuo.

¹H-NMR (400 MHz; CD₃OD): δ 7.90 (s, 1H), 7.74 (d, 2H), 7.62 (d, 2H), 7.45 (d, 2H), 7.24 (d, 2H), 7.17-7.10 (m, 4H), 4.42-4.38 (m, 1H), 4.11 (d, 2H), 3.45-3.27 (m, 6H), 2.97 (t, 2H), 1.72-1.64 (m, 2H), 1.39-1.30 (m, 8H), 0.89 (t, 3H).

Example 3

(S)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-dimethylamino-propan-2-ol dihydrochloride



A solution of 2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-4-[4-((S)-1-oxiranylmethoxy)-phenyl]-1H-imidazole (from intermediate A3) in 3 mL of dimethylamine in THF (2M) was stirred at 76° C. for 1 h in a microwave reactor. Upon completion (determined by LC/MS), the reaction was

22


evaporated in vacuo and purified by silica gel flash column chromatography using a gradient of EtOAc to 96% EtOAc/(2M NH₃/MeOH) as an eluent to afford (S)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-dimethylamino-propan-2-ol.

The resultant free base was converted to the corresponding dihydrochloride salt by dissolution in 1 mL of DCM and 3 mL of HCl/dioxane (4.0 M) and removal of solvent in vacuo.

¹H-NMR (400 MHz; CD₃OD): δ 7.90 (s, 1H), 7.73 (d, 2H), 7.61 (d, 2H), 7.44 (d, 2H), 7.23 (d, 2H), 7.16-7.10 (m, 4H), 4.38 (m, 1H), 4.09 (d, 2H), 3.37 (d, 2H), 2.98-2.96 (m, 8H), 1.71-1.63 (m, 2H), 1.37-1.30 (m, 2H), 0.88 (t, 3H).

Example 4

(S)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-diethylamino-propan-2-ol dihydrochloride




A solution of 2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-4-[4-((S)-1-oxiranylmethoxy)-phenyl]-1H-imidazole (from intermediate A3) in 1 mL of diethylamine and 2 mL of THF was stirred at 76° C. for 1 h in a microwave reactor. Upon completion (determined by LC/MS), the reaction was evaporated in vacuo and purified by silica gel flash column chromatography using a gradient of EtOAc to 96% EtOAc/(2M NH₃/MeOH) as an eluent to afford (S)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-diethylamino-propan-2-ol.

The resultant free base was converted to the corresponding dihydrochloride salt by dissolution in 1 mL of DCM and 3 mL of HCl/dioxane (4.0 M) and removal of solvent in vacuo.

¹H-NMR (400 MHz; CD₃OD): δ 7.90 (s, 1H), 7.73 (d, 2H), 7.61 (d, 2H), 7.45 (d, 2H), 7.24 (d, 2H), 7.16-7.10 (m, 4H), 4.42-4.37 (m, 1H), 4.10 (d, 2H), 3.42-3.26 (m, 6H), 2.96 (t, 2H), 1.71-1.63 (m, 2H), 1.38-1.31 (m, 8H), 0.88 (t, 3H).

Example 5

(R)-1-(4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-dimethylamino-propan-2-ol dihydrochloride



A solution of 1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-4-[4-((R)-1-oxiranylmethoxy)-phenyl]-1H-imidazole (~100 mg, ~0.20 mmol, from intermediate B2) in

23


3 mL of dimethylamine in THF (2M) was stirred at 60° C. overnight in a teflon-capped vial. Upon completion (determined by LC/MS), the reaction was dried in vacuo and purified by silica gel flash column chromatography using a gradient of EtOAc to 4% ammonia/MeOH (2.0M) in EtOAc as an eluent to afford (R)-1-(4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-dimethylamino-propan-2-ol.

The resultant free base was converted to the corresponding dihydrochloride salt by dissolution in 1 mL of DCM and 3 mL of HCl/dioxane (4.0 M) and removal of solvent in vacuo.

¹H-NMR (400 MHz; CD₃OD): δ 8.05 (s, 1H), 7.75 (d, 2H), 7.65 (d, 2H), 7.44 (d, 2H), 7.23 (d, 2H), 7.16-7.10 (m, 4H), 4.69 (s, 2H), 4.42-4.36 (m, 1H), 4.11 (d, 2H), 3.60 (q, 2H), 3.37 (d, 2H), 2.99 (s, 3H), 2.96 (s, 3H) 1.20 (t, 3H).

Example 6

(S)-1-(4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-dimethylamino-propan-2-ol dihydrochloride




A solution of 1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-4-[4-((S)-1-oxiranylmethoxy)-phenyl]-1H-imidazole (~100 mg, ~0.20 mmol, from intermediate B3) in 3 mL of dimethylamine in THF (2M) was stirred at 60° C. overnight in a teflon-capped vial. Upon completion (determined by LC/MS), the reaction was dried in vacuo and purified by silica gel flash column chromatography using a gradient of EtOAc to 4% ammonia/MeOH (2.0M) in EtOAc as an eluent to afford (S)-1-(4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-dimethylamino-propan-2-ol.

The resultant free base was converted to the corresponding dihydrochloride salt by dissolution in 1 mL of DCM and 3 mL of HCl/dioxane (4.0 M) and removal of solvent in vacuo.

¹H-NMR (400 MHz; CD₃OD): δ 8.05 (s, 1H), 7.76 (d, 2H), 7.65 (d, 2H), 7.44 (d, 2H), 7.23 (d, 2H), 7.16-7.10 (m, 4H), 4.69 (s, 2H), 4.42-4.36 (m, 1H), 4.10 (d, 2H), 3.60 (q, 2H), 3.37 (d, 2H), 2.99 (s, 3H), 2.96 (s, 3H) 1.20 (t, 3H).

Example 7

(R)-1-(4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-diethylamino-propan-2-ol dihydrochloride



24


A solution of 1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-4-[4-((R)-1-oxiranylmethoxy)-phenyl]-1H-imidazole (~100 mg, ~0.20 mmol, from intermediate B2) in 1 mL of diethylamine and 2 mL of THF was stirred at 60° C. overnight in a teflon-capped vial. Upon completion (determined by LC/MS), the reaction was dried in vacuo and purified by silica gel flash column chromatography using a gradient of EtOAc to 4% ammonia/MeOH (2.0M) in EtOAc as an eluent to afford (R)-1-(4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-diethylamino-propan-2-ol.

The resultant free base was converted to the corresponding dihydrochloride salt by dissolution in 1 mL of DCM and 3 mL of HCl/dioxane (4.0 M) and removal of solvent in vacuo.

¹H-NMR (400 MHz; CD₃OD): δ 8.06 (s, 1H), 7.76 (d, 2H), 7.66 (d, 2H), 7.43 (d, 2H), 7.22 (d, 2H), 7.16 (d, 2H), 7.10 (d, 2H), 4.69 (s, 2H), 4.42-4.36 (m, 1H), 4.11 (d, 2H), 3.62-3.56 (q, 2H), 3.41-3.24 (m, 6H), 1.36 (t, 6H), 1.19 (t, 3H).

Example 8

(S)-1-(4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-diethylamino-propan-2-ol dihydrochloride




A solution of 1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-4-[4-((S)-1-oxiranylmethoxy)-phenyl]-1H-imidazole (~100 mg, ~0.20 mmol, from intermediate B3) in 1 mL of diethylamine and 2 mL of THF was stirred at 60° C. overnight in a teflon-capped vial. Upon completion (determined by LC/MS), the reaction was dried in vacuo and purified by silica gel flash column chromatography using a gradient of EtOAc to 4% ammonia/MeOH (2.0M) in EtOAc as an eluent to afford (S)-1-(4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-diethylamino-propan-2-ol.

The resultant free base was converted to the corresponding dihydrochloride salt by dissolution in 1 mL of DCM and 3 mL of HCl/dioxane (4.0 M) and removal of solvent in vacuo.

¹H-NMR (400 MHz; CD₃OD): δ 8.06 (s, 1H), 7.75 (d, 2H), 7.65 (d, 2H), 7.43 (d, 2H), 7.22 (d, 2H), 7.16 (d, 2H), 7.10 (d, 2H), 4.69 (s, 2H), 4.42-4.36 (m, 1H), 4.11 (d, 2H), 3.62-3.56 (q, 2H), 3.41-3.24 (m, 6H), 1.36 (t, 6H), 1.19 (t, 3H).

Example 9

(R)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-methylamino-propan-2-ol dihydrochloride



25


A solution of 2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-4-[4-((R)-1-oxiranylmethoxy)-phenyl]-1H-imidazole (50 mg, 0.11 mmol, from intermediate A2) in 4 mL of methylamine in MeOH (2M) was stirred at 60° C. overnight in a teflon-capped vial. Upon completion (determined by LC/MS), the reaction was dried in vacuo and purified by silica gel flash column chromatography using a gradient of EtOAc to 4% ammonia/MeOH (2.0M) in EtOAc as an eluent to afford (R)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-methylamino-propan-2-ol.

The resultant free base was converted to the corresponding dihydrochloride salt by dissolution in 1 mL of DCM and 3 mL of HCl/dioxane (4.0 M) and removal of solvent in vacuo.

¹H-NMR (400 MHz; CD₃OD): δ 7.90 (s, 1H), 7.73 (d, 2H), 7.61 (d, 2H), 7.43 (d, 2H), 7.23 (d, 2H), 7.15-7.10 (m, 4H), 4.34-4.24 (m, 1H), 4.14-4.06 (m, 2H), 3.30-3.16 (m, 2H), 2.96 (t, 2H), 2.76 (s, 3H), 1.70-1.63 (m, 2H), 1.40-1.28 (m, 2H), 0.87 (t, 3H).

Example 10

(S)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-methylamino-propan-2-ol dihydrochloride




A solution of 2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-4-[4-((S)-1-oxiranylmethoxy)-phenyl]-1H-imidazole (50 mg, 0.11 mmol, from intermediate A3) in 4 mL of methylamine in MeOH (2M) was stirred at 60° C. overnight in a teflon-capped vial. Upon completion (determined by LC/MS), the reaction was dried in vacuo and purified by silica gel flash column chromatography using a gradient of EtOAc to 4% ammonia/MeOH (2.0M) in EtOAc as an eluent to afford (S)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-methylamino-propan-2-ol.

The resultant free base was converted to the corresponding dihydrochloride salt by dissolution in 1 mL of DCM and 3 mL of HCl/dioxane (4.0 M) and removal of solvent in vacuo.

¹H-NMR (400 MHz; CD₃OD): δ 7.90 (s, 1H), 7.73 (d, 2H), 7.61 (d, 2H), 7.43 (d, 2H), 7.23 (d, 2H), 7.15-7.10 (m, 4H), 4.34-4.24 (m, 1H), 4.12-4.06 (m, 2H), 3.30-3.16 (m, 2H), 2.96 (t, 2H), 2.76 (s, 3H), 1.70-1.63 (m, 2H), 1.40-1.28 (m, 2H), 0.87 (t, 3H).

Example 11

(R)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-ethylamino-propan-2-ol dihydrochloride



26


A solution of 2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-4-[4-((R)-1-oxiranylmethoxy)-phenyl]-1H-imidazole (50 mg, 0.11 mmol, from intermediate A2) in 4 mL of ethylamine in MeOH (2M) was stirred at 60° C. overnight in a teflon-capped vial. Upon completion (determined by LC/MS), the reaction was dried in vacuo and purified by silica gel flash column chromatography using a gradient of EtOAc to 4% ammonia/MeOH (2.0M) in EtOAc as an eluent to afford (R)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-ethylamino-propan-2-ol.

The resultant free base was converted to the corresponding dihydrochloride salt by dissolution in 1 mL of DCM and 3 mL of HCl/dioxane (4.0 M) and removal of solvent in vacuo.

¹H-NMR (400 MHz; CD₃OD): δ 7.90 (s, 1H), 7.73 (d, 2H), 7.61 (d, 2H), 7.44 (d, 2H), 7.23 (d, 2H), 7.15-7.10 (m, 4H), 4.34-4.24 (m, 1H), 4.14-4.06 (m, 2H), 3.34-3.28 (m, 1H), 3.19-3.11 (m, 3H), 2.96 (t, 2H), 1.71-1.63 (m, 2H), 1.38-1.30 (m, 5H), 0.88 (t, 3H).

Example 12

(S)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-ethylamino-propan-2-ol dihydrochloride




A solution of 2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-4-[4-((S)-1-oxiranylmethoxy)-phenyl]-1H-imidazole (50 mg, 0.11 mmol, from intermediate A3) in 4 mL of ethylamine in MeOH (2M) was stirred at 60° C. overnight in a teflon-capped vial. Upon completion (determined by LC/MS), the reaction was dried in vacuo and purified by silica gel flash column chromatography using a gradient of EtOAc to 4% ammonia/MeOH (2.0M) in EtOAc as an eluent to afford (S)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-ethylamino-propan-2-ol.

The resultant free base was converted to the corresponding dihydrochloride salt by dissolution in 1 mL of DCM and 3 mL of HCl/dioxane (4.0 M) and removal of solvent in vacuo.

¹H-NMR (400 MHz; CD₃OD): δ 7.90 (s, 1H), 7.73 (d, 2H), 7.61 (d, 2H), 7.44 (d, 2H), 7.23 (d, 2H), 7.16-7.10 (m, 4H), 4.34-4.24 (m, 1H), 4.14-4.06 (m, 2H), 3.34-3.28 (m, 1H), 3.18-3.11 (m, 3H), 2.96 (t, 2H), 1.71-1.63 (m, 2H), 1.38-1.29 (m, 5H), 0.88 (t, 3H).

Example 13

(R)-1-(4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-methylamino-propan-2-ol dihydrochloride



27


A solution of 1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-4-[4-((R)-1-oxiranylmethoxy)-phenyl]-1H-imidazole (50 mg, 0.11 mmol, from intermediate B2) in 4 mL of methylamine in MeOH (2M) was stirred at 60° C. overnight in a teflon-capped vial. Upon completion (determined by LC/MS), the reaction was dried in vacuo and purified by silica gel flash column chromatography using a gradient of EtOAc to 4% ammonia/MeOH (2.0M) in EtOAc as an eluent to afford (R)-1-(4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-methylamino-propan-2-ol.

The resultant free base was converted to the corresponding dihydrochloride salt by dissolution in 1 mL of DCM and 3 mL of HCl/dioxane (4.0 M) and removal of solvent in vacuo.

¹H-NMR (400 MHz; CD₃OD): δ 8.06 (s, 1H), 7.75 (d, 2H), 7.65 (d, 2H), 7.43 (d, 2H), 7.23 (d, 2H), 7.16-7.10 (m, 4H), 4.69 (s, 2H), 4.31-4.26 (m, 1H), 4.13-4.07 (m, 2H), 3.60 (q, 2H), 3.32-3.28 (m, 1H), 3.21-3.15 (m, 1H), 2.76 (s, 3H), 1.19 (t, 3H).

Example 14

(S)-1-(4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-methylamino-propan-2-ol dihydrochloride




A solution of 1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-4-[4-((S)-1-oxiranylmethoxy)-phenyl]-1H-imidazole (50 mg, 0.11 mmol, from intermediate B3) in 4 mL of methylamine in MeOH (2M) was stirred at 60° C. overnight in a teflon-capped vial. Upon completion (determined by LC/MS), the reaction was dried in vacuo and purified by silica gel flash column chromatography using a gradient of EtOAc to 4% ammonia/MeOH (2.0M) in EtOAc as an eluent to afford (S)-1-(4-{1-[4-(4-Chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-methylamino-propan-2-ol.

The resultant free base was converted to the corresponding dihydrochloride salt by dissolution in 1 mL of DCM and 3 mL of HCl/dioxane (4.0 M) and removal of solvent in vacuo.

¹H-NMR (400 MHz; CD₃OD): δ 8.06 (s, 1H), 7.75 (d, 2H), 7.65 (d, 2H), 7.44 (d, 2H), 7.23 (d, 2H), 7.16-7.10 (m, 4H), 4.69 (s, 2H), 4.31-4.26 (m, 1H), 4.13-4.07 (m, 2H), 3.60 (q, 2H), 3.32-3.28 (m, 1H), 3.21-3.15 (m, 1H), 2.76 (s, 3H), 1.20 (t, 3H).

Example 15

(R)-1-(4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-ethylamino-propan-2-ol dihydrochloride



28


A solution of 1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-4-[4-((R)-1-oxiranylmethoxy)-phenyl]-1H-imidazole (50 mg, 0.11 mmol, from intermediate B2) in 4 mL of ethylamine in MeOH (2M) was stirred at 60° C. overnight in a teflon-capped vial. Upon completion (determined by LC/MS), the reaction was dried in vacuo and purified by silica gel flash column chromatography using a gradient of EtOAc to 4% ammonia/MeOH (2.0M) in EtOAc as an eluent to afford (R)-1-(4-{1-[4-(4-Chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-ethylamino-propan-2-ol.

The resultant free base was converted to the corresponding dihydrochloride salt by dissolution in 1 mL of DCM and 3 mL of HCl/dioxane (4.0 M) and removal of solvent in vacuo.

¹H-NMR (400 MHz; CD₃OD): δ 8.02 (s, 1H), 7.75 (d, 2H), 7.64 (d, 2H), 7.44 (d, 2H), 7.23 (d, 2H), 7.16-7.10 (m, 4H), 4.67 (s, 2H), 4.29-4.25 (m, 1H), 4.13-4.06 (m, 2H), 3.60 (q, 2H), 3.32-3.28 (m, 1H), 3.19-3.11 (m, 3H), 1.35 (t, 3H), 1.20 (t, 3H).

Example 16

(S)-1-(4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-ethylamino-propan-2-ol dihydrochloride




A solution of 1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-4-[4-((S)-1-oxiranylmethoxy)-phenyl]-1H-imidazole (50 mg, 0.11 mmol, from intermediate B3) in 4 mL of ethylamine in MeOH (2M) was stirred at 60° C. overnight in a teflon-capped vial. Upon completion (determined by LC/MS), the reaction was dried in vacuo and purified by silica gel flash column chromatography using a gradient of EtOAc to 4% ammonia/MeOH (2.0M) in EtOAc as an eluent to afford (S)-1-(4-{1-[4-(4-Chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-ethylamino-propan-2-ol.

The resultant free base was converted to the corresponding dihydrochloride salt by dissolution in 1 mL of DCM and 3 mL of HCl/dioxane (4.0 M) and removal of solvent in vacuo.

¹H-NMR (400 MHz; CD₃OD): δ 8.04 (s, 1H), 7.75 (d, 2H), 7.65 (d, 2H), 7.44 (d, 2H), 7.23 (d, 2H), 7.16-7.10 (m, 4H), 4.68 (s, 2H), 4.29-4.25 (m, 1H), 4.13-4.06 (m, 2H), 3.60 (q, 2H), 3.32-3.28 (m, 1H), 3.19-3.11 (m, 3H), 1.35 (t, 3H), 1.20 (t, 3H).

Example Z

[3-(4-{2-butyl-1-[4-(4-chlorophenoxy)-phenyl]-1H-imidazole-4-yl}-phenoxy)-propyl]-diethylamine dihydrochloride



Example Z may be prepared according to the method described in PCT publication number WO 2003/075921 for Example 406.

Biological Assay

The following assay method may be used to identify compounds of Formula (I) or pharmaceutically acceptable salts thereof which are useful as inhibitors of binding of physiological RAGE ligands, such as S100b and β -amyloid, to RAGE.

S100b, β -amyloid, or CML (500 ng/100 μ L/well) in 100 mM sodium bicarbonate/sodium carbonate buffer (pH 9.8) is loaded onto the wells of a NUNC Maxisorp flat bottom 96-well microtitre plate. The plate is incubated at 4° C. overnight. The wells are aspirated and treated with 50 mM imidazole buffer saline (pH 7.2) (with 5 mM $\text{CaCl}_2/\text{MgCl}_2$) containing 1% bovine serum albumin (BSA) (300 μ L/well) for 1 h at RT. The wells are aspirated.

Test compounds are dissolved in nanopure water (concentration: 10-100 μ M). DMSO may be used as co-solvent. 25 μ L of test compound solution in 4% DMSO is added, along with 75 μ L sRAGE (6 nM FAC) to each well and samples are incubated for 1 h at 37° C. The wells are washed several times with 155 mM NaCl pH 7.2 buffer saline and are soaked for several seconds between each wash.

Non-radioactive detection is performed by adding:

10 μ L Biotinylated goat F(ab')₂ Anti-mouse IgG. (8.0 \times 10⁻⁴ mg/mL, FAC), 5 μ L Alk-phos-Streptavidin (3 \times 10⁻³ mg/mL FAC), 0.42 μ L per 5 mL Monoclonal antibody for sRAGE (FAC 6.0 \times 10⁻³ mg/mL) to 5 mL 50 mM imidazole buffer saline (pH 7.2) containing 0.2% bovine serum albumin and 5 mM CaCl_2 . The mixture is incubated for 30 minutes at RT.

100 μ L of complex is added to each well and incubation is allowed to proceed at rt for 1 h. Wells are washed several times with wash buffer and soaked several seconds between each wash. 100 μ L 1 mg/mL (pNPP) in 1 M diethanolamine (pH adjusted to 9.8 with HCl) is added. Color is allowed to develop in the dark for 30 min to 1 h at rt. The reaction is quenched with 10 μ L of stop solution (0.5-1.0 N NaOH in 50% ethanol) and the absorbance is measured spectrophotometrically with a microplate reader at 405 nm.

The Examples 1-16 (hydrochloride salt form) were tested according to the assay method described above, employing S100b or β -amyloid as the RAGE ligand, and were found to possess IC₅₀ shown below. IC₅₀ (μ M) of in the ELISA assay represents the concentration of compound at which 50% signal has been inhibited.

Example	IC ₅₀ (β -amyloid) (μ M)	IC ₅₀ (S100b) (μ M)
1	0.85	0.66
2	0.76	0.55
3	0.80	0.84
4	0.65	0.54

-continued

Example	IC ₅₀ (β -amyloid) (μ M)	IC ₅₀ (S100b) (μ M)
5	1.02	0.71
6	0.78	0.77
7	1.17	1.05
8	1.26	0.80
9	1.59	1.13
10	1.32	1.14
11	1.02	0.81
12	1.19	0.98
13	2.16	4.61
14	2.37	4.56
15	2.47	3.14
16	1.55	3.13

Pharmacokinetics

Pharmacokinetic screening in rats was performed on various compounds to measure brain and plasma concentrations at 6 hour time point.

The parameters for the pharmacokinetic protocol were as follows.

Amount of compound: 5 mg/kg

Species: Rat; Strain: Sprague Dawley; Sex: Male

Average body weight at dose: weight ranged from 271 to 423 grams

Average age at dose: age ranged from 9 to 14 weeks

Diet Status Overnight fasting

Number of Animals (n) for each time point: 2

Dosing: Oral (PO)

Formulation: 2% Tween 80 in distilled water

Each formulation was administered once by oral gavage. The dose volume was 5 mL/kg for all animals. The actual volume administered to each animal was calculated and adjusted based on the most recent body weight.

Blood samples (approximately 300 μ L whole blood) at (1, 2, and 4 h) was collected from each animal via tail vein except for terminal blood samples. Terminal blood (6 h) samples were collected via cardiac puncture. All samples were collected into tubes containing lithium heparin (Multivette 600 LH-Gel, Sarstedt, Newton, N.C., USA). Following collection, the tubes were placed in refrigerator (maximum 30 minutes) or until centrifugation under refrigeration (at 2 to 8° C.) at 1500g for approximately 15 minutes. Each harvested plasma sample was then transferred into 1.2 mL polypropylene tubes, on the 96-Well Plate according to the 96-Well Plate plasma sample map and kept in freezer. Plasma samples were then analyzed for test substances.

Brain samples were collected immediately after the animals were euthanized at designated time points. Brain samples were rinsed with saline, blotted dry, and weighed. Brain samples were placed into individual containers and kept in freezer (-20° C.). Brain samples were then analyzed for test articles.

After analysis, all the plasma results are reported as ng/mL and brain sample results are reported as ng/g. In the table below, "ND" stands for not determined and "NA" stands for not applicable.

Ex.	Brain (ng/g)	Plasma (ng/mL)	B/P Ratio	R ¹	R ²	Q ¹	Config
Z	697	92	7.7	-CH ₃ CH ₂	-CH ₃ CH ₂	butyl	NA
1	626	18	34.1	-CH ₃	-CH ₃	butyl	R
2	718	24	30.6	-CH ₃ CH ₂	-CH ₃ CH ₂	butyl	R
3	1120	48	23.3	-CH ₃	-CH ₃	butyl	S
4	610	74	8.8	-CH ₃ CH ₂	-CH ₃ CH ₂	butyl	S
5	3325	200	16.7	-CH ₃	-CH ₃	ethoxymethyl	R
6	3905	155	25.3	-CH ₃	-CH ₃	ethoxymethyl	S


Ex.	Brain (ng/g)	Plasma (ng/mL)	B/P Ratio	R ¹	R ²	Q ¹	Config
7	1385	153	9.1	—CH ₃ CH ₂	—CH ₃	ethoxymethyl	R
8	2705	137	19.6	—CH ₃ CH ₂	—CH ₃	ethoxymethyl	S
9	537	76	7.2	H	—CH ₃	butyl	R
10	212	74	2.9	H	—CH ₃	butyl	S
11	343	72	4.8	H	—CH ₃ CH ₂	butyl	R
12	540	124	4.5	H	—CH ₃ CH ₂	butyl	S
13	ND	ND	ND	H	—CH ₃	ethoxymethyl	R
14	ND	ND	ND	H	—CH ₃	ethoxymethyl	S
15	ND	ND	ND	H	—CH ₃ CH ₂	ethoxymethyl	R
16	ND	ND	ND	H	—CH ₃ CH ₂	ethoxymethyl	S

The specific pharmacological responses observed may vary according to and depending on the particular active compound selected or whether there are present pharmaceutical carriers, as well as the type of formulation and mode of administration employed, and such expected variations or differences in the results are contemplated in accordance with practice of the present invention.

While the invention has been described and illustrated with reference to certain preferred embodiments thereof, those skilled in the art will appreciate that various changes, modifications and substitutions can be made therein without departing from the spirit and scope of the invention. For example, effective dosages other than the preferred dosages as set forth herein may be applicable as a consequence of variations in the responsiveness of the mammal being treated for RAGE-mediated disease(s). Likewise, the specific pharmacological responses observed may vary according to and depending on the particular active compound selected or whether there are present pharmaceutical carriers, as well as the type of formulation and mode of administration employed, and such expected variations or differences in the results are contemplated in accordance with the objects and practices of the present invention.

We claim:

1. A method for treating Alzheimer's disease comprising administering to a subject a compound of Formula (I) or a pharmaceutically acceptable salt thereof,



wherein

R¹ and R² are independently selected from the group consisting of: —CH₃, —CH₂CH₃, —CH(CH₃)₂, and —CH₂CH₂CH₃; and

Q¹ is selected from the group consisting of —CH₂OCH₂CH₃ and —CH₂CH₂CH₂CH₃.

2. The method of claim 1, wherein the compound is (R)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-dimethylamino-propan-2-ol or a pharmaceutically acceptable salt thereof.

3. The method of claim 1, wherein the compound is (R)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-diethylamino-propan-2-ol or a pharmaceutically acceptable salt thereof.

4. The method of claim 1, wherein the compound is (S)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-dimethylamino-propan-2-ol or a pharmaceutically acceptable salt thereof.

5. The method of claim 1, wherein the compound is (S)-1-(4-{2-butyl-1-[4-(4-chloro-phenoxy)-phenyl]-1H-imidazol-4-yl}-phenoxy)-3-diethylamino-propan-2-ol or a pharmaceutically acceptable salt thereof.

6. The method of claim 1, wherein the compound is (R)-1-(4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-dimethylamino-propan-2-ol or a pharmaceutically acceptable salt thereof.

7. The method of claim 1, wherein the compound is (S)-1-(4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-dimethylamino-propan-2-ol or a pharmaceutically acceptable salt thereof.

8. The method of claim 1, wherein the compound is (R)-1-(4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-diethylamino-propan-2-ol or a pharmaceutically acceptable salt thereof.

9. The method of claim 1, wherein the compound is (S)-1-(4-{1-[4-(4-chloro-phenoxy)-phenyl]-2-ethoxymethyl-1H-imidazol-4-yl}-phenoxy)-3-diethylamino-propan-2-ol or a pharmaceutically acceptable salt thereof.

* * * * *