

US009695453B2

(12) **United States Patent**
Corsa et al.

(10) **Patent No.:** **US 9,695,453 B2**
(45) **Date of Patent:** ***Jul. 4, 2017**

(54) **PROCESS FOR THE PRODUCTION OF
HYALURONIC ACID IN *ESCHERICHIA COLI*
OR *BACILLUS MEGATERIUM***

(71) Applicant: **FIDIA FARMACEUTICI S.P.A.**,
Abano Terme (IT)

(72) Inventors: **Vincenza Corsa**, Albano Terme (IT);
Alessandro Negro, Padua (IT); **Sonia
Bisicchia**, Abano Terme (IT)

(73) Assignee: **FIDIA FARMACEUTICI S.P.A.**,
Abano Terme (IT)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

This patent is subject to a terminal dis-
claimer.

(21) Appl. No.: **14/884,274**

(22) Filed: **Oct. 15, 2015**

(65) **Prior Publication Data**

US 2016/0032337 A1 Feb. 4, 2016

Related U.S. Application Data

(63) Continuation of application No. 13/821,953, filed as
application No. PCT/EP2011/065641 on Apr. 15,
2013, now Pat. No. 9,163,270.

(30) **Foreign Application Priority Data**

Sep. 9, 2010 (IT) MI2010A1641

(51) **Int. Cl.**

C12P 19/26 (2006.01)
C12P 19/04 (2006.01)
C08B 37/08 (2006.01)
C12N 9/04 (2006.01)
C12N 9/10 (2006.01)
C12N 9/12 (2006.01)
C12N 9/92 (2006.01)
C12N 15/75 (2006.01)

(52) **U.S. Cl.**

CPC **C12P 19/04** (2013.01); **C08B 37/0072**
(2013.01); **C12N 9/0006** (2013.01); **C12N**
9/1051 (2013.01); **C12N 9/1241** (2013.01);
C12N 9/92 (2013.01); **C12N 15/75** (2013.01);
C12P 19/26 (2013.01); **C12Y 101/01022**
(2013.01); **C12Y 204/01212** (2013.01); **C12Y**
207/07009 (2013.01); **C12Y 503/01009**
(2013.01)

(58) **Field of Classification Search**

CPC **C12Y 101/01022**; **C12Y 204/01212**; **C12Y**
207/07009; **C12Y 503/01009**

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,955,310 A 9/1999 Widner et al.
7,811,806 B2 10/2010 Sloma et al.
9,163,270 B2 * 10/2015 Corsa C08B 37/0072
2003/0175902 A1 9/2003 Sloma et al.

OTHER PUBLICATIONS

Chien, L. et al., "Enhanced hyaluronic acid production in *Bacillus subtilis* by coexpressing bacterial hemoglobin", *Biotechnology Progress*, Sep. 2007, vol. 23, No. 5, pp. 1017-1022.

Crater et al., "Molecular Characterization of hasC from an Operon Required for Hyaluronic Acid Synthesis in Group A Streptococci," *JBC*, Dec. 1995, vol. 270, No. 48, pp. 28676-28680.

Dougherty et al. "Molecular Characterization of hasB from an Operon Required for Hyaluronic Acid Synthesis in Group A Streptococci," *JBC*, Dec. 1993, vol. 268, No. 10, pp. 7118-7124-28680.

Duenas et al., "Synthesis of a eukaryotic virus protein in a prokaryotic viral-cell system: production of the adenovirus type 2 fiber shaft fragment by a tightly regulated T7POL-M13 expression system," *J. Virological Methods*, Jan. 1999, vol. 79, pp. 121-131.

Gamer, M. et al., "A T7 RNA polymerase-dependent gene expression system for *Bacillus megaterium*", *Applied Microbiology and Biotechnology*, Springer, Berlin, DE, Mar. 24, 2009, vol. 82, No. 6, pp. 1195-1203.

Kang, Y. et al., "One step engineering of T7-expression strains for protein production: Increasing the host-range of the T7-expression system", *Protein Expression and Purification*, Academic Press, San Diego, CA, US, Sep. 8, 2007, vol. 55, No. 2, pp. 325-333.

Mao, Z. et al., "A recombinant *E. coli* bioprocess for hyaluronan synthesis", *Applied Microbiology and Biotechnology*, Springer, Berlin, DE, Mar. 24, 2009, vol. 84, No. 1, pp. 63-69.

Mathur et al., "Biochemical characterization of recombinant phosphoglucose isomerase of *Mycobacterium tuberculosis*," *Biochem Biophys Res Commun.*, Elsevier, Sep. 2005, vol. 337, pp. 626-632.

Studier, F. W. et al., "Use of T7 RNA polymerase to direct expression of cloned genes", *Methods in Enzymology*, Academic Press Inc., San Diego, CA, US, Jan. 1, 1990, vol. 185, pp. 60-89.

Widner, B. et al., "Hyaluronic acid production in *Bacillus subtilis*", *Applied and Environmental Microbiology*, American Society for Microbiology, US, Jul. 1, 2005, vol. 71, No. 1, pp. 3747-3752.

Yu, H. et al., "Metabolic engineering of *Escherichia coli* for biosynthesis of hyaluronic acid", *Metabolic Engineering*, Academic Press, US, Dec. 24, 2007, vol. 10, No. 1, pp. 24-32.

* cited by examiner

Primary Examiner — Karen Cochrane Carlson


(74) *Attorney, Agent, or Firm* — Birch, Stewart, Kolasch & Birch, LLP

(57) **ABSTRACT**

A method of producing hyaluronic acid (HA) in *Escherichia coli* and *Bacillus megaterium* through episomal plasmid vectors wherein the gene is under the control of strong promoter T7, preferably under the control of strong promoter T7 of bacteriophage T7, and a system for the selection of stable bacterial strains producing high levels of hyaluronic acid, are provided.

32 Claims, 6 Drawing Sheets


HAS1 e TUAD expression in E.coli TOP-10



Cells incorporating plasmide HT01 are larger than cells with BS5 (difficulty to grow)

(Cells incorporating BS5 are more yellow than parental cells)

Fig. 1



TnsD expression in E coli BL21 DE3

Fig. 2

Constitutive expression of Hyaluronan synthase (Streptococcus) in E. coli




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8




Fig. 9




Fig. 10

**PROCESS FOR THE PRODUCTION OF
HYALURONIC ACID IN *ESCHERICHIA COLI*
OR *BACILLUS MEGATERIUM***

This application is a Continuation of copending application Ser. No. 13/821,953, filed on Apr. 15, 2013, which was filed as the National Phase of PCT International Application No. PCT/EP2011/065641 on Sep. 9, 2011, which claims the benefit under 35 U.S.C. §119(a) to Patent Application No. MI2010A001641, filed in Italy on Sep. 9, 2010, all of which are hereby expressly incorporated by reference into the present application.

SUBJECT OF THE INVENTION

The present invention discloses a process for the production of hyaluronic acid (HA) in *Escherichia coli* and *Bacillus megaterium* through episomal plasmid vectors wherein the gene is under the control of a strong T7 promoter, preferably under the control of a strong T7 promoter of bacteriophage T7, and a system for the selection of stable bacterial strains producing high levels of hyaluronic acid.

FIELD OF INVENTION

Hyaluronic acid is a natural linear polysaccharide which consists of alternating β -1-4 D-glucuronic acid and β -1-3 N-acetyl glucosamine. Hyaluronic acid is part of the glycosaminoglycan family, and can reach a molecular weight of 10^7 Da, with approx. 300000 repeating saccharide units. It is widely distributed in the connective tissue and extracellular matrix in the epithelium of eukaryotic organisms, where it is located on the cell surface, but can also be synthesised in some prokaryotic organisms, such as those of the *Streptococcus* family. Glycosaminoglycans are ideal joint lubricants, but also perform many other functional roles in tissue repair, cell motility, adhesion and development, cancer and angiogenesis. Products based on hyaluronic acid have been developed on the basis of these important characteristics, and are used in orthopaedics, rheumatology and dermatology.

The most common natural sources of HA include rooster combs, the classic material from which HA is extracted, and some bacteria, especially those belonging to the *Streptococcus* family. All these different sources present numerous disadvantages: hyaluronic acid obtained from rooster combs can, for example, cause allergies in humans because it is of avian origin, while HA from bacterial sources must be free of all the toxins normally present in those bacteria which can cause possibly serious immune/inflammatory reactions. The current industrial HA purification processes therefore comprise many different steps, with a consequent increase in the final costs of manufacturing the raw material.

There is consequently a strongly felt need for alternative sources that eliminate all the adverse events described, while maintaining reasonable manufacturing costs. In recent years, biosynthesis pathways for the synthesis of hyaluronic acid have been included in detail in numerous organisms. While the genes required for hyaluronic acid synthesis which are present in eukaryotic organisms are distributed throughout the genome, in bacterial systems said genes are often present and organised in operons. For example, in *Streptococcus equi* the operon for hyaluronic acid comprises 5 genes: hasA, hasB, hasC, hasD and hasE. Sometimes, however, the genes are present in two operons: in *Streptococcus equisimilis* one operon with genes hasA, hasB and hasC is present, and another with genes hasC, hasD and hasE. The genes

homologous with hasB, hasC, hasD and hasE of the Streptococci are present in many organisms, and synthesise the enzymes necessary for the synthesis of hyaluronic acid precursors D-glucuronic acid and N acetyl-D glucosamine, which are also the essential constituents of the bacterial walls. In the case of streptococci, hyaluronan synthase (hasA, which is present in the plasma membrane) is the key enzyme for the final synthesis of hyaluronic acid because it performs two functions: it catalyses the union of D-glucuronic acid and N-acetyl-D-glucosamine, and transports the chain of newly-formed hyaluronic acid out of the cell. The study of the enzymes responsible for hyaluronic acid synthesis has allowed the development of recombinant systems in various organisms, such as *Bacillus subtilis*, *Lactococcus lactis*, *Escherichia coli* and *Agrobacterium radiobacter*. The first organism engineered to produce hyaluronic acid was *B. subtilis*, through cloning in its chromosome of an operon that carries the hasA gene from *Streptococcus* (which is missing in *Bacillus*), with the tuaD and gtaB genes of *Bacillus* (corresponding to hasB and hasC of *Streptococcus*), under the control of a constitutive promoter (US2003/175902). In this way a biosynthesis pathway was organised in operons similar to those of *Streptococcus equi*, one of the major natural producers of hyaluronic acid. However, the system thus perfected leads to the industrial production of a hyaluronic acid with a weight average molecular weight of less than 1 MDA, with very low manufacturing yields.

The system of expression of hyaluronic acid according to the present invention uses bacteria of the strains *Bacillus Megaterium* and *Escherichia coli*.

Bacillus Megaterium is an aerobic gram-positive bacterium, which was described over 100 years ago. Its large size (1 μ m, i.e. 100 times larger than *E. coli* in both vegetative and spore-forming form) has made it very popular for morphological analysis studies. This bacterium can contain many different types of plasmids; the plasmid DNA can be transferred by protoplast transformation obtained by treatment with polyethylene glycol, and they all work extremely well, with excellent structural stability. The bacterium can be transduced with phages, and the frequency of transformation can reach 10^6 transformants per μ g of DNA. Several hundred mutants are currently available, which cover various biosynthesis pathways: catabolism, division, sporulation, germination, antibiotic resistance and recombination.

No less than seven plasmids have been found in different strains of *B. megaterium*, with sizes ranging from 5.4 to 165 kb. The genomes of two strains (DSM319:EMBL, accession number CP001983, and QM B1551:EMBL, accession number CP001982) and those of the seven natural plasmids are now available. Although it is considered to be a bacterium present in soil, *B. megaterium* has been found in various ecological niches such as dried meat, seawater and fish. *B. megaterium* is able to grow in various carbon sources, including slaughter waste and industrial syrups with a broad spectrum of sugars (62 of the 95 tested), which include carboxylic acids like acetate. *B. megaterium* can be cultured at high density, up to 80 g of dry weight per litre. Considerable knowledge has been obtained of various recombinant enzymes with different industrial applications which can be secreted in this organism, such as α -amylase, β -amylase, penicillin amidase, neutral protease and β -glucanase. Particularly important are amylases, used in the bread-making industry, glucose dehydrogenase, used industrially for the production of NADH and as a biosensor, and penicillin

amidase, used to generate new synthetic antibiotics. Finally, *B. megaterium* is the major source of vitamin B12.

DETAILED DESCRIPTION OF THE INVENTION

The present invention discloses and claims a process for the production of hyaluronic acid (HA) in high industrial yields in *Bacillus megaterium* and *Escherichia coli* through episomal plasmid vectors wherein the genes for the synthesis of the enzymes required for HA production, are under the control of the strong T7 promoter, preferably under the control of the strong T7 promoter of bacteriophage T7, and a system for the selection of engineered, stable bacterial strains producing high amounts of hyaluronic acid having well defined weight average molecular weights (in the following also indicated as MW).

In order to produce recombinant proteins (in this case the enzymatic proteins required for the synthesis of HA) efficiently, systems which use highly controllable strong promoters need to be designed. The invention discloses a process for the transformation of the above-disclosed bacteria, using a very efficient system for the control of the transcription of the genes introduced, as the gene of interest is placed under the control of the promoter dependent on T7 RNA polymerase.

During construction in *E. coli* of the vectors expressing hyaluronic acid in the form of plasmids, it was discovered that the genes thus introduced (which are responsible for synthesis of the hyaluronic acid-producing enzymes) are cell-toxic when their transduction control is a strong constitutive promoter. In fact, in *E. coli* transformed with genes *hasA* and *tuaD*, gene transduction of *hasA* alone leads to a great reduction in the D-glucuronic acid precursors required to constitute the bacterial wall, with the result that the cell dies, whereas gene transduction of *tuaD* alone generates uncontrolled synthesis of D-glucuronic acid which, by acidifying the bacterium and depriving it of glucose (its precursor), causes its death. Conversely, the transduction of both genes by bacterial polymerases leads to the activation of the two enzymes at different times, because they require different construction times with different procedures and sites of action (for example, *hasA* is a transmembrane protein with different domains crossing it, so a much longer time is needed for its synthesis and correct folding). The cell can only survive if balanced quantities of the precursor enzymes and the enzyme necessary for hyaluronic acid synthesis are present. In this case, the excess D-glucuronic acid, which is toxic at high levels in the cell, is used by hyaluronan synthase (*hasA*) which, combining it with glucosamine, incorporates it in the nascent hyaluronic acid and exports it from the cell, thus keeping the cell alive.

Consequently, although both *hasA* and *tuaD* are necessary for the synthesis of hyaluronic acid, it is essential for the two genes to work in concert, leaving the cell the time required to:

produce D-glucuronic acid at non-toxic levels and trigger the transcription of the *hasA* gene in such a way that the latter is able to dispose of the high levels of D-glucuronic acid as they accumulate in the cell.

In the present invention, the problems described above have been solved by

placing the plasmid genes, necessary for the synthesis of the above disclosed enzymes, under the control of a T7 promoter, preferably under the control of the T7 promoter of bacteriophage T7, dependent on T7 RNA polymerase, which uses repressor Xy1R in *B. megate-*

rium (and *lac* in *E. coli*) for its induction. The T7 promoter of bacteriophage T7 is dependent on the presence of T7 polymerase, so the HA synthesis genes placed under its control can only be transcribed by T7 RNA polymerase, not by the action of the polymerases naturally present in the bacterium;

perfecting a system of selection of stable, engineered and secreting *B. megaterium* strains and *E. Coli* strains, preferably of viable, engineered and secreting *B. megaterium* strains, wherein the enzymes necessary for the HA synthesis, are present in "balanced" amounts, thus non toxic for the cell.

It is therefore object of the present invention a process for the preparation of hyaluronic acid in *Escherichia coli* or *Bacillus megaterium*, preferably in *B. megaterium*, comprising the following steps:

(a) culture of bacterial host cells of *Escherichia coli* or *Bacillus megaterium*, preferably of host cells of *B. megaterium*, transformed in a stable way with the T7 RNA polymerase system under conditions suitable for the production of hyaluronic acid in the presence of isopropyl- β -thiogalactopyranoside (IPTG) or xylose respectively as inducers, wherein said bacterial host cells are characterised by being further transformed with:

- (i) at least one episomal plasmid vector comprising a sequence coding for the enzyme hyaluronan synthase and a sequence coding for the enzyme UDP-glucose dehydrogenase in tandem under the control of the strong inducible T7 promoter, preferably under the control of the T7 promoter of bacteriophage T7; or
- (ii) at least one episomal plasmid vector comprising a sequence coding for the enzyme hyaluronate synthase, a sequence coding for the enzyme UDP-glucose dehydrogenase, a sequence coding for the enzyme UDP-glucose pyrophosphorylase and a sequence coding for the enzyme glucose 6 phosphate isomerase, under the control of the strong inducible T7, preferably under the control of the T7 promoter of bacteriophage T7;

(b) recovery of hyaluronic acid from the culture medium, wherein such bacterial host cells of *Escherichia coli* or *Bacillus megaterium* transformed in a stable way with the T7 RNA polymerase system and with plasmid vector (i) or (ii) able to produce hyaluronic acid of step a) are pre-selected in the plate on IPTG or xylose gradient respectively.

The Applicant preferably used *B. megaterium* (preferably pertaining to QMB1551 or DSM319 strains), transformed with the T7 RNA polymerase system, for its subsequent transformation with the episomal plasmid containing the genes for HA synthesis, as it presents various advantages as host for the expression of heterologous DNA:

- the HA produced is easily secreted;
- it is free of exotoxins and endotoxins, unlike gram-negative bacteria;
- it does not contain any alkaline protease, and consequently does not induce the breakdown of the protein produced;
- it is structurally very stable by comparison with recombinant plasmids: *B. megaterium* can contain a much larger number of episomal plasmids than *Bacillus Subtilis*, which are more stable; *B. megaterium* can also support much larger inserts than *E. coli* and *B. subtilis*, and this characteristic is very important when, as in the case of the present invention, a long metabolic pathway like that of hyaluronic acid is to be engineered.

The T7 RNA polymerase system transferred to *B. megaterium* (and to *E. coli*, preferably to *E. coli* BL21 DE3 strain)

controls the expression of the genes responsible for synthesis of the HA biosynthesis pathway (cloned in episomal plasmids), and guarantees

very high activity and selectivity of gene transcription; a consequent very high production of the recombinant proteins required for the synthesis of hyaluronic acid.

The final yield of the desired product will be very high: much higher than that obtained with *B. subtilis*, where the operon system is cloned on the chromosome of the bacterium, and is under the control of non-inducible constitutive promoters.

In fact, the T7 RNA polymerase system described above is inducible: it is introduced artificially into the bacterium and activated by the Applicant by adding substances like IPTG (for *E. coli* in quantities of between 0.1 mM and 10 mM, preferably between 0.4 and 1 mM) or xylose (for *B. Megaterium* in quantities of between 0.1% and 10%, preferably between 0.5% and 1% w/v); in their presence, the inducer bonds to the repressor, modifying its configuration, and the repressor then detaches from the promoter, allowing the polymerases of the bacterium to transduce the gene for synthesis of T7 RNA polymerase. The latter, in turn, can only activate the gene transcription of the genes placed under the control of a T7 promoter. In this way the synthesis of the whole biosynthesis process for the production of HA can be controlled. The system is so efficient in that a single polymerase is dedicated to the gene of interest, and the RNA polymerase of the bacterium is not involved. With this methodology, the cell protein synthesis system is saturated, so that the proteins of interest are obtained in amounts to 50% or more of the total proteins.

Further, as demonstrated in the following by the Applicant, by modulating the fermentation times, the Applicant can obtain the production of high amounts of HA with specific weight average molecular weights, comprised in a range of from 100 KD to above 2 MD. More particularly, when the process according to the invention uses bacterial host cells of *B. megaterium* and fermentation time is comprised of from 80 to 160 hours, it is possible to obtain HA having a weight average MW comprised in the range 100-500 KD; when fermentation time is comprised of from 40 to 80 hours, it is possible to obtain HA having a weight average MW comprised in the range 500-1000 KD; when fermentation time is comprised of from 12 to 40 hours, it is possible to obtain HA having a weight average MW comprised in the range $1 \times 10^6 - 3 \times 10^6$ D.

In a preferred embodiment of the present invention, the sequence coding for the enzyme hyaluronan synthase (hasA) is obtained from a *Streptococcus* strain, preferably from *Streptococcus zooepidemicus*, and the sequences coding for enzymes UDP-glucose dehydrogenase (hasB or tuaD), UDP-glucose pyrophosphorylase (hasC or gtaB) and glucose 6 phosphate isomerase (hasE or pgi), are derived from *B. subtilis*.

According to a particularly preferred embodiment of the present invention, the sequences coding for enzymes hyaluronan synthase, UDP-glucose dehydrogenase, UDP-glucose pyrophosphorylase and glucose 6 phosphate isomerase include an upstream Shine-Dalgarno sequence.

Even more preferably, said plasmid vector (i) comprises or consists of the nucleotide sequence as defined in SEQ ID NO:1 or in SEQ ID NO:2.

The subsequent purification of the HA secreted will be extremely simple, with the result that the industrial production process will be much cheaper than the process according to the state of the art.

Specifically, *E. coli* strains BL21 DE3 (Stratagene, Calif., USA) have T7 RNA polymerase cloned in the chromosome of the bacterium under the control of the inducible promoter lac. It can be induced with IPTG for the transcription of the T7 RNA polymerase gene. At this point, the T7 RNA polymerase produced can transcribe the genes under its control.

A similar system has also been engineered in *B. megaterium*. In this case the system uses two plasmids: the first leads to the synthesis of the enzymatic protein T7 RNA polymerase, and the second (engineered) to that of the messenger of the gene (or genes) of interest, under the control of the T7 promoter of bacteriophage T7. The first plasmid, pT7-RNAP (MoBiTec), derives from plasmid pBM100 264 (MoBiTec), which replicates in *B. megaterium* QM B1551 (MoBiTec) and also contains the replication origin of *E. coli*, resistance to ampicillin and chloramphenicol and the promoter for xylose PXY1A, and its repressor Xy1R, which control the synthesis of T7 RNA polymerase, whose gene sequence is in the same plasmid. The plasmid for synthesis of recombinant proteins, pPT7 (MoBiTec), derives from *B. cereus* and leads to a replication origin of *B. megaterium* and resistance to ampicillin and chloramphenicol, and a replication origin for *E. coli* and the promoter T7 controlled by T7 RNA polymerase.

When the protein of interest is to be synthesised, xylose is added to the cells, and activates its promoter by detaching the repressor. The promoter, freed, then allows the polymerase of the bacterium to transcribe the gene for synthesis of the T7 RNA polymerase enzyme which, moving onto the T7 promoter of the other plasmid, transcribes its gene of interest, namely the genes required for HA synthesis. The system is highly efficient, because a single polymerase is dedicated to the transcription of the gene of interest, and the multiple copies of the two plasmids ensure that the transcript levels are extremely high.

A further object of the present invention are plasmid vectors, containing the two genes hasA and tuaD or the four genes hasA, tuaD, gtaB and pgi (corresponding to hasE), under the control of T7 promoter of RNA polymerase of bacteriophage T7, which are suitable to allow the production in *B. megaterium* and/or in *E. coli*, preferably in *B. megaterium*, of hyaluronic acid in high yield, according to the methodology described above. Preferably, the sequences coding for the hyaluronan synthase enzyme, UDP-glucose dehydrogenase, UDP-glucose pyrophosphorylase and glucose 6 phosphate isomerase include an upstream Shine-Dalgarno sequence. These vectors can also be constructed so as contain any other gene relating to the biosynthesis of hyaluronic acid.

Unlike those available to date, the starting plasmid is small, which allows engineering of the entire hyaluronic acid biosynthesis pathway (i.e. the four genes hasA, tuaD, gtaB and pgi) in a single plasmid, which is herein referred to as pPT7hasAtuaDgtaBpgi, making the present invention economically advantageous and successfully applicable on an industrial scale. In a preferred embodiment of the present invention the plasmid vector is pPT7hasAtuaD (SEQ ID NO:1) or pPT7hasAtuaDgtaBpgi (SEQ ID NO:2).

The present invention also relates to a method and relative system for the production/construction of bacterial strains, transformed with plasmid containing the entire hyaluronic acid biosynthesis pathway, with the 2 genes or 4 genes, and the selection of stable, viable, replicating and HA-secreting bacterial strains with high yield.

Said method of construction of engineered strain with the 2 genes or 4 genes plasmid vector for the HA synthesis comprises the following steps:

Cloning of the *tuaD* gene (UDP-glucose dehydrogenase) from *Bacillus Subtilis*,

Cloning of the *hasA* gene (hyaluronan synthase) from *Streptococcus zooepidemicus*,

Construction of the plasmid pGEM4hasA,

Construction of a plasmid with the *tuaD* gene following *hasA*,

Cloning of the *hasA-tuaD* gene in the plasmid for *B. megaterium* pPT7: pPT7hasAtuaD;

the process for the construction of the 4 genes route proceeds with the following steps:

Cloning of the *gtaB* gene: construction of the plasmid pGEM4hasA-gtaB,

Cloning of the *pgi* gene from *Bacillus Subtilis*,

Construction of plasmid pPT7hasAtuaDgtaBpgi, which is referred to as pT7hyal,

Transformation of plasmids pPT7hasAtuaD and pPT7hasAtuaDgtaBpgi into *Bacillus megaterium* or *E. coli*, preferably in *Bacillus megaterium*,

Selection of hyaluronic acid-secreting cells by xylose gradient for *Bacillus megaterium* or IPTG gradient for *E. coli*,

Selection of stable, viable, replicant and secreting high amounts of HA cells.

A further object of the invention is therefore a system for the selection of transfected, secreting, viable cells: the IPTG gradient allows the selection of transfected, viable cells, capable of replication and above all, secreting HA with high yields.

The present invention will be now disclosed by way of example but not of limitation, according to preferred embodiments with particular reference to the attached figures, wherein:

FIG. 1 shows a comparison in plates between the growth of cells *E. coli* TOP10, incorporating plasmid pHT01 (control) and cells *E. coli* TOP10, incorporating pBS5 (*hasA+tuaD*);

FIG. 2 shows the gel analysis of the expression of gene *tuaD* in *E. coli* BL21 DE3;

FIG. 3 shows the analysis in gel electrophoresis of the constitutive expression of hyaluronan synthase (Street) in *E. coli*; the encoded protein designated SeHAS is 417 amino acids long (calculated molecular weight 47,778; calculated PI 9.1) and is the smallest member of the HAS family identified thus far; the enzyme migrates anomalously fast in SDS polyacrylamide gel electrophoresis (about 42000 Da);

FIG. 4 shows the plasmid map pPT7 comprising the promoter and the terminator of T7 RNA polymerase of bacteriophage T7; the replication origin of *Coli* and *Megaterium*; ampicillin resistance gene; tetracycline resistance gene;

FIG. 5 shows the restriction map of plasmid PT7hyal;

FIG. 6 shows the analysis by SDS-page of cell lysates of *E. coli* BL21 DE3 to verify the presence of proteins that lead to the synthesis of hyaluronic acid;

FIG. 7 shows the comparison of the production of HA in plate between colonies of *E. coli* BL21 DE3 transformed with the plasmid pPT7 (colony control), pPT7hasAtuaD (colony 6) and pPT7hasAtuaDgtaBpgi (pT7Hyal—Colony 2) after 24 hours of growth at 37° C., in the presence of IPTG;

FIG. 8 shows the results of plating assays for the selection of cells able to express high levels of hyaluronic acid in the presence or absence of IPTG;

FIG. 9 shows the results of plating assays in the presence of IPTG to test the degree of survival of cells capable of producing HA;

FIG. 10 shows the carbazole analysis of the precipitates of HA in the test tube.

The following examples describe the various steps required for the embodiment of the invention, by way of example but not of limitation.

Example 1

Cloning of the *tuaD* Gene (UDP-Glucose Dehydrogenase) from *Bacillus subtilis*

The sequence of the *tuaD* gene, which is 9300 bp long in *B. subtilis*, is present in the databases under the access number AF015609 in the system which codes for the teichuronic acid operon and comprises eight genes, *tuaA*-*CDEFGH*. In the present case the gene of interest *tuaD* falls between bases 3582-4984 bp. Software analysis for restriction enzymes indicates that the restriction sites *ClaI*, *EcoRI*, *PstI*, *HindIII* and *SphI* are present, and therefore cannot be used for cloning. The start codon is not a methionine but a valine; in the present invention, it was replaced with the codon for methionine, which is much more efficient in the transduction of the protein. Two oligonucleotide primers with the following sequence were used to recover this sequence:

(SEQ ID NO: 3)
5'atgaaaaaatagctgtcattggaacag 3'
and

(SEQ ID NO: 4)
5'ttataaattgctggtcccaagtct 3'.

The genomic DNA from *B. subtilis* strain 168 (ACTT 23857D-5) was obtained with the Qiagen extraction kit. With 32 cycles of PCR, using DNA from *B. subtilis* as template and the two said oligonucleotides, an amplificate of the expected molecular weight was obtained. The amplificate obtained was tested for the presence of restriction enzyme *EcoRI*. After cutting with this enzyme in 1% agarose gel, two bands of DNA weighing 470 bp and 920 bp were present, which correspond to those expected. To clone the *tuaD* gene in an expression vector, two other oligonucleotides with the following sequence were synthesised:

(SEQ ID NO: 5)
5'gctggatccatgaaaaaatagctgtcattgg 3'
and

(SEQ ID NO: 6)
5'ctcgtagcttataaattgacgcttcccaag 3'

in order to insert said sequence between the restriction sites *BamHI* and *NheI* in the expression vector, plasmid pRSET B (INVITROGEN).

A Shine-Dalgarno (SD) sequence needs to be introduced into gene *tuaD* upstream of the 5' end of the gene to allow efficient recognition by the bacterial RNA polymerase. For this purpose the DNA was amplified with the following oligonucleotide primers:

(SEQ ID NO: 7)
5'cgacatatgaaaaaatagctgtcattgg 3'

-continued

and

(SEQ ID NO: 8)
5' ctcgctagcttataaattgacgcttcccaag 3'.

Two restriction sites NdeI and NheI are present in said primers at 5', which allow their cloning in vector pRSET B between the same sites. A sequence SD, consequently present upstream of restriction site NdeI of plasmid pRSET B, is particularly efficient and necessary for the RNA polymerase in order to synthesise the protein. Restriction site XbaI, which will be required for the subsequent clonings, is also present even before said sequence. The vector created, pRSET B, was therefore called pRSETuaD.

Thus in this plasmid, the sequence coding for tuaD falls between restriction sites NdeI and NheI; restriction site XbaI, which is necessary for the subsequent cloning, is present before and upstream of said plasmid, and other restriction sites, including BamHI--BglII--XhoI, are present behind the tuaD gene.

The diagram below summarises the sites of interest present in plasmid pRSETuaD

XbaI--NdeI-----tuaD-----NheI--
BamHI--BglII-XhoI

The plasmid described is an expression vector functioning not only in *B. megaterium* but also in *E. coli*, because the gene is under the control of T7 promoter of bacteriophage T7; if it is transformed into bacterial cells BL21 DE3, which are able to transcribe T7 RNA polymerase, it therefore enables them to express the tuaD gene. After induction with 1 mM of IPTG the cells in *E. coli* are able to produce the protein of the expected molecular weight, but not hyaluronic acid. The construction is particularly efficient because the level of expression is very high. The sizes of the colonies which carry plasmid pRSETuaD are tiny compared with the control cells (FIG. 1), which demonstrates the toxicity of the tuaD gene. This cloning is difficult precisely because it is apparently difficult for the colonies to grow; the particularly high level of enzyme UDP-glucose dehydrogenase probably drains the cell glucose because it is required for the formation of the hyaluronic acid precursor. The cells in which the synthesis of tuaD is induced with IPTG are therefore no longer able to survive for a long time, so the gene product is toxic.

In conclusion, the tuaD gene was isolated and cloned in a plasmid, and the sequence proved correct. The gene expressed in *E. coli* is able to produce a protein of the expected molecular weight (54 kDa, FIG. 2), which is toxic to the cell. However, these cells are unable to produce significant quantities of hyaluronic acid, as hyaluronan synthase (hasA) is lacking.

Example 2

Cloning of the hasA (Hyaluronan Synthase) Gene from *Streptococcus zooepidemicus*

The gene sequence for hyaluronan synthase is present in the databases under the access number AY173078, and is 3552 bp long; the sequence coding for the protein is between bases 1 and 1254. The restriction sites HindIII and StuI are present in this sequence, and therefore cannot be used for cloning, but can be used to verify the cloning. Two oligonucleotides for use with PCR were designed and synthesised to recover the coding sequence:

(SEQ ID NO: 9)
5'atgagaacattaaaaaacctcataac 3'
and

(SEQ ID NO: 10)
5'taataatTTTTTACGTGTTCCCCAG 3'

The genomic DNA from the bacterium *Streptococcus zooepidemicus* was recovered with the Qiagen extraction kit. The 1254 bp coding sequence was recovered with PCR. The expected amplicate of the correct dimensions was controlled with restriction enzyme HindIII, and gave rise to two bands of approx. 100 bp and 1150 bp which correspond to the expected cut.

Example 3

Construction of the Plasmid pGEM4hasA

Two other oligonucleotides with the following sequence were created to clone the hasA sequence in plasmid pGEM4Z:

(SEQ ID NO: 11)
5' ggaggatccatgagaacattaaaaaacctcat 3'
and

(SEQ ID NO: 12)
5' cagtctagattataataatTTTTTACGTGTCC 3'

The BamHI restriction site was created in the first oligonucleotide close to 5', and the XbaI restriction site was created in the second oligonucleotide, again at 5'. The amplicate obtained through these two oligonucleotides was cloned between restriction sites BamHI and XbaI in plasmid pGEM4Z (PROMEGA) between the same sites to give plasmid pGEM4hasA.

The DNA sequence between said two restriction sites was analysed with an ABI 7000 sequencer, proved correct, and is identical to the one published.

HindIII-BamHI-----hasA-----XbaI-Sall

The plasmid was checked for expression of the recombinant protein in *E. coli* and presented a molecular weight of approx. 42 kDa, which agrees with the weight reported for the protein in the literature, although it has a theoretical molecular weight of 47.778 kDa (FIG. 3).

The cloning of hasA from *streptococcus* was therefore also demonstrated in terms of protein expression. The plasmid is unable to produce significant quantities of hyaluronic acid because it lacks the tuaD gene.

Example 4

Construction of a Plasmid with the tuaD Gene Following hasA

With this construction, the hasA gene is placed in tandem with the tuaD gene. For this purpose, plasmid pGEM4hasA, which already contains the hasA gene, is used as vector. The plasmid was cut with XbaI and Sall, and the tuaD gene sequence from plasmid pRSETuaD was cut with XbaI and XhoI and cloned in the same sites (Xho I and Sall are compatible)

pGEM4hasA
HindIII-BamHI-----hasA-----XbaI-Sall
pRSE tuaD
XbaI--NdeI-----tuaD-----NheI--
BamHI--BglII-XhoI

11

the following final sequence being obtained:
HindIII-BamHI-----hasA-----XbaI--NdeI-----
tuaD-----NheI-BamHI--BglI-XhoI

Example 5

Cloning of the hasA-tuaD Gene in the Plasmid
pPT7 for *B. megaterium*

This plasmid pPT7 (MoBiTec) contains two origins of replication, one for *E. coli* and one for *B. megaterium*, and can therefore be propagated in both bacteria. It also contains resistance to the antibiotics ampicillin and tetracycline, which can be used for *E. coli* and *B. megaterium* respectively, and the recognition sequence for T7 RNA polymerase, namely the promoter dependent on T7 RNA polymerase of bacteriophage T7 followed by its terminator.

The plasmid contains restriction site BsrGI with the sequence tgtaca a few bases after the Shine-Dalgarno

12

sequence, and a site BamHI (ggatcc) after the initial methionine. Two oligonucleotides were synthesised for the cloning so as to create the following two restriction sites at the end:

(SEQ ID NO: 13)
5'GCTTGATACATGAGAACATTAAAAACCTCA 3'

(SEQ ID NO: 14)
5'AGGGATCCTTATAAATTGACGCTTCCCAAG 3'

i.e. BsrGI and BamHI upstream and downstream of genes hasA and tuaD respectively. The 2698 bp amplicon obtained was cut with the restriction enzymes BsrGI and BamHI and cloned in the same restriction sites as plasmid pPT7 to obtain plasmid pPT7hasAtuaD (FIG. 4).

The complete sequence of this plasmid, called pPT7hasAtuaD, was analysed, and is set out below:

0 CTTTTAGGTTCTAAATCGTGTCTTTCTTGAATTGTGCTGTTTATCCTTTACCTTGTC
60 TACAAACCCCTTAAAAACGTTTTTAAAGGCTTTTAAGCCGTCTGTACGTTTCCTTAAGGCC
120 AAATTAATACGACTCACTATAGGGAGACCACAACGGTTTCCGAATATTAATTAACCAAG

Bsp1407I

180 GAGGTGAAATGTACAATGAGAACATTAAAAACCTCATAACTGTTGTGGCCTTTAGTATT
1 M R T L K N L I T V V A F S I

HindIII

240 TTTTGGGTACTGTTGATTTACGTCAATGTTTATCTCTTTGGTGCTAAAGGAAGCTTGTC

1 F W V L L I Y V N V Y L F G A K G S L S

300 ATTTATGGCTTTTGGCTGATAGCTTACCTATTAGTCAAATGTCCTTATCCTTTTTTTAC

1 I Y G F L L I A Y L L V K M S L S F F Y

360 AAGCCATTTAAGGAAGGGCTGGGCAATATAAGGTTGCAGCCATTATTCCTCTTATAAC

1 K P F K G R A G Q Y K V A A I I P S Y N

420 GAAGATGCTGAGTCATTGCTAGAGACCTTAAAAAGTGTTCAGCAGCAAACCTATCCCCTA

1 E D A E S L L E T L K S V Q Q Q T Y P L

480 GCAGAAATTTATGTTGTTGACGATGGAAGTGTGATGAGACAGGTATTAAGCGCATTGAA

1 A E I Y V V D D G S A D E T G I K R I E

540 GACTATGTGCGTGACACTGGTGACCTATCAAGCAATGTCATTGTTACCGGTCAGAAAA

1 D Y V R D T G D L S S N V I V H R S E K

600 AATCAAGGAAAGCGTCATGCACAGGCCTGGGCCTTTGAAAGATCAGACGCTGATGCTTTT

1 N Q G K R H A Q A W A F E R S D A D V F

660 TTGACCGTTGACTCAGATACTTATATCTACCCTGATGCTTTAGAGGAGTTGTTAAAAACC

1 L T V D S D T Y I Y P D A L E E L L K T

720 TTTAATGACCCAACCTGTTTTTGTGCGACGGGTCACCTTAATGTCAGAAATAGACAAACC

1 F N D P T V F A A T G H L N V R N R Q T

780 AATCTCTTAACACGCTTGACAGATATTCGCTATGATAATGCTTTTGGCGTTGAACGAGCT

1 N L L T R L T D I R Y D N A F G V E R A

840 GCCCAATCCGTTACAGGTAATATTCTCGTTTGCTCAGGCCCGCTTAGCGTTTACAGACGC

-continued

1 A Q S V T G N I L V C S G P L S V Y R R
 900 GAGGTGGTTGTTCCCTAACATAGATAGATACATCAACCAGACCTTCCTGGGTATTCCTGTA
 1 E V V V P N I D R Y I N Q T F L G I P V
 960 AGTATCGGTGATGACAGGTGCTTGACCAACTATGCAACTGATTTAGGAAAGACTGTTTAT
 1 S I G D D R C L T N Y A T D L G K T V Y
 1020 CAATCCACTGCTAAATGTATTACAGATGTTCTGACAAGATGTCTACTTACTTGAAGCAG
 1 Q S T A K C I T D V P D K M S T Y L K Q
 1080 CAAAACCGCTGGAACAAGTCCTTCTTTAGAGAGTCCATTATTTCTGTTAAGAAAATCATG
 1 Q N R W N K S F F R E S T I S V K K I M
 1140 AACAATCCTTTTGTAGCCCTATGGACCATACTTGAGGTGTCTATGTTTATGATGCTTGTT
 1 N N P F V A L T I L E E V S M F M M L V
 1200 TATTCTGTGGTGGATTTCTTTGTAGGCAATGTCAGAGAATTTGATTGGCTCAGGGTTTGT
 1 Y S V V D F F V G N V R E F D W L R V L
 1260 GCCTTTCTGGTGATTATCTTCATTGTTGCTCTTTGTCGTAATATCACTATATGCTTAAG
 1 A F L V I I F I V A L C R N I H Y M L K
 1320 CACCCGCTGTCCTTCTTGTTATCTCCGTTTTATGGGGTACTGCTTTGTTTGTCTACAGC
 1 H P L S F L L S P F Y G V L L C L S Y S
 1380 CCTTGAAATGTATTCTCTTTTTACTATTAGAAATGCTGACTGGGGAACACGTAAAAAAT
 1 P

 XbaI NdeI
 1440 TATTATAATCTAGAAATAATTTTGTTTAACTTTAAGAAGGAGATATACATATGAAAAAAA
 3 M K K I
 1500 TAGCTGTCATTGGAACAGGTTATGTAGGACTCGTATCAGGCACTTGCTTTGCGGAGATCG
 3 A V I G T G Y V G L V S G T C F A E I G

 EcoRV ClaI
 1560 GCAATAAAGTTGTTTGTGTGATATCGATGAATCAAAAATCAGAAGCCTGAAAAATGGGG
 3 N K V V C C D I D E S K I R S L K N G V
 1620 TAATCCCAATCTATGAACCAGGGCTTGACAGACTTAGTTGAAAAAATGTGCTGGATCAGC
 3 I P I Y E P G L A D L V E K N V L D Q R

 EcoRV
 1680 GCCTGACCTTTACGAACGATATCCCGTCTGCCATTGCGGCCTCAGATATTATTATATTG
 3 L T F T N D I P S A I R A S D I I Y I A
 1740 CAGTCGGAACGCCTATGTCCAAAACAGGTGAAGCTGATTTAACGTACGTCAAAGCGGCGG
 3 V G T P M S K T G E A D L T Y V K A A A
 1800 CGAAAACAATCGGTGAGCATCTTAACGGCTACAAAAGTGATCGTAAATAAAAGCACAGTCC
 3 K T I G E H L N G Y K V I V N K S T V P
 1860 CGGTTGGAACAGGGAAACTGGTGCATCTATCGTTCAAAAAGCCTCAAAGGGGAGATACT
 3 V G T G K L V Q S I V Q K A S K G R Y S

 EcoRI
 1920 CATTTGATGTTGTATCTAACCTGAATTCCTTCGGGAAGGGTCAGCGATTCATGACACGA

-continued

3 F D V V S N P E F L R E G S A I H D T M
1980 TGAATATGGAGCGTGCCGTGATTGGTTCAACAAGTCATAAAGCCGCTGCCATCATTGAGG

3 N M E R A V I G S T S H K A A A I I E E
2040 AACTTCATCAGCCATTCCATGCTCCTGTCAATTAACAAACCTAGAAAGTGCAGAAATGA

3 L H Q P F H A P V I K T N L E S A E M I
EcoRV

2100 TTAAATACGCCGGAATGCATTTCTGGCGACAAAGATTTCTTTATCAACGATATCGCAA
3 K Y A A N A F L A T K I S F I N D I A N

2160 ACATTTGTGAGCGAGTCGGCGCAGACGTTTCAAAAGTTGCTGATGGTGTGGTCTTGACA
3 I C E R V G A D V S K V A D G V G L D S

2220 GCCGTATCGGCAGAAAGTTCTTAAAGCTGGTATTGGATTTCGGCGGTTTCATGTTTTCCAA
3 R I G R K F L K A G I G F G G S C F P K

2280 AGGATACAACCGCGCTGCTTCAAATCGAAAATCGGCAGGCTATCCATTCAAGCTCATCG
3 D T T A L L Q I A K S A G Y P F K L I E

2340 AAGCTGTCATTGAAACGAACGAAAAGCAGCGTGTTCATATTGTAGATAAACTTTTGACTG
3 A V I E T N E K Q R V H I V D K L L T V

2400 TTATGGGAAGCGTCAAAGGGAGAACCATTTTCAGTCCTGGGATTAGCCTTCAAACCGAATA
3 M G S V K G R T I S V L G L A F K P N T

PstI

2460 CGAACGATGTGAGATCCGCTCCAGCGCTTGATATTATCCCAATGCTGCAGCAGCTGGGCG
3 N D V R S A P A L D I I P M L Q Q L G A

HindIII

2520 CCCATGTAAAAGCATAACGATCCGATTGCTATTCTGAAGCTTCAGCGATCCTTGGCGAAC
3 H V K A Y D P I A I P E A S A I L G E Q

SphI

2580 AGGTCGAGTATTACACAGATGTGTATGCTGCGATGGAAGACACTGATGCATGCCTGATTT
3 V E Y Y T D V Y A A M E D T D A C L I L

2640 TAACGGATTGGCCGGAAGTGAAAGAAATGGAGCTTGTAAGTGAACCCCTTAAAC
3 T D W P E V K E M E L V K V K T L L K Q

2700 AGCCAGTCATCATTGACGGCAGAAATTTATTTTCACTTGAAGAGATGCAGGCAGCCGGAT
3 P V I I D G R N L F S L E E M Q A A G Y

2760 ACATTTATCACTCTATCGGCCGTCCCGCTGTTTCGGGGAACGGAACCCCTGACAAGTATT
3 I Y H S I G R P A V R G T E P S D K Y F

BamHI

2820 TTCCGGGCTTGCCGCTTGAAGAATTGGCTAAAGACTTGGGAAGCGTCAATTTATAAGGAT
3 P G L P L E E L A K D L G S V N L

SphI

(SEQ ID NO: 1)

2880 CCGGCCGATGCCGGCTAATCGCGACCGGTTAACTAGCATAACCCCTTGGGGCCTCTAAA
2940 CGGGTCTTGAGGGGTTTTTGTAAAGGAGGAACCTATATCCGGTCCAAGAATTGGAGCCA

3000 ATCAATTCTGCGGAGAACTGTGAATGCGCAAACCAACCCTTGGCAGAACATATCCATCG

-continued

3060 CGTCCGCCATCTCCAGCAGCCGCGACGCGGCATCTCGGGCCGCGTTGCTGGCGTTTTTC
 3120 CATAGGCTCCGCCCCCTGACGAGCATCACAAAAATCGACGCTCAAGTCAGAGGTGGCGA
 3180 AACCCGACAGGACTATAAAGATACCAGGCGTTTCCCCCTGGAAGCTCCCTCGTGCCTCT
 3240 CCTGTTCCGACCCTGCCGCTTACCGGATACCTGTCCGCCTTCTCCCTTCGGGAAGCGTG
 3300 GCGCTTTCTCATAGCTCACGCTGTAGGTATCTCAGTTCGGTGTAGGTCGTTTCGCTCCAAG
 3360 CTGGGCTGTGTGCACGAACCCCCGTTTCAGCCCGACCGCTGCGCCTTATCCGGTAACAT
 3420 CGTCTTGAGTCCAACCCGGTAAGACGACTTATCGCCACTGGCAGCAGCCACTGGTAAC
 3480 AGGATTAGCAGAGCGAGGTATGTAGGCGGTGCTACAGAGTCTTGAAGTGGTGGCCTAAC
 3540 TACGGCTACACTAGAAGGACAGTATTTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTTC
 3600 GGAAAAAGAGTTGGTAGCTCTTGATCCGGCAAACAAACCACCGCTGGTAGCGGTGGTTTT
 3660 TTTGTTTGAAGCAGCAGATTACGCGCAGAAAAAAGGATCTCAAGAAGATCCTTTGATC
 3720 TTTTCTACGGGGTCTGACGCTCAGTGAACGAAAACCTCACGTTAAGGGATTTTGGTCATG
 3780 AGATTATCAAAAAGGATCTTACCCTAGATCCTTTTAAATTAATAAATGAAGTTTAAATCA
 3840 ATCTAAAGTATATATGAGTAAACTTGGTCTGACAGTTACCAATGCTTAATCAGTGAGGCA
 3900 CCTATCTCAGCGATCTGTCTATTTCTGTTTCATCCATAGTTGCCCTGACTCCCCGTCGTGTAG
 3960 ATAACCTACGATACGGGAGGGCTTACCATCTGGCCCCAGTGCTGCAATGATACCGCGAGAC
 4020 CCACGCTCACCGGCTCCAGATTTATCAGCAATAAACAGCCAGCCGGAAGGGCCGAGCGC
 4080 AGAAGTGGTCTGCAACTTTATCCGCCTCCATCCAGTCTATTAATTGTTGCCGGAAGCT
 Pst I
 4140 AGAGTAAGTAGTTCGCCAGTTAATAGTTTGGCGAACGTTGTTGCCATTGCTGCAGGCATC
 Hpy99 I
 4200 GTGGTGTACGCTCGTCGTTTGGTATGGCTTCATTTCAGCTCCGGTTCCTAACGATCAAGG
 4260 CGAGTTACATGATCCCCATGTTGTGCAAAAAGCGGTTAGCTCCTTCGGTCCCTCGATC
 4320 GTTGTGAGAAGTAAGTTGGCCGCGAGTGTATCACTCATGGTTATGGCAGCACTGCATAAT
 Sca I
 4380 TCTCTTACTGTGTCATGCCATCCGTAAGATGCTTTTCTGTGACTGGTGAGTACTCAACCAAG
 4440 TCATTCTGAGAATAGTGTATGCGGCGACCGAGTTGCTCTTGCCCGGCGTCAACACGGGAT
 4500 AATACCGCGCCACATAGCAGAACTTTAAAAGTGCTCATCATTTGAAAAACGTTCTTCGGGG
 4560 CGAAAACCTCAAGGATCTTACCGCTGTTGAGATCCAGTTCGATGTAACCCACTCGTGCA
 4620 CCCAACTGATCTTCAGCATCTTTTACTTTTACCAGCGTTTCTGGGTGAGCAAAAACAGGA
 4680 AGGCAAAATGCCGCAAAAAGGGAATAAGGGCGACACGGAAATGTTGAATACTCATACTC
 4740 TTCCTTTTCAATATTATTGAAGCATTTATCAGGGTTATTGTCTCATGAGCGGATACATA
 4800 TTTGAATGATTTAGAAAAATAAACAAATAGGGGTTCCGCGCACATTTCCCGAAAAGTG
 4860 CCACCTGACGTCTAAGAAACCATTATTTATCATGACATTAACCTATAAAAATAGGCGTATC
 EcoRI
 4920 ACGAGGCCCTTTCGTCTTCAAGAAATCCTGTTATAAAAAAGGATCAATTTTGAACCTC
 4980 TCCCAAAGTTGATCCCTTAACGATTTAGAAATCCCTTTGAGAATGTTTATATACATTCAA
 —
 5040 GGTAACCAGCCAATAATGACAATGATTCCTGAAAAAGTAATAACAAATTACTATACAG
 5100 ATAAGTTGACTGATCAACTTCCATAGGTAACAACCTTTGATCAAGTAAGGGTATGGATAA
 5160 TAAACCACCTACAATTGCAATACCTGTTCCCTCTGATAAAAAGCTGGTAAAGTTAAGCAA

-continued

5220 ACTCATTCCAGCACCAGCTTCCTGCTGTTTCAAGCTACTTGAAACAATTGTTGATATAAC
 5280 TGTTTTGGTGAACGAAAGCCCACCTAAAACAATACGATTATAATTGTCATGAACCATGA
 5340 TGTTGTTTCTAAAAGAAAGGAAGCAGTTAAAAGCTAACGAAAGAAATGTAACCTCCGAT
 5400 GTTTAACACGTATAAAGGACCTCTTCTATCAACAAGTATCCCACCAATGTAGCCGAAAAT

ScaI

5460 AATGACACTCATTGTTCCAGGGAAAATAATTACACTTCCGATTTCCGGCAGTACTTAGCTG
 5520 GTGAACATCTTTCATCATATAAGGAACCATAGAGACAAACCTGCTACTGTTCCAAATAT
 5580 AATCCCCCACAAAGAACTCCAATCATAAAAGGTATATTTTCCCTAATCCGGGATCAAC
 5640 AAAAGGATCTGTTACTTTCTGATATGTTTTACAAATATCAGGAATGACAGCACGCTAAC
 5700 GATAAGAAAAGAAATGCTATATGATGTTGTAAACAACATAAAAAATACAATGCCTACAGA

EcoRV

5760 CATTAGTATAAATTCCTTTGATATCAAAATGACCTTTTATCCTTACTTCTTTCTTTAATAA
 5820 TTTCATAAGAAACGGAACAGTGATAATTGTTATCATAGGAATGAGTAGAAGATAGGACCA
 5880 ATGAATATAATGGGCTATCATTCCACCAATCGCTGGACCGACTCCTTCTCCCATGGCTAC

ClaI

5940 TATCGATCCAATAAGACCAAATGCTTTACCCCTATTTTCTTTGGAATATAGCGCGCAAC
 6000 TACAACCATACGAGTGCTGGAAATGCAGCTGCACCAGCCCTTGAATAAACGAGCCAT
 6060 AATAAGTAAGGAAAAGAAAGAATGGCCAACAACCCAATTACCGACCCGAAACAATTTAT
 6120 TATAATTCAAATAGGAGTAACCTTTGATGCCTAATTGATCAGATAGCTTTCCATATAC
 6180 AGCTGTTCCAATGGAAAAGGTTAACATAAAGGCTGTGTTACCCAGTTTGTACTCGCAGG
 6240 TGGTTTATTAATAATCATTGCAATATCAGGTAATGAGACGTTCAAAACCATTTCAATTA
 6300 TACGCTAAAAAAGATAAAATGCAAAGCCAAATTAATAATTTGGTTGTGTCGTAAATTCGA
 6360 TTGTGAATAGGATGTATTACATTTACCCCTCCAATAATGAGGGCAGACGTAGTTTATAG
 6420 GGTAAATGATACGCTTCCCTCTTTAATTGAACCCGTGTACATTATTACACTTCATAAT
 6480 TAATTCCTCCTAAACTTGATTAACATTTTACCACATATAAACTAAGTTTTAAATTCAG
 6540 TATTTCACTACTTATAACAATATGGCCCGTTTGTGAACTACTCTTTAATAAAATAAT
 6600 TTTTCCGTTCCCAATTCACATTGCAATAATAGAAAATCCATCTTCATCGGCTTTTTCGT
 6660 CATCATCTGTATGAATCAAATCGCCTTCTTCTGTGTCATCAAGGTTAATTTTTTATGTA
 6720 TTTCTTTAACAACCACCATAGGAGATTAACCTTTTACGGTGTAACCTTCTCCAAAT
 6780 CAGACAAACGTTTCAAATCTTTTCTTCATCATCGGTCATAAAATCCGTATCCTTTACAG
 6840 GATATTTGCAGTTTCGTCAATTGCCGATTGTATATCCGATTATATTTATTTTTCGGTC
 6900 GAATCATTTGAACTTTTACATTTGGATCATAGTCTAATTTCAATGCTTTTTCCAAAAT
 6960 GAATCCATTGTTTTGATTCACGTAGTTTTCTGTATTCTTAAAAAAGTTGGTTCCACAC
 7020 ATACCAATACATGCATGTGCTGATATAAGAATTATCTTTATTTTATTGTCACTTCCG
 7080 TTGCACGCATAAAACCAACAAGATTTTTATTAATTTTTTATATGCATCATTCCGGCGAA
 7140 ATCCTTGAGCCATATCTGACAACTCTTATTTAATTCCTCGCCATCATAAACATTTTTAA
 7200 CTGTTAATGTGAGAAACAACCAACGAAGTGTGGCTTTTGTTTAATAACTTCAGCAACAA
 7260 CCTTTTGTGACTGAATGCCATGTTTCATTGCTCTCCTCCAGTTGCACATTGGACAAAGCC
 7320 TGGATTTACAAAACCACTCGATACAACCTTTCTTTTCGCTGTTTCACGATTTTGTATT
 7380 ACTCTAATATTTACAGACAATCTTTTACTCTTTACGCTTTTTAAATTCAGAATATGCA

-continued

7440 GAAGTTCAAAGTAATCAACATTAGCGATTTTCTTTTCTCTCCATGGTCTCACTTTTCCAC
 7500 TTTTGTCTTGTCCTAAACCCCTTGATTTTTCATCTGAATAAATGCTACTATTAGGAC
 7560 ACATAATATTTAAAGAAACCCCATCTATTTAGTTATTTGTTTGGTCACTTATAACTTTA
 7620 ACAGATGGGGTTTTTCTGTGCAACCAATTTAAGGGTTTTCAATACTTTAAACACATAC
 7680 ATACCAACTTCAACGCACCTTTCAGCAACTAAAATAAAAAATGACGTTATTTCTATATG
 7740 TATCAAGATAAGAAAGAACAAGTTCAAAACCATCAAAAAAGACACCTTTTCAGGTGCTT
 7800 TTTTATTTTATAAACTCATTCCTGATCTCGACTTCGTTCTTTTTTACCTCTCGGTTA
 7860 TGAGTTAGTTCAAATTCGTT

The plasmid has a molecular weight of 7880 bp and contains the various genes responsible for hyaluronic acid synthesis under the control of strong T7 promoter of bacteriophage T7. The hasA synthase sequence from *Streptococcus equi* falls between bases 196 and 1383, and that of the tuaD gene between bases 1430 and 2873.

Example 6

Cloning of the gtaB Gene (UDP-Glc
Pyrophosphorylase)

The gtaB gene from *Bacillus Subtilis* was recovered from the bacterial genome as above, and through two oligonucleotides having the following sequence:

(SEQ ID NO: 15)
 5' ATGTCTAGAATAATAAGGAAGGTGCCTTTTAAATGAA 3'

(SEQ ID NO: 16)
 5' CTCTCGAGCTAGCTTAGATTTCTTCTTTGTTTAGTAAAG 3'

The amplified product of 925 bp was cut with XbaI and XhoI and cloned in plasmid pGEM4hasA in the same restriction sites; plasmid pGEMhasA-gtaB is obtained in this way.

Example 7

Cloning of the Pgi Gene from *Bacillus subtilis* in
Plasmid pRSET B

The pgi gene (glucose 6 phosphate isomerase, also called phosphoglucoisomerase pgi, corresponding to hasE from *S.*

zooepidemicus) was recovered from the bacterial genome as described above with these two oligonucleotides

(SEQ ID NO: 17)
 5' TACATATGACGCATGTACGCTTGACTACTCCAAAAG 3'

(SEQ ID NO: 18)
 5' ATGCTAGCTCATTTATAATCTTCCAGACGTTTTTCAAG 3'

and PCR, and cloned after cutting with restriction enzymes NdeI and NheI in plasmid pRSETB between the same restriction sites. Plasmid pRSEpgi is obtained in this way. It places the pgi gene under the control of a T7 promoter, and when it is transferred to cells of *E. coli* BL21 DE3 it produces the protein of the expected molecular weight. This plasmid was cut with XbaI and PstI, and the 1340 bp fragment was cloned in plasmid pGEMhasA-gtaB between sites NheI and PstI. Restriction site Xba, like NheI, is lost after cloning. In this way the pgi gene is placed behind the gtaB gene. The plasmid, called pGEM hasA-gtaB-pgi, was cut with XbaI and XhoI, and the fragment which contains the sequences coding for gtaB and pgi was cloned in plasmid pRSETuaD between the same sites. The plasmid obtained was called pRSETuaD-gtaB-pgi.

The latter was cut with XbaI and BamHI and the fragment which contains the sequence coding for tuaD, gtaB and pgi was cloned in plasmid pPT7hasAtuaD between the same sites to obtain plasmid pPT7hasAtuaDgtaBpgi, which we will call pT7hyal.

The sequence of plasmid pT7hyal is shown below

0 CTTTTTAGGTTCTAAATCGTGTCTTTTCTTGGAAATGTGCTGTTTTATCCTTTACCTTGTC
 60 TACAAACCCCTTAAAAACGTTTTTAAAGGCTTTTAAAGCCGTCTGTACGTTCCCTTAAGGCG
 120 AAATTAATACGACTCACTATAGGGAGACCACAACGGTTTCCCGAATATTAATTAACCAAG

Bsp1407I

180 GAGGTGAAATGTACAATGAGAACATTAAAAACCTCATAACTGTTGTGGCCTTTAGTATT

1 M R T L K N L I T V V A F S I

HindIII

240 TTTTGGTACTGTTGATTTACGTCAATGTTTATCTCTTTGGTGTCTAAAGGAAGCTTGTC

1 F W V L L I Y V N V Y L F G A K G S L S

300 ATTTATGGCTTTTGTGATAGCTTACCTATTAGTCAAAATGTCCTTATCCTTTTTTTTAC

1 I Y G F L L I A Y L L V K M S L S F F Y

-continued

360 AAGCCATTTAAGGGAAGGGCTGGGCAATATAAGGTTGCAGCCATTATTCCCTCTTATAAC
 1 K P F K G R A G Q Y K V A A I I P S Y N
 420 GAAGATGCTGAGTCATTGCTAGAGACCTTAAAAAGTGTTCAGCAGCAAACCTATCCCCTA
 1 E D A E S L L E T L K S V Q Q Q T Y P L
 480 GCAGAAATTTATGTTGTTGACGATGGAAGTGTGATGAGACAGGTATTAAGCGCATTGAA
 1 A E I Y V V D D G S A D E T G I K R I E
 540 GACTATGTGCGTGACACTGGTGACCTATCAAGCAATGTCATTGTTACCGGTCAGAAAAA
 1 D Y V R D T G D L S S N V I V H R S E K
 600 AATCAAGGAAAGCGTCATGCACAGGCCTGGGCCTTTGAAAGATCAGACGCTGATGTCTTT
 1 N Q G K R H A Q A W A F E R S D A D V F
 660 TTGACCGTTGACTCAGATACTTATATCTACCCTGATGCTTTAGAGGAGTTGTTAAAAACC
 1 L T V D S D T Y I Y P D L E E E L L K T
 720 TTTAATGACCCAACTGTTTTTGTGCGACGGGTACCTTAATGTCAGAAATAGACAAACC
 1 F N D P T V F A A T G H L N V R N R Q T
 780 AATCTCTTAACACGCTTGACAGATATTCGCTATGATAATGCTTTTGGCGTTGAACGAGCT
 1 N L L T R L T D I R Y D N A F G V E R A
 840 GCCCAATCCGTTACAGGTAATATTCTCGTTTGTCTCAGGCCCGCTTAGCGTTTACAGACGC
 1 A Q S V T G N I L V C S G P L S V Y R R
 900 GAGGTGGTTGTTCCCTAACATAGATAGATACATCAACCAGACCTTCTGGGTATTCCTGTA
 1 E V V V P N I D R Y I N Q T F L G I P V
 960 AGTATCGGTGATGACAGGTGCTTGACCAACTATGCAACTGATTTAGGAAAGACTGTTTAT
 1 S I G D D R C L T N Y A T D L G K T V Y
 1020 CAATCCACTGCTAAATGTATTACAGATGTTCTGACAAGATGTCTACTTACTTGAAGCAG
 1 Q S T A K C I T D V P D K M S T Y L K Q
 1080 CAAAACCGCTGGAACAAGTCCTTCTTTAGAGAGTCCATTATTTCTGTTAAGAAAATCATG
 1 Q N R W N K S F F R E S I I S V K K I M
 1140 AACAATCCTTTTGTAGCCCTATGGACCATACTTGAGGTGTCTATGTTTATGATGCTTGT
 1 N N P F V A L W T I L E V S M F M M L V
 1200 TATTCTGTGGTGGATTTCTTTGTAGGCAATGTCAGAGAATTTGATTGGCTCAGGGTTTTG
 1 Y S V V D F F V G N V R E F D W L R V L
 1260 GCCTTTCTGGTGATTATCTTCATTGTTGCTCTTTGTCGTAATATTCATATATGCTTAAG
 1 A F L V I I F I V A L C R N I H Y M L K
 1320 CACCCGCTGTCCTTCTTGTATCTCCGTTTTATGGGGTACTGCTTTGTTTGTCTACAGC
 1 H P L S F L L S P F Y G V L L C L S Y S
 1380 CCTTGAAATGTATTCTCTTTTTACTATTAGAAATGCTGACTGGGGAACACGTAAAAAAT
 1 P

XbaI

NdeI

1440 TATTATAATCTAGAAATAATTTGTTTAACTTTAAGAAGGAGATATACATATGAAAAAA
 3 M K K I

-continued

1500 TAGCTGTCATTGGAACAGGTTATGTAGGACTCGTATCAGGCACTTGCTTTGCGGAGATCG
 3 A V I G T G Y V G L V S G T C F A E I G

EcoRV

ClaI

1560 GCAATAAAGTTGTTTGTGTGATATCGATGAATCAAAAATCAGAAGCCTGAAAAATGGGG
 3 N K V V C C D I D E S K I R S L K N G V

1620 TAATCCCAATCTATGAACCAGGGCTTGCAGACTTAGTTGAAAAAATGTGCTGGATCAGC
 3 I P I Y E P G L A D L V E K N V L D Q R

EcoRV

1680 GCCTGACCTTTACGAACGATATCCCGTCTGCCATTCGGGCCTCAGATATTATTTATATTG
 3 L T F T N D I P S A I R A S D I I Y I A

1740 CAGTCGGAACGCCTATGTCCAAAACAGGTGAAGCTGATTTAACGTACGTCAAAGCGGCGG
 3 V G T P M S K T G E A D L T Y V K A A A

1800 CGAAAACAATCGGTGAGCATCTTAACGGCTACAAAAGTGATCGTAAATAAAAGCACAGTCC
 3 K T I G E H L N G Y K V I V N K S T V P

1860 CGGTTGGAACAGGGAAACTGGTGCAATCTATCGTTCAAAAAGCCTCAAAGGGGAGATACT
 3 V G T G K L V Q S I V Q K A S K G R Y S

EcoRI

1920 CATTTGATGTTGTATCTAACCCGTAATTCCTTCGGGAAGGGTCAGCGATTCATGACACGA
 3 F D V V S N P E F L R E G S A I H D T M

1980 TGAATATGGAGCGTGCCGTGATTGGTTCAACAAGTCATAAAGCCCTGCCATCATTGAGG
 3 N M E R A V I G S T S H K A A A I I E E

2040 AACTTCATCAGCCATTCCATGCTCCTGTCAATTAACAACCTAGAAAAGTGCAGAAATGA
 3 L H Q P F H A P V I K T N L E S A E M I

EcoRV

2100 TTAAATACGCCGGAATGCATTTCTGGCGACAAAGATTTCTTTATCAACGATATCGCAA
 3 K Y A A N A F L A T K I S F I N D I A N

2160 ACATTTGTGAGCGAGTCGGCGCAGACGTTTCAAAAAGTTGCTGATGGTGTGGTCTTGACA
 3 I C E R G A D D V S K V A D G V G L D S

2220 GCCGTATCGGCAGAAAGTTCCTTAAAGCTGGTATTGGATTCCGGCGGTTTCATGTTTTCCAA
 3 R I G R K F L K G I G G F G G S C F P K

2280 AGGATACAACCGCGCTGCTTCAAATCGCAAAATCGGCAGGCTATCCATTCAAGCTCATCG
 3 D T T A L L Q I A K S A G Y P F K L I E

2340 AAGCTGTCATTGAAACGAACGAAAAGCAGCGTGTTTCATATGTAGATAAACTTTTACTG
 3 A V I E T N E K Q R V H I V D K L L T V

2400 TTATGGGAAGCGTCAAAGGGAGAACCATTTTCAGTCCTGGGATTAGCCTTCAAACCGAATA
 3 M G S V K G R T I S V L G L A F K P N T

PstI

2460 CGAACGATGTGAGATCCGCTCCAGCGCTTGATATTATCCCAATGCTGCAGCAGCTGGGCG
 3 N D V R S A P A L D I I P M L Q Q L G A

HindIII

2520 CCCATGTAAAAGCATACGATCCGATTGCTATTCTGAAGCTTCAGCGATCCTTGGCGAAC
 3 H V K A Y D P I A I P E A S A I L G E Q

SphI

2580 AGGTCGAGTATTACACAGATGTGTATGCTGCGATGGAAGACACTGATGCATGCCTGATTT
 3 V E Y Y T D V Y A A M E D T D A C L I L

2640 TAACGGATTGGCCGGAAGTGAAGAAATGGAGCTTGAAAAGTGAAAACCTCTTAAAAC
 3 T D W P E V K E M E L V K V K T L L K Q

2700 AGCCAGTCATCATTGACGGCAGAAATTTATTTTCACTTGAAGAGATGCAGGCAGCCGGAT
 3 P V I I D G R N L F S L E E M Q A A G Y

2760 ACATTTATCACTCTATCGGCCGTCCTGTTTCGGGGAACGGAACCTCTGACAAGTATT
 3 I Y H S I G R P A V R G T E P S D K Y F

2820 TTCCGGGCTTGCCGCTTGAAGAATTGGCTAAAGACTTGGGAAGCGTCAATTTATAAGCTA
 3 P G L P L E E L A K D L G S V N L

2880 GAATAATAAGGAAGGTGCCTTTTAAATGAAAAAGTACGTAAAGCCATAATTCCAGCAGC
 2 M K K V R K A I I P A A

2940 AGGCTTAGGAACACGTTTTCTTCCGGCTACGAAAGCAATGCCGAAAGAAATGCTTCCTAT
 2 G L G T R F L P A T K A M P K E M L P I

3000 CGTTGATAAACCTACCATTCAATACATAATTGAAGAAGCTGTTGAAGCCGGTATTGAAGA
 2 V D K P T I Q Y I I E E A V E A G I E D

3060 TATTATTATCGTAACAGGAAAAAGCAAGCGTGCGATTGAGGATCATTTTGATTACTCTCC
 2 I I I V T G K S K R A I E D H F D Y S P

3120 TGAGCTTGAAAGAAACCTAGAAGAAAAAGGAAAACTGAGCTGCTTGAAAAAGTGAAAAA
 2 E L E R N L E E K G K T E L L E K V K K

3180 GGCTTCTAACCTGGCTGACATTCACTATATCCGCCAAAAAGAACCTAAAGGTCTCGGACA
 2 A S N L A D I H Y I R Q K E P K G L G H

3240 TGCTGTCTGGTGCGCACGCAACTTTATCGGCGATGAGCCGTTTGGGTTACTGCTTGGTGA
 2 A V W C A R N F I G D E P F A V L L G D

3300 CGATATTGTTTCAAGGCTGAAACTCCAGGGTTGCGCCAATTAATGGATGAATATGAAAAAC
 2 D I V Q A E T P G L R Q L M D E Y E K T

3360 ACTTTCTTCTATTATCGGTGTTTCAAGGTTGCGCGAAGAAGAAACACACCGCTACGGCAT
 2 L S S I I G V Q Q V P E E E T H R Y G I

3420 TATTGACCCGCTGACAAGTGAAGGCCCGGTTATCAGGTGAAAAACTTCGTTGAAAAACC
 2 I D P L T S E G R R Y Q V K N F V E K P

3480 GCCTAAAGGCACAGCACCTTCTAATCTTGCCATCTTAGGCCGTTACGTATTCACGCCTGA
 2 P K G T A P S N L A I L G R Y V F T P E

Bg1II

3540 GATCTTCATGTATTTAGAAGAGCAGCAGGTTGGCGCCGGCGAGAAATTCAGCTCACAGA
 2 I F M Y L E E Q Q V G A G G E I Q L T D

3600 CGCCATTCAAAGCTGAATGAAATCAAAGAGTGTGCTTACGATTTTGAAGCAAGCG
 2 A I Q K L N E I Q R V F A Y D F E G K R

-continued

3660 TTATGATGTTGGTGAAAAGCTCGGCTTTATCACAACAACCTTTGAATTTGCGATGCAGGA
 2 Y D V G E K L G F I T T T L E F A M Q D
 3720 TAAAGAGCTTCGCGATCAGCTCGTTCCATTTATGGAAGGTTTACTAAACAAGAAGAAAT
 2 K E L R D Q L V P F M E G L L N K E E I
 NdeI
 3780 CTAAGCTAGAAATAATTTTGTTTAACTTTAAGAAGGAGATATACATATGACGCATGTACG
 2 M T H V R
 3840 CTTGACTACTCCAAAAGCGTTGACTTTCTTTCCAACGGAACATGAACTTACATACCTGCG
 2 L T T P K A L T F F P T E H E L T Y L R
 3900 GGACTTTGTAAAAACAGCACACCATAATATCCATGAGAAAACAGGCGGGCAGCGATTT
 2 D F V K T A H H N I H E K T G A G S D F
 EcoRI
 3960 TCTAGGCTGGGTGGACCTCCCTGAACATTATGATAAAGAAGAATTCGCGCGCATCCAAAA
 2 L G W V D L P E H Y D K E E F A R I Q K
 4020 AAGCGCGGAAAAAATCCAATCTGACTCTGATGTCTTGCTTGTGTCGGCATCGGCGGTTTC
 2 S A E K I Q S D S D V L L V V G I G G S
 4080 TTATCTTGGAGCGGGCAGCGATTGAAGCGCTGAATCACGCGTTTTATAACACTTTGCC
 2 Y L G A R A A I E A L N H A F Y N T L P
 4140 AAAAGCCAAACCGGCAATCCGCAAGTCATTTTAACTTCTCTATTAATGTGATTTCTAA
 2 K A K R G N P Q V I F N F S I N V I S K
 HindIII
 4200 ATCAGGTACGACAACTGAACCTGCAATCGCTTTCCGTATTTTCCGCAAGCTTCTTGAAGA
 2 S G T T T E P A I A F R I F R K L L E E
 4260 GAAATACGGTAAAGAAGAAGCGAAAGCGGATTTATGCAACAACCTGATAAAGAGCGCGG
 2 K Y G K E E A K A R I Y A T T D K E R G
 4320 CGCATTA AAAACGCTTTCTAACGAAGAAGGCTTTGAATCATTGTAATTCCTGACGATGT
 2 A L K T L S N E E G F E S F V I P D D V
 4380 CGGCGGCCGTTATTTCAGTTTTAACAGCTGTAGGTCTCTTGCCGATTGCTGTGACGGCGT
 2 G G R Y S V L T A V G L L P I A V S G V
 4440 CAACATTGACGACATGATGAAAGGCGCCCTGGATGCGAGCAAAGATTTTGCAACATCTGA
 2 N I D D M M K G A L D A S K D F A T S E
 4500 ACTGGAAGATAACCCAGCATACCAATATGCGGTTGTTTCGCAATGTCCTTTATAATAAGGG
 2 L E D N P A Y Q Y A V V R N V L Y N K G
 4560 CAAAACAATTGAAATGCTCATCAACTACGAACCGGCGCTTCAATACTTTGCGGAATGGTG
 2 K T I E M L I N Y E P A L Q Y F A E W W
 4620 GAAGCAGCTGTTTCGGAGAAAGCGAAGGAAAGATGAGAAGGCGATTTATCCTTCTTCAGC
 2 K Q L F G E S E G K D E K G I Y P S S A
 4680 GAACTATCAACAGACCTTCATTCTTTAGGCCAGTATGTACAAGAAGGCCGAGAGATTT
 2 N Y S T D L H S L G Q Y V Q E G R R D L
 4740 ATTCGAAACGGTCCTGAACGTAGAGAAGCCTAAACATGAACTGACAATTGAGGAAGCGGA

-continued

2 F E T V L N V E K P K H E L T I E E A D
 4800 TAACGATCTTGACGGCTTGAAC TATTTAGCCGGTAAAAC TGTGATTTGTTAACAAAA
 2 N D L D G L N Y L A G K T V D F V N K K
 4860 AGCATTCCAAGGTACAATGCTTGCCCATACAGACGGAATGTTCCGAACTTAATCGTTAA
 2 A F Q G T M L A H T D G N V P N L I V N
 4920 CATTCTGAGCTGAATGCATATACTTTTGGATACCTTGTATATTTCTTCGAAAAAGCCTG
 2 I P E L N A Y T F G Y L V Y F F E K A C
 4980 CGCGATGAGCGGTTACCTCCTTGGCGTCAATCCGTTTGACCAGCCTGGTGTAGAAGCGTA
 2 A M S G Y L L G V N P F D Q P G V E A Y
 5040 TAAAGTCAATATGTTTGCCTTACTCGGCAAACCTGGCTTTGAAGAGAAAAAGCAGAGCT
 2 K V N M F A L L G K P G F E E K K A E L

NheI

5100 TGAAAAACGTCTGGAAGATTATAAATGAGCTAGCATGACTGGTGGACAGCAAATGGGTCG
 2 E K R L E D Y K

BamHI KpnI

SphI AgeI

(SEQ ID NO: 2)

5160 GGATCTGTACGACGATGACGATAAGGATCCGGTACCGGCCGCATGCCGGCTAATCGCGAC
 5220 CGGTAACTAGCATAACCCCTTGGGGCCTCTAACCGGTCTTGAGGGGTTTTTTGCTAAA
 5280 GGAGGAACTATATCCGGTCCAAGAATTGGAGCCAATCAATCTTGCGGAGAACTGTGAAT
 5340 GCGCAAACCAACCCTTGGCAGAACATATCCATCGCGTCCGCCATCTCCAGCAGCCGCACG
 5400 CGGCGCATCTCGGGCCGCGTTGCTGGCGTTTTTCCATAGGCTCCGCCCCCTGACGAGCA
 5460 TCACAAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACTATAAAGATACCA
 5520 GGCCTTTCCCCCTGGAAGCTCCCTCGTGCCTCTCCTGTTCCGACCCTGCCGCTTACCGG
 5580 ATACCTGTCCGCTTTCTCCCTTCGGGAAGCGTGGCGCTTTCTCATAGCTCACGCTGTAG
 5640 GTATCTCAGTTCGGTGTAGGTCGTTCCCTCCAAGCTGGGCTGTGTGCACGAACCCCCGT
 5700 TCAGCCCGACCGCTGCGCCTTATCCGGTAACTATCGTCTTGAGTCCAACCCGGTAAGACA
 5760 CGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATGTAGG
 5820 CGGTGTACAGAGTTCTTGAAGTGGTGGCCTAACTACGGCTACACTAGAAGGACAGTATT
 5880 TGGTATCTGCGCTCTGCTGAAGCCAGTTACCTTCGAAAAAGAGTTGGTAGCTCTTGATC
 5940 CGGCAAACAAACCACCGCTGGTAGCGGTGGTTTTTTTTGTTGCAAGCAGCAGATTACGCG
 6000 CAGAAAAAAGGATCTCAAGAAGATCCTTTGATCTTTCTACGGGTCTGACGCTCAGTG
 6060 GAACGAAAACCTCACGTTAAGGGATTTTGGTCATGAGATTATCAAAAAGGATCTTCACCTA
 6120 GATCCTTTTAAATTAATAATGAAGTTTTAAATCAATCTAAAGTATATATGAGTAAACTTG
 6180 GTCGTACAGTTACCAATGCTTAATCAGTGAGGCACCTATCTCAGCGATCTGCTATTTTCG
 6240 TTCATCCATAGTTGCCTGACTCCCCGTCGTGTAGATAACTACGATACGGGAGGGCTTACC
 6300 ATCTGGCCCCAGTGTGCAATGATACCGCGAGACCCACGCTCACCGGCTCCAGATTTATC
 6360 AGCAATAAACCCAGCCAGCCGGAAGGGCCGAGCGCAGAAGTGGTCTGCAACTTTATCCGC
 6420 CTCCATCCAGTCTATTAATTGTTGCCGGAAGCTAGAGTAAGTAGTTCCGCGTAAATAG

PstI

6480 TTTGCGCAACGTTGTTGCCATTGCTGCAGGCATCGTGGTGTACGCTCGTCGTTTGGTAT
 6540 GGCTTCATTCAGCTCCGGTTCCCAACGATCAAGGCGAGTTACATGATCCCCATGTTGTG

- continued

6600 CAAAAAGCGGTTAGCTCCTTCGGTCCTCCGATCGTTGTCAGAAGTAAGTTGGCCGCAGT
 6660 GTTATCACTCATGGTTATGGCAGCACTGCATAATTCTTACTGTCATGCCATCCGTAAG

ScaI

6720 ATGCTTTTCTGTGACTGGTGAGTACTCAACCAAGTCATTCTGAGAATAGTGTATGCGGCG
 6780 ACCGAGTTGCTCTTGCCCGCGTCAACACGGGATAATACCGCGCCACATAGCAGAACTTT
 6840 AAAAGTGCTCATCATTGGAAAACGTTCTTCGGGGCGAAAACCTCAAGGATCTTACCGCT
 6900 GTTGAGATCCAGTTCGATGTAACCACTCGTGCACCCAACGATCTTCAGCATCTTTTAC
 6960 TTTCACCAGCGTTTCTGGGTGAGCAAAAACAGGAAGGCAAAATGCCGCAAAAAGGGAAAT
 7020 AAGGGCGACACGGAAATGTTGAATACTCATACTCTTCCTTTTCAATATTATTGAAGCAT
 7080 TTATCAGGGTTATTGTCTCATGAGCGGATACATATTTGAATGTATTTAGAAAATAAACA
 7140 AATAGGGTTCCGCGCACATTTCCCGAAAAGTGCCACCTGACGTCTAAGAAACCATTAT

EcoRI

7200 TATCATGACATTAACCTATAAAAAATAGGCGTATCACGAGGCCCTTTCGTCTTCAAGAATT
 7260 CCTGTTATAAAAAAAGGATCAATTTTGAACCTCTCTCCAAAGTTGATCCCTTAACGATTT
 7320 AGAAATCCCTTTGAGAATGTTTATATACATTCAAGGTAACCAGCCAACATAATGACAATGA
 7380 TTCTGAAAAAAGTAATAACAAATTAATAACAGATAAGTTGACTGATCAACTTCCATAG
 7440 GTAACAACCTTTGATCAAGTAAGGGTATGGATAATAAACCACTACAATTGCAATACCTG
 7500 TTCCCTCTGATAAAAAGCTGGTAAAGTTAAGCAAACCTATTCCAGCACAGCTTCCTGCT
 7560 GTTTCAGCTACTTGAAACAATTGTTGATATAACTGTTTTGGTGAACGAAAGCCCACCTA
 7620 AAACAAATACGATTATAATTGTATGAACCATGATGTTGTTTCTAAAAGAAAGGAAGCAG
 7680 TTAAAAAGCTAACAGAAAGAAATGTAACCTCCGATGTTTAAACACGTATAAAGGACCTCTTC
 7740 TATCAACAAGTATCCCACCAATGTAGCCGAAAATAATGACACTCATTGTTCCAGGGAAAA

ScaI

7800 TAATTACACTTCCGATTTCCGCGACTACTTAGCTGGTGAACATCTTTCATCATATAAGGAA
 7860 CCATAGAGACAAACCCTGCTACTGTTCCAAATATAATTCCCCACAAAGAACTCCAATCA
 7920 TAAAAGGTATATTTTTCCCTAATCCGGGATCAACAAAAGGATCTGTTACTTTCTGATAT
 7980 GTTTTACAAATATCAGGAATGACAGCACGCTAACGATAAGAAAAGAAATGCTATATGATG

EcoRV

8040 TTGTAAACAACATAAAAAATACAATGCCTACAGACATTAGTATAATTCCTTTGATATCAA
 8100 AATGACCTTTTATCCTTACTTCTTTCTTTAATAATTTTATAAGAAACGGAACAGTGATAA
 8160 TTGTTATCATAGGAATGAGTAGAAGATAGGACCAATGAATATAATGGGCTATCATTCCAC
 8220 CAATCGCTGGACCGACTCCTTCTCCCATGGCTACTATCGATCCAATAAGACCAAATGCTT
 8280 TACCCCTATTTTCTTTGGAATATAGCGCGCAACTACAACCATTACGAGTGCTGGAAATG
 8340 CAGCTGCACCAGCCCCTTGAATAAAACGAGCCATAATAAGTAAGGAAAAGAAAGAAATGGC
 8400 CAACAAACCAATTACCGACCCGAAACAATTTATTATAATTCAAATAGGAGTAACCTTT
 8460 TGATGCCAATTGATCAGATAGCTTCCATATACAGCTGTTCCAATGGAAAAGGTTAACA
 8520 TAAAGGCTGTGTTACCCAGTTTGTACTCGCAGGTGGTTTATTAAAATCATTGCAATAT
 8580 CAGGTAATGAGACGTTCAAACCATTTTCAATTAATACGCTAAAAAAGATAAAAATGCAA
 8640 GCCAAATTAATAATTTGGTTGTGTCGTAATTCGATTGTGAATAGGATGTATTCACATTT
 8700 ACCCTCAATAATGAGGGCAGACGTAGTTTATAGGGTTAATGATACGCTTCCCTCTTTTA

- continued

8760 ATTGAACCCGTGTTACATTACACTTCATAATTAATTCCTCCTAAACTTGATTA AAC
8820 ATTTTACCACATATAAACTAAGTTTTAAATTCAGTATTTTCATCACTTATACAACAATATG
8880 GCCCGTTTGTGAACTACTCTTTAATAAAATAATTTTTCCGTTCCCAATTCACATTGCA
8940 ATAATAGAAAATCCATCTTCATCGGCTTTTTTCGTATCATCTGTATGAATCAAATCGCCT
9000 TCTTCTGTGTCATCAAGGTTTAAATTTTTTATGTATTTCTTTTAAACAAACCACCATAGGAG
9060 ATTAACCTTTTACGGTGTAACCTTCCTCCAAATCAGACAAACGTTTCAAATTCCTTTTCT
9120 TCATCATCGGTATAAAATCCGTATCCTTTACAGGATATTTTGCAGTTTCGTCAATTGCC
9180 GATTGTATATCCGATTTATATTTATTTTTCCGTCGAATCATTGAACTTTTACATTTGGA
9240 TCATAGTCTAATTTTATTGCCTTTTTCCAAAATTGAATCCATGTTTTTGTTCACGTAG
9300 TTTTCTGTATTCTTAAAATAAGTTGGTTCCACACATACCAATACATGCATGTGCTGATTA
9360 TAAGAATTATCTTTATTATTATTGTCACTTCCGTTGCACGCATAAAACCAACAAGATTT
9420 TTATTAATTTTTTTATATTGCATCATTCGGCGAAATCCTTGAGCCATATCTGACAAACTC
9480 TTATTTAATCTTCGCCATCATAAACATTTTTAACTGTTAATGTGAGAAACAACCAACGA
9540 ACTGTTGGCTTTTGTTTAATAACTTCAGCAACAACCTTTTGTGACTGAATGCCATGTTTC
9600 ATTGCTCTCCTCCAGTTGCACATTGGACAAAGCCTGGATTTACAAAACCACTCGATAC
9660 AACTTTCTTTCGCCTGTTTCACGATTTTGTTTATACTCTAATATTTCAGCACAATCTTTT
9720 ACTCTTTCAGCCTTTTTAAATTCAGAATATGCAGAAGTTCAAAGTAATCAACATTAGCG
9780 ATTTTCTTTCTCTCCATGGTCTCACTTTTCCACTTTTTGTCTTGTCCACTAAAACCCTT
9840 GATTTTTTCATCTGAATAAATGCTACTATTAGGACACATAATATTAAGAAACCCCATC
9900 TATTTAGTTATTTGTTTGGTCACTTATAACTTTAACAGATGGGGTTTTTCTGTGCAACCA
9960 ATTTTAAGGGTTTTCAATACTTTAAAACACATACATACCAACACTTCAACGCACCTTTCA
10020 GCAACTAAAATAAAAATGACGTTATTTCTATATGTATCAAGATAAGAAAGAACAAGTTCA
10080 AAACCATCAAAAAAGACACCTTTTCAGGTGCTTTTTTTATTTTATAAACTCATTCCCTG
10140 ATCTCGACTTCGTTCTTTTTTTTACCTCTCGGTTATGAGTTAGTTCAAATTCGTT

40

This plasmid has a molecular weight of 10194 bp and contains the various genes responsible for hyaluronic acid synthesis under the control of a strong T7 promoter of bacteriophage T7. The hasA sequence from *Streptococcus equi* is included between bases 196 and 1383, that of the tuaD gene between bases 1430 and 2873, the sequence coding for gtaB between bases 2905 and 3781, and that for pgi between bases 3824 and 5125.

Example 8

Restriction map of plasmid pT7hyal

When plasmid pT7hyal is cut with restriction enzymes it gives rise to a restriction map which corresponds to that expected after sequencing. In column 1 of FIG. 5, it is shown that the cutting with enzyme EcoRI gives rise to three bands 4900 bp, 3240 bp and 2020 bp from plasmid; in column 2 it is shown that the cutting with EcoRI and HindIII gives rise to six bands 3290 bp, 2950 bp, 1660 bp, 1400 bp, 610 bp, and 290 bp; in column 3 it is shown that the cutting with HindIII gives rise to three bands 6240 bp, 2240 bp and 1690 bp, and in column 4 it is shown that, the cutting with restriction enzyme XbaI, single site gives rise to a single band (FIG. 5).

Example 9

Check on Synthesis of Proteins which Lead to Hyaluronic Acid Synthesis

The two plasmids pPT7hasAtuaD and pPT7hasAtuaDgtaBpgi (pT7Hyal) were transformed into bacterial cells of *E. coli* BL21 DE3. After induction with IPTG, the cells were lysed, and the sample obtained was loaded into an SDS-PAGE to test for the presence of the proteins which lead to hyaluronic acid synthesis (FIG. 6). The preparation in column 2 corresponds to cells which carried plasmid pPT7hasAtuaD: as shown in FIG. 6, compared with the control colonies in column 1, column 2 presents a protein with a molecular weight of 54 kDa which corresponds to tuaD, and a protein with a weight of 42 kDa which corresponds to hasA. The samples in column 7 and 8 which carry plasmid pPT7hasAtuaDgtaBpgi produce, compared with control colonies 5 and 6, a protein with a molecular weight of 54 kDa which corresponds to tuaD, a protein with a molecular weight of 51 kDa which corresponds to pgi, a protein with a weight of 42 kDa which corresponds to hasA, and a protein with a molecular weight of 32 kDa which corresponds to gtaB. In conclusion, both plasmids produce the proteins of the expected molecular weight required for hyaluronic acid synthesis.

Example 10

Synthesis of Hyaluronic Acid in *E. coli* and Selection by IPTG Gradient

Plasmids pPT7 (control colony), pPT7hasAtuaD (colony 6) and pPT7hasAtuaDgtaBpgi (pT7Hyal—colony 2) were transformed into bacterial cells BL21 DE3. After 24 hours' growth at 37° C., the colonies were analysed for the production of hyaluronic acid. In solution, the cells which carry plasmids pPT7hasAtuaD (colony 6) or plasmids pPT7hasAtuaDgtaBpgi (colony 2) grow much more slowly, and after induction with IPTG only produce low levels of hyaluronic acid. The cells were then plated in the presence of IPTG (FIG. 7).

The control colonies that carry plasmid pPT7 (and no hyaluronic acid synthesis gene) grow more easily, and are larger and flatter, than colony 6 and colony 2, in which the bacteria are engaged in producing hyaluronic acid; in fact, colonies 2 and 6 are shinier than the control as they produce hyaluronic acid. To select cells able to express high levels of hyaluronic acid, the cells were plated in the presence or absence of IPTG (FIG. 8). In the presence of IPTG the majority of the colonies die, and only some survive, especially those close to the IPTG gradient formed. These cells were selected and replated in the presence of IPTG to establish their survival rate (FIG. 9): all of them remained alive, maintaining their HA synthesis capacity.

The above statements are demonstrated by the fact that the cells of colonies 6 and 2 were cultured in solution for 48 hours in the presence of IPTG and 1% saccharose. 1 ml of this bacterial culture was centrifuged to obtain the precipitate, and the bacterial precipitate was then lysed in the presence of 0.1% SDS for 10 minutes. After adding 2 volumes of absolute ethyl alcohol, the result was as shown in FIG. 10.

As will be seen, only colonies 6 and 2 give rise to a hyaluronic acid precipitate (which was tested with the carbazole test).

Example 11

Transformation of Plasmids pPT7hasAtuaD and pPT7hasAtuaDgtaBpgi into *Bacillus megaterium*

The *B. megaterium* used in the present invention is already pre-transformed with plasmid pT7-RNAP (QM B1551 MoBiTec) (this plasmid is able to replicate in both *E. coli* and *B. megaterium* because it contains two origins of replication which allow its propagation in both bacteria). It also contains resistance to ampicillin and chloramphenicol, which can be used for *E. coli* and *B. megaterium* respectively. The plasmid contains the sequence able to code for T7 RNA polymerase under the control of the inducible promoter for xylose, and also contains the repressor for the xylose promoter; if the cells are maintained in the absence of xylose, they are therefore unable to transcribe T7 RNA polymerase.

For the transformation of this bacterium it was necessary to remove its bacterial wall to obtain the protoplasts to use for the transformation. To remove the bacterial wall, 50 ml of LB medium were introduced into a 300 ml Erlenmeyer flask, and 1 ml of *Bacillus megaterium* grown overnight under aerobic conditions was added. When the cell density at OD578 reached the value of 1, the cells were centrifuged at 4500 rpm for 15 minutes. The cells were then suspended in 5 ml of 17.5 g/L of Antibiotic Medium no. 3, 500 mM

saccharose, 20 mM sodium maleinate and 20 mM MgCl₂ pH6 (buffer SMMP). 50 ml of lysozyme 1 mg/ml in SMMP buffer were added and the mixture was maintained at 37° C. for 60 minutes, so as to remove the cell wall; the cells were then gently centrifuged at 1300 rpm for 10 minutes. The bacterial cells were then suspended in 5 ml of fresh SMMP buffer without stirring, as the protoplasts are sensitive to physical stress. This washing was repeated once more. After suspension, the protoplasts were ready to be used directly for the transformation or to be frozen at -80° C. in SMMP, which contains 15% glycerol. However, the transformations are much more efficient when the protoplasts are freshly prepared. For the transformation, 500 µl of protoplast suspension were mixed with 1 µg of plasmid DNA pPT7hasAtuaD or pPT7hasAtuaDgtaBpgi; 1.5 ml of PEG-P (40% w/v PEG6000 in 1×SMM) were then added, and the mixture was placed at ambient temperature for 2 minutes. 5 ml of SMMP were added, and the tubes were gently mixed by rotation.

The bacteria were centrifuged gently at 3000 rpm for 10 minutes at ambient temperature. The supernatant was discarded, and the almost invisible sediment contained the bacteria of interest. 500 µl of SMMP was added to the bacteria, which were then incubated for 90 minutes at 37° C. under slow stirring, at a maximum of 10 rpm; 2.5 ml of CR5 top agar were then prepared in sterile tubes in a hot bath at 43° C.

The CR5 top agar was prepared by mixing two components:

- a) 51.5 g of saccharose, 3.25 g of MOPS and 0.33 g of NaOH in 250 ml of H₂O pH7.3, sterilised by filtration
- b) 2.0 g of agar, 0.1 g of casaminoacids, 5 g of yeast extract and 142.5 ml of H₂O.

After autoclaving for 20 minutes, the two ingredients, cooled to 50° C., were mixed together.

After growth, 100 µl of the above disclosed cell preparation were added to 2.5 ml of top agar, mixed gently by rotating with both hands, and deposited on a pre-heated plate containing the antibiotic (4.5 µg/ml of chloramphenicol and 10 µg/ml of tetracycline). The mixture was incubated overnight at 37° C.; the colonies resulting larger or smaller, depending on their access to air.

Example 12

Expression of Hyaluronic Acid in *B. megaterium*

The transformed *B. megaterium* cells were cultured in LB medium with tetracycline and chloramphenicol up to an optical density at 578 nm of 0.4 at 37° C. The induction was conducted with the addition of 0.5% of D-xylose (w/v), followed by incubation at 37° C. The optical density of the bacteria was read every 30 minutes until the optical density at 600 nm reached 1.5; at this point the cells reached the steady state. These cells, as in the case of *E. coli*, are unable to produce hyaluronic acid directly after induction.

Example 13

System for the Selection of Hyaluronic Acid Secreting Cells

To obtain *B. megaterium* cells able to produce hyaluronic acid, the plate selection system presented for *E. coli* was employed, using xylose as inductor instead of IPTG. The cells which produce high levels of hyaluronic acid in the plate were then selected. Those cells survive, and can be

cultured. The supernatant contains the hyaluronic acid produced (its presence is confirmed by carbazole analysis when it is precipitated with two volumes of ethanol).

Example 14

Fermentation of transformed *B. megaterium* cells Selected on Gradient

Bacillus megaterium cells transformed with two genes pPT7hasAtuaD plasmid or with four genes pPT7hasAtuaDgtaBpgi plasmid, and selected on xylose gradient were cultured in a 20 l fermenter in 5 l of MM++ medium and glucose or saccharose as carbon source.

Xylose was added as inductor after the start of fermentation.

In the following some fermentation processes for the production of HA are illustrated, said processes mainly differing because of:

- the starting source of carbon;
- the added feed (glucose or saccharose);
- the fermentation temperature (the temperature can be established in a range of from 20 to 38° C., preferably of from 25 to 35° C.);
- time of fermentation.

Culture media used:

LB broth (Miller), pH 7

MM++ (Minimal Medium Bs), pH 7, containing per liter:
1 g (NH₄)₂ HPO₄; 1 g NH₄NO₃; 2.5 g K₂HPO₄; 2.5 g KH₂PO₄; 0.2 g MgSO₄·7H₂O; 0.01 g FeSO₄·7H₂O; 0.007 g MnSO₄·7H₂O.

Example 14a

Production of HA Having a Weight Average MW Comprised in the Range of 100-500 KD

The bacterial strain *B. Megaterium* (QM B1551), transfected with the plasmid pPT7hasAtuaDgtaBpgi selected on xylose gradient 0.5% w/v, as described in Example 13, was used.

Procedure: a single colony resistant to xylose was inoculated into 5 ml of sterile LB medium containing 5 mg/l of tetracycline and the inductor. The culture was grown at 37° C., under stirring at 200 rpm.

After 8 hours, 50 µl of this culture were inoculated into a flask containing 50 ml of the medium mentioned above (containing the inductor), and it was made to grow under the same conditions described above.

Subsequently, spent further 14-16 hours, 2 ml of this culture were inoculated into a flask containing 500 ml of the medium above, and it was made to grow under the same conditions until reaching a D.O.^{600nm} of 0.6-0.8.

500 ml of the culture thus obtained were then inoculated in the fermenter containing MM++ medium, and the fermentation conditions involved maintaining the culture under stirring at 600 rpm, aeration with 20-24 litres of air/min, a temperature of 37° C. (the temperature of fermentation can be established in a range between 25° C. and 38° C.), and a pH of 6.9 to 7.1. The initial source of carbon was 2% saccharose.

After 4 hours of fermentation, a 2% saccharose supply was added. At 24 hours of fermentation, xylose was added to a final concentration of 0.5%; this induction proceeded for 4 hours; at the end, 10% saccharose was added in steps.

At the end of fermentation (130 hours), the bacterial culture was discharged and centrifuged at 7500 rpm at 8° C. for 20 minutes.

The fermentation broth thus obtained, clarified as free of the cellular component, was analyzed to determine the concentration of HA with the carbazole method (Bitter and Muir, 1962, *Anal. Biochem.* 4:330-334).

Results: The analysis resulted in a concentration of HA of 3.5 g/l.

Determination of weight average molecular weight MW:

For its analysis it was used the method of the intrinsic viscosity (as described in Terbojevich et al., *Carbohydr. Res.* 1986, 363-377, incorporated herein by reference).

Results: the analyzed HA sample showed a weight average molecular weight MW in the range of 100-300 KD.

Example 14b

Production of HA Having a Weight Average MW Comprised in the Range of 1×10⁶-2×10⁶ D

The bacterial strain *B. Megaterium* (QM B1551), transfected with the two genes plasmid pPT7hasAtuaD and with the four genes plasmid pPT7hasAtuaDgtaBpgi, selected on xylose gradient, as described in Example 13, was used.

Procedure: for each plasmid which was used, a single colony resistant to xylose was processed as indicated in example 14a. The initial source of carbon was saccharose at 2%: in this example the further supply was glucose (further experimental tests showed that it can be substituted with equal or lower amounts of saccharose). The fermentation conditions were the same as those used in example 14a with the only difference of the fermentation temperature: 25° C.

The culture media used for the fermentation were those disclosed according to example 14a.

At the end of the process (ended after 24 hours), the fermentation broth was analyzed to determine the concentration of HA with the carbazole method.

Results: *B. Megaterium* (QM B1551), transfected with the two genes plasmid pPT7hasAtuaD: the analysis resulted in a concentration of HA of 2.5 g/l;

B. Megaterium (QM B1551), transfected with the four genes plasmid pPT7hasAtuaDgtaBpgi: the analysis resulted in a concentration of HA of 3.2 g/l;

Determination of weight average molecular weight MW: For its analysis it was used the method of the intrinsic viscosity as indicated in the previous example 14a.

Results: the analyzed HA sample produced by *B. Megaterium* transfected with the two genes plasmid showed a weight average molecular weight MW in the range of 1.3×10⁶-1.7×10⁶D;

the analyzed HA sample produced by *B. Megaterium* transfected with the four genes plasmid showed a weight average molecular weight MW in the range of 1.6×10⁶-2×10⁶D.

The system engineered in *B. megaterium* is inducible, so the fermentation process can be continued by stimulating the production of HA to obtain the desired weight average molecular weight MW; fermentation times between 80 and 160 hours result in a medium-low weight average molecular weight MW, comprised in the range between 100-500 KD, fermentation times between 40 and 80 hours result in a weight average molecular weight in the range between 500-1000 KD, fermentation times between 12 and 40 hours result in a weight average molecular weight MW in the range 1×10⁶-3×10⁶ D.

41

With the experiments and the results obtained above, the Applicant has demonstrated to have perfected a system of production of HA in *B. megaterium* by plasmid vectors by: engineering of 2 genes (or 4 genes) plasmid vectors for the synthesis of enzymes needed for the production of said polysaccharide, whose gene control is placed under the control of strong promoter T7 of bacteriophage T7;

42

perfecting a system of selection of these transfected strains for the production of stable, viable, replicating and HA secreting strains;
creating an inducible system of HA production, thus controllable both in order to obtain high concentrations of HA and for the production of said polysaccharide at different weight average molecular weight MW.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 18

<210> SEQ ID NO 1

<211> LENGTH: 7880

<212> TYPE: DNA

<213> ORGANISM: artificial

<220> FEATURE:

<223> OTHER INFORMATION: Plasmid containing hasA and tuaD gene under the control of promoter T7

<400> SEQUENCE: 1

```

ctttttaggt tctaaatcgt gtttttcttg gaattgtgct gttttatcct ttaccttgtc      60
tacaaacccc ttaaaaacgt ttttaaaggc ttttaagccg tctgtacggt ccttaaggcg      120
aaattaatac gactcactat agggagacca caacggtttc ccgaatatta attaaccaag      180
gaggtgaaat gtacaatgag aacattaaaa aacctcataa ctggtgtggc ctttagtatt      240
ttttgggtac tggtgattta cgtcaatggt tatctctttg gtgctaaagg aagcttgtca      300
atztatggct ttttgetgat agcttaccta ttagtcaaaa tgccttatac ctttttttac      360
aagccattta aggaagggc tgggcaatat aaggttgcag ccattattcc ctcttataac      420
gaagatgctg agtcattgct agagacctta aaaagtgttc agcagcaaac ctatccccta      480
gcagaaattt atggtgttga cgatggaagt gctgatgaga caggtattaa gcgcattgaa      540
gactatgtgc gtgacactgg tgacctatca agcaatgtca ttgttcaccg gtcagaaaaa      600
aatcaaggaa agcgtcatgc acaggcctgg gcctttgaaa gatcagacgc tgatgtcttt      660
ttgaccggtg actcagatac ttatatctac cctgatgctt tagaggagt gttaaaaacc      720
tttaatgacc caactgtttt tgctgcgacg ggtcacctta atgtcagaaa tagacaaacc      780
aatctcttaa cacgcttgac agatattcgc tatgataatg cttttggcgt tgaacgagct      840
gccaatccg ttacaggtaa tattctcggt tgctcaggcc cgcttagcgt ttacagacgc      900
gaggtgggtg ttcctaacat agatagatac atcaaccaga ccttcctggg tattcctgta      960
agtatcggtg atgacagggt cttgaccaac tatgcaactg atttaggaaa gactgtttat     1020
caatccactg ctaaagtgtat tacagatggt cctgacaaga tgtctactta cttgaagcag     1080
caaaaccgct ggaacaagtc cttctttaga gagtccatta tttctgttaa gaaaatcatg     1140
aacaatcctt ttgtagccct atggaccata cttgagggtg ctatgtttat gatgcttggt     1200
tattctgtgg tggatttctt tgtaggcaat gtcagagaat ttgattggct cagggttttg     1260
gcctttctgg tgattatctt cattgttgct ctttgtcgta atattcacta tatgcttaag     1320
caccgctgt ccttcttggt atctccggtt tatgggggtac tgctttggtt gtccctacagc     1380
ccttgaatt gtattctctt tttactatta gaaatgctga ctggggaaca cgtaaaaaat     1440
tattataatc tagaaataat tttgtttaac ttaagaagg agatatacat atgaaaaaaa     1500
tagctgtcat tggaacagggt tatgtaggac tcgtatcagg cacttgcttt gggagatcg     1560
gcaataaagt tgtttgctgt gatatcgatg aatcaaaaat cagaagcctg aaaaatgggg     1620
taatcccaat ctatgaacca gggcttgcag acttagttga aaaaaatgtg ctggatcagc     1680

```


-continued

gcctgacctt tacgaacgat atcccgtctg ccattcgggc ctcagatatt atttatattg	1740
cagtcggaac gcctatgtcc aaaacaggtg aagctgattt aacgtacgtc aaagcggcgg	1800
cgaaaacaat cggtagcat cttaacggct acaaagtgat cgtaaataaa agcacagtcc	1860
cggttggaac agggaaactg gtgcaatcta tcgttcaaaa agcctcaaag gggagatact	1920
catttgatgt tgtatctaac cctgaattcc ttcgggaagg gtcagcgatt catgacacga	1980
tgaatatgga gcgtgccgtg attggttcaa caagtcataa agccgctgcc atcattgagg	2040
aacttcatca gccattccat gctcctgtca ttaaaacaaa cctagaaagt gcagaaatga	2100
ttaaatacgc cgcaatgca tttctggcga caaagatttc ctttatcaac gatatcgcaa	2160
acatttgatga gcgagtcggc gcagacgttt caaaagttgc tgatgggtgtt ggtcttgaca	2220
gccgtatcgg cagaaagttc cttaaagctg gtattggatt cggcggttca tgttttccaa	2280
aggatacaac cgcgctgctt caaatcgcaa aatcggcagg ctatccattc aagctcatcg	2340
aagctgtcat tgaacgaac gaaaagcagc gtgttcatat tgtagataaa cttttgactg	2400
ttatgggaag cgtcaaaggg agaaccattt cagtccctggg attagccttc aaaccgaata	2460
cgaacgatgt gagatccgct ccagcgttg atattatccc aatgctgcag cagctgggcg	2520
cccatgtaaa agcatacgat ccgattgcta ttctgaagc ttcagcgatc cttggcgaac	2580
aggtcgagta ttacacagat gtgtatgctg cgatggaaga cactgatgca tgctgattt	2640
taacggattg gccggaagtg aaagaaatgg agcttgtaa agtgaaaacc ctcttaaaac	2700
agccagtcat cattgacggc agaaatttat tttcacttga agagatgcag gcagccggat	2760
acatttatca ctctatcggc cgtcccgtg ttcggggaac ggaaccctct gacaagtatt	2820
ttccgggctt gccgctgaa gaattggcta aagacttggg aagcgtcaat ttataaggat	2880
ccggccgcat gccggcta at cgcgaccggt taactagcat aacccttgg gcctctaaa	2940
cgggtcttga ggggtttttt gctaaaggag gaactatata cggccaaga attggagcca	3000
atcaattctt gcggagaact gtgaatgctc aaaccaacc ttggcagaac atatccatcg	3060
cgcccgcat ctccagcagc cgcacgcggc gcatctcggg ccgcttctgt ggcgtttttc	3120
cataggctcc gccccctga cgagcatcac aaaaatcgac gctcaagtca gaggtggcga	3180
aaccgcagag gactataaag ataccaggcg tttccccctg gaagctcct cgtgcgctct	3240
cctgttccga ccctgccgt taccggatac ctgtccgct ttctccctc ggaagcgtg	3300
gcgctttctc atagctcacg ctgtaggtat ctcagttcgg tgtaggtcgt tcgctccaag	3360
ctgggctgtg tgcacgaacc cccggttcag cccgaccgct gcgccttata cggtaactat	3420
cgtcttgagt ccaaccggt aagacacgac ttatcgccac tggcagcagc cactggtaac	3480
aggattagca gagcggaggta tggtaggggt gctacagagt tcttgaagtg gtggcctaac	3540
tacggctaca ctagaaggac agtatttggg atctgcgctc tgctgaagcc agttacctc	3600
ggaaaaagag ttggtagctc ttgatccggc aaacaaacca ccgctggtag cggtggtttt	3660
tttgtttgca agcagcagat tacgcgaga aaaaaaggat ctcaagaaga tcctttgatc	3720
ttttctacgg ggtctgacgc tcagtggaac gaaaactcac gttaaggat tttggtcatg	3780
agattatcaa aaaggatctt cacctagatc cttttaaatt aaaaatgaag ttttaaatca	3840
atctaaagta tatatgagta aacttggct gacagttacc aatgcttaat cagtgaggca	3900
cctatctcag cgatctgtct atttcgttca tccatagttg cctgactccc cgtcgtgtag	3960
ataactacga tacgggaggg cttaccatct ggccccagt ctgcaatgat accgcgagac	4020

-continued

ccacgctcac	cggctccaga	tttatcagca	ataaaccagc	cagccggaag	ggccgagcgc	4080
agaagtggtc	ctgcaacttt	atccgcctcc	atccagtcta	ttaattgttg	ccgggaagct	4140
agagtaagta	gttcgccagt	taatagtttg	cgcaacgttg	ttgccattgc	tgcaggcatc	4200
gtggtgtcac	gctcgtcgtt	tggtatggct	tcattcagct	ccggttccca	acgatcaagg	4260
cgagttacat	gatccccat	gttgtgcaaa	aaagcgggta	gctccttcgg	tccctccgatc	4320
gttgtcagaa	gtaagttggc	cgcagtgtta	tactcatgg	ttatggcagc	actgcataat	4380
tctcttactg	tcatgccatc	cgtaagatgc	ttttctgtga	ctgggtgagta	ctcaaccaag	4440
tcattctgag	aatagtgtat	gcgccgaccg	agttgctctt	gcccggcgtc	aacacgggat	4500
aataccgcgc	cacatagcag	aactttaaaa	gtgctcatca	ttggaaaacg	ttcttcgggg	4560
cgaaaactct	caaggatctt	accgctgttg	agatccagtt	cgatgtaacc	cactcgtgca	4620
cccaactgat	cttcagcatc	ttttactttc	accagcgttt	ctgggtgagc	aaaaacagga	4680
aggcaaatg	ccgcaaaaa	gggaataagg	gcgacacgga	aatggtgaat	actcatactc	4740
ttcctttttc	aatattattg	aagcatttat	cagggttatt	gtctcatgag	cggatacata	4800
tttgaatgta	tttagaaaa	taaacaata	ggggttccgc	gcacatttcc	ccgaaaagtg	4860
ccacctgacg	tctaagaaac	cattattatc	atgacattaa	cctataaaaa	taggcgtatc	4920
acgaggccct	ttcgtcttca	agaattcctg	ttataaaaa	aggatcaatt	ttgaactctc	4980
tcccaaagtt	gatcccttaa	cgatttagaa	atcccttga	gaatgtttat	atacattcaa	5040
ggtaaccagc	caactaatga	caatgattcc	tgaaaaagt	aataacaaat	tactatacag	5100
ataagttgac	tgatcaactt	ccataggtaa	caaccttga	tcaagtaagg	gtatggataa	5160
taaaccacct	acaattgcaa	tacctgttcc	ctctgataaa	aagctggtaa	agttaagcaa	5220
actcattcca	gcaccagctt	cctgctgttt	caagctactt	gaaacaattg	ttgatataac	5280
tgttttgggtg	aacgaaagcc	cacctaaaac	aaatacgatt	ataattgtca	tgaaccatga	5340
tgttgtttct	aaaagaaagg	aagcagttaa	aaagctaaca	gaaagaaatg	taactccgat	5400
gtttaacacg	tataaaggac	ctcttctatc	aacaagtatc	ccaccaatgt	agccgaaaat	5460
aatgacactc	attgttccag	ggaaaataat	tacacttccg	atttcggcag	tacttagctg	5520
gtgaacatct	ttcatcatat	aaggaacct	agagacaaac	cctgctactg	ttccaaatat	5580
aattccccca	caaagaactc	caatcataaa	aggtatattt	ttccctaate	cgggatcaac	5640
aaaaggatct	gttactttcc	tgatatgttt	tacaaatate	aggaatgaca	gcacgctaac	5700
gataagaaaa	gaaatgctat	atgatgtttg	aaacaacata	aaaaatacaa	tgcttacaga	5760
cattagtata	attcctttga	tatcaaaatg	accttttate	cttacttctt	tctttaataa	5820
tttcataaga	aacggaacag	tgataattgt	tatcatagga	atgagtagaa	gataggacca	5880
atgaatataa	tgggctatca	ttccaccaat	cgctggaccg	actccttctc	ccatggctac	5940
tatcgatcca	ataagaccaa	atgctttacc	cctattttcc	tttggaatat	agcgcgcaac	6000
tacaaccatt	acgagtgtctg	gaaatgcagc	tgaccagccc	ccttgaataa	aacgagccat	6060
aataagtaag	gaaaagaaag	aatggccaac	aaaccaat	accgaccgga	aacaatttat	6120
tataattcca	aataggagta	accttttgat	gcctaattga	tcagatagct	ttccatatac	6180
agctgttcca	atggaaaagg	ttaacataaa	ggctgtgttc	accagtttg	tactcgcagg	6240
tggtttatta	aatcatttg	caatcagcag	taatgagacg	ttcaaacca	tttcatttaa	6300
tacgctaaaa	aaagataaaa	tgcaaagcca	aattaaaatt	tggttgtgtc	gtaaattcga	6360
ttgtgaatag	gatgtattca	catttcaccc	tccaataatg	agggcagacg	tagtttatag	6420

-continued

```

ggttaatgat acgcttcctt cttttaattg aaccctgtta cattcattac acttcataat 6480
taattcctcc taaacttgat taaaacattt taccacatat aaactaagtt ttaaattcag 6540
tatttcatca cttatacaac aatatggccc gtttgttgaa ctactcttta ataaaataat 6600
ttttccggtc ccaattccac attgcaataa tagaaaatcc atcttcatcg gctttttcgt 6660
catcatctgt atgaatcaaa tgccttctt ctgtgtcatc aaggtttaat tttttatgta 6720
tttcttttaa caaaccacca taggagatta accttttaag gtgtaaacct tcctccaaat 6780
cagacaaacg tttcaaattc ttttcttcat catcggtcac aaaatccgta tcctttacag 6840
gatattttgc agtttctgca attgccgatt gtatatccga tttatattta tttttcggtc 6900
gaatcatttg aacttttaca tttggatcat agtctaattt cattgccttt ttccaaaatt 6960
gaatccattg tttttgattc acgtagtttt ctgtattctt aaaataagtt ggttccacac 7020
ataccaatac atgcatgtgc tgattataag aattatcttt attatttatt gtcacttccg 7080
ttgcacgcat aaaaccaaca agattttttat taattttttt atattgcatc attcggcgaa 7140
atccttgagc catatctgac aaactcttat ttaattcttc gccatcataa acatttttaa 7200
ctgttaatgt gagaaacaac caacgaactg ttggcttttg ttaataact tcagcaacaa 7260
ccttttgtga ctgaatgcca tgtttcattg ctctctcca gttgcacatt ggacaaagcc 7320
tggatttaca aaaccacact cgatacaact ttctttcgcc tgtttcacga ttttgtttat 7380
actctaatat ttcagcacia tcttttactc tttcagcctt tttaaattca agaatatgca 7440
gaagttcaaa gtaatcaaca ttagcgattt tcttttctct ccatggcttc acttttccac 7500
tttttgtctt gtccactaaa acccttgatt tttcatctga ataatgcta ctattaggac 7560
acataatatt aaaagaaacc cccatctatt tagttatttg tttggtcact tataacttta 7620
acagatgggg tttttctgtg caaccaattt taagggtttt caatacttta aaacacatac 7680
ataccaacac ttcaacgcac ctttcagcaa ctaaaataaa aatgacgta tttctatatg 7740
tatcaagata agaaagaaca agttcaaac catcaaaaa agacaccttt tcaggtgctt 7800
tttttatttt ataaactcat tccctgatct cgacttcggt ctttttttac ctctcggtta 7860
tgagttagtt caaattcgtt 7880

```

<210> SEQ ID NO 2

<211> LENGTH: 10194

<212> TYPE: DNA

<213> ORGANISM: artificial

<220> FEATURE:

<223> OTHER INFORMATION: Plasmid containing hasA, tuaD, gtaB, pgi gene under the control of T7 promoter

<400> SEQUENCE: 2

```

cttttttaggt tctaaatcgt gtttttcttg gaattgtgct gttttatcct ttaccttgtc 60
tacaaacccc ttaaaaacgt ttttaaagcc ttttaagccg tctgtacggt ccttaaggcg 120
aaattaatac gactcactat agggagacca caacggtttc ccgaatatta attaaccaag 180
gaggtgaaat gtacaatgag aacattaaaa aacctcataa ctggttggtg ctttagtatt 240
ttttgggtac tgttgattta cgtcaatggt tatctctttg gtgctaaagg aagcttgctca 300
atztatggct ttttctgat agcttaccta ttagtcaaaa tgctccttct ctttttttac 360
aagccattta aggaagggc tgggcaatat aaggttgcag ccattattcc ctcttataac 420
gaagatgctg agtcattgct agagacctta aaaagtgttc agcagcaaac ctatccccta 480
gcagaaattt atgttggtga cgatggaagt gctgatgaga caggtattaa gcgcattgaa 540

```

-continued

gactatgtgc	gtgacactgg	tgacctatca	agcaatgtca	ttgttcaccg	gtcagaaaa	600
aatcaaggaa	agcgatcatgc	acaggcctgg	gcctttgaaa	gatcagacgc	tgatgtcttt	660
ttgaccgttg	actcagatac	ttatatctac	cctgatgctt	tagaggagtt	gttaaaaaacc	720
tttaatgacc	caactgtttt	tgctgcgacg	ggcacctta	atgtcagaaa	tagacaaaacc	780
aatctcttaa	cacgcttgac	agatattcgc	tatgataatg	cttttggcgt	tgaacgagct	840
gccaatccg	ttacaggtaa	tattctcggt	tgctcaggcc	cgcttagcgt	ttacagacgc	900
gaggtgggtg	ttcctaacad	agatagatac	atcaaccaga	ccttctctggg	tattctctgta	960
agtatcgggtg	atgacaggtg	cttgaccaac	tatgcaactg	atntagaaa	gactgtttat	1020
caatccactg	ctaaatgtat	tacagatggt	cctgacaaga	tgtctactta	cttgaagcag	1080
caaaacogct	ggaacaagtc	cttctttaga	gagtccatta	tttctgtaa	gaaaatcatg	1140
aacaatcctt	ttgtagccct	atggaccata	cttgaggtgt	ctatgtttat	gatgcttggt	1200
tattctgtgg	tggatttctt	tgtaggcaat	gtcagagaat	ttgattggct	cagggttttg	1260
gcctttctgg	tgattatctt	cattgttgct	ctttgtcgta	atattcacta	tatgcttaag	1320
caccogctgt	ccttcttggt	atctccggtt	tatgggttac	tgctttgttt	gtcctacagc	1380
ccttgaaatt	gtattctctt	ttactatta	gaaatgctga	ctggggaaca	cgtaaaaaat	1440
tattataatc	tagaaataat	tttgtttaac	tttaagaagg	agatatacat	atgaaaaaaaa	1500
tagctgtcat	tggaacaggt	tatgtaggac	tcgtatcagg	cacttgcttt	gcggagatcg	1560
gcaataaagt	tgtttgctgt	gatatcgatg	aatcaaaaat	cagaagcctg	aaaaatgggg	1620
taatcccaat	ctatgaacca	gggcttgacg	acttagttga	aaaaaatgtg	ctggatcagc	1680
gcctgacctt	tacgaacgat	atcccgtctg	ccattcgggc	ctcagatatt	atztatattg	1740
cagtcggaac	gcctatgtcc	aaaacaggtg	aagctgattt	aacgtacgtc	aaagcggcgg	1800
cgaaaaaat	cggtgagcat	cttaacggct	acaaagtgat	cgtaataaaa	agcacagtcc	1860
cggttggaa	agggaaactg	gtgcaatcta	tcgttcaaaa	agcctcaaag	gggagatact	1920
catttgatgt	tgtatctaac	cctgaattcc	ttcgggaagg	gtcagegatt	catgacacga	1980
tgaatatgga	gcggtccgtg	attggttcaa	caagtcataa	agccgctgcc	atcattgagg	2040
aacttcatca	gccattccat	gctcctgtca	ttaaaacaaa	cctagaaagt	gcagaaatga	2100
ttaaatcgc	cgcaatgca	tttctggcga	caaagatttc	ctttatcaac	gatatcgcaa	2160
acatttgatg	gagagtcggc	gcagacgttt	caaaagttgc	tgatgggtgt	ggtcttgaca	2220
gccgtatcgg	cagaaagtcc	cttaaaagctg	gtattggatt	cggcgggttc	tgttttccaa	2280
aggatacaac	cgcgctgctt	caaatcgcaa	aatcggcagg	ctatccattc	aaagctcatcg	2340
aagctgtcat	tgaaacgaac	gaaaagcagc	gtgttcatat	tgtagataaa	cttttgactg	2400
ttatgggaag	cgtcaaaggg	agaaccattt	cagtccctggg	attagccttc	aaaccgaata	2460
cgaacgatgt	gagatccgct	ccagcgtttg	atattatccc	aatgctgcag	cagctgggcg	2520
cccattgaaa	agcatacgat	ccgattgcta	ttcctgaagc	ttcagcgtac	cttggcgaac	2580
aggtcagata	ttacacagat	gtgtatgctg	cgatggaaga	cactgatgca	tgcttgattt	2640
taacggattg	gccggaagtg	aaagaaatgg	agcttgtaaa	agtgaaaacc	ctcttaaac	2700
agccagtcac	cattgacggc	agaaatttat	ttcacttga	agagatgcag	gcagccggat	2760
acatttatca	ctctatcggc	cgtcccgtcg	ttcggggaac	ggaaccctct	gacaagtatt	2820
ttccgggctt	gccgcttgaa	gaattggcta	aagacttggg	aagcgtcaat	ttataagcta	2880

-continued

gaataataag	gaaggtgcct	tttaaatgaa	aaaagtacgt	aaagccataa	ttccagcagc	2940
aggcttagga	acacgttttc	ttccggctac	gaaagcaatg	ccgaaagaaa	tgcttcctat	3000
cgttgataaa	cctaccattc	aatacataat	tgaagaagct	gttgaagccg	gtattgaaga	3060
tattattatc	gtaacaggaa	aaagcaagcg	tgcgattgag	gatcattttg	attactctcc	3120
tgagcttgaa	agaaacctag	aagaaaaagg	aaaaactgag	ctgcttgaaa	aagtgaaaaa	3180
ggcttctaac	ctggctgaca	ttcactatat	ccgcaaaaaa	gaacctaaag	gtctcggaca	3240
tgctgtctgg	tgcgacgca	actttatcgg	cgatgagccg	tttgccgtac	tgcttggtga	3300
cgatattggt	caggctgaaa	ctccaggggt	gcgccaatta	atggatgaat	atgaaaaaac	3360
actttcttct	attatcggtg	ttcagcaggt	gccgaagaa	gaaacacacc	gctacggcat	3420
tattgaccgc	ctgacaagtg	aaggccgccc	ttatcaggtg	aaaaacttcg	ttgaaaaacc	3480
gcctaaaggc	acagcacctt	ctaactctgc	catcttaggc	cgttacgtat	tcacgcctga	3540
gatcttcag	tatttagaag	agcagcaggt	tgccgcccgc	ggagaaattc	agctcacaga	3600
cgccattcaa	aagctgaatg	aaattcaaag	agtgtttgct	tacgattttg	aaggcaagcg	3660
ttatgatggt	ggtgaaaagc	tcggctttat	cacaacaact	cttgaatttg	cgatgcagga	3720
taaagagctt	cgcgatcagc	tcggtccatt	tatggaaggt	ttactaaaca	aagaagaat	3780
ctaagctaga	aataattttg	tttaacttta	agaaggagat	atacatatga	cgcatgtacg	3840
cttgactact	ccaaaagcgt	tgactttctt	tccaacggaa	catgaactta	catacctgcg	3900
ggactttgta	aaaacagcac	accataatat	ccatgagaaa	acaggcgcgg	gcagcgattt	3960
tctaggctgg	gtggacctcc	ctgaacatta	tgataaagaa	gaattcgcgc	gcatccaaaa	4020
aagcgcggaa	aaaatccaat	ctgactctga	tgtcttgctt	gttgtcggca	tcggcggttc	4080
ttatcttgga	gcgcgggcag	cgattgaagc	gctgaatcac	gcgttttata	acactttgcc	4140
aaaagccaaa	cgcggaatc	cgcaagtcac	tttaacttc	tctattaatg	tgatttctaa	4200
atcaggtacg	acaactgaac	ctgcaatcgc	ttccgtatt	ttccgcaagc	ttcttgaaga	4260
gaaatacggg	aaagaagaag	cgaaagcgcg	gatttatgca	acaactgata	aagagcgcgg	4320
cgcatataaa	acgctttcta	acgaagaagg	ctttgaaatc	ttcgtaattc	ctgacgatgt	4380
cgccggccgt	tattcagttt	taacagctgt	aggtctcttg	ccgattgctg	tcagcggcgt	4440
caacattgac	gacatgatga	aaggcgcctt	ggatgcgagc	aaagattttg	caacatctga	4500
actggaagat	aaccagcat	accaatatgc	ggtgttccgc	aatgtccttt	ataataaggg	4560
caaaaacaatt	gaaatgetca	tcaactacga	accggcgtt	caatactttg	cggaatggtg	4620
gaagcagctg	ttcggagaaa	gcgaagggaa	agatgagaag	ggcatttatc	cttcttcagc	4680
gaactattca	acagaccttc	attctttagg	ccagtatgta	caagaaggcc	gcagagattt	4740
attcgaaacg	gtcctgaacg	tagagaagcc	taaacadgaa	ctgacaattg	aggaagcggg	4800
taacgatctt	gacggcttga	actatttagc	cggtaaaact	gttgatttcg	ttaacaaaaa	4860
agcattccaa	ggtacaatgc	ttgccatac	agacggaaat	gttccgaact	taatcgtaa	4920
cattcctgag	ctgaatgcat	atacttttgg	ataccttgta	tatttcttcg	aaaaagcctg	4980
cgcgatgagc	ggttacctcc	ttggcgtaa	tccgtttgac	cagcctggtg	tagaagcgta	5040
taaagtcaat	atgtttgctg	tactcggcaa	acctggcttt	gaagagaaaa	aagcagagct	5100
tgaaaaacgt	ctggaagatt	ataaatgagc	tagcatgact	ggtggacagc	aaatgggtcg	5160
ggatctgtac	gacgatgacg	ataaggatcc	ggtaccggcc	gcatgccggc	taatcgcgac	5220
cggttaacta	gcataacccc	ttggggcctc	taaacgggtc	ttgaggggtt	ttttgctaaa	5280

-continued

ggaggaacta	tatccggtcc	agaattgga	gccaatcaat	tcttgcgag	aactgtgaat	5340
gcgcaaacca	acccttggca	gaacatatcc	atcgcgccg	ccatctccag	cagccgcacg	5400
cggcgcatct	cgggcccgt	tgctggcggt	tttccatagg	ctccgcccc	ctgacgagca	5460
tcacaaaaat	cgacgctcaa	gtcagagggtg	gcgaaacccg	acaggactat	aaagatacca	5520
ggcgtttccc	cctggaagct	ccctcggtcg	ctctcctggt	ccgaccctgc	cgcttacccg	5580
atacctgtcc	gcctttctcc	cttcgggaag	cgtagcgctt	tctcatagct	cacgctgtag	5640
gtatctcagt	tcgggtgtagg	tcggtcgctc	caagctgggc	tgtgtgcacg	aaccccccg	5700
tcagcccagc	cgctgcccct	tatccggtaa	ctatcgctct	gagccaacc	cggtaaagaca	5760
cgacttatcg	ccactggcag	cagccactgg	taacaggatt	agcagagcga	ggtatgtagg	5820
cggtgctaca	gagttcttga	agtgggtggc	taactacggc	tacactagaa	ggacagtatt	5880
tggtatctgc	gctctgctga	agccagttac	cttcggaaaa	agagttggta	gctcttgatc	5940
cggcaaacaa	accaccgctg	gtagcgggtg	tttttttgtt	tgcaagcagc	agattacgcg	6000
cagaaaaaaaa	ggatctcaag	aagatccttt	gatcttttct	acggggtctg	acgctcagtg	6060
gaacgaaaac	tcacgttaag	ggattttggt	catgagatta	tcaaaaagga	tcttcaccta	6120
gatcctttta	aattaaaaat	gaagttttaa	atcaatctaa	agtatatatg	agtaaacttg	6180
gtctgacagt	taccaatgct	taatcagtga	ggcacctatc	tcagcgatct	gtctatttctg	6240
ttcatccata	gttgctgac	tccccgtcgt	gtagataact	acgatacggg	agggcttacc	6300
atctggcccc	agtgtgcaa	tgataccgcg	agaccacgc	tcaccggctc	cagatttatc	6360
agcaataaac	cagccagccg	gaagggccga	gcgcagaagt	ggtcctgcaa	ctttatccgc	6420
ctccatccag	tctattaatt	gttgcccggga	agctagagta	agtagttcgc	cagttaatag	6480
ttgcgcaac	gttggtgcca	ttgctgcagg	catcgtgggtg	tcacgctcgt	cgtttggtat	6540
ggcttcattc	agctccggtt	cccaacgatc	aaggcgagtt	acatgatccc	ccatggtgtg	6600
caaaaaagcg	gtagctcct	tcggtcctcc	gatcgttgtc	agaagtaagt	tgcccgcagt	6660
ggtatcactc	atggttatgg	cagcactgca	taattctctt	actgtcatgc	catccgtaag	6720
atgcttttct	gtgactggtg	agtactcaac	caagtcattc	tgagaatagt	gtatgcggcg	6780
accgagttgc	tcttgcccgg	cgtaacacg	ggataatacc	gcgccacata	gcagaacttt	6840
aaaagtgctc	atcattggaa	aacgttcttc	ggggcgaaaa	ctctcaagga	tcttaccgct	6900
ggtgagatcc	agttcgatgt	aaccactcgt	tgaccccaac	tgatcttcag	catcttttac	6960
tttaccagc	gtttctgggt	gagcaaaaa	aggaaggcaa	aatgccgcaa	aaaagggaa	7020
aagggcgaca	cggaaatggt	gaatactcat	actcttctct	tttcaatatt	attgaagcat	7080
ttatcagggg	tattgtctca	tgagcggata	catatttgaa	tgtatttaga	aaaataaaca	7140
aatagggggt	ccgcgcacat	ttccccgaaa	agtgccacct	gacgtctaag	aaaccattat	7200
tatcatgaca	ttaacctata	aaaataggcg	tatcacgagg	ccctttcgtc	ttcaagaatt	7260
cctgttataa	aaaaaggatc	aattttgaac	tctctcccaa	agttgatccc	ttaacgattt	7320
agaaatccct	ttgagaatgt	ttatatacat	tcaaggtaac	cagccaacta	atgacaatga	7380
ttcctgaaaa	aagtaataac	aaattactat	acagataagt	tgactgatca	acttccatag	7440
gtaacaacct	ttgatcaagt	aagggtatgg	ataataaacc	acctacaatt	gcaataacctg	7500
ttccctctga	taaaagctg	gtaaagttaa	gcaaacctcat	tccagcacca	gcttctctgct	7560
gtttcaagct	acttgaaca	attggtgata	taactgtttt	ggtgaacgaa	agcccaccta	7620

-continued

aaacaaatac	gattataatt	gtcatgaacc	atgatgttgt	ttctaaaaga	aaggaagcag	7680
ttaaaaagct	aacagaaaga	aatgtaactc	cgatgtttaa	cacgtataaa	ggacctcttc	7740
tatcaacaag	tatcccacca	atgtagccga	aaataatgac	actcattggt	ccagggaaaa	7800
taattacact	tccgatttcg	gcagtactta	gctggtgaac	atctttcatc	atataaggaa	7860
ccatagagac	aaacctgct	actgttccaa	atataattcc	cccacaaaga	actccaatca	7920
taaaaggat	atctttccct	aatccgggat	caacaaaagg	atctgttact	ttcctgatat	7980
gttttacaaa	tatcaggaat	gacagcacgc	taacgataag	aaaagaaatg	ctatatgatg	8040
ttgtaacaa	cataaaaaat	acaatgccta	cagacattag	tataattcct	ttgatatcaa	8100
aatgaccttt	tatccttact	tctttcttta	ataatttcat	aagaaacgga	acagtgataa	8160
ttgttatcat	aggaatgagt	agaagatagg	accaatgaat	ataatgggct	atcattccac	8220
caatcgctgg	accgactcct	tctcccatgg	ctactatoga	tccaataaga	ccaaatgctt	8280
taccctatt	ttcctttgga	atatagcgcg	caactacaac	cattacgagt	gctggaaatg	8340
cagctgcacc	agccccttga	ataaaacgag	ccataataag	taaggaaaag	aaagaatggc	8400
caacaaacc	aattaccgac	ccgaaacaat	ttattataat	tccaaatagg	agtaaccttt	8460
tgatgcctaa	ttgatcagat	agctttccat	atacagctgt	tccaatggaa	aaggttaaca	8520
taaaggctgt	gttcaccag	ttgtactcg	cagggtggtt	attaaaatca	tttgcaatat	8580
caggtaatga	gacgttcaaa	accatttcat	ttaatacgct	aaaaaaagat	aaaatgcaaa	8640
gccaaattaa	aatttggttg	tgtcgtaaat	tcgattgtga	ataggatgta	ttcacatttc	8700
accctccaat	aatgagggca	gacgtagttt	atagggttaa	tgatagcctt	ccctctttta	8760
attgaacct	gttacattca	ttacacttca	taattaatc	ctcctaaact	tgattaaaac	8820
attttaccac	atataaacta	agttttaaat	tcagtatttc	atcacttata	caacaatatg	8880
gcccgtttgt	tgaactactc	tttaataaaa	taatttttcc	gttcccaatt	ccacattgca	8940
ataatagaaa	atccatcttc	atcggctttt	tcgtcatcat	ctgtatgaat	caaatcgctt	9000
tcttctgtgt	catcaagggt	taatttttta	tgtatttctt	ttaacaaacc	accataggag	9060
attaaccttt	tacgggtgta	accttctctc	aaatcagaca	aacgtttcaa	attcttttct	9120
tcatcatcgg	tcataaaatc	cgtatccttt	acaggatatt	ttgcagtttc	gtcaattgcc	9180
gattgtatat	ccgatttata	tttatttttc	ggtcgaatca	tttgaacttt	tacatttgga	9240
tcatagteta	atctcattgc	ctttttccaa	aattgaatcc	attgtttttg	attcacgtag	9300
ttttctgtat	tcttaaaata	agttggttcc	acacatacca	atacatgcat	gtgctgatta	9360
taagaattat	ctttattatt	tattgtcact	tccgttgcac	gcataaaacc	aacaagattt	9420
ttattaattt	ttttatattg	catcattegg	cgaaatcctt	gagccatata	tgacaaactc	9480
ttatttaatt	cttcgccatc	ataaacattt	ttaactgtta	atgtgagaaa	caaccaacga	9540
actgttggct	tttgtttaat	aacttcagca	acaacctttt	gtgactgaat	gccatgtttc	9600
attgctctcc	tccagttgca	cattggacaa	agcctggatt	tacaaaacca	cactcgatac	9660
aactttcttt	cgctgtttc	acgattttgt	ttatacteta	atatttcagc	acaatctttt	9720
actctttcag	cctttttaaa	ttcaagaata	tgcagaagtt	caaagtaatc	aacattagcg	9780
atcttctttt	ctctccatgg	tctcactttt	ccactttttg	tcttgtccac	taaaacctt	9840
gatttttcat	ctgaataaat	gctactatta	ggacacataa	tattaaaaga	aacccccatc	9900
tatttagtta	tttgtttgg	cacttataac	tttaacagat	ggggtttttc	tgtgcaacca	9960
attttaaggg	ttttcaatac	tttaaacac	atacatacca	acacttcaac	gcacctttca	10020

-continued

gcaactaaaa taaaaatgac gttatttcta tatgtatcaa gataagaaag aacaagttca 10080
aaaccatcaa aaaaagacac cttttcaggt gcttttttta ttttataaac tcattccctg 10140
atctcgactt cgttcttttt ttacctctcg gttatgagtt agttcaaatt cgtt 10194

<210> SEQ ID NO 3
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: tuaD gene amplification primer

<400> SEQUENCE: 3

atgaaaaaat agctgtcatt ggaacag 27

<210> SEQ ID NO 4
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: tuaD gene amplification primer

<400> SEQUENCE: 4

ttataaattg tcggtcccaa gtct 24

<210> SEQ ID NO 5
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: tuaD gene cloning primer

<400> SEQUENCE: 5

gctggatcca tgaaaaaata gctgtcattg g 31

<210> SEQ ID NO 6
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: tuaD gene cloning primer

<400> SEQUENCE: 6

ctcgctagct tataaattga cggttcccaa g 31

<210> SEQ ID NO 7
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: Amplification primer for the introduction of a Shine-Dalgarno sequence in the tuaDgene

<400> SEQUENCE: 7

cgacatatga aaaaatagct gtcattgg 28

<210> SEQ ID NO 8
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: artificial
<220> FEATURE:
<223> OTHER INFORMATION: Amplification primer for the introduction of a Shine-Dalgarno sequence in the tuaDgene

<400> SEQUENCE: 8

-continued

ctcgctagct tataaattga cgcttcccaa g 31

<210> SEQ ID NO 9
 <211> LENGTH: 26
 <212> TYPE: DNA
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: hasA gene amplification primer

<400> SEQUENCE: 9

atgagaacat taaaaaacct cataac 26

<210> SEQ ID NO 10
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: hasA gene amplification primer

<400> SEQUENCE: 10

taataatttt ttacgtgttc cccag 25

<210> SEQ ID NO 11
 <211> LENGTH: 32
 <212> TYPE: DNA
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: hasA cloning primer

<400> SEQUENCE: 11

ggagatcca tgagaacatt aaaaaacctc at 32

<210> SEQ ID NO 12
 <211> LENGTH: 31
 <212> TYPE: DNA
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: hasA cloning primer

<400> SEQUENCE: 12

cagtctagat tataataatt tttacgtgtc c 31

<210> SEQ ID NO 13
 <211> LENGTH: 30
 <212> TYPE: DNA
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Cloning primer for the introduction of BsrGI e
 BamHI sites upstream and downstream the hasA and tuaDgene

<400> SEQUENCE: 13

gcttgatcat gagaacatta aaaaacctca 30

<210> SEQ ID NO 14
 <211> LENGTH: 30
 <212> TYPE: DNA
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Cloning primer for the introduction of BsrGI e
 BamHI sites upstream and downstream the hasA and tuaDgene

<400> SEQUENCE: 14

aggatcctt ataaattgac gcttcccaag 30

<210> SEQ ID NO 15
 <211> LENGTH: 37

-continued

<212> TYPE: DNA
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: gtaB cloning primer

<400> SEQUENCE: 15

atgtctagaa taataaggaa ggtgcctttt aaatgaa 37

<210> SEQ ID NO 16
 <211> LENGTH: 39
 <212> TYPE: DNA
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: gtaB cloning primer

<400> SEQUENCE: 16

ctctcgagct agcttagatt tcttctttgt ttagtaaag 39

<210> SEQ ID NO 17
 <211> LENGTH: 36
 <212> TYPE: DNA
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: pgi cloning primer

<400> SEQUENCE: 17

tacatatgac gcatgtacgc ttgactactc caaaag 36

<210> SEQ ID NO 18
 <211> LENGTH: 38
 <212> TYPE: DNA
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: pgi cloning primer

<400> SEQUENCE: 18

atgctagctc atttataatc ttccagacgt ttttcaag 38

The invention claimed is:

1. A process for the preparation of hyaluronic acid in *Bacillus megaterium*, comprising the following steps:

(a) culturing bacterial host cells of *Bacillus megaterium*, transformed in a stable way with the T7 RNA polymerase system under conditions suitable for the production of hyaluronic acid in the presence of xylose as an inducer, wherein said bacterial host cells are characterized by being further transformed with:

(i) at least one episomal plasmid vector comprising a sequence coding for the enzyme hyaluronan synthase and a sequence coding for the enzyme UDP-glucose dehydrogenase in tandem under the control of the strong inducible T7 promoter; or

(ii) at least one episomal plasmid vector comprising a sequence coding for the enzyme hyaluronate synthase, a sequence coding for the enzyme UDP-glucose dehydrogenase, a sequence coding for the enzyme UDP-glucose pyrophosphorylase and a sequence coding for the enzyme glucose 6 phosphate isomerase, under the control of the strong inducible T7 promoter;

(b) recovering hyaluronic acid from the culture medium, wherein such bacterial host cells of *Bacillus megaterium* transformed in a stable way with the T7 RNA polymerase system and with plasmid vector (i) or (ii) able to produce hyaluronic acid of step a) are pre-selected on a xylose gradient.

40 2. The process according to claim 1, wherein the xylose inducer is added to a concentration of between 0.1% and 10% w/v.

45 3. The process of claim 2, in which the xylose inducer is added to a concentration of between 0.5% and 1% w/v.

4. The process according to 1, wherein said bacterial host cells of *Bacillus megaterium* transformed with the T7 RNA polymerase system belong to *B. megaterium* strain QM B1551 or DSM319.

50 5. The process according to claim 1, wherein the sequence coding for the enzyme hyaluronan synthase (hasA) is obtained from a strain of *Streptococcus*, and the sequences coding for enzymes UDP-glucose dehydrogenase (hasB or tuaD), UDP-glucose pyrophosphorylase (gtaB) and glucose 6 phosphate isomerase (pgi or hasE) are obtained from *Bacillus subtilis*.

6. The process according to claim 1, in which the sequences coding for the enzyme UDP-glucose dehydrogenase, hyaluronan synthase, UDP-glucose pyrophosphorylase and glucose 6 phosphate isomerase are operatively linked to an upstream Shine-Dalgarno sequence.

7. The process according to claim 1, wherein said plasmid vector (i) comprises the nucleotide sequence of SEQ ID NO:1.

65 8. The process according to claim 1, wherein said plasmid vector (ii) comprises the nucleotide sequence of SEQ ID NO:2.

9. The process according to claim 1, wherein the fermentation time is in the range between 80 and 160 hours and the product HA has a weight average molecular weight in the range 100-500 KDa.

10. The process according to claim 1, wherein the fermentation time is in the range between 40 and 80 hours and the product HA has a weight average molecular weight in the range 500-1000 KDa.

11. The process according to claim 1, wherein the fermentation time is in the range between 12 and 40 hours and the product HA has a weight average molecular weight in the range $1 \times 10^6 - 3 \times 10^6$ D.

12. A process for the preparation of hyaluronic acid in *Escherichia coli*, comprising the following steps:

(a) culturing bacterial host cells of *Escherichia coli*, transformed in a stable way with the T7 RNA polymerase system under conditions suitable for the production of hyaluronic acid in the presence of isopropyl- β -thio-galactopyranoside (IPTG) as an inductor, wherein said bacterial host cells are characterised by being further transformed with:

(i) at least one episomal plasmid vector comprising a sequence coding for the enzyme hyaluronan synthase and a sequence coding for the enzyme UDP-glucose dehydrogenase in tandem under the control of the strong inducible T7 promoter; or

(ii) at least one episomal plasmid vector comprising a sequence coding for the enzyme hyaluronate synthase, a sequence coding for the enzyme UDP-glucose dehydrogenase, a sequence coding for the enzyme UDP-glucose pyrophosphorylase and a sequence coding for the enzyme glucose 6 phosphate isomerase, under the control of the strong inducible T7 promoter;

(b) recovering hyaluronic acid from the culture medium, wherein such bacterial host cells of *Escherichia coli* transformed in a stable way with the T7 RNA polymerase system and with plasmid vector (i) or (ii) able to produce hyaluronic acid of step a) are pre-selected on an IPTG gradient.

13. The process according to claim 12, wherein the IPTG inducer is added to a concentration of between 0.1 mM and 10 mM.

14. The process of claim 13, in which the IPTG inducer is added to a concentration of between 0.4 mM to 1.0 mM.

15. The process according to claim 12, wherein said bacterial host cells of *Escherichia coli* transformed with the T7 RNA polymerase system belong to *E. coli* strain BL21 DE3.

16. The process according to claim 12, wherein the sequence coding for the enzyme hyaluronan synthase (hasA) is obtained from a strain of *Streptococcus*, and the sequences coding for enzymes UDP-glucose dehydrogenase (hasB or tuaD), UDP-glucose pyrophosphorylase (gtaB) and glucose 6 phosphate isomerase (pgi or hasE) are obtained from *Bacillus subtilis*.

17. The process according to claim 12, in which the sequences coding for the enzyme UDP-glucose dehydrogenase, hyaluronan synthase, UDP-glucose pyrophosphorylase and glucose 6 phosphate isomerase are operatively linked to an upstream Shine-Dalgarno sequence.

18. The process according to claim 12, wherein said plasmid vector (i) comprises the nucleotide sequence of SEQ ID NO:1.

19. The process according to claim 12, wherein said plasmid vector (ii) comprises the nucleotide sequence of SEQ ID NO:2.

20. A plasmid vector comprising a strong inducible bacteriophage T7 promoter operationally linked to a sequence coding for a hyaluronan synthase enzyme and a sequence coding for a UDP-glucose dehydrogenase enzyme in tandem.

21. The plasmid vector according to claim 20, wherein said sequence coding for the enzyme hyaluronan synthase is a hasA gene from *Streptococcus zooepidemicus*, and said sequence coding for the enzyme UDP-glucose dehydrogenase is a tuaD gene from *Bacillus subtilis*.

22. The plasmid vector according to claim 21, comprising the nucleotide sequence SEQ ID NO:1.

23. A plasmid vector comprising a strong inducible bacteriophage T7 promoter operationally linked to a sequence coding for a hyaluronate synthase enzyme, a sequence coding for a UDP-glucose dehydrogenase enzyme, a sequence coding for a UDP-glucose pyrophosphorylase enzyme and a sequence coding for a glucose 6 phosphate isomerase enzyme.

24. The plasmid vector according to claim 23, wherein said sequence coding for the enzyme hyaluronan synthase is a hasA gene from *Streptococcus zooepidemicus*, said sequence coding for the enzyme UDP-glucose dehydrogenase is a tuaD gene from *Bacillus subtilis*, said sequence coding for the enzyme UDP-glucose pyrophosphorylase is a gtaB gene from *Bacillus subtilis* and said sequence coding for the enzyme glucose 6 phosphate isomerase is a pgi gene from *Bacillus subtilis*.

25. The plasmid vector according to claim 24, comprising the nucleotide sequence SEQ ID NO:2.

26. The plasmid vector according to claim 23, wherein the sequence coding for the enzyme UDP-glucose dehydrogenase, hyaluronan synthase, UDP-glucose pyrophosphorylase and glucose 6 phosphate isomerase are operatively linked to an upstream Shine-Dalgarno sequence.

27. A recombinant host bacterial cell belonging to the genus *Bacillus* previously transformed with the T7 RNA polymerase system, comprising at least one plasmid vector according to claim 20.

28. The recombinant host bacterial cell according to claim 27, which is *Bacillus megaterium*.

29. A method for obtaining recombinant host bacterial cells, according to claim 27, which are capable of producing high levels of hyaluronic acid, comprising selecting bacterial host cells transformed with a plasmid vector comprising a strong inducible bacteriophage T7 promoter operationally linked to a sequence coding for a hyaluronan synthase enzyme and a sequence coding for a UDP-glucose dehydrogenase enzyme in tandem and transformed with a T7 RNA polymerase system, on a xylose gradient.

30. A recombinant host bacterial cell belonging to the genus *Escherichia* previously transformed with the T7 RNA polymerase system, comprising at least one plasmid vector according to claim 20.

31. The recombinant host bacterial cell according to claim 30, which is *Escherichia coli*.

32. A method for obtaining recombinant host bacterial cells, according to claim 30, which are capable of producing high levels of hyaluronic acid, comprising selecting bacterial host cells transformed with a plasmid vector comprising a strong inducible bacteriophage T7 promoter operationally linked to a sequence coding for a hyaluronan synthase enzyme and a sequence coding for a UDP-glucose dehydrogenase enzyme in tandem and transformed with a T7 RNA polymerase system, on an IPTG gradient.

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 9,695,453 B2
APPLICATION NO. : 14/884274
DATED : July 4, 2017
INVENTOR(S) : Vincenza Corsa et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

On the Title Page

At item (72), Inventors, change “**Vincenza Corsa, Albano Terme (IT)**” to --**Vincenza Corsa, Abano Terme (IT)**--.

At item (63), **Related U.S. Application Data**, change “Continuation of application No. 13/821,953, filed as application No. PCT/EP2011/065641 on Apr. 15, 2013, now Pat. No. 9,163,270.” to --Continuation of application No. 13/821,953, filed as application No. PCT/EP2011/065641 on Sep. 9, 2011, new Pat No. 9,163,270.--.

Signed and Sealed this
Twenty-fourth Day of October, 2017



Joseph Matal

*Performing the Functions and Duties of the
Under Secretary of Commerce for Intellectual Property and
Director of the United States Patent and Trademark Office*