

US009725701B2

(12) **United States Patent**
McLaren et al.

(10) **Patent No.:** **US 9,725,701 B2**
(45) **Date of Patent:** **Aug. 8, 2017**

(54) **PROKARYTOIC-TYPE ISOCITRATE DEHYDROGENASE AND ITS APPLICATION FOR IMPROVING NITROGEN UTILIZATION IN TRANSGENIC PLANTS**

(71) Applicant: **Iowa Corn Promotion Board,**
Johnston, IA (US)

(72) Inventors: **James McLaren,** Chesterfield, MO
(US); **Brian Vande Berg,** Morrisville,
NC (US)

(73) Assignee: **Iowa Corn Promotion Board,**
Johnston, IA (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 966 days.

(21) Appl. No.: **13/833,247**

(22) Filed: **Mar. 15, 2013**

(65) **Prior Publication Data**

US 2014/0283224 A1 Sep. 18, 2014

(51) **Int. Cl.**
C12N 15/29 (2006.01)
C12N 15/82 (2006.01)
C12N 9/04 (2006.01)

(52) **U.S. Cl.**
CPC **C12N 9/0006** (2013.01); **C12N 15/8261**
(2013.01)

(58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

6,121,513 A	9/2000	Zhang et al.	
6,204,039 B1 *	3/2001	Allen	C12N 9/0006 435/190
6,562,958 B1	5/2003	Breton et al.	
6,673,910 B1	1/2004	Breton	
6,995,250 B1	2/2006	Hirano et al.	
7,214,786 B2	5/2007	Kovalic et al.	
7,504,111 B2	3/2009	Fontana et al.	
7,517,684 B2	4/2009	Rubinfeld et al.	
7,569,389 B2	8/2009	Feldmann et al.	
7,630,836 B2	12/2009	Omura et al.	
7,777,097 B2	8/2010	Glazebrook et al.	
7,862,827 B2	1/2011	Giuliani et al.	
7,867,704 B2	1/2011	Kapur et al.	
8,247,650 B2	8/2012	Hershey et al.	
2011/0252503 A1 *	10/2011	McLaren	C12N 9/1217 800/278

FOREIGN PATENT DOCUMENTS

CN	101421295	4/2009
CN	102711445	10/2012
WO	9928479	6/1999
WO	2011056769	5/2011

OTHER PUBLICATIONS

Lemaitre et al. (Plant Physiology, Jul. 2007, vol. 144, pp. 1546-1558).*

Sahara et al. (Biosci. Biotechnol. Biochem. 66:489-500(2002)).*

Guo et al (2004, Proc. Natl. Acad. Sci. USA 101: 9205-9210).*

Liu et al. (J. Plant Biol. (2010) 53:107-112).*

Chen, Ridong, "Plant NADP-Dependent Isocitrate Dehydrogenases are Predominantly Localized in the Cystol", "Planta", 1998, pp. 280-285, vol. 207, Published in: US.

Foyer, C. et al., "Respiration and Nitrogen Assimilation: Targeting Mitochondria-associates Metabolism as a Means to Enhance Nitrogen Use Efficiency", "Journal of Experimental Biology", 2011, pp. 1467-1482, vol. 62, No. 4.

Igamberdiev, A. et al., "Regulation of NAD- and NADP-dependent Isocitrate Dehydrogenases by Reduction Levels of Pyridine Nucleotides in Mitochondria and Cytosol of Pea Leaves", "Biochim Biophys Acta", Sep. 30, 2003, pp. 117-125, vol. 1606.

Leterrier, M. et al., "Cytosolic NADP-Isocitrate Dehydrogenase of Pea Plants: Genomic Clone Characterization and Fuctional Analysis Under Abiotic Stress Conditions", "Free Radical Research", Feb. 1, 2007, pp. 191-199, vol. 41, No. 2.

Sienkiewicz-Porzucek, A. et al., "Mild Reductions in Mitochondrial NAD-Dependent Isocitrate Dehydrogenase Activity Result in Altered Nitrate Assimaltion and Pigmentation But Do Not Impact Growth", "Molecular Plant", 2009, pp. 156-173, vol. 3, No. 1.

Sahara et al., "Cloning, Sequencing, and Expression of a Gene Encoding the Monomeric Isocitrate Dehydrogenase of the Nitrogen-fixing Bacterium", 2002, pp. 589-500, vol. 66, No. 3, Publisher: Bioscience Biotechnology Biochemistry, Published in: Sapporo, Japan.

Sahara et al., "Isocitrate dehydrogenase", "GenBank Abstract", Jul. 27, 2011.

Popova et al., "Citrate and isocitrate in plant metabolism", "Biochimica et Biophysica Ada", May 27, 1998, pp. 307-325, vol. 1364, No. 3, Published in: NL.

Anonymous, "UNIPROT:P16100", "http://bis/exam/dbfetch.jsp?id=UNIPROTP16100", Jan. 23, 2007.

* cited by examiner

Primary Examiner — Shubo (Joe) Zhou

Assistant Examiner — Charles Logsdon

(74) *Attorney, Agent, or Firm* — Davis, Brown, Koehn,
Shors & Roberts, P.C.; Kent A. Herink; Matthew W.
Coryell

(57) **ABSTRACT**

The present invention relates to transgenic plants that have increased nitrogen use efficiency, stress tolerance, and/or alleviating a limitation such that yield is increased, or a combination of these and that have been transformed using a novel vector construct including a synthetic isocitrate dehydrogenase (icdh) gene that modulates nitrogen use in plants. The invention also relates to stacking the icdh gene with other exogenous or heterologous genes that modulate nitrogen use in the plant, including a N-acetylglutamate kinase gene. The invention also relates to methods of expressing in plants the nucleic acid molecules corresponding to the nucleic acid sequences that modulate nitrogen use in plants or are modulated by nitrogen conditions.

23 Claims, 2 Drawing Sheets




FIG. 1




FIG. 2

**PROKARYOTIC-TYPE ISOCITRATE
DEHYDROGENASE AND ITS APPLICATION
FOR IMPROVING NITROGEN UTILIZATION
IN TRANSGENIC PLANTS**

REFERENCE TO THE SEQUENCE LISTING

Applicant hereby incorporates by reference material submitted via Web-EFS in the following file:

File Name	Date Created	Size
13-833247_5T25.txt	August 27, 2013	16 kb

FIELD OF THE INVENTION

The invention relates generally to plants with improved nitrogen utilization and stress tolerance, more specifically, to heterologous expression of an isocitrate dehydrogenase (ICDH) enzyme in plants, including the overexpression and characterization of a prokaryotic-based isocitrate dehydrogenase that improves stress tolerance and nitrogen uptake, metabolism or both. The invention also includes stacking of the *icdh* gene with one or more other transgenes to improve nitrogen utilization and/or stress tolerance.

BACKGROUND OF THE INVENTION

Plants require nitrogen during their vegetative and reproductive growth phases. Nitrogen is made available to the plant through soil mineralization, the application of nitrogen fertilizer, or both. It has been estimated, however, that between 50 and 70 percent of nitrogen applied to crops is lost from the plant-soil system [Peoples, M. B. et al., "Minimizing Gaseous Losses of Nitrogen," In *Nitrogen Fertilizer in the Environment* (Bacon, P. E., ed.) Marcel Dekker, pp. 565-606 (1995)]. Nitrogen is one of the most expensive plant nutrients to supply, nitrogen fertilizer is not always available at a reasonable cost, and excessive application of nitrogen fertilizer can result in environmental challenges. Corn is an example of an agronomically important plant that often requires nitrogen fertilizers to perform at its genetic potential.

Native ICDH can exist in the mitochondria, chloroplast and cytosol, with each having a different physiological impact although the catalytic action may be similar. In general, ICDH1 is found in the cytosol and ICDH2 is found in the chloroplast.

For co-factor reducing power, ICDH can use either nicotinamide adenine dinucleotide (NAD⁺) or nicotinamide adenine dinucleotide phosphate (NADP⁺), depending on which metabolic pathway it is active. Some publications indicate that the main function of ICDH may be to generate reducing power (NADH, NADPH) for other metabolic reactions, for example, in the β -oxidation of unsaturated fatty acids. Other theories include the suggestion that the reaction product, 2-oxoglutarate (OG), could be used to support amino acid synthesis via the GOGAT cycle (Hodges, M. Enzyme redundancy and the importance of 2-oxoglutarate in plant ammonium assimilation. *J. Exp. Botany* (2002), 53, 905). In addition, the over-expression of the ICDH enzyme in a stack with another gene or genes may allow the effective utilization of the additional carbon skeletons. A previous study of transgenic tobacco plants that overexpressed a mitochondrial *icdh* gene was focused on redox pathways and

did not mention nor evaluate any possible impact on nitrogen utilization (Gray, G., Villarimo, A., Whitehead, C., McIntosh, L. Transgenic Tobacco (*Nicotiana tabacum* L.) Plants with Increased Expression Levels of Mitochondrial NADP⁺-dependent Isocitrate Dehydrogenase: Evidence Implicating this Enzyme in the Redox Activation of the Alternative Oxidase, *Plant and Cell Physiology* 2004; 45, 1413-1425).

Regulation of NAD- and NADP-dependent isocitrate dehydrogenases (NAD-ICDH, EC 1.1.1.41 and NADP-ICDH, EC 1.1.1.42) is complex due to expression, substrates, compartments and post-translational regulation. While it is unclear which ICDH version generates OG for amino acids, any such OG would have to be in, or enter, the chloroplast where nitrogen is assimilated into amino acids. The literature suggests that plant cytosolic versions of ICDH are homodimers with subunits of approximately 47 kD. Mitochondrial ICDH is suspected to have more subunits. Bacterial versions of ICDH may be monomeric and have been considered to overcome the typical regulation of expression and function that occurs with plant ICDH in plants, that is, phosphorylation may inactivate the homodimer.

Cytosolic NADP-specific ICDH catalyzes the conversion of citrate to oxoglutarate. One strategy is to design a construct containing a gene encoding a monomeric prokaryotic-type isocitrate dehydrogenase gene (*icdh*), and to direct overexpression of ICDH in the cytoplasm of plants. The expressed ICDH enzyme will enhance the plant's ability to utilize available nitrogen via an enhanced flow of carbon into the nitrogen assimilatory mechanism. Here, we describe the overexpression and characterization of a synthetic *icdh* gene based on selection from among bacterial *icdh* sequences and optimized for expression in corn, and the stacking of a *icdh* genes with other transgenes.

SUMMARY OF THE INVENTION

The present invention relates to transgenic plants that have increased nitrogen use efficiency, stress tolerance, or both, that have been transformed using a novel vector construct including an *icdh* nucleic acid sequence that modulates nitrogen use in plants. A variety of *icdh* nucleic acid sequences were identified for use with the present invention from the several bacterial and plant genomic sequencing projects that have been archived in public databases from which sequences that encode ICDH enzymes with robust activity could be selected. These candidate *icdh* sequences were then screened to deselect those that had a relatively high content of poly A regions, which can be inhibitory to expression in plants. The sequence chosen to exemplify these *icdh* sequences was then codon-optimized for expression in maize (SEQ ID No 1). The invention also includes stacking an *icdh* gene with one or more heterologous genes so as to induce the over-expression of the ICDH enzyme in combination with nitrogen assimilatory enzymes. The invention also relates to isolated vectors for transforming plants and to antibodies for detecting expression of the nucleotide sequence(s) of interest in the transformed plants. The invention also relates to methods of expressing in plants the nucleic acid molecules corresponding to the nucleic acid sequences that modulate nitrogen use in plants.

Specifically, vectors for transforming plants and bacterial cells have been constructed using the nucleotide sequences SEQ ID NO: 1 and 3, as well as combinations, variants, fragments, and complements thereof. These vectors include a 5' DNA promoter sequence and a 3' terminator sequence,

wherein the nucleic acid sequence, the DNA promoter sequence, and the terminator sequence are operatively coupled to permit transcription of the nucleotide sequence. In some embodiments, the promoter sequence may be a constitutive plant promoter or a tissue specific promoter.

The invention also includes polyclonal antibodies, comprising polyclonal antibodies to a polypeptide encoded by nucleotide sequences SEQ ID NO: 1 and 3 and combinations thereof.

The invention also includes plants transformed with a nucleotide sequences SEQ ID NO: 1 and 3, as well as combinations, variants and fragments thereof. The plant is selected from the group consisting of corn (maize), sorghum, wheat, sunflower, tomato, crucifers, peppers, potato, cotton, rice, soybean, sugarbeet, sugarcane, tobacco, barley, and oilseed rape, *Brassica* sp., alfalfa, rye, millet, safflower, peanuts, sweet potato, cassava, coffee, coconut, pineapple, citrus trees, cocoa, tea, banana, avocado, fig, guava, mango, olive, papaya, cashew, macadamia, almond, oats, vegetables, grasses (such as turf grasses, forage grasses, or pasture grasses), ornamentals, trees (such as fruit trees, nut trees, pulp trees, oil palms) and conifers. The invention also includes a component part of such plants, plant seed produced from such plants, and a plant seed transformed with a vector construct of the present invention.

The invention also includes a host cell transformed with a nucleotide sequence selected from SEQ ID NO: 1 and 3, and combinations thereof. The host cell may be a bacterial cell or a plant cell.

The invention also includes a method of expressing a nucleic acid molecule that modulates nitrogen in a plant, said method comprising the steps of providing a transgenic plant or plant seed transformed with a vector construct according to the present invention, and growing the transgenic plant or a plant grown from the transgenic plant seed under conditions effective to express the nucleic acid molecule in said transgenic plant or said plant grown from the transgenic plant seed. Growing of the transgenic plant is effective in increasing nitrogen uptake of said transgenic plant or said plant grown from the transgenic plant seed, and/or in increasing efficiency of nitrogen utilization of said transgenic plant or said plant grown from the transgenic plant seed, and/or alleviating a limitation such that yield is increased in said transgenic plant or said plant grown from the transgenic plant seed. The invention also includes the foregoing methods wherein a transgenic plant is provided or a transgenic seed is provided. The invention also includes the foregoing method wherein the plant is selected from the group consisting of corn (maize), sorghum, wheat, sunflower, tomato, crucifers, peppers, potato, cotton, rice, soybean, sugarbeet, sugarcane, tobacco, barley, and oilseed rape, *Brassica* sp., alfalfa, rye, millet, safflower, peanuts, sweet potato, cassava, coffee, coconut, pineapple, citrus trees, cocoa, tea, banana, avocado, fig, guava, mango, olive, papaya, cashew, macadamia, almond, oats, vegetables, grasses (such as turf grasses, forage grasses, or pasture grasses), ornamentals, trees (such as fruit trees, nut trees, pulp trees, oil palms) and conifers.

The invention also includes a method of improving the stress tolerance of a plant by expressing a nucleic acid molecule modulated by nitrogen in a plant, said method comprising the steps of providing a transgenic plant or plant seed transformed with a vector construct according to the present invention and growing the transgenic plant or a plant grown from the transgenic plant seed under conditions

effective to express the nucleic acid molecule in said transgenic plant or said plant grown from the transgenic plant seed.

The invention also includes a method of altering the morphology of a plant by expressing a nucleic acid molecule modulated by nitrogen in a plant, said method comprising the steps of providing a transgenic plant or plant seed transformed with a vector construct according to the present invention and growing the transgenic plant or a plant grown from the transgenic plant seed under conditions effective to express the nucleic acid molecule in said transgenic plant or said plant grown from the transgenic plant seed.

The invention also includes a vector construct, comprising a nucleotide sequence encoding the ICDH amino acid sequence including SEQ ID NO: 2 and 4, and combinations thereof, a 5' DNA promoter sequence, and a 3' terminator sequence, wherein the nucleotide sequence, the DNA promoter sequence, and the terminator sequence are operatively coupled to permit transcription of the nucleotide sequence.

The invention also includes a vector construct comprising a nucleotide sequence that modulates nitrogen in a plant, wherein said nucleotide sequence is selected from SEQ ID NO: 1 and 3, and combinations thereof; a nucleotide sequence having at least 85% sequence identity to the corresponding nucleotide sequence of SEQ ID NO: 1 and 3, and combinations thereof, wherein said nucleotide sequence modulates nitrogen in a plant; a nucleotide sequence selected from those encoding the ICDH amino acid sequences SEQ ID NO: 2 and 4, and combinations thereof; and, a nucleotide sequence encoding an amino acid sequence having at least 85% sequence identity to the amino acid sequence of SEQ ID NO: 2 and 4, and combinations thereof; wherein said nucleotide sequence modulates nitrogen in a plant, wherein said construct further comprises a 5' DNA promoter sequence and a 3' terminator sequence, wherein the nucleotide sequence, the DNA promoter sequence, and the terminator sequence are operatively coupled to permit transcription of the nucleotide sequence.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 is a vector map for the plasmid pMDO8901, wherein the Main elements in the plasmid (clockwise from top) are: Right Border, ScUbi4 promoter, 5' UTR exon, intron, *icdh* gene, 35S terminator, ScUbi4 promoter, 5' UTR exon, intron, chloroplast transit peptide from EPSPS, *nagk* gene, 35S terminator, ScUbi4 promoter, 5' UTR exon, intron, chloroplast transit peptide from EPSPS, glyphosate tolerance SM (GRG23ac35), 35S terminator, Left Border.

FIG. 2 is a vector map for the plasmid pMDO8902, wherein the main elements in the plasmid (clockwise from top) are: Right Border, ScUbi4 promoter, 5' UTR exon, intron, *icdh* gene, 35S terminator, ScUbi4 promoter, 5' UTR exon, intron, chloroplast transit peptide from EPSPS, *nagk* gene, 35S terminator, ScUbi4 promoter, 5' UTR exon, intron, glyphosate tolerance SM, 35S terminator, Left Border.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

The development of plant varieties that use nitrogen more efficiently will reduce the need for excessive inputs of nitrogen, save production costs for farmers, benefit farmers in developing countries who do not have access to fertilizer inputs, and reduce environmental contamination associated with the application of excessive nitrogen fertilizers. One

approach that has been used in the development of plant varieties with improved nitrogen utilization relies on conventional plant breeding techniques. However, such approaches have had variable success due to lack of specification in the genetic recombination.

There is a need to develop plant cultivars that absorb and use nitrogen more efficiently. Plant scientists have adopted the shorthand term nitrogen use efficiency (NUE), and a variety of methods of measuring and evaluating NUE have been developed [Craswell, E. T. and Godwin, D. C. (1984) The efficiency of nitrogen fertilizers applied to cereals grown in different climates. In *Advances in Plant Nutrition* (Vol. 1) (Tinker, P. B. and Lauchli, A., eds), pp. 1-55, Praeger Publishers; Steenbjerg, F. and Jakobsen, S. T. (1963) Plant nutrition and yield curves. *Soil Sci.* 95, 69-90; Siddiqi, M. Y. and Glass, D. M. (1981) Utilization index: a modified approach to the estimation and comparison of nutrient utilization efficiency in plants. *J. Plant Nutr.* 4, 289-302; Moll, R. H. et al. (1982) Analysis and interpretation of factors which contribute to efficiency of nitrogen utilization. *Agron. J.* 74, 562-564]. There are differences in the specific definitions, and context of use. For example, some definitions are based on total biomass while others are based on the weight of grain yielded. Another set of definitions uses the efficiency of extracting nitrogen from the soil. The efficiency with which applied nitrogen is used to improve grain yield may be measured by agronomic efficiency (AE), the product of physiological efficiency and utilization efficiency, or NUE_g which is the product of uptake efficiency and utilization efficiency. Other definitions take physiological factors into account.

As used in this specification, the term nitrogen use efficiency, or NUE, is defined to include a measurable change in any of the main nitrogen metabolic pool sizes in the assimilation pathways (for example, may include a measurable change in one or more of the following: nitrate, nitrite, ammonia, glutamic acid, aspartic acid, glutamine, asparagine, lysine, leucine, threonine, methionine, glycine, tryptophan, tyrosine, total protein content of a plant part, total nitrogen content of a plant part, and/or chlorophyll content), or where the plant is shown to provide the same or elevated biomass or harvestable yield at lower nitrogen fertilization levels, or where the plant is shown to provide elevated biomass or harvestable yields at the same nitrogen fertilization levels when compared to a plant that has not been transformed with a nitrogen-modulating nucleic acid construct of the invention. A "measurable change" can include an increase or a decrease in the amount of any component ("metabolic pool") of the nitrogen assimilation pathway. A change can include either a decrease or an increase in one or more metabolic pools in the pathway, or a decrease in one or more pools with a concomitant increase in one or more other pool(s), such as when one intermediate in the nitrogen assimilation pathway is being utilized for the purpose of generating another intermediate or product of the pathway. For example, in the conversion of glutamate to glutamine, the level of glutamate may decrease while the level of glutamine may increase. Thus, while not being bound by any particular theory or mechanism, any change in one or more of these pools indicates that nitrogen is being utilized more efficiently by the plant.

An increase in nitrogen utilization efficiency can be associated with about a 5%, about a 10%, 15%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 125%, 150%, about a 200% or greater measurable change in any of the main nitrogen metabolic pool sizes in the assimilation pathway. In one embodiment, the transgenic plants of the

invention have an increased nitrogen uptake from the environment when compared to a plant that does not contain a nitrogen-modulating sequence of the invention. By "nitrogen modulating sequence" it is intended to mean a nucleotide or amino acid sequence that modulates NUE, by way of non-limiting example: either by generating an enzyme that impacts NUE, or by generating a protein that interacts with the components involved in NUE, or by generating a protein that impacts the internal homeostatic signal cascade regulating NUE, or by a combination of these mechanisms that results in a measurable change in N uptake, N assimilation, N metabolism, N transport, N utilization, N storage, or a combinations of these. The present invention further provides a method of improving stress tolerance in a plant by expressing one or more nitrogen-modulating nucleotide sequences within the plant. In one embodiment, the nitrogen-modulating nucleotide sequence is SEQ ID NO: 1, or variants and fragments thereof. In another embodiment, the nitrogen-modulating nucleotide sequence is a nucleotide sequence that encodes SEQ ID NO: 2, or variants and fragments thereof. In another embodiment, the nitrogen-modulating nucleotide sequence is a nucleotide sequence that encodes SEQ ID NO 1 plus SEQ ID NO: 2, or variants and fragments thereof, respectively.

As used herein, the term "stress" or "stress condition" refers to the exposure of a plant, plant cell, or the like, to a physical, environmental, biological or chemical agent or condition that has an adverse effect on metabolism, growth, development, propagation and/or survival of the plant (collectively "growth"). A stress can be imposed on a plant due, for example, to an environmental factor such as water (e.g., flooding, drought, dehydration), anaerobic conditions (e.g., a low level of oxygen), abnormal osmotic conditions, salinity or temperature (e.g., hot/heat, cold, freezing, frost), a deficiency of nutrients such as nitrogen, phosphate, potassium, sulfur, micronutrient, or exposure to pollutants, or by a hormone, second messenger or other molecule. Anaerobic stress, for example, is due to a reduction in oxygen levels (hypoxia or anoxia) sufficient to produce a stress response. A flooding stress can be due to prolonged or transient immersion of a plant, plant part, tissue or isolated cell in a liquid medium such as occurs during monsoon, wet season, flash flooding or excessive irrigation of plants, or the like. A cold stress or heat stress can occur due to a decrease or increase, respectively, in the temperature from the optimum range of growth temperatures for a particular plant species. Such optimum growth temperature ranges are readily determined or known to those skilled in the art. Dehydration stress can be induced by the loss of water, reduced turgor, or reduced water content of a cell, tissue, organ or whole plant. Drought stress can be induced by or associated with the deprivation of water or reduced supply of water to a cell, tissue, organ or organism. Saline stress (salt stress) can be associated with or induced by a perturbation in the osmotic potential of the intracellular or extracellular environment of a cell. Osmotic stress also can be associated with or induced by a change, for example, in the concentration of molecules in the intracellular or extracellular environment of a plant cell, particularly where the molecules cannot be partitioned across the plant cell membrane.

An improvement in stress tolerance can be assessed by any quantitative or qualitative measure of plant performance under a given stress condition and is relative to the performance of a plant grown under the same stress conditions that has not been transformed with a nitrogen-modulating sequence of the invention. Thus, the plants may exhibit improved nitrogen contents, altered amino acid or protein

compositions, altered carbohydrate composition, altered oil composition, vigorous growth characteristics, increased vegetative yields or better seed yields and qualities. These plants may be identified by examining any of following parameters: 1) the rate of growth, measured in terms of rate of increase in fresh or dry weight; 2) vegetative yield of the mature plant, in terms of fresh or dry weight; 3) the seed or fruit yield; 4) the seed or fruit weight; 5) the total nitrogen content of the plant; 6) the total nitrogen content of the fruit or seed; 7) the free amino acid content of the plant; 8) the free amino acid content of the fruit or seed; 9) the total protein content of the plant; 10) the total protein content of the fruit or seed; 11) measurable change in carbohydrates or oils. The procedures and methods for examining these parameters are well known to those skilled in the art. These methods may involve enzymatic assays and immunoassays to measure enzyme/protein levels; assays to measure the amino acid composition, free amino acid pool or total nitrogen content of various plant tissues; measurement of growth rates in terms of fresh weight gains over time; or measurement of plant yield in terms of total dry weight and/or total seed weight.

Transformation of Bacterial or Plant Cells

Provided herein are novel nucleotide sequences that modulate nitrogen utilization efficiency in plants. Also provided are amino acid sequences of the proteins of the invention, that may be nitrogen-modulating or modulated by nitrogen concentration.

The nitrogen-modulating nucleotide sequences of the invention may be modified to obtain or enhance expression in plant cells. The nitrogen-modulating sequences of the invention may be provided in expression cassettes for expression in the plant of interest. "Plant expression cassette" includes DNA constructs that are capable of resulting in the expression of a protein from an open reading frame in a plant cell. The cassette will include in the 5'-3' direction of transcription, a transcriptional initiation region (i.e., promoter) operably-linked to a DNA sequence of the invention, and a transcriptional and translational termination region (i.e., termination region) functional in plants. The cassette may additionally contain at least one additional gene to be co-transformed into the organism, such as a selectable marker gene or a stacked gene of different function. Alternatively, the additional gene(s) can be provided on multiple expression cassettes. Such an expression cassette is provided with a plurality of restriction sites for insertion of the nitrogen-modulating sequence to be under the transcriptional regulation of the regulatory regions.

By "promoter" is intended a nucleic acid sequence that functions to direct transcription of a downstream coding sequence. The promoter, together with other transcriptional and translational regulatory nucleic acid sequences (also termed as "control sequences"), are necessary for the expression of a DNA sequence of interest. Preferably, the promoter is one that is known to stimulate transcription in the organism into which the nucleotide sequence of the invention is being introduced.

The promoter may be native or analogous, or foreign or heterologous, to the plant host and/or to the DNA sequence of the invention. Additionally, the promoter may be the natural sequence or alternatively a synthetic sequence. Where the promoter is "native" or "homologous" to the plant host, it is intended that the promoter is found in the native plant into which the promoter is introduced. Where the promoter is "foreign" or "heterologous" to the DNA sequence of the invention, it is intended that the promoter is not the native or naturally occurring promoter for the

operably linked DNA sequence of the invention. "Heterologous" generally refers to the nucleic acid sequences that are not endogenous to the cell or part of the native genome in which they are present, and have been added to the cell by infection, transfection, microinjection, electroporation, microprojection, or the like. By "operably linked" is intended a functional linkage between a promoter and a second sequence, wherein the promoter sequence initiates and mediates transcription of the DNA sequence corresponding to the second sequence. Generally, "operably linked" means that the nucleic acid sequences being linked are contiguous, including exons and introns and, where necessary to join two protein coding regions, contiguous and in the same reading frame.

In one embodiment, the promoter is a constitutive promoter. Suitable constitutive promoters for use in plants include: the promoters from plant viruses, such as the peanut chlorotic streak caulimovirus (PC1SV) promoter (U.S. Pat. No. 5,850,019); the 35S promoter from cauliflower mosaic virus (CaMV) (Odell et al. (1985) *Nature* 313:810-812); promoters of *Chlorella* virus methyltransferase genes (U.S. Pat. No. 5,563,328) and the full-length transcript promoter from figwort mosaic virus (FMV) (U.S. Pat. No. 5,378,619); the promoters from such genes as rice actin (McElroy et al. (1990) *Plant Cell* 2:163-171); ubiquitin (Christensen et al. (1989) *Plant Mol. Biol.* 12:619-632 and Christensen et al. (1992) *Plant Mol. Biol.* 18:675-689), including the TrpPro5 promoter (U.S. patent application Ser. No. 10/377,318; filed Mar. 16, 2005); pEMU (Last et al. (1991) *Theor. Appl. Genet.* 81:581-588); MAS (Velten et al. (1984) *EMBO J.* 3:2723-2730); maize H3 histone (Lepetit et al. (1992) *Mol. Gen. Genet.* 231:276-285 and Atanassova et al. (1992) *Plant J.* 2(3):291-300); *Brassica napus* ALS3 (PCT application WO 97/41228); and promoters of various *Agrobacterium* genes (see U.S. Pat. Nos. 4,771,002; 5,102,796; 5,182,200; and 5,428,147).

In another embodiment, the promoter is a tissue-specific promoter. A list of commonly-used tissue-specific promoters can be found in Reviewed in Moore et al. (2006) *Plant J.* 45(4):651-683, which is herein incorporated by reference in its entirety.

Often, such constructs will also contain 5' and 3' untranslated regions. Such constructs may contain a "signal sequence" or "leader sequence" to facilitate co-translational or post-translational transport of the peptide of interest to certain intracellular structures such as the chloroplast (or other plastid), endoplasmic reticulum, or Golgi apparatus, or to be secreted. For example, the gene can be engineered to contain a signal peptide to facilitate transfer of the peptide to the endoplasmic reticulum. By "signal sequence" is intended a sequence that is known or suspected to result in co-translational or post-translational peptide transport across the cell membrane. In eukaryotes, this typically involves secretion into the Golgi apparatus, with some resulting glycosylation. By "leader sequence" is intended any sequence that when translated, results in an amino acid sequence sufficient to trigger co-translational transport of the peptide chain to a sub-cellular organelle. Thus, this includes leader sequences targeting transport and/or glycosylation by passage into the endoplasmic reticulum, passage to vacuoles, plastids including chloroplasts, mitochondria, and the like. It may also be preferable to engineer the plant expression cassette to contain an intron, such that mRNA processing of the intron is required for expression.

By "3' untranslated region" is intended a nucleotide sequence located downstream of a coding sequence. Polyadenylation signal sequences and other sequences encoding

regulatory signals capable of affecting the addition of poly-adenylic acid tracts to the 3' end of the mRNA precursor are 3' untranslated regions. By "5' untranslated region" is intended a nucleotide sequence located upstream of a coding sequence.

Other upstream or downstream untranslated elements include enhancers. Enhancers are nucleotide sequences that act to increase the expression of a promoter region. Enhancers are well known in the art and include, but are not limited to, the SV40 enhancer region and the 35S enhancer element.

The termination region may be native with the transcriptional initiation region, may be native with the nitrogen-modulating sequence of the present invention, or may be derived from another source. Convenient termination regions are available from the Ti-plasmid of *A. tumefaciens*, such as the octopine synthase and nopaline synthase termination regions, or the potato proteinase inhibitor II sequence (PinII) as described in Liu et al. (2004) *Acta Biochim Biophys Sin* 36(8):553-558. See also Guerineau et al. (1991) *Mol. Gen. Genet.* 262:141-144; Proudfoot (1991) *Cell* 64:671-674; Sanfacon et al. (1991) *Genes Dev.* 5:141-149; Mogen et al. (1990) *Plant Cell* 2:1261-1272; Munroe et al. (1990) *Gene* 91:151-158; Ballas et al. (1989) *Nucleic Acids Res.* 17:7891-7903; and Joshi et al. (1987) *Nucleic Acid Res.* 15:9627-9639.

Where appropriate, the gene(s) may be optimized for increased expression in the transformed host cell. That is, the genes can be synthesized using host cell-preferred codons for improved expression, or may be synthesized using codons at a host-preferred codon usage frequency. Generally, the GC content of the gene will be increased. See, for example, Campbell and Gowri (1990) *Plant Physiol.* 92:1-11 for a discussion of host-preferred codon usage. Methods are known in the art for synthesizing host-preferred genes. See, for example, U.S. Pat. Nos. 6,320,100; 6,075,185; 5,380,831; and 5,436,391, U.S. Published Application Nos. 20040005600 and 20010003849, and Murray et al. (1989) *Nucleic Acids Res.* 17:477-498, herein incorporated by reference.

In one embodiment, the nucleic acids of interest are targeted to the chloroplast for expression. In this manner, where the nucleic acid of interest is not directly inserted into the chloroplast, the expression cassette will additionally contain a nucleic acid encoding a transit peptide to direct the gene product of interest to the chloroplasts. Such transit peptides are known in the art. See, for example, Von Heijne et al. (1991) *Plant Mol. Biol. Rep.* 9:104-126; Clark et al. (1989) *J. Biol. Chem.* 264:17544-17550; Della-Cioppa et al. (1987) *Plant Physiol.* 84:965-968; Romer et al. (1993) *Biochem. Biophys. Res. Commun.* 196:1414-1421; and Shah et al. (1986) *Science* 233:478-481.

The nucleic acids of interest to be targeted to the chloroplast may be optimized for expression in the chloroplast to account for differences in codon usage between the plant nucleus and this organelle. In this manner, the nucleic acids of interest may be synthesized using chloroplast-preferred codons. See, for example, U.S. Pat. No. 5,380,831, herein incorporated by reference.

Typically this "plant expression cassette" will be inserted into a "plant transformation vector." By "transformation vector" is intended a DNA molecule that is necessary for efficient transformation of a cell. Such a molecule may consist of one or more expression cassettes, and may be organized into more than one "vector" DNA molecule. For example, binary vectors are plant transformation vectors that utilize two non-contiguous DNA vectors to encode all requisite cis- and trans-acting functions for transformation

of plant cells (Hellens and Mullineaux (2000) *Trends in Plant Science* 5:446-451). "Vector" refers to a nucleic acid construct designed for transfer between different host cells. "Expression vector" refers to a vector that has the ability to incorporate, integrate and express heterologous DNA sequences or fragments in a foreign cell.

This plant transformation vector may be comprised of one or more DNA vectors needed for achieving plant transformation. For example, it is a common practice in the art to utilize plant transformation vectors that are comprised of more than one contiguous DNA segment. These vectors are often referred to in the art as "binary vectors." Binary vectors as well as vectors with helper plasmids are most often used for *Agrobacterium*-mediated transformation, where the size and complexity of DNA segments needed to achieve efficient transformation is quite large, and it is advantageous to separate functions onto separate DNA molecules. Binary vectors typically contain a plasmid vector that contains the cis-acting sequences required for T-DNA transfer (such as left border and right border), a selectable marker that is engineered to be capable of expression in a plant cell, and a "nucleotide sequence of interest" (a nucleotide sequence engineered to be capable of expression in a plant cell for which generation of transgenic plants is desired). Also present on this plasmid vector are sequences required for bacterial replication. The cis-acting sequences are arranged in a fashion to allow efficient transfer into plant cells and expression therein. For example, the selectable marker gene and the gene of interest are located between the left and right borders. Often a second plasmid vector contains the trans-acting factors that mediate T-DNA transfer from *Agrobacterium* to plant cells. This plasmid often contains the virulence functions (Vir genes) that allow infection of plant cells by *Agrobacterium*, and transfer of DNA by cleavage at border sequences and vir-mediated DNA transfer, as is understood in the art (Hellens and Mullineaux (2000) *Trends in Plant Science*, 5:446-451). Several types of *Agrobacterium* strains (e.g. LBA4404, GV3101, EHA101, EHA105, etc.) can be used for plant transformation. The second plasmid vector is not necessary for transforming the plants by other methods such as microprojection, microinjection, electroporation, polyethylene glycol, etc.

Altered or Improved Variants Useful in the Constructs of the Invention

It is recognized that nucleotide and amino acid sequences useful in the present invention may be altered by various methods, and that these alterations may result in sequences encoding proteins with amino acid sequences different than that encoded by the nitrogen-modulating sequences disclosed herein.

Nucleotide sequences useful in the present invention include the sequences set forth in SEQ ID NO: 1 and 3, and combinations, variants, fragments, and complements thereof. As used herein, the term "nucleotide sequence" or "nucleic acid molecule" is intended to include DNA molecules (e.g., cDNA or genomic DNA) and RNA molecules (e.g., mRNA) and analogs of the DNA or RNA generated using nucleotide analogs. The nucleic acid molecules can be single-stranded or double-stranded, but preferably are double-stranded DNA. By "complement" is intended a nucleotide sequence that is sufficiently complementary to a given nucleotide sequence such that it can hybridize to the given nucleotide sequence to thereby form a stable duplex. The corresponding amino acid sequences for the nitrogen-modulating proteins encoded by these nucleotide sequences are set forth in SEQ ID NO: 2 and 4, as well as combinations, variants and fragments thereof. The invention also

encompasses the use of nucleic acid molecules comprising nucleotide sequences encoding partial-length nitrogen-modulating proteins, and complements thereof.

Nucleic acid molecules that are fragments of these nitrogen-modulating nucleotide sequences are also useful in the present invention. By “fragment” is intended a portion of a nucleotide sequence encoding a nitrogen-modulating protein. A fragment of a nucleotide sequence may encode a biologically active portion of a nitrogen-modulating protein, or it may be a fragment that can be used as a hybridization probe or PCR primer using methods disclosed below. Nucleic acid molecules that are fragments of a nitrogen-modulating nucleotide sequence comprise at least about 15, 20, 50, 75, 100, 200, 300, 350, or at least about 400 contiguous nucleotides, or up to the number of nucleotides present in a full-length nitrogen-modulating nucleotide sequence disclosed herein depending upon the intended use. By “contiguous” nucleotides is intended nucleotide residues that are immediately adjacent to one another.

Polypeptides that are fragments of these nitrogen-modulating polypeptides are also useful in the present invention. By “fragment” is intended a portion of an amino acid sequence encoding a nitrogen-modulating protein as set forth SEQ ID NO: 2 and/or 4, and that retains nitrogen utilization efficiency. A biologically active portion of a nitrogen-modulating protein can be a polypeptide that is, for example, 10, 25, 50, 100, 125, 150, 175, 200, 250, 300, 350, 400 or more amino acids in length. Such biologically active portions can be prepared by recombinant techniques and evaluated for nitrogen utilization efficiency. As used here, a fragment comprises at least 8 contiguous amino acids of SEQ ID NO: 2 and/or 4. The invention encompasses other fragments, however, such as any fragment in the protein greater than about 10, 20, 30, 50, 100, 150, 200, 250, 300, 350, or 400 amino acids.

The invention also encompasses the use of variant nucleic acid molecules, or variant amino acid sequences, in the methods and compositions of the inventions. “Variants” of the nitrogen-modulating nucleotide sequences include those sequences that encode a nitrogen-modulating protein disclosed herein but that differ conservatively because of the degeneracy of the genetic code, as well as those that are sufficiently identical as discussed above. Naturally occurring allelic variants can be identified with the use of well-known molecular biology techniques, such as polymerase chain reaction (PCR) and hybridization techniques as outlined below. Variant nucleotide sequences also include synthetically derived nucleotide sequences that have been generated, for example, by using site-directed mutagenesis but which still encode the nitrogen-modulating proteins disclosed in the present invention as discussed below. Variant proteins useful in the present invention are biologically active, that is they retain the desired biological activity of the native protein, that is, nitrogen utilization efficiency and/or improved stress tolerance.

By “variants” is intended proteins or polypeptides having an amino acid sequence that is at least about 60%, 65%, about 70%, 75%, 80%, 85%, or 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequences of SEQ ID NO: 2 and/or 4. Variants also include polypeptides encoded by a nucleic acid molecule that hybridizes to a nucleic acid molecule of SEQ ID NO: 1 and/or 3, or a complement thereof, under stringent conditions. Variants include polypeptides that differ in amino acid sequence due to mutagenesis. Variant proteins encompassed by the present invention are biologically active, that is they continue to possess the desired biological activity of the

native protein, that is, retain nitrogen utilization efficiency and/or improved stress tolerance.

Preferred nitrogen-modulating proteins useful in the present invention are encoded by a nucleotide sequence sufficiently identical to a nucleotide sequence of SEQ ID NO: 1 and/or 3. The term “sufficiently identical” is intended an amino acid or nucleotide sequence that has at least about 60% or 65% sequence identity, about 70% or 75% sequence identity, about 80% or 85% sequence identity, or about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% sequence identity compared to a reference sequence using one of the alignment programs described herein using standard parameters. One of skill in the art will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning, and the like.

To determine the percent identity of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e., percent identity = number of identical positions / total number of positions (e.g., overlapping positions) × 100). In one embodiment, the two sequences are the same length. The percent identity between two sequences can be determined using techniques similar to those described below, with or without allowing gaps. In calculating percent identity, typically exact matches are counted.

The determination of percent identity between two sequences can be accomplished using a mathematical algorithm. A nonlimiting example of a mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul (1990) *Proc. Natl. Acad. Sci. USA* 87:2264, modified as in Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-5877. Such an algorithm is incorporated into the BLASTN® and BLASTX® programs of Altschul et al. (1990) *J. Mol. Biol.* 215:403. BLAST® nucleotide searches can be performed with the BLASTN® program, score=100, wordlength=12, to obtain nucleotide sequences homologous to nitrogen-modulating nucleic acid molecules of the invention. BLAST® protein searches can be performed with the BLASTX® program, score=50, wordlength=3, to obtain amino acid sequences homologous to nitrogen-modulating protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST® can be utilized as described in Altschul et al. (1997) *Nucleic Acids Res.* 25:3389. Alternatively, PSI-Blast® can be used to perform an iterated search that detects distant relationships between molecules. See Altschul et al. (1997) *supra*. When utilizing BLAST®, Gapped BLAST®, and PSI-Blast® programs, the default parameters of the respective programs (e.g., BLASTX® and BLASTN®) can be used. Another non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the ClustalW algorithm (Higgins et al. (1994) *Nucleic Acids Res.* 22:4673-4680). ClustalW compares sequences and aligns the entirety of the amino acid or DNA sequence, and thus can provide data about the sequence conservation of the entire amino acid sequence. The ClustalW algorithm is used in several commercially available DNA/amino acid analysis software packages, such as the ALIGNX® module of the Vector NTI® Program Suite (Invitrogen Corporation, Carlsbad, Calif.). After alignment of amino acid sequences with ClustalW, the percent amino acid identity can be assessed. A non-limiting example of a

software program useful for analysis of ClustalW alignments is GENEDOCT™. GENEDOCT™ (Karl Nicholas) allows assessment of amino acid (or DNA) similarity and identity between multiple proteins. Another non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller (1988) *CABIOS* 4:11-17. Such an algorithm is incorporated into the ALIGN® program (version 2.0), which is part of the GCG sequence alignment software package (available from Accelrys, Inc., 9865 Scranton Rd., San Diego, Calif., USA). When utilizing the ALIGN® program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used.

A preferred program is GAP version 10, which used the algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443-453. GAP Version 10 may be used with the following parameters: % identity and % similarity for a nucleotide sequence using GAP Weight of 50 and Length Weight of 3, and the nwsgapdna.cmp scoring matrix; % identity and % similarity for an amino acid sequence using GAP Weight of 8 and Length Weight of 2, and the BLOSUM62 Scoring Matrix. Equivalent programs may also be used. By “equivalent program” is intended any sequence comparison program that, for any two sequences in question, generates an alignment having identical nucleotide or amino acid residue matches and an identical percent sequence identity when compared to the corresponding alignment generated by GAP Version 10.

The skilled artisan will further appreciate that changes can be introduced by mutation into the nucleotide sequences of the invention thereby leading to changes in the amino acid sequence of the encoded nitrogen-modulating protein, without altering the biological activity of the protein. Thus, variant isolated nucleic acid molecules can be created by introducing one or more nucleotide substitutions, additions, or deletions into the corresponding nucleotide sequence disclosed herein, such that one or more amino acid substitutions, additions or deletions are introduced into the encoded protein. Mutations can be introduced by standard techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. Such variant nucleotide sequences are also encompassed by the present invention.

For example, conservative amino acid substitutions may be made at one or more predicted, preferably nonessential amino acid residues. A “nonessential” amino acid residue is a residue that can be altered from the wild-type sequence of a nitrogen-modulating protein without altering the biological activity, whereas an “essential” amino acid residue is required for biological activity. A “conservative amino acid substitution” is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Amino acid substitutions may be made in nonconserved regions that retain function. In general, such substitutions would not be made for conserved amino acid residues, or for amino acid residues residing within a conserved motif, where such residues are essential for protein activity. However, one of skill in the art would understand that functional

variants may have minor conserved or nonconserved alterations in the conserved residues. Examples of residues that are conserved and that may be essential for protein activity include, for example, residues that are identical between all proteins contained in an alignment of similar or related sequences known to be involved in nitrogen assimilation. Examples of residues that are conserved but that may allow conservative amino acid substitutions and still retain activity include, for example, residues that have only conservative substitutions between all proteins contained in an alignment of similar or related sequences known to be involved in nitrogen assimilation.

Alternatively, variant nucleotide sequences can be made by introducing mutations randomly along all or part of the coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for ability to confer nitrogen utilization efficiency to identify mutants that retain activity. Following mutagenesis, the encoded protein can be expressed recombinantly, and the activity of the protein can be determined using standard assay techniques.

Using methods such as PCR, hybridization, and the like, corresponding nitrogen-modulating sequences can be identified, such sequences having substantial identity to the sequences of the invention. See, for example, Sambrook J., and Russell, D. W. (2001) *Molecular Cloning: A Laboratory Manual*. (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.) and Innis, et al. (1990) *PCR Protocols: A Guide to Methods and Applications* (Academic Press, NY). In a hybridization method, all or part of the nitrogen-modulating nucleotide sequence can be used to screen cDNA or genomic libraries. Methods for construction of such cDNA and genomic libraries are generally known in the art and are disclosed in Sambrook and Russell, 2001, supra.

Variants and fragments of the nucleotide or amino acid sequences of the present invention generally will encode protein fragments that retain the biological activity of the full-length nitrogen-modulating protein; i.e., retain nitrogen utilization efficiency. By “retains nitrogen utilization efficiency” is intended that the variant or fragment will have at least about 30%, at least about 50%, at least about 70%, or at least about 80% of the nitrogen utilization efficiency and/or stress tolerance of the full-length nitrogen-modulating protein disclosed herein as SEQ ID NO: 2 and/or 4, or the full-length nitrogen-modulating nucleotide sequence disclosed herein as SEQ ID NO: 1 and/or 3. Methods for monitoring nitrogen utilization efficiency include detecting a change in any of the main nitrogen metabolic pool sizes in the assimilation pathways (for example, a measurable change in nitrate, nitrite, ammonia, glutamic acid, aspartic acid, glutamine, asparagine, lysine, leucine, threonine, methionine, glycine, tryptophan, tyrosine, total protein content of a plant part, total nitrogen content of a plant part, and/or chlorophyll content) or detecting the ability of a plant to provide the same or elevated yield at lower nitrogen fertilization levels, or the ability of a plant to provide elevated yields at the same nitrogen fertilization levels when compared to a plant that does not contain or express a nitrogen-modulating sequence of the invention. The designation of “same” or “lower” nitrogen fertilization levels refers to the level of nitrogen generally applied to a plant not expressing a nitrogen-modulating sequence of the invention. Sufficient nitrogen levels are known in the art for the majority, if not all, plant varieties of interest. Additional guidance may be found in, for example, Hewitt (1966) *Sand and Water Culture Methods Used in the Study of Plant Nutrition*, 2nd ed., Farnham Royal (Bucks), Commonwealth

Agricultural Bureaux; and, Hewitt (1975) *Plant Mineral Nutrition*, London, English University Press.

The polypeptide sequences useful in the present invention may be altered in various ways including amino acid substitutions, deletions, truncations, and insertions. Methods for such manipulations are generally known in the art. For example, amino acid sequence variants of the nitrogen-modulating proteins disclosed herein can be prepared by mutations in the nucleotide sequences. This may also be accomplished by one of several forms of mutagenesis and/or in directed evolution. In some aspects, the changes encoded in the amino acid sequence will not substantially affect function of the protein. Such variants will possess the desired nitrogen utilization efficiency. However, it is understood that the ability of the nitrogen-modulating sequences of the invention to alter or improve nitrogen utilization may be further improved by one use of such techniques upon the compositions of this invention. For example, one may express the nucleotide sequences disclosed herein in host cells that exhibit high rates of base misincorporation during DNA replication, such as XL-1 Red (Stratagene, La Jolla, Calif.). After propagation in such strains, one can isolate the DNA (for example by preparing plasmid DNA, or by amplifying by PCR and cloning the resulting PCR fragment into a vector), transform it into plants as described elsewhere herein, and measure nitrogen utilization efficiency.

Alternatively, alterations may be made to the protein sequence of many proteins at the amino or carboxy terminus without substantially affecting activity. This can include insertions, deletions, or alterations introduced by modern molecular methods, such as PCR, including PCR amplifications that alter or extend the protein coding sequence by virtue of inclusion of amino acid encoding sequences in the oligonucleotides utilized in the PCR amplification. Alternatively, the protein sequences added can include entire protein-coding sequences, such as those used commonly in the art to generate protein fusions. Such fusion proteins are often used to (1) increase expression of a protein of interest, (2) introduce a binding domain, enzymatic activity, or epitope to facilitate either protein purification, protein detection, or other experimental uses known in the art, or, (3) target secretion or translation of a protein to a subcellular organelle, such as the periplasmic space of gram-negative bacteria, or the endoplasmic reticulum of eukaryotic cells, the latter of which often results in glycosylation of the protein.

Variant nucleotide and amino acid sequences of the present invention also encompass sequences derived from mutagenic and recombinogenic procedures such as DNA shuffling. With such a procedure, one or more different nitrogen-modulating protein coding regions can be used to create a new nitrogen-modulating protein possessing the desired properties. In this manner, libraries of recombinant polynucleotides are generated from a population of related sequence polynucleotides comprising sequence regions that have substantial sequence identity and can be homologously recombined in vitro or in vivo. For example, using this approach, sequence motifs encoding a domain of interest may be shuffled between the nitrogen-modulating sequence useful in the present invention and other known nitrogen-modulating sequences to obtain a new sequence coding for a protein with an improved property of interest, such as improved nitrogen utilization. Strategies for such DNA shuffling are known in the art. See, for example, Stemmer (1994) *Proc. Natl. Acad. Sci. USA* 91:10747-10751; Stemmer (1994) *Nature* 370:389-391; Cramer et al. (1997) *Nature Biotech.* 15:436-438; Moore et al. (1997) *J. Mol. Biol.* 272:336-347; Zhang et al. (1997) *Proc. Natl. Acad. Sci.*

USA 94:4504-4509; Cramer et al. (1998) *Nature* 391:288-291; and U.S. Pat. Nos. 5,605,793 and 5,837,458.

Plant Transformation

Methods of the invention involve introducing one or more nitrogen-modulating nucleotide sequences into a plant. In some embodiments, only one of the nitrogen-modulating sequences disclosed herein is introduced into the plant. In other embodiments, at least 2, at least 3, at least 4, or more of the sequences are introduced. Where multiple sequences are introduced, each of the nucleotide sequences is non-identical. Two nucleotide sequences are considered non-identical if they differ in at least one nucleotide position. Thus, non-identical nucleotide sequences include two or more different nucleotide sequences that each encodes the same amino acid sequence (e.g., one or more has been optimized for expression in the plant), as well as two or more different nucleotide sequences that encode at least two different amino acid sequences.

By "introducing" it is intended to present to the plant one or more constructs comprising the one or more nitrogen-modulating sequences in such a manner that the construct(s) gain(s) access to the interior of a cell of the plant. The methods of the invention do not require that a particular method for introducing a nucleotide construct to a plant is used, only that the nucleotide construct(s) gain(s) access to the interior of at least one cell of the plant. Methods for introducing nucleotide constructs into plants are known in the art including, but not limited to, stable transformation methods, transient transformation methods, and virus-mediated methods.

In general, plant transformation methods involve transferring heterologous DNA into target plant cells (e.g. immature or mature embryos, suspension cultures, undifferentiated callus, protoplasts, etc.), followed by applying a maximum threshold level of appropriate selection (depending on the selectable marker gene) to recover the transformed plant cells from a group of untransformed cell mass. Explants are typically transferred to a fresh supply of the same medium and cultured routinely. Subsequently, the transformed cells are differentiated into shoots after placing on regeneration medium supplemented with a maximum threshold level of selecting agent (i.e., antibiotics, such as spectinomycin and kanamycin). The shoots are then transferred to a selective rooting medium for recovering rooted shoot or plantlet. The transgenic plantlet then grow into mature plant and produce fertile seeds (e.g. Hiei et al. (1994) *The Plant Journal* 6:271-282; Ishida et al. (1996) *Nature Biotechnology* 14:745-750). Explants are typically transferred to a fresh supply of the same medium and cultured routinely. A general description of the techniques and methods for generating transgenic plants are found in Ayres and Park (1994) *Critical Reviews in Plant Science* 13:219-239 and Bommineni and Jauhar (1997) *Maydica* 42:107-120. Since the transformed material contains many cells, both transformed and non-transformed cells are present in any piece of subjected target callus or tissue or group of cells. The ability to kill non-transformed cells and allow transformed cells to proliferate results in transformed plant cultures. Often, the ability to remove non-transformed cells is a limitation to rapid recovery of transformed plant cells and successful generation of transgenic plants. Molecular and biochemical methods can then be used to confirm the presence of the integrated heterologous gene of interest in the genome of transgenic plant.

Generation of transgenic plants may be performed by one of several methods, including but not limited to introduction of heterologous DNA by *Agrobacterium* into plant cells

(*Agrobacterium*-mediated transformation), bombardment of plant cells with heterologous foreign DNA adhered to particles, and various other non-particle direct-mediated methods (e.g. Hiei et al. (1994) *The Plant Journal* 6:271-282; Ishida et al. (1996) *Nature Biotechnology* 14:745-750; Ayres and Park (1994) *Critical Reviews in Plant Science* 13:219-239; Bommineni and Jauhar (1997) *Maydica* 42:107-120) to transfer DNA.

Methods for transformation of chloroplasts are known in the art. See, for example, Svab et al. (1990) *Proc. Natl. Acad. Sci. USA* 87:8526-8530; Svab and Maliga (1993) *Proc. Natl. Acad. Sci. USA* 90:913-917; Svab and Maliga (1993) *EMBO J.* 12:601-606. The method relies on particle gun delivery of DNA containing a selectable marker and targeting of the DNA to the plastid genome through homologous recombination. Additionally, plastid transformation can be accomplished by transactivation of a silent plastid-borne transgene by tissue-preferred expression of a nuclear-encoded and plastid-directed RNA polymerase. Such a system has been reported in McBride et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:7301-7305.

Transformation of bacterial cells is accomplished by one of several techniques known in the art, including but not limited to electroporation or chemical transformation (see, for example, Ausubel, ed. (1994) *Current Protocols in Molecular Biology*, John Wiley and Sons, Inc., Indianapolis, Ind.). Markers conferring resistance to toxic substances are useful in identifying transformed cells (having taken up and expressed the test DNA) from non-transformed cells (those not containing or not expressing the test DNA).

In one aspect of the invention, the nucleotide sequences of the invention are useful as markers to assess transformation of bacterial or plant cells. In this manner, transformation is assessed by monitoring nitrogen utilization efficiency as described above.

Transformation of plant cells can be accomplished in similar fashion. By "plant" is intended whole plants, or component parts including plant organs (e.g., leaves, stems, roots, etc.), seeds, plant cells, propagules, embryos and progeny of the same. Plant cells can be differentiated or undifferentiated (e.g. callus, suspension culture cells, protoplasts, leaf cells, root cells, phloem cells, pollen). "Transgenic plants" or "transformed plants" or "stably transformed" plants or cells or tissues refer to plants that have incorporated or integrated exogenous nucleic acid sequences or DNA fragments into the plant cell. By "stable transformation" is intended that the nucleotide construct introduced into a plant integrates into the genome of the plant and is capable of being inherited by progeny thereof.

The cells that have been transformed may be grown into plants in accordance with conventional ways. See, for example, McCormick et al. (1986) *Plant Cell Reports* 5:81-84. These plants may then be grown, and either pollinated with the same transformed strain or different strains, and the resulting hybrid having constitutive expression of the desired phenotypic characteristic identified. Two or more generations may be grown to ensure that expression of the desired phenotypic characteristic is stably maintained and inherited and then seeds harvested to ensure expression of the desired phenotypic characteristic has been achieved. In this manner, the present invention provides transformed seed (also referred to as "transgenic seed") having a nucleotide construct of the invention, for example, an expression cassette of the invention, stably incorporated into their genome. Methods to Increase Plant Yield by Modulating Nitrogen Utilization

Methods for increasing plant yield are provided. The methods comprise introducing into a plant or plant cell a nitrogen-modulating nucleotide sequence disclosed herein such that an increase in nitrogen utilization efficiency corresponds to an increase in plant yield. As defined herein, the "yield" of the plant refers to the quality and/or quantity of biomass, and/or harvestable yield, produced by the plant. By "biomass" is intended any measured plant product (e.g., any component part of a plant, such as seed, stalk, root, grain, leaf, etc.). An increase in biomass production is any improvement in the yield of the measured plant product. An increase in harvestable yield is a higher weight of a plant component that is easily collected using known harvest methods, or an increase in the compositional amount of a compound of interest in the harvested part: a nonlimiting example, being the amount of an amino acid, such as lysine, that is harvested per unit land area. Increasing plant yield or harvestable yield has several commercial applications. For example, increasing plant leaf biomass may increase the yield of leafy vegetables for human or animal consumption. Additionally, increasing leaf biomass can be used to increase production of plant-derived pharmaceutical or industrial products. An increase in yield can comprise any statistically significant increase including, but not limited to, at least a 1% increase, at least a 3% increase, at least a 5% increase, at least a 10% increase, at least a 20% increase, at least a 30%, at least a 50%, at least a 70%, at least a 100% or a greater increase in plant yield compared to the yield of a plant into which a nucleotide sequence that modulates use of nitrogen of the invention has not been introduced.

Plants

The present invention may be used for transformation of any plant species, including, but not limited to, monocots and dicots. Examples of plants of interest include, but are not limited to, corn (maize), sorghum, wheat, sunflower, tomato, crucifers, peppers, potato, cotton, rice, soybean, sugarbeet, sugarcane, tobacco, barley, and oilseed rape, *Brassica* sp., alfalfa, rye, millet, safflower, peanuts, sweet potato, cassava, coffee, coconut, pineapple, citrus trees, cocoa, tea, banana, avocado, fig, guava, mango, olive, papaya, cashew, macadamia, almond, oats, vegetables, grasses (such as turf grasses, forage grasses, or pasture grasses), ornamentals, trees (such as fruit trees, nut trees, pulp trees, oil palms) and conifers.

Vegetables include, but are not limited to, onions, tomatoes, lettuce, green beans, lima beans, peas, and members of the genus *Curcumis* such as cucumber, cantaloupe, and muskmelon. Ornamentals include, but are not limited to, azalea, hydrangea, hibiscus, roses, tulips, daffodils, petunias, carnation, poinsettia, and chrysanthemum. Preferably, plants of the present invention are crop plants (for example, maize, sorghum, wheat, sunflower, tomato, crucifers, peppers, potato, cotton, rice, soybean, sugarbeet, sugarcane, tobacco, barley, oilseed rape, etc.).

This invention is particularly suitable for any member of the monocot plant family including, but not limited to, maize, rice, barley, oats, wheat, sorghum, rye, sugarcane, pineapple, yams, onion, banana, coconut, and dates.

Evaluation of Plant Transformation

Following introduction of heterologous foreign DNA into plant cells, the transformation or integration of heterologous gene in the plant genome is confirmed by various methods such as analysis of nucleic acids, proteins and metabolites associated with the integrated gene.

PCR analysis is a rapid method to screen transformed cells, tissue or shoots for the presence of incorporated nucleotide sequences at the earlier stage before transplanting

into the soil (Sambrook and Russell (2001) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.). PCR is carried out using oligonucleotide primers specific to the gene of interest or *Agrobacterium* vector background, etc.

Plant transformation may be confirmed by Southern blot analysis of genomic DNA (Sambrook and Russell, 2001, supra). In general, total DNA is extracted from the transformant, digested with appropriate restriction enzymes, fractionated in an agarose gel and transferred to a nitrocellulose or nylon membrane. The membrane or "blot" is then probed with, for example, radiolabeled ^{32}P target DNA fragments to confirm the integration of the introduced gene in the plant genome according to standard techniques (Sambrook and Russell, 2001, supra).

In Northern analysis, RNA is isolated from specific tissues of transformant, fractionated in a formaldehyde agarose gel, blotted onto a nylon filter according to standard procedures that are routinely used in the art (Sambrook and Russell, 2001, supra). Expression of RNA encoded by the nucleotide sequence of the invention is then tested by hybridizing the filter to a radioactive probe derived from a polynucleotide of the invention, by methods known in the art (Sambrook and Russell, 2001, supra).

Western blot and biochemical assays and the like may be carried out on the transgenic plants to determine the presence of protein encoded by the nitrogen-modulating gene by standard procedures (Sambrook and Russell, 2001, supra) using antibodies that bind to one or more epitopes present on the nitrogen-modulating protein. For example, the polyclonal antibodies generated by the methods of the present invention can be used to detect the presence of a nitrogen-modulating protein.

Antibodies

Antibodies to the polypeptides useful in the present invention, or to variants or fragments thereof, are also encompassed. Methods for producing antibodies are well known in the art (see, for example, Harlow and Lane (1988) *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.; U.S. Pat. No. 4,196,265).

EXPERIMENTAL

I. icdh Gene

Materials and Methods

Using genomic tools, multiple searches were made among the plant and bacterial icdh genes in Genbank. The target focus was on cytosolic, monomeric, NADP⁺-dependent versions, and with a particular "AT" pattern in the polyA signals that would not be inhibitory to expression in plants. This work eventually led to a sequence from an *Azotobacter* species that had a fit with the criteria. The sequence of the selected bacterial isocitrate dehydrogenase gene (icdh) was codon optimized for expression in maize. A synthetic gene encoding an ICDH enzyme was generated (SEQ ID NO: 1):

ATGAGCACCCCAAGATCATCTACACCTTGACAGATGAGGCGCCGGCGCT
GGCCACCTACAGCTTGCTGCCCATCATCAAGGCTTCACTGGAAGCTCAG
GCATTGCTGTGGAACAAGGACATCTCCCTTGCTGGAAGGCTGATCGCC
ACCTTCCAGAATATTTGACAGACACCAGAAGATCTCTGATGATCTTGC
TGAGCTGGGGAAGCTGGCCACCAGCCAGATGCCAACATCATCAAGCTGC

-continued

CAAACATCTCTGCTTCAGTTCCCTCAGCTGAAGGCCCGCCATCAAGGAACTC
CAGCAGCAAGGCTACAAGCTGCCAGATTATCCAGAAGAACCAAAAACAGA
5 CACAGAGAAGGATGTCAAGGCAAGATATGACAAGATCAAGGGCAGCGCCG
TCAACCCCGTGCTGAGAGAAGGAAATTCAGACCGCCGCGCGCGCTCTCC
GTCAAGAACTATGCAAGGAAGCATCCTCACAAGATGGGCGCCTGGAGCGC
10 CGACAGCAAGAGCCATGTTGCTCACATGGACAATGGAGATTTCTATGGAT
CAGAGAAGGCGCGCTGATTGGTGCTCCTGGAAGCGTCAAGATTGAGCTG
ATCGCCAAGGATGGAAGCAGCACCGTGCTGAAGGCCAAGACATCAGTTCA
15 AGCTGGAGAGATCATCGACAGCTCCGTGATGAGCAAGAATGCTCTGAGGA
ACTTCATTGCTGCCGAGATTGAAGATGCCAAGAAGCAAGGAGTGCTGCTC
TCCGTCCACCTCAAGGCCACCATGATGAAGGTTTTCAGATCCCATCATGTT
TGGCCAGATTGTTTCAGAGTTCTACAAGGATGCTCTCACCAGCATGCTG
20 AGGTGCTGAAGCAGATTGGATTTGATGTCAACAATGGCATTGGAGATCTC
TATGCAAGGATCAAGACCTACCAGAAGCAAAGCAGAAGGAGATTGAAGC
TGACATCCAAGCTGTTTATGCTCAAAGGCCGAGCTGGCAATGGTGAACA
25 GCGACAAGGGCATCACCACCTCCATGTTCTTCTGATGTCATCGTCGAC
GCCTCCATGCCGGCCATGATCAGAGATTTCAGGGAAGATGTGGGGCCAGA
TGGCAAGCTGCATGACACCAAGGCCGTCATCCAGATCGCTGCTATGCTG
30 GCGTCTACCAGGTGGTGATTGAAGATTGCAAGCAGCATGGCGCCTTCGAC
CCAACAACAATGGGCTCAGTTCCAAATGTTGGGCTGATGGCGCAGAAGGC
AGAAGAATATGGAAGCCATGACAAGACCTTTCAGATCCCTGCTGATGGCG
35 TCGTCCGCGTCACTGATGAAAGCGGCAAGCTGCTGCTGGAGCAATCAGTG
GAAGCTGGAGACATCTGGAGGATGTGCCAAGCAAAGGATGCTCCCATCCA
AGATTGGGTGAAGCTCGCCGTCAACAGGGCGCGCCACCAACACGCCGG
40 CGGTGTTCTGGCTGGACCCAGCAAGGGCTCATGATGCTCAGGTGATCGCC
AAGGTGGAGAGATATCTAAAGGATTATGACACCTCCGGCCTGGACATCAG
GATCTTGTGCGCGGTGGAAGCAACAAGGTTCTCCTTGGCAAGGATCAGAG
45 AAGGAAAGGACACCATCTCAGTGACAGGAAATGTGCTGAGGGACTACCTC
ACCGACCTCTTCCCATCATGGAGCTGGGCACCTCCGCCAAGATGCTCTC
CATTGTTCTCTGATGAGCGGCGGCGCCTCTTTGAACTGGAGCTGGAG
50 GATCAGCGCCCAAGCATGTTTCAAGGATTCCTGGAAGAAGGCTACCTCAGA
TGGGACAGCCTTGAGAGTTTCTGGCGCTCGCCGCTCCTTGGAGCATCT
TGGAATGCCATAAGAACCACAAAGGCGCTGGTGCTGGCCTCCACCTAG
55 ATCAAGCTACTGGCAAGATCCTGGACAACAACAAGAGCCCAGCAAGGAAG
GTTGGTGAGATCGACAACAGAGGAAGCCACTTCTACCTGGCGCTCTACTG
GGCTCAAGCTCTTGTGCTCAAACAGAGGACAAGGAGCTACAAGCTCAGT
60 TCACCGGCATTGCCAAGGCGCTGACAGACAATGAAACAAAAATGTTTGA
GAGCTGGCTGCTGCTCAAGGAAAGCCGGTGGACATTGCTGGCTACTACCA
TCCAAACACCGACCTCACCAGCAAGGCCATCAGGCCATCTGCCACCTTCA
65 ATGCTGCTCTGGCGCGCTGGCATAGTAAGG

The *icdh* DNA sequence shown above encodes the following ICDH protein sequence (SEQ ID NO:2) (741 amino acids):

```
mstpkiiytl tdeapalaty sllpiikaft gssgiavetr
dislagrlia tfpeyltdtq kisddlaelg klattpdani
iklpnisasv pqlkaakel qqqgyklpdy peepktdtek
dvkarydkik gsavnpvlre gnsdrrapls vknyarkhph
kmgawsadsk shvahmdngd fygsekaali gapgsvkiel
iakdgsstvl kaktsvqage iidssvmskn alrnfiaaei
edakkqgvll svhlkatmmk vsdpimfgqi vsefykdalt
khaevlkqig fdvnmigdl yariktlpea kqkeieadiq
avyaqrpqla mvnsdkgitn lhvpsdvivd asmpamirds
gkmwgpdgkl hdtkavipdr cyagvyqvvi edckqhgaft
pttmgsvpnv glmaqkaeey gshdktfqip adgvrvrtde
sgklleqsv eagdiwrncq akdapiqdwv klavnrarat
ntpavfwldp arandaqvia kverylkdyd tsgldirils
pveatrfsla riregkdtis vtgnvirdyl tdlfpimelg
tsakmlsivp lmsggglfet gaggsapkhv qqfleegylyr
wdslgeflal aaslehlgn yknpkalvla stldqatgki
ldnnkspark vgeidnrgh fylalywaqa laaqtedkel
qaqftgiaka ltdnetkivg elaaaqgkp diagyypnt
dltskairps atfnaalapl a
```

Vector Construction for Overexpression of ICDH

The open reading frame described in the previous section was introduced into a vector for plant expression. The vector also contains a gene encoding a glyphosate tolerant EPSPS enzyme (GRG23ace5) that was used as the selectable marker during maize transformation. Expression of each of these genes was controlled by the ScUbi4 promoter to produce robust expression in maize. A vector map of this vector, denominated pMDO8901, is shown in FIG. 1.

Plant Transformation

The pMDO8901 vector was used to carry out *Agrobacterium*-mediated transformation of maize. Following vector construction and transformation of *Agrobacterium*, the vectors were confirmed by Southern blot by methods known in the art. Positive *Agrobacterium* strains that passed these tests were then grown on a solid medium to produce cell counts for large-scale transformation experiments.

The vector pMDO8901 was introduced into an *Agrobacterium tumefaciens* strain by electroporation. The formation

of the recombinant vector, pMDO8901, was confirmed by Southern blot hybridization of this *Agrobacterium* strain. The selection agent for these experiments was glyphosate

The *Agrobacterium* strain harboring the cointegrate can be used to transform plants, for example, by the PureIntro method (Japan Tobacco, Inc.).

Western Blot Analysis

Expression of *icdh* in these plants was examined by generating antibodies that bind specifically to the ICDH protein. Briefly, the *icdh* gene was subcloned into the vector pRSF1b (Novagen) to allow overexpression of the ICDH protein in *E. coli* following IPTG induction. The vector also introduces a 6xHis tag at the N-terminus of the protein. Following protein overexpression, the ICDH protein was purified by cobalt column chromatography and the identity of the purified protein was confirmed by N-terminal sequencing. The purified protein was then used to immunize rabbits, with serum collection beginning 42 days after immunization.

Next, the ICDH antiserum was used to assess protein expression in the transgenic maize plants by Western blot analysis. Leaf samples were taken from individual plants following 4 weeks of growth in the greenhouse, and protein extracts were prepared by grinding the plant material in water. Protein concentration in each extract was determined by Bradford assay, and 25 ug of each extract was separated on polyacrylamide gels with a 4-12% gradient. The separated proteins were transferred to nitrocellulose and then probed with the rabbit antiserum at a 1:5000 dilution. Following wash steps, the nitrocellulose was contacted with goat anti-rabbit conjugated with horseradish peroxidase (1:10,000 dilution), and antibody complexes were visualized using ECL detection reagents (GE Healthcare). At the T0 stage most *icdh* events were found to be expressing ICDH. Two events had particularly good expression levels and were promoted to the T1 stage.

Maize Nitrogen Analysis

A series of assays that quantify nitrogen intermediates in plants have been developed. These nitrogen assay methods are described in a previous patent filing (WO 2008/051608 "Plants with improved nitrogen utilization and stress tolerance"). These assays were utilized here to analyze a total of 10 transgenic plants containing the *icdh* gene. Each of the plants was sampled (leaf) following 4 weeks of growth in soil in a greenhouse. These leaf samples were processed to determine their nitrate, asparagine, glutamine, aspartic acid, glutamic acid, ammonium, total amino acid, chlorophyll, and total protein levels. Included alongside in the analysis were plants that were transformed with a construct containing only the GRG23ace5 selectable marker (no *icdh* nor *nagk*). These plants were likewise sampled at 4 weeks and are referred to as "non-GOI" plants. The results of the nitrogen assays carried out on both types of plants are shown below in Table 1.

TABLE 1

Nitrogen levels, ICDH vs. non-GOI maize events, 4 weeks after transfer to soil									
Plant #	GOI	Aspartic Acid ($\mu\text{g/g}$)	Glutamic Acid ($\mu\text{g/g}$)	Asparagine ($\mu\text{g/g}$)	Glutamine ($\mu\text{g/g}$)	Total Chlorophyll (a + b) (mg/g)	Ammonium ($\mu\text{g/g}$)	Total Amino Acids (mg/g)	Total Protein (mg/g)
15091	ICDH	177.3	1240.2	1692.0	5175.0	0.086	755.795	285.830	16.990
15092	ICDH	451.7	1161.9	1898.3	2055.2	0.007	533.265	234.620	25.471
15093	ICDH	82.9	962.6	450.0	1344.0	0.061	502.607	157.230	24.621
15094	ICDH	201.3	1417.4	148.9	2144.1	0.074	670.780	166.334	19.411
15095	ICDH	174.9	1176.2	1403.1	3009.6	0.066	714.197	196.974	22.753
15096	ICDH	51.2	1435.5	2006.0	2630.9	0.052	626.751	302.167	20.094

TABLE 1-continued

Nitrogen levels, ICDH vs. non-GOI maize events, 4 weeks after transfer to soil									
Plant #	GOI	Aspartic Acid ($\mu\text{g/g}$)	Glutamic Acid ($\mu\text{g/g}$)	Asparagine ($\mu\text{g/g}$)	Glutamine ($\mu\text{g/g}$)	Total Chlorophyll (a + b) (mg/g)	Ammonium ($\mu\text{g/g}$)	Total Amino Acids (mg/g)	Total Protein (mg/g)
15097	ICDH	213.6	1213.7	1364.2	2357.3	0.054	634.482	207.681	24.317
15098	ICDH	256.4	1762.7	3595.3	5148.4	0.357	1822.651	448.316	42.052
15099	ICDH	82.4	848.0	1946.7	2503.9	0.064	590.783	201.124	19.159
15100	ICDH	208.4	1562.0	2226.4	4765.9	0.061	1140.701	303.099	19.038
15101	ICDH	167.6	914.3	1184.2	2873.7	0.049	542.457	267.801	18.433
15122	ICDH	269.2	1081.6	1424.1	6786.6	0.061	574.964	265.796	35.271
15123	ICDH	322.1	1713.5	1625.9	7988.8	0.062	24.671	375.517	23.669
15124	ICDH	296.5	1262.3	1138.7	6023.9	0.077	367.674	237.945	29.045
15125	ICDH	387.2	2006.6	1652.1	7789.2	0.079	926.448	289.497	30.555
IC 1 Control	non-GOI	537.8	596.3	2335.0	3516.7	0.154	984.302	362.225	28.635
IC 2 Control	non-GOI	101.6	473.7	1490.3	2485.0	0.157	505.712	264.265	34.564
IC 3 Control	non-GOI	153.6	914.9	1799.3	2191.5	0.066	534.599	297.756	34.019
IC 4 Control	non-GOI	59.7	526.2	313.0	659.3	0.082	347.337	183.701	24.077
Avg		213.2	627.8	1484.4	2213.1	0.114	984.3	277.0	30.3
Std Dev		219.8	197.9	855.4	1181.5	0.048	273.6	74.3	5.0

20

These data demonstrate that the synthetic gene we designed encodes a functional ICDH enzyme.

Maize Plants Containing Icdh Gene Showed Differences Over Controls

Events 15122 and 15125 were selected and progressed onto the T1 stage. The T1 plants were sampled and evaluated as described previously. The results are set out in Tables 2-4.

TABLE 2

T0 Event Promoted to T1 Testing Due to Elevated Glutamine									
15122 Plant #	GOI	Aspartic Acid ($\mu\text{g/g}$)	Glutamic Acid ($\mu\text{g/g}$)	Asparagine ($\mu\text{g/g}$)	Glutamine ($\mu\text{g/g}$)	Total Chlorophyll (a + b) (mg/g)	Ammonium ($\mu\text{g/g}$)	Total Amino Acids (mg/g)	Total Protein (mg/g)
2	ICDH	331	2537	575	2551	0.197	566	229	18.6
3	ICDH	456	2299	647	2105	0.144	477	254	18.1
5	ICDH	563	2880	220	2106	0.450	294	170	12.9
7	ICDH	1151	2350	671	2628	0.243	351	217	22.5
8	ICDH	664	2707	466	3826	0.115	413	202	14.0
9	ICDH	446	3155	89	1156	0.148	406	163	27.9
Avg		602	2655	445	2396	0.216	418	206	19.0
Std Dev		292	328	240	875	0.123	95	35	5.6

TABLE 3

T0 Event Promoted to T1 Testing Due to Elevated Aspartic Acid, Glutamic Acid and Glutamine									
15122 Plant #	GOI	Aspartic Acid ($\mu\text{g/g}$)	Glutamic Acid ($\mu\text{g/g}$)	Asparagine ($\mu\text{g/g}$)	Glutamine ($\mu\text{g/g}$)	Total Chlorophyll (a + b) (mg/g)	Ammonium ($\mu\text{g/g}$)	Total Amino Acids (mg/g)	Total Protein (mg/g)
1	ICDH	478	2929	140	3961	0.166	609	204	21.6
2	ICDH	827	1912	855	4177	0.207	391	256	19.3
3	ICDH	838	2406	416	4364	0.157	613	251	14.2
4	ICDH	865	24410	606	6268	0.284	376	192	28.2
5	ICDH	391	2564	167	2447	0.184	535	221	17.3
10	ICDH	328	3010	376	2128	0.209	417	345	14.0
Avg		621	6205	427	3891	0.201	490	245	19.1
Std Dev		248	8927	271	1494	0.046	109	55	5.3

TABLE 4

Controls									
Control Plant #	GOI	Aspartic Acid (µg/g)	Glutamic Acid (µg/g)	Asparagine (µg/g)	Glutamine (µg/g)	Total Chlorophyll (a + b) (mg/g)	Ammonium (µg/g)	Total Amino Acids (mg/g)	Total Protein (mg/g)
1		253	2351	1759	2412	0.298	567	235	22.0
2		739	2154	3488	3989	0.089	452	226	17.4
3		297	1767	373	2532	0.107	408	197	21.2
4		905	2233	941	3237	0.145	314	190	17.2
5		394	3037	3346	5262	0.155	504	252	19.5
6		789	2535	4166	9084	0.133	459	295	28.9
Avg		563	2346	2346	4419	0.154	451	232	21.0
Std Dev		281	424	1538	2516	0.074	86	38	4.3

The data for T1 plants indicated lower asparagine levels for the transgenic events (both the average and for most plants). Individual plants had results that differed from the control; for example, event 15122 plant #5, had higher glutamic acid and higher chlorophyll but lower amino acids and lower protein levels compared to the controls.

In Event 15125, plant #4 had much a higher level of glutamic acid. The recorded number was so high as to appear that it might be an outlier or a sampling error. However, the same plant had higher chlorophyll and higher protein levels compared to the control (the chlorophyll and protein measurements are taken from a separate sample than the glutamine), therefore all the elevated levels could not have been due to the same experimental error, even if one assumed there was an error for glutamine.

Overall the data indicates some potential effects on the measured pool sizes (e.g. lower asparagine) and others for individual plants (e.g. chlorophyll). Lower asparagine could be due to more N incorporation into other pools resulting in less N for internal transport (via the transport amino acid, asparagine).

II. icdh Gene+nagk Gene

Vector Construction for Overexpression of Bacterial icdh+nagk

The ICDH vector pMDO8901 was further modified to add a gene encoding an arginine-insensitive N-acetylglutamate kinase (NAGK) protein. This new vector (pMDO8902; FIG. 2) is designed to introduce overexpression of both ICDH and NAGK proteins to further improve nitrogen utilization in plants.

The DNA sequence encoding NAGK, SEQ ID NO: 3 set out below, includes a chloroplast transit peptide (CTP) obtained from the 5' end of the enolpyruvyl shikimate phosphate synthase (EPSPS) gene from the algae *Chlamydomonas reinhardtii*. The sequence of the CTP is distinguished from the nagk gene in that the nagk gene is shown in boldface type:

SEQ ID NO: 3

ATGCAGCTGCTCAACCAGCGGCAGGCGCTGCGGCTGGGAAGAAGCTCCGC
CAGCAAGAACCAGCAGGTGGCGCCGCTGGCATCAAGCCGGCAAGCAGCC
TCTCCGTCTCCGCCTCCTCCGTGGCGCCGGCGCCGCTCGGGCGCCG
GCCGGCGCCGGCGCCCGCGCGCTGGTGGTGCAGCGCTCCGCCACCAAGGA
GAAGGIGGAGGAGCTCACCATCCAGATGCTGCATGAGGTGATGGTATCA
AGTGGCGCGGCAGCATGCTGGAGCAGCTGCCGGAGAGCTTCTACAACAAG

-continued

CTGGCGACGCTGCAAGCAGAAGGAAGCATCGTCATTGTTTCATGGAGG
20 AGGGCCGGCCATCAACCAGATGCTGGAGCAGCTGAAGATTGAGCCAACT
TCTCAAATGGGCTGAGGGTGACAGATGAGCCAACAATGCAAGCTGTGGAG
ATGGTGCTCTCAGGGCCATCAACAAGCTGGTGGTGGAGGAAGCTGCTGCA
25 CGCCGGCGCAAGGCATGGGGCTCAGCGGCGTGGATGGAAGCCTGCTGC
AAGCTGTTGAGAAGACTCAAGGCCTCGGCTGGTGGGCAGCATCACCGTG
GTGGATCAAGCGCGCTCCAGCTGCTGCTGAGCAATGGCTACATCCCGGT
GGTGTCTCCCATCGCCGTCTCAGAAGATGGAAGAACAAGATACAACTGCA
30 ACGCCGACACCGTCCGCCGGCGCCATTGCTTCAGCTCTCGGCGCCAAGCAG
CTGCTGATGCTCACTGATGTTCTGGCATCTGGGCAGAAAATGAGCTGGG
AGAGAAGCAGCTGCTGCCGACGGTGACAACAGAAGATATTCAGCTGATGA
35 TGAAGAACCAGATCATCACCGCGGCATGATCCCCAAGGTGCAAGCGGCG
CTGGATGCTCTAGCTCAAGGAGTTCAAGAAAGTGGTATCTGCAAAGGAGA
AGCTGAGACGCTGGACGGCGTGGTGAAGGGCATGGCCGTCCGCACCTCCA
40 TCTCCGCCGAGATGAGCAGAGGACAAGATTCTCAAGCCTTCATCAGCAAC
AAGGTGTGAGG

This amino acid sequence of NAGK (after removal of the CTP), SEQ ID NO: 4, is shown here (277 amino acids):

SEQ ID NO: 4:

MLHEVMVIKC GGSMLQLPE SFYNKLATLQ AEGRSIVIVH
GGGPAINQML EQLKIEPTFS NGLRVTDEPT MQAVEMVLSG
50 PINKLVVRKL LHAGGKAWGL SGVDGSLLOA VEKTOGLGLV
GSITVVDQAP LQLLLSNGYI PVVSPIAVSE DGRTRYNCNA
DTVAGAIASA LGAKQLMLT DVPGIWAENE LGEKQLLPTV
55 TTEDIQLMMK NQIITGMIP KVQAALDALA QGVQEVVICK
GEAETLDGVV KGMVGTSSIS AEMSRGQDSQ AFISNKV

Maize Transformation with Vector pMDO8902

The plant vectors pMDO8902 was transformed into *Agrobacterium* and subsequently entered into plant transformation experiments, as previously described, to introduce the stacked genes into the maize genome. The selection agent for these experiments was glyphosate.

Western Blot Analysis

As before, the generated polyclonal antibodies were used in Western blots to determine if the transgenic protein was being expressed. At the T0 stage most icdh events were

found to be expressing ICDH. Two events were promoted to the T1 stage and each of these had good expression levels. Nitrogen Assays, T0 Events

A series of assays that quantify nitrogen intermediates in plants have been developed. These nitrogen assay methods are described in the previous section. Briefly, each of the plants was sampled (leaf) following 4 weeks of growth in soil in a greenhouse. These leaf samples were processed to

determine their nitrate, asparagine, glutamine, aspartic acid, glutamic acid, ammonium, total amino acid, chlorophyll and total protein levels. Included alongside in the analysis were plants that were transformed with a construct containing only the glyphosate selectable marker (no icdh, nagk, genes) and are referred to as control "non-GOI" plants.

The T0 data is shown in Table 5.

TABLE 5

Nitrogen levels, ICDH + NAGK vs. non-GOI maize events, 4 weeks after transfer to soil									
Plant #	GOI	Aspartic Acid ($\mu\text{g/g}$)	Glutamic Acid ($\mu\text{g/g}$)	Asparagine ($\mu\text{g/g}$)	Glutamine ($\mu\text{g/g}$)	Total Chlorophyll (a + b) (mg/g)	Ammonium ($\mu\text{g/g}$)	Total Amino Acids (mg/g)	Total Protein (mg/g)
15102	ICDH + NAGK	146.0	785.7	1230.0	1464.9	0.092	754.605	216.736	24.521
15103	ICDH + NAGK	187.1	372.6	973.5	3279.8	0.127	836.415	268.941	43.110
15104	ICDH + NAGK	318.6	905.9	2202.9	7399.3	0.370	1087.857	605.138	53.679
15105	ICDH + NAGK	264.6	1082.6	2941.9	11934.4	0.134	865.201	440.129	25.618
15106	ICDH + NAGK	194.3	438.0	1766.1	10241.0	0.144	724.948	390.051	43.614
15107	ICDH + NAGK	122.4	481.7	1683.7	7154.5	0.126	582.000	282.809	30.601
15108	ICDH + NAGK	222.0	27.8	1839.7	4568.3	0.152	700.611	295.594	35.903
15109	ICDH + NAGK	466.7	603.5	1699.4	8891.9	0.281	851.859	347.156	28.974
15110	ICDH + NAGK	124.8	475.6	996.7	6484.5	0.114	654.786	236.433	37.768
IC 1 Control	non-GOI	537.8	596.3	2335.0	3516.7	0.154	984.3	362.2	28.6
IC 2 Control	non-GOI	101.6	473.7	1490.3	2485.0	0.157	505.7	264.3	34.6
IC 3 Control	non-GOI	153.6	914.9	1799.3	2191.5	0.066	534.6	297.8	34.0
IC 4 Control	non-GOI	59.7	526.2	313.0	659.3	0.082	347.3	183.7	24.1
Avg		213.2	627.8	1484.4	2213.1	0.114	593.0	277.0	30.3
Std Dev		219.8	197.9	855.4	1181.5	0.048	273.6	74.3	5.0

Compared to the control (and standard deviation), one T0 event (15109) had higher levels of aspartic acid. Several events had higher glutamine levels, higher total amino acids and higher protein compared to the control. From these results the events 15105 (elevated Glu, Asp, Gln, total AAs) and 15106 (elevated Gln, total AAs, protein) were selected for T1 stage evaluations.

T1 Results

For each of the selected events, six T1 plants were produced, grown and sampled as described previously. Assays were performed and the results are shown in Table 6.

TABLE 6

Nitrogen levels, ICDH + NAGK vs. non-GOI maize events, 4 weeks after transfer to soil									
Plant #	GOI	Aspartic Acid ($\mu\text{g/g}$)	Glutamic Acid ($\mu\text{g/g}$)	Asparagine ($\mu\text{g/g}$)	Glutamine ($\mu\text{g/g}$)	Total Chlorophyll (a + b) (mg/g)	Ammonium ($\mu\text{g/g}$)	Total Amino Acids (mg/g)	Total Protein (mg/g)
15105, #5	ICDH + NAGK	947	2789	1269	13837	0.544	930	555	18.7
15105, #13	ICDH + NAGK	827	1809	298	3163	0.312	619	339	22.0
15105, #14	ICDH + NAGK	430	1825	371	2818	0.384	611	511	17.6
15105, #15	ICDH + NAGK	785	3149	335	6217	0.214	460	423	21.7
15105, #17	ICDH + NAGK	274	1006	388	2632	0.609	425	368	19.9
15105, #18	ICDH + NAGK	453	2030	320	2523	0.233	485	247	17.1
Avg		619	2101	497	5198	0.383	588	407	19.5
Std Dev		442	766	380	4455	0.163	185	114	2.1
15106, #5	ICDH + NAGK	722	3797	3858	10471	0.674	963	425	17.9
15106, #9	ICDH + NAGK	184	2469	549	2556	0.279	600	260	13.9
15106, #10	ICDH + NAGK	373	1408	3319	10395	0.234	773	458	21.0
15106, #11	ICDH + NAGK	1463	4102	4382	9148	0.383	637	632	22.1
15106, #13	ICDH + NAGK	544	3437	3294	5693	0.223	884	416	23.3
15106, #17	ICDH + NAGK	591	2331	2790	5572	0.255	815	501	30.4
Avg		646	2924	3032	7306	0.341	779	449	21.4
Std Dev		442	1027	1333	3198	0.173	140	121	5.5
Control 1		253	2351	1759	2412	0.298	567	235	22.0
Control 2		739	2154	3488	3989	0.089	452	226	17.4
Control 3		297	1767	373	2532	0.107	408	197	21.2
Control 4		905	2233	941	3237	0.145	314	190	17.2
Control 5		394	3037	3346	5262	0.155	504	252	19.5
Control 6		789	2535	4166	9084	0.133	459	295	28.9
Avg		563	2346	2346	4419	0.154	451	232	21.0
Std Dev		281	424	1538	2516	0.074	86	38	4.3

Compared to the control average (and Standard Deviation), the averages showed that 15105 had lower asparagine, higher chlorophyll, and higher total amino acids; 15106 had higher glutamate, higher chlorophyll, and higher total amino acids.

Since the stack ICDH+NAGK appeared to have several positive effects, especially in the later nitrogen-containing metabolites (e.g. chlorophyll, total amino acids), we used previous data on NAGK effects (U.S. patent application Ser. No. 12/916,854, filed Nov. 1, 2010, and incorporated herein in its entirety by this reference) and compared that to ICDH alone and ICDH+NAGK. The results are shown in Table 7 (data are % of control).

TABLE 7

Data on NAGK Alone, ICDH Alone and ICDH + NAGK								
	Aspartic Acid (µg/g)	Glutamic Acid (µg/g)	Asparagine (µg/g)	Glutamine (µg/g)	Total Chlorophyll (a + b) (mg/g)	Ammonium (µg/g)	Total Amino Acids (mg/g)	Total Protein (mg/g)
NAGK Alone	162%	155%	191%	140%	90%	120%	124%	105%
ICDH Alone	105%	111%	18%	65%	131%	102%	98%	87%
ICDH + NAGK	112%	107%	75%	141%	234%	152%	184%	97%

We excluded the one plant (15125 #4) from the ICDH values since it seemed to be a high outlier. The results show a pattern where: (1) NAGK alone tends to impact the intermediate N-metabolites (Asp, Asn, Glu, Gln); (2) ICDH alone impacts chlorophyll; and (3) ICDH+NAGK shows impacts on chlorophyll and total amino acids. This stack effect indicates that N was assimilated and moved towards end-products to a greater extent than with either gene alone.

The foregoing description and drawings comprise illustrative embodiments of the present inventions. The foregoing embodiments and the methods described herein may vary based on the ability, experience, and preference of those skilled in the art. Merely listing the steps of the method in a certain order does not constitute any limitation on the order of the steps of the method. The foregoing description and drawings merely explain and illustrate the invention, and the invention is not limited thereto. Those skilled in the art who have the disclosure before them will be able to make modifications and variations therein without departing from the scope of the invention.

SEQUENCE LISTING

SEQ ID NO: 1
 ATGAGCACCCCAAGATCATCTACACCTTGACAGATGAGGCGCCGCGCT
 GGCCACCTACAGCTTGCTGCCATCATCAAGGCTTCTACTGGAAGCTCAG
 GCATTGCTGTGGAACAAGGGACATCTCCCTTGCTGGAAGGCTGATCGCC
 ACCTTCCCAGAATATTTGACAGACACCCAGAAGATCTCTGATGATCTTGC
 TGAGCTGGGGAAGCTGGCCACCACGCCAGATGCCAACATCATCAAGCTGC
 CAAACATCTCTGCTTCAAGTTCCTCAGCTGAAGGCCGCCATCAAGGAACTC
 CAGCAGCAAGGCTACAAGCTGCCAGATTATCCAGAAGAACAAAAACAGA
 CACAGAGAAGGATGTCAAGGCAAGATATGACAAGATCAAGGGCAGCGCCG

-continued

TCAACCCCGTGTGAGAGAAGGAAATTCAGACCGCCGCGCCGCTCTCC
 GTCAAGAACTATGCAAGGAAGCATCTCACAAAGATGGGCGCCTGGAGCGC
 5 CGACAGCAAGAGCCATGTTGCTCACATGGACAATGGAGATTTCTATGGAT
 CAGAGAAGGCGCGCTGATTGGTGTCTCTGGAAGCGTCAAGATTGAGCTG
 ATCGCCAAGGATGGAAGCAGCACCGTGTGAAGGCCAAGACATCAGTTCA
 10 AGCTGGAGAGATCATCGACAGCTCCGTGATGAGCAAGAATGCTCTGAGGA
 ACTTCATTGCTGCCGAGATTGAAGATGCCAAGAAGCAAGGAGTGTCTCTC

-continued

30 TCCGTCCACCTCAAGGCCACCATGATGAAGGTTTCAGATCCCATCATGTT
 TGGCCAGATTGTTTCAGAGTTCTACAAGGATGCTCTCACCAGCATGCTG
 AGGTGCTGAAGCAGATTGGATTTGATGTCAACAATGGCATTGGAGATCTC
 35 TATGCAAGGATCAAGACCTACCAGAAGCAAAGCAGAAGGAGATTGAAGC
 TGACATCCAAGCTGTTTATGCTCAAAGGCCGAGCTGGCAATGGTGAACA
 GCGACAAGGGCATCACCAACCTCCATGTTTCTTCTGATGTCATCGTCGAC
 40 GCCTCCATGCCGCGCCATGATCAGAGATTCAGGGAAGATGTGGGGCCAGA
 TGGCAAGCTGCATGACACCAAGGCCGTCATCCCAGATCGCTGCTATGCTG
 GCGTCTACCAGGTGGTGTGATTGAAGATTGCAAGCAGCATGGCGCCTTCGAC
 45 CCAACAACAATGGGCTCAGTTCCAAATGTTGGGCTGATGGCGCAGAAGGC
 AGAAGAATATGGAAGCCATGACAAGACCTTTCAGATCCCTGCTGATGGCG
 TCGTCCGCGTCACTGATGAAAGCGGCAAGCTGCTGCTGGAGCAATCAGTG
 50 GAAGCTGGAGACATCTGGAGGATGTGCCAAGCAAAGGATGCTCCCATCCA
 AGATTGGGTGAAGCTCGCCGTCAACAGGGCGCGCCACCAACACGCCGG
 CGGTGTTCTGGCTGGACCCAGCAAGGGCTCATGATGCTCAGGTGATCGCC
 55 AAGGTGGAGAGATATCTAAAGGATTATGACACCTCCGGCTGGACATCAG
 GATCTTGTGCGCGGTGGAAGCAACAAGGTTCTCCTTGGCAAGGATCAGAG
 AAGGAAAGGACACCATCTCAGTGACAGGAAATGTGCTGAGGGACTACCTC
 ACCGACCTCTTCCCATCATGGAGCTGGGCACCTCCGCCAAGATGCTCTC
 60 CATTGTTCTCTGATGAGCGGCGCGCCTCTTTGAAACTGGAGCTGGAG
 GATCAGCGCCCAAGCATGTTTCAAGGTTCTTGAAGAAGGCTACCTCAGA
 TGGGACAGCCTTGGAGAGTTCTTGGCGCTCGCCGCTCCTTGGAGCATCT
 65 TGGAAATGCCTACAAGAACCCAAAGGCGCTGGTGTGCTGGCTCCACCTAG

-continued

ATCAAGCTACTGGCAAGATCCTGGACAACAACAAGAGCCCAGCAAGGAAG
 GTTGGTGAGATCGACAACAGAGGAAGCCACTTCTACCTGGCGCTCTACTG
 GGCTCAAGCTCTTGCTGCTCAAACAGAGGACAAGGAGCTACAAGCTCAGT
 TCACCGGCATTGCCAAGGCGCTGACAGACAATGAAACAAAAATTGTTGGA
 GAGCTGGCTGCTGCTCAAGGAAAGCCGGTGGACATTGCTGGCTACTACCA
 TCCAAACACCGACCTCACCAGCAAGGCCATCAGGCCATCTGCCACCTTCA
 ATGCTGCTCTGGCGCCGCTGGCATAGTAAGG

SEQ ID NO: 2

MSTPKIIYTL TDEAPALATY SLLPIIKAFT GSSGIAVETR
 DISLAGRLIA TFPEYLTDTQ KISDDLAE LG KLATTPDANI
 IKLPNISASV PQLKAAIKEL QOQGYKLPDY PEEPKTDEK
 DVKARYDKIK GSAVNPVRE GNSDRRAPLS VKNYARKHPH
 KMGAWSADSK SHVAHMDNGD FYGSEKAALI GAPGSVKIEL
 IAKDGSSTVL KAKTSVQAGE IIDSSVMSKN ALRNFIAAEI
 EDAKQGVLL SVHLKATMMK VSDPIMFGQI VSEFYKDALT
 KHAEVLKQIG FDVNNGIGDL YARIKTLPEA KQKEIEADIQ
 AVYAQRPLA MVNSDKGITN LHVPSDVIVD ASMPAMIRDS
 GKMWGPDKL HDTKAVIPDR CYAGVYQVVI EDCKQHGAFF
 PTTMGSVPNV GLMAQKAEY GSHDKTFQIP ADGVVVRTDE
 SGKLLLEQSV EAGDIWRMCQ AKDAPIQDWV KLA VNRARAT
 NTPAVFWLDP ARAHDAQVIA KVERYLKDYD TSGLDIRILS
 PVEATRFSLA RIREGKDTIS VTGNVLRDYL TDLFPIMELG
 TSAKMLSIVP LMSGGLFET GAGGSAPKHV QQFLEEGYLR
 WDSLGEFLAL AASLEHLGNA YKNPKALVLA STLDQATGKI
 LDNNKSPARK VGEIDNRGSH FYLALYWAQA LAAQTEDKEL
 QAQFTGIACA LTDNETKIVG ELAAAQGKPV DIAGYYHPNT
 DLTSKAIRPS ATFNAALAPL A

SEQ ID NO: 3

ATGCAGCTGCTCAACCAGCGGCAGGCGCTGCGGCTGGGAAGAAGCTCCGC

-continued

CAGCAAGAACCAGCAGGTGGCGCCGCTGGCATCAAGGCCGGCAAGCAGCC
 TCTCCGTCCTCCGCTCCTCCGTGGCGCCGGCGCCGGCCTGCTCGGCGCCG
 5 GCCGGCGCCGGCCCGCGCCGCTGGTGGTGC GCGCCTCCGCCACCAAGGA
 GAAGGTGGAGGAGCTCACCATCCAGATGCTGCATGAGGTGATGGTGATCA
 AGTGC GCGGCAGCATGCTGGAGCAGCTGCCGGAGAGCTTCTACAACAAG
 10 CTGGCGACGCTGCAAGCAGAAGGAAGAAGCATCGTCATTGTTTCATGGAGG
 AGGGCCGGCCATCAACCAGATGCTGGAGCAGCTGAAGATTGAGCCAACCT
 TCTCAAATGGGCTGAGGGTGACAGATGAGCCAACAATGCAAGCTGTGGAG
 15 ATGGTGCTCTCAGGGCCATCAACAAGCTGGTGGTGAGGAAGCTGCTGCA
 CGCCGGCGCAAGGCATGGGGCTCAGCGGCGTGATGGAAGCCTGCTGC
 AAGCTGTTGAGAAGACTCAAGGCCTCGGCTGGTGGCAGCATCACCGTG
 GTGGATCAAGCGCCGCTCCAGCTGCTGCTGAGCAATGGCTACATCCCGGT
 20 GGTGTCTCCCATCGCCGCTCAGAAGATGGAAGAACAAGATACA AACTGCA
 ACGCCGACACCGTCGCCGGCGCCATTGCTTCAGCTCTCGGCGCCAAGCAG
 CTGCTGATGCTCACTGATGTTCTCGGCATCTGGGCAGAAAATGAGCTGGG
 25 AGAGAAGCAGCTGCTGCCGACGGTGACAACAGAAGATATTCAGCTGATGA
 TGAAGAACCAGATCATCACCGCGGCATGATCCCCAAGGTGCAAGCGGCG
 CTGGATGCTCTAGCTCAAGGAGTTCAAGAAGTGGTGATCTGCAAAGGAGA
 30 AGCTGAGACGCTGGACGGCGTGGTGAAGGGCATGGCCGTCGGCACCTCCA
 TCTCCGCCGAGATGAGCAGAGGACAAGATTCTCAAGCCTTCATCAGCAAC
 AAGGTGTGAGG

SEQ ID NO: 4

MLHEVMVIK GGSMLQLPE SFYNKLATLQ AEGRSIVIVH
 GGGPAINQML EQLKIEPTFS NGLRVTDEPT MQAVEMVLSG
 40 PINKLVVRKL LHAGGKAWGL SGVDGSLLOA VEKTQGLGLV
 GSITVVDQAP LQLLLSNGYI PVSPIAVSE DGRTRYNCNA
 DTVAGAIASA LGAKQLMLT DVPGIWAENE LGEKQLLPTV
 45 TTEDIQLMMK NQIITGGMIP KVQAALDALA QGVQEVVICK
 GEAETLDGVV KGMAVGTSIS AEMSRGQDSQ AFISNKV

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 4

<210> SEQ ID NO 1

<211> LENGTH: 2231

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 1

atgagcacc ccaagatcat ctacaccttg acagatgagg cgccggcgct ggccacctac 60
 agcttgctgc ccatcatcaa ggctttcact ggaagctcag gcattgctgt ggaaacaagg 120
 gacatctccc ttgctggaag gctgatcgcc accttcccag aatatttgac agacaccag 180

-continued

```

aagatctctg atgatcttgc tgagctgggg aagctggcca ccacgccaga tgccaacatc 240
atcaagctgc caaacatctc tgcttcagtt cctcagctga aggccgcat caaggaactc 300
cagcagcaag gctacaagct gccagattat ccagaagaac caaaaacaga cacagagaag 360
gatgtcaagg caagatatga caagatcaag ggcagcgccg tcaaccccggt gctgagagaa 420
ggaaattcag accgcccgcg gccgctctcc gtcaagaact atgcaaggaa gcatcctcac 480
aagatgggcg cctggagcgc cgacagcaag agccatgttg ctcacatgga caatggagat 540
ttctatggat cagagaaggc ggcgctgatt ggtgctctcg gaagcgtcaa gattgagctg 600
atcgccaagg atggaagcag caccgtgctg aaggccaaga catcagttca agctggagag 660
atcatcgaca gctccgtgat gagcaagaat gctctgagga acttcattgc tgccgagatt 720
gaagatgcca agaagcaagg agtgctgctc tccgtccacc tcaaggccac catgatgaag 780
gtttcagatc ccatcatgtt tggccagatt gttcagagt tctacaagga tgctctcacc 840
aagcatgctg aggtgctgaa gcagattgga tttgatgtca acaatggcat tggagatctc 900
tatgcaagga tcaagaccct accagaagca aagcagaagg agattgaagc tgacatccaa 960
gctgtttatg ctcaaaggcc gcagctggca atggtgaaca gcgacaaggg catcaccaac 1020
ctccatgttc cttctgatgt catcgtcgac gcctccatgc cggccatgat cagagattca 1080
gggaagatgt gggggccaga tggcaagctg catgacacca aggccgtcat cccagatcgc 1140
tgctatgctg gcgtctacca ggtggtgatt gaagattgca agcagcatgg cgccttcgac 1200
ccaacaacaa tgggctcagt tccaaatggt gggctgatgg cgcagaaggc agaagaatat 1260
ggaagccatg acaagacctt tcagatccct gctgatggcg tcgtccgctg cactgatgaa 1320
agcggcaagc tgctgctgga gcaatcagtg gaagctggag acatctggag gatgtgcaa 1380
gcaaaggatg ctcccatcca agattgggtg aagctcgccg tcaacagggc gcgcgccacc 1440
aacacgccgg cgggtgtctg gctggacca gcaagggctc atgatgctca ggtgatcgcc 1500
aaggtggaga gatatctaaa ggattatgac acctccggcc tggacatcag gatcttgtcg 1560
ccggtggaag caacaaggtt ctcttggca aggatcagag aaggaaagga caccatctca 1620
gtgacaggaa atgtgctgag ggactacctc accgacctct tccccatcat ggagctgggc 1680
acctccgcca agatgctctc cattgttctc ctgatgagcg gcggcggcct ctttgaaact 1740
ggagctggag gatcagcgcc caagcatggt cagcagttcc tggagaagg ctacctcaga 1800
tgggacagcc ttggagagt cctggcgctc gccgcctcct tggagcatct tggaaatgcc 1860
tacaagaacc caaaggcgt ggtgctggcc tccaccctag atcaagctac tggcaagatc 1920
ctggacaaca acaagagccc agcaaggaag gttggtgaga tcgacaacag aggaagccac 1980
ttctacctgg cgctctactg ggctcaagct cttgctgctc aaacagagga caaggagcta 2040
caagctcagt tcaccggcat tgccaaggcg ctgacagaca atgaaacaaa aattggtgga 2100
gagctggctg ctgctcaagg aaagccggtg gacattgctg gctactacca tccaaacacc 2160
gacctacca gcaaggccat caggccatct gccaccttca atgctgctct ggcgcccgtg 2220
gcatagtaag g 2231

```

```

<210> SEQ ID NO 2
<211> LENGTH: 741
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

```

```

<400> SEQUENCE: 2

```

-continued

Met Ser Thr Pro Lys Ile Ile Tyr Thr Leu Thr Asp Glu Ala Pro Ala
 1 5 10 15
 Leu Ala Thr Tyr Ser Leu Leu Pro Ile Ile Lys Ala Phe Thr Gly Ser
 20 25 30
 Ser Gly Ile Ala Val Glu Thr Arg Asp Ile Ser Leu Ala Gly Arg Leu
 35 40 45
 Ile Ala Thr Phe Pro Glu Tyr Leu Thr Asp Thr Gln Lys Ile Ser Asp
 50 55 60
 Asp Leu Ala Glu Leu Gly Lys Leu Ala Thr Thr Pro Asp Ala Asn Ile
 65 70 75 80
 Ile Lys Leu Pro Asn Ile Ser Ala Ser Val Pro Gln Leu Lys Ala Ala
 85 90 95
 Ile Lys Glu Leu Gln Gln Gln Gly Tyr Lys Leu Pro Asp Tyr Pro Glu
 100 105 110
 Glu Pro Lys Thr Asp Thr Glu Lys Asp Val Lys Ala Arg Tyr Asp Lys
 115 120 125
 Ile Lys Gly Ser Ala Val Asn Pro Val Leu Arg Glu Gly Asn Ser Asp
 130 135 140
 Arg Arg Ala Pro Leu Ser Val Lys Asn Tyr Ala Arg Lys His Pro His
 145 150 155 160
 Lys Met Gly Ala Trp Ser Ala Asp Ser Lys Ser His Val Ala His Met
 165 170 175
 Asp Asn Gly Asp Phe Tyr Gly Ser Glu Lys Ala Ala Leu Ile Gly Ala
 180 185 190
 Pro Gly Ser Val Lys Ile Glu Leu Ile Ala Lys Asp Gly Ser Ser Thr
 195 200 205
 Val Leu Lys Ala Lys Thr Ser Val Gln Ala Gly Glu Ile Ile Asp Ser
 210 215 220
 Ser Val Met Ser Lys Asn Ala Leu Arg Asn Phe Ile Ala Ala Glu Ile
 225 230 235 240
 Glu Asp Ala Lys Lys Gln Gly Val Leu Leu Ser Val His Leu Lys Ala
 245 250 255
 Thr Met Met Lys Val Ser Asp Pro Ile Met Phe Gly Gln Ile Val Ser
 260 265 270
 Glu Phe Tyr Lys Asp Ala Leu Thr Lys His Ala Glu Val Leu Lys Gln
 275 280 285
 Ile Gly Phe Asp Val Asn Asn Gly Ile Gly Asp Leu Tyr Ala Arg Ile
 290 295 300
 Lys Thr Leu Pro Glu Ala Lys Gln Lys Glu Ile Glu Ala Asp Ile Gln
 305 310 315 320
 Ala Val Tyr Ala Gln Arg Pro Gln Leu Ala Met Val Asn Ser Asp Lys
 325 330 335
 Gly Ile Thr Asn Leu His Val Pro Ser Asp Val Ile Val Asp Ala Ser
 340 345 350
 Met Pro Ala Met Ile Arg Asp Ser Gly Lys Met Trp Gly Pro Asp Gly
 355 360 365
 Lys Leu His Asp Thr Lys Ala Val Ile Pro Asp Arg Cys Tyr Ala Gly
 370 375 380
 Val Tyr Gln Val Val Ile Glu Asp Cys Lys Gln His Gly Ala Phe Asp
 385 390 395 400
 Pro Thr Thr Met Gly Ser Val Pro Asn Val Gly Leu Met Ala Gln Lys
 405 410 415

-continued

Ala Glu Glu Tyr Gly Ser His Asp Lys Thr Phe Gln Ile Pro Ala Asp
420 425 430

Gly Val Val Arg Val Thr Asp Glu Ser Gly Lys Leu Leu Leu Glu Gln
435 440 445

Ser Val Glu Ala Gly Asp Ile Trp Arg Met Cys Gln Ala Lys Asp Ala
450 455 460

Pro Ile Gln Asp Trp Val Lys Leu Ala Val Asn Arg Ala Arg Ala Thr
465 470 475 480

Asn Thr Pro Ala Val Phe Trp Leu Asp Pro Ala Arg Ala His Asp Ala
485 490 495

Gln Val Ile Ala Lys Val Glu Arg Tyr Leu Lys Asp Tyr Asp Thr Ser
500 505 510

Gly Leu Asp Ile Arg Ile Leu Ser Pro Val Glu Ala Thr Arg Phe Ser
515 520 525

Leu Ala Arg Ile Arg Glu Gly Lys Asp Thr Ile Ser Val Thr Gly Asn
530 535 540

Val Leu Arg Asp Tyr Leu Thr Asp Leu Phe Pro Ile Met Glu Leu Gly
545 550 555 560

Thr Ser Ala Lys Met Leu Ser Ile Val Pro Leu Met Ser Gly Gly Gly
565 570 575

Leu Phe Glu Thr Gly Ala Gly Gly Ser Ala Pro Lys His Val Gln Gln
580 585 590

Phe Leu Glu Glu Gly Tyr Leu Arg Trp Asp Ser Leu Gly Glu Phe Leu
595 600 605

Ala Leu Ala Ala Ser Leu Glu His Leu Gly Asn Ala Tyr Lys Asn Pro
610 615 620

Lys Ala Leu Val Leu Ala Ser Thr Leu Asp Gln Ala Thr Gly Lys Ile
625 630 635 640

Leu Asp Asn Asn Lys Ser Pro Ala Arg Lys Val Gly Glu Ile Asp Asn
645 650 655

Arg Gly Ser His Phe Tyr Leu Ala Leu Tyr Trp Ala Gln Ala Leu Ala
660 665 670

Ala Gln Thr Glu Asp Lys Glu Leu Gln Ala Gln Phe Thr Gly Ile Ala
675 680 685

Lys Ala Leu Thr Asp Asn Glu Thr Lys Ile Val Gly Glu Leu Ala Ala
690 695 700

Ala Gln Gly Lys Pro Val Asp Ile Ala Gly Tyr Tyr His Pro Asn Thr
705 710 715 720

Asp Leu Thr Ser Lys Ala Ile Arg Pro Ser Ala Thr Phe Asn Ala Ala
725 730 735

Leu Ala Pro Leu Ala
740

<210> SEQ ID NO 3
 <211> LENGTH: 1061
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 3

atgcagctgc tcaaccagcg gcagcgctg cggtgggaa gaagctccgc cagcaagaac 60
 cagcaggtgg cgccgctggc atcaaggccg gcaagcagcc tctccgtctc cgctctctcc 120
 gtggcgccgg cgccggcctg ctgggcgccc gccggcgcgc gccgcccgc cgtggtggtg 180

-continued

```

cgcgctccg ccaccaagga gaaggtggag gagctcacca tccagatgct gcatgaggtg 240
atggtgatca agtgcggcgg cagcatgctg gagcagctgc cggagagctt ctacaacaag 300
ctggcgagcg tgcaagcaga aggaagaagc atcgtcattg ttcattggagg agggccggcc 360
atcaaccaga tgctggagca gctgaagatt gagccaacct tctcaaatgg gctgaggggtg 420
acagatgagc caacaatgca agctgtggag atggtgctct cagggcccat caacaagctg 480
gtggtgagga agctgctgca cgccggcggc aaggcatggg gcctcagcgg cgtggatgga 540
agcctgctgc aagctgttga gaagactcaa ggctcggcc tggggggcag catcacctg 600
gtggatcaag cgccgctcca gctgctgctg agcaatggct acatcccggg ggtgtctccc 660
atcgccgtct cagaagatgg aagaacaaga tacaactgca acgcccacac cgtcgccggc 720
gccattgctt cagctctcgg cgccaagcag ctgctgatgc tcaactgatgt tcctggcatc 780
tgggcagaaa atgagctggg agagaagcag ctgctgccga cggtgacaac agaagatatt 840
cagctgatga tgaagaacca gatcatcacc ggccgatga tccccaaagt gcaagcggcg 900
ctggatgctc tagctcaagg agttcaagaa gtggtgatct gcaaaggaga agctgagacg 960
ctggacggcg tggatgaagg catggccgctc ggcacctcca tctccgccga gatgagcaga 1020
ggacaagatt ctcaagcctt catcagcaac aaggtgtgag g 1061

```

```

<210> SEQ ID NO 4
<211> LENGTH: 277
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct

```

<400> SEQUENCE: 4

```

Met Leu His Glu Val Met Val Ile Lys Cys Gly Gly Ser Met Leu Glu
1          5          10          15

Gln Leu Pro Glu Ser Phe Tyr Asn Lys Leu Ala Thr Leu Gln Ala Glu
20          25          30

Gly Arg Ser Ile Val Ile Val His Gly Gly Gly Pro Ala Ile Asn Gln
35          40          45

Met Leu Glu Gln Leu Lys Ile Glu Pro Thr Phe Ser Asn Gly Leu Arg
50          55          60

Val Thr Asp Glu Pro Thr Met Gln Ala Val Glu Met Val Leu Ser Gly
65          70          75          80

Pro Ile Asn Lys Leu Val Val Arg Lys Leu Leu His Ala Gly Gly Lys
85          90          95

Ala Trp Gly Leu Ser Gly Val Asp Gly Ser Leu Leu Gln Ala Val Glu
100         105         110

Lys Thr Gln Gly Leu Gly Leu Val Gly Ser Ile Thr Val Val Asp Gln
115        120        125

Ala Pro Leu Gln Leu Leu Leu Ser Asn Gly Tyr Ile Pro Val Val Ser
130        135        140

Pro Ile Ala Val Ser Glu Asp Gly Arg Thr Arg Tyr Asn Cys Asn Ala
145        150        155        160

Asp Thr Val Ala Gly Ala Ile Ala Ser Ala Leu Gly Ala Lys Gln Leu
165        170        175

Leu Met Leu Thr Asp Val Pro Gly Ile Trp Ala Glu Asn Glu Leu Gly
180        185        190

Glu Lys Gln Leu Leu Pro Thr Val Thr Thr Glu Asp Ile Gln Leu Met
195        200        205

```

-continued

Met	Lys	Asn	Gln	Ile	Ile	Thr	Gly	Gly	Met	Ile	Pro	Lys	Val	Gln	Ala
	210					215					220				
Ala	Leu	Asp	Ala	Leu	Ala	Gln	Gly	Val	Gln	Glu	Val	Val	Ile	Cys	Lys
225					230				235					240	
Gly	Glu	Ala	Glu	Thr	Leu	Asp	Gly	Val	Val	Lys	Gly	Met	Ala	Val	Gly
				245						250				255	
Thr	Ser	Ile	Ser	Ala	Glu	Met	Ser	Arg	Gly	Gln	Asp	Ser	Gln	Ala	Phe
			260					265						270	
Ile	Ser	Asn	Lys	Val											
			275												

We claim:

1. A construct comprising a nucleotide sequence selected from the group consisting of:

- a) the nucleotide sequence SEQ ID NO: 1; and
- b) a nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 2; and, operably linked to a heterologous 5' DNA plant promoter.

2. An expression vector comprising a nucleotide sequence selected from the group consisting of:

- a) the nucleotide sequence SEQ ID NO: 1; and
 - b) a nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 2;
- and an operably linked 5' DNA plant promoter sequence; wherein said nucleotide sequence modulates nitrogen use in a plant.

3. The expression vector according to claim 2, further comprising a 3' terminator sequence, wherein the nucleotide sequence, the DNA promoter sequence, and the terminator sequence are operatively coupled to permit transcription of the nucleotide sequence.

4. The expression vector according to claim 3, wherein the promoter sequence is selected from the group consisting of constitutive plant promoters and tissue specific promoters.

5. A plant, transformed with the construct of claim 1; wherein said nucleotide sequence modulates nitrogen use in a plant.

6. The plant according to claim 5, wherein the plant is selected from the group consisting of corn (maize); sorghum; wheat; sunflower; tomato; crucifers; peppers; potato; cotton; rice; soybean; sugarbeet; sugarcane; tobacco; barley; and oilseed rape; *Brassica* sp.; alfalfa; rye; millet; safflower; peanuts; sweet potato; cassava; coffee; coconut; pineapple; cocoa; tea; banana; avocado; fig; guava; mango; olive; papaya; cashew; macadamia; almond; oats; vegetables; grasses; vegetables, including but not limited to, onions, tomatoes, lettuce, green beans, lima beans, peas, and members of the genus *Cucumis* such as cucumber, cantaloupe, and muskmelon; ornamentals, including, but are not limited to, azalea, hydrangea, hibiscus, roses, tulips, daffodils, petunias, carnation, poinsettia, and chrysanthemum; pulp trees; oil palm; and conifers.

7. A component part of the plant of claim 6 that contains the construct, and further comprises a nucleotide sequence selected from the group consisting of: the nucleotide sequence SEQ ID NO: 3; and a nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 4.

8. A plant seed produced from the plant of claim 6 that contains the construct, and further comprises a second nucleotide sequence selected from the group consisting of:

the nucleotide sequence SEQ ID NO: 3; and a nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 4.

9. A plant seed transformed with the vector of claim 2.

10. A host cell, transformed with the construct of claim 1; wherein said nucleotide sequence modulates nitrogen use in a plant.

11. The host cell of claim 10, wherein said host cell further comprises at least a second nucleotide sequence selected from the nucleotide sequences SEQ ID NO: 3 and a nucleotide sequence encoding the amino acid sequences of SEQ ID NO: 4.

12. The host cell according to claim 10, wherein the host cell is selected from the group consisting of bacterial cells and plant cells.

13. A vector construct, comprising:

a) a first nucleotide sequence encoding an amino acid sequence selected from the group consisting of:

- i) the nucleotide sequences SEQ ID NO: 1;
- ii) a nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 2;

wherein said nucleotide sequence modulates nitrogen use in a plant;

b) a 5' DNA plant promoter sequence; and,

c) a 3' terminator sequence, wherein the nucleotide sequence, the DNA promoter sequence, and the terminator sequence are operatively linked to permit transcription of the nucleotide sequence.

14. The vector construct according to claim 13, further comprising a second nucleotide sequence selected from the group consisting of: the nucleotide sequence SEQ ID NO: 3; and a nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 4.

15. A method of expressing a nucleic acid molecule that modulates nitrogen in a plant, said method comprising the steps of providing a transgenic plant or plant seed transformed with a vector construct according to claim 2, and growing the transgenic plant or a plant grown from the transgenic plant seed under conditions effective to express the nucleic acid molecule in said transgenic plant or said plant grown from the transgenic plant seed.

16. The method according to claim 15, wherein expression of the nucleic acid molecule is effective in alleviating a limitation such that yield is increased in said transgenic plant or said plant grown from the transgenic plant seed.

17. The method according to claim 15, wherein expression of the nucleic acid molecule is effective in increasing efficiency of nitrogen utilization of said transgenic plant or said plant grown from the transgenic plant seed.

18. The method according to claim 15, wherein the plant is selected from the group consisting of corn (maize);

sorghum; wheat; sunflower; tomato; crucifers; peppers; potato; cotton; rice; soybean; sugarbeet; sugarcane; tobacco; barley; and oilseed rape; *Brassica* sp.; alfalfa; rye; millet; safflower; peanuts; sweet potato; cassava; coffee; coconut; pineapple; cocoa; tea; banana; avocado; fig; guava; mango; olive; papaya; cashew; macadamia; almond; oats; vegetables; grasses; vegetables, including but not limited to, onions, tomatoes, lettuce, green beans, lima beans, peas, and members of the genus *Cucumis* such as cucumber, cantaloupe, and muskmelon; ornamentals, including, but are not limited to, azalea, hydrangea, hibiscus, roses, tulips, daffodils, petunias, carnation, poinsettia, and chrysanthemum; pulp trees; oil palm; and conifers.

19. The method according to claim 15, wherein expression of the nucleic acid molecule is effective in improving the stress tolerance of said transgenic plant or said plant grown from the transgenic plant seed.

20. The method according to claim 15, wherein expression of the nucleic acid molecule is effective in altering the

morphology of said transgenic plant or said plant grown from the transgenic plant seed.

21. A transgenic corn plant, comprising a corn plant transformed with a nucleotide sequence comprising a gene encoding a cytosolic ICDH enzyme and at least an NAGK gene that improve nitrogen utilization efficiency in corn wherein the transformed corn plant has increased chlorophyll over the starting corn plant.

22. A transgenic corn plant, comprising a corn plant transformed with a nucleotide sequence comprising a gene encoding a cytosolic ICDH enzyme and at least an nagk gene that improve nitrogen utilization efficiency in corn wherein the transformed corn plant has increased total amino acids over the starting corn plant.

23. A method of improving the grain number of a corn plant, comprising the steps of introducing into the genome of the plant a cytosolic icdh gene and an nagk gene that improves nitrogen utilization efficiency and growing the transformed plant to produce grain.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 9,725,701 B2
APPLICATION NO. : 13/833247
DATED : August 8, 2017
INVENTOR(S) : McLaren et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

In the Title:

There is a typo in the title. It should read: “A **Prokaryotic**-type Isocitrate Dehydrogenase and Its Application for Improving Nitrogen Utilization in Transgenic Plants”

Signed and Sealed this
Twenty-fourth Day of October, 2017



Joseph Matal

*Performing the Functions and Duties of the
Under Secretary of Commerce for Intellectual Property and
Director of the United States Patent and Trademark Office*