



US009803249B2

(12) **United States Patent**
Taylor et al.

(10) **Patent No.:** **US 9,803,249 B2**
(45) **Date of Patent:** ***Oct. 31, 2017**

(54) **DETECTING NEOPLASM**

(56) **References Cited**

(71) Applicant: **Mayo Foundation for Medical Education and Research**, Rochester, MN (US)

U.S. PATENT DOCUMENTS

(72) Inventors: **William R. Taylor**, Lake City, MN (US); **Jonathan J. Harrington**, Madison, WI (US); **Patrick S. Quint**, Kasson, MN (US); **Hongzhi Zou**, Middleton, WI (US); **Harold S. Bergen, III**, Spring Valley, MN (US); **David I. Smith**, Rochester, MN (US); **David A. Ahlquist**, Rochester, MN (US)

6,143,529	A	11/2000	Lapidus et al.
8,673,555	B2	3/2014	Taylor
2007/0172823	A1	7/2007	Steinberg
2007/0202513	A1	8/2007	Shuber
2016/0168643	A1	6/2016	Ahlquist

FOREIGN PATENT DOCUMENTS

WO	2001/042781	6/2001
WO	2003/087390	10/2003
WO	2006/094149	9/2006
WO	06/119434	11/2006
WO	2007/134779	11/2007

OTHER PUBLICATIONS

(73) Assignee: **MAYO FOUNDATION FOR MEDICAL EDUCATION AND RESEARCH**, Rochester, MN (US)

Kraunz et al (British Journal of Cancer, 2005, 93: 949-952).*

Dai et al (Neoplasia, 2001, 3(4): 314-323).*

Horikoshi et al (Leuk Res, 1994, 18(9): Abstract).*

Ahlquist et al., 2000, "Colorectal cancer screening by detection of altered human DNA in stool: feasibility of a multitarget assay panel", Gastroenterology, 119:1219.

Boynton et al., 2003, "DNA integrity as a potential marker for stool-based detection of colorectal cancer", Clin. Chem., 49:1058.

Davies, 2005, "Somatic mutations of the protein kinase gene family in human lung cancer", Cancer Res. 65(17):7591-5.

Greenman et al., 2007, "Patterns of somatic mutation in human cancer genomes", Nature, 446(7132):153-8.

Sjoblom et al., 2006, "The consensus coding sequences of human breast and colorectal cancers", Science, 314(5797):268-74.

Soussi and Lozano, 2005, "p53 mutation heterogeneity in cancer", Biochem. Biophys. Res. Commun. 331(3):834-42.

Stephens et al., 2005, "A screen of the complete protein kinase gene family identifies diverse patterns of somatic mutations in human breast cancer", Nat. Genet. 37(6):590-2.

Wilm et al., 1996, "Femtomole sequencing of proteins from polyacrylamide gels by nano-electrospray mass spectrometry", Nature, 379:466-469.

Wood et al., 2007, "The genomic landscapes of human breast and colorectal cancers", Science, 318(5853):1108-13.

Zou et al., 2006, "A sensitive method to quantify human long DNA in stool: relevance to colorectal cancer screening", Cancer Epidemiol. Biomarkers Prev, 15(6):1115.

Zou et al., 2007, "Highly methylated genes in colorectal neoplasia: implications for screening", Cancer Epidemiol. Biomarkers Prev, 16(12):2686-2696.

Zou et al., 2008, "High detection rates of colorectal neoplasia by stool DNA testing with a novel digital melt curve assay", Gastroenterology 136(2):459-70.

Cavestro, G., et al., "Role of faecal elastase 1 in pancreatic cancer: A pilot study," Pancreas, Raven press, new York, NY, US, vol. 29, No. 4, Nov. 5, 2004, pp. 349-350.

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **15/467,739**

(22) Filed: **Mar. 23, 2017**

(65) **Prior Publication Data**

US 2017/0198361 A1 Jul. 13, 2017

Related U.S. Application Data

(63) Continuation of application No. 15/197,105, filed on Jun. 29, 2016, now Pat. No. 9,632,093, which is a continuation of application No. 14/827,013, filed on Aug. 14, 2015, now Pat. No. 9,399,800, which is a continuation of application No. 14/168,552, filed on Jan. 30, 2014, now Pat. No. 9,121,070, which is a continuation of application No. 12/866,558, filed as application No. PCT/US2009/033793 on Feb. 11, 2009, now Pat. No. 8,673,555.

(60) Provisional application No. 61/029,221, filed on Feb. 15, 2008.

(51) **Int. Cl.**
C12Q 1/68 (2006.01)
G01N 33/574 (2006.01)

(52) **U.S. Cl.**
CPC **C12Q 1/6886** (2013.01); **G01N 33/57419** (2013.01); **C12Q 2600/154** (2013.01); **C12Q 2600/156** (2013.01); **C12Q 2600/158** (2013.01); **G01N 2333/928** (2013.01); **G01N 2333/966** (2013.01); **G01N 2560/00** (2013.01)

(58) **Field of Classification Search**
None
See application file for complete search history.

(Continued)

Primary Examiner — Sean E Aeder
(74) *Attorney, Agent, or Firm* — Casimir Jones, SC

(57) **ABSTRACT**

This document relates to methods and materials for detecting premalignant and malignant neoplasms. For example, methods and materials for determining whether or not a stool sample from a mammal contains nucleic acid markers or polypeptide markers of a neoplasm are provided.

5 Claims, 12 Drawing Sheets

(56)

References Cited

OTHER PUBLICATIONS

Shimada, Shimya, et al., "Pancreatic elastase IIIA and its variants are expressed in pancreatic carcinoma cells," *International Journal of Molecular Medicine*, Spandidos Publications, GR, vol. 10, No. 5, Nov. 1, 2002.

Haug, Ulrike, et al., "Mutant-enriched PCR and allele-specific hybridization reaction to detect K-ras mutations in stool DNA: High prevalence in a large sample of older adults," *Clinical Chemistry*, vol. 53, No. 4, Apr. 2007.

Naruse, S., et al., "Fecal pancreatic elastase: A reproducible marker for severe exocrine pancreatic insufficiency," *Journal of Gastroenterology* 200609 JP LNKD, vol. 41, No. 9, Sep. 2006.

Jiang X, et al., "T1102 Detection of Colorectal Neoplasia by Stool DNA Testing: High Discrimination with Multi-Marker Quantitation," *Gastroenterology*, Elsevier, Philadelphia, PA, vol. 134, No. 4, Apr. 1, 2008, pp. A-484.

Hanley, Robert, et al., "DNA Integrity Assay: A Plasma-Based Screening Tool for the Detection of Prostate Cancer," *Clin. Cancer Res.* vol. 12 No. 15 pp. 4569-4574, Aug. 1, 2006.

Koinuma et al. Screening for genomic fragments that are methylated specifically in colorectal carcinoma with a methylated MLH1 promoter (*Carcinogenesis*, 2005, 26(12): 2078-2085).

Etzioni et al The Case for Early Detection (*Nature Reviews*, Apr. 2003, 3: internet pp. 1-10).

Mercer Use of Multiple Markers to Enhance Clinical Utility (*Immunol Ser.*, 1990, 53:39-54).

Wang, et al. Usefulness of p53 gene mutations in the supernatant of bile for diagnosis of biliary tract carcinoma: comparison with K-ras mutation (*J. Gastroenterol*, 2002, 37:831-839).

Kraunz et al., (*British Journal of Cancer*, 2005, 93:949-952).

Nelson et al. (*Cancer Research*, 1999, 59: 4570-4573).

Montiero et al. (*J. Clin Microbiol*, 1997, 35(4): 995-998).

* cited by examiner

FIG. 1




FIG. 2




FIG. 3




FIG. 4





FIG. 5

A. K-ras mutation (5571G>T,12G>V)



B. APC mutation (102457delC)




FIG. 6




FIG. 7




FIG. 8




FIG. 9





FIG. 10






FIG. 12



DETECTING NEOPLASM**CROSS-REFERENCE TO RELATED APPLICATIONS**

This application is a continuation of U.S. patent application Ser. No. 15/197,105, filed Jun. 29, 2016, which is a continuation of U.S. patent application Ser. No. 14/827,013, now U.S. Pat. No. 9,309,800, which is a continuation of U.S. patent application Ser. No. 14/168,552, now U.S. Pat. No. 9,121,070, which is a continuation of U.S. patent application Ser. No. 12/866,558, now U.S. Pat. No. 8,673,555, which is a Section 371 U.S. national stage entry of international Patent Application No. PCT/US2009/033793, filed Feb. 11, 2009, which claims priority to U.S. Provisional Patent Application No. 61/029,221, filed Feb. 15, 2008, the contents of which are hereby incorporated by reference in their entireties.

BACKGROUND

1. Technical Field

This document relates to methods and materials involved in detecting premalignant and malignant neoplasms (e.g., colorectal and pancreatic cancer).

2. Background Information

About half of all cancer deaths in the United States result from aero-digestive cancer. For example, of the estimated annual cancer deaths, about 25 percent result from lung cancer; about 10 percent result from colorectal cancer; about 6 percent result from pancreas cancer; about 3 percent result from stomach cancer; and about 3 percent result from esophagus cancer. In addition, over 7 percent of the annual cancer deaths result from other aero-digestive cancers such as naso-oro-pharyngeal, bile duct, gall bladder, and small bowel cancers.

SUMMARY

This document relates to methods and materials for detecting premalignant and malignant neoplasms (e.g., colorectal and pancreatic cancer). For example, this document provides methods and materials that can be used to determine whether a sample (e.g., a stool sample) from a mammal contains a marker for a premalignant and malignant neoplasm such as a marker from a colonic or supracolonic aero-digestive neoplasm located in the mammal. The detection of such a marker in a sample from a mammal can allow a clinician to diagnose cancer at an early stage. In addition, the analysis of a sample such as a stool sample can be much less invasive than other types of diagnostic techniques such as endoscopy.

This document is based, in part, on the discovery of particular nucleic acid markers, polypeptide markers, and combinations of markers present in a biological sample (e.g., a stool sample) that can be used to detect a neoplasm located, for example, in a mammal's small intestine, gall bladder, bile duct, pancreas, liver, stomach, esophagus, lung, or naso-oro-pharyngeal airways. For example, as described herein, stool can be analyzed to identify mammals having cancer. Once a particular mammal is determined to have stool containing a neoplasm-specific marker or collection of markers, additional cancer screening techniques can be used to identify the location and nature of the neoplasm. For example, a stool sample can be analyzed to determine that the patient has a neoplasm, while magnetic resonance imaging (MRI), endoscopic analysis, and tissue biopsy tech-

niques can be used to identify the location and nature of the neoplasm. In some cases, a combination of markers can be used to identify the location and nature of the neoplasm without additional cancer screening techniques such as MRI, endoscopic analysis, and tissue biopsy techniques.

In general, one aspect of this document features a method of detecting pancreatic cancer in a mammal. The method comprises, or consists essentially of determining the ratio of an elastase 3A polypeptide to a pancreatic alpha-amylase polypeptide present within a stool sample. The presence of a ratio greater than about 0.5 indicates that the mammal has pancreatic cancer. The presence of a ratio less than about 0.5 indicates that the mammal does not have pancreatic cancer.

In another aspect, this document features a method of detecting pancreatic cancer in a mammal. The method comprises or consists essentially of determining the level of an elastase 3A polypeptide in a stool sample from the mammal. The presence of an increased level of an elastase 3A polypeptide, when compared to a normal control level, is indicative of pancreatic cancer in the mammal.

In another aspect, this document features a method of detecting pancreatic cancer in a mammal. The method comprises, or consists essentially of, determining the level of a carboxypeptidase B polypeptide in a stool sample from the mammal. An increase in the level of a carboxypeptidase B polypeptide, when compared to a normal control level, is indicative of pancreatic cancer in the mammal.

In another aspect, this document features a method of detecting pancreatic cancer in a mammal. The method comprises, or consists essentially of, determining whether or not a stool sample from the mammal comprises a ratio of a carboxypeptidase B polypeptide to a carboxypeptidase A2 polypeptide that is greater than about 0.5. The presence of the ratio greater than about 0.5 indicates that the mammal has pancreatic cancer.

In another aspect, this document features a method of detecting cancer or pre-cancer in a mammal. The method comprises, or consists essentially of, determining whether or not a stool sample from the mammal has an increase in the number of DNA fragments less than 200 base pairs in length, as compared to a normal control. The presence of the increase in the number of DNA fragments less than 200 base pairs in length indicates that the mammal has cancer or pre-cancer. The DNA fragments can be less than 70 base pairs in length.

In another aspect, this document features a method of detecting colorectal cancer or pre-cancer in a mammal. The method comprises, or consists essentially of, determining whether or not a stool sample from the mammal has an elevated K-ras (Kirsten rat sarcoma-2 viral (v-Ki-ras2) oncogene homolog (GenBank accession no. NM_033360; gil34485724)) mutation score, an elevated BMP3 (bone morphogenetic protein 3 (GenBank accession no. M22491; gil179505)) methylation status, and an elevated level of human DNA as compared to a normal control. The presence of the elevated K-ras mutation score, elevated BMP3 methylation status, and elevated level of human DNA level indicates that the mammal has colorectal cancer or pre-cancer. The K-ras mutation score can be measured by digital melt curve analysts. The K-ras mutation score can be measured by quantitative allele specific PCR.

In another aspect, this document features a method of detecting aero-digestive cancer or pre-cancer in a mammal. The method comprises, or consists essentially of, determining whether or not a stool sample from the mammal has an elevated K-ras mutation score, an elevated BMP3 methylation status, and an elevated level of human DNA as

compared to a normal control. The presence of the elevated K-ras mutation score, elevated BMP3 methylation status, and elevated level of human DNA level indicates that the mammal has aero-digestive cancer or pre-cancer. The K-ras mutation score can be measured by digital melt curve analysis. The K-ras mutation score can be measured by quantitative allele-specific PCR. The method can further comprise determining whether or not a stool sample from the mammal has an elevated APC mutation score. The APC mutation score can be measured by digital melt curve analysis.

In another aspect, this document features a method of detecting aero-digestive cancer or pre-cancer in a mammal. The method comprises, or consists essentially of, determining whether or not the mammal has at least one mutation in six nucleic acids selected from the group consisting of p16, p53, k-ras, APC (adenomatosis polyposis coli tumor suppressor (GenBank accession no. NM_000038; gi|189011564)), SMAD4 (SMAD family member 4 (GenBank accession no. NM_005359; gi|195963400)), EGFR (epidermal growth factor receptor (GenBank accession no. NM_005228 gi|41327737)), CTNNB1 (catenin (cadherin-associated protein), beta 1 (88 kD) (GenBank accession no. X87838; gi|1154853)), and BRAF (B-Raf proto-oncogene serine/threonine-protein kinase (p94) (GenBank accession no. NM_004333; gi|187608632)) nucleic acids. The presence of at least one mutation in each of the six nucleic acids indicates that the mammal has aero-digestive cancer or pre-cancer. The method can further comprise determining whether or not a stool sample from the mammal has an elevated level of a carboxypeptidase B polypeptide as compared to a normal control. The presence of the elevated level of a carboxypeptidase B polypeptide indicates that the mammal has aerodigestive cancer or pre-cancer in the mammal. The method can further comprise determining whether or not a stool sample from the mammal has an elevated amount of DNA fragments less than 70 base pairs in length as compared to a normal control. The presence of the elevated amount of DNA fragments less than 70 base pairs in length indicates that the mammal has aero-digestive cancer or pre-cancer. The method can further comprise determining whether or not a stool sample from the mammal has an elevated amount of DNA fragments greater than 100 base pairs in length as compared to normal controls. The presence of the elevated amount of DNA fragments greater than 100 base pairs in length indicates that the mammal has aero-digestive cancer or pre-cancer. The method can further comprise determining whether or not a stool sample from the mammal has an elevated BMP3 methylation status. The elevated BMP3 methylation status level indicates that the mammal has aero-digestive cancer or pre-cancer. The determining step can comprise using digital melt curve analysis.

In another aspect, this document features a method of detecting aero-digestive cancer or pre-cancer in a mammal. The method comprises, or consists essentially of, measuring mutations in a matrix marker panel in a stool sample. The marker panel can comprise measuring DNA mutations in p16, p53, k-ras, APC, SMAD4, EGFR, CTNNB1, and BRAF nucleic acids. The presence of a mutation in each of the nucleic acids is indicative of the presence of aero-digestive cancer or pre-cancer in a mammal.

In another aspect, this document features a method of detecting aero-digestive cancer in a mammal. The method comprises, or consists essentially of, determining whether or not the methylation status of an ALX4 (aristaless-like homeobox 4 (GenBank accession no. AF294629, gi|108637481)) nucleic acid in a stool sample from the

mammal is elevated, as compared to a normal control. The presence of an elevated ALX4 methylation status indicates the presence of aero-digestive cancer in the mammal.

In another aspect, this document features a method of diagnosing pancreatic cancer in a mammal. The method comprises, or consists essentially of, obtaining a stool sample from the mammal, determining the ratio of an elastase 3A polypeptide to a pancreatic alpha-amylase polypeptide present within a stool sample, and communicating a diagnosis of pancreatic cancer if the ratio is greater than about 0.5, thereby diagnosing the mammal with pancreatic cancer.

In another aspect, this document features a method of diagnosing a mammal with pancreatic cancer. The method comprises, or consists essentially of, obtaining a stool sample from the mammal, measuring mutations in a matrix marker panel of nucleic acids present in the sample, determining the ratio of a carboxypeptidase B polypeptide to a carboxypeptidase A2 polypeptide present within the sample, and communicating a diagnosis of pancreatic cancer or pre-cancer if a mutation is detected in each of the marker panel nucleic acids and the ratio is greater than 0.5, thereby diagnosing the mammal. The matrix marker panel comprises or consists essentially of p16, p53, k-ras, APC, SMAD4, EGFR, CTNNB1, and BRAF nucleic acids.

In another aspect, this document features method of diagnosing a mammal with colorectal cancer. The method comprises, or consists essentially of, obtaining a stool sample from the mammal, detecting mutations in a matrix marker panel comprising of p16, p53, k-ras, APC, SMAD4, EGFR, CTNNB1, and BRAF nucleic acids in DNA present within the sample, measuring the level of a serotransferrin polypeptide present within the sample, and communicating a diagnosis of colorectal cancer or pre-cancer if a mutation is detected in each of the nucleic acids and the level of a serotransferrin polypeptide is elevated as compared to a reference level, thereby diagnosing the mammal.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention pertains. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

The details of one or more embodiments of the invention are set forth in the accompanying drawings and the description below. Other features, objects, and advantages of the invention will be apparent from the description and from the claims.

DESCRIPTION OF THE DRAWINGS

FIG. 1: Adjusted Cut-off Levels with Quantitative Stool Markers to Achieve 95% Specificity across Age and Gender using the Q-LEAD Model. Solid line for women, dotted line for men.

FIG. 2: Sensitive and specific detection of pancreatic cancer by fecal ratio of carboxypeptidase B:carboxypeptidase A2. Note that ratio in stools from patients with colorectal cancer is no different from ratios with healthy controls.

5

FIG. 3: Elastase levels quantified in stools from patients with pancreatic cancer, patients with colorectal cancer, and healthy controls.

FIG. 4: Ratio of elastase 3A:pancreatic alpha amylase differentiates patients with pancreatic cancer from patients with colorectal cancer and from healthy controls.

FIG. 5: Digital Melt Curve to detect mutations by targeted gene scanning (temperature (x-axis) v. temperature-normalized fluorescence (y-axis)). Eight pairs of primers, which amplify 100-350 bp gene fragments, were used to scan K-ras and APC genes and detect mutations (substitution and deletion mutations, respectively) at 1% mutant/wild type ratio.

FIG. 6: Quantitative detection of low abundance mutations by Digital Melt Curve Assay. Varying Mutant: Wild-type ratios of K-ras and APC gene mixtures were prepared and assayed blindly by Digital Melt Curve.

FIG. 7: High analytical sensitivity by Digital Melt Curve (temperature (x-axis) v. temperature-normalized fluorescence (y-axis)). To test the detection limit of digital melt curve (DMC) assay, mutant copies were spiked in wild-type copies at 0.1, 0.5, 1, 5, and 10% dilutions. DMC could detect up to 0.1% mutant/wild-type level when 1000 copies were dispersed to one 96-well plate. The numbers of positive wells increased proportionally when spiked mutant copies were increased. A pair of primers that amplify 248 bp K-ras gene fragment were used as an example here. Primers that amplify-119 bp K-ras gene, 162 bp APC gene, and 346 bp APC genes were also used to test the detection limit and quantitative property of DMC.

FIG. 8: Superior screen detection of colorectal precancerous polyps by Digital Melt Curve (DMC). Histogram compares sensitivity by DMC with that by common fecal occult blood tests (Hemoccult and HemoccultSENSA) and by the commercial stool DNA test (PreGenPlus, Exact Sciences). Detection by DMC was significantly better than by any other test ($p < 0.05$).

FIG. 9: Distributions of short fragment human DNA (short DNA) and long fragment human DNA (long DNA) in stools from patients with normal colonoscopy, large precancerous adenomas, and colorectal cancer. Human DNA quantified by an assay of Alu repeats. Short DNA represents 45 bp fragment amplification products, and long DNA represents 245 bp amplification products.

FIG. 10: Stool distributions of short and long DNA in patients with pancreatic cancer and in healthy controls. Short DNA represents 45 bp fragment amplification products, and long DNA represents 245 bp amplification products.

FIG. 11: Methylated BMP3 gene in stool for detection of colorectal neoplasia. Methylated BMP3 was blindly quantified in stools from patients with colorectal cancers, precancerous adenomas, and normal individuals with real-time methylation-specific PCR. Each circle represents a stool sample.

FIG. 12: Frequency of Specific Base Changes in Colorectal Tumors.

DETAILED DESCRIPTION

This document provides methods and materials related to detecting a neoplasm in a mammal (e.g., a human). For example, this document provides methods and materials for using nucleic acid markers, polypeptide markers, and combinations of markers present in a biological sample (e.g., a stool sample) to detect a neoplasm in a mammal. Such a neoplasm can be a cancer or precancer in the head and neck,

6

lungs and airways, esophagus, stomach, pancreas, bile ducts, small bowel, or colorectum. It will be appreciated that the methods and materials provided herein can be used to detect neoplasm markers in a mammal having a combination of different neoplasms. For example, the methods and materials provided herein can be used to detect nucleic acid and polypeptide markers in a human having lung and stomach neoplasms.

In some cases, the methods and materials provided herein can be used to quantify multiple markers in biological samples, stool) to yield high sensitivity for detection of lesions (e.g., neoplasms), while preserving high specificity. Such methods can include, for example, a logistic model that adjusts specificity cut-offs based on age, gender, or other variables in a target population to be tested or screened.

In some cases, the methods and materials provided herein can be used to determine whether a mammal (e.g., a human) has colorectal cancer or pancreatic cancer. For example, serotransferin, methylated BMP3, and mutant BRAF markers in stool can be used to identify a mammal as likely having colorectal cancer, while mutant p16, carboxypeptidase B/A, and elastase 2A markers can be used to identify a mammal as likely having pancreatic cancer.

Any suitable method can be used to detect a nucleic acid marker in a mammalian stool sample. For example, such methods can involve isolating DNA from a stool sample, separating out one or more particular DNAs from the total DNA, subjecting the DNAs to bisulfite treatment, and determining whether the separated DNAs are abnormally methylated (e.g., hypermethylated or hypomethylated). In some cases, such methods can involve isolating DNA from a stool sample and determining the presence or absence of DNA having a particular size (e.g., short DNA). It is noted that a single stool sample can be analyzed for one nucleic acid marker or for multiple nucleic acid markers. For example, a stool sample can be analyzed using assays that detect a panel of different nucleic acid markers. In addition, multiple stool samples can be collected from a single mammal and analyzed as described herein.

Nucleic acid can be isolated from a stool sample using, for example, a kit such as the QIAamp DNA Stool Mini Kit (Qiagen Inc., Valencia, Calif.). In addition, nucleic acid can be isolated from a stool sample using the following procedure: (1) homogenizing samples in an excess volume ($>1:7$ w:v) of a stool stability buffer (0.5M Tris pH 9.0, 150 mM EDTA, 10 mM NaCl) by shaking or mechanical mixing; (2) centrifuging a 10 gram stool equivalent of each sample to remove all particulate matter; (3) adding 1 of 100 $\mu\text{g}/\mu\text{L}$, RNase A to the supernatant and incubating at 37° C. for 1 hour; (4) precipitating total nucleic acid with $1/10$ volume 3M NaAc and an equal volume isopropanol; and (5) centrifuging and then resuspending the DNA pellet in TE (0.01 M Tris pH 7.4, 0.001 M EDTA). U.S. Pat. Nos. 5,670,325; 5,741,650, 5,928,870; 5,952,178, and 6,020,137 also describe various methods that can be used to prepare and analyze stool samples.

One or more specific nucleic acid fragments can be purified from a nucleic acid preparation using, for example, a modified sequence-specific hybrid capture technique (see e.g., Ahlquist et al. (2000) *Gastroenterology*, 119:1219-1227). Such a protocol can involve: (1) adding 300 μL of sample preparation to an equal volume of a 6 M guanidine isothiocyanate solution containing 20 pmol biotinylated oligonucleotides (obtained from, for example, Midland Certified Reagent Co., Midland, Tex.) with sequences specific for the DNA fragments to be analyzed; (2) incubating for two hours at 25° C.; (3) adding streptavidin coated magnetic

beads to the solution and incubating for an additional hour at room temperature; (4) washing the bead/hybrid capture complexes four times with IX B+W buffer (1 M NaCl, 0.01 M Tris-HCl pH 7.2, 0.001 M EDTA, 0.1% Tween) 20); and (5) eluting the sequence specific captured DNA into 35 μ L L-TE (1 mM Tris pH 7.4, 0.1 M EDTA) by heat denaturation of the bead/hybrid capture complexes. Any other suitable technique also can be used to isolate specific nucleic acid fragments.

Nucleic acid can be subjected to bisulfite treatment to convert unmethylated cytosine residues to uracil residues, while leaving any 5-methylcytosine residues unchanged. A bisulfite reaction can be performed using, for example, standard techniques: (1) denaturing approximately 1 μ g of genomic DNA (the amount of DNA can be less when using micro-dissected DNA specimens) for 15 minutes at 45° C. with 2 N NaOH; (2) incubating with 0.1 M hydroquinone and 3.6 M sodium bisulfite (pH 5.0) at 55° C. for 4-12 hours; (3) purifying the DNA from the reaction mixture using standard (e.g., commercially-available) DNA miniprep columns or other standard techniques for DNA purification; (4) resuspending the purified DNA sample in 55 μ L water and adding 5 μ L 3 N NaOH for a desulfonation reaction that typically is performed at 40° C. for 5-10 minutes; (5) precipitating the DNA sample with ethanol, washing the DNA, and resuspending the DNA in an appropriate volume of water. Bisulfite conversion of cytosine residues to uracil also can be achieved using other methods (e.g., the CpGenome™ DNA Modification Kit from Serologicals Corp., Norcross, Ga.).

Any appropriate method can be used to determine whether a particular DNA is hypermethylated or hypomethylated. Standard PCR techniques, for example, can be used to determine which residues are methylated, since unmethylated cytosines converted to uracil are replaced by thymidine residues during PCR. PCR reactions can contain, for example, 10 μ L of captured DNA that either has or has not been treated with sodium bisulfite, IX PCR buffer, 0.2 mM dNTPs, 0.5 μ M sequence specific primers (e.g., primers flanking a CpG island within the captured DNA), and 5 units DNA polymerase (e.g., Amplitaq DNA polymerase from PE Applied Biosystems, Norwalk, Conn.) in a total volume of 50 μ L. A typical PCR protocol can include, for example, an initial denaturation step at 94° C. for 5 min, 40 amplification cycles consisting of 1 minute at 94° C., 1 minute at 60° C., and 1 minute at 72° C., and a final extension step at 72° C. for 5 minutes.

To analyze which residues within a captured DNA are methylated, the sequences of PCR products corresponding to samples treated with and without sodium bisulfite can be compared. The sequence from the untreated DNA will reveal the positions of all cytosine residues within the PCR product. Cytosines that were methylated will be converted to thymidine residues in the sequence of the bisulfite-treated DNA, while residues that were not methylated will be unaffected by bisulfate treatment.

Purified nucleic acid fragments from a stool sample or samples can be analyzed to determine the presence or absence of one or more somatic mutations. Mutations can be single base changes, short insertion/deletions, or combinations thereof. Methods of analysis can include conventional Sanger based sequencing, pyrosequencing, next generation sequencing, single molecule sequencing, and sequencing by synthesis. In some cases, mutational status can be determined by digital PCR followed by high resolution melting curve analysis. In other cases, allele specific primers or probes in conjunction with amplification methods can be

used to detect specific mutations in stool DNA. The mutational signature can comprise not only the event of a base or sequence change in a specific gene, but also the location of the change within the gene, whether it is coding, non-coding, synonymous or non-synonymous, a transversion or transition, and the dinucleotide sequence upstream and downstream from the alteration.

In some cases, a sample can be assessed for the presence or absence of a polypeptide marker. For example, any appropriate method can be used to assess a stool sample for a polypeptide marker indicative of a neoplasm. For example, a stool sample can be used in assays designed to detect one or more polypeptide markers. Appropriate methods such as those described elsewhere (Aebersold and Mann, *Nature*, 422:198-207 (2003) and McDonald and Yates, *Dis. Markers*, 18:99-105 (2002)) can be adapted or designed to detect polypeptides in a stool. For example, single-reaction monitoring using a TSQ mass spectrometer can specifically target polypeptides in a stool sample. High resolution instruments like the LTQ-FT or LTQ orbitrap can be used to detect polypeptides present in a stool sample.

The term “increased level” as used herein with respect to the level of an elastase 3A polypeptide is any level that is above a median elastase 3A polypeptide level in a stool sample from a random population of mammals (e.g., a random population of 10, 20, 30, 40, 50, 100, or 500 mammals) that do not have an aero-digestive cancer. Elevated polypeptide levels of an elastase 3A polypeptide can be any level provided that the level is greater than a corresponding reference level. For example, an elevated level of an elastase 3A polypeptide can be 0.5, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more fold greater than the reference level of elastase 3A polypeptide in a normal sample. It is noted that a reference level can be any amount. For example, a reference level for an elastase 3A polypeptide can be zero. In some cases, an increased level of an elastase 3A polypeptide can be any detectable level of an elastase 3A polypeptide in a stool sample.

The term “increased level” as used herein with respect to the level of a carboxypeptidase B polypeptide level is any level that is above a median carboxypeptidase B polypeptide level in a stool sample from a random population of mammals (e.g., a random population of 10, 20, 30, 40, 50, 100, or 500 mammals) that do not have an aero-digestive cancer. Elevated polypeptide levels of carboxypeptidase B polypeptide can be any level provided that the level is greater than a corresponding reference level. For example, an elevated level of carboxypeptidase B polypeptide can be 0.5, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more fold greater than the reference level carboxypeptidase B polypeptide observed in a normal stool sample. It is noted that a reference level can be any amount. For example, a reference level for a carboxypeptidase B polypeptide can be zero. In some cases, an increased level of a carboxypeptidase B polypeptide can be any detectable level of a carboxypeptidase B polypeptide in a stool sample.

The term “increased level” as used herein with respect to the level of DNA fragments less than about 200 or less than about 70 base pairs in length is any level that is above a median level of DNA fragments, less than about 200 or less than about 70 base pairs in length in a stool sample from a random population of mammals (e.g., a random population of 10, 20, 30, 40, 50, 100, or 500 mammals) that do not have an aero-digestive cancer. In some cases, an increased level of DNA fragments less than about 200 or less than about 70 base pairs in length can be any detectable level of DNA

fragments less than about 200 or less than about 70 base pairs in length in a stool sample.

The term “elevated methylation” as used herein with respect to the methylation status of a BMP3 or ALX nucleic acid is any methylation level that is above a median methylation level in a stool sample from a random population of mammals (e.g., a random population of 10, 20, 30, 40, 50, 100, or 500 mammals) that do not have an aero-digestive cancer. Elevated levels of BMP3 or ALX methylation can be any level provided that the level is greater than a corresponding reference level. For example, an elevated level of BMP3 or ALX methylation can be 0.5, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more fold greater than the reference level methylation observed in a normal stool sample. It is noted that a reference level can be any amount.

The term “elevated mutation score” as used herein with respect to detected mutations in a matrix panel of particular nucleic acid markers is any mutation score that is above a median mutation score in a stool sample from a random population of mammals (e.g., a random population of 10, 20, 30, 40, 50, 100, or 500 mammals) that do not have an aero-digestive cancer. An elevated mutation score in a matrix panel of particular nucleic acid markers can be any score provided that the score is greater than a corresponding reference score. For example, an elevated score of K-ras or APC mutations can be 0.5, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more fold greater than the reference score of K-ras or APC mutations observed in a normal stool sample it is noted that a reference score can be any amount.

In some cases, a ratio of particular polypeptide markers can be determined and used to identify a mammal having an aero-digestive cancer (e.g., a colorectal cancer or a pancreatic cancer). For example, a ratio provided herein (e.g., the ratio of carboxypeptidase B polypeptide levels to carboxypeptidase A2 polypeptide levels) to can be used as described herein to identify a mammal having a particular neoplasm (e.g., pancreatic cancer).

In some cases, a matrix marker panel can be used to identify mammals having an aero-digestive cancer (e.g., a colorectal cancer or a pancreatic cancer). In some cases, such panel also can identify the location of the aero-digestive cancer. Such a panel can include nucleic acid markers, polypeptide markers, and combinations thereof and can provide information about a mutated marker gene, the mutated region of the marker gene, and/or type of mutation. For example, data can be analyzed using a statistical model to predict tumor site (e.g., anatomical location or tissue of origin) based on inputs from sequencing data (such as by specific nucleic acid or combination of nucleic acids mutated, specific mutational location on a nucleic acid, and nature of mutation insertion, deletion, transition, or transversion) or by any combination thereof) and/or data from polypeptide or other types of markers. For example, a Site of Tumor Estimate (SITE) model can be used to predict tumor site using a matrix panel of markers that are present to variable extent across tumors.

In some cases, data can be analyzed using quantified markers to create a logistic model, which can have both high sensitivity and high specificity. For example, a logistic model can also incorporate population variables like gender and age to adjust cut-off levels for test positivity and thereby optimize assay performance in a screening setting. In some cases, a Quantitative Logistic to Enhance Accurate Detection (Q-LEAD) Model can be used with any marker class or combination of markers as long as they can be quantified.

This document also provides methods and materials to assist medical or research professionals in determining

whether or not a mammal has an aero-digestive cancer. Medical professionals can be, for example, doctors, nurses, medical laboratory technologists, and pharmacists. Research professionals can be, for example, principle investigators, research technicians, postdoctoral trainees, and graduate students. A professional can be assisted by (1) determining the ratio of particular polypeptide markers in a stool sample, and (2) communicating information about the ratio to that professional, for example. In some cases, a professional can be assisted by (1) determining the level of human DNA, the methylation status of genes such as BMP3, and the mutation score of genes such as APC and K-ras, and (2) communicating information about the level of DNA, the methylation status of particular genes, and the mutation score of particular genes to the professional in some cases, a professional can be assisted by (1) detecting mutations in cancer-related genes such as K-ras, p53, APC, p16, EGFR, CTNNB1, BRAF, and SMAD4, as a matrix marker panel, and (2) communicating information regarding the mutations to the professional.

After the ratio of particular polypeptide markers, or presence of particular nucleic acid markers in a stool sample is reported, a medical professional can take one or more actions that can affect patient care. For example, a medical professional can record the results in a patient’s medical record. In some cases, a medical professional can record a diagnosis of an aero-digestive cancer, or otherwise transform the patient’s medical record, to reflect the patient’s medical condition. In some cases, a medical professional can review and evaluate a patient’s entire medical record, and assess multiple treatment strategies, for clinical intervention of a patient’s condition. In some cases, a medical professional can record a tumor site prediction with the reported imitations. In some cases, a medical professional can request a determination of ratio of particular polypeptide markers to predict tumor site. In some cases, a medical professional can review and evaluate a patient’s entire medical record and assess multiple treatment strategies, for clinical intervention of a patient’s condition.

A medical professional can initiate or modify treatment of an aero-digestive cancer after receiving information regarding a ratio of particular polypeptide markers or the presence of nucleic acid markers in a patient’s stool sample. In some cases, a medical professional can compare previous reports and the recently communicated ratio of particular polypeptide markers, or presence of nucleic acid markers, and recommend a change in therapy. In some cases, a medical professional can enroll a patient in a clinical trial for novel therapeutic intervention of an aero-digestive cancer. In some cases, a medical professional can elect waiting to begin therapy until the patient’s symptoms require clinical intervention.

A medical professional can communicate the ratio of particular polypeptide markers to a patient or a patient’s family. In some cases, a medical professional can provide a patient and/or a patient’s family with information regarding aerodigestive cancers, including treatment options, prognosis, and referrals to specialists, e.g., oncologists and/or radiologists. In some cases, a medical professional can provide a copy of a patient’s medical records to communicate the ratio of particular polypeptide markers to a specialist.

A research professional can apply information regarding a subject’s ratio of particular polypeptide markers to advance aerodigestive cancer research. For example, a researcher can compile data on the ratio of particular polypeptide markers, and/or presence of particular nucleic acid

markers, with information regarding the efficacy of a drug for treatment of aero-digestive cancer to identify an effective treatment. In some cases, a research professional can obtain a subject's ratio of particular polypeptide markers, and/or determine the presence of particular nucleic acid markers to evaluate a subject's enrollment, or continued participation in a research study or clinical trial. In some cases, a research professional can classify the severity of a subject's condition, based on the ratio of particular polypeptide markers and/or the levels of particular nucleic acid markers. In some cases, a research professional can communicate a subject's ratio of particular polypeptide markers, and/or the presence of particular nucleic acid markers to a medical professional. In some cases, a research professional can refer a subject to a medical professional for clinical assessment of an aero-digestive cancer, and treatment of an aero-digestive cancer.

Any appropriate method can be used to communicate information to another person (e.g., a professional). For example, information can be given directly or indirectly to a professional. For example, a laboratory technician can input the ratio of particular polypeptide markers and/or particular nucleic acid markers into a computer-based record. In some cases, information is communicated by making a physical alteration to medical or research records. For example, a medical professional can make a permanent notation or flag a medical record for communicating a diagnosis to other medical professionals reviewing the record. In addition, any type of communication can be used to communicate the information. For example, mail, e-mail, telephone, and face-to-face interactions can be used. The information also can be communicated to a professional by making that information electronically available to the professional. For example, the information can be communicated to a professional by placing the information on a computer database such that the professional can access the information. In addition, the information can be communicated to a hospital, clinic, or research facility serving as an agent for the professional.

The invention will be further described in the following examples, which do not limit the scope of the invention described in the claims.

EXAMPLES

Example 1—Multi-Marker Quantitation and a Q-LEAD Model

Most approaches at marker detection in stool have been qualitative. When such qualitative approaches are applied to assay of multiple markers (targeting multiple markers is required with neoplasm detection due molecular heterogeneity), sensitivity is achieved at the expense of compounded non-specificity. Non-specificity can lead to prohibitive programmatic cost with population screening due to the expensive and unnecessary evaluations of false-positive tests. However, if markers are quantified, then a logistic model can be created to achieve both high sensitivity and high specificity. Such a logistic model can also incorporate population variables like gender and age to adjust cut-off levels for test positivity and thereby optimize assay performance in a screening setting (FIG. 1). This Quantitative Logistic to Enhance Accurate Detection (Q-LEAD) Model can be used with any marker class or combination of markers as long as they can be quantified.

A combination of more than one marker was undertaken to achieve the desired sensitivity and specificity for cancer detection. Binary regression methods predicting disease as a

function of diagnostic tests estimate the optimal combination of the tests for classifying a subject as diseased or not. McIntosh and Pepe, *Biometrics* 58: 657-664 (2002). A logistic regression model can assess the relationship between a binary dependent response variable such as presence or absence of disease and one or more independent predictor variables. The independent predictors may be qualitative (e.g., binary) or quantitative (e.g., a continuous endpoint). In the Q-LEAD model, the independent predictors can include such biological markers as K-ras, BMP3 and DNA concentration, and others. Importantly, the model incorporates the demographic variables of gender and age, as we have observed that both age and gender influence molecular marker levels in stool. As average stool marker levels increase with age and male gender, failure to adjust for these variables would yield suboptimal specificity in Men and elderly persons tested. Coefficients are estimated from the sample data for each term in the model. The result of the model is a risk score for each subject. Cutoffs for predicting disease state from this risk score can be determined in order to maximize sensitivity and specificity of the marker combinations for predicting disease as desired. The inclusion of demographic variables allows these cutoffs to be determined as a function of age and gender.

As an application of the Q-LEAD Model, the following was performed to evaluate a quantitative stool DNA assay approach targeting three informative markers for the detection of colorectal neoplasia. Subjects included 34 with colorectal cancer, 20 with adenomas >1 cm, and 26 with normal colonoscopy. Subjects added a DNA stabilization buffer with stool collection, and stools were frozen at -80° C. within 72 hours. From thawed stool aliquots, crude DNA was extracted by standard methods, and target genes were enriched by sequence capture. K-ras mutation score, methylation of BMP3 gene, and concentration of human DNA (245 bp length) were respectively quantified by a digital melt curve assay, real-time methylation-specific PCR and real-time Alu PCR, respectively. Assays were performed blinded. A logistic model, which incorporates three markers and gender, was constructed to analyze discrimination by combined markers.

Age medians were 60 for patients with colorectal cancer, 66 for those with adenomas, and 61 for normal controls; and male/female distributions were 23/11, 9/11, and 10/16, respectively. Detection rates of colorectal neoplasms were determined by individual quantitative markers at specificity cut-offs of 96 percent and by combined markers (Table 1). Discrimination by combined markers was calculated using a qualitative binomial method (each marker considered as positive or negative based on individual 96 percent specificity) and by the Q-LEAD model (sensitivity data shown at overall specificity of 96 percent).

TABLE 1

Specificity and sensitivity of cancer markers.				
Specificity	Sensitivity			
	Cancers	Adenomas	Both	
Individual Markers				
k-ras mutation	96%	42%	32%	38%
BMP3 methylation	96%	45%	32%	40%
DNA concentration	96%	65%	40%	56%

13

TABLE 1-continued

Specificity and sensitivity of cancer markers.				
Specificity	Sensitivity			
	Cancers	Adenomas	Both	
Combined Markers				
Qualitative method	88%	90%	58%	78%
Q-LEAD Model	96%	90%	47%	76%

By quantitative assay and multivariable analysis of an informative marker panel, stool DNA testing can achieve high sensitivity while preserving high specificity for detection of colorectal neoplasia. The particular three-marker combination of mutant K-ras, BMP3 methylation, and human DNA concentration represents a complementary, high-yield panel.

The above data set and additional data were analyzed as follows. A quantitative stool DNA assay approach targeting four informative markers for use in the detection of colorectal cancer and advanced adenoma was evaluated. Subjects comprised 74 patients with colorectal cancer, 27 with an adenoma >1 cm, and 100 with normal colonoscopy. Stools were collected with a stabilization buffer before or >1 week after colonoscopy and were frozen at -80 °C within 24 hours of collection. From thawed stool aliquots, crude DNA was extracted as described above, and target genes were enriched by sequence-specific capture. Human DNA concentration, K-ras and APC mutation scores, and BMP3 methylation were sensitively quantified by real-time Alu PCR, a digital melt curve assay (Zou et al., *Gastroenterology*, "High Detection Rates of Colorectal Neoplasia by Stool DNA Fasting With a Novel Digital Melt Curve Assay," (2008)), and real-time methylation-specific PCR, respectively. Assays were performed blindly. Sensitivities and specificities of single makers and their combinations were analyzed.

Age medians were 61 for patients with colorectal cancer, 67 for those with adenomas, and 59 for normal controls; and, male/female ratios were 52/22, 15/12, and 37/63, respectively. The table displays detection rates of colorectal neoplasms by individual quantitative markers at specificities of 90% and by combined markers at two specificities. (Table 2) Data in this table represent a training set and have not been adjusted for age and gender. Yet, it is clear that the full panel of Alu, K-ras, APC, and BMP3 detected more neoplasms than any individual marker, $p < 0.05$. At 90% specificity, the full panel detects more adenomas >3 cm (90%, 9/10) than <3 cm (47%, 8/17), $p < 0.05$, and more colorectal cancers at

14

stages III-IV (89%, 40/45) than at stages I-II (69%, 20/29) $p < 0.05$. Neoplasm detection rates were not affected by tumor location.

TABLE 2

Specificity and sensitivity of a four marker panel					
Specificity	Sensitivity				
	Colorectal cancers		Adenomas		
	I-II	III-IV	≤3 cm	>3 cm	
Individual Markers					
APC mutation	90%	38%	40%	47%	50%
k-ras mutation	90%	46%	42%	24%	50%
BMP3 methylation	90%	36%	38%	12%	30%
DNA concentration	90%	52%	76%	41%	50%
Combined Markers					
	90%	69%	89%	47%	90%

In conclusion, a quantitative stool DNA assay system that incorporates a stabilization buffer with specimen collection, high analytical sensitivity, and a panel of broadly informative markers can achieve high detection rates of both colorectal cancers and advanced adenoma.

Example 2—SITE Model and Matrix Marker Panel

A statistical model (Site of Tumor Estimate (SITE)) can be used to predict tumor site (e.g., anatomical location or tissue of origin) based on inputs from sequencing data (such as by specific nucleic acid or combination of nucleic acids mutated, specific mutational location on a nucleic acid, and nature of mutation insertion, deletion, transition, or transversion) or by any combination thereof) and/or data from polypeptide or other types of markers.

A matrix marker panel was developed to include eight cancer-related genes: K-ras, p53, APC, p16, EGFR, CTNNB1, BRAF, and SMAD4. The mutation frequencies of these genes were tabulated against the six major aerodigestive cancers based on literature or public database reviews and on actual sequencing observations (Table 3). Literature frequencies were derived from the COSMIC somatic mutation database, review articles, and texts.

TABLE 3

Matrix Panel of Markers by Tumor Site										
AD Cancer Site	N	p16	p53	K-ras	APC	SMAD4	EGFR	CTNNB1	BRAF	Total Unique
Literature										
Colorectal		<5%	50-75%	40%	85%	14%	NA	13%	20%	
Pancreatic		85-100%	50-60%	80-90%	10-40%	30%	NA	3-8%	<5%	
Lung		15-25%	25-75%	20-40%	5%	7%	30%	6%	<5%	
Bile Duct		15-60%	30-60%	40%	30-40%	17%	NA	1%	14%	
Gastric		5-30%	20-50%	10%	20-60%	NA	<1%	30-50%	<5%	
Esophageal		5-90%	40-90%	5-12%	5-60%	NA	NA	<1%	<5%	

TABLE 3-continued

Matrix Panel of Markers by Tumor Site										
AD Cancer Site	N	p16	p53	K-ras	APC	SMAD4	EGFR	CTNNB1	BRAF	Total Unique
Actual (non-dbsNP)										
Colorectal	57	5%	47%	26%	75%	25%	12%	2%	30%	98%
Pancreatic	24	29%	17%	62%	54%	8%	8%	4%	8%	83%
Lung	56	9%	57%	9%	16%	14%	14%	2%	4%	77%
Bile Duct	15	13%	27%	13%	20%	13%	0	0	7%	67%
Gastric	23	17%	22%	4%	35%	17%	4%	4%	0	65%
Esophageal	24	4%	46%	4%	33%	17%	4%	0	4%	79%

Some of the frequencies include other genetic alterations than simply single base changes and small insertions/deletions such as methylation events, large homozygous deletions, and copy number changes. Such alterations would not be reflected in the actual frequency table. Actual frequencies were derived by sequencing coding and flanking gene regions from 245 patient tissue samples reflecting the spectrum of aero-digestive cancers. Only non-synonymous and splice site alterations were tabulated. When specific mutational hot-spot sites were able to be identified for particular genes, only those sites were analyzed.

The matrix panel includes markers that are present to variable extent across these tumors so that their aggregate use achieves high overall sensitivity and allows prediction of tumor site using the SITE Model. 70% of tumors harbored one or more mutations from the eight gene panel. Some gene mutations, like those associated with p16, are common in tumors above the colon but are rare for those in the colon. Mutant K-ras is frequent with colorectal and pancreatic cancers but infrequent in the other cancers. Mutations in EGFR clustered with lung and colorectal tumors and mutations in SMAD4 clustered with stomach and colorectal tumors. Genes such as p53, are commonly mutated across many different types of cancers, but specific mutational locations or types of mutations within p53 and other genes differ between tumor site (e.g., Greenman et al., *Nature*, 446(7132):153-8 (2007); Soussi and Lozano, *Biochem. Biophys. Res. Commun.*, 331(3):834-42 (2005); Stephens et al. *Nat. Genet.*, 37(6):590-2 (2005); Sjoblom et al., *Science*, 314(5797):268-74 (2006); Wood et al., *Science*, 318(5853):1108-13 (2007); and Davies, *Cancer Res.*, 65(17):7591-5 (2005)) and can be factored in to the SITE Model to predict tumor site. Single base substitutions were the most common type of mutation throughout the panel and those that predicted colorectal tumors included C-G and A-T transversions (FIG. 12). Other tumor sites had similarly unique base change profiles. (Table 4). Insertion/deletions mutations were most common with colorectal tumors, particularly adenomas.

TABLE 4

Specific Base Change Fractions in AD Tumors						
Tumor	C > T (G > A)	T > C (A > G)	G > C (C > G)	A > T (T > A)	G > T (C > A)	T > G (A > C)
Head and Neck	0.38	0.12		0.38		0.12
Esophageal	0.8	0.07			0.13	
Lung	0.3	0.11	0.02	0.13	0.34	0.09
Stomach	0.5	0.25		0.17	0.08	
Pancreas	0.41	0.15	0.04	0.07	0.33	
Bile Duct	0.5		0.12	0.25	0.12	

TABLE 4-continued

Specific Base Change Fractions in AD Tumors						
Tumor	C > T (G > A)	T > C (A > G)	G > C (C > G)	A > T (T > A)	G > T (C > A)	T > G (A > C)
CRA	0.34	0.05	0.11	0.11	0.34	0.03
CRC	0.4	0.05	0.02	0.24	0.26	0.03

Polypeptide markers found in stool, such as by proteomic approaches, can also be used to detect aero-digestive neoplasms and predict tumor site. The following was performed to identify and explore candidate polypeptide markers in stool for the discriminate detection of pancreatic cancer. Subjects included 16 cases with pancreatic cancer, 10 disease controls (colorectal cancer), and 24 healthy controls. Whole stools were collected and frozen promptly in aliquots at -80° C. Thawed aliquots were centrifuged, and the aqueous supernatant from each was analyzed. Polypeptides were separated by 1-D electrophoresis, excised from gels, and digested for mass spectrometric analysis using an LTQ-Orbitrap. Data outputs were searched using Mascot, Sequest, and X! Tandem programs against an updated Swissprot database that included all cataloged species. Unique peptide counts and ratio calculations were performed using Scaffold software.

Median age for pancreatic cancer cases was 67, for colorectal cancer controls 63, and for healthy controls 62; and male/female distributions were 9/7, 6/4, and 9/15, respectively. Using shotgun-proteomic techniques on stools, two pancreatic enzymes (carboxypeptidases B and A2) were conspicuous, as unique spectral counts of the former were commonly elevated with pancreatic cancer and of the latter commonly decreased. Considered together as the ratio of carboxypeptidase B/carboxypeptidase A2, pancreatic cancer cases were almost completely separated from colorectal cancer and healthy control groups. Median ratios were 0.9, 0.2, and 0.3, respectively. At a specificity cut-off for the carboxypeptidase B/A2 ratio at 100% (i.e., ratios from normal control and colorectal cancer stools all below cut-off), sensitivity for pancreatic cancer was 86 percent (FIG. 2). Only two pancreatic cancers were misclassified.

These results demonstrate that a stool assay of polypeptide markers can be a feasible non-invasive approach to the detection pancreatic cancer. These results also demonstrate that multivariable analysis of specific polypeptide ratios can be used.

In addition, polypeptide markers unique to colorectal neoplasms were identified (Table 5). For example, serotransferrin was found in stools from patients with colorectal cancer but not in those with pancreatic cancer. These markers when considered as part of a matrix panel contribute both

to overall sensitivity for tumor detection and help discriminate colorectal from pancreatic cancer.

TABLE 5

Positive Stool Findings.		
	Carboxypeptidase B/A2*	Serotransferrin
Colorectal Cancer	0	60%
Pancreatic Cancer	86%	0
Normal controls	0	0

*ratio >0.75 considered positive

Another polypeptide in stool that is pancreatic cancer specific is elastase 3A. Methods and Results demonstrating this are as follows:

Stool Preparation

Samples were collected in phosphate buffered saline and either dropped off in clinic or mailed in collection tub. Samples were homogenized and frozen within 72 hours after receipt. Frozen stools were diluted 1:3 w:v in PBS (Roche, Cat#1666789). Diluted stools were stomached in a filter bag (Brinkman, BA6041/STR 177x305 mm) for 60 seconds on control setting and spun at 10,000 rpms for 30 minutes. Following an additional 10 minute spin at 14,000 rpm, the supernatant was filtered through a 0.45- μ m syringe filter and analyzed. Total protein present in stool was quantitated using a Bradford Protein Assay kit (Pierce).

1-Dimensional Electrophoresis

Stool supernatants were diluted 1:1 in Leammli-BME buffer and run on a 10.5-14% gradient gel. Vertical slices were cut from 250 kDa to 15 kDa and in-gel digested using methods described elsewhere (e.g., Wilm et al., *Nature*, 379:466-469 (1996)). Bands were destained, dehydrated, digested in trypsin, extracted, and for MS analysis.

Mass Spectrometry

Lyophilized samples were reconstituted and injected with a flow of 500 nL/min and a 75 minute gradient from 5-90% 98% acetonitrile. MS was performed in data dependent mode to switch automatically between MS and MS² acquisition on the three most abundant ions. Survey scans were acquired with resolution $r=60,000$ at 40 m/z using FWHM with a target accumulation of 10^6 counts. An isolation width of 2.5 m/z was applied. Exclusion mass width was 0.6 m/z on low end and 1.5 m/z on high end. All acquisition and method development was performed using Xcaliher version 2.0.

Database Searching All ms/ms

Samples were analyzed using Mascot (Matrix Science, London, UK; version 2.1.03), Sequest (ThermoFinnigan, San Jose, Calif.; version 27, rev. 12) and X! Tandem (World Wide Web at "thegpm.org" version 2006.09.15.3). Mascot and X! Tandem were searched with a fragment ion mass tolerance of 0.80 Da and a parent ion tolerance of 10.0 PPM. Sequest was searched with a fragment ion mass tolerance of 1.00 Da. Nitration of tyrosine was specified in Mascot as a variable modification.

Criteria for Polypeptide Identification

Scaffold (version Scaffold-01_06_06, Proteome Software Inc., Portland, Oreg.) was used to validate MS/MS based polypeptide identifications. Peptide identifications were accepted if they could be established at greater than 95.0% probability as specified by the Peptide Prophet algorithm (Keller et al, *Anal. Chem.*, 74(20):5383-92 (2002)). Polypeptide identifications were accepted if they could be established at greater than 99.0 percent probability and contained at least two identified peptides. Polypeptide probabilities

were assigned by the Protein Prophet algorithm (Nesvizhskii, *Anal. Chem.*, 75(17):4646-58 (2003)). Polypeptides that contained similar peptides and could not be differentiated based on MS/MS analysis alone were grouped to satisfy the principles of parsimony.

Specific to Elastase SA Ratio Determination

Ratios of elastase 3A were determined using spectral counts for each polypeptide. Ratios were determined by dividing the number of unique peptides of elastase 3A (determined using a composite ID from database search modules Mascot, XTandem, and Seaquest and compiled in Scaffold) by the number of unique peptides from another polypeptide such as pancreatic alpha-amylase.

Results

The concentration of a specific pancreatic enzyme, elastase 3A, was consistently found to be elevated in the fecal supernatant of patients with pancreatic cancer as compared to normal controls or patients with non-pancreatic cancer (FIG. 3). These findings indicate that fecal concentration of elastase 3A is an accurate marker for pancreatic cancer. In addition, the ratio of elastase 3A against other pancreatic enzymes (or other stable fecal polypeptides) was found to be especially discriminant for pancreatic cancer and obviates the need to determine absolute elastase 3A concentrations (FIG. 4). While mass spectrometry was used to make these observations, elastase 3A levels and ratios including elastase 3A can be measured using other methods as well.

Example 3—Digital Melt Curve Assay for Scanning Mutations

A sensitive, rapid, and affordable method for scanning mutations in bodily fluids at high-throughput was developed. A melt curve assay is a post-PCR technique that can be used to scan for mutations in PCR amplicons. Mutations in PCR products can be detected by changes in the shape of the melting curve (heterozygote from mutant sample) compared to a reference sample (homozygote from wild-type sample) (FIG. 5). Melt curve assay can scan all mutations in a DNA fragment <400 bp in less than 10 minutes, rather than individually targeting single mutations. Regular melt curve assays can detect mutations down to a limit of 5% mutant: wild-type and, thus, are not sensitive enough to detect mutations in many biological samples. For instance, in stool, an analytical sensitivity of 1% or less is required in order to detect precancerous polyps or small early stage cancers. Importantly, a quantitative score can be given to density of target mutations (FIG. 6).

Digital PCR can augment the sensitivity of PCR to detect low abundance mutations. Gene copies can be diluted and distributed into 96 wells of a plate to increase the percentage of mutant copy to wild-type copies in certain wells. For example, if a stool DNA sample contains only 1% of mutant BRAF copies compared to wild-type copies, distributing 300 copies of BRAF gene into a 96-well plate can lead to three wells with an average mutant ratio of 33 percent (1:3). After PCR amplification, these three wells with mutant copies can be detected by sequencing or other approaches. Since digital PCR requires PCR on a whole 96-well plate and 96 sequencings (or other approaches) for each target, it can be slow and costly.

The concept of digital melt curve assay is to combine the scanning ability and speed of high resolution melt curve assay with the sensitivity of digital PCR. Miniaturizing and

automating this technology dramatically lowers per assay cost and achieves high-throughput necessary for population screening.

The following procedure was used to perform a digital melt curve assay. To prepare a DNA sample, gene target fragments (e.g., BRAF, K-ras, APC, p16, etc.) were captured from stool DNA using a sequence-specific capture method and were quantified with real-time PCR. About 200 to 2000 gene copies were mixed in tube with all PCR reagents. An average of 2 to 20 copies (variable) were distributed to each well of a 96-well plate. PCR amplification was performed using specific primers on the plate (e.g., one target per plate). Final concentrations of PCR mastermix for Digital Melt

Curve assays in a 96-well plate (500 μ L dispersed to 96 wells with each well containing 5 μ L) were as follows: 2 \times pfx amplification buffer (Invitrogen), 0.3 mM each dNTP, 200 nM forward primer, 200 nM reverse primer, 1 mM MgSO₄, 0.02 unit/ μ L Platinum® pfx polymerase (Invitrogen), and 0.1 unit/ μ L LcGreen+dye (Idaho Tech). A high resolution melt curve assay was used to identify the wells with mutant copies. Sequencing was optionally performed to confirm 1 to 2 representative wells.

In some cases, emulsion PCR can be used in place of digital PCR. In such cases, each lipid drop can become a tiny PCR reactor of one single molecule of gene.

TABLE 6

Sequence Specific Capture Probes and DNA Primer					
Gene	Target Region	Capture Probe/ Primer	Oligo Sequence (5'→3')	SEQ ID No.	
KRAS	Condons 12/13	Probe	GTGGACGAATATGATCCAACAATAGAGGTAAATCTTG	1	
		Primer 1	AGGCCTGCTGAAAATGACTG	2	
	Condons 12/13	Primer 2	TTGTTGGATCATATTCGTCCAC	3	
			TAAGCCTGCTGAAAATGAC	4	
		Primer 3	ATCAAAGAATGGTCTGCAC	5	
	Condons 12/13	Primer 3	CGTCTGCAGTCAACTGGAATTT	6	
			TGTATCGTCAAGGCACCTTGC	7	
	Condons 12/13	Primer 4	CTTAAGCGTCGATGGAGGAG	8	
TTGTTGGATCATATTCGTCCAC			3		
BRAF	V600E	Probe	CCAGACAACCTGTTCAAACCTGATGGGACCCACTCCATC	9	
		Primer	CCACAAAATGGATCCAGACA	10	
	V600E		TGCTTGCTCTGATAGGAAAATG	11	
APC	MCR	Probe 1	CAGATAGCCCTGGACAAACCATGCCACCAAGCAGAAG	12	
		Probe 2	TTCCAGCAGTGTACAGCACCCCTAGAACCAAATCCAG	13	
		Probe 3	ATGACAATGGGAATGAAACAGAATCAGAGCAGCCTAAAG	14	
	Condons 1286-1346	Primer 1	TTCATATCATCTTTTGTTCATCAGC	15	
			CGCTCCTGAAGAAAATTCAA	16	
	Condons 1346-1367	Primer 2	TGCAGGGTTCTAGTTTATCTTCA	17	
			CTGGCAATCGAACGACTCTC	18	
	Condons 1394-1480	Primer 3	CAGGAGACCCCACTCATGTT	19	
			TGGCAAAATGTAATAAAGTATCAGC	20	
	Condons 1450-1489	Primer 4	CATGCCACCAAGCAGAAGTA	21	
			CACTCAGGCTGGATGAACAA	22	
	Condon 1554	Primer 5	GAGCCTCGATGAGCCATTTA	23	
			TCAATATCATCATCTGAATCATC	24	
	102457delC	Primer 6	GTGAACCATGCAGTGGAATG	25	
			ACTTCTCGCTTGGTTTGGAGC	26	
	102457delC	Primer 7	CAGGAGACCCCACTCATGTT	19	
CATGGTTTGTCCAGGGCTAT			27		
102457delC	Primer 8	GTGAACCATGCAGTGGAATG	25		
		AGCATCTGGAAGAACCCTGGA	28		
TP53	Exon 4	Probe	AAGACCCAGGTCCAGATGAAGCTCCAGAATGCCAGA	29	
		Primer	CCCTTCCCAGAAAACCTACC	30	
	Exon 5	Probe	GCCAGGCATTGAAGTCTCAT	31	
			CATGGCCATCTACAAGCAGTACAGCACATGACGGAG	32	
		Primer	CACTTGTGCCCTGACTTTCA	33	
	Exon 6	Probe	AACCAGCCCTGTCGTCTCT	34	
			AGTGAAGGAAATTTGCGTGTGGAGTATTTGGATGAC	35	
	Exon 7	Primer	CAGGCCTCTGATTCCTCACT	36	
			CTTAACCCCTCCTCCCAGAG	37	
	Exon 8	Probe	ATGTGTAACAGTTCCCTGCATGGGCGGCATGAACCGGA	38	
			C'TTGGCCTGTGTTATCTCC	39	
	Exon 8	Primer	GGGTCAGAGGCAAGCAGA	40	
			CGCACAGAGGAAGAGAATCTCCGCAAGAAAGGGGAGC	41	
	Exon 8	Primer	GGGAGTAGATGGAGCCTGGT	42	
			GCTTCTTGTCTGCTTGCTT	43	

Example 4—Sensitive Detection of Mutations Using a Digital Melt Curve Assay

The following was performed to develop a quantitative method for scanning gene mutations and to evaluate the sensitivity of the quantitative method for detecting target

mutations in stool. A digital melt curve assay was designed by combining digital PCR to a modified melt curve assay. Target genes in low concentration were PCR amplified with a saturated DNA dye, LcGreen+, in a 96-well plate. Each well contained a small number of gene copies, which allowed high mutation/wild-type ratios in some wells that

were then detected by melt curve scanning using a Light-Scanner. Mutations were scored based on the number of wells containing mutant copies in a 96-well plate. To test sensitivity, mutant genes were spiked into a wild-type pool at 0.1, 0.5, 1, 5, and 10% dilutions, and analyzed using digital melt curve assay with 250-1000 gene copies per 96-well plate. This method was then applied in the stool detection of APC, p53, K-ras, and BRAF mutations from 48 patients known to have mutations in one of these genes in matched tumor tissue. Subjects included 9 patients with pancreatic cancer, 31 with colorectal cancer, and 8 with colorectal adenoma >1 cm. All mutations detected by digital melt curve were further confirmed by Sanger sequencing.

The digital melt curve assay detected as few as 0.1% mutant copies for amplicons <350 bp using one 96-well plate (FIG. 7), compared to the detection limit for regular melt curve of ≥ 5 percent. Each mutation scanning took 8-10 minutes with this manual approach. Mutations of APC, p53, K-ras, and BRAF genes were all successfully scanned with digital melt curve in quantitative fashion. Tissue-confirmed mutations were detected from matched stools in 88 percent (42/48) of patients with gastrointestinal neoplasms, including 89 percent with pancreatic cancer, 90 percent with colorectal cancer, and 75 percent with colorectal adenoma >1 cm.

These results demonstrate that a digital melt curve assay can be a highly sensitive approach for detecting mutations in stool, and that it has potential for diagnostic application with both upper and lower gastrointestinal neoplasms.

Example 5—Using a Digital Melt Curve Assay to Detect Adenomas

Archived stools were used to evaluate a digital melt curve assay of DNA markers for detection of advanced adenomas and to compare the accuracy of the digital melt curve assay with that of occult blood testing and a commercial DNA marker assay method (EXACT Sciences). Average risk subjects collected stools without a preservative buffer and mailed them to central processing laboratories for banking and blinded stool testing by Hemocult, HemocultSENSA, and DNA marker assay. All subjects underwent a colonoscopy, and tissue from advanced adenomas was archived. Archival stools were selected from the 27 patients with a colorectal adenoma >1 cm found to harbor mutant K-ras on tissue analyses and from the first 25 age and gender matched subjects with normal colonoscopy. Standard methods were used to extract crude DNA from fecal aliquots, and K-ras gene was enriched by sequence capture. Mutations in the K-ras gene were quantified by a digital melt curve assay based on the number of wells containing mutant gene copies in a 96-well plate and confirmed by sequencing.

Median age with adenomas was 67 and controls 71; and males/females were 12/15 and 13/14, respectively. Median adenoma size was 1.5 cm (range 1-3 cm). Based on a cut-off of >3 wells with mutant K-ras, the digital melt curve assay yielded an overall sensitivity of 59 percent for adenomas with a specificity of 92 percent; sensitivity for adenomas >2 cm was 80 percent (8/10) and for those <2 cm was 47 percent (8/17), $p=0.1$. In these same stools, overall adenoma detection rates were 7 percent by Hemocult, 15 percent by HemocultSENSA, and 26 percent by the EXACT Sciences K-ras assay ($p<0.05$ for each vs. digital melt curve) (FIG. 8). Respective specificities were 92 percent, 92 percent, and 100 percent.

These results demonstrate that an analytically-sensitive digital melt curve assay method can be used to detect a

majority of advanced colorectal adenomas and improve yield over current stool test approaches.

Example 6—Short DNA as a Cancer Marker

Free human DNA is present in all human stools and arises from cells shed (exfoliated) from the normal surface (mucosa) of the aero-digestive tract (mouth/throat, lungs, and all digestive organs) and from tumors or other lesions that may be present. It has been generally accepted that “long DNA” in stool reflects that presence of colorectal and other aero-digestive tumors, in that cells exfoliated from cancers do not undergo typical cell death (apoptosis) which would shorten DNA. Specifically, because DNA from apoptotic cells would be broken down to fragment lengths shorter than 100 bp, long DNA was defined as being longer than 100 bp. Indeed, levels of long DNA were elevated in stools from patients with colorectal and other cancers as compared to those from healthy controls (Zou et al., *Cancer Epidemiol. Biomarkers Prev.* 15: 115 (2006); Ahlquist et al., *Gastroenterology*, 119:1219 (2000); and Boynton et al., *Clin. Chem.*, 49:1058 (2003)). As such, long DNA in stool can serve as a marker for colorectal and other tumors.

“Short DNA” (i.e., <100 bp in length), however, was found to be as or more discriminant than long DNA as a tumor marker in stool for detection of both colorectal (FIG. 9) and pancreatic (FIG. 10) neoplasia.

Briefly, methods and materials similar to those described elsewhere were used to detect short DNA present in stool samples (Zou et al., *Cancer Epidemiol. Biomarkers Prev.*, 15(6): 1115 (2006)). Total DNA was extracted by isopropanol precipitation from 19 blinded stool samples. 9 pancreatic adenocarcinoma, and 10 age/gender matched normals. The DNA pellets were taken up in 8 mL of 10-fold diluted TE, pH 8. The Alu sequence consists of conserved regions and variable regions. In the putative consensus Alu sequence, the conserved regions are the 25-bp span between nucleotide positions 23 and 47 and the 16-bp span between nucleotide positions 245 and 260. Although primers can be designed in any part of the Alu sequences, for more effectively amplifying Alu sequences, the PCR primers are preferably completely or partially (at least the 3'-regions of the primers) located in the conserved regions. Primers specific for the human Alu sequences were used to amplify fragments of differing lengths inside Alu repeats. The sequences were as follows:

Ampli- con size	Primer Sequences
245 bp	Forward Primer: 5'-ACGCCTGTAATCCCAGCACTT-3' (SEQ ID NO: 44) Reverse Primer: 5'-TCGCCAGGCTGGAGTGCA-3' (SEQ ID NO: 45)
130 bp	Forward Primer: 5'-TGGTGAAACCCCGTCTCTAC-3' (SEQ ID NO: 46) Reverse Primer: 5'-CTCACTGCAACCTCCACCTC-3' (SEQ ID NO: 47)
45 bp	Forward Primer: 5'-TGGTGAAACCCCGTCTCTAC-3' (SEQ ID NO: 46) Reverse Primer: 5'-CGCccGGCTAATTTTTGTAT-3' (SEQ ID No: 48)

Stool DNA was diluted 1:5 with Ix Tris-EDTA buffer (pH 7.5) for PCR amplification. Tris-EDTA buffer-diluted stool DNA (14) was amplified in a total volume of 25 μ L

containing Ix iQ SYBR Green Supermix (Bio-Rad, Hercules, Calif.), 200 nmol/L each primer under the following conditions: 95° C. for 3 minutes followed by 40 cycles of 95° C., 60° C., and 72° C. for 30 seconds each. A standard curve was created for each plate by amplifying 10-fold serially diluted human genomic DNA samples (Novagen, Madison, Wis.). Melting curve analysis was made after each PCR to guarantee that only one product was amplified for all samples.

Amplification was carried out in 96-well plates in an iCycler (Bio-Rad). Each plate consisted of stool DNA samples and multiple positive and negative controls. Each assay was done in duplicate.

The following was performed to compare long DNA (245 bp) and short (45 bp) human DNA in stool for detection of upper and lower GI neoplasms, and to assess the effect of GI tumor site on human DNA levels in stool. Subjects included 33 patients with colorectal cancer, 20 with colorectal adenomas >1 cm, 13 with pancreatic cancer, and 33 colonoscopically-normal controls. Subjects added a preservative buffer to stools at time of collection to prevent post-defecation bacterial metabolism of DNA, and stools were frozen within 8 hours at -80° C. Using a validated quantitative assay for human DNA (Zou et al. *Epidemiol. Biomarkers Prev.*, 15:1115 (2006)), 245 bp and 45 bp Alu sequences were amplified from all stools in blinded fashion. Sensitivities for long and short DNA were based on 97 percent specificity cut-offs.

Age medians were 60, 66, 69, and 62 for colorectal cancer, colorectal adenoma, pancreatic cancer, and control groups, respectively, and male/female distributions were 22/11, 9/11, 9/4, 11/21, respectively. In stools from neoplasm and control groups, amplification products were quantitatively greater for short DNA versus long DNA. Respective sensitivities by long and short DNA were 66 percent and 62 percent with the 29 distal colorectal neoplasms, 46 percent and 46 percent with the 24 proximal colorectal neoplasms, and 15 percent and 31 percent (p=0.16) with the 13 pancreatic cancers. By Wilcoxon Rank-Stun test, effect of neoplasm site on detection rates was significant for both long DNA (p=0.004) and short DNA (p=0.02). Among colorectal neoplasms, respective sensitivities by long and short DNA were 48 percent and 52 percent with lesions <3 cm, 63 percent and 63 percent with those >3 cm, 64 percent and 61 percent with cancers, and 35 percent and 45 percent with adenomas.

These results demonstrate that short and long DNA can be comparably sensitive for stool detection of GI neoplasms. However, detection rates vary with tumor site, being greatest with the most distal lesions and lowest with the most proximal ones. These results were consistent with substantial luminal degradation of DNA exfoliated from more proximal GI neoplasms.

It was also demonstrated that mutant gene markers in stool can be detected to a greater extent if amplicon size is

less than 70 bp, consistent with luminal degradation. Thus, short DNA can serve as a marker per se and as the target size for mutation detection.

Example 7—Use of Fecal Methylated BMP3 as a Neoplasia Marker

Stools from patients with colorectal tumors were found to contain significantly elevated amounts of methylated BMP3 gene copies, but those from normal individuals were found to contain none or only trace amounts. When fecal methylated BMP3 was assayed with an appropriate amplification method, colorectal cancers and premalignant adenomas were specifically detected (FIG. 11). Fecal methylated BMP3 detected a higher percentage of proximal colon tumors than distal tumors, so it can be combined with markers for distal colorectal tumors to create complementary marker panels. Fecal methylated BMP3 was very specific with few false-positive reactions.

Similar results can be obtained using other genes and methods such as those described elsewhere (Zou et al., *Cancer Epidemiol. Biomarkers Prev.*, 16(12):2686 (2007)).

Example 8—Detecting Aero-Digestive Cancers by Stool DNA Testing

Tissue samples from patients with confirmed acre-digestive tumors were extracted and sequenced to assess the presence or absence of somatic gene alterations. Germline DNA from the same patients were used as controls. Once an alteration was confirmed, a matched stool sample was tested for that alteration. Two separate methods were utilized to detect the mutation in stool: Allele specific PCR and digital melt curve analysis. For both methods, we focused on amplifying the shortest fragments possible (>100 bp) that have been shown to contain higher levels of the mutant sequence.

Digital Melt Curve (DMC)

We studied 138 patients (69 cases with a GI neoplasm and 69 age/sex-matched asymptomatic controls with normal colonoscopy) by first, identifying a mutation in neoplasm tissue, and then determining if that specific mutation could be detected in stool from that individual. Stools were collected with a stability buffer and frozen at -80° C. until assayed.

Genes commonly mutated in GI neoplasms (TP53, KRAS, APC, CDH1, CTNNB1, BRAF, SMAD4, and P16) were sequenced from DNA extracted from tumor tissue, to identify a target mutation for each case. Target genes were isolated by hybrid capture (Table 7) and the tissue-confirmed somatic mutations were assayed in stool by the digital melt curve method, as described in Example 1. Mutations detected in stool were confirmed by sequencing. Assays were performed blinded.

TABLE 7

Sequence Specific Capture Probes and Primers for AD Cancer Mutation Detection		
MUTATION IN TISSUE	CAPTURE PROBE	SEQ ID No.
12487C > CT: 167Q > Q/X	ATGGCCATCTACAAGCAGTCATAGCACATGACGGAGGTTGT	49
102447_102450het_delTTGGT	AGAGTGAACCATGCAGTGGAAAAGTGGCATTATAAGCCC	50
12410G > GA, 141C > C/Y	TTTGCCAACTGGCCAAGACCTACCCCTGTGCAGCTGTG	51
102678het_delA	CAGATGCTGATACTTTATTACTTTTGCCACGGAAAGTACT	52
102594_102598het_delAGAGA	AAAGCACCTACTGCTGAAAGAGAGTGGACCTAAGCAAG	53
102644_102646het_insG	ATGCTGCAGTTCCAGAGGGTCCAGGTTCTTCCAGATGC	54

TABLE 7-continued

Sequence Specific Capture Probes and Primers for AD Cancer Mutation Detection		
102594_102595het_delAG	TAAAGCACCTACTGCTGAAAAGAGAGAGTGGACCTAAGCAAG	55
102106het_delT	CACAGGAAGCAGATTCTGCAATACCCCTGCAAATAGCA	56
102442het_delT	TTCAGAGTGAACCATGCAGGGAATGGTAAGTGGCATTAT	57
apc 102494C > CT: 1429Q > Q/X	TCCAGATAGCCCTGGATAAACCATGCCACCAAG	58
apc 102557C > CT: 1450R > RA	CTCAAACAGCTCAAACCAAGTGAGAAGTACCTAAAAATAAA	59
apc 102140het_delA	AGCAGAAATAAAAGAAAAGTTGGAAGTACCTAAAAATAAA	60
apc 102494C > CT: 1429Q > Q/X	TCCAGATAGCCCTGGATAAACCATGCCACCAAG	58
apc 102554het_delA	CTCAAACAGCTCAAACAGCGAGAAGTACCTAAA	61
tp53 E5 12647A > AG: 193H > H/R	TCTGGCCCTCCTCAGCGTCTTATCCGAGTGGAAAG	62
tp53 E5 12742G > GA	CCTATGAGCCGCTGAGATCTGGTTTGAACCTGGG	63
tp53 E5 12706C > CT: 213R > R/X	ATGACAGAAACACTTTTTGACATAGTGTGGTGGTG	64
tp53 E4 12712A > AG: 215S > S/G	GAAACACTTTTTCGACATGGTGTGGTGGTGCCCTAT	65
tp53 E4 12388T > TC: 134F > F/L	CTGCCCTCAACAAGATGCTTTGCCAACTGGCCAAG	66
tp53 E3 11606G > GA: 125T > T/T	AAGTCTGTGACTTGCACAGTTCAGTTGCCCTGAGGG	67
tp53 E6 13379C > CT: 248R > R/W	GCATGGGCGGCATGAACCTGGAGGCCATCCTCACC	68
tp53 12E3 11326A > AC (splice site)	TCTTTTCACCCATCTACCGTCCCTCCCTTGCCGTCCC	69
tp53 E6 13412G > GT: 259D > D/Y	CCATCATCACACTGGAATACTCCAGGTCCAGGAGCC	70
tp53 E4 12449G > GT: 154G > GN	ACCCCGCCCGTCCACCCGCGTCC	71
tp53 E7 13872G > GT, 298E > E/X	GGAACAGCTTTGAGGTGTGTGTGTGTGCCTGTCTC	72
APC 102843C > CG: 1545S > S/X	TCAGAGCAGCCTAAAGAATGAAATGAAAACCAAGAGAAA	73
tp53 E4 12392G > GT, 135C > C/F	CCTCAACAAGATGTTTTTCCAACTGGCCAAGACCT	74
APC 102557C > CT: 1450R > R/X	CTCAAACAGCTCAAACCAAGTGAGAAGTACCTAAAAATAAA	59
tp53 E7 13819G > T: 280R > I	CCTGTCTGGGATAGACCCGCGCAC	75
tp53 13E4 11326A > AC (splice site)	TCTTTTCACCCATCTACCGTCCCTCCCTTGCCGTCCC	69
tp53 E7 13412G > GT: 259D > D/Y	CCATCATCACACTGGAATACTCCAGGTCCAGGAGCC	70
tp53 E5 12449g > GT: 154G > G/V	ACCCCGCCCGTCCACCCGCGTCC	71
tp53 E8 13872G > Gt: 298E > E/X	GGAGCCTCACCCTAGCTGCCCCCAGG	76
tp53 E8 13813C > CG, 278P > P/R	GTGTTTGTGCCTGTCTGTTGGAGAGACCCGCG	77
tp53 E8 13851A > AT, 291K > K/X	GGAAGAGAATCTCCGCTAGAAAGGGGAGCCTCA	78
smad4 E2 19049G > GA, 18118A > A/A	GTTAAATATTGTGAGTATGCATTTGACTTAAAATGTGATAG	79
tp53 E8 13777G > GA: 266G > G/E	TAGTGGTAATCTACTGGAACGGAACAGCTTTGAGGTG	80
tp53 E6 12653T > TC: 195I > I/T	CCTCCTCAGCATCTTACCCGAGTGAAGGAAAT	81
tp53 E7 13379C > CT: 248R > R/W	GCATGGGCGGCATGAACCTGGAGGCCATCCTCACC	68
tp53 E6 12647A > AG: 193H > H/R	TCTGGCCCTCCTCAGCGTCTTATCCGAGTGGAAAG	62
tp53 E6 12712A > AG: 215S > S/G	GAAACACTTTTTCGACATGGTGTGGTGGTGCCCTAT	65
tp53 E8 13872G > GT, 298E > E/X	GGAGCCTCACCCTAGCTGCCCCCAGG	76
tp53 E7 13370G > GA: 245G > G/S	AGTTCCTGCATGGGCAGCATGAACCCGAGGC	82
tp53 E4 11580het_delG	CTGGGCTTCTTGATTCTGGACAGCCAAGTCTGTGA	83
tp53 E5 12524A > AG, 179H > H/R	TGCCCCACCGTGAGCGCTGC	84
>tp53 E6 12661G > GT, 198E > E/X	TCAGCATCTTATCCGAGTGTAAAGAAATTTGCGTGTGGA	85
tp53 E8 13872G > GT, 298E > E/X	GGAGCCTCACCCTAGCTGCCCCCAGG	76
apc 102494C > CT: 1429Q > Q/X	TCCAGATAGCCCTGGATAAACCATGCCACCAAG	58
apc 102557C > CT: 1450R > R/X	CTCAAACAGCTCAAACCAAGTGAGAAGTACCTAAAAATAAA	59
apc 102140het_delA	AGCAGAAATAAAAGAAAAGTTGGAAGTACCTAAAAATAAA	60
apc 102494C > CT: 1429Q > Q/X	TCCAGATAGCCCTGGATAAACCATGCCACCAAG	58
apc 102134G > GT: 1309E > E/X	TGCAAATAGCAGAAATAAAATAAAAGATTGGAAGTGGTCA	86
apc102554het_delA	CTCAAACAGCTCAAACAGCGAGAAGTACCTAAA	61
apc 102852het_insA	CTAAAGAATCAAATGAAAACCAAGAGAAAGAGGCAGAA	87
Kras 5571G > GA: 12G > G/D	GTGGTAGTTGGAGCTGATGGCGTAGGCAAGAGT	88
tp53 E4 12392G > GA: 135C > C/Y	CCTCAACAAGATGTTTTACCAACTGGCCAAGACCT	89
tp53 E5 12655C > CT: 196R > R/X	CCTCCTCAGCATCTTATCTGAGTGGAAAGGAAATTTGC	90
tp53 E6 13350G > GA: 238C > C/Y	TCCACTACAACCTACATGTATAACAGTTCTTGCATGGG	91
tp53 E6 13420G > GA	CACTGGAAGACTCCAGATCAGGAGCCACTTGCC	92
tp53 E5 12712A > AG: 215S > S/G	GAAACACTTTTTCGACATGGTGTGGTGGTGCCCTAT	65
Kras 5571G > GA: 12G > G/D	GTGGTAGTTGGAGCTGATGGCGTAGGCAAGAGT	88
P16 (ink4a) E1 19638A > AT	GGAGAGGGGGAGTGCAGGCAGCGGG	93
P16 (ink4a) E2 23353G > GT: 447D > DY	CCCAACTGCGCCTACCCCGCCACTC	94
P16 (ink4a) E1 19638A > AT	GGAGAGGGGGAGTGCAGGCAGCGGG	93
p16(ink4a) E2 23402het_delT	GCCCGGGAGGGCTCCTGGACACGCTG	95
p16(ink4a) E2 23403C > CA: 484F > F/	GCCCGGGAGGGCTTACTGGACACGCTGGT	96
ctnbn1 25541het_delT	CAATGGGTGCATATCACAGATTCTTTTTTTTTAAATTAAGTAACA	97
cdh 1 E9 76435het_delA	TCTTATCTCAAAGAACAACAAAAAGAGGAATCCTTTAG	98
cdh1 E1 743_744het_insAGCCCTGCGCCCA	GCGCCAGCCCTGCGCCCATTCCTC	99
cdh 1 E13 8685386854het_insA	AAGTAAGTCCAGCTGGCAAAGTACTCAGCCTTTGACTT	100
cdh 1 E14 91472C > CT: 751N > N/N	AGGATGACACCCGGGACAATGTTTATTACTATGATGAAG	101
cdh 1 E 15	TTTTTCTCAAAGGACTGACGCTCGGCCTGAAGT	102
92868_92896het delTTGACTTGA		
GCCAGCTGCACAGGGGCCTG		
cdh 1 E4 71669*het_delA	CAAGCAGAATTGCTCACTTTCCCAACTCCTCTCC	103
cdh 1 E7 74926G > GA: 289A > A/T	GGTCACAGCCACAGACACGGACGATGATGTGAA	104
cdh 1 E1 736_742het_delTGGCCTC	AGCCCTGCGCCCTTCTCTCCCG	105
p16(ink4a) E1 19638A > AT	GGAGAGGGGGAGTGCAGGCAGCGGG	93
tp53 E4 12365A > AG: 126Y > Y/C	TTCCTCTTCTACAGTGTCCCTGCCCCTCAAC	106
tp53 E4 12548G > GA	TGCTCAGATAGCGATGATGAGCAGCTGGGGCTG	107
p16(ink4a) E1 19810T > TG: 491I > I/S	GGTCGGAGGGCCGAGCCAGGTGGGTAGA	108
tp53 E7 13757G > GA	GCTTCTCTTTCTATCCTAAGTAGTGGTAATCTACTGG	109
tp53 E7 13815G > GC: 279G > G/R	TTGTGCCTGTCTCCTCGGAGAGACCCGCG	110
tp53 E7 13816G > GA: 279G > G/E	TGTGCCTGTCTGAGAGAGACCCGCGC	111

TABLE 7-continued

Sequence Specific Capture Probes and Primers for AD Cancer Mutation Detection		
MUTATION IN TISSUE	SENSE PRIMER 1 (5' TO 3')	SEQ ID No.
tp53 E5 12365A > AC, 126Y > Y/S	TTCCTCTTCTACAGTCCTCCCTGCCCTCAAC	112
tp53 E5 12491A > AT, 168H > HL	TCTACAAGCAGTCACAGTCATGACGGAGGTTGTGGA	113
		113
		113
kras 5570G > GC, 12G > G/R	GTGGTAGTTGGAGCTGATGGCGTAGGCAAGAGT	88
tp53 E7 13370G > GA, 245G > G/S	AGTTCCTGCATGGGCAGCATGAACCGGAGGC	82
apc 102864_102865het_delAG	AAATGAAAACCAAGAGAAAGGCAGAAAAAATATTGATTC	114
tp53 E5 12386T > TC, 133M > M/T	CCTGCCCTCAACAAGACGTTTGGCCAACTGGCC	115
cdh1 E15 93059G > GA	GCTCATCTCTAAGCTCAGGAAGAGTTGTGTCAAAAATGAGA	116
tp53 E8 13798G > GA: 273R > R/H	CGGAACAGCTTTGAGGTGCATGTTTGTGCCTGTCTGGG	117
p53 E6, 12698_12701het_delAC (1 or 2 AC repeats)	TGGAAGGAAATTTGCGTGTGGAGTATTTGGATGACAG	118
P53 E8, 13824C > CT, 282R > R/W	TGTCCTGGGAGAGACTGGCGCACAGAGGAAGAGAAT	119
APC 102151G > GA, 1314R > R/R	AAGAAAAGATTGGAAC TAGATCAGCTGAAGATCCTGTG	120
P53 ES 12457 G > G/T	CGCCCGGCACCCCGCTTCCGCGCCATGGCCA	121
p53 E813812C > CG, 278P > P/A	GTGTTGTGCCTGTGCTGGGAGAGACCGGCG	122
APC 102686het_delA	AGGTTCTTCCAGATGCTGATACTTTATTACATTTTGC	123
APC het_delAG between 102594_102603 (1 of 5 AG repeats)	GCGAGAAGTACCTAAAAATAAAGCACCTACTGCTGAA	124
APC 102240C > CA, 1344S > S/X	CAGGGTTCTAGTTTATCTTAAGAATCAGCCAGGCACA	125
102676102680delACATT	CCAGATGCTGATACTTTATTTTGGCCACGGAAAGTACTC	126
12487C > CT: 167Q > Q/X	ATGGCCATCTACAAGCAGTCATAGCACATGACGGAGGTTGT	49
102447_102450het_delTGGT	AGAGTGAACCATGCAGTGGAAAAGTGGCATTATAAGCCC	50
12410G > GA, 141C > C/Y	TTTGCCAACTGGCCAAGACTACCCCTGTGCAGCTGTG	51
102678het_delA	CAGATGCTGATACTTTATTACTTTTGGCCACGGAAAGTACT	52
102594_102598het_delAGAGA	AAAGCACCTACTGCTGAAAGAGAGTGGACCTAAGCAAG	127
102776A > AT: 1523R > R/X	ATGACAATGGGAATGAAACAGAATCAGAGCAGCCTAAAG	14
102644_102645het_insG	ATGCTGCAGTTCAGAGGGTCCAGGTTCTTCCAGATGC	54
102594_102595het_delAG	TAAAGCACCTACTGCTGAAAAGAGAGAGTGGACCTAAGCAAG	55
102106het_delT	CACAGGAAGCAGATTCCTGCAATACCCCTGCAATAGCA	56
102442het_delT	TTCAGAGTGAACCATGCAGGGAATGGTAAGTGGCATTAT	57
apc 102494C > CT: 1429Q > Q/X	TCCAGATAGCCCTGGATAAACCATGCCACCAAG	58
apc 102140het_delA	AGCAGAAATAAAAAGAAAAGTTGGAAGTGGTGCAGCTGA	60
apc 102554het_delA	CTCAAACAGCTCAAACCGAGGAGTACCTAAA	61
12487C > CT: 167Q > Q/X	AGTACTCCCCTGCCCTCAAC	128
102447_102450het_delTGGT	TTTGAGAGTCGTTTCGATTGC	129
12410G > GA, 141C > C/Y	AGTACTCCCCTGCCCTCAAC	128
102678het_delA	TCCAGGTTCTTCCAGATGCT	130
102594_102598het_delAGAGA	AGCTCAAACCAAGCGAGAAG	131
102644_102646het_insG	GGACCTAAGCAAGCTGCAGTA	132
102594_102595het_delAG	AGCTCAAACCAAGCGAGAAG	131
102106het_delT	CAGACGACACAGGAAGCAGA	133
102442het_delT	TTTGAGAGTCGTTTCGATTGC	129
apc 102494C > CT: 1429Q > Q/X	GTGAACCATGCAGTGGAAATG	25
apc 102557C > CT: 1450R > RA	ACCACCTCCTCAAACAGCTC	134
apc 102140het_delA	CAGACGACACAGGAAGCAGA	133
apc 102494C > CT: 1429Q > Q/X	GTGAACCATGCAGTGGAAATG	25
apc 102554het_delA	CATGCCACCAAGCAGAAGTA	21
tp53 E5 12647A > AG: 193H > H/R	CAGGCCTCTGATTCTCTACT	36
tp53 E5 12742G > GA	CATAGTGTGGTGGTGCCTTA	135
tp53 E5 12706C > CT: 213R > R/X	GTGGAAGGAAATTTGCGTGT	136
tp53 E4 12712A > AG: 215S > S/G	GTGGAAGGAAATTTGCGTGT	136
tp53 E4 12388T > TC: 134F > F/L	TGTTCACTTGTGCCCTGACT	137
tp53 E3 11606G > GA: 125T > T/T	GTCTGGGCTTCTTGCAATTCT	138
tp53 E6 13379C > CT: 248R > R/W	TGGCTCTGACTGTACCACCA	139
tp53 12E3 11326A > AC (splice site)	ACCTGGTCTCTGACTGCTC	140
tp53 E6 13412G > GT: 259D > D/Y	CCTCACCATCATCACACTGG	141
tp53 E4 12449G > GT: 154G > GN	GTGCAGCTGTGGGTTGATT	142
tp53 E7 13872G > GT, 298E > E/X	GGAAGAGAATCTCCGCAAGA	143
APC 102843C > CG: 1545S > S/X	ATGCCCTCAGTTCAGGAAAA	144
tp53 E4 12392G > GT, 135C > C/F	TGCCCTGACTTTCAACTCTGT	145
APC 102557C > CT: 1450R > R/X	ACCACCTCCTCAAACAGCTC	134
tp53 E7 13819G > T: 280R > I	CTACTGGGACGGAACAGCTT	146
tp53 13E4 11326A > AC (splice site)	ACCTGGTCTCTGACTGCTC	140
tp53 E7 13412G > GT: 259D > D/Y	CCTCACCATCATCACACTGG	41
tp53 E5 12449g > GT: 154G > G/V	GTGCAGCTGTGGGTTGATT	142
tp53 E8 13872G > Gt: 298E > E/X	GGAAGAGAATCTCCGCAAGA	143
tp53 E8 13813C > CG, 278P > P/R	CTACTGGGACGGAACAGCTT	146
tp53 E8 13851A > AT, 291K > K/X	GCGCACAGAGGAAGAGAATC	147
smad4 E2 19049G > GA, 18118A > A/A	AGGTGCCCTGATCTTCACAA	148
tp53 E8 13777G > GA: 266G > G/E	TTTCCTTACTGCCTCTTGCTTC	149
tp53 E6 12653T > TC: 195I > I/T	CAGGCCTCTGATTCTCTACT	36
tp53 E7 13379C > CT: 248R > R/W	TGGCTCTGACTGTACCACCA	139
tp53 E6 12647A > AG: 193H > H/R	CAGGCCTCTGATTCTCTACT	36
tp53 E6 12712A > AG: 215S > S/G	GTGGAAGGAAATTTGCGTGT	136

TABLE 7-continued

Sequence Specific Capture Probes and Primers for AD Cancer Mutation Detection		
tp53 E8 13872G > GT, 298E > E/X	GGAAGAGAATCTCCGCAAGA	143
tp53 E7 13370G > GA: 245G > G/S	TGGCTCTGACTGTACCACCA	139
tp53 E4 11580het_delG	CCCTTCCCAGAAAACCTACC	30
tp53 E5 12524A > AG, 179H > H/R	TGGCCATCTACAAGCAGTCA	150
>tp53 E6 12661G > GT, 198E > E/X	CAGGCCTCTGATTCCTCACT	36
tp53 E8 13872G > GT, 298E > E/X	GGAAGAGAATCTCCGCAAGA	143
apc 102494C > CT: 1429Q > Q/X	CAGGAGACCCCACTCATGTT	19
apc 102557C > CT: 1450R > R/X	CAGGAGACCCCACTCATGTT	19
apc 102140het_delA	TTCATTATCATCTTTGTCATCAGC	15
apc 102494C > CT: 1429Q > Q/X	CAGGAGACCCCACTCATGTT	19
apc 102134G > GT: 1309E > E/X	TTCATTATCATCTTTGTCATCAGC	15
apc102554het_delA	CAGGAGACCCCACTCATGTT	19
apc 102852het_insA	GAGCCTCGATGAGCCATTTA	23
Kras 5571G > GA: 12G > G/D	AGGCCTGCTGAAAATGACTG	2
tp53 E4 12392G > GA: 135C > C/Y	TGTTCACTTGTGCCCTGACT	137
tp53 E5 12655C > CT: 196R > R/X	CAGGCCTCTGATTCCTCACT	36
tp53 E6 13350G > GA: 238C > C/Y	TGGCTCTGACTGTACCACCA	139
tp53 E6 13420G > GA	CCTCACCATCATCACACTGG	141
tp53 E5 12712A > AG: 215S > S/G	GTGGAAGGAAATTTGCGTGT	136
Kras 5571G > GA: 12G > G/D	AGGCCTGCTGAAAATGACTG	2
P16 (ink4a) E1 19638A > AT	AGCCAGTCAGCCGAAGG	151
P16 (ink4a) E2 23353G > GT: 447D > DY	CACCCCTGGCTCTGACCAT	152
P16 (ink4a) E1 19638A > AT	AGCCAGTCAGCCGAAGG	151
p16(ink4a) E2 23402het_delT	GACCCCGCCACTCTCAC	153
p16(ink4a) E2 23403C > CA: 484F > F/ ctnbl 25541het_delT	GACCCCGCCACTCTCAC	153
cdh 1 E9 76435het_delA	ATATTTCAATGGGTTCATATCACAG	154
cdh1 E1 743_744het_insAGCCCTGCGCCCA	GCCATGATCGCTCAAATACA	155
cdh 1 E13 8685386854het_insA	ACTTGGGAGGGACGCATT	156
cdh 1 E14 91472C > CT: 751N > N/N	CATTCTGGGGATTCTTGGAG	157
cdh 1 E 15	CTGTTTCTTCGGAGGAGAGC	158
92868_92896het delTTGACTTGA	TTCCTACTCTTCATTGTACTTCAACC	159
GCCAGCTGCACAGGGCCTG		
cdh 1 E4 71669*het_delA	CGTTTCTGGAATCCAAGCAG	160
cdh 1 E7 74926G > GA: 289A > A/T	CCAGGAACCTCTGTGATGGA	161
cdh 1 E1 736_742het_delTGGCCTC	ACTTGGGAGGGACGCATT	156
p16(ink4a) E1 19638A > AT	AGCCAGTCAGCCGAAGG	151
tp53 E4 12365A > AG: 126Y > Y/C	CACCTGTGCCCTGACTTTCA	33
tp53 E4 12548G > GA	CACATGACGGAGGTTGTGAG	162
p16(ink4a) E1 19810T > TG: 491I > I/S	TTCCAATTCCCCTGCAA	163
tp53 E7 13757G > GA	GGGACAGGTAGGACCTGATTT	164
tp53 E7 13815G > GC: 279G > G/R	CTACTGGGACGGAACAGCTT	146
tp53 E7 13816G > GA: 279G > G/E	CTACTGGGACGGAACAGCTT	146
tp53 E5 12365A > AC, 126Y > Y/S	CACCTGTGCCCTGACTTTCA	33
tp53 E5 12491A > AT, 168H > HL	TGGCCATCTACAAGCAGTCA	150
	TGGCCATCTACAAGCAGTCA	150
	TGGCCATCTACAAGCAGTCA	150
	AGGCCTGCTGAAAATGACTG	2
kras 5570G > GC, 12G > G/R	TGGCTCTGACTGTACCACCA	139
tp53 E7 13370G > GA, 245G > G/S	TGACAATGGGAATGAAACAGA	165
apc 102864_102865het_delAG	TGTTCACTTGTGCCCTGACT	137
tp53 E5 12386T > TC, 133M > M/T	CCAAAGCATGGCTCATCTCTA	205
cdh1 E15 93059G > GA	CTACTGGGACGGAACAGCTT	146
tp53 E8 13798G > GA: 273R > R/H	GTGGAAGGAAATTTGCGTGT	136
p53 E6, 12698_12701het_delAC (1 or 2 AC repeats)	CTACTGGGACGGAACAGCTT	146
P53 E8, 13824C > CT, 282R > R/W	CAGACGACACAGGAAGCAGA	133
APC 102151G > GA, 1314R > R/R	GTGACGCTGTGGGTTGATT	142
P53 E5 12457 G > G/T	CTACTGGGACGGAACAGCTT	146
p53 E813812C > CG, 278P > P/A	CTGCAGTTCAGAGGGTCCAG	210
APC 102686het_delA	AGCTCAAACCAAGCGAGAAG	131
APC het_delAG between 102594_102603 (1 of 5 AG repeats)		
APC 102240C > CA, 1344S > S/X	CCCTAGAACCAAATCCAGCA	166
102676102680delACATT	CTGCAGTTCAGAGGGTCCAG	167
12487C > CT: 167Q > Q/X	AGTACTCCCCTGCCCTCAAC	128
102447_102450het_delTGGT	TTTGAGAGTCGTTTCGATTGC	129
12410G > GA, 141C > C/Y	AGTACTCCCCTGCCCTCAAC	128
102678het_delA	TCCAGGTTCTTCCAGATGCT	130
102594_102598het_delAGAGA	AGCTCAAACCAAGCGAGAAG	131
102776A > AT: 1523R > R/X	TTTGCCACGGAAGTACTCC	168
102644_102645het_insG	GGACCTAAGCAAGCTGCAGTA	132
102594_102595het_delAG	AGCTCAAACCAAGCGAGAAG	131
102106het_delT	CAGACGACACAGGAAGCAGA	133
102442het_delT	TTTGAGAGTCGTTTCGATTGC	129
apc 102494C > CT: 1429Q > Q/X	GTGAACCATGCAGTGGAAATG	25
apc 102140het_delA	CAGACGACACAGGAAGCAGA	133
apc 102554het_delA	CATGCCACCAAGCAGAAGTA	21

TABLE 7-continued

Sequence Specific Capture Probes and Primers for AD Cancer Mutation Detection		
MUTATION IN TISSUE	ANTISENSE PRIMER 2 (5' TO 3')	SEQ ID No.
12487C > CT: 167Q > Q/X	CTCACAACCTCCGTCATGTG	169
102447_102450het_delTGGT	CATGGTTTGTCCAGGGCTAT	27
12410G > GA, 141C > C/Y	CTCCGTCATGTGCTGTGACT	170
102678het_delA	CACTCAGGCTGGATGAACAA	22
102594_102598het_delAGAGA	AGCATCTGGAAGAACCCTGGA	28
102644_102646het_insG	CACTCAGGCTGGATGAACAA	22
102594_102595het_delAG	AGCATCTGGAAGAACCCTGGA	28
102106het_delT	TGCTGGATTTGGTTCTAGGG	171
102442het_delT	CATGGTTTGTCCAGGGCTAT	27
apc 102494C > CT: 1429Q > Q/X	AGCTGTTTGGAGAGGTGGTG	172
apc 102557C > CT: 1450R > RA	GCAGCTTGCTTAGGTCCACT	173
apc 102140het_delA	TGCTGGATTTGGTTCTAGGG	171
apc 102494C > CT: 1429Q > Q/X	AGCTGTTTGGAGAGGTGGTG	172
apc 102554het_delA	GCAGCTTGCTTAGGTCCACT	173
tp53 E5 12647A > AG: 193H > H/R	ACACGCAAATTTCCCTCCAC	174
tp53 E5 12742G > GA	AACCACCCTTAACCCCTCCT	175
tp53 E5 12706C > CT: 213R > R/X	CAGTTGCAAACCAGACCTCA	176
tp53 E4 12712A > AG: 215S > S/G	CAGTTGCAAACCAGACCTCA	176
tp53 E4 12388T > TC: 134F > F/L	GCAGGCTTTGGCCAGTTG	177
tp53 E3 11606G > GA: 125T > T/T	GCCAGGCATTGAAGTCTCAT	31
tp53 E6 13379C > CT: 248R > R/W	CCAGTGTGATGATGGTGAGG	178
tp53 12E3 11326A > AC (splice site)	GGGGACAGCATCAAATCATC	179
tp53 E6 13412G > GT: 259D > D/Y	GGGTCAGAGGCAAGCAGA	40
tp53 E4 12449G > GT: 154G > GN	CTCCGTCATGTGCTGTGACT	170
tp53 E7 13872G > GT, 298E > E/X	GCTTCTTGCTCTTGCTTGCTT	43
APC 102843C > CG: 1545S > S/X	TTTTTCTGCCTCTTTCTCTTGG	180
tp53 E4 12392G > GT, 135C > C/F	CTGCACAGGGCAGGTCTT	181
APC 102557C > CT: 1450R > R/X	GCAGCTTGCTTAGGTCCACT	173
tp53 E7 13819G > T: 280R > I	GCGGAGATTCTCTTCTCTCTG	182
tp53 13E4 11326A > AC (splice site)	GGGGACAGCATCAAATCATC	179
tp53 E7 13412G > GT: 259D > D/Y	GGGTCAGAGGCAAGCAGA	40
tp53 E5 12449g > GT: 154G > G/V	CTCCGTCATGTGCTGTGACT	170
tp53 E8 13872G > Gt: 298E > E/X	GCTTCTTGCTCTTGCTTGCTT	43
tp53 E8 13813C > CG, 278P > P/R	GCGGAGATTCTCTTCTCTCTG	182
tp53 E8 13851A > AT, 291K > K/X	TTCTTGCTCTTGCTTGCTTACC	183
smad4 E2 19049G > GA, 18118A > A/A	TGGATTACACAGACACTATCACA	184
tp53 E8 13777G > GA: 266G > G/E	CACAAACACGCACCTCAAAG	185
tp53 E6 12653T > TC: 195I > I/T	ACACGCAAATTTCCCTCCAC	174
tp53 E7 13379C > CT: 248R > R/W	CCAGTGTGATGATGGTGAGG	178
tp53 E6 12647A > AG: 193H > H/R	ACACGCAAATTTCCCTCCAC	174
tp53 E6 12712A > AG: 215S > S/G	CAGTTGCAAACCAGACCTCA	76
tp53 E8 13872G > GT, 298E > E/X	GCTTCTTGCTCTTGCTTGCTT	43
tp53 E7 13370G > GA: 245G > G/S	CCAGTGTGATGATGGTGAGG	178
tp53 E4 11580het_delG	ACTGACCGTGCAAGTCACAG	186
tp53 E5 12524A > AG, 179H > H/R	CTGCTCACCATCGCTATCTG	187
>tp53 E6 12661G > GT, 198E > E/X	CCAAATACTCCACACGCAAA	188
tp53 E8 13872G > GT, 298E > E/X	GCTTCTTGCTCTTGCTTGCTT	43
apc 102494C > CT: 1429Q > Q/X	TGGCAAATGTAATAAAGTATCAGC	20
apc 102557C > CT: 1450R > R/X	TGGCAAATGTAATAAAGTATCAGC	20
apc 102140het_delA	CGCTCCTGAAGAAAATTCAA	16
apc 102494C > CT: 1429Q > Q/X	TGGCAAATGTAATAAAGTATCAGC	20
apc 102134G > GT: 1309E > E/X	CGCTCCTGAAGAAAATTCAA	16
apc102554het_delA	TGGCAAATGTAATAAAGTATCAGC	20
apc 102852het_insA	TCAATATCATCATCATCTGAATCATC	24
Kras 5571G > GA: 12G > G/D	TTGTTGGATCATATTCGTCCAC	3
tp53 E4 12392G > GA: 135C > C/Y	GCAGGCTTTGGCCAGTTG	177
tp53 E5 12655C > CT: 196R > R/X	ACACGCAAATTTCCCTCCAC	174
tp53 E6 13350G > GA: 238C > C/Y	CCAGTGTGATGATGGTGAGG	178
tp53 E6 13420G > GA	GGGTCAGAGGCAAGCAGA	40
tp53 E5 12712A > AG: 215S > S/G	CAGTTGCAAACCAGACCTCA	176
Kras 5571G > GA: 12G > G/D	TTGTTGGATCATATTCGTCCAC	3
P16 (ink4a) E1 19638A > AT	GAGGGGCTGGCTGGTC	189
P16 (ink4a) E2 23353G > GT: 447D > DY	GGGTCGGGTGAGAGTGG	190
P16 (ink4a) E1 19638A > AT	GAGGGGCTGGCTGGTC	189
p16(ink4a) E2 23402het_delT	CAGCTCCTCAGCCAGGTC	191
p16(ink4a) E2 23403C > CA: 484F > F/	CAGCTCCTCAGCCAGGTC	191
ctnnb1 25541het_delT	TCAAATCAGCTATAAATACGAAACA	192
cdh 1 E9 76435het_delA	TCTCAGGGGGCTAAAGGATT	193
cdh1 E1 743_744het_insAGCCCTGCGCCCA	GAAGAAGGGAAGCGGTGAC	194
cdh 1 E13 8685386854het_insA	GGAAATAAACCTCCTCCATTTT	195
cdh 1 E14 91472C > CT: 751N > N/N	CCGCCCTCTTCTTCATCATA	196
cdh 1 E 15	TGCAACGTCGTTACGAGTCA	197
92868_92896het delTTGACTTGA		
GCCAGCTGCACAGGGGCTG		

TABLE 7-continued

Sequence Specific Capture Probes and Primers for AD Cancer Mutation Detection			
cdh 1 E4 71669*het_delA	GCAGCTGATGGGAGGAATAA	198	
cdh 1 E7 74926G > GA: 289A > A/T	TGAGGATGGTGTAAAGCGATG	199	
cdh 1 E1 736_742het_delTGCGCCC	GAAGAAGGGAAGCGGTGAC	194	
p16(ink4a) E1 19638A > AT	CTCACAACTCCGTCATGTG	169	
tp53 E4 12365A > AG: 126Y > Y/C	GCCAGTTGGCAAAACATCT	200	
tp53 E4 12548G > GA	AACCAGCCCTGTCGTCCT	34	
p16(ink4a) E1 19810T > TG: 491I > I/S	CCCACGCACCGAATAGT	201	
tp53 E7 13757G > GA	AGCTGTTCCGTCCCAGTAGA	202	
tp53 E7 13815G > GC: 279G > G/R	GCGGAGATTCTTCTCCTCTG	182	
tp53 E7 13816G > GA: 279G > G/E	GCGGAGATTCTTCTCCTCTG	182	
tp53 E5 12365A > AC, 126Y > Y/S	GCCAGTTGGCAAAACATCT	200	
tp53 E5 12491A > AT, 168H > HL	CTGCTCACCATCGCTATCTG	187	
	TCACCATCGCTATCTGAGCA	203	
	AACCAGCCCTGTCGTCCT	34	
	TTGTTGGATCATATTCGTCCAC	3	
kras 5570G > GC, 12G > G/R	CCAGTGTGATGATGGTGAGG	178	
tp53 E7 13370G > GA, 245G > G/S	GGTCTTTTTCAGAATCAATAGTTTT	204	
apc 102864_102865het_delAG	GCAAGTCTTGGCCAGTTG	177	
tp53 E5 12386T > TC, 133M > M/T	CTCAGGCAAGCTGAAAACAT	206	
cdh1 E15 93059G > GA	GCGGAGATTCTTCTCCTCTG	182	
tp53 E8 13798G > GA: 273R > R/H	GCCAGATTCTTCTCCTCTG	182	
p53 E6, 12698_12701het_delAC (1 or 2 AC repeats)	AGCTGTTTGAGGAGGTGGTG	172	
P53 E8, 13824C > CT, 282R > R/W	GCGGAGATTCTTCTCCTCTG	182	
APC 102151G > GA, 1314R > R/R	GTGACACTGCTGGAACCTCG	207	
P53 ES 12457 G > G/T	CTCCGTCATGTGCTGTGACT	170	
p53 E813812C > CG, 278P > P/A	GCGGAGATTCTTCTCCTCTG	182	
APC 102686het_delA	CACTCAGGCTGGATGAACAA	22	
APC het_delAG between 102594_102603 (1 of 5 AG repeats)	AGCATCTGGAAGAACCCTGGA	28	
APC 102240C > CA, 1344S > S/X	TGTCTGAGCACCACCTTTTGG	208	
102676102680delACATT	CACTCAGGCTGGATGAACAA	22	
12487C > CT: 167Q > Q/X	CTCACAACTCCGTCATGTG	169	
102447_102450het_delTGGT	CATGGTTTGTCAGGGCTAT	27	
12410G > GA, 141C > C/Y	CTCCGTCATGTGCTGTGACT	170	
102678het_delA	CACTCAGGCTGGATGAACAA	22	
102594_102598het_delAGAGA	AGCATCTGGAAGAACCCTGGA	28	
102776A > AT: 1523R > R/X	TTTCTGAACTGGAGGCATT	209	
102644_102645het_insG	CACTCAGGCTGGATGAACAA	22	
102594_102595het_delAG	AGCATCTGGAAGAACCCTGGA	28	
102106het_delT	TGCTGGATTTGGTTCTAGGG	171	
102442het_delT	CATGGTTTGTCAGGGCTAT	27	
apc 102494C > CT: 1429Q > Q/X	AGCTGTTTGAGGAGGTGGTG	172	
apc 102140het_delA	TGCTGGATTTGGTTCTAGGG	171	
apc 102554het_delA	GCAGCTTGCTTAGGTCCACT	173	

Target imitations were not detected in control stools. Target mutations were detected in stools from 68% (47/69) of patients with a GI neoplasm. Specifically, target mutations were detected in stools from 71% (36/51) of patients with cancer [40% (2/5) with oropharyngeal, 65% (11/17) with esophageal, 100% (4/4) with gastric, 55% (6/11) with pancreatic, 75% (3/4) with biliary or gallbladder, and 100%

(10/10) with colorectal] and from 61% (11/18) with precancers [100% (2/2) with pancreatic intraductular papillary mucinous neoplasia and 56% (9/16) with colorectal advanced adenoma]. Mutant copies in genes recovered from stool averaged 0.4% (range 0.05-13.4%) for supracolonial and 1.4% (0.1-15.6%) for colorectal neoplasms, $p=0.004$ (Table 8).

TABLE 8

Digital Melt Curve Detection of Validated Mutations in AD Cancer Patient Stool								
#	ID	Site	Age	Gender	Tissue Mutation	Stool Detection	Mutation Frequency %	Normal Control
1	1163	Head/Neck(pharynx)	73	M	tp53	YES	0.8	Neg
2	1250	Head/Neck(pharynx)	49	M	tp53	NO		Neg
3	1295	Head/Neck(pharynx)	47	F	tp53	NO		Neg
4	1391	Head/Neck	65	M	tp53 TP53	NO (both)		Neg
5	1427	Head/Neck	60	M	tp53 tp53	YES(p53-1), No (p53-2)	0.05	Neg
1	745	Esophagus	84	F	tp53	YES	0.4	Neg
2	769	Esophagus	56	F	tp53	YES	0.4	Neg
3	782	Esophagus	55	M	tp53	NO		Neg
4	789	Esophagus	61	M	tp53	YES	1.6	Neg
5	819	Esophagus	53	M	tp53	YES	0.2	Neg
6	873	Esophagus	61	M	tp53	YES	0.2	Neg
7	906	Esophagus	55	M	APC	YES	0.8	Neg
8	1049	Esophagus	57	M	tp53	NO		Neg
9	1064	Esophagus	72	F	tp53	NO		Neg
10	1067	Esophagus	72	M	tp53	YES	0.7	Neg

TABLE 8-continued

Digital Melt Curve Detection of Validated Mutations in AD Cancer Patient Stool									
#	ID	Site	Age	Gender	Tissue Mutation		Stool Detection	Mutation Frequency %	Normal Control
11	1103	Esophagus	78	M	tp53		NO		Neg
12	1199	Esophagus	66	M	tp53		YES	0.5	NEG
13	1307	Esophagus	51	M	tp53		NO		NEG
14	1373	Esophagus	76	M	tp53		YES	0.5	NEG
15	1414	Esophagus	66	M	tp53		YES	0.1	NEG
16	1448	Esophagus	82	M	tp53		NO		NEG
17	1072	Esophagus			tp53		YES	0.4	NEG
1	798	Stomach	81	M	cdh1		YES	13.2	NEG
3	1221	Stomach	55	M	cdh1	cdh1	YES(both)	8,1.3	NEG
4	1224	Stomach	75	F	smad4	cdh1	YES(smad4),No(CDH1)	0.2	NEG
5	1402	Stomach	56	M	APC	tp53	YES (p53)	0.1	NEG
1	848	Gall Bladder	67	M	tp53		YES	0.1	NEG
2	1315	Gall Bladder	57	F	tp53		YES	1.4	NEG
1	1043	Bile Duct	51	F	APC		NO		NEG
2	1554	Bile Duct	77	M	cdh1		YES	13.4	NEG
1	757	Pancreatic Cancer in situ	78	M	K-ras		YES	0.2	NEG
2	1349	Pancreatic Cancer in situ	64	M	K-ms		YES	0.2	NEG
1	839	Pancreas	69	F	tp53		YES	0.2	NEG
2	1204	Pancreas	65	F	p16		NO		NEG
3	1253	Pancreas	63	F	K-ras	tp53	Yes(k-ras), No(p53)	2	NEG
4	1400	Pancreas	71	F	tp53	K-ras	No (both)		NEG
5	1547	Pancreas	77	F	tp53		NO		NEG
6	1217	Pancreas			K-ras		NO		NEG
7	1073	Pancreas			K-ras		NO		NEG
8	532	Pancreas			K-ras		YES	1	NEG
9	1592	Pancreas			K-ras	P53	YES (both)	0.3	NEG
10	1695	Pancreas			K-ras		YES	0.2	NEG
11	1058	Pancreas			K-ras	P53 APC	YES(K-ras)	0.2	NEG
1	438	Colorectal Cancer	78	F	APC		YES	1.2	NEG
2	446	Colorectal Cancer	74	M	BRAF		YES	0.4	NEG
3	529	Colorectal Cancer	46	M	K-RAS		YES	1	NEG
4	489	Colorectal Cancer	73	M	K-RAS		YES	2.6	NEG
5	549	Colorectal Cancer	79	M	BRAF		YES	1.6	NEG
6	551	Colorectal Cancer	69	M	K-RAS		YES	5.8	NEG
7	584	Colorectal Cancer	68	M	K-RAS		YES	1.4	NEG
8	894	Colorectal Cancer	57	M	P53	APC	YES(p53,APC)	1.6,5	NEG
9	998	Colorectal Cancer	45	F	APC	KRAS	YES(K-ras,APC)	0.6,0.8	NEG
10	1009	Colorectal Adenoma	65	F	P53		YES	12.9	NEG
1	513	Colorectal Adenoma	65	F	APC		YES	0.1	NEG
2	546	Colorectal Adenoma	61	M	APC		NO		NEG
3	547	Colorectal Adenoma	52	F	APC		NO		NEG
4	568	Colorectal Adenoma	52	M	APC		YES	7.8	NEG
5	578	Colorectal Adenoma	71	F	APC		NO		NEG
6	590	Colorectal Adenoma	54	F	APC		YES	3.2	NEG
7	701	Colorectal Adenoma	72	F	APC		NO		NEG
8	855	Colorectal Adenoma	75	M	K-RAS		YES	0.4	NEG
9	860	Colorectal Adenoma	53	M	APC		YES	15.6	NEG
10	900	Colorectal Adenoma	64	F	APC	K-RAS	No(both)		NEG
11	962	Colorectal Adenoma	56	M	K-RAS		Yes	1	NEG
12	965	Colorectal Adenoma	82	M	APC	K-RAS	No (both)		NEG
13	991	Colorectal Adenoma	79	M	APC	K-RAS	YES(K-ras), No(APC)	0.2	NEG
14	1135	Colorectal Adenoma	59	M	K-RAS		YES	13	NEG
15	1231	Colorectal Adenoma	50	M	APC		NO		NEG
16	1559	Colorectal Adenoma			K-RAS		YES	1	NEG

55

We also performed an initial pilot study with 10 stool samples from patients with confirmed bile duct cancers to determine if DMC technology could detect mutations in k-ras, a well characterized gene known be mutated in this population. K-ras mutations were detected in stools for 3/10 or 4/10 bile duct cancers (depending on mutation score of 5 or 3, respectively) (Table 9). As K-ras is mutant in 30-40% of bile duct cancers, these results indicate that the detection assay is picking up the appropriate proportion of cancer samples.

TABLE 9

K-ras mutation scores for patients with bile duct cancer.				
Sample #	Pathology	K-ras Mutation Score	Mutation Detected	
			A	B
520	BD Cancer	0		
528	BD Cancer	0		
559	BD Cancer	0		
558	BD Cancer	1	Codon 12	GAT

60

65

TABLE 9-continued

K-ras mutation scores for patients with bile duct cancer.				
Sample		K-ras Mutation	Mutation Detected	
#	Pathology	Score	A	B
515	BD Cancer	2	Codon 12 GAT	
543	BD Cancer	2	Codon 13 GAC	
806	BD Cancer	3	Codon 12 TGT	
539	BD Cancer	5	Codon 12 GAT	Codon 13 GGA
512	BD Cancer	6	Codon 12 GAT	Codon 12 GAT
725	BD Cancer	25	Codon 13 GAC	Codon 12 GAT; Codon 12 GTT

Allele Specific PCR

The allele specific-PCR assay was a modified version of a previously published method (e.g., Cha et al., Mismatch Amplification Mutation Assay (MAMA): Application to the c-H-ras Gene PCR Methods and Applications, 2:14-20 (1992) Cold Spring Harbor Laboratory). TP53 gene fragments were captured from stool DNA samples with probes specific to mutations identified in the matched tissue (Table 7). Copy numbers were assessed by qPCR. Samples were adjusted to 10,000 fragments each and amplified with allele specific primer sets.

Sample	F Primer	R Primer
A743	GACAGAAACACTTTAT (SEQ ID No: 211)	CGGCTCATAGGG (SEQ ID No: 217)
A848	ACACTTTTCGACAAG (SEQ ID No: 212)	AAACCAGACCTCAG (SEQ ID No: 218)
A789	CCTCAACAAGATAC (SEQ ID No: 213)	CAGCTGCACAGG (SEQ ID No: 219)
A782	GCCGCCTGAAA (SEQ ID No: 214)	AGACCCCAGTTGC (SEQ ID No: 220)
A873	GCGGCATGAAAT (SEQ ID No: 215)	TTCCAGTGTGATGAT (SEQ ID No: 221)
A769	CCCCTCCTCAGAG (SEQ ID No: 216)	CTTCCACTCGGATAA (SEQ ID No: 222)

The forward primer in each case is specific for each TP53 mutation.

Esophagus and Stomach

Targeting mutations found in esophageal cancers or those from gastroesophageal junction (on p53, APC, or K-ras), the same mutation was detected by allele-specific PCR in matched stools from five of five (100%) cancers but in none of the controls (Table 10). The threshold cycle (Ct), designates the PCR cycle at which the product enters the exponential phase of amplification.

Gallbladder

Targeting a mutation confirmed in a gallbladder cancer, the same mutation was found in the matched stool from that patient using allele-specific PCR (Table 10).

TABLE 10

Quantitative Mutant Allele Specific-PCR Results for Matched Aero-digestive Cancers				
Sample	Gene	# fragments	Ct	
A769	esophageal/gastric cancer	p53	10K	71
N	normal	p53	10K	>80

TABLE 10-continued

Quantitative Mutant Allele Specific-PCR Results for Matched Aero-digestive Cancers				
Sample	Gene	# fragments	Ct	
A782	esophageal/gastric cancer	p53	10K	38.8
N	normal	p53	10K	44.1
A745	esophageal/gastric cancer	p53	30K	42.5
N	normal	p53	30K	45.6
A873	esophageal/gastric cancer	p53	30K	37.8
N	normal	p53	30K	40.4
A848	gall bladder cancer	p53	10K	22.4
N	normal	p53	10K	36.3
A789	esophageal/gastric cancer	p53	10K	25.9
N	normal	p53	10K	28.7

Example 9—Candidate Stool Polypeptide Markers Identified for Colorectal Cancer and Precancerous Adenomas

The following list of polypeptides were identified by a statistical analysis model using all data generated from mass spectral of fecal protein extracts: β 2-macroglobulin, complement C3 protein, serotransferrin, haptoglobin, carbonic anhydrase 1, xaa-pro dipeptidase, leukocyte elastase inhibitor, hemoglobin, glucose-6-phosphate, and catalase. This list of polypeptides is in order of difference from normal. Thus, the mean spectral abundance for β 2-macroglobulin is most different from normal for cancer and adenoma.

The statistical significance of relative poly peptide abundance between normal, adenoma, and colorectal cancer (CRC) was obtained using normalized spectral count data from a zero inflated poisson regression model as an offset term in the protein specific differential expression analysis. The differential expression analysis also incorporated the zero inflated poisson regression model. Polypeptides were then ranked according to their statistical significance and whether the expression profile followed the clinically relevant pattern of Normal<Adenoma<CRC. Using a rule that a positive test required that any three of top six markers be positive, the sensitivity and specificity of this panel were both 100% in a training set.

The listed polypeptides can be used individually or in any combination to detect colorectal cancer or precancerous adenomas.

Example 10—Identification of Polypeptide Markers for Pancreatic Cancer

Potential polypeptide markers for pancreatic cancer prediction were identified. Utilizing a Scaffold (Proteome Software) side-by-side comparison of spectral abundances, ratios of the spectral counts of carboxypeptidase B (CBPB1_HUMAN) and Carboxypeptidase A1 (CBPA1_HUMAN) were compared. A value for carboxypeptidase B/A1 of 0.7 or higher was predictive of pancreatic cancer (in normal stools, an average ratio of 2:3 B/A1 was observed). Specificity for training data was 100% with a sensitivity of 88%, while sensitivity from a validation set was 82% at the same specificity.

Example 11—Use of Fecal Methylated ALX4 as a Neoplasia Marker

Stools from patients with colorectal tumors were found to contain significantly elevated amounts of methylated ALX4 gene copies, but those from normal individuals were found to contain none or only trace amounts. When fecal methylated ALX4 was assayed with an appropriate amplification method, colorectal cancers and premalignant adenomas were specifically detected. At 90% specificity, fecal methylated ALX4 detected 59% colorectal cancer and 54%

pre-malignant adenomas, allowing for the detection of both colorectal cancer and premalignant adenomas.

OTHER EMBODIMENTS

It is to be understood that while the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 222

<210> SEQ ID NO 1

<211> LENGTH: 37

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 1

gtggacgaat atgatccaac aatagaggta aatcttg

37

<210> SEQ ID NO 2

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 2

aggcctgctg aaaatgactg

20

<210> SEQ ID NO 3

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 3

ttggttgatc atattcgtcc ac

22

<210> SEQ ID NO 4

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 4

taaggcctgc tgaaaatgac

20

<210> SEQ ID NO 5

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 5

atcaaagaat ggtcctgcac

20

-continued

<210> SEQ ID NO 6
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 6

cgctctgcagt caactggaat tt 22

<210> SEQ ID NO 7
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 7

tgtatcgtca aggcactctt gc 22

<210> SEQ ID NO 8
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 8

cttaagcgtc gatggaggag 20

<210> SEQ ID NO 9
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 9

ccagacaact gttcaaactg atgggaccca ctccatc 37

<210> SEQ ID NO 10
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 10

ccacaaaatg gatccagaca 20

<210> SEQ ID NO 11
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 11

tgcttgctct gataggaaaa tg 22

<210> SEQ ID NO 12
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

-continued

<400> SEQUENCE: 12
cagatagccc tggacaaacc atgccaccaa gcagaag 37

<210> SEQ ID NO 13
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 13
ttccagcagt gtcacagcac cctagaacca aatccag 37

<210> SEQ ID NO 14
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 14
atgacaatgg gaatgaaaca gaatcagagc agcctaaag 39

<210> SEQ ID NO 15
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 15
ttcattatca tctttgtcat cagc 24

<210> SEQ ID NO 16
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 16
cgctcctgaa gaaaattcaa 20

<210> SEQ ID NO 17
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 17
tgcagggttc tagtttatct tca 23

<210> SEQ ID NO 18
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 18
ctggcaatcg aacgactctc 20

<210> SEQ ID NO 19
<211> LENGTH: 20

-continued

<212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 19

 caggagaccc cactcatggt 20

 <210> SEQ ID NO 20
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 20

 tggcaaatg taataaagta tcagc 25

 <210> SEQ ID NO 21
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 21

 catgccacca agcagaagta 20

 <210> SEQ ID NO 22
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 22

 cactcaggct ggatgaacaa 20

 <210> SEQ ID NO 23
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 23

 gagcctcgat gagccattta 20

 <210> SEQ ID NO 24
 <211> LENGTH: 26
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 24

 tcaatatcat catcatctga atcatc 26

 <210> SEQ ID NO 25
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 25

-continued

 gtgaacctg cagtggatg 20

<210> SEQ ID NO 26
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 26

acttctcgt tggtttgagc 20

<210> SEQ ID NO 27
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 27

catggtttgt ccaggctat 20

<210> SEQ ID NO 28
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 28

agcatctgga agaactgga 20

<210> SEQ ID NO 29
 <211> LENGTH: 37
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 29

aagaccagg tccagatgaa gctcccagaa tgccaga 37

<210> SEQ ID NO 30
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 30

cccttcccag aaaactacc 20

<210> SEQ ID NO 31
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 31

gccaggcatt gaagtctcat 20

<210> SEQ ID NO 32
 <211> LENGTH: 37
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 32

 catggccatc tacaagcagt cacagcacat gacggag 37

<210> SEQ ID NO 33
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 33

 cacttgtagc ctgactttca 20

<210> SEQ ID NO 34
 <211> LENGTH: 19
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 34

 aaccagccct gtcgtctct 19

<210> SEQ ID NO 35
 <211> LENGTH: 37
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 35

 agtggaggga aattgctg tggagtattt ggatgac 37

<210> SEQ ID NO 36
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 36

 caggcctctg attcctcact 20

<210> SEQ ID NO 37
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 37

 cttaaccct cctcccagag 20

<210> SEQ ID NO 38
 <211> LENGTH: 37
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 38

 atgtgtaaca gttcctgcat gggcggcatg aaccgga 37

-continued

<210> SEQ ID NO 39
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 39

cttgggcctg tggtatctcc 20

<210> SEQ ID NO 40
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 40

gggtcagagg caagcaga 18

<210> SEQ ID NO 41
 <211> LENGTH: 37
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 41

cgcacagagg aagagaatct ccgcaagaaa ggggagc 37

<210> SEQ ID NO 42
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 42

gggagtagat ggagcctggt 20

<210> SEQ ID NO 43
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 43

gcttcttgtc ctgcttgctt 20

<210> SEQ ID NO 44
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 44

acgcctgtaa tcccagcact t 21

<210> SEQ ID NO 45
 <211> LENGTH: 19
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

-continued

<400> SEQUENCE: 45
 tcgcccaggc tggagtgca 19

<210> SEQ ID NO 46
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 46
 tggtgaaacc ccgtctctac 20

<210> SEQ ID NO 47
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 47
 ctactgcaa cctccacctc 20

<210> SEQ ID NO 48
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 48
 cgcccggcta atttttgtat 20

<210> SEQ ID NO 49
 <211> LENGTH: 41
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 49
 atggccatct acaagcagtc atagcacatg acggaggttg t 41

<210> SEQ ID NO 50
 <211> LENGTH: 39
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 50
 agagtgaacc atgcagtgga aaagtggcat tataagccc 39

<210> SEQ ID NO 51
 <211> LENGTH: 37
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 51
 tttgccaact ggccaagacc taccctgtgc agctgtg 37

<210> SEQ ID NO 52

-continued

<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 52

cagatgctga tactttatta cttttgccac ggaaagtact 40

<210> SEQ ID NO 53
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 53

aaagcaccta ctgctgaaag agagtggacc taagcaag 38

<210> SEQ ID NO 54
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 54

atgctgcagt tcagaggggt ccaggttctt ccagatgc 38

<210> SEQ ID NO 55
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 55

taaagcacct actgctgaaa agagagagtg gacctaagca ag 42

<210> SEQ ID NO 56
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 56

cacaggaagc agattctgca ataccctgca aatagca 37

<210> SEQ ID NO 57
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 57

ttcagagtga accatgcagg gaatggtaag tggcattat 39

<210> SEQ ID NO 58
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 58

-continued

tccagatagc cctggataaa ccatgccacc aag 33

<210> SEQ ID NO 59
 <211> LENGTH: 41
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide
 <400> SEQUENCE: 59

ctcaaacagc tcaaaccaag tgagaagtac ctaaaaataa a 41

<210> SEQ ID NO 60
 <211> LENGTH: 38
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide
 <400> SEQUENCE: 60

agcagaaata aaagaaaagt tggaactagg tcagctga 38

<210> SEQ ID NO 61
 <211> LENGTH: 34
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide
 <400> SEQUENCE: 61

ctcaaacagc tcaaaccagc gagaagtacc taaa 34

<210> SEQ ID NO 62
 <211> LENGTH: 35
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide
 <400> SEQUENCE: 62

tctggcccct cctcagcgtc ttatccgagt ggaag 35

<210> SEQ ID NO 63
 <211> LENGTH: 35
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide
 <400> SEQUENCE: 63

cctatgagcc gcctgagatc tggtttgcaa ctggg 35

<210> SEQ ID NO 64
 <211> LENGTH: 35
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide
 <400> SEQUENCE: 64

atgacagaaa cactttttga catagtgtgg tgggtg 35

<210> SEQ ID NO 65
 <211> LENGTH: 35
 <212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 65

 gaaacacttt tcgacatggt gtgggtggtgc cctat 35

<210> SEQ ID NO 66
 <211> LENGTH: 35
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 66

 ctgccctcaa caagatgctt tgccaactgg ccaag 35

<210> SEQ ID NO 67
 <211> LENGTH: 35
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 67

 aagtctgtga cttgcacagt cagttgccct gaggg 35

<210> SEQ ID NO 68
 <211> LENGTH: 35
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 68

 gcatgggagg catgaactgg aggcccatcc tcacc 35

<210> SEQ ID NO 69
 <211> LENGTH: 35
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 69

 tcttttcacc catctaccgt cccccttgcc gtccc 35

<210> SEQ ID NO 70
 <211> LENGTH: 35
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 70

 ccatcatcac actggaatac tccaggtcag gagcc 35

<210> SEQ ID NO 71
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 71

 acccccgcc gtcaccgcg tcc 23

-continued

<210> SEQ ID NO 72
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 72

ggaacagctt tgaggtgtgt gtttgtgcct gtcct 35

<210> SEQ ID NO 73
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 73

tcagagcagc ctaaagaatg aatgaaaac caagagaaa 39

<210> SEQ ID NO 74
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 74

cctcaacaag atgtttttcc aactggccaa gacct 35

<210> SEQ ID NO 75
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 75

cctgtcctgg gatagaccgg cgcac 25

<210> SEQ ID NO 76
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 76

ggagcctcac cactagctgc ccccagg 27

<210> SEQ ID NO 77
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 77

gtgtttgtgc ctgtcgtggg agagaccggc g 31

<210> SEQ ID NO 78
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 78

ggaagagaat ctccgctaga aaggggagcc tca 33

<210> SEQ ID NO 79
 <211> LENGTH: 41
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 79

gttaaatatt gtcagtatgc atttgactta aaatgtgata g 41

<210> SEQ ID NO 80
 <211> LENGTH: 37
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 80

tagtggaat ctactggaac ggaacagctt tgaggtg 37

<210> SEQ ID NO 81
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 81

cctcctcagc atcttaccg agtggaagga aat 33

<210> SEQ ID NO 82
 <211> LENGTH: 31
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 82

agttcctgca tgggcagcat gaaccggagg c 31

<210> SEQ ID NO 83
 <211> LENGTH: 36
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 83

ctgggcttct tgcattctgg acagccaagt ctgtga 36

<210> SEQ ID NO 84
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 84

tgccccacc gtgagcgctg c 21

-continued

<210> SEQ ID NO 85
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 85

tcagcatcctt atccgagtgt aaggaaattt gcgtgtgga 39

<210> SEQ ID NO 86
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 86

tgcaaatagc agaaataaaa taaaagattg gaactaggtc a 41

<210> SEQ ID NO 87
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 87

ctaaagaatc aatgaaaaa ccaagagaaa gaggcagaa 39

<210> SEQ ID NO 88
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 88

gtggtagttg gagctgatgg cgtaggcaag agt 33

<210> SEQ ID NO 89
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 89

cctcaacaag atgttttacc aactggccaa gacct 35

<210> SEQ ID NO 90
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 90

cctcctcagc atcttatctg agtggagga aatttgc 37

<210> SEQ ID NO 91
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

-continued

<400> SEQUENCE: 91
tccactacaa ctacatgtat aacagttcct gcatggg 37

<210> SEQ ID NO 92
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 92
cactggaaga ctccagatca ggagccactt gcc 33

<210> SEQ ID NO 93
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 93
ggagaggggg agtgcaggca gcggg 25

<210> SEQ ID NO 94
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 94
cccaactgcg cctaccccg cactc 25

<210> SEQ ID NO 95
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 95
gcccgggagg gctcctggac acgctg 26

<210> SEQ ID NO 96
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 96
gcccgggagg gcttactgga cacgctggt 29

<210> SEQ ID NO 97
<211> LENGTH: 44
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 97
caatgggtca tatcacagat tctttttttt aaattaaagt aaca 44

<210> SEQ ID NO 98
<211> LENGTH: 40

-continued

<212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 98

 tcttatctca aaagaacaac aaaaaagagg aatcctttag 40

<210> SEQ ID NO 99
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 99

 gcgcccagcc ctgcgccc at tctc 25

<210> SEQ ID NO 100
 <211> LENGTH: 39
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 100

 aagtaagtcc agctggcaaa gtgactcagc ctttgactt 39

<210> SEQ ID NO 101
 <211> LENGTH: 39
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 101

 aggatgacac ccgggacaat gtttattact atgatgaag 39

<210> SEQ ID NO 102
 <211> LENGTH: 36
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 102

 ttttttctcc aaaggactga cgctcggcct gaagtg 36

<210> SEQ ID NO 103
 <211> LENGTH: 34
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 103

 caagcagaat tgctcacttt cccaactcct ctcc 34

<210> SEQ ID NO 104
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 104

-continued

 ggtcacagcc acagacacgg acgatgatgt gaa 33

<210> SEQ ID NO 105
 <211> LENGTH: 24
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 105

agccctgcgc cccttctct cccg 24

<210> SEQ ID NO 106
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 106

ttcctcttcc tacagtgtc ccctgccctc aac 33

<210> SEQ ID NO 107
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 107

tgctcagata gcgatgatga gcagctgggg ctg 33

<210> SEQ ID NO 108
 <211> LENGTH: 27
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 108

ggtcggaggc cgagccaggt gggtaga 27

<210> SEQ ID NO 109
 <211> LENGTH: 39
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 109

gcttctcttt tcctatccta agtagtggtgta atctactgg 39

<210> SEQ ID NO 110
 <211> LENGTH: 27
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 110

ttgtgcctgt cctcggagag accggcg 27

<210> SEQ ID NO 111
 <211> LENGTH: 27
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 111

 tgtgcctgtc ctgagagaga ccggcgc 27

<210> SEQ ID NO 112
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 112

 ttctcttcc tacagtcctc ccctgccctc aac 33

<210> SEQ ID NO 113
 <211> LENGTH: 37
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 113

 tctacaagca gtcacagctc atgacggagg ttgtgga 37

<210> SEQ ID NO 114
 <211> LENGTH: 40
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 114

 aaatgaaaac caagagaaag gcagaaaaa ctattgattc 40

<210> SEQ ID NO 115
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 115

 cctgccctca acaagacgtt ttgccactg gcc 33

<210> SEQ ID NO 116
 <211> LENGTH: 41
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 116

 gctcatctct aagctcagga agagttgtgt caaaaatgag a 41

<210> SEQ ID NO 117
 <211> LENGTH: 39
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 117

 cggaacagct ttgagtgca tgtttgtgcc tgcctggg 39

-continued

<210> SEQ ID NO 118
 <211> LENGTH: 37
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 118

 tggaaaggaaa tttgcgtgtg gagtatttgg atgacag 37

<210> SEQ ID NO 119
 <211> LENGTH: 36
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 119

 tgtcctggga gagactggcg cacagaggaa gagaat 36

<210> SEQ ID NO 120
 <211> LENGTH: 38
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 120

 aagaaaagat tggaaactaga tcagctgaag atcctgtg 38

<210> SEQ ID NO 121
 <211> LENGTH: 30
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 121

 cgcccggcac ccgcttccgc gccatggcca 30

<210> SEQ ID NO 122
 <211> LENGTH: 31
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 122

 gtgtttgtgc ctgtgctggg agagaccggc g 31

<210> SEQ ID NO 123
 <211> LENGTH: 37
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 123

 aggttcttcc agatgctgat actttattac attttgc 37

<210> SEQ ID NO 124
 <211> LENGTH: 37
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

-continued

<400> SEQUENCE: 124
gcgagaagta cctaaaaata aagcacctac tgctgaa 37

<210> SEQ ID NO 125
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 125
cagggttcta gtttatctta agaatcagcc aggcaca 37

<210> SEQ ID NO 126
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 126
ccagatgctg atactttatt ttgccacgga aagtactc 38

<210> SEQ ID NO 127
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 127
aaagcaccta ctgctgaaag agagtggacc taagcaag 38

<210> SEQ ID NO 128
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 128
agtactcccc tgccctcaac 20

<210> SEQ ID NO 129
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 129
ttgagagtc gttcgattgc 20

<210> SEQ ID NO 130
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 130
tccaggttct tccagatget 20

<210> SEQ ID NO 131

-continued

<211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 131

 agctcaaacc aagcgagaag 20

<210> SEQ ID NO 132
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 132

 ggacctaaagc aagctgcagt a 21

<210> SEQ ID NO 133
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 133

 cagacgacac aggaagcaga 20

<210> SEQ ID NO 134
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 134

 accacctcct caaacagctc 20

<210> SEQ ID NO 135
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 135

 catagtgtgg tggcgccta 20

<210> SEQ ID NO 136
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 136

 gtggaaggaa atttgcgtgt 20

<210> SEQ ID NO 137
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 137

-continued

tggtcacttg tgcctgact	20
<p><210> SEQ ID NO 138 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetically generated oligonucleotide</p>	
<400> SEQUENCE: 138	
gtctgggctt cttgcattct	20
<p><210> SEQ ID NO 139 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetically generated oligonucleotide</p>	
<400> SEQUENCE: 139	
tggtctgac tgtaccacca	20
<p><210> SEQ ID NO 140 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetically generated oligonucleotide</p>	
<400> SEQUENCE: 140	
acctggtcct ctgactgctc	20
<p><210> SEQ ID NO 141 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetically generated oligonucleotide</p>	
<400> SEQUENCE: 141	
cctcaccatc atcacactgg	20
<p><210> SEQ ID NO 142 <211> LENGTH: 19 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetically generated oligonucleotide</p>	
<400> SEQUENCE: 142	
gtgcagctgt gggttgatt	19
<p><210> SEQ ID NO 143 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetically generated oligonucleotide</p>	
<400> SEQUENCE: 143	
ggaagagaat ctccgcaaga	20
<p><210> SEQ ID NO 144 <211> LENGTH: 20 <212> TYPE: DNA</p>	

-continued

<213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 144

 atgcctccag ttcaggaaaa 20

<210> SEQ ID NO 145
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 145

 tgcctgact ttcaactctg t 21

<210> SEQ ID NO 146
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 146

 ctactgggac ggaacagctt 20

<210> SEQ ID NO 147
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 147

 gcgcacagag gaagagaatc 20

<210> SEQ ID NO 148
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 148

 agtggcctg atcttcacaa 20

<210> SEQ ID NO 149
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 149

 tttccttact gcctcttget tc 22

<210> SEQ ID NO 150
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 150

 tggccatcta caagcagtca 20

-continued

<210> SEQ ID NO 151
 <211> LENGTH: 17
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 151

 agccagtcag ccgaagg 17

<210> SEQ ID NO 152
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 152

 caccctggct ctgacat 18

<210> SEQ ID NO 153
 <211> LENGTH: 17
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 153

 gaccccgcca ctctcac 17

<210> SEQ ID NO 154
 <211> LENGTH: 24
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 154

 atatttcaat gggcatatc acag 24

<210> SEQ ID NO 155
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 155

 gccatgatcg ctcaaataca 20

<210> SEQ ID NO 156
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 156

 acttgcgagg gacgcatt 18

<210> SEQ ID NO 157
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:

-continued

<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 157

cattctgggg attcttgag 20

<210> SEQ ID NO 158
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 158

ctgtttcttc ggaggagac 20

<210> SEQ ID NO 159
 <211> LENGTH: 26
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 159

ttcctactct tcattgtact tcaacc 26

<210> SEQ ID NO 160
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 160

cgtttctgga atccaagcag 20

<210> SEQ ID NO 161
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 161

ccaggaacct ctgtgatgga 20

<210> SEQ ID NO 162
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 162

cacatgacgg aggttgtag 20

<210> SEQ ID NO 163
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 163

ttccaattcc cctgcaaa 18

-continued

<210> SEQ ID NO 164
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 164

 gggacaggta ggacctgatt t 21

<210> SEQ ID NO 165
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 165

 tgacaatggg aatgaaacag a 21

<210> SEQ ID NO 166
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 166

 ccctagaacc aaatccagca 20

<210> SEQ ID NO 167
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 167

 ctgcagttca gaggtccag 20

<210> SEQ ID NO 168
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 168

 tttgccacgg aaagtactcc 20

<210> SEQ ID NO 169
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 169

 ctcacaacct ccgtcatgtg 20

<210> SEQ ID NO 170
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

-continued

<400> SEQUENCE: 170
ctccgtcatg tgctgtgact 20

<210> SEQ ID NO 171
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 171
tgctggattt ggttctaggg 20

<210> SEQ ID NO 172
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 172
agctgtttga ggaggtggtg 20

<210> SEQ ID NO 173
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 173
gcagcttgct taggtccact 20

<210> SEQ ID NO 174
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 174
acacgcaaat ttccttccac 20

<210> SEQ ID NO 175
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 175
aaccaccctt aaccctcct 20

<210> SEQ ID NO 176
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 176
cagttgcaaa ccagacctca 20

<210> SEQ ID NO 177
<211> LENGTH: 18

-continued

<212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 177

 gcaggtcttg gccagttg 18

 <210> SEQ ID NO 178
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 178

 ccagtgtgat gatggtgagg 20

 <210> SEQ ID NO 179
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 179

 ggggacagca tcaaatcatc 20

 <210> SEQ ID NO 180
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 180

 tttttctgcc tctttctctt gg 22

 <210> SEQ ID NO 181
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 181

 ctgcacaggg caggtctt 18

 <210> SEQ ID NO 182
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 182

 gcggagattc tcttctctg 20

 <210> SEQ ID NO 183
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 183

-continued

ttcttgctct gcttgcttac c	21
<p><210> SEQ ID NO 184 <211> LENGTH: 24 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetically generated oligonucleotide</p>	
<400> SEQUENCE: 184	
tggattcaca cagacactat caca	24
<p><210> SEQ ID NO 185 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetically generated oligonucleotide</p>	
<400> SEQUENCE: 185	
cacaaacacg cacctcaaag	20
<p><210> SEQ ID NO 186 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetically generated oligonucleotide</p>	
<400> SEQUENCE: 186	
actgaccgtg caagtcacag	20
<p><210> SEQ ID NO 187 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetically generated oligonucleotide</p>	
<400> SEQUENCE: 187	
ctgctcacca tcgctatctg	20
<p><210> SEQ ID NO 188 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetically generated oligonucleotide</p>	
<400> SEQUENCE: 188	
ccaaatactc cacacgcaaa	20
<p><210> SEQ ID NO 189 <211> LENGTH: 16 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetically generated oligonucleotide</p>	
<400> SEQUENCE: 189	
gaggggctgg ctggtc	16
<p><210> SEQ ID NO 190 <211> LENGTH: 17 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence</p>	

-continued

<220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 190

 gggtcgggtg agagtgg 17

<210> SEQ ID NO 191
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 191

 cagctcctca gccaggtc 18

<210> SEQ ID NO 192
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 192

 tcaaatcagc tataaatagc aaaca 25

<210> SEQ ID NO 193
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 193

 tctcaggggg ctaaaggatt 20

<210> SEQ ID NO 194
 <211> LENGTH: 19
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 194

 gaagaagga agcgggtgac 19

<210> SEQ ID NO 195
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 195

 ggaaataaac ctctccatt ttt 23

<210> SEQ ID NO 196
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 196

 ccgcctcctt cttcatcata 20

-continued

<210> SEQ ID NO 197
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 197

tgcaacgtcg ttacgagtca 20

<210> SEQ ID NO 198
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 198

gcagctgatg ggaggaataa 20

<210> SEQ ID NO 199
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 199

tgaggatggt gtaagc gatg 20

<210> SEQ ID NO 200
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 200

gccagttggc aaaacatct 19

<210> SEQ ID NO 201
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 201

cccaacgcac cgaatagt 18

<210> SEQ ID NO 202
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 202

agctgttccg tcccagtaga 20

<210> SEQ ID NO 203
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

-continued

<400> SEQUENCE: 203
tcaccatcgc tatctgagca 20

<210> SEQ ID NO 204
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 204
ggtccttttc agaatcaata gtttt 25

<210> SEQ ID NO 205
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 205
ccaaagcatg gctcatctct a 21

<210> SEQ ID NO 206
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 206
ctcaggcaag ctgaaaacat 20

<210> SEQ ID NO 207
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 207
gtgacactgc tggaaactcg 20

<210> SEQ ID NO 208
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 208
tgtctgagca ccacttttgg 20

<210> SEQ ID NO 209
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 209
tttctgaac tggaggcatt 20

<210> SEQ ID NO 210

-continued

<211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 210

 ctgcagttca gaggtccag 20

<210> SEQ ID NO 211
 <211> LENGTH: 16
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 211

 gacagaaaca ctttat 16

<210> SEQ ID NO 212
 <211> LENGTH: 15
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 212

 acacttttcg acaag 15

<210> SEQ ID NO 213
 <211> LENGTH: 14
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 213

 cctcaacaag atac 14

<210> SEQ ID NO 214
 <211> LENGTH: 11
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 214

 gccgctgaa a 11

<210> SEQ ID NO 215
 <211> LENGTH: 12
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 215

 gcggcatgaa at 12

<210> SEQ ID NO 216
 <211> LENGTH: 13
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

 <400> SEQUENCE: 216

-continued

cccctcctca gag 13

<210> SEQ ID NO 217
 <211> LENGTH: 12
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 217

cggctcatag gg 12

<210> SEQ ID NO 218
 <211> LENGTH: 14
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 218

aaaccagacc tcag 14

<210> SEQ ID NO 219
 <211> LENGTH: 12
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 219

cagctgcaca gg 12

<210> SEQ ID NO 220
 <211> LENGTH: 13
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 220

agaccccagt tgc 13

<210> SEQ ID NO 221
 <211> LENGTH: 15
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 221

ttccagtgtg atgat 15

<210> SEQ ID NO 222
 <211> LENGTH: 15
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetically generated oligonucleotide

<400> SEQUENCE: 222

cttccactcg gataa 15

107

What is claimed is:

1. A method for characterizing a human biological sample comprising:

- (a) measuring a K-ras mutation score from a human biological sample, wherein said measuring of said K-ras mutation score is measured by digital melt curve analysis;
- (b) measuring a methylation level of a CpG site of from the human biological sample;
- (c) determining whether or not the human biological sample has an elevated K-ras mutation score and an elevated BMP3 methylation status as compared to a control from a subject that does not have premalignant or malignant colorectal cancer.

2. The method of claim **1**, wherein the human biological sample is a stool sample.

3. A method for characterizing a human stool sample comprising:

108

(a) measuring a K-ras mutation score from a human stool sample;

(b) measuring a methylation level of a CpG site of BMP3 from the human stool sample;

(c) determining whether or not the stool biological sample has an elevated K-ras mutation score and an elevated BMP3 methylation status as compared to a control from a subject that does not have premalignant or malignant colorectal cancer.

4. The method of claim **3**, wherein said measuring of said K-ras mutation score is measured by digital melt curve analysis.

5. The method of claim **3**, wherein said measuring of said K-ras mutation score is measured by quantitative allele-specific PCR.

* * * * *