



US009908908B2

(12) **United States Patent**
Zhang et al.

(10) **Patent No.:** **US 9,908,908 B2**
(45) **Date of Patent:** **Mar. 6, 2018**

(54) **TENOFOVIR PRODRUG AND
PHARMACEUTICAL USES THEREOF**

(71) Applicant: **Jiangsu Hansoh Pharmaceutical Co.,
Ltd., Lianyungang, Jiangsu (CN)**

(72) Inventors: **Fuyao Zhang, Shanghai (CN); Dong
Wei, Shanghai (CN)**

(73) Assignee: **Jiangsu Hansoh Pharmaceutical Co.,
Ltd., Lianyungang, Jiangsu (CN)**

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **14/423,182**

(22) PCT Filed: **Jul. 10, 2013**

(86) PCT No.: **PCT/CN2013/079123**

§ 371 (c)(1),

(2) Date: **Feb. 23, 2015**

(87) PCT Pub. No.: **WO2014/032481**

PCT Pub. Date: **Mar. 6, 2014**

(65) **Prior Publication Data**

US 2015/0225433 A1 Aug. 13, 2015

(30) **Foreign Application Priority Data**

Aug. 30, 2012 (CN) 2012 1 0315565

Feb. 1, 2013 (CN) 2013 1 0041647

(51) **Int. Cl.**

C07D 403/02 (2006.01)

A61K 31/4188 (2006.01)

A61K 31/4985 (2006.01)

A61K 31/675 (2006.01)

C07F 9/6561 (2006.01)

(52) **U.S. Cl.**

CPC **C07F 9/65616** (2013.01); **A61K 31/675**
(2013.01)

(58) **Field of Classification Search**

CPC **C07D 403/02; C07F 9/65616; A61K**
31/4188; A61K 31/4985; A61K 31/675

USPC **514/810; 544/244**

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,278,153 A 1/1994 Biller
5,545,750 A 8/1996 Kempf et al.
5,670,675 A 9/1997 Kempf et al.
5,717,095 A 2/1998 Arimilli et al.
5,731,432 A 3/1998 Edon et al.
5,733,788 A 3/1998 Bischofberger
5,789,394 A 8/1998 Nguyen-Ba et al.
5,798,340 A 8/1998 Bischofberger et al.
5,854,228 A 12/1998 Webb, II et al.
5,874,577 A 2/1999 Chen et al.
5,886,179 A 3/1999 Arimilli et al.

5,977,061 A 11/1999 Holy et al.
6,005,107 A 12/1999 Nguyen-Ba et al.
6,228,861 B1 5/2001 Nagarathnam et al.
6,331,543 B1 12/2001 Garvey et al.
6,455,513 B1 9/2002 McGuigan et al.
6,465,649 B1 10/2002 Gutierrez et al.
6,489,476 B1 12/2002 Dang et al.
6,653,296 B1 11/2003 Holy et al.
6,756,360 B1 6/2004 Erion et al.
7,300,924 B2 11/2007 Boojamra et al.
7,390,791 B2 6/2008 Becker et al.
7,439,350 B2 10/2008 Bischofberger et al.
7,951,787 B2 5/2011 McGuigan
8,835,630 B1 9/2014 Hostetler et al.
8,946,243 B2 2/2015 Margolis et al.
2002/0035155 A1 3/2002 Kaddurah-Daouk
2003/0050251 A1 3/2003 Semple et al.
2003/0144283 A1 7/2003 Coleman et al.
2003/0236227 A1 12/2003 Druzgala et al.
2004/0002475 A1 1/2004 Hakimelahi
2005/0080078 A1 4/2005 Aquila et al.
2005/0148623 A1 7/2005 DeGoey et al.

(Continued)

FOREIGN PATENT DOCUMENTS

CN 1425680 A 6/2003
CN 1465582 A 1/2004

(Continued)

OTHER PUBLICATIONS

Cihlar, Tomas "Amidate Prodrugs of Nucleotide Phosphonates:
From Design to In vivo Proof of Concept" Collect. Symp. Series
(2008), 10: pp. 45-53.*

(Continued)


Primary Examiner — Matthew Coughlin

Assistant Examiner — Sagar Patel

(74) *Attorney, Agent, or Firm* — Panitch Schwarze
Belisario & Nadel LLP

(57) **ABSTRACT**

The invention relates to a tenofovir prodrug and pharma-
ceutical uses thereof. In particular, the invention relates to a
compound as shown in general formula (I):



and its isomer, pharmaceutically acceptable salt, hydrate or
solvate, as well as their uses in preparing drugs for treating
viral infectious diseases, especially HIV infection, hepatitis
B, or diseases caused by hepatitis B virus, wherein the
definition of each substituent in the general formula (I) is as
defined in the specification.

16 Claims, No Drawings

(56)

References Cited

U.S. PATENT DOCUMENTS

2005/0215612 A1 9/2005 Kuo et al.
 2006/0030545 A1 2/2006 Cheng et al.
 2006/0046981 A1 3/2006 Shibata
 2009/0163712 A1 6/2009 Guo et al.
 2013/0210757 A1 8/2013 Huang et al.
 2014/0256684 A1 9/2014 Beard et al.
 2014/0256685 A1 9/2014 Beard et al.
 2014/0288025 A1 9/2014 Milne et al.
 2014/0315790 A1 10/2014 Bushell et al.

FOREIGN PATENT DOCUMENTS

CN 1523028 A 8/2004
 CN 1583769 A 2/2005
 CN 1634943 A 7/2005
 CN 1796393 A 7/2006
 CN 1810816 A 8/2006
 CN 1966514 A 5/2007
 CN 1984640 A 6/2007
 CN 1986553 A 6/2007
 CN 101003550 A 7/2007
 CN 101066980 A 11/2007
 CN 101066981 A 11/2007
 CN 101085785 A 12/2007
 CN 101089004 A 12/2007
 CN 101143879 A 3/2008
 CN 101190927 A 6/2008
 CN 101190935 A 6/2008
 CN 101230065 A 7/2008
 CN 101278938 A 10/2008
 CN 101279987 A 10/2008
 CN 101307076 A 11/2008
 CN 101418017 A 4/2009
 CN 101450954 A 6/2009
 CN 101463045 A 6/2009
 CN 101475594 A 7/2009
 CN 101531679 A 9/2009
 CN 101531680 A 9/2009
 CN 101574356 A 11/2009
 CN 101585854 A 11/2009
 CN 101591357 A 12/2009
 CN 101648974 A 2/2010
 CN 101659676 A 3/2010
 CN 101712692 A 5/2010
 CN 101759722 A 6/2010
 CN 101781334 A 7/2010
 CN 101781335 A 7/2010
 CN 101870713 A 10/2010
 CN 1947796 B 11/2010
 CN 101906119 A 12/2010
 CN 101928302 A 12/2010
 CN 101948485 A 1/2011
 CN 101985454 A 3/2011
 CN 102040626 A 5/2011
 CN 102060876 A 5/2011
 CN 102070670 A 5/2011
 CN 102093417 A 6/2011
 CN 102276667 A 12/2011
 CN 102485229 A 6/2012
 CN 103845345 A 6/2014
 CN 103923124 A 7/2014
 CN 104072539 A 10/2014
 CN 104558035 A 4/2015
 CN 104628773 A 5/2015
 EP 00206459 A2 12/1986
 EP 269947 A1 6/1988
 EP 270885 A1 6/1988
 EP 402646 A1 12/1990
 EP 481214 A1 4/1992
 EP 745893 A2 12/1996
 EP 829493 A2 3/1998
 EP 853084 A2 7/1998
 EP 989122 A1 3/2000
 EP 0996430 A2 5/2000
 EP 1065207 A1 1/2001

EP 1627902 A1 2/2006
 JP 2001031690 A 2/2001
 JP 2001031691 A 2/2001
 KR 2012025678 A 3/2012
 WO 9119721 A1 12/1991
 WO 9209611 A1 6/1992
 WO 9319075 A1 9/1993
 WO 9403467 A2 2/1994
 WO 9404143 A1 3/1994
 WO 9418200 A1 8/1994
 WO 9507920 A1 3/1995
 WO 9623801 A2 8/1996
 WO 9629336 A1 9/1996
 WO 9633200 A1 10/1996
 WO 9713507 A1 4/1997
 WO 9717969 A1 5/1997
 WO 9724361 A1 7/1997
 WO 9735855 A1 10/1997
 WO 9735990 A2 10/1997
 WO 98004569 A1 2/1998
 WO 9811121 A1 3/1998
 WO 9842758 A1 10/1998
 WO 1999/005150 A1 2/1999
 WO 9941277 A1 8/1999
 WO 9945016 A2 9/1999
 WO 9962871 A1 12/1999
 WO 2000014095 A1 3/2000
 WO 2000018775 A1 4/2000
 WO 2000038666 A2 7/2000
 WO 2000075236 A1 12/2000
 WO 2000076505 A1 12/2000
 WO 2001032636 A1 5/2001
 WO 2001049688 A1 7/2001
 WO 2001066553 A2 9/2001
 WO 2001098344 A2 12/2001
 WO 0208241 A2 1/2002
 WO 200204475 A1 1/2002
 WO 2002003978 A2 1/2002
 WO 2002008241 A2 1/2002
 WO 2002012366 A1 2/2002
 WO 2002020475 A2 3/2002
 WO 2002022572 A2 3/2002
 WO 2002094185 A2 11/2002
 WO 2002096892 12/2002
 WO 2003006628 1/2003
 WO 2003016274 A2 2/2003
 WO 2003020773 A2 3/2003
 WO 2003024943 A2 3/2003
 WO 03053989 A1 7/2003
 WO 2003087298 A2 10/2003
 WO 2003091264 A2 11/2003
 WO 2003095665 11/2003
 WO 2004024681 A2 3/2004
 WO 2004029064 A1 4/2004
 WO 2004037161 A2 5/2004
 WO 2004041167 A2 5/2004
 WO 2004064845 A1 8/2004
 WO 2004096285 A2 11/2004
 WO 2004101579 A2 11/2004
 WO 2005012324 A2 2/2005
 WO 2005016942 A1 2/2005
 WO 2005044279 A1 5/2005
 WO 2005044757 A1 5/2005
 WO 2005058330 A1 6/2005
 WO 2005066189 A1 7/2005
 WO 2005079812 A1 9/2005
 WO 2005085183 A1 9/2005
 WO 2005087788 A2 9/2005
 WO 2005123729 A1 12/2005
 WO 2006013203 A2 2/2006
 WO 2006017044 A2 2/2006
 WO 2006059357 A2 6/2006
 WO 2006102594 A1 9/2006
 WO 2006114064 A2 11/2006
 WO 2006114065 A2 11/2006
 WO 2006114492 A1 11/2006
 WO 2006130217 A2 12/2006
 WO 2006133632 A1 12/2006
 WO 2007002808 A1 1/2007

(56)

References Cited

FOREIGN PATENT DOCUMENTS

WO 2007002912 A2 1/2007
 WO 2007009236 A1 1/2007
 WO 2007013085 A1 2/2007
 WO 2007013086 A1 2/2007
 WO 2007014352 A2 2/2007
 WO 2007044565 A2 4/2007
 WO 2002057288 A1 5/2007
 WO 2007084157 A2 7/2007
 WO 2007/095269 A2 8/2007
 WO 2007101371 A1 9/2007
 WO 2007111994 A2 10/2007
 WO 2007128815 A1 11/2007
 WO 2007130783 A2 11/2007
 WO 2007134457 A1 11/2007
 WO 2007136650 A2 11/2007
 WO 2007143823 A1 12/2007
 WO 2007143824 A1 12/2007
 WO 2008005555 A1 1/2008
 WO 2008007392 A2 1/2008
 WO 2008010985 A2 1/2008
 WO 2008047345 A1 4/2008
 WO 2008056264 A2 5/2008
 WO 2008082601 A2 7/2008
 WO 2008088147 A1 7/2008
 WO 2008093173 A1 8/2008
 WO 2008096369 A2 8/2008
 WO 2008104408 A2 9/2008
 WO 2008134578 A2 11/2008
 WO 2008140302 A1 11/2008
 WO 2008143500 A1 11/2008
 WO 2008157657 A1 12/2008
 WO 2009005693 A1 1/2009
 WO 2009023718 A2 2/2009
 WO 2009/052050 A1 4/2009
 WO 2009046573 A1 4/2009
 WO 2009055289 A2 4/2009
 WO 2009064174 A1 5/2009
 WO 2009074351 A2 6/2009
 WO 2009077671 A1 6/2009
 WO 2009094190 A2 7/2009
 WO 2009105513 A2 8/2009
 WO 2009130437 A1 10/2009
 WO 2010026603 A2 3/2010
 WO 2010068708 A2 6/2010
 WO 2010081082 A2 7/2010
 WO 2010091386 A2 8/2010
 WO 2010108995 A1 9/2010
 WO 2010125200 A1 11/2010
 WO 2010142761 A1 12/2010
 WO 2011005672 A2 1/2011
 WO 2011017253 A1 2/2011
 WO 2011031628 A1 3/2011
 WO 2011038207 A1 3/2011
 WO 2011053812 A1 5/2011
 WO 2011069688 A1 6/2011
 WO 2011100698 A2 8/2011
 WO 2011111074 A2 9/2011
 WO 2011128374 A1 10/2011
 WO 2011130557 A2 10/2011
 WO 2011133963 A2 10/2011
 WO 2011150360 A1 12/2011
 WO 2012079035 A1 6/2012
 WO 2012098255 A1 7/2012
 WO 2012116050 A2 8/2012
 WO 2012117219 A1 9/2012
 WO 2012142671 A1 10/2012
 WO 2012149236 A1 11/2012
 WO 2012149567 A1 11/2012
 WO 2012154698 A2 11/2012
 WO 2013019874 A1 2/2013
 WO 2013052094 A2 4/2013
 WO 2013078277 A1 5/2013

WO 2013078288 A1 5/2013
 WO 2013090420 A2 6/2013
 WO 2013095684 A1 6/2013
 WO 2013115916 A1 8/2013
 WO 2013116720 A1 8/2013
 WO 2013116730 A1 8/2013
 WO 2013167743 A1 11/2013
 WO 2013175231 A1 11/2013
 WO 2013187978 A1 12/2013
 WO 2014032481 A1 3/2014
 WO 2014033688 A1 3/2014
 WO 2014068265 A1 5/2014
 WO 2014070771 A1 5/2014
 WO 2014088923 A1 6/2014
 WO 2014099578 A1 6/2014
 WO 2014177464 A2 11/2014
 WO 2014183462 A1 11/2014
 WO 2015040640 A2 3/2015
 WO 2015079240 A1 6/2015
 WO 2015079241 A1 6/2015

OTHER PUBLICATIONS

Lee, W., G. He, E. Eisenberg, T. Cihlar, S. Swaminathan, A. Mulato, and K. Cundy "Selective Intracellular Activation of a Novel Prodrug of the Human Immunodeficiency Virus Reverse Transcriptase Inhibitor Tenofovir leads to Preferential Distribution and Accumulation in Lymphatic Tissue" *Antimicro. Agents Chemother.* (2005), 49 (5), pp. 1898-1906.*
 Birkus, G., N. Kutty, G. He, A. Mulato, W. Lee, M. McDermott, and T. Cihlar "Activation of 9-[(R)-2-[[[(S)-[[[(S)-1-(Isopropoxycarbonyl)ethyl]amino]phenoxyphosphinyl]-methoxy]propyl]adenine (GS-7340) and Other Tenofovir Phosphonoamidate Prodrugs by Human Proteases" *Mol. Pharmacol.* (2008), 74(1), pp. 92-100.*
 Int'l Search Report dated Oct. 17, 2013 in Int'l Application No. PCT/CN2013/079123.
Journal of Medicinal Chemistry (2011), 54(16), 5680-5693 (Tyrosine-based 1-(S)-[3-hydroxy-2-(phosphonomethoxy)propyl]cytosine and -adenine ((S)-HPMPC and (S)-HPMPA) prodrugs: Synthesis, stability, antiviral activity, and in vivo transport studies) (Abstract Only).
Nucleosides, Nucleotides & Nucleic Acids (2001), 20(4-7), 621-628 (Practical synthesis, separation, and stereochemical assignment of the PMPA pro-drug GS-7340) (Abstract Only).
Bioorganic & Medicinal Chemistry Letters (2001), 11(8), 1053-1056 (Synthesis and evaluation of novel amidate prodrugs of PMEAs and PMPA) (Abstract Only).
International Journal of Antimicrobial Agents (1999), 12(1), 53-61 (Inhibition of the in vitro growth of *Plasmodium falciparum* by acyclic nucleoside phosphonates) (Abstract Only).
Drug Metabolism and Disposition (1997), 25(3), 362-366 (Pharmacokinetics and metabolism of selected prodrugs of PMEAs in rats).
Nucleosides & Nucleotides (1995), 14(3-5), 767-770 (In vivo antiretroviral efficacy of oral bis(POM)-PMEA, the bis(pivaloyloxymethyl)prodrug of 9-(2-phosphonylmethoxyethyl)adenine (PMEA)) (Abstract Only).
Journal of Medicinal Chemistry (1994), 37(12), 1857-64 (Synthesis, Oral Bioavailability Determination, and in vitro Evaluation of Prodrugs of the Antiviral Agent 9-[2-(Phosphonomethoxy)ethyl]adenine (PMEA)) (Abstract Only).
 Balzarini et al. *Federation of European Biochemical Societies*, 410 (1997) 324-328.
 Perrone et al. *J. Med. Chem.* (2007) 50, 1840-1849.
 Cahard et al. *Mini-Reviews in Medicinal Chemistry* (2004) 4, 371-381.
 Pertusati et al. *Antiviral Chem and Chemotherapy* (2012) 22, 181-203.

* cited by examiner

1

TENOFOVIR PRODRUG AND PHARMACEUTICAL USES THEREOF

CROSS-REFERENCE TO RELATED APPLICATIONS


This application is a Section 371 of International Application No. PCT/CN2013/079123, filed on Jul. 10, 2013, which was published in the Chinese language on Mar. 6, 2014, under International Publication No. WO 2014/032481 A1, and the disclosure of which is incorporated herein by reference.

FIELD OF THE INVENTION


The present invention relates to a tenofovir prodrug and a stereoisomer, pharmaceutically acceptable salt, hydrate, or solvate thereof, as well as their medical use.

BACKGROUND OF THE INVENTION


Hepatitis B virus (HBV) is a kind of DNA virus that causes human acute or chronic hepatitis. As HBV infection is a direct cause of serious liver disease including cirrhosis and hepatocellular carcinoma in humans, hepatitis B is a major threat to human health. HBV-DNA (deoxyribonucleic acid) is the core of HBV and the basis of viral replication. Nucleotides can inhibit viral polymerases by competitive binding to the natural deoxyribose substrate directly, and terminate the DNA chain by inserting DNA. Thus, nucleotides, such as Cidofovir, Adefovir, Lamivudine, and Tenofovir, are the main drugs for treating hepatitis B. Tenofovir is a novel nucleotide reverse transcriptase inhibitor, which is effective against a variety of viruses for the treatment of viral infections. As the phosphate group is a dianion at physiological pH, tenofovir has poor cell membrane permeability, low bioavailability, and dose-dependent renal toxicity, which limits its therapeutic effect. Thus, tenofovir must be prepared into a phosphonate prodrug form via various technical means, such as esterification and salification, for clinical applications. For example, Tenofovir disoproxil fumarate developed by Gilead Sciences Inc. is the first generation oral active tenofovir prodrug for the treatment of HIV infection and hepatitis B.



Tenofovir



2



Tenofovir disoproxil fumarate


As Tenofovir disoproxil fumarate is highly sensitive to serum enzyme mediated hydrolysis reactions, its drug concentration cannot be effectively increased at the active site. Moreover, two equivalents of potentially toxic formaldehyde are released upon metabolism, and side effects such as lactic acidosis, severe hepatomegaly and lipodystrophy have been found during clinical use. In order to improve the stability of tenofovir prodrug in plasma and reduce the metabolite-tenofovir concentration to reduce drug toxicity, many pharmaceutical companies are conducting research and development on the next generation of tenofovir prodrugs, and have made some progress. Some new prodrugs have been in phase I/II clinical studies. For example, International Patent Application WO0208241 discloses a kind of natural amino acid (monosubstituted) synthesized tenofovir phosphonamidate prodrug (e.g., GS-7340), and International Patent Application WO2009105513 discloses a kind of novel tenofovir phosphate bisamide prodrug. Compared to tenofovir phosphodiester, these novel prodrugs have improved plasma stability, thereby increasing the cumulative concentration of the active metabolite-tenofovir in peripheral blood mononuclear cells (PBMCs) and the therapeutic effect. For example, the total concentration of the active ingredient produced by GS-7340 in PBMCs is 10 times greater than tenofovir disoproxil, and 30 times greater than tenofovir. However, GS-7340 has certain degradation in plasma and 1-2% of the metabolite-tenofovir can be detected in plasma. So it is inevitable that GS-7340 has toxicity as a side effect generated by tenofovir disoproxil, which results in drug safety problems. Thus, it is important to further develop a tenofovir prodrug with high efficacy and low toxicity. With the purpose of further improving the stability of tenofovir disoproxil in plasma, the present invention synthesizes a series of tenofovir phosphonamidate prodrugs with disubstituted amino acids. This kind of prodrug is proven to be very stable in plasma and no metabolite-tenofovir is found in plasma. On the other hand, the concentration of active metabolite-tenofovir in PBMCs is increased significantly compared with GS-7340. Thus, the present invention makes it possible to provide a new generation of tenofovir prodrugs with high efficacy and low toxicity.

SUMMARY OF THE INVENTION

Surprisingly, the inventors found a series of compounds, which have higher efficacy and lower toxicity than the prior art. Compared to GS-7340, the compounds according to the present invention are stable enough in plasma, and the metabolite-tenofovir is completely undetectable in plasma. On the other hand, the concentration of tenofovir is greatly improved in PBMCs. Such a result is totally unexpected for those skilled in the art.

The present invention relates to a compound of general formula (I) and a stereoisomer, pharmaceutically acceptable salt, hydrate, or solvate thereof:

3



wherein:

R^1 and R^2 are C_{1-6} alkyl respectively, or R^1 and R^2 together with the attached carbon atom form a C_{3-7} cycloalkyl;

R^3 is hydrogen, C_{1-6} alkyl, substituted or unsubstituted C_{6-10} aryl or 6- to 10-membered heteroaryl;



Ar is substituted or unsubstituted C_{6-10} aryl or 6- to 10-membered heteroaryl.

The compound of general formula (I) according to the present invention can be used as a prodrug of tenofovir. This prodrug is stable in plasma and the concentration of active metabolite-tenofovir in PBMCs is improved significantly compared to that of GS-7340.

In the compound of general formula (I) according to the present invention, the phosphorus atom is chiral and the configuration is S or R, or a mixture of R and S.

In an embodiment of the present invention, the compound of general formula (I) or the stereoisomer, pharmaceutically acceptable salt, hydrate or solvate thereof, wherein the stereoisomer comprises a tautomer, cis-trans isomer, conformational isomer, mesomer or enantiomeric or diastereomeric optical isomer.

In a preferred embodiment of the present invention, compounds having the following structures are provided, but the compound of general formula (I) according to the present invention is not limited to the following structures:




4

-continued

(I)


5



10


15

20




25

30



35

40




45

50

55

Ib

60



65

Ic

Id


Ie

If


Ig

5


-continued




5h



5i




5j



5k


In another preferred embodiment of the present invention, chiral compounds with the following structures are disclosed, but the compound of general formula (I) according to the present invention is not limited to the following structures:




5l

6


-continued




6h




6i



6j



6k



6l

Ib1

Ic1


Id1

Ie1


If1

7


-continued




Ig1




Ih1



Ii1




Ij1



Ik1

The compound of general formula (I) according to the present invention can be prepared according to the following method:

8



5

10

(II)

15

20

25

30

wherein the compound of general formula (II) can be prepared from tenofovir according to the method of Chinese Patent ZL01813161.1 as well as other conventional methods in the art.

Chiral isomer (II) can be separated from a mixture of isomers (I') with a reverse phase column or chiral column.

35

40

45

50

55

60

65

70

75

80

85

90

95

100

105

110

115

120

125

130

135

140

145

150

155

160

165

170

175

180

185

190

195

200

205

210

215

220

225

230

235

240

245

250

255

260

265

270

275

280

285

290

295

300

305

310

315

320

325

330

335

340

345

350

355

360

365

370

375

380

385

390

395

400

405

410

415

420

425

430

435

440

445

450

455

460

465

470

475

480

485

490

495

500

505

510

515

520

525

530

535

540

545

550

555

560

565

570

575

580

585

590

595

600

605

610

615

620

625

630

635

640

645

650

655

660

665

670

675

680

685

690

695

700

705

710

715

720

725

730

735

740

745

750

755

760

765

770

775

780

785

790

795

800

805

810

815

820

825

830

835

840

845

850

855

860

865

870

875

880

885

890

895

900

905

910

915

920

925

930

935

940

945

950

955

960

965

970

975

980

985

990

995

1000

1005

1010

1015

1020

1025

1030

1035

1040

1045

1050

1055

1060

1065

1070

1075

1080

1085

1090

1095

1100

1105

1110

1115

1120

1125

1130

1135

1140

1145

1150

1155

1160

1165

1170

1175

1180

1185

1190

1195

1200

1205

1210

1215

1220

1225

1230

1235

1240

1245

1250

1255

1260

1265

1270

1275

1280

1285

1290

1295

1300

1305

1310

1315

1320

1325

1330

1335

1340

1345

1350

1355

1360

1365

1370

1375

1380

1385

1390

1395

1400

1405

1410

1415

1420

1425

1430

1435

1440

1445

1450

1455

1460

1465

1470

1475

1480

1485

1490

1495

1500

1505

1510

1515

1520

1525

1530

1535

1540

1545

1550

1555

1560

1565

1570

1575

1580

1585

1590

1595

1600

1605

1610

1615

1620

Unless otherwise stated, the terms in the present invention have the following meanings.

The term "alkyl" means saturated aliphatic hydrocarbon groups comprising a straight or branched chain having 1 to 6 carbon atoms. Examples of alkyl include, but are not limited to, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, t-butyl, sec-butyl, n-pentyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 2,2-dimethylpropyl, 1-ethylpropyl, 2-methylbutyl, 3-methylbutyl, n-hexyl, 1-ethyl-2-methylpropyl, 1,1,2-trimethylpropyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 2,2-dimethylbutyl, 1,3-dimethylbutyl, 2-ethylbutyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 2,3-dimethylbutyl and the like. Alkyl can be substituted or unsubstituted. When substituted, the substituent can be substituted at any possible attachment point, and the substituent is preferably one or more groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, halogen, thiol, hydroxyl, nitro, cyano, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, cycloalkoxy, heterocycloalkoxy, cycloalkylthio, heterocycloalkylthio, and oxo.

The term "cycloalkyl" means a saturated or partially unsaturated monocycle or polycycle hydrocarbon substituent, which comprises 3 to 7 carbon atoms. Examples of monocyclic cycloalkyl include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, cyclohexadienyl, cycloheptyl, cycloheptatrienyl and the like. Examples of polycyclic cycloalkyl include, but are not limited to, spiro-ring cycloalkyl, fused-ring cycloalkyl and bridged-ring cycloalkyl. The cycloalkyl can be substituted or unsubstituted. When substituted, the substituent is preferably one or more groups independently selected from the group consisting of alkyl, alkoxy, halogen, hydroxy, nitro, cyano, cycloalkyl, heterocycloalkyl, aryl, and heteroaryl.

The term "aryl" means a 6 to 10 membered full-carbon monocycle or fused polycycle (i.e., rings which share adjacent pairs of carbon atoms) and polycycle having a conjugated π electron system (i.e., rings having adjacent pairs of carbon atoms), such as phenyl and naphthyl. The aryl can be substituted or unsubstituted. When substituted, the substituent is preferably one or more groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, halogen, thiol, hydroxyl, nitro, cyano, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, cycloalkoxy, heterocycloalkoxy, cycloalkylthio, and heterocycloalkylthio.

The term "heteroaryl" means a heteroaromatic system of 6 to 10 ring atoms, preferably 5 to 6 ring atoms, which comprises one, two, three or four heteroatoms including O, S or N, such as pyridinyl, or pyrimidinyl. "Heteroaryl" can be optionally substituted or unsubstituted. When substituted, the substituent is preferably one or more groups independently selected from the group consisting of alkyl, alkenyl, alkynyl, alkoxy, alkylthio, alkylamino, halogen, thiol, hydroxyl, nitro, cyano, cycloalkyl, heterocycloalkyl, aryl, heteroaryl, cycloalkoxy, heterocycloalkoxy, cycloalkylthio, and heterocycloalkylthio.

DETAILED DESCRIPTION OF THE INVENTION

The present invention will be illustrated by the following examples which enable those skilled in the art to understand the present invention more clearly. The examples are merely


for illustrating the technical solutions of the present invention and should not be considered as limiting the scope of the invention.

Example 1

Step 1:


Trimethylchlorosilane (6.0 g) was added dropwise to a solution of phenol (5 g) and triethylamine (10.1 mL) in dichloromethane (150 mL) at 0° C. After addition, the reaction mixture was stirred for 18 hours after the temperature was raised to 20° C. The white solid was removed and washed with dichloromethane. The filtrate was combined and the solvent was evaporated to give phenoxy trimethylsilane (4.2 g) as a colorless oil.

Step 2:




DMF (0.1 mL) and dichlorosulfoxide (0.73 g) were added to a suspension of tenofovir (1 g, purchased from Suzhou Henderson Pharmaceutical Co., Ltd.) in sulfolane (2.5 mL) at 70° C., and then the temperature was raised to 100° C. The reaction mixture was stirred at 100° C. for 1.5 hours until a clear solution was obtained. Then, phenoxy trimethylsilane (0.70 g) was added rapidly, and the mixture was left to stir at 100° C. for another 1.5 hours. Then, the solvent was evaporated to give a viscous yellow oil. The oil was dissolved in methanol and adjusted to pH 3 with 45% aqueous potassium hydroxide. The precipitate was filtered and dried to give a white powder solid IIa (0.7 g). MS (m/z) 363.96 (MH+).

Step 3:



11

-continued



Ia

DMF (0.1 mL) and dichlorosulfoxide (343 mg) were added to a mixture of compound IIa (600 mg) in sulfolane (1 mL) at 60° C. The mixture was then stirred at 60° C. for 30 minutes until a clear solution was obtained. The resulting solution was added to a solution of amino acid ester IIIa (750 mg, purchased from Shanghai Darui Fine Chemicals Co., Ltd.) and diisopropylamine (452 mg) in dichloromethane (7 mL) at 0° C. The mixture was stirred at 20° C. for 2 hour, and then washed with 5% aqueous sodium dihydrogen phosphate and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated to give a yellow oil crude product, which was purified via column chromatography to give an oil product Ia (150 mg).

¹H-NMR (400 MHz, CDCl₃) δ 8.34 (m, 1H), 8.05 (m, 1H), 7.36~6.95 (m, 5H), 6.49 (b, 2H), 6.22~5.84 (m, 1H), 5.01 (m, 1H), 4.42 (m, 1H), 4.40~3.60 (m, 3H), 1.52~1.18 (m, 15H). MS (m/z) 491.13 (MH⁺).


The Preparation of Chiral Compounds Ia1 and Ia2:

Method 1: Non-Chiral Column Preparation

The crude product Ia (150 mg) was separated via preparative HPLC (preparative column: Waters Symmetry C18, Mobile phase: A: 0.02% aqueous phosphoric acid; B: methanol) to give compound Ia1 (50 mg, retention time: 50.65 min): MS (m/z) 491.17 (MH⁺) and compound Ia2 (61 mg, retention time: 47.57 min): MS (m/z) 491.10 (MH⁺).

Method 2: Chiral Column Preparation

The crude product Ia (150 mg) was separated via preparative HPLC (preparative column: Chiralpak AS-H, mobile phase: A: n-hexane; B: ethanol) to give compound Ia1 (62 mg, retention time: 6.53 min) and compound Ia2 (78 mg, retention time: 6.11 min).




Ia1

12


-continued

Ia2



15

Example 2



20

25

30

35

40


45

50

55

60

65




Ib

DMF (0.1 mL) and dichlorosulfoxide (343 mg) were added to a mixture of compound IIa (600 mg) in sulfolane (1 mL) at 60° C. The mixture was then stirred at 60° C. for 30 minutes until a clear solution was obtained. The resulting solution was added to a solution of amino acid ester IIIb (760 mg, purchased from Shanghai Darui Fine Chemicals Co., Ltd.) and diisopropylamine (452 mg) in dichloromethane (7 mL) at 0° C. The mixture was stirred at 20° C. for 2 hour, and then washed with 5% aqueous sodium dihydrogen phosphate and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated to give a yellow oil crude product, which was purified via column chromatography to give an oil product Ib (221 mg).

¹H-NMR (400 MHz, CDCl₃) δ 8.38 (m, 1H), 8.01 (m, 1H), 7.34~6.95 (m, 5H), 6.48~6.18 (m, 1H), 5.84 (b, 2H), 5.01~4.82 (m, 1H), 4.42 (m, 1H), 4.2~03.60 (m, 5H), 2.68 (m, 1H), 1.41~1.10 (m, 12H).

The crude product Ib (100 mg) was separated via preparative HPLC (preparative column: Chiralpak AS-H, mobile phase: A: n-hexane; B: ethanol) to give compound Ib1 (35 mg). MS (m/z) 489.26 (MH⁺).

13




Example 3

Step 1:

Trimethylchlorosilane (6.3 g) was added dropwise to a solution of p-chlorophenol (5 g) and triethylamine (10.8 mL) in dichloromethane (150 mL) at 0° C. After addition, the reaction mixture was stirred for 18 hours after the temperature was raised to 20° C. The solvent was evaporated to give p-chlorophenoxy trimethylsilane (5.1 g) as a colorless oil.


Step 2:



DMF (0.1 mL) and dichlorosulfoxide (0.73 g) were added to a suspension of tenofovir (1 g) in sulfolane (2.5 mL) at 70° C., and then the temperature was raised to 100° C. The reaction mixture was stirred at 100° C. for 1.5 hours until a clear solution was obtained. Then, p-chlorophenoxy trimethylsilane (0.77 g) was added rapidly, and the mixture was left to stir at 100° C. for 1.5 hours. The solvent was evaporated to give a viscous yellow oil. The oil was dissolved in methanol and then adjusted to pH 3 with 45% aqueous potassium hydroxide. The precipitate was filtered and dried to give a white powder solid IIc (800 mg). MS (m/z) 398.05 (MH⁺).

14

Step 3:




DMF (0.1 mL) and dichlorosulfoxide (343 mg) were added to a mixture of compound IIc (600 mg) in sulfolane (1 mL) at 60° C. The mixture was then stirred at 60° C. for 30 minutes until a clear solution was obtained. The resulting solution was added to a solution of amino acid ester IIIa (731 mg) and diisopropylamine (452 mg) in dichloromethane (7 mL) at 0° C. The mixture was stirred at 20° C. for 2 hours, and then washed with 5% aqueous sodium dihydrogen phosphate and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated to give a yellow oil crude product which was purified via column chromatography to give an oil product Ic (121 mg).

¹H-NMR (400 MHz, CDCl₃) δ 8.35 (m, 1H), 8.01 (m, 1H), 7.28 (m, 1H), 7.22 (m, 1H), 7.15~7.13 (m, 1H), 6.94 (m, 1H), 5.88 (b, 2H), 5.07 (m, 2H), 4.42 (m, 1H), 4.21 (m, 1H), 3.90~3.81 (m, 2H), 3.71~3.54 (m, 1H), 1.56~1.24 (m, 15H).

The crude product Ic (70 mg) was separated by preparative HPLC (preparative column: Chiralpak AS-H, mobile phase: A: n-hexane; B: ethanol) to give compound Ic1 (21 mg). MS (m/z) 525.26 (MH⁺).

Ic1



Example 4


Step 1:

Trimethylchlorosilane (6.3 g) was added dropwise to a solution of p-methoxyphenol (5 g) and triethylamine (10.8 mL) in dichloromethane (150 mL) at 0° C. After addition,

15


the reaction mixture was stirred for 18 hours after the temperature was raised to 20° C. The solvent was evaporated to give p-methoxyphenoxy trimethylsilane (4.7 g) as a colorless oil.

Step 2:



DMF (0.1 mL) and dichlorosulfoxide (0.73 g) were added to a suspension of tenofovir (1 g) in sulfolane (2.5 mL) at 70° C., and then the temperature was raised to 100° C. The reaction mixture was stirred at 100° C. for 1.5 hours until a clear solution was obtained. Then, p-methoxyphenoxy trimethylsilane (0.75 g) was added rapidly and the mixture was left to stir at 100° C. for 1.5 hours. The solvent was evaporated to give a viscous yellow oil. The oil was dissolved in methanol and then adjusted to pH 3 with 45% aqueous potassium hydroxide. The precipitate was filtered and dried to give a white powder solid IId (600 mg). MS (m/z) 394.11 (MH⁺).

Step 3:




16

DMF (0.1 mL) and dichlorosulfoxide (181 mg, 1.52 mmol) were added to a mixture of compound IId (300 mg) in sulfolane (1 mL) at 60° C. The mixture was then stirred at 60° C. for 30 minutes until a clear solution was obtained. The resulting solution was added to a solution of amino acid ester IIIa (386 mg) and diisopropylamine (343 mg) in dichloromethane (5 mL) at 0° C. The mixture was stirred at 20° C. for 2 hours, and then washed with 5% aqueous sodium dihydrogen phosphate and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated to give a yellow oil crude product, and then purified via column chromatography to give an oil product Id (40 mg). ¹H-NMR (400 MHz, CDCl₃) δ 8.35 (m, 1H), 8.04 (m, 1H), 7.12~6.85 (m, 4H), 5.86 (b, 2H), 5.06 (m, 1H), 4.42 (m, 1H), 4.18 (m, 1H), 4.08~3.94 (m, 3H), 3.82 (m, 3H), 3.7~73.61 (m, 1H), 1.55~1.17 (m, 15H).

The crude product Id (30 mg) was separated by preparative HPLC (preparative column: Chiralpak AS-H, mobile phase: A: n-hexane; B: ethanol) to give 12 mg compound Id1. MS (m/z) 521.23 (MH⁺).


Id1



Example 5


Compounds Ie and Ie1 were prepared according to the same preparation method as that of compounds Ic and Ic1.

Ie



17

-continued





Ie: $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 8.27 (m, 1H), 8.04 (s, 1H), 7.96 (m, 1H), 7.84 (m, 1H), 7.62 (m, 1H), 7.52~7.33 (m, 4H), 5.78 (b, 2H), 5.04~4.98 (m, 1H), 4.38~3.71 (m, 6H), 1.57~1.06 (m, 15H).

Ie1: MS (m/z) 541.11.

Example 6

Compounds If and If1 were prepared according to the same preparation method as that of compounds Ic and Ic1.



If: $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 8.33 (m, 1H), 8.02 (s, 1H), 7.81~7.66 (m, 4H), 7.49~7.41 (m, 2H), 7.31~7.06 (m, 1H), 5.72 (b, 2H), 5.06~4.99 (m, 1H), 4.43~4.35 (m, 1H), 4.19~3.91 (m, 4H), 3.74~3.65 (m, 1H), 1.57~1.20 (m, 15H).


If1: MS (m/z) 541.10.

Example 7

Compounds Ih and Ih1 were prepared according to the same preparation method as that of compounds Ic and Ic1.

18

Ie1




Ih

5

10

15

20



Ih1

25

Ih: $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 8.33 (m, 1H), 7.95 (m, 1H), 7.00~6.95 (m, 3H), 5.83 (b, 2H), 5.05~4.99 (m, 2H), 4.35~4.31 (m, 1H), 4.23~4.17 (m, 1H), 4.01~3.83 (m, 3H), 3.80~3.77 (m, 1H), 2.35 (s, 3H), 2.31 (s, 3H), 1.33~1.19 (m, 15H).



30

Ih1: MS (m/z) 519.15.

If

Example 8

Compounds Ii and Ii1 were prepared according to the same preparation method as that of compounds Ic and Ic1.



If: $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 8.33 (m, 1H), 8.02 (s, 1H), 7.81~7.66 (m, 4H), 7.49~7.41 (m, 2H), 7.31~7.06 (m, 1H), 5.72 (b, 2H), 5.06~4.99 (m, 1H), 4.43~4.35 (m, 1H), 4.19~3.91 (m, 4H), 3.74~3.65 (m, 1H), 1.57~1.20 (m, 15H).

If1: MS (m/z) 541.10.


Example 7

Compounds Ih and Ih1 were prepared according to the same preparation method as that of compounds Ic and Ic1.

If

Example 8

Compounds Ii and Ii1 were prepared according to the same preparation method as that of compounds Ic and Ic1.




Ii

40

45

50

55



Ii1

65


Ii: $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 8.36 (m, 1H), 8.00 (m, 1H), 7.17 (m, 1H), 7.02 (m, 1H), 6.97 (m, 2H), 5.71 (b, 2H), 5.06 (m, 1H), 4.43 (m, 1H), 4.20 (m, 1H), 4.06~3.84 (m, 3H), 3.72~3.61 (m, 1H), 1.56~1.22 (m, 15H).

Ii1: MS (m/z) 509.25.

19

Example 9

Compounds Ij and Ij1 were prepared according to the same preparation method as that of compounds Ic and Ic1.



Ij: ¹H-NMR (400 MHz, CDCl₃) δ 8.32 (m, 1H), 8.06 (s, 1H), 7.58 (m, 2H), 7.52 (m, 2H), 5.89 (b, 2H), 5.02~4.96 (m, 1H), 4.43~4.36 (m, 2H), 4.04~3.91 (m, 4H), 1.58~1.23 (m, 15H).

Ij1: MS (m/z) 559.08.

Example 10 Preparation of Fumarate of Compound Ia1

Compound Ia1 (480 mg), fumaric acid (120 mg) and acetonitrile were added sequentially to a single-necked flask at 20° C. The mixture was warmed to 60° C. and stirred at this temperature until the solid was completely dissolved. Stirring was continued for another 5 minutes, and then the solution was cooled to 20° C. and filtered to obtain the fumarate of compound Ia1 as a white granular solid (490 mg).

¹H NMR (400 MHz, D₂O) δ 7.21 (m, 2H), 7.11 (m, 1H), 6.67 (m, 2H), 6.57 (s, 2H), 4.77 (m, 1H), 4.29 (m, 1H), 4.17 (m, 1H), 4.06 (m, 1H), 3.93 (m, 1H), 1.07 (m, 6H), 1.21 (m, 9H).

Example 11 Antiviral Experiment

1. Study of In Vitro Anti-Hepatitis B Virus Activity

HepG 2.2.15 cell was used as the hepatitis B virus (HBV) vehicle to determine the inhibition effect of the compounds on DNA-replication of HBV.

Test method: HepG 2.2.15 cells were seeded into a 96-well culture plate. Various dilutions of test samples and positive control were added, respectively, after 24 hours, in which a cell control well was set. The medium was replaced with culture containing various dilutions of test samples after 72 hours. The supernatants and HepG 2.2.15 cells were collected after 6 days of culture. The HBV DNA-replication was tested via the dot blot method, and IC₅₀ was calculated (the results are shown in Table 1).

20

2. Cytotoxicity Test

Test method: HepG 2.2.15 cells were seeded into a 96-well culture plate and various dilutions of test samples and positive control were added, respectively. CellTiter-Blue (Promega, Catalog #G8081) was added for 6 days of culture. Fluorescence reading was counted with Flexstation 3 to calculate CC₅₀ (the results were shown in Table 1).

TABLE 1

HBV inhibition rate and cytotoxicity results of the compounds		
Compound	IC ₅₀ (nM)	Toxicity CC ₅₀ (nM)
Positive control GS-7171(Mixture)	35	>10000
Positive control GS-7340 (Chiral)	14	>10000
Ia (Mixture)	21	>10000
Ia1 (Chiral)	5.0	>10000
Ia2 (Chiral)	32	>10000
Ib1 (Chiral)	15	>10000
Ic1 (Chiral)	5.3	>10000
Id1 (Chiral)	6.2	>10000
Ie1 (Chiral)	5.7	>10000
If1 (Chiral)	7.5	>10000
Ih1 (Chiral)	5.9	>10000
Ii1 (Chiral)	8.3	>10000
Ij1 (Chiral)	7.0	>10000

Positive control is GS-7171 and GS-7340, which are disclosed in Examples 2 and 3 of Chinese Patent ZL01813161.1. GS-7171 can be resolved into diastereomers GS-7340 and GS-7339, in which GS7340 has the better efficacy.

3. Conclusion

Experimental results show that the compounds Ia1, Ib1, Ic1, Id1, Ie1, If1, Ih1, and Ij1 have a significant inhibitory effect on HBV-DNA replication without cytotoxicity, in which the inhibitory effects on HBV-DNA replication of compounds Ia1, Ic1, Id1, Ie1, If1, Ih1, Ii1 and Ij1 were better than that of the positive control GS7340.

Example 12 Stability Test in Acidic Medium and Simulated Gastric Juice

1. Materials, Reagents and Manufacturers

Name	Content	Manufacturer
Pepsin	1:3000	Shanghai Runjie Chemical Reagent Co., Ltd.
HCl	36%	Jiangsu Qiangsheng Chemical Co., Ltd.
Ammonium acetate	98.0%	Sinopharm Chemical Reagent Co., Ltd.
GS-7340	98.7%	UniTris Biopharma Co. Ltd.
Ia1	98.7%	UniTris Biopharma Co. Ltd.

2. Preparation of Reagents

2.1 Hydrochloric Acid Solution (pH 2.0)

4.5 mL of 36% hydrochloric acid was transferred into a 1 L volumetric flask and diluted with water to 1 L to prepare a stock solution. Then 10 mL of the above solution was transferred into a 50 mL volumetric flask and diluted with water to 50 mL to prepare the hydrochloric acid solution with pH 2.0.

2.2 Simulated Gastric Juice (pH 2.0)

10 ml of the stock solution and 500.0 mg of pepsin were transferred into a 50 mL volumetric flask and diluted with water to 50 ml, which was then subjected to ultrasound to dissolve the pepsin (the solution was not clear now) and then filtered to give a clear solution as simulated gastric juice.

21

2.3 Preparation of the Sample Solution

2.3.1 Hydrochloric Acid Solution of GS-7340

5.0 mg of GS-7340 were transferred into a 5 mL volumetric flask and mixed with 2.5 mL of isopropyl alcohol to dissolve GS-7340, and then the hydrochloric acid solution (pH 2.0) was added to 5 mL. The solution was shaken well and filtered for using.

2.3.2 Simulated Gastric Juice of GS-7340

5.0 mg of GS-7340 were transferred into a 5 mL volumetric flask and mixed with 2.5 mL of isopropyl alcohol to dissolve GS-7340, and then the simulated gastric juice was added to 5 mL. The solution was shaken well and filtered for using.

2.3.3 Hydrochloric Acid Solution of Compound Ia1

5.0 mg of compound Ia1 were transferred into a 5 mL volumetric flask and mixed with 2.5 mL of isopropyl alcohol to dissolve compound Ia1, and then the hydrochloric acid solution (pH 2.0) was added to 5 mL. The solution was shaken well and filtered for using.

2.3.4 Simulated Gastric Juice of Compound Ia1

5.0 mg of compound Ia1 were transferred into a 5 mL volumetric flask and mixed with 2.5 mL of isopropyl alcohol to dissolve compound Ia1, and then the simulated gastric juice was added to 5 mL. The solution was shaken well and filtered for using.

2.4 Sampling

The prepared sample was filled into the chromatography vial as the initial sample and immediately injected. Meanwhile, the rest of the samples were put into a 37° C. incubator, and were injected into the HPLC system after 6 hours.

The stability results of compound Ia1 and GS-7340 in an acid medium and simulated gastric juice are shown in Table 2.

TABLE 2

The stability results of compound Ia1 and GS-7340 in an acid medium and simulated gastric juice						
Sample	Acid medium (pH = 2)			Simulated gastric juice		
	Initial purity	Purity after 6 hours	Change value	Initial purity	Purity after 6 hours	Change value
GS-7340	98.7%	95.5%	3.2%	98.7%	94.4%	4.3%
Ia1	98.7%	98.1%	0.6%	98.7%	98.2%	0.5%

3. Conclusion

Experimental results show that the stability of tenofovir phosphoramidate prodrug (Ia1) having a disubstituted amino acid is significantly improved compared to tenofovir phosphoramidate prodrug (GS-7340) having a monosubstituted amino acid.

Example 13: Metabolic Stability in Fresh Human Whole Blood and Distribution Test in PBMCs of Tenofovir Prodrug

1. Materials

Compound: Ia1 and GS-7340

2. Test Method

Different tenofovir prodrugs were incubated together with fresh human whole blood at 37° C. Plasma and PBMCs were separated, respectively, from the whole blood (Ficoll density gradient centrifugation method) after 1 hour and 2 hours incubation to determine the concentrations of prototype drug and metabolite-tenofovir in plasma and PBMCs. The

22

PBMCs were counted via cell counter and each PBMC was treated as 200 fL to calculate intracellular drug concentration.

3. Plasma/PBMC Sample Treatment

20 µL of internal standard solution (400 ng/mL SN-38 solution), 5.04, methanol-water (50:50, v/v), and 200 µL acetonitrile were added to 100 µL plasma sample or PBMC sample, respectively. The mixture was mixed by vortex for 1 min and centrifuged for 5 min (14000 rpm). 20 µL supernatant and 180 µL mobile phase were mixed by vortex for 1 min, and 10 µL of the above mixture was injected into LC/MS/MS for analysis.

The results of the metabolic stability in fresh human whole blood and the distribution test in PBMCs of tenofovir prodrug are shown in Table 3.

TABLE 3

The results of the metabolic stability in fresh human whole blood and the distribution test in PBMCs cells of tenofovir prodrug					
Incubation duration	Analyte	GS-7340		Ia1	
		Prototype concentration (µM)	PAMA* concentration (µM)	Prototype concentration (µM)	PAMA* concentration (µM)
0 hour	Initial concentration	16.9	0	12.4	0
1 hour	In Plasma	14.5	0.71	12.3	0
	In PBMCs	2448	199	1528	320
2 hour	In Plasma	13.0	1.37	12.2	0
	In PBMCs	2357	207	1282	546

*PAMA is the active metabolite-tenofovir of prodrug

4. Conclusion

As can be seen from Table 3, certain active metabolite-tenofovir was detected in plasma for the positive control GS-7340 after incubation with fresh human whole blood, and the active metabolite-tenofovir released in plasma increased significantly over the incubation time. However, no active metabolite-tenofovir was detected in plasma for the compound Ia1 of the present invention after incubation with fresh human whole blood, and the active metabolite-tenofovir was not always detected over the incubation time, which illustrates that the stability of compound Ia1 in plasma was significantly better than the positive control GS-7340. Therefore, compound Ia1 of the present invention has significant advantages in reducing toxic side effects resulting from tenofovir in plasma compared with the positive control GS-7340.

It also can be seen from Table 3 that the concentration of the active metabolite-tenofovir in peripheral blood mononuclear cells (PBMCs) for compound Ia1 of the present invention increased significantly over the incubation time, while the concentration of the active metabolite-tenofovir in peripheral blood mononuclear cells (PBMCs) for compound GS-7340 was almost the same. The concentration of active metabolite-tenofovir in PBMCs for compound Ia1 was about three times of that for the positive control of GS-7340 after incubation for 2 hours. Thus, compound Ia1 of the present invention has significant advantages in terms of therapeutic effect compared with the positive control GS-7340.

Example 14: Anti-AIDS (HIV) Virus Test

1. Objective: To evaluate anti-HIV activity and cytotoxicity, and EC50 and CC50 value of three compounds.

2. Materials and Method

2.1 Materials

Compound: Compound Ia1, GS-7340, and tenofovir disoproxil

RPMI medium (Invitrogen 21969-035)

DMEM medium (Invitrogen 21969-035)

Glutamic acid 200 mM (Invitrogen 25030)

Fetal bovine serum ((Invitrogen 16000-044)

Penicillin/streptomycin ((Invitrogen 15140-122)

DPBS buffer (Invitrogen 14190-094)

Trypsin-EDTA (Invitrogen 25200)

Trypan blue Sigma T8154

DMSO Sigma D2650

MUG Biochemika 69590

2.2 Test Method

1) MT-2 cells were infected with HIV-1 (IIIb) to form multiplicity of infection (MOI) 0.01 TCID₅₀ per cell.

2) The mixture of virus and cells was incubated in a 384-well plate for 3 days.

3) Cells for the cytotoxic detection were incubated in a 384-well plate for 3 days.

4) The supernatant was transferred to a new plate and incubated with reporter cells (HeLa) for 24 hours.

5) Detecting beta-GAL activity to evaluate anti-HIV activity.

6) Luminescent signal of the cells free of virus was detected to evaluate cytotoxicity after incubation for 3 days.

7) The antiviral activity and cytotoxicity were calculated according to the following equation:

$$\text{Antiviral activity (\%)} = 100 - \frac{(\text{Detection value} - \text{Maximum value})}{(\text{Minimum value} - \text{Maximum value})} * 100$$

$$\text{Cytotoxicity (\%)} = 100 - \frac{(\text{Detection value} - \text{Maximum value})}{(\text{Minimum value} - \text{Maximum value})} * 100$$

8) EC₅₀ and CC₅₀ were calculated with Fit Curve of Graphpad Prism 5 (the results are shown in Table 4).

TABLE 4

The results of anti-HIV activity and cytotoxicity		
Compounds	EC ₅₀ (nM)	Toxicity CC ₅₀ (mM)
Positive control (GS-7340)	12	>150
Positive control (Tenofovir Disoproxil)	11	16.9
Ia1 (chiral)	8	>150


3. Conclusion

The above results show that compound Ia1 has a strong inhibitory effect on HIV virus with no cytotoxicity.

The present invention has been described and illustrated by specific embodiments. Certain modifications and equivalent variations are apparent for those skilled in this art and are included within the scope of the present invention.

The invention claimed is:

1. A chiral compound of formula (I) or a stereoisomer, pharmaceutically acceptable salt, hydrate, or solvate thereof:



wherein:




R¹ and R² are each methyl, or R¹ and R² together with the attached carbon atom form a C₃ cycloalkyl;

R³ is hydrogen, C₁₋₆ alkyl, substituted or unsubstituted C₆₋₁₀ aryl, or 6- to 10-membered heteroaryl; and

Ar is substituted or unsubstituted C₆₋₁₀ aryl, or 6- to 10-membered heteroaryl.






2. The chiral compound of formula (I) according to claim 1, or the stereoisomer, pharmaceutically acceptable salt, hydrate, or solvate thereof, wherein the stereoisomer is selected from the group consisting of a tautomer, cis-trans isomer, conformational isomer, mesomer, an enantiomeric optical isomer, and a diastereomeric optical isomer.

3. The chiral compound of formula (I) according to claim 1, or the pharmaceutically acceptable salt thereof, wherein the compound is:






25

-continued




26

-continued



4. A chiral compound of formula (Ia1) or a pharmaceutically acceptable salt thereof:



5. A pharmaceutical composition comprising the chiral compound of formula (I) or the stereoisomer, pharmaceutically acceptable salt, hydrate, or solvate thereof according to claim 1 and a pharmaceutically acceptable carrier.

6. The pharmaceutical composition according to claim 5, wherein the pharmaceutically acceptable carrier is selected from the group consisting of water for injection, lyophilized powder excipient, and oral preparation excipient.

7. A pharmaceutical composition comprising the chiral compound of formula (Ia1) or the pharmaceutically acceptable salt thereof according to claim 4 and a pharmaceutically acceptable carrier.

8. The pharmaceutical composition according to claim 7, wherein the pharmaceutically acceptable carrier is selected

27

from the group consisting of water for injection, lyophilized powder excipient, and oral preparation excipient.

9. A pharmaceutical composition comprising the chiral compound of formula (I) or the pharmaceutically acceptable salt thereof according to claim 3 and a pharmaceutically acceptable carrier.

10. A method of treating a viral infection in a subject in need thereof, the method comprising administering to the subject the pharmaceutical composition according to claim 5.

11. The method according to claim 10, wherein the viral infection is selected from the group consisting of HIV infection, hepatitis B, and diseases caused by hepatitis B virus.

12. A method of treating a viral infection in a subject in need thereof, the method comprising administering to the subject the pharmaceutical composition according to claim 7.

28

13. The method according to claim 12, wherein the viral infection is selected from the group consisting of HIV infection, hepatitis B, and diseases caused by hepatitis B virus.

14. A method of treating a viral infection in a subject in need thereof, the method comprising administering to the subject the pharmaceutical composition according to claim 9.

15. The method according to claim 14, wherein the viral infection is selected from the group consisting of HIV infection, hepatitis B, and diseases caused by hepatitis B virus.

16. The chiral compound according to claim 1, or the stereoisomer, pharmaceutically acceptable salt, hydrate, or solvate thereof, wherein R¹ and R² are each methyl.

* * * * *