



US009909152B2

(12) **United States Patent**  
**Dai et al.**

(10) **Patent No.:** **US 9,909,152 B2**  
(45) **Date of Patent:** **Mar. 6, 2018**

(54) **ENHANCED ITACONIC ACID PRODUCTION  
IN *ASPERGILLUS* WITH INCREASED *LAEA*  
EXPRESSION**

2010/0062485 A1 3/2010 Kang et al.  
2013/0137150 A1 5/2013 Dai et al.  
2015/0232892 A1 8/2015 Dai et al.

(71) Applicant: **Battelle Memorial Institute**, Richland,  
WA (US)

OTHER PUBLICATIONS

(72) Inventors: **Ziyu Dai**, Richland, WA (US); **Scott E.  
Baker**, Richland, WA (US)

Sadowski et al., *Current Opinion in Structural Biology* 19:357-362,  
2009.\*

(73) Assignee: **Battelle Memorial Institute**, Richland,  
WA (US)

Kubodera et al., *Biosci. Biotechnol. Biochem.* 66(2):404-406,  
2002.\*

(\* ) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 102 days.

Bentley and Thiessen, "Biosynthesis of Itaconic Acid in *Aspergillus  
Terreus*," *J Biol Chem* 226:703-720, 1957.

(21) Appl. No.: **14/928,511**

Aebi et al., "Cloning and Characterization of the ALG3 Gene of  
*Saccharomyces cerevisiae*," *Glycobiol.* 6:439-444, 1996.

(22) Filed: **Oct. 30, 2015**

Andersen et al., "Comparative Genomics of Citric-Acid-Producing  
*Aspergillus niger* ATCC 1015 Versus Enzyme-Producing CBS  
513.88," *Genome Res.* 21:885-897, 2011.

(65) **Prior Publication Data**

US 2016/0046967 A1 Feb. 18, 2016

Apweiler et al., "On the Frequency of Protein Glycosylation, as  
Deduced from Analysis of the Swiss-PROT Database," *Biochim.  
Biophys. Acta* 1473:4-8, 1999.

**Related U.S. Application Data**

(63) Continuation-in-part of application No. 14/703,499,  
filed on May 4, 2015, now Pat. No. 9,206,450, which  
is a continuation-in-part of application No.  
13/691,396, filed on Nov. 30, 2012, now Pat. No.  
9,023,637.

Baba et al., "Identification and Characterization of *Penicillium  
citrinum* VeA and LaeA as Global Regulators for ML-236B Pro-  
duction," *Curr. Genet.* 58:1-11, 2012.

(60) Provisional application No. 61/565,018, filed on Nov.  
30, 2011.

Bayram et al., "VelB/VeA/LaeA Complex Coordinates Light Signal  
with Fungal Development and Secondary Metabolism," *Science*  
320:1504-1506, 2008.

(51) **Int. Cl.**

**C12P 7/46** (2006.01)  
**C12P 7/44** (2006.01)  
**C12N 1/15** (2006.01)  
**C12N 1/14** (2006.01)  
**C12P 7/48** (2006.01)  
**C07K 14/38** (2006.01)  
**C12N 9/10** (2006.01)

Bayram et al., "LaeA Control of Velvet Family Regulatory Proteins  
for Light-Dependent Development and Fungal Cell-Type Specif-  
icity," *PLoS Genet.* 6:e1001226, 2010.

(52) **U.S. Cl.**

CPC ..... **C12P 7/44** (2013.01); **C07K 14/38**  
(2013.01); **C12N 1/14** (2013.01); **C12N**  
**9/1051** (2013.01); **C12P 7/48** (2013.01); **C12Y**  
**204/01258** (2013.01)

Bok and Keller, "LaeA, a Regulator of Secondary Metabolism in  
*Aspergillus* spp.," *Eukaryot. Cell* 3:527-535, 2004.

(58) **Field of Classification Search**

CPC ... **C12P 7/44**; **C12P 7/48**; **C07K 14/38**; **C12Y**  
**204/01258**; **C12N 9/1051**; **C12N 1/14**  
See application file for complete search history.

Bok et al., "Secondary Metabolic Gene Cluster Silencing in  
*Aspergillus nidulans*," *Mol. Microbiol.* 61:1636-1645, 2006.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,532,148 A 7/1996 Datta et al.  
9,023,637 B2 5/2015 Dai et al.  
2004/0058872 A1 3/2004 Keller et al.  
2009/0124014 A1\* 5/2009 Berg ..... C12N 15/815  
435/471

Bok et al., "Chromatin-Level Regulation of Biosynthetic Gene  
Clusters," *Nat. Chem. Biol.* 5:462-464, 2009.

Bouhired et al., "Accurate Prediction of the *Aspergillus nidulans*  
Terrequinone Gene Cluster Boundaries Using the Transcriptional  
Regulator LaeA," *Fungal Genet. Biol.* 44:1134-1145, 2007.

Bowman et al., "Mutational Analysis of the  
Glycosylphosphatidylinositol (GPI) Anchor Pathway Demonstrates  
that GPI-Anchored Proteins Are Required for Cell Wall Biogenesis  
and Normal Hyphal Growth in *Neurospora crassa*," *Eukaryot. Cell*  
5:587-600, 2006.

Branden et al., "Introduction to Protein Structure," Garland Pub-  
lishing Inc., New York, p. 247, 1991.

Butchko et al., "Lae1 Regulates Expression of Multiple Secondary  
Metabolite Gene Clusters in *Fusarium verticillioides*," *Fungal  
Genet. Biol.* 49:602-612, 2012.

(Continued)

Primary Examiner — Delia M Ramirez

(74) Attorney, Agent, or Firm — Klarquist Sparkman,  
LLP

(57) **ABSTRACT**

Fungi, such as *Aspergillus niger*, having a dolichyl-P-Man:  
Man(5)GlcNAc(2)-PP-dolichyl mannosyltransferase (Alg3)  
gene genetic inactivation, increased expression of a loss of  
aflR expression A (LaeA), or both, are described. In some  
examples, such mutants have several phenotypes, including  
an increased production of citric acid relative to the parental  
strain. Methods of using the disclosed fungi to make citric  
acid are also described, as are compositions and kits includ-  
ing the disclosed fungi. Further described are *Aspergillus*  
*terreus* fungi overexpressing the LaeA gene and the use of  
such fungi for the production of itaconic acid.

**16 Claims, 27 Drawing Sheets**

(56)

## References Cited

## OTHER PUBLICATIONS

- Calvo, "The VeA Regulatory System and its Role in Morphological and Chemical Development in Fungi," *Fungal Genet. Biol.* 45:1053-1061, 2008.
- Dai et al., "The Galphal Protein Regulates *Aspergillus niger* Growth and Development and the Citric Acid Production in Response to Nitrogen Sources," Abstracts of the General Meeting of the American Society for Microbiology, vol. 108, Jun. 5, 2008, pp. 0-024, US ISSN: 1060-2011.
- Davidson et al., "Functional Analysis of the ALG3 Gene Encoding the Dol-P-Man: Man<sub>5</sub>G1cNAc<sub>2</sub>-PP-Dol Mannosyltransferase Enzyme of *P. pastoris*," *Glycobiology* 14:399-407, 2004.
- Dellaporta et al., "A Plant DNA Miniprep: Version II," *Plant Mol. Biol. Rep.* 1:19-21, 1983.
- Denecke et al., "Congenital Disorder of Glycosylation Type Id: Clinical Phenotype, Molecular Analysis, Prenatal Diagnosis, and Glycosylation of Fetal Proteins," *Pediatr. Res.* 58:248-253, 2005.
- de Oliveira and de Graaff, "Proteomics of Industrial Fungi: Trends and Insights for Biotechnology," *Appl. Microbiol. Biotechnol.* 89:225-237, 2011.
- Deshpande et al., "Protein Glycosylation Pathways in Filamentous Fungi," *Glycobiology* 18:626-637, 2008.
- Georgianna et al., "Beyond Aflatoxin: Four Distinct Expression Patterns and Functional Roles Associated with *Aspergillus flavus* Secondary Metabolism Gene Clusters," *Mol. Plant Pathol.* 11:213-226, 2010.
- Gerngross, "Advances in the Production of Human Therapeutic Proteins in Yeasts and Filamentous Fungi," *Nat. Biotechnol.* 22:1409-1414, 2004.
- Geysens et al., "Genomics of Protein Folding in the Endoplasmic Reticulum, Secretion Stress and Glycosylation in the Aspergilli," *Fungal Genet. Biol.* 46:S121-S140, 2009.
- Haltiwanger and Lowe, "Role of Glycosylation in Development," *Ann. Rev. Biochem.* 73:491-537, 2004.
- Huang et al., Improving itaconic acid production through genetic engineering of an industrial *Aspergillus terreus* strain, *Microb Cell Fact* 13:119, 2014.
- Jacobs et al., "Effective Lead Selection for Improved Protein Production in *Aspergillus niger* Based on Integrated Genomics," *Fungal Genet. Biol.* 46:S141-S152, 2009.
- Kainz et al., "N-Glycan Modification in *Aspergillus* Species," *Appl. Environ. Microbiol.* 74:1076-1086, 2008.
- Kale et al., "Requirement of LaeA for Secondary Metabolism and Sclerotial Production in *Aspergillus flavus*," *Fungal Genet. Biol.* 45:1422-1429, 2008.
- Kajiura et al., "*Arabidopsis thaliana* ALG3 Mutant Synthesizes Immature Oligosaccharides in the ER and Accumulates Unique N-Glycans," *Glycobiology* 20:736-751, 2010.
- Keller et al., "LaeA, A Global Regulator of *Aspergillus* Toxins," *Med. Mycol.* 44:S83-S85, 2006.
- Kim et al., "Centralized Modularity of N-Linked Glycosylation Pathways in Mammalian Cells," *PLoS One* 4:e7317, 2009.
- Körner et al., "Carbohydrate Deficient Glycoprotein Syndrome Type IV: Deficiency of Dolichyl-P-Man:Man<sub>5</sub>G1cNAc<sub>2</sub>-PP-dolichyl Mannosyltransferase," *EMBO J.* 18:6816-6822, 1999.
- Kornfeld and Kornfeld, "Assembly of Asparagine-Linked Oligosaccharides," *Annu. Rev. Biochem.* 54:631-664, 1985.
- Kotz et al., "Approaching the Secrets of N-Glycosylation in *Aspergillus fumigatus*: Characterization of the AfOch1 Protein," *PLoS One* 5:e15729, 2010.
- Kozak, "Initiation of translation in prokaryotes and eukaryotes," *Gene* 234:187-208, 1999.
- Kranz et al., "CDG-Id in Two Siblings With Partially Different Phenotypes," *Am. J. Med. Genet.* 143A:1414-1420, 2007.
- Kukuruzinska et al., "Protein Glycosylation in Yeast," *Annu. Rev. Biochem.* 56:915-944, 1987.
- Maddi and Free, "α-1,6-Mannosylation of N-Linked Oligosaccharide Present on Cell Wall Proteins is Required for Their Incorporation into the Cell Wall in the Filamentous Fungus *Neurospora crassa*," *Eukaryot. Cell* 9:1766-1775, 2010.
- Magnuson and Lasure, "Organic Acid Production by Filamentous Fungi," in *Advances in Fungal Biotechnology for Industry, Agriculture, and Medicine*, Lange and Lange (eds.), pp. 307-340, Kluwer Academic/Plenum Publishers, 2004.
- Manthri et al., "Deletion of the TbALG3 Gene Demonstrates Site-Specific NGlycosylation and N-Glycan Processing in *Trypanosoma brucei*," *Glycobiology* 18:367-383, 2008.
- Maras et al., "Molecular Cloning and Enzymatic Characterization of a *Trichoderma reesei* 1,2-α-D-mannosidase," *J. Biotechnol.* 77:255-263, 2000.
- Nam et al., "The Effects of Culture Conditions on the Glycosylation of Secreted Human Placental Alkaline Phosphatase Produced in Chinese Hamster Ovary Cells," *Biotech. Bioeng.* 100:1178-1192, 2008.
- Nevalainen et al., "Heterologous Protein Expression in Filamentous Fungi," *Trends Biotechnol.* 23:468-474, 2005.
- Oda et al., "*Aspergillus oryzae* laeA Regulates Kojic Acid Synthesis Genes," *Biosci. Biotechnol. Biochem.* 75:1832-1834, 2011.
- Pang et al., "Human Sperm Binding is Mediated by the Sialyl-Lewis<sup>x</sup> Oligosaccharide on the Zona Pellucida," *Science* 333:1761-1764, 2011.
- Pel et al., "Genome Sequencing and Analysis of the Versatile Cell Factory *Aspergillus niger* CBS 513.88," *Nature Biotechnol.* 25:221-231, 2007.
- Punt et al., "Filamentous Fungi as Cell Factories for Heterologous Protein Production," *Trends Biotechnol.* 20:200-206, 2002.
- Ramamoorthy et al., "veA-Dependent RNA-pol II Transcription Elongation Factor-Like Protein, RtfA, is Associated with Secondary Metabolism and Morphological Development in *Aspergillus nidulans*," *Mol. Microbiol.* 85:795-814, 2012.
- Reyes-Dominguez et al., "Heterochromatic Marks are Associated with the Repression of Secondary Metabolism Clusters in *Aspergillus nidulans*," *Mol. Microbiol.* 76:1376-1386, 2010.
- Roze et al., "Volatile Profiling Reveals Intracellular Metabolic Changes in *Aspergillus parasiticus*: veA Regulates Branched Chain Amino Acid and Ethanol Metabolism," *BMC Biochem.* 11:33, 2010.
- Sakai et al., "Heterologous Expression System in *Aspergillus oryzae* for Fungal Biosynthetic Gene Clusters of Secondary Metabolites," *Appl. Microbiol. Biotechnol.* 93:2011-2022, 2012.
- Sauer et al., "Microbial Production of Organic Acids: Expanding the Markets," *Trends Biotechnol.* 26:100-108, 2008.
- Schollen et al., "CDG-Id Caused by Homozygosity for an ALG3 Mutation Due to Segmental Maternal Isodisomy UPD3(q21.3-pter)," *Eur. J. Med. Genet.* 48:153-158, 2005.
- Schuster et al., "On the Safety of *Aspergillus niger*—A Review," *Appl. Microbiol. Biotechnol.* 59:426-435, 2002.
- Seffernick et al., "Melamine Deaminase and Atrazine Chlorohydrolase: 98 Percent Identical but Functionally Different," *J Bacteriol* 183(8):2405-2410, 2001.
- Silberstein and Gilmore, "Biochemistry, Molecular Biology, and Genetics of the Oligosaccharyltransferase," *FASEB J.* 10:849-858, 1996.
- Soukup et al., "Overexpression of the *Aspergillus nidulans* Histone 4 Acetyltransferase EsaA Increases Activation of Secondary Metabolite Production," *Mol. Microbiol.* 86:314-330, 2012.
- Sousa et al., "The ARO4 gene of *Candida albicans* encodes a tyrosine-sensitive DAHP synthase: evolution, functional conservation and phenotype of Aro3p-, Aro4p-deficient mutants," *Microbiology* 148(Pt5):1291-1303, 2002.
- Stibler et al., "Carbohydrate-Deficient Glycoprotein Syndrome—A Fourth Subtype," *Neuropediatrics* 26: 235-237, 1995.
- Sun et al., "Congenital Disorder of Glycosylation Id Presenting with Hyperinsulinemic Hypoglycemia and Islet Cell Hyperplasia," *J. Clin. Endocrinol. Metab.* 90:4371-4375, 2005.
- Trombetta and Parodi, "Quality Control and Protein Folding in the Secretory Pathway," *Ann. Rev. Cell Dev. Biol.* 19:649-676, 2003.
- Tsang et al., "Analytical and Computational Approaches to Define the *Aspergillus niger* Secretome," *Fungal Genet. Biol.* 46:S153-S160, 2009.
- Uniprot ALG3\_ASPNC [online] Nov. 2, 2010 [retrieved Jan. 9, 2013], available on the internet <URL: [www.uniprot.org/uniprot/A2RA94.txt?version=27](http://www.uniprot.org/uniprot/A2RA94.txt?version=27)>.




(56)

**References Cited**

OTHER PUBLICATIONS

- Witkowski et al., "Conversion of a  $\beta$ -Ketoacyl Synthase to a Malonyl Decarboxylase by Replacement of the Active-Site Cysteine with Glutamine," *Biochemistry* 38:11643-11650, 1999.
- Xing et al., "Molecular Cloning and Characterization of the Global Regulator LaeA in *Penicillium citrinum*," *Biotechnol. Lett.* 32:1733-1737, 2010.
- Yan and Lennarz, "Unraveling the Mechanism of Protein N-glycosylation," *J. Biol. Chem.* 280:3121-3124, 2005.
- Yu et al., "Conservation of Structure and Function of the Aflatoxin Regulatory Gene aflR from *Aspergillus nidulans* and *A. flavus*" *Curr. Genet.* 29:549-555, 1996.
- Zhou et al., "Global analysis of gene transcription regulation in prokaryotes," *Cell Mol Life Sci* 63(19-20):2260-2290, 2006.

\* cited by examiner






FIG. 2

FIG. 3A

*11414kusa*

*alg3Δ*




FIG. 3B










FIG. 4

CM

PDA









FIG. 5

FIG. 6A *11414kuaA* *Alg3Δ*  
8 hrs after inoculation



15 hrs after inoculation





FIG. 6B






FIG. 7

Transgene fragment for complemented transformation




FIG. 8A




FIG. 8B

# FIG. 9

GENE ID: 5996303 AOR\_1\_556094 | dolichyl-P-Man:Man(5)GlcNAc(2)-PP-dolichyl mannosyltransferase [Aspergillus oryzae RIB40] (10 or fewer PubMed links)

Score = 137 bits (74), Expect = 2e-28  
Identities = 400/553 (72%), Gaps = 40/553 (7%)  
Strand=Plus/Plus

Query	801	CAGGTTTTACTCGCGATACCGTTCCTACAAAACAACCCGGCGGGGTATC-TCTCGCGGGC	859
Sbjct	643	CAGGTTCTACTAGCGATTCCCTTCCTACAGGGTAAACCCCATAGGATA-CGTCGCGCGGGC	701
Query	860	GTTTCGAGCTAACCAGACAGTTCATGTTTAAATGGACAGTCAATTGGAGATTTGTTGGCGA	919
Sbjct	702	CTTTGAGTTGACTAGACAGTTTATGTTCAAATGGACTGTCAAATTGGAGGTTTGTGGGEGA	761
Query	920	AGAAGTATTCTIAT-CTAAGAGCTTTTCCCTGGCATTGCT-GGCCGTCCACATTTGTGCTG	977
Sbjct	762	AGACTTGTTCCTATCCAAACAG-TTTTCTCTAGCCTTACTAGG-TTTCATATTTTCTG	819
Query	978	CTAGGCG-CT-TTTGCCGTCACTGGTTGGCTGA-GATAC-ICCAGG-TCTAGCTTGCCTG	1032
Sbjct	820	CT-GG-GATTATTTGTTACCACAGGCTGG-ITACG-GCCGT-CAGGATCTAACGTCCCTG	874
Query	1033	-CGTTCAITTCGGAATCTGCTAGC-GGGTCGACATCGCACAGT-GTCCCTCCCAAACCCT	1089
Sbjct	875	AC-ITCCTCCGGAGCCTACT-CCAAGGACGCCAACGCACCCTGGT-GCTTTCTAAGTCTT	931
Query	1090	ACATCATGAGCGTGATGCTCTCGTCTCTGACAG-TTGGCTTGTGTGCGCAAGGTCCCTT	1148
Sbjct	932	TCATAATGACCGTGATGTTGACATCGCTGGC-GATCGGGTGTGTGCGCAAGGTCCCTT	990
Query	1149	CATTACCAATTCCTCGCCTACCTCTCCTGGGCGACACCCTT-CCTCCTCTGGCGCGCAGG	1207
Sbjct	991	CATTACCAATTCCTTGCCTATCTCTCCTGGGCTACGCC-TTGCCTTCTCTGGCGGGCTCG	1049
Query	1208	GTTTCATCCAATC-TTGCTGTAC-CCTATCTGGGCTA-TGCAAGAGTGGGCTTGAACA-	1263
Sbjct	1050	GCTCCATCCGATCCTTA-TATATGCG-ATCTGGGC-ACTACAGGAGTGGGCTTGAATGT	1106
Query	1264	CATTCCCAGCACCAAC-CTCAGTTCCATCATT-GTTGTCCTCTCACTTGCTACCCAGAG	1321
Sbjct	1107	C-TACCCAAGCACCAATGC-CAGTTCT-TCGCTCGTTGTCTTCTCACTTGCTGTTTACAG-G	1162
Query	1322	TTT-CGGCGTCCCT 1333	
Sbjct	1163	TTTTCGGTGTCCCT 1175	



FIG. 10B

\*\*+--+\*\*\*\*\*+\*+.\*+\*\*\*\*\*+. . . .+ . . .\* . . .+ . . .+\*\*\*\*\* . . \* .  
*Aspergillus niger* 322) VGLLCARSLHYQFFAYLSWATPFLLWRAGFHPI---LLYLIWAMQEWAWNTFPSTNLSSI  
*Aspergillus nid.* 322) VGLLCARSLHYQFFAYLSWATPFLLWQAGYHPI---LVYALWLWQEWAWNVYPSTNLSSA  
*Fusarium oxy.* 341) IGLLFARSLHYQFYAYLAWAIPYLLWRATEDPV---IVAIIWAAQEWAWNVYPSTDLSSY  
*Neurospora cra.* 349) VGLLFARSLHYQFYAYVAWSTPFLLWRAGLHPV---LVYLLWAVHEWAWNVFPSTPASSA  
*Saccharomyces cer.* 342) IGMCFSRSLHYQFYVWYFHTLPYLLWGGVKKLARLLRVLIILGLIELSWNTYPSTNYSSL  
*Arabidopsis tha.* 345) IGIVFARSLHYQFYVWYFHTLPYLLWRIPF-PT--WLRLIMFLGIELCWNVYPSTPSSSG  
*Homo sap.* 343) IGICFSRSLHYQFYVWYFHTLPYLLWAMPARWLTHLLRLLVLGLIELSWNTYPSTSCSSA

.+ . . . . .+\*+ . . . . . . . . . .+ . . .+ . . .-  
*Aspergillus niger* 379) IVVLSLATQSGVLANASASA-FYTMRSNPSGKEHNQ--  
*Aspergillus nid.* 379) AVVLLLCAQVLGVLVNRDRA-FPSSPPTPKAKQHVQ--  
*Fusarium oxy.* 398) IAVNTMLATVVLVYLGTARR-AVPAPAAQVGNVDDKKNK  
*Neurospora cra.* 406) VVVGVLGVTVAGVWFGAREEwEPCMKSSSKKEEAAMR·  
*Saccharomyces cer.* 402) SLHVCHLIILLCLWLNPNPA-SPSHRSENKAKSH-----  
*Arabidopsis tha.* 402) LLLCLHLIILVGLWLAPSVD-PYQLKEHPKSQIHKKA-  
*Homo sap.* 403) ALHICHAVILLQLWLGFPQPF--PKSTQHSK-KAH-----

# FIG. 11

Score = 279 bits (713), Expect = 2e-87, Method: Compositional matrix adjust.  
 Identities = 159/405 (39%), Positives = 239/405 (59%), Gaps = 6/405 (1%)

Query	12	FNPRHTKWMAPLLVLGDAFLCALIIWKVPYTEIDWATYMQQISLYLSGERDYTLIRGSTG	71
		F P +T + +L + + +I KV YTEIDW YM ++ ++G DYT ++G TG	
Sbjct	31	FKPEYILLVTAFLWFLEIAINIWVIQKVSYTEIDWKAYMDEVEGVINGTYDYTLKGDG	90
Query	72	PLVYPAAHVYSYALYHLLTDEGRDIFFGQILFAVLYLITLVVVLCCY-RQSGAPPYLLPL	130
		PLVYPA VY +T LY+LTD G +I GQ +FAV YLI L++V+ Y R PPY+	
Sbjct	91	PLVYPAGFVYIFTGLYYLTDHGHNIRLQYVFAVSYLINLLVMRIYHRTRKVPYVFFF	150
Query	131	LVL-SKRLHSVYVLRLEFNDGLAALAMWVAILLFMNRKWTAAVAVWSTGVAIKMTLLLLAP	189
		+ S R+HS+++LRLFND +A + + AI LF++ +WE A++S V++KM +LL AP	
Sbjct	151	ICCASYRIHSIFILRLFNDPVMMLCFCAINLPLDGRWTLCCALYSLAVSVKMNVLFFAP	210
Query	190	AIAVVTVLSLSLGPVGLGVLAVLVQVLLAIPFLQNNPAGYLSRAFELTRQFMFKWTVNW	249
		+ + * L ++ L ++Q++L +PFL NP GY+SRAF+L RQF+EKWTVNW	
Sbjct	211	GLLFLLLCEFGWLKTLPRALCAVIQLVGLPFLVNPVGYVSRAFDLGRQFLFKWTVNW	270
Query	250	RFVGEVFLSKSFLALLAVHIVLLGFAVTVGLRYSRSSLPAFIRNLLAGRHRTVSLPK	309
		RF+ E+VFL++ F LALL HI L FA+ W R S SS+ +++ + +	
Sbjct	271	RFLPEDVFLNRYFHLALLLAHITILLLFALKRWKR-SGSSIWTILKDPSEKETAHKVNA	329
Query	310	PYIMSVMLSSLTVGLLCARSLHYQFFAYLSWATPELLWRAGFHP---ILLYLIWAMQEWA	366
		++ ++ +S +G+ +RSLHYQF+ + P+LLW G +L LI + E +	
Sbjct	330	DQMVLIIFTSNFIGMCFSRSLHYQFYVWYFHTLPYLLWSGGVKKLARLLRVLILGLIELS	389
Query	367	WNTFPSTNLSSIIIVVLSLATQSFGLANSASAFYTMRSNPSGKEH	411
		WNT+PSIN SS+ + + + N A + RS K H	
Sbjct	390	WNTYPSYSSLSLHVCHLIILLCLWLNPNPASPSHRSENKAKSH	434



# FIG. 12

**A. *nidulans***

Query 1 MTSPAHNHYSYHSEPTSSDRGRSRQNSDAMDIOQSI TERE PATR-----YAVAGGPAPWN 53  
M SP N+YSY S D GR SRQNSDAMDI IT +EP Y GGPA +  
Sbjct 14 MASPNNRNNYSYQGI ESYDSGRSRQNSDAMD I HVIT AQEPPREPPDNDPYDGHGCPAGTS 73

**A. *niger***

Query 54 RNGSPSMSPINSE RNQFHEENGR TYHGFR RGM YFLPCDEQE QDRLDIFHKLFTVARVSES 113  
P R F+EENGR TYHG+RRG+Y LPCDEQE QDRLDIFHKLFTVAR+SES

Sbjct 74 HYSKPP-----NRWLFYEENGR TYHGYRRGVYPLPCDEQE QDRLDIFHKLFTVARMSES 127

Query 114 LIYAPHPTNGRFLDLGCGTGIWAI EVANKY PDAFVAGVDLAPIQPPNHPKNCEFYAPFDF 173  
LIYAPHP NGRFLDLGCGTGIWAI+VA+KYP+AFVAGVDLAPIQPPNHP NCEFYAPFDF

Sbjct 128 LIYAPHPPNGRFLDLGCGTGIWAI DV AHKYPNAFVAGVDLAPIQPPNHPDNCEFYAPFDF 187

Query 174 EAPWAMGEDSWDLIHLQMGCGSVMGWPNLYRRIFAHLRPGAWFEQVEIDFEPRCDDRSLD 233  
EAPW +GE+SWDLIHLQMGCGSV+GW NLY+RI HL+PGAWFEQVEIDFEPRCDDRSL+

Sbjct 188 EAPWTLGENSWDLIHLQMGCGSVLGWQNLYKRILRHLPQAWFEQVEIDFEPRCDDRSLN 247

Query 234 GTALRHWDCLKQATAETMRPIAHSSRDTIKDLQDAGFTEIDHQIVGLPLNPWHQDEHER 293  
G ALR WY LKQAT +TMRPIAHSSRDTI+ L++ACFT+IDHQ+VGLPLNPWH+DEHE+

Sbjct 248 GLALREWYQYLKQATQDTMRPIAHSSRDTIRHLEEAGFTQIDHQMVG LPLNPWHRDEHEQ 307

Query 294 KVARWYNLAVSESEIENLSLAPFSRVYRWPLERIQQLAADVKSEAFNKEIHAYNILHIYQA 353  
KVARWYNLA+SESIE LSLAPFSR++ W L+RI+Q+ A+VKS+AFNKEIHAYNILHIYQA

Sbjct 308 KVARWYNLAISESEIETLSLAPFSRIFHWDLDRIRQITAEVKSQAFNKEIHAYNILHIYQA 367

Query 354 RKP 356

RKP

Sbjct 368 RKP 370

FIG. 13




FIG. 14




FIG. 15A. Oligo primers:PTR5F/PTR3R





FIG. 15B. Oligo primers:LaeA5F/TRP3R






FIG. 16

FIG. 17A

**laeA deletion construct:**




FIG. 17B

**laeA complementation construct:**





FIG. 18

*laeA* gene deletion



Bλ

Clone-1

Clone-2

Clone-3

Clone-4

Clone-5

Clone-6

Clone-7

Clone-8

Clone-9

parent

FIG. 19






FIG. 20






FIG. 22A




FIG. 22B




FIG. 23




FIG. 24A




FIG. 24B





FIG. 25



## ENHANCED ITACONIC ACID PRODUCTION IN *ASPERGILLUS* WITH INCREASED *LAEA* EXPRESSION

### CROSS-REFERENCE TO RELATED APPLICATIONS

This is a continuation-in-part of U.S. patent application Ser. No. 14/703,499, filed May 4, 2015, issued as U.S. Pat. No. 9,206,450 on Dec. 8, 2015, which is a continuation-in-part of U.S. application Ser. No. 13/691,396, filed Nov. 30, 2012, issued as U.S. Pat. No. 9,023,637 on May 5, 2015, which claims the benefit of U.S. Provisional Application No. 61/565,018, filed Nov. 30, 2011. The above-referenced applications are herein incorporated by reference in their entirety.

### ACKNOWLEDGMENT OF GOVERNMENT SUPPORT

This invention was made with government support under contract number DE-AC05-76RL01830 awarded by the U.S. Department of Energy. The government has certain rights in the invention.

### FIELD

This disclosure concerns recombinant *Aspergillus terreus* fungi that are genetically enhanced to increase the expression levels of the loss of aflR expression A (LaeA) gene, which results in a significant increase in itaconic acid production. This disclosure further concerns methods of using these fungi to produce itaconic acid.

### BACKGROUND

Filamentous fungi, such as *Aspergillus niger*, are well known for their industrial applications in protein and chemical productions. They are used to produce a wide variety of products ranging from human therapeutics, glycosyl hydrolases to specialty chemicals (Punt et al., *Trends Biotechnol* 20(5):200-206, 2002; Schuster et al., *Appl Microbiol Biotechnol* 59(4-5):426-435, 2002; Gerngross, *Nat Biotechnol* 22(11):1409-1414, 2004; Nevalainen et al., *Trends Biotechnol* 23(9):468-474, 2005; Sauer et al., *Trends Biotechnol* 26(2):100-8, 2008; Magnuson and Lasure (2004). "Organic acid production by filamentous fungi." *Advances in fungal biotechnology for industry, agriculture, and medicine*, pages 307-340). Some of industrial *A. niger* strains are capable of growing on solutions of glucose or sucrose in excess of 20% (w/v) and converting approximately 90% of the supplied carbohydrate to citric acid. These remarkable properties are the reason that *A. niger* has been used to produce citric acid for more 80 years and is currently the primary source of commercial citric acid production (Magnuson and Lasure (2004). "Organic acid production by filamentous fungi." *Advances in fungal biotechnology for industry, agriculture, and medicine*, pages 307-340).

The maximum product output in fermentation processes is the result of optimal metabolic pathways and cellular formation, which are influenced by endogenous and exogenous factors. Cellular metabolisms are tightly controlled and highly interconnected, and are regulated spatially and temporally at different levels, such as transcription, post-transcription, translation, and post-translation. Therefore, different approaches have been explored to understand the regulatory mechanisms of metabolic processes and cellular

formation for maximizing the product output in filamentous fungi. For example, comparative genomics was used to examine citric-acid-producing versus enzyme-producing *A. niger* strains (Andersen et al., *Genome Res.* 21(6): 885-97, 2011), proteomics was used to examine filamentous fungi related to enzymes or organic acid production (de Oliveira and de Graaff, *Appl. Microbiol. Biotechnol.* 89(2): 225-37, 2011), or combination of both genomics and proteomics were used to examine enzyme production (Jacobs et al., *Fungal Genetics and Biology* 46(1, Supplement):S141-S152, 2009). Although these studies examined the potential involvement of selected genes and proteins in optimizing production of organic acids or proteins in filamentous fungi, methods for altering the complex post-translation modifications (such as N-glycosylation of cellular proteins) for signal transduction, cellular formation and metabolism at different growth and development stages, which may affect product output, have not been examined.

Protein glycosylation is a ubiquitous and structurally diverse form of post translation modification, which occurs at all domains of life. More than two-thirds of eukaryotic proteins are predicted to be glycosylated (Apweiler et al., *Biochim Biophys Acta* 1473(1):4-8, 1999). N- and O-linked protein glycosylation are common types of protein glycosylation, occurring mainly on the asparagine (N) and serine/threonine (S/T) residues, respectively. N-linked glycosylation has been implicated in many biochemical and cellular processes, including protein secretion, stability and translocation, maintenance of cell structure, receptor-ligand interactions and cell signaling, cell-cell recognition, pathogen infection, and host defense in various organisms (Haltiwanger and Lowe, *Ann. Rev. Biochem.* 73(1):491-537, 2004; Dellaporta et al., *Plant Mol. Biol. Reporter* 1(4):19-21, 1983; Nam et al., *Biotech. Bioengineer.* 100(6): 1178-1192, 2008; Trombetta and Parodi, *Ann. Rev. Cell Dev. Biol.* 19(1):649-676, 2003; Tsang et al., *Fungal Genetics and Biology* 46(1): S153-S160, 2009; Pang et al., *Science*, 333(6050):1761-4, 2011).

N-glycosylation is highly complex and has been extensively studied in mammalian systems (Yan and Lennarz, *J. Biol. Chem.* 280(5):3121, 2005; Silberstein and Gilmore, *FASEB J.* 10(8): 849, 1996; Kornfeld and Kornfeld, *Annu. Rev. Biochem.* 54:631-664, 2005; Kim et al., *PLoS ONE* 4(10): e7317, 2009, 2009) and yeast (Kukuruzinska et al., *Annu. Rev. Biochem.* 56(1):915-944, 1987). The protein N-glycosylation pathways in filamentous fungi have also been identified (Deshpande et al., *Glycobiology* 18(8):626-637, 2008; Geysens et al., *Fungal Genetics and Biology* 46(1, Supplement): S121-S140, 2009) on the basis of the known genomic sequences. Several genes involved in N-glycosylation have been studied in filamentous fungi (Kotz et al., *PLoS ONE* 5(12):e15729, 2010; Kainz et al., *Appl Environ Microbiol* 74(4):1076-86, 2008; Maras et al., *J. Biotechnol.* 77(2-3):255-63, 2000; Maddi and Free, *Eukaryot Cell* 9(11):1766-75, 2010; Bowman et al., *Eukaryotic Cell* 5(3):587-600, 2006). In these studies, the effects of gene deletion on N-linked glycan patterns formation, the cell wall formation, overall protein secretion and/or the phenotypic changes were demonstrated.

Alg3 is localized in the ER and catalyzes the initial transfer of a mannose residue from dolichol pyrophosphate-mannose to lipid-linked Man5GlcNAc2-PP-Dol on the ER luminal side. It is involved in the early N-glycan synthesis in eukaryotes for the assembly of a Glc3Man9GlcNAc2 core oligosaccharide that is linked to the lipid carrier dolichol pyrophosphate. The Alg3 gene and its functions have been identified and studied in *S. cerevisiae*, *P. pastoris*, *T. brucei*,

*A. thaliana*, and human (Aebi et al., *Glycobiol.* 6(4):439-444, 1996; Korner et al., *EMBO J.* 18(23): 6816-6822, 1999; Davidson et al., *Glycobiology* 14(5):399-407, 2004; Manthri et al., *Glycobiol.* 18(5):367-83, 2008; Kajiura et al., *Glycobiol.* 20(6):736-51, 2010). In these studies, the Alg3 mutants exhibited a unique structural profile in the glycoproteins, such as Man3GlcNAc2, Man4GlcNAc2, Man5GlcNAc2, GlcMan5GlcNAc2, and Glc3Man5GlcNAc2, which affected the overall N-glycosylation by incomplete utilization of N-linked glycosites in glycoproteins. No obvious growth phenotype was observed in those Alg3Δ mutants of *S. cerevisiae*, *P. pastoris*, *T. brucei*, and plant except that the Alg3 defect in human caused severe diseases such as profound psychomotor delay, optic atrophy, acquired microcephaly, iris obovomas and hypsarrhythmia (Stibler et al., *Neuropediatrics* 26(5): 235-7, 1995; Sun et al., *J. Clin. Endocrinol. Metab.* 90(7):4371-5, 2005; Schollen et al., *Eur. J. Med. Genet.* 48(2):153-158, 2005; Kranz et al., *Am. J. Med. Genet.* 143A(13):1414-20, 2007; Denecke et al., *Pediatr. Res.* 58(2):248-53, 2005).

LaeA, a global regulator gene for the secondary metabolism, was first identified in *A. nidulans* through complementing the aflR deficient mutants (Bok and Keller, *Eukaryot Cell* 3:527-535, 2004). Deletion of LaeA gene inhibits the expression of secondary metabolic gene clusters, such as sterigmatocystin, penicillin, and lovastatin, but has no effect on spore production in *A. nidulans*. The LaeA that was confirmed as a nuclear protein and a putative methyltransferase does not involve in gene clusters for nutrient utilization (Bok et al., *Mol Microbiol* 61:1636-45, 2006). Furthermore, the role of LaeA in secondary metabolism was confirmed in *Aspergillus flavus* and *Aspergillus oryzae* (Kale et al., *Fungal Genet. Biol.* 45:1422-9, 2008; Oda et al., *Biosci Biotechnol Biochem* 75:1832-4, 2011). Evidence indicates that LaeA reverses gene repression at the level of the heterochromatin state (Reyes-Dominguez et al., *Molecular Microbiology* 76:1376-86, 2010). LaeA is a component of the heterotrimeric VeA/VelB/LaeA protein complex (Bayram et al., *Science Signalling* 320:1504, 2008), which involves in the acetylation signal transduction for secondary metabolite production in *A. nidulans* (Soukup et al., *Mol. Microbiol.*, 86(2):314-30, 2012). The veA/VelB/LaeA complex may coordinately respond to environmental cues (Ramamoorthy et al., *Mol. Microbiol.*, 85(4):795-814, 2012) and has a role in fungal morphology (Calvo, *Fungal Genetics and Biology* 45:1053-61, 2008). LaeA may direct the formation of the VelB-VosA and VelB-VelA-LaeA complexes, control veA modification and protein levels, and be involved in light regulation of growth and development (Bayram et al., *PLoS genetics* 6: e1001226, 2010).

### SUMMARY

Provided herein are isolated *Aspergillus terreus* fungi transformed with a heterologous nucleic acid molecule comprising an *Aspergillus* species gene. Also provided is a method of making itaconic acid by culturing an isolated *A. terreus* fungus transformed with a heterologous nucleic acid molecule comprising an *Aspergillus* species LaeA gene under conditions that permit the fungus to make itaconic acid. In some embodiments, the heterologous nucleic acid construct includes a heterologous LaeA gene, a heterologous promoter, a heterologous transcription terminator, and/or a heterologous selective marker gene. In non-limiting examples, the heterologous *Aspergillus* species LaeA gene is an *A. fumigatus* laeA gene, an *A. nidulans* laeA gene, an *A. niger* LaeA gene, or an *A. oryzae* laeA gene.

Although the current commercial conversion rate of carbohydrate to citric acid in *A. niger* is more than eighty to ninety percent, further improvement in production of citric acid and other metabolites is desirable. This disclosure describes the role of the dolichyl-P-Man:Man(5)GlcNAc(2)-PP-dolichyl mannosyltransferase gene ( $\alpha$ -1,3-mannosyltransferase, Alg3) on the spore germination, filamentous growth, sporulation, and production of citric acid in *Aspergillus niger*. In addition, the role of LaeA in citric acid production by its over-expression is shown, alone or in combination with an Alg3Δ mutant background.

Based on these observations, provided herein are isolated fungi (such as filamentous fungi) having a gene inactivation (also referred to herein as a gene deletion) of a dolichyl-P-Man:Man(5)GlcNAc(2)-PP-dolichyl mannosyltransferase (Alg3) gene (referred to herein as Alg3Δ strains), a gene enhancement (e.g., overexpression) of a LaeA gene (referred to herein as upregulated LaeA strains), or both. Any strain of fungi can be used, such as a filamentous fungi, for example *Aspergillus niger* (*A. niger*) or particular strains thereof (for example *A. niger* strain 11414 or 11414KusA). In particular examples, an Alg3Δ strain exhibits one or more of the following characteristics: slower growth on citric acid production (CAP) medium, complete medium (CM) or potato dextrose agar (PDA) medium; earlier spore germination and a higher germination rate in CAP medium; delayed spore germination in CM or PDA medium; reduced sporulation on complete medium; or combinations thereof. In some examples, such increases or decreases are relative to *A. niger* strain 11414KusA grown under the same conditions. The combination of Alg3Δ and over-expression of LaeA resulted in some improvement of sporulation on CM.

In particular examples, such Alg3Δ strains, up-regulated LaeA strains, or Alg3Δ-upregulated LaeA strains, produce more citric acid when grown in CAP medium, such as at least 20%, at least 50%, or at least 70% more than *A. niger* strain 11414KusA under identical growing conditions after at least 4 days, at least 5 days or at least 10 days. Thus, one strategy to increase citric acid production is to reduce the carbohydrate consumption for protein glycosylation and cellular formation, as altering protein glycosylation can augment the carbohydrate flux into citric acid production in *A. niger*.

Also provided herein are compositions (such as fermentation broth) and kits that include a fungal Alg3Δ strain, up-regulated LaeA strain, or Alg3Δ-upregulated LaeA strain.

Also provided herein are methods of making citric acid using the disclosed fungal Alg3Δ strains, up-regulated LaeA strains, and Alg3Δ-upregulated LaeA strains. For example, such a method can include culturing an isolated Alg3Δ fungus, up-regulated LaeA fungus, or Alg3Δ-upregulated LaeA fungus, under conditions that permit the fungus to make citric acid, thereby making citric acid. For example, the Alg3Δ fungus, up-regulated LaeA fungus, or Alg3Δ-upregulated LaeA fungus, can be cultured in CAP medium. In some examples, the method further includes isolating the citric acid produced.

The foregoing and other objects and features of the disclosure will become more apparent from the following detailed description, which proceeds with reference to the accompanying figures.

### BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1A is a schematic drawing showing a restriction map of the 10.9 kb fragment containing the *A. niger* Alg3 gene.



## 5

FIG. 1B is a schematic drawing showing the introduction of the hyg-selective marker, which was flanked by the upstream and downstream DNA sequences of Alg3. Integration of the linear molecules by homologous recombination replaces Alg3 with hph in the chromosome.

FIG. 1C is a digital image of a Southern blot showing the genomic DNA hybridization of parent and Alg3Δ strains. One parent and two selected Alg3Δ strains are shown, which have the correct enzyme restriction pattern.

FIG. 2 is a series of digital images showing the Alg3Δ and parent 11414kusA strains grown on agar plates of complete medium (CM), potato dextrose (PDA), and minimal medium (MM) at 30° C. for 15 hours or 48 hours. Parent strain (panel A), Alg3Δ strain (panel B), and both parent and Alg3Δ strains (panel C) grown on complete medium plate. Parent strain (panel D), Alg3Δ strain (panel E), and both parent and Alg3Δ strains (panel F) were grown on PDA agar plates. Parent strain (panel G), Alg3Δ strain (panel H), and both parent and Alg3Δ strains (panel I) were grown on MM agar plates.

FIG. 3A is a series of digital images showing the spore germination of parent 11414kusA and Alg3Δ strains in liquid cultures of complete medium (CM), potato dextrose medium (PDA), and minimal medium (MM) at 30° C. for 7 hours. Panels A and B are germinated spores in CM liquid culture. Panels C and D are germinated spores in PDA liquid culture. Panels E and F are germinated spores in MM liquid culture. The left panels for the parent 11414kusA strain and the right panels for the Alg3Δ strain.

FIG. 3B provides graphs showing the time courses of the percentage of spore germination of parent 11414kusA and Alg3Δ strains grown in the liquid cultures of complete medium (CM; top graph), potato dextrose medium (PDA; middle graph), or minimal medium (MM; bottom graph). The solid filled cycle is the parent 11414kusA strain and open cycle is the Alg3Δ strain.

FIG. 4 shows a series of digital images (panels A and D) and stereo microscopy digital images (panels B, C, E and F) of parent and Alg3Δ strains grown on agar plates of complete medium (CM) and potato dextrose (PDA) at 30° C. for 4 days.

FIG. 5 provides stereo microscopy digital images parent 11414kusA and Alg3Δ strains grown on citric acid production (CAP) medium plates at different pH levels for 27 hours. The left panels for parent strain 11414kusA and right panels for the Alg3Δ strain.

FIGS. 6A and 6B show spore germination of parent 11414kusA and Alg3Δ strains in citric acid production (CAP) liquid medium. (FIG. 6A, top left and bottom left panel) Inverted microscopic images for parent 11414kusA strain. (FIG. 6A, top right and bottom right panel) Inverted microscopic images for Alg3Δ strain. (FIG. 6A, top panels) Strains grown in CAP liquid culture at 30° C. for 8 hours. (FIG. 6A, bottom panels) Strains grown in CAP liquid culture at 30° C. for 15 hours. (FIG. 6B) Time course of the spore germination rate (%) of parent and Alg3Δ strains grown in CAP liquid culture. The solid cycle for parent 11414kusA strain and open cycle for the Alg3Δ strain.

FIG. 7 is a graph showing the time course of citric acid production by parent 11414kusA and Alg3Δ strains in the liquid culture of citric acid production at 30° C. and 200 rpm.

FIGS. 8A and 8B show (FIG. 8A) a schematic diagram showing the construction of the transgene used to complement the Alg3Δ mutant, and (FIG. 8B) a graph showing the citric acid production by parent strain 11414kusA

## 6

(kusA), Alg3Δ mutant (Alg3) and Alg3Δ mutant complemented with Alg3 gene (cAlg3) after growth at 30° C., 200 rpm for 12 days

FIG. 9 shows an alignment of Alg3 nucleic acid sequences from *A. niger* (top strand, nucleotides 1986-2518 of SEQ ID NO: 1) and *A. oryzae* (bottom strand, nucleotides 643-1175 of SEQ ID NO: 3).

FIGS. 10A and 10B show an alignment of Alg3 protein sequences from *A. niger* (SEQ ID NO: 2), *A. nidulans* (SEQ ID NO: 31), *Fusarium oxysporum* (SEQ ID NO: 32), *Neurospora crassa* (SEQ ID NO: 33), *S. cerevisiae* (SEQ ID NO: 34), *Arabidopsis thaliana* (SEQ ID NO: 35), and *Homo sapiens* (SEQ ID NO: 36). The signs at the top of the alignment show: ‘-’ the average weight of column pair exchanges is less than weight matrix mean value; ‘.’ is less than mean value plus one SD; ‘+’ is less than mean value plus two SD; and ‘\*’ is more than mean value plus two SD.

FIG. 11 shows an alignment of Alg3 protein sequences from *A. niger* (top strand, amino acids 12-411 of SEQ ID NO: 2) and *S. cerevisiae* (bottom strand, amino acids 31-434 of SEQ ID NO: 34).

FIG. 12 shows an alignment of LaeA protein sequences from *A. nidulans* (top strand, amino acids 14-372 of SEQ ID NO: 41) and *A. niger* (bottom strand, amino acids 14-370 of SEQ ID NO: 59).

FIG. 13 is a schematic diagram showing the construction of pPTRpGPDALaeA plasmid vector. Both the glyceraldehyde 3-phosphate dehydrogenase (gpdA) promoter and LaeA (loss of aflR expression A) gene coding sequence of genomic DNA were isolated by overlap PCR from pAN7-1 plasmid vector and *A. nidulans* with additions of HindIII restriction enzyme sites at 5'-end of gpdA promoter and 3'-end of LaeA gene, confirmed by DNA sequence, and ligated into pPTR1 plasmid vector at HindIII restriction enzyme site.

FIG. 14 is a schematic illustrating a plasmid vector pRS426-LaeA, which contains the upstream region of pyrG gene of *A. niger*, the coding region of LaeA gene under the control of gpdA promoter and transcriptional terminator of trpC gene from *A. nidulans*, the pyrithiamine resistance (ptrA) gene from *A. oryzae*, and the downstream region of pyrG gene of *A. niger*. The unique restriction enzyme PmeI site was introduced at the both end of transgene expression fragment.

FIGS. 15A and 15B show the results of polymerase chain reaction (PCR) analysis of LaeA gene insertion in the transgenic *A. niger* genome of heterologous expression of *A. nidulans* LaeA gene. (FIG. 15A) PCR products of *A. oryzae* ptrA gene detected in selected single spore colony isolate (SCI) of LaeA gene transgenic mutants and parent kusA and Alg3Δ are control strains. (FIG. 15B) PCR products of transgene expression DNA fragment including the gpdA promoter, LaeA coding region and trpC gene transcriptional terminator. The SCI-1 to SCI-4 is the individual single spore colony of LaeA gene transgenic mutants and parent kusA and Alg3Δ are control strains. Lambda DNA marker is the restriction fragment of BstEII restriction enzyme.

FIG. 16 is a bar graph showing the results of citric acid production after 10 days in culture of parent strain (kusA), alg3Δ mutant (Alg3), and over-expression of LaeA gene in alg3Δ (LaeA-1 and LaeA-2) mutants. The data for each strain is the average of three replicates.

FIG. 17A is a schematic diagram showing the construction of the pBSK-LaeA deletion plasmid vector for *A. niger*. The DNA fragments of the 5' and 3' ends of *A. niger* LaeA (lack of aflR expression) gene and the hph (hygromycin phosphotransferase) expression cassette were isolated from

*A. niger* genomic DNA (LaeA) or pCB1003 plasmid vector DNA by PCR with oligonucleotide pairs 1 and 2; 3 and 4; and 5 and 6. The PCR DNA fragments were assembled together by Gibson assembly cloning kit in pBSK backbone vector prepared by HindIII/PstI double digestion. The plasmid DNA fragment containing the LaeA gene deletion cassette was prepared by restriction enzyme digestion with restriction endonucleases of HindIII and XbaI and used for *A. niger* transformation.

FIG. 17B is a schematic diagram showing the construction of the pRS426-*A. niger* LaeA complementation plasmid vector for *A. niger*. The LaeA gene containing both its promoter and transcriptional terminator was isolated by PCR with oligonucleotide pair 9 and 10 from *A. niger* genomic DNA. The PCR fragment was cloned into the plasmid vector pRSB426-LaeA prepared by restriction endonuclease HindIII digestion and Klenow treatment with blunt-end ligation. The new plasmid DNA vector contains the upstream region of the pyrG gene of *A. niger*, the entire region of *A. niger* LaeA gene, the transcriptional terminator of the trpC gene from *A. nidulans*, the pyrithiamine resistance (ptrA) gene from *A. oryzae*, and the downstream region of the pyrG gene of *A. niger*. The unique restriction endonuclease PmeI site was introduced at both ends of the transgene expression fragment.

FIG. 18 shows a digital image of the results of PCR analysis of the LaeA gene deletion in the transgenic *A. niger* genome by homologous recombination at the pyrG locus. The PCR products corresponding to the 5' end of the *A. niger* LaeA gene and part of the hph expression cassette were amplified with oligonucleotide pair 7 and 8, and were detected in selected single spore colony isolate (SCI) of LaeA gene deletion transgenic mutants.

FIG. 19 shows a digital image of the results of PCR analysis of the LaeA gene complementation (cLaeA) in the genetic background of LaeA deletion mutant or *A. nidulans* LaeA over-expression (LaeA) in the genetic background of 11414kusA of those transgenic *A. niger* genome by homologous recombination. PCR products of *A. oryzae* ptrA gene detected in selected single spore colony isolate (SCI) of LaeA gene transgenic mutants and parent kusA (negative) and plasmid DNA of pRS426-laeA (positive) represent control DNA.

FIG. 20 is a bar graph showing the results of citric acid production after 5 days in culture of the parent strain (kusA), the LaeAΔ mutant, *A. niger* LaeA gene complementation in the LaeAΔ mutant (cLaeAΔ), and over-expression of *A. nidulans* LaeA gene in the parent (KusA) strain. The data for each strain is the average of at least three biological replicates.

FIG. 21A is a diagram illustrating the LaeA deletion construct in which the LaeA coding region was replaced by the bacterial hygromycin B phosphotransferase (hph) gene. Oligonucleotides 1546, 108, 107 and 1553 (see Table 6) were used for LaeA gene deletion confirmation in *A. terreus*.

FIG. 21B shows the genomic PCR confirmation of selected LaeA deletion mutants with oligonucleotide pair 1546/107 or 108/1553.

FIG. 22A is a diagram illustrating LaeA complementation in *A. terreus* LaeA deletion mutants. The complementation transgene fragment was targeted upstream of the LaeA gene and the bacterial bleomycin resistance gene (ble) was used for transgenic strain selection. The insertion of LaeA complementation was confirmed by oligonucleotide pair 1556/107.

FIG. 22B shows gel electrophoresis of genomic PCR product corresponding to ble and part of hph.

FIG. 23 is a graph showing itaconic acid production in selected strains of parent, LaeA deletion (LaeAΔ) and LaeA complementation (cLaeAΔ) mutants after 4 days of culture in itaconic acid production medium.

FIG. 24A shows a diagram of the transgene over-expression cassette of the *A. nidulans* LaeA gene under the control of the *A. nidulans* gpdA promoter and TrpC terminator with a locus target of 2.5 kb of the *A. terreus* LaeA gene. The pyrithiamine resistance gene (ptrA) of *A. oryzae* was used for transgene expression selection. Oligonucleotides 1563, 1198 and 1564 (see Table 6) were used for genomic PCR to confirm the gene insertion.

FIG. 24B shows gel electrophoresis of PCR products corresponding to the oligonucleotide pairs 1198/1564 and 1563/1564 in selected transgenic strains.

FIG. 25 is a bar graph showing itaconic acid production in selected *A. terreus* transgenic strains with the *A. nidulans* LaeA over-expression transgene after 5 days of culture in itaconic acid production medium.

#### SEQUENCE LISTING

The nucleic and amino acid sequences listed in the accompanying sequence listing are shown using standard letter abbreviations for nucleotide bases, and three letter code for amino acids, as defined in 37 C.F.R. 1.822. Only one strand of each nucleic acid sequence is shown, but the complementary strand is understood as included by any reference to the displayed strand. The Sequence Listing is submitted as an ASCII text file, created on Oct. 19, 2015, 157 KB, which is incorporated by reference herein. In the accompanying sequence listing:

SEQ ID NOS: 1 and 2 are exemplary Alg3 nucleic acid and protein sequences, respectively, from *A. niger*.

SEQ ID NOS: 3 and 4 are exemplary Alg3 nucleic acid and protein sequences, respectively, from *A. oryzae*.

SEQ ID NOS: 5-30 show exemplary primer sequences.

SEQ ID NOS: 31-36 are exemplary Alg3 protein sequences from *A. nidulans*, *Fusarium oxysporum*, *Arabidopsis thaliana*, *Neurospora crassa*, *S. cerevisiae*, and *Homo sapiens*, respectively.

SEQ ID NO: 37 is an exemplary *Aspergillus nidulans* glyceraldehyde 3-phosphate dehydrogenase (gpdA) promoter sequence.

SEQ ID NOS: 38 and 39 are exemplary forward and reverse primers, respectively, that can be used to isolate or amplify an *A. nidulans* gpdA promoter.

SEQ ID NOS: 40 and 41 are exemplary *Aspergillus nidulans* methyltransferase (LaeA) coding and protein sequences, respectively.

SEQ ID NOS: 42 and 43 are exemplary forward and reverse primers, respectively, that can be used to isolate or amplify an *A. nidulans* LaeA sequence.

SEQ ID NO: 44 is the nucleic acid sequence of the pGPDA-LaeA fragment described in FIG. 13 and Example 5.

SEQ ID NO: 45 is the upstream region of *A. niger* pyrG gene.

SEQ ID NO: 46 is the trpC transcriptional terminator of *A. nidulans*.

SEQ ID NO: 47 is the pyrithiamine resistance gene (ptrA) of *Aspergillus oryzae*.

SEQ ID NO: 48 is the downstream region of *A. niger* pyrG gene.

SEQ ID NOS: 49 and 50 are exemplary forward and reverse primers, respectively, that can be used to isolate or amplify an upstream region of *A. niger* pyrG.

SEQ ID NOS: 51 and 52 are exemplary forward and reverse primers, respectively, that can be used to isolate or amplify the *trpC* transcriptional terminator of *A. nidulans*.

SEQ ID NOS: 53 and 54 are exemplary forward and reverse primers, respectively, that can be used to isolate or amplify *ptrA* of *Aspergillus oryzae*.

SEQ ID NOS: 55 and 56 are exemplary forward and reverse primers, respectively, that can be used to isolate or a downstream region of *A. niger pyrG*.

SEQ ID NO: 57 is the nucleic acid sequence of the transgene fragment described in FIG. 13 and Example 6.

SEQ ID NOS: 58 and 59 are exemplary *Aspergillus niger* *LaeA* coding and protein sequences, respectively.

SEQ ID NO: 60 is the nucleic acid sequence of the transgene fragment used to complement the *alg3Δ* mutant with the original *alg3* gene at *pyrG* locus.

SEQ ID NO: 61 is the nucleic acid sequence of the 5' end of the *A. niger LaeA* gene.

SEQ ID NO: 62 is the nucleic acid sequence of the 3' end of the *A. niger LaeA* gene.

SEQ ID NO: 63 is the nucleic acid sequence of the *hph* expression cassette.

SEQ ID NO: 64 is the nucleic acid sequence of the *A. niger LaeA* gene.

SEQ ID NO: 65 is the nucleic acid sequence of the *LaeA* deletion cassette.

SEQ ID NO: 66 is the nucleic acid sequence of the *LaeA* complementation construct.

SEQ ID NOS: 67-100 are oligonucleotide primers.

SEQ ID NO: 101 is the nucleotide sequence of the 1546/1548 fragment.

SEQ ID NO: 102 is the nucleotide sequence of the 1549/1550 fragment.

SEQ ID NO: 103 is the nucleotide sequence of the 1551/1553 fragment.

SEQ ID NO: 104 is the nucleotide sequence of the 1547/1552 fragment.

SEQ ID NO: 105 is the nucleotide sequence of the 1554/1555 fragment.

SEQ ID NO: 106 is the nucleotide sequence of the 1556/1557 fragment.

SEQ ID NO: 107 is the nucleotide sequence of the 1558/1559 fragment.

SEQ ID NO: 108 is the nucleotide sequence of the 1561/1562 fragment.

SEQ ID NO: 109 is the nucleotide sequence of the 1563/1564 fragment.

SEQ ID NO: 110 is the nucleotide sequence of the 1565/1566 fragment.

#### DETAILED DESCRIPTION

Unless otherwise explained, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. Definitions of common terms in molecular biology may be found in Benjamin Lewin, *Genes V*, published by Oxford University Press, 1994 (ISBN 0-19-854287-9); Kendrew et al. (eds.), *The Encyclopedia of Molecular Biology*, published by Blackwell Science Ltd., 1994 (ISBN 0-632-02182-9); and Robert A. Meyers (ed.), *Molecular Biology and Biotechnology: a Comprehensive Desk Reference*, published by VCH Publishers, Inc., 1995 (ISBN 1-56081-569-8).

The singular terms “a,” “an,” and “the” include plural referents unless context clearly indicates otherwise. Similarly, the word “or” is intended to include “and” unless the

context clearly indicates otherwise. Hence “comprising A or B” means including A, or B, or A and B. It is further to be understood that all base sizes or amino acid sizes, and all molecular weight or molecular mass values, given for nucleic acids or polypeptides are approximate, and are provided for description. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present disclosure, suitable methods and materials are described below. All references and GENBANK™ Accession numbers mentioned herein are incorporated by reference (the sequence available on Nov. 30, 2011). The materials, methods, and examples are illustrative only and not intended to be limiting.

In order to facilitate review of the various embodiments of the disclosure, the following explanations of specific terms are provided:

**Alg3** (dolichyl-P-Man:Man(5)GlcNAc(2)-PP-dolichyl mannosyltransferase): Also known as asparagine-linked glycosylation 3 and  $\alpha$ -1,3-mannosyltransferase. Alg3 encodes an enzyme which catalyzes the addition of the first dol-p-man derived mannose in an  $\alpha$ -1,3 linkage to Man5GlcNAc2-PP-Dol. The term Alg3 (or Alg3) includes any Alg3 gene (such as a fungal Alg3 sequence), cDNA, mRNA, or protein, that is an Alg3 involved in catalyzing the addition of the first dol-p-man derived mannose in an  $\alpha$ -1,3 linkage to Man5GlcNAc2-PP-Dol, and when genetically inactivated results in a fungus that has an ability to produce more citric acid than the parent strain (such as at least 20%, at least 30%, at least 50%, at least 60%, or at least 70% more than a parent strain under the same growing conditions).

Alg3 sequences are publicly available for many species of *Aspergillus*. For example, GENBANK™ Accession Nos: XM\_001823992.2 and XP\_001824044 disclose *Aspergillus oryzae* RIB40 Alg3 nucleic acid and protein sequences, respectively; GENBANK™ Accession Nos: XM\_001398659.2 and XP\_001398696.2 disclose *Aspergillus niger* CBS 513.88 Alg3 nucleic acid and protein sequences, respectively (SEQ ID NOS: 1 and 2); and GENBANK™ Accession Nos: XM\_748359.1 and XP\_753452 disclose *Aspergillus fumigatus* Af293 Alg3 nucleic acid and protein sequences, respectively. Additional exemplary Alg3 sequences are provided in SEQ ID NOS: 1-4 and 31-36. However, one skilled in the art will appreciate that in some examples, an Alg3 sequence can include variant sequences (such as allelic variants and homologs) that retain Alg3 activity but when genetically inactivated in *Aspergillus* results in a fungus that has an ability to produce more citric acid than the parent strain (such as at least 20%, at least 30%, at least 50%, at least 60%, or at least 70% more under the same growing conditions).

***Aspergillus terreus***: A fungus found in soil and vegetation. *A. terreus* is found worldwide, but is more prevalent in warmer climates, such as tropical and subtropical regions. This fungus is commonly used in industry to produce important organic acids, such as itaconic acid and cis-aconitic acid, and was the initial source for the drug mevastatin (lovastatin), a drug for lowering serum cholesterol (Bennett, “An Overview of the Genus *Aspergillus*,” pages 1-17, *Aspergillus Molecular Biology and Genomics*, edited by Masayuki Machida and Katsuyi Gomi, Caister, Academic Press, 2010).

**Detectable**: Capable of having an existence or presence ascertained. For example, production of citric acid is detectable if the signal generated is strong enough to be measurable.

**Genetic enhancement or up-regulation**: When used in reference to the expression of a nucleic acid molecule, such

as a gene, refers to any process which results in an increase in production of a gene product. A gene product can be RNA (such as mRNA, rRNA, tRNA, and structural RNA) or protein. Examples of processes that increase transcription include those that facilitate formation of a transcription initiation complex, those that increase transcription initiation rate, those that increase transcription elongation rate, those that increase processivity of transcription and those that relieve transcriptional repression (for example by blocking the binding of a transcriptional repressor). Gene up-regulation can include inhibition of repression as well as stimulation of expression above an existing level. Examples of processes that increase translation include those that increase translational initiation, those that increase translational elongation and those that increase mRNA stability. In one example, additional copies of genes are introduced into a cell in order to increase expression of that gene in the resulting transgenic cell.

Gene up-regulation includes any detectable increase in the production of a gene product. In certain examples, production of a gene product increases by at least 1.5-fold, at least 2-fold, or at least 5-fold), such as LaeA. For example, a genetic enhancement of a LaeA gene in *Aspergillus* (e.g., *A. niger* or *A. terreus*) results in an *Aspergillus* strain having increased levels of the LaeA protein relative to the parent strain, which can increase the ability of the fungus to produce more citric acid or itaconic acid. Genetic enhancement is also referred to herein as “enhancing or increasing expression.”

Genetic inactivation or down-regulation: When used in reference to the expression of a nucleic acid molecule, such as a gene, refers to any process which results in a decrease in production of a gene product. A gene product can be RNA (such as mRNA, rRNA, tRNA, and structural RNA) or protein. Therefore, gene down-regulation or deactivation includes processes that decrease transcription of a gene or translation of mRNA.

For example, a mutation, such as a substitution, partial or complete deletion, insertion, or other variation, can be made to a gene sequence that significantly reduces (and in some cases eliminates) production of the gene product or renders the gene product substantially or completely non-functional. For example, a genetic inactivation of an Alg3 gene in *Aspergillus* (e.g., *A. niger*) results in *Aspergillus* having a non-functional or non-existent Alg3 protein, which results in an ability of the fungus to produce more citric acid. Genetic inactivation is also referred to herein as “functional deletion”.

Heterologous: Derived from separate genetic sources or species. For example, an LaeA gene that is heterologous to *A. terreus* is an LaeA gene from a species other than *A. terreus*.

Isolated: To be significantly separated from other agents. An “isolated” biological component (such as a nucleic acid molecule or protein) has been substantially separated, produced apart from, or purified away from other biological components in the cell of the organism in which the component occurs, for example, other chromosomal and extra-chromosomal DNA and RNA, and proteins. Nucleic acid molecules and proteins which have been “isolated” include nucleic acid molecules and proteins purified by standard purification methods. The term also embraces nucleic acid molecules and proteins prepared by recombinant expression in a host cell as well as chemically synthesized proteins and nucleic acids. Samples of isolated biological components include samples of the biological component wherein the

biological component represents greater than 90% (for example, greater than 95%, such as greater than 98%) of the sample.

An “isolated” microorganism (such as an Alg3Δ strain of *Aspergillus*) has been substantially separated or purified away from microorganisms of different types, strains, or species. Microorganisms can be isolated by a variety of techniques, including serial dilution and culturing and resistance to certain chemicals.

Itaconic acid: An unsaturated dicarbonic acid; also known as 2-methylidenebutanedioic acid. Itaconic acid can be produced by the fermentation of carbohydrates using *Aspergillus terreus*. Itaconic acid can be used as a monomer for the production of a number of different products, including resins, plastics, acrylate latexes, paints and synthetic fibers.

Itaconic acid production medium: Media containing in grams per liter (g/l): glucose, 100; ammonium sulfate, 2.36; potassium phosphate dibase, 0.11; magnesium sulfate heptahydrate, 2.08; calcium chloride dihydrate, 0.13; sodium chloride, 0.074; copper sulfate pentahydrate  $2 \times 10^{-4}$ ; ferrous sulfate heptahydrate,  $5.5 \times 10^{-3}$ ; manganese chloride tetrahydrate,  $7 \times 10^{-4}$ ; and zinc sulfate heptahydrate,  $1.3 \times 10^{-3}$ .

LaeA (loss of afiR expression A): LaeA encodes a protein which regulates secondary metabolite production in *Aspergillus*. The term LaeA (or LaeA) includes any LaeA gene (such as a fungal LaeA sequence), cDNA, mRNA, or protein, that is an LaeA involved in secondary metabolite production. In some embodiments, when LaeA expression is increased, for example in combination with a genetically inactivated Alg3 gene, it results in a fungus that has an ability to produce more citric acid than the parent strain (such as at least 20%, at least 30%, at least 40%, 50%, at least 60%, or at least 70% more than a parent strain under the same growing conditions). In some embodiments, when LaeA expression is increased, for example in *A. terreus*, it results in a fungus that has an ability to produce more itaconic acid than the parent strain (such as at least 20%, at least 30%, at least 40%, 50%, at least 60%, or at least 70% more than a parent strain under the same growing conditions).

LaeA sequences are publicly available for many species of *Aspergillus*. For example, GENBANK™ Accession Nos: AB267276 and BAF74528.1 disclose *Aspergillus oryzae* LaeA nucleic acid and protein sequences, respectively; GENBANK™ Accession No. EHA27020.1 discloses an exemplary *Aspergillus niger* ATCC1015 LaeA protein sequence, a parent strain of 11414kusA (other exemplary sequences are provided in SEQ ID NOS: 58 and 59); GENBANK™ Accession No: CBF88745 discloses an *Aspergillus nidulans* LaeA protein sequence; and GENBANK™ Accession Nos: AY422723 and AAR01218 disclose *Aspergillus fumigatus* LaeA nucleic acid and protein sequences, respectively. Additional exemplary LaeA sequences are provided in SEQ ID NOS: 40-41 and 58-59.

However, one skilled in the art will appreciate that in some examples, an LaeA sequence can include variant sequences (such as allelic variants and homologs) that retain LaeA activity and when genetically up-regulated in *Aspergillus* (for example with addition of copy or in combination with Alg3Δ) results in a fungus that has an ability to produce more citric acid than the parent strain (such as at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, or at least 70% more under the same growing conditions).

Mutation: A change in a nucleic acid sequence (such as a gene sequence) or amino acid sequence, for example as compared to a nucleic acid or amino acid sequence present in a wild-type or native organism. In particular examples, a

mutation is introduced into an Alg3 gene in *Aspergillus*. Mutations can occur spontaneously, or can be introduced, for example using molecular biology methods. In particular examples, a mutation includes one or more nucleotide substitutions, deletions, insertions, or combinations thereof. In particular examples, the presence of one or more mutations in a gene can significantly inactivate that gene.

**Operably linked:** A first nucleic acid sequence is operably linked to a second nucleic acid sequence where the first nucleic acid sequence is joined in a functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Generally, operably linked DNA sequences are contiguous and, where necessary to join two protein-coding regions, in the same reading frame.

**Promoter:** An array of nucleic acid sequences that governs transcription of a nucleic acid. A promoter includes necessary nucleic acid sequences to initiate gene transcription. A promoter also optionally includes distal enhancer or repressor elements which can be located as much as several thousand base pairs from the start site of transcription.

**Recombinant:** A recombinant nucleic acid molecule or protein is one that has a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two otherwise separated segments of sequence. In particular examples, this artificial combination is accomplished by chemical synthesis or by the artificial manipulation of isolated segments of nucleic acids, for example, by genetic engineering techniques such as those described in Sambrook et al. (ed.), *Molecular Cloning: A Laboratory Manual*, 3d ed., vol. 1-3, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 2001. The term recombinant includes nucleic acid molecules that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid molecule.

**Sequence identity/similarity:** The identity/similarity between two or more nucleic acid sequences, or two or more amino acid sequences, is expressed in terms of the identity or similarity between the sequences. Sequence identity can be measured in terms of percentage identity; the higher the percentage, the more identical the sequences are. Sequence similarity can be measured in terms of percentage similarity (which takes into account conservative amino acid substitutions); the higher the percentage, the more similar the sequences are.

Methods of alignment of sequences for comparison are well known in the art. Various programs and alignment algorithms are described in: Smith & Waterman, *Adv. Appl. Math.* 2:482, 1981; Needleman & Wunsch, *J. Mol. Biol.* 48:443, 1970; Pearson & Lipman, *Proc. Natl. Acad. Sci. USA* 85:2444, 1988; Higgins & Sharp, *Gene*, 73:237-44, 1988; Higgins & Sharp, *CABIOS* 5:151-3, 1989; Corpet et al., *Nuc. Acids Res.* 16:10881-90, 1988; Huang et al. *Computer Appls. in the Biosciences* 8, 155-65, 1992; and Pearson et al., *Meth. Mol. Bio.* 24:307-31, 1994. Altschul et al., *J. Mol. Biol.* 215:403-10, 1990, presents a detailed consideration of sequence alignment methods and homology calculations.

The NCBI Basic Local Alignment Search Tool (BLAST) (Altschul et al., *J. Mol. Biol.* 215:403-10, 1990) is available from several sources, including the National Center for Biological Information (NCBI, National Library of Medicine, Building 38A, Room 8N805, Bethesda, Md. 20894) and on the Internet, for use in connection with the sequence analysis programs blastp, blastn, blastx, tblastn and tblastx. Additional information can be found at the NCBI web site.

BLASTN is used to compare nucleic acid sequences, while BLASTP is used to compare amino acid sequences. To compare two nucleic acid sequences, the options can be set as follows: -i is set to a file containing the first nucleic acid sequence to be compared (e.g., C:\seq1.txt); -j is set to a file containing the second nucleic acid sequence to be compared (e.g., C:\seq2.txt); -p is set to blastn; -o is set to any desired file name (e.g., C:\output.txt); -q is set to -1; -r is set to 2; and all other options are left at their default setting. For example, the following command can be used to generate an output file containing a comparison between two sequences: C:\B12seq -i c:\seq1.txt -j c:\seq2.txt -p blastn -o c:\output.txt -q -1 -r 2.

To compare two amino acid sequences, the options of B12seq can be set as follows: -i is set to a file containing the first amino acid sequence to be compared (e.g., C:\seq1.txt); -j is set to a file containing the second amino acid sequence to be compared (e.g., C:\seq2.txt); -p is set to blastp; -o is set to any desired file name (e.g., C:\output.txt); and all other options are left at their default setting. For example, the following command can be used to generate an output file containing a comparison between two amino acid sequences: C:\B12seq -i c:\seq1.txt -j c:\seq2.txt -p blastp -o c:\output.txt. If the two compared sequences share homology, then the designated output file will present those regions of homology as aligned sequences. If the two compared sequences do not share homology, then the designated output file will not present aligned sequences.

Once aligned, the number of matches is determined by counting the number of positions where an identical nucleotide or amino acid residue is presented in both sequences. The percent sequence identity is determined by dividing the number of matches either by the length of the sequence set forth in the identified sequence, or by an articulated length (e.g., 100 consecutive nucleotides or amino acid residues from a sequence set forth in an identified sequence), followed by multiplying the resulting value by 100. For example, a nucleic acid sequence that has 1166 matches when aligned with a test sequence having 1554 nucleotides is 75.0 percent identical to the test sequence (i.e.,  $1166 \div 1554 * 100 = 75.0$ ). The percent sequence identity value is rounded to the nearest tenth. For example, 75.11, 75.12, 75.13, and 75.14 are rounded down to 75.1, while 75.15, 75.16, 75.17, 75.18, and 75.19 are rounded up to 75.2. The length value will always be an integer. In another example, a target sequence containing a 20-nucleotide region that aligns with 20 consecutive nucleotides from an identified sequence as follows contains a region that shares 75 percent sequence identity to that identified sequence (i.e.,  $15 \div 20 * 100 = 75$ ).

For comparisons of amino acid sequences of greater than about 30 amino acids, the Blast 2 sequences function is employed using the default BLOSUM62 matrix set to default parameters, (gap existence cost of 11, and a per residue gap cost of 1). Homologs are typically characterized by possession of at least 70% sequence identity counted over the full-length alignment with an amino acid sequence using the NCBI Basic Blast 2.0, gapped blastp with databases such as the nr or swissprot database. Queries searched with the blastn program are filtered with DUST (Hancock and Armstrong, 1994, *Comput. Appl. Biosci.* 10:67-70). Other programs use SEG. In addition, a manual alignment can be performed. Proteins with even greater similarity will show increasing percentage identities when assessed by this method, such as at least 75%, 80%, 85%, 90%, 95%, or 99% sequence identity.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode identical or similar (conserved) amino acid sequences, due to the degeneracy of the genetic code. Changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid molecules that all encode substantially the same protein. Such homologous nucleic acid sequences can, for example, possess at least 60%, 70%, 80%, 90%, 95%, 98%, or 99% sequence identity determined by this method.

One of skill in the art will appreciate that these sequence identity ranges are provided for guidance only; it is possible that strongly significant homologs could be obtained that fall outside the ranges provided.

Transformed: A cell, such as a fungal cell, into which a nucleic acid molecule has been introduced, for example by molecular biology methods known in the art. As used herein, the term transformation encompasses all techniques by which a nucleic acid molecule might be introduced into such a cell, including, but not limited to transfection with viral vectors, conjugation, transformation with plasmid vectors, and introduction of naked DNA by chemical-mediated, electroporation, lipofection, and biolistic particle delivery.

#### Overview

Disclosed herein is the finding that overexpression of the LaeA gene in *A. terreus* results in a significant increase in production of itaconic acid. In particular, *A. terreus* transformed with a nucleic acid construct comprising an *Aspergillus* species LaeA gene produces significantly more itaconic acid relative to *A. terreus* that is not transformed with the construct.

Providing herein are isolated *A. terreus* fungi transformed with a heterologous nucleic acid construct comprising an *Aspergillus* species LaeA (loss of aflR expression A) gene. Expression of LaeA is increased in the transformed fungus compared to an *A. terreus* fungus that is not transformed with the heterologous nucleic acid construct.

In some embodiments, the heterologous nucleic acid construct comprises a heterologous LaeA gene, a heterologous promoter, a heterologous transcription terminator, a heterologous selective marker gene, or any combination thereof.

In some embodiments, the *Aspergillus* species LaeA gene is a homologous LaeA gene (i.e. an *A. terreus* LaeA gene). In other embodiments, the LaeA gene is a heterologous LaeA gene, such as an LaeA gene from *A. nidulans*, *A. niger*, *A. oryzae*, *A. fumigatus* or another *Aspergillus* species. In some examples, the *A. nidulans* LaeA gene encodes a protein having at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% sequence identity to the amino acid sequence of SEQ ID NO: 41. In non-limiting examples, the *A. nidulans* LaeA gene encodes a protein comprising the amino acid sequence of SEQ ID NO: 41. In some examples, the *A. niger* LaeA gene encodes a protein having at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identity to the amino acid sequence of SEQ ID NO: 59. In non-limiting examples, the *A. niger* LaeA gene encodes a protein comprising the amino acid sequence of SEQ ID NO: 59.

In some embodiments, the heterologous nucleic acid molecule comprises a promoter operably linked to an *Aspergillus* species LaeA gene, a transcription terminator and/or a selective marker gene. In particular examples, the promoter is a heterologous promoter, such as, but not limited to, the *A. nidulans* gpdA promoter, or any other promoter that is

functional in *Aspergillus* (e.g., the actA, ubi4, Arsa-7, A-37, Brsa-109, gpdA, pyrG or tefl $\alpha$  promoters from *A. niger*). In other particular examples, the *Aspergillus* species LaeA gene is a heterologous LaeA gene, such as, but not limited to, an *A. nidulans*, *A. niger*, *A. oryzae*, or *A. fumigatus* LaeA gene. In other particular examples, the transcription terminator is a heterologous transcription terminator, such as, but not limited to, the *A. nidulans* TrpC transcription terminator. As an alternative to trpC, other transcriptional terminators can be used, such as promoters which include a transcriptional terminator (e.g., actA, Arsa7, Arsa-37, gpdA, pyrG, tefl $\alpha$  or ubi4). In yet other particular examples, the selective marker is a heterologous selective marker gene, such as a heterologous antibiotic resistance gene. In specific examples, selective marker gene is an *A. oryzae* pyrithiamine resistance (ptrA) gene. Other selective marker genes include, for example, the hygromycin B phosphotransferase (hph) gene, the bar gene (which confers glufosinate resistance), or bleomycin resistance gene (ble), or auxotrophic markers, such as pyrG.

In specific non-limiting examples, the heterologous nucleic acid molecule comprises the nucleotide sequence of SEQ ID NO: 109.

Further provided herein is a method of making itaconic acid. In some embodiments, the includes culturing an isolated *A. terreus* fungus transformed with a heterologous nucleic acid molecule comprising an *Aspergillus* species LaeA gene under conditions that permit the fungus to make itaconic acid. Expression of LaeA is increased in the transformed fungus compared to an *A. terreus* fungus that is not transformed with the heterologous nucleic acid construct. In some embodiments, the fungus is cultured in itaconic acid production medium, which contains grams per liter (g/l): glucose, 100; ammonium sulfate, 2.36; potassium phosphate dibase, 0.11; magnesium sulfate heptahydrate, 2.08; calcium chloride dihydrate, 0.13; sodium chloride, 0.074; copper sulfate pentahydrate  $2 \times 10^{-4}$ ; ferrous sulfate heptahydrate,  $5.5 \times 10^{-3}$ ; manganese chloride tetrahydrate,  $7 \times 10^{-4}$ ; and zinc sulfate heptahydrate,  $1.3 \times 10^{-3}$ .

In some embodiments of the methods of making itaconic acid, the heterologous nucleic acid construct comprises a heterologous LaeA gene, a heterologous promoter, a heterologous transcription terminator, a heterologous selective marker gene, or any combination thereof.

In some embodiments of the methods, the *Aspergillus* species LaeA gene is a homologous LaeA gene (i.e. an *A. terreus* LaeA gene). In other embodiments, the LaeA gene is a heterologous LaeA gene, such as an LaeA gene from *A. nidulans*, *A. niger*, *A. oryzae*, *A. fumigatus* or another *Aspergillus* species. In some examples, the *A. nidulans* LaeA gene encodes a protein having at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% sequence identity to the amino acid sequence of SEQ ID NO: 41. In non-limiting examples, the *A. nidulans* LaeA gene encodes a protein comprising the amino acid sequence of SEQ ID NO: 41. In some examples, the *A. niger* LaeA gene encodes a protein having at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identity to the amino acid sequence of SEQ ID NO: 59. In non-limiting examples, the *A. niger* LaeA gene encodes a protein comprising the amino acid sequence of SEQ ID NO: 59.

In some embodiments of the method, the heterologous nucleic acid molecule comprises a promoter operably linked to an *Aspergillus* species LaeA gene, a transcription terminator and/or a selective marker gene. In particular examples, the promoter is a heterologous promoter, such as, but not

limited to, the *A. nidulans* *gpdA* promoter. In other particular examples, the *Aspergillus* species *LaeA* gene is a heterologous *LaeA* gene, such as, but not limited to, an *A. nidulans*, *A. niger*, *A. oryzae*, or *A. fumigatus* *LaeA* gene. In other particular examples, the transcription terminator is a heterologous transcription terminator, such as, but not limited to, the *A. nidulans* *TrpC* transcription terminator. In yet other particular examples, the selective marker is a heterologous selective marker gene, such as a heterologous antibiotic resistance gene. In specific examples, the selective marker gene is an *A. oryzae* pyrithiamine resistance (*ptrA*) gene. In specific non-limiting examples, the heterologous nucleic acid molecule comprises the nucleotide sequence of SEQ ID NO: 109.

Also provided by the present disclosure are compositions that include isolated *A. terreus* fungi transformed with a heterologous nucleic acid construct. The compositions may further include growth medium. Further provided by the present disclosure are kits that include isolated *A. terreus* fungi transformed with a heterologous nucleic acid construct, such as a kit that includes a medium for culturing, storing, or growing the fungus, and/or antibiotics (for selection).

This disclosure also provides the first demonstration that genetic inactivation of *Alg3*, a gene involved in protein N-linked glycosylation, can result in substantial improvement of citric acid production in *A. niger*, while the total biomass is similar to the parent strain. The core oligosaccharide *Glc3Man9GlcNAc2* is synthesized by a series of membrane-bound glycosyltransferases, which begins on the cytoplasmic side of the membrane of the endoplasmic reticulum (ER) and flips into the luminal side of the ER membrane to complete its synthesis. The lipid-linked core *Glc3Man9GlcNAc2* is subsequently transferred to a nascent protein in the ER, where the glycoproteins are folded and then shuttled to the Golgi for additional, but divergent processing. The *Alg3* gene encodes the enzyme  $\alpha$ -1,3-mannosyltransferase that converts *Man5GlcNAc2-Dol-PP* to *Man6GlcNAc2-Dol-PP* on the ER membrane of the luminal side. Provided herein is a homolog of *Saccharomyces cerevisiae* *Alg3* identified from *Aspergillus niger* (e.g., see SEQ ID NOS: 1 and 2).

It is shown herein that genetic inactivation of *Alg3* in *A. niger* resulted in a significant reduction of growth on complete medium (CM) and potato dextrose agar medium (PDA), but no effect on minimal medium (MM). The *Alg3* deletion also caused the substantial reduction in spore production of *A. niger* on CM, but no significant change on the PDA. When the spores were germinated in CM or PDA liquid culture medium, the *Alg3* $\Delta$  strain showed pronounced delay in spore germination. This growth phenotype is similar to the mutants with defects in signal transduction pathways observed in *A. nidulans* and *A. niger* (Fillinger et al., *Mol. Microbiol.* 44(4):1001-16, 2002; Saudohar et al., *Microbiol.* 148(8):2635-45, 2002; Xue et al., *Eukaryot Cell* 3(2):557-60, 2004). Deletion of *pkaA*, *cycaA* or *schA/pkaA* in *A. nidulans* substantially reduces its growth on CM medium plates and spore germination rate in MM liquid culture medium (Fillinger et al., *Mol. Microbiol.* 44(4):1001-1016, 2002) and similar growth phenotypes were observed in the strains with the deletion of *pkaR*, *pkaC* or double deletion of *pkaR/pkaC* in *A. niger* (Saudohar et al., *Microbiol.* 148(8):2635-2645, 2002). However, functional deletion of the MAP kinase *SakA* in *A. fumigatus* delays the spore germination in liquid CM, but stimulates spore germination in MM liquid medium (Xue et al., *Eukaryot Cell* 3(2): 557-560, 2004).

Furthermore, the *Alg3* deletion reduced the overall growth on citric acid production (CAP) medium plates at different pHs. In contrast, the *Alg3* deletion triggered early spore germination and substantially improved spore germination rate in CAP liquid culture medium. Citric acid production in CAP liquid culture medium was significantly improved in *A. niger*. When the *alg3* $\Delta$  mutant was complemented with the original *alg3* gene at *pyrG* locus (FIG. 8A; SEQ ID NO: 60), its transcription levels was similar to parent strain with the cycle threshold (Ct) values about 23, while the Ct value for *alg3* $\Delta$  mutant was 30.8 in CAP liquid culture conditions, which was determined by real-time reverse-transcription PCR. Consequently, the citric acid production in the resulted complemented mutant strains was similar to the parent strain as shown in FIG. 8B. The results shown herein demonstrate the involvement of *Alg3* on the growth and development and citric acid production in *A. niger*.

It is proposed that inactivation of *Alg3* influences the N-glycosylation of those proteins involving in signal transduction pathways. The N-glycosylation consensus sequence (N-glycosite) for N-glycosylation in those proteins from the signal transduction pathways was observed. Most of those proteins contained 1 to 7 N-glycosites, such as, 6 N-glycosites found in *sskB* (map kinase kinase), 7 in *Ste11/SteC*, 5 in *acyA*, 5 in *rgsA*, 6 in *rgsC*, 4 in *gprA*, 4 in *pkaC2*, 4 in *flbA* and 3 in *G $\beta$* . Comparison of these results with previous studies indicates that the effects of the *Alg3* deletion on spore germination and growth may be regulated by altering the N-glycosylation in those proteins involved in signal transduction pathways in *A. niger*.

When the *Alg3* $\Delta$  strain was grown on CM medium, spore production of *Alg3* $\Delta$  mutants was dramatically reduced as compared to the parent strain, while maintaining a similar level when grown on PDA medium. This phenotype of sporulation production may be influenced by both endogenous and exogenous factors. For example, protein glycosylation was greatly influenced by culture conditions in filamentous fungi, such as fully glycosylated *Ce17A* only isolated from MM culture medium (Stals et al., *Glycobiology* 14(8):725-737, 2004). In addition, higher amounts of proteases were secreted by the *Alg3* $\Delta$  strain than the parent in liquid MM culture supplemented 1 g/l yeast extract, which further influenced nutrient uptakes, cellular formation and overall N-glycosylation. This would alter the yield and N-glycosylation in G protein system in *A. niger*, where G protein signaling is crucial for detection of major environmental stimuli for food acquisition, asexual sporulation, and spore germination (Chang et al., *Genetics* 167(3):305, 2004; Li et al., *Annu. Rev. Microbiol.* 61:423-452, 2007).

The spores of parent strain germinated more slowly and had a lower germination rate than the *Alg3* $\Delta$  strain in CAP liquid culture medium, which contains limited nitrogen source (3.1 g/l of  $\text{NH}_4\text{NO}_3$ ), similar to MM. A similar phenotype was observed when the stress activating kinase, a MAP kinase, was deleted in *A. fumigatus* (*SakA* $\Delta$  strain) and grown in MM liquid culture medium (Xue et al., *Eukaryot Cell* 3(2): 557-560, 2004). The spore germination of *SakA* $\Delta$  strain was dramatically influenced by nitrogen sources. For example, similar rates of spore germination between parent and *SakA* $\Delta$  strains were observed on MM containing 10 mM  $\text{NH}_4\text{Cl}$  or 10 mM Pro, while the spore germination rates of *SakA* $\Delta$  strain was much higher than the parent strain in the MM culture medium containing 10 mM  $\text{NaNO}_3$ ,  $\text{NaNO}_2$ , or Phe. In addition, the CAP medium contains high level of glucose and low pH, which contributes additional stresses to *A. niger* growth. Although the

Alg3Δ strain had earlier and higher germination in CAP medium, its biomass formation was less than the parent strain at early stages. The dried biomass yields for both parent and Alg3Δ strains were similar after growth in CAP medium for four and half days. However, more citric acid was produced by the Alg3Δ strain than the parent strain. This indicates more glucose was directly converted to citric acid by influence citric acid metabolism and reduction of glucose consumption for complex N-glycan formation and sequentially for other cellular metabolisms.

This disclosure also provides the first demonstration that genetic inactivation of Alg3, in combination with an increase in expression of the loss of aflR expression A (LaeA) gene, can result in substantial improvement of citric acid production in *A. niger*. It is proposed that increased expression of LaeA can also improve citric acid production in *A. niger* or other filamentous fungi. To increase expression of LaeA in fungal cells, a transgene was generated and expressed in *A. niger* as follows. The LaeA gene of *A. nidulans* was operably controlled by glyceraldehyde 3-phosphate dehydrogenase (gpdA) promoter and trpC transcriptional terminator (TrpC) of *A. nidulans*. This chimeric gene was flanked with the upstream of *A. niger* pyrG gene, the pyrithiamine resistance (ptrA) gene of *A. oryzae* and the downstream of *A. niger* pyrG gene. The transgene expression fragment containing the chimeric gene was used to transform the protoplasts of alg3Δ mutants of *A. niger*.

The present disclosure also describes the generation of an *A. niger* strain having a deletion of the LaeA gene (LaeAΔ), as well as an *A. niger* LaeAΔ strain complemented with a transgene encoding LaeA (c1LaeAΔ) at the pyrG locus. These strains were used to evaluate the role of LaeA expression on citric acid production in *Aspergillus*. The data disclosed herein demonstrates that deletion of the LaeA gene in *A. niger* (LaeAΔ) results in a loss of citric acid production in culture. When the original *A. niger* LaeA gene was used for complementation in the LaeAΔ mutant (c1LaeAΔ) at the pyrG locus, citric acid production was partially recovered, which indicates the importance of the chromosomal location of LaeA gene. When *A. nidulans* LaeA was over-expressed in the *A. niger* parent strain, citric acid production was higher than the parent strain. These results indicate that LaeA enhances citric acid production in *A. niger*.

In summary, the deletion of Alg3, increasing expression of LaeA, or both, can be used to increase citric acid production in fungi (such as filamentous fungi, e.g., *A. niger*). In addition, deletion of Alg3 alters the overall N-glycosylation and further influences the spore germination, filamentous growth, sporulation and other organic acid production in *A. niger*.

#### Alg3Δ Fungi

The present disclosure provides isolated fungi having its Alg3 gene inactivated, wherein such inactivation results in increased citric acid production by the fungi. Such fungi are referred to herein as Alg3Δ fungi. It is disclosed herein that genetic inactivation of Alg3 results in *Aspergillus* fungi that can increase citric acid production as compared to *Aspergillus* having a native Alg3 sequence.

Contemplated herein are isolated fungi containing a genetic inactivation of a dolichyl-P-Man:Man(5)GlcNAc(2)-PP-dolichyl mannosyltransferase gene (Alg3). Any fungus can be used, such as any genus or variety of *Aspergillus*. In particular examples, the disclosed *Aspergillus* fungus is *A. niger*, such as *Aspergillus niger* strain 11414 (American Type Culture Collection (ATCC) No. 11414; NRRL 2270);

1015 (ATCC No. 1015; NRRL 328, CBS 113.46); NRRL 3 (ATCC No. 9029, CBS 120.49, N400); NRRL 3122 (ATCC No. 22343); or 11414KusA-. In other specific examples, the *Aspergillus* is *A. aculeatus*, *A. awamori*, *A. carbonarius*, *A. wentii*, *A. foetidus*, *A. oryzae*, *A. terreus*, or *A. fumigatus*.

In addition, any method for genetic inactivation can be used, as long as the expression of the gene is significantly reduced or eliminated, or the function of the expressed protein is significantly reduced or eliminated. In particular examples, the Alg3 gene is genetically inactivated by complete or partial deletion mutation or by insertional mutation. In some examples genetic inactivation need not be 100% genetic inactivation. In some embodiments, genetic inactivation refers to at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, or at least 95% gene or protein inactivation. The term "reduced" or "decreased" as used herein with respect to a cell and a particular gene or protein activity refers to a lower level of activity than that measured in a comparable cell of the same species. For example, a particular fungi lacking Alg3 activity has reduced Alg3 activity if a comparable fungi not having an Alg3 genetic inactivation has detectable Alg3 activity.

Alg3 sequences are disclosed herein and others are publicly available, for example from GENBANK™ or EMBL. In some examples, the Alg3 gene functionally deleted encodes a protein having at least 80%, at least 90%, at least 95%, at least 97%, or at least 98% sequence identity to SEQ ID NO: 2, 4, 31, 32, 33, 34, 35, or 36. In some examples, the Alg3 gene functionally deleted comprises at least 80%, at least 90%, at least 95%, at least 97%, or at least 98% sequence identity to SEQ ID NO: 1 or 3 or nucleotides 1186-2582 of SEQ ID NO: 1.

The inactivation of Alg3 results in many phenotypes in the fungi. For example, Alg3Δ mutants can have one or more of the following phenotypes: slower growth on citric acid production (CAP) medium, earlier spore germination in CAP medium (for example germination in at least 3 hours, at least 4 hours, or at least 5 hours after inoculation, such as within 3 hours of inoculation), increased spore germination rate in CAP medium, increased citric acid production in CAP medium, slower growth on complete medium (CM) or potato dextrose (PDA) medium, delay initiation of spore germination in CM or PDA medium, reduced sporulation on CM, or combinations thereof.

Such changes (such as increases or decreases) can be relative to a fungi having a wild-type Alg3 gene, such as a parental strain (e.g., *A. niger* strain 11414KusA), grown under the same conditions as the Alg3Δ mutant. In some examples, an increased germination rate is germination of at least 20%, at least 25%, or at least 30% of the spores from an Alg3Δ fungus have germinated 8 hours after inoculation in CAP medium (such as 20% to 35%, such as 32%), as compared to no more than 20%, no more than 15%, or no more than 10% (such as 5 to 15%, or 10%) for *A. niger* strain 11414KusA. In some examples, an increased germination rate is germination of at least 80%, at least 85%, or at least 90% of the spores from an Alg3Δ fungus have germinated 15 hours after inoculation in CAP medium (such as 80% to 95%, such as 90%), as compared to no more than 60%, no more than 65%, or no more than 75% (such as 55 to 65%, or 60%) for *A. niger* strain 11414KusA. In some examples, increased citric acid production in CAP medium is an increase of at least 10%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 65%, or at least 70%, by an Alg3Δ fungus as compared to *A. niger* strain 11414KusA. In some examples, reduced sporulation on complete medium is a reduction of sporulation by at least



20%, at least 30%, at least 40%, at least 50%, or at least 60%, (such as a 40% to 60% reduction) by an Alg34 fungus as compared to *A. niger* strain 11414KusA.

One skilled in the art will appreciate that additional genes can also be inactivated, wherein the additional genes may or may not provide additional enhancement of citric acid production to the fungus. In one example KusA (e.g., GENBANK™ Accession No. EF061656) is also genetically inactivated.

Also provided by the present disclosure are compositions that include isolated Alg3Δ fungi, such as a growth medium. Also provided by the present disclosure are kits that include isolated Alg3Δ fungi, such as a kit that includes a medium for culturing, storing, or growing the fungus. Exemplary mediums include solid medium (such as those containing agar, for example CM, PDA or MM) and liquid media (such as a fermentation broth, such as CM, MM, or CAP medium).

#### A. Methods of Functionally Deleting Genes

As used herein, an “inactivated” or “functionally deleted” gene means that the gene has been mutated, for example by insertion, deletion, or substitution (or combinations thereof) of one or more nucleotides such that the mutation substantially reduces (and in some cases abolishes) expression or biological activity of the encoded gene product. The mutation can act through affecting transcription or translation of the gene or its mRNA, or the mutation can affect the polypeptide product itself in such a way as to render it substantially inactive.

Genetic inactivation of one or more genes (which in some examples is also referred to as functional deletion) can be performed using any conventional method known in the art. In one example, a strain of *Aspergillus* is transformed with a vector which has the effect of down-regulating or otherwise inactivating an Alg3 gene. This can be done by mutating control elements such as promoters and the like which control gene expression, by mutating the coding region of the gene so that any protein expressed is substantially inactive, or by deleting the Alg3 gene entirely. For example, an Alg3 gene can be functionally deleted by complete or partial deletion mutation (for example by deleting a portion of the coding region of the gene) or by insertional mutation (for example by inserting a sequence of nucleotides into the coding region of the gene, such as a sequence of about 1-5000 nucleotides). Thus, the disclosure in some examples provides transformed fungi that include at least one exogenous nucleic acid molecule which genetically inactivates an Alg3 gene (such as a nucleic acid sequence encoding SEQ ID NO: 2 or 4). In one example, such a transformed cell produces more citric acid, for example relative to a comparable fungus with a native Alg3 sequence.

In particular examples, an insertional mutation includes introduction of a sequence that is in multiples of three bases (e.g., a sequence of 3, 9, 12, or 15 nucleotides) to reduce the possibility that the insertion will be polar on downstream genes. For example, insertion or deletion of even a single nucleotide that causes a frame shift in the open reading frame, which in turn can cause premature termination of the encoded Alg3 polypeptide or expression of a substantially inactive polypeptide. Mutations can also be generated through insertion of foreign gene sequences, for example the insertion of a gene encoding antibiotic resistance (such as hygromycin or bleomycin).

In one example, genetic inactivation is achieved by deletion of a portion of the coding region of the Alg3 gene. For example, some, most (such as at least 50%) or virtually the entire coding region can be deleted. In particular examples, about 5% to about 100% of the gene is deleted, such as at

least 20% of the gene, at least 40% of the gene, at least 75% of the gene, or at least 90% of the Alg3 gene.

Deletion mutants can be constructed using any of a number of techniques known in the art. In one example, allelic exchange is employed to genetically inactivate one or more genes in *Aspergillus*. A specific example of such a method is described in Example 2 below.

In one example, a strategy using counterselectable markers can be employed which has been utilized to delete genes. For a review, see Reyrat et al. (*Infec. Immun.* 66:4011-4017, 1998). In this technique, a double selection strategy is employed wherein a plasmid is constructed encoding both a selectable and counterselectable marker, with flanking DNA sequences derived from both sides of the desired deletion. The selectable marker is used to select for fungi in which the plasmid has integrated into the genome in the appropriate location and manner. The counterselectable marker is used to select for the very small percentage of fungi that have spontaneously eliminated the integrated plasmid. A fraction of these fungi will then contain only the desired deletion with no other foreign DNA present.

In another technique, the cre-lox system is used for site specific recombination of DNA (for example see Steiger et al., *Appl. Environ. Microbiol.* 77(1):114, 2011). The system includes 34 base pair lox sequences that are recognized by the bacterial cre recombinase gene. If the lox sites are present in the DNA in an appropriate orientation, DNA flanked by the lox sites will be excised by the cre recombinase, resulting in the deletion of all sequences except for one remaining copy of the lox sequence. Using standard recombination techniques, the targeted gene of interest (e.g., Alg3) can be deleted in the *Aspergillus* genome and to replace it with a selectable marker (for example a gene coding for kanamycin resistance) that is flanked by the lox sites. Transient expression (by electroporation of a suicide plasmid containing the cre gene under control of a promoter that functions in *Aspergillus*) of the cre recombinase should result in efficient elimination of the lox flanked marker. This process will produce a mutant containing the desired deletion mutation and one copy of the lox sequence.

In another method, an Alg3 gene sequence in the *Aspergillus* genome is replaced with a marker gene, such as green fluorescent protein, β-galactosidase, or luciferase. In this technique, DNA segments flanking a desired deletion are prepared by PCR and cloned into a suicide (non-replicating) vector for *Aspergillus*. An expression cassette, containing a promoter active in *Aspergillus* and the appropriate marker gene, is cloned between the flanking sequences. The plasmid is introduced into wild-type *Aspergillus*. Fungi that incorporate and express the marker gene are isolated and examined for the appropriate recombination event (replacement of the wild type Alg3 gene with the marker gene).

Thus, for example, a fungal cell can be engineered to have a disrupted Alg3 gene using common mutagenesis or knock-out technology. (Methods in Yeast Genetics (1997 edition), Adams, Gottschling, Kaiser, and Sterns, Cold Spring Harbor Press, 1998; Datsenko and Wanner, *Proc. Natl. Acad. Sci. USA* 97: 6640-5, 2000; and Dai et al., *Appl. Environ. Microbiol.* 70(4):2474-85, 2004). Alternatively, antisense technology can be used to reduce or eliminate the activity of Alg3. For example, a fungal cell can be engineered to contain a cDNA that encodes an antisense molecule that prevents Alg3 from being translated. The term “antisense molecule” encompasses any nucleic acid molecule or nucleic acid analog (e.g., peptide nucleic acids) that contains a sequence that corresponds to the coding strand of an endogenous Alg3 gene. An antisense molecule also can have

flanking sequences (e.g., regulatory sequences). Thus, anti-sense molecules can be ribozymes or antisense oligonucleotides. A ribozyme can have any general structure including, without limitation, hairpin, hammerhead, or axehead structures, provided the molecule cleaves RNA. Further, gene silencing can be used to reduce the activity of Alg3.

#### B. Measuring Gene Inactivation

A fungus having an inactivated Alg3 gene can be identified using any method known in the art. For example, PCR and nucleic acid hybridization techniques, such as Northern and Southern analysis, can be used to confirm that a fungus has an inactivated Alg3 gene. Alternatively, real-time reverse transcription PCR (qRT-PCR) can be used for detection and quantification of targeted messenger RNA, such as mRNA of Alg3 gene in the parent and mutant strains as grown at the same culture conditions. Immunohistochemical and biochemical techniques can also be used to determine if a cell expresses Alg3 by detecting the expression of the Alg3 peptide encoded by *Alga*. For example, an antibody having specificity for Alg3 can be used to determine whether or not a particular fungus contains a functional nucleic acid encoding Alg3 protein. Further, biochemical techniques can be used to determine if a cell contains a particular gene inactivation by detecting a product produced as a result of the expression of the peptide. For example, structural determination of N-glycans excised from glycoproteins can indicate that a fungal cell contains an inactivated Alg3 gene. In addition, measurements of sporulation, germination, secondary metabolite production, and citric acid production can be measured using the methods described herein.

#### C. Measuring Citric Acid Production

Methods of determining whether a genetic inactivation of Alg3 in *Aspergillus* increases citric acid production, for example relative to the same strain with a native Alg3 sequence (such as a parental strain), are routine in the art. Although particular examples are disclosed herein, the methods are not limiting.

For example, production of citric acid by *Aspergillus* (such as an Alg3Δ strain) can be measured using a spectrophotometric assay. In one example citric acid production can be determined with an endpoint spectrophotometric enzyme assay (for example see, Bergmeyer, H. U. 1985. *Metabolites 2*: tri- and dicarboxylic acids, purines, pyrimidines and derivatives, coenzymes, inorganic compounds, p. 5-10. In *Citric acids*. VCH Publishers, Weinheim, Germany). Citric acid can also be measured by liquid chromatography (LC) or high-performance liquid chromatography (HPLC) methods.

#### D. Alg3 Sequences

Alg3 protein and nucleic acid sequences are publicly available and specific examples are provided herein. In addition, Alg3 sequences can be identified using routine molecular biology methods.

Examples of Alg3 nucleic acid sequences shown in SEQ ID NOS: 1 and 3. However, the disclosure also encompasses variants of SEQ ID NOS: 1 and 3 which retain the ability to encode an Alg3 protein. One skilled in the art will understand that variant Alg3 nucleic acid sequences can be inactivated. Variant sequences may contain a single insertion, a single deletion, a single substitution, multiple insertions, multiple deletions, multiple substitutions, or any combination thereof (e.g., single deletion together with multiple insertions). In addition, the degeneracy of the code permits multiple nucleic acid sequences to encode the same protein. For example, FIG. 9 shows an alignment of Alg3 nucleic acid sequences from *A. niger* (nucleotides 1986-2518 of SEQ ID NO: 1) and *A. oryzae* (nucleotides 643-1175 of SEQ ID NO: 3), which permits one to identify nucleotides that

can tolerate substitution (e.g., those that are not conserved between species) and those that may not (e.g., those that are conserved between species). Such nucleic acid molecules can share at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to any known Alg3 nucleic acid sequence, such as SEQ ID NO: 1 or 3 or nucleotides 1186-1306, 1393-1916 and 1989-2582 of SEQ ID NO: 1.

Examples of Alg3 protein sequences shown in SEQ ID NOS: 2, 4, 31, 32, 33, 34, 35, and 36. However, the disclosure also encompasses variants SEQ ID NOS: 2, 4, 31, 32, 33, 34, 35, and 36 which retain Alg3 activity. One skilled in the art will understand that variant Alg3 enzyme sequences can be inactivated. Variant sequences may contain a single insertion, a single deletion, a single substitution, multiple insertions, multiple deletions, multiple substitutions, or any combination thereof (e.g., single deletion together with multiple insertions). Such polypeptides share at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to an Alg3 sequence, such as SEQ ID NO: 2, 4, 31, 32, 33, 34, 35, or 36.

Variant sequences can be identified, for example by aligning known Alg3 sequences. For example, FIGS. 10A and 10B show the alignment of seven different Alg3 sequences from different organisms. In addition, FIG. 11 shows a detailed alignment of Alg3 protein sequences from *A. niger* (amino acids 12-411 of SEQ ID NO: 2) and *S. cerevisiae*, indicating amino acids that are identical, conserved (+) or not conserved (space). Based on these alignments, variants of Alg3 sequences can be identified. For example, amino acid residues that are conserved between organisms are ones that should not be substituted (such as amino acids M50, T70, Y81, Q100 and D150 based on the numbering for *A. niger*), while amino acid residues that are not conserved between organisms are ones likely to tolerate substitution (such as amino acids R8, L160, 5395 and N405 based on the numbering for *A. niger*). Similarly, amino acid positions in FIGS. 10A and 10B indicated with different amino acids at the same position are ones likely to tolerate substitution, while positions with the same amino acid (\*) are not.

In some examples, an Alg3 sequence that is to be genetically inactivated encodes or includes one or more conservative amino acid substitutions. A conservative amino acid substitution is a substitution of one amino acid (such as one found in a native sequence) for another amino acid having similar biochemical properties. Typically, conservative substitutions have little to no impact on the activity of a resulting peptide. In one example, an Alg3 sequence (such as any of SEQ ID NOS: 2, 4, 31, 32, 33, 34, 35, or 36) includes one or more amino acid substitutions (for example at 1, 2, 5 or 10 residues). Examples of amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative substitutions include: Ser for Ala; Lys for Arg; Gln or His for Asn; Glu for Asp; Ser for Cys; Asn for Gln; Asp for Glu; Pro for Gly; Asn or Gln for His; Leu or Val for Ile; Ile or Val for Leu; Arg or Gln for Lys; Leu or Ile for Met; Met, Leu or Tyr for Phe; Thr for Ser; Ser for Thr; Tyr for Trp; Trp or Phe for Tyr; and Ile or Leu for Val. Further information about conservative substitutions can be found in, among other locations in, Ben-Bassat et al., (*J. Bacteriol.* 169:751-7, 1987), O'Regan et al., (*Gene* 77:237-51, 1989), Sahin-Toth et al., (*Protein Sci.* 3:240-7, 1994), Hochuli et al., (*Bio/Technology* 6:1321-5, 1988), WO 00/67796 (Curd et al.) and in standard textbooks of genetics and molecular biology.

The Alg3 gene inactivated in a fungus, in particular examples, includes a sequence that encodes an Alg3 protein having at least at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to an Alg3 sequence, such as SEQ ID NO: 2, 4, 31, 32, 33, 34, 35, or 36, wherein the protein can catalyze the addition of the first dol-p-man derived mannose in an  $\alpha$ -1,3 linkage to Man5GlcNAc2-PP-Dol. In a specific example, the Alg3 gene inactivated in a fungus encodes an Alg3 protein shown in SEQ ID NO: 2, 4, 31, 32, 33, 34, 35, or 36.

The Alg3 gene that is to be inactivated in a fungus, in particular examples, includes a sequence having at least at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to an Alg3 nucleic acid sequence, such as SEQ ID NO: 1 or 3 or nucleotides 1186-1306, 1393-1916 and 1989-2582 of SEQ ID NO: 1, and encode an Alg3 protein that can catalyze the addition of the first dol-p-man derived mannose in an  $\alpha$ -1,3 linkage to Man5GlcNAc2-PP-Dol. In a specific example, the Alg3 gene inactivated in a fungus is shown in SEQ ID NO: 2 or 4.

One skilled in the art will appreciate that additional Alg3 sequences can be identified using any method such as those described herein. For example, Alg3 nucleic acid molecules that encode an Alg3 protein can be identified and obtained using common molecular cloning or chemical nucleic acid synthesis procedures and techniques, including PCR. In addition, standard nucleic acid sequencing techniques and software programs that translate nucleic acid sequences into amino acid sequences based on the genetic code can be used to determine whether or not a particular nucleic acid has any sequence homology with known Alg3 sequences. Sequence alignment software such as MEGALIGN (DNASTAR, Madison, Wis., 1997) can be used to compare various sequences.

In addition, nucleic acid hybridization techniques can be used to identify and obtain a nucleic acid molecule that encodes an Alg3 protein. Briefly, any known Alg3 nucleic acid molecule, or fragment thereof, can be used as a probe to identify similar nucleic acid molecules by hybridization under conditions of moderate to high stringency. Such similar nucleic acid molecules then can be isolated, sequenced, and analyzed to determine whether the encoded protein is an Alg3 protein.

Any method can be used to introduce an exogenous nucleic acid molecule into a fungal cell, for example to genetically inactivate Alg3. For example, chemical mediated-protoplast transformation, electroporation, *Agrobacterium*-mediated transformation, fusion of protoplasts, and biolistic delivery are common methods for introducing nucleic acid into fungal cells. (See, e.g., Ito et al., *J. Bacteriol.* 153:163-8, 1983; Durrens et al., *Curr. Genet.* 18:7-12, 1990; Sambrook et al., *Molecular cloning: A laboratory manual*, Cold Spring Harbour Laboratory Press, New York, USA, third edition, 2001; and Becker and Guarente, *Methods in Enzymology* 194:182-7, 1991. An exogenous nucleic acid molecule contained within a particular cell of the disclosure can be maintained within that cell in any form. For example, exogenous nucleic acid molecules can be integrated into the genome of the cell or maintained in an episomal state. That is, a cell can be a stable or transient transformant.

#### Fungi with Increased LaeA Expression or Increased LaeA Expression and Alg3 Deletion

The present disclosure provides isolated fungi having increased LaeA expression, wherein such increased expres-

sion or activity (for example in combination with an Alg3 functional inactivation, Alg3 $\Delta$ ) results in increased citric acid production by the fungi. Such fungi are referred to herein as increased LaeA fungal strains. It is disclosed herein that increased expression of LaeA (for example in combination with genetic inactivation of Alg3, Alg3 $\Delta$ ) results in *Aspergillus* fungi that can increase citric acid production as compared to *Aspergillus* having native levels of expression.

Contemplated herein are isolated fungi having increased LaeA activity/expression, for example in combination with a genetic inactivation of Alg3. Any fungus can be used, such as any genus or variety of *Aspergillus*. In particular examples, the *Aspergillus* fungus is *A. niger*, such as *Aspergillus niger* strain 11414 (American Type Culture Collection (ATCC) No. 11414; NRRL 2270); 1015 (ATCC No. 1015; NRRL 328, CBS 113.46); NRRL 3 (ATCC No. 9029, CBS 120.49, N400); NRRL 3122 (ATCC No. 22343); or 11414KusA-. In other specific examples, the *Aspergillus* is *A. aculeatus*, *A. awamori*, *A. carbonarius*, *A. wentii*, *A. foetidus*, *A. fumigatus*, *A. oryzae*, or *A. terreus*.

The present disclosure also provides isolated *A. terreus* fungi having increased LaeA expression, wherein such increased expression or activity results in increased itaconic acid production by the fungi. It is disclosed herein that increased expression of LaeA results in *A. terreus* fungi that can increase itaconic acid production as compared to *A. terreus* having native levels of LaeA expression. Contemplated herein are isolated *A. terreus* fungi transformed with a nucleic acid construct comprising an *Aspergillus* species LaeA gene. The LaeA gene can be from any species of *Aspergillus*. In some examples, the LaeA gene is an *A. niger*, *A. nidulans*, *A. aculeatus*, *A. awamori*, *A. carbonarius*, *A. wentii*, *A. foetidus*, *A. fumigatus*, *A. oryzae*, or *A. terreus* LaeA gene.

Any method for genetic enhancement or up-regulation can be used, as long as the expression of the gene and/or gene product is significantly increased, or the function of the expressed protein is significantly increased. In particular examples, LaeA gene expression is up-regulated by transformation of the fungi with one or more copies of a LaeA coding or genomic sequence (which can be a native or non-native LaeA sequence). In some embodiments, up-regulation refers to an increase in gene or protein expression of at least 20%, at least 40%, at least 50%, at least 100%, at least 150%, at least 200%, at least 300%, or at least 500%, for example relative to the parental fungal strain without the additional copies of an LaeA gene. The term "increased" or "up-regulated" as used herein with respect to a cell and a particular gene or protein activity refers to a higher level of activity than that measured in a comparable cell of the same species. For example, a particular fungi having increased or up-regulated LaeA activity has increased LaeA activity if a comparable fungi having native LaeA activity has less detectable LaeA activity (for example as measured by gene or protein expression).

LaeA sequences are disclosed herein and others are publicly available, for example from GENBANK™ or EMBL. In some examples, the LaeA gene upregulated encodes a protein having at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% sequence identity to SEQ ID NO: 41 or SEQ ID NO: 59. In some examples, the LaeA gene upregulated comprises at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% sequence identity to SEQ ID NO: 40 (e.g., nt 1-236 and 367-1252 of SEQ ID NO: 41) or SEQ ID NO: 58 (e.g., nt 1-230 and 373-1267 of SEQ ID NO: 58).

Increasing LaeA activity (for example in combination with genetic inactivation of Alg3) results in many phenotypes in the fungi. For example, such recombinant fungi exhibit increased citric acid production in CAP medium. Such increases can be relative to a fungi having a native or wild-type level of LaeA (or LaeA and Alg3) gene or protein expression, such as a parental strain (e.g., *A. niger* strain 11414KusA), grown under the same conditions as the fungi with increased LaeA activity (or increased LaeA activity and decreased Alg3 activity). In some examples, increased citric acid production in CAP medium is an increase of at least 20%, at least 30%, at least 50%, at least 60%, at least 65%, or at least 70%, by such a recombinant fungus as compared to *A. niger* strain 11414KusA. In some examples, recombinant fungi with increased LaeA activity (for example in combination with genetic inactivation of Alg3) have increased sporulation relative to *A. niger* Alg3Δ on MM, accumulate red color pigments to *A. niger* strain 11414KusA on complete medium, or both. In other examples, increasing LaeA activity in *A. terreus* fungi results in an increase in itaconic acid production in itaconic acid production media. Such increases can be relative to *A. terreus* fungi having a native or wild-type level of LaeA gene or protein expression, such as a parental strain *A. terreus* strain grown under the same conditions as the fungi with increased LaeA activity. In some examples, increased itaconic acid production is an increase of at least 20%, at least 30%, at least 50%, at least 60%, at least 65%, or at least 70%, as compared to a parental strain.

One skilled in the art will appreciate that additional genes can also be inactivated or upregulated, wherein the additional genes may or may not provide additional enhancement of citric acid production or itaconic acid production to the fungus. In one example KusA (e.g., GENBANK™ Accession No. EF061656) is also genetically inactivated.

Also provided by the present disclosure are compositions that include isolated LaeA up-regulated fungi, such as a growth medium. Also provided by the present disclosure are kits that include isolated LaeA up-regulated fungi, such as a kit that includes a medium for culturing, storing, or growing the fungus. Exemplary mediums include solid medium (such as those containing agar, for example CM, PDA or MM) and liquid media (such as a fermentation broth, such as CM, MM, or CAP medium).

#### A. Methods of Up-Regulating Gene and/or Protein Expression

As used herein, an “activated” or “up-regulated” gene means that expression of the gene or gene product (e.g., protein) has been up-regulated, for example by introduction of additional copies of the appropriate gene or coding sequence into the fungus (or other common molecular biology methods), such that the introduce nucleic acid sequence is expressed, resulting in increased expression or biological activity of the encoded gene product.

Increasing expression of one or more genes (which in some examples is also referred to as up-regulation) can be performed using any conventional method known in the art. In one example, a strain of *Aspergillus* is transformed with a vector which has the effect of up-regulating or otherwise activating a LaeA gene (such as a native or non-native LaeA gene). This can be done by introducing one or more LaeA coding sequences (such as a gene sequence), whose expression is controlled by elements such as promoters and the like which control gene expression, by introducing a nucleic acid sequence which itself (or its encoded protein) can increase LaeA protein activity in the fungus, or by introducing another molecule (such as a protein or antibody) increases

LaeA protein activity in the fungus. For example, a LaeA gene can be up-regulated by introduction of a vector that includes one or more LaeA sequences (such as 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 LaeA sequences or copies of such sequences) into the desired fungus. In some examples, such LaeA sequences are from different fungal species, can be multiple copies from a single species, or combinations thereof, such as LaeA sequences from at least 2, 3, 4, 5, 6, 7, 8, 9, or 10 different fungal species. In some examples, the LaeA sequence(s) introduced into the fungus is optimized for codon usage. Thus, the disclosure in some examples provides transformed fungi that include at least one exogenous nucleic acid molecule which includes a LaeA gene or coding sequence (such as a nucleic acid sequence encoding SEQ ID NOs: 41 or 59), for example in combination with Alg3Δ. In one example, such transformed cells produce more citric acid, for example relative to a comparable fungus with a native LaeA sequence (or a native LaeA sequence combined with a native Alg3 sequence).

In another technique, the cre-lox system is used for site specific recombination of DNA (for example see Steiger et al., *Appl. Environ. Microbiol.* 77(1):114, 2011). The system includes 34 base pair lox sequences that are recognized by the bacterial cre recombinase gene. If the lox sites are present in the DNA in an appropriate orientation, DNA flanked by the lox sites will be excised by the cre recombinase, resulting in the deletion of all sequences except for one remaining copy of the lox sequence. Using standard recombination techniques, the targeted gene of interest (e.g., LaeA) can be deleted in the *Aspergillus* genome and replaced with one or more copies of a non-native LaeA sequence (for example in *A. niger*, replacing one or both *A. niger* LaeA sequences with one or more, or combination of, LaeA sequences from *A. nidulans*, *A. flavus*, *fusarium oxysporum*, *penicillium chrysogenum*, which have high secondary metabolite production) flanked by the lox sites. Transient expression (by electroporation of a suicide plasmid containing the cre gene under control of a promoter that functions in *Aspergillus*) of the cre recombinase should result in efficient elimination of the lox flanked marker. This process will produce a fungus containing the desired insertion mutation and one copy of the lox sequence.

In one example, one or more LaeA genes are introduced into fugal cells by chemical mediated proteoplast transformation in combination of yeast-gap repairing method for transgene expression construction.

In one example, a transgene is generated and expressed in the desired fungal cell, such as an Alg3Δ cell, to increase LaeA expression. For example, such a transgene can include a LaeA genomic or cDNA sequence (such as one having at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to any known LaeA sequence, such as SEQ ID NO: 40 or 58), for example operably linked to a promoter, such as a glyceraldehyde 3-phosphate dehydrogenase (gpdA) promoter or other promoter, such as one that has high activity in CAP culture medium, for example a polyubiquitin promoter, Arsa-7, and A-37 from *A. niger*. In one example, the promoter has at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 37. In one example, the promoter comprises or consists of the sequence shown in SEQ ID NO: 37. In some examples, the transgene further includes pyrG upstream and downstream sequences (for example that are at the 5'- and 3'-end, respectively, of the transgene). The pyrG gene in *A. niger* is mutated and has lost its original functions. Thus, other non-essential gene loci can be used as long as it

is not influenced by the native neighbor genes. In one example, the *pyrG* upstream and downstream sequences have at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 45 and 48, respectively. In one example, the *pyrG* upstream and downstream sequences comprise or consist of the sequence shown in SEQ ID NO: 45 or 48, respectively. In some examples, the transgene further includes a *trpC* transcriptional terminator sequence of *A. nidulans*, for example downstream of the *LaeA* sequence. As an alternative to *trpC*, other transcriptional terminators can be used, such as promoters which include a transcriptional terminators (e.g., *ArsA7*, *Arsa-37*, polyubiquitin (*ubi4*)). In one example, the *trpC* transcriptional terminator has at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 46. In one example, the *trpC* transcriptional terminator comprises or consists of the sequence shown in SEQ ID NO: 46. In some examples, the transgene further includes a *ptrA* sequence, for example downstream of the *trpC* transcriptional terminator sequence. As an alternative to *ptrA*, the bleomycin gene or *bar* gene can be used. In one example, the *ptrA* sequence has at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 47. In one example, the *ptrA* sequence comprises or consists of the sequence shown in SEQ ID NO: 47. In one example, the transgene comprises a sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 97%, at least 98%, or at least 99% sequence identity to SEQ ID NO: 44 or 57. In one example, the transgene comprises or consists of the sequence shown in SEQ ID NO: 44 or 57.

Thus, for example, a fungal cell can be engineered to have increased copies of *LaeA* using common recombinant technology methods.

#### B. Measuring Gene Activation or Up-Regulation

A fungus having an activated or up-regulated *LaeA* gene can be identified using any method known in the art. For example, PCR and nucleic acid hybridization techniques, such as Northern, RT-PCR, and Southern analysis, can be used to confirm that a fungus has an up-regulated *LaeA* gene, such as an increase in the *LaeA* copy number. Immunohistochemical and biochemical techniques can also be used to determine if a cell expresses *LaeA* by detecting the expression of the *LaeA* peptide encoded by *LaeA*. For example, an antibody having specificity for *LaeA* can be used to determine whether or not a particular fungus has increased *LaeA* protein expression. Further, biochemical techniques can be used to determine if a cell has increased *LaeA* expression by detecting a product produced as a result of the expression of the peptide. For example, measurement of secondary metabolites can indicate that a fungal cell contains an up-regulated *LaeA* gene. In addition, measurements of citric acid production can be measured using the methods described herein.

#### C. Measuring Citric Acid Production

Methods of determining whether a genetic up-regulation of *LaeA* (alone or in combination with inactivation of *Alg3*) in *Aspergillus* increases citric acid production, for example relative to the same strain with a native *LaeA* sequence, *Alg3* sequence, or both (such as a parental strain), are routine in the art. Although particular examples are disclosed herein (see above and in the examples below), the methods are not limiting.

#### D. Measuring Itaconic Acid Production

Methods of determining whether a genetic up-regulation (e.g. overexpression) of *LaeA* in *Aspergillus terreus*

increases itaconic acid production, for example relative to the same strain that does not overexpress *LaeA* (such as a parental strain), are routine in the art. Although particular examples are disclosed herein (see Example 14), the methods are not limiting. For example, production of itaconic acid by *A. terreus* can be performed as previously described by Dickman (*Analytical Chem* 24: 1064-1066, 1952), or using the modified method described in Example 14.

#### E. *LaeA* Sequences

*LaeA* protein and nucleic acid sequences are publicly available and specific examples are provided herein. In addition, *LaeA* sequences can be identified using routine molecular biology methods.

Examples of *LaeA* nucleic acid sequences shown in SEQ ID NOs: 40 and 58. However, the disclosure also encompasses variants of SEQ ID NOs: 40 and 58 (such as the coding regions nt 1-236 and 367-1252 of SEQ ID NO: 41 and nt 1-230 and 373-1267 of SEQ ID NO: 58) which retain the ability to encode a *LaeA* protein. One skilled in the art will understand that variant *LaeA* nucleic acid sequences can be used to increase expression of *LaeA*. Variant sequences may contain a single insertion, a single deletion, a single substitution, multiple insertions, multiple deletions, multiple substitutions, or any combination thereof (e.g., single deletion together with multiple insertions). In addition, the degeneracy of the code permits multiple nucleic acid sequences to encode the same protein. Thus, in one example, a *LaeA* sequence having at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to any known *LaeA* sequence, such as SEQ ID NO: 40 or SEQ ID NO: 58 (such as the coding regions nt 1-236 and 367-1252 of SEQ ID NO: 41 and nt 1-230 and 373-1267 of SEQ ID NO: 58) can be expressed in a fungal cell to increase *LaeA* expression in the fungal cell.

For example, FIG. 12 shows an alignment of *LaeA* protein sequences from *A. nidulans* (aa 14-372 of SEQ ID NO: 41) and *A. niger* (aa 14-370 of SEQ ID NO: 59), which permits one to identify amino acids that can tolerate substitution (e.g., those that are not conserved between species) and those that may not (e.g., those that are conserved between species). Based on these alignments, variants of *LaeA* sequences can be identified. For example, amino acid residues that are conserved between organisms are ones that should not be substituted (such as amino acids S16, M42, and P52 based on the numbering for *A. nidulans*), while amino acid residues that are not conserved between organisms are ones likely to tolerate substitution (such as amino acids T15, S44, and N325 based on the numbering for *A. nidulans*). Similarly, amino acid positions in FIG. 12 indicated with a space are ones likely to tolerate substitution, while positions with the same amino acid are not.

Such protein molecules can share at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to any known *LaeA* nucleic acid sequence, such as SEQ ID NOs: 41 and 59, and such variants can be used to increase *LaeA* activity in a fungal cell. One skilled in the art will understand that variant *LaeA* enzyme sequences can be used to increase *LaeA* activity in a fungal cell. Variant sequences may contain a single insertion, a single deletion, a single substitution, multiple insertions, multiple deletions, multiple substitutions, or any combination thereof (e.g., single deletion together with multiple insertions).

In some examples, a *LaeA* sequence whose expression is to be up-regulated encodes or includes one or more conservative amino acid substitutions. A conservative amino acid

substitution is a substitution of one amino acid (such as one found in a native sequence) for another amino acid having similar biochemical properties. Typically, conservative substitutions have little to no impact on the activity of a resulting peptide. In one example, a LaeA sequence (such as any of SEQ ID NOs: 41 and 59) includes one or more amino acid substitutions (for example at 1, 2, 5 or 10 residues). Examples of amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative substitutions include those discussed above for Alg3.

The LaeA gene up-regulated in a fungus, in particular examples, includes a sequence that encodes a LaeA protein having at least at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to an LaeA sequence, such as SEQ ID NO: 41 or SEQ ID NO: 59, wherein the protein can regulate secondary metabolite production in *Aspergillus*. In a specific example, the LaeA gene up-regulated in a fungus encodes a LaeA protein shown in SEQ ID NO: 41 or 59.

The LaeA gene up-regulated in a fungus, in particular examples, includes a sequence having at least at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% sequence identity to a LaeA nucleic acid sequence, such as SEQ ID NO: 40 or SEQ ID NO: 58 (or to the coding regions nt 1-236 and 367-1252 of SEQ ID NO: 41 or nt 1-230 and 373-1267 of SEQ ID NO: 58), and encodes a LaeA protein which can regulate secondary metabolite production in *Aspergillus*. In a specific example, the LaeA gene upregulated in a fungus is shown in SEQ ID NO: 40 or SEQ ID NO: 58 (or includes the coding regions nt 1-236 and 367-1252 of SEQ ID NO: 41 or nt 1-230 and 373-1267 of SEQ ID NO: 58).

One skilled in the art will appreciate that additional LaeA sequences can be identified and obtained using any method such as those described herein. For example, LaeA nucleic acid molecules that encode a LaeA protein can be identified and obtained using common molecular cloning or chemical nucleic acid synthesis procedures and techniques, including PCR. In addition, standard nucleic acid sequencing techniques and software programs that translate nucleic acid sequences into amino acid sequences based on the genetic code can be used to determine whether or not a particular nucleic acid has any sequence homology with known LaeA sequences. Sequence alignment software such as MEGA-LIGN (DNASTAR, Madison, Wis., 1997) can be used to compare various sequences.

In addition, nucleic acid hybridization techniques can be used to identify and obtain a nucleic acid molecule that encodes a LaeA protein. Briefly, any known LaeA nucleic acid molecule, or fragment thereof, can be used as a probe to identify similar nucleic acid molecules by hybridization under conditions of moderate to high stringency. Such similar nucleic acid molecules then can be isolated, sequenced, and analyzed to determine whether the encoded protein is a LaeA protein. The gene specific oligonucleotide pair can also be designed, synthesized and used for real-time RT-PCR to quantify the LaeA gene transcription level.

Any method can be used to introduce an exogenous nucleic acid molecule into a fungal cell, for example to genetically enhance LaeA expression. For example, chemical mediated-protoplast transformation, electroporation, *Agrobacterium*-mediated transformation, fusion of protoplasts, and biolistic delivery are common methods for introducing nucleic acid into fungal cells. (See, e.g., Ito et al., *J. Bacteriol.* 153:163-8, 1983; Durrens et al., *Curr. Genet.* 18:7-12, 1990; Sambrook et al., *Molecular cloning: A labo-*

ratory manual, Cold Spring Harbour Laboratory Press, New York, USA, second edition, 1989; and Becker and Guarente, *Methods in Enzymology* 194:182-7, 1991). An exogenous nucleic acid molecule contained within a particular cell of the disclosure can be maintained within that cell in any form. For example, exogenous nucleic acid molecules can be integrated into the genome of the cell or maintained in an episomal state. That is, a cell can be a stable or transient transformant.

#### F. LaeA Deletion and Complementation

Disclosed herein is a LaeA deletion cassette for generating a fungus, such an *Aspergillus* species fungus, having a deletion in the LaeA gene. A schematic of the LaeA deletion cassette is shown in FIG. 17A. To generate the deletion cassette, DNA fragments comprising the 5' and 3' ends of the *A. niger* LaeA gene (SEQ ID NOs: 61 and 62, respectively) were isolated from *A. niger* genomic DNA and the hph expression cassette (SEQ ID NO: 63) (hph) was isolated from pCB1003 plasmid vector DNA by PCR. The PCR DNA fragments were assembled in the pBSK backbone vector to produce the LaeA deletion cassette (SEQ ID NO: 65).

Provided herein is a nucleic acid molecule comprising a nucleotide sequence at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to SEQ ID NO: 65. In some embodiments, the nucleic acid molecule comprises or consists of the nucleotide sequence of SEQ ID NO: 65. Further provided is an isolated fungus transformed with a nucleic acid molecule comprising a nucleotide sequence at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to SEQ ID NO: 65. In some embodiments, the isolated fungus is transformed with a nucleic acid molecule comprising or consisting of the nucleotide sequence of SEQ ID NO: 65. In some embodiments, the fungus is a species of *Aspergillus*, such as *A. niger*.

Further provided herein are isolated fungi (such as filamentous fungi) having a gene inactivation (also referred to as a gene deletion) of a LaeA gene. Any strain of fungi can be used, such as a filamentous fungi, for example *A. niger* or particular strains thereof (for example *A. niger* strain 11414 or 11414KusA). In particular examples, the LaeA gene is genetically inactivated by complete or partial deletion mutation or by insertional mutation. In some examples genetic inactivation need not be 100% genetic inactivation. In some embodiments, genetic inactivation refers to at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, or at least 95% gene or protein inactivation. The term "reduced" or "decreased" as used herein with respect to a cell and a particular gene or protein activity refers to a lower level of activity than that measured in a comparable cell of the same species. For example, a particular fungi lacking LaeA activity has reduced LaeA activity if a comparable fungi not having an LaeA genetic inactivation has detectable LaeA activity. In some embodiments, the isolated fungi having a gene inactivation are generated using the LaeA deletion construct set forth herein as SEQ ID NO: 65.

Also provided herein is a LaeA complementation construct for complementing expression of LaeA in fungi having a deleted LaeA gene. A schematic of the LaeA complementation construct is shown in FIG. 17B. To generate the complementation construct, the LaeA gene containing both its promoter and transcriptional terminator was isolated by PCR from *A. niger* genomic DNA. The PCR fragment was cloned into the plasmid vector pRSB426-LaeA. The new plasmid DNA vector contained the upstream region of the

pyrG gene of *A. niger*, the entire region of *A. niger* LaeA gene, the transcriptional terminator of the trpC gene from *A. nidulans*, the pyrithiamine resistance (ptrA) gene from *A. oryzae*, and the downstream region of the pyrG gene of *A. niger* (SEQ ID NO: 66).

Provided herein is a nucleic acid molecule comprising a nucleotide sequence at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to SEQ ID NO: 66. In some embodiments, the nucleic acid molecule comprises or consists of the nucleotide sequence of SEQ ID NO: 66. Further provided is an isolated fungus transformed with a nucleic acid molecule comprising a nucleotide sequence at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to SEQ ID NO: 66. In some embodiments, the isolated fungus is transformed with a nucleic acid molecule comprising or consisting of the nucleotide sequence of SEQ ID NO: 66. In some embodiments, the fungus is a species of *Aspergillus*, such as *A. niger*.

Further provided herein are isolated fungi (such as filamentous fungi) having a gene inactivation (also referred to as a gene deletion) of a LaeA gene and further transformed by a nucleic acid construct comprising a nucleotide sequence at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% identical to SEQ ID NO: 66. In some embodiments, the isolated fungus is transformed with a nucleic acid molecule comprising or consisting of the nucleotide sequence of SEQ ID NO: 66. In some embodiments, the fungus is a species of *Aspergillus*, such as *A. niger*. Any strain of fungi can be used, such as a filamentous fungi, for example *A. niger* or particular strains thereof (for example *A. niger* strain 11414 or 11414KusA).

#### Production of Citric Acid Using Alg3Δ Mutants, Fungi with Increased LaeA Expression, or Both

The fungi provided herein, namely Alg3Δ fungi, up-regulated LaeA fungi, and fungi with both Alg3Δ and up-regulated LaeA, can be used to produce citric acid, as well as derivatives thereof such as hydroxycitric acid (for example for medical applications). Such fungi can be from any species, such as *Aspergillus* or *Rhizopus* cells. For example, the disclosure provides methods of making citric acid, which can include culturing Alg3Δ fungi, up-regulated LaeA fungi, or fungi with both Alg3Δ and up-regulated LaeA, under conditions that permit the fungus to make citric acid, for example in CAP medium.

Citric acid (2-hydroxy-propane-1,2,3-tricarboxylic acid) combines a pleasant taste with low toxicity and palatability and is a ubiquitous food additive. It is also able to complex heavy metal ions, like iron and copper, and is therefore applied in the stabilization of oils and fats or ascorbic acid during metal ion-catalyzed oxidation reactions. Consequently, it is today one of the bulk products produced by fermentation, most of which occurs with the fungus *Aspergillus niger*, although a small portion is also produced by fermentation with yeast, such as *Candida oleophila* and *Candida lipolytica*.

Citric acid production generally requires a unique combination of several unusual nutrient conditions (e.g., excessive concentrations of carbon source, H<sup>+</sup>, and dissolved oxygen, or suboptimal concentrations of certain trace metals and phosphate), which synergistically influence the yield of citric acid. Table 1 below shows the environmental parameters that influence citric acid accumulation.

TABLE 1

Parameters that influence citric acid accumulation by <i>A. niger</i>	
Parameter	Requirement for citric acid accumulation
Carbon source concentration	Higher than 50 g/l
Carbon source type	Enable rapid catabolism
Nitrogen source	Consumption leads to some decrease in pH
Phosphate concentration	Suboptimal
Aeration	In excess
Trace metal ions	Limiting, especially Mn <sup>2+</sup>
pH	Below pH 3

Methods of making citric acid, which can include culturing Alg3Δ fungi, up-regulated LaeA fungi, or fungi with both Alg3Δ and up-regulated LaeA, under conditions that permit the fungus to make citric acid, are provided. In general, the culture media and/or culture conditions can be such that the fungi grow to an adequate density and produce citric acid efficiently. In one example the Alg3Δ fungi, up-regulated LaeA fungi, or fungi with both Alg3Δ and up-regulated LaeA, are cultured or grown in a liquid medium that includes sucrose and/or glucose as the carbon source, for example at a concentration of at least 50 g/liter, such as at least 100 g/l, or at least 140 g/l. Thus, a fungus within the scope of the disclosure in some examples can utilize a variety of carbon sources. In one example the Alg3Δ fungi, up-regulated LaeA fungi, or fungi with both Alg3Δ and up-regulated LaeA, are cultured or grown in a liquid medium that includes a very small amount of manganese, such as less than 100 parts per billion (ppb), less than 50 ppb, less than 20 ppb, less than 15 ppb, for example 5 ppb to 15 ppb or 10 ppb to 15 ppb, such as 5, 10, 13, 15 or 20 ppb. In one example the Alg3Δ fungi, up-regulated LaeA fungi, or fungi with both Alg3Δ and up-regulated LaeA, are cultured or grown in a liquid medium having an initial pH of less than 3, such as less than 2.5, for example about pH 1.8 to 3, 1.8 to 2.5, 1.8 to 2.2, 1.9 to 2.1, for example pH 1.8, 1.9, 2, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8 or 2.9. In some examples the Alg3Δ fungi, up-regulated LaeA fungi, or fungi with both Alg3Δ and up-regulated LaeA, are cultured or grown in a liquid medium at about 25 to 35° C. (such as 28 to 32° C., or 30° C.) with rotation of 180 to 300 rpm.

In a specific example, the Alg3Δ fungi, up-regulated LaeA fungi, or fungi with both Alg3Δ and up-regulated LaeA, are grown in citric acid production (CAP) medium. In a specific example, the CAP medium includes 140 g of glucose/liter, 3.1 g of NH<sub>4</sub>NO<sub>3</sub>/liter, 0.15 g of KH<sub>2</sub>PO<sub>4</sub>/liter, 0.15 g of NaCl/liter, 2.2 g of MgSO<sub>4</sub> 7H<sub>2</sub>O/liter, 6.6 mg of ZnSO<sub>4</sub> 7H<sub>2</sub>O/liter, and 0.1 mg of FeCl<sub>3</sub>/liter adjusted to about pH 2 with 4 M H<sub>2</sub>SO<sub>4</sub>. Cations can be removed from the glucose solution by ion exchange on Dowex 50W-X8, 100/200-mesh, H cation exchange resin (Fisher Scientific, Pittsburgh, Pa.) prior to adding the other nutrient components. The manganese concentration in the medium can be adjusted by the addition of appropriate volumes of a stock solution of MnCl<sub>2</sub> 4H<sub>2</sub>O (10 mM). In one example, the manganese concentration is less than 50 ppb, such as less than 20 ppb, for example 5 to 15 ppb, such as 10 ppb.

Methods of culturing *Aspergillus* to enable citric acid production are well known in the art. In one example, the fungi are grown in culture containers (such as baffled flasks, and in some examples are silanized (5% solution of dichlorodimethylsilane in heptane (Sigma, St. Louis, Mo.)). The Alg3Δ fungi, up-regulated LaeA fungi, or fungi with both Alg3Δ and up-regulated LaeA, provided herein can be grown in CAP media containing low amounts of Mn<sup>2+</sup>(e.g.,

10 ppb) at 30° C. with rotation (e.g., 200 to 250 rpm) for at least 3 days (e.g., 3 to 7 days). Each culture container is inoculated with spores (such as at least 10<sup>6</sup> spores/ml) and incubated for at least 12 or at least 15 hours at 30° C. and 200 to 250 rpm to obtain properly pelleted morphology.

In one example, the Alg3Δ fungi, up-regulated LaeA fungi, or fungi with both Alg3Δ and up-regulated LaeA, produce more citric acid than a corresponding fungus with wild-type Alga. In specific examples, the Alg3Δ fungi, up-regulated LaeA fungi, or fungi with both Alg3Δ and up-regulated LaeA, produce at least 25 g/l of citric acid (for example at least 30 g/l, at least 32 g/l, at least 35 g/l, at least 40 g/l, at least 42 g/l, at least 45 g/l, at least 50 g/l, at least 52 g/l or at least 55 g/l), for example after at least 4 days (such as at least 5 days, at least 6 days, at least 7 days, at least 8 days, or at least 10 days, such as after 4 to 6 days, 8 to 10 days, or 4 to 5 days) when grown in CAP medium at 30° C. with 200 rpm shaking.

In some examples, the method further includes isolating the citric acid made by the Alg3Δ fungi, up-regulated LaeA fungi, or fungi with both Alg3Δ and up-regulated LaeA. Once produced, any method can be used to isolate the citric acid. For example, common separation techniques can be used to remove the fungal biomass from the culture medium, and common isolation procedures (e.g., filtration, distillation, precipitation, electro dialysis, and ion-exchange procedures) can be used to obtain the citric acid from the broth (such as a fungi-free broth). In addition, the citric acid can be isolated from the culture medium after the citric acid production phase has been terminated.

### Example 1

#### Materials and Methods

This example describes methods used in the experiments described in Examples 2-5 below.

#### Strains and Media.

The *Escherichia coli* strains Top10 and *Saccharomyces cerevisiae* strain YVH10 were used as hosts for routine cloning and gap repair experiments. *A. niger* strain ATCC 11414 (American Type Culture Collection, Rockville, Md.), was grown on potato dextrose agar plates (PDA) and complete medium (CM) agar plates at 30° C. for culture maintenance and spore preparation, respectively. The mutant strain *Aspergillus niger* 11414KusA was generated by the deletion of kusA in *A. niger* strain 11414 by the replacement with *A. fumigatus* pyrG gene, which encodes the ortholog of the ku70 protein that involves in the non-homologous end joining pathway of DNA repair for the integration of a DNA fragment into the genome in other eukaryotes, and was confirmed by Southern blotting analysis. The 11414kusA strain with high rate of homologous replacement was mainly used as a parent strain. The cultures on PDA or complete medium (CM) agar plates were incubated for four days at 30° C. and the spores were harvested by washing with sterile 0.8% Tween 80 (polyoxyethylenesorbitan monooleate). The CM medium contains 20 g of D-glucose/liter, 5 g yeast extract/liter, 2 g trypticase peptone/liter, 1 g casamino acids/liter, 6 g NaNO<sub>3</sub>/liter, 0.52 g KCl/liter, 0.52 g MgSO<sub>4</sub>·7H<sub>2</sub>O/liter, 1.52 g KH<sub>2</sub>PO<sub>4</sub>/liter, 36.7 mg ZnSO<sub>4</sub>·7H<sub>2</sub>O/liter, 18.3 mg H<sub>3</sub>BO<sub>3</sub>/liter, 8.3 mg MnCl<sub>2</sub>·4H<sub>2</sub>O/liter, 8.3 mg FeSO<sub>4</sub>·7H<sub>2</sub>O, 2.8 mg CoCl<sub>2</sub>·6H<sub>2</sub>O/liter, 2.7 mg CuSO<sub>4</sub>·5H<sub>2</sub>O/liter, 2.5 mg Na<sub>2</sub>MoO<sub>4</sub>·2H<sub>2</sub>O/liter, 83.3 mg Na<sub>2</sub>EDTA/liter, 1 mg biotin/liter, 1 mg pyridoxin/liter, 1 mg thiamine/liter, 1 mg riboflavin/liter, 1 mg p-aminobenzoic acid/liter and 1 mg nicotinic acid/liter. The PDA medium

contains 4 g/liter potato starch and 20 g/liter dextrose. Conidia were enumerated with a hemacytometer. Aliquots of the resulting spore suspension (1×10<sup>9</sup> spores/ml) were used to inoculate baffled-flask liquid cultures. The citric acid production (CAP) medium contained 140 g/l of glucose, 3.1 g/l NH<sub>4</sub>NO<sub>3</sub>, 0.15 g/l KH<sub>2</sub>PO<sub>4</sub>, 0.15 g/l NaCl, 2.2 g/l MgSO<sub>4</sub>·7H<sub>2</sub>O, 6.6 mg/l ZnSO<sub>4</sub>·7H<sub>2</sub>O, and 0.1 mg/l FeCl<sub>3</sub> adjusted to pH 2.1 with 4 M H<sub>2</sub>SO<sub>4</sub>. Cations were removed from the glucose solution by ion-exchange on Dowex 50W-X8, 100-200 mesh, H cation exchange resin (Fisher Scientific, Pittsburgh, Pa.) prior to adding the other nutrient components.

#### Culture Methods.

Glass baffled-flasks of 250 ml or 1000 ml were silanized by rinsing in a 5% solution of dichlorodimethylsilane in heptane (Sigma, St. Louis, Mo.) to minimize leaching of metals. For citric acid production tests, 1×10<sup>6</sup> spores/ml of parent or mutant strains were grown in 80 ml CAP media containing 10 ppb Mn<sup>2+</sup> in 250 ml baffled flasks or 220 ml CAP media in 1000 ml baffled flasks at 30° C. and 200 rpm. Samples for citric acid analysis were taken at intervals. The biomass of transgenic clones and parent stain were prepared from 2 ml CM station cultures with proper antibiotics and grown in 16×125 mm glass culture-tubes at 30° C. without shaking. The biomass formed on the surface of the culture medium was collected, frozen immediately in liquid nitrogen and dried in the lyophilizer.

#### Dried Biomass Measurement.

After proper cultivation, the cell mass from citric acid production culture was collected by centrifugation at room temperature and 4500×g for 5 min in Sorvall floor centrifuge with swinging-bucket rotor. The cell mass was then transferred onto the Whatman Grade No 1 filter paper or left in centrifuge tubes for freeze-drying. The biomass was then dried in high temperature oven at 80° C. or freeze-dried in the lyophilizer. Prior to being used, the centrifuge tube or Whatman filter paper was weighted and re-weighted after the biomass was completely dried.

Total Genomic DNA Isolation for PCR and Southern Blotting Analysis.

Total genomic DNA was isolated from *A. niger* according to the SDS extraction method described previously by Dellaporta et al. (*Plant Molecular Biology Reporter* 1(4): 19-21, 1983) with some modifications. Briefly, fungal biomass from 2 ml station cultures was looped and transferred into a 1.5 ml microcentrifuge tube. A needle size hole on the cap was punched with 18 gauge needle. The tube was immediately frozen in liquid N<sub>2</sub> for 5 minutes and biomass in the tube was dried in a VirTis benchtop manifold freeze dryer (SP Scientific, Gardiner, N.Y.) overnight. The dried biomass and two 3.5 mm diameter glass beads were transferred into the 2 ml polypropylene microvial, where biomass was pulverized into fine power with Mini-Beadbeater-8 (Bio Spec Products Inc., Bartlesville, Okla.) for one minute. Then, 500 μl of 60° C. extraction buffer and 80 μl of 15% SDS were added into the microcentrifuge tube and incubated at 65° C. for at least 30 minutes with occasionally swirling to mix. Two-hundred microliters of 5M potassium acetate was added, mixed and incubated on ice for 30 minutes. The supernatant was collected by centrifugation at 12,000 g for 10 minutes at 4° C. and transferred into the new microcentrifuge tube. The total nucleic acids were precipitated with 780 μl of 2-propanol for 30 minutes at -20° C. and centrifuged at 12,000 g for 10 minutes. The nucleic acids were re-suspended in 200 μl 50TE buffer containing 2 RNase (10 μg/μl stock solution) and incubated in Eppendorf thermomixer at 50° C. and 500 rpm for 30 minutes. The proteins



and cell debris was removed by being added and well mixed 20  $\mu$ l 3M sodium acetate and equal volume of phenol: chloroform and centrifuged at 15,000 g for 5 minutes. The supernatant was transferred to new DNase-free microcentrifuge tube containing 220  $\mu$ l of 2-propanol, mixed well and incubated at room temperature for 5 minutes. The genomic DNA was pelleted by centrifugation at 15,000 g for 10 minutes and washed with 500  $\mu$ l of 70% ethanol. The genomic DNA was re-suspended in 80  $\mu$ l 10 mM TrisHCl (pH8.0) buffer and determined with Qubit fluorometer (Invitrogen, Carisbad, Calif.). One microgram of total genomic DNA was digested with restriction endonuclease BamH and SacII. The genomic DNA fragments were separated in 1% agarose gel electrophoretically and transferred onto the zeta-probe membrane (BioRad) with alkaline capillary transfer method. A 3.8 kb genomic DNA fragment containing the Alg3 sequence was used for preparation of the biotin-labeled probe. The genomic DNA in Zeta-probe membrane was hybridized with the biotin-labeled probe overnight in 60° C. hybridization oven. The genomic DNA on hybridized membrane was visualized with North2South chemiluminescent detection kit (Pierce Protein Research Products, Rockford, Ill.).

## Spore Production and Germination.

The spore production on the PDA or CM agar plates described above was excised with plastic closures of culture tubes in 27 mm diameter and transferred into the 50 ml centrifuge tubes containing 25 ml 0.8% tween 80. The spores were released from the agar surface by scraping with plastic loops and vortexed with vortex mixer at top speed. The spores were diluted properly and enumerated with a hemacytometer. The spore production in a unit area (cm<sup>2</sup>) was determined. For spore germination, 1 $\times$ 10<sup>5</sup> spores per well were added into each well of 24 well Schwarz sensoplate and incubated in the microscopic incubator with temperature control at 30° C. The spore germination was automatically imaged hourly for 24 hours through the Olympus inverted system microscope (Olympus America Inc., Center Valley, Pa., USA). The spore germination was visualized with Adobe Photoshop CSS (San Jose, Calif.) and counted manually.

## Citric Acid Measurements.

Citric acid concentrations were determined with an endpoint spectrophotometric enzyme assay as described in the instruction from the manufacturer (R-Biopharm AG/Roche, Darmstadt, Germany) with a proper dilution.

Table 2 shows oligonucleotides used in the methods.

TABLE 2

Oligonucleotides		
Name	Sequence	Product size (kb)
Alg3A construction		
Alg3-ForScr	CGGTTTCCCTTCAGTTCCAGT (SEQ ID NO: 5)	
Alg3-1	GTAACGCCAGGGTTTTCCAGTCACGACGCATAACTTCTCTCCCCTCC (SEQ ID NO: 6)	1.06
Alg3-2	ATCCACTTAACGTTACTGAAATCTCCAAC <u>TTCATGGACACACAGACC</u> (SEQ ID NO: 7)	
Hph-F	GGTCTGTGTGTGCCATGAAGTTGGAGATTTAGTAACGTTAAGTGGAT (SEQ ID NO: 8)	1.49
Hph-R	GCTACTACTGATCCCTCTGCGTCGGAGACAGAAGATGATATTGAAGGAG (SEQ ID NO: 9)	1.03
Alg3-3	CTCCTTCAATATCATCTTCTGTCTCCGACGCAGAGGGATCAGTAGTAGC (SEQ ID NO: 10)	
Alg3-4	GCGGATAACAATTTACACAGGAAACAGCCGTGAGAGGTTTGTAGTACG (SEQ ID NO: 11)	
Alg3-RevScr	AAGCTGAGAGCGACATCTTCA (SEQ ID NO: 12)	
hyg-RevScr	GTA <u>CTTCTACACAGCCATCGGTCCA</u> (SEQ ID NO: 13)	
hyg-ForScr	GTA <u>CTTCTACACAGCCATCGGTCCA</u> (SEQ ID NO: 14)	
Alg3A + Alg3 construction		
pryGScr	TCTGCTGTCTTGCATGAGGTCCTT (SEQ ID NO: 15)	
pyrGScr	Agcgtaggacaaggctcgtctctgt (SEQ ID NO: 16)	2.34
5-pyrG5F	GTAACGCCAGGGTTTTCCAGTCACGACG <u>tttaaac</u> ATGCATCATTCTCCGCTTTGT (SEQ ID NO: 17)	1.69
5-pyrG3R	agaaagagtcacoggtcacGacatcgccaatcacctcaatcac (SEQ ID NO: 18)	1.47 1.47
ble5F	gtgattgaggtgattggcgatgtCgtgaccoggtgactctttct (SEQ ID NO: 19)	1.23

TABLE 2-continued

Oligonucleotides		
Name	Sequence	Product size (kb)
Ble3R	<b>TCCAACCTTGTAGCAACCAAAGCTTCGAGCGTCCCAAAACCT</b> (SEQ ID NO: 20)	
Alg3-5F1	<u>AGGTTTTGGGACGCTCGAAGCTTTGGTTGCTACAAGTTGGA</u> (SEQ ID NO: 21)	
Alg3-3R1	TCAAGTAGAGCACAGCAAATAGTATCTGA (SEQ ID NO: 22)	
Alg3-5F2	<b>TCAGATACTATTGCTGTGCTCTACTTGA</b> (SEQ ID NO: 23)	
Alg3-3R2	ttgatccttgtgccacaccaTCCTACGTGGTCATCGATACCA (SEQ ID NO: 24)	
3-pyrG5F	<u>TGGTATCGATGACCACGTAGGA</u> tggtgtggcacaaggatcaa (SEQ ID NO: 25)	
3-pyrG3R	GCGGATAACAATTTACACAGGAAACAGCgtttaactgtgccagtcaa ttgtccgaagt (SEQ ID NO: 26)	
Alg3Seq-1	TACAGACGCGGTACGCATGT (SEQ ID NO: 27)	
Alg3seq-2	TGCTATTGTCCACAGATACCGAGA (SEQ ID NO: 28)	
Alg3seq-3	GAGCTAACCAGACAGTTCATGT (SEQ ID NO: 29)	
Alg3seq-4	Tcgtcgtaccgcattgatcct (SEQ ID NO: 30)	

## Example 2

Genetic Inactivation of Alg3 in *A. niger*

This example describes methods used to genetically clone and then inactivate Alg3 in *A. niger* strain 11414KusA. Based on these teachings, one skilled in the art will appreciate that Alg3 can be similarly inactivated in other strains of *Aspergillus*.

Alg3 has been identified and characterized in *Arabidopsis thaliana*, *Homo sapiens*, *Pichia pastoris*, *Trypanosoma brucei* and *Saccharomyces cerevisiae* (see for example Korner et al., *EMBO J.* 18(23): 6816-6822, 1999; Davidson et al., *Glycobiology* 14(5):399-407, 2004; Manthri et al., *Glycobiology* 18(5): 367-383, 2008; Kajiura et al., *Glycobiology* 20(6):736-751, 2010). A database search based on the amino acid sequence of *S. cerevisiae* Alg3 identified a putative  $\alpha$ -1,3-mannosyltransferase gene in JGI (DOE Joint Genome Institute)-*A. niger* genome database (jgilAspni5142720). The *A. niger* Alg3 gene contains two introns and its 1400 bp open reading frame (nt 1186-1306, 1393-1916 and 1989-2582 of SEQ ID NO: 1) encodes a protein consisting of 413 amino acids (SEQ ID NO: 2), which contains one potential N-glycosite at the amino acid position 374. The predicted Alg3 amino acid sequence has 39% sequence identity to the *S. cerevisiae* Alg3.

Alg3 was functionally inactivated in *A. niger* using a gene deletion vector constructed by yeast gap repairing approach. The 5'- and 3'-end of the hygromycin marker (hph) gene was flanked with about 1 kb upstream and downstream fragments of Alg3 coding region that were isolated by PCR from *A. niger* genomic DNA. The DNA sequence of the upstream and downstream fragments was confirmed by DNA sequencing analysis. The Alg3 in *A. niger* was deleted by homologous replacement with hygromycin marker (hph) gene in the kusA deletion background of *A. niger*, where the kusA gene, encoding the ortholog of the Ku70 protein in other eukary-

30

otes, was deleted for dramatically improved homologous integration efficiency. FIGS. 1A and 1B show the predicted restriction enzyme digestion patterns of genomic DNA of the parent and mutant strains with BamHI and SacII. FIG. 1C shows the Southern blotting analysis of the digested genomic DNA of parent and mutant strains. The results confirm that the Alg3 coding region in *A. niger* was replaced by the hygromycin selection marker gene (hph) in the Alg3 $\Delta$  strains.

## Example 3

Effects of Alg3 Deletion on *A. niger* Growth and Development

This example describes methods used to determine the effect genetically inactivating Alg3 in *A. niger*.

It was previously demonstrated that the deletion of Alg3 in different organisms causes underglycosylation, but no obvious phenotype changes were observed at the selected culture condition in those studies (Aebi et al., *Glycobiology* 6(4):439-444, 1996; Korner et al., *EMBO J.* 18(23): 6816-6822, 1999; Davidson et al., *Glycobiology* 14(5):399-407, 2004; Manthri et al., *Glycobiology* 18(5): 367-383, 2008; Kajiura et al., *Glycobiology* 20(6):736-751, 2010).

The effects of the Alg3 deletion were examined on CM, PDA and MM plates. As exhibited in FIGS. 2A-I, the Alg3 $\Delta$  strain grew much slower than the parent strain when grown on either CM or PDA medium plate, but there was no significant difference between the Alg3 $\Delta$  mutants and parent strain when grown on the MM medium plate. When both Alg3 $\Delta$  strain and parent strain were grown in the liquid culture of CM and PDA, the initiation of spore germination of Alg3 $\Delta$  strain was pronouncedly delayed (FIG. 3A), but the spore germination rate was not affected by the Alg3 deletion (FIG. 3B). These results demonstrate that deletion of Alg3 has significant effects on *A. niger* growth on nutrient rich media.

65

The effects of Alg3 deletion on spore production on both CM and PDA plates were also examined. The Alg3 deletion had a substantial reduction of sporulation on CM medium plate, while no obvious difference was exhibited on PDA plate (FIG. 4). The spore production at given area was enumerated with hemocytometer (Table 3). The average spore production of Alg3Δ was  $2.64 \times 10^7$  spores/cm<sup>2</sup>, about 40% of parent strain ( $6.44 \times 10^7$  spores/cm<sup>2</sup>) on CM medium plates, while average spore production per a square millimeter was similar between Alg3Δ mutant ( $7.72 \times 10^7$  spores/cm<sup>2</sup>) and parent strains ( $7.89 \times 10^7$  spores/cm<sup>2</sup>) on PDA medium plates.

TABLE 3

Strain	CM pates	PDA
11414-kusA	$6.44 \pm 1.24$	$7.72 \pm 1.78$
Alg3Δ	$2.64 \pm 0.49$	$7.89 \pm 1.18$

## Example 4

## Effects of Alg3 Deletion on Spore Germination, Growth and Citric Acid Production

This example describes methods used to measure spore germination, growth, and citric acid production in the Alg3 Δ *A. niger* strain generated in Example 1. Based on these teachings, one skilled in the art will appreciate that spore germination, growth, and citric acid production can be similarly measured in other Alg3Δ strains of *Aspergillus*.

*A. niger* strain ATCC11414 is a strain developed for industrial production of citric acid. *A. niger* morphology plays a role in citric acid production. The fungal morphology affect overall molecular regulation in response to the endogenous and exogenous factors, which include the regulations of transcription, post-transcription, translation and post-translation. Therefore, the effects of Alg3 deletion on *A. niger* growth on CAP agar plates at different pHs or in CAP liquid culture conditions was determined.

FIG. 5 shows the effects of Alg3 deletion in fungi grown on CAP medium plates at different pH conditions after 28 hours in culture. When the Alg3Δ strain was grown on the CAP medium plates at pH 1.8, its growth was relatively slower than the parent strain, where its colonies were much smaller than the parent strain. At pH 2.1, the Alg3Δ strain formed a less tight pellet than that of parent strain. Growth of the Alg3Δ strain on the CAP medium at pH 4.5 and pH 7.0 was affected more profoundly than that of parent strain, where the Alg3Δ strain formed thinner layer of biomass on the agar plates than that of parent strain. These results show that the Alg3 deletion reduces the normal growth of *A. niger* at different levels on CAP culture medium plates at different pHs.

The spore germination of the Alg3Δ and parent strains in CAP liquid culture medium was also examined by using automated microscopic imaging, enumerating the germination manually in a same visual unit area, and expressing spore germination as a percentage of total spores at the same visual unit area. FIG. 6 shows the different dynamics of spore germination between the Alg3Δ and parent strains. The spore germination of Alg3Δ strain began as early as 3 hours after spore inoculation, while the parent strain was not initiated until 6 hours after spore inoculation. After 8 hours

inoculation, more than 32% spores of Alg3Δ mutant germinated, while only 10% of parent strain spores did (FIG. 6A, top panels; and FIG. 6B). After 15 hours growth in CAP liquid medium, more than 90% spores of Alg3Δ strain germinated, while only 50% spores of parent strain did (FIG. 6A, bottom panels; and FIG. 6B). After 24 hours of growth in the liquid culture, about 75% spores of parent strain germinated, while the Alg3Δ strain achieved 94% of germination rate (FIG. 6B). The Alg3 deletion leads to earlier germination and a higher germination rate than parent strain.

The effect of Alg3 deletion on citric acid production was determined in CAP flask cultures. FIG. 7 shows the time course of citric acid production in CAP liquid medium. The yield of citric acid production was similar between the Alg3Δ mutant and parent strain in 2 days culture. After 3 days culture, the average citric acid production by Alg3Δ mutants was 8.8 g/l citric acid, while the parent strain only produced 5.8 g/l. After 4.5 days of culture, the parent strain accumulated 18.8 g/l citric acid and the Alg3Δ mutant produced 33.3 g/l citric acid (more than 70% higher than the parent strain). Thus, the Alg3 deletion substantially improves the citric acid production in *A. niger*.

The effect of Alg3 deletion on citric acid production was also examined by complementation of its original gene into the alg3Δ mutant. FIG. 8B shows the citric acid production in CAP liquid medium after 10 days of culture. The yield of citric acid production was similar between the alg3Δ complemented (cAlg3Δ) mutant and parent strain, but much lower than the alg3Δ mutant in 10 days culture. After 10 days culture, the average citric acid production by Alg3Δ mutants was 46.1 g/l citric acid, while the parent and calg3Δ strain only produced 34.8 and 29.4 g/l, respectively.

## Example 5

## pPTRpGPDALaeA Plasmid Vector Construction

This example describes methods used to generate the pPTRpGPDALaeA plasmid vector (FIG. 13). One skilled in the art will appreciate that although gpdA and LaeA sequences were used from *Aspergillus nidulans*, one skilled in the art will appreciate that variants of these sequences can be used in the fungi and methods provided herein, such as gpdA and LaeA sequences from other *Aspergillus* species. In one example, a gpdA sequence having at least 80%, at least 90%, at least 95%, or at least 98% sequence identity to SEQ ID NO: 37 is used in the fungi and methods provided herein.

The *Aspergillus nidulans* glyceraldehyde 3-phosphate dehydrogenase (gpdA) promoter (SEQ ID NO: 37) was isolated from the pAN8-1 plasmid DNA using the primer set gpdA5F/gpdA3R (gpdA5F: CGCAGATCTC AAGCTG-TAAG GATTCGGCA SEQ ID NO: 38; gpdA3R: CAC-CGGGCC ATCTCAAACA TTGTGATGTC TGCT-CAAGCG SEQ ID NO: 39) and the LaeA coding sequence of genomic DNA from *A. nidulans* (SEQ ID NO: 40) obtained by PCR using LaeA5F/LaeA3R (LaeA5F: CGCT-TGAGCA GACATCACAA TGTTTGAGAT GGGCCCG-GTG; SEQ ID NO: 42; LaeA3R: CGCAGATCTG AGGAT-TATGA GAAGGGAGC; SEQ ID NO: 43).

The DNA fragment of pGPDA and LaeA was filled together by overlap PCR and a HindIII restriction enzyme site was introduced at both 5'- and 3'-end of the DNA fragment. The DNA fragment (SEQ ID NO: 44) and pPTR1 plasmid DNA were cut with Hind III and ligated together by a quick DNA ligation kit at 25° C. for 30 min. The ligated plasmid DNA was transferred into the Top10 *E. coli* competent cells by lithium acetate mediated transformation. The

## 43

transformed bacterial colonies were screened for the DNA fragment insertion by PCR with the primers *gpdA5F* (SEQ ID NO: 38) and *LaeA3R* (SEQ ID NO: 43). The plasmid DNA for the selected transformed colonies was prepared for restriction enzyme confirmation and further expression vector construction.

## Example 6

## pRS426-LaeA Vector Construction

This example describes methods used to generate a transgene containing *A. niger* *LaeA* (FIG. 14).

PCR was performed to isolate DNA fragments of *A. niger* *pyrG* upstream region (SEQ ID NO: 45), *trpC* transcriptional terminator of *A. nidulans* (SEQ ID NO: 46), pyrithiamine resistance gene (*ptrA*) of *Aspergillus oryzae* (SEQ ID NO: 47), and *A. niger* *pyrG* downstream region (SEQ ID NO: 48), using primers *pyrGU5F/PTRU3R* (*pyrGU5F*: GTAACGCCAG GGTTCCTCCA GTCACGACGT TTAAACATGC ATCATTCTCC CGCTTTGT, SEQ ID NO: 49; *pyrGU3R*: TGCCGAAATC CTTACAGCTT GAAGCTTCAT CGCCAATCAC CTCAATCAC, SEQ ID NO: 50), *Trp5F/Trp3R* (*Trp5F*: AGCTCCCTTC TCATAATCCT CAAGCTTGGA CCGATGGCTG TGTAGAAGT, SEQ ID NO: 51; *Trp3R*: CGTAATCAAT TGCCCGTCTG TCAGAGAGCG GATTCCTCAG TCTCGT; SEQ ID NO: 52), *PTR5F/PTR3R* (*PTR5F*: ACGAGACTGA GGAATCCGCT CTCTGACAGA CGGGCAATTG ATTACG, SEQ ID NO: 53; *PTR3R*: ACAGCAGTGC TTATCTGCGA TGACGAGCCG CTCTTGCATC TTTGT, SEQ ID NO: 54) and *PyrGD5F/PTRD3R* (*pyrGD5F*: ACAAAGATGC AAGAGCGGCT CGTCATCGCA GATAAGCACT GCTGT; SEQ ID NO: 55, *pyrGD3R*: TGAGACGCTG TTTCACCGAG TACATCGCCA ATCACCTCAA TCAC, SEQ ID NO: 56), respectively.

As shown in FIG. 14, the DNA fragment of pGDPALaeA (SEQ ID NO: 44) was isolated from pPTRpGDPALaeA (FIG. 13) by HindIII digestion. The yeast gap repairing vector pRS426 was double digested with restriction enzyme HindIII and XhoI. Hundred nanograms of each DNA fragment generated from PCR (i.e., SEQ ID NOs: 45-56) or restriction enzyme digestions were used for *S. cerevisiae* transformation. The gap repairing plasmid DNA in the total *S. cerevisiae* genomic DNA was isolated by transferred into the Top10 *E. coli* cells.

The transformed plasmid DNA was confirmed by PCR and digested with PmeI. The PmeI DNA fragment (SEQ ID NO: 57) was used for *A. niger* transformation.

One skilled in the art will appreciate that although the *pyrG* upstream and downstream sequences, *trpC* transcriptional terminator sequence, and *ptrA* sequence used were from particular organisms, one skilled in the art will appreciate that variants of these sequences can be used in the fungi and methods provided herein, such as those from other *Aspergillus* species. In one example, *pyrG* upstream and downstream sequences having at least 80%, at least 90%, at least 95%, or at least 98% sequence identity to SEQ ID NO: 45 and 48 are used in the fungi and methods provided herein. In one example, a *trpC* transcriptional terminator sequence having at least 80%, at least 90%, at least 95%, or at least 98% sequence identity to SEQ ID NO: 46 is used in the fungi and methods provided herein. In one example, a *ptrA* sequence having at least 80%, at least 90%, at least 95%, or

## 44

at least 98% sequence identity to SEQ ID NO: 47 is used in the fungi and methods provided herein.

## Example 7

Expression of pGDPALaeA-Containing Transgene in *A. niger* Alg3Δ

This example describes methods used to introduce pGPDALaeA (SEQ ID NO: 57) into *A. niger*.

The originally transformed *A. niger* colonies were picked from the minimal medium plates with 0.1 μg/ml pyrithiamine hydrobromide selection on minimal medium agar plates without thiamine supplementation. The single spore colonies were picked for spore production after the initial transformant spores were grown on the same selection medium plates. The biomass was harvested from the single spore colony isolates grown in minimal medium with pyrithiamine selection and dried in the VirTis bench top freeze dryer. The genomic DNA was prepared for PCR confirmation of pGDPALaeA insertion in transgenic *A. niger*. As shown in FIG. 15A, the primer set of *PTR5F* (SEQ ID NO: 53) and *PTR3R* (SEQ ID NO: 54) was used to confirm the presence of *A. oryzae* pyrithiamine resistance gene (*ptrA*) in transgenic *A. niger* with the expected size of 2 kb PCR DNA fragment. As shown in FIG. 15B, the primer set *LaeA5F* (SEQ ID NO: 42) and *TRP3R* (SEQ ID NO: 52) was used to demonstrate that the transgene *A. nidulans* *LaeA* was under the control of *gdpA* promoter and *trpC* transcriptional terminator of *A. nidulans* with the expected 3.4 kb PCR fragment size. The genomic DNA of parent strain and the plasmid DNA of transgene vector carrying the pGDPALaeA fragment were used for negative and positive references.

## Example 8

## Increased Production of Citric Acid

This example describes methods used to demonstrate that citric acid production was increased in the presence of increased expression of *LaeA*, alone or in combination with deletion of *Alg3*.

Citric acid was produced as described in Example 1.

As shown in FIG. 16, of citric acid production was increased in the *Alg3Δ* mutant (*Alg3*), and in the mutants over-expressing *LaeA* in *Alg3Δ* (*LaeA-1* and *LaeA-2*), as compared to the parent strain (*kusA*).

## Example 9

## pBSK-LaeA a Plasmid Vector Construction

This example describes methods used to generate the pBSK-LaeAΔ plasmid vector (FIG. 17A). The 5' (SEQ ID NO: 61) and 3' (SEQ ID NO: 62) ends of the *Aspergillus niger* *LaeA* gene were isolated from *A. niger* genomic DNA by PCR using oligonucleotide sets D1 & D2, and D5 & D6 (see Table 4 below). The *hph* expression cassette was isolated from plasmid vector DNA of pCB1003 (SEQ ID NO: 63) by PCR using the oligonucleotide sets D3 & D4. The DNA fragments were assembled together into the backbone plasmid vector of pBSK linearized with restriction endonucleases of both HindIII and PstI using the Gibson assembly cloning kit. The assembled plasmid DNA was transferred into the Top10 *E. coli* competent cells by lithium acetate mediated transformation. The transformed bacterial

## 45

colonies were screened for the DNA fragment insertion by restriction endonuclease digestion of PvuII and XhoI. The plasmid DNA for the transformed colonies was prepared and digested with endonucleases of HindIII and XbaI for further LaeA deletion in *A. niger*.

TABLE 4

Oligonucleotide Primers for <i>A. niger</i> LaeA Deletion		
Name	Sequence	SEQ ID NO:
D1	gtcgcggtatcgataAGCTTCAAGACAGC GGCTGCAA	67
D2	gccgaccgTGGTAGAGATACAGGGGTTTC	68
D3	ctctaccaCGGTCCGCATCTACTCTATTC	69
D4	gcgactgaGCTGGAGCTAGTGGAGGT	70
D5	gctccagcTCAGTCGCATCTTTCAGT	71
D6	atcccccggtgcaTGGTAGGCTGTCT CAAGG	72
SC7	ACTCGCAGCAGAGATGCCATCT	73
SC8	CGTTATGTTTATCGGCACCTTTCAT	74

## Example 10

pRS426-*A. niger* LaeA Complementation Plasmid  
Vector Construction

This example describes methods used to generate the pBSK-LaeAΔ plasmid vector (FIG. 17B). The entire LaeA gene (2.34 kb; SEQ ID NO: 64) containing 740 bp of promoter region and 330 bp of transcriptional terminator region was isolated from *A. niger* genomic DNA by PCR using oligonucleotide set CP9 & CP10 (see Table 5 below). The DNA fragment was ligated into the plasmid DNA of pRS426-*A. nidulans* LaeA over-expression vector at a HindIII restriction endonuclease site after the pGPDALaeA fragment in the vector was removed by agarose gel separation. The assembled plasmid DNA was transferred into the Top10 *E. coli* competent cells by lithium acetate mediated transformation. The transformed bacterial colonies were screened for the DNA fragment insertion by restriction endonuclease digestion of BamHI and XhoI. The plasmid DNA for the transformed colonies was prepared and digested with PmeI restriction endonuclease for *A. niger* LaeA expression at pyrG locus to complement the LaeA deletion in *A. niger* LaeAΔ mutants.

TABLE 5

Oligonucleotide Primers for <i>A. niger</i> LaeA Complementation		
Name	Sequence	SEQ ID NO:
CP9	aggtgattggcgatgaTGCCGTTTCAGC TGCTCTGC	75
CP10	acagccatcggtccaTCTCTCCTCGTA ACGCCTG	76
SC11	ggatagaatcgggtgccgctgatct	77
SC12	gagaacctggcaccgaaggt	78

## 46

## Example 11

Deletion of *A. niger* LaeA Gene in *A. niger*

This example describes methods used to introduce the HindIII/XbaI DNA fragment contain the LaeA deletion cassette (SEQ ID NO: 65) into *A. niger*.

The originally transformed *A. niger* colonies were picked from the minimal medium plates with 100 μg/ml Hygromycin B selection on minimal medium agar plates. The single spore colonies were picked for spore production after the initial transformant spores were grown on the same selection medium plates. The biomass was harvested from the single spore colony isolates grown in complete medium with Hygromycin B selection and dried in the VirTis bench top freeze dryer. The genomic DNA was prepared for PCR confirmation of the replacement of LaeA gene coding region in transgenic *A. niger*. As shown in FIG. 18, the primer set SC7 and SC8 (see Table 4) was used to confirm the hph replacement of the LaeA gene in transgenic *A. niger* with the expected 2 kb PCR DNA fragment. The genomic DNA of parent strain was used for negative and positive references.

## Example 12

Expression of pGDPALaeA-Containing Transgene  
in *A. niger*

This example describes methods used to introduce the *A. niger* LaeA gene into the pyrG locus in the *A. niger* LaeAΔ mutant to complement the loss of LaeA function with PmeI DNA fragment (SEQ ID NO: 66).

The originally transformed *A. niger* colonies were picked from the minimal medium plates with 0.1 μg/ml pyrithiamine hydrobromide selection on minimal medium agar plates without thiamine supplementation. The single spore colonies were picked for spore production after the initial transformant spores were grown on the same selection medium plates. The biomass was harvested from the single spore colony isolates grown in minimal medium with pyrithiamine selection and dried in the VirTis bench top freeze dryer. The genomic DNA was prepared for PCR confirmation of LaeA gene insertion in transgenic *A. niger*. As shown in FIG. 19, the primer set of PTR5F (SEQ ID NO: 53) and PTR3R (SEQ ID NO: 54) was used to confirm the presence of *A. oryzae* pyrithiamine resistance gene (ptrA) in transgenic *A. niger* with the expected size of 2 kb PCR DNA fragment. The genomic DNA of the parent strain and the plasmid DNA of the transgene vector carrying the *A. oryzae* pyrithiamine resistance gene (ptrA) fragment were used for negative and positive references.

## Example 13

The Effects of LaeA Gene Deletion and  
Complementation and Over-Expression of *A. nidulans*  
LaeA Gene on Citric Acid Production

This example describes methods used to demonstrate that citric acid production in *A. niger* mutant strains was significantly influenced by perturbing the LaeA gene expression in *A. niger*. At least three individual clones per mutant strain were selected for spore production on complete medium plates at 30° C. for 4 days. The spores were counted with a hemocytometer. Citric acid production culture was initiated by inoculation with a 1×10<sup>6</sup> spore/ml of 75 ml citric acid production culture medium in a 250 ml baffled Erlenmeyer

glass flask siliconized with a 5% solution of dichlorodimethylsilane in hexane solvent. The culture was maintained at 30° C. and 220 rpm in shaker incubator for 5 days. One microliter of culture was harvested and briefly spun down in a microcentrifuge at full speed. The supernatants of cultures were diluted 200-fold with dH<sub>2</sub>O prior to citric acid measurement. The citric acid in the supernatant was quantified with the citric acid assay kit from R-Biopharm AG. The detailed procedure for the citric acid assay was performed essentially according to the manufacturer's description. The results in FIG. 20 show that deletion of the *LaeA* gene in *A. niger* (*LaeAΔ*) led to loss of citric acid production in citric acid production culture. When the original *A. niger* *LaeA* gene was used for complementation in the *LaeAΔ* mutant (*cLaeAΔ*) at the *pyrG* locus, the citric acid production was partially recovered, which indicates the importance of chromosome location of *LaeA* gene. When *A. nidulans* *LaeA* was over-expressed in the *A. niger* parent strain (*LaeA*), the citric acid production was higher than the parent strain. This indicates that the *LaeA* is involved in citric acid production in *A. niger*.

#### Example 14

##### Enhancement of Itaconic Production by *LaeA* Complementation *A. terreus* Growth Conditions

All strains were maintained on potato dextrose agar (PDA), which is composed of 39 g/L of Difco Potato Dextrose Agar (BD, USA). Spore inocula were grown on PDA and harvested after 5 days. The transformant was selected for either hygromycin or bleomycin resistance on minimum media (MM, 50 ml 20× nitrate salts, 1 ml 1000× trace elements, 1 ml 1000× vitamin stock, 10 g glucose, pH 6.5, 100 μg/ml hygromycin or 250 μg/ml bleomycin). The 20×NO<sub>3</sub> salts contains in one liter H<sub>2</sub>O: Na<sub>2</sub>NO<sub>3</sub>, 120 g; KCL, 10.4 g; MgSO<sub>4</sub>·7H<sub>2</sub>O, 10.4 g; KH<sub>2</sub>PO<sub>4</sub>, 30.4 g. The 1000× vitamin solution (yellow) contains in 100 ml H<sub>2</sub>O: biotin, 0.01 g; pyridoxine-HCl, 0.01 g; thiamine-HCl, 0.01 g; riboflavin, 0.01 g; para-amino benzoic acid, 0.01 g; and nicotinic acid (or niacin), 0.01 g, filtered and stored at 4° C. The 1000× trace elements contains per 100 ml H<sub>2</sub>O: ZnSO<sub>4</sub>·7H<sub>2</sub>O, 2.2 g; H<sub>3</sub>BO<sub>3</sub>, 1.1 g; MnCl<sub>2</sub>·4H<sub>2</sub>O, 0.5 g; FeSO<sub>4</sub>·7H<sub>2</sub>O, 0.5 g; CoCl<sub>2</sub>·6H<sub>2</sub>O, 0.17 g; CuSO<sub>4</sub>·5H<sub>2</sub>O, 0.16 g; Na<sub>2</sub>MoO<sub>4</sub>·2H<sub>2</sub>O, 0.15 g; Na<sub>2</sub>EDTA, 5 g, which is added in the order, boil and cool to room temperature and adjusted the pH to 6.5 with KOH. For pyrithiamine marker selection, the thiamine-HCl was eliminated from minimum media.

About 1×10<sup>6</sup> spores/ml were used to inoculate 50 ml of itaconic acid production media in 250 ml glass flask. The itaconic acid production medium essentially followed the description of Riscaldati et al. (*J Biotechnol* 83:219-230, 2000). Itaconic acid production medium contains in grams per liter (g/l): glucose, 100; ammonium sulfate, 2.36; potassium phosphate dibase, 0.11; magnesium sulfate heptahydrate, 2.08; calcium chloride dihydrate, 0.13; sodium chloride, 0.074; copper sulfate pentahydrate 2×10<sup>-4</sup>; ferrous sulfate heptahydrate, 5.5×10<sup>-3</sup>; manganese chloride tetrahydrate, 7×10<sup>-4</sup>; and zinc sulfate heptahydrate, 1.3×10<sup>-3</sup>. The flask cultures were shaken at 30° C. and 150 rpm. At least three biological replicates for each strain were used for each experiment.

##### Itaconic Acid Measurement

The itaconic acid method followed the previous description by Dickman (*Analytical Chem* 24: 1064-1066, 1952) with some modification. The reaction volume was scaled

down to 1.4 ml for each reaction. Ten to 250 μl of culture supernatant or itaconic acid standard dissolved in dH<sub>2</sub>O were used. The proper amount of culture supernatant or itaconic acid standard was added into 690 to 930 μl dH<sub>2</sub>O in 1.5 ml microcentrifuge tube. Then, 60 μl of freshly prepared metaphosphoric acid (0.425 g/ml) was added into the tube, mixed well, and incubated at 4° C. for 10 minutes. Next, 400 μl of freshly prepared potassium permanganate (5 mM) from stock solution (100 mM) was added, mixed well and incubated in the dark at room temperature for 10 minutes. Finally, the absorbency of the reaction mixture was determined at 530 nm spectrophotometrically.

##### *LaeA* Deletion Construct

The DNA fragments corresponding to the upstream and downstream regions of the *LaeA* gene in *A. terreus* and *hph* were isolated with oligonucleotide pairs 1546/1548, 1551/1553 and 1549/1550, respectively (see Table 6 for oligonucleotide sequences). These three DNA fragments (SEQ ID NOs: 89-91) were fused together by double-joint PCR (Yu, et al., *Fungal Genetics and Biology* 41: 973-981, 2004). The whole *LaeA* deletion fragment was isolated by PCR with oligonucleotide pair 1547/1552 (see Table 6 and SEQ ID NO: 104). FIG. 21A depicts the *LaeA* deletion construct. FIG. 21B shows genomic PCR confirmation of selected *LaeA* deletion mutants with oligonucleotide pair 1546/107 or 108/1553.

##### *LaeA* Complementation Construct

The *LaeA* complementation construct was prepared for targeting upstream of the *LaeA* locus in *A. terreus* using the Gibson Assembly method. The DNA fragments of the whole *LaeA* gene, *ble* selection marker gene, and *hph* gene used for gene deletion were isolated by PCR with oligonucleotide pairs 1554/1555, 1556/1557, and 1558/1559, respectively (see Table 6). In order to isolate the whole transgene cassette, an *Xba*I restriction endonuclease site was introduced into the upstream fragment of 1554/1555. The three DNA fragments (SEQ ID NOs: 107-109) were then assembled into the cloning vector of pBluescript SK(-). FIG. 22A is a diagram illustrating the *LaeA* complementation constructs. The insertion of *LaeA* complementation was confirmed by oligonucleotide pair 1556/107. FIG. 22B shows gel electrophoresis of genomic PCR products corresponding to *ble* and part of *hph*.

##### *LaeA* Over-Expression Construct

The *Aspergillus nidulans* *LaeA* gene under the control of *A. nidulans* *gpdA* promoter and *TrpC* transcription terminator, and the selection marker gene of pyrithiamine resistance gene (*ptrA*) of *Aspergillus oryzae*, were isolated from the transgene expression construct used for *A. niger* transformation (as described in Example 6) with oligonucleotide pair 1563 and 1564 (Table 6). Due to the extremely high homologous replacement rate in *A. terreus*, the *laeA* over-expression transgene cassette was designed to replace the 127 bp genomic DNA fragment of the non-gene region located at 1571 bp upstream in *A. terreus* *laeA* coding gene sequence. The corresponding DNA fragments flanked to the both ends of *LaeA* expression cassette for homologous replacement were isolated by PCR with oligonucleotide pairs 1561/1562 and 1565/1566 (Table 6). The three DNA fragments (SEQ ID NOs: 96-98) were fused together in pBluescript SK(-) by Gibson Assembly. The restriction endonuclease site for *Pme*I was introduced into the 5' end of the upstream region fragment of 1561/1562 (SEQ ID NO: 96) and the 3' end of the downstream region fragment of 1565/1566 (SEQ ID NO: 110), which enables the isolation of whole transgene expression construct. FIG. 24A is a diagram of the transgene over-expression cassette of the *A.*

*nidulans* LaeA gene. Oligonucleotides 1563, 1198 and 1564 (see Table 6) were used for genomic PCR to confirm the gene insertion. FIG. 24B shows gel electrophoresis of PCR products corresponding to the oligonucleotide pairs 1198/1564 and 1563/1564 in selected transgenic strains.

#### *A. terreus* Protoplast Isolation and Transformation

*A. terreus* ( $10^6$  conidia/ml) was inoculated into 100 ml of complete media containing in g/l: D-glucose, 10; peptone, 2; yeast extract, 1; casamino acids, 1; 20× nitrate salts, 50 ml; trace elements, 1 ml; vitamin solution, 1 ml (pH 6.5) or MM without thiamine-HCl. The inoculated culture was shaken at 150 rpm and 30° C. for 16-22 hours. The mycelia were harvested by filtering the culture through Miracloth and rinsed well with sterile dH<sub>2</sub>O. The mycelia (mass of approximately several bean sizes) were digested with 20 ml of 20 mg/ml lysing enzyme (L1412, Sigma) that was dissolved in osmotic washing buffer (0.5 M KCl, 10 mM sodium phosphate, pH 5.8) for approximately 2-3 hours at 80 rpm and 30° C. Protoplasts were filtered through sterile miracloth into a 50 ml sterile centrifuge tube and centrifuged for 10 minutes at 1000×g and 4° C. Protoplasts were washed twice with 20 ml washing solution (0.6M KCl, 0.1 M Tris-HCl, pH 7.0) and once in 10 ml conditioning solution (0.6M KCl, 50 mM CaCl<sub>2</sub>, 10 mM Tris-HCl, pH 7.5).

For transformation, 1-2 µg DNA was added to  $2 \times 10^7$  protoplasts in 0.1 ml conditioning solution. After addition of 25 µl PEG solution, the protoplasts were incubated on ice for 20 minutes. An additional 500 µl PEG solution was carefully added, mixed and incubated on ice for additional 5 minutes. Finally, 1 ml cold conditioning solution was added and mixed by inverting the tube several times. About 300-500 µl of transformed protoplasts were pipetted into 12 ml melted MM agar (MM+0.6M KCl+1.5% agar+antibiotics) tubes at 50° C. The transformed protoplasts were mixed well by inverting the tubes 3-4 times and poured directly onto the 100 mm petri dishes. The petri dishes were incubated overnight at 30° C. An additional 8 ml of regular MM agar with proper antibiotics was overlaid and further incubated at 30° C. for approximately 3-4 days. The transformed colonies were picked and further grown on 2 ml MM agar gel for spore production.

TABLE 6

Oligonucleotides		
Oligo	DNA sequence	SEQ ID NO:
107	GTACTTCTACACAGCCATCGGTCCA	79
108	CGTTATGTTTATCGGCACTTTGCAT	80
1546	ACAGGTA CTCCATCTTGACTGGT	81
1547	<u>TCTcctccaacgtccgatct</u>	82
1548	acctccactagctccagcaagccgaacaga ggtaaagacga	83
1549	tcgtctttacctctgttcggcttgctggag ctagtggaggtca	84
1550	taccaacgtgagaccattttTCTcggtcggc atctactctattcct	85

TABLE 6-continued

Oligonucleotides		
Oligo	DNA sequence	SEQ ID NO:
5	1551 <u>aggaatagagtagatgccgaccgAGAAaat</u> ggtcgcacgttggt a	86
	1552 AAGCGTCTCTTTCCTGGGTCTT	87
10	1553 TGCCAGTTCTGTTGGACATCTCT	88
	1554 tcgaggtcgacggtatcgat atctagaACA GGTACTTCCATCTTG TAC	89
	1555 ggtcactgtTCCTGGCAGCTGACATTG	90
15	1556 ctgccaggaACAGTGACCGGTGACTCT	91
	1557 aagcagcagatACGACCGTTGATCTGCTTG	92
	1558 acggtcgtATCTGCTGCTTGGTGAC	93
20	1559 actagtgatccccgggctgcagCGGTCCG GCATCTACTCTATTC	94
	1561 cgaggtcgacggtatcgataGTTTAAACCT CCCAGGTACCGACTAAC	95
25	1562 ctcaatcacagATCATGTTTGGGTGGGTTC	96
	1563 aaacatgatcTGTGATTGAGGTGATTGGCG	97
	1564 ctctgtgectACAGCAGTGCTTATCTGC GATG	98
30	1565 gcaactgctgtAGGCACAGTAACAGGTAG GTAGACAG	99
	1566 agtggatccccgggctgcaGTTTAAACTC CGCACCACGAAAGCAACT	100

#### Results

Itaconic acid production in selected LaeA deletion (LaeAΔ) and LaeA complementation (cLaeAΔ) mutants of *A. terreus* was tested after four days of culture in itaconic acid production medium. The parental *A. terreus* strain was also tested as a control. As shown in FIG. 23, deletion of LaeA significantly decreased the production of itaconic acid. However, complementation of the LaeA gene in *A. terreus* enabled itaconic acid production to similar levels observed in the *A. terreus* parent.

An additional study was performed to test the effect of overexpression of the LaeA gene in *A. terreus*. FIG. 25 is a graph showing itaconic acid production in selected *A. terreus* transgenic strains having the *A. nidulans* LaeA overexpression transgene following five days of culture in itaconic acid production medium. Also shown in FIG. 25 is the percentage of itaconic acid production relative to the parent strain. The results demonstrated that overexpression of LaeA in *A. terreus* led to a significant increase in itaconic acid production relative to itaconic acid production in the parental strain.

In view of the many possible embodiments to which the principles of the disclosed invention may be applied, it should be recognized that the illustrated embodiments are only examples of the disclosure and should not be taken as limiting the scope of the invention. Rather, the scope of the invention is defined by the following claims. We therefore claim as our invention all that comes within the scope and spirit of these claims.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 110

<210> SEQ ID NO 1  
 <211> LENGTH: 3888  
 <212> TYPE: DNA  
 <213> ORGANISM: Aspergillus niger  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (1186)..(1306)  
 <220> FEATURE:  
 <221> NAME/KEY: Intron  
 <222> LOCATION: (1307)..(1392)  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (1393)..(1916)  
 <220> FEATURE:  
 <221> NAME/KEY: Intron  
 <222> LOCATION: (1917)..(1988)  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (1989)..(2582)

<400> SEQUENCE: 1

```

tctgacattg aactacagtc ataacttctc tcccctcccc ggcttccccg gtogtccttc      60
tatcgccgcc cccccgcta tccattgggtg aggccatgct atgattgggg cgctctacct      120
tgctgtcat taagtttttg tgctttgggtt gctacaaggt tggactattc ctttcttttg      180
cattgttctt ttacaccgaa actaacgcat gcccgcggcc tagtgaacgc atcgtcgtta      240
ccgtcacctt ccgctccgca taaatcttct ccttaattgc tcacatccac atttcaacat      300
gccgcccgat ctacatcccc ggtcacggtc aacgtcctcc ctcttcgccc ggacgctcct      360
ggcttccttg gtcgtcgtcg gctaccaca tgtctttccc tgtccccccc cgcgtcgcac      420
attcgccgat tctgagatga taatgtcggc agatgggcaa cctatccaga gaatccgcag      480
acgacgtcgg aaagacgaag aactccttgg ccaggatggc aatccgctcg gccagacgca      540
gcctgcagcc gatgaagagg tgtctacggt cctacaattg gaagaagaag cacagagatt      600
ggccaaggca ggccacgagt gtctgttccc caagcctcga gggattcttg gcgaattgct      660
gggttttacg agtagtggag gtatttcgac atcaacgacg acacaggcac aacagggttg      720
agagggtcgg taaccgacaa ttggaaagca aggaggactc aatcaaggct aaaataggct      780
ttgagcatgt acagcgtgaa ggaactgtgg attatatcac gaaataagcc atgaggcagt      840
cgtgttggct tcgacggagg ttggcttcgg aaattttcgg ccgggcacca aatccgaccc      900
atggcagcaa tatacccatc tttgtatcga tagtgtactt acaaaaactg tctactatat      960
gattatgcat atgcgattaa atacaactct caattgatgc acaattcgcc tcaacttcta     1020
tggtaacgaa cccacctgtt ctgcagacat gcggccgcgc ggtctgtgtg tgtccatgaa     1080
cgtaatccc aaacgggaca gctctcattg gctctccggc acgaagaggc caccgcacat     1140
ctacagctgt agaagaaagt agctggttca acaaccgcat gcaag atg gac tgg atg     1197
                                     Met Asp Trp Met
                                     1

cgc cta att cgc gat ttg tgt ttc aat ccc cga cac aca aaa tgg atg     1245
Arg Leu Ile Arg Asp Leu Cys Phe Asn Pro Arg His Thr Lys Trp Met
5          10          15          20

gct ccg ctc ctg gtc ctg ggt gac gct ttc ctc tgc gcg ctg atc atc     1293
Ala Pro Leu Leu Val Leu Gly Asp Ala Phe Leu Cys Ala Leu Ile Ile
          25          30          35

tgg aaa gtg ccc t gtaaggctac agctaagctc cgttcacacc cttttgcgac     1346
Trp Lys Val Pro
          40

```



-continued

aagtgaagca atgccactaa cctagccccg ttgctattgt ccacag at acc gag	1400
Tyr Thr Glu	
att gac tgg gcc acg tac atg caa caa ata tcg ctt tat ttg tca gga	1448
Ile Asp Trp Ala Thr Tyr Met Gln Gln Ile Ser Leu Tyr Leu Ser Gly	
45 50 55	
gaa cgc gat tat act ctc atc aga gga tca acc ggt ccc ctt gtc tac	1496
Glu Arg Asp Tyr Thr Leu Ile Arg Gly Ser Thr Gly Pro Leu Val Tyr	
60 65 70 75	
ccg gcc gcc cat gta tac agt tat acg gcc ctc tac cat ctc acc gat	1544
Pro Ala Ala His Val Tyr Ser Tyr Thr Ala Leu Tyr His Leu Thr Asp	
80 85 90	
gag ggg cgc gat att ttc ttc ggt cag ata cta ttt gct gtg ctc tac	1592
Glu Gly Arg Asp Ile Phe Phe Gly Gln Ile Leu Phe Ala Val Leu Tyr	
95 100 105	
ttg atc acg ctg gtg gtt gtg ctg tgc tgt tat aga cag tcg ggt gct	1640
Leu Ile Thr Leu Val Val Val Leu Cys Cys Tyr Arg Gln Ser Gly Ala	
110 115 120	
ccg ccg tac ttg ctt ccg ctg ctg gtc ctt tcc aag aga ctt cac agc	1688
Pro Pro Tyr Leu Leu Pro Leu Leu Val Leu Ser Lys Arg Leu His Ser	
125 130 135	
gtt tat gtc ctg cgt ctg ttc aat gat ggc ttg gcg gcg ctg gcg atg	1736
Val Tyr Val Leu Arg Leu Phe Asn Asp Gly Leu Ala Ala Leu Ala Met	
140 145 150 155	
tgg gtt gcc att ctg tta ttc atg aat cgg aag tgg acg gct gcg gtc	1784
Trp Val Ala Ile Leu Leu Phe Met Asn Arg Lys Trp Thr Ala Ala Val	
160 165 170	
gca gtg tgg tct act ggt gtt gcg att aag atg aca ctg ttg ctg ctg	1832
Ala Val Trp Ser Thr Gly Val Ala Ile Lys Met Thr Leu Leu Leu Leu	
175 180 185	
gcc ccg gct att gct gtg gtc acg gtg ctt agt ctg tcg ctt ggt cct	1880
Ala Pro Ala Ile Ala Val Val Thr Val Leu Ser Leu Ser Leu Gly Pro	
190 195 200	
agc gtg ggg ctg ggg gtt ctg gcg gtg ctt gtc cag gtaggttccc	1926
Ser Val Gly Leu Gly Val Leu Ala Val Leu Val Gln	
205 210 215	
atgaggctgt agggttggcc aaaggcaatt tgtgtgaaga cttgtctgac attgaactac	1986
ag gtt tta ctc gcg ata ccg ttc cta caa aac aac ccg gcg ggg tat	2033
Val Leu Leu Ala Ile Pro Phe Leu Gln Asn Asn Pro Ala Gly Tyr	
220 225 230	
ctc tcg cgg gcg ttc gag cta acc aga cag ttc atg ttt aaa tgg aca	2081
Leu Ser Arg Ala Phe Glu Leu Thr Arg Gln Phe Met Phe Lys Trp Thr	
235 240 245	
gtc aat tgg aga ttt gtt ggc gaa gaa gta ttc tta tct aag agc ttt	2129
Val Asn Trp Arg Phe Val Gly Glu Glu Val Phe Leu Ser Lys Ser Phe	
250 255 260	
tcc ctg gca ttg ctg gcc gtc cac att gtg ctg cta ggc gct ttt gcc	2177
Ser Leu Ala Leu Leu Ala Val His Ile Val Leu Leu Gly Ala Phe Ala	
265 270 275	
gtc act ggt tgg ctg aga tac tcc agg tct agc ttg cct gcg ttc att	2225
Val Thr Gly Trp Leu Arg Tyr Ser Arg Ser Ser Leu Pro Ala Phe Ile	
280 285 290	
ccg aat ctg cta gcg ggt cga cat cgc aca gtg tcc ctc ccc aaa ccc	2273
Arg Asn Leu Leu Ala Gly Arg His Arg Thr Val Ser Leu Pro Lys Pro	
295 300 305 310	
tac atc atg agc gtg atg ctc tcg tct ctg aca gtt ggc ttg ttg tgc	2321
Tyr Ile Met Ser Val Met Leu Ser Ser Leu Thr Val Gly Leu Leu Cys	
315 320 325	
gca agg tcc ctt cat tac caa ttc ttc gcc tac ctc tcc tgg gcg aca	2369
Ala Arg Ser Leu His Tyr Gln Phe Phe Ala Tyr Leu Ser Trp Ala Thr	

-continued

330	335	340	
ccc ttc ctc ctc tgg cgc gca ggg ttt cat cca atc ttg ctg tac ctt			2417
Pro Phe Leu Leu Trp Arg Ala Gly Phe His Pro Ile Leu Leu Tyr Leu			
345	350	355	
atc tgg gct atg caa gag tgg gct tgg aac aca ttc ccc agc acc aac			2465
Ile Trp Ala Met Gln Glu Trp Ala Trp Asn Thr Phe Pro Ser Thr Asn			
360	365	370	
ctc agt tcc atc att gtt gtc ctc tca ctt gct acc cag agt ttc ggc			2513
Leu Ser Ser Ile Ile Val Val Leu Ser Leu Ala Thr Gln Ser Phe Gly			
375	380	385	390
gtc ctt gcg aat agt gcc agc gcc ttt tat acc atg cgt tcg aac cct			2561
Val Leu Ala Asn Ser Ala Ser Ala Phe Tyr Thr Met Arg Ser Asn Pro			
395	400	405	
agc ggt aaa gag cat aac caa tagaagtgac acccggccag tatcgagatc			2612
Ser Gly Lys Glu His Asn Gln			
410			
gggctgtgac aggtgcatcg ataatcgcaa tcagtcttgt acccatgaga atccctgaaa			2672
aagtaagact gctctgtcag gtagtccatt gcccatgcca taggttcgga cgcctaaagg			2732
atcaatcaag atgccaatca agcatccgac tcatcggaag aaggcatctt gccgacattg			2792
gactcatcct cttcgtccga gtcgtcggcg acaacagcag cttgcttagc gaactccttt			2852
ggcctgcaga gggatcagta gtagccccag gcaccgcgat tgaggggatcc actcaccaaa			2912
acggggcttt gcgccacgct ttctcatcac gaacgatggt gactatttct cccttgagct			2972
tgttgtcgcg gccctcaaga gcattggtat cgatgaccac gtaggaacct cgcttgatcc			3032
agatggtcga tcggaagcgg gcagggagct ccacaaaac cgactccttc gagggaagtt			3092
ccaccgagta gatgttgttt ccggtcgect tgatagcccc ggcaattaga tggccctgag			3152
acagctcatc cggcgggaac atggtctctt ccgccgtagc gagaaccttg cgcctagggg			3212
gacccatttc taacgtatac gcaagtgggt cagcgggaag tagtgtcgat ctaaattggg			3272
atgaacacca aagagactga agaaagagtg agcaaaatgc gagaaaccgt cgactgcgga			3332
gagttttact cgaatgaaga aattcgggcg gcagaaaatc cccgggtggg agtgtgggtc			3392
ccccatccgc ttctttgcca atctcgtctc ctctctctc tcctctccgc catctacgga			3452
gtagagctcc agtactacta tctacttacg gccgtgctca tatccattgg ttgtgaggaa			3512
tgtggcatct tagttggcta cacagtgcac tacacaatcc atcgaatcaa ggtcttctctg			3572
gcgagtcagt tcccctgcac aatgtttggc ctaaggcggga catctagacc acttgacctc			3632
gtgtttttgg gcccttctgc cggcatttgg agtggatgta tcatatcgtg aaacttgcat			3692
gtgtctgatt ggcttccggc ttttatctct ttgaacttcg ctactgggtg gggttcgcgg			3752
agtgatattc cgtcaagct aattcacctc catggatatt aataatgacc ccgacacggt			3812
ggggatcccc aaacctccgt aaaagaatcg tttatcctgg ggtctcggac agatttcgta			3872
ctacaaacct ctcacg			3888

<210> SEQ ID NO 2  
 <211> LENGTH: 413  
 <212> TYPE: PRT  
 <213> ORGANISM: Aspergillus niger

<400> SEQUENCE: 2

Met Asp Trp Met Arg Leu Ile Arg Asp Leu Cys Phe Asn Pro Arg His															
1			5				10						15		
Thr Lys Trp Met Ala Pro Leu Leu Val Leu Gly Asp Ala Phe Leu Cys															
			20				25						30		

-continued

Ala Leu Ile Ile Trp Lys Val Pro Tyr Thr Glu Ile Asp Trp Ala Thr  
 35 40 45  
 Tyr Met Gln Gln Ile Ser Leu Tyr Leu Ser Gly Glu Arg Asp Tyr Thr  
 50 55 60  
 Leu Ile Arg Gly Ser Thr Gly Pro Leu Val Tyr Pro Ala Ala His Val  
 65 70 75 80  
 Tyr Ser Tyr Thr Ala Leu Tyr His Leu Thr Asp Glu Gly Arg Asp Ile  
 85 90 95  
 Phe Phe Gly Gln Ile Leu Phe Ala Val Leu Tyr Leu Ile Thr Leu Val  
 100 105 110  
 Val Val Leu Cys Cys Tyr Arg Gln Ser Gly Ala Pro Pro Tyr Leu Leu  
 115 120 125  
 Pro Leu Leu Val Leu Ser Lys Arg Leu His Ser Val Tyr Val Leu Arg  
 130 135 140  
 Leu Phe Asn Asp Gly Leu Ala Ala Leu Ala Met Trp Val Ala Ile Leu  
 145 150 155 160  
 Leu Phe Met Asn Arg Lys Trp Thr Ala Ala Val Ala Val Trp Ser Thr  
 165 170 175  
 Gly Val Ala Ile Lys Met Thr Leu Leu Leu Ala Pro Ala Ile Ala  
 180 185 190  
 Val Val Thr Val Leu Ser Leu Ser Leu Gly Pro Ser Val Gly Leu Gly  
 195 200 205  
 Val Leu Ala Val Leu Val Gln Val Leu Leu Ala Ile Pro Phe Leu Gln  
 210 215 220  
 Asn Asn Pro Ala Gly Tyr Leu Ser Arg Ala Phe Glu Leu Thr Arg Gln  
 225 230 235 240  
 Phe Met Phe Lys Trp Thr Val Asn Trp Arg Phe Val Gly Glu Glu Val  
 245 250 255  
 Phe Leu Ser Lys Ser Phe Ser Leu Ala Leu Leu Ala Val His Ile Val  
 260 265 270  
 Leu Leu Gly Ala Phe Ala Val Thr Gly Trp Leu Arg Tyr Ser Arg Ser  
 275 280 285  
 Ser Leu Pro Ala Phe Ile Arg Asn Leu Leu Ala Gly Arg His Arg Thr  
 290 295 300  
 Val Ser Leu Pro Lys Pro Tyr Ile Met Ser Val Met Leu Ser Ser Leu  
 305 310 315 320  
 Thr Val Gly Leu Leu Cys Ala Arg Ser Leu His Tyr Gln Phe Phe Ala  
 325 330 335  
 Tyr Leu Ser Trp Ala Thr Pro Phe Leu Leu Trp Arg Ala Gly Phe His  
 340 345 350  
 Pro Ile Leu Leu Tyr Leu Ile Trp Ala Met Gln Glu Trp Ala Trp Asn  
 355 360 365  
 Thr Phe Pro Ser Thr Asn Leu Ser Ser Ile Ile Val Val Leu Ser Leu  
 370 375 380  
 Ala Thr Gln Ser Phe Gly Val Leu Ala Asn Ser Ala Ser Ala Phe Tyr  
 385 390 395 400  
 Thr Met Arg Ser Asn Pro Ser Gly Lys Glu His Asn Gln  
 405 410

<210> SEQ ID NO 3  
 <211> LENGTH: 1923  
 <212> TYPE: DNA  
 <213> ORGANISM: Aspergillus oryzae  
 <220> FEATURE:

-continued

---

<221> NAME/KEY: CDS  
 <222> LOCATION: (1)..(1242)

<400> SEQUENCE: 3

atg gag ttg aag cac ttc atc cac gaa ctc tgc cta aac ccc aga cat	48
Met Glu Leu Lys His Phe Ile His Glu Leu Cys Leu Asn Pro Arg His	
1 5 10 15	
aca aaa tgg att gca ccg ctt ctt gtc ata ggc gat gcc ttc cta tgt	96
Thr Lys Trp Ile Ala Pro Leu Leu Val Ile Gly Asp Ala Phe Leu Cys	
20 25 30	
gct ctt atc atc tgg aag atc cca tat act gag atc gac tgg acg acg	144
Ala Leu Ile Ile Trp Lys Ile Pro Tyr Thr Glu Ile Asp Trp Thr Thr	
35 40 45	
tac atg cag cag ata gcg ctt tac atc tct ggc gaa cgt gac tat acc	192
Tyr Met Gln Gln Ile Ala Leu Tyr Ile Ser Gly Glu Arg Asp Tyr Thr	
50 55 60	
ctg atc aag ggg tcc act gga ccc ctt gta tac ccg gcc gcc cac gta	240
Leu Ile Lys Gly Ser Thr Gly Pro Leu Val Tyr Pro Ala Ala His Val	
65 70 75 80	
tat agc tac atg gca ctc tat cac cta aca gat gaa ggc cga gat att	288
Tyr Ser Tyr Met Ala Leu Tyr His Leu Thr Asp Glu Gly Arg Asp Ile	
85 90 95	
ctt ttc ggg cag ata tta ttt gct gtc ctt tac ctt gtc acg cta gca	336
Leu Phe Gly Gln Ile Leu Phe Ala Val Leu Tyr Leu Val Thr Leu Ala	
100 105 110	
gtt gtg atg gtt tgc tac agg cag tca ggt gcc cct ccg tac ctg ttt	384
Val Val Met Val Cys Tyr Arg Gln Ser Gly Ala Pro Pro Tyr Leu Phe	
115 120 125	
cct ctt ctt gtc ctt tcc aag cgg ctc cac agt gtc ttt gtt ttg cgc	432
Pro Leu Leu Val Leu Ser Lys Arg Leu His Ser Val Phe Val Leu Arg	
130 135 140	
ctt ttc aat gat ggc ctc gcg gtc tgt gcc atg tgg ata gcg att ctg	480
Leu Phe Asn Asp Gly Leu Ala Val Cys Ala Met Trp Ile Ala Ile Leu	
145 150 155 160	
ctc ttc cag aat aag aaa tgg acg gct ggt gtt acg gcc tgg act gtt	528
Leu Phe Gln Asn Lys Lys Trp Thr Ala Gly Val Thr Ala Trp Thr Val	
165 170 175	
ggg gtt ggc att aag atg acg tta ctg ctc ctt gcg cca gcc att gcg	576
Gly Val Gly Ile Lys Met Thr Leu Leu Leu Leu Ala Pro Ala Ile Ala	
180 185 190	
gtg gta act gtg ctc agc ctt tcc ctc gtg cct agt att cga ctt gga	624
Val Val Thr Val Leu Ser Leu Ser Leu Val Pro Ser Ile Arg Leu Gly	
195 200 205	
att cta gca ttg ctc att cag gtt cta cta gcg att ccc ttc cta cag	672
Ile Leu Ala Leu Leu Ile Gln Val Leu Leu Ala Ile Pro Phe Leu Gln	
210 215 220	
ggt aac ccc ata gga tac gtc gcg cgg gcc ttt gag ttg act aga cag	720
Gly Asn Pro Ile Gly Tyr Val Ala Arg Ala Phe Glu Leu Thr Arg Gln	
225 230 235 240	
ttt atg ttc aaa tgg act gtc aat tgg agg ttt gtg ggt gaa gac ttg	768
Phe Met Phe Lys Trp Thr Val Asn Trp Arg Phe Val Gly Glu Asp Leu	
245 250 255	
ttc cta tcc aaa cag ttt tct cta gcc tta cta ggt ttg cat att ttt	816
Phe Leu Ser Lys Gln Phe Ser Leu Ala Leu Leu Gly Leu His Ile Phe	
260 265 270	
ctg ctg gga tta ttt gtt acc aca ggc tgg tta cgg ccg tca gga tct	864
Leu Leu Gly Leu Phe Val Thr Thr Gly Trp Leu Arg Pro Ser Gly Ser	
275 280 285	
aac gtc cct gac ttc ctc cgg agc cta ctc caa gga cgc caa cgc acc	912
Asn Val Pro Asp Phe Leu Arg Ser Leu Leu Gln Gly Arg Gln Arg Thr	

-continued

290	295	300	
gtg gtg ctt tct aag tct ttc ata atg acc gtg atg ttg aca tcg ctg			960
Val Val Leu Ser Lys Ser Phe Ile Met Thr Val Met Leu Thr Ser Leu			
305	310	315	320
gcg atc ggg ttg ttg tgc gca agg tcc ctt cat tac caa ttc ttt gcc			1008
Ala Ile Gly Leu Leu Cys Ala Arg Ser Leu His Tyr Gln Phe Phe Ala			
	325	330	335
tat ctc tcc tgg gct acg cct tgc ctt ctc tgg cgg gct cgg ctc cat			1056
Tyr Leu Ser Trp Ala Thr Pro Cys Leu Leu Trp Arg Ala Arg Leu His			
	340	345	350
ccg atc ctt ata tat gcg atc tgg gca cta cag gag tgg gct tgg aat			1104
Pro Ile Leu Ile Tyr Ala Ile Trp Ala Leu Gln Glu Trp Ala Trp Asn			
	355	360	365
gtc tac cca agc acc aat gcc agt tct tcg gtc gtt gtc ttc tca ctt			1152
Val Tyr Pro Ser Thr Asn Ala Ser Ser Ser Val Val Val Phe Ser Leu			
	370	375	380
gct gtt cag gtt ttc ggt gtc ctc ctc aat agc aga aac gca ctg agc			1200
Ala Val Gln Val Phe Gly Val Leu Leu Asn Ser Arg Asn Ala Leu Ser			
	385	390	395
gat gcg cct ccg aga cgc aaa gga aag gag cac atc cag tga			1242
Asp Ala Pro Pro Arg Arg Lys Gly Lys Glu His Ile Gln			
	405	410	
taatgagcca ctcataggct ccaaatttat cgttcccttc attgttttca tcaattggtc			1302
ctcgtgggttc acattaaatg tagacgcacg tgcattggat acagcgatca ccatacgtcg			1362
attagttcaa gcatcagagt cgcccgaggg aggcattcttg cccacatttg actcttctc			1422
atcctctgaa tccgaagcaa caaccgtaga ttgcttgacg aattccttcg gcctaattgga			1482
gccttgccgc aaactttctc atcccgaacg atattaatga ttccccagc aagcttattg			1542
tcccggctct caagagcgtt ggtatcgacc acgacgtagg agttgcgttt catccagatt			1602
cttgaacgga agcgggaagg tagttcgacc aaaaccgtgt cttttgatgg cagttcgacc			1662
acataataat tgttccccgt tgctttgatg actcgcgcga tttgctgtcc ctggtgaagc			1722
tcatctggag gggcatagt ttcttccgct gttgcgagta ccttgcgcgg tgggtgggcc			1782
atgatggat attgcaattg ggatctcttt tcttggaaacc taaattgtcg caatctctta			1842
gatgctgtct tgtatgaagc aattgctgat cagataagct gtggtgaggg gtagtcttat			1902
tgtgtttgat agcggagggg t			1923
<p>&lt;210&gt; SEQ ID NO 4                  &lt;211&gt; LENGTH: 413                  &lt;212&gt; TYPE: PRT                  &lt;213&gt; ORGANISM: Aspergillus oryzae</p>			
<p>&lt;400&gt; SEQUENCE: 4</p>			
Met Glu Leu Lys His Phe Ile His Glu Leu Cys Leu Asn Pro Arg His			
1	5	10	15
Thr Lys Trp Ile Ala Pro Leu Leu Val Ile Gly Asp Ala Phe Leu Cys			
	20	25	30
Ala Leu Ile Ile Trp Lys Ile Pro Tyr Thr Glu Ile Asp Trp Thr Thr			
	35	40	45
Tyr Met Gln Gln Ile Ala Leu Tyr Ile Ser Gly Glu Arg Asp Tyr Thr			
	50	55	60
Leu Ile Lys Gly Ser Thr Gly Pro Leu Val Tyr Pro Ala Ala His Val			
	65	70	75
Tyr Ser Tyr Met Ala Leu Tyr His Leu Thr Asp Glu Gly Arg Asp Ile			
	85	90	95

-continued

Leu Phe Gly Gln Ile Leu Phe Ala Val Leu Tyr Leu Val Thr Leu Ala  
 100 105 110  
 Val Val Met Val Cys Tyr Arg Gln Ser Gly Ala Pro Pro Tyr Leu Phe  
 115 120 125  
 Pro Leu Leu Val Leu Ser Lys Arg Leu His Ser Val Phe Val Leu Arg  
 130 135 140  
 Leu Phe Asn Asp Gly Leu Ala Val Cys Ala Met Trp Ile Ala Ile Leu  
 145 150 155 160  
 Leu Phe Gln Asn Lys Lys Trp Thr Ala Gly Val Thr Ala Trp Thr Val  
 165 170 175  
 Gly Val Gly Ile Lys Met Thr Leu Leu Leu Ala Pro Ala Ile Ala  
 180 185 190  
 Val Val Thr Val Leu Ser Leu Ser Leu Val Pro Ser Ile Arg Leu Gly  
 195 200 205  
 Ile Leu Ala Leu Leu Ile Gln Val Leu Leu Ala Ile Pro Phe Leu Gln  
 210 215 220  
 Gly Asn Pro Ile Gly Tyr Val Ala Arg Ala Phe Glu Leu Thr Arg Gln  
 225 230 235 240  
 Phe Met Phe Lys Trp Thr Val Asn Trp Arg Phe Val Gly Glu Asp Leu  
 245 250 255  
 Phe Leu Ser Lys Gln Phe Ser Leu Ala Leu Leu Gly Leu His Ile Phe  
 260 265 270  
 Leu Leu Gly Leu Phe Val Thr Thr Gly Trp Leu Arg Pro Ser Gly Ser  
 275 280 285  
 Asn Val Pro Asp Phe Leu Arg Ser Leu Leu Gln Gly Arg Gln Arg Thr  
 290 295 300  
 Val Val Leu Ser Lys Ser Phe Ile Met Thr Val Met Leu Thr Ser Leu  
 305 310 315 320  
 Ala Ile Gly Leu Leu Cys Ala Arg Ser Leu His Tyr Gln Phe Phe Ala  
 325 330 335  
 Tyr Leu Ser Trp Ala Thr Pro Cys Leu Leu Trp Arg Ala Arg Leu His  
 340 345 350  
 Pro Ile Leu Ile Tyr Ala Ile Trp Ala Leu Gln Glu Trp Ala Trp Asn  
 355 360 365  
 Val Tyr Pro Ser Thr Asn Ala Ser Ser Ser Val Val Val Phe Ser Leu  
 370 375 380  
 Ala Val Gln Val Phe Gly Val Leu Leu Asn Ser Arg Asn Ala Leu Ser  
 385 390 395 400  
 Asp Ala Pro Pro Arg Arg Lys Gly Lys Glu His Ile Gln  
 405 410

<210> SEQ ID NO 5  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Alg3-ForScr primer

<400> SEQUENCE: 5

cggtttccct tcagtttcca gt

22

<210> SEQ ID NO 6  
 <211> LENGTH: 49  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:

-continued

---

<223> OTHER INFORMATION: Alg3-1 primer

<400> SEQUENCE: 6

gtaacgccag ggttttccca gtcacgacgt cataacttct ctcccctcc 49

<210> SEQ ID NO 7  
 <211> LENGTH: 49  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Alg3-2 primer

<400> SEQUENCE: 7

atccacttaa cgttactgaa atctccaact tcatggacac acacagacc 49

<210> SEQ ID NO 8  
 <211> LENGTH: 49  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Hph-F primer

<400> SEQUENCE: 8

ggtctgtgtg tgtccatgaa gttggagatt tcagtaacgt taagtggat 49

<210> SEQ ID NO 9  
 <211> LENGTH: 49  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Hph-R primer

<400> SEQUENCE: 9

gctactactg atccctctgc gtcggagaca gaagatgata ttgaaggag 49

<210> SEQ ID NO 10  
 <211> LENGTH: 49  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Alg3-3 primer

<400> SEQUENCE: 10

ctccttcaat atcatcttct gtctccgacg cagagggatc agtagtagc 49

<210> SEQ ID NO 11  
 <211> LENGTH: 49  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Alg3-4 primer

<400> SEQUENCE: 11

gcgataaca atttcacaca ggaaacagcc gtgagagggt ttagtagc 49

<210> SEQ ID NO 12  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Alg3-RevScr primer

<400> SEQUENCE: 12

aagctgagag cgacatcttc a 21

-continued

---

<210> SEQ ID NO 13  
 <211> LENGTH: 25  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: hyg-RevScr primer  
  
 <400> SEQUENCE: 13  
  
 gtacttctac acagccatcg gtcca 25

<210> SEQ ID NO 14  
 <211> LENGTH: 25  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: hyg-ForScr primer  
  
 <400> SEQUENCE: 14  
  
 gtacttctac acagccatcg gtcca 25

<210> SEQ ID NO 15  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: pryGScr primer  
  
 <400> SEQUENCE: 15  
  
 tctgctgtct tgcattgaggt cctt 24

<210> SEQ ID NO 16  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: primer  
  
 <400> SEQUENCE: 16  
  
 agcgtaggac aaggtcgtct ctgt 24

<210> SEQ ID NO 17  
 <211> LENGTH: 58  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: 5-pyrG5F primer  
  
 <400> SEQUENCE: 17  
  
 gtaacgccag ggttttccca gtcacgacgt ttaaacaatgc atcattctcc cgttttgt 58

<210> SEQ ID NO 18  
 <211> LENGTH: 43  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: 5-pyrG3R primer  
  
 <400> SEQUENCE: 18  
  
 agaaagagtc accggtcacg acatcgccaa tcacctcaat cac 43

<210> SEQ ID NO 19  
 <211> LENGTH: 43  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: ble5F primer



-continued

---

<400> SEQUENCE: 19  
 gtgattgagg tgattggcga tgtcgtgacc ggtgactctt tct 43

<210> SEQ ID NO 20  
 <211> LENGTH: 42  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Ble3R primer

<400> SEQUENCE: 20  
 tccaaccttg tagcaaccaa agcttcgagc gtcccaaac ct 42

<210> SEQ ID NO 21  
 <211> LENGTH: 42  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Alg3-5F1 primer

<400> SEQUENCE: 21  
 aggttttggg acgctcgaag ctttggttgc tacaaggttg ga 42

<210> SEQ ID NO 22  
 <211> LENGTH: 29  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Alg3-3R1 primer

<400> SEQUENCE: 22  
 tcaagtagag cacagcaaat agtatctga 29

<210> SEQ ID NO 23  
 <211> LENGTH: 29  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Alg3-5F2 primer

<400> SEQUENCE: 23  
 tcagatacta ttgctgtgc tctacttga 29

<210> SEQ ID NO 24  
 <211> LENGTH: 42  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Alg3-3R2 primer

<400> SEQUENCE: 24  
 ttgatccttg tgccacacca tctacgtgg tcatcgatac ca 42

<210> SEQ ID NO 25  
 <211> LENGTH: 42  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: 3-pyrG5F primer

<400> SEQUENCE: 25  
 tggatcgat gaccagtag gatggtgtgg cacaaggatc aa 42

<210> SEQ ID NO 26  
 <211> LENGTH: 60

-continued

---

<212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: 3=pyrG3R primer  
  
 <400> SEQUENCE: 26  
  
 gcggataaca atttcacaca ggaaacagcg tttaaactgt gccagtcaat tgcccgaagt 60

<210> SEQ ID NO 27  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Alg3seq-1 primer  
  
 <400> SEQUENCE: 27  
  
 tacagacgcg tgtacgcatg t 21

<210> SEQ ID NO 28  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Alg3seq-2 primer  
  
 <400> SEQUENCE: 28  
  
 tgctattgtc cacagatacc gaga 24

<210> SEQ ID NO 29  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Alg3seq-3 primer  
  
 <400> SEQUENCE: 29  
  
 gagctaacca gacagttcat gt 22

<210> SEQ ID NO 30  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Alg3seq-4 primer  
  
 <400> SEQUENCE: 30  
  
 tcgtcgtagc gcattgatcc t 21

<210> SEQ ID NO 31  
 <211> LENGTH: 413  
 <212> TYPE: PRT  
 <213> ORGANISM: Aspergillus nidulans  
  
 <400> SEQUENCE: 31  
  
 Met Ala Leu Thr Asp Leu Val Ser Gly Leu Cys Ser Asn Pro Lys His  
 1 5 10 15  
  
 Thr Lys Trp Ile Ala Pro Ile Leu Asn Ile Ala Asp Gly Leu Leu Cys  
 20 25 30  
  
 Ala Phe Ile Ile Trp Lys Val Pro Tyr Thr Glu Ile Asp Trp Thr Thr  
 35 40 45  
  
 Tyr Met Gln Gln Val Lys Leu Tyr Leu Ser Gly Glu Arg Asp Tyr Thr  
 50 55 60  
  
 Leu Ile Lys Gly Ser Thr Gly Pro Leu Val Tyr Pro Ala Ala His Val  
 65 70 75 80

-continued

Tyr Ser Tyr Ser Leu Phe His His Leu Thr Asp Glu Gly Arg Asp Ile  
 85 90 95  
 Val Phe Gly Gln Ile Ile Phe Ala Phe Leu Tyr Leu Ile Cys Leu Thr  
 100 105 110  
 Val Val Met Ala Cys Tyr Arg Arg Val Gly Ala Pro Pro Tyr Leu Phe  
 115 120 125  
 Pro Leu Leu Val Leu Ser Lys Arg Leu His Ser Val Tyr Met Leu Arg  
 130 135 140  
 Leu Phe Asn Asp Gly Leu Ala Ala Leu Ala Met Trp Gly Ser Ile Trp  
 145 150 155 160  
 Leu Phe Ile Asn Arg Lys Trp Thr Pro Ala Val Val Leu Trp Ser Leu  
 165 170 175  
 Gly Leu Gly Val Lys Met Thr Leu Ile Leu Leu Val Pro Ala Val Met  
 180 185 190  
 Val Val Leu Ala Leu Ser Leu Asp Ile Gly Arg Cys Ile Arg Leu Ala  
 195 200 205  
 Gly Leu Ala Leu Gly Ile Gln Ile Leu Leu Ala Ile Pro Phe Leu Lys  
 210 215 220  
 Thr Asn Pro Ser Gly Tyr Phe Glu Arg Ala Phe Glu Phe Gly Arg Gln  
 225 230 235 240  
 Phe Met Phe Lys Trp Thr Val Asn Trp Arg Phe Val Gly Glu Asp Ile  
 245 250 255  
 Phe Leu Ser Lys Gly Phe Trp Ala Gly Leu Ile Val Leu His Leu Leu  
 260 265 270  
 Ile Leu Val Val Leu Gly Phe Thr Cys Phe Leu Asn Pro Ser Gly Thr  
 275 280 285  
 Ser Leu Pro Asp Phe Ala Gly Arg Phe Leu Thr Gly Gln His Arg Gly  
 290 295 300  
 Ile Ala Leu His Pro Ser Phe Ile Met Ser Ala Leu Leu Thr Ser Leu  
 305 310 315 320  
 Ser Val Gly Leu Leu Cys Ala Arg Ser Leu His Tyr Gln Phe Phe Ala  
 325 330 335  
 Tyr Leu Ser Trp Ala Thr Pro Phe Leu Leu Trp Gln Ala Gly Tyr His  
 340 345 350  
 Pro Ile Leu Val Tyr Ala Leu Trp Leu Val Gln Glu Trp Ala Trp Asn  
 355 360 365  
 Val Tyr Pro Ser Thr Asn Leu Ser Ser Ala Ala Val Val Leu Leu Leu  
 370 375 380  
 Gly Ala Gln Val Leu Gly Val Leu Val Asn Arg Asp Arg Ala Phe Pro  
 385 390 395 400  
 Ser Ser Pro Pro Thr Pro Lys Ala Lys Gln His Val Gln  
 405 410

<210> SEQ ID NO 32  
 <211> LENGTH: 434  
 <212> TYPE: PRT  
 <213> ORGANISM: Fusarium oxysporum

<400> SEQUENCE: 32

Met Pro Glu Ser Ala Ser Gly Thr Leu Ser Gln Gly Val Arg Phe Leu  
 1 5 10 15  
 Arg Asn Val Leu Asn Gly Arg His Ala Leu Ser Lys Leu Ile Pro Ile  
 20 25 30  
 Ala Leu Trp Leu Val Asp Ala Leu Gly Cys Gly Leu Ile Ile Trp Lys

-continued

35					40					45					
Ile	Pro	Tyr	Thr	Glu	Ile	Asp	Trp	Val	Ala	Tyr	Met	Gln	Gln	Ile	Ser
	50					55					60				
Gln	Phe	Val	Ser	Gly	Glu	Arg	Asp	Tyr	Thr	Lys	Met	Glu	Gly	Asp	Thr
	65					70					75				80
Gly	Pro	Leu	Val	Tyr	Pro	Ala	Ala	His	Val	Tyr	Thr	Tyr	Thr	Gly	Leu
				85					90					95	
Tyr	Tyr	Ile	Thr	Asp	Lys	Gly	Thr	Asn	Ile	Leu	Leu	Ala	Gln	Gln	Ile
			100					105					110		
Phe	Ala	Val	Leu	Tyr	Met	Ala	Thr	Leu	Ala	Val	Val	Met	Leu	Cys	Tyr
		115					120					125			
Trp	Lys	Ala	Lys	Val	Pro	Pro	Tyr	Met	Phe	Ile	Phe	Leu	Ile	Ala	Ser
	130					135					140				
Lys	Arg	Leu	His	Ser	Leu	Phe	Val	Leu	Arg	Cys	Phe	Asn	Asp	Cys	Phe
	145					150					155				160
Ala	Val	Phe	Phe	Leu	Trp	Leu	Thr	Ile	Phe	Leu	Phe	Gln	Arg	Arg	Gln
				165					170					175	
Trp	Thr	Val	Gly	Ser	Leu	Val	Tyr	Ser	Trp	Gly	Leu	Gly	Ile	Lys	Met
			180					185					190		
Ser	Leu	Leu	Leu	Val	Leu	Pro	Ala	Ile	Gly	Val	Ile	Leu	Phe	Leu	Gly
		195					200					205			
Arg	Gly	Leu	Trp	Pro	Ser	Leu	Arg	Leu	Ala	Trp	Leu	Met	Ala	Gln	Ile
	210					215					220				
Gln	Phe	Ala	Ile	Gly	Leu	Pro	Phe	Ile	Thr	Lys	Asn	Pro	Arg	Gly	Tyr
	225					230					235				240
Ala	Ala	Arg	Ala	Phe	Glu	Leu	Ser	Arg	Gln	Phe	Gln	Phe	Lys	Trp	Thr
				245					250					255	
Val	Asn	Trp	Arg	Met	Leu	Gly	Glu	Glu	Val	Phe	Leu	Ser	Lys	Tyr	Phe
			260					265					270		
Ala	Leu	Ser	Leu	Leu	Ala	Cys	His	Ile	Leu	Val	Leu	Leu	Ile	Phe	Ile
		275					280					285			
Ser	Lys	Arg	Trp	Ile	Gln	Pro	Thr	Gly	Arg	Ser	Leu	Tyr	Asp	Leu	Ile
	290					295					300				
Pro	Ser	Phe	Leu	Arg	Leu	Lys	Ser	Pro	Phe	Thr	Met	Gln	Glu	Gln	Leu
	305					310					315				320
Arg	Ile	Ser	His	Tyr	Val	Thr	Pro	Glu	Tyr	Ala	Met	Thr	Thr	Met	Leu
			325						330					335	
Thr	Ala	Asn	Leu	Ile	Gly	Leu	Leu	Phe	Ala	Arg	Ser	Leu	His	Tyr	Gln
			340					345					350		
Phe	Tyr	Ala	Tyr	Leu	Ala	Trp	Ala	Thr	Pro	Tyr	Leu	Leu	Trp	Arg	Ala
		355					360					365			
Thr	Glu	Asp	Pro	Val	Ile	Val	Ala	Ile	Ile	Trp	Ala	Ala	Gln	Glu	Trp
	370					375					380				
Ala	Trp	Asn	Val	Tyr	Pro	Ser	Thr	Asp	Leu	Ser	Ser	Thr	Ile	Ala	Val
	385					390					395				400
Asn	Thr	Met	Leu	Ala	Thr	Val	Val	Leu	Val	Tyr	Leu	Gly	Thr	Ala	Arg
			405						410					415	
Arg	Ala	Val	Pro	Ala	Pro	Ala	Ala	Gln	Val	Gly	Asn	Val	Asp	Asp	Lys
		420					425						430		
Asn	Lys														

&lt;210&gt; SEQ ID NO 33

&lt;211&gt; LENGTH: 502

-continued

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Neurospora crassa

&lt;400&gt; SEQUENCE: 33

Met Ala Ala Pro Ser Ser Arg Pro Glu Ser Asn Pro Pro Leu Tyr Lys  
 1 5 10 15  
 Gln Ala Leu Asp Phe Ala Leu Asp Val Ala Asn Gly Arg His Ala Leu  
 20 25 30  
 Ser Lys Leu Ile Pro Pro Ala Leu Phe Leu Val Asp Ala Leu Leu Cys  
 35 40 45  
 Gly Leu Ile Ile Trp Lys Val Pro Tyr Thr Glu Ile Asp Trp Ala Ala  
 50 55 60  
 Tyr Met Glu Gln Val Ser Gln Ile Leu Ser Gly Glu Arg Asp Tyr Thr  
 65 70 75 80  
 Lys Val Arg Gly Gly Thr Gly Pro Leu Val Tyr Pro Ala Ala His Val  
 85 90 95  
 Tyr Ile Tyr Thr Gly Leu Tyr His Leu Thr Asp Glu Gly Arg Asn Ile  
 100 105 110  
 Leu Leu Ala Gln Gln Leu Phe Ala Gly Leu Tyr Met Val Thr Leu Ala  
 115 120 125  
 Val Val Met Gly Cys Tyr Trp Gln Ala Lys Ala Pro Pro Tyr Leu Phe  
 130 135 140  
 Pro Leu Leu Thr Leu Ser Lys Arg Leu His Ser Ile Phe Val Leu Arg  
 145 150 155 160  
 Cys Phe Asn Asp Cys Phe Ala Val Leu Phe Leu Trp Leu Ala Ile Phe  
 165 170 175  
 Phe Phe Gln Arg Arg Asn Trp Gln Ala Gly Ala Leu Leu Tyr Thr Leu  
 180 185 190  
 Gly Leu Gly Val Lys Met Thr Leu Leu Ser Leu Pro Ala Val Gly  
 195 200 205  
 Ile Val Leu Phe Leu Gly Ser Gly Ser Phe Val Thr Thr Leu Gln Leu  
 210 215 220  
 Val Ala Thr Met Gly Leu Val Gln Ile Ala Ile Gly Leu Pro Phe Ile  
 225 230 235 240  
 Thr Lys Asn Pro Arg Gly Tyr Ala Ala Arg Ala Phe Glu Leu Ser Arg  
 245 250 255  
 Gln Phe Gln Phe Lys Trp Thr Val Asn Trp Arg Met Leu Gly Glu Glu  
 260 265 270  
 Val Phe Leu Ser Lys Tyr Phe Ala Leu Ser Leu Leu Ala Cys His Ile  
 275 280 285  
 Leu Val Leu Leu Ile Leu Ile Gly Val Pro Phe Leu Ala His Tyr Pro  
 290 295 300  
 Thr Glu Tyr Leu Ser Arg Ala Phe Glu Leu Ser Arg Gln Phe Phe Phe  
 305 310 315 320  
 Lys Trp Thr Val Asn Trp Arg Phe Val Gly Glu Glu Ile Phe Leu Ser  
 325 330 335  
 Lys Gly Phe Ala Leu Thr Leu Leu Ala Leu His Val Leu Val Leu Gly  
 340 345 350  
 Ile Phe Ile Thr Thr Arg Trp Ile Lys Pro Ala Arg Lys Ser Leu Val  
 355 360 365  
 Gln Leu Ile Ser Pro Val Leu Leu Ala Gly Lys Pro Pro Leu Thr Val  
 370 375 380  
 Pro Glu His Arg Ala Ala Ala Arg Asp Val Thr Pro Arg Tyr Ile Met  
 385 390 395 400

-continued

Thr Thr Ile Leu Ser Ala Asn Ala Val Gly Leu Leu Phe Ala Arg Ser  
 405 410 415  
 Leu His Tyr Gln Phe Tyr Ala Tyr Val Ala Trp Ser Thr Pro Phe Leu  
 420 425 430  
 Leu Trp Arg Ala Gly Leu His Pro Val Leu Val Tyr Leu Leu Trp Ala  
 435 440 445  
 Val His Glu Trp Ala Trp Asn Val Phe Pro Ser Thr Pro Ala Ser Ser  
 450 455 460  
 Ala Val Val Val Gly Val Leu Gly Val Thr Val Ala Gly Val Trp Phe  
 465 470 475 480  
 Gly Ala Arg Glu Glu Trp Glu Pro Gly Met Lys Ser Ser Ser Lys Lys  
 485 490 495  
 Glu Glu Ala Ala Met Arg  
 500

<210> SEQ ID NO 34  
 <211> LENGTH: 434  
 <212> TYPE: PRT  
 <213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 34

Met Ala Gly Gly Lys Lys Lys Ser Ser Thr Ala Pro Ser Arg Phe Gln  
 1 5 10 15  
 Lys Thr Leu Ser Ser Ile Trp Gln Asp Lys His Thr Val Leu Phe Lys  
 20 25 30  
 Pro Glu Tyr Thr Leu Leu Val Thr Ala Val Leu Trp Phe Leu Glu Ile  
 35 40 45  
 Ala Ile Asn Ile Trp Val Ile Gln Lys Val Ser Tyr Thr Glu Ile Asp  
 50 55 60  
 Trp Lys Ala Tyr Met Asp Glu Val Glu Gly Val Ile Asn Gly Thr Tyr  
 65 70 75 80  
 Asp Tyr Thr Gln Leu Lys Gly Asp Thr Gly Pro Leu Val Tyr Pro Ala  
 85 90 95  
 Gly Phe Val Tyr Ile Phe Thr Gly Leu Tyr Tyr Leu Thr Asp His Gly  
 100 105 110  
 His Asn Ile Arg Leu Gly Gln Tyr Val Phe Ala Val Ser Tyr Leu Ile  
 115 120 125  
 Asn Leu Leu Leu Val Met Arg Ile Tyr His Arg Thr Lys Lys Val Pro  
 130 135 140  
 Pro Tyr Val Phe Phe Phe Ile Cys Cys Ala Ser Tyr Arg Ile His Ser  
 145 150 155 160  
 Ile Phe Ile Leu Arg Leu Phe Asn Asp Pro Val Ala Met Met Leu Cys  
 165 170 175  
 Phe Gly Ala Ile Asn Leu Phe Leu Asp Gly Arg Trp Thr Leu Gly Cys  
 180 185 190  
 Ala Leu Tyr Ser Leu Ala Val Ser Val Lys Met Asn Val Leu Leu Phe  
 195 200 205  
 Ala Pro Gly Leu Leu Phe Leu Leu Leu Cys Glu Phe Gly Leu Trp Lys  
 210 215 220  
 Thr Leu Pro Arg Leu Ala Leu Cys Ala Val Ile Gln Leu Leu Val Gly  
 225 230 235 240  
 Leu Pro Phe Leu Ile Thr Tyr Pro Val Ser Tyr Ile Ala Asn Ala Phe  
 245 250 255  
 Asp Leu Gly Arg Val Phe Ile His Phe Trp Ser Val Asn Phe Lys Phe

-continued

260					265					270					
Val	Pro	Glu	Arg	Val	Phe	Val	Ser	Lys	Glu	Phe	Ala	Val	Cys	Leu	Leu
		275					280					285			
Ile	Ala	His	Leu	Phe	Leu	Leu	Val	Ala	Phe	Ala	Leu	Lys	Arg	Trp	Lys
		290					295					300			
Arg	Ser	Gly	Ser	Ser	Ile	Trp	Thr	Ile	Leu	Lys	Asp	Pro	Ser	Glu	Arg
		305					310					315			
Lys	Glu	Thr	Ala	His	Lys	Val	Asn	Ala	Asp	Gln	Met	Val	Leu	Ile	Leu
				325					330					335	
Phe	Thr	Ser	Asn	Phe	Ile	Gly	Met	Cys	Phe	Ser	Arg	Ser	Leu	His	Tyr
			340					345					350		
Gln	Phe	Tyr	Val	Trp	Tyr	Phe	His	Thr	Leu	Pro	Tyr	Leu	Leu	Trp	Ser
		355					360					365			
Gly	Gly	Val	Lys	Lys	Leu	Ala	Arg	Leu	Leu	Arg	Val	Leu	Ile	Leu	Gly
		370					375					380			
Leu	Ile	Glu	Leu	Ser	Trp	Asn	Thr	Tyr	Pro	Ser	Thr	Asn	Tyr	Ser	Ser
		385					390					395			
Leu	Ser	Leu	His	Val	Cys	His	Leu	Ile	Ile	Leu	Leu	Cys	Leu	Trp	Leu
			405						410					415	
Asn	Pro	Asn	Pro	Ala	Ser	Pro	Ser	His	Arg	Ser	Glu	Asn	Lys	Ala	Lys
			420					425					430		

Ser His

&lt;210&gt; SEQ ID NO 35

&lt;211&gt; LENGTH: 437

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Arabidopsis thaliana

&lt;400&gt; SEQUENCE: 35

Met	Ala	Gly	Ala	Ser	Ser	Pro	Ala	Ser	Leu	Arg	Ala	Ser	Arg	Ser	Arg
				5					10					15	
Arg	Leu	Gly	Lys	Glu	Thr	Asn	Arg	Ser	Asp	Leu	Phe	Lys	Lys	Pro	Ala
			20					25					30		
Val	Pro	Phe	Ala	Phe	Ala	Leu	Ile	Leu	Ala	Asp	Ala	Ile	Leu	Val	Ala
		35					40					45			
Leu	Ile	Ile	Ala	Tyr	Val	Pro	Tyr	Thr	Lys	Ile	Asp	Trp	Asp	Ala	Tyr
		50					55					60			
Met	Ser	Gln	Val	Ser	Gly	Phe	Leu	Gly	Gly	Glu	Arg	Asp	Tyr	Gly	Asn
				70								75			80
Leu	Lys	Gly	Asp	Thr	Gly	Pro	Leu	Val	Tyr	Pro	Ala	Gly	Phe	Leu	Tyr
				85					90					95	
Val	Tyr	Ser	Ala	Val	Gln	Asn	Leu	Thr	Gly	Gly	Glu	Val	Tyr	Pro	Ala
			100					105					110		
Gln	Ile	Leu	Phe	Gly	Val	Leu	Tyr	Ile	Val	Asn	Leu	Gly	Ile	Val	Leu
		115					120					125			
Ile	Ile	Tyr	Val	Lys	Thr	Asp	Val	Pro	Trp	Trp	Ala	Leu	Ser	Leu	Leu
		130					135					140			
Cys	Leu	Ser	Lys	Arg	Ile	His	Ser	Ile	Phe	Val	Leu	Arg	Leu	Phe	Asn
				150					155					160	
Asp	Cys	Phe	Ala	Met	Thr	Leu	Leu	His	Ala	Ser	Met	Ala	Leu	Phe	Leu
				165					170					175	
Tyr	Arg	Lys	Trp	His	Leu	Gly	Met	Leu	Val	Phe	Ser	Gly	Ala	Val	Ser
			180					185					190		
Val	Lys	Met	Asn	Val	Leu	Leu	Tyr	Ala	Pro	Thr	Leu	Leu	Leu	Leu	Leu





-continued

Ala	Thr	Leu	Leu	Leu	Val	Phe	Leu	Ile	Tyr	His	Gln	Thr	Cys	Lys	Val
130						135					140				
Pro	Pro	Phe	Val	Phe	Phe	Phe	Met	Cys	Cys	Ala	Ser	Tyr	Arg	Val	His
145				150						155					160
Ser	Ile	Phe	Val	Leu	Arg	Leu	Phe	Asn	Asp	Pro	Val	Ala	Met	Val	Leu
				165					170					175	
Leu	Phe	Leu	Ser	Ile	Asn	Leu	Leu	Leu	Ala	Gln	Arg	Trp	Gly	Trp	Gly
			180					185					190		
Cys	Cys	Phe	Phe	Ser	Leu	Ala	Val	Ser	Val	Lys	Met	Asn	Val	Leu	Leu
		195					200					205			
Phe	Ala	Pro	Gly	Leu	Leu	Phe	Leu	Leu	Leu	Thr	Gln	Phe	Gly	Phe	Arg
210						215					220				
Gly	Ala	Leu	Pro	Lys	Leu	Gly	Ile	Cys	Ala	Gly	Leu	Gln	Val	Val	Leu
225					230					235					240
Gly	Leu	Pro	Phe	Leu	Leu	Glu	Asn	Pro	Ser	Gly	Tyr	Leu	Ser	Arg	Ser
				245					250					255	
Phe	Asp	Leu	Gly	Arg	Gln	Phe	Leu	Phe	His	Trp	Thr	Val	Asn	Trp	Arg
			260					265					270		
Phe	Leu	Pro	Glu	Ala	Leu	Phe	Leu	His	Arg	Ala	Phe	His	Leu	Ala	Leu
		275					280					285			
Leu	Thr	Ala	His	Leu	Thr	Leu	Leu	Leu	Leu	Phe	Ala	Leu	Cys	Arg	Trp
		290				295					300				
His	Arg	Thr	Gly	Glu	Ser	Ile	Leu	Ser	Leu	Leu	Arg	Asp	Pro	Ser	Lys
305					310					315					320
Arg	Lys	Val	Pro	Pro	Gln	Pro	Leu	Thr	Pro	Asn	Gln	Ile	Val	Ser	Thr
				325					330					335	
Leu	Phe	Thr	Ser	Asn	Phe	Ile	Gly	Ile	Cys	Phe	Ser	Arg	Ser	Leu	His
			340					345					350		
Tyr	Gln	Phe	Tyr	Val	Trp	Tyr	Phe	His	Thr	Leu	Pro	Tyr	Leu	Leu	Trp
		355					360					365			
Ala	Met	Pro	Ala	Arg	Trp	Leu	Thr	His	Leu	Leu	Arg	Leu	Leu	Val	Leu
		370				375					380				
Gly	Leu	Ile	Glu	Leu	Ser	Trp	Asn	Thr	Tyr	Pro	Ser	Thr	Ser	Cys	Ser
385					390					395					400
Ser	Ala	Ala	Leu	His	Ile	Cys	His	Ala	Val	Ile	Leu	Leu	Gln	Leu	Trp
				405					410					415	
Leu	Gly	Pro	Gln	Pro	Phe	Pro	Lys	Ser	Thr	Gln	His	Ser	Lys	Lys	Ala
			420					425					430		

His

&lt;210&gt; SEQ ID NO 37

&lt;211&gt; LENGTH: 1291

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Aspergillus nidulans

&lt;400&gt; SEQUENCE: 37

aagcttcaag ctgtaaggat ttcggcacgg ctacggaaga cggagaagcc caccttcagt	60
ggactcgagt accatttaat tctattttgtg tttgatcgag acctaataca gccctacaa	120
cgaccatcaa agtcgtatag ctaccagtga ggaagtggac tcaaatcgac ttcagcaaca	180
tctcctggat aaactttaag cctaaactat acagaataag atgggtggaga gcttataccg	240
agctcccaaa tctgtccaga tcatggttga cgggtgctg gatcttcta tagaatcatc	300
cttattcgtt gacctagctg attctggagt gaccagagg gtcattgactt gagcctaaaa	360

-continued

---

```

tccgccgect ccaccatttg tagaaaaatg tgacgaactc gtgagctctg tacagtgacc 420
ggtgactcct tctggcatgc ggagagacgg acggacgcag agagaagggc tgagtaataa 480
gcgccactgc gccagacagc tctggcggct ctgaggtgca gtggatgatt attaatccgg 540
gaccggccgc ccctccgcc cgaagtggaa aggttgggtg gccctcgtt gaccaagaat 600
ctattgcac atcggagaat atggagcttc atcgaatcac cggcagtaag cgaaggagaa 660
tgtgaagcca ggggtgtata gccgtcggcg aatagcatg ccattaacct aggtacagaa 720
gtccaattgc ttccgatctg gtaaaagatt cacgagatag taccttctcc gaagtaggta 780
gagcgagtac ccggcgcgta agctccctaa ttggccatc cggcatctgt agggcgtcca 840
aatatcgtgc ctctcctgct ttgcccgggtg tatgaaaccg gaaaggccgc tcaggagctg 900
gccagcggcg cagaccgga acacaagctg gcagtcgacc catccggtgc tctgcactcg 960
acctgctgag gtcctcagt ccctggtagg cagctttgcc ccgtctgtcc gcccggtgtg 1020
tcggcggggg tgacaaggtc gttgcgtcag tccaacattt gttgccatat tttcctgctc 1080
tccccaccag ctgctctttt cttttctctt tcttttccca tcttcagtat attcatcttc 1140
ccatccaaga acctttattt ccctaagta agtactttgc tacatccata ctccatcctt 1200
cccatccctt attccttga acctttcagt tcgagctttc ccacttcac gcagcttgac 1260
taacagctac cccgcttgag cagacatcac a 1291

```

```

<210> SEQ ID NO 38
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: gpdA5F primer

```

```

<400> SEQUENCE: 38

```

```

cgcagatctc aagctgtaag gatttcggca 30

```

```

<210> SEQ ID NO 39
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: gpdA3R primer

```

```

<400> SEQUENCE: 39

```

```

caccgggccc atctcaaaca ttgtgatgtc tgctcaagcg 40

```

```

<210> SEQ ID NO 40
<211> LENGTH: 1255
<212> TYPE: DNA
<213> ORGANISM: Aspergillus nidulans
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(236)
<220> FEATURE:
<221> NAME/KEY: Intron
<222> LOCATION: (237)..(366)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (367)..(1252)

```

```

<400> SEQUENCE: 40

```

```

atg ttt gag atg ggc ccg gtg gga act cgt ctc ccc gcc atg acc tct 48
Met Phe Glu Met Gly Pro Val Gly Thr Arg Leu Pro Ala Met Thr Ser
1 5 10 15

```

```

cca gcg cac aac cac tac agc tac cac tct ccc acc tcc agc gac aga 96
Pro Ala His Asn His Tyr Ser Tyr His Ser Pro Thr Ser Ser Asp Arg

```

-continued

	20	25	30	
ggc cgg tca agg cag aac tcg gat gcc atg gac atc cag tcc atc act				144
Gly Arg Ser Arg Gln Asn Ser Asp Ala Met Asp Ile Gln Ser Ile Thr				
	35	40	45	
gaa cga gag ccg gcg acc aga tac gcg gtt gcg ggc ggc cct gcg ccc				192
Glu Arg Glu Pro Ala Thr Arg Tyr Ala Val Ala Gly Gly Pro Ala Pro				
	50	55	60	
tgg aat cgc aac ggg tct ccg agc atg agc cct atg tat agc aa				236
Trp Asn Arg Asn Gly Ser Pro Ser Met Ser Pro Met Tyr Ser Asn				
	65	70	75	
gtacatctct cttaccctc cgtttctttc tgcttttcta ccaccccatc cctctttcca				296
gtctgagtc aggcttggtc cgcttgaagt ggctaatgtg atcctegtct tctctcttcc				356
tgtgttttag c aat tcc gag cga aac cag ttt cat gaa gag aac gga cgc				406
Asn Ser Glu Arg Asn Gln Phe His Glu Glu Asn Gly Arg				
	80	85	90	
acc tac cat ggc ttt cgc agg gga atg tat ttt ctt ccg tgc gat gag				454
Thr Tyr His Gly Phe Arg Arg Gly Met Tyr Phe Leu Pro Cys Asp Glu				
	95	100	105	
caa gaa cag gat cgc ctc gac atc ttc cat aag cta ttc acg gta gcg				502
Gln Glu Gln Asp Arg Leu Asp Ile Phe His Lys Leu Phe Thr Val Ala				
	110	115	120	
cgg gta tcg gag agt ctg atc tac gcg ccc cat cca acc aac ggc cgg				550
Arg Val Ser Glu Ser Leu Ile Tyr Ala Pro His Pro Thr Asn Gly Arg				
	125	130	135	140
ttt ctg gac cta gga tgt gga act ggt atc tgg gcg atc gag gta gcg				598
Phe Leu Asp Leu Gly Cys Gly Thr Gly Ile Trp Ala Ile Glu Val Ala				
	145	150	155	
aac aag tac cct gat gcg ttt gtc gct ggt gtg gat ttg gct cct att				646
Asn Lys Tyr Pro Asp Ala Phe Val Ala Gly Val Asp Leu Ala Pro Ile				
	160	165	170	
cag cct ccg aac cac ccg aag aac tgc gag ttc tac gcg ccc ttc gac				694
Gln Pro Pro Asn His Pro Lys Asn Cys Glu Phe Tyr Ala Pro Phe Asp				
	175	180	185	
ttc gaa gcg cca tgg gcc atg ggg gag gat tcc tgg gat cta atc cat				742
Phe Glu Ala Pro Trp Ala Met Gly Glu Asp Ser Trp Asp Leu Ile His				
	190	195	200	
ctg cag atg ggt tgc ggt agt gtc atg ggc tgg cca aac ttg tat cga				790
Leu Gln Met Gly Cys Gly Ser Val Met Gly Trp Pro Asn Leu Tyr Arg				
	205	210	215	220
agg ata ttc gca cat ctc cgt ccc ggt gcc tgg ttt gag cag gtt gag				838
Arg Ile Phe Ala His Leu Arg Pro Gly Ala Trp Phe Glu Gln Val Glu				
	225	230	235	
atc gat ttc gag cct cga tgt gat gat cgg tca cta gat gga acg gca				886
Ile Asp Phe Glu Pro Arg Cys Asp Asp Arg Ser Leu Asp Gly Thr Ala				
	240	245	250	
ttg cgg cat tgg tac gat tgt ctt aaa cag gcg aca gca gag acc atg				934
Leu Arg His Trp Tyr Asp Cys Leu Lys Gln Ala Thr Ala Glu Thr Met				
	255	260	265	
cgg cca atc gcc cat agc tcc cgc gat aca ata aaa gac ctg cag gac				982
Arg Pro Ile Ala His Ser Ser Arg Asp Thr Ile Lys Asp Leu Gln Asp				
	270	275	280	
gct ggg ttc acg gag atc gac cat caa ata gtg gga ctc ccg ctc aac				1030
Ala Gly Phe Thr Glu Ile Asp His Gln Ile Val Gly Leu Pro Leu Asn				
	285	290	295	300
ccg tgg cat cag gac gaa cac gag cgg aag gtg gca cgt tgg tat aac				1078
Pro Trp His Gln Asp Glu His Glu Arg Lys Val Ala Arg Trp Tyr Asn				
	305	310	315	
ctg gcc gtc tca gag agc atc gaa aac ctc agt ctg gct ccc ttc agt				1126

-continued

---

Leu Ala Val Ser Glu Ser Ile Glu Asn Leu Ser Leu Ala Pro Phe Ser  
                   320                                  325                                  330

cgt gtc tat cgc tgg ccc ctg gag aga atc cag caa ctc gcc gca gat 1174  
 Arg Val Tyr Arg Trp Pro Leu Glu Arg Ile Gln Gln Leu Ala Ala Asp  
                   335                                  340                                  345

gtg aag tcc gaa gca ttc aac aaa gag atc cat gcc tac aat ata ctg 1222  
 Val Lys Ser Glu Ala Phe Asn Lys Glu Ile His Ala Tyr Asn Ile Leu  
                   350                                  355                                  360

cac ata tac cag gct agg aaa cca tta aga taa 1255  
 His Ile Tyr Gln Ala Arg Lys Pro Leu Arg  
                   365                                  370

<210> SEQ ID NO 41  
 <211> LENGTH: 374  
 <212> TYPE: PRT  
 <213> ORGANISM: Aspergillus nidulans

<400> SEQUENCE: 41

Met Phe Glu Met Gly Pro Val Gly Thr Arg Leu Pro Ala Met Thr Ser  
 1                  5                                  10                                  15

Pro Ala His Asn His Tyr Ser Tyr His Ser Pro Thr Ser Ser Asp Arg  
                   20                                  25                                  30

Gly Arg Ser Arg Gln Asn Ser Asp Ala Met Asp Ile Gln Ser Ile Thr  
                   35                                  40                                  45

Glu Arg Glu Pro Ala Thr Arg Tyr Ala Val Ala Gly Gly Pro Ala Pro  
                   50                                  55                                  60

Trp Asn Arg Asn Gly Ser Pro Ser Met Ser Pro Met Tyr Ser Asn Asn  
 65                                  70                                  75                                  80

Ser Glu Arg Asn Gln Phe His Glu Glu Asn Gly Arg Thr Tyr His Gly  
                   85                                  90                                  95

Phe Arg Arg Gly Met Tyr Phe Leu Pro Cys Asp Glu Gln Glu Gln Asp  
                   100                                  105                                  110

Arg Leu Asp Ile Phe His Lys Leu Phe Thr Val Ala Arg Val Ser Glu  
                   115                                  120                                  125

Ser Leu Ile Tyr Ala Pro His Pro Thr Asn Gly Arg Phe Leu Asp Leu  
                   130                                  135                                  140

Gly Cys Gly Thr Gly Ile Trp Ala Ile Glu Val Ala Asn Lys Tyr Pro  
 145                                  150                                  155                                  160

Asp Ala Phe Val Ala Gly Val Asp Leu Ala Pro Ile Gln Pro Pro Asn  
                   165                                  170                                  175

His Pro Lys Asn Cys Glu Phe Tyr Ala Pro Phe Asp Phe Glu Ala Pro  
                   180                                  185                                  190

Trp Ala Met Gly Glu Asp Ser Trp Asp Leu Ile His Leu Gln Met Gly  
                   195                                  200                                  205

Cys Gly Ser Val Met Gly Trp Pro Asn Leu Tyr Arg Arg Ile Phe Ala  
                   210                                  215                                  220

His Leu Arg Pro Gly Ala Trp Phe Glu Gln Val Glu Ile Asp Phe Glu  
 225                                  230                                  235                                  240

Pro Arg Cys Asp Asp Arg Ser Leu Asp Gly Thr Ala Leu Arg His Trp  
                   245                                  250                                  255

Tyr Asp Cys Leu Lys Gln Ala Thr Ala Glu Thr Met Arg Pro Ile Ala  
                   260                                  265                                  270

His Ser Ser Arg Asp Thr Ile Lys Asp Leu Gln Asp Ala Gly Phe Thr  
                   275                                  280                                  285

Glu Ile Asp His Gln Ile Val Gly Leu Pro Leu Asn Pro Trp His Gln  
                   290                                  295                                  300

-continued

Asp Glu His Glu Arg Lys Val Ala Arg Trp Tyr Asn Leu Ala Val Ser  
 305 310 315 320

Glu Ser Ile Glu Asn Leu Ser Leu Ala Pro Phe Ser Arg Val Tyr Arg  
 325 330 335

Trp Pro Leu Glu Arg Ile Gln Gln Leu Ala Ala Asp Val Lys Ser Glu  
 340 345 350

Ala Phe Asn Lys Glu Ile His Ala Tyr Asn Ile Leu His Ile Tyr Gln  
 355 360 365

Ala Arg Lys Pro Leu Arg  
 370

<210> SEQ ID NO 42  
 <211> LENGTH: 40  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: LaeA5F primer

<400> SEQUENCE: 42

cgcttgagca gacatcacia tgtttgagat gggcccggtg 40

<210> SEQ ID NO 43  
 <211> LENGTH: 29  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: LaeA3R primer

<400> SEQUENCE: 43

cgcagatctg aggattatga gaagggagc 29

<210> SEQ ID NO 44  
 <211> LENGTH: 2866  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: pGPDA promoter LeaA fragment

<400> SEQUENCE: 44

aagcttcaag ctgtaaggat ttcggcacgg ctacggaaga cggagaagcc caccttcagt 60  
 ggactcgagt accatttaat tctatttgtg tttgatcgag acctaataca gcccctacia 120  
 cgaccatcaa agtcgtatag ctaccagtga ggaagtggac tcaaatcgac ttcagcaaca 180  
 tctcctggat aaactttaag cctaaactat acagaataag atgggtggaga gcttataccg 240  
 agtcccaaaa tctgtccaga tcatggttga cgggtgctg gatcttcta tagaatcatc 300  
 cttattcggt gacctagctg attctggagt gaccagagg gtcattgact gagcctaaaa 360  
 tccgcccct ccaccatttg tagaaaaatg tgacgaactc gtgagctctg tacagtgacc 420  
 ggtgactctt tctggcatgc ggagagacgg acggacgcag agagaagggc tgagtaataa 480  
 gcgccactgc gccagacagc tctggcggct ctgaggtgca gtggatgatt attaattcgg 540  
 gaccggccgc cctccgccc cgaagtggaa aggtcgggtg gcccctcgtt gaccaagaat 600  
 ctattgcatc atcggagaat atggagcttc atcgaatcac cggcagtaag cgaaggagaa 660  
 tgtgaagcca ggggtgtata gccgtcggcg aatagcatg ccattaacct aggtacagaa 720  
 gtccaattgc ttcgatctg gtaaaagatt cacgagatag taccttctcc gaagtaggta 780  
 gagcgagtac cggcgcgta agtccctaa ttggccatc cggcatctgt agggcgtcca 840  
 aatattcgtc ctctcctgct ttgcccgtg tatgaaaccg gaaaggccgc tcaggagctg 900

-continued

---

```

gccagcggcg cagaccggga acacaagctg gcagtcgacc catccggtgc tctgcactcg 960
acctgctgag gtccctcagt ccctggtagg cagctttgcc ccgtctgtcc gcccggtgtg 1020
tcggcggggg tgacaaggtc gttgcgtcag tccaacattt gttgccatat tttcctgctc 1080
tccccaccag ctgctctttt cttttctctt tcttttccca tcttcagtat attcatcttc 1140
ccatccaaga acctttattt ccctaagta agtactttgc tacatccata ctccatcctt 1200
cccatccctt attcctttga acctttcagt tcgagctttc ccacttcacg gcagcttgac 1260
taacagctac cccgcttgag cagacatcac aatgtttgag atgggcccgg tgggaactcg 1320
tctccccgcc atgacctctc cagcgcacaa cactacagc taccactctc ccacctccag 1380
cgacagaggc cgggtcaaggc agaactcgga tgccatggac atccagtcca tcaactgaacg 1440
agagccggcg accagatacg cggttgcggg cggccctgcg ccctggaatc gcaacgggtc 1500
tccgagcatg agccctatgt atagcaagta catctctctt acccctccgt ttctttctgc 1560
ttttctacca ccccatccct ctttccagtc tgagtccagg cttgttccgc ttgaagtggc 1620
taatgtgatc ctgctcttct ctctttctgt gttttagcaa ttccgagcga aaccagtttc 1680
atgaagagaa cggacgcacc taccatggct ttcgcagggg aatgtatttt cttccgtgcg 1740
atgagcaaga acaggatcgc ctgcacatct tccataagct attcacggta gcgcggtat 1800
cggagagtct gatctacgcg ccccatccaa ccaacggccg gtttctggac ctaggatgtg 1860
gaactgggat ctgggcgatc gaggtagcga acaagtaccg tgatgcgttt gtcgctgggtg 1920
tggatttggc tcctattcag cctccgaacc acccgaagaa ctgcgagttc tacgcgccct 1980
tcgacttcga agcgcctatg gccatggggg aggattcctg ggatctaata catctgcaga 2040
tgggttgccg tagtgtcatg ggctggccaa acttgatcgc aaggatattc gcacatctcc 2100
gtcccgggtc ctggtttgag caggttgaga tcgatttoga gcctcgatgt gatgatcgg 2160
cactagatgg aacggcattg cggcattggt acgattgtct taaacaggcg acagcagaga 2220
ccatgcggcc aatcgcccat agctcccgcg atacaataaa agacctgcag gacgctgggt 2280
tcacggagat cgaccatcaa atagtgggac tcccgtcaa cccgtggcat caggacgaac 2340
acgagcggaa ggtggcacgt tggataaacc tggccgtctc agagagcatc gaaaacctca 2400
gtctggctcc cttcagtcgt gtctatcgtt ggcccctgga gagaatccag caactcgccg 2460
cagatgtgaa gtccgaagca ttcaacaaag agatccatgc ctacaatata ctgcacatat 2520
accaggctag gaaaccatta agataagagc aaaaggcgac cacatccagg aacgcaaac 2580
gaaaaggagg aaaactgcta gcgcaagttt atgtcacgct ggcacacgcc cagccatcag 2640
aaatctcaac agcgaagtt atgaaccgca tcaaccgagt atgaacgaca attcgtccat 2700
cacacacctc tcggttcctc tcgcaggccc agcatggcgc cctatcaacc tgctttacga 2760
cgtcgtatat actggcgaag taccctctct atctactctg gcgctctaga taccgtgaag 2820
atgcagacaa aattggccga gctcccttct cataatcctc aagctt 2866

```

&lt;210&gt; SEQ ID NO 45

&lt;211&gt; LENGTH: 1279

&lt;212&gt; TYPE: DNA

<213> ORGANISM: *Aspergillus niger*

&lt;400&gt; SEQUENCE: 45

```

atgcatcatt ctcccgtttt gtttttgggc ccaaactaac cgagtaggtg tggcatttgc 60
gggcatgatg tttcaactac cgctgatcat tatcaccgcc ccattagaga agatccaaga 120

```

-continued

---

```

ccctactggg aaggtgatag gcaattccat tttctgggtt agtttttgtc ttgtcggcca 180
gcctttggga gctttgctgt acttctttgc ctggcaagcg aagtatggca gtgtgagccg 240
aatgtgaatg taaaagcacg cacgtgtccg ctgtttgtca tagatgtaaa taaatgccaa 300
caacttcagc cttttttga aaagcaaagc aaccgaagta aacgatcctg taccatcagc 360
gctcctcaca atggaatcct ttagatgttt ctgttccatt catcttgctt actgcaatgt 420
tcttttgcg tttgactaat tctccggatg ttgaatggca acgctgtcgg cgtcgggtct 480
tcagggatcc gccaaggatg ctctggatcc gcatccggcc gctcttgccg cccatcaatc 540
gcccgactat aaatcgaact actttcggca tcttctagac ttctaatac cgcctagtca 600
tagcagatcc aagctgagaa caccacaagt aaatatcacc catcatgctt accctgaccg 660
tcctgaaaa ctacgggatg gtgccaattc tacaattcct tgcagacaat gccattctcc 720
ccatgaagtc tgatgctaac tctctgcag ctctgtcatt gccgtcgtc tgggtgcat 780
ccccgtcctg agcttcgtcc atggcgccgt cgtgtctcgt ctccgcaagg aagctgattg 840
cccctaccct cactgctatg cgaccgtaga gcagtgcaag accaacgtaa gccaacctca 900
cacaacagc attcctcgag ctaacataca ttccgaaccg tgcagcccaa ggccgagcag 960
ttcaactgcg ctacgcgcgc tcatgccaac ttcttgaga actccagcca aactatgctc 1020
ttcctcctgg tagctggact gaagtacccc cagtggcgga ctggcctcgg aagcatctgg 1080
gtcctcggtc gctcactgtt cctttacgga tatgtgtact ccggcaagcc gcgggggtcgc 1140
ggtcgtttgt acggcagctt ctacttgctt gcacagggag ctctctgggg cttgacgtct 1200
tttgagttg cgagggagt gatttcctac ttctaagttt ggactgaatc cgtggtgtga 1260
ttgaggtgat tggcgatgt 1279

```

```

<210> SEQ ID NO 46
<211> LENGTH: 472
<212> TYPE: DNA
<213> ORGANISM: Aspergillus nidulans
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (361)..(361)
<223> OTHER INFORMATION: n is a, c, g, or t

```

```

<400> SEQUENCE: 46

```

```

ggaccgatgg ctgtgtagaa gtactcgcgc atagtggaaa ccgacgcccc agcactcgtc 60
cgagggcaaaa ggaatagagt agatgccgac cgcgggatcc acttaacgtt actgaaatca 120
tcaaacagct tgacgaatct ggatataaga tcgttgggtg cgatgtcagc tccggagtgt 180
agacaaatgg tgttcaggat ctcgataaga tacgttcatt tgtccaagca gcaaagagtg 240
ccttctagtg atttaatagc tccatgtcaa caagaataaa acgcgttttc gggtttacct 300
cttcagata cagctcatct gcaatgcatt aatgcattga ctgcaaccta gtaacgcctt 360
ncaggctccg gcgaagagaa gaatagctta gcagagctat tttcattttc gggagacgag 420
atcaagcaga tcaacgggtc tcaagagacc tacgagactg aggaatccgc tc 472

```

```

<210> SEQ ID NO 47
<211> LENGTH: 2005
<212> TYPE: DNA
<213> ORGANISM: Aspergillus oryzae

```

```

<400> SEQUENCE: 47

```

```

tctgacagac gggcaattga ttacgggatc ccattggtaa cgaaatgtaa aagctaggag 60
atcgtccgcc gatgtcagga tgatttcact tgtttcttgt ccggtcacc ggtcaaagct 120

```

-continued

---

```

aaagaggagc aaaaggaacg gatagaatcg ggtgccgctg atctatacgg tatagtgccc 180
ttatcacggt gactcaaccc atgctattta actcaacccc tccttctgaa ccccaccatc 240
ttcttccttt tcctctcatc ccacacaatt ctctatctca gatttgaatt ccaaaagtcc 300
tcggacgaaa ctgaacaagt cttectcect tcgataaacc tttggtgatt ggaataactg 360
accatcttct atagttcca aaccaaccga caatgtaaat aactcctcg attagcctc 420
tagagggcat acgatggaag tcatggaata cttttggctg gactctcaca atgatcaagg 480
tatcttaggt aacgtctttg gcgtgggccc gtgttcgctc ccagtcacg atgcattcac 540
atgccctccc taagctggc cctagactct aggatcctag tctagaagga catggcatcg 600
atggactggg ttcgttctga gattatacgg ctaaaacttg atctggataa taccagcgaa 660
aagggtcatg cttctctcg ttcttctgt tgatggaatg gctaacagat gatagtcatt 720
gcaacttgaa acatgtctcc tccagctgcc atctacgaac ccaactgtggc cgctaccggc 780
ctcaagggtta aggtcgtggt ttctgagacc gtcccgttg agggagcttc tcagaccaag 840
ctggttgacc atttcggtgg caagtgggac gagttcaagt tcgcccctat ccgcgaaagc 900
caggtctctc gtgccatgac cagacgttac tttgaggacc tggacaagta cgctgaaagt 960
gacgttgca ttgttggtgc tggttcctgc ggtctgagca ctgcgtacgt cttggccaag 1020
gctcgtccgg acctgaagat tgctatcgtc gaggccagcg tctctcctgg tcagtgtcc 1080
atgatggatt gccttgact cagctttccg gaactaacgt gcaatagggt gcggtgcctg 1140
gttgggtggc caactctttt ctgctatggt catgcgccgt cccgcggaag tcttctgaa 1200
cgagctgggt gttccttacg aagaggacgc aaaccccaac tacgttgtcg tcaagcacgc 1260
ctccctgttt acctcgacac tcatgtcgaa ggttctctcc tccccaatg tcaagctctt 1320
caatgctacc gctggtgagg acttgatcac ccgtccgacc gagaacggca acccccagat 1380
tgctgggtgt gtcgtcaact ggacgctggt cacccttcc cagcatgatc actcctgcat 1440
ggaccccaac actatcaacg ctctgtcat catcagtacc actggtcacg atgggcccatt 1500
cggcgccttc tgtgcaagc gcttggtgtc catgggcagc gtcgacaagc taggtggcat 1560
gcgtggtctc gacatgaact cggccgagga tgccatcgtc aagaacaccc gcgaggttac 1620
taagggttg ataacggcg gtatggagct gtctgaaatt gatggctta accgcatggg 1680
ccctaccttc ggtgccatgg ttctcagtgg tgtcaaggct gccgaggagg cattgaaggt 1740
gttcgacgag cgtcagcgcg agtggtgctga gtaaagtact cactaccga atgggttcag 1800
tgcatgaacc ggatttgtct tacggctctt gacgataggg gaatgatgat tatgtgatag 1860
ttctgagatt tgaatgaact cgtagctcg taatccacat gcatatgtaa atggctgtgt 1920
cccgtatgta acggtggggc attctagaat aattatgtgt aacaagaaag acagtataat 1980
acaaacaaag atgcaagagc ggctc 2005

```

&lt;210&gt; SEQ ID NO 48

&lt;211&gt; LENGTH: 1052

&lt;212&gt; TYPE: DNA

<213> ORGANISM: *Aspergillus niger*

&lt;400&gt; SEQUENCE: 48

```

gtcatcgcag ataagcactg ctgtcttgca tccaagtacg cgtcagcaga aatacgggac 60
ttccgaaagt atatggcaaa attaaagaac ttgactctcc agcaatgttt tgccctgacc 120
gtcgctaaaa cgttactacc cctatacccg tctgtttgct ccagcccag gcattaggtc 180

```



-continued

---

tgactgacag cacggcgcca tgcgggcttg ggacgccatg tccgtcgcgt gataaggggt	240
gatccatgca gctactatcc ttccatcggt ccattcccat ccttgctcta tctccatcct	300
tgaaacttta ctagttagt tggatgctcg agcttgctct cggctactcc gtccaatgga	360
taagaccccg atgccggtcc tcattggctc ccagctggta tcgcccacac cttcgtgtga	420
tcgcctctct gcttcccctc atcatcatta ctaactagta catccaaaag ccatcccagt	480
gcttcccctc acccttgccc aagacattcc aagtgggctc tcggctggaa aacatggacc	540
cattggttcc atcgataagc tagctcctcg tccgttacc cagattgata ccagataaca	600
ttgaccagcg gcttatcacc gaggtctgcg ggtgagacc cccctgacac aagttagata	660
aaagaaactc gcctcattgt gcttccgatg ggtcgggatg acgagccttc ggaaagagct	720
ggcgcctctt taaaggggac agctgtcgcc aagtgtgaa attctccgat aactactaac	780
aatctctccc ttccttcccg ctactgtggt caccaaatca actctctttt ctggccaag	840
atctaactatg gcggatgaga agactgaaaa gtctccccc ccatgacgag tggatgagga	900
gactggcaca acagaggaaa ttgacccgac aatggcaaac catacgaagg atgcagacga	960
ggcactggcg gtcttcgaag acctccatgg tgaagtcac acacttgatg aggagacaaa	1020
caaaaggata cttcggacaa ttgactggca ca	1052

<210> SEQ ID NO 49  
 <211> LENGTH: 58  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: pyrGU5F primer

<400> SEQUENCE: 49

gtaacgccag ggttttcca gtcacgacgt ttaaacatgc atcattctcc cgctttgt	58
--	----

<210> SEQ ID NO 50  
 <211> LENGTH: 49  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: pyrGU3R primer

<400> SEQUENCE: 50

tgccgaaatc cttacagctt gaagcttcat cgccaatcac ctcaatcac	49
---	----

<210> SEQ ID NO 51  
 <211> LENGTH: 49  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Trp5F primer

<400> SEQUENCE: 51

agctcccttc tcataatcct caagcttggc ccatggcttg tgtagaagt	49
---	----

<210> SEQ ID NO 52  
 <211> LENGTH: 46  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Trp3R primer

<400> SEQUENCE: 52

cgtaatcaat tgcccgtctg tcagagagcg gattcctcag tctcgt	46
--	----

-continued

---

<210> SEQ ID NO 53  
 <211> LENGTH: 46  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: PTR5F primer  
  
 <400> SEQUENCE: 53  
  
 acgagactga ggaatccgct ctctgacaga cgggcaattg attacg 46

<210> SEQ ID NO 54  
 <211> LENGTH: 45  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: PTR3R primer  
  
 <400> SEQUENCE: 54  
  
 acagcagtgc ttatctgcca tgacgagccg ctcttgcatc tttgt 45

<210> SEQ ID NO 55  
 <211> LENGTH: 45  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: PyrGD5F primer  
  
 <400> SEQUENCE: 55  
  
 acaaagatgc aagagcggct cgatcatgca gataagcact gctgt 45

<210> SEQ ID NO 56  
 <211> LENGTH: 44  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: PyrGD3R primer  
  
 <400> SEQUENCE: 56  
  
 tgagacgctg tttcaccgag tacatcgcca atcacctcaa tcac 44

<210> SEQ ID NO 57  
 <211> LENGTH: 7689  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: transgene  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (4513)..(4513)  
 <223> OTHER INFORMATION: n is a, c, g, or t  
  
 <400> SEQUENCE: 57  
  
 gtttaaacat gcatcattct cccgctttgt ttttgggccc aaactaaccg agtaggtgtg 60  
 gcatttgccg gcatgatgtt tcaactaccg ctgatcatta tcaccgcccc attagagaag 120  
 atccaagacc ctactgggaa ggtgatagcc aattccattt tctgggtag tttttgtctt 180  
 gtcggccagc ctttgggagc tttgctgtac ttctttgcoct ggcaagcgaa gtatggcagt 240  
 gtgagccgaa tgtgaatgta aaagcacgca cgtgtccgct gtttgtcata gatgtaaata 300  
 aatgccaaca acttcagcca ttttttgaag agcaaagcaa ccgaagtaaa cgatcctgta 360  
 ccatcagcgc tctcacaat ggaatctttt agatgtttct gttccattca tcttgcttac 420  
 tgcaatgttc ttttcgcgtt tgactaattc tccggatggt gaatggcaac gctgtcggcg 480  
 tcgggtcttc agggatccgc caaggatgct ctggatccgc atccggccgc tcttgccgcc 540

-continued

catcaatcgc	ccgactataa	atcgaactac	tttcggcatc	ttctagactt	cctaataaccg	600
cctagtcata	gcagattcaa	gctgagaaca	ccacaagtaa	atatcaccca	tcatgcttac	660
cctgaccgtc	cctgaaaact	acgggtatgt	gccaatctta	caattccttg	cagacaatgc	720
cattctcccc	atgaagtctg	atgctaacta	tctgcagct	ctgtcattgc	cgctcgtctg	780
ggtgccatcc	ccgtcctgag	cttcgtccat	ggcgccgtcg	tgtctcgtct	ccgcaaggaa	840
gctgattgcc	cctaccctca	ctgctatgcg	accgtagagc	agtgcaagac	caacgtaagc	900
caacctcaca	caaacaggat	tctcagagct	aacatacatt	ccgaaccgtg	cagcccaagg	960
ccgagcagtt	caactgcgct	cagcgcgctc	atgccaaactt	ccttgagaac	tccagccaaa	1020
ctatgctctt	cctcctggta	gctggactga	agtacccccca	gttggcgact	ggcctcggaa	1080
gcatctgggt	cctcggtcgc	tcaactgttcc	tttacggata	tgtgtactcc	ggcaagccgc	1140
ggggtcgcgg	tcgtttgtag	ggcagcttct	acttgcttgc	acagggagct	ctctggggct	1200
tgacgtcttt	tggagttgcg	agggagttga	tttctactt	ctaagtttgg	actgaatccg	1260
tgggtgatt	gaggtgattg	gcgatgaagc	ttcaagctgt	aaggatttcg	gcacggctac	1320
ggaagacgga	gaagcccacc	ttcagtggac	tcgagtacca	tttaattcta	tttgtgtttg	1380
atcgagacct	aatacagccc	ctacaacgac	catcaaagtc	gtatagctac	cagtgaggaa	1440
gtggactcaa	atcgacttca	gcaacatctc	ctggataaac	tttaagccta	aactatacag	1500
aataagatgg	tggagagctt	ataccgagct	cccaaactcg	tccagatcat	ggttgaccgg	1560
tgcttgatc	ttctataga	atcatcctta	ttcgttgacc	tagctgattc	tggagtgacc	1620
cagagggcca	tgacttgagc	ctaaaatccg	ccgcctccac	cattttaga	aaaatgtgac	1680
gaactcgtga	gctctgtaca	gtgaccgggtg	actctttctg	gcatgcggag	agacggacgg	1740
acgcagagag	aagggctgag	taataagcgc	cactgcgcca	gacagctctg	gcggctctga	1800
ggtgcagtgg	atgattatta	atccgggacc	ggccgcccct	ccgccccgaa	gtggaaaggc	1860
tgggtgccc	ctcgttgacc	aagaatctat	tgcacatcg	gagaatatgg	agcttcatcg	1920
aatcaccggc	agtaagcga	ggagaatgtg	aagccagggg	tgtatagccg	tcggcgaaat	1980
agcatgccat	taacctaggt	acagaagtcc	aattgcttcc	gatctggtaa	aagattcacg	2040
agatagtacc	ttctccgaag	taggtagagc	gagtaccogg	cgcgtaagct	ccctaattgg	2100
cccatccggc	atctgtaggg	cgtccaaata	tcgtgcctct	cctgctttgc	ccggtgatg	2160
aaaccggaaa	ggccgctcag	gagctggcca	gcggcgcaga	ccgggaacac	aagctggcag	2220
tcgaccatc	ccgtgctctg	cactcgacct	gctgaggtcc	ctcagtcctt	ggtaggcagc	2280
tttccccgt	ctgtccgccc	ggtgtgtcgg	cggggtgac	aaggctcgtt	cgctcagtcca	2340
acatttggtg	ccatattttc	ctgctctccc	caccagctgc	tcttttctt	tctctttctt	2400
ttccatctt	cagtatattc	atcttcccat	ccaagaacct	ttatttcccc	taagtaagta	2460
ctttgctaca	tccatactcc	atccttccca	tcccttattc	ctttgaacct	ttcagttcga	2520
gctttcccac	ttcatcgcag	cttgactaac	agctaccccg	cttgagcaga	catcacaatg	2580
tttgagatgg	gcccgggtggg	aactcgtctc	cccgccatga	cctctccagc	gcacaaccac	2640
tacagctacc	actctcccac	ctccagcgac	agaggccggg	caaggcagaa	ctcggatgcc	2700
atggacatcc	agtcacac	tgaacgagag	ccggcgacca	gatacgcggg	tgcggggcggc	2760
cctgcgccct	ggaatcgcaa	cgggtctccg	agcatgagcc	ctatgtatag	caagtacatc	2820
tctcttacc	ctccgtttct	ttctgctttt	ctaccacccc	atccctcttt	ccagtctgag	2880
tccaggttg	ttccgcttga	agtggcta	gtgatcctcg	tcttctctct	ttctgtgttt	2940

-continued

---

tagcaattcc	gagcgaaacc	agtttcatga	agagaacgga	cgcacctacc	atggctttcg	3000
caggggaatg	tattttcttc	cgtgcgatga	gcaagaacag	gatcgectcg	acatcttcca	3060
taagctattc	acggtagcgc	gggtatcgga	gagtctgatc	tacgcgcccc	atccaaccaa	3120
cggccggttt	ctggacctag	gatgtggaac	tggatatctgg	gcgatcgagg	tagcgaacaa	3180
gtacctgat	gcgtttgtcg	ctgggtgga	tttggctcct	attcagcctc	cgaaccaccc	3240
gaagaactgc	gagttctacg	cgcccttcga	cttcgaagcg	ccatgggcca	tgggggagga	3300
ttcctgggat	ctaataccatc	tgcagatggg	ttgcggtagt	gtcatgggct	ggccaaactt	3360
gtatcgaagg	atattcgcac	atctccgtcc	cggtgccctgg	tttgagcagg	ttgagatcga	3420
tttcgagcct	cgatgtgatg	atcggtcact	agatggaacg	gcattgcggc	attggtacga	3480
ttgtcttaaa	caggcgacag	cagagaccat	gcggccaatc	gcccatagct	cccgcgatac	3540
aataaaagac	ctgcaggacg	ctgggttcac	ggagatcgac	catcaaatag	tgggactccc	3600
gctcaaccgg	tggcatcagg	acgaacacga	gcggaagggtg	gcacgttggg	ataacctggc	3660
cgtctcagag	agcatcgaaa	acctcagtct	ggctcccttc	agtcgtgtct	atcgctggcc	3720
cctggagaga	atccagcaac	tgcgccgaga	tgtgaagtcc	gaagcattca	acaagagat	3780
ccatgcctac	aataactgc	acataacca	ggctaggaaa	ccattaagat	aagagcaaaa	3840
ggcgaccaca	tccaggaacg	caaaacgaaa	aggaggaaaa	ctgctagcgc	aagtttatgt	3900
cacgctggca	cacgcccagc	catcagaaat	ctcaacagcg	aaagttatga	accgcatcaa	3960
ccgagtatga	acgacaattc	gtccatcaca	cacccttcgg	ttcctctcgc	aggcccagca	4020
tggcgcccta	tcaacctgct	ttacgacgtc	gtatatactg	gcgaagtatc	ctctctatct	4080
actctggcgc	tctagatacc	gtgaagatgc	agacaaaatt	ggccgagctc	ccttctcata	4140
atcctcaagc	ttggaccgat	ggctgtgtag	aagtactcgc	cgatagtgga	aaccgacgcc	4200
ccagcactcg	tccgagggca	aaggaataga	gtagatgccg	accgcgggat	ccacttaacg	4260
ttactgaaat	catcaaacag	cttgacgaat	ctggatataa	gatcgttggg	gtcgatgtca	4320
gctccggagt	tgagacaaat	gggtttcagg	atctcgataa	gatacgttca	tttgtccaag	4380
cagcaaagag	tgccttctag	tgatttaata	gctccatgtc	aacaagaata	aaacgcgttt	4440
tgggtttac	ctcttcaga	tacagctcat	ctgcaatgca	ttaatgcatt	gactgcaacc	4500
tagtaacgcc	ttncaggctc	cggcgaagag	aagaatagct	tagcagagct	atthtcattt	4560
tccggagacg	agatcaagca	gatcaacggg	cgtcaagaga	cctacgagac	tgaggaatcc	4620
gctctctgac	agacgggcaa	ttgattacgg	gatcccattg	gtaacgaaat	gtaaaagcta	4680
ggagatcgtc	cgccgatgtc	aggatgattt	cacttgtttc	ttgtccggct	caccgggtcaa	4740
agctaaagag	gagcaaaagg	aacggataga	atcgggtgcc	gctgatctat	acggtatagt	4800
gcccttatca	cgttgactca	acctatgcta	tttaactcaa	cccctccttc	tgaacccac	4860
catcttcttc	cttttctct	catcccacac	aattctctat	ctcagatttg	aattccaaaa	4920
gtcctcggac	gaaactgaac	aagtcttctc	cccttcgata	aacctttggg	gattggaata	4980
actgaccatc	ttctatagtt	cccaaaccac	ccgacaatgt	aaatacactc	ctcgattagc	5040
cctctagagg	gcatacgatg	gaagtcatgg	aatacttttg	gctggactct	cacaatgatc	5100
aaggtatctt	aggtaacgtc	tttggcgtgg	gccggtgttc	gttcccagtc	atcgatgcat	5160
tcacatgccc	tccctaagct	gggccctaga	ctctaggatc	ctagtctaga	aggacatggc	5220
atcgatggac	tgggttcggt	ctgagattat	acggctaaaa	cttgatctgg	ataataccag	5280

-continued

---

cgaaaaggg	catgccttct	ctcgttcttc	ctggtgatgg	aatggctaac	agatgatagt	5340
cattgcaact	tgaaacatgt	ctcctccagc	tgccatctac	gaaccactg	tggccgctac	5400
cggcctcaag	ggtaaggtcg	tggtttctga	gaccgtcccc	gttgaggag	cttctcagac	5460
caagctggtg	gaccatttcg	gtggcaagtg	ggacgagttc	aagtctgccc	ctatccgca	5520
aagccaggtc	tctcgtgcca	tgaccagacg	ttactttgag	gacctggaca	agtacgctga	5580
aagtgaogtt	gtcattgttg	gtgctgggtc	ctgctgctcg	agcactgctg	acgtcttggc	5640
caaggctcgt	ccggacctga	agattgctat	ctcctgagcc	agcgtctctc	ctggctcagta	5700
gtccatgatg	gattgccttg	cactcagctt	tccggaacta	acgtgcaata	ggtggcgggtg	5760
cctggttggg	tggccaactc	ttttctgcta	tggctcatgcg	ccgtcccgcg	gaagtcttcc	5820
tgaacgagct	gggtgttctt	tacgaagagg	acgcaaacc	caactacgtt	gtcgtcaagc	5880
acgcctccct	gtttacctcg	acactcatgt	cgaaggttct	ctccttcccc	aatgtcaagc	5940
tcttcaatgc	taccgctgtt	gaggacttga	tcaccctgct	gaccgagaac	ggcaaccccc	6000
agattgctgg	tgttgcctgc	aactggacgc	tggctaccct	tcaccacgat	gatcactcct	6060
gcatggacct	caaacctatc	aacgctcctg	tcatcatcag	taccactggt	cacgatgggc	6120
cattcggcgc	cttctgtgcg	aagcgttgg	tgtccatggg	cagcgtcagc	aagctaggtg	6180
gcatgcgtgg	tctcgacatg	aactcggccg	aggatgccat	cgtcaagaac	accgcgaggg	6240
ttactaaggg	cttgataatc	ggcggatgg	agctgtctga	aattgatggc	tttaaccgca	6300
tgggcccctac	cttcgggtgcc	atggttctca	gtggtgtcaa	ggctgccgag	gaggcattga	6360
aggtgttcga	cgagcgtcag	cgcgagtgtg	ctgagtaa	gactcactac	ccgaatgggt	6420
tcagtgcattg	aaccggattt	gtcttacggg	ctttgacgat	aggggaatga	tgattatgtg	6480
atagttctga	gatttgaatg	aactcgttag	ctcgtaatcc	acatgcatat	gtaaattggct	6540
gtgtcccgtg	tgtaacggtg	gggcattcta	gaataattat	gtgtaacaag	aaagacagta	6600
taatacaaac	aaagatgcaa	gagcggctcg	tcatcgcaga	taagcactgc	tgtcttgcatt	6660
ccaagtccagc	gtcagcagaa	atacgggact	tccgaaagta	tatggcaaaa	ttaaagaact	6720
tgactctcca	gcaatgtttt	gccctgaccg	tcgctaaaac	gttactaccc	ctatacccg	6780
ctgtttgtcc	cagcccagag	cattaggtct	gactgacagc	acggcgccat	gcgggcttgg	6840
gacgccatgt	ccgtcgcgtg	ataaggggtg	atccatgcag	ctactatcct	tccatcgttc	6900
cattcccattc	cttgcctat	ctccatcctt	gaaactttac	tagtttagtt	ggatgctcga	6960
gcttgccttc	ggctactccg	tccaatggat	aagaccccga	tgccggctct	cattgggtctc	7020
cagctgggat	cgccccaac	ttcgtgtgat	cgctctctg	cttcccctca	tcatcattac	7080
taactagtac	atccaaaagc	catcccagtg	cttcccctca	cccttgccc	agacattcca	7140
agtgggcctt	cggctggaaa	acatggacct	attggttcca	tcgataagct	agctcctcgt	7200
ccgttaccct	agattgatac	cagataacat	tgaccagcgg	cttatcaccg	aggtctgcgg	7260
gtgagacccc	ccctgcgaca	agttagataa	aagaaaactg	cctcattgtg	cttccgatgg	7320
ggtcggatga	cgagccttcg	gaaagagctg	gcgcctcttt	aaaggggaca	gctgtcgcca	7380
agttgtgaaa	ttctccgata	actactaaca	atctctccct	tccttcccgc	tactgtggtc	7440
accaaataca	ctctcttttc	toggccaaga	tctaactatg	cggatgagaa	gactgaaaag	7500
tctccccac	cgatgacggt	ggatgaggag	actggcacia	cagaggaaat	tgacccgaca	7560
atggcaaagc	atacgaagga	tgacagcag	gcaactggcg	tcttcgaaga	cctccatggt	7620
gaagtcatca	cacttgatga	ggagacaaac	aaaaggatac	ttcggacaat	tgactggcac	7680

-continued

```

agtttaaac 7689

<210> SEQ ID NO 58
<211> LENGTH: 1270
<212> TYPE: DNA
<213> ORGANISM: Aspergillus niger
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(230)
<220> FEATURE:
<221> NAME/KEY: Intron
<222> LOCATION: (231)..(372)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (373)..(1267)

<400> SEQUENCE: 58

atg ttt gaa atc agc cga ctt ttg cat cag cca att act atg gct tcg 48
Met Phe Glu Ile Ser Arg Leu Leu His Gln Pro Ile Thr Met Ala Ser
1 5 10 15

ccg aat cgc aat aac tac agc tac caa ggg ata gaa tcc tat gat tcc 96
Pro Asn Arg Asn Asn Tyr Ser Tyr Gln Gly Ile Glu Ser Tyr Asp Ser
20 25 30

ggc cgt tcc agg caa aac tcg gat gct atg gac att cac gtc att acg 144
Gly Arg Ser Arg Gln Asn Ser Asp Ala Met Asp Ile His Val Ile Thr
35 40 45

gcc caa gaa cct cct cga gaa ccc ccg gac aac aac gat cct tat gat 192
Ala Gln Glu Pro Pro Arg Glu Pro Pro Asp Asn Asn Asp Pro Tyr Asp
50 55 60

ggc cat ggg ggt cca gct ggg act agc cat tat agc aa gtacttctcc 240
Gly His Gly Gly Pro Ala Gly Thr Ser His Tyr Ser Lys
65 70 75

cttctcatac tctgcacccc acgtaccccg caaaatccct ttttctcatg ccgtgcaaat 300

atcacactta tttctacaac taccgggcca ctaattcagg gaactttctt ttccggttgt 360

cgtttaatct ag g cct cca aac aga tgg ctc ttc tat gaa gaa aat ggg 409
Pro Pro Asn Arg Trp Leu Phe Tyr Glu Glu Asn Gly
80 85

cga aca tat cat gga tat cgc aga gga gtt tac ccg ctg cca tgc gat 457
Arg Thr Tyr His Gly Tyr Arg Arg Gly Val Tyr Pro Leu Pro Cys Asp
90 95 100 105

gaa cag gaa cag gac cgt ctc gat atc ttc cat aaa ctg ttc aca gta 505
Glu Gln Glu Gln Asp Arg Leu Asp Ile Phe His Lys Leu Phe Thr Val
110 115 120

gca cgg atg tcc gag agc tta atc tac gca cct cac ccc cca aat ggt 553
Ala Arg Met Ser Glu Ser Leu Ile Tyr Ala Pro His Pro Pro Asn Gly
125 130 135

cga ttc cta gat ctg ggg tgc ggc act ggg atc tgg gcc att gat gta 601
Arg Phe Leu Asp Leu Gly Cys Gly Thr Gly Ile Trp Ala Ile Asp Val
140 145 150

gcc cac aag tat ccc aat gct ttc gtt gct gga gta gat cta gca cct 649
Ala His Lys Tyr Pro Asn Ala Phe Val Ala Gly Val Asp Leu Ala Pro
155 160 165

ata cag cct ccc aac cac ccc gat aac tgc gag ttc tat gca cct ttt 697
Ile Gln Pro Pro Asn His Pro Asp Asn Cys Glu Phe Tyr Ala Pro Phe
170 175 180 185

gac ttt gag gcg cca tgg acg ctt ggg gaa aat tct tgg gat ctc att 745
Asp Phe Glu Ala Pro Trp Thr Leu Gly Glu Asn Ser Trp Asp Leu Ile
190 195 200

cat cta cag atg ggt tgc ggc agt gtt ctg ggc tgg cag aat ctc tac 793
His Leu Gln Met Gly Cys Gly Ser Val Leu Gly Trp Gln Asn Leu Tyr
205 210 215

```

-continued

---

```

aag cga atc tta agg cat ctt cag cct ggg gca tgg ttt gaa cag gtg      841
Lys Arg Ile Leu Arg His Leu Gln Pro Gly Ala Trp Phe Glu Gln Val
      220                      225                      230

gaa ata gat ttc gaa ccc cgc tgc gat gat cgc tcc ctg aat gga ctg      889
Glu Ile Asp Phe Glu Pro Arg Cys Asp Asp Arg Ser Leu Asn Gly Leu
      235                      240                      245

gca ctc cgg gag tgg tac cag tac ctg aag cag gcg aca caa gat aca      937
Ala Leu Arg Glu Trp Tyr Gln Tyr Leu Lys Gln Ala Thr Gln Asp Thr
      250                      255                      260                      265

atg cga ccc ata gcg cac agc tcg cgg gat acc atc aga cac ctt gag      985
Met Arg Pro Ile Ala His Ser Ser Arg Asp Thr Ile Arg His Leu Glu
      270                      275                      280

gag gca ggc ttt acc cag atc gac cat cag atg gtg ggg ctg cct ctc      1033
Glu Ala Gly Phe Thr Gln Ile Asp His Gln Met Val Gly Leu Pro Leu
      285                      290                      295

aac cct tgg cac cgt gat gaa cat gag cag aag gta gcc cgt tgg tat      1081
Asn Pro Trp His Arg Asp Glu His Glu Gln Lys Val Ala Arg Trp Tyr
      300                      305                      310

aac ctc gca atc tct gag agt atc gag acg ctc agc ctc gcc cct ttc      1129
Asn Leu Ala Ile Ser Glu Ser Ile Glu Thr Leu Ser Leu Ala Pro Phe
      315                      320                      325

agt cgc atc ttt cac tgg gat ctg gat agg atc aga cag atc aca gcg      1177
Ser Arg Ile Phe His Trp Asp Leu Asp Arg Ile Arg Gln Ile Thr Ala
      330                      335                      340                      345

gag gtc aag tca caa gcc ttc aac aag gaa atc cac gct tac aat atc      1225
Glu Val Lys Ser Gln Ala Phe Asn Lys Glu Ile His Ala Tyr Asn Ile
      350                      355                      360

tta cat ata tac cag gca cgg aag cgg ggc ggc cca tca ctt tga      1270
Leu His Ile Tyr Gln Ala Arg Lys Pro Gly Gly Pro Ser Leu
      365                      370                      375

```

&lt;210&gt; SEQ ID NO 59

&lt;211&gt; LENGTH: 375

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Aspergillus niger

&lt;400&gt; SEQUENCE: 59

```

Met Phe Glu Ile Ser Arg Leu Leu His Gln Pro Ile Thr Met Ala Ser
1                      5                      10                      15

Pro Asn Arg Asn Asn Tyr Ser Tyr Gln Gly Ile Glu Ser Tyr Asp Ser
      20                      25                      30

Gly Arg Ser Arg Gln Asn Ser Asp Ala Met Asp Ile His Val Ile Thr
      35                      40                      45

Ala Gln Glu Pro Pro Arg Glu Pro Pro Asp Asn Asn Asp Pro Tyr Asp
      50                      55                      60

Gly His Gly Gly Pro Ala Gly Thr Ser His Tyr Ser Lys Pro Pro Asn
      65                      70                      75                      80

Arg Trp Leu Phe Tyr Glu Glu Asn Gly Arg Thr Tyr His Gly Tyr Arg
      85                      90                      95

Arg Gly Val Tyr Pro Leu Pro Cys Asp Glu Gln Glu Gln Asp Arg Leu
      100                     105                     110

Asp Ile Phe His Lys Leu Phe Thr Val Ala Arg Met Ser Glu Ser Leu
      115                     120                     125

Ile Tyr Ala Pro His Pro Pro Asn Gly Arg Phe Leu Asp Leu Gly Cys
      130                     135                     140

Gly Thr Gly Ile Trp Ala Ile Asp Val Ala His Lys Tyr Pro Asn Ala
      145                     150                     155                     160

```

-continued

Phe Val Ala Gly Val Asp Leu Ala Pro Ile Gln Pro Pro Asn His Pro  
 165 170 175  
 Asp Asn Cys Glu Phe Tyr Ala Pro Phe Asp Phe Glu Ala Pro Trp Thr  
 180 185 190  
 Leu Gly Glu Asn Ser Trp Asp Leu Ile His Leu Gln Met Gly Cys Gly  
 195 200 205  
 Ser Val Leu Gly Trp Gln Asn Leu Tyr Lys Arg Ile Leu Arg His Leu  
 210 215 220  
 Gln Pro Gly Ala Trp Phe Glu Gln Val Glu Ile Asp Phe Glu Pro Arg  
 225 230 235 240  
 Cys Asp Asp Arg Ser Leu Asn Gly Leu Ala Leu Arg Glu Trp Tyr Gln  
 245 250 255  
 Tyr Leu Lys Gln Ala Thr Gln Asp Thr Met Arg Pro Ile Ala His Ser  
 260 265 270  
 Ser Arg Asp Thr Ile Arg His Leu Glu Glu Ala Gly Phe Thr Gln Ile  
 275 280 285  
 Asp His Gln Met Val Gly Leu Pro Leu Asn Pro Trp His Arg Asp Glu  
 290 295 300  
 His Glu Gln Lys Val Ala Arg Trp Tyr Asn Leu Ala Ile Ser Glu Ser  
 305 310 315 320  
 Ile Glu Thr Leu Ser Leu Ala Pro Phe Ser Arg Ile Phe His Trp Asp  
 325 330 335  
 Leu Asp Arg Ile Arg Gln Ile Thr Ala Glu Val Lys Ser Gln Ala Phe  
 340 345 350  
 Asn Lys Glu Ile His Ala Tyr Asn Ile Leu His Ile Tyr Gln Ala Arg  
 355 360 365  
 Lys Pro Gly Gly Pro Ser Leu  
 370 375

<210> SEQ ID NO 60  
 <211> LENGTH: 6411  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: transgene to complement alg3delta mutant

<400> SEQUENCE: 60

gtttaaacat gcatcattct cccgctttgt ttttgggccc aaactaaccg agtaggtgtg 60  
 gcatttgccg gcatgatggt tcaactaccg ctgatcatta tcaccgcccc attagagaag 120  
 atccaagacc ctactgggaa ggtgataggc aattccattt tctgggtag tttttgtctt 180  
 gtcggccagc ctttgggagc tttgctgtac ttctttgctt ggcaagcgaa gtatggcagt 240  
 gtgagccgaa tgtgaatgta aaagcacgca cgtgtccgct gtttgtcata gatgtaaata 300  
 aatgccaaca acttcagcca ttttttgaag agcaaagcaa ccgaagtaaa cgatcctgta 360  
 ccatcagcgc tcctcacaat ggaatctttt agatgtttct gttccattca tcttgcttac 420  
 tgcaatgttc ttttcgctt tgactaattc tccggatggt gaatggcaac gctgtcggcg 480  
 tcgggtcttc agggatccgc caaggatgct ctggatccgc atccggccgc tcttgccgcc 540  
 catcaatcgc ccgactataa atcgaactac tttcggcatc ttctagactt cctaataaccg 600  
 cctagtcata gcagattcaa gctgagaaca ccacaagtaa atatcaccca tcatgcttac 660  
 cctgaccgtc cctgaaaact acgggtatgt gccaaattcta caattccttg cagacaatgc 720  
 catttcccc atgaagtctg atgctaacta tcctgcagct ctgtcattgc cgtcgtctctg 780  
 ggtgccatcc ccgtcctgag cttcgtccat ggcgcccgtc tgtctcgtct ccgcaaggaa 840



-continued

---

gctgattgcc	cctaccctca	ctgctatgcg	accgtagagc	agtgcaagac	caacgtaagc	900
caacctcaca	caaacaggat	tcctcgagct	aacatacatt	cgaaccgtg	cagcccaagg	960
ccgagcagtt	caactgcgct	cagcgcgctc	atgccaaactt	ccttgagaac	tccagccaaa	1020
ctatgctctt	cctcctggta	gctggactga	agtacccccca	gttggcgact	ggcctcggaa	1080
gcatctgggt	cctcggtcgc	tcaactgttcc	tttacggata	tgtgtactcc	ggcaagccgc	1140
ggggtcgcgg	tcgtttgtac	ggcagcttct	acttgcttgc	acagggagct	ctctggggct	1200
tgacgtcttt	tggagttgcg	agggagttga	tttcctactt	ctaagtttgg	actgaatccg	1260
tggtgtgatt	gaggtgattg	gcgatgtcgt	gaccggtgac	tctttctggc	atgcccagag	1320
acggacggac	gcagagagaa	gggctgagta	ataagccact	ggccagacag	ctctggcggc	1380
tctgaggtgc	agtggatgat	tattaatccg	ggaccggccg	cccctccgcc	ccgaagtgga	1440
aaggtcggg	tgcccctcgt	tgaccaagaa	tctattgcat	catcggagaa	tatggagctt	1500
catcgaatca	ccggcagtaa	gcgaaggaga	atgtgaagcc	aggggtgtat	agccgtcggc	1560
gaaatagcat	gccattaacc	taggtacaga	agtccaattg	cttccgatct	ggtaaaagat	1620
tcacgagata	gtaccttctc	cgaagtaggt	agagcgagta	cccggcgcgt	aagctcccta	1680
attggcccat	ccggcatctg	tagggcgctc	aaatatcgtg	cctctcctgc	tttgcccggg	1740
gtatgaaacc	ggaaaggccg	ctcaggagct	ggccagccgg	gcagaccggg	aacacaagct	1800
ggcagtcgac	ccatccggg	ctctgcactc	gacctgctga	ggccctcag	tcctggtag	1860
gcagctttgc	cccgtctgtc	cgcccgggtg	gtcggcgggg	ttgacaaggt	cggtgcgtca	1920
gtccaacatt	tgttgccata	ttttcctgct	ctccccacca	gctgctcttt	tcttttctct	1980
ttcttttccc	atcttcagta	tattcatctt	ccatccaag	aacctttatt	tcccctaagt	2040
aagtactttg	ctacatccat	actccatcct	tcccatccct	tattcctttg	aacctttcag	2100
ttcgagcttt	cccacttcat	cgcagcttga	ctaacagcta	ccccgcttga	gcagacatca	2160
ccatggccaa	gttgaccagt	gccgttccgg	tgctcaccgc	gcgcgacgtc	gccggagcgg	2220
tcgagttctg	gaccgaccgg	ctcgggttct	cccgggactt	cgtggaggac	gacttcgccc	2280
gtgtggtccg	ggacgacgtg	accctgttca	tcagcgcggt	ccaggaccag	gtggtgccgg	2340
acaacaccct	ggcctgggtg	tgggtgcgcg	gcctggacga	gctgtacgcc	gagtggtcgg	2400
aggtcgtgtc	cacgaacttc	cgggacgcct	ccgggccggc	catgaccgag	atcggcagac	2460
agccgtgggg	gcgggagttc	gccctgcgcg	accggccggg	caactgctg	cacttcgtgg	2520
ccgaggagca	ggactgaccg	acgccgacca	acaccgcccg	tccgacggcg	gcccacgggt	2580
cccaggagct	tgagatccag	gagcaggact	gaccgacgcc	gaccaacacc	gccggctcca	2640
cgcggccccga	cgggtccgag	gcctcggaga	tccgtcccc	ttttcctttg	tcgatatcat	2700
gtaattagtt	atgtcacgct	tacattcacg	ccctcccccc	acatccgctc	taaccgaaaa	2760
ggaaggagtt	agacaacctg	aagtctaggt	ccctatttat	ttttttatag	ttatgttagt	2820
attaagaacg	ttatttatat	ttcaaatttt	tctttttttt	ctgtacagac	gcgtgtacgc	2880
atgtaacatt	atactgaaaa	ccttgcttga	gaaggttttg	ggacgctcga	agccgatgaa	2940
gaggtgtcta	cgttcctaca	attggaagaa	gaagcacaga	gattggccaa	ggcaggccac	3000
gagtgtcctg	ttcccagcc	tcgagggatt	ctgggcgaat	tgctgggttt	tacgagtagt	3060
ggaggtattt	cgacatcaac	gacgacacag	gcacaacagg	ttggagaggg	tcggtaacccg	3120
acaattggaa	agcaaggagg	actcaatcaa	ggctaaaata	ggctttgagc	atgtacagcg	3180

-continued

tgaaggaact	gtggattata	tcacgaaata	agccatgagg	cagtcgtgtt	ggcttcgacg	3240
gaggttggt	tcggaaattt	tcggccgggc	accaaaccg	acccatggca	gcaatatacc	3300
catctttgta	tcgatagtgt	acttacaaaa	actgtctact	atatgattat	gcatatgcga	3360
ttaaatacaa	ctctcaattg	atgcacaatt	cgctcaact	tctatggtaa	cgaaccacc	3420
tgttctgcag	acatgcggcc	gcgcggtctg	tgtgtgtcca	tgaacgctaa	tcccaaacgg	3480
gacagctctc	attggctctc	cggcacgaag	aggccaccg	acatctacag	ctgtagaaga	3540
aagtagctgg	ttcaacaacc	gcatgcaaga	tggactggat	gcgcctaatt	cgcgatttgt	3600
gtttcaatcc	ccgacacaca	aatggatgg	ctccgctcct	ggcctgggt	gacgctttcc	3660
tctgcgcgct	gatcatctgg	aaagtgcct	gtaaggctac	agctaagctc	cgttcacacc	3720
cttttgcgac	aagtgaagca	atgccactaa	cctagccccg	ttgctattgt	ccacagatac	3780
cgagattgac	tgggccacgt	acatgcaaca	aatatcgctt	tatttgcag	gagaacgcga	3840
ttatactctc	atcagaggat	caaccggctc	ccttgtctac	ccggccgccc	atgtatacag	3900
ttatacggcc	ctctaccatc	tcaccgatga	ggggcgcgat	atcttcttcg	gtcagatact	3960
atctgtctg	ctctacttga	tcacgctgg	ggtgtgctg	tgctgttata	gacagtcggg	4020
tgctccgccc	tacttgcttc	cgctgctgg	cctttccaag	agacttcaca	gcgtttatgt	4080
cctgcgctg	ttcaatgatg	gcttggcggc	gctggcgatg	tgggttgcca	ttctgttatt	4140
catgaatcgg	aagtggacgg	ctgcggctgc	agtgtggtct	actggtgttg	cgattaagat	4200
gacactgctg	ctgctggccc	cgctattgc	tgtggtcacg	gtgcttagtc	tgctgcttgg	4260
tcctagcgtg	gggctggggg	ttctggcgg	gctgtccag	gtaggttccc	atgaggctgt	4320
agggttgccc	aaaggcaatt	tgtgtgaaga	cttgtctgac	attgaactac	aggttttact	4380
cgcgataccg	ttcctacaaa	acaaccggc	gggtatctc	tcgcgggcgt	tcgagctaac	4440
cagacagttc	atgtttaa	ggacagtcaa	ttggagattt	gttggcgaag	aagtattctt	4500
atctaagagc	tttccctgg	cattgctggc	cgccacatt	gtgctgctag	gcgcttttgc	4560
cgctactgg	tggctgagat	actccaggtc	tagcttgcc	gcgttcattc	ggaatctgct	4620
agcgggtcga	catcgcacag	tgtccctccc	caaaccctac	atcatgagcg	tgatgctctc	4680
gtctctgaca	gttggcttgt	tgtgcgcaag	gtccctcat	taccaattct	tcgctacct	4740
ctcctggg	acacccttc	tcctctggc	cgagggtt	catccaatct	tgctgtacct	4800
tatctgggct	atgcaagagt	ggccttgaa	cacattcccc	agcaccaacc	tcagttccat	4860
cattgttgct	ctctcaactg	ctaccagag	ttcggcgctc	cttgcgaata	gtgccagcgc	4920
ctttataacc	atgcgttcga	accctagcgg	taaagagcat	aaccaataga	agtgacaccc	4980
ggccagtatc	gagatcggc	tgtgacaggt	gcatcgataa	tcgcaatcag	tcttgtaccc	5040
atgagaatcc	ctgaaaaagt	aagactgctc	tgtcaggtag	tccattgccc	atgcgatagg	5100
ttcggacgcc	taaaggatca	atcaagatgc	caatcaagca	tccgactcat	cggaagaagg	5160
catcttgccg	acattggact	catcctcttc	gtccgagtcg	tcggcgacaa	cagcagcttg	5220
cttagcgatg	gtgtggcaca	aggatcaatg	cggtacgacg	atctgatgca	gataagcagg	5280
ctgcgaagta	gtaactcttg	cgtagagaaa	atggcgacgg	gtggctgata	agggcgggtga	5340
taagcttaat	tgtcatcgca	gataagcact	gctgtcttgc	atccaagtca	gcgtcagcag	5400
aaatacggga	cttccgaaag	tatatggcaa	aattaaagaa	cttgactctc	cagcaatggt	5460
ttgcctgac	cgctgctaaa	acgttactac	ccctataccc	gtctgtttgt	cccagcccga	5520
ggcattaggt	ctgactgaca	gcacggcgcc	atgcgggctt	gggacgcat	gtccgtcgcg	5580

-continued

---

```

tgataagggg tgatccatgc agctactatc cttccatcgt tccattccca tccctgtcct 5640
atctccatcc ttgaaacttt actagtttag ttggatgctc gagcttctc tcggctactc 5700
cgtccaatgg ataagacccc gatgccggtc ctcatggctc tccagctggg atcgccccaa 5760
ccttcgtgtg atcgccctc tgcttcccct catcatcatt actaactagt acatccaaaa 5820
gccatcccag tgcttcccct cacccttgcc caagacattc caagtgggcc ttcggctgga 5880
aaacatggac ccattggttc catcgataag ctagctctc gtccgttacc ccagattgat 5940
accagataac attgaccagc ggcttatcac cgaggtctgc ggggtgagacc ccccctgcga 6000
caagttagat aaaagaaact cgcctcattg tgcttccgat ggggtcggat gacgagcctt 6060
cggaaagagc tggcgctct ttaaagggga cagctgtcgc caagttgtga aattctccga 6120
taactactaa caatctctcc cttccttccc gctactgtgg tcaccaaata aactctcttt 6180
tctcgcccaa gatctaacat ggcggtatgag aagactgaaa agtctcccc accgatgacg 6240
gtggatgagg agactggcac aacagaggaa attgaccoga caatggcaaa gcatacgaag 6300
gatgcagacg aggcactggc ggtcttcgaa gacctccatg gtgaagtcac cacacttgat 6360
gaggagacaa acaaaaggat acttcggaca attgactggc acagtttaaa c 6411

```

```

<210> SEQ ID NO 61
<211> LENGTH: 899
<212> TYPE: DNA
<213> ORGANISM: Aspergillus niger

```

```

<400> SEQUENCE: 61

```

```

tcaagacagc ggctgcaaca accttaaagg cagggttget ttgctttttg ccttctttcc 60
tctgcctctc attccacccc cctctctctc tccccctctc ccgctgtccg aatggacttg 120
cacttcoggg atcggatttt cctttctga tccgatgatc cagtccaatt gacctaacctc 180
tcgatcgtcc tccccgatc catggttggt ctttagtgcg ctctctctc gtaacgcctg 240
acacctgctg ctgttatcga taccctattt attatttatt atttattttt tttccctatt 300
ttcctatgcc gctgcccgtga ccgctacttt tctcttttgt tctctcccc ccgacggggt 360
cctgtccttg cggggaggag acttttttcc ccctacctat ttagtatcct cattcctcct 420
tgtgcctcgg ccgagatttg gagatactct tctccaccgt gtgtgactgt ggggtgtgcgc 480
atgcacgggt gtgctgcct ctttgctctc atacaactct cctgggttgg ggcaaattta 540
caccattgtg gactttgat cattcttact tactatcctt ggctttttgc tctccggctt 600
tccactctcc tcttcttcc aatcctttta cccctctctg gccactgatt cattccgagg 660
aggatacccc cagtttagtag ttttggtggt gccgtctttt cgtctgatca gcttttcaat 720
ccaatcattg gctcaatttg cggcgccgaa ctccaacttc ctctatgtgc cacctgactt 780
acgattcccc gatatcacct gcagcctgca tacaccctgt tggctaacat tggcgtttta 840
cgttaccgctc acgaagaacg ccggtctctt tgattgaacg aaccctgta tctctacca 899

```

```

<210> SEQ ID NO 62
<211> LENGTH: 1012
<212> TYPE: DNA
<213> ORGANISM: Aspergillus niger

```

```

<400> SEQUENCE: 62

```

```

tcagtcgcat ctttactgg gatctggata ggatcagaca gatcacagcg gaggtcaagt 60
cacaagcctt caacaaggaa atccacgctt acaatatctt acatatatac caggcacgga 120

```

-continued

---

```

agccgggccc cccatcactt tgaaagtaca gagaaaatta cggcagtgcc ggctgtaatg 180
cagatttcat tccggatacc catatgcagc cttttgtggg agggccatcc tacctgttct 240
cttctttttg ttcattccat ttttttctcg atgaggatta gtgacgacca attccatctc 300
cttgacggga tcatactgaa atgcttatac accaaagcga gcaaagccca caaaaccatc 360
actggacttg aactcgtacc aacactgcta gtttgacgca agaatgccag gaacgcagac 420
aggcttatct tcattgtgtt attgcctact ttagtgacga cagctgaacg gcacaaaggg 480
acgcgaagta tattatcaat cacaccatgt ggattgttac cgtcgcaaaa gatactgctc 540
gaggttttaa ataatagaag cacttcaaga agatgagggtg ccggttgcgg gactcgaaga 600
agtggccgac gactataaag taccagattt gccaggatcc aaaagaagct accaatctca 660
tgttgcatth ttgaaagcc ctgtaattca ggccggctga actgcggaga tcccagaatc 720
atggcagttg actgagtact tgctgtgacg ggccctaaac atcagccatt ccatccgaac 780
tcaaaactct ctgatgaccg tggcggctct gtggcacagc caccgataat tttttttggt 840
ggcacttatt aatgaattag atgagaaaga agcgaagaac cgagggcttg aacaaacgca 900
aagccctcca aatggttact attgaacttt ttgctctccc accgaaactc ccaactgatca 960
tgtaccgtaa gtatatcgaa aaatatgcca tgtcccttga gacagcctac ca 1012

```

&lt;210&gt; SEQ ID NO 63

&lt;211&gt; LENGTH: 1360

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic polynucleotide (hph expression cassette)

&lt;400&gt; SEQUENCE: 63

```

gctggagcta gtggaggtca acacatcaat gctatthttgg tttagtcgtc caggcggatc 60
acaaaatttg tgctgthtga caagatggth catttaggca actggtcaga tcagcccact 120
tgtaagcagt agcggcggcg ctgcaagtgt gactcttatt agcagacagg aacgaggaca 180
ttattatcat ctgctgcttg gtgcacgata acttgthgctt ttgtcaagca aggtaagtga 240
acgaccgggt cataccttct taagttcgcc cttcctccct ttatttcaga ttcaatctga 300
cttacctatt ctaccaagc atccaaatgc ctgaaactca cgcgacgtct gtcgagaagt 360
ttctgatcga aaagttcgac agcgtctccg acctgatgca gctctcggag ggccaagaat 420
ctcgtgctth cagcttcgat gtaggagggc gtggatatgt cctgcccggta aatagctgcg 480
ccgatggtht ctacaaagat cgthtatgtht atcggcactt tgcacggcc gcgctcccga 540
ttccggaagt gcttgacatt ggggagthtca gcgagagcct gacctattgc atctcccgcc 600
gtgcacaggg tgthcacgtg caagacctgc ctgaaaccga actgcccgtt gthctggagc 660
cggthcggga ggccatggat gcgatcgtg cggccgatct tagccagacg agcgggthctg 720
gcccattcgg accgcaagga atcggthcaat aactacatg gcgthgattc atatgthcga 780
thgctgatcc ccatgthgat cactggcaaa ctgthgatgga cgacaccgtc agthcgtccg 840
thcgcagggc thctgatgag ctgatgctth gggccgagga ctgccccgaa gthcggcacc 900
thcgtcacgc ggattthcgc thcaacaatg thctgacgga caatggccgc ataacagcgg 960
thattgactg gacgagggcg atgthcgggg attthcaata cgaggtcggc aacatcttht 1020
thctggagggc gthgthgthct thgatggagc agcagacgcg ctactthcag cggagghatc 1080
cggagctthc aggatcggcg cgctcccggg cgtatathct ccgcatthgt ththgaccaac 1140

```

-continued

---

tctatcagag	cttggttgac	ggcaatttcg	atgatgcagc	ttgggcgag	ggtcgatgag	1200
acgcaatcgt	ccgatccgga	gccgggactg	tcgggcttac	acaaatcgcc	cgcagaagcg	1260
cggccgtctg	gaccgatggc	tgtgtagaag	tactcgccga	tagtggaaac	cgacgcccc	1320
gcactcgtcc	gagggcaaag	gaatagagta	gatgccgacc			1360

&lt;210&gt; SEQ ID NO 64

&lt;211&gt; LENGTH: 2340

&lt;212&gt; TYPE: DNA

<213> ORGANISM: *Aspergillus niger*

&lt;400&gt; SEQUENCE: 64

tctctcctcg	taacgcctga	cacctgctgc	tgttatcgat	accctattta	ttatttatta	60
tttatttttt	ttccctattt	tcctatgccg	ctgccgtgac	cgctactttt	ctcttttggt	120
ctctcccccc	cgacgggttc	ctgtccttgc	ggggaggaga	cttttttccc	cctacctatt	180
tagtatactc	attcctcctt	gtgcctcggc	cgagatttgg	agatactctt	ctccaccgtg	240
tgtgactgtg	gggtgtgcga	tgcacgggtg	tgcgtgcctc	tttgctctca	tacaactctc	300
ctgggttggg	gcaaatttac	accattgtgg	actttggatc	attcttactt	actatccttg	360
gctttttgct	ctccggcttt	ccactctcct	ccttcttcca	atccttttac	cccttctcgg	420
ccactgattc	attccgcgga	ggatacccc	agttagtagt	tttgtggttg	ccgtcttttc	480
gtctgatcag	cttttcaatc	caatcattgg	ctcaatttgc	ggcgccgaac	tccaacttcc	540
tctatgtgcc	acctgactta	cgattccccg	atatcacctg	cagcctgcat	acacctggtt	600
ggtaacatt	ggcgttttac	gttaccgtca	cgaagaacgc	cggtctcttt	gattgaacga	660
acctctgtat	ctctaccata	ccttgagcag	gaaaagtcga	atctcttccc	agcgaaccga	720
tccttggaact	tcagggttcc	atgtttgaaa	tcagccgact	tttgcacag	ccaattacta	780
tggcttccgc	gaatcgcaat	aactacagct	accaagggat	agaatcctat	gattccggcc	840
gttccaggca	aaactcggat	gctatggaca	ttcacgtcat	tacggcccaa	gaacctctc	900
gagaaccccc	ggacaacaac	gatccttatg	atggccatgg	gggtccagct	gggactagcc	960
attatagcaa	gtacttctcc	cttctcatac	tctgcacccc	acgtaccccc	caaaatccct	1020
ttttctcatg	ccgtgcaaat	atcacactta	tttctacaac	taccgggcca	ctaattcagg	1080
gaactttctt	ttccgttggt	cgtttaatct	aggcctccaa	acagatggct	cttctatgaa	1140
gaaaatgggc	gaacatatca	tggatatcgc	agaggagttt	acctcgctgc	atgcatgaa	1200
caggaaacagg	accgtctcga	tatcttccat	aaactgttca	cagtagcacg	gatgtccgag	1260
agcttaatct	acgcacctca	ccccccaaat	ggtcgattcc	tagatctggg	gtgcggcact	1320
gggatctggg	ccattgatgt	agcccacaag	tatcccgaatg	ctttcggtgc	tggagtagat	1380
ctagcaccta	tacagcctcc	caaccacccc	gataactgag	agttctatgc	accttttgac	1440
tttgaggcgc	catggacgct	tggggaaaat	tcttgggatc	tcattcatct	acagatgggt	1500
tgcggcagtg	ttctgggctg	gcagaatctc	tacaagcgaa	tcttaaggca	tcttcagcct	1560
ggggcatggg	ttgaacaggt	ggaaatagat	ttcgaacccc	gctgcgatga	tcgctccctg	1620
aatggactgg	cactccggga	gtggtaccag	tacctgaagc	aggcgacaca	agatacaatg	1680
cgacccatag	cgcacagctc	gcgggatacc	atcagacacc	ttgaggaggc	aggctttacc	1740
cagatcgacc	atcagatggt	ggggctgect	ctcaaccctt	ggcaccgtga	tgaacatgag	1800
cagaaggtag	cccgttggtg	taacctcgca	atctctgaga	gtatcgagac	gctcagcctc	1860
gcccctttca	gtcgcactct	tcactgggat	ctggatagga	tcagacagat	cacagcggag	1920

-continued

---

```

gtcaagtcac aagccttcaa caaggaaatc cacgcttaca atatcttaca tatataccag 1980
gcacggaagc cgggeggccc atcactttga aagtacagag aaaattacgg cagtggcggc 2040
tgtaatgcag atttcattcc ggatacccat atgcagcctt ttgtgggagg gccatcctac 2100
ctgttctctt ctttttgttc attccatttt tttctcgatg aggattagtg acgaccaatt 2160
ccatctcctt gacgggatca tactgaaatg cttatacacc aaagcgagca aagcccacaa 2220
aaccatcact ggacttgaac tcgtaccaac actgctagtt tgacgcaaga atgccaggaa 2280
cgcagacagg cttatcttca ttgtgttatt gcctacttta gtgcagacag ctgaacggca 2340

```

&lt;210&gt; SEQ ID NO 65

&lt;211&gt; LENGTH: 3306

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic polynucleotide (laeA deletion cassette)

&lt;400&gt; SEQUENCE: 65

```

aagcttcaag acagcggctg caacaacctt aaaggcaggg ttgctttgct ttttgccttc 60
tttctctgct cctctattcc acccccctc tctcttccc cctctccgct gtccgaatgg 120
acttgcaact cgggatcgg attttccctt tctgatccga tgatccagtc caattgacct 180
aactctcgat cgtcctccc cgatccatgg ttgttcttta gtcgcctctc tctctgtaac 240
gcctgacacc tgctgctgtt atcgataccc tatttattat ttattattta ttttttttcc 300
ctattttctt atgccgctgc cgtgaccgct acttttctct tttgttctct ccccccgac 360
gggttctctg ccttgccggg aggagacttt tttccccta cctatttagt atcctcattc 420
ctccttgctc ctggccgag atttgagat actcttctcc accgtgtgtg actgtgggtg 480
tgcgatgca cgggtgtgcg tgectctttg ctctcataca actctcctgg gttggggcaa 540
atttacacca ttgtggactt tggatcattc ttacttacta tccttggctt tttgctctcc 600
ggctttccac tctcctcctt cttccaatcc ttttaccct tctcggccac tgattcattc 660
cgcggaggat acccccagtt agtagttttg tggttgctt cttttcgtct gatcagcttt 720
tcaatccaat cattggctca atttgccgcg ccgaactcca acttcctcta tgtgccacct 780
gacttacgat tccccgatat cacctgcagc ctgcatacac cctggttggct aacattggcg 840
ttttacgtta ccgtcacgaa gaacgccggg ctctttgatt gaacgaacc ctgtatctct 900
accacggctg gcatctactc tattcctttg ccctcggagc agtgctgggg cgtcgggttc 960
cactatcggc gagtacttct acacagccat cggtcagac ggcgcgctt ctgcgggcca 1020
tttgtgtacg cccgacagtc ccgctccgg atcggacgat tgcgtcgcat cgacctgcg 1080
cccaagctgc atcatcgaat ttgccgtaa ccaagctctg atagagttgg tcaagaccaa 1140
tgcggagcat atacgcccg aggccggcg atcctgcaag ctccggatgc ctccgctcga 1200
agtagcggct ctgctgctcc atacaagcca accacggcct ccagaagaag atgttggcga 1260
cctcgtattg ggaatccccg aacatcgctt cgtccagtc aatgaccgct gttatgcggc 1320
cattgtccgt caggacattg ttggagccga aatccgctg cacgaggtgc cggacttcgg 1380
ggcagtcctc ggcccaaagc atcagctcat cgagagcctg cgcgacggac gcaactgacg 1440
tgtcgtccat cacagtttgc cagtataca catggggatc agcaatcgcg catatgaaat 1500
cacgccatgt agtgattga ccgattcctt gcggtccgaa tgggcccgaac ccgctcgtct 1560
ggctaagatc ggccgagcg atcgcattca tggcctccgc gaccggctcc agaacagcgg 1620

```

-continued

---

```

gcagttcggg ttcaggcagg tcttgcaacg tgacaccctg tgcacggcgg gagatgcaat 1680
aggtcaggct ctcgctgaac tccccaatgt caagcacttc cggaatcggg agcgcgggccc 1740
atgcaaagtg ccgataaaca taacgatctt tgtagaaacc atcggcgcag ctatttacc 1800
gcaggacata tccacgccct cctacatcga agctgaaagc acgagattct tcgccctccg 1860
agagctgcat caggtcggag acgctgtcga acttttcgat cagaaacttc tcgacagacg 1920
tcgcggtgag ttcaggcatt tggatgcttg ggtagaatag gtaagtcaga ttgaatctga 1980
aataaagggg ggaagggcga acttaagaag gtatgaccgg gtcgttcact taccttgctt 2040
gacaaacgca caagttatcg tgcaccaagc agcagatgat aataatgtcc tcgttcctgt 2100
ctgctaataa gagtcacact tcgagcggcc cgcctactgc ttacaagtgg gctgatctga 2160
ccagttgcct aatgaacca tcttgcaaaa cgacacaaat tttgtgatcc gcctggacga 2220
ctaaacccaa atagcattga tgtgttgacc tccactagct ccagctcagt cgcacttttc 2280
actgggatct ggataggatc agacagatca cagcggaggc caagtcacaa gccttcaaca 2340
aggaaatcca cgcttacaat atcttacata tataccaggc acggaagccg ggccggccat 2400
cactttgaaa gtacagagaa aattacggca gtgccggctg taatgcagat ttcattccgg 2460
ataccatata gcagcctttt gtgggagggc catcctacct gttctctctt tttgttcat 2520
tccatttttt tctcgtatgag gattagtgc gaccaattcc atctccttga cgggatcata 2580
ctgaaatgct tatacaccia agcagcaaaa gccacaaaaa ccatcactgg acttgaactc 2640
gtaccaacac tgctagttagt acgcaagaat gccaggaacg cagacaggct tatcttcatt 2700
gtgttattgc ctactttagt gcagacagct gaacggcaca aagggacgcg aagtatatta 2760
tcaatcacac catgtggatt gttaccgtcg caaaagatac tgctcgaggt tttaaataat 2820
agaagcactt caagaagatg aggtgccgtt gtcgggactc gaagaagtgg ccgacgacta 2880
taaagtacca gatttgccag gatccaaaag aagctaccaa tctcatgttg catttttgga 2940
aagccctgta attcaggccg gctgaactgc ggagatccca gaatcatggc agttgactga 3000
gtacttgctg tgacgggccc taaacatcag ccattccatc cgaactcaaa actctctgat 3060
gaccgtggcg gctctgtggc acagccaccg ataatttttt ttggtggcac ttattaatga 3120
attagatgag aaagaagcga agaaccgagg gcttgaacaa acgcaaagcc ctccaaatgg 3180
ttactattga actttttgct ctcccaccga aactcccact gatcatgtac cgtaagtata 3240
tcgaaaaata tgccatgtcc cttgagacag cctaccatgc agcccggggg atccactagt 3300
tctaga 3306

```

&lt;210&gt; SEQ ID NO 66

&lt;211&gt; LENGTH: 7163

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic polynucleotide (LaeA complementation construct)

&lt;400&gt; SEQUENCE: 66

```

gtttaaacad gcatcattct cccgctttgt tttggggccc aaactaacgg agtaggtgtg 60
gcatttgccg gcatgatgtt tcaactaccg ctgatcatta tcaccgcccc attagagaag 120
atccaagacc ctactgggaa ggtgataggc aattccattt tctgggtag tttttgtctt 180
gtcggccagc ctttgggagc tttgctgtac ttctttgctt ggcaagcgaa gtatggcagt 240
gtgagccgaa tgtgaatgta aaagcacgca cgtgtccgct gtttgcata gatgtaata 300

```

-continued

---

aatgccaaca	acttcagcca	ttttttgaaa	agcaaagcaa	ccgaagtaaa	cgatcctgta	360
ccatcagcgc	tcctcacaat	ggaatctttt	agatgtttct	gttccattca	tcttgcttac	420
tgcaatgttc	ttttcgcggt	tgactaattc	tccggatggt	gaatggcaac	gctgtcggcg	480
tcgggtcttc	agggatccgc	caaggatgct	ctggatccgc	atccggccgc	tcttgcgccc	540
catcaatcgc	ccgactataa	atcgaactac	tttcggcacc	ttctagactt	cctaataaccg	600
cctagtcata	gcagattcaa	gctgagaaca	ccacaagtaa	atatcaccca	tcatgcttac	660
cctgaccgtc	cctgaaaact	acgggtatgt	gccaaattcta	caattccttg	cagacaatgc	720
cattctcccc	atgaagtctg	atgctaacta	tcttcagact	ctgtcattgc	cgtcgctctg	780
ggtgccatcc	ccgtcctgag	cttcgtccat	ggcgccgctg	tgtctcgtct	ccgcaaggaa	840
gctgattgcc	cctaccctca	ctgctatgcg	accgtagagc	agtgcaagac	caacgtaagc	900
caacctcaca	caaacaggat	tcctcgagct	aacatacatt	ccgaaccgtg	cagcccaagg	960
ccgagcagtt	caactgcgct	cagcgcgctc	atgccaaact	ccttgagaac	tccagccaaa	1020
ctatgctctt	cctcctggta	gctggactga	agtaccccc	gttggecact	ggcctcggaa	1080
gcatctgggt	cctcggtcgc	tcaactgttc	tttacggata	tgtgtactcc	ggcaagccgc	1140
ggggtcgcgg	tcgttgttac	ggcagcttct	acttgcctgc	acagggagct	ctctggggct	1200
tgacgtcttt	tggagttgcg	agggagttga	tttcctactt	ctaagtttgg	actgaatccg	1260
tgggtgtgatt	gaggtgattg	gcgatgatgc	cgttcagctg	tctgcactaa	agtaggcaat	1320
aacacaatga	agataagcct	gtctgcgctc	ctggcattct	tgcgtcaaac	tagcagtgtt	1380
ggtacgagtt	caagtccagt	gatggttttg	tgggctttgc	tcgctttggt	gtataagcat	1440
ttcagtatga	tcccgtcaag	gagatggaat	tggctgtcac	taatcctcat	cgagaaaaaa	1500
atggaatgaa	caaaaagaag	agaacaggta	ggatggccct	cccacaaaag	gctgcatatg	1560
ggtatccgga	atgaaatctg	cattacagcc	ggcaactgcc	taattttctc	tgtactttca	1620
aagtgatggg	ccgcccggct	tccgtgcctg	gtatatatgt	aagatattgt	aagcgtggat	1680
ttccttggtg	aaggcttggt	acttgacctc	cgctgtgac	tgtctgatcc	tatccagatc	1740
ccagtgaaag	atgcgactga	aaggggagag	gctgagcgtc	tcgatactct	cagagattgc	1800
gaggttatac	caacgggcta	ccttctgctc	atgttcatca	cggtgccaaag	ggttgagagg	1860
cagccccacc	atctgatggt	cgatctgggt	aaagcctgcc	tcctcaaggt	gtctgatggt	1920
atcccgcgag	ctgtgcgcta	tgggtcgcac	tgtatcttgt	gtcgcctgct	tcaggactctg	1980
gtaccactcc	cggagtgcca	gtccattcag	ggagcgatca	tcgcagcggg	gttcgaaatc	2040
tatttccacc	tgttcaaacc	atgccccagg	ctgaagatgc	cttaagattc	gcttgtagag	2100
attctgccag	cccagaacac	tgcgcgaacc	catctgtaga	tgaatgagat	cccagaatt	2160
ttccccaaag	gtccatggcg	cctcaaagtc	aaaaggtgca	tagaactcgc	agttatcggg	2220
gtggttggga	ggctgtatag	gtgctagatc	tactccagca	acgaaagcat	tgggatactt	2280
gtgggctaca	tcaatggccc	agatcccagt	gccgcacccc	agatctagga	atcgaccatt	2340
tggggggtga	ggtgcgtaga	ttaagctctc	ggacatccgt	gctactgtga	acagtttatg	2400
gaagatatcg	agacggtcct	gttcctgttc	atcgcattgc	agcgggtaaa	ctcctctgcg	2460
atatccatga	tatgttcgcc	cattttcttc	atagaagagc	catctgtttg	gaggcctaga	2520
ttaaaccgaac	aacggaaaag	aaagttccct	gaattagctg	cccggtagtt	gtagaaataa	2580
gtgtgatatt	tgcacggcat	gagaaaaagg	gattttgccc	ggtacgtggg	gtgcagagta	2640



-continued

---

tgagaagggga	gaagtacttg	ctataatggc	tagtcccagc	tggacccccca	tggccatcat	2700
aaggatcggt	gttgtccggg	ggttctcgag	gaggttcttg	ggccgtaatg	acgtgaatgt	2760
ccatagcatc	cgagttttgc	ctggaacggc	cggaatcata	ggattctatc	ccttggtagc	2820
tgtagttatt	gcgattcggc	gaagccatag	taattggctg	atgcaaaagt	cggtcgattt	2880
caaacatgga	accctgaagt	ccaaggatcg	gttcgctggg	aagagattcg	acttttctctg	2940
ctcaaggtat	ggtagagata	caggggttcg	ttcaatcaaa	gagaccggcg	ttcttcgtga	3000
cggtaacgta	aaacgccaat	gtagccaac	aggggtgatg	caggctgcag	gtgatatcgg	3060
ggaatcgtaa	gtcaggtggc	acatagagga	agttggagtt	cggcgccgca	aattgagcca	3120
atgattggat	tgaaaagctg	atcagacgaa	aagacggcaa	ccacaaaact	actaactggg	3180
ggtatcctcc	gcggaatgaa	tcagtggccg	agaaggggta	aaaggattgg	aagaaggagg	3240
agagtggaaa	gccggagagc	aaaaagccaa	ggatagtaag	taagaatgat	ccaaagtcca	3300
caatggtgta	aatttgcccc	aaccaggag	agttgtatga	gagcaaagag	gcacgcacac	3360
ccgtgcatgc	gcacaccac	agtcacacac	ggtggagaag	agtatctcca	aatctcggcc	3420
gaggcacaag	gaggaatgag	gatactaaat	aggtaggggg	aaaaaagtct	cctccccgca	3480
aggacaggaa	cccgtcgggg	gggagagaac	aaaagagaaa	agtagcggtc	acggcagcgg	3540
cataggaaaa	tagggaaaaa	aaataaataa	taaataataa	atagggtatc	gataacagca	3600
gcaggtgtca	ggcgttacga	ggagagatgg	accgatggct	gtgtagaagt	actcgccgat	3660
agtggaaaacc	gacgccccag	cactcgtccg	agggcaaagg	aatagagtag	atgccgaccg	3720
cgggatccac	ttaacgttac	tgaaatcatc	aaacagcttg	acgaatctgg	atataagatc	3780
gttggtgtcg	atgtcagctc	cggagttgag	acaaatggtg	ttcaggatct	cgataagata	3840
cgttcatttg	tccaagcagc	aaagagtgcc	ttctagtgat	ttaatagctc	catgtcaaca	3900
agaataaaac	gcgttttcgg	gtttacctct	tccagataca	gctcatctgc	aatgcattaa	3960
tgattgact	gcaacctagt	aacgccttca	ggctccggcg	aagagaagaa	tagcttagca	4020
gagctatfff	cattttcggg	agacgagatc	aagcagatca	acggctcgtca	agagacctac	4080
gagactgagg	aatccgctct	ctgacagacg	ggcaattgat	tacgggatcc	cattggtaac	4140
gaaatgtaaa	agctaggaga	tcgtccggcg	atgtcaggat	gatttcactt	gtttcttgtc	4200
cggctcaccg	gtcaaagcta	aagaggagca	aaaggaaacg	atagaatcgg	gtgccgctga	4260
tctatacggg	atagtgcctt	tatcacggtg	actcaaccca	tgctatttaa	ctcaaccctt	4320
ccttctgaac	cccaccatct	tcttcttttt	cctctcatcc	cacacaattc	tctatctcag	4380
atftgaattc	caaaagtcct	cggacgaaac	tgaacaagtc	ttcctccctt	cgataaacct	4440
ttggtgattg	gaataactga	ccatcttcta	tagttcccaa	accaaccgac	aatgtaaata	4500
cactcctcga	ttagccctct	agagggcata	cgatggaagt	catggaatac	ttttggctgg	4560
actctcacia	tgatcaaggt	atcttaggta	acgtctttgg	cgtgggcccg	tgttcgttcc	4620
cagtcacoga	tgcattcaca	tgccctccct	aagctggggc	ctagacteta	ggatcctagt	4680
ctagaaggac	atggcatcga	tggactgggt	tcgttctgag	attatacggc	taaaacttga	4740
tctggataat	accagcgaac	agggatcatg	cttctctcgt	tcttctgtt	gatggaatgg	4800
ctaacagatg	atagtcatgt	caacttgaac	catgtctcct	ccagctgcca	tctacgaacc	4860
cactgtggcc	gctaccggcc	tcaagggtaa	ggctcgtggt	tctgagaccg	tccccgttga	4920
gggagcttct	cagaccaagc	tggtggacca	tttcggtggc	aagtgggacg	agttcaagtt	4980
cgccctatc	cgcgaaagcc	aggtctctcg	tgccatgacc	agacgttact	ttgaggacct	5040

-continued

---

```

ggacaagtac gctgaaagtg acgttgatcat tgttgggtgct ggttcctgcg gtctgagcac 5100
tgcgtacgctc ttggccaagg ctcgctccgga cctgaagatt gctatcgctcg aggccagcgt 5160
ctctcctgggt cagtagtcca tgatggattg ccttgcactc agctttccgg aactaacgtg 5220
caataggtgg cggtgccctgg ttgggtggcc aactcttttc tgctatggtc atgcgccgctc 5280
ccgcggaagt cttcctgaac gagctgggtg ttccttacga agaggacgca aaccccaact 5340
acgttgctcg caagcacgcc tccctgttta cctcgacact catgtcgaag gttctctcct 5400
tccccaatgt caagctcttc aatgctaccg ctggtgagga cttgatcacc cgtccgaccg 5460
agaacggcaa cccccagatt gctgggtgtg tcgtcaactg gacgctggtc acccttcacc 5520
acgatgatca ctctgcatg gaccccaaca ctatcaacgc tctgtcatc atcagtacca 5580
ctggtcacga tgggccattc ggcgccttct gtgcgaagcg cttgggtgtcc atgggcagcg 5640
tcgacaagct aggtggcatg cgtgggtctcg acatgaactc ggccgaggat gccatcgta 5700
agaacacccg cgaggttact aagggcttga taatcgccgg tatggagctg tctgaaattg 5760
atggctttaa ccgcatgggc cctaccttcg gtgccatggt tctcagtgggt gtcaaggctg 5820
ccgaggaggc attgaaggtg ttcgacgagc gtcagcgcga gtgtgctgag taaatgactc 5880
actaccgaa tgggttcagt gcatgaaccg gatttgtctt acggctcttg acgatagggg 5940
aatgatgatt atgtgatagt tctgagattt gaatgaactc gttagctcgt aatccacatg 6000
catatgtaaa tggctgtgtc ccgatgtaa cgggtgggca ttctagaata attatgtgta 6060
acaagaaaga cagtataata caaacaaga tgcaagagcg gctcgtcatc gcagataagc 6120
actgctgtct tgcattcaag tcagcgtcag cagaaatacg ggacttccga aagtatatgg 6180
caaaattaa gaacttgact ctccagcaat gttttgcct gaccgtcgtc aaaacgttac 6240
taccctata cccgtctgtt tgtcccagcc cgaggcatta ggtctgactg acagcacggc 6300
gccatgcggg cttgggacgc catgtccgct gcgtgataag ggttgatcca tgcagctact 6360
atccttccat cgttccattc ccaccttgt cctatctcca tccttgaaac tttactagtt 6420
tagttggatg ctcgagcttg ctctcgcta ctccgtccaa tggataagac cccgatgccg 6480
gtcctcattg gtctccagct ggtatcgccc caacctcgt gtgatcgctc ctctgcttcc 6540
cctcatcacc attactaact agtacatcca aaagccatcc cagtgtctcc cctcacctt 6600
gccaagaca ttccaagtgg gccttcggct ggaaaacatg gaccattgg ttccatcgat 6660
aagctagctc ctcgctcgtt accccagatt gataccagat aacattgacc agcggcttat 6720
caccgaggtc tgcgggtgag aacccccctg cgacaagtta gataaaagaa actcgcctca 6780
ttgtgcttcc gatgggtcg gatgacgagc ctcggaaag agctggcgcc tctttaagg 6840
ggacagctgt cgccaagttg tgaaattctc cgataactac taacaatctc tcccttctt 6900
cccgtactg tggtcaccaa atcaactctc tttctcggc caagatctaa catggcggat 6960
gagaagactg aaaagtctcc cccaccgatg acggtggatg aggagactgg cacaacagag 7020
gaaattgacc cgacaatggc aaagcatagc aaggatgcag acgaggcact ggcggctctc 7080
gaagacctcc atggtgaagt catcacactt gatgaggaga caaacaag gatacttcgg 7140
acaattgact ggcacagttt aac 7163

```

&lt;210&gt; SEQ ID NO 67

&lt;211&gt; LENGTH: 38

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

-continued

---

<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 67

gtcgcagcgtg tgcgataagct tcaagacagc ggctgcaa 38

<210> SEQ ID NO 68  
 <211> LENGTH: 28  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 68

gccgaccgtg gtagagatac aggggttc 28

<210> SEQ ID NO 69  
 <211> LENGTH: 29  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 69

ctctaccacg gtcggcatct actctattc 29

<210> SEQ ID NO 70  
 <211> LENGTH: 26  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 70

gcgactgagc tggagctagt ggaggt 26

<210> SEQ ID NO 71  
 <211> LENGTH: 27  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 71

gctccagctc agtcgcatct ttcactg 27

<210> SEQ ID NO 72  
 <211> LENGTH: 33  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 72

atcccccggtg ctgcatgta ggctgtctca agg 33

<210> SEQ ID NO 73  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 73

actcgcagca gagatgcat ct 22

-continued

---

<210> SEQ ID NO 74  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 74

cgttatgttt atcggcactt tgcac 25

<210> SEQ ID NO 75  
<211> LENGTH: 34  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 75

aggtgattgg cgatgatgcc gttcagctgt ctgc 34

<210> SEQ ID NO 76  
<211> LENGTH: 34  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 76

acagccatcg gtccatctct cctcgtaacg cctg 34

<210> SEQ ID NO 77  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 77

ggatagaatc gggtgccgct gatct 25

<210> SEQ ID NO 78  
<211> LENGTH: 21  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 78

gagaaccatg gcaccgaagg t 21

<210> SEQ ID NO 79  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 79

gtacttctac acagccatcg gtcca 25

<210> SEQ ID NO 80  
<211> LENGTH: 25  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Synthetic oligonucleotide

-continued

---

<400> SEQUENCE: 80  
 cgttatgttt atcggcactt tgcatt 25

<210> SEQ ID NO 81  
 <211> LENGTH: 25  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 81  
 acaggtactt ccatcttgta ctggt 25

<210> SEQ ID NO 82  
 <211> LENGTH: 20  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 82  
 tctcctccaa cgtccgatct 20

<210> SEQ ID NO 83  
 <211> LENGTH: 41  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 83  
 acctccacta gctccagcaa gccgaacaga ggtaaagacg a 41

<210> SEQ ID NO 84  
 <211> LENGTH: 43  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 84  
 tcgtctttac ctctgttcgg cttgctggag ctagtggagg tca 43

<210> SEQ ID NO 85  
 <211> LENGTH: 45  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 85  
 taccaacgtg cgaccatttt ctcggtcggc atctactcta ttct 45

<210> SEQ ID NO 86  
 <211> LENGTH: 45  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 86  
 aggaatagag tagatgccga ccgagaaaat ggtcgcacgt tggta 45

<210> SEQ ID NO 87  
 <211> LENGTH: 22

-continued

---

<212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
  
 <400> SEQUENCE: 87  
  
 aagcgtctct ttctgggctc tt 22

<210> SEQ ID NO 88  
 <211> LENGTH: 23  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
  
 <400> SEQUENCE: 88  
  
 tgccagttct gttggacatc tct 23

<210> SEQ ID NO 89  
 <211> LENGTH: 48  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
  
 <400> SEQUENCE: 89  
  
 tcgaggtcga cggtatcgat atctagaaca ggtacttcca tcttgtac 48

<210> SEQ ID NO 90  
 <211> LENGTH: 27  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
  
 <400> SEQUENCE: 90  
  
 ggtcactggt cctggcagct gacattg 27

<210> SEQ ID NO 91  
 <211> LENGTH: 27  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
  
 <400> SEQUENCE: 91  
  
 ctgccaggaa cagtgaccgg tgactct 27

<210> SEQ ID NO 92  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
  
 <400> SEQUENCE: 92  
  
 aagcagcaga tacgaccgtt gatctgcttg 30

<210> SEQ ID NO 93  
 <211> LENGTH: 26  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide  
  
 <400> SEQUENCE: 93

-continued

---

acggtcgtat ctgctgcttg gtgcac 26

<210> SEQ ID NO 94  
 <211> LENGTH: 45  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 94

actagtggat cccccgggct gcagcggctg gcactactc tattu 45

<210> SEQ ID NO 95  
 <211> LENGTH: 47  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 95

cgaggctgac ggtatcgata gtttaaacct cccaggtacc gactaac 47

<210> SEQ ID NO 96  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 96

ctcaatcaca gatcatgttt gggtagggttc 30

<210> SEQ ID NO 97  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 97

aaacatgatc tgtgattgag gtgattggcg 30

<210> SEQ ID NO 98  
 <211> LENGTH: 32  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 98

ctctgtgctt acagcagtgc ttatctgcca tg 32

<210> SEQ ID NO 99  
 <211> LENGTH: 38  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 99

gcactgctgt aggcacagag taacaggtag gtagacag 38

<210> SEQ ID NO 100  
 <211> LENGTH: 48  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence

-continued

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic oligonucleotide

&lt;400&gt; SEQUENCE: 100

agtggatccc ccgggctgca gtttaaactc ccacgcacga aagcaact 48

&lt;210&gt; SEQ ID NO 101

&lt;211&gt; LENGTH: 1102

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic polynucleotide (1546/1548 fragment)

&lt;400&gt; SEQUENCE: 101

acaggtactt ccactctgta ctggtaagtc ccgtccgtac tttaaataca ggggacctcc 60

cgtacgtcag gtacctatgt catgattctt cttctccccg gcattaagat ttgtttcttc 120

ttattaatta tcactactat tattcctgca gcaggaacgc cccctatcct ctctccaac 180

gtccgatcta tcgtgtggtt cgagtcgtgc tgttcacaca cacgcacgca cacacacact 240

ctcaaacata tctgccccct cttgccccct catccctcca gaaccgagat tcacgggtgcc 300

aactctcttt tccttggtgc tcctatcccc tttaaaaaaa aagaggtacg cagaaccgaa 360

cccaatcgac ctccgcccgc taattaagtg ggattctctg gccccaggg atcggtctct 420

tccttccatc ctggcgctcg aaaccagacg tcccaatcac cggagctgac cctggatcgg 480

cctctgatcc atggctagt tgcctgaca tttgtcggcc gttaaagccg tgggacctgc 540

gcccagtaca taaaccgcct gcgtcgtgac accgcctctt tggccgcgct tcccgtctac 600

ttctctogac gcctccgcta ttttcatcgc ggcgctcacc gcggcgccca tcccattgcg 660

ccgcagttgc tttcgtgctt gggattttgg tttccgtttc ttgtcttttt ctcttgggcg 720

ttcgtctcat cccgacctgc tccacggctt gattgattta tttgattttc aattttgtct 780

aaagtgcctt gcatcagacc gtccctcaac tccgaattcg tcttgtcgga acacctccag 840

ctgcctcgcc cgtccccgcg ctccacctcc gagcaatttc gcatgaggcg cagtcagcct 900

gatatttgat tgaacgcctt gctttcgtcc attgccagge cctcagactt gtctgtggcc 960

gcctgttata caggacggcc cttcggcttc tgatctatat cctgatctcc tgggtgttctt 1020

cattatcgac agctctcccc ggttcggctt cggcttctct ccggcattcg aactgctttc 1080

gtctttacct ctgttcggct tg 1102

&lt;210&gt; SEQ ID NO 102

&lt;211&gt; LENGTH: 1361

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic polynucleotide (1549/1550 fragment)

&lt;400&gt; SEQUENCE: 102

gctggagcta gtggaggcca acacatcaat gctattttgg tttagtcgtc caggcggatc 60

acaaaatttg tgtcgtttga caagatgggt catttaggca actggtcaga tcagcccact 120

tgtaagcagt agcggcggcg ctccaagtgt gactcttatt agcagacagg aacgaggaca 180

ttattatcat ctgctgcttg gtgcacgata acttgtcgtt ttgtcaagca aggtaagtga 240

acgaccgggt cataccttct taagtccgcc cttcctccct ttatttcaga ttcaatctga 300

cttacctatt ctacccaagc atccaaatgc ctgaactcac cgcgacgtct gtcgagaagt 360

ttctgatcga aaagtccgac agcgtctccg acctgatgca gctctcggag ggccaagaat 420



-continued

---

```

ctcgtgcttt cagcttcgat gtaggagggc gtggatatgt cctgcccggta aatagctgcg 480
ccgatggttt ctacaaagat cgttatgttt atcggcactt tgcateggcc gcgctcccga 540
ttccggaagt gcttgacatt ggggagttca gcgagagcct gacctattgc atctcccgcc 600
gtgcacaggg tgtcacgttg caagacctgc ctgaaaccga actgcccgtt gttctggagc 660
cggtcgaggga ggccatggat gcgatcgctg cggccgatct tagccagacg agcgggttcg 720
gcccattcgg accgcaagga atcgggtcaat aactacatg gcgtgatttc atatgcgcga 780
ttgctgatcc ccatgtgtat cactggcaaa ctgtgatgga cgacaccgtc agtgcgtccg 840
tcgcgaggc tctcgtatgag ctgatgcttt gggccgagga ctgccccgaa gtccggcacc 900
tcgtgcacgc ggatttcggc tccaacaatg tctgacgga caatggccgc ataacagcgg 960
tcattgactg gagcggaggc atgttcgggg attcccaata cgaggtcgcc aacatcttct 1020
tctggaggcc gtggttggt tgtatggagc agcagacgcg ctacttcgag cggaggcatc 1080
cggagcttgc aggatcgccg cgctccggg cgtatatgct ccgcattggt cttgaccaac 1140
tctatcagag cttggttgac ggcaatttcg atgatgcagc ttgggcgcag ggtcgtatgc 1200
acgcaatcgt ccgatccgga gccgggactg tcgggcgtac acaaatcgcc cgcagaagcg 1260
cggccgtctg gaccgatggc tgtgtagaag tactcgccga tagtggaac cgacgcccc 1320
gcaactcgtcc gagggcaaaag gaatagagta gatgccgacc g 1361

```

&lt;210&gt; SEQ ID NO 103

&lt;211&gt; LENGTH: 1300

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic polynucleotide (1551/1553 fragment)

&lt;400&gt; SEQUENCE: 103

```

aaatggctgc acgttggtac catctcgcca ttacggagag tatcgaaacc ctcaagtctgg 60
ccccgtttag cggggttttc ggttgggata gggaaaaaat ccggcgtatc gcgtctgagg 120
tcaggtcgga ggctttcaac aaggacatcc atgcctataa cattttacat atctaccaag 180
ctcggaaacc tctgtcaac tgacaaccag cgatcaacta gcgagagacc gagaacatca 240
aagtcgogct aggatttctt gtccaagagt ggataatgga gtccctgttt cgctcctctc 300
ttcccttggc tccacgagcg actgcgatcg agggtaactc tccgtagtac ctgtgctatc 360
gcggcctcat actggccttg gaccccagca atgctgatgc attctggtag ccgaaccgtg 420
caacgcactg gctgcgacgt ccagatattc acgacgtctc gcgggagctt ctcaactttgc 480
gtatatgggt ctaaagtgag tgtctgcccg ctgtgttcaa tcacgctgca atgtcagctg 540
ccaggatgaa agactataca aacaagctct gctcaactta taccaattct ttcgctttca 600
cgaaaggatt aagagcgagc ccgcggaatt tcccatgggt ggttgaccct gcgtcagaga 660
gcggttaatt gagattgacc tgtcttgaat ctctccgctc tcgtaccttc aaaacatgct 720
cctcaacttc ccttgacat gcggctgtag ttgagacgca tctgcctctc caccgacgg 780
ctctgggctt attttgattg cccatattta acggaccctg tgaaacactc gtactgtgtc 840
gccacgagtt tggaaccgga gataccctag aactacatc tactgatgaa ggaaaaataa 900
atcgaatcaa ctaaaacaag ctgaatctcc cttgtcctat tccttctatt gggccgacgg 960
agactattcc ggctatacaa ttgatacttc ataagatgcy tgtctacata cgcgagtaag 1020
atacagaaac acaaacacca accaccccc aagcccaccg aaacgcccac aaagaccag 1080
gaaagagacg ctttcgaaca tcgcataata atagcgggat gcaccgggtt ccccaaaact 1140

```

-continued

---

```

catcgccggc gacccttctg acgagacgtg ttattcttgc cactcgccgt atcttcataa 1200
tcccgaccag cgccgggact gcgccttcgc atggccccg ggccctaccg gacaatatcc 1260
tcttcaactct cactatcaga gatgtccaac agaactggca 1300

```

&lt;210&gt; SEQ ID NO 104

&lt;211&gt; LENGTH: 3389

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic polynucleotide (1547/1552 fragment)

&lt;400&gt; SEQUENCE: 104

```

tctcctccaa cgtccgatct atcgtgtggt tcgagtcgtg ctgttcacac acacgcacgc 60
acacacacac tctcaaacat atctgcccc tcttgccct ccatccctcc agaaccggag 120
ttcaagggtgc caactctctt ttcttgtgt ctctatccc gtttaaaaa aaagaggtac 180
gcagaaccga acccaatcga cctccgccga ctaattaagt gggattctct ggccccagg 240
gatcggcttc ttcttccat cctgcggtc gaaaccagac gtcccaatca ccggagctga 300
ccctggatcg gcctctgatc catggctagt ttgccctgac atttgtcgc cgttaaagcc 360
gtgggacctg cgcccagtac ataaaccgcc tgcgtcgtga caccgctct ttggccgcgc 420
ttcccgctca ctttcttga cgctccggt atttcatcg cggcgctcat cgccggcccc 480
atcccattgc gccgcagttg ctttcgtgcg tgggattttg gtttccggtt cttgtctttt 540
tctcttgggc gttcgtctca tcccagctg ctccacggct tgattgattt atttgatttt 600
caattttgtc taaagtcgcc tgcacagac cgtcctccaa ctccgaattc gtcttgcg 660
aacacctcca gctgcctcgc ccgtccccc cctcaccatc cgagcaattt cgcgatgaggc 720
gcagtcagcc tgatattga ttgaacgcct tgcttctcgc cattgccagg ccctcagact 780
tgtctgtggc cgctgttat ccaggacggc ccttcgggtc ctgatctata tctgatctc 840
ctggtgttct tcattatcga cagctctccc cggttcggtc tcggcttgtc tccggcattc 900
gaactgcttt cgtctttacc tctgttcggc ttgtcggagc tagtggaggt caacacatca 960
atgctatttt ggtttagtcg tccaggcgga tcacaaaatt tgtgtcgttt gacaagatgg 1020
ttcatttagg caactggtca gatcagccca cttgtaagca gtacggcgcg cgtcgaagt 1080
gtgactctta ttagcagaca ggaacgagga cattattatc atctgctgct tgggtgcacga 1140
taacttgtgc gtttgtcaag caaggtaagt gaacgacccg gtcatacctt ctttaagttcg 1200
cccttctccc ctttatttca gattcaatct gacttaccta ttctacccaa gcatccaaat 1260
gcctgaactc accgcgacgt ctgtcgagaa gtttctgatc gaaaagttcg acagcgtctc 1320
cgacctgatg cagctctcgg agggcgaaga atctcgtgct ttcagcttcg atgtaggagg 1380
gcgtggatat gtctgcggg taaatagctg cgcgatggg ttctacaaag atcgttatgt 1440
ttatcggcac tttgcatcgg ccgcgctccc gattccggaa gtgcttgaca ttggggagtt 1500
cagcgagagc ctgacctatt gcatctccc ccgtgcacag ggtgtcacgt tgcaagacct 1560
gcctgaaacc gaactgcccg ctgttctgga gccggtcgcg gaggccatgg atgcgatcgc 1620
tgcggccgat cttagccaga cgagcgggtt cggccattc ggaccgcaag gaatcgggtca 1680
atacactaca tggegtgatt tcatatgcgc gattgctgat ccccatgtgt atcaactggca 1740
aactgtgatg gacgacaccg tcagtgcgtc cgtcgcgcag gctctcgatg agctgatgct 1800
ttgggcccag gactgcccc aagtccggca cctcgtgcac gcggatttcg gctccaacaa 1860

```

-continued

---

tgtcctgacg gacaatggcc gcataacagc ggtcattgac tggagcagag cgatgttcgg	1920
ggattcccaa tacgaggtcg ccaacatctt cttctggagg ccgtgggttg cttgtatgga	1980
gcagcagacg cgctacttcg agcggaggca tccggagctt gcaggatcgc cgcgcctccg	2040
ggcgtatatg ctccgcattg gtcttgacca actctatcag agcttggttg acggcaattt	2100
cgatgatgca gcttggggcg agggtcgatg cgacgcaatc gtccgatccg gagccgggac	2160
tgtcggggcg acacaaatcg cccgcagaag cgcggccgtc tggaccgatg gctgtgtaga	2220
agtactcgcc gatagtggaa accgacgccc cagcactcgt ccgagggcaa aggaatagag	2280
tagatgccga ccgagaaaat ggtcgcacgt tggtagcacc tcgccattac ggagagtacc	2340
gaaaccctca gtctggcccc gtttagccgg gttttcgggt gggatagga aaaaatccgg	2400
cgtatcgcgt ctgaggtcag gtcggaggct ttcaacaagg acatccatgc ctataacatt	2460
ttacatatct accaagctcg gaagcctcct gtcaactgac aaccagcgt caactagcgc	2520
gagaccgaga acatcaaagt cgcgctagga tttctgtcc aagagtggat aatggagtcc	2580
ctgtttcgtc cctctctcc cttggctcca cgacgcactg cgatcgaggg taactctccg	2640
tagtacctgt gctatcgcgg cctcactctg gccttgacc ccagcaatgc tgatgcattc	2700
tggtagccga accgtgcaac gcaactggctg cgacgtccag atattcacga cgtctcgcgg	2760
gagctttcca ctttgcgtat atgggtctaa agtgagtgtc tgcccgtgt gttcaatcac	2820
gctgcaatgt cagctgccag gatgaaagac tatacaaca agctctgctc aactataacc	2880
aattctttcg ctttcacgaa aggattaaga gcgagcccgc ggaatttccc atgggtggtt	2940
gaccctgcgt cagagagcgg ttaattgaga ttgacctgtc ttgaatctct ccgctctcgt	3000
accttcaaaa catgtccctc aacttccctt gcacatgcgg ctgtagttga gacgcactg	3060
cctctccacc cgacggctct gggcttattt tgattgcca tatttaacgg acccgttgaa	3120
acactcgtac tgtgtcgcca cgagtttggg acccgagata ccctagacac tacatctact	3180
gatgaaggaa aaataaatcg aatcaactaa aacaagctga atctcccttg tcctattcct	3240
tctattgggc cgacggagac tattccggct atacaattga tacttcataa gatgcgtgtc	3300
tacatacgcg agtaagatac agaaacacaa acaccaacca ccccaaaagc ccaccgaaac	3360
gccgacaaag acccaggaaa gagacgctt	3389

&lt;210&gt; SEQ ID NO 105

&lt;211&gt; LENGTH: 2678

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic polynucleotide (1554/1555 fragment)

&lt;400&gt; SEQUENCE: 105

acaggtactt ccatcttgta ctggtaagtc ccgtccgtac tttaaataca ggggacctcc	60
cgtaacgtcag gtacctatgt catgattctt cttctccccg gcattaagat ttgtttcttc	120
ttattaatta tcatcactat tattcctgca gcaggaaagc cccctatcct ctctccaac	180
gtccgatcta tcgtgtggtt cgagtcgtgc tgttcacaca cacgcacgca cacacacact	240
ctcaaacata tctgccccct cttgccccct catccctcca gaacccgagt tcacgggtgcc	300
aactctcttt tccttgtgtc toctatcccc tttaaaaaaa aagaggtacg cagaaccgaa	360
cccaatcgac ctccgcccgc taattaagtg ggattctctg gccccaggg atcggttctt	420
tccttccatc ctgcccgtcg aaaccagacg tcccaatcac cggagctgac cctggatcgg	480
cctctgatcc atggctagtt tgccctgaca tttgtcggcc gttaaagccg tgggacctgc	540

-continued

---

```

gccagtaga taaaccgct ggcgcgtgac accgcctctt tggccgcgct tcccgtctac 600
tttcttcgac gcctccgtta ttttcacgca ggcgcctcgc ggggcgccc tcccattgca 660
ccgcagttgc tttcgtgctt gggattttgg tttccgtttc ttgtcttttt ctcttggggc 720
ttcgtctcat cccgacctgc tccacggctt gattgattta tttgattttc aattttgtct 780
aaagtcgctt gcatcagacc gtctccaac tccgaattcg tcttgctgga acacctccag 840
ctgcctcgcc cgtccccgc ctcaccatcc gagcaatttc gcatgaggcg cagtcagcct 900
gatatttgat tgaacgcctt gctttcgtcc attgccaggc cctcagactt gtctgtggcc 960
gcctgttata caggacggcc cttcggcttc tgatctatat cctgatctcc tgggtgttctt 1020
cattatcgac agctctcccc ggttcggctt cggcttctct cgggcattcg aactgctttc 1080
gtctttacct ctgttcggct tgttgctgga gagggtcccg cattgactcg gaggcctcat 1140
gtttgggaac ggacacggcc cgggacaacc tatcaccatg acccctccgt cttacaataa 1200
gtttggctcc cagtcgtcgg cttctggccg ctcgagaacc aattccgatg ctatggacat 1260
ccacgtcata acggatcggg acttggtgca acgagaacat aatcctggct acagcaattg 1320
gacgaacaat ggttccccat cgatatacac caagtaattc acccccggcc cctttttttc 1380
ccattctatg tggattgttc taccagttc acttttctct tgggtgatt aatcttctat 1440
cccaggagtc cagaaaagca gtactatgaa gaaaatggac ggctgtacca tgcgtatcga 1500
aaaggagtct atatgctacc atgcgatgac gaggagcagg atgcctcga tcttttccac 1560
aagttgttca ccgaggcgag ggtatctgat ggtctgattt acgcacctca cccgagaaac 1620
ggtcagttct tggatttggg atgtgggaca gggatctggg caattgacgt ggccaacaaa 1680
taccocgatg ctttcgttgt cggggtatg cttgctccga tccagcccc aaaccacccc 1740
aagaactgca aattttacgc tctttctgat tacgaaagcc cctgggcat ggggtgaagac 1800
tcctgggact tgatccacct ccaaatgggc tgcggaagcg tggtaggatg gccaaacctc 1860
taccgaagga tcttcgccc ccttcgaccg ggtgcttggg ttgagcaagt ggagatcgat 1920
ttcgaacccc gttgtgacga ccgatccttg ggacggactg gcgcttcgca aatggtacca 1980
atactgaaa cgagcaacc aggagacctt gcgaccaatc gccacaact cccgggaaac 2040
catcaaagat ttgcaggacg ccggcttcac tgagatcgat catcagatgg ttggattgcc 2100
cttgaatcct tggcaccagg acgagcacga gaaaatggtc gcacgttggg accatctcgc 2160
cattacggag agtatcgaac ccctcagctt ggccccgttt agccgggttt tcggttggga 2220
tagggaaaaa atccggcgta tcgctctgca ggtcaggctg gaggctttca acaaggacat 2280
ccatgcctat aacattttac atatctacca agctcggaag cctcctgtca actgacaacc 2340
agcgatcaac tagcgcgaga ccgagaacat caaagtcgca ctaggatttc ttgtccaaga 2400
gtggataatg gagtcctgt ttcgctctc tcttccttg gctccacgag cgactgcat 2460
cgagggtaac tctcgtagt acctgtgcta tcgcgccctc atactggcct tggaccccag 2520
caatgctgat gcattctggg agccgaaccg tgcaacgcac tggctgcat gtccagatat 2580
tcacgacgca tcgcgggagc ttctcacttt gcgtatatgg gtctaaagt agtgtctgcc 2640
cgctgtgttc aatcacgctg caatgtcagc tgccagga 2678

```

&lt;210&gt; SEQ ID NO 106

&lt;211&gt; LENGTH: 1651

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

-continued

---

<223> OTHER INFORMATION: Synthetic polynucleotide (1556/1557 fragment)

&lt;400&gt; SEQUENCE: 106

```

acagtgaccg gtgactcttt ctggcatgcg gagagacgga cggacgcaga gagaagggct    60
gagtaataag ccactggcca gacagctctg gcggctctga ggtgcagtgg atgattatta    120
atccgggacc ggccgcccct ccgccccgaa gtggaaaggc tgggtgtgccc ctcgttgacc    180
aagaatctat tgcacatcg  gagaatatgg agcttcatcg aatcacccgc agtaagcgaa    240
ggagaatgtg aagccagggg  tgtatagccg tcggcgaaat agcatgccat taacctaggt    300
acagaagtcc aattgcttcc gatctggtaa aagattcacg agatagtacc ttctccgaag    360
taggtagagc gagtaccggg  cgcgtaagct ccctaattgg cccatccggc atctgtaggg    420
cgtccaaata tcgtgectct cctgctttgc ccggtgatg aaaccgaaa ggccgctcag    480
gagctggcca gcggcgcaga ccgggaacac aagctggcag tcgaccatc  cggtgctctg    540
cactcgacct gctgaggtcc ctcagtccct ggtaggcagc tttgccccgt ctgtccgccc    600
ggtgtgtcgg cggggttgac aaggctgttg cgtcagtcca acatttgttg ccatattttc    660
ctgctctccc caccagctgc tcttttcttt tctctttctt ttcccatctt cagtatattc    720
atcttcccat ccaagaacct ttatttcccc taagtaagta ctttgctaca tccatactcc    780
atccttccca tcccttattc ctttgaacct ttcagttoga gctttccac  ttcacgcag    840
cttgactaac agtaccctcg cttgagcaga catcaccatg gccaaagtga ccagtgccgt    900
tccggtgctc accgcgcgcg acgtcgcggg agcggctcag ttctggaccg accggctcgg    960
gttctcccgg gacttcgtgg aggacgactt cgcgggtgtg gtccgggacg acgtgaccct   1020
gttcatcagc gcggtccagg accaggtggg gccggacaac accctggcct ggggtgtgggt   1080
gcgcggcctg gacgagctgt acgccgagtg gtcggaggtc gtgtccacga acttccggga   1140
cgctccggg  ccggccatga ccgagatcgg cgagcagccg tgggggcggg agttcgccct   1200
gcgcgacccg gccggcaact gegtgcactt cgtggccgag gagcaggact gaccgacgcc   1260
gaccaacacc gccggtccga cggcggccca cgggtcccag gagcttgaga tccacttaac   1320
gttactgaaa tcatcaaaac gcttgacgaa tctggatata agatcgttgg tgtcgatgtc   1380
agctccggag ttgagacaaa tgggtgttcag gatctcgata agatacgttc atttgtccaa   1440
gcagcaaaga gtgccttcta gtgatttaat agctccatgt caacaagaat aaaacgcgtt   1500
ttcgggttta cctcttccag atacagctca tctgcaatgc attaatgcat tgactgcaac   1560
ctagtaacgc cttacaggct ccggcgaaga gaagaatagc ttagcagagc tattttcatt   1620
ttcgggagac gagatcaagc agatcaacgg t                                     1651

```

&lt;210&gt; SEQ ID NO 107

&lt;211&gt; LENGTH: 1362

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic polynucleotide (1558/1559 fragment)

&lt;400&gt; SEQUENCE: 107

```

tgctggagct agtggaggtc aacacatcaa tgctatthttg gtttagtcgt ccaggcggat    60
cacaaaattt gtgtcgtttg acaagatggg tcatttaggc aactggctcag atcagcccac    120
ttgtaagcag tagcggcggc gctcgaagtg tgactcttat tagcagacag gaacgaggac    180
attattatca tctgctgctt ggtgcacgat aacttgtgcg tttgtcaagc aaggtaagtg    240
aacgacccgg tcataccttc ttaagttcgc ccttctccc tttatttcag attcaatctg    300

```

-continued

---

```

acttacctat tctaccaag catccaaatg cctgaactca ccgcgacgtc tgtcgagaag 360
tttctgatcg aaaagttcga cagcgtctcc gacctgatgc agctctcgga gggcgaagaa 420
tctcgtgctt tcagcttcga tgtaggaggg cgtggatagc tcctgcgggt aaatagctgc 480
gccgatgggt tctacaaaga tcgttatggt taccggcact ttgcatcggc cgcgctcccc 540
attccggaag tgcttgacat tggggagttc agcgagagcc tgacctattg catctccccg 600
cgtgcacagg gtgtcacggt gcaagacctg cctgaaaccg aactgccccg tgttctggag 660
ccggctcgcg aggccatgga tgcgatcgct gcggccgatc ttagccagac gagcgggttc 720
ggcccattcg gaccgcaagg aatcgggtcaa taccctacat ggcgtgattt catatgcgcg 780
attgctgatc cccatgtgta tcaactggcaa actgtgatgg acgacaccgt cagtgcgtcc 840
gtcgcgcagg ctctcgatga gctgatgctt tgggcccagg actgccccga agtccggcac 900
ctcgtgcacg cggatttcgg ctccaacaat gtctgacgg acaatggccg cataacagcg 960
gtcattgact ggagcggagg gatgttcggg gattcccaat acgaggtcgc caacatcttc 1020
ttctggaggc cgtggttggc ttgtatggag cagcagacgc gctacttcca gcgaggcat 1080
ccggagcttg caggatcgcc gcgcctccgg gcgtatatgc tccgcattgg tcttgaccaa 1140
ctctatcaga gcttgggtga cggcaatttc gatgatgcag cttggggcga gggtcgatgc 1200
gacgcaatcg tccgatccgg agccgggact gtcggggcga cacaatcgc ccgcagaagc 1260
gcggccgtct ggaccgatgg ctgtgtagaa gtactcggcg atagtggaaa ccgacgcccc 1320
agcactcgtc cgagggcaaa ggaatagagt agatgccgac cg 1362

```

&lt;210&gt; SEQ ID NO 108

&lt;211&gt; LENGTH: 1017

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic polynucleotide (1561/1562 fragment)

&lt;400&gt; SEQUENCE: 108

```

acctcccagg taccgactaa ctagatctag tagcacctgt actaccttac taaggttcct 60
agaatgtgct ccgtaacatt cgtcccaaac taaggtagct gtgggaggcg ggccgtcacc 120
accggaagt cactttcatc gtcagcaaca gtggatcatt tcctagaaag cgtttggcat 180
gtttttcttt ttctttcca tttttttta ccaacaaaac cctcctactg gccggagtcc 240
cgtcgtgccg agaatgagcg gcagttaccg agaagagtaa tcgcacggag tcctatcagc 300
aacgaggaga ttgttgacc catctgattg aacgttaact tctatggagt acaggtagctg 360
ttggcttagg agtctcccct acgctcccga atctctctg actccagagt gcttgtaac 420
atgatatggt ctctgttca ctctcccatt ctgcaggacg gaatgatttt ccgcatgcgg 480
gcgcgctatt cacacgagtt gattagtttt aatctatcgg acgaaagggt atcgttcatg 540
ttggtctggg ttaagttccg tcaaatagtt tcccgcact tggttctggg cgacagtgat 600
tgacgccagg cccgagcact tgggtagtcc gttaggaaa aaagggggga gaaataaata 660
tagaaaaga aaagagaaa agaaattaa aaaaaaggg ccggtgacag atagataaat 720
gtcaccgaag gagcggatgt acggacgacg tacaggtagc cgacatgcat accctttata 780
catgaatgga atgtactccg tacatatacc tagcgatata tcgcctaccg attactccgt 840
aatctatata ggaatgggtga gacacaagaa tagaacaact tcgtccacca agatccatcc 900
atggctggga tcgccccgc aaccaatgac tctggctccg aagcgccta ccttctcggg 960

```

-continued

---

gagatggcgg ttctccagg acctttcggg ttcggacgaa cccacccaaa catgatc 1017

&lt;210&gt; SEQ ID NO 109

&lt;211&gt; LENGTH: 5390

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic polynucleotide (1563/1564 fragment)

&lt;400&gt; SEQUENCE: 109

tgtgattgag gtgattggcg atgaagcttc aagctgtaag gatttcggca cggctacgga 60  
agacggagaa gcccaccttc agtggactcg agtaccattt aattctattt gtgtttgatc 120  
gagacctaata acagccccta caacgaccat caaagtcgta tagctaccag tgaggaagtg 180  
gactcaaatac gacttcagca acatctcctg gataaacttt aagcctaaac tatacagaat 240  
aagatgggtgg agagcttata ccgagctccc aaatctgtcc agatcatggt tgaccgggtgc 300  
ctggatcttc ctatagaatc atccttattc gttgacctag ctgattctgg agtgaccacg 360  
agggatcatga cttgagccta aaatccgccc cctccaccat ttgtagaaaa atgtgacgaa 420  
ctcgtgagct ctgtacagt accgggtgact ctttctggca tgcggagaga cggaccggacg 480  
cagagagaag ggctgagtaa taagcgccac tgcgcccagac agctctggcg gctctgaggt 540  
gcagtggatg attattaatc cgggaccggc cgcctctccg ccccgaagtg gaaaggctgg 600  
tgtgcccctc gttgaccaag aatctattgc atcatcggag aatatggagc ttcacgcaat 660  
caccggcagt aagcgaagga gaatgtgaag ccaggggtgt atagccgtcg gcgaaatagc 720  
atgccattaa cctaggtaca gaagtccaat tgcttccgat ctggtaaaag attcacgaga 780  
tagtaccttc tccgaagtag gtagagcgag taccggcgcg gtaagctccc taattggccc 840  
atccggcacc tgtagggcgt ccaaatatcg tgcctctcct gctttgcccg gtgtatgaaa 900  
ccgaaaggc cgctcaggag ctggccagcg gcgacagacc ggaacacaag ctggcagtcg 960  
accatccgg tgctctgac tegacctgct gaggtccctc agtccctggt aggcagcttt 1020  
gccccgtctg tccgcccggg gtgtcggcgg ggttgacaag gtcggtgctg cagtccaaca 1080  
tttgttgcca tttttctctg ctctccccac cagctgctct tttcttttct ctttcttttc 1140  
ccatcttcag tatattcatc ttcccatcca agaaccttta tttcccctaa gtaagtactt 1200  
tgctacatcc atactccatc cttcccatcc cttattcctt tgaacctttc agttcgagct 1260  
ttcccacttc atcgagctt gactaacagc taccocgctt gagcagacat cacaatgttt 1320  
gagatgggccc cgggtgggaa tcgtctcccc gccatgacct ctccagcgca caaccactac 1380  
agctaccact ctcccactc cagcgacaga ggccggtcaa ggcagaactc ggatgccatg 1440  
gacatccagt ccatcactga acgagagccg gcgaccagat acgcggttgc gggcggccct 1500  
gcgcccctgga atcgcaacgg gtctccgagc atgagcccta tgtatagcaa gtacatctct 1560  
cttaccctc cgtttctttc tgcttttcta ccaccccatc cctctttcca gtctgagtc 1620  
aggcttggtc cgcttgaagt ggctaagtgt atcctcgtct tctctctttc tgtgttttag 1680  
caattccgag cgaaaccagt ttcattgaaga gaacggagcg acctaccatg gctttcgcag 1740  
gggaatgtat tttcttccgt gcgatgagca agaacaggat cgcctcgaca tcttccataa 1800  
gctattcacg gtagcgcggg tatcggagag tctgatctac gcgccccatc caaccaacgg 1860  
ccggtttctg gacctaggat gtggaactgg tatctggggc atcgaggtag cgaacaagta 1920  
ccctgatgag tttgtcgtg gtgtggattt ggctcctatt cagcctccga accaccgaa 1980  
gaactgcgag ttctacgag ccttcgactt cgaagcgcca tgggcatggt gggaggattc 2040

-continued

---

ctgggatcta	atccatctgc	agatggggtg	cggtagtgtc	atgggctggc	caaacttgta	2100
tccaaggata	ttcgacatc	tccgtcccgg	tgcttggttt	gagcagggtg	agatcgattt	2160
cgagcctcga	tgtgatgatc	ggtcactaga	tggaaaggca	ttgcggcatt	ggtacgattg	2220
tcttaaacag	gcgacagcag	agaccatgcg	gccaatcgcc	catagctccc	gcgatacaat	2280
aaaagacctg	caggacgctg	ggttcacgga	gatcgaccat	caaatagtgg	gactcccgct	2340
caacccgtgg	catcaggacg	aacacgagcg	gaaggtggca	cgttggtata	acctggccgt	2400
ctcagagagc	atcgaaaacc	tcagtctggc	tcccttcagt	cggtgtctatc	gctggcccct	2460
ggagagaatc	cagcaactcg	ccgcagatgt	gaagtccgaa	gcattcaaca	aagagatcca	2520
tgctacaat	atactgcaca	tataccaggc	taggaaacca	ttaagataag	agcaaaaggc	2580
gaccacatcc	aggaacgcaa	aacgaaaagg	aggaaaactg	ctagcgcaag	tttatgtcac	2640
gctggcacac	gcccagccat	cagaaatctc	aacacgga	ggtatgaacc	gcatcaaccg	2700
agtatgaacg	acaattcgtc	catcacacac	ccttcggttc	ctctcgcagg	cccagcatgg	2760
cgccctatca	acctgcttta	cgacgtcgta	tatactggcg	aagtatcctc	tctatctact	2820
ctggcgctct	agataccgtg	aagatgcaga	caaaattggc	cgagctccct	tctcataatc	2880
ctcaagcttg	gaccgatggc	tgtgtagaag	tactcgccga	tagtggaaac	cgacgcccc	2940
gcactcgtcc	gagggcaaag	gaatagagta	gatgccgacc	gcgggatcca	cttaacgtta	3000
ctgaaatcat	caaacagctt	gacgaatctg	gatataagat	cgttggtgtc	gatgtcagct	3060
ccggagttga	gacaaatggt	gttcaggatc	tcgataagat	acgttcattt	gtccaagcag	3120
caaagagtgc	cttctagtga	tttaatagct	ccatgtcaac	aagaataaaa	cgcgttttcg	3180
ggtttacctc	ttccagatac	agctcatctg	caatgcatta	atgcattgac	tgcaacctag	3240
taacgcctta	caggctccgg	cgaagagaag	aatagcttag	cagagctatt	ttcattttcg	3300
ggagacgaga	tcaagcagat	caacggtcgt	caagagacct	acgagactga	ggaatccgct	3360
ctctgacaga	cgggcaattg	attacgggat	cccattggta	acgaaatgta	aaagctagga	3420
gatcgtccgc	cgatgtcagg	atgatttcac	ttgtttcttg	tccggctcac	cggtcaaagc	3480
taaagaggag	caaaaggaac	ggatagaatc	gggtgcccgt	gatctatagc	gtatagtgcc	3540
cttatcacgt	tgactcaacc	catgctatct	aactcaaccc	ctccttctga	acccaccat	3600
cttcttctct	ttcctctcat	cccacacaat	tctctatctc	agatttgaat	tccaaaagtc	3660
ctcggacgaa	actgaacaag	tcttctctcc	ttcgataaac	ctttggtgat	tggaataact	3720
gaccatcttc	tatagttccc	aaaccaaccg	acaatgtaaa	tacactctc	gattagccct	3780
ctagagggca	tacgatggaa	gtcatggaat	acttttggtc	ggactctcac	aatgatcaag	3840
gtatcttagg	taacgtcttt	ggcgtgggcc	ggtgttcggt	cccagtcctc	gatgcattca	3900
catgccctcc	ctaagctggg	ccctagactc	taggatccta	gtctagaagg	acatggcatc	3960
gatggactgg	gttcgttctg	agattatacg	gctaaaactt	gatctggata	ataccagcga	4020
aaagggatcat	gccttctctc	gttcttctctg	ttgatggaat	ggctaacaga	tgatagtcat	4080
tgcaacttga	aacatgtctc	ctccagctgc	catctacgaa	cccactgtgg	ccgctaccgg	4140
cctcaagggg	aaggctcgtg	tttctgagac	cgccccggt	gagggagctt	ctcagaccaa	4200
gctgttgga	catttcggtg	gcaagtggga	cgagttcaag	ttegcccta	tccgcaaaag	4260
ccaggtctct	cgtgccatga	ccagacgtta	ctttgaggac	ctggacaagt	acgctgaaag	4320
tgacgttgtc	attgttggtg	ctggttctctg	cggtctgagc	actgcgtacg	tcttgccaa	4380



-continued

---

```

ggctcgccg gacctgaaga ttgctatcgt cgaggccagc gtctctcctg gtcagtagtc 4440
catgatggat tgccttgac tcagctttcc ggaactaacg tgcaataggt ggcgggcct 4500
ggttgggtgg ccaactcttt tctgctatgg tcatgcgccg tcccgcggaa gtcttctga 4560
acgagctggg tgttccttac gaagaggacg caaaccccaa ctacgttgtc gtcaagcacg 4620
cctccctggt tacctcgaca ctcatgtcga aggttctctc ctcccccaat gtcaagctct 4680
tcaatgctac cgctgttgag gacttgatca cccgtccgac cgagaacggc aacccccaga 4740
ttgctgggtg tgcgtcaac tggacgctgg tcacccttca ccacgatgat cactcctgca 4800
tggaccccaa cactatcaac gctcctgtca tcatcagtac cactggtcac gatgggcat 4860
tcggcgccct ctgtgcgaag cgcttgggtg ccatgggcag cgtcgacaag ctaggtggca 4920
tgcgtggctc cgacatgaac tcggccgagg atgccatcgt caagaacacc cgcgaggtta 4980
ctaagggctt gataatcggc ggtatggagc tgtctgaaat tgatggcttt aaccgcatgg 5040
gcctacctt cggtgccatg gttctcagtg gtgtcaaggc tgccgaggag gcattgaagg 5100
tgttcgacga gcgtcagcgc gagtgtgctg agtaaatgac tcactaccg aatgggttca 5160
gtgcatgaac cggatttgc ttacggtctt tgacgatagg ggaatgatga ttatgtgata 5220
gttctgagat ttgaatgaac tcgtagctc gtaatccaca tgcatatgta aatggctgtg 5280
tcccgatgt aacgggtggg cattctagaa taattatgtg taacaagaaa gacagtataa 5340
tacaacaaa gatgcaagag cggctcgtca tcgcagataa gactgctgt 5390

```

&lt;210&gt; SEQ ID NO 110

&lt;211&gt; LENGTH: 1078

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Synthetic polynucleotide (1565/1566 fragment)

&lt;400&gt; SEQUENCE: 110

```

aggcacagag taacaggtag gtagacagtc gattgattgt aaatatcgtt attctacagt 60
tctcagaaat cgccccgctg ctttctgggc ttctattctc tctctctctc tctttttttt 120
tttttttttt ttccctggtt ttctaatttt ttttttctgg tccaatttct cccaccccat 180
ttgtttcccg gcccgtggaa cccaacgggg agcaagaaag actagactac tagactaggt 240
aactacctgt agctgggtgc taataataat attcacggcg gcaagacagc catggcctta 300
taaaccttcc tgtcatcctt accgctgacc tagtattatt attattatat gtggtgaggt 360
cgtacctagg tacctgggag cttaaagaga tactacaggt acttccatct tgtactggta 420
agtcccgctc gtactttaa tacaggggac ctcccgtagc tcaggtacct atgtcatgat 480
tcttctctc cccggcatta agatttgttt cttcttatta attatcatca ctattattcc 540
tgcagcagga acgcccccta tcctctctc caacgtccga tctatcgtgt gtttcgagtc 600
gtgctgttca cacacacgca cgcacacaca cactctcaaa catatctgcc cctcttgcc 660
ctccatccc tccagaacc gagttcacgg tgccaactct ctttctctg tgtctctat 720
cccgtttaa aaaaagagg tacgcagaac cgaacccaat cgacctccgc cgactaatta 780
agtgggattc tctggcccc agggatcggc ttcttcttc catcctgagg ctgaaacca 840
gacgtcccaa tcaccggagc tgaccctgga tcggcctctg atccatggct agtttgcct 900
gacatttgc gcccgtaaa gccgtgggac ctgcgccag tacataaacc gcctgcgtcg 960

```

-continued

---

```

tgacaccgcc tctttggcgc cgtttcccgt ctacttttctt cgacgcctcc gttattttca 1020
tcgcggcgct cctcggcgcg cccatcccat tgcgcgcgag ttgctttcgt gcgtggga 1078

```

---

The invention claimed is:

**1.** An isolated *Aspergillus terreus* fungus transformed with a heterologous nucleic acid construct comprising an *Aspergillus* species LaeA (loss of aflR expression A) gene, wherein the LaeA gene encodes a LaeA protein comprising an amino acid sequence which is at least 80% identical to the amino acid sequence of SEQ ID NO: 41 or SEQ ID NO: 59 and the heterologous nucleic acid construct is inserted upstream of the coding region of the endogenous LaeA gene, and wherein expression of the *Aspergillus* species LaeA gene is increased in the transformed fungus compared to an *A. terreus* fungus that is not transformed with the heterologous nucleic acid construct.

**2.** The isolated *A. terreus* fungus of claim 1, wherein the heterologous nucleic acid construct further comprises a heterologous promoter, a heterologous transcription terminator, a heterologous selective marker gene, or any combination thereof.

**3.** The isolated *A. terreus* fungus of claim 1, wherein the *Aspergillus* species LaeA gene encodes a protein comprising the amino acid sequence of SEQ ID NO: 41.

**4.** The isolated *A. terreus* fungus of claim 1, wherein the *Aspergillus* species LaeA gene encodes a protein comprising the amino acid sequence of SEQ ID NO: 59.

**5.** The isolated *A. terreus* fungus of claim 1, wherein the heterologous nucleic acid construct comprises a promoter operably linked to the *Aspergillus* species LaeA gene, a transcription terminator and a selective marker gene.

**6.** The isolated *A. terreus* fungus of claim 5, wherein:  
the promoter comprises the *A. nidulans* gpdA promoter;  
the *Aspergillus* species LaeA gene is an *A. nidulans* LaeA gene;  
the transcription terminator comprises the *A. nidulans* TrpC transcription terminator; or  
the selective marker gene comprises the *A. oryzae* pyrithiamine resistance (ptrA) gene.

**7.** The isolated *A. terreus* fungus of claim 6, wherein the heterologous nucleic acid construct comprises the nucleotide sequence of SEQ ID NO: 109.

**8.** A method of making itaconic acid, comprising culturing the isolated *A. terreus* fungus of claim 1 under conditions that permit the fungus to make itaconic acid, thereby making itaconic acid.

**9.** The method of claim 8, wherein the fungus is cultured in itaconic acid production media.

**10.** The method of claim 8, wherein the heterologous nucleic acid construct further comprises a heterologous promoter, a heterologous transcription terminator, a heterologous selective marker gene, or any combination thereof.

**11.** The method of claim 10, wherein the *Aspergillus* species LaeA gene in the heterologous nucleic acid construct is an *A. nidulans* or an *A. niger* LaeA gene.

**12.** The method of claim 9, wherein the *Aspergillus* species LaeA gene encodes a protein having an amino acid sequence at least 80% identical to the amino acid sequence of SEQ ID NO: 41 or SEQ ID NO: 59.

**13.** The method of claim 12, wherein the *Aspergillus* species LaeA gene encodes a protein having the amino acid sequence of SEQ ID NO: 41 or the amino acid sequence of SEQ ID NO: 59.

**14.** The method of claim 8, wherein the heterologous nucleic acid construct comprises a promoter operably linked to the *Aspergillus* species LaeA gene, a transcription terminator and a selective marker gene.

**15.** The method of claim 14, wherein:  
the promoter comprises the *A. nidulans* gpdA promoter;  
the *Aspergillus* species LaeA gene is an *A. nidulans* or an *A. niger* LaeA gene;  
the transcription terminator comprises the *A. nidulans* TrpC transcription terminator; or  
the selective marker gene comprises the *A. oryzae* pyrithiamine resistance (ptrA) gene.

**16.** The method of claim 15, wherein the heterologous nucleic acid construct comprises the nucleotide sequence of SEQ ID NO: 109.

\* \* \* \* \*