


Jan. 19, 1937.

R. H. McKEE

Plant Pat. 225

HYBRID POPLAR

Original Filed Aug. 26, 1930



Inventor
RALPH H. McKEE

By his Attorney

C. R. Parker, Jr.

UNITED STATES PATENT OFFICE

225

HYBRID POPLAR

Ralph H. McKee, New York, N. Y., assignor to
McKee Poplar Forestation, Inc., New York,
N. Y., a corporation of New York

Original application August 26, 1930, Serial No.
477,979. Divided and this application January
28, 1935, Serial No. 3,858. Renewed July 25,
1936

1 Claim. (Cl. 47—59)

This invention relates to a new and distinct
hybrid poplar and more particularly to a hybrid
poplar which has a very rapid rate of growth,
which is substantially immune to the ordinary
diseases to which poplars are susceptible, and
the wood of which is particularly suitable for use
in making paper pulp.

This is a division of my copending application
Serial No. 477,979, filed August 26, 1930.

As is well known, poplar wood is one of the
most valuable woods for use in the making of
wood pulp. At the present time there are known
in the United States and foreign countries, ac-
cording to Illick (1924) and to Sargent (1922),
approximately thirty varieties of poplar and five
commonly recognized hybrid poplars. Dode
(1905) lists 110 species but the accuracy of this
is questioned. Only three species of these are
used commercially for pulpwood. The common-
ly recognized hybrids include the Lombardy pop-
lar (male), Regenerata poplar (female), Carolina
poplar (Eugenei) (male) and Robusta poplar
(male), the parent trees of which are not defi-
nitely known. The fifth of the hybrid poplars
known as *P. (Populus) generosa* which was first
obtained in England in 1912, had as its male
parent *P. trichocarpa* and as its female parent
P. balsamifera virginiana (sometimes termed *P.*
angulata). As stated, these hybrids were known
prior to the present invention and are disclaimed
as forming no part of the invention.

Due to the increasing consumption of poplar
wood in cellulose industries, and the resulting
decreasing supply, it is obviously desirable to pro-
vide abundant sources of poplar wood for future
consumption. However, several disadvantages
attend the reforestation of land with ordinary
poplars. For example, ordinary poplar species
used commercially, such as *P. tremuloides* and
P. grandidentata, require many years of growth
to provide trees of suitable size for supplying
wood for use in making wood pulp or for lumber,
e. g. a tree 12 inches in diameter will ordinarily
be 80 years or more old. In addition, for maxi-
mum growth of ordinary poplars, a rich soil
or the use of fertilizers is required. Further,
ordinary poplars are susceptible to numerous
common tree diseases, such as Cytospora. Final-
ly, the length of fiber of the wood of ordinary
poplars is not so great as might be desired for
the making of paper pulp, so that a portion of
longer fiber pulp (such as that from spruce)
is often added to the poplar pulp to give the paper
the desired strength.

The present invention overcomes all of the

foregoing disadvantages and provides a new va-
riety of hybrid poplar which not only grows
quickly in ordinary soil but is particularly suit-
able for supplying wood of high grade for lum-
ber and for use in making wood pulp.

When two species of the *Populus* genus are
crossed seeds are obtained which when grown
produce seedlings of varying types. Some of
these seedlings will be dwarfs, some will grow at
the same rate as their parents, and others will
grow so rapidly that, relatively speaking, they
partake of the nature of giants. The present
invention is concerned with the propagating of
those seedlings which are giants, i. e. which have
a rapid growth rate. In general, it is found that
the varieties which are particularly useful are
those whose dimension is more than twice the
corresponding dimension of the parent trees.
By dimension is meant the height of the tree at
the same age, as well as the diameter of the tree
at the same age. In some cases, I have found
that the hybrid tree produced has a relative
dimension as great as three and one-half times
the corresponding dimension of the parent trees.

In order to obtain seedlings of the desired char-
acteristics as set forth above I have grown ap-
proximately 16,000 seedlings obtained by cross-
ing different varieties of poplars and determined
which of these hybrid seedlings possessed the
characteristics of the desired tree. The better
of these seedlings have been propagated vege-
tatively and the desired characteristics were
found to continue and show in the trees so pro-
duced vegetatively. This, of course, is frequent-
ly not true of subsequent generations grown from
seed after a hybrid crossing. A portion of this
work has been described in the *Journal of Hered-*
ity, vol. 24, pages 216-229, in 1933.

Of course, one could not be certain of obtaining
trees of the desired characteristics no matter
how much work was done. The uncertainty of
results is always present in attempting to get by
hybridization a particular type of tree.ordi-
narily in hybridization one simply makes numer-
ous crosses and picks out the one most interest-
ing. In this particular case, however, applicant
was searching for trees of a particular type and,
in general, discarded the hybrids produced unless
they partook of the nature of the type for which
he was searching. The present application is for
a hybrid in which the male parent was
P. trichocarpa and the female parent *P. angulata*.

The method which I have employed in pro-
ducing the seedlings of the hybrid poplar set
forth above consisted in tying paper sacks over

the clusters of flowers on the female trees before the flowers had opened in order to prevent normal pollination. Later, when the flowers had opened, pollen was obtained from the flowers of the selected male tree. The pollen was taken at once to the female tree and, by means of a wad of cotton, dusted over the female flowers. The sacks were immediately replaced to prevent normal pollination. After about two weeks the paper sacks were replaced by green cheesecloth sacks.

Approximately six weeks later when the seeds had ripened, they were removed, and the cotton carefully picked off and the seed planted in damp sand. When the seeds had germinated and the little trees had reached a height of about two inches, the small trees were transferred to larger containers and separated somewhat. Later, the seedlings were individually planted. At the end of the following summer the seedlings of desired characteristics were selected for propagation.

I have found that by simply taking the cuttings and making a hole approximately eight inches in depth in the damp ground by means of a sharp stick, putting the lower end of a cutting twelve or fourteen inches in length in the hole, and pressing the ground back around the cutting, fully 95 per cent. of the cuttings of the new hybrids will grow. As is known to those skilled in the art, this vitality is not shown by their parents. Indeed, in the case of the common poplar (*P. tremuloides*) for example, not more than one or two in a thousand will grow under the same circumstances.

The general method of propagation applicable to the new hybrid poplar constituting the present invention is as follows: From the seedling tree produced, cuttings of the new wood (the preceding summer's growth or the growth of two summers) are made after the leaves have fallen in the fall. These cuttings may be taken, generally speaking, during the period from November to March. The cutting from the small tree is then divided into pieces approximately from ten to fifteen inches in length. These cuttings are stored in an ice house or similar cool room, but are not allowed to dry out, until the time for planting is at hand. The cuttings may be planted either in the fall before the ground has become frozen or, as is preferable, in the spring shortly after the frost has left the ground and before the weeds have started to grow vigorously. The cutting is inserted in the ground, where it will begin to grow with considerable rapidity. I have found that, in the latitude of New York, with average summer weather, the cutting will grow into a tree of about six feet in height by the 15th of September. Under the same circumstances, the pure species which were the parents of this same hybrid will not reach a height of more than two or three feet. The growth of this new hybrid poplar in succeeding years is about the same as during the first year, that is, a gain of about six feet in height and approximately an inch in diameter per year may be counted upon.

I have found that from one cutting there will be obtained in one summer a five to eight foot tree. By cutting this tree into cuttings of say fourteen inches in length, there may be obtained from six to ten cuttings suitable for planting the following year, and from the old stalk there will come up a number of shoots so that during the following year there will be obtained not only the six to ten cuttings (each capable of providing six to ten new cuttings for propagation) but also

from the original stalk sufficient material to furnish another fifteen to twenty cuttings.

As will be apparent, if the tree is not cut up it will grow to be a mature tree and can be expected to increase in diameter roughly an inch per year and in height about six feet per year. The exact increase will be determined by the length of the growing season and by the supply of rain during the particular summer in question. I have found that the new hybrid poplar above referred to thrives on sunshine and rain but requires little in the way of rich soil or fertilizers. Analyses of the wood show that this hybrid poplar, contrary to most trees, does not take up from the soil much more than a trace of nitrogen, potash or phosphates. In fact, attempts to secure the usual effects of fertilizer have failed to show more than about 10 per cent. increase in growth due to fertilizer treatment. In no event has the rate of growth of the herein described hybrid poplar been increased 20 per cent by the use of fertilizer.

While the above describes the method always used in the early work of propagation of this new hybrid, later work has shown that the method of the McKee Patent No. 1,943,030, Propagation of trees from cuttings, is applicable so that all of the tree except the leaves can be used for propagation and thus a much more rapid multiplication of available trees be provided.

A tree which had been grown from a cutting for a single summer was sawed off even with the ground in the fall in order to make cuttings for propagation. From the one year root there grew the succeeding summer several shoots, a number of which reached the height of 9 ft. 6 in. in the one season's growth. Such trees show an annual ring of about $\frac{5}{8}$ in. Of the hybrid poplars known previously *P. eugenei* is generally considered as the most rapid growing but it shows annual rings of only about $\frac{1}{8}$ in. The wild poplars, such as *P. tremuloides*, which are now customarily used for lumber and pulpwood purposes show an annual ring of about $\frac{1}{11}$ in. It is, accordingly, evident that the new hybrid described herein has a surprising rate of growth in that it is much more rapid in growing than the pure species, ordinary aspen, *P. tremuloides*, which is considered the fastest growing commercial wood of the northern forests, or the hybrid poplars grown in lumber plantations in Italy and France, *P. eugenei*, which is considered the fastest growing of the hybrid poplars hitherto known. It is to be noted further that the hardness and density of the wood of the present hybrid as compared with *P. tremuloides* and *P. eugenei* is about the same. Rapid growing trees often give soft and light woods.

A study of the chromosome numbers as shown in the root system from growing cuttings was made. The stainings were by the Flemming and Heidenhain stains. In the original drawing the magnification is approximately 3400 times and the cell represented is from the cortical zone of a root tip stained with Heidenhain stain. The nucleus is in the equatorial plate stage of mitosis and shows 38 chromosomes.

Stems

The stems are round at the base becoming angular toward the tip. They are glabrous, grayish brown at the base and green toward the tip. The lenticels are round to oval at the base, measuring 0.7 to 1.5 mm. in diameter, and are

linear toward the tip, measuring 0.6 to 3.0 mm. long.

Leaves

The infantile leaves are ovate to lanceolate, generally rounded or very slightly cordate at the base, with tips acute. The blades are usually greenish but frequently are more or less suffused with a bright red pigment which overshadows the green. This reddish tinge appears especially in young terminal leaves of an older shoot. The first leaves of a cutting usually are bright green. The margins are pubescent and are finely crenate with 8 to 12 crenations per centimeter. The veins are prominent and bright red in color. The blades measure 2.1 to 2.8 cm. wide by 2.8 to 3.7 cm. long. The petioles are red, pubescent and measure 1.5 to 1.8 cm. long.

The mature leaves are ovate, measuring 7.0 to 17.5 cm. wide by 8.7 to 25 cm. long. The bases are truncate to cordate and the tips acute to acuminate. The blades are bright green, glabrous above and below with margins sparsely pubescent and crenate. The crenations occur 3.5 to 4.0 per centimeter, and are cuspidate with cusps brown. The midrib is yellow to bright red. The large veins are usually yellow, sometimes slightly tinged with pale red pigment. The petioles are green to red, ridged, often flattened, lenticellate, and measure 3.5 to 7.5 cm. long. The stipules are triangular, pale green, becoming darker, hispid, viscid, and fugacious.

Buds

The buds are broadly triangular to lanceolate and measure 2.0 to 6.0 mm. long. They are pale to reddish brown, glossy, viscid, and appressed.

Leaf scars

The leaf scars are broad, triangular, decurrent, and the bundle scars are prominent, keeled by prominent ridges which appear corky in older stems.

Resistance to disease

This new hybrid poplar is not attacked by the common poplar diseases. For example, it seems to be entirely immune to *Cytospora*, the most common disease of both the wild and cultivated poplars. It is susceptible to *Melampsora medusae* in Maine but it has not been attacked by this rust in Missouri, Wisconsin, or New York. This is the red rust that has proven to be a pest to most poplars growing in America, i. e., those designated under the names of *P. tremuloides*, *grandidentata*, *angulata*, *balsamifera*, *deltoides*, *wislizeni*, *occidentalis*, *dilatata*, and *condicans*. European poplars are similarly affected.

One would like to have available the flower and seed characteristics but obviously they can not be given as it would require a tree 15 to 20 years old to get these characters in true form. It was felt that it was safe not to attempt to get flowers and seed growth by grafting on mature trees as such growth is often affected by the hormones present in the sap of the grafted tree.

A study of the root producing hormones along the lines of the work done by Wend brings out that this hybrid has an exceptionally high percentage of such hormones.

The leaves of the young poplar hybrid cov-

ered by this application are surprisingly large, being often 9 in. in length and 8 in. in breadth. Such leaves are shown by trees in their first or second year of growth from cuttings. An older tree, say 5 years old, shows leaves which are approximately 4 in. by 3¼ in.

The characteristics of this hybrid are:

- (1) Fiber length of over 0.70 mm. in first year's growth.
- (2) Growth not increased 20% by fertilizer.
- (3) Rapid growth in diameter, e. g. ⅝ in. ring.
- (4) Rapid growth in height.
- (5) Relatively large proportion of the wood in the bole.
- (6) Relatively few branches.
- (7) Large leaves.
- (8) Chromosomes 38.
- (9) Resistance to disease.
- (10) Rapid propagation.

There are scarcely any other characteristics that could be named which, from the viewpoint of those engaged in the art of growing trees for pulpwood or lumber, could be as distinctive as the major ones of those listed above, i. e., length of fiber, speed of growth, and nearly entire lack of fertilizer effect.

Further evidence of the hardy strength of this hybrid is the fact that young trees, 12 to 14 in. high, have survived submersion for as much as a week under flood waters and continued with growth unimpaired.

The new hybrid poplar which I have developed shows not only a more rapid rate of growth than the parent trees but also a greater freedom from the ordinary diseases to which poplars are susceptible. In addition, this hybrid poplar shows an increased length of fiber as compared with that of the parent trees. As is obvious, this greater length of fiber is important, not only because it provides a wood of greater mechanical strength, but even more so because it provides a wood which when used for the making of paper is found to give materially stronger paper. As is well known, greater length of fiber is a decided advantage from the paper making standpoint. The ordinary poplar used for paper making, known as *P. tremuloides*, has, in the case of a one year old plant sampled a foot from the ground, a fiber length averaging about 0.54 mm. whereas the new hybrid set forth above has fiber length under the same conditions of about 0.74 mm. These figures are for small trees in a single summer's growth. Older trees have correspondingly longer length fibers; that is, the same approximate ratios are kept up.

From the foregoing it will be seen that I have developed, from the parents *P. trichocarpa* and *P. angulata*, a new hybrid poplar which is particularly suitable for supplying wood for lumber and for use in making wood pulp and which, because of the rapid growth rate of the trees and freedom from disease, is particularly valuable for reforestation purposes.

I claim:

A hybrid poplar as described, characterized by its very rapid rate of growth, its great fiber length, its large leaves, its disease resistance, and its ability to thrive and maintain substantially its maximum rate of growth in the absence of fertilizer.

RALPH H. MCKEE.