
Nov. 1, 1949.

Plant Pat. 889

ROSE PLANT

Filed July 22, 1948



Svend Poulsen
By Hobbstokh
attorneys

# UNITED STATES PATENT OFFICE

889

#### ROSE PLANT

Svend Poulsen, Copenhagen, Denmark, assignor to The Conard-Pyle Corporation, West Grove, Pa., a corporation of Pennsylvania

Application July 22, 1948, Serial No. 40,069

10

40

1 Claim. (Cl. 47—61)

The present invention relates to a new and distinct variety of rose plant of the hybrid polyantha class characterized as to novelty dominantly by its adaptability for both cut-flowers and garden decoration, by the purity of the color 5 of its flowers and their excellent form and large size for varieties of this class, combined with its exceptionally vigorous growing habit over a long blooming period and extraordinary freedom of flowering.

Asexual reproduction shows that these characteristics hold true through succeeding propagations.

In the drawing are shown specimens of this new variety, illustrating the flowers at different 15 stages of development from bud to full blown blossoms, together with specimens of young and mature foliage.

The following is a detailed description of the new variety, all references to color being in ac- 20 cordance with Ridgway's Color Standards and Nomenclature, and ordinary dictionary definition.

Type: Hardy; bush; for cut flowers and garden 25 decoration.

Class: Hybrid polyantha.

Breeding: Seedling.

Seed parent.—A seedling of "Mrs. Cutbush." Pollen parent.—"Edina."

Propagation: Holds its distinguishing characteristics through succeeding propagations by budding.

### Flower

Locality where grown: West Grove, Pennsyl- 35 vania.

Flowers borne: Several to stem in cluster, on normal, medium long stems.

Quantity of bloom: Abundant outdoors and in greenhouse.

Continuity: Continuous.

Fragrance: Slight to moderate; old China rose fragrance.

Bud:

green; medium rough; numerous minute red prickles.

Before calyx breaks.—Size — Medium. Form—pointed; with foliaceous appendages on the surface of the bud; with slen- 50 der, entire foliaceous parts extending beyond the tip of the bud equal to one-half or more of its length.

As calyx breaks.—Color—White.

As first petal opens.—Size—Medium small. 55

Form—elliptical, pointed. Color—white (sometimes faint blush). Outside: White flushed Light Viridine Yellow, Plate V, at base; inside: White delicately veined Light Viridine Yellow to Pale Viridine Yellow, Plate V.

Opening.—Opens up well.

Bloom: Size, when fully open—large; 3 inches to  $3\frac{1}{4}$  inches.

Petalage.—Double; 30 to 50 or more petals; arranged regularly with petaloids in center.

Form.—Cupped at first; becoming open cupped; petals remaining at first loosely cupped, outer row of petals first, rolled outward; becoming later, at maturity. loosely cupped.

Petals: Firm and silky; with inside like white silk crepe and outside more shiny.

Shape.—Outside—rounded heart shape; scalloped with one notch usually, sometimes a compound notch. Intermediateobovate; emarginate scalloped with one notch; less deeply notched than outside petals. Inside—elongated heart shape, scalloped with one notch, with some irregular petaloids.

This description of a newly opened flower was made from a rose grown in a greenhouse in the month of March at West Grove, Pa.

Color.—Outside petal.—outside surface: White with trace of Martius Yellow, Plate IV, at base, variable with angle of light; inside surface: same as outside surface. Intermediate petal — outside surface: White with trace of Martius Yellow, Plate IV, at base, variable with angle of light; inside surface: same as outside surface. Inner petal—outside surface: White with trace of Martius Yellow, Plate IV, at base, variable with angle of light; inside surface: same as outside surface.

This description was made from a rose that Peduncle.—Long; slender; erect; medium  $_{45}$  was open for three days in a greenhouse in the month of March, at West Grove, Pa.

> Color.—Outside petal.—outside surface: White with faint trace of Martius Yellow. Plate IV, at base of petals, variable with angle of light; inside surface: same as outside surface. Inside petal-outside surface: White with faint trace of Martius Yellow, Plate IV, at base of petals, variable with angle of light; inside surface; same as outside surface.

General color effect.—Newly opened flower— White with luminous glow. Three days open—White with faint luminous glow at center.

Behavior .- Drop off cleanly.

Flower longevity.—On bush in greenhouse— 10 to 14 days in March. Cut roses grown in greenhouse kept at living room temperature—5 to 6 days in March.

## Reproductive organs

Stamens: Many; arranged irregularly about pistils, mixed with petaloids.

Filaments: Medium length; yellow; most with anthers.

Anthers: Small; yellow-brown.

loosely bunched.

Stigma: Pale yellow green.

Ovaries: All enclosed in calyx.

Sepals: Moderately long; recurved; many curled. Color-Biscay Green (Plate XVII), sometimes lined or touched variably with Sorghum Brown (Plate XXXIX) at tip.

### Plant

Foliage:

Leaves.--Compound of seven leaflets; abundant; medium size; moderately glossy.

Leastets.—Shape — rounded obovate with apex obtuse; base rounded obtuse; margin serrate.

Color.—Mature—upper surface: Yew Green, Plate XXXI; under surface: Jade Green, 35 Plate XXXI. Young-upper surface: Cress

Green, Plate XXXI; under surface—Light Cress Green, Plate XXXI.

Rachis.--Medium heavy. Upper side--minutely prickly; under side—sparsely thorny.

Stipules.-Long; narrow; with medium long points, turning out at varying angles of less than 45° or about; finely serrate.

Disease.—Resistant.

10 Growth:

15 -

Habit.—Bushy; upright; branched.

Growth.—Vigorous.

Canes.—Medium heavy.

Main stems.—Bright green. Thorns—several: long; hooked downward; with long, narrow base. Prickles and hairs-none.

Pistils: Medium number.

Branches.—Bright green. Thorns—Few to Styles: Even; medium long; medium heavy; several; medium long; hooked downward; with narrow base. Prickles and hairsnone.

New shoots.—Bright green. Thorns, prickles, and hairs—none.

I claim:

A new and distinct variety of rose plant of the 25 hybrid polyantha class, characterized as to novelty by its adaptability for use as cut-flowers and garden decoration; its exceptionally vigorous growing habit over a long blooming period: the extraordinary freedom of flowering; the purity of the color of its flowers, and the excellent form and large size of said flowers, substantially as shown and described.

SVEND POULSEN.

No references cited.