


June 12, 1973

W. H. JESSEL, JR., ET AL

Plant Pat. 3,358

CHRYSANTHEMUM PLANT

Filed Aug. 25, 1971



1

3,358

CHRYSANTHEMUM PLANT

Walter H. Jessel, Jr., Doylestown, and William E. Duffett,
Akron, Ohio, assignors to Yoder Brothers, Inc., Bar-
berton Ohio

Filed Aug. 25, 1971, Ser. No. 175,011

Int. Cl. A01h 5/00

U.S. Cl. Pkt.—77

1 Claim

The present invention comprises a new and distinct
cultivar of chrysanthemum plant which is a seedling of
#60228-2×62966-1. Both parents are unpatented and
are identified for breeding purposes by the above num-
bers to distinguish the same from other parents in the
same program.

The new cultivar is similar in many respects to the
unpatented but commercially well-known cultivar Nep-
tune, having the same characteristics as Neptune of:

- (1) A year-round recommended flowering period.
- (2) Similar flower size.
- (3) Decorative flower form and white flower color.
- (4) Prominent stipules.

The new cultivar is distinguished from Neptune by the
following characteristics, when grown under comparable
conditions:

- (1) Response is approximately 4-10 days earlier de-
pending on the season of the year.
- (2) A more vigorous variety which can be grown
under medium rather than short treatment as Neptune
normally is.
- (3) Stronger, heavier stems.
- (4) No pinking of the flowers with a cool finish of
60° or less.
- (5) No longitudinal petal roll under low light and
also high temperature finishes of 70°-75° or above, which
conditions cause a reduction in flower size and con-
tributes to an aged appearance on Neptune.
- (6) No open center or significant petal reduction
under low light or low temperature conditions.

2

(7) A wider, more spreading habit, when both varie-
ties are grown and flowered with their recommended re-
sponse and treatments.

(8) Averages one more break per plant.

(9) Foliage approximately one-quarter inch shorter
in length.

(10) A lighter foliage color.

(11) Foliage less indented with less serration.

The new cultivar was selected from the progeny of the
designated cross and when asexually reproduced by cut-
tings at Barberton, Ohio, has been found to retain its
distinctive characteristics through successive propagations.

The new cultivar was selected from a flowering block
of the parent cultivar at Barberton, Ohio; and when asex-
ually reproduced by cuttings at Barberton, Ohio, has
been found to retain its distinctive characteristics through
successive propagations.

The new cultivar when grown in the vicinity of Bar-
berton, Ohio, has a response period of approximately 8
weeks, and the following detailed description is based on
observations made of the new cultivar in a greenhouse
in Barberton, Ohio. The response time, blooming period,
color, and total vigor may vary significantly with vary-
ing environmental conditions such as temperature, day
length, and light intensity. Suggested flowering in the
northern and western United States, and coastal Califor-
nia is from January through December; Florida, Novem-
ber through May.

The accompanying drawing illustrates the new cultivar,
with the color being as nearly true as possible with color
photographs of this type.

In the description which follows, color references are
to the Munsell Color Book, 1963 edition, and compari-
sons are made with the cultivar Neptune. Where only a
single value or description appears from the new cultivar,
the same value or description applies also for the com-
parison cultivar Neptune.

Botanical classification: *Chrysanthemum morifolium*

	Windsong	Neptune
Bloom:		
Size.....	Approximately 4"	
Fully expanded.....	Approximately 5"	
Borne.....	Terminal on disbudded plants	
Stems.....	Very strong	Strong.
Form.....	Decorative	
Permanence.....	10-14 days	
Color:		
Center of flower.....	White, 10Y9/1	Streaked pink on white 2.5RP5/12 with a cool finish or 60° or less to white, 7.5Y9/2 at finishing temperatures above this level.
Base of petals.....	White, N9.5/	
Inside of petals.....	do	
Reverse of petals.....	do	
Tonality from a distance.....	White	
Discoloration.....	do	White streaked pink, 2.5RP5/12 on N9.5/ with a cool finish of 60° or less to white, 7.5Y9/2 at finishing temperature above this level.
Petals:		
Texture.....	Smooth	
Appearance and form.....	Open at base, broad shallow keel, slightly recurving near apex, tapering to a blunt point.	Slightly open at base keeled at mid point, sev- eral ridges on bottom, 5/16" wide.
Arrangement.....	Composite, whorled on a single receptacle	
Persistence.....	Resist shatter	
Fragrance.....	Typical chrysanthemum	
Reproductive organs:		
Stamen, anthers.....	None	None to 25-30.
Pollen.....	do	None to scant.
Arrangement.....	Clustered in center of flower, if present	
Styles.....	Present ray florets	Present both ray and disc florets.
Length.....	Short	
Ovaries.....	At the base of petal attached to receptacle	
Plant:		
Form.....	Herbaceous	Upright.
Growth.....	Spreading	Approximately 12" when grown with 9 week short treatment as defined in the Yode, timetables.
Height.....	Approximately 11" when grown with 8 week medium treatment as defined in the Yoder timetables.	Approximately 5 1/2" when grown with 9 week short treatment as defined in the Yoder timetables.
Spread.....	do	

	Windsong	Neptune
Foliage:		
Top side.....	Dark green, 7.5 GY3/4.	Very dark green, 5 GY2/4.
Size.....	3 3/4" long, 2 1/4" wide.	4" long, 2 1/2" wide.
Quantity.....	Numerous.	
Shape.....	Spatulate lobed.	
Texture.....	Smooth.	
Ribs and veins.....	Prominent.	
Edge.....	Moderately indented.	Indented.
Serration.....	Slight.	Moderate.
Under side.....	Light green, 5 GY5/4.	Light green, 10 GY5/4.
Stipules.....	Prominent.	

We claim:

1. A new and distinct cultivar of chrysanthemum characterized particularly as to uniqueness when compared to the cultivar Neptune by its response which is approximately 4-10 days earlier depending on the season of the year, a more vigorous variety which can be grown under medium rather than short treatment, stronger, heavier stems, no pinking of the flowers with a cool finish of 60° or less, no longitudinal petal roll under low light and also high temperature finishes of 70°-75° or above, no open center or significant petal reduction

under low light or low temperature conditions, a wider, more spreading habit when both varieties are grown and flowered with their recommended response and treatments, average of one more break per plant, foliage approximately one-quarter inch shorter in length, a lighter foliage color, with the foliage being less indented with less serration.

No references cited.

ROBERT E. BAGWILL, Primary Examiner

UNITED STATES PATENT OFFICE
CERTIFICATE OF CORRECTION

Patent No. PP-3,358

Dated June 12, 1973

Inventor(s) WALTER H. JESSEL, JR. and WILLIAM E. DUFFETT

It is certified that error appears in the above-identified patent and that said Letters Patent are hereby corrected as shown below:

In column 2, line 3, "teratments" should be --treatments--;
line 8, "idented" should be --indented--; line 36, "from"
should be --for--.

In the tabulation, opposite "Color: , Center of flower",
under "Neptune", "finish or 60° or less" should read --finish
of 60° or less--.

Signed and sealed this 1st day of October 1974.

(SEAL)

Attest:

McCOY M. GIBSON JR.
Attesting Officer

C. MARSHALL DANN
Commissioner of Patents