C. B. COWARD ET AL

CHRYSANTHEMUM PLANT

Filed March 20, 1974



1

3,741 CHRYSANTHEMUM PLANT

Charles B. Coward, Fort Myers, Fla., and Walter H. Jessel, Jr., Doylestown, and William E. Duffett, Akron, Ohio, assignors to Charles B. Coward, Fort Myers, Fla. Filed Mar. 20, 1974, Ser. No. 453,034
Int. Cl. A01h 5/00

U.S. Cl. Plt.—81

1 Claim

The present invention comprises a new and distinct 10 cultivar of chrysanthemum plant known by the varietal name Beloved which is a sport of the cultivar known in the United States as Belair, which forms the subject matter of U.S. Plant Pat. No. 3,184, issued May 23, 1972.

The new cultivar is similar in many respects to Belair, 15 having the same characteristics of:

- 1. Use as a cut spray.
- 2. Vigor and general growth habit.
- 3. Flowering response.
- 4. Flower size, form and form retention.

The new cultivar is distinguished from Belair by its more intense lavender color when grown under comparable conditions. The color of Belair is more lavender pink. 25

The new cultivar is also similar in many respects to Belrose, disclosed in our pending application Ser. No. 416,175, filed Nov. 15, 1973. Belrose is also a sport of Belair, with the flower color of Belrose being a noticeably darker purple than the lavender color of Beloved.

The new cultivar was discovered at Fort Myers, Florida, and when asexually reproduced by cuttings at Barberton, Ohio, has been found to retain its distinctive characteristics through successive propagations.

The new cultivar, when grown in the vicinity of Barberton, Ohio, has a response period of approximately 9 weeks, and the following detailed description is based on observations made of the new cultivar in a greenhouse in Barberton, Ohio. The response time, blooming period, color, and total vigor may vary significantly with varying environmental conditions such as temperature, day length, and light intensity. Suggested flowering in the northern United States is from April through November 15; southeastern United States, March 15 through December 15; southwestern United States, March 15 through December, and southern Florida under outdoor conditions November through April.

The accompanying photographic drawing shows the unique characteristics of the new cultivar, the color being 50 as nearly true as possible with color illustrations of this type. The drawing is a comparison photograph, with the new cultivar Beloved appearing at the lower left, the cultivar Belair in the lower right, and the cultivar Belrose in the middle. Belrose is a noticeably darker purple as will 55 be apparent from the drawing.

Color references in the following description are to the Munsell Book of Color, 1963 edition, and color comparisons are made with the cultivar Belair. Where only a single value or description for the color appears in the 60 column under Beloved, the same value or description applies as well to Belair.

BOTANICAL CLASSIFICATION: Chrysanthemum morifolium

BLOOM:

size—2¾"
fully expanded—3¼"

2

borne—terminal cluster of 8-12 on elongated pedicel according to culture stems—elongated pedicels 6"-10" in length branching from main stem form—decorative permanence—7-10 days

	Beloved	Belair
color:		
center of	light yellow-green,	
flower.	2.5GY8/6.	
base of petals	do	
inside of petals.	lavender, 2.5 RP6/10 over light lavender, 2.5 RP8/4.	lavender pink, 5RP8/4 over pale lavender, 5RP9/1.
reverse of petals.	greyed lavender, 2.5RP6/8 over pale lavender, 2.5RP9/2.	light lavender pink, 5RP9/2 over pale lavender, 5RP9/1.
tonality from a distance.	lavender	lavender pink.
discoloration	lavender, 2.5 RP6/10 over light lavender, 2.5 RP8/4.	lavender pink, 5RP8/4 over pale lavender, 5RP9/1.

20

PETALS: texture—smooth

appearance and form—slightly open at base, keeled at mid point, tapering to a blunt point

arrangement—composite, whorled on a single receptacle

persistence—resist shatter

fragrance—typical chrysanthemum

REPRODUCTIVE ORGANS:

stamen, anthers—none to 20, more or less depending on culture

pollen—none to scant

arrangement—clustered in center of flower, if present styles—present in both ray and disc floret

length—short

ovaries—at the base of petal attached to receptacle PLANT:

form—herbaceous

growth—upright

height—approximately 36"-42" given 1-2 long day weeks with a response of 9 weeks, and 10-11 week total crop time, as defined in the timetables of Yoder Brothers, Inc., Barberton, Ohio, for the various seasons of the year

spread—none when grown single stem to slight when pinched

FOLIAGE:

top side—green, 7.5GY3/4 size—7''-9" long, 4"-5" wide

quantity—numerous

shape—spatulate lobed

texture—coarse

ribs and veins—prominent

edge—deeply indented

serration—slight

under side—greyed green, 5GY4/4

stipules—prominent at base of petiole

We claim:

1. A new and distinct cultivar of chrysanthemum characterized particularly as to its uniqueness when compared to the parent cultivar Belair by its more intense lavender coloration.

No reference cited.

ROBERT E. BAGWILL, Primary Examiner