

[54] CACTACEAE PLANT NAMED RUDOLPH
 [75] Inventor: Harry Higaki, San Mateo, Calif.
 [73] Assignee: Bay City Flower, Co., Half Moon Bay, Calif.
 [21] Appl. No.: 851,789
 [22] Filed: Apr. 14, 1986
 [51] Int. Cl. A01H 5/00
 [52] U.S. Cl. Plt./88
 [58] Field of Search Plt./88

Primary Examiner—Robert E. Bagwill
 Attorney, Agent, or Firm—James R. Cypher

EXEMPLARY CLAIM

A new and distinct plant variety of the Cactaceae fam-

ily obtained through cross-pollination of *Schlumbergera truncata* "Christmas Charm" and *Schlumbergera truncata* "Maria", distinguished from its parents and other known related varieties by a growth habit which combines a "reddish" colored bloom, a more vigorous growth rate than both parents, breaking propensity comparable to its paternal parent and better than its maternal parent, vigorous free budding and free flowering comparable to its paternal parent and better than its maternal parent, an upright and dense appearance and considerable resistance to nutrient deficiencies and fungus type diseases.

1 Drawing Sheet

1

BACKGROUND OF THE INVENTION

The invention relates to a new and distinct plant variety of the Cactaceae family which has been named *Schlumbergera truncata* "Rudolph" by the inventor.

Certain plant varieties of the Cactaceae family, which are among those of the *truncata* species of the *Schlumbergera* genus, tend to bloom in the months of November and December in the Northern Hemisphere. Because of their blooming time they appear in the market primarily during the Thanksgiving and Christmas seasons, and hence the common name Christmas Cactus. During the Christmas season, a "reddish" colored bloom is often very desirable. There are many "reddish" colored varieties available on the market, but none of the varieties combine the characteristics of having a fast growth rate, good breaking, and free budding and flowering. The "reddish" varieties that are available are either slow growing with good breaking, and free budding and flowering characteristics, or fast growing with poor breaking, and low budding and flowering characteristics.

The main objective of the invention has been to develop a variety of the Cactaceae family with a "reddish" colored bloom that has a fast growth rate, a vigorous breaking propensity and is free budding and free flowering. Another objective has been to develop a variety that also has an upright and dense appearance, and has suitable resistance to nutrient deficiencies and fungus type diseases.

The objectives of the invention have been obtained by the development of the new variety. The new variety was developed in a nursery located in Half Moon Bay, Calif., as a hybrid obtained through the cross-pollination of *Schlumbergera truncata* "Christmas Charm" and *Schlumbergera truncata* "Maria". "Christmas Charm" being the paternal parent and "Maria" the maternal parent. The seeds taken from the fertilized seed pod were cultivated at the Half Moon Bay nursery and, after observation of the seedlings, the hybridized plant of the new variety was selected and asexually reproduced by the inventor at the nursery by the propagation of stem cuttings taken from the original hybrid plant.

2

Through successive propagations, it has been recognized that specimens of the new variety resemble the parent varieties, but are distinguishable from the parents and other related varieties known to me, by a growth habit which is clearly visible in the specimens propagated and grown under nursery conditions at the Half Moon Bay location, as combining the following characteristics:

1. A "reddish" colored bloom.
2. A faster growth rate than both parents.
3. Breaking propensity comparable to the paternal parent and better than the maternal parent.
4. A vigorous free budding and flowering habit comparable to the paternal parent and better than the maternal parent.
5. An upright and dense appearance.
6. Considerable resistance to nutrient deficiencies and fungus type diseases.

BRIEF DESCRIPTION OF THE DRAWING

The accompanying drawings serve, by color photographic means, to illustrate the new plant variety and wherein:

FIG. 1 is a color photograph of a potted plant of the new variety illustrating the overall appearance and form of the plant; and

FIG. 2 is a color photograph showing a fully opened bloom of the new variety together with buds of the new variety.

DETAILED DESCRIPTION OF THE DRAWING

The following is a detailed description of the new variety with colors and hues being named in accord with the Horticultural Colour Chart issued by the British Colour Council in collaboration with The Royal Horticultural Society (two volume set issued separately in 1938 and 1942). The following description is based on observations of well fertilized plants of 16-18 months of age from initial propagation which were grown under 50-70% shaded polyhouse nursery conditions in Half Moon Bay, Calif., where temperatures average from 55° to 65° F. during the summer months, and from 45° to 55° F. during the winter months.

DETAILED PLANT DESCRIPTION

I. Name: *Schlumbergera truncata*.

II. Parentage:

Maternal.—*Schlumbergera truncata* "Maria" (*Zygocactus truncatus* "Maria"). 5

Paternal.—*Schlumbergera truncata* "Christmas Charm" (*Zygocactus truncatus* "Christmas Charm").

III. Classification: 10

Botanic.—(Britton and Rose, *The Cactaceae*, Constable and Co., Ltd., London, 1937, Vol. IV). (Bailey and Bailey and the staff of the Bailey Hortorium, Hortus Third, 1976). 15

(1) Family:	Cactaceae	
(2) Tribe:	Cereeae	
(3) Sub-tribe:	Epiphyllanae	
(4) Genus:	<i>Schlumbergera</i>	
(5) Species:	<i>Truncata</i> (Haw.) Moran	20

Commercial.—Thanksgiving cactus.

IV. Form: Terrestrial, shade-loving, succulent, leafless plant with jointed and branched stems. 25

V. Stems:

A. *General*.—Irregular with usually mono-chotomous to di-chotomous branching of mostly upright, adventitiously rootable, irregularly strongly revolute to flattened phylloclades that have a prominent midrib and prominently toothed lateral wings. 30

B. *Phylloclades*.—(1) *General*: Elongated with strong revolute character of the youngest phylloclades with the overall orientation that of a convex surface for one side of a phylloclade and a corresponding concave surface of the opposing side of the phylloclade, the phylloclades becoming mostly flattened with age. Phylloclades with transversely elongate, areole bearing truncated apex, with inwardly tapering basal wing margins that merge through a broad, usually pointed basal juncture with the phylloclade below and with an axillary areole associated with each tooth. (2) *Midrib*: *General* — Extends longitudinally of phylloclade and continuously through joints with laterally tapering cortex at wing insertions. Pith surrounding vascular bundles that branch and provide lateral extensions of the vascular system to marginal teeth. *Texture*: Smooth, waxy epidermis with wax in small embedded scales and becoming corky in basal stem areas with age. *Size* (18 mos) — *Length*: Usually between 47 and 57 mm with the average for respective plant specimens being usually between 50 and 54 mm. *Thickness*: usually between 1.5 and 4.5 mm with the average for respective plant specimens being between 2.2 and 3.7 mm. *Color* (at maturity): commonly spinach green (0960/2) to (0960/3). (3) *Wings*. — *general shape*: Generally strongly revolute when young becoming flattened with age from midrib cortex to tooth insertions with thinning taper toward margins. *Margins*: toothed. *Texture*: succulent to slightly coriaceous with smooth, waxy epidermis having wax arranged in small embedded scales and becoming corky in basal plant areas with age. *Size* (18 mo). — *Center thickness*: usually between 0.8 and 1.8 mm. *Width* (measured from phylloclade 65

axis to most offset lateral areole): usually between 12.5 and 21.2 mm. *Color* (at maturity): commonly spinach green (0960) to (0960/1). (4) *Teeth*. — *General*: Generally flattened and tapering along margins from wing insertions to an apex having a hyaline, pointed spine with random bending. *Adaxial marginal shape*: generally straight for larger teeth to distinctly concave for smaller teeth. *Adaxial marginal shape*: irregular, varying from being distinctly convex to generally straight or with the distal terminus being slightly convex. *Orientation*: usually projects distally of the phylloclade base in an alternate arrangement. *Margins*: entire. *Texture*: succulent to coriaceous with smooth, waxy epidermis having wax in small embedded scales and becoming corky in basal plant areas with age. *Number*: usually 6 to 8 per phylloclade. *Size* (18 mo). — *Center thickness*: usually between 0.5 and 1.2 mm. *Areole to apex dimension* (adaxial marginal side): usually between 0.2 and 5.2 mm for teeth located distally of basal teeth. *Color* (at maturity): commonly Scheeles green (860/3). (5) *Areoles*: terminal areole — Large elongated oval-shape with several acicular bristles and a dense mat of multi-cellular hairs and several buds that may mature into either new phylloclades of flowers. The opposing ends of the areoles are located adjacent to subsidiary areoles which are in turn located at the axils of teeth that are located at the distal end of the phylloclade. *Axillary areoles*: acicular bristles without glochidia but having numerous short colorless or light brown, multicellular hairs.

VI. Buds: Unarmored, ovid and chlorophyllous when first emerging.

VII. Flowers:

A. *General*.—Sessile, zygomorphic, usually in pairs, occasionally solitary or in clusters ranging up to four, terminal, perfect, epigynous with double hypanthium and tepals having aspiral emergence as a perianth provided with a sepaloid series of free tepals, a tube laminating series of tepals, and a tube forming series of united tepals.

B. *Sepaloid series*.—(1) *General*: free tepals inserted on top of the ovary. (2) *Shape*: deltoid in outer members of the whorl and with the inner members being irregular in shape varying from elliptic to slightly oblanceolate, tips of tepals are acute in outer members of the whorl and acuminate for inner member of the whorl. *Margins* are entire or with sparse irregular teeth for the apical portion of the inner members of the whorl. (3) *Texture*: succulent and glabrous outer whorl members and grading inwardly in the whorl to thin blades with fleshy basal areas. (4) *Number*: usually 9 or 10 (5) *Size* (at full bloom): base-tip dimension — usually less than 30 mm, range for longest tepal of the sepaloid series between 25 and 30 mm. *Maximum width dimension* — less than 15 mm, range of maximum between 11 and 15 mm. (6) *Color*: outer whorl members — Varies with position of tepal within the whorl. Smallest outer whorl members with a uniform field that in color is of a violet-red hue. The larger outer whorl members are characterized by

violet-red basal areas merging distally with red
 hued marginal and apical zones. The outer whorl
 members are basally magenta to magenta rose,
 (27/2) to (027/2), with the marginal and distal
 zones being spiraea red (025/2) with the transi- 5
 tion zone of color fusion between the basal and
 marginal areas being rose bengal to spiraea red,
 (25/3 to 025/3). Inner whorl members have basal
 areas dominated by magenta (27/3) intensifying 10
 to magenta (27/2) immediately distal to the basal
 area. Marginal areas are dominated by rose ben-
 gal (25/1) (25/2) intergrading to geranium lake
 (20/1) (20/2) as the marginal and the area imme-
 diately distal to the basal area merge. (7) Orienta-
 tion at full bloom: generally acute to slightly 15
 reflexed.

C. *Tube laminating series*.—(1) General: tepals in-
 serted on ovary and basally united below the
 throat as outer laminations on the perianth tube
 and with progressively greater amounts of basal 20
 fusion inwardly in the whorl. (2) Shape: grading
 inwardly in whorl with progressively longer
 base-tip dimensions and with oblanceolate to
 spatulate tepals and with the free portion of the
 blade being lanceolate to slightly oblanceolate, 25
 tip narrow to broadly acute generally with a
 short apiculate ultimate apex. Margins mostly
 entire at base with very small, sparse and irregu-
 lar teeth on the margin from near the middle of
 the tepal to the apex. (3) Texture: succulent and 30
 fleshy basal area, thin and soft blades. (4) Num-
 ber: usually 4 to 6. Size (at full bloom): Base-tip
 usually between 26 and 56 mm. Maximum width
 dimensions: usually between 13 and 17 mm. (5)
 Color: tepals with magenta basal areas (27/3) 35
 distally becoming more intensely colored (27/2).
 The marginal area of the blades of the tube lami-
 nating series is rose bengal (25/1) (25/2), inter-
 grading to geranium lake (20/1) (20/2) as the
 marginal and distal basal areas merge. 40

D. *Tube forming series*.—(1) General: tepals basally
 united to form perianth tube that is inserted on
 ovary and equipped with a vestigial carina at the
 throat. (2) Shape: perianth tube — Enlongatd
 and ellipsoidal in cross-section. Blades elliptic to 45
 slightly oblanceolate, tips acute to slightly acu-
 minate. Margins basally entire, apically with
 small irregular teeth. Carina: vestigial at throat
 of perianth tube. (3) Texture: perianth tube —
 thick, succulent, slightly ribbed at point of lateral 50
 fusion of tepals. Blades: proximally fleshy, dis-
 tally thin and soft. Carina: fleshy. (4) Blade num-
 ber: usually 8. (5) Size (at full bloom): perianth
 tube — base to keel length. Usually between 32
 and 36 mm along axis of tube with average 55
 length difference between measurements along
 dorsal and ventral sides for respective specimens
 usually between 2.0 and 4.5 mm. Internal major
 axis (at throat): usually between 8 and 11 mm
 when measured perpendicular to axis of perianth 60
 tube. Internal minor axis (at throat): usually be-
 tween 4 and 8 mm when measured perpendicular
 to axis of perianth tube. Blades — length (keel to
 tip): usually between 25 and 33 mm. width (maxi-
 mum): usually between 12 and 15 mm. (6) Color 65
 (at full bloom): perianth tube — usually translu-
 cent white with lines marking lateral fusion of
 tepals light magenta (27/3). Blades — Tepal

blades with marginal blade areas which in color
 are dominated by rose bengal (red) to geranium
 lake (red) that merges with a violet-red (ma-
 genta) for the basal blade area that extends dis-
 tally of the throat. The basal area magenta (27/3)
 distally becomes geranium lake (20/1) and
 merges with the marginal area rose bengal
 (25/1). Carina: characterized by an intense fuch-
 sia purple (28/1) to strong peony purple (729/2).
 Orientation at full bloom: reflexed.

E. *Androecium (stamens)*.—(1) General: numerous
 exerted and diadelphous stamens with one
 group having filaments basally fused to the peri-
 anth tube and the other group having filaments
 basally united to form a nectary housing, thin
 annulus around the style and which is provided
 with a thin, slightly deflexed, irregularly toothed
 margin at the throat of the annulus. (2) Stamen
 number: tube attached group — usually between
 18 and 22. Basally united group — usually be-
 tween 18 and 22. (3) Filament general: translu-
 cent white and glabrous with anther connective.
 Shape: long, slender, gradually tapering from
 base to anther connective. Texture: glabrous and
 smooth. Color: translucent white. Size (at full
 bloom) length: tube attached group — usually
 between 37 and 45 mm. Basally united group:
 usually between 39 and 45 mm. Diameter: usu-
 ally between 0.2 and 0.3 mm intermediate the
 opposing ends. (4) Anthers generally: adnate
 with four longitudinally dehiscent pollen sacs.
 Shape: elongated. Texture: waxy. Color (post
 dehiscence, pollen color): usually greenish-yel-
 low, uranium green (63/2).

F. *Gynoecium (pistil)*.—(1) General: compound,
 parietal placentation with a united style sur-
 rounded by an annular diffuse nectary at its in-
 sertion. (2) Style general: stout and inserted in
 ovary. Shape: elongated cylindrical and gener-
 ally tapering. Texture: fleshy and glabrous.
 Color: proximal end is pure magenta (27) gradu-
 ally changing to fuchsia purple (28) at the distal
 end of the style. Size (at full bloom): Length:
 usually about 56 mm. Diameter: usually between
 0.6 and 0.8 mm intermediate the opposing ends.
 (3) Stigma — General: exerted and erect with
 usually 6 or 7 inner marginally adhering lobes.
 Shape: elongated and tapering toward lobe tips
 and having relatively blunt apices. Texture:
 fleshy with short glutinous hairs. Color: area
 proximal to point of attachment of stigma to
 style persian rose (628/1) distally decreasing in
 intensity to (628/2) and (628/3) at the tips of the
 stigma lobes. Size (lobe length at full bloom):
 usually between 2.8 and 4.2 mm. (4) Ovary Gen-
 eral: epigynous with thin epidermis and distally
 located concavity and with single cavity usually
 having 6 or 7 carpels with numerous ovules.
 Shape: terete to ovoid, broadening from inser-
 tion to floral end. Texture: succulent and glabrous
 with thin epidermis. Color: mostly translucent
 white or very light green-yellow or chartreuse
 (663/3) with portion immediately distal to point
 of attachment to phylloclade rhodamine pink
 (527/3) and portion immediately bordering at-
 tachment of tepals magenta (27/3). Size (at full
 bloom) — Length (insertion to concavity base):
 usually between 7 and 10 mm. Major axis (distal

end of concavity): usually between 7 and 8 mm.
 Minor axis (distal end of concavity): usually between 6 and 8 mm.

The following is a general description of a flower of the new plant variety grown at Half Moon Bay, Calif., under greenhouse nursery conditions.

Bloom life: 8 Days.

Sepaloid series of tepals:

- (1) *Number*.—10.
- (2) *Size (at full bloom)*.—Maximum base-tip dimension: 26 mm. Minimum base-tip dimension: 3 mm. Maximum width dimension: 12 mm.
- (3) *Color*.—Most outer whorl basally magenta (27/2) to magenta rose (027/2), marginal and distal areas spiraea red (025/2) with transition zone of rose bengal (25/3) to spiraea red (025/3) between basal and marginal areas. Inner whorl members basally magenta (27/3) becoming more intensely colored (27/2) distally, marginal zones rose bengal (25/1) (252) with transition zone of geranium lake (20/1) (20/2).

Tube laminating series:

- (1) *Number*.—6.
- (2) *Size (at full bloom)*.—Maximum base-tip dimension: 56 mm. Minimum base-tip dimension: 26 mm. Maximum width dimension: 13 mm. Minimum width dimension: 10 mm.
- (3) *Color*.—Basally magenta (27/3) (27/2), marginal blade area rose bengal (25/1) (25/2).

Tube forming series:

- (1) *Number*.—8.
- (2) *Size (at full bloom)*.—Perianth tube: Base to throat length 34 mm. Interior major axis (at throat): 10 mm. Interior minor axis (at throat): 6 mm. Blades: maximum length (throat to tip) 28 mm. Minimum length (throat to tip): 26 mm. Maximum width: 14 mm. Minimum width: 12 mm.
- (3) *Color*.—Perianth tube: translucent white with the lines of lateral fusion of tepals light magenta (27/3). Blades: basally magenta (27/3) distally becoming geranium lake (20/1), marginal areas are rose bengal (25/1). Carina: Fuchsia purple (28/1) to peony purple (729/2).

Androecium:

- (1) *Stamen number*.—Tube attached group: 90. Basally united group: 19.
- (2) *Filaments*.—Color: translucent white. Size (at full bloom): Length — tube attached group: 43 mm (avg.). Basally united group: 41.5 mm (avg.). Diameter: 0.25 mm (avg.).
- (3) *Anthers*.—Color (post dehiscence): Uranium green (63/2).

Gynoecium:

- (1) *Style*.—Color: Proximal end pure magenta (27), distally becoming fuchsia purple (28). Size (at full bloom): Length — 56 mm. Diameter — 0.7 mm intermediate the two opposing ends.
- (2) *Stigma*.—Color: Proximally persian rose (628/1) decreasing in intensity distally to (628/2) or (628/3). Size (lobe length.) 3.5 mm (avg.).
- (3) *Ovary*.—Color: Translucent white to very light green-yellow or chartreuse green (663/3) with zone immediately distal to attachment with phylloclade rhodamine pink (527/3) and the portion

immediately bordering attachment of tepals magenta (27/3). Length (insertion to concavity base) — 10 mm. Major axis (distal end of concavity) — 8 mm. Minor axis (distal end of concavity) — 7 mm.

The following is a general description of a specimen of the new plant variety that was grown from the propagation of a single phylloclade in a nursery at Half Moon Bay, Calif.

Age from initial propagation: 18 months.

Branches from propagated cutting: 7.

Total number of phylloclades grown from cutting: 77.

General:

Branch No.	No. of Phylloclades	Max. Length	No. of Tips
1	28	219 mm	11
2	24	197 mm	11
3	14	223 mm	5
4	6	175 mm	2
5	3	76 mm	2
6	1	29 mm	1
7	1	19 mm	1

Midribs:

Branch No.	Length (avg.)	Thickness (avg.)
1	47.9 mm	3.3 mm
2	42.8 mm	2.9 mm
3	49.8 mm	3.4 mm
4	45.8 mm	2.2 mm
5	39.7 mm	2.0 mm
6	32.0 mm	1.2 mm
7	19.0 mm	1.7 mm

Wings:

Branch No.	Center Thickness (avg.)	Max. Width (avg.)
1	1.4 mm	35.2 mm
2	1.2 mm	32.4 mm
3	1.1 mm	30.2 mm
4	1.0 mm	31.3 mm
5	0.9 mm	25.3 mm
6	0.8 mm	10.0 mm
7	1.2 mm	8.0 mm

Teeth:

Branch No.	No./Phylloclade (avg.)	Center Thickness (avg.)	Areole to Apex Dimension (avg.)
1	6.5	0.7 mm	2.7 mm
2	7.0	0.6 mm	3.2 mm
3	6.4	0.6 mm	3.7 mm
4	6.3	0.5 mm	3.7 mm
5	6.0	0.5 mm	2.1 mm
6	6.0	0.6 mm	0.7 mm
7	5.0	0.7 mm	0.5 mm

Phylloclade color: Spinach green (0960) (0960/1) to Scheeles green (860/3).

I claim:

1. The new and distinct hybrid plant variety of the Cactaceae family substantially as herein shown and described.

* * * * *



FIGURE 1



FIGURE 2