## United States Patent [19]

## Kientzler

[11] Patent Number: Plant 6,925 [45] Date of Patent: Jul. 18, 1989

[54]	IMPATIENS PLANT NAMED MIMAS		Primary Examiner—Robert E. Bagwill
[75]	Inventor:	Ludwig Kientzler, Bad Kreuznach, Fed. Rep. of Germany	Attorney, Agent, or Firm—Foley & Lardner, Schwartz, Jeffery, Schwaab, Mack, Blumenthal & Evans
[73]	Assignee:	Paul Ecke, Jr., Encinitas, Calif.	[57] ABSTRACT
[21]	Appl. No.:	119,843	An Impatiens plant named Mimas having many large red-purple flowers and dark green foliage, upright growth habit, early flowering, and a floriferous habit.
[22]	Filed:	Nov. 12, 1987	
[51] [52]	Int. Cl. <sup>4</sup>		growth habit, carry howering, and a hornerous matrice
[58]			1 Drawing Sheet

1

The present invention relates to a new and distinct cultivar of Impatiens plant known by the cultivar name Mimas, and botanically known as Impatiens.

Mimas was developed by applicant through a controlled breeding program in Bad Kreuznach, Federal 5 Republic of Germany. Both the male and female parents are unknown at this time. Mimas was discovered and selected as one flowering plant within the progeny of the cross by applicant in a controlled environment in Bad Kreuznach, Federal Republic of Germany.

The first act of asexual reproduction of Mimas was accomplished when terminal or stem cuttings were taken from the initial selection by applicant in a controlled environment in Bad Kreuznach, Federal Republic of Germany. Horticultural examination of selected 15 plants of Mimas has demonstrated that the combination of characteristics as herein disclosed for Mimas are firmly fixed and are retained through successive generations of asexual reproduction.

Mimas has not been observed under all possible envi- 20 ronmental conditions. The phenotype may vary significantly with variations in environment such as temperature, light intensity, soil conditions, nutrient feeding, and day length.

The following characteristics distinguish the new 25 Impatiens from both its parent varieties and other cultivated Impatiens of this type known and used in the floriculture industry within the knowledge of applicant:

- 1. Unique radiant lilac flowers.
- 2. Very floriferous with many large flowers above the dark green foliage.
- 3. Early flowering.
- 4. The growth habit is upright and self branching, giving a nice full appearance; medium height.

The accompanying photographs illustrate a perspective view of the overall appearance of the new cultivar. Sheet 1 comprises the color photograph in which the

precise flower color is not uniformly depicted.

Sheet 2 is the same photograph as appears on Sheet 1, but is in black and white with three flowers being in color. The color closely corresponds to the color values noted below.

The following is a detailed description of Mimas 45 based on plants produced under commercial practice in Encinitas, Calif. Unrooted cuttings were rooted in March under intermittent mist, and the plants subsequently transplanted to 6" pots. Flowering commenced in July, and a plant nicely proportioned for a 6" pot was

2

ready five or six weeks later. Most of the data which follows was collected between September 1-5 in Encinitas, Calif. The measurements and description of foliage were taken from fully expanded leaves.

Color references are made to The Royal Horticultural Society Colour Chart except where general terms of ordinary dictionary significance are used. Color chart correlations were made indoors but under natural light conditions, with the leaves and flowers being compared to the R.H.S. Colour Chart near a west-facing window.

Parentage: Seedling from cross pollination, with both the male and female parents being unknown at this time.

Propagation:

Type cutting.—Tip cutting. Stems 2-3 cm long will develop to 4-5 cm after 21 days in propagation. Time to initiate roots.—8-10 days at 73° C. Nicely developed root mass in 18-21 days.

Rooting habit.—Numerous, fibrous adventitious roots from the stem.

Plant description:

35

Form.—Symmetrical, bush shaped, flowering herb, with self-branching characteristics giving the plant a full appearance.

Habit of growth.—Vigorous, self-branching, medium habit, producing whorls of leaves and flowers on branched stems. Growth is indeterminate and flowering is continuous.

Foliage description.—Leaves are simple. Lower leaves on stem are opposite. Higher leaves in whorls of 5. Leaf shape: Lanceolate to ovate, tip acuminate, base acute. Leaf blade size: 7-9 cm×3-3.5-4 cm. Leaf margin: Ciliate and lightly serrated. Leaf texture: Upper surface: glabrous. Under surface: glabrous. Leaf color: Dark green, with some yellow variegation along the mid-vein of older leaves. Upper surface: Darker than 139A. Under surface: Near 197A. Venation: Pinnate.

Flowering description:

Flowering habits.—Very floriferous, flowering continuously. Flowers develop progressively around the whorl of leaves, taking 5-7 days from buds which show color to bloom. Flowers are single and large, lasting for 2-3 weeks.

Natural flowering season.—Flowering is indeterminate and occurs throughout the year. Quantity of

flowers increases with increasing light intensity and duration.

Flower buds.—Ellipsoidal and covered with 3 sepals. A reddish approximately 5.5 cm long spur originates from the base of the major sepal. Two 5 additional, rudimentary sepals are fused to back of the top petal.

Flowers borne.—Singly, on pedicels 5 cm long, pedicels reddish in color.

Quantity of flowers.—Floriferous. Two flowers per 10 leaf occurring progressively around the whorl of leaves so that tight buds to mature blooms are visible at the same time in large numbers.

Petals.—Number of petals: Five (5). Shape: Top named I petal heart shaped with wide base, all other pet- 15 thereof. als heart shaped with pointed base. All petals

overlap. Color: Radiant red-purple. Upper surface: Near 74A. Under surface: 67B. Size of flowers: 5 cm across.

Reproductive organs.—Stamens: Five (5) with the lower stamen being shorter than the other stamens. Stamens united in an asymmetrical tube surrounding the ovary. Anthers: Hooded, cream fused with red. Pollen: Cream color. Stigma: Five pointed star, colorless. Styles: Very short, reddish in color. Ovary: 5 celled, 4–5 mm. long until fertilized, green in color.

1. A new and distinct cultivar of Impatiens plant named Mimas, as illustrated and described, and parts thereof.

\* \* \* \*

20

25

30

35

40

45

50

55

60

