

[54] NECTARINE — EARLY RED JIM

[75] Inventor: James W. Taylor, Dinuba, Calif.

[73] Assignee: Ito Packing Co., Inc., Reedley, Calif.

[21] Appl. No.: 283,990

[22] Filed: Dec. 13, 1988

[51] Int. Cl.<sup>4</sup> ..... A01H 5/00

[52] U.S. Cl. .... Plt./41

[58] Field of Search ..... Plt. 41

Primary Examiner—James R. Feyrer

Attorney, Agent, or Firm—Dennis B. Haase

[57] ABSTRACT

A new and distinct variety of nectarine tree characterized by a yellow fleshed, cling stone fruit, which is attractive in appearance and having excellent sugar content and flavor, and, further, having the characteristic and advantage of maturing in the early part of the late season, thereby becoming marketable at a time when comparable fruit is likely to be less available.

1 Drawing Sheet

1

BACKGROUND OF THE VARIETY

The present invention relates to a new and distinct variety of nectarine tree, which I refer to as "Early Red Jim", displaying a yellow fleshed, cling stone fruit, which is both attractive in appearance and having the characteristics of maturing in the late season period. This variety was developed from a continued fruit breeding program.

ORIGINAL AND ASEXUAL REPRODUCTION OF THE NEW VARIETY

This novel nectarine was derived from an ongoing fruit breeding program in the improvement of nectarines to enhance quality and market acceptability. The flowers of the Red Jim variety (U.S. Plant Pat. No. 4,518) were emasculated and pollinated with pollen from the variety May Grand (U.S. Plant Pat. No. 2,794). Seeds produced from this method were stratified and grown to a height of about 18 inches. These seedlings were bud grafted into dehorned orchard trees in the experimental orchard of Ito Packing Co., Inc., Reedley, Calif., for testing and selection. Each clone was budded into several limbs to make sure that it propagated true to type.

SUMMARY OF THE VARIETY

The instant variety developed through the above method was selected because of its uniquely deep red color of the skin, good flavor, aroma and texture, which contributes to its exceptional eating quality. The novel nectarine of the present invention has a much more attractive red color than other varieties ripening at the same period, and a maturity of approximately three weeks prior to a similar variety, which is the subject matter of a copending application.

BRIEF DESCRIPTION OF THE DRAWING

The accompanying drawing illustrates typical specimens of the fruit and foliage of my new variety as grown in the San Joaquin Valley of California. Two specimens are shown, one of which is in the side elevation and the other from the apex view; yet another specimen in a section side elevation illustrating the internal texture and color of the flesh and pit.

DETAILED DESCRIPTION

The following is a detailed description of my new variety. Each color reference is with respect to the

2

Maerz and Paul Dictionary of Colors, except in instances where terminology having generally accepted meaning is employed.

- 5 Parentage: Red Jim × May Grand F1.
- Propagation: Maintains its distinguishing characteristics through several bud grafts.
- Locality where grown and observed: Near Reedley, County of Fresno, Calif.

TREE

- 10 Size: Large, spreading, open, vase form.
- Vigor: Vigorous, productive.
- Regularity of bearing: Regular bearer.
- 15 Trunk: Medium size; medium texture.
- Leaves:
  - Color.—Top side, 32B10; underside, 311B.
  - Size.—Average length — 4.8 inches; average width of 1.2 inch; medium thickness.
  - Petiole.—Medium length; medium thickness.
  - Glandular characteristics.—Average number — four; opposite; medium size; reniform; position — usually two on at the base of the leaf blade near the attachment of the petiol and occasionally one on the petiole near the leaf blade.
  - Stipules.—Wanting.
- 25 Flower bud: Medium size.
- Flower: Large, Showy.
  - Color.—Pink.
  - 30 Anthers.—Red.
- Time of full bloom: With variety Red Jim (U.S. Plant Pat. No. 4,518).

FRUIT

- 35 Maturity: When described, firm ripe, July 25, 1988.
- Date of first picking: July 20, 1988.
- Date of last picking: July 30, 1988.
- Size: Average axial length 3.0 inches; average transverse width in suture plane 2 7/8 inches.
- Axial diameter.—About 2 7/8 inches.
- Form: Essentially round but compressed at the base. Often slightly larger on right half but may be larger on right half from the ventral suture.
- 45 Tendency to crack.—None observed.
- Suture.—Shallow in lower one-third of fruit but deepens in the upper one-third of the fruit to the apex. The apex being below the contour of the fruit. Dorsal suture present but very shallow



except near the apex where it deepens to meet ventral suture.  
*Ventral surface.*—Round.  
*Cavity.*—Round, medium depth.  
*Base.*—Flattened slightly.  
*Apex.*—Smooth with little or no point, recessed with the suture.

Skin:

*Thickness.*—Medium thick; medium texture.  
*Tendency to crack.*—None observed.  
*Bloom.*—Wanting.  
*Color.*—Base color yellow (10L6) but only around the base of the fruit where it is pressed against the fruit. Lighter portions are light orange (10LB) heavily speckled with red (6L4) over most of the surface. Much of the surface of red color (6L3) which deepens to a darker red (6L4) as the fruit matures.

Flesh:

*Texture.*—Very firm, meaty.  
*Color.*—Yellow with slight redish tinge (9J2). Considerable red around the pit extending through much of the flesh (5E6).  
*Juice.*—Average for cling stone nectarine.  
*Fibers.*—Considerable around pit with many running out into the flesh.  
*Ripening.*—Even. Stays firm on the tree for a long time after ripening.  
*Flavor.*—Good balance of acid and sugar. Sweeter than many nectarines at this season with soluble solids of 14% (refractometer).  
*Aroma.*—Very pronounced.  
*Eating quality.*—Excellent.

Stone:

*Adherence to flesh.*—Cling stone.

*Size.*—Average thickness, 11/16 inch; average width, 1 1/8 inch; average length, 1 3/8 inch.  
*Form.*—Oval.  
*Hilum.*—Small, oblong.  
*Dorsal edge.*—Shallow grooves.  
*Ventral edge.*—Narrow, very little wing toward the apex.  
*Surface.*—Irregular broad grooves, deeper than in most varieties.  
*Color.*—Brown (7L12) shaded to redish brown throughout (47L1).  
*Tendency to split.*—Seldom.

Use: Fresh market.

Keeping quality: Excellent.

Shipping quality: Excellent.

Resistance to disease: Similar to other glanded varieties of nectarines.

Although this novel variety of plum tree possesses the described characteristics as a result of the growing conditions in Fresno County, Calif., in the central portion of the San Joaquin Valley, it is to be understood that variations of the usual magnitude in characteristics incident to growing conditions, fertilization, pruning and pest control are to be expected.

Having thus described and illustrated my new variety of plum tree, what is claimed as new and described to be secured by Letters Patent is:

1. A new and distinct variety of nectarine tree with fruit of yellow flesh, cling stone type, substantially as herein shown and described, having a maturity period in the early part of the late season, and having fruit of good sugar content, the fruit being large with skin of a deep red color over much of the exterior, and flesh of a firm texture with good flavor.

\* \* \* \* \*

40

45

50

55

60

65

**U.S. Patent**

**Mar. 13, 1990**

**Plant 7,186**

