

United States Patent [19] Kientzler

- **IMPATIENS PLANT NAMED GRENADA** [54]
- Ludwig Kientzler, Gensingen, [75] Inventor: Germany
- [73] Assignee: Paul Ecke Ranch, Inc., Encinitas, Calif.
- [21] Appl. No.: 280,204
- [22] Filed: Jul. 25, 1994

Plant 9,343 **Patent Number:** [11] Oct. 24, 1995 **Date of Patent:** [45]

Primary Examiner—James R. Feyrer Attorney, Agent, or Firm—Foley & Lardner

[57] ABSTRACT

A new and distinct cultivar of New Guinea Impatiens named Grenada, particularly characterized by compact growth habit with excellent branching characteristics, large, coral pink flowers, early flowering and very floriferous habits, dark green foliage, and suitability to 4 inch and 6 inch pots, and 8 inch hanging basket cultures.

[51]	Int. Cl. ⁶
[52]	U.S. Cl
[58]	Field of Search

1 Drawing Sheet

The present invention relates to a new and distinct cultivar of plant known as New Guinea Impatiens. The new cultivar is known by the cultivar name Grenada, and was developed by the inventor Ludwig Kientzler in Gensingen, Federal Republic of Germany by crossing the cultivar designated QT 435 (seed parent) with the cultivar BD 22 (pollen parent). Both parents are proprietary cultivars used in the breeding program.

Asexual reproduction by terminal (stem tip) cuttings taken by me or under my supervision at Gensingen, Federal Republic of Germany, has shown that the unique features of this new Impatiens are stabilized and are reproduced true to type in successive propagations.

The following characteristics distinguish the new Impatiens from both its parent varieties and other cultivars of this ¹⁵ general type known and used in the floriculture industry:

ordinary dictionary significance are used. Parentage: Cross between QT 435 (seed) and BD 22 (pollen) Asexual reproduction:

- A. Cutting type.—Tip, with stems 2–3 cm long and developing to 4-5 cm after 21 days in propagation.
- B. Time to initiate roots.—8–10 days at 23° C.; nicely developed root mass in 18–21 days.
- C. Rooting habit.-Numerous, fibrous, adventitious roots from the stem.

Plant description:

- A. Form.—Symmetrical, bush shaped, flowering herb, with self-branching characteristics giving the plant a full appearance. Pinching is not recommended since it does not enhance branching and only delays flowering by removing flower buds.

1. Compact growth habit with excellent branching characteristics. Plants in 16.5 cm pots were 18 cm tall and 33 cm wide, eight weeks after first flowering.

2. Large, coral pink flowers. Open flowers measure up to 206.0 cm in diameter.

3. Early flowering and very floriferous. Plants in 16.5 cm pots were in flower 6 weeks after planting.

4. Dark green foliage.

5. Well suited both to 4" pot and 6" pot, and 8" hanging ²⁵ basket cultures.

Grenada is similar in many respects to Melissa, disclosed in U.S. Plant Pat. No. 7,837. The flower color of Grenada is a coral pink, similar in color to Melissa. The leaf color of Granada is similar to Melissa. Granada has a more compact ³⁰ growth habit, has large flowers and blooms earlier than Melissa. Reference is made to Chart A at the end of the specification which compares certain characteristics of Grenada and Melissa.

The accompanying colored photograph is a top perspec-³⁵ tive view of the new cultivar, showing color as true as it is reasonably possible to obtain in a colored reproduction of this type. The following is a detailed description of my new Impatiens cultivar based on plants grown under commercial ⁴⁰ practice in Encinitas, Calif. Plants were started as rooted cuttings and were transplanted in mid-November into 16.5 cm (6.5 inch) pots, one pot per plant. By mid-January, plants were growing vigorously and were in flower. The values, measurements and observations noted below were taken 45 from plants in full bloom in mid-February.

- B. Habit of growth.-Vigorous, self-branching habit, producing whorls of leaves and flowers. Growth is indeterminate and flowering is continuous.
- C. Foliage description.—Leaves are simple. Lower leaves on stems in whorls of 3-4. Higher on stem, leaves in whorls of 6-8. 1. Leaf shape: Broadly elliptic to ovate. Leaf blade tips are acuminate; bases of immature leaves are acute, mature leaves, attenuate. 2. Leaf blade size: Mature leaves 13–14 cm long and 4.5-5 cm wide. 3. Petiole length: 2.5-3 cm. 4. Leaf Margin: Ciliated and finely serrated. 5. Leaf texture: Slightly rugose. a. Upper surface: Glabrous. b. Under surface: Glabrous. 6. Leaf color: Dark green with a reddish midvein. a. Upper surface: Darker than 139A. b. Under surface: Greyed purple, near 183A. 7. Venation: Pinnate.

Flowering description:

A. Flowering habits.--Very floriferous. Flowering is continuous. Flowers develop progressively around the whorl of leaves, taking 5-7 days from buds which show color, to bloom. Flowers are single and large, lasting for 2-3 weeks.

Color references are made to the Royal Horticultural Society Colour Chart (RHS), except where general terms of B. Natural flowering season.—Flowering is indeterminate and occurs throughout the year. Quantity of flowers increases with increasing light intensity and duration. However, floriferousness may wane during hot summer days in temperate climates.

- C. Flower buds.—Ellipsoidal and covered with 3 sepals plus rudimentary sepals fused into the under surface of the superior petal. A spur originates from the base of the inferior sepal. Spur is reddish and 4.5-5 cm long.
- D. *Flowers borne*.—Singly, on red pedicels about 4–4.5 cm long.

Plant 9,343

3

- E. Quantity of flowers.—One flower per leaf. Flowers occur progressively around the whorl of leaves so that tight buds to mature flowers are visible at the same time.
- F. Petals.—Petals open nearly perpendicular to the 5 pedicel so that the plane of the flower surface is nearly flat. 1. Number of petals: Five (5) petals, all overlapping. 2. Shape: All petals are_heart-shaped. Superior petal has a broad base. Other petals have a pointed base. 3. Color: Coral pink, with a purplish- 10 _ red eye. a. Upper surface: Between 41B and 41C. Purplish-red eye near 61B. b. Under surface: Between 43C and 43D. 4. Flower size: Up to 6.0 cm in diameter. G. Reproductive organs.—Flowers are monoecious. 1. 15 Stamens: Five (5), broad and fused to form a tube around the ovary; red in color. 2. Anther: Hooded and cream colored. 3. Pollen: Cream colored. 4. Stigma: Five pointed star, colorless. 5. Styles: Very short. 6. Ovary: 5-celled, 4-5 mm long until fertil- 20 ized. Grows to 1 cm or longer after fertilization;

purple.

Resistance to disease: Good resistance to common stem and root diseases, Rhizoctonia and Pythium. Botrytis may be a problem in humid or mist propagation environments, but no greater than with other New Guinea Impatiens cultivars.

4

CHART A

	Grenada	Melissa
Flower Color	RHS 41 B-C	RHS 43 C-D
Plant Height	18 cm	25 cm
(chows the not)		

Plant Width	33 cm	48 cm	
Weeks to flower	6 wks	8 wks	
Flower Diameter	6 cm	5.5 cm	

I claim:

1. A new and distinct cultivar of New Guinea Impatiens named Grenada, as illustrated and described.

* * * * *

.

.

. 1

.

. .

· . · ·

U.S. Patent Oct. 24, 1995

Plant 9,343

. . . .

. . .

•

.





.

. .

1. ¹. -.



· · ·

.

· · ·

· .

.

· · .

.

.

• .