



US00PP10080P

United States Patent [19] Meier et al.

[11] Patent Number: Plant 10,080
[45] Date of Patent: Oct. 21, 1997

[54] 'BA 77 700' KENTUCKY BLUEGRASS

[75] Inventors: Virgil D. Meier; Eugene W. Mayer, both of Marysville, Ohio; J. Kevin Turner, Salem, Oreg.

[73] Assignee: OMS Investments, Inc., Wilmington, Del.

[21] Appl. No.: 680,167

[22] Filed: Jul. 15, 1996

[51] Int. Cl.⁶ A01H 5/00

[52] U.S. Cl. Plt./90.2

[58] Field of Search Plt./90.2

[56] **References Cited**

U.S. PATENT DOCUMENTS

P.P. 3,156	5/1972	Fuchigami et al.	Plt./90.2
P.P. 3,186	5/1972	Barenbrug et al.	Plt./90.2
P.P. 4,336	11/1978	Mayer et al.	Plt./90.2
P.P. 6,280	9/1988	Meier et al.	Plt./90.2

P.P. 6,537	1/1989	Meier et al.	Plt./90.2
P.P. 6,538	1/1989	Meier et al.	Plt./90.2
P.P. 6,585	2/1989	Meier et al.	Plt./90.2
P.P. 7,831	3/1992	Meier et al.	Plt./90.2
P.P. 8,490	12/1993	Meier et al.	Plt./90.2
P.P. 9,036	1/1995	Meier et al.	Plt./90.2
P.P. 9,209	7/1995	Meier et al.	Plt./90.2
P.P. 9,611	7/1996	Meier et al.	Plt./90.2
P.P. 9,848	4/1997	Meier et al.	Plt./90.2

Primary Examiner—James R. Feyrer
Attorney, Agent, or Firm—Jones, Day, Reavis & Pogue

[57] **ABSTRACT**

A variety of Kentucky Bluegrass having a medium to high level of resistance to a broad spectrum of serious diseases, including leaf spot, dollar spot, rusts, and necrotic ring spot; an attractive green color throughout the growing season; good early spring greening; medium to high quality turf formation under a wide variety of environmental conditions; a moderately wide blade; and a high level of seed yielding capacity.

1 Drawing Sheet

1

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to a new and distinct variety of *Poa pretensis* that has been designated Ba 77-700 Kentucky Bluegrass.

2. Description of Related Art

Kentucky Bluegrasses have been disclosed in U.S. Plant Pat. No. 3,156, issued May 9, 1972; U.S. Plant Pat. No. 3,186, issued May 23, 1972; U.S. Plant Pat. No. 4,336, issued Nov. 28, 1978; U.S. Plant Pat. No. 6,280, issued Sep. 6, 1988; U.S. Plant Pat. Nos. 6,537 and 6,538, issued on Jan. 17, 1989; U.S. Plant Pat. No. 6,585, issued Feb. 7, 1989; U.S. Plant Pat. No. 7,831, issued Mar. 17, 1992; U.S. Plant Pat. No. 8,490, issued Dec. 7, 1993; U.S. Plant Pat. No. 9,036, issued Jan. 3, 1995; U.S. Plant Pat. No. 9,209 issued Jul. 18, 1995; and pending U.S. plant pataent application Ser. No. 08/453,864, filed May 30, 1995 now U.S. Plant Pat. No. 9,611, Ser. No. 08/532,995, filed Sep. 22, 1995, now U.S. Plant Pat. No. 9,848 and Ser. No. 08/604,763 filed Feb. 23, 1996, as well as U.S. plant patent application Ser. No. 08/680,168, filed coextensively herewith.

SUMMARY OF THE VARIETY

Ba 77-700 plant material was selected from an old turf area on a farmstead in Union County, Ohio. After testing and observing this variety, it was determined to be a distinct variety, and it was asexually propagated by rhizomes, tillers and disseminules.

Seed of Ba 77-700 Kentucky Bluegrass was produced first at Marysville, Ohio and later at Gervais, Oreg. This seed was used to plant turf performance evaluation trials and later seed production fields. Asexual production of Ba 77-700 by propagules (tillers and rhizomes) and by disseminules (modified caryopses produced by apomixis) has consistently produced progeny plants indistinguishable from the mother plant. The apomixis level of Ba 77-700 is approximately 95% based upon examining seedling characteristics of

2

approximately 100 to 150 seedlings from different crop years in a growth chamber.

Ba 77-700 has a number of highly desirable characteristics including a medium to high level of resistance to Drechslera spp. that causes leaf spot, melting out and crown rot, Puccinia spp. that causes several types of rust infections, Schlerotinia homoeocarpa that causes dollar spot and Leptosphaeria korrae that causes necrotic ring spot. Ba 77-700 has an attractive leafy turf type growth habit; moderately wide leaf blades and an attractive dark green color which can be maintained throughout the entire growing season. Ba 77-700 demonstrates good early spring greening and winter color under mild winter conditions. Ba 77-700 is an overall good turfgrass performer as evidenced by medium to high scores for quality and color. Ba 77-700 has a high seed yield potential in the bluegrass seed production region of the northwestern United States.

In comparison with a number of other Kentucky Bluegrass varieties, Ba 77-700 seed is above average in length; and is average in width and rachilla length. The panicle is average in length; below average in width and whorl numbers and above average in branch numbers for both the lower and third whorls. Ba 77-700 has average size spikelets and glumes and below average number of florets. The peduncle is significantly longer and wider than many other grasses. The culm and top internode are of average length but the number of nodes on the culm are significantly lower.

Ba 77-700 has a flag leaf that is average in length, above average in width and significantly thicker than many other Kentucky Bluegrass varieties. The ligule and leaf margin hair are significantly longer and more abundant than many other Kentucky Bluegrass varieties. The vegetative leaf is average in length and width but above average in thickness. The ligule of the vegetative leaf is significantly shorter than many other grasses and the sheath color is above average.

BRIEF DESCRIPTION OF THE DRAWING

FIG. 1 is a Ba 77-700 Kentucky Bluegrass panicle; FIG. 2 is a Ba 77-700 Kentucky Bluegrass seed; and

FIG. 3 is a Ba 77-700 Kentucky Bluegrass plant during anthesis.

DETAILED DESCRIPTION OF THE VARIETY

77-700 Kentucky Bluegrass (*Poa pratensis* L.) is perennial with creeping rhizomes forming a dense turf. When plants overwinter in the field under freezing temperatures and are then brought into the greenhouse during late winter to continue growth undisturbed by clipping under moderate temperatures (60°–80° F.), culms are erect averaging 39.7 cm in length and 2.9 nodes per culm and the uppermost internode averages 9.7 cm in length. The peduncle averages 26.9 cm in length and 0.82 mm in width. The leaf averages 4.3 cm in length, 3.7 mm in width and 0.22 mm in thickness and the ligule 1.3 mm in length. The vegetative leaf averages 18.6 cm in length, 3.8 mm in width, 0.341 mm in thickness and a ligule length of 0.21 mm.

The panicle averages 77.5 mm in length, 56.1 mm in width, and 8.7 whorls. The lowest whorl averages 4.0 branches and the third whorl from the bottom of the panicle averages 3.5 branches. A spikelet in the lowest whorl averages 4.6 mm in length, 2.4 mm in width, 3.3 florets and the outer glume and inner glume average 2.7 and 3.3 mm in length, respectively. A spikelet from the third whorl from the bottom of the panicle averages 4.8 mm in length, 2.6 mm in width, 3.4 florets, and the outer glume and inner glume averages 2.8 and 3.3 mm in length, respectively.

After the seed has been conditioned, the lemma has a generally smooth keel with occasional short hairs and a few long fine hairs at the base. The conditioned seed of Ba 77-700 averages 3.05 mm in length, 0.88 mm in width, a rachilla length of 0.82 mm and an average quantity of lemma hairs. The seed count for Ba 77-700 averages approximately 1,065,922 seeds per pound.

Comparisons of Ba 77-700 with other Kentucky Bluegrass varieties in terms of seed numbers per pound are shown in Table 1 as follows:

TABLE 1

Comparison of Seed Numbers Per Pound of Ba 77-700 and Other Kentucky Bluegrass Varieties at Marysville, OH After Conditioning

Variety	Seeds Per Pound
Ba77-700	1,065,922
Abbey	1,003,037
Adelphi	1,383,976
Allure	1,359,914
America	1,659,824
Ascot	1,039,000
Baron	1,051,693
Bristol	1,270,821
Buckingham	978,000
Cannon	1,171,000
Chateau	1,300,105
Coventry	1,246,200
Eclipse	1,335,668
Fairfax	1,250,997
Georgetown	1,431,000
Gnome	1,017,641
Kelly	921,166
Marquis	1,054,642
Midnight	1,227,000
Nassau	1,127,130
Ram I	1,214,000
Touchdown	1,211,000
Victa	1,038,298
Viva	1,025,586

Ba 77-700 differs significantly morphologically from many other Kentucky Bluegrass varieties in regard to the following characteristics: (1) peduncle length and width; (2) culm nodes; (3) flag leaf thickness, ligule length and leaf margin hairs and (4) vegetative leaf ligule length.

Since environmental conditions such as soil and climate may influence morphological characteristics to some extent, comparisons of morphological characteristics of Ba 77-700 were made with other Kentucky Bluegrass varieties under like conditions and the comparisons are set forth in the following Tables 2–10:

TABLE 2

Morphological Comparison of Seed and Rachilla Measurements and Lemma Hair of Ba 77-700 and Other Kentucky Bluegrass Varieties at Marysville, OH After Conditioning

Variety	Seed		Rachilla mm	Lemma* Hair
	Length mm	Width mm		
Ba 77-700	3.05	0.88	0.82	3.8
Abbey	2.97	0.89	0.80	3.8
Adelphi	2.70	0.84	0.65	4.4
Allure	2.78	0.86	0.70	4.7
America	2.40	0.68	0.68	2.8
Ascot	2.99	0.85	0.82	3.9
Baron	3.08	0.81	0.71	5.0
Bristol	2.94	0.88	0.73	4.3
Buckingham	3.23	0.89	0.88	4.2
Cannon	3.00	0.94	0.86	4.9
Chateau	2.81	0.86	0.71	4.5
Coventry	2.71	0.81	0.70	4.0
Eclipse	2.77	0.83	0.68	3.5
Fairfax	2.77	0.89	0.68	4.4
Georgetown	2.94	0.82	0.74	4.9
Gnome	2.78	0.83	0.75	4.1
Kelly	3.07	0.89	0.75	4.2
Marquis	2.97	0.87	0.83	4.7
Midnight	2.94	0.76	0.78	5.7
Nassau	3.07	0.86	0.68	3.4
Ram I	3.23	0.89	0.80	6.0
Touchdown	2.93	0.88	0.71	4.6
Victa	3.00	0.80	0.82	3.5
Viva	3.04	0.91	0.80	4.4
LSD (.05)	0.16	0.05	0.13	0.86

*Rating Scale: 0–9; 9 = longest row of hairs

TABLE 3

Morphological Comparison of Panicles, Whorl Number and Whorl Branches of Ba 77-700 and Other Kentucky Bluegrass Varieties in the Greenhouse at Marysville, OH

Variety	Panicles			Number of Branches		
	Nodding*	Length mm	Width mm	Whorl Number	Lowest Whorl	Third Whorl
Ba 77-700	1.2	77.5	56.1	8.7	4.0	3.5
Abbey	1.6	80.4	59.3	9.3	3.8	3.3
Adelphi	1.8	106.8	76.4	10.0	3.6	3.0
Allure	2.2	60.8	47.6	8.4	3.9	3.0
America	2.3	67.7	51.1	8.8	3.3	3.5
Ascot	2.4	77.1	66.0	9.2	2.3	2.1
Baron	2.2	92.6	71.0	10.0	3.4	2.8
Bristol	2.0	85.5	61.5	8.4	2.8	2.7
Buckingham	2.0	88.8	62.0	9.3	2.1	2.1
Cannon	1.1	80.2	55.3	9.8	4.1	3.9
Chateau	2.9	65.2	57.3	8.5	3.3	2.8
Coventry	2.5	64.0	54.2	8.4	3.3	2.5
Eclipse	1.3	89.2	74.0	10.7	3.3	2.1
Fairfax	1.5	67.4	46.6	8.8	3.3	3.1

TABLE 3-continued

Morphological Comparison of Panicles, Whorl Number and Whorl Branches of Ba 77-279 and Other Kentucky Bluegrass Varieties in the Greenhouse at Marysville, OH						
Variety	Panicles				Number of Branches	
	Nodding*	Length mm	Width mm	Whorl Number	Lowest Whorl	Third Whorl
Georgetown	1.0	80.0	57.0	7.4	2.1	2.6
Gnome	1.1	80.6	56.0	10.6	4.6	3.9
Kelly	2.0	88.0	70.2	9.8	4.6	3.7
Marquis	1.1	82.0	63.0	10.3	3.9	3.6
Midnight	2.0	75.8	48.4	7.1	2.9	3.0
Nassau	2.2	91.2	68.6	10.0	2.5	2.1
Ram I	1.3	67.7	47.0	7.3	3.0	3.2
Touchdown	1.1	73.1	75.0	7.6	2.1	2.3
Victa	1.7	74.9	58.3	10.0	4.7	3.5
Viva	1.6	88.8	68.4	8.8	4.3	3.7
LSD (.05)	0.68	7.35	8.62	0.66	0.72	0.50

*Rate Scale: 1-9; 9 = most nodding

TABLE 4

Morphological Comparison of Spikelets and Florets of Ba 77-700 and Other Kentucky Bluegrass Varieties in the Greenhouse at Marysville, OH

Variety	Spikelet*				Number of Florets	
	Lowest Whorl		Third Whorl		Lowest Whorl	Third Whorl
	Length mm	Width mm	Length mm	Width mm		
Ba 77-700	4.6	2.4	4.8	2.6	3.3	3.4
Abbey	4.4	2.2	4.5	2.2	3.0	3.0
Adelphi	4.6	2.2	4.8	2.2	4.4	4.4
Allure	4.3	2.2	4.3	2.2	3.4	3.2
America	4.6	2.2	4.4	2.3	4.5	4.3
Ascot	5.4	2.6	5.3	2.9	3.1	3.1
Baron	5.4	2.8	5.4	3.1	4.0	4.2
Bristol	4.9	2.4	5.0	2.5	4.2	4.5
Buckingham	5.6	3.2	5.6	3.3	4.1	4.1
Cannon	4.7	2.4	4.6	2.6	2.7	2.7
Chateau	4.4	2.4	4.4	2.4	3.4	3.5
Coventry	4.4	2.2	4.5	2.4	3.2	3.5
Eclipse	4.6	2.4	4.6	2.4	3.6	3.6
Fairfax	4.6	2.3	4.7	2.4	3.2	3.3
Georgetown	5.0	2.3	5.0	2.6	4.8	4.9
Gnome	4.6	2.5	4.6	2.9	3.2	3.2
Kelly	5.1	2.5	5.2	2.8	3.7	3.8
Marquis	4.3	2.2	4.4	2.3	3.1	2.9
Midnight	5.3	2.5	5.5	2.4	4.4	4.7
Nassau	4.7	2.6	4.8	3.0	4.2	4.3
Ram I	5.4	2.6	5.3	2.8	3.6	3.4
Touchdown	5.1	2.9	4.8	2.8	4.1	4.0
Victa	4.5	2.5	4.5	2.4	3.3	3.2
Viva	4.6	2.5	4.7	2.6	2.6	2.8
LSD (.05)	0.41	0.35	0.38	0.35	0.58	0.53

*Microscope measurements

TABLE 5

Morphological Comparison of Glumes of Ba 77-700 and Other Kentucky Bluegrass Varieties in the Greenhouse at Marysville, OH

Variety	Glume Length (mm)				Glume Hairs*			
	Outer		Inner		Lowest Whorl			
	Lowest Whorl	Third Whorl	Lowest Whorl	Third Whorl	Outer	Inner	Outer	Inner
Ba 77-700	2.7	2.8	3.3	3.3	2.4	2.2	1.9	2.1
Abbey	2.7	2.7	3.1	3.1	0.6	0.9	1.0	1.0
Adelphi	2.7	2.6	3.0	3.1	0.6	2.7	1.1	1.9
Allure	2.6	2.7	3.0	3.0	2.5	3.0	2.5	2.5
America	2.1	2.1	2.5	2.5	0.2	0.5	0.3	0.3
Ascot	3.7	3.7	4.1	4.0	2.1	3.0	2.0	2.3
Baron	3.1	3.2	3.6	3.7	4.8	4.4	4.2	4.6
Bristol	2.8	3.0	3.3	3.4	1.0	2.2	0.8	2.2
Buckingham	3.5	3.3	3.9	3.8	2.6	3.2	2.2	2.8
Cannon	2.8	2.9	3.2	3.1	1.7	2.3	2.1	2.3
Chateau	2.8	2.9	3.1	3.2	1.6	2.6	2.0	1.8
Coventry	2.7	2.8	3.1	3.1	1.5	2.7	1.6	2.4
Eclipse	3.0	3.0	3.4	3.4	1.5	1.9	1.8	1.9
Fairfax	2.8	2.9	3.1	3.2	2.2	2.2	2.1	3.1
Georgetown	2.8	2.8	3.1	3.1	1.7	2.7	1.2	2.5
Gnome	2.8	2.9	3.3	3.3	1.0	1.9	1.2	1.6
Kelly	3.0	3.1	3.4	3.5	1.3	1.5	1.1	0.8
Marquis	2.8	2.7	3.1	3.1	0.9	1.1	1.7	1.8
Midnight	2.6	2.6	3.0	3.1	1.8	2.8	1.0	2.4
Nassau	2.6	2.7	2.9	3.0	2.1	2.6	4.2	4.4
Ram I	2.9	3.0	3.6	3.5	1.1	2.3	0.7	1.2
Touchdown	3.3	3.3	3.8	3.8	0.7	2.1	1.1	1.4
Victa	2.8	2.7	3.1	3.1	0.8	1.1	1.1	1.3
Viva	2.8	2.8	3.2	3.1	1.7	1.3	1.3	2.0
LSD (.05)	0.23	0.22	0.22	0.23	0.9	0.9	0.9	0.9

*Ratings Scale: 0-9; 9 - most hairs

TABLE 6

Morphological Comparison of Flag Leaves of Ba 77-700 and Other Kentucky Bluegrass Varieties in the Greenhouse at Marysville, OH

Variety	Length cm	Width mm	Thickness mm	Leaf Angle ^{1/}	Leaf Curve ^{2/}
Ba 77-700	4.3	3.7	0.22	2.0	4.6
Abbey	3.8	3.4	0.14	2.7	4.1
Adelphi	5.7	3.4	0.18	5.2	5.0
Allure	3.8	2.9	0.21	1.0	4.9
America	4.3	2.6	0.21	4.6	5.0
Ascot	4.7	3.5	0.29	2.1	4.5
Baron	7.0	4.2	0.15	9.2	4.4
Bristol	4.3	3.2	0.21	2.3	5.0
Buckingham	5.3	3.4	0.23	1.4	5.0
Cannon	4.3	3.5	0.22	1.9	5.0
Chateau	4.0	2.7	0.19	3.4	2.0
Coventry	3.7	3.0	0.14	0.7	4.3
Eclipse	3.9	3.2	0.19	8.7	5.0
Fairfax	4.3	3.2	0.20	1.0	4.9
Georgetown	5.7	2.8	0.22	1.9	5.0
Gnome	4.5	3.5	0.19	4.3	5.0
Kelly	4.8	4.1	0.18	3.1	4.4
Marquis	5.5	3.6	0.24	2.7	4.6
Midnight	3.8	2.6	0.19	1.1	5.0
Nassau	6.0	3.7	0.19	5.9	4.7
Ram I	3.5	3.1	0.19	1.0	4.8

TABLE 6-continued

Morphological Comparison of Flag Leaves of Ba 77-700 and Other Kentucky Bluegrass Varieties in the Greenhouse at Marysville, OH					
Variety	Length cm	Width mm	Thickness mm	Leaf Angle ^{1/}	Leaf Curve ^{2/}
Touchdown	4.1	2.7	0.16	3.4	4.1
Victa	3.9	3.7	0.27	2.1	4.6
Viva	4.5	3.6	0.18	5.1	4.4
LSD (.05)	1.0	0.4	0.02	2.5	0.4

^{1/}Degrees from the stem^{2/}Rating Scale: 1-9; 1 = curves up; 5 = no curve; 9 = curves down

TABLE 7

Morphological Comparison of Flag Leaves (Ligule Length, Hairs on the Leaf Margin and Ligule, and Sheath Color) of Ba 77-700 and Other Kentucky Bluegrass Varieties in the Greenhouse at Marysville, OH.

Variety	Ligule Length (mm)	Leaf Margin	Hairs ^{1/}	
			Li- gule	Sheath Color ^{2/}
Ba 77-700	1.3	1.5	2.3	1.8
Abbey	1.3	0.8	1.6	0.2
Adelphi	1.1	1.8	1.6	0.4
Allure	0.8	0.4	2.3	2.0
America	0.6	0.4	0.7	1.9
Ascot	1.5	0.6	0.7	0.9
Baron	1.2	1.0	3.8	2.6
Bristol	0.6	0.4	1.9	2.0
Buckingham	1.2	0.5	2.1	2.0
Cannon	1.3	0.4	3.6	1.5
Chateau	0.9	1.1	3.1	0.3
Coventry	0.9	0.8	1.7	2.0
Eclipse	1.1	1.0	1.0	0.2
Fairfax	1.0	0.1	2.5	1.7
Georgetown	0.6	0.6	2.0	2.0
Gnome	0.8	0.9	3.2	1.5
Kelly	1.5	1.1	2.5	2.2
Marquis	0.8	0.5	3.8	1.5
Midnight	0.4	0.4	0.3	2.0
Nassau	1.1	2.6	2.4	0.8
Ram I	0.7	0.5	1.3	2.0
Touchdown	1.1	1.0	1.0	7.7
Victa	1.3	1.0	0.8	0.5
Viva	1.5	0.4	2.2	0.5
LSD (.05)	0.17	0.47	0.7	0.76

^{1/}Rating Scale: 0-9; 0 = none; 9 = many^{2/}Rating Scale: 0-9; 9 = dark purple

TABLE 8

Morphological Comparison of Peduncles, Culms, Node Numbers Per Culm and Internode Length of Ba 77-700 and Other Kentucky Bluegrass Varieties in the Greenhouse at Marysville, OH

Variety	Peduncle Length cm	Peduncle Width mm	Culm Length cm	Nodes Per Culm	Top Internode Length (cm)
Ba 77-700	26.9	0.82	39.7	2.9	9.7
Abbey	23.2	0.65	41.3	4.1	11.0
Adelphi	24.5	0.66	40.0	4.0	10.4
Allure	17.9	0.61	28.0	3.1	7.7
America	16.2	0.62	31.8	3.1	11.0
Ascot	20.0	0.64	36.3	3.0	12.7
Baron	34.0	0.70	52.8	4.2	12.8
Bristol	22.9	0.73	43.5	4.0	12.6
Buckingham	24.7	0.71	44.9	3.7	13.4
Cannon	24.3	0.81	39.5	3.1	10.8
Chateau	20.7	0.61	38.9	4.4	9.9
Coventry	19.6	0.49	34.7	4.5	7.9

TABLE 8-continued

Morphological Comparison of Peduncles, Culms, Node Numbers Per Culm and Internode Length of Ba 77-700 and Other Kentucky Bluegrass Varieties in the Greenhouse at Marysville, OH

Variety	Peduncle Length cm	Peduncle Width mm	Culm Length cm	Nodes Per Culm	Top Internode Length (cm)
Eclipse	23.7	0.66	39.2	4.9	9.9
Fairfax	21.0	0.60	31.9	3.2	7.2
Georgetown	21.7	0.68	37.5	3.2	11.1
Gnome	19.4	0.84	35.3	3.1	9.2
Kelly	29.4	0.75	46.6	3.6	11.8
Marquis	27.1	0.83	34.1	3.4	7.8
Midnight	20.5	0.72	29.5	2.8	7.6
Nassau	25.0	0.66	36.0	3.9	7.4
Ram I	22.2	0.69	30.5	2.5	6.6
Touchdown	23.6	0.52	38.0	4.0	8.4
Victa	21.2	0.81	37.9	3.7	10.3
Viva	26.4	0.70	42.7	4.1	9.2
LSD (.05)	3.2	0.08	3.3	0.4	1.9

TABLE 9

Morphological Comparison of Vegetative Leaves (Length, Width, Thickness, Angle and Curvature) of Ba 77-700 and Other Kentucky Bluegrass Varieties in the Greenhouse at Marysville, OH

Variety	Length cm	Width mm	Thickness mm	Leaf	
				Angle ^{1/}	Curve ^{2/}
Ba 77-700	18.6	3.8	0.34	61.6	7.7
Abbey	20.7	3.6	0.28	51.3	7.4
Adelphi	19.4	3.7	0.28	67.0	4.6
Allure	18.1	3.5	0.30	53.0	6.2
America	21.4	3.1	0.32	66.0	6.4
Ascot	17.4	3.4	0.35	67.9	5.5
Baron	18.2	4.3	0.29	35.0	4.4
Bristol	18.2	3.7	0.32	65.5	6.9
Buckingham	18.9	4.5	0.36	50.9	5.0
Cannon	22.1	3.3	0.30	62.5	8.2
Chateau	23.1	3.3	0.33	39.7	5.4
Coventry	23.4	3.6	0.25	63.0	7.3
Eclipse	20.3	4.4	0.30	65.5	5.4
Fairfax	21.9	3.5	0.28	51.7	8.3
Georgetown	19.5	3.6	0.29	56.7	5.8
Gnome	18.2	2.9	0.29	43.2	6.6
Kelly	22.8	3.9	0.30	55.5	7.0
Marquis	19.7	3.7	0.38	46.4	7.8
Midnight	19.5	3.5	0.31	63.3	5.0
Nassau	16.3	3.7	0.29	63.2	4.9
Ram I	19.1	3.6	0.27	70.8	6.5
Touchdown	16.7	3.7	0.24	51.3	4.0
Victa	21.5	3.6	0.36	56.3	6.7
Viva	22.9	3.8	0.40	53.0	7.1
LSD (.05)	3.50	0.43	0.04	15.9	1.5

^{1/}Degrees from the stem^{2/}Rating scale: 1-9; 1 = curves up; 5 = no curve; 9 = curves down

TABLE 10

Morphological Comparison of Vegetative Leaves (Ligule Length, Hairs on the Leaf Margin and Ligule, Sheath Roughness and Color) of Ba 77-700 and Other Kentucky Bluegrass Varieties in the Greenhouse at Marysville, OH

Variety	Ligule Length mm	Hairs ^{1/}			
		Leaf Ligule	Leaf Margin	Sheath Roughness ^{2/}	Sheath Color ^{3/}
Ba 77-700	0.21	3.1	2.0	7.3	1.4
Abbey	0.32	4.4	2.8	7.7	0.0

TABLE 10-continued

Morphological Comparison of Vegetative Leaves (Ligule Length, Hairs on the Leaf Margin and Ligule, Sheath Roughness and Color) of Ba 77-700 and Other Kentucky Bluegrass Varieties in the Greenhouse at Marysville, OH

Variety	Ligule Length mm	Hairs ^{1/}			Sheath Roughness ^{2/}	Sheath Color ^{3/}
		Leaf Ligule	Leaf Margin	Sheath		
Adelphi	0.34	2.6	3.4	8.8	0.0	
Allure	0.14	2.0	1.6	8.3	1.9	
America	0.12	1.7	2.1	8.4	2.0	
Ascot	0.22	1.5	2.0	8.8	0.2	
Baron	0.33	4.6	3.6	7.4	0.4	
Bristol	0.12	2.6	3.0	8.9	2.0	
Buckingham	0.14	3.3	2.6	8.7	2.0	
Cannon	0.16	3.2	0.7	8.2	2.0	
Chateau	0.22	3.3	3.1	7.8	0.5	
Coventry	0.32	3.1	2.4	8.1	0.3	
Eclipse	0.38	2.8	2.6	9.0	0.2	
Fairfax	0.14	2.7	1.4	8.1	1.7	
Georgetown	0.14	2.5	2.4	8.5	2.0	
Gnome	0.23	1.6	0.8	7.5	1.4	
Kelly	0.34	4.6	2.4	8.8	0.0	
Marquis	0.21	3.6	1.1	6.8	1.1	
Midnight	0.10	1.3	1.6	7.7	2.0	
Nassau	0.34	2.5	3.3	8.5	1.4	
Ram I	0.09	1.8	2.0	7.7	2.0	
Touchdown	0.29	1.4	1.3	7.9	8.0	
Victa	0.26	2.3	1.7	7.9	0.1	
Viva	0.23	5.2	4.2	8.5	0.0	
LSD (.05)	0.04	0.79	0.70	0.89	0.57	

^{1/}Rating Scale: 0-9; 0 = none; 9 = many

^{2/}Rating Scale: 1-9; 1 = rough; 9 = smooth

^{3/}Rating Scale: 0-9; 0 = no color; 9 = dark purple color

Ba 77-700 has performed well as exhibited by medium to high turf quality ratings throughout the United States in comparison to other Kentucky Bluegrass varieties. Ba 77-700 has an attractive green color which can be maintained throughout the growing season.

With regard to a comparative analysis conducted for purposes of determining color of Ba 77-700 plants relative to other Kentucky Bluegrass varieties, readings were taken of the vegetative color of Ba 77-700 during mid-October while the turf was actively growing with adequate nutrient and water availability. The readings were taken in full sun with several actively growing leaves being compared, one at a time, utilizing color chips from the Munsell Book of Color as a reference. On this basis, the color of Ba 77-700 was determined to be 5 GY 4/6. During the same time period, the color of similar leaves of other Kentucky Bluegrass varieties were determined by the same procedure to be as follows: Victa — 5 GY 4/4; Bristol — 7.5 GY 4/4; Ascot — 5 GY 4/4; Midnight — 7.5 GY 4/4; Abbey — 5 GY 4/6; and Buckingham — 7.5 GY 4/4. However, it should be noted that the general apparent color of turf does not always correlate directly with the color of the individual actively growing leaves within the turf and that turf color varies with nutrient level and time of year with some varieties being darker or lighter green depending on such factors.

Ba 77-700 demonstrates good early spring greening and winter color in mild winter climates.

Comparisons of Ba 77-700 with other varieties for quality, genetic color, spring greening and winter color, living ground cover, leaf texture and seedling vigor are set forth hereinafter in Tables 11-17:

TABLE 11

A Comparison of Quality of Ba 77-700 and Other Kentucky Bluegrass Varieties at Five (5) Test Locations in the U.S.

Variety	Locations				
	Fort Collins, Co	Lexington, Kentucky	Adelphia, NJ	Kingston, RI	Beltsville, MD
Ba 77-700	6.3	6.5	4.8	5.7	6.5
A-34	5.2	5.8	5.2	5.6	6.7
Abbey	6.3	6.2	4.7	5.7	6.3
Able I	6.3	6.9	6.2	5.8	5.9
Allure	5.9	5.4	4.9	5.3	6.5
Ascot	6.7	6.5	5.9	5.7	6.0
Banff	6.1	5.8	5.6	5.0	5.2
Baron	5.3	6.0	4.6	5.7	6.5
Buckingham	5.6	5.9	5.0	4.6	5.3
Challenger	6.1	6.6	5.4	5.2	6.6
Cannon	6.1	6.0	4.9	5.4	6.4
Classic	5.8	5.8	5.5	5.4	5.6
Coventry	6.1	5.6	4.7	5.0	6.6
Eclipse	6.6	6.5	6.3	6.1	6.8
Estate	5.8	4.8	5.0	5.2	6.3
Fairfax	5.9	5.1	5.1	5.3	6.8
Georgetown	5.8	5.9	5.1	5.3	5.3
Glade	6.1	6.4	5.1	5.8	6.7
Gnome	5.9	6.2	4.6	5.4	6.5
Haga	5.6	6.0	4.8	5.2	6.0
Kelly	5.9	6.1	4.5	5.2	6.4
Kenblue	3.7	5.0	2.4	4.9	5.4
Liberty	6.2	5.6	5.7	5.3	5.9
Marquis	6.2	6.9	4.5	5.6	6.4
Merion	5.1	5.9	4.0	5.4	5.7
Merit	6.1	6.5	4.2	5.7	6.3
Midnight	7.1	8.2	6.7	5.1	7.5
Monopoly	4.7	5.5	5.0	5.5	5.9
Nassau	5.8	6.5	4.7	5.2	4.9
Ram I	6.1	5.0	4.6	5.7	6.9
South Dakota	4.1	4.3	2.3	4.6	5.2
Touchdown	6.4	5.8	5.3	5.2	6.4
Viva	6.1	6.3	4.2	5.9	6.3
LSD (.05)	0.7	1.0	0.7	0.7	0.8

Quality Rating Scale: 1-9; 9 = ideal turf

TABLE 12

A Comparison of Genetic Color of Ba 77-700 and Other Kentucky Bluegrass Varieties in Three Test (A-C) Conducted at Ten (10) Locations in the US

Variety	TESTS (Annual Means)		
	A	B	C
Ba 77-700	5.7	5.5	5.8
A-34	5.0	5.4	4.6
Abbey	5.6	5.9	5.8
Able I	6.6	6.8	6.8
Allure	5.7	5.2	5.6
Ascot	6.9	6.8	7.2
Banff	5.1	6.0	5.5
Baron	6.0	6.0	5.9
Buckingham	7.0	7.2	7.2
Cannon	6.0	5.7	6.4
Challenger	6.0	6.2	5.9
Classic	5.2	5.9	5.2
Coventry	5.9	5.6	5.8
Eclipse	6.3	6.6	6.4
Estate	5.8	5.3	5.3
Fairfax	6.0	5.4	5.4
Georgetown	5.2	6.1	5.3
Glade	6.7	6.3	6.7
Gnome	6.0	6.1	5.8
Haga	5.1	5.8	5.3
Kelly	5.6	5.7	6.1

TABLE 12-continued

A Comparison of Genetic Color of Ba 77-700 and Other Kentucky Bluegrass Varieties in Three Test (A-C) Conducted at Ten (10) Locations in the US

Variety	TESTS (Annual Means)		
	A	B	C
Kenblue	4.6	5.0	5.3
Liberty	5.9	4.0	6.1
Marquis	6.1	6.1	6.4
Merion	5.7	5.8	5.4
Merit	5.7	5.7	6.2
Midnight	7.1	7.3	7.7
Monopoly	4.8	4.9	4.8
Nassau	6.0	6.6	6.5
Ram I	6.7	5.9	6.6
South Dakota	4.5	4.8	4.7
Touchdown	5.5	5.5	6.3
Viva	5.5	5.5	5.9
LSD (.05)	0.5	0.4	0.7

Rating Scale: 1-9; 9 = dark green.

TABLE 13

A Comparison of Spring Greenup of Ba 77-700 and Other Kentucky Bluegrass Varieties at Martinsville, New Jersey

Variety	Spring Greenup
Ba 77-700	6.0
A-34	6.0
Abbey	5.3
Able I	6.3
Allure	4.3
Ascot	3.7
Banff	3.7
Baron	4.0
Buckingham	5.0
Cannon	5.3
Challenger	5.3
Classic	5.3
Coventry	4.3
Eclipse	4.7
Estate	5.0
Fairfax	4.0
Georgetown	4.7
Glade	4.0
Gnome	4.7
Haga	5.7
Kelly	4.0
Kenblue	3.0
Liberty	4.7
Marquis	5.0
Merion	3.7
Merit	4.0
Midnight	2.7
Monopoly	4.7
Nassau	5.3
Ram I	5.0
South Dakota	4.0
Touchdown	5.7
Viva	5.0
LSD (.05)	1.8

Spring Greenup Rating Scale: 1-9; 9 = Completely green

TABLE 14

A Comparison of Winter Color of Ba 77-700 and Other Kentucky Bluegrass Varieties in Two Tests (A-B).

Variety	A ^{1/}	B ^{2/}
Ba 77-700	5.0	5.0
A-34	6.7	2.0
Abbey	1.7	6.3
Able I	6.3	3.3
Allure	6.0	6.3
Ascot	4.7	4.3
Banff	9.0	3.7
Baron	2.3	5.7
Buckingham	5.3	6.3
Cannon	2.3	6.0
Challenger	7.3	4.3
Classic	9.0	3.3
Coventry	7.0	3.7
Eclipse	4.7	3.7
Estate	5.7	5.0
Fairfax	6.0	4.5
Georgetown	8.0	3.3
Glade	1.7	3.7
Gnome	1.0	5.7
Haga	9.0	3.0
Kelly	2.3	5.7
Kenblue	1.3	3.0
Liberty	7.3	3.3
Marquis	2.0	5.0
Merion	6.7	3.3
Merit	3.7	6.0
Midnight	3.0	1.7
Monopoly	4.3	2.7
Nassau	7.7	5.0
Ram I	1.7	3.7
South Dakota	2.3	4.0
Touchdown	3.0	2.3
Viva	2.3	4.0
LSD (.05)	1.9	2.1

Winter Color Rating Scale: 1-9; 9 = Complete color retention

^{1/}Location of Test A: North Brunswick, NJ

^{2/}Location of Test B: Post Falls, ID

TABLE 15

A Comparison of Percent Living Ground Cover in the Fall of Ba 77-700 and Other Kentucky Bluegrass Varieties at Haymarket, VA

Variety	% Ground Cover
Ba 77-700	70.0
A-34	30.0
Abbey	68.3
Able I	28.3
Allure	83.3
Ascot	11.7
Banff	21.7
Baron	71.0
Buckingham	30.0
Cannon	75.0
Challenger	60.0
Classic	31.7
Coventry	65.0
Eclipse	48.3
Estate	71.7
Fairfax	70.0
Georgetown	33.3
Glade	68.3
Gnome	70.0
Haga	53.3
Kelly	83.3
Kenblue	94.3
Liberty	33.3
Marquis	88.3
Merion	81.7

TABLE 15-continued

A Comparison of Percent Living Ground Cover in the Fall of Ba 77-700 and Other Kentucky Bluegrass Varieties at Haymarket, VA

Variety	% Ground Cover
Merit	71.7
Midnight	38.3
Monopoly	85.0
Nassau	36.7
Ram I	65.0
South Dakota	86.0
Touchdown	33.3
Viva	85.0
LSD (.05)	29.9

TABLE 16

A Comparison of Leaf Texture of Ba 77-700 and Other Kentucky Bluegrass Varieties in Two Tests (A-B)

Variety	A ^{1/}	B ^{2/}
Ba 77-700	4.0	4.0
Abbey	5.0	4.3
Able I	5.7	6.3
Allure	4.3	4.0
Ascot	5.3	6.0
A-34	4.3	5.2
Banff	5.3	5.2
Baron	5.0	4.2
Buckingham	4.0	3.7
Cannon	4.0	4.0
Challenger	5.0	4.5
Classic	5.3	5.3
Coventry	5.0	4.2
Eclipse	5.7	5.7
Estate	4.3	3.8
Fairfax	4.7	4.0
Georgetown	6.0	4.8
Glade	5.0	4.6
Gnome	5.3	4.2
Haga	5.7	5.2
Kelly	4.3	4.0
Kenblue	6.0	5.8
Liberty	5.0	4.7
Marquis	4.3	4.2
Merion	4.0	4.2
Merit	4.7	4.0
Midnight	6.7	6.0
Monopoly	5.0	4.5
Nassau	5.0	4.3
Ram I	5.0	4.8
South Dakota	3.7	4.8
Touchdown	5.3	5.8
Viva	4.3	4.0
LSD (.05)	0.9	0.8

Leaf Texture Rating Scale: 1-9; 9 = very fine

^{1/}Test Location: Martinsville, NJ^{2/}Test Location: Martinsville, NJ; and pooled data from Halsey, Hubbard and Gervais, OR

TABLE 17

A Comparison of Seeding Vigor of Ba 77-700 and Other Kentucky Bluegrass Varieties at Six Locations in The U.S. and Canada^{1/}

Variety	Seedling Vigor (Mean)
Ba 77-700	6.8
A-34	6.0
Abbey	6.4
Able I	6.3

TABLE 17-continued

A Comparison of Seeding Vigor of Ba 77-700 and Other Kentucky Bluegrass Varieties at Six Locations in The U.S. and Canada^{1/}

Variety	Seedling Vigor (Mean)
Allure	6.2
Ascot	6.2
Banff	7.7
Baron	6.7
Buckingham	5.3
Cannon	6.3
Challenger	5.6
Classic	7.1
Coventry	6.1
Eclipse	6.1
Estate	6.1
Fairfax	6.4
Georgetown	7.0
Glade	6.7
Gnome	6.3
Haga	7.1
Kelly	6.6
Kenblue	7.4
Liberty	6.4
Marquis	6.3
Merion	2.6
Merit	6.4
Midnight	5.4
Monopoly	7.1
Nassau	6.4
Ram I	6.5
South Dakota	6.7
Touchdown	5.8
Viva	6.3
LSD (.05)	0.7

Rating Scale: 1-9; 9 = Maximum vigor.

^{1/}Locations: Fort Collins, CO; Post Falls, ID; North Brunswick, NJ; Adelphia, NJ; Richmond Hills, Ontario; pooled data from Halsey, Hubbard and Gervais, OR.

Turf diseases are one of the major causes of inconsistent and poor turf performance. Ba 77-700 has been found to have a medium to high level of resistance to leaf spot and melting out caused by *Drechslera poae* (formerly called *Helminthosporium vagans*). Ba 77-700 furthermore has been found to have a medium to high level of resistance to dollar spot caused by *Sclerotinia homoeocarpa* and to several rust diseases caused by *Puccinia* spp. as well as to necrotic ring spot caused by *Leptosphaeria korrae*.

Comparisons of disease incidence of Ba 77-700 as compared with other Kentucky Bluegrass varieties in regard to leaf spot, dollar spot, rusts and necrotic ring spot are presented in Tables 18-23 as follows:

TABLE 18

A Comparison of Leaf Spot Incidence in Ba 77-700 and Other Kentucky Bluegrass Varieties in Two Tests (A-B) Conducted at Four Locations in the U.S.

Variety	Means	
	Test A ^{1/}	Test B ^{2/}
Ba77-700	7.2	5.7
A-34	7.2	5.4
Abbey	7.8	5.1
Able I	7.8	7.1
Allure	7.7	5.8
Ascot	8.1	7.4
Banff	4.8	5.9
Baron	7.1	5.4

TABLE 18-continued

A Comparison of Leaf Spot Incidence in Ba 77-700 and Other Kentucky Bluegrass Varieties in Two Tests (A-B) Conducted at Four Locations in the U.S.

Variety	Means	
	Test A ^{1/}	Test B ^{2/}
Buckingham	8.3	6.8
Cannon	8.0	5.8
Challenger	8.0	6.9
Classic	7.8	5.4
Coventry	6.8	5.8
Eclipse	7.9	7.4
Estate	7.9	5.9
Fairfax	6.6	4.9
Georgetown	8.0	6.3
Glade	4.8	4.8
Gnome	7.6	5.7
Haga	7.2	5.9
Kelly	7.2	5.1
Kenblue	1.2	1.8
Liberty	8.1	6.8
Marquis	7.4	5.6
Merion	8.0	7.3
Merit	7.4	5.4
Midnight	8.1	6.6
Monopoly	7.1	5.2
Nassau	8.2	6.2
Ram I	3.4	4.9
South Dakota	2.2	1.8
Touchdown	7.4	6.3
Viva	7.4	5.1
LSD (.05)	0.9	0.7

Rating Scale: 1-9; 9 = no disease

^{1/}Locations: Post Falls, ID; North Brunswick and Adelphia, NJ

^{2/}Locations: Post Falls, ID; North Brunswick and Adelphia, NJ; Marysville, OH

TABLE 19

A Comparison of Dollar Spot Disease Incidence of Ba 77-700 and Other Kentucky Bluegrass Varieties in Two Tests Conducted in the U.S.

Variety	Means	
	Test A ^{1/}	Test B ^{2/}
Ba 77-700	7.0	7.0
A-34	6.7	6.0
Abbey	6.8	6.3
Able I	6.8	6.1
Allure	6.4	5.3
Ascot	7.8	7.0
Banff	6.4	6.2
Baron	6.9	6.8
Buckingham	7.9	4.8
Canon	6.6	6.4
Challenger	6.4	4.7
Classic	7.4	6.3
Coventry	5.9	5.9
Eclipse	7.5	7.2
Estate	5.3	5.6
Fairfax	5.8	6.0
Georgetown	6.9	6.1
Glade	7.0	6.4
Gnome	6.7	6.1
Haga	6.3	6.2
Kelly	7.3	6.3
Kenblue	6.8	6.8
Liberty	7.3	6.8
Marquis	6.8	6.7
Merion	6.9	6.1
Merit	6.8	6.7
Midnight	8.0	6.6
Monopoly	7.1	6.7

TABLE 19-continued

A Comparison of Dollar Spot Disease Incidence of Ba 77-700 and Other Kentucky Bluegrass Varieties in Two Tests Conducted in the U.S.

Variety	Means	
	Test A ^{1/}	Test B ^{2/}
Nassau	7.4	6.7
Ram I	6.0	4.9
South Dakota	7.3	5.8
Touchdown	6.8	6.1
Viva	7.1	6.2
LSD (.05)	1.1	0.8

Rating Scale: 1-9; 9 = no disease.

^{1/}Locations: Urbana, IL; Adelphia, NJ; Kingston, RI; Haymarket, VA

^{2/}Locations: Urbana, IL; Silver Spring, MD; Kingston, RI

TABLE 20

A Comparison of Stem Rust Disease Incidence in Ba 77-700 and Other Kentucky Bluegrass Varieties at Adelphia, NJ

Variety	Stem Rust
Ba 77-700	7.3
A-34	4.3
Abbey	7.0
Able I	3.3
Allure	6.3
Ascot	7.3
Banff	7.7
Baron	7.0
Buckingham	5.7
Cannon	7.3
Challenger	4.0
Classic	8.0
Coventry	6.7
Eclipse	4.0
Estate	6.3
Fairfax	6.3
Georgetown	8.7
Glade	7.7
Gnome	7.0
Haga	8.0
Kelly	8.0
Kenblue	5.7
Liberty	7.7
Marquis	6.7
Merion	1.0
Merit	6.3
Midnight	8.0
Monopoly	6.3
Nassau	8.0
Ram I	8.0
South Dakota	4.3
Touchdown	2.3
Viva	6.7
LSD (.05)	1.4

Rating Scale: 1-9; 9 = no disease

TABLE 21

A Comparison of Leaf Rust Disease Incidence in Ba 77-700 and Other Kentucky Bluegrass Varieties at Kingston, R.I.

Variety	Leaf Rust
Ba 77-700	8.0
A-34	6.0
Abbey	6.3
Able I	7.0
Allure	6.0
Ascot	6.0
Banff	7.0

TABLE 21-continued

A Comparison of Leaf Rust Disease Incidence in Ba 77-700 and Other Kentucky Bluegrass Varieties at Kingston, R.I.	
Variety	Leaf Rust
Baron	8.3
Buckingham	6.7
Cannon	6.3
Challenger	5.7
Classic	8.7
Coventry	7.0
Eclipse	5.7
Estate	6.3
Fairfax	6.0
Georgetown	7.7
Glade	7.0
Gnome	6.7
Haga	7.0
Kelly	5.0
Kenblue	6.7
Liberty	7.3
Marquis	6.3
Merion	2.3
Merit	6.0
Midnight	6.0
Monopoly	5.3
Nassau	7.0
Ram I	6.7
South Dakota	6.7
Touchdown	3.7
Viva	5.7
LSD (.05)	2.4

Rating Scale: 1-9; 9 = no disease.

TABLE 22

A Comparison of Stripe Rust Disease Incidence in Ba 77-700 and Other Kentucky Bluegrass Varieties in Tests Conducted at Three Locations ^{1/}	
Variety	Stripe Rust
Ba 77-700	5.0
A-34	5.0
Abbey	4.3
Able I	6.0
Allure	4.3
Ascot	6.0
Banff	6.0
Baron	6.0
Buckingham	5.7
Cannon	4.7
Challenger	5.3
Classic	6.3
Coventry	5.0
Eclipse	6.0
Estate	5.7
Fairfax	5.7
Georgetown	6.3
Glade	3.0
Gnome	4.7
Haga	6.3
Kelly	5.0
Kenblue	3.3
Liberty	5.3
Marquis	5.3
Merion	3.7
Merit	5.0
Midnight	4.7
Monopoly	5.3
Nassau	5.3
Ram I	6.0
South Dakota	4.0

TABLE 22-continued

A Comparison of Stripe Rust Disease Incidence in Ba 77-700 and Other Kentucky Bluegrass Varieties in Tests Conducted at Three Locations ^{1/}	
Variety	Stripe Rust
Touchdown	3.0
Viva	5.0
LSD (.05)	1.2

Rating Scale: 1-9; 9 = no disease

^{1/}Pooled data from Halsey, Hubbard and Gervais, Oregon.

TABLE 23

A Comparison of Necrotic Ring Spot Incidence in Ba 77-700 and Other Kentucky Bluegrass Varieties at Post Falls, ID	
Variety	Necrotic Ring Spot
Ba 77-700	7.3
A-34	6.0
Abbey	7.3
Able I	7.3
Allure	3.3
Ascot	8.0
Banff	7.7
Baron	7.3
Buckingham	8.7
Cannon	7.0
Challenger	8.3
Classic	6.7
Coventry	5.0
Eclipse	7.7
Estate	4.7
Fairfax	3.5
Georgetown	7.7
Glade	8.3
Gnome	7.3
Haga	7.7
Kelly	7.3
Kenblue	5.7
Liberty	6.7
Marquis	7.7
Merion	5.3
Merit	7.7
Midnight	8.3
Monopoly	6.7
Nassau	7.7
Ram I	7.0
South Dakota	7.0
Touchdown	6.3
Viva	7.7
LSD (.05)	1.9

Rating Scale: 1-9; 9 = no disease

Ba 77-700 has shown good drought tolerance and seed yielding capability. Comparisons for drought tolerance and seed yields of Ba 77-700 with other Kentucky Bluegrasses are presented in Tables 24-26.

TABLE 24

A Comparison of Drought Tolerance of Ba 77-700 and Other Kentucky Bluegrass Varieties at Haymarket, VA	
Variety	Drought Tolerance
Ba77-700	5.3
A-34	3.0
Abbey	4.3
Able I	3.3
Allure	4.0
Ascot	4.0
Banff	3.3

TABLE 24-continued

A Comparison of Drought Tolerance of Ba 77-700 and Other Kentucky Bluegrass Varieties at Haymarket, VA	
Variety	Drought Tolerance
Baron	4.0
Buckingham	4.0
Cannon	4.0
Challenger	3.3
Classic	3.7
Coventry	3.7
Eclipse	4.3
Estate	4.3
Fairfax	4.3
Georgetown	3.7
Glade	5.0
Gnome	3.7
Haga	3.0
Kelly	4.0
Kenblue	4.0
Liberty	3.0
Marquis	4.7
Merion	3.7
Merit	4.7
Midnight	3.7
Monopoly	4.0
Nassau	3.3
Ram I	4.3
South Dakota	4.0
Touchdown	4.0
Viva	4.7
LSD (.05)	1.1

Dormancy Rating Scale: 1-9; 9 = no dormancy.

TABLE 25

A Comparison of Drought Recovery of Ba 77-700 and Other Kentucky Bluegrass Varieties in Two Tests (A-B)		
Variety	Test A ^{1/}	Test B ^{2/}
Ba 77-700	4.8	3.0
A-34	3.3	1.0
Abbey	4.5	2.7
Able I	4.3	1.3
Allure	4.7	5.3
Ascot	3.2	1.0
Banff	3.5	1.0
Baron	5.5	4.0
Buckingham	2.8	2.0
Cannon	4.3	2.7
Challenger	3.5	3.3
Classic	4.2	1.3
Coventry	4.7	2.7
Eclipse	3.3	2.0
Estate	5.5	3.0
Fairfax	5.0	3.0
Georgetown	3.5	1.0
Glade	4.2	2.3
Gnome	4.8	2.7

TABLE 25-continued

A Comparison of Drought Recovery of Ba 77-700 and Other Kentucky Bluegrass Varieties in Two Tests (A-B)		
Variety	Test A ^{1/}	Test B ^{2/}
Haga	4.0	2.7
Kelly	4.5	4.3
Kenblue	4.8	4.7
Liberty	4.2	1.7
Marquis	4.2	4.7
Merion	3.5	4.3
Merit	5.7	3.7
Midnight	4.5	1.0
Monopoly	4.7	5.3
Nassau	5.0	2.3
Ram I	4.3	3.0
South Dakota	3.7	3.3
Touchdown	4.3	1.3
Viva	4.8	3.7
LSD (.05)	1.2	1.8

Drought Recovery Rating Scale: 1-9; 9 = Complete recovery
^{1/}Locations: Carbondale, IL; and pooled data from Halsey, Hubbard and Gervais, OR
^{2/}Locations: Haymarket, VA

TABLE 26

A Comparison of Seed Yield in Pounds Per Acre of Ba 77-700 and Other Kentucky Bluegrass Varieties in Two (2) Tests		
Variety	Test A ^{1/}	Test B ^{2/}
Ba 77-700	1495	1569
Abbey	—	1549
Allure	859	—
Bristol	862	—
Buckingham	865	748
Cannon	—	1551
Coventry	—	1046
Fairfax	—	1159
Victa	1342	—
Viva	1382	1612
LSD (.05)	216	160

^{1/}Test Location: Gervais, OR
^{2/}Test Location: La Grande, OR

What is claimed is:

1. A variety of Kentucky Bluegrass plant, substantially as shown and described, characterized by a medium to high level of resistance to leaf spot, dollar spot, rusts, and necrotic ring spot; an attractive green color throughout the growing season; good early spring greening; medium to high quality turf formation under a wide variety of environmental conditions; a moderately wide blade; and a high level of seed yielding capacity.

* * * * *

FIG. 1



FIG. 2



FIG. 3

