



US00PP10095P

United States Patent [19]  
Henny

[11] Patent Number: Plant 10,095  
[45] Date of Patent: Nov. 4, 1997

[54] MANDEVILLA: JUMBO BLUSH  
[75] Inventor: Richard J. Henny, Tavares, Fla.  
[73] Assignee: Gem Ornamentals, Inc., Tavares, Fla.  
[21] Appl. No.: 637,121  
[22] Filed: Apr. 24, 1996  
[51] Int. Cl.<sup>6</sup> ..... A01H 5/00  
[52] U.S. Cl. .... Plt./54.1  
[58] Field of Search ..... Plt./54.1

[56] References Cited  
U.S. PATENT DOCUMENTS

P.P. 8,479 11/1993 Henny ..... Plt./54.1  
P.P. 8,842 7/1994 Henny ..... Plt./54.1

OTHER PUBLICATIONS  
Bailey, L. H., et al., "Mandevilla", *Hortus Third*, MacMillan  
Publ. Co., Inc., N.Y. (1976) p. 711.

Graf, A. B., (Ed.), "Mandevilla", *Exotica*, Roehrs Co.,  
Rutherford, N.J., (1967), p. 1651.

Huxley, A., et al. (Eds.), "Mandevilla", *The New Royal  
Horticultural Society Dictionary of Gardening* (1992), vol.  
3, p. 187.

Primary Examiner—James R. Feyrer  
Attorney, Agent, or Firm—Godfrey & Kahn, S.C.

[57] ABSTRACT

A new and distinct variety of Mandevilla characterized by  
large funnelform to trumpet-shaped flowers that open pale  
pink, fade to white within one to three days after anthesis  
and remain on the plant an additional two to three days.  
Flower color is highlighted by a bright red-purple throat that  
becomes somewhat lighter as the flower ages.

1 Drawing Sheet

1

BACKGROUND INFORMATION

The present invention relates to a new and distinct hybrid  
of Mandevilla that originated from self-pollination of  
Mandevilla 'Alice Du Pont', a member of the family Apo-  
cynaceae. The parent is believed to be an interspecific  
hybrid; the seed parent of which was *Mandevilla splendens*.  
This new Mandevilla is the result of deliberate pollinations  
performed by the inventor in a greenhouse at Gem Orna-  
mentals in Tavares, Fla. in 1990. This new variety was  
selected out for further evaluation because it had large, pale  
pink flowers with a bright red-purple throat. The flowers  
faded to white with age. The new variety has been repro-  
duced by asexual propagation (cuttings) at the same loca-  
tion. Each of the progeny exhibits identical characteristics to  
the original mother plant establishing this hybrid as repro-  
ducible and true to type.

This new hybrid is unlike any Mandevilla commercially  
available nor like any of those described by Graf or Bailey  
(see references) as evidenced by the following unique com-  
binations of characteristics.

1. Large funnelform to trumpet-shaped flowers that open  
pale pink, fade to white within one to three days after  
anthesis and remain on the plant an additional two to three  
days.

2. Flower color is highlighted by a bright red-purple throat  
that becomes somewhat lighter as the flower ages.

3. Elliptic-ovate shaped, leathery, shiny dark green and  
evergreen leaves.

DESCRIPTION OF THE DRAWING

This new variety of Mandevilla is illustrated by the  
accompanying photographic drawing.

FIG. 1 is a photographic print in full color of a six-month  
old plant illustrating its trumpet-shaped, pale-pink to white  
flowers with red-purple throats.

2

DETAILED PLANT DESCRIPTION

The following is a detailed description of the new  
Mandevilla plant based on observations made at the nursery  
of Gem Ornamentals, Inc. in Lake County, Tavares, Fla. The  
color determinations and comparisons are all based on The  
R.H.S. Colour Chart published by The Royal Horticultural  
Society of England in association with the Flower Council  
of Holland. Colors are approximate as color depends on  
horticultural practices such as temperature, light level and  
fertilization rate, among others.

Growth habit: The growth habit is a woody, counter-  
clockwise twining vine with fleshy bidentate to tetraden-  
tate (2–4 toothed) nodal appendages; opposite, elliptic to  
obovate leaves, and large funnelform light pink flowers  
borne on alternate, axillary racemes.

Stems: Stems terete, olive green (Yellow Green Group 144A  
to B) turning brown in age (new growth often tinged with  
reddish-brown), minutely tomentose, hairs initially  
colorless, brownish in age.

Foliage:

Type.—Evergreen, herbaceous.

Leaf arrangement.—Opposite.

Shape.—Elliptic to obovate, rarely ovate in small  
leaves.

Length.—(7.8) 12–18 cm.

Width.—(3.4) 4.5–6.9 cm.

Apex.—Acuminate.

Base.—Cordate.

Margin.—Entire.

Texture.—Leathery, rugose (tissue between lateral  
veins raised).

Pubescence.—Minutely tomentose throughout, hairs  
colorless, erect.

Petiole.—Channeled adaxially, minutely tomentose,  
0.5–1.0 cm long.

Color.—Upper leaf surface glittering Yellow-Green  
Group 139A to 135A; midrib and lateral veins dull  
Yellow-Green Group 143C. Lower leaf surface



opaque Yellow-Green Group 147B. Midrib below Yellow-Green Group 145D.

#### Inflorescence:

*Type*.—Simple raceme, apparently indeterminate.

*Position*.—Lateral, axillary, one per node.

*Peduncle*.—5.5–11.6 cm long, minutely papillose.

*Bracts*.—Narrowly triangular, herbaceous, 0.5–1.0 cm long, 0.2–0.4 cm wide.

*Pedicel*.—Twisted, minutely papillose, 1.3–1.8 cm long.

*Color*.—Peduncle and rachis of inflorescence — R.H.S. Fan 3 Yellow Green Group 144A and 144B. Bract—R.H.S. Fan 3 Yellow Green Group 144B with tip and apical margins R.H.S. Red-Purple Group 60A. Pedicel — R.H.S. Fan 3 Yellow Green Group 144B remaining yellow green until the flower falls.

*Number of flowers*.—8 to 9 (or more) produced per raceme, alternate but occasionally subopposite. One to three flowers per raceme are open at one time.

*Inflorescence length*.—10–30 cm (or more) long, longer than subtending leaves, increasing with age.

Flowers produced throughout the flowering season of March to November at Tavares, Fla.

#### Flowers:

*Structure*.—Bisexual, calyx of 5 basally connate sepals, corolla of five parts, sympetalous, basal portion a straight tube proper the corolla widening above into a campanulate throat and spreading corolla lobes, stamens epipetalous, filaments bearded, anthers connivent, bilobed at base, adhering to stigma.

*Calyx lobes*.—Ovate, acuminate at tip, thickened medially and at base, minutely papillose.

*Corolla tube*.—Straight, minutely papillose, 1.5–1.8 cm long, 0.4–0.5 cm in diameter at base.

*Corolla throat*.—Bell-shaped, 3.4–3.6 cm long, 2.5–3.0 cm wide at the opening.

*Corolla width*.—12.0–14.5 cm.

*Corolla lobes*.—Very strongly unequal sided (dimitate), apex obtuse, slightly excentric, broadly spatulate-elliptic to obovate. The left hand side greatly reduced  $\frac{1}{2}$  ovate, the right hand side broadly elliptic to broadly ovate; glabrous.

*Fragrance*.—The light scent of petunia is noticeable in the morning, being most pronounced at mid-day and fading later in the afternoon.

*Color*.—At opening, the outer surface of the tube proper is light green (R.H.S. Fan 3 Yellow Green Group 145B) to cream-green (R.H.S. Fan 3 Yellow Green Group 145D). The throat is pink (R.H.S. Fan 2 Red-Purple Group 65D). Along the inner surface of the throat, the basal  $\frac{1}{4}$  is yellow-green (R.H.S. Fan 3 Yellow Green Group 150B) and the apical  $\frac{3}{4}$  of the throat is dark pink (R.H.S. Fan 2 Red-Purple Group 64C), darker over the venation. The outer surface of the corolla lobes are pink (R.H.S. Fan 2 Red-Purple Group 65B to 65D) toward the center, darkening at the apex to dark pink (R.H.S. Fan 2 Red-Purple Group 64C).

The inner surface of the corolla lobes are pink (R.H.S. Fan 2 Red-Purple Group 65C) darkening at the apex to dark pink (R.H.S. Fan 2 Red-Purple

Group 64C). A broad triangular region of dark pink (R.H.S. Fan 2 Red-Purple Group 64C) extends about 1.0 cm up the lobe.

As the flower matures, the outer surfaces of the throat and corolla lobe fade to pale pink (paler than R.H.S. Fan 2 Red-Purple Group 62D). On the inside surface, the center of the petal lobe becomes white (R.H.S. Fan 4 White Group 155C). The deep pink (R.H.S. Fan 2 Red-Purple Group 64C) fades from the base of the petal lobe and persists only in the inner throat. The yellow-green color (R.H.S. Fan 3 Yellow-Green Group 150B) is absent in the throat of a mature flower. The broad triangular region of dark pink (R.H.S. Fan 2 Red-Purple Group 64C) extending about 1.0 cm up the lobe fades to pale pink (R.H.S. Fan 2 Red-Purple Group 65C) the same as the petal surface.

#### Overall size and growth habit:

*Size*.—The ultimate size of the new variety is unknown as no plant has reached full maturity. This new hybrid has grown between 1.0 to 1.5 m in 2 months under greenhouse conditions in the summer in Tavares, Fla.

*Hardiness*.—The new variety has tolerated temperatures to 35° F. but is killed at 32° F.

*Fertility*.—The new variety is both male and female fertile.

#### Additional comments:

*Differences from Mandevilla 'Alice du Pont'*.—Mandevilla 'Alice du Pont' has flowers that open light pink and become very dark pink with age. In contrast, flowers of the new variety open light pink and fade to white. Average corolla width of the new variety (12–14 cm) is wider than 'Alice du Pont' (9.5–11.5 cm).

*Differences from Mandevilla 'White Delite'*.—Mandevilla 'White Delite' has smaller flowers than the new variety. The corolla width of 'White Delite' is 8–10 cm thus narrower than the new variety corolla width of 12–14 cm. The new variety has a very prominent red-purple coloration in the throat. Mandevilla 'White Delite' lacks any red-purple coloration in the throat.

#### REFERENCES

- Bailey, L.H. and E.Z. Bailey, 1976, *Hortus Third*, MacMillan, New York.  
Graf, A.B., 1978, *Tropica—Color Cyclopedia of Exotic Plants and Trees*, Roehrs Co., East Rutherford, N.J.

#### I claim:

1. A new and distinct variety of Mandevilla plant substantially as shown and described, characterized by:

- a) large funnelform to trumpet-shaped flowers that open pale pink, fade to white within one to three days after anthesis and remain on the plant an additional two to three days;
- b) a flower color highlighted by a bright red-purple throat that becomes somewhat lighter as the flower ages; and
- c) elliptic-ovate shaped, leathery, shiny dark green and evergreen leaves.

\* \* \* \* \*

FIG. 1

