



US00PP10461P

# United States Patent [19]

Shaw

[11] **Patent Number: Plant 10,461**

[45] **Date of Patent: Jun. 23, 1998**

[54] **STRAWBERRY PLANT NAMED 'GAVIOTA'**

P.P. 8,746 5/1994 Izhar et al. .... Plt./49

[75] Inventor: **Douglas V. Shaw**, Davis, Calif.

[73] Assignee: **The Regents of the University of California**, Oakland, Calif.

*Primary Examiner*—James R. Feyrer  
*Assistant Examiner*—Kent L. Bell  
*Attorney, Agent, or Firm*—Burns, Doane, Swecker & Mathis, L.L.P.

[21] Appl. No.: **747,507**

[22] Filed: **Nov. 12, 1996**

[51] **Int. Cl.**<sup>6</sup> ..... **A01H 5/00**

[52] **U.S. Cl.** ..... **Plt./49**

[58] **Field of Search** ..... **Plt./49, 48**

## [57] **ABSTRACT**

'C205' is a new and distinct cultivar of strawberry plant of the short-day type which produces greater quantities of late-season fruit than the cultivars 'Chandler' and 'Camarosa' when grown in central California. Fruiting plants of 'C205' are smaller and more compact, more open, more erect, and less vigorous than plants of 'Chandler' or 'Camarosa'.

## [56] **References Cited**

### U.S. PATENT DOCUMENTS

P.P. 8,660 3/1994 Voth et al. .... Plt./49

## **3 Drawing Sheets**

### **1**

#### RELATED APPLICATIONS

There are no related applications.

#### FIELD OF THE INVENTION

This invention relates to a new and distinctive short-day type strawberry cultivar designated as 'C205', which resulted from a cross performed in 1991 between advanced selections Cal 87.112-6 and Cal 88.270-1. The cultivar is botanically identified as *F. xananassa Duch.*

'C205' was first fruited at the University of California Wolfskill Experimental Orchard, near Winters, Calif. in 1992 where it was selected, originally designated Cal 91.248-2, and propagated asexually by runners. Following selecting and during testing the plant of this disclosure was designated 'CN205'. With the decision that this plant was to be released, it was given the name 'Gaviota' for the purposes of introduction into commerce and for international registration and recognition. Asexual propagules from this original source have been tested at the Watsonville Strawberry Research Facility, the South Coast Research and Extension Center, and to a limited extent in grower fields starting in 1994.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 depicts the general flowering and fruiting characteristics of the plant.

FIG. 2 depicts a typical mature leaf during late spring.

FIG. 3 depicts representative mid-season fruit.

#### DETAILED DESCRIPTION OF THE INVENTION

'C205' is typical of short-day strawberry cultivars and produces fruit over an extended period when treated appropriately in arid, subtropical climates. The production pattern for 'C205' is similar to that for 'Chandler' (U.S. Plant Pat. No. 5,262) and 'Camarosa' (U.S. Plant Pat. No. 8,708), although it produces greater quantities of late-season fruit than either cultivar in central California, and is somewhat later to initiate fruiting than 'Camarosa'. 'C205' will be of special interest for winter plantings, where 'Chandler' and 'Camarosa' have been successful, and in summer plantings

### **2**

where 'Pajaro' (U.S. Plant Pat. No. 4,538) has been successful.

5 Plants and foliage: Fruiting plants of 'C205' are smaller and more compact, more open, more erect, and less vigorous than plants of 'Chandler' or 'Camarosa'. Comparative statistics for foliar characters near mid-season are given for the three cultivars in Table 1. Individual leaflets for 'C205' are larger than for 'Chandler' or 'Camarosa', and somewhat more rounded. Leaves (including petioles) are similar in length and broader than those of 'Chandler', and shorter and broader than those of 'Camarosa'. Petioles are thicker and more stiff than those of 'Chandler' and are similar to those of 'Camarosa'. Also, the upper surface of leaves for 'C205' are darker than either 'Chandler' or 'Camarosa' and the lower surfaces are somewhat lighter (Table 2). Leaves of 'C205' have variable leaf convexity, are generally more concave than leaves of 'Chandler' or 'Camarosa', and have fewer and more rounded serrations than the comparison cultivars.

10 Isozymes in leaf extracts: 'C205' has been classified for three isozyme systems using starch gel electrophoresis (Table 3): Phosphoglucosomerase (PGI), Leucine Aminopeptides (LAP), and Phosphoglucosomutase (PGM). It is distinguishable from 'Chandler' but not from 'Camarosa' using this methodology. For electrophoretic procedures see: J. Amer. Soc. Hort. Sci. 106:684-687.

15 Disease and pest reaction: 'C205' is moderately susceptible to common leaf spot (*Ramularia tulasnei*) and Verticillium wilt (*Verticillium dahliae*), relatively resistance to powdery mildew (*sphaerotheca macolaris*) and Anthracnose crown rot (*Colletotrichum acutatum*). When treated properly, it has tolerance to two-spotted spidermites (*Tetranychus urticae*) equal or greater than 'Chandler' and 'Camarosa'. 'C205' is tolerant to strawberry viruses encountered in California.

TABLE 1

Foliar characteristics for 'C205', 'Chandler', and 'Camarosa'			
Foliar Character	Cultivar		
	'Chandler'	'Camarosa'	'C205'
<u>Mid-tier leaflet</u>			
<u>Length (mm)</u>			
mean	64.1	68.7	71.7
range	57–69	60–83	66–85
<u>Width (mm)</u>			
mean	61.8	64.2	65.7
range	55–66	56–80	61–74
<u>Mid-tier leaf</u>			
<u>Length (mm)</u>			
mean	110.0	131.9	122.1
range	100–122	115–146	115–134
<u>Width (mm)</u>			
mean	119.5	121.5	119.2
range	109–135	110–147	108–145
# Leaflets/leaf	3	3	3
Leaf convexity	most concave, some flat/convex	flat–convex, some concave	convex, few flat or concave
<u>Serrations</u>			
number	med–many	few	few to moderate
shape	semipointed	rounded, some semipointed	rounded to semipointed
Leaf pubescence	moderate	light–moderate	moderate
Petiole pubescence			
density	mod.–heavy	heavy	heavy
direction	perpendicular	perpendicular to acropetal	perpendicular

Flowering, fruiting, fruit, and production characteristics: Comparative statistics for flower and fruit characters near mid-season are given for 'C205', 'Chandler' and 'Camarosa' in Table 4. The primary flowers for 'C205' are similar in size to 'Chandler' and 'Camarosa'; the sepals are somewhat larger than for 'Chandler' and substantially smaller than for 'Camarosa'. The calyx for 'C205' varies from slightly indented to slightly necked, and each primary flower has 4–6 petals. The fruit shape for 'C205' can vary but is typically a rounded conic, and is easily distinguished from 'Chandler' (flat conic, with some long conic) and 'Camarosa' (shortened flat conic). External and internal fruit color for 'C205' is darker than 'Chandler' and slightly darker than 'Camarosa' (Table 2). Achenes vary from yellow to dark red, and are even with the first surface or occasionally slightly extruded. 'C205' has been tested under a variety of cultural regimes, and optimal performance is obtained when nursery treatments and nutritional programs similar to those for 'Chandler' and 'Camarosa' are used. In general, 'C205' is less adapted to very early season planting but less sensitive to excess chilling than 'Camarosa'. 'C205' retains excellent fruit quality in summer planting systems.

When treated with appropriate planting regimes, 'C205' has larger fruit and produces greater yields than 'Chandler' (Table 5); 'C205' has lower yield but equal or larger sized fruit than 'Camarosa'. 'C205' is similar to 'Chandler' in its production pattern, somewhat later to initiate production than 'Camarosa', and produces substantially more late-season fruit than either comparison cultivar

(with conventional winter planting in central California). Commercial appearance ratings have been comparable to or better than those for 'Chandler' and 'Camarosa'. Fruit for 'C205' is substantially firmer than fruit and 'Chandler' but slightly less firm than 'Camarosa'. Subjectively, 'C205' has very good flavor, somewhat less aromatic than 'Chandler', but with better balance and texture than 'Camarosa'. The fruit will be outstanding for both fresh market and processing, and will be useful for home garden purposes.

TABLE 2

Foliar and fruit color characteristics for 'C205', 'Chandler', and 'Camarosa'			
Color Character	Cultivar		
	'Chandler'	'Camarosa'	'C205'
<u>Leaf color (CIELAB)</u>			
<u>Adaxial</u>			
<u>L*</u>			
mean	33.4	33.2	30.2
range	30.2–37.8	30.9–34.3	28.8–31.8
<u>a*</u>			
mean	–7.5	–7.2	–6.3
range	–6.7–8.6	–5.6–8.9	–5.6–7.5
<u>b*</u>			
mean	16.9	14.7	11.1
range	14.7–21.2	13.4–17.7	9.5–13.8
Munsell	5GY 5/6	2.5GY 5/5	2.5GY 5/5
<u>Abaxial</u>			
<u>L*</u>			
mean	48.9	48.7	49.1
range	47.1–50.0	47.3–50.0	46.9–50.3
<u>a*</u>			
mean	–7.6	–7.6	–7.8
range	–6.3–8.2	–7.3–8.3	–7.3–8.2
<u>b*</u>			
mean	21.1	20.7	18.6
range	18.3–23.6	19.8–21.8	16.8–20.1
Munsell	5GY 4/3	5GY 4/3	5GY 4/3
<u>Fruit color (CIELAB)</u>			
<u>External</u>			
<u>L*</u>			
mean	24.0	25.7	23.9
range	22.6–25.7	22.3–28.0	22.1–25.1
<u>a*</u>			
mean	26.9	22.0	23.9
range	23.9–29.2	16.7–28.0	21.4–25.1
<u>b*</u>			
mean	13.9	26.8	12.4
range	10.0–16.3	24.5–28.5	9.5–16.9
Munsell	5R 5/13	2.5R 4/10	5R 4/12
<u>Internal</u>			
<u>L*</u>			
mean	45.5	44.6	48.7
range	42.0–49.9	38.4–52.0	45.5–51.3
<u>a*</u>			
mean	39.0	39.6	33.1
range	32.6–43.1	34.3–41.5	30.3–36.0



TABLE 2-continued

Foliar and fruit color characteristics for 'C205', 'Chandler', and 'Camarosa'			
Color	Cultivar		
	'Chandler'	'Camarosa'	'C205'
Character			
b*			
—			
mean	29.4	26.7	24.2
range	24.7–33.8	24.5–28.5	21.2–27.2
Munsell	5R 6/11	7.5R 5/13	5R 5/13

\*CIELAB is the abbreviation of the international color system known as "Commission Internationale De L'Eclairage" 1978. Recommendations on uniform color spaces - - - color difference equations, psychometric color terms, Supplement No. 2 to CIE Publication No. 15. PARIS.

TABLE 3

Isozyme phenotypes for 'C205', 'Chandler', and 'Camarosa'			
locus	Cultivar		
	'Chandler'	'Camarosa'	'C205'
PGI	A1	A2	A2
LAP	B3	B3	B3
PGM	C1	C1	C1

TABLE 4

Flower and fruit characters for 'C205', 'Chandler', and 'Camarosa'			
Character	Cultivar		
	'Chandler'	'Camarosa'	'C205'
# petals			
mean	5.8	5.5	5.1
range	5–7	5–7	4–6
Flower position	even to	most even	exposed

TABLE 4-continued

Flower and fruit characters for 'C205', 'Chandler', and 'Camarosa'			
Character	Cultivar		
	'Chandler'	'Camarosa'	'C205'
(relative to foliage)	exposed	some exposed	
Calyx diam. (mm)			
mean	39.6	64.3	43.2
range	33–44	40–77	35–55
Corolla diam. (mm)			
mean	38.9	35.9	37.7
range	37–42	31–40	35–41
Fruit shape			
length/width			
ratio	1.10	1.15	1.10
range	1.00–1.43	0.96–1.19	1.08–1.12
subjective	conic to flat conic	flat conic to flat	mostly rounded conic, some flat conic
Calyx position	slight neck	event to slight indent	even to slight indent
Seed position	even	even to slight indent	even

TABLE 5

Performance of 'C205', 'Chandler', and 'Camarosa' evaluated at the Watsonville Research Facility in 1995 and 1996. All plants for these trials were harvested from Macdoel on October 17, and transplanted after one week supplemental storage. Harvest was initiated in early April and continued through the first week of September. (52" 2-row beds, 17,300 plants/acre, 100 grams/plant = 316.3 crates/acre)

Item	Yield (g/plant)	Appearance Score	Size (g/fruit)	Firmness
'Chandler'	1,372	3.1	22.9	6.1
'Camarosa'	1,820	3.7	26.3	7.8
'C205'	1,443	3.7	27.4	7.3

We claim:

1. The new and distinct cultivar of strawberry plant substantially as herein described and illustrated.

\* \* \* \* \*





Figure 1.






Figure 2.






Figure 3.