

US00PP10645P

Plant 10,645

Oct. 13, 1998

United States Patent

Kristiansen

Primary Examiner—James R. Feyrer

Patent Number:

Date of Patent:

[11]

[45]

Assistant Examiner—Elizabeth C. Kemmerer Inventor: Kell Kristiansen, Årslev, Denmark Attorney, Agent, or Firm—C. A. Whealy

ABSTRACT [57] Assignee: Asterklubben, Arslev, Denmark

A distinct cultivar of Aster plant named 'Patricia Viking', Appl. No.: 772,719 characterized by its rapid growth rate and moderate vigor;

Filed: Dec. 23, 1996

ASTER PLANT NAMED 'PATRICIA VIKING'

conditions; daisy-type inflorescences with purple ray florets

and yellow centers; and good post-production longevity. Plt./68.1 U.S. Cl.

1 Drawing Sheet

The present invention relates to a new and distinct cultivar of pot Aster plant, botanically known as Aster novi-belgii and referred to by the cultivar name Patricia Viking.

The new cultivar is a product of a planned breeding program conducted by the inventor in Arslev, Denmark. The objective of the breeding program was to create new Aster cultivars having desirable inflorescence colors, uniform plant habit and rapid growth rate.

The new cultivar originated from a cross made in 1994, by the inventor of the cultivar Patricia Ballard (not patented) as the male or pollen parent with the cultivar Royal Ruby (not 10 patented) as the female or seed parent.

The cultivar Patricia Viking was discovered and selected by the inventor as a flowering plant within the progeny of the stated cross in a controlled environment in Arslev, Denmark. Asexual reproduction of the new cultivar by terminal cut- 15 tings taken at Arslev, Denmark, has shown that the unique features of this Aster are stable and reproduced true to type in successive generations of asexual reproduction.

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Patricia 20' Viking'. These characteristics in combination distinguish 'Patricia Viking' as a new and distinct cultivar:

- 1. Rapid growth rate and moderate vigor.
- 2. Moderate branching.
- 3. Freely flowering even under winter conditions.
- 4. Daisy-type inflorescences with purple ray florets and yellow centers.
 - 5. Good post-production longevity.

The cultivar Patricia Viking has not been observed under all possible environmental conditions. The phenotype may ³⁰ vary somewhat with variations in environment such as temperature, daylength and light intensity, without, however, any variance in genotype.

In comparison to plants of the new Aster, plants of the male parent, the cultivar Patricia Ballard, differ in ray floret 35 color. In comparison to plants of the new Aster, plants of the female parent, the cultivar Royal Ruby, are taller, differ in ray floret color, and have poor postproduction longevity.

The accompanying colored photographs illustrate the overall appearance of the new cultivar, showing the colors as 40 true as it is reasonably possible to obtain in colored reproductions of this type.

The first photograph comprises a top perspective view of typical flowering plant of the new Aster.

The second photograph comprises a top perspective view ⁴⁵ of a typical inflorescence of the new Aster. Floret and foliage colors in these photographs may vary from the actual ray and foliage colors due to light reflectance.

In the following description, color references are made to The Royal Horticultural Society Colour Chart except where

general terms of ordinary dictionary significance are used. The following observations and measurements describe plants grown in Arslev, Denmark, under glass with average day and night temperatures of 20 C. Plants received long day/short night treatments for the first six weeks, then short day/long night until flowering. Measurements and numerical values represent ranges or averages for six typical flowering

Botanical classification: Aster novi-belgii cultivar Patricia Viking.

Commercial classification: Pot Aster.

Parentage:

plants.

Male or pollen parent.—Aster novi-belgii Patricia Ballard (not patented).

Female or seed parent.—Aster novi-belgii Royal Ruby (not patented).

Propagation:

Type.—Terminal tip cuttings.

Time to rooting.—Summer: About 12 days at a temperature of 21 C. Winter: About 13 days at a temperature of 21 C.

Rooting habit.—Fine, fibrous and well-branched. Plant description:

Appearance.—Perennial herbaceous decorative container plant. Stems upright to outward, inverted triangle, and moderate branching. Rapid growth rate and moderate vigor. Appropriate for 10 to 15-cm containers. Usually about 11 weeks are required to produce a flowering plant.

Plant height.—13 to 16 cm.

Plant width.—25 to 30 cm.

Stem diameter.—2 to 5 mm.

Foliage description.—Arrangement: Distichous. Shape: Elliptic with acute apex and auriculate base, margin entire. Size: Length: 2 to 3.5 cm. Width: 5 to 10 mm. Texture: Glabrous, leathery. Color: Abaxial surface: 137A. Adaxial surface: 137C.

Flowering description:

Appearance.—Daisy inflorescence form. Flat inflorescences borne on terminals above foliage, arising from leaf axils. Disc and ray florets arranged acropetally on a capitulum.

Flowering response.—Under natural conditions, plant flower in the late summer/autumn in the Northern Hemisphere. At other times of the year, inflorescence initiation and development can be induced under short day/long night conditions.

Post-production longevity.—Good, inflorescences last about three to four weeks on the plant. Inflorescences persistent.

moderate branching; freely flowering even under winter

3

Quantity of Inflorescences.—Inflorescences form at every leaf axil. Freely flowering, usually 5 to 8 inflorescences per flowering stem.

Inflorescence size.—Diameter: 2.6 to 3 cm. Depth (height): 1 to 1.5 cm. Diameter of disc: 5 to 9 mm.

Bud.—Shape: Ovoid. Size: Length: 8 to 10 mm. Diameter: 6 to 8 mm.

Ray florets.—Quantity: 100 to 150 per inflorescence. Number of whorls: More than two. Shape: Narrowly obovate with obtuse apex and entire margin. Size: Length: 1.4 to 1.6 cm. Width: 1 to 2 mm. Texture: Satiny, smooth and glabrous. Aspect: Flat to slightly concave. Orientation: Horizontal to peduncle. Color: Mature, abaxial surface: 82B/82C. Mature, adaxial surface: 82D.

Disc florets.—Quantity: 30 to 45 per inflorescence. Shape: Funnel. Length: 1 to 1.2 cm. Color, apex of corolla: Immature: Dark purple. Mature: Almost transparent.

4

Peduncle.—Aspect: Angled about 45° to the stem.
Length: 2.5 to 3 cm. Texture: Glabrous. Color: 137A.
Reproductive organs.—Androecium: Present on disc florets only. Anther position: Even with apex of disc corolla. Anther size: About 2 mm. Anther color: 14C.
Pollen: Moderate, 14C in color. Gynoecium: Present on both ray and disc florets. Stigma position: Above the anthers. Stigma shape: Bifurcate. Stigma color:

Disease resistance: No known Aster diseases observed to date on plants grown under commercial greenhouse conditions.

Seed production: Seed production has not been observed.

It is claimed:

4C.

1. A new and distinct cultivar of Aster plant named 'Patricia Viking', as illustrated and described.

* * * *



