



US00PP10826P

United States Patent [19]
Andel

[11] Patent Number: Plant 10,826
[45] Date of Patent: Mar. 16, 1999

[54] ALSTROEMERIA PLANT NAMED ‘STALYA’
[75] Inventor: Jacob van Andel, Aalsmeer,
Netherlands
[73] Assignee: Van Staaveren b.v., Aalsmeer,
Netherlands
[21] Appl. No.: 881,825
[22] Filed: Jun. 19, 1997
[51] Int. Cl.⁶ A01H 5/00
[52] U.S. Cl. Plt./87.1
[58] Field of Search Plt./87.1

[56] References Cited
PUBLICATIONS
Huxley et al. (Eds.) “Alstroemeria”, The New R.H.S Dic-
tionary of Gardening, The Stockton Press New York. vol. 1
pp. 141–148, 1992.
Primary Examiner—Howard J. Locker
Assistant Examiner—Melissa L. Kimball
Attorney, Agent, or Firm—Christie, Parker & Hale, LLP
[57] ABSTRACT
A new and distinct Alstroemeria plant variety producing
flowers of good quality and attractive coloration.
1 Drawing Sheet

1

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct variety
of Alstroemeria originated as a seedling by crossing as seed
parent an unreleased, unpatented seedling identified as
88T1104-01 and as pollen parent an unreleased, unpatented
seedling identified as 87F1459-01 from among my collec-
tion of Alstroemeria seedlings maintained under controlled
conditions in a greenhouse at Van Staaveren B.V., Aalsmeer,
The Netherlands, for developmental purposes. The varietal
denomination of the new variety is ‘Stalya’.
The first act of asexual reproduction of ‘Stalya’ was in
Aalsmeer, The Netherlands, by propagation by dividing
rhizomes. Asexual reproduction by dividing rhizomes
through successive generations in Aalsmeer, The Nether-
lands, has demonstrated that the combination of character-
istics as herein disclosed for ‘Stalya’ are firmly fixed and
retained through successive generations.
‘Stalya’ has not been observed under all possible envi-
ronmental conditions. Phenotypic expression may vary with
variations in environment such as temperature, light inten-
sity, day length and growing and cultural conditions.

SUMMARY OF THE INVENTION

Flowers of the new variety of Alstroemeria, ‘Stalya’ are
characterized by unusual coloration, as described herein, and
good quality.

DESCRIPTION OF ILLUSTRATION

The new variety of Alstroemeria hybrid is illustrated in
the accompanying illustration, which shows typical flower
characteristics, with colors being as nearly true as is possible
for illustrations of this type.

DETAILED DESCRIPTION OF THE NEW
VARIETY

The following is a detailed description of the new cultivar
of Alstroemeria hybrid as observed in a greenhouse in
Aalsmeer, The Netherlands. Color designations indicated
are in accordance with The Royal Horticultural Society
Colour Chart by color plate designations.

Botanical classification:
Family.—Liliceae.
Genus.—Alstroemeria sp.
Commercial.—Alstroemeria hybrid.

2

Origin: Seedling.
Plant
Form: Bush.
Shape: Upright.
Height: About 150 to 250 cm.
Internode length: About 1 to 8 cm.
Growth rate: Fast; the length of time (number of days) taken
from young plants (rooted rootstocks, the stage of devel-
opment in which the varieties are sold) to flowering stage
under commercial greenhouse conditions is 80–90 days.
Main stem or cane length: About 200 cm, pinching not
required.
Growth: Vigorous.
Strength: Strong.
Foliage:
Quantity.—Abundant.
Number of leaves.—About 35 to 45 per stem.
Size of leaf.—Length: About 15 to 20 cm. Width: About
3 to 5 cm.
Shape of leaf.—Elliptical.
Margin type.—Entire.
Texture.—Glossy.
Color.—Upper side: Near 137A. Underside: Near
137C.
Rhizomes.—Yes. Color: Near 155D. Size: About 10 to
30 cm.
Inflorescence—Flower/Floret
Blooming habit: Concurrent; Spent umbels not required to
be pinched to maintain the claimed plant in a continuous
blooming condition.
Blooms: Profusely.
Spring.—Yes.
Fall.—Yes.
Size:
Diameter.—About 6 cm.
Depth.—About 5 cm.
Borne: Singly.
Shape or form:
When bloom first opens.—Cup.
When bloom matures: Cup.
Fragrance: None.
Lasting quality:
On plant.—About 3 weeks.
As cut flower.—About 14 to 16 days.
Number of tepals.—6.

Arrangement.—2 concentric circles of 3 tepals.
Form.—Obovate.
Margin type.—Straight.
Apex.—Pointed.
Texture.—Soft.
Appearance.—Velvety.
Color (summer/fall):
Outer tepal/floret.—Body near 31B/53B/58C; base near 31C; reverse side near 53B/58C.
Inside tepal/floret.**13** Body near 7A/31B/53B/58C; base near 7A; reverse side near 7A/31B/53B/58C.
Spots: The two inner lateral tepals have 20–40 spots, color near 185A. The inner median tepal has 0–15 spots, color near 185A.
Peduncle or flower stem:
Length.—About 10 to 15 cm.
Color.—Near 137B.
Strength.—Strong.
Upright.—Upright, not hairy.
Discoloration after full bloom.—None.
Bloom affect by wet or hot weather.—No.
Persistence: No.
Disease resistance: Unknown.

Bud

Size:
Diameter.—About 1 to 2 cm.
Depth or length.—About 2 to 4 cm.
Form or shape: Pear-shaped, long.
Rate of opening: Fast.
Color of tepals/florets:
When tepals begin to unfurl.—Near 7A/31B/53B/58C.
When tepals first divide.—Near 7A/31B/53B/58C.

Peduncle or flower stem:
Length.—About 6 to 8 cm.
Hairy.—No.
Strength or character: Erect.
Color: Near 137B to 137C.

Reproductive Organs

Stamen:
Number.—6 .
Arrangement.—Around the style, side-by-side.
Anthers:
Size.—About 8 to 10 mm.
Color .—Near 167A.
Filaments:
Length.—About 5 cm.
Color.—Near 41C.
Pollen:
Color.—Near 167A.
Pistils.—1.
Styles:
Length.—About 4 to 5 cm.
Color.—Near 41C.
Stigmas:
Color.—Near 41D.

Fruit

Fertile: Yes.
Shape: Round.
Color at maturity: Near 163A.
I claim:
1. A new variety of *Alstroemeria* plant substantially as herein described and shown.

* * * * *

