



US00PP11060P

United States Patent [19]  
Dümmen

[11] Patent Number: Plant 11,060  
[45] Date of Patent: Sep. 14, 1999

[54] NEW GUINEA IMPATIENS PLANT NAMED  
‘DUERIREST’  
[75] Inventor: Marga Dümmen, Rheinberg, Germany  
[73] Assignee: Dümmen Jungpflanzenkulturen,  
Rheinberg, Germany  
[21] Appl. No.: 09/038,768  
[22] Filed: Mar. 11, 1998  
[51] Int. Cl.<sup>6</sup> ..... A01H 5/00  
[52] U.S. Cl. .... Plt./318

[58] Field of Search ..... Plt./317, 318  
Primary Examiner—Howard J. Locker  
Attorney, Agent, or Firm—C. A. Whealy

[57] ABSTRACT  
A new and distinct variety of New Guinea Impatiens plant  
named ‘Duerirest’, characterized by its light pink and red  
bi-colored flowers; floriferousness; freely and basally  
branching plant habit; and dark green non variegated foli-  
age.

1 Drawing Sheet

1

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct variety  
of New Guinea Impatiens plant, botanically known as *Impa-  
tiens hawkeri*, and hereinafter referred to by the name  
‘Duerirest’. The new variety is being marketed under the  
name Riviera Red Star.  
The new variety is a product of a planned breeding  
program conducted by the inventor in Rheinberg, Germany.  
The objective of the breeding program is to develop New  
Guinea Impatiens that are freely branching; early flowering;  
and that have desirable flower and leaf colors.  
The new cultivar originated from a cross made by the  
inventor of the variety ‘Pago Pago’ (disclosed in U.S. Plant  
Pat. No. 9,143) as the male, or pollen, parent with the variety  
‘Sarabande’ (not patented) as the female, or seed, parent.  
‘Duerirest’ was discovered and selected in 1995 by the  
inventor as a single flowering plant within the progeny of the  
stated cross in a controlled environment in Rheinberg,  
Germany.  
Asexual reproduction of the new variety by terminal  
cuttings taken at Rheinberg, Germany, has shown that the  
unique features of this new variety are stable and reproduced  
true to type in successive generations of asexual reproduc-  
tion.

BRIEF SUMMARY OF THE INVENTION

The following traits have been repeatedly observed and  
are determined to be the unique characteristics of ‘Duerir-  
est’. These characteristics in combination distinguish ‘Duer-  
irest’ as a new and distinct variety:  
1. Light pink and red bi-colored flowers.  
2. Freely flowering.  
3. Dense and bushy plant habit.  
4. Freely and basally branching plant habit.  
5. Dark green non-variegated foliage.  
The new variety can be compared to the male parent, the  
variety ‘Pago Pago’. However, in side-by-side comparisons  
conducted by the inventor in Rheinberg, Germany, plants of  
the new variety differed from plants of ‘Pago Pago’, in the  
following characteristics:  
1. Plants of the new variety have a faster growth rate than  
plants of ‘Pago Pago’.  
2. Plants of the new variety are taller and have longer  
internodes than plants of ‘Pago Pago’.  
3. Plants of the new variety are much more freely branch-  
ing than plants of ‘Pago Pago’.

2

4. Plants of the new variety are twice as floriferous as  
plants of ‘Pago Pago’.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the  
overall appearance of the new variety, showing the colors as  
true as it is reasonably possible to obtain in colored repro-  
ductions of this type. The photograph comprises a side  
perspective view of a typical plant of ‘Duerirest’ grown in a  
12-cm container. Flower and foliage colors may appear  
different from the actual colors due to light reflectance.

DETAILED BOTANICAL DESCRIPTION

Plants of ‘Duerirest’ have not been observed under all  
possible environmental conditions. The phenotype may vary  
somewhat with variations in environment such as  
temperature, daylength and light intensity, without,  
however, any variance in genotype. The following observa-  
tions and measurements describe plants grown in 12-cm  
containers during the spring in Rheinberg, Germany, under  
commercial practice in a glass-covered greenhouse with day  
and night temperatures about 18° C. and light levels about  
45 thousand lux.  
In the following description, color references are made to  
The Royal Horticultural Society Colour Chart except where  
general terms of ordinary dictionary significance are used.  
Ps  
Botanical classification: *Impatiens hawkeri* ‘Duerirest’.  
Parentage:  
Male, or pollen, parent.—*Impatiens hawkeri* ‘Pago  
Pago’.  
Female, or seed, parent.—*Impatiens hawkeri* ‘Sara-  
bande’.  
Propagation:  
Type cutting.—Terminal cuttings.  
Time to initiate roots.—Summer: About 10 days at 22°  
C. Winter: About 12 days at 22° C.  
Time to develop roots.—Summer: About 21 days at 22°  
C. Winter: About 26 days at 22° C.  
Rooting habit.—Freely branching.  
Plant description:  
General appearance.—Rounded; dense and bushy;  
very freely and basally branching; upright and  
spreading; rapid growth rate and vigorous; suitable  
for 10 to 16-cm pots.  
Plant height.—About 22 cm from soil level to top of  
plant plane.

*Lateral branches*.—Length: About 15.5 cm. Diameter: About 4.1 mm. Internode length: About 5.5 cm. Color: 138A.

*Foliage description*.—Quantity of leaves per lateral branch: About 18. Shape: Elliptic with acuminate tip and attenuate base. Length: About 6.6 cm. Width: About 2.5 cm. Texture: Velvety, smooth and glabrous. Margin: Serrulate. Color: Young foliage, upper surface: 147A. Young foliage, lower surface: 139A. Mature foliage, upper surface: 132A. Mature foliage, lower surface: 53A. Veins, upper surface: 186A. Veins, lower surface: 53A. Petiole: Length: About 1.7 cm. Diameter: About 1.9 mm. Color: 136C.

Flower description:

*Flowering habit*.—Freely flowering.

*Natural flowering season*.—Year-round. Plants flower about 8 weeks after planting.

*Flowers borne*.—Flower buds develop in apical leaf axils. Open flowers are displayed above the foliage.

*Quantity of inflorescences*.—Freely flowering, usually about 85 buds and flowers per plant.

*Flower shape*.—Rounded and flat in aspect.

*Flower diameter*.—About 5.1 cm.

*Flower depth (height)*.—About 5.8 mm.

*Petals*.—Shape: Reniform with emarginate apex and acute base. Quantity, arrangement: 5 petals overlapping. Aspect: Flat. Length: About 2.8 cm. Width:

About 2.6 cm. Texture: Smooth, shiny, and glabrous. Margin: Entire. Color: When opening, upper surface: Background color, light pink, 49B to 52D; red, 41A, star; and dark red, 53C, eye. When opening, lower surface: 41A/41B. Fully opened, upper surface: Background color, light pink, 49B to 52D; red, 41A, star; and dark red, 53C, eye. Fully opened, lower surface: 41A/41B. Upper surface fading to: Background color, light pink, 52D, to white, 155D.

*Peduncle*.—Angle: Erect. Length: About 3.5 cm. Color: 29A.

*Flower bud*.—Shape: Ovoid with spur. Length: About 1.6 cm. Diameter: About 8.5 mm. Color: 33A.

*Spur*.—Shape: Needle-shaped, curved at end. Length: About 5 cm. Color: 22A.

*Reproductive organs*.—Androecium: Stamen number: 5. Anther shape: Ovate. Anther size: About 4 mm. Anther color: 49A. Pollen color: 29D. Amount of pollen: Moderate. Gynoecium: Pistil length: About 4.9 mm. Style color: 146A. Stigma color: 129B.

Disease resistance: No fungal, bacterial nor viral problems observed.

Seed development: Seed production is very rarely observed.

It is claimed:

1. A new and distinct variety of New Guinea Impatiens plant named 'Duerirest', as illustrated and described.

\* \* \* \* \*



