



US00PP11292P

United States Patent [19] Stravers

[11] Patent Number: Plant 11,292
[45] Date of Patent: Mar. 14, 2000

[54] GERBERA PLANT NAMED 'TEREROS'

[75] Inventor: Lambertus Johannes Maria Stravers,
Kudelstaart, Netherlands

[73] Assignee: Terra Nigra Holding B.V., De Kwakel,
Netherlands

[21] Appl. No.: 09/048,199

[22] Filed: Mar. 26, 1998

[51] Int. Cl.⁷ A01H 5/00

[52] U.S. Cl. Plt./357

[58] Field of Search Plt./357

[56] References Cited

U.S. PATENT DOCUMENTS

P.P. 7,846 3/1992 Stravers Plt./357

OTHER PUBLICATIONS

Huxley, Anthony, ed., The New RHS Dictionary of Garden-
ing, Stockton Press, New York, p. 402, 1992.

Primary Examiner—Howard J. Locker
Assistant Examiner—Wendy Anne Baker

[57] ABSTRACT

A new and distinct cultivar of Gerbera plant named
'Tereros', as illustrated and described, characterized by its
semi-double type, the special combination of the colors
yellow and orange throughout the flower in nice contrast to
the small brown/purple disc, the light yellow styles and
anthers, an overall flower diameter of 100 mm and with
flowers of extreme good quality.

1 Drawing Sheet

1

BACKGROUND OF THE INVENTION

The present invention comprises a new and distinct cul-
tivar of *Gerbera jamesonii*, referred to by the cultivar name
'Tereros'.

'Tereros' was originated from a hybridization program in
De Kwakel, The Netherlands in 1993. The female parent
was the unnamed seedling '92.008' (unpatented) and the
male parent was 'Sundance' (unpatented). The female parent
'92.008' was completely yellow, semi-double and had a
distinct black center. The male parent differs from the new
invention because of its single type flower. The present
invention comprises an improvement of the vase-life of the
male parent. The new cultivar was selected by me from the
progeny of the stated parentage on or about February 1994.
The first asexual reproduction of 'Tereros' was accom-
plished when vegetative cuttings were taken on February
1995 in De Kwakel. The new cultivar is presently being
propagated by cuttings and tissue culture. Horticultural
examination of selected units initiated in 1997 has demon-
strated that the combination of characteristics as herein
disclosed for 'Tereros' are firmly fixed and are retained
through successive generations of asexual reproduction.

'Tereros' has not been observed under all possible envi-
ronmental conditions. The phenotype may vary significantly
with variations in environment such as temperature, light
intensity and day length. The following observations, mea-
surements and comparisons describe plants grown in De
Kwakel, The Netherlands, under greenhouse conditions
which closely approximate those generally used in commer-
cial practice. The following traits have been repeatedly
observed and are determined to be basic characteristics of
'Tereros', which in combination distinguish this Gerbera
from its parents and all other varieties of which I am aware:

1. Type: Semi-double.
2. Color of ray floret: Yellow and orange.
3. Color of disc before opening of disc florets: Brown/
purple, R.H.S. 187A; more intense in center.
4. Color of perianth lobe: Dark green, R.H.S. 137A.
5. Diameter of flower head: Approx. 90–100 mm.

Of the many commercial cultivars known to me, there is
no cultivar similar in comparison to 'Tereros'.

2

BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying photographic drawings show typical
inflorescence and foliage characteristics of the new cultivar
with the colors being shown as nearly true as is reasonably
possible to attain in illustrations of this type.

The top photograph illustrates the flower head pictured
across face.

The photograph at the bottom illustrates a typical leaf of
the present invention.

BOTANICAL DESCRIPTION OF THE PLANT

Botanical: *Gerbera jamesonii* cv. 'Tereros'.

In the following description, color references are made to
The Royal Horticultural Society Colour Chart (R.H.S.). The
color values were determined at 15.30 p.m. on Jan. 26, 1998
under artificial light at De Kwakel.

The following traits have been repeatedly observed and
are determined to be basic characteristics of 'Tereros', which
in combination distinguish this Gerbera as a new and distinct
cultivar.

INFLORESCENCE

A. Flowerhead:

Type.—Semi-double.

Diameter.—Medium (approx. 93 mm).

Color (general tonality from a distance of 3 meters).—
Yellow and orange.

Shape.—Incurving funnel-shaped.

Involucre.—Height from point of attachment of involu-
cre to top of flower head: low (approx. 28 mm).
Height: low (approx. 14 mm). Diameter: small-
medium (approx. 38 mm). Number of bracts:
medium (approx. 65). Color: green, R.H.S. 137A.
Longitudinal axis of inner rows: straight. Anthocya-
nin: present. Pubescence: dense.

Ray florets.—Number: medium (57). Shape: broad obo-
vate. Longitudinal axis outer row: incurving. Lon-
gitudinal axis inner row: incurving-straight. Longi-
tudinal axis of ray female floret: reflexing.

Outer ray floret.—Cross section: flat. Length: short (approx. 40 mm). Width: medium (approx. 8–10 mm). Longitudinal folding: medium. Angle of apex: obtuse. Shape of apex: rounded. Incisions of apex: 1. Depth of incision: medium. Color (topside): R.H.S. 12A+R.H.S. 28B at edges. Color (bottom side): R.H.S. 1A. Color distribution on inner side: uniform. Edge of different color: present (R.H.S. 28B). Striation: absent. Claw spot: absent.

B. Disc florets:

Disc diameter.—Small (approx. 18 mm).

Color (immature, bottom).—White-green (50% at bottom, R.H.S. 154C).

Color (immature, top).—Orange (50% at top, R.H.S. 30A).

Main color upperside corolla.—Female flowers: 70% orange, 30% yellow (R.H.S. 32A, R.H.S. 12A). Male flowers: 50% yellow, 50% orange (R.H.S. 3B, R.H.S. 32A).

C. Reproductive organs:

Style.—Main color distal part: light yellow (R.H.S. 4B).

Stigma.—Main color: light yellow (R.H.S. 4B).

Anthers.—Main color: light yellow (R.H.S. 7A). Color of top relative to other parts is identical. Longitudinal stripes are absent. Intensity of anthocyanin coloration is absent.

Pappas.—Main color: purple (R.H.S. 187B). Color of top relative to other parts is identical. Level of top relative to closed disc florets: above (1–2 mm in center).

Fertility.—Fertility as well as the seedsetting is good.

D. Peduncle:

Length.—Medium (approx. 64 cm).

Cross section.—Elliptic.

Tendency to fasciation.—Absent.

Thickness.—Medium.

Strength.—Strong.

Pubescence.—Medium.

Color.—Medium green (R.H.S. 144A).

Anthocyanin coloration.—At base: absent or very weak (R.H.S. 152C). At top: absent.

Involucral bracts.—Absent.

PLANT

A. General appearance:

Height.—40 cm (excluding any flowers).

B. Foliage:

Leaf blade.—Length: long (approx. 32±1 cm). Width: medium (approx. 14–16 cm). Thickness: medium. Blistering: medium. Pubescence: On upper side (midrib excluded): sparse. Depth of cuts or incisions in leaf: Basal part: deep. Central part: medium. Distal part: medium. Color: Upper side of the leaf blade: medium green (R.H.S. 137A). Bottom side of the leaf blade: R.H.S. 138B. Glossiness on upper side: strong. Angle of apex: acute. Shape of apex: pointed. Margin of lobes: serrate. Number of lobes: approx. 8. Extensions of margin: medium.

Petiole.—Petiole length: medium (approx. 21 cm). Color of petiole: R.H.S. 144A. Petiole anthocyanin coloration: weak.

C. Disease resistance: No special disease resistance.

OTHER CHARACTERISTICS

An important characteristic for the variety 'Tereros' is the very sturdy flower.

I claim:

1. A new and distinct cultivar of Gerbera plant named 'Tereros', substantially as herein shown and described, characterized particularly as to novelty by its characteristics enumerated above.

* * * * *

