



US00PP11375P

United States Patent [19]  
Dümmen

[11] Patent Number: Plant 11,375  
[45] Date of Patent: May 9, 2000

[54] NEW GUINEA IMPATIENS PLANT NAMED  
‘DUEPETREST’  
[75] Inventor: Marga Dümmen, Rheinberg, Germany  
[73] Assignee: Dummen Jungpflanzenkulturen,  
Rheinberg, Germany  
[21] Appl. No.: 09/103,176  
[22] Filed: Jun. 22, 1998  
[51] Int. Cl.<sup>7</sup> ..... A01H 5/00  
[52] U.S. Cl. .... Plt./318  
[58] Field of Search ..... Plt./318

[56] References Cited  
U.S. PATENT DOCUMENTS  
P.P. 9,143 5/1995 Kientzler ..... Plt./318  
P.P. 10,302 3/1998 Kientzler ..... Plt./318

OTHER PUBLICATIONS

GTITM UPOVROM Citation for ‘Duepetrest’ as Per QZ  
PBR 960330; Mar. 4, 1996.  
GTITM UPOVROM Citation for ‘Duepetrest’ as Per JP  
PBR 9854; May 6, 1997.

Primary Examiner—Howard J. Locker  
Assistant Examiner—Kent L. Bell  
Attorney, Agent, or Firm—C. A. Whealy

[57] ABSTRACT  
A new and distinct variety of New Guinea Impatiens plant  
named ‘Duepetrest’, characterized by its soft red and light  
lavender bi-colored petal color; floriferousness; dense and  
bushy plant habit; freely and basally branching plant habit;  
and dark green non-variegated foliage.

1 Drawing Sheet

1

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct variety  
of New Guinea Impatiens plant, botanically known as *Impa-*  
*tiens hawkeri*, and hereinafter referred to by the name  
‘Duepetrest’. The new variety is marketed under the trade  
name Red Fox Petticoat Red Star.  
The new variety is a product of a planned breeding  
program conducted by the Inventor in Rheinberg, Germany.  
The objective of the breeding program is to develop New  
Guinea Impatiens that are freely branching; compact; early  
flowering; and that have desirable flower and leaf colors.  
The new cultivar originated from a cross made by the  
Inventor of two unidentified proprietary New Guinea Impa-  
tiens seedling selections.  
‘Duepetrest’ was discovered and selected in 1994 by the  
Inventor as a single flowering plant within the progeny of the  
stated cross in a controlled environment in Rheinberg,  
Germany.  
Asexual reproduction of the new variety by terminal  
cuttings taken at Rheinberg, Germany, has shown that the  
unique features of this new variety are stable and reproduced  
true to type in successive generations of asexual reproduc-  
tion.  
BRIEF SUMMARY OF THE INVENTION  
The following traits have been repeatedly observed and  
are determined to be the unique characteristics of ‘Duepe-  
trest’. These characteristics in combination distinguish  
‘Duepetrest’ as a new and distinct variety:  
1. Soft red and light lavender bi-colored petal color.  
2. Freely flowering.  
3. Dense and bushy plant habit.  
4. Freely and basally branching plant habit.  
5. Dark green non-variegated foliage.  
The new variety can be compared to the commercial  
cultivar ‘Pago Pago’ (disclosed in U.S. Plant Pat. No. 9,143).  
However, in side-by-side comparisons conducted by the  
Inventor in Rheinberg, Germany, plants of the new variety  
are much more compact and have more incised flower

2

petals. In addition, petals of plants of the new New Guinea  
Impatiens have a more pink red central stripe whereas petals  
of plants of the cultivar ‘Pago Pago’ have a more orange red  
stripe.  
BRIEF DESCRIPTION OF THE PHOTOGRAPH  
The accompanying colored photograph illustrates the  
overall appearance of the new variety, showing the colors as  
true as it is reasonably possible to obtain in colored repro-  
ductions of this type. The photograph comprises a side  
perspective view of a typical plant of ‘Duepetrest’. Flower  
and foliage colors may appear different from the actual  
colors due to light reflectance.  
DETAILED BOTANICAL DESCRIPTION  
Plants of ‘Duepetrest’ have not been observed under all  
possible environmental conditions. The phenotype may vary  
somewhat with variations in environment such as  
temperature, daylength and light intensity, without,  
however, any variance in genotype. The following observa-  
tions and measurements describe plants grown in 12-cm  
containers during the spring in Rheinberg, Germany, under  
commercial practice in a glass-covered greenhouse with day  
temperatures about 22° C., night temperatures about 18° C.  
and light levels about 42.5 thousand lux.  
In the following description, color references are made to  
The Royal Horticultural Society Colour Chart except where  
general terms of ordinary dictionary significance are used.  
Botanical classification: *Impatiens hawkeri* ‘Duepetrest’.  
Parentage:  
Male, or pollen, parent.—Unidentified proprietary  
*Impatiens hawkeri* seedling selection.  
Female, or seed, parent.—Unidentified proprietary  
*Impatiens hawkeri* seedling selection.  
Propagation:  
Type cutting.—Terminal cuttings.  
Time to initiate roots.—Summer: About 10 days at 22°  
C. Winter: About 12 days at 22° C.  
Time to develop roots.—Summer: About 21 days at 22°  
C. Winter: About 26 days at 22° C.

*Rooting habit.*—Fine, freely branching.

Plant description:

*General appearance.*—Rounded; compact; full, dense and bushy; very freely and basally branching; upright and spreading; moderate growth rate; suitable for 10 to 15-cm pots.

*Plant height.*—About 15.5 cm from soil level to top of plant plane.

*Lateral branches.*—Quantity: About 17. Length: About 13.4 cm. Diameter: About 5.1 mm. Internode length: About 3.75 cm. Color: 138A.

*Foliage description.*—Quantity of leaves per lateral branch: About 26. Shape: Elliptic with acuminate tip and attenuate base. Length: About 11 cm. Width: About 4 cm. Texture: Smooth, glabrous. Margin: Serrulate. Color: Young foliage, upper surface: 139A. Young foliage, lower surface: 59A. Mature foliage, upper surface: 139A. Mature foliage, lower surface: 59A. Veins, mature foliage, upper surface: 59B. Veins, mature foliage, lower surface: 136A. Petiole: Length: About 2.8 cm. Diameter: About 2.9 mm. Color: 133C.

Flower description:

*Flowering habit.*—Freely flowering. Flowers not persistent.

*Natural flowering season.*—Year-round. Plants flower about 8 weeks after planting. Flowers typically last less than one week depending on environmental and cultural conditions.

*Flowers borne.*—Flower buds develop in apical leaf axils. Open flowers are displayed above the foliage.

*Quantity of inflorescences.*—Freely flowering, usually about 100 flower buds and open flowers per plant.

*Flower shape.*—Oval.

*Flower diameter.*—About 5.4 cm.

*Flower depth (height).*—About 1 cm.

*Petals.*—Shape: Obcordate with emarginate apex and acute base. Quantity, arrangement: Five petals overlapping. Aspect: Flat. Length: About 3 cm. Width: About 2.5 cm. Texture: Smooth, satiny, glabrous. Margin: Entire. Color: When opening, upper surface: 56C; 40C central stripe and central area on upper petal; 54B base. When opening, lower surface: 39A. Fully opened, upper surface: 75C; 52A central stripe and central area on upper petal; 57C base. Fully opened, lower surface: 39A.

*Peduncle.*—Angle: Erect. Strength: Slender, but strong. Length: About 4 cm. Color: 134B.

*Flower bud.*—Shape: Ovoid with spur. Length: About 1.8 cm. Diameter: About 9 mm. Color: 39A.

*Spur.*—Shape: Needle-shaped, curved at end. Length: About 5.4 cm. Color: 29A-29C.

*Reproductive organs.*—Androecium: Stamen number: 5. Anther shape: Oval. Anther size: About 4 mm. Anther color: 54C. Pollen color: 27C. Amount of pollen: Abundant. Gynoecium: Pistil length: About 7.8 mm. Style color: 131A. Ovary color: 131A.

Disease resistance: Resistance to pathogens common to New Guinea Impatiens has not been observed.

Seed development: Seed production has not been observed.

It is claimed:

1. A new and distinct variety of New Guinea Impatiens plant named ‘Duepetrest’, as illustrated and described.

\* \* \* \* \*

