

Plant 11,486

Aug. 22, 2000

# United States Patent [19]

Trees

- [54] NEW GUINEA IMPATIENS PLANT NAMED 'BFP-796'
- [75] Inventor: Scott C. Trees, Nipomo, Calif.
- [73] Assignee: Ball FloraPlant, a division of Ball Horticultural Company, West Chicago, Ill.
- [21] Appl. No.: **09/112,648**
- [22] Filed: Jul. 9, 1998

[56] **References Cited** U.S. PATENT DOCUMENTS

**Patent Number:** 

**Date of Patent:** 

[11]

[45]

P.P. 10,108 11/1997 Trees ..... Plt./318 Primary Examiner—Howard J. Locker Assistant Examiner—Kent L. Bell Attorney, Agent, or Firm—C. A. Whealy [57] ABSTRACT

A new and distinct cultivar of New Guinea Impatiens plant named 'BFP-796', characterized by its large and numerous orange-colored flowers; upright and mounded plant habit;

[51]	Int. Cl. <sup>7</sup> A	01H 5/00
[52]	U.S. Cl.	. Plt./318
[58]	Field of Search	Plt./318

short internodes; good basal branching; and dark green leaves.

**1 Drawing Sheet** 

### BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of New Guinea Impatiens plant, botanically known as *Impatiens hawkeri*, and hereinafter referred to by the cultivar name 'BFP-796'.

The new New Guinea Impatiens is a product of a planned breeding program conducted by the Inventor in Arroyo Grande, Calif. The objective of the breeding program was to develop new cultivars with excellent basal branching, 10 upright and medium growth habit, numerous large flowers, and interesting flower and foliage colors.

The new cultivar originated from a cross made by the Inventor to two unidentified proprietary *Impatiens hawkeri* selections. The cultivar 'BFP-796' was discovered and 15 selected by the Inventor as a flowering plant within the progeny of the stated cross in a controlled environment in Arroyo Grande, Calif., on Sep. 15, 1996. 2

leaves of plants of the new New Guinea Impatiens will maintain solid dark green color.

3. Flowers of plants of the new New Guinea Impatiens are more orange in color than flowers of plants of the cultivar 'BSR-195 Salmon'.

4. Plants of the new New Guinea Impatiens have longer peduncles than plants of the cultivar 'BSR-195 Salmon'.

A detailed comparison of plants of the cultivar 'BFP-796' and the cultivar 'BSR-195 Salmon' appears in Chart A at the end of the specification.

#### BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproduction of this type. The photograph comprises a top perspective view of a typical plant of 'BFP-796'. Flower and foliage colors in the photograph appear lighter than the actual colors due to light reflectance.

Asexual reproduction of the new cultivar by terminal cuttings taken at Arroyo Grande, Calif., has shown that the <sup>20</sup> unique features of this new Impatiens are stable and reproduced true to type in successive generations of asexual reproduction.

#### SUMMARY OF THE INVENTION

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'BFP-796'. These characteristics in combination distinguish 'BFP-796' as a new and distinct cultivar:

- 1. Large and numerous orange-colored flowers.
- 2. Upright and mounded plant habit.
- 3. Short internodes.
- 4. Good basal branching.
- 5. Dark green leaves.
- The new New Guinea Impatiens can be compared to the

#### DETAILED BOTANICAL DESCRIPTION

The cultivar 'BFP-796' has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and light intensity, without, however, any variance in genotype. The following observations, measurements and comparisons describe plants grown in 10-cm pots in Arroyo 30 Grande, Calif., under commercial practice in a glass-covered greenhouse with day temperatures about 22° C. and night temperatures about 18° C.

In the following description, color references are made to The Royal Horticultural Society Colour Chart except where general terms of ordinary dictionary significance are used.

Botanical classification: *Impatiens hawkeri* cultivar 'BFP-796'.

commercial cultivar, 'BSR-195 Salmon' (disclosed in U.S. Plant Pat. No. 8,870). However, in side-by-side comparisons conducted by the Inventor in Arroyo Grande, Calif., plants 40 of the new New Guinea Impatiens differ from plants of the cultivar 'BSR-195 Salmon' in the following characteristics:

1. Plants of the new New Guinea Impatiens are more compact and have shorter lateral branches than plants of the cultivar 'BSR-195 Salmon'. 45

2. Under high light conditions, leaves of plants of the cultivar 'BSR-195 Salmon' will become variegated whereas

Parentage:

- Male parent.—Unidentified proprietary Impatiens hawkeri selection.
- Female parent.—Unidentified proprietary Impatiens hawkeri selection.

Propagation:

*Type cutting.*—Terminal cuttings. *Time to initiate roots.*—About 7 days at temperatures about 18° C.

### Plant 11,486

#### 3

*Time to develop roots.*—About 21 days at temperatures about 18° C.

*Rooting habit.*—Fibrous, fine, freely branching. Plant description:

*Plant form.*—Upright and mounded.

Growth and branching habit.—Moderately vigorous. Good basal branching with about six lateral branches at the base, dense and bushy growth. Appropriate for 10 and 12.5-cm containers.

Plant height.—About 12.6 cm.

*Crop time.*—About eight weeks are required to produce a finished flowering plant in a 10-cm container from a rooted cutting.

#### 4

- *Peduncles.*—Length: About 5.6 cm. Angle: About 45° to the lateral branch. Strength: Moderately strong. Color: 144D.
- *Reproductive organs.*—Androecium: Stamen number: Five, anthers fused, filaments free. Anther shape: Obovate. Anther color: 12D. Amount of pollen: Moderate. Pollen color: 13D. Gynoecium: Fiveloculate fused. Gynoecium color: 143C.
- Disease resistance: Under commercial conditions, resistance to pathogens has not been observed.

Seed development: Seed production has not been observed.

#### CHART A

- Lateral branches.—Length: About 6 cm. Diameter: About 6 mm. Internode length: About 2 cm. Color: 144C.
- Foliage description.—Leaves simple, generally symmetrical, abundant, in whorls or opposite. Quantity of leaves per lateral branch: About nine. Length: About 9 cm. Width: About 3.2 cm. Shape: Lanceolate with acuminate apex, attenuate base and ciliate and serrate margin. Texture: Rugose, slightly pubescent. Color: Fully expanded foliage, upper surface: 137A. Fully expanded foliage, lower surface: 147B. Venation, upper surface: 145C. Venation, lower surface: 145C. Petiole: Length: About 1.5 cm. Diameter: About 3 mm. Color: 145C.

Flower description:

*Flower type and habit.*—Large orange-colored flowers. Freely and continuously flowering. Flowers arise from leaf axils. Usually about nine flowers and flower buds per lateral branch. Flowers positioned mostly above the foliage and typically face outward to upward. Flowers generally flat and rounded. Flow-

CHART A			
CHARACTERISTIC	'BFP-796'	'BSR-195 SALMON'	
PLANT HEIGHT	About 12.6 cm	About 13.5 cm	
LATERAL BRANCH	About 6 cm	About 7 cm	
LENGTH			
LATERAL BRANCH	About 6 mm	about 4 mm	
DIAMETER STEM COLOR	144C	144D with 183D	
STEM COLOR	1440	near nodes	
QUANTITY OF LEAVES	About 9	About 11	
PER LATERAL			
BRANCH			
LEAF LENGTH	About 9 cm	About 8 cm	
LEAF WIDTH	About 3.2 cm	About 3.3 cm	
LEAF COLOR, UPPER	137A	139A	
SURFACE	1470	1200	
LEAF COLOR, LOWER SURFACE	147B	138B	
LEAF COLOR, UPPER	137A	139A at margins,	
SURFACE UNDER	137 <b>A</b>	center yellow, 7A	
HIGH LIGHT			
LEAF VEIN COLOR,	145C	144C	
UPPER & LOWER			
SURFACES			
PETIOLE COLOR	145C	144C	
FLOWER DIAMETER	About 7.5 cm	About 7.2 cm	
PETAL COLOR, WHEN	40 <b>A</b>	43A/44A	
OPENING, UPPER SURFACE			
PETAL COLOR, WHEN	41B	47C	
OPENING, LOWER	1110		
SURFACE			
PETAL COLOR,	40B with $51A$	46C	
OPENED, UPPER	at base		
SURFACE	<b>2010</b>		
PETAL COLOR,	38B	46D	
OPENED, LOWER SURFACE			
PETAL COLOR, FADING	39 A	47C	
TO			
PEDUNCLE LENGTH	About 5.6 cm	About 4.5 cm	
FLOWER BUD COLOR	41B	47D	

ers persistent.

*Flowering season.*—Year-round under greenhouse conditions. In the garden, flowering is continuous from spring until fall.

*Flower longevity.*—About 3 or 4 days.

Flower size.—Diameter: About 7.5 cm. Depth: About 1.5 cm.

*Flower buds.*—Length: About 2.5 cm. Diameter: About 1.5 cm. Shape: Ovoid, pointed. Color: 41B.

- *Petals.*—Quantity: Five, overlapping. Length: About 3 cm. Width: About 3.5 cm. Shape: Obcordate with emarginate apex, attenuate base and entire margin. Texture: smooth, iridescent. Color: When opening, upper surface: 40A. When opening, lower surface: 41B. Fully opened, upper surface: 40B with 51A at base. Fully opened, lower surface: 38B. Upper surface fading to: 39A.
- Spur.—Length: About 6.6 cm. Shape: Narrow and curved. Color: 53C.

It is claimed:

1. A new and distinct cultivar of New Guinea Impatiens plant named 'BFP-796', as illustrated and described.

### **U.S. Patent**

· · · . · ·

. .

. . . . . . . . . . 

. . . . . . . .

. . . . . . . . . -----.....

· - · ·

. . . .

. . . . . . . . ·· · · · · · ·: :-: :-:

. . . . . . . . . .

· · · · · · · · · · · · · · · · · ·

#### Aug. 22, 2000

## Plant 11,486

۰.

.

· · .

••••••••••• 

.... . . . . . . . . . . . . .

. . . . . . . . . . . . . . . . . . .

•••••••

. .. . . . . . . . . .. . . . .... . . . . . . . . . . . . . . . . .... ... • . . . . . . . . . . . . . . . . .... ..... . . . . . . . . . . . . . . . . .

### . . . .

. . .

# 

· · · · · · · · · · . . . . . .

.....

·... ...

· · · · · · · . . . . . . . . . . .

. .

.

. . . . . . . . . . . . . . . . .

. . . . . . .

•••••••••••

.

•

• . . . . . . . . . . · · · · · · · · . . . . •••••

. . . . . . . .

- . . . · -. . . .

. : . : ' . . . . . . . . . . . ... ...