



US00PP11535P

# United States Patent [19]

## Kientzler

[11] Patent Number: Plant 11,535  
[45] Date of Patent: Oct. 3, 2000

[54] NEW GUINEA IMPATIENS PLANT NAMED 'KIPAB'

[75] Inventor: **Ludwig Kientzler**, Gensingen, Germany

[73] Assignee: **Paul Ecke Ranch**, Encinitas, Calif.

[21] Appl. No.: **09/093,166**

[22] Filed: **Jun. 8, 1998**

[51] Int. Cl.<sup>7</sup> ..... A01H 5/00

[52] U.S. Cl. ..... Plt./318

[58] Field of Search ..... Plt./318

[56]

### References Cited

#### U.S. PATENT DOCUMENTS

P.P. 10,858 4/1999 Kientzler ..... Plt./318  
P.P. 10,957 6/1999 Bull ..... Plt./318

Primary Examiner—Howard J. Locker

Assistant Examiner—Kent L. Bell

Attorney, Agent, or Firm—C. A. Whealy

[57]

### ABSTRACT

A new and distinct cultivar of New Guinea Impatiens plant named 'Kipab', characterized by its small and numerous pink-colored flowers; early flowering; upright and relatively compact growth habit; freely branching plant habit; and dark green, slightly shiny non-variegated leaves.

### 1 Drawing Sheet

2

by-side comparisons in Encinitas, Calif. under commercial practice, plants of the new New Guinea Impatiens differed from plants of the cultivar 'Grenada' in the following characteristics:

5 1. Plants of the new New Guinea Impatiens have much shorter internodes than plants of the cultivar 'Grenada'.

2. Leaves of plants of the new New Guinea Impatiens are shorter, narrower and less shiny than leaves of plants of the cultivar 'Grenada'.

10 3. The lower surfaces of leaves of plants of the new New Guinea Impatiens are green whereas the lower surfaces of leaves of plants of the cultivar 'Grenada' are dark purple.

4. Leaf petioles of plants of the new New Guinea Impatiens are longer and much lighter pink than leaf petioles of plants of the cultivar 'Grenada'.

15 5. Flowers of plants of the new New Guinea Impatiens are smaller than flowers of plants of the cultivar 'Grenada'.

6. Flower color of plants of the new New Guinea Impatiens is lighter and less salmon pink than flower color of plants of the cultivar 'Grenada'.

20 7. Plants of the new New Guinea Impatiens flower earlier than plants of the cultivar 'Grenada'.

25 A detailed comparison of plants of the new New Guinea Impatiens and the cultivar 'Grenada' appears in Chart A at the end of the specification.

### BRIEF DESCRIPTION OF THE PHOTOGRAPHS

30 The accompanying colored photographs illustrate the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type.

35 The photograph at the top of the sheet comprises a side perspective view of a typical plant of 'Kipab'.

40 The photograph at the bottom of the sheet comprises a close-up view of a typical flower and leaves of 'Kipab' (left) and 'Grenada' (right). Flower and foliage colors in the photographs may differ from the actual colors due to light reflectance.

### DETAILED BOTANICAL DESCRIPTION

45 The cultivar Kipab has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as tempera-

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Kipab'. These characteristics in combination distinguish 'Kipab' as a new and distinct cultivar:

1. Small and numerous pink-colored flowers.
2. Early flowering.
3. Upright and relatively compact growth habit.
4. Freely branching plant habit.
5. Dark green, slightly shiny non-variegated leaves.

Plants of the new New Guinea Impatiens are more compact and have darker pink flowers than plants of the male parent, MF 352. Compared to plants of the female parent, MF 291, plants of the new New Guinea Impatiens have greener foliage, and lighter pink and more rounded flowers.

Plants of the new New Guinea Impatiens are similar in flower color to the New Guinea Impatiens cultivar 'Grenada' (disclosed in U.S. Plant Pat. No. 9,343). However in side-

# Plant 11,535

**3**

ture and light intensity, without, however, any variance in genotype. The following observations, measurements and comparisons describe plants grown in 10-cm pots in Encinitas, Calif., under commercial practice in a fiberglass-covered greenhouse with day temperatures ranging from 23 to 29° C. and night temperatures ranging from 16 to 18° C. and light levels of about 3,000 footcandles.

In the following description, color references are made to The Royal Horticultural Society Colour Chart except where general terms of ordinary dictionary significance are used.

**Classification:**

*Botanical.*—*Impatiens hawkeri* cultivar ‘Kipab’.

*Commercial.*—New Guinea Impatiens cultivar ‘Kipab’.

**Parentage:**

*Male parent.*—Proprietary selection of *Impatiens hawkeri* identified as MF 352.

*Female parent.*—Proprietary selection of *Impatiens hawkeri* identified as MF 291.

**Propagation:**

*Type cutting.*—Terminal cuttings.

*Time to initiate roots.*—Summer: About 14 days with 21° C. soil temperature. Winter: About 18 days with 21° C. soil temperature.

*Time to develop roots.*—Summer: About 21 days with 21° C. soil temperature. Winter: About 24 days with 21° C. soil temperature.

**Plant description:**

*Plant form.*—Upright, rounded and mounding.

*Growth habit.*—Moderately vigorous. Moderate branching, dense and bushy growth. Appropriate for 10 and 12.5-cm containers.

*Crop time.*—From planting of a rooted cutting, about six to eight weeks are required to produce a finished flowering plant in a 12.5-cm container.

*Plant size.*—Height: About 15.5 cm. Width or spread: About 22 cm.

*Lateral branches.*—Quantity: About six. Length: About 9 cm. Diameter: About 7 mm. Internode length: About 2.75 cm. Color: 59B.

*Foliage description.*—Leaves simple, generally symmetrical, abundant, opposite or in whorls of three, horizontal to plant and flat in aspect. Length: About 10.5 cm. Width: About 4.2 cm. Shape: Elliptic with acuminate apex, attenuate base and serrulate margin with ciliation. Texture: Smooth, slightly shiny. Color: Young foliage, upper surface: Brighter green than 147A. Young foliage, lower surface: 147C. Fully expanded foliage, upper surface: 147A. Fully expanded foliage, lower surface: 147C. Venation, upper surface: 148D. Venation, lower surface: 51D. Petiole: Length: About 4 cm. Diameter: About 3.5 mm. Color: 51C.

**Flower description:**

*Flower type and habit.*—Small pink-colored flowers. Freely and continuously flowering. Flowers arise from leaf axils. Usually about seven or eight flowers per lateral branch. Flowers positioned above the foliage and face slightly outward. Flowers flat and rounded triangular. Flowers persistent.

*Flowering season.*—Year-round under greenhouse conditions. In the garden, flowering is continuous from

**4**

spring until fall. Flowers typically last less than one week depending on environmental and cultural conditions.

*Flower size.*—Length: About 4 cm. Width: About 3.5 cm. Depth: About 4 mm.

*Flower buds.*—Length: About 1.8 cm. Diameter: About 1.2 cm. Shape: Ovoid. Color: 54C/54D.

*Petals.*—Quantity: Five. Length: Upper petal: About 2 cm. Middle petals: About 1.8 cm. Lower petals: About 2.5 cm. Width: Upper petal: About 3 cm. Middle petals: About 1.8 cm. Lower petals: About 2.6 cm. Shape: Cordate with emarginate apex, acute base and entire margin. Texture: Smooth, iridescent. Color: When opening, upper surface: 55B. When opening, lower surface: 55C. Fully opened, upper surface: 55B/55C with distinct eye, 58B to 57A at attachment. Fully opened, lower surface: 55D. Fading to: 55D with distinct eye, 57A.

*Spur.*—Length: About 4.1 cm. Shape: Narrow and curved. Color: 51A.

*Peduncles.*—Length: About 4.7 cm. Angle: Erect and slightly outward. Strength: Moderately strong. Color: 50B/50C.

*Reproductive organs.*—Androecium: Stamen number: Five, anthers fused, filaments free. Anther shape: Obovate. Anther size: 3 mm by 4 mm. Anther color: 4D to white. Amount of pollen: Moderate. Pollen color: Cream. Gynoecium: Five-loculate fused. Gynoecium length: About 3 mm. Gynoecium color: 145A.

**Disease resistance:** Under commercial conditions, resistance to pathogens has not been observed.

**Seed development:** Seed production has not been observed.

<u>CHART A</u>		
CHARACTERISTICS	‘KIPAB’	‘GRENADA’
INTERNODE LENGTH	About 2.75 cm	About 4 cm
LEAF LENGTH	About 10.5 cm	About 11.75 cm
LEAF WIDTH	About 4.2 cm	About 5 cm
LEAF COLOR, FULLY EXPANDED, LOWER SURFACE	147C	59A/59B
VENATION COLOR, UPPER SURFACE	148D	60A
VENATION COLOR, LOWER SURFACE	51D	59A
PETIOLE LENGTH (length by width)	About 4 cm by 3.5 cm	About 5 cm by 5 cm
SPUR LENGTH	About 4.1 cm	About 4.3 cm
PETAL COLOR, WHEN OPENING, UPPER SURFACE	55B	47D
PETAL COLOR, OPENED, UPPER SURFACE	55B/55C Eye, 58B/57A	48C Eye, 57A
PETAL COLOR, OPENED, LOWER SURFACE	55D	48D
PEDUNCLE COLOR	50B/50C	47A/47B

It is claimed:

1. A new and distinct cultivar of New Guinea Impatiens plant named ‘Kipab’, as illustrated and described.

\* \* \* \* \*

**U.S. Patent**

**Oct. 3, 2000**

**Plant 11,535**

