US00PP11700P

United States Patent [19] Klemm

- NEW GUINEA IMPATIENS PLANT NAMED [54] 'MIAMI'
- Siegfried Klemm, Stuttgart, Germany [75] Inventor:
- Assignee: Klemm + Sohn, Stuttgart, Germany [73]
- Appl. No.: 09/213,449 [21]
- Dec. 17, 1998 [22] Filed:
- Int. Cl.⁷ [51] A01H 5/00 [52] **US CL Plt /318**

Plant 11,700 **Patent Number:** [11] **Date of Patent:** Dec. 19, 2000 [45]

Primary Examiner—Howard J. Locker Assistant Examiner—Michelle Kizilkaya Attorney, Agent, or Firm—C. A. Whealy

[57] ABSTRACT

A new and distinct cultivar of New Guinea Impatiens plant named 'Miami', characterized by its numerous large red purple-colored flowers that are held above the foliage; uniform and rounded growth habit; medium plant height; freely branching, dense and bushy plant habit; and glossy, dark green leaves.

$\begin{bmatrix} J^{2} \end{bmatrix}$	U.S. CI	••••••••••••••••••	111./310
[58]	Field of Search	•••••••••••••••••••••••••••••••••••••••	Plt./318

1 Drawing Sheet

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of New Guinea Impatiens plant, botanically known as Impatiens hawkeri, and hereinafter referred to by the cultivar name Miami.

The new cultivar is a product of a planned breeding program conducted by the Inventor in Stuttgart, Germany. The objective of the breeding program was to develop varieties with good branching, uniform and rounded growth 10 habit, medium height, early flowering, floriferousness, bright flower color, and large round flowers.

The new cultivar originated from a cross made by the Inventor of the New Guinea Impatiens selection identified as K 127, not patented, as the male, or pollen parent, with the 15 nonpatented New Guinea Impatiens cultivar Hongkong as the female, or seed parent.

Plants of the new Impatiens are similar in flower color to plants of the nonpatented New Guinea Impatiens cultivar Kampala. However, in side-by-side comparisons conducted in Stuttgart, Germany, plants of the new Impatiens have a slightly lighter flower color, larger flowers and more uniform plant growth than plants of the cultivar Kampala.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. The photograph comprises a top perspective view of a typical plant of 'Miami'. Flower and foliage colors in the photographs may differ from the actual colors due to light reflectance.

The cultivar Miami was discovered and selected by the Inventor in 1994 as a flowering plant within the progeny of 20 the stated cross in a controlled environment in Stuttgart, Germany.

Asexual reproduction of the new cultivar by terminal cuttings taken at Stuttgart, Germany, has shown that the unique features of this new Impatiens are stable and repro-25 duced true to type in successive generations of asexual reproduction.

SUMMARY OF THE INVENTION

-30 The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Miami'. These characteristics in combination distinguish 'Miami' as a new and distinct cultivar:

1. Numerous large red purple-colored flowers that are 35 held above the foliage.

2. Uniform and rounded growth habit.

3. Medium plant height.

DETAILED BOTANICAL DESCRIPTION

Plants of the cultivar Miami have not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and light intensity, without, however, any variance in genotype. The following observations, measurements and comparisons describe plants grown in Stuttgart, Germany, under greenhouse and outdoor production practices.

In the following description, color references are made to The Royal Horticultural Society Colour Chart except where general terms of ordinary dictionary significance are used.

Classification:

Botanical.—Impatiens hawkeri cultivar Miami.

Commercial.—New Guinea Impatiens cultivar Miami. Parentage:

Male parent.—Impatiens hawkeri selection identified as K 127.

Female parent.—Impatiens hawkeri cultivar Hongkong, not patented. 40 Propagation:

4. Freely branching, dense and bushy plant habit. 5. Glossy, dark green leaves.

Compared to plants of the male parent, the salmon pink selection identified as K 127, plants of the new Impatiens differ in flower color, are more compact, more uniform, more upright, and have larger and more round flowers. ⁴⁵ Compared to plants of the female parent, the violet cultivar Hongkong, plants of the new Impatiens have a lighter flower color, are more uniform, have larger leaves, and have larger and more round flowers.

Type cutting.—Terminal cuttings.

Time to initiate and develop roots.—About 16 days. Plant description:

Crop time (precocity).—From rooted cuttings to flowering, about 80 to 90 days are required. *Plant form.*—Uniform and rounded growth habit, plants appear almost spherical; upright yet spreadıng.

Plant 11,700

3

Growth habit.—Moderately vigorous. Freely branching, dense and bushy growth. Appropriate for 10 to 25-cm containers.

- *Plant size*.—Height, about 15.5 cm; spread, about 27 cm.
- Lateral stems.—Quantity: About 10 basal branches; each basal branch with about 2 lateral stems. Length: About 15.5 cm.
- Foliage description.—Medium-sized leaves; simple, generally symmetrical, moderate in quantity, opposite or in whorls, usually flat. Shape: Oblonglanceolate; margin, shallow serration. Texture: Smooth, glossy. Color: Upper surface: A mixture between 136A and 147A; venation, 145C. Lower surface: 148D, venation, main veins, 181C, and lateral veins, 146B. Petiole length: About 2 cm.

4

Flower buds.—Length: About 1.2 cm. Diameter: About 7 mm. Shape: Ovoid. Color: 78B.

- Petals.—Quantity: Five. Shape: Cordate with acute base and entire margin. Texture: Satiny, smooth.
 Color: Fully opened, upper surface: Red purple, 74A; base or "eye", 46B. Color may fade slightly with subsequent development. Fully opened, lower surface: A mixture between 74B and 74C.
- Sepals.—Three, lower sepal spurred. Color, lateral sepals: Inside: 138C. Outside: 148B. Color, spurred sepal: Inside: 65B. Outside: Center, 53A; surround-ing area, 65B.

Flower description:

- Flower type and habit.—Large red purple-colored flowers. Early and freely flowering; usually about 30 flowers per plant. Single flowers arranged in clusters arising from leaf axils. Flowers positioned above the foliage and face upward and outward. Flower shape is rounded. Flowers not persistent.
- *Flowering season.*—Year-round under greenhouse conditions. In the garden, flowering is continuous from spring until fall.
- Flower longevity.—About 10 to 14 days on the plant; about 1 to 2 days as a cut flower.
- Flower size.—Diameter: About 6.5 cm. Depth: About 1.4 cm.

- Peduncles.—Length: About 3.8 cm. Angle: About 45° to stem. Strength: Very stiff, strong. Color: Redbrown.
- Reproductive organs.—Androecium: Stamen number: Five, anthers fused, filaments free. Anther shape: Obovate. Pollen color: Cream. Gynoecium: Fiveloculate fused. Style length: About 4 mm. Style color: Green. Stigma color: Cream.
 Fruit.—Green, pulpy explosive capsule.
 Seed.—Brown, about 2 to 3 mm in length.
- Disease resistance: Under commercial conditions, good resistance to Botrytis has been observed.

It is claimed:

1. A new and distinct cultivar of New Guinea Impatiens plant named 'Miami', as illustrated and described.

* * * * *

U.S. Patent

Dec. 19, 2000

Plant 11,700

