



US00PP11818P2

(12) **United States Plant Patent**  
**VandenBerg**

(10) **Patent No.:** **US PP11,818 P2**

(45) **Date of Patent:** **Mar. 13, 2001**

(54) **CHRYSANTHEMUM PLANT NAME**  
**'YOBETH'**

(75) **Inventor:** **Cornelis P. VandenBerg**, Salinas, CA  
(US)

(73) **Assignee:** **Yoder Brothers, Inc.**, Barberton, OH  
(US)

(\* ) **Notice:** Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

(21) **Appl. No.:** **09/224,727**

(22) **Filed:** **Jan. 4, 1999**

(51) **Int. Cl.<sup>7</sup>** ..... **A01H 5/00**

(52) **U.S. Cl.** ..... **Plt./287; Plt./286**

(58) **Field of Search** ..... **Plt./287, 286**

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

P.P. 10,212 \* 1/1998 Wain ..... Plt./287

\* cited by examiner

*Primary Examiner*—Howard J. Locker

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A distinct cultivar of Chrysanthemum plant named 'Yobeth',  
characterized by its uniformly mounded plant habit;  
decorative-type inflorescences that are about 3.6 cm in  
diameter; attractive reddish purple-colored ray florets and;  
numerous inflorescences per plant.

**1 Drawing Sheet**

**1**

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct cultivar  
of Chrysanthemum plant, botanically known as *Dendran-  
thema grandiflora* and referred to by the cultivar name  
Yobeth.

The new cultivar is a product of a planned breeding  
program conducted by the Inventor in Salinas, Calif. The  
objective of the breeding program is to create new garden-  
type Chrysanthemum cultivars having inflorescences with  
desirable inflorescence forms and floret colors and good  
garden performance.

The new cultivar originated from a cross made by the  
Inventor in November, 1991, of the *Dendranthema grandiflora*  
cultivar Bravo (U.S. Plant Pat. No. 6,888) as the  
female, or seed, parent with an unidentified proprietary  
seedling selection as the male, or pollen parent.

The cultivar Yobeth was discovered and selected by the  
Inventor as a flowering plant within the progeny of the  
cross in a controlled environment in Salinas, Calif., in  
October, 1992. The selection of this plant was based on its  
desirable inflorescence form and ray floret color.

Asexual reproduction of the new cultivar by terminal  
cuttings taken in a controlled environment in Salinas, Calif.,  
has shown that the unique features of this new Chrysanthemum  
are stable and reproduced true to type in successive  
generations.

**SUMMARY OF THE INVENTION**

The cultivar Yobeth has not been observed under all  
possible environment conditions. The phenotype may vary  
somewhat with variations in environment such as  
temperature, daylength and light intensity, without,  
however, any variance in genotype.

The following traits have been repeatedly observed and  
are determined to be the unique characteristics of 'Yobeth'.  
These characteristics in combination distinguish 'Yobeth' as  
a new and distinct cultivar:

1. Uniformly mounded plant habit.
2. Relatively small decorative-type inflorescences that are  
about 3.6 cm in diameter.

**2**

3. Attractive reddish purple-colored ray florets.

4. Numerous inflorescences per plant.

The new Chrysanthemum is similar to the parent cultivar  
Bravo. However in side-by-side comparisons under com-  
mercial practice, plants of the new Chrysanthemum differed  
from plants of the cultivar Bravo in the following charac-  
teristics:

1. Plants of the new Chrysanthemum are fuller and more  
dense than plants of the cultivar Bravo.
2. Plants of the new Chrysanthemum flower about one  
week later than plants of the cultivar Bravo grown  
under natural season conditions.
3. Ray florets of plants of the new Chrysanthemum are  
reddish purple in color whereas ray florets of plants of  
the cultivar Bravo are red in color.

**BRIEF DESCRIPTION OF THE PHOTOGRAPHS**

The accompanying photographs illustrate the overall  
appearance of the new cultivar. The photograph at the top of  
the sheet comprises a side perspective view of a typical  
flowering plant of 'Yobeth'. The photograph at the bottom of  
the sheet comprises a close-up view of typical inflorescences  
of the cultivar 'Yobeth'. These photographs show the colors  
as true as it is reasonably possible to obtain in colored  
reproductions of this type. Floret and foliage colors in the  
photographs may differ from the actual colors due to light  
reflectance.

**DETAILED BOTANICAL DESCRIPTION**

In the following description, color references are made to  
The Royal Horticultural Society Colour Chart except where  
general terms of ordinary dictionary significance are used.  
The following observations and measurements describe  
plants grown in Leamington, Ontario, Canada, under con-  
ditions which approximate those generally used in commer-  
cial garden Chrysanthemum production. One rooted cutting  
was planted in a 15-cm container on Jul. 20, 1998 and plants  
were grown outdoors under natural season conditions. Mea-  
surements and numerical values represent averages for typi-  
cal flowering containers.

Botanical classification: *Dendranthema grandiflora* cultivar Yobeth.

Commercial classification: Decorative-type garden chrysanthemum.

Parentage:

*Female or seed parent.*—*Dendranthema grandiflora* cultivar Bravo, disclosed in U.S. Plant Pat. No. 6,888.

*Male or pollen parent.*—Unidentified proprietary seedling selection.

Propagation:

*Type.*—Terminal tip cuttings.

*Time to rooting.*—Seven to ten days with soil temperatures of 21° C.

*Rooting habit.*—Fine, fibrous and well-branched.

Plant description:

*Appearance.*—Perennial herbaceous decorative-type garden Chrysanthemum. Inverted triangle. Stems initially upright, then outwardly spreading giving a uniformly mounded appearance to the plant. Freely branching with lateral branches potentially developing at every node, when pinched, about 10 laterals develop.

*Plant height.*—About 30 cm.

*Plant spread.*—About 43 cm.

*Foliage description.*—Leaf arrangement: Alternate. Length: About 4.7 cm. Width: About 3.5 cm. Apex: Mucronate. Base: Attenuate. Margin: Palmately lobed, sinuses parallel. Texture: Upper and lower surfaces sparsely pubescent. Veins prominent on lower surface. Petiole length: About 1.7 cm. Petiole diameter: About 2 mm. Color: Young foliage upper surface: 147A. Young foliage lower surface: 147B. Mature foliage upper surface: 147A. Mature foliage lower surface: 147B. Venation upper surface: 147A to 147B. Venation lower surface: 147B.

Inflorescence description:

*Appearance.*—Decorative-type inflorescence form with oblong-shaped ray florets. Inflorescences borne on terminals above foliage, arising from leaf axils. Disk and ray florets arranged acropetally on a capitulum. One inflorescence per terminal with numerous inflorescences per plant, about 10 per lateral stem.

*Flowering response.*—Under natural season conditions, plant flowers in mid to late September in the Northern Hemisphere about 73 days after planting, and flower for at least three weeks depending on weather conditions.

*Inflorescence bud. (Before showing color).*—Height: About 4 mm. Diameter: About 6 mm. Phyllary color: Close to 141A.

*Inflorescence size.*—Diameter: About 3.6 cm. Depth (height): About 1.1 cm. Diameter of disc: About 3 mm.

*Ray florets.*—Shape: Oblong. Length: About 1.6 cm. Width: About 5 mm. Apex: Dentate. Margin: Entire. Texture: Smooth, glabrous, satiny. Orientation: Initially upright, then horizontal. Number of ray florets per inflorescence: Typically more than 100. Color: When opening: Upper surface: 59A, white at base. Lower surface: 59B, white at base. Opened inflorescence: Upper surface: 59B, fading to 59C to 77C to 77D; white at base. Lower surface: 77C to 77D, white at base.

*Disc florets.*—Shape: Tubular, apex dentate. Length: About 3.5 mm. Width: Apex: About 1 mm. Base: About 1 mm. Number of disc florets per inflorescence: Typically about 12. Color: Immature: 154A. Mature: Apex: 9A. Mid-section and base: White.

*Peduncle.*—Aspect: Flexible, angled about 45° to the stem. Length: First peduncle: About 3.4 cm. Fourth peduncle: About 8.5 cm. Diameter: About 1.5 mm. Texture: Pubescent. Color: Close to 147A.

*Reproductive organs.*—Androecium: Present on disc florets only. Anther color: 9A. Pollen: Scare, 9A. Gynoecium: Present on both ray and disc florets.

Disease resistance: Resistance to known Chrysanthemum diseases has not been observed on plants grown under commercial production conditions.

Seed production: Seed production has not been observed.

It is claimed:

1. A new and distinct cultivar of Chrysanthemum plant named 'Yobeth', as illustrated and described.

\* \* \* \* \*

