



US00PP12466P2

(12) **United States Plant Patent**
Akerboom

(10) **Patent No.:** **US PP12,466 P2**

(45) **Date of Patent:** **Mar. 19, 2002**

(54) **ASTER PLANT NAMED 'KARMIJN MILKA'**

PP10,624 P * 9/1998 Akerboom Plt./355

(75) Inventor: **Petrus J. Akerboom**, Ter Aar (NL)

OTHER PUBLICATIONS

(73) Assignee: **De Nachtvliinder B.V.**, Ter Aar (NL)

UPOV-ROM GTITM Computer Database 2000/02, GTI
JOUVE Retrieval Software, citations for 'Karmijn Milka',
May 2000.*

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

* cited by examiner

(21) Appl. No.: **09/263,153**

Primary Examiner—Bruce R. Campell

(22) Filed: **Mar. 5, 1999**

Assistant Examiner—Wendy C Baker

(51) **Int. Cl.**⁷ **A01H 5/00**

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(52) **U.S. Cl.** **Plt./355**

(57) **ABSTRACT**

(58) **Field of Search** Plt./355

A distinct cultivar of Aster plant named 'Karmijn Milka',
characterized by its freely and uniform flowering; decorative
inflorescence form with purple ray florets; and good post-
production longevity.

(56) **References Cited**

1 Drawing Sheet

U.S. PATENT DOCUMENTS

PP10,602 P * 9/1998 Akerboom Plt./355

1

2

BACKGROUND OF THE INVENTION

The present Invention relates to a new and distinct culti-
var of Aster plant, botanically known as *Aster novi-belgii*
and referred to by the cultivar name Karmijn Milka.

The cultivar Karmijn Milka has not been observed under
all possible environmental conditions; The phenotype may
vary somewhat with variations in environment such as
temperature, daylength and light intensity, without,
5 however, any variance in genotype.

The new Aster is a whole plant mutation of the commer-
cial *Aster novi-belgii* cultivar Milka, disclosed in U.S. Plant
Pat. No. 10,602. The new Aster was discovered and selected
by the Inventor in a controlled environment in Ter Aar, The
Netherlands, as a single flowering plant among a population
of plants of the parent cultivar Milka in June, 1994. This
single plant consistently formed inflorescences having
darker purple ray florets compared to the lighter purple ray
florets of the cultivar Milka.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the
overall appearance of the new Aster, showing the colors as
true as it is reasonably possible to obtain in colored repro-
ductions of this type. Colors in the photograph may differ
slightly from the color values cited in the detailed botanical
description which accurately describe the colors of the new
Aster. The photograph comprises a side perspective view of
a typical cut flower stem of the new Aster.

Asexual reproduction of the new Aster by terminal cut-
tings taken at Ter Aar, The Netherlands, has shown that the
unique features of this new Aster are stable and reproduced
true to type in successive generations.

SUMMARY OF THE INVENTION

DETAILED BOTANICAL DESCRIPTION

The following traits have been repeatedly observed and
are determined to be the unique characteristics of 'Karmijn
Milka'. These characteristics in combination distinguish
'Karmijn Milka' as a new and distinct cultivar:

In the following description, color references are made to
The Royal Horticultural Society Colour Chart except where
general terms of ordinary dictionary significance are used.
The following observations and measurements describe
plants grown as cut flowers during the summer in Ter Aar,
The Netherlands, in a glass-covered greenhouse with aver-
age day temperatures of 17 to 35° C. and night temperatures
of 14 to 20° C. Plants received long day/short night
treatments, 18 hours light, for seven weeks and then were
forced into flower with short day/long night treatments, 12
hours light.

1. Freely and uniform flowering.
2. Decorative inflorescence form with purple ray florets.
3. Good post-production longevity.

Plants of the new Aster can be compared to plants of the
sibling cultivar Pink Milka (U.S. Plant patent application
Ser. No. 09/263,152). Plants of the new Aster have narrowly
elliptic and purple ray florets whereas plants of the cultivar
Pink Milka have narrowly obovate and light pink ray florets.

Botanical classification: *Aster novi-belgii* cultivar Karmijn
Milka.

Commercial classification: Cut flower Aster; can also be
produced as a potted plant.

Plants of the new Aster can also be compared to plants of
the sibling cultivar Dark Milka (U.S. Plant patent applica-
tion Ser. No. 09/263,154). Plants of the new Aster have
lighter purple ray florets than plants of the cultivar Dark
Milka.

Parentage: Whole plant mutation of *Aster novi-belgii* culti-
var Milka, disclosed in U.S. Plant Pat. No. 10,602.

Propagation:

Type.—Terminal tip cuttings.

Time to initiate roots.—About 8 to 10 days at a temperature of 18 to 23° C.

Time to develop roots.—Summer: About 16 to 20 days at a temperature of 23° C. Winter: About 20 to 25 days at a temperature of 18° C.

Root description.—Fine, fibrous and well-branched.

Plant description:

Appearance.—Herbaceous cut flower. Stems upright; medium plant height at flowering. Freely and uniformly flowering; purple-colored decorative inflorescences.

Crop time.—About 12.5 to 13 weeks from planting to harvest of cut flowering stems.

Branching habit.—Freely branching, typically more than 25 lateral branches per plant.

Growth rate.—Moderate to slow.

Plant height.—About 100 to 110 cm.

Plant width.—About 35 cm.

Lateral branch length.—At the top of the flowering stem: About 1 mm. At the middle of the flowering stem: About 18 to 30 cm. At the bottom of the flowering stem: About 2 to 12.5 cm.

Lateral branch diameter.—At the top of the flowering stem: About 1.2 mm. At the middle of the flowering stem: About 1.4 to 1.9 mm. At the bottom of the flowering stem: About 1.9 to 2.3 mm.

Internode length.—5th node from apex: About 1.4 to 1.8 cm. 10th node from apex: About 1.9 to 2.5 cm. 15th node from apex: About 2.3 to 2.6 cm. 20th node from apex: About 2 to 3.2 cm.

Stem color.—Green, 144A to 146A, little to no anthocyanin at internodes.

Stem texture.—Slightly pubescent.

Foliage description.—Arrangement: Alternate. Quantity: About 45 to 48 per lateral stem. Shape: Narrowly elliptic; apex acute; base attenuate, sessile; margin serrate. Size: At the top of the flowering stem: About 1.1 to 1.6 cm in length and about 2 to 2.4 mm in width. At the middle of the flowering stem: About 14.1 to 15.7 cm in length and about 1.3 to 1.7 cm in width. At the bottom of the flowering stem: About 16.8 to 19.8 cm in length and about 2 to 2.5 cm in width. Texture: Pubescent on upper surface, leathery. Color: Young leaves, upper surface: 147A. Young leaves, lower surface: Slightly lighter than 137C. Fully expanded leaves, upper surface: 147A. Fully expanded leaves, lower surface: 146B. Venation, upper and lower surfaces: 145C.

Flowering description:

Appearance.—Decorative inflorescence form. Inflorescences held on wiry peduncles, arising from leaf axils; inflorescences face upright. Disc and ray florets arranged acropetally on a capitulum.

Flowering response.—Under natural conditions, plants flower in the late summer/autumn. At other times of the year, inflorescence initiation and development can be induced under short day/long night conditions. Response time is about 5.5 to 6 weeks.

Post-production longevity.—Good, inflorescences last about 3.5 to 4.5 weeks on the plant; about 2 to 3 weeks as a cut flower without postproduction treatments. Inflorescences persistent.

Quantity of inflorescences.—Inflorescences form at every leaf axil. Freely flowering, usually about 54 to 78 inflorescences per flowering stem.

Fragrance.—None.

Inflorescence size.—Diameter: About 3.5 cm. Depth (height): About 9.5 to 14 mm. Disc diameter: About 8 mm.

Inflorescence bud.—Shape: Cylindrical. Length: About 7 to 8 mm. Diameter: About 7 to 8 mm. Color: 137B.

Ray florets.—Quantity of ray florets per inflorescence: About 122 to 152. Shape: Narrowly elliptic; apex rounded; base attenuate; margin entire. Length: About 1.1 to 1.3 cm. Width: About 1.2 to 1.9 mm. Texture: Satiny, smooth and glabrous. Color: When opening, upper surface: 82B. When opening, lower surface: 82A to 82B. Mature, upper surface: 82B; fading to 82D. Mature, lower surface: 82B.

Disc florets.—Quantity: About 107 per inflorescence. Shape: Tubular. Length: About 6 mm. Diameter: At apex, about 2 mm; at base, about 1 mm. Color: Immature: 145A to 154A. Mature: 9A.

Peduncle.—Strength: Strong. Length: Apical peduncle: About 4 to 6 mm. Fourth peduncle: About 2 to 3 mm. Seventh peduncle: About 2 mm. Color: 146A to 147A.

Reproductive organs.—Androecium: Not present on ray florets. Gynoecium: Pistil length: About 3.5 to 4.5 mm. Stigma color: Yellow, 10C to 10D. Style length: About 2 mm.

Seed development.—Not observed.

Disease resistance: Plants of the new Aster have not been observed to be resistant to pathogens common to Asters, however plants of the new Aster appear to be less sensitive to Powdery Mildew than other known cultivars of Aster.

It is claimed:

1. A new and distinct cultivar of Aster plant named 'Karmijn Milka', as illustrated and described.

* * * * *

