



US00PP12476P2

(12) **United States Plant Patent**
Trees

(10) **Patent No.:** **US PP12,476 P2**
(45) **Date of Patent:** **Mar. 19, 2002**

(54) **GERANIUM PLANT NAMED ‘BALFANFLAI’**

(75) **Inventor:** **Scott C. Trees**, Shell Beach, CA (US)

(73) **Assignee:** **Ball FloraPlant, A Division of Ball Horticultural Co.**, West Chicago, IL (US)

(*) **Notice:** Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) **Appl. No.:** **09/822,008**

(22) **Filed:** **Mar. 31, 2001**

(51) **Int. Cl.⁷** **A01H 5/00**

(52) **U.S. Cl.** **Plt./330**

(58) **Field of Search** **Plt./330**

Primary Examiner—Bruce R. Campell
Assistant Examiner—Kent L. Bell
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of Geranium plant named ‘Balfanflai’, characterized by its upright and mounded growth habit; freely basal branching habit; dark green leaves with darker green zonation pattern; and light red-colored semi-double flowers.

1 Drawing Sheet

1

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of Geranium plant, botanically known as *Pelargonium×hortorum*, and hereinafter referred to by the name ‘Balfanflai’.

The new Geranium is a product of a planned breeding program conducted by the Inventor in Arroyo Grande, Calif. The objective of the breeding program was to develop new cultivars with freely-branching and vigorous growth habit, dark green-colored foliage, uniform flowering, and interesting flower and foliage colors.

The new Geranium originated from a cross made by the Inventor of the proprietary *Pelargonium×hortorum* selection 7788-35 as the female, or seed, parent with the *Pelargonium×hortorum* cultivar ‘Isabell’, disclosed in U.S. Plant Pat. No. 7,080, as the male, or pollen, parent. The cultivar ‘Balfanflai’ was discovered and selected by the Inventor as a flowering plant within the progeny from this cross in a controlled environment in Arroyo Grande, Calif., in 1998.

Asexual reproduction of the new cultivar by terminal cuttings taken at Arroyo Grande, Calif., has shown that the unique features of this new Geranium are stable and reproduced true to type in successive generations of asexual reproduction.

SUMMARY OF THE INVENTION

The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘Balfanflai’. These characteristics in combination distinguish ‘Balfanflai’ as a new and distinct cultivar:

1. Upright and mounded growth habit.
2. Freely basal branching habit.
3. Dark green leaves with darker green zonation pattern.
4. Light red-colored semi-double flowers.

Plants of the new Geranium differ primarily from plants of the female parent selection in flower color and flower form. Plants of the new Geranium differ from plants of the male parent, the cultivar ‘Isabell’, in the following characteristics:

1. Plants of the new Geranium are less vigorous and have shorter internodes than plants of the cultivar ‘Isabell’.

2

2. Plants of the new Geranium have smaller umbels than plants of the cultivar ‘Isabell’.

3. Plants of the new Geranium have shorter peduncles than plants of the cultivar ‘Isabell’.

The new Geranium can be compared to the cultivar, ‘Tango Dark Red’, not patented. However, in side-by-side comparisons conducted in Arroyo Grande, Calif., plants of the new Geranium differ from plants of the cultivar ‘Tango Dark Red’ in the following characteristics:

1. Plants of the new Geranium are taller and more vigorous than plants of the cultivar ‘Tango Dark Red’.
2. Plants of the new Geranium have shorter internodes than plants of the cultivar ‘Tango Dark Red’.
3. Plants of the new Geranium have larger flowers than plants of the cultivar ‘Tango Dark Red’.
4. Plants of the new Geranium have longer peduncles than plants of the cultivar ‘Tango Dark Red’.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. The photograph comprises a side perspective view of a typical flowering plant of ‘Balfanflai’. Flower and foliage colors in the photograph may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new Geranium.

DETAILED BOTANICAL DESCRIPTION

The cultivar ‘Balfanflai’ has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and light intensity, without, however, any variance in genotype. The following observations, measurements and comparisons describe plants grown 10-cm pots in West Chicago, Ill., under commercial practice in a polycarbonate-covered greenhouse with day temperatures about 20 to 22° C., night temperatures about 18 to 20° C. and light levels about 2,500 to 3,000 foot-candles. Plants used for the photograph and description were about 8 to 10 weeks from

planting rooted cuttings. In the following description, color references are made to The Royal Horticultural Society Colour Chart except where general terms of ordinary dictionary significance are used.

Botanical classification: *Pelargonium x hortorum* cultivar 'Balfanflai'.

Parentage:

Female parent.—Proprietary *Pelargonium x hortorum* selection 7788-35.

Male parent.—*Pelargonium x hortorum* cultivar 'Isabell', disclosed in U.S. Plant Pat. No. 7,080.

Propagation:

Type cutting.—Terminal cuttings.

Time to initiate roots.—About 7 days at 18° C.

Time to develop roots.—About 21 days at 18° C.

Root description.—Fibrous, freely branching.

Plant description:

General appearance.—Upright and mounded growth habit. Appropriate for 10-cm and larger containers.

Crop time.—Starting with a rooted cutting, about 8 to 10 weeks are required to produced a finished flowering plant in 10-cm container.

Growth and branching habit.—Vigorous and freely basal branching with about 5 lateral branches at the base. Pinching, that is, removal of terminal apices, is typically not required.

Plant height (to top of foliage).—About 13 cm.

Plant width.—About 19.2 cm.

Lateral branches.—Length: About 11 cm. Internode length: About 7 mm. Color: 143C.

Foliage description.—Leaves simple, generally symmetrical, abundant, opposite. Quantity of leaves per lateral branch: About 18. Length: About 4.4 cm. Width: About 7.7 cm. Shape: Reniform. Apex: Rounded. Base: Cordate. Margin: Crenate, undulate. Venation: Palmate. Texture: Velvety; upper surface pubescent and pubescence along veins on lower surface. Color: Young foliage, upper surface: Closest to 139A; slight zonation pattern, 139A in color and located on the outer half of the leaf. Young foliage, lower surface: Closest to 137C. Fully expanded foliage, upper surface: Slightly lighter than 139A; slight zonation pattern, 139A in color, about 2 cm in width, and located on the outer half of the leaf. Fully expanded foliage, lower surface: Closest to 137B. Venation, upper and lower surfaces: Closest to 144A. Petiole: Length: About 9.5 cm. Diameter: About 2 mm. Color: Closest to 144A.

Flower description:

Flower arrangement and type.—Light red-colored flowers arranged in hemispherical umbels arising

from apical leaf axils. Umbels displayed above the foliage. At full flowering, usually about 3 open umbels and about 3 developing umbels per plant. Flowers semi-double and rounded in shape. Umbels persistent. Flowers not fragrant.

Flowering season.—Year-round under greenhouse conditions. In the garden, flowering is continuous from spring until fall.

Umbels.—Number of flowers per umbel: About 8. Diameter: About 7.6 cm. Height: About 4.6 cm. Flower diameter: About 4.3 cm.

Flower buds.—Length: About 1.2 cm. Diameter: About 7 mm. Shape: Ovoid. Color: When showing color, between 44A and 45A.

Petals/petaloids.—Quantity: About 6 to 9 petals and about 1 or 2 petaloids per flower. Petaloids variable in size and shape. Arrangement: Imbricate. Petal length: Upper petals: About 2.7 cm. Lower petals: About 2.5 cm. Petal width: Upper petals: About 2.5 cm. Lower petals: About 2.2 cm. Petal shape: Obovate. Petal/petaloid apex: Rounded. Petal/petaloid base: Attenuate. Petal/petaloid margin: Entire. Petal/petaloid texture: Smooth. Petal/petaloid color: Upper surface: Closest to 43A, venation on upper petals, 46B; iridescent. Lower surface: Closest to 40A.

Sepals.—Quantity: About 5, not imbricate. Length: About 9 mm. Width: About 4 mm. Shape: Ensiform. Apex: Narrowly acute. Margin: Entire. Texture: Densely pubescent. Color, upper and lower surfaces: 183B, towards apex, 143A.

Peduncle (umbel stem).—Length: About 12.7 cm. Angle: Erect. Strength: Strong. Texture: Pubescent. Color: Closest to 144A overlain with 183B.

Pedicel (individual flower stem).—Length: About 2.4 cm. Angle: Erect. Strength: Strong. Texture: Pubescent. Color: Closest to 183A.

Reproductive organs.—Androeceum: Anther quantity: About 5. Anther size: About 1 mm. Anther color: 186A. Pollen amount: Moderate. Pollen color: Closest to 25A. Gynoecium: Pistil quantity: One. Pistil length: About 9 mm. Stigma shape: Five-parted, star-shaped. Stigma color: Closest to 45B. Style length: About 3 mm. Style color: 45A.

Seed.—Not observed.

Disease resistance: Resistance to pathogens common to *Pelargonium* has not been observed.

It is claimed:

1. A new and distinct cultivar of *Geranium* plant named 'Belfanflai', as herein illustrated and described.

* * * * *

