



US00PP12508P2

(12) United States Plant Patent
Bessho et al.**(10) Patent No.: US PP12,508 P2****(45) Date of Patent: Apr. 2, 2002**

- (54) **CALIBRACHOA PLANT NAMED 'KAKEGAWA S8'**
- (75) Inventors: **Masao Bessho**, Kakegawa; **Toshimi Ohga**, Shizuoka-Prefecture, both of (JP)
- (73) Assignee: **Sakata Seed Corporation**, Yokohama (JP)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.
- (21) Appl. No.: **09/369,952**
- (22) Filed: **Aug. 4, 1999**
- (51) **Int. Cl.**⁷ **A01H 5/00**
- (52) **U.S. Cl.** **Plt./263**
- (58) **Field of Search** **Plt./263, 356**

- (56) **References Cited**
PUBLICATIONS
- UPOV-ROM GTITM Computer Database 2000/04 GTI JOUVE Retrieval Software.*
- * cited by examiner
- Primary Examiner*—Bruce R. Campell
Assistant Examiner—A. Para
- (74) *Attorney, Agent, or Firm*—James R. Cypher; Charles R. Cypher

(57) **ABSTRACT**

A new Calibrachoa plant which combines a decumbent and compact growth habit with large funnellform flowers, having mostly pink lobes and throat. The plant grows vigorously, branching only moderately. The petals are united and each has a fissure and a slight indent. Fully-opened flowers have a diameter between 3.1–3.6 cm. The throat and limbs of the petals are mostly light pink (R.H.S. 74C). The tube has partially white portions and partially yellow (R.H.S. 7B) portions. Typically, young flowers will close under low light and low temperature conditions such as late in the day and at night, giving them their generally funnellform appearance.

1 Drawing Sheet**1****BACKGROUND**

This invention relates to a new and distinct cultivar of Calibrachoa plant, hereinafter referred to by the name 'Kakegawa S8'. 'Kakegawa S8' is a hybrid. In 1996, the inventors Masao Bessho and Tosimi Ohga crossed a variety obtained from a commercial market in South America with a patented hybrid at a commercial nursery in Kakegawa, Japan. The F₁ progeny of this cross were found to have desirable characteristics, and the present invention was selected from this generation. The plant has been asexually reproduced through a number of generations since that time and its distinguishing characteristics have remain firmly fixed.

The Calibrachoa genus is a relatively new designation for a number of species that were formerly considered part of the Petunia genus.

The Petunia genus was originally established in 1803 by A. L. Jussieu, who described both *P. parviflora* Juss. and *P. nyctaginiflora* Juss. as type species. Using a non-horticultural system that selected the first mentioned species as the type species (lectotype), N. L. Britton and H. A. Brown declared *P. parviflora* Juss. as the type species for Petunia in 1913.

During the 1980's and 1990, H. J. Wijsman published a series of articles regarding the ancestry of *P. hybrida* (Hook.) Vilm., the Garden Petunia, and the inter-relationship of several species classified as Petunia. These studies discovered that *P. hybrida* (Hook.) Vilm. and its ancestral species, *P. nyctaginiflora* Juss. (= *P. axillaris* (Lam.) B.S.P.) and *P. violacea* Lindley (= *P. integrifolia* (Hook.) Schinz & Thellung.), possessed 14 pairs of chromosomes while several other species, including *P. parviflora* Juss., possessed 18 pairs of chromosomes. Since *P. parviflora* Juss. was the lectotype species for the Petunia genus, Wijsman and J. H. de Jong proposed transferring the 14 chromosome species to

2

the genus Stimoryne. Horticulturists opposed reclassifying the Garden Petunia and in 1986, Wijsman proposed the alternative of making *P. nyctaginiflora* Juss. the lectotype species for Petunia and transferring the 18 chromosome species to another genus. The I. N. G. Committee adopted this proposal. By 1990 Wijsman had transferred several species, including *P. parviflora* Juss. (= *C. parviflora*) to Calibrachoa, originally established by Llave and Lexarza in 1825. *Calibrachoa parviflora* (= *C. mexicana* la Lave & Lexarza) is now the type species for the genus Calibrachoa.

Classification of the current Petunia and Calibrachoa species is still in progress. New species are also being identified. At present, Calibrachoa can be distinguished from Petunia based on the higher chromosome number, chromosome morphology, plant branching habit and type of flower bud aestivation.

To summarize a few of the more easily identified distinguishing characteristics: whereas Petunia species have 2n=14 chromosomes, Calibrachoa species have 2n=18 chromosomes. Whereas Petunia species bear a flower peduncle and one new stem from a node, Calibrachoa bear a flower peduncle and up to three stems. Finally, Petunia species have a cochlear corolla bud. A single outermost petal covers the other four petals which are radially folded and terminally contorted. Calibrachoa flower buds are flat with all five petals linearly folded. The two lower petals are fused together and form a cover around the three other petals.

Further information on the transfer of certain species of Petunia to Calibrachoa can found in the following articles which are incorporated by reference herein. H. J. Wijsman, *On the Interrelationships of Certain Species of Petunia VI New names for the species of Calibrachoa formerly included into Petunia (Solanaceae)*, Acta. Bot. Neerl. 39(1), 101–102 (March, 1990). H. J. Wijsman and J. H. de Jong, *On the Interrelationships of Certain Species of Petunia IV Hybridization between P. Linearis and P. Calycina and Nomencla-*

torial Consequences in the Petunia Group, Acta. Bot. Neerl. 34, 337–349 (August, 1985). H. J. Wijsman, J. H. de Jong and T. M. Pedersen, *On the Interrelationships of Certain Species of Petunia III The Position of P. Linearis and P. Calycina*, Acta. Bot. Neerl. 32(4), 323–332, (August, 1983). Toshio Ando, Yoshiro Ueda, and Goro Hashimoto, *Historical Survey and Present Status of Systematics in the Genus Petunia Jussieu (Solanaceae)*, Technical Bulletin of Faculty of Horticulture, Chiba University, No. 45 (1992).

The new variety's female parent is a member of the Calibrachoa genus. No species determination of the female parent has been made. The female or seed parent is known to the inventors by its breeder code—C-10A. Parent C-10A was found in a commercial market in South America. It is characterized by white funnel-shaped flowers having a purple ring around the base of the throat and a decumbent growth habit.

The male parent was patented in the United States on May 6, 1997, by the same inventors. Its U.S. patent number is U.S. Plant Pat. No. 9,885. The inventors have named it 'Liricashower Blue'. Like the seed parent, its botanical classification has not been carried to the species level. In U.S. Plant Pat. No. 9,885, the genus for 'Liricashower Blue' was identified as Petunia. Since that time, as discussed above, the inventors have learned that the genus Petunia has been split by the I.N.G., and this particular variety, because of its chromosome number and bud aestivation is more accurately characterized as a member of the Calibrachoa genus. 'Liricashower Blue' is itself a hybrid plant, the result of a cross between plants obtained from commercial markets in South America.

The distinguishing characteristics of the parent 'Liricashower Blue' include bright funnel-shaped flowers, a decumbent growth habit and a large profusion of blooms. With regard to the color of the flowers, specifically, the throat and limbs of the petals are violet (R.H.S. 88A at maturity) and the tube is white to light green-yellow.

The initial cross-pollination of the parents, resulting in F₁ generation seed, was made in October of 1996. This F₁ seed was sown and yielded 20 plants. From these 20 plants, three plants were selected for their appealing flower color and decumbent growth habit. In August of 1997, the three selected plant lines were vegetatively propagated and tested for ease of reproducibility and stability of traits. The present variety was selected from the three for its vigorous growth, decumbent habit and light pink flower color. 'Kakegawa S8' has a pigmentation pattern similar to its pollen parent, 'Liricashower Blue'.

In December of 1997, cuttings of 'Kakegawa S8' were sent to a commercial nursery in Salinas, Calif. During the spring and summer of 1998, plants were grown under the direction and supervision of the inventors for evaluation of stability of the line's desired traits. Plants were grown in hanging pots at the commercial nursery in Salinas, Calif., and in cultivated fields at the commercial nursery in Kakegawa, Japan. Vegetative propagation in Japan was done in Kakegawa, Japan. The terminal 1.0 to 1.5 inches of an actively growing stem is excised. The basal half is stripped of leaves and dipped in a 1:19 dilution of Dip-N-Gro rooting solution (solution:water). The dipped portion of the stem is then inserted into moist peat-based soilless plant growing media. Soil trays with 1.0 inch diameter by 1.5 inch deep cells are used. The cuttings are kept in a warm greenhouse under a clear plastic tent with occasional misting from an automatic system. After about four weeks the cuttings have grown roots, bound together the soil as a root ball and can

be transplanted to pots. Two generations of successive propagation (December 1997 and December 1998) were performed between the final selection and collecting data for the application. Vegetative propagation in Salinas, Calif. was performed in greenhouses at the Sakata Seed America Plug & Propagation facility. It takes approximately four weeks to produce a root cutting. 'Kakegawa S8' was determined by the inventors to have its characteristics, as herein described, firmly fixed.

'Kakegawa S8' is a new variety of Calibrachoa plant having a decumbent and compact growth habit. The plant grows vigorously, branches only moderately, and makes an excellent hanging basket. Its flowers are funnel-shaped. The petals are united and each has a fissure and a slight indent. The flowers are axillary and solitary. Fully-opened flowers have a diameter between 3.1–3.5 cm. The throat and limbs of the petals are light pink (R.H.S. 74C, red-purple group). The tube is alternately white or yellow (R.H.S. 7B, yellow group). The plant is very resistant to rain, heat and drought. The plant grows and flowers best under low soil pH conditions (pH 5–6). Typically, young flowers will close under low light and low temperature conditions such as late in the day and at night, giving them their generally funnelform appearance.

The new variety is distinguished from other Calibrachoa plants by the combination of its characteristics. It combines a unique flower color with large flowers, a decumbent, vigorous growth habit with moderate branching, and a tendency for its flowers to close in the evening.

The closest commercial cultivar of which the inventors are aware is the petunia-like plant named 'Million Bells Trailing Pink'. The distinguishing characteristics which differentiate 'Kakegawa S8' from 'Million Bells Trailing Pink' are:

	'Kakegawa S8'	'Million Bells Trailing Pink'
Branching	Moderate branching	Abundant branching
Growth Habit	Decumbent	Mounding
Flower Color	Throat and limbs of the petals are light pink (R.H.S. 74C).	Bluish Pink
Tube Color	Partially white and partially yellow (R.H.S. 7B).	Pale yellow with purple veins.
Flower Diameter	31–35 mm	Smaller
Blooming Habit	Closes in the evening	Stays open all day

BRIEF DESCRIPTION OF DRAWINGS

The accompanying drawings serve by color photographic means to illustrate the new plant variety, Kakegawa S8. The colors are represented as true as possible using conventional photographic procedures.

FIG. 1 is a close-up view of multiple blooms illustrating the color of the flowers and the plant's decumbent, compact growth habit; and large profusion of blooms.

FIG. 2 is a view of the new cultivar after growing for several weeks in a hanging basket.

DETAILED DESCRIPTION OF THE NEW PLANT

The following description is based on observations and measurements of plants grown in pots in Salinas, Calif. The

plants were grown indoors. The plants received fertilizer water (constant liquid feeding) with an 18-8-8 feeding level. The plants had been growing for about 5 months since they had been transferred to their pot as rooted cuttings. The average growing temperature is 65 degrees Fahrenheit at night, and 75 degrees Fahrenheit in the day. Color designations were made according to The Royal Horticultural Society Colour Chart published by The Royal Horticultural Society of London, England.

Origin: Seedling

Parentage:

Female parent.—Unpatented and unnamed commercial market plant from South America, breeder code C-10A.

Male parent.—'Liricashower Blue' U.S. Plant Pat. No. 9,885.

Classification:

Family.—Solanaceae.

Genus.—Calibrachoa.

Botanical.—Unknown member of the Calibrachoa genus.

Commercial.—'Kakegawa S8'.

Plant:

Growth habit.—Decumbent.

Plant height.—8 cm.

Spreading area of plant.—15 cm. in all directions from the edge of the pot result from plants grown in a 41 cm. diameter hanging basket.

Type.—Perennial.

Disease resistance.—The plant is susceptible to Botrytis, powdery mildew, various stem and root rots, and certain viruses, like Tobacco Mosaic Virus and Impatiens Necrotic Spot Virus. Plants can be infested with aphids, leafminer, whitefly and various Lepitopdera.

Blooming.—Mature plants growing in a 41 cm. Diameter hanging basket can have in excess of 700 open flowers at any given time. The inflorescence is solitary.

Stem:

Thickness.—Main stem 2.0–2.5 mm, lateral stems 1.0–1.5 mm.

Color.—Yellowish green (R.H.S. 144B).

Pubescence.—Stem pubescence is moderate in density and short. The pubescence is colorless and bulbous at the tip.

Branching.—Moderate. One flower peduncle and three break stems at each node.

Length of internode.—20–25 mm.

Leaf:

Form.—Alternate, elliptical with mucronate tips and entire margins.

Length (average).—22 mm.

Width (average).—7 mm.

Thickness.—0.5–0.8 mm.

Color.—The adaxial surface color of the leaves is R.H.S. 137B. The abaxial surface color of the leaves is R.H.S. 138B.

Pubescence.—Pubescence is short, colorless and bulbous at the tip.

Flower:

Form.—Flowers are axillary and solitary. Flowers are bisexual. Calyx, corolla and androecium are all 5-parted. Calyx is persistent. Corolla is funnelform with 5 united petals. As flowers age the throat and limbs of the flower flare more abruptly. Petals have broad limbs, each petal is fissured with a slight indent.

Diameter of corolla.—31–36 mm when fully open.

Tube length.—14 mm.

Throat diameter.—7 mm.

Color of petals.—Young flowers: Light Pink (R.H.S. 75C).

Color of petals in full bloom.—When fully mature: the adaxial surface color of the limbs is R.H.S. N74A, the adaxial surface color of the throat is R.H.S. 79C, and the adaxial surface color of the tube is R.H.S. 7B at base fading to 1D at the throat; the abaxial surface color of the limbs and throat is R.H.S. 77C, and the abaxial surface color of the tube is R.H.S. 8D; and the color of the flower petal vein is R.H.S. 79C.

Reproductive organs.—Androecium consisting of five stamens, 2 with long filaments bending upward over the pistil and 3 with short filaments, anthers are yellow and filaments are white. The color of the stamens is 160A. One pistil. Ovary is superior. Stigma is oval with a septation running through the middle on the long axis. The color of the ovary, pistil, style and a stigma is R.H.S. 140C.

Blooming habit.—Young flowers close in the evening and under low-light conditions.

Fragrance.—The flowers of this plant do not possess any fragrance.

Sepals.—There are five sepals on each flower; the adaxial surface color of the sepal is R.H.S. 137B and the abaxial surface color of the sepal is 137B. Sepal pubescence is moderate in density, short, clear and bulbous at the tip. The sepals are fused at their base. Where they are not fused they taper in a lanceolate manner to an acute pointed tip.

Seeds/fruit: No seeds or fruit are produced.

Blooming season.—Plants flower April through October in Salinas, Calif.

Bud:

Aestivation.—Buds are flat with all five petals linearly folded and the two lower petals forming a cover around the three other petals. All the petals are fused together.

Chromosome number: 2n=18.

We claim:

1. A new and distinct variety of Calibrachoa plant as illustrated and described.

* * * * *



FIGURE 1




FIGURE 2