



US00PP12817P2

(12) **United States Plant Patent**
Gilford et al.

(10) **Patent No.:** **US PP12,817 P2**

(45) **Date of Patent:** **Jul. 30, 2002**

(54) **STRAWBERRY PLANT NAMED**
'MARATHON'

(75) Inventors: **Kristie L. Gilford**, Dover, FL (US);
Bruce D. Mowrey, Watsonville, CA
(US); **Jorge Campos Garcia**, Plant
City, FL (US); **Thomas M. Sjulín**,
Aromas; **Larry T. Kodama**, Freedom,
both of CA (US)

(73) Assignee: **Driscoll Strawberry Associates, Inc.**,
Watsonville, CA (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **09/396,213**

(22) Filed: **Sep. 15, 1999**

(30) **Foreign Application Priority Data**

Mar. 30, 1999 (CA) 99-1603

(51) **Int. Cl.**⁷ **A01H 5/00**

(52) **U.S. Cl.** **Plt./209**

(58) **Field of Search** **Plt./208, 209**

Primary Examiner—Bruce R. Campell

Assistant Examiner—Michelle Kizilkaya

(74) *Attorney, Agent, or Firm*—Pennie & Edmonds LLP

(57) **ABSTRACT**

This invention relates to a new and distinct variety of
strawberry named 'Marathon', botanically identified as
Fragaria×ananassa. The closest known varieties are 'Key
Largo' (U.S. Plant Pat. No. 8,649), 'Captiva' (U.S. Plant Pat.
No. 11,277, allowed) and 'Mirador' (U.S. Plant Pat. No.
11,279, allowed). The new variety is a partially everbearing
fruit bearing variety. The variety is distinguished from 'Key
Largo', 'Captiva' and 'Mirador' by its strong plant vigor,
concave to flat cross section of the leaf, medium leaf
blistering, same size diameter of inner to outer calyx,
broader than longer fruit ratio length/width, large fruit size
and cordate fruit shape.

2 Drawing Sheets

1

1. BACKGROUND OF THE INVENTION

The new variety originated as a result of a controlled cross
between the strawberry plants 'Key Largo' (U.S. Plant Pat.
No. 8,649) and 'L3' (unpatented proprietary) in an ongoing
breeding program, and was discovered as a seedling in a
controlled breeding plot. The original seedling of the new
cultivar was asexually propagated by stolons in Shasta
County, Calif. Propagules were transplanted to a controlled
breeding plot in Valrico, Hillsborough County, Fla. where
the new cultivar first fruited in December 1994 and was
selected for further evaluation. 'Marathon' was subsequently
asexually propagated and underwent further testing in the
area of Valrico, Fla. from 1995–1999. This propagation and
testing has demonstrated that the combination of traits
disclosed herein which characterize the new variety are fixed
and retained true to type through successive generations as
asexual reproduction.

2. SUMMARY OF THE INVENTION

The present invention relates to a new and distinct variety
of strawberry named 'Marathon'. The variety is botanically
identified as *Fragaria×ananassa*. The new variety is distin-
guished from other varieties by a number of characteristics
as set forth in Tables 1, 2 and 3.

3. COMPARISON TO CLOSEST VARIETIES

The varieties which we believe to be closest to 'Marathon'
from those known to us are 'Key Largo' (U.S. Plant Pat. No.
8,649), 'Captiva' (U.S. Plant Pat. No. 11,277, allowed) and
'Mirador' (U.S. Plant Pat. No. 11,279, allowed). There are
several characteristics of 'Marathon' that are different from,
or not possessed by 'Key Largo', 'Captiva' or 'Mirador'.
Table 1 provides information on the plant and fruit charac-
teristics of the new variety 'Marathon' compared with char-
acteristics of 'Key Largo', 'Captiva' and 'Mirador'. Table 2

2

provides additional information of the plant and fruit char-
acteristics of the new variety 'Marathon' compared with
characteristics of the varieties 'Captiva', 'Key Largo' and
'Mirador'. Table 3 provides isozyme characteristics of the
new variety as compared to the varieties 'Key Largo',
'Captiva' and 'Mirador'.

4. BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying photographs show typical specimens
of the new variety, including fruit, foliage and flowers, in
color as nearly true as it is reasonably possible to make in
color illustrations of these characteristics.

FIG. 1 shows the fruit in longitudinal section, illustrating
the typical flesh coloration.

FIG. 2 shows the upper surface of two typical folioles of
the new variety.

FIG. 3 shows the flower and reproductive organs of the
new variety, as well as the size and position of the petals and
sepals and the underside of the calyx.

FIG. 4 shows typical whole fruit.

5. DESCRIPTION OF THE NEW VARIETY

The following detailed description of the new variety is
based upon observations taken of plants and fruit grown in
Valrico, Fla., U.S.A. Observations of 'Marathon', 'Key
Largo', 'Captiva' and 'Mirador' were taken in side by side
comparisons in 1999. This description is in accordance with
UPOV terminology. Color designations, color descriptions,
and other phenotypical descriptions may deviate from the
stated values and descriptions depending upon variation in
environmental, seasonal, climatic and cultural conditions.
Colors are described and the closest matching color ratings
are provided using The Royal Horticultural Society (R.H.S.)
Colour Chart.

5.1 Propagation

The new variety is principally propagated by way of stolons. Although propagation by stolons is presently preferred, other known methods of propagating strawberry plants may be employed.

5.2 Characteristics of the New Variety

Information the new variety is presented in Tables 1, 2 and 3. In the tables, the flowers described are secondary flowers except where indicated. Petal color of 'Marathon' is white which is not referenced in The Royal Horticultural Society Colour Chart. The fruit described is the secondary fruit on one year old plants. Fruit and flower measurements are an average of both primary and secondary fruit and flowers.

Plants and Foliage

Plants of 'Marathon' are of strong vigor while those of 'Key Largo', 'Captiva' and 'Mirador' are of medium vigor. The average height of 'Marathon' is 21.0 cm, the average diameter is 37.9 cm, and the average number of crowns/plant is 3.8. The upper leaf surface of 'Marathon' is dark green with an R.H.S. color rating of 137B, whereas the color of three comparison varieties have an R.H.S. color rating of 137A. The underside of the leaves of 'Marathon' has a R.H.S. color rating of 138B, while 'Key Largo', 'Captiva' or 'Mirador' have an R.H.S. color rating of 139C. The shape of leaves of 'Marathon' in cross section is concave while those of 'Key Largo' and 'Captiva' and 'Mirador' are slightly concave. The leaves of 'Marathon' exhibit medium interveinal leaf-blistering. 'Marathon' has three leaflets and the leaves are of medium glossiness. The shape of the base of the terminal leaflet is slightly oblique and the teeth of the terminal leaflet are obtuse. The average number of teeth per terminal leaflet is 25.9. The average length and width of the terminal leaflet are 10.6 cm and 11.3 cm, respectively.

The petiole of 'Marathon' has medium pubescence. The pose of the petiole hairs is outwards compared to that of 'Captiva' and 'Key Largo' which is upwards. The average petiole length is 10.4 cm. The petiole color has an R.H.S. color rating of 141D.

The bract frequency is 30% and is typically single compared to that of 'Captiva' and 'Mirador' for which the bracts occur with a 60% frequency. The stipule length and width are 3.7 cm and 2.3 cm, respectively.

The stipule pubescence is medium. Marathon has a medium number to many stolons that are medium to thick in thickness with medium to strong anthocyanin coloration and sparse pubescence.

Flower and Fruit Characteristics

The position of the 'Marathon' inflorescence is beneath the foliage compared to that of 'Mirador' which is above. The flowers are large in size. The average petal length and width are 1.3 cm and 1.4 cm, respectively. The average flower diameter is 4.1 cm. The average calyx diameter is 3.7 cm.

The diameter of 'Marathon's' calyx is larger than the corolla while that of 'Mirador' is the same size. On secondary flowers, the diameter of the inner calyx is the same as the outer calyx. On secondary flowers with 5 to 6 petals, the petals are overlapping. The petal length-to-width ratio on secondary flowers is as broad as they are long compared to that of 'Mirador' which are broader than long.

The fruiting trusses are long in length with an average length of 18.5 cm from the crown to the base of the terminal flower or fruit. The attitude at first picking is prostrate compared to the semi-erect 'Key Largo' and erect 'Mirador'.

Observations of 'Marathon' fruit were taken of secondary fruit on one year old plants. The fruits are predominantly cordate in shape. The fruit ratio of length to maximum width is as broad as long. The fruit are large in size. The average length and width of the fruit are 4.3 cm and 4.3 cm, respectively. There is a slight difference in fruit shape between the primary and secondary fruits. The band without achenes is narrow. The fruit surface has a weak unevenness.

The fruit skin color is an even orange-red with an R.H.S. color rating of 42A and strong glossiness.

The insertion of the achenes is below the surface of the fruit as compared to that of 'Captiva' and 'Mirador' which are level with the surface.

The calyx is set above the fruit and the pose of the calyx segments is spreading. The calyx diameter is larger than the fruit diameter. The adherence of the calyx to the fruit is medium.

The fruit flesh is medium firm when fully ripe. The color of the fruit flesh is medium red with an R.H.S. color rating of 42C. The fruit flesh color is slightly uneven, and the fruit flesh color distribution is marginal and central. The fruit has a medium sized hollow center.

The fruit has a medium sweetness with medium acidity and fine textures.

The time of flowering when 50% of the plants are at first flower is early. Initial flowering occurs in late-October to early-November.

The harvest maturity when 50% of the plants possess ripe fruit is early. Season of harvest is from late-November through early-April. 'Marathon' is a partially everbearing variety.

The average number of stamens is 28.

Anthocyanin coloration is red to purple.

The average number of pedicels/peduncle is 2.3. Average diameter of peduncle is 4.5 mm. Average length of peduncle is 5.8 cm. Average pedicel diameter is 2.7 mm. Average length of pedicels is 11.8 cm. Average petiole diameter is 3.8 mm.

Insertion of achenes is below surface. Achene color is yellow to red (12A to 46A).

Average sepal number is 10.9 and sepal length is 1.5 cm.

Percent brix is 8.3. Amount of pollen produced is abundant and fertile.

TABLE 1

	DETAILED COMPARISON OF 'MARATHON', KEY LARGO, 'CAPTIVA' AND 'MIRADOR'			
	MARATHON	CAPTIVA	KEY LARGO	MIRADOR
<u>Plant Characteristics</u>				
Avg. Height of Plant	21.0 cm	13.0 cm	18.3 cm	20.0 cm
Avg. Spread of Plant	37.9 cm	33.1 cm	39.2 cm	35.0 cm
Avg. Number of Crowns	3.8	4.4	4.0	3.0

TABLE 1-continued

DETAILED COMPARISON OF 'MARATHON', KEY LARGO, 'CAPTIVA' AND 'MIRADOR'				
	MARATHON	CAPTIVA	KEY LARGO	MIRADOR
<u>Leaf Characteristics</u>				
Terminal Leaflet Width	11.3 cm	7.3 cm	8.9 cm	9.0 cm
Terminal Leaflet Length	10.6 cm	7.2 cm	8.5 cm	8.3 cm
Terminal Leaflet Length/Width Ratio	0.94	0.99	0.96	0.92
Number of Teeth/Terminal Leaflet	25.9	25.4	23.8	20.0
Petiole Length	10.4	8.1	10.4	12.8
Bract Frequency	30% Typically Single	60% Typically Single	30% Typically in pairs	60% Typically Single
Stipule Length	3.7	3.6	3.7	3.6
Stipule Width	2.3	1.8	1.9	2.2
<u>Flower Characteristics</u>				
Petal Width	1.4 cm	1.4 cm	1.5 cm	1.6 cm
Petal Length	1.3 cm	1.4 cm	1.5 cm	1.5 cm
Petal Length/Width Ratio	0.93	1.00	1.00	0.94
Flower Diameter	4.1 cm	3.6 cm	3.9 cm	4.0 cm
Calyx Diameter	3.7 cm	4.3 cm	5.1 cm	4.0 cm
<u>Fruit Characteristics</u>				
Fruit Width	4.3 cm	3.6 cm	3.5 cm	3.5 cm
Fruit Length	4.3 cm	4.1 cm	4.2 cm	3.5 cm
Fruit Length/Width Ratio	1.00	1.14	1.20	1.00
Average Berry Weight (g)	26.5	17.8	22.4	16.3
Fruit Skin Color	Orange Red 42A	Orange Red 45A	Dark Red 46B	Red 45B
Fruit Flesh Color	Light Red 42C	Light Red 49C	Light Red 44C	Pale Rose 43B
Total Yield (g/plant)	638	369	277	551

TABLE 2

CHARACTERISTICS OF 'MARATHON', 'KEY LARGO', 'CAPTIVA', AND 'MIRADOR'				
	MARATHON	CAPTIVA	KEY LARGO	MIRADOR
<u>Plant</u>				
Habit	Flat globuse	Flat Globuse	Globuse	Globuse
Density	Medium	Medium	Medium	Open
Vigor	Strong	Medium	Medium	Medium
<u>Leaf</u>				
Color of upper side	Dark Green 137B	Dark Green 137A	Dark Green 137A	Dark Green 137A

TABLE 2-continued

CHARACTERISTICS OF 'MARATHON', 'KEY LARGO', 'CAPTIVA', AND 'MIRADOR'				
	MARATHON	CAPTIVA	KEY LARGO	MIRADOR
<u>Color or under side</u>				
Color or under side	Light Green 138B	Light Green 139C	Light Green 139C	Light Green 139C
<u>Shape in cross section</u>				
Shape in cross section	Concave	Slightly Concave	Slightly Concave	Slightly Concave
<u>Interveinal blistering</u>				
Interveinal blistering	Medium	Weak	Weak	Weak
<u>Glossiness</u>				
Glossiness	Medium	Medium	Medium	Medium
<u>Number of leaflets</u>				
Number of leaflets	3	3	3	3
<u>Terminal leaflet margin profile</u>				
Terminal leaflet margin profile	Revolute	Revolute	Revolute	Revolute
<u>Terminal leaflet shape of base</u>				
Terminal leaflet shape of base	Slightly oblique	Obtuse	Slightly oblique	Slightly oblique
<u>Terminal leaflet shape of teeth</u>				
Terminal leaflet shape of teeth	Obtuse	Obtuse	Rounded	Rounded
<u>Terminal leaflet average number of teeth</u>				
Terminal leaflet average number of teeth	25.9	25.4	23.8	20.0
<u>Petiole pubescence</u>				
Petiole pubescence	Medium	Medium	Medium	Medium
<u>Petiole pose of hairs</u>				
Petiole pose of hairs	Outwards	Upwards	Upwards	Outwards
<u>Petiole Color</u>				
Petiole Color	141D	141C	141C	141C
<u>Stipule Pubescence</u>				
Stipule Pubescence	Medium	Medium	Medium	Medium
<u>Bracts frequency on petioles</u>				
Bracts frequency on petioles	30% Typically Single	60% Typically Single	30% Typically in pairs	60% Typically Single
<u>Stolon</u>				
Number	Medium to many	Medium to many	—	Many
<u>Anthocyanin coloration</u>				
Anthocyanin coloration	Medium to strong	Strong	—	Medium to strong
<u>Thickness</u>				
Thickness	Medium to thick	Medium to thick	—	Thin to medium
<u>Pubescence</u>				
Pubescence	Sparse	Sparse to medium	—	Sparse to medium
<u>Inflorescence</u>				
Position relative to foliage	Beneath	Beneath	Beneath	Above
<u>Flower size</u>				
Flower size	Large	Medium	Large	Small
<u>Diameter of calyx relative to corolla</u>				
Diameter of calyx relative to corolla	Larger	Larger	Much larger	Same size
<u>Diameter of inner calyx relative to outer</u>				
Diameter of inner calyx relative to outer	Same size	Smaller	Smaller	Smaller
<u>Spacing of petals</u>				
Spacing of petals	Overlapping	Overlapping	Overlapping	Overlapping
<u>Fruiting Truss</u>				
Attitude at first picking	Prostrate	Prostate	Semi-erect	Erect
Length	18.5 cm	9.4 cm	16.6 cm	17.3 cm
<u>Fruit</u>				
Predominant shape	Cordate	Conical	Conical	Conical
<u>Difference in shapes between primary and secondary fruits</u>				
Difference in shapes between primary and secondary fruits	Slight	Slight	Slight	Slight

TABLE 2-continued

CHARACTERISTICS OF 'MARATHON', 'KEY LARGO', 'CAPTIVA', AND 'MIRADOR'				
	MAR- ATHON	CAPTIVA	KEY LARGO	MIRADOR
Band without achenes	Narrow	Narrow	Narrow	Narrow
Unevenness of surface	Weak	Weak	Weak	Weak
Skin color	Orange Red 42A	Orange red 45A	Dark red 46B	Red 45B
Evenness of color	Even	Slightly uneven	Even	slightly uneven
Glossiness	Strong	Medium	Strong	Medium
Insertion of achenes	Below surface	Level with surface	Below surface	Level with surface
Insertion of calyx	Above fruit	Above fruit	Above fruit	Above fruit
Pose of the calyx segments	Spreading	Spreading	Reflexed	Spreading
Size of calyx in relation to fruit	Larger	Larger	Larger	Larger
Firmness of flesh	Medium	Firm	Firm	Firm
Color of flesh	Light Red 42C	Light Red 49C	Light Red 44B	Pale Rose 43B
Evenness of flesh color	Slightly uneven	Slightly uneven	Slightly uneven	Slightly uneven
Distribution of flesh color	Marginal and central	Marginal and central	Marginal and central	Only marginal
Hollow center size	Medium	Small	Medium	Medium
Sweetness	Medium	Medium	Medium	Medium
Texture when tasted	Fine	Fine	Medium	Fine
Acidity	Medium	Medium	Medium	Medium

5.3 Insect and Disease Resistance and Susceptibility

The 'Marathon' variety is moderately susceptible to *Xanthomonas fragariae* and Botrytis fruit rot. The variety is susceptible to Powdery Mildew and Verticillium wilt. The

'Marathon' variety is also susceptible to injury by the two-spotted spider mite (*Tetranychus urticae*), lygus bug (*Lygus hesperus*) and Aphis spp. (Aphids). 'Marathon' is moderately resistant to the Strawberry Mottle Virus.

5.4 ISOZYME ANALYSIS

In addition to the morphological description above, the new cultivar 'Marathon' has been analyzed to obtain an indication of its genetic makeup to provide further means for identifying the new variety and distinguishing it from some other somewhat similar and/or related strawberry varieties. Specifically, leaf samples of 'Marathon', 'Key Largo', 'Captiva' and 'Mirador' were analyzed by electrophoresis for isozyme patterns of the enzymes phosphoglucosomerase (PGI), leucine aminopeptidase (LAP) and phosphoglucomutase (PGM). See *J. Amer. Soc. Hort. Sci.* 106:684-687. Isozyme characterization of the four varieties is presented in Table 3, with the letters representing the banding patterns for each enzyme as designated in the above-identified article.

TABLE 3

ISOZYME ANALYSIS FOR 'MARATHON', 'CAPTIVA', 'KEY LARGO' AND 'MIRADOR'				
Locus	MARATHON	CAPTIVA	KEY LARGO	MIRADOR
PGI	A1	A2	A1	A3
LAP	B3	B3	B3	B3*
PGM	C4	C2	C4	C4

*With a slower migration band staining more weakly than typical B3 pattern.

What is claimed is:

1. A new and distinct variety of strawberry plant, substantially as shown and described.

* * * * *



FIG. 1




FIG. 2




FIG. 3



FIG. 4