



US00PP13180P2

**(12) United States Plant Patent
Strope****(10) Patent No.: US PP13,180 P2****(45) Date of Patent: Nov. 5, 2002****(54) NEW GUINEA IMPATIENS PLANT NAMED
'BALCEBSAFO'****(75) Inventor: Kerry M. Strope, Pismo Beach, CA
(US)****(73) Assignee: Ball FloraPlant, a division of Ball
Horticultural corporation, West
Chicago, IL (US)****(*) Notice:** Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.**(21) Appl. No.: 09/772,824****(22) Filed: Jan. 30, 2001****(51) Int. Cl.⁷ A01H 5/00****(52) U.S. Cl. Plt./318****(58) Field of Search Plt./318***Primary Examiner*—Bruce R. Campell*Assistant Examiner*—June Hwu**(74) Attorney, Agent, or Firm**—Wood, Phillips, Katz, Clark
& Mortimer**(57) ABSTRACT**A new and distinct cultivar of New Guinea impatiens plant
named Balcebsafo, characterized by its large light salmon
flowers, dark green foliage, compact upright and mounded
habit and excellent basal branching.**1 Drawing Sheet****1****BOTANICAL CLASSIFICATION***Impatiens hawkeri* cultivar 'Balcebsafo'.**BACKGROUND OF INVENTION**The present invention comprises a new and distinct New
Guinea impatiens plant, botanically known as *Impatiens*
hawkeri, and hereinafter referred to by the cultivar name
'Balcebsafo'.The new cultivar was developed by the inventor through
a controlled breeding program during 1997 at Arroyo
Grande, Calif. The objective of the breeding program was to
develop cultivars with numerous large flowers, compact to
moderate growth habit and excellent basal branching.The female (seed) parent of 'Balcebsafo' was the propri-
etary *Impatiens hawkeri* cultivar designated '1199' (non-
patented), characterized by its compact habit, lavender flow-
ers and dark green foliage. The male (pollen) parent of
'Balcebsafo' was the *Impatiens hawkeri* cultivar 'Toga'
(U.S. Plant Pat. No. 10,304) characterized by its compact
habit, lavender flowers and dark green foliage. 'Balcebsafo'
was discovered and selected in December 1997 by the
inventor as a flowering plant within the progeny of the stated
cross.Asexual reproduction of the new cultivar by terminal
cuttings has demonstrated that the characteristics of the new
cultivar as herein described are firmly fixed and are retained
through successive generations of such asexual propagation.
The new cultivar was asexually reproduced in Arroyo
Grande, Calif. and West Chicago, Ill.**SUMMARY OF INVENTION**

It was found that the cultivar of the present invention:

- (a) exhibits large round light salmon flowers,
- (b) forms dark green foliage,
- (c) exhibits a good basal branching character, and
- (d) exhibits a medium upright mounded growth habit.

The new cultivar of the present invention can be com-
pared to 'Celebration Deep Coral' (U.S. Plant Pat. No.
9,214). In side by side comparisons, plants of the new
cultivar differ from 'Celebration Deep Coral' by the follow-**2**ing characteristics: it is found that the new cultivar is more
compact, has longer internodes, longer petioles and lighter
flower color. The new cultivar can also be compared to
'Celebrette Peach' (U.S. Plant Pat. No. 10,108). In side by
side comparisons it is found that the new cultivar is more
compact, has shorter peduncles, larger, pinker flowers,
longer internodes and different foliage coloration. A detailed
comparison can be found below in Table A.**TABLE A**

CHARACTERISITIC	'BALCEBSAFO'	CELEBRETTE PEACH
PLANT HEIGHT	18.4 cm	21 cm
STEM COLOR	185A	53B
BRANCH LENGTH	16.2 cm	18.3 cm
INTERNODE LENGTH	6.9 cm	2.6 cm
MATURE LEAF LENGTH × WIDTH	8.2 cm × 3.1 cm	9 cm × 3.4 cm
MATURE LEAF COLOR-UPPER SURFACE	139A WITH VENATION OF LIGHT GREEN	136A WITH VENATION OF 63A
MATURE LEAF COLOR-LOWER SURFACES	191 A WITH VENATION OF 184B	183C WITH VENATION OF 185B
PETIOLE LENGTH	1.2 cm	1.5 cm
FULLY OPENED PETAL COLOR- UPPER SURFACE	48B GRADUALLY FADING THROUGH 48C, 48D AND 39D TO BASE OF 51B	SUPERIOR PETAL: 41A AT EDGE; 63B AT BASE. OTHER PETALS: 41A AT EDGE, FADING THROUGH 63D TO 63A AT BASE
PEDUNCLE LENGTH	4.2 cm	5.4 cm
PEDUNCLE COLOR	53A	144D
FLOWER DIAMETER	6.4 cm	5.5 cm
SPUR COLOR	53A	53C
SPUR LENGTH	5.6 cm	4.3 cm
POLLEN COLOR	19B	11C

CHARACTERISITIC**CELEBRATION DEEP
CORAL**

PLANT HEIGHT	27.1 cm
STEM COLOR	187D
BRANCH LENGTH	23.3 cm
INTERNODE LENGTH	5.8 cm
MATURE LEAF LENGTH × WIDTH	8.4 cm × 3.5 cm

TABLE A-continued

MATURE LEAF	187 A WITH MIDVEIN OF 45B
COLOR-UPPER SURFACE MATURE LEAF	187C
COLOR-LOWER SURFACES	
PETIOLE LENGTH	.9 cm
FULLY OPENED	50A
PETAL COLOR- UPPER SURFACE	
PEDUNCLE LENGTH	4.9 cm
PEDUNCLE COLOR	53D
FLOWER DIAMETER	6.8 cm
SPUR COLOR	53C
SPUR LENGTH	5.2 cm
POLLEN COLOR	19B

BRIEF DESCRIPTION OF PHOTOGRAPH

The accompanying photographs show as nearly true as it is reasonably possible to make the same in color illustrations of this type, typical flower and foliage characteristics of the new cultivar. The plants were grown in a greenhouse at West Chicago, Ill., U.S.A.

DETAILED DESCRIPTION

The 'Balcebsafo' cultivar has not been observed under all possible environmental conditions to date. Accordingly, it is possible that the phenotype may vary somewhat with variations in the environment, such as temperature, light intensity, and day length.

The chart used in the identification of colors described herein is the R.H.S. Colour Chart of The Royal Horticultural Society, London, England. The color values were determined on Aug. 30, 2000 between 1:00 and 3:00 p.m. under natural daylight conditions. The plants were ten (10) weeks old and were produced from cuttings taken from stock plants and were grown in 10 cm pots under greenhouse conditions comparable to those used in commercial practice while utilizing a soilless growth medium and maintaining temperatures of approximately 72° F. during the day and approximately 65° F. during the night.

Parentage: Proprietary seedling '1199'×'Toga'.

Propagation:

Type cutting.—Terminal tip.

Time to initiate roots.—Approximately 7–14 days with the shorter times generally being experienced in the summer and the longer times in the winter.

Time to develop roots.—Approximately 21 days.

Rooting habit.—Fibrous, branching.

Plant description:

Habit of growth.—Moderately vigorous. A mature plant, 8 to 10 weeks after planting of a rooted cutting, measures approximately 18.4 cm in height and approximately 33.4 cm in width.

Form.—Upright and mounded.

Foliage.—Leaves are simple, lanceolate, acuminate apex, attenuate base, serrate and ciliate margin. Texture: smooth. Size of mature foliage: approximately

8.2 cm in length, approximately 3.1 cm in width. Color of mature foliage: upper surface is 139A with venation of 144D; lower surface is 191A with venation of 184B. Petiole: Length: approximately 1.2 cm. Color: 184C.

Stem.—Color: 185A. Internode length: approximately 6.9 cm.

Flower description:

Flowering habit.—Freely flowering.

Natural flowering season.—Year round in greenhouse environment. Flowering is continuous from spring until fall in the garden.

Bud shape, size, color.—Shape: Ovate. Length: 1 cm. Color: 60B.

Flower longevity.—Approximately seven days.

Flowers borne.—Above foliage arising from leaf axils.

Flower color.—Upper surface of petals is 48B at edges gradually fading through 48C, 48D and 39D to bases of 51B; lower surface is between 40A and 50C with midvein of 51B.

Quantity of flowers.—Approximately 2 flowers and 5 buds per stem.

Petals.—Number: Five. Shape: Obovate, superior petal has broad base, other four petals have narrow base. Approximate size: Superior petal width, 4.1 cm; superior petal length, 2.8 cm. Lateral petal width, 2.7 cm; lateral petal length, 2.8 cm; lower petals width, 3.1 cm, lower petal length, 3.4 cm.

Flower diameter.—Approximately 6.4 cm.

Spur.—Origin: from inferior sepal; Length: Approximately 5.6 cm. Color: 53A. Shape: Very Slightly Curved.

Peduncle.—Length: Approximately 4.2 cm. Color: 53A.

Flower buds.—Length: approximately 2 cm, width: approximately 1.2 cm when first showing color.

Reproductive organs.—Androecium: five stamens, anthers are fused together forming one organ that surrounds the pistil. Generally, the anthers shed pollen prior to the stigma becoming receptive, pollen color is 19B. Gynoecium: the stigma color is 154D the style color is 144A and the ovary color is 144A.

Anther shape, size, color.—Anther Shape: Hooded, Anther size: 2.2 mm. Anther color: 20B.

Pistil length, stigma color.—Pistil Length: 3 mm. Stigma color: 70B.

Seed production.—Seed production has not been observed.

Disease resistance: Resistance to pathogens has not been observed.

I claim:

1. A new and distinct cultivar of New Guinea Impatiens plant named 'Balcebsafo' substantially as herein shown and described, which:

- exhibits attractive, large round light salmon flowers,
- forms dark green foliage,
- exhibits a good basal branching character, and exhibits a medium upright mounded growth habit.

* * * * *

