



US00PP13229P2

(12) **United States Plant Patent**
Hammett

(10) **Patent No.:** **US PP13,229 P2**

(45) **Date of Patent:** **Nov. 12, 2002**

(54) **NEMESIA PLANT NAMED ‘NEMLI’**

(75) **Inventor:** **Keith Richard William Hammett,**
Auckland (NZ)

(73) **Assignee:** **Seaview Nurseries Ltd.,** Manurewa
(NZ)

(*) **Notice:** Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) **Appl. No.:** **09/968,327**

(22) **Filed:** **Sep. 30, 2001**

(51) **Int. Cl.⁷** **A01H 5/00**

(52) **U.S. Cl.** **Plt./263**

(58) **Field of Search** **Plt./263**

(56) **References Cited**

U.S. PATENT DOCUMENTS

PP12,014 P2 * 7/2001 Hubbard Plt./263

* cited by examiner

Primary Examiner—Kent L. Bell

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Nemesia* plant named ‘Nemli’,
characterized by its upright, outwardly spreading and com-
pact plant habit; freely branching habit; small green leaves;
and numerous light purple-colored flowers.

1 Drawing Sheet

1

**BOTANICAL CLASSIFICATION/CULTIVAR
DESIGNATION**

Nemesia caerula cultivar ‘Nemli’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Nemesia* plant, botanically known as *Nemesia caerula*
and referred to by the name ‘Nemli’.

The new *Nemesia* is a product of a planned breeding
program conducted by the Inventor in Auckland, New
Zealand. The objective of the program is to create new
compact *Nemesia* cultivars with numerous flowers and
unique flower colors.

The new *Nemesia* originated from a cross by the Inventor
of the *Nemesia caerula* cultivar ‘Pearl’, not patented, as the
female, or seed, parent with the *Nemesia caerula* cultivar
‘Delphi’, not patented, as the male, or pollen, parent on or
about Oct. 30, 1998. The cultivar ‘Nemli’ was discovered
and selected by the Inventor as a flowering plant within the
progeny of the stated cross in a controlled environment in
Auckland, New Zealand on or about Apr. 16, 1999.

Asexual reproduction of the new *Nemesia* by terminal
cuttings taken in a controlled environment in Auckland,
New Zealand since April, 1999, has shown that the unique
features of this new *Nemesia* are stable and are reproduced
true to type in successive generations.

SUMMARY OF THE INVENTION

The new *Nemesia* has not been observed under all pos-
sible environmental conditions. The phenotype may vary
somewhat with variations in environment such as
temperature, daylength and light intensity, without,
however, any variance in genotype.

The following characteristics have been repeatedly
observed and are determined to be basic characteristics of
‘Nemli’ and distinguish ‘Nemli’ as a new and distinct
cultivar:

1. Upright, outwardly spreading and compact plant habit.
2. Freely branching habit.

2

3. Small green leaves.

4. Numerous light purple-colored flowers.

Plants of the new *Nemesia* differ primarily from plants of
the female parent, the cultivar ‘Pearl’, in the following
characteristics:

1. Plants of the the new *Nemesia* are more compact than
plants of the cultivar ‘Pearl’.

2. Plants of the new *Nemesia* are more freely flowering
than plants of the cultivar ‘Pearl’.

3. Plants of the new *Nemesia* have light purple-colored
flowers whereas plants of the cultivar ‘Pearl’ have pale
pink-colored flowers.

Plants of the new *Nemesia* differ primarily from plants of
the male parent, the cultivar ‘Delphi’, in flower color.

Plants of the new *Nemesia* can be compared to plants of
the cultivar ‘Blueberry Sachet’, not patented. In side-by-side
comparisons conducted in Encinitas, Calif., plants of the
new *Nemesia* differed from plants of the cultivar ‘Blueberry
Sachet’ in the following characteristics:

1. Plants of the new *Nemesia* were larger and more
outwardly spreading than plants of the cultivar ‘Blue-
berry Sachet’.

2. Plants of the new *Nemesia* had lighter green foliage
than plants of the cultivar ‘Blueberry Sachet’.

3. Plants of the new *Nemesia* had light purple-colored
flowers whereas plants of the cultivar ‘Blueberry
Sachet’ had blue-colored flowers.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the
overall appearance of the new *Nemesia*, showing the colors
as true as it is reasonably possible to obtain in colored
reproductions of this type. Colors in the photographs may
differ slightly from the color values cited in the detailed
botanical description which accurately describe the colors of
the new *Nemesia*.

The photograph at the top of the sheet comprises a side
perspective view of a typical potted plant of ‘Nemli’ that was
about 14 weeks old.

The photograph at the bottom of the sheet comprises a close-up view of typical leaves and flowers of 'Nemli'.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observations and averaged measurements describe plants grown in Encinitas, Calif., in a polyethylene-covered greenhouse during the spring with day temperatures about 24° C., night temperatures about 19° C., and light levels about 4,000 foot-candles. Plants were grown for 14 weeks in 16.5-cm containers with one plant per container and were pinched. Color references are made to The Royal Horticultural Society Colour Chart except where general terms of ordinary dictionary significance are used.

Parentage:

Female parent.—*Nemesia caerulea* cultivar 'Pearl', not patented.

Male parent.—*Nemesia caerulea* cultivar 'Delphi', not patented.

Propagation:

Type.—By vegetative cuttings.

Time to initiate roots.—About 14 days at 20° C.

Time to develop roots.—About 30 days at 20° C.

Root description.—Fibrous, fine.

Plant description:

General appearance.—Upright, outwardly spreading and compact plant habit. Freely branching, typically about 30 laterals with numerous secondary laterals developing after pinching. Numerous light purple-colored zygomorphic flowers. Moderately vigorous.

Plant height.—About 28 cm.

Plant diameter or spread.—About 40 cm.

Lateral branches.—Appearance: Slender and filamentous; square in cross-section. Length: About 26 cm. Diameter: About 3 mm. Internode length: About 2.5 to 6 cm. Strength: Moderately strong, wiry; with development, lateral branches tend to bend with weight of flowers. Texture: Glabrous. Color: 144A.

Foliage description.—Arrangement: Opposite, simple. Quantity: About 12 to 14 per lateral branch. Shape: Elliptic. Apex: Broadly acute. Base: Acute to rounded. Length: About 3 cm. Width: About 1.8 cm. Margin: Serrate. Texture: Glabrous. Venation pattern: Pinnate, arcuate. Frgrance: Pungent, grassy. Petiole length: About 5 mm. Petiole diameter: About 2 mm. Color: Young leaves, upper surface: 147A. Young leaves, lower surface: 147B. Fully expanded leaves, upper surface: 147B. Fully expanded leaves, lower surface 147C. Venation, upper surface: 147B. Venation, lower surface: 147C. Petiole: 147B.

Flowering description:

Arrangement/appearance.—Zygomorphic solitary flowers arranged on terminal racemes; flowering acropetally towards apex. Flowers bilabiate with nectar spur. Flowers face outward. Flowers last about four to five days on the plant. Flowers not persistent.

Natural flowering season.—Natural flowering season is spring to fall; flowering continuous during this period.

Quantity of flowers.—Freely flowering with about 15 flowers and flower buds per raceme at one time.

Fragrance.—Faint, spicy and sweet.

Inflorescence length.—About 11.5 cm.

Inflorescence diameter.—About 2 cm.

Flower length.—About 1 cm.

Flower width.—About 1.5 cm.

Flower depth, including nectar spur.—About 1 cm.

Nectar spur length.—About 7 mm.

Flower buds.—Shape: Ovoid with spur. Length: About 6 mm. Diameter: About 5 mm. Color: 85C to 85D.

Petals.—Arrangement/shape: Five petals total. Four upper petals are fused at base to form an upright lobed and arched banner lip; lower petal modified into a larger lip with nectar spur and convex oval protuberance which serves as pollinator nectar guide and landing platform. Apex: Rounded. Margin: Entire. Length: Upper lip petals: About 7 mm. Lower lip petal: About 9 mm. Width: Upper lip petals: Lateral two petals: About 5 mm Center two petals: About 4 mm. Lower lip petal: About 1.2 cm. Texture: Smooth, velvety. Color: When opening, upper surface: 87A. When opening, lower surface: 76B. Fully opened, upper surface: 88C; color fading to 84A towards the margin and 85A towards the center with subsequent development. Fully opened, lower surface: 76C to 76D. Nectar guide: 9B. Nectar spur: 155B.

Sepals.—Quantity: Five-parted, star-shaped calyx. Shape: Elliptic. Apex: Acute. Base: Fused. Margin: Entire. Length: About 2 mm. Diameter: About 1.5 mm. Texture, upper and lower surfaces: Pubescent. Color: Upper surface: 144A. Lower surface: 147B.

Peduncle.—Length: About 11.5 cm. Diameter: About 1.5 mm. Strength: Strong. Angle: Upright to 45° from the stem. Color: 144C.

Pedicle.—Length: About 8 mm. Diameter: Less than 1 mm. Strength: Slender, but hold flowers outward. Angle: About 45° from the stem. Color: 144C.

Androecium.—Stamen number: Four per flower. Anther shape: Oval. Anther size: Less than 1 mm. Anther color: 6A. Amount of pollen: Scarce. Pollen color: 6A.

Gynoecium.—Pistil number: One per flower. Stigma shape: Flattened. Pistil length: About 2 mm. Style length: About 1 mm. Style color: 144D. Stigma shape: Rounded. Stigma color: 144D. Ovary color: 144B.

Seed/fruit.—Seed and fruit production has not been observed on plants of the new *Nemesia*.

Disease/pest resistance: Plants of the new *Nemesia* have not been observed to be resistant to pathogens or pests common to *Nemesias*.

Temperature tolerance: Plants of the new *Nemesia* have been observed to be tolerant to temperatures ranging from 4° to 32° C.

It is claimed:

1. A new and distinct cultivar of *Nemesia* plant named 'Nemli', as illustrated and described.

* * * * *

