



US00PP13385P2

(12) **United States Plant Patent**
Sekiguchi

(10) **Patent No.:** **US PP13,385 P2**

(45) **Date of Patent:** **Dec. 17, 2002**

(54) **OSTEOSPERMUM PLANT NAMED**
'SEIDACRE'

(76) **Inventor:** **Masayuki Sekiguchi**, 497
Kita-shiroda-cho, Maebashi-shi,
Gunma-ken (JP)

(*) **Notice:** Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) **Appl. No.:** **09/996,362**

(22) **Filed:** **Nov. 30, 2001**

(51) **Int. Cl.⁷** **A01H 5/00**

(52) **U.S. Cl.** **Plt./360**

(58) **Field of Search** **Plt./360**

Primary Examiner—Bruce R. Campell

Assistant Examiner—A. Para

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A distinct cultivar of *Osteospermum* plant named 'Seidacre',
characterized by its uniformly mounded and outwardly
spreading plant habit; freely branching growth habit; freely
flowering habit; and pale yellow-colored ray florets and dark
purple-tipped disc florets.

1 Drawing Sheet

1

**BOTANICAL CLASSIFICATION/CULTIVAR
DESIGNATION**

Osteospermum ecklonis cultivar Seidacre.

BACKGROUND OF THE INVENTION

The present Invention relates to a new and distinct culti-
var of *Osteospermum* plant, botanically known as
Osteospermum ecklonis, and hereinafter referred to by the
name 'Seidacre'.

The new *Osteospermum* is a product of a planned breed-
ing program conducted by the Inventor in Gunma-den,
Japan. The objective of the breeding program is to create
new *Osteospermum* cultivars with interesting floret colors.

The new *Osteospermum* originated from a cross made by
the Inventor in April, 1986 of two unidentified selections of
Osteospermum ecklonis, not patented. The new *Osteosper-*
mum was discovered and selected by the Inventor as a single
flowering plant within the progeny of the stated cross grown
in a controlled environment in Gunma-den, Japan.

Asexual reproduction of the new *Osteospermum* by ter-
minal vegetative cuttings was first conducted in Gunma-den,
Japan on Jun. 1, 1991. Asexual reproduction by cuttings has
shown that the unique features of this new *Osteospermum*
are stable and reproduced true to type in successive genera-
tions.

SUMMARY OF THE INVENTION

The cultivar Seidacre has not been observed under all
possible environmental conditions. The phenotype may vary
somewhat with variations in environment such as
temperature, daylength and light intensity without, however,
any variance in genotype.

The following traits have been repeatedly observed and
are determined to be the unique characteristics of 'Seidacre'.
These characteristics in combination distinguish 'Seidacre'
as a new and distinct *Osteospermum*:

1. Compact, mounded and outwardly spreading plant habit.
2. Freely branching growth habit.
3. Freely flowering habit.

2

4. Pale yellow-colored ray florets and dark purple-tipped
disc florets.

Plants of the new *Osteospermum* differ from plants of the
female parent, an unidentified selection, in the following
characteristics:

1. Plants of the new *Osteospermum* are shorter than plants
of the female parent.
2. Plants of the new *Osteospermum* flower earlier than
plants of the female parent.
3. Ray florets of plants of the new *Osteospermum* are pale
yellow in color whereas ray florets of plants of the female
parent are white in color.

Plants of the new *Osteospermum* differ from plants of the
male parent, the unidentified *Osteospermum* selection, pri-
marily in ray floret coloration.

Plants of the new *Osteospermum* are similar in ray floret
coloration to plants of the cultivar Cape Daisy Maputo, not
patented. However, plants of the new *Osteospermum* are
more outwardly spreading and more freely flowering than
plants of the cultivar Cape Daisy Maputo.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the
overall appearance of the new *Osteospermum* showing the
colors as true as it is reasonably possible to obtain in colored
reproductions of this type. Colors in the photographs may
differ from the color values cited in the detailed botanical
description which accurately describe the colors of the new
Osteospermum.

The photograph at the top of the sheet comprises a side
perspective view of three typical flowering plants of 'Sei-
dacre' grown in a 22-cm container.

The photograph at the bottom of the sheet is a close-up
view of typical developing inflorescence buds, typical devel-
oping inflorescences, upper and lower surfaces of typical
fully opened inflorescences, and upper and lower surfaces of
typical leaves of 'Seidacre'.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observa-
tions and measurements describe plants grown in Bonsall,
Calif., in an outdoor nursery during the late spring and early

summer under full sun conditions with day temperatures ranging from 18 to 35° C. and night temperatures ranging from 4 to 18° C. After planting rooted cuttings, plants were grown for about eight weeks in 22-cm containers with three plants per container and were pinched one time. Color references are made to The Royal Horticultural Society Colour Chart, 1995 edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Osteospermum ecklonis* cultivar Seidacre.

Parentage:

Female, or seed, parent.—Unidentified *Osteospermum ecklonis* selection, not patented.

Male, or pollen, parent.—Unidentified *Osteospermum ecklonis* selection, not patented.

Propagation:

Type.—Terminal vegetative cuttings.

Time to initiate rooting.—Summer: About 12 days at 25° C. Winter: About 20 days at 15° C.

Time to develop roots.—Summer: About 20 days at 25° C. Winter: About 27 days at 15° C.

Root description.—Fine, fibrous and well-branched.

Plant description:

Appearance.—Perennial herbaceous container and garden plant. Compact and mounded plant habit; initially upright and then outwardly spreading. Freely branching, about nine lateral branches develop after pinching; dense and full plants. Moderately vigorous growth habit.

Plant height.—About 22 cm.

Plant width or area of spread.—All three plants, about 25 cm; individual plants, about 15 cm.

Lateral branches.—Length: About 15 cm. Diameter: About 3 mm. Internode length: About 1.25 cm. Aspect: Upright to outward. Strength: Strong. Texture: Pubescent. Color: 143C.

Foliage description.—Arrangement: Alternate; simple. Number of leaves per lateral branch: About 15. Length: About 4.25 cm. Width: About 1.1 cm. Shape: Elliptic, strap-like. Apex: Acute. Base: Attenuate. Margin: Entire with occasional tiny serrations. Venation pattern: Pinnate. Texture: Smooth, glabrous. Color: Young foliage, upper surface: 143A. Young foliage, lower surface: 143C. Fully expanded foliage, upper surface: 147B. Fully expanded foliage, lower surface: 147C. Venation, upper and lower surfaces: 147C. Petiole: Length: About 1.1 cm. Diameter: About 5 mm. Texture, upper and lower surfaces: Glabrous. Color, upper and lower surfaces: 144B.

Inflorescence description:

Appearance.—Terminal and axillary inflorescences held above and beyond the foliage on moderately strong peduncles. Composite inflorescence form, radially symmetrical, with ligulate-shaped ray florets and disc florets massed at the center; ray and disc florets arranged acropetally on a capitulum. Inflorescences persistent. Inflorescences face upright or outward.

Flowering response.—Plants flower continuous and freely from the spring through the fall.

Postproduction longevity.—Inflorescences maintain good color and substance for about one week on the plant when grown in an outdoor environment.

Quantity of inflorescences.—Freely flowering; at one time, about nine open inflorescences and inflorescence buds per lateral stem.

Fragrance.—None detected.

Inflorescence bud (at stage of showing color).—

Length: About 1.5 cm. Diameter: About 7 mm.

Shape: Ovoid. Color, ray florets, lower or outer surface: 2A.

Inflorescence size.—Diameter: About 5.25 cm. Depth (height): About 2 cm. Disc diameter: About 1 cm.

Ray florets.—Length: About 3.5 cm. Width: About 7 mm. Shape: Ligulate. Apex: Slightly emarginate. Base: Acute. Margin: Entire. Texture: Velvety. Orientation: Initially upright then about 45 to 60° from vertical. Number of ray florets per inflorescence: About 16 in a single whorl. Color: When opening, upper surface: 10C. When opening, lower surface: 10A. Fully opened, upper surface: Main color, pale yellow, 10A to 10B; towards base, a white, 155D, band; below white band, a band of purple, 78A; below purple band at very base, 90A; main color fades to 10B with subsequent development. Fully opened, lower surface: Ground color, 12B, overlain with fine longitudinal stripes, 199B to 199C.

Disc florets.—Shape: Tubular, elongated. Apex: Five-pointed. Length: About 8 mm. Width: At apex: About 2 mm. At base: About 1 mm. Number of disc florets per inflorescence: About 45. Color: Immature: 10C; apex, 86A. Mature: 1D; apex, 83A.

Phyllaries.—Length: About 1.5 cm. Diameter: About 1.5 mm. Shape: Linear. Apex: Narrowly acute. Base: Fused. Margin: Entire. Texture: Coarse, pubescent. Number per inflorescence: About 16 in a single whorl. Color: Upper surface: 146A. Lower surface: 146C.

Peduncles.—Length, terminal peduncle: About 4.5 cm. Length, second peduncle: About 4 cm. Length, third peduncle: About 3 cm. Diameter: About 2 mm. Angle: Terminal peduncles, erect; secondary and tertiary peduncles, about 45° from vertical. Strength: Strong. Texture: Pubescent. Color: 146B.

Reproductive organs.—Androecium: Present on disc florets only. Stamen number: Five per floret; fused around style. Anther shape: Oblong. Anther length: About 1 mm. Anther color: 83A. Pollen amount: Moderate. Pollen color: 23A. Gynoecium: Present on both ray and disc florets. Pistil number: One per floret. Pistil length: About 7 mm. Stigma shape: Two-parted. Stigma color: 83A. Style length: About 5 mm. Style color: 84C. Ovary color: 145A.

Seed/fruit.—Seed and/or fruit production has not been observed.

Disease/pest resistance. Resistance to pathogens and pests common to *Osteospermums* has not been observed on plants grown under commercial greenhouse or outdoor conditions.

Temperature tolerance. Plants of the new *Osteospermum* have been observed to tolerate temperatures from 4 to 35° C.

It is claimed:

1. A new and distinct cultivar of *Osteospermum* plant named 'Seidacre', as illustrated and described.

* * * * *

