



US00PP13410P3

(12) **United States Plant Patent**
Strope

(10) **Patent No.:** **US PP13,410 P3**

(45) **Date of Patent:** **Dec. 24, 2002**

(54) **NEW GUINEA IMPATIENS PLANT NAMED**
'BALCELBUPI'

(75) **Inventor:** **Kerry M. Strope**, Pismo Beach, CA
(US)

(73) **Assignee:** **Ball FloraPlant, a division of Ball**
Horticultural Company, West
Chicago, IL (US)

(*) **Notice:** Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) **Appl. No.:** **09/772,807**

(22) **Filed:** **Jan. 30, 2001**

(65) **Prior Publication Data**

US 2002/0133859 P1 Sep. 19, 2002

(51) **Int. Cl.⁷** **A01H 5/00**

(52) **U.S. Cl.** **Plt./318**

(58) **Field of Search** **Plt./318**

Primary Examiner—Bruce R. Campell

Assistant Examiner—June Hwu

(74) *Attorney, Agent, or Firm*—Wood, Phillips, Katz, Clark
& Mortimer

(57) **ABSTRACT**

A new and distinct New Guinea impatiens cultivar named
Balcelbupi, characterized by its large blush pink flowers,
dark green foliage, compact upright and mounded habit and
excellent basal branching.

1 Drawing Sheet

1

BACKGROUND OF INVENTION

The present invention comprises a new and distinct New
Guinea impatiens plant, botanically known as *Impatiens*
hawkeri, and hereinafter referred to by the cultivar name
'Balcelbupi'.

The new cultivar was developed by the inventor in a
controlled breeding program during 1997 at Arroyo Grande,
Calif. The objective of the breeding program was to develop
new cultivars with numerous, large flowers, excellent basal
branching and upright medium growth habit.

The female (seed) parent of 'Balcelbupi' was 'Moorea'
(U.S. Plant Pat. No. 9,147) characterized by its compact
growth habit, pure white flowers and medium green foliage.
The male (pollen) parent of 'Balcelbupi' was the proprietary
Impatiens hawkeri cultivar designated '50-4' (non-patented)
characterized by its compact habit, pink flowers and dark
green foliage. 'Balcelbupi' was discovered and selected as a
flowering plant within the progeny of the stated cross in
December of 1997.

Asexual reproduction of the new cultivar by terminal
cuttings has demonstrated that the characteristics of the new
cultivar as herein described are firmly fixed and retained
through successive generations of such asexual propagation.
The new cultivar was asexually reproduced in Arroyo
Grande, Calif. and West Chicago, Ill.

SUMMARY OF INVENTION

It was found that the cultivar of the present invention:

- (a) Exhibits large round blush pink flowers,
- (b) Forms dark green foliage,
- (c) Exhibits a good basal branching character, and
- (d) Exhibits a medium, upright mounded growth habit.

The new cultivar of the present invention can be compared
to both its female parent 'Moorea' and Celebration
Blush White (U.S. Plant Pat. No. 8,538). When compared to
'Moorea' it is found that the new cultivar has larger flowers,
smaller leaves, longer internodes and shorter spur. When
compared to Celebration Blush White, the new cultivar is

2

more compact, has shorter internodes, smaller leaves and
larger flowers of a darker pink color. A detailed comparison
can be found in Table A, below.

Distinctive Characteristics:

Characteristic	'Goldenglossy'	'Sherwoodii'	'Conti'	'Sunrise'
Height (Mature)	3-4'	3-4'	1 1/2'-2'	3-4'
Width (Mature)	4-5'	4-5'	4-5'	4-5'
Growth Habit	Compact	Compact	Ground cover	Compact
Leaf Length	3/4-1 1/4"	3/4-1 1/4"	3/4-1"	3/4-1"
Leaf Width	3/8-5/8"	3/8-5/8"	3/8-5/8"	5/16-1/2"
Leaf Variegation (Immature)	Yellow Group 10B	None	Greyed-Yellow	Greyed-Yellow
Leaf Variegation (Mature)	Yellow-Green G.144A	None	Greyed-Yellow G.160C	Greyed-Yellow G.160A

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying photographs show as nearly true as it
is reasonably possible to make the same in color illustrations
of this type, typical flower and foliage characteristics of the
new cultivar. The plants were grown in a greenhouse at West
Chicago, Ill., U.S.A.

DETAILED DESCRIPTION

The 'Balcelbupi' cultivar has not been observed under all
possible environmental conditions to date. Accordingly, it is
possible that the phenotype may vary somewhat with variations
in the environment, such as temperature, light intensity,
and day length.

The chart used in the identification of colors described
herein is The R.H.S. Colour Chart of The Royal Horticultural
Society, London, England. The color values were determined
on Aug. 30, 2000 between 1:00 and 1:45 p.m. under natural
daylight conditions. The following measur-

ments and describe plants ten (10) weeks old and produced from cuttings taken from stock plants and grown in 10 cm pots under greenhouse conditions comparable to those used in commercial practice while utilizing a soilless growth medium and maintaining temperatures of approximately 72° F. during the day and approximately 65° F. during the night.

Classification:

Botanical.—*Impatiens hawkeri* cultivar ‘Balcelbupi’.

Parentage: ‘Moorea’ (seed)×Proprietary seedling ‘50-4’ (pollen).

Propagation:

Type cutting.—Terminal tip.

Time to initiate roots.—Approximately 7–14 days with the shorter times generally being experienced in the summer and the longer times in the winter.

Time to develop roots.—Approximately 21 days.

Rooting habit.—Fibrous, branching.

Plant description:

Habit of growth.—Moderately vigorous with good basal branching. A mature plant, 8 to 10 weeks after the planting of a rooted cutting, commonly measures approximately 18.4 cm in height and approximately 29 cm in width.

Plant form.—Upright and mounded.

Foliage.—Shape: Lanceolate; acuminate apex; attenuate base; ciliate and serrate margin. Texture: Smooth, slightly pubescent on upper surface only. Size of mature foliage: Approximately 6.8 cm in length; approximately 1.9 cm in width. Color of mature foliage: Upper surface is 139A, with mid vein lighter than 53D; Lower surface is 137C with veins of 53C. Petiole: Length: approximately 6 mm, Color: 53D.

Stem.—Color 178A. Internode length: Approximately 5.2 cm.

Flower description:

Flowering habit.—Freely flowering.

Natural flowering season.—Year round in greenhouse environment. Flowering is continuous from spring until fall in the garden.

Bud shape, color.—Bud Shape: Ovate. Color: 55C.

Flowers borne.—Above foliage arising from leaf axils.

Quantity of flowers.—Approximately 1 flower and 4 buds per stem.

Flower size.—Approximately 7.1 cm in diameter.

Flower buds.—Length: approximately 2.1 cm. Width: approximately 1.3 cm.

Petals.—Number: Five, overlapping. Size: Superior petal length — 3.3 cm, superior petal width — 5.2 cm. Lateral petal length — 3.7 cm, lateral petal width — 3.4 cm. Lower petal length — 4.2 cm, lower petal width — 4.3 cm. Shape: obovate with emarginate apex attenuate base and entire margin. Texture: smooth and iridescent.

Flower color.—Light Pink. Upper surface of petals has a background of 73D with fine venation of 73B overall and bases of 73A forming an “eye”; lower surface has background of 73D, fine venation overall of 73B, midveins and bases of 61B.

Spur.—Length: Approximately 4.7 cm. Shape: Sharply curved. Color: 60B.

Peduncles.—Length approximately 4.3 cm. Color: 178B.

Reproductive organs.—Androecium: Five stamens. Anther size: Approximately 2.5 mm. Color: 39D. Anthers are fused together forming one organ that surrounds the pistil. Generally, the anthers shed pollen prior to the stigma becoming receptive. Pollen color: 19D. Gynoecium: Pistil number: One. Stigma length: >1 mm. Stigma color: 154D. Style length: 1 mm. Style color: 144A. Ovary length: 4 mm. Ovary color: 144A.

Seed development.—Seed production has not been observed.

Disease resistance.—Resistance to pathogens has not been observed.

I claim:

1. A new and distinct cultivar of New Guinea impatiens plant named ‘Balcelbupi’ substantially as herein shown and described, which:

- (a) Exhibits large round blush pink flowers,
- (b) Forms dark green foliage,
- (c) Exhibits a good basal branching character, and
- (d) Exhibits an upright mounded growth habit.

* * * * *

