



US00PP13446P2

(12) **United States Plant Patent**
Tristram

(10) **Patent No.: US PP13,446 P2**
(45) **Date of Patent: Dec. 31, 2002**

(54) **SCABIOSA PLANT NAMED ‘WALMINIBLUE’**
(76) **Inventor: David Ralph Tristram**, Old Rectory,
Arundel, Binsted, W. Sussex (GB),
BN18 0LL
(*) **Notice:** Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.
(21) **Appl. No.: 09/962,382**
(22) **Filed: Sep. 25, 2001**
(51) **Int. Cl.⁷ A01H 5/00**

(52) **U.S. Cl. Plt./263**
(58) **Field of Search Plt./263**
Primary Examiner—Bruce R. Campell
Assistant Examiner—June Hwu
(74) *Attorney, Agent, or Firm*—Mark P. Bourgeois

(57) **ABSTRACT**
A new cultivar of *Scabiosa columbaria* named ‘Walm-
inibblue’ that is characterized by its small plant size, compact
plant habit, proportionately sized, intensely colored lavender
blue flowers, prolific and continuous flowering, and sterility.

2 Drawing Sheets

1

**CROSS-REFERENCE TO RELATED
APPLICATIONS**

The present application is co-pending with another appli-
cation entitled *Scabiosa Plant Named ‘Walminipink’* (appli-
cation Ser. No. 09/962,381).

LATIN NAME OF THE GENUS AND SPECIES

Scabiosa columbaria.

VARIETY DENOMINATON

‘Walminibblue’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Scabiosa* plant, botanically known as *Scabiosa colum-
baria* and will be referred to hereafter by its cultivar name
‘Walminibblue’. *Scabiosa columbaria* is a hardy perennial
grown for landscape use.

The new cultivar, ‘Walminibblue’, was discovered in 1988
by the inventor at his nursery in West Sussex, England.
‘Walminibblue’ was discovered as a naturally occurring
mutation, amongst a large crop of *Scabiosa columbaria*
‘Butterfly Blue’ (un-patented). The new invention was dis-
covered as a whole plant, however it originated as a single
cutting taken from ‘Butterfly Blue’. Several selections were
discovered and observed before ‘Walminibblue’ was selected
as unique and desirable for its characteristics of plant size,
plant habit, flower color, and sterility.

The new variety of *Scabiosa* can be characterized by its
smaller size: the foliage dimensions, diameter of the flower
heads and the height of the plant are less in comparison to
the parent plant, ‘Butterfly Blue’. The new invention has
never set fruit and all flowers observed have been self-
sterile. Near self-sterility is a characteristic of ‘Butterfly
Blue’ and the inventor’s *Scabiosa* variety ‘Pink Mist’ (U.S.
Plant Pat. No. 8,957), however, it is not a characteristic of
any other commercial cultivars of *Scabiosa columbaria*
known to the inventor. The flower color of ‘Walminibblue’ is
deeper and more blue in comparison to ‘Butterfly Blue’. The
new cultivar is similar to both ‘Butterfly Blue’ and ‘Pink
Mist’ in that it has a compact habit and has a continuous and
prolific flowering habit.

2

Asexual reproduction of the new cultivar was first accom-
plished by taking cuttings in West Sussex, England by the
inventor in 1988. The characteristics of this cultivar have
been determined to be stable and are reproduced true to type
in successive generations both by vegetative cuttings and by
tissue culture.

SUMMARY OF THE INVENTION

The following traits have been repeatedly observed and
are determined to be the basic characteristics of the new
cultivar. These attributes in combination distinguish the new
invention from wild forms of *Scabiosa columbaria* and
distinguish it as a new and distinct cultivar. ‘Walminibblue’
has not been tested under all possible conditions and phe-
notypic differences may be observed with variations in
environmental, climatic, and cultural conditions:

1. ‘Walminibblue’ is smaller in height and has less of a
spread in comparison to ‘Butterfly Blue’. ‘Walminibblue’
attains 10 to 18 cm in height and 10 to 20 cm in width while
‘Butterfly Blue’ attains 15 to 30 cm in height and 17 to 30
cm in width, dependent on growing conditions.

2. The plant habit of ‘Walminibblue’ is compact with a tight
rosette of basal growth.

3. The flowers of ‘Walminibblue’ are smaller than those of
Butterfly Blue, ranging from 2.5 to 4.3 cm in diameter while
the flower size of ‘Butterfly Blue’ ranges from 4.0 to 6.0 cm
in diameter. This range in flower size is found on a single
plant.

4. The flowers of ‘Walminibblue’ are lavender blue tending
towards pure blue and are deeper in color than ‘Butterfly
Blue’. The flowers of ‘Walminibblue’ are typically 92A
whereas those of ‘Butterfly Blue’ are typically 91A–91B.

5. ‘Walminibblue’ blooms continuously and profusely from
early spring until frost. Typically, there are 10–12 flower
heads showing color at one time.

6. The flowers of ‘Walminibblue’ are presumed to be
sterile, as fruit set has not been observed to date.

BRIEF DESCRIPTION OF THE DRAWING

The photograph on Sheet 1 is of a 6 month-old plant of
‘Walminibblue’ as grown in a one-gallon container. The plant
was grown in a greenhouse in Encinitas, Calif. under natural

lighting and with an average daytime temperature of 75° F. and an average nighttime temperature of 70° F.

The photograph on Sheet 2 is a close-up of recently opened flowers. The colors in these photographs are as accurate as could be obtained by conventional photography. The flower color is pinker in the photographs due to the difficulty in accurately capturing blue tones with photographic film.

BOTANICAL DESCRIPTION OF THE PLANT

The following is a detailed botanical description of the new cultivar as grown under greenhouse conditions in Encinitas, Calif. The 6-month old plant was grown under natural lighting with an average daytime temperature of 75° F. and an average nighttime temperature of 70° C. and a constant feed of 20-10-20 fertilizer at a rate of 175 ppm Nitrogen. General plant descriptions describe a plant as observed in the garden in West Sussex, England for 6 years. The color determination is in accordance with The R.H.S. Colour Chart of The Royal Horticultural Society, London, England, except where general color terms of ordinary dictionary significance are used.

Botanical classification: *Scabiosa columbaria* 'Walminibblue'.

Market classification: Hardy perennial for landscape use.

Parentage: Naturally occurring mutation of *Scabiosa columbaria* 'Butterfly Blue' (not patented).

Plant description:

Blooming period.—Natural flowering season is from early spring until frost in the Northern hemisphere.

Plant habit.—Dense, mounded rosette with upright flowering stems.

Height and spread.—10 to 18 cm in height in bloom (bloom stalks vary in height) and 10 to 20 cm in spread depending on growing conditions.

Hardiness.—Zone 3–9.

Type.—Hardy herbaceous perennial.

Propagation.—By terminal tip cuttings and division.

Root system.—Fibrous, branching.

Time to initiate roots.—10 to 15 days in the summer months at temperatures of 64° F. to 72° F. and 15 to 20 days in the winter months at temperatures of 53° F. to 60° F.

Vigor.—A rooted cutting will establish roots and initiate flowers in a six-inch container after 15 weeks at an average temperature of 75° F. and constant fertilizer feed of 20-10-20 at a rate of 175 ppm Nitrogen.

Culture.—Well drained soils of average fertility in full sun.

Diseases and pests.—Powdery mildew is the only known disease known to be a problem in the garden.

Foliage description (characteristics that differ in basal and flowering stem leaves): Basal leaves: Simple, opposite arrangement, obovate to oblanceolate, entire to coarsely lobed with lobes and apex rounded, 5.0 to 10 cm in length and 1.5 to 2.5 cm in width.

Flowering stem leaves: Simple, obovate, pinnatifid to pinnatisect with up to seven pinnae on each side, leaflets are opposite or alternately arranged, lobes are ovate to lanceolate, apices of leaf and leaflets are acute. Stem leaves are 5.0 to 12.0 cm in length and 2.0 to 3.0 in width.

Foliage description (characteristics that are similar for the basal and flowering stem leaves):

Venation.—Not prominent, with the exception of the mid-rib on the lower portion of the basal leaves, color 139D (upper and lower surface).

Surface.—Pubescent on upper and lower surfaces.

Color.—Upper surface, mature and immature leaves, 137A. Lower surface, mature and immature leaves, 137C.

Flowering stem description (peduncle):

Shape.—Round.

Size.—1.5–3.0 mm in diameter in width, 10 to 18 cm in height with the longer stems occurring when temperatures are warmer, above 75° F.

Surface.—Pubescent.

Color.—138B.

Branching.—Initially none, but as the flower stem elongates, lateral branches are produced, each with a single flower head.

Flower description:

Type.—Terminal flat heads, regular to slightly irregular, round, numerous with outer florets that are larger than the inner florets.

Fragrance.—None.

Size.—2.5 to 4.3 cm in diameter cm in diameter, approximately 1.0 cm in depth.

Lastingness of the flowers.—Approximately one week, not self cleaning; bristled calyx and involucre bracts are persistent.

Buds.—Dome-shaped, emerge 143D and 5.0 mm in diameter, gradually changing to the petal color 92A, with a diameter of 1.0 cm.

Involucre bracts.—10–15 in a single row, held horizontally, linear-lanceolate, apex acute, length ranges from 5.0 mm to 12.0 mm, width ranges from 1.5 mm to 3.0 mm, base of bract is 145A, portion of bract extended beyond flower head is 137A, pubescent.

Calyx.—Densely packed to form a receptacle-like structure. Cupular, composed of 5 united sepals, 144 C, 1.5 mm in width and 2.0 mm in height with a bristle, 202A, extending 2.0 mm in length beyond each sepal. Entire Calyx is persistent.

Corolla.—Zygomorphic, five petals, partially united near base. Outer florets are 8 to 15 mm in length and 4 to 6 mm in width, approximately 15–30 ray florets per inflorescence, color 92A that fades to 91A with maturity on both surfaces. Inner florets are 4–7 mm in length and 2–3 mm in width, approximately 50–120 disc florets per inflorescence, color 92A on both surfaces, fading to 91A and then fading to 159D at maturity.

Reproductive organs.—Stamens; 4, partially fused, can be included or extend up to 4 mm above corolla in early stages of flowering and then are not visible, 68D in color, no pollen was observed to date.

Pistil.—One, 159D, clearly visible in later stage of flowering, extending up to 6.0 mm beyond corolla, superior ovary.

Fruit/seed.—Fruit production has not been observed to date, presumed to be sterile.

I claim:

1. A new and distinct cultivar of *Scabiosa* plant named 'Walminibblue' as described and illustrated.

* * * * *



